

Ke Li's lemma for quantum hypothesis testing in general von Neumann algebras

Yan Pautrat and Simeng Wang

Université Paris-Saclay, CNRS, Laboratoire de Mathématiques d'Orsay
91405 Orsay, France

January 14, 2021

Abstract

A lemma stated by Ke Li in [9] has been used in e.g. [3, 8, 12, 14, 15] for various tasks in quantum hypothesis testing, data compression with quantum side information or quantum key distribution. This lemma was originally proven in finite dimension, with a direct extension to type I von Neumann algebras. Here we show that the use of modular theory allows to give more transparent meaning to the objects constructed by the lemma, and to prove it for general von Neumann algebras. This yields immediate generalizations of e.g. [3].

1 Introduction

Quantum hypothesis testing is concerned with the situation where one considers a von Neumann algebra \mathcal{M} , equipped with a state which is either ρ or σ ; this uncertainty in the nature of the state makes sense in particular when \mathcal{M} is viewed as modeling the observable quantities of a quantum system, and the physical state of this system — itself modeled by a state in the mathematical sense — is only known to be one or the other. A natural task is then to try and determine which state is the true one by producing, in physical terms, an experimental measurement procedure such that, depending on the measurement outcome, one will conclude that the actual state is ρ or σ .

In the model of orthodox quantum mechanics, a measurement procedure is determined by a self-adjoint element X of the observable algebra \mathcal{M} , and this X is simply called an *observable*. Non-trivial measurements will have a random outcome; the set of possible outcomes is exactly the spectrum of that element, and if ω is the actual state of the system and we denote by ξ_X the spectral measure of X , then the probability distribution for the measurement outcomes is $\rho \circ \xi_X$. In the situation described above it will suffice to consider an observable with spectrum $\{0, 1\}$, that is, an orthogonal projector of \mathcal{M} . We will take advantage of this simplification

and continue this preliminary discussion assuming that the observable describing the discriminating experience, is an orthogonal projector T , which we call a *test*.

Suppose the decision rule is that, if the measurement outcome is 0, then the observer concludes that the true state is ρ , and if the measurement outcome is 1, then they conclude that the true state is σ . There are two ways in which this conclusion can be wrong: either the actual state was ρ and the measurement outcome was 1, or the actual state was σ and the measurement outcome was 0. Applying the rules determining the distribution of the measurement outcomes shows that the former occurs with probability $\rho(T)$, and the latter with probability $\sigma(\text{Id} - T)$. Assuming that the possible states ρ and σ are fixed, we denote these two types of error

$$\alpha(T) := \rho(T) \quad \beta(T) := \sigma(\text{Id} - T). \quad (1)$$

If, to the experimenter's knowledge, both ρ and σ may be the state of the system, then one typically wishes to make both $\alpha(T)$ and $\beta(T)$ small. There are, however, various ways in which this can be done, and we postpone the corresponding discussion to section 4.

The result we are concerned with gives a test T such that $\alpha(T)$ and $\beta(T)$ satisfy a pair of upper bounds. To state it, let us assume that \mathcal{M} is a finite-dimensional matrix algebra: $\mathcal{M} = \mathcal{B}(\mathbb{C}^n)$, and denote by ϱ and ς the density matrices associated with the states ρ and σ , that is:

$$\rho = \text{tr}(\varrho \cdot) \quad \sigma = \text{tr}(\varsigma \cdot). \quad (2)$$

Assume for simplicity that ρ and σ are faithful states, or equivalently that ϱ, ς are invertible matrices. Consider then the vector space $\mathcal{B}(\mathbb{C}^n)$, on which we define the vectors

$$\Omega_\rho = \varrho^{1/2} \quad \Omega_\sigma = \varsigma^{1/2}$$

and the operator

$$\Delta_{\rho|\sigma} : X \mapsto \varrho X \varsigma^{-1}.$$

If $\mathcal{B}(\mathbb{C}^n)$ is equipped with the scalar product $\langle X, Y \rangle = \text{tr}(X^* Y)$ then $\Delta_{\rho|\sigma}$ is self-adjoint. It then holds that for any $\epsilon > 0$ there exists a test T such that

$$\alpha(T) \leq \epsilon \quad \beta(T) \leq \langle \Omega_\sigma, \mathbb{1}_{(\epsilon, +\infty)}(\Delta_{\rho|\sigma}) \Omega_\sigma \rangle. \quad (3)$$

This result was first proven in [9], as a technical step towards the ‘‘second order Stein’s lemma’’ which we discuss in section 4. It is generally quoted as ‘‘Ke Li’s lemma’’ for quantum hypothesis testing, even though it is not the only result of Ke Li relevant to this field. The statement as written above, however, does not appear in [9] and in particular, there is no mention of the operator $\Delta_{\rho|\sigma}$ in that article. The result was later reformulated in [3] (with the same goal of proving the second order Stein’s lemma) to involve the operator $\Delta_{\rho|\sigma}$, which the reader may already have recognized to be the finite-dimensional instance of a *relative modular operator*. From then on it was expected that a proof written solely in terms of modular theory on $\mathcal{B}(\mathbb{C}^n)$ should be possible, in which case, according to a standard rule of thumb discussed in [7] it

should extend with minimal effort to general von Neumann algebras. It turns out that such an extension holds for any two faithful normal states ρ, σ on a σ -finite von Neumann algebra (yielding an extension of the second order Stein's lemma as stated in [3], see section 4). To state this extension we need, however, to recall how the relative modular operator $\Delta_{\rho|\sigma}$ is defined in the general case, and we therefore postpone its statement to Section 3.

Even though our proof is written in the language of modular theory, it did not proceed from a simple "modular translation" of Ke Li's finite dimensional proof, as we could not find such a thing. Note that until now, no extension of inequalities (3) was found, except for the extension to separable type I von Neumann algebras in [11], in which case ρ and σ were still of the form (2) and one could simply proceed by finite-dimensional approximations of ϱ and ς .

Let us recall shortly the definition of T in the proof of (3) as in [3, 9] in order to underline its connection with modular theory, and motivate our definition of T below. Consider spectral decompositions of ϱ, ς :

$$\varrho = \sum_{x=1}^n \lambda_x |a_x\rangle\langle a_x| \quad \varsigma = \sum_{y=1}^n \mu_y |b_y\rangle\langle b_y|$$

where $(\lambda_x)_x$ and $(\mu_y)_y$ are labeled in nondecreasing order, and both families $(a_x)_x$ and $(b_y)_y$ are orthonormal bases of \mathbb{C}^n . Define for any $y = 1, \dots, n$:

$$Q_y = \mathbb{1}_{[0, \epsilon\mu_y]}(\varrho) = \sum_{x=1}^n \mathbb{1}_{\lambda_x \leq \epsilon\mu_y} |a_x\rangle\langle a_x|,$$

so that $(Q_y)_y$ is a nondecreasing family of projectors, and

$$\xi_y = Q_y b_y = \sum_{x=1}^n \mathbb{1}_{\lambda_x \leq \epsilon\mu_y} \langle a_x, b_y \rangle a_x.$$

Ke Li then defines his test as the orthogonal projector T_{KL} onto the vector space spanned by the $\xi_y, y = 1, \dots, n$.

For X an operator on \mathbb{C}^n denote now by L_X the operator on $\mathcal{B}(\mathbb{C}^n)$ acting by $Y \mapsto XY$, and let J be the antilinear involution $X \mapsto X^*$ on $\mathcal{B}(\mathbb{C}^n)$. Notice then that $(JL_X J)Y = YX^*$ and let $\mathcal{M}, \mathcal{M}'$ be the sets of operators of the form L_X or $JL_X J$ respectively. We then have:

$$\Delta_{\rho|\sigma} = \sum_{x,y=1}^n \lambda_x \mu_y^{-1} L_{|a_x\rangle\langle a_x|} J L_{|b_y\rangle\langle b_y|} J$$

so that

$$\mathbb{1}_{(0, \epsilon]}(\Delta_{\rho|\sigma}) = \sum_{x,y=1}^n \mathbb{1}_{\lambda_x \leq \epsilon\mu_y} L_{|a_x\rangle\langle a_x|} J L_{|b_y\rangle\langle b_y|} J = \sum_{y=1}^n L_{Q_y} J L_{|b_y\rangle\langle b_y|} J$$

and

$$\mathbb{1}_{(0,\epsilon]}(\Delta_{\rho|\sigma})\Omega_\sigma = \sum_{x,y=1}^n \mathbb{1}_{\lambda_x \leq \epsilon \mu_y} \mu_y^{1/2} \langle a_x, b_y \rangle |a_x\rangle\langle b_y|. \quad (4)$$

It is then immediate to remark that $\mathbb{1}_{(0,\epsilon]}(\Delta_{\rho|\sigma})\Omega_\sigma b_y = \mu_y \xi_y$ for any $y = 1, \dots, n$, so that

$$\text{Ran}(\mathbb{1}_{(0,\epsilon]}(\Delta_{\rho|\sigma})\Omega_\sigma) = \text{span}\{\xi_y, y = 1, \dots, n\}$$

where the last equation is due to the faithfulness of σ . Last, remark that the range $\text{Ran}(\mathbb{1}_{(0,\epsilon]}(\Delta_{\rho|\sigma})\Omega_\sigma)$ is the same as $\mathcal{M}'\mathbb{1}_{(0,\epsilon]}(\Delta_{\rho|\sigma})\Omega_\sigma$. Therefore, T_{KL} is equivalently defined by the fact that $L_{T_{\text{KL}}}$ is the orthogonal projection on $\mathcal{M}'\mathbb{1}_{(0,\epsilon]}(\Delta_{\rho|\sigma})\Omega_\sigma$. Since all quantities appearing in this last sentence are well-defined in the standard representation of any von Neumann algebra (see section 2), this gives a satisfactory starting point for our proof of the extension of Ke Li's lemma.

The structure of the paper is as follows. In section 2 we recall the elements of the modular theory of von Neumann algebras required by the statement of our result. In section 3 we give our result and its proof. In section 4 we discuss the merits of our result and its possible applications.

Acknowledgements YP wishes to thank Nilanjana Datta and Cambyse Rouzé for initiating the pleasant collaboration that led to [3] and the present endeavors, Ke Li and Magdalena Musat for instructive discussions and most of all Yoshiko Ogata who helped chase the modular in Ke Li's original proof. YP was supported by ANR grant "NonStops" ANR-17-CE40-0006. SW was partially supported by a public grant as part of the FMJH and ANR grant "ANCG" ANR-19-CE40-0002.

2 Modular theory and Haagerup L^p spaces

We recall first a few notions on von Neumann algebras and relative modular operators. Suggested references are [2], and for a pedagogical introduction we recommend section 2 (and in particular sections 2.11 and 2.12) of [7]. We then move on to describe a few properties of Haagerup's L^p spaces following [5] and [13].

Let \mathcal{M} be a von Neumann algebra, i.e. a $*$ -algebra of bounded linear operators on some Hilbert space, that is closed in the weak operator topology and contains the identity. The set of operators commuting with all elements of \mathcal{M} is called the commutant of \mathcal{M} , and denoted \mathcal{M}' ; von Neumann's bicommutant theorem then states that a $*$ -algebra of bounded linear operators containing the identity is a von Neumann algebra if and only if $\mathcal{M} = \mathcal{M}''$, the latter being the bicommutant $(\mathcal{M}')'$ of \mathcal{M} .

The notion of nonnegative operators induces a partial order on \mathcal{M} : by definition, $X \leq Y$ if $Y - X$ is a nonnegative operator. A map on \mathcal{M} such that the null operator is the only nonnegative operator mapped to the zero element is called *faithful*. Linear forms ω on \mathcal{M} with the additional regularity property that

$\omega(\limsup_n X_n) = \limsup_n \omega(X_n)$ if X_n is an increasing sequence of self-adjoint operators are called *normal*. A linear form which maps nonnegative operators to nonnegative scalars is called *positive*. A positive linear form ω from \mathcal{M} to $[0, +\infty]$ will be called a *weight*. A weight ω such that the set of nonnegative elements A with $\omega(A) < +\infty$ is weakly dense is called *semifinite*. A weight ω with the property that $\omega(\text{Id}) = 1$ is called a *state* (and necessarily takes values in $[0, +\infty)$). A weight or state with the property that $\omega(XY) = \omega(YX)$ for all X, Y in \mathcal{M} is called *tracial*, or a *trace*.

A standard representation of some von Neumann algebra \mathcal{M} is a quadruple $(\pi, \mathcal{H}, \mathcal{H}^+, \text{J})$ where \mathcal{H} is a Hilbert space (with scalar product denoted $\langle \cdot, \cdot \rangle$), π is a faithful morphism of $*$ -algebras from \mathcal{M} to $\mathcal{B}(\mathcal{H})$, \mathcal{H}^+ is a self-dual cone of \mathcal{H} (i.e. the set of ϕ in \mathcal{H} such that $\langle \phi, \psi \rangle \geq 0$ for all ψ in \mathcal{H}^+ is \mathcal{H}^+ itself), J is an anti-unitary involution of \mathcal{H} , and these different objects satisfy the following properties:

- $\text{J}\mathcal{M}\text{J} = \mathcal{M}'$,
- $\text{J}X\text{J} = X^*$ for X in $\mathcal{M} \cap \mathcal{M}'$,
- $\text{J}\psi = \psi$ for ψ in \mathcal{H}^+ ,
- $\text{J}X\text{J}X\mathcal{H}^+ \subset \mathcal{H}^+$ for $X \in \mathcal{H}$.

A standard representation of a von Neumann algebra \mathcal{M} always exists (we will describe one of them below), and if $(\pi_1, \mathcal{H}_1, \mathcal{H}_1^+, \text{J}_1)$ and $(\pi_2, \mathcal{H}_2, \mathcal{H}_2^+, \text{J}_2)$ are two standard representations, then there exists a unitary operator $U : \mathcal{H}_1 \rightarrow \mathcal{H}_2$ such that $U\pi_1(X)U^* = \pi_2(X)$ for all $X \in \mathcal{M}$, $U\text{J}_1U^* = \text{J}_2$ and $U\mathcal{H}_1^+ = \mathcal{H}_2^+$.

We will now describe the modular structure associated with a von Neumann algebra \mathcal{M} : fix a standard representation $(\pi, \mathcal{H}, \mathcal{H}_+, \text{J})$ of \mathcal{M} . Then for any normal state ω on \mathcal{M} there exists a unique Ω_ω in \mathcal{H}_+ such that

$$\omega(X) = \langle \Omega_\omega, \pi(X)\Omega_\omega \rangle \text{ for any } X \in \mathcal{M}. \quad (5)$$

This Ω_ω is cyclic in the sense that the closure of $\pi(\mathcal{M})\mathcal{H}$ is \mathcal{H} itself. If in addition ω is faithful, then Ω_ω is separating in $\pi(\mathcal{M})$, that is, $\pi(X)\Omega_\omega = 0$ if and only if $X = 0$. We continue by fixing two normal faithful states ρ and σ . We can then define a densely defined operator $S_{\rho|\sigma}$ by

$$S_{\rho|\sigma}X\Omega_\sigma = X^*\Omega_\rho.$$

This operator turns out to be closable, and its closure $\overline{S}_{\rho|\sigma}$ has polar decomposition

$$\overline{S}_{\rho|\sigma} = \text{J}\Delta_{\rho|\sigma}^{1/2}$$

where $\Delta_{\rho|\sigma}$ is a positive-definite operator on \mathcal{H} , which is in general unbounded.

We are now in a position to state our main result, but will first introduce additional elements required for our proof. The reader may wish to take a peek at Theorem 1 in section 3.

To prove the result we consider the Haagerup L^p -spaces associated with \mathcal{M} . We will not need the detailed construction of these spaces and only recall their properties

(see chapter 2 of [13] for a complete presentation; in the rest of this section all numbered references point to those notes, with e.g. Theorem 2.7 meaning Theorem 7 of chapter 2 of [13]). An important role will be played by a distinguished von Neumann algebra \mathcal{N} , equipped with a normal semifinite faithful trace τ . This algebra \mathcal{N} then acts on the Hilbert space $L^2(\mathbb{R}, \mathcal{H})$ which we simply denote \mathcal{K} , for its nature will not matter. For a definition of this algebra \mathcal{N} , see the beginning of chapter 2 of [13]: between saying too much and saying too little, it seems more convenient to say too little. This \mathcal{N} is such that there exists a faithful normal representation of \mathcal{M} as a sub-von Neumann algebra of \mathcal{N} ; in order to spare ourselves an additional notation for this representation, we simply assume that \mathcal{M} is realized as a subalgebra of \mathcal{N} , and therefore acts on $\mathcal{K} = L^2(\mathbb{R}, \mathcal{H})$. We say that a densely defined unbounded operator on \mathcal{K} is measurable with respect to (\mathcal{N}, τ) if for every $\delta > 0$ there exists a projection P in \mathcal{N} such that the range of P is included in the domain of \mathcal{N} , and $\tau(\text{Id} - P) < \delta$. We say that an unbounded operator on \mathcal{K} is affiliated with \mathcal{N} if any Borel bounded functional of either $\frac{1}{2}(K + K^*)$ or $\frac{1}{2i}(K - K^*)$ is an element of \mathcal{N} . Then for all $1 \leq p \leq \infty$, there exists a space $L^p(\mathcal{M})$ which is a subspace of the set of those measurable operators with respect to (\mathcal{N}, τ) which are affiliated with \mathcal{N} , which satisfies the following properties. For all $1 \leq p < \infty$, an operator X on \mathcal{K} is in $L^p(\mathcal{M})$ if and only its polar decomposition $X = U|X|$ satisfies $U \in \mathcal{M}$ and $|X|^p \in L^1(\mathcal{M})$. Moreover there is a faithful positive linear functional $\text{tr} : L^1(\mathcal{M}) \rightarrow \mathbb{C}$ such that

$$\text{tr}(XY) = \text{tr}(YX) \text{ for all } X \in L^p(\mathcal{M}), Y \in L^q(\mathcal{M}) \text{ with } 1/p + 1/q = 1.$$

If we define $\|X\|_p = \text{tr}(|X|^p)^{1/p}$ then the closed product of measurable operators on these spaces satisfies the Hölder inequality, and in particular we have $L^p(\mathcal{M})L^q(\mathcal{M}) \subset L^r(\mathcal{M})$ for any $p, q, r \in [1, +\infty]$ with $1/r = 1/p + 1/q$. This induces a natural Hilbert space structure on $L^2(\mathcal{M})$ by taking $p = q = 2$. In addition (Theorem 2.7 and Proposition 2.10), $\mathcal{M} = L^\infty(\mathcal{M})$ and the normal states ρ, σ on \mathcal{M} are of the form

$$\rho = \text{tr}(\varrho \cdot) \quad \sigma = \text{tr}(\varsigma \cdot) \quad (6)$$

where ϱ, ς are positive elements of $L^1(\mathcal{M})$ with $\text{tr}(\sigma) = 1$. This of course echoes (2) and will allow us to mimic some of the ‘‘spatial’’ arguments in Ke Li’s proof. Taking the values of p, q, r appropriately, we obtain a faithful representation of \mathcal{M} on $L^2(\mathcal{M})$ as

$$\pi(X)Y = XY, \quad X \in \mathcal{M}, Y \in L^2(\mathcal{M}).$$

Denote $J : X \mapsto X^*$ and $L^2(\mathcal{M})_+$ the subset of $L^2(\mathcal{M})$ made of nonnegative operators. Then $(\pi, L^2(\mathcal{M}), L^2(\mathcal{M})_+, J)$ is a standard form of \mathcal{M} (see Theorem 2.36). Last, we will use the crucial fact that $L^p(\mathcal{M})$ is isometrically isomorphic to a subspace of the tracial weak L^p -space on (\mathcal{R}, τ) , and more precisely we have (Lemma 2.5):

$$\text{tr}(X) = \tau(\mathbb{1}_{(1, \infty)}(X)) \text{ for any } X \in L^1(\mathcal{M})_+. \quad (7)$$

3 Our result and its proof

We first state the general form of Ke Li’s lemma. As discussed in section 2, the vector Ω_σ and the operator $\Delta_{\rho|\sigma}$ are meant in any standard representation of \mathcal{M} .

THEOREM 1. Let \mathcal{M} be a von Neumann algebra, and ρ, σ be two faithful normal states on \mathcal{M} . For any $\epsilon > 0$ there exists a test $T \in \mathcal{M}$ such that

$$\alpha(T) \leq \epsilon \quad \beta(T) \leq \langle \Omega_\sigma, \mathbb{1}_{(\epsilon, +\infty)}(\Delta_{\rho|\sigma})\Omega_\sigma \rangle.$$

Remark 2. Our assumption that ρ and σ are faithful is unessential, simplifies the proof and the description of $\Delta_{\rho|\sigma}$ in section 2, and can be dropped by considering an approximation of general ρ, σ by faithful states.

The rest of this section is dedicated to the proof of Theorem 1. We consider the standard representation $(\pi, L^2(\mathcal{M}), L^2(\mathcal{M})_+, \mathbb{J})$ of \mathcal{M} that we discussed in the second part of section 2. Once again we simply write \mathcal{M} for $\pi(\mathcal{M})$, so that \mathcal{M} will act by left-multiplication on $L^2(\mathcal{M})$.

Let \mathfrak{P} be the following orthogonal projection operator on $L^2(\mathcal{M})$:

$$\mathfrak{P} = \mathbb{1}_{(0, \epsilon]}(\Delta_{\rho|\sigma}).$$

We recall that if X is an element of $L^2(\mathcal{M})$ then its *support* in \mathcal{M} , which we denote by $\ell(X)$ (with an ℓ for “left support”), is defined as the smallest orthogonal projector P in \mathcal{M} such that $PX = X$. It is immediate to show that this projector has range $\overline{\mathcal{M}'X}$. Similarly we define the support of $X \in L^2(\mathcal{M})$ in \mathcal{M}' , which we denote by $r(X)$ (with an r for “right support”) as the smallest orthogonal projector P' in \mathcal{M}' such that $P'X = X$.

First define our test T as the *support* in \mathcal{M} of $\mathfrak{P}(\Omega_\sigma)$. By definition $\mathfrak{P}(\Omega_\sigma)$ is an element of $L^2(\mathcal{M})$, and therefore a densely defined closed operator on \mathcal{K} which is affiliated with \mathcal{N} ; the polar decomposition of $\mathfrak{P}(\Omega_\sigma)$ as an operator on \mathcal{K} has partial isometry U which from the properties of $L^2(\mathcal{M})$ belongs to \mathcal{M} . Then UU^* is by definition the smallest orthogonal projector P in $\mathcal{B}(\mathcal{K})$ such that $P\mathfrak{P}(\Omega_\sigma) = \mathfrak{P}(\Omega_\sigma)$, and belongs to \mathcal{M} . Therefore T is equivalently defined as the orthogonal projector onto $\mathcal{M}'\mathfrak{P}(\Omega_\sigma)$, as UU^* , or as left-multiplication by the orthogonal projector onto the closed range of $\mathfrak{P}(\Omega_\sigma)$ when the latter is viewed as an operator on \mathcal{K} . Note that by definition, $T \in \mathcal{M}$.

From the first expression for T we have

$$\sigma(\text{Id} - T) = \|\Omega_\sigma - T\Omega_\sigma\|^2 \leq \|\Omega_\sigma - \mathfrak{P}(\Omega_\sigma)\|^2 = \langle \Omega_\sigma, \mathbb{1}_{(\epsilon, +\infty)}(\Delta_{\rho|\sigma})\Omega_\sigma \rangle$$

where:

- the first equality follows from (5) and the fact that $\text{Id} - T$ is a self-adjoint projection,
- the inequality follows from the inequality $\|\Psi - T\Psi\| \leq \|\Psi - \Phi\|$, valid for any Ψ in $L^2(\mathcal{M})$ and $\Phi \in \text{Ran } T$ by definition of an orthogonal projection, and the fact that $\mathfrak{P}(\Omega_\sigma)$ is an element of $\mathcal{M}'\mathfrak{P}(\Omega_\sigma) \subset \text{Ran } T$,
- the second equality follows from (5) and the fact that $\text{Id} - \mathfrak{P}$ is a self-adjoint projection.

This proves the second bound in Theorem 1, and we now move on to prove the first bound.

Let $\varrho = \int_0^1 \lambda \, d\mathfrak{e}_\rho(\lambda)$ and $\varsigma = \int_0^1 \mu \, d\mathfrak{e}_\sigma(\mu)$ be spectral decompositions of ϱ, ς in \mathcal{K} . Then from the definition in the second part of section 2,

$$\Delta_{\rho|\sigma} = \varrho \mathbf{J} \varsigma^{-1} \mathbf{J} \text{ on } \mathcal{M}\Omega_\sigma$$

so that, denoting $\tilde{\mathfrak{e}}_\sigma := \mathbf{J} \mathfrak{e}_\sigma \mathbf{J}$,

$$\Delta_{\rho|\sigma} = \int_{\mathbb{R}_+^2} \lambda \mu^{-1} \, d\mathfrak{e}_\rho(\lambda) \, d\tilde{\mathfrak{e}}_\sigma(\mu)$$

at least on $\mathcal{M}\Omega_\sigma$, and since the latter integral expression defines a closed operator, equality holds everywhere. Approximating $\mathbf{1}_{(0,\epsilon]}$ by polynomials weakly, it is easy to see that

$$\mathfrak{P} = \int_{\mathbb{R}_+^2} \mathbf{1}_{(0,\epsilon]}(\lambda \mu^{-1}) \, d\mathfrak{e}_\rho(\lambda) \, d\tilde{\mathfrak{e}}_\sigma(\mu).$$

Applying Fubini's theorem twice, we have

$$\begin{aligned} \mathfrak{P}(\Omega_\sigma) &= \int_{\mathbb{R}_+^2} \mathbf{1}_{(0,\epsilon]}(\lambda \mu^{-1}) \mu^{1/2} \, d\mathfrak{e}_\rho(\lambda) \, d\tilde{\mathfrak{e}}_\sigma(\mu) \\ &= \int_0^\infty \mu^{1/2} \mathbf{1}_{(0,\epsilon\mu]}(\varrho) \, d\tilde{\mathfrak{e}}_\sigma(\mu). \end{aligned} \quad (8)$$

We now prove a regularity lemma that will allow us to approximate T thanks to Riemann sums approximating the expression (8) for $\mathfrak{P}(\Omega_\sigma)$. Assume therefore that $(X_n)_n$ is a sequence of bounded operators on \mathcal{K} that converges to X strongly, and let p_n (respectively p) be the support of X_n (respectively X) as an operator on \mathcal{K} . For any ξ in \mathcal{K} we have using $p_n X_n \xi = X_n \xi$ and $p X \xi = X \xi$:

$$\|p_n p X \xi - p X \xi\| \leq \|p_n X \xi - p_n X_n \xi\| + \|p_n X_n \xi - X \xi\| \leq 2 \|X_n \xi - X \xi\|$$

and the upper bound goes to zero as $n \rightarrow \infty$. In particular, $p_n p$ converges to p strongly, which further implies that $(1-p)p_n p$ and $p p_n (1-p)$ converge to 0 weakly. Now consider P_n (respectively P) the operator of left-multiplication by p_n (respectively p) on $L^2(\mathcal{M})$. By the normality of ρ , we get

$$\begin{aligned} \rho(P) &= \lim_n \rho(P P_n P) \\ &= \lim_n \rho(P P_n P) + \lim_n \rho((1-P) P_n P) + \lim_n \rho(P P_n (1-P)) \\ &= \lim_n (\rho(P_n) - \rho((1-P) P_n (1-P))) \\ &\leq \limsup_n \rho(P_n). \end{aligned} \quad (9)$$

Recall now that our test T is defined as the support $\ell(X)$ of $X := \mathfrak{P}(\Omega_\sigma)$. Let $(X_n)_n$ be a sequence of Riemann sums of the form

$$X_n = \sum_i \mu_i^{1/2} \mathbb{1}_{[0, \epsilon \mu_i]}(\varrho) (\tilde{e}_\sigma(\mu_i) - \tilde{e}_\sigma(\mu_{i-1}))$$

that converges strongly to X , and let T_n be the support $\ell(X_n)$ of X_n . From (9), we have $\rho(T) \leq \limsup_n \rho(T_n)$ and we therefore consider $\rho(T_n)$. Obviously

$$T_n \leq \sum_i \ell(\mathbb{1}_{[0, \epsilon \mu_i]}(\varrho) (\tilde{e}_\sigma(\mu_i) - \tilde{e}_\sigma(\mu_{i-1}))) =: T'_n,$$

where we omitted the scalar coefficients $\lambda_i^{1/2}$ since X and its multiple λX must have the same range for any $X \in L^2(\mathcal{M})$. Note that $\mathbb{1}_{[0, \epsilon \mu_i]}(\varrho)$ is a projection whose range contains that of $\ell(\mathbb{1}_{[0, \epsilon \mu_i]}(\varrho) (\tilde{e}_\sigma(\mu_i) - \tilde{e}_\sigma(\mu_{i-1})))$, so we write

$$\ell(\mathbb{1}_{[0, \epsilon \mu_i]}(\varrho) (\tilde{e}_\sigma(\mu_i) - \tilde{e}_\sigma(\mu_{i-1}))) = \mathbb{1}_{[0, \epsilon \mu_i]}(\varrho) \ell(\mathbb{1}_{[0, \epsilon \mu_i]}(\varrho) (\tilde{e}_\sigma(\mu_i) - \tilde{e}_\sigma(\mu_{i-1}))).$$

Note that (see e.g. [4])

$$\tau(\mathbb{1}_{(1, \infty)}(X)) \leq \tau(\mathbb{1}_{(1, \infty)}(Y)) \quad \text{if } 0 \leq X \leq Y.$$

Thus together with (7),

$$\begin{aligned} \text{tr}(\varrho T'_n) &= \sum_i \text{tr}(\varrho \mathbb{1}_{[0, \epsilon \mu_i]}(\varrho) \ell(\mathbb{1}_{[0, \epsilon \mu_i]}(\varrho) (\tilde{e}_\sigma(\mu_i) - \tilde{e}_\sigma(\mu_{i-1})))) \\ &= \sum_i \tau\left(\mathbb{1}_{(1, \infty)}\left(\varrho \mathbb{1}_{[0, \epsilon \mu_i]}(\varrho) \ell(\mathbb{1}_{[0, \epsilon \mu_i]}(\varrho) (\tilde{e}_\sigma(\mu_i) - \tilde{e}_\sigma(\mu_{i-1})))\right)\right) \\ &\leq \sum_i \tau\left(\mathbb{1}_{(1, \infty)}\left(\epsilon \mu_i \ell(\mathbb{1}_{[0, \epsilon \mu_i]}(\varrho) (\tilde{e}_\sigma(\mu_i) - \tilde{e}_\sigma(\mu_{i-1})))\right)\right) \\ &\leq \sum_i \mathbb{1}_{(1, \infty)}(\epsilon \mu_i) \tau\left(\ell(\mathbb{1}_{[0, \epsilon \mu_i]}(\varrho) (\tilde{e}_\sigma(\mu_i) - \tilde{e}_\sigma(\mu_{i-1})))\right). \end{aligned}$$

Note that $p_{i,1} := \mathbb{1}_{[0, \epsilon \mu_i]}(\varrho)$ and $p_{i,2} := \tilde{e}_\sigma(\mu_i) - \tilde{e}_\sigma(\mu_{i-1})$ are projections. In this case it is folklore that $\ell(p_{i,1} p_{i,2})$ is equivalent to $r(p_{i,1} p_{i,2})$, i.e. that there exists a unitary U in \mathcal{M} such that $\ell(p_{i,1} p_{i,2}) = UU^*$ and $r(p_{i,1} p_{i,2}) = U^*U$, and

$$r(p_{i,1} p_{i,2}) = p_{i,1} \vee p_{i,2}^\perp - p_{i,2}^\perp \leq 1 - p_{i,2}^\perp = p_{i,2}.$$

In other words, $\ell(p_{i,1} p_{i,2})$ is equivalent to a subprojection of $p_{i,2}$, whence $\tau(\ell(p_{i,1} p_{i,2})) \leq \tau(p_{i,2})$. So the previous inequality reads

$$\text{tr}(\varrho T'_n) \leq \tau\left(\sum_i \mathbb{1}_{(1, \infty)}(\epsilon \mu_i) \tilde{e}_\sigma(\mu_i) - \tilde{e}_\sigma(\mu_{i-1})\right)$$

Taking the limit of Riemann sums and using (7) again, we get

$$\text{tr}(\varrho T) \leq \tau\left(\int \mathbb{1}_{(1, \infty)}(\epsilon \mu) d\tilde{e}_\sigma(\mu)\right) = \tau(\mathbb{1}_{(1, \infty)}(\epsilon \varsigma)) = \text{tr}(\epsilon \varsigma) = \epsilon.$$

The proof is complete. \square

4 Applications and comparison with the Neyman–Pearson test

Let us now return to the practical task of interest, which is of discriminating between the states ρ and σ . As we mentioned in the introduction, one will typically try to make both error probabilities $\alpha(T)$ and $\beta(T)$ small (recall that $\alpha(T)$ and $\beta(T)$ are defined by (1)). However, it is expected that there is a tradeoff between one error probability and the other. One must therefore make more precise in what sense we want to “make both $\alpha(T)$ and $\beta(T)$ ” small. One possible sense is natural if we assume that prior probabilities p and $q := 1 - p$ can be assigned to the states ρ and σ respectively. One may then decide to minimize the quantity $p\alpha(T) + q\beta(T)$. This is the realm of symmetric hypothesis testing, as opposed to asymmetric hypothesis testing where one wishes e.g. to minimize $\alpha(T)$ under a specific constraint on $\beta(T)$.

To state the relevant result for symmetric hypothesis testing we define a specific test T_{NP} called the *Neyman–Pearson* test. In the finite-dimensional case this test is defined as

$$T_{\text{NP}} := \ell(\mathbb{1}_{\mathbb{R}_+}(p\varrho - q\varsigma)).$$

In the general case the definition is formally similar:

$$T_{\text{NP}} := \ell((p\rho - q\sigma)_+)$$

but here one needs to rely on the (unique) Jordan decomposition

$$p\rho - q\sigma = (p\rho - q\sigma)_+ - (p\rho - q\sigma)_-$$

with $(p\rho - q\sigma)_\pm$ two positive normal forms on \mathcal{M} . We will freely denote T_{NP} as $T_{\text{NP}}(p, q)$ when we need to emphasize the dependency on p, q .

The following result is known as the quantum Neyman–Pearson lemma and was proved in [1] in the finite-dimensional case and in [6] in the general case:

$$p\alpha(T_{\text{NP}}) + q\beta(T_{\text{NP}}) = \inf_T (p\alpha(T) + q\beta(T)) \quad (10)$$

where the infimum is indifferently over the set of orthogonal projections of \mathcal{M} , or over the set of elements T of \mathcal{M} satisfying $0 \leq T \leq \text{Id}$. The associated relevant upper bound was also proven in [1, 6] and is called the *Chernoff bound*:

$$p\alpha(T_{\text{NP}}) + q\beta(T_{\text{NP}}) \leq \inf_{s \in [0, 1]} p^s q^{1-s} \langle \Omega_\sigma, \Delta_{\rho|\sigma}^s \Omega_\sigma \rangle. \quad (11)$$

Note that the Chernoff bound (with an additional lower bound, see [6, 10]) is sufficient to prove the standard Stein’s lemma and Hoeffding bounds for asymmetric hypothesis testing, see [6].

Denote by T_{KL} (or, again, by $T_{\text{KL}}(\epsilon)$ when we need to emphasize the dependence on ϵ) the test constructed in section 3 (we confess a slight inconsistency in our choice of notation, as the present T_{KL} is the $L_{T_{\text{KL}}}$ of section 1). A first interesting observation is that, in the case where \mathcal{M} is commutative, or when ϱ and ς commute, then

$$T_{\text{KL}}(\epsilon) = T_{\text{NP}}\left(\frac{1}{1 + \epsilon}, \frac{\epsilon}{1 + \epsilon}\right).$$

This can be easily seen in the finite-dimensional case, i.e. when $\mathcal{M} = \{L_X, X \in \mathcal{B}(\mathbb{C}^n)\}$ as described in section 1 (and using the same notation). Using the fact that the families $(a_x)_x$ and $(b_y)_y$ are the same (up to permutation) when ϱ and ς commute, expression (4) takes the form

$$\mathbb{1}_{(\epsilon, +\infty)}(\Delta_{\rho|\sigma})\Omega_\sigma = \sum_{x=1}^n \mathbb{1}_{\lambda_x > \epsilon \mu_x} \mu_x^{1/2} |a_x\rangle\langle a_x|. \quad (12)$$

On the other hand, when $p = \frac{1}{1+\epsilon}$,

$$\mathbb{1}_{\mathbb{R}_+}(p\varrho - q\varsigma) = \sum_{x=1}^n \mathbb{1}_{\lambda_x > \epsilon \mu_x} |a_x\rangle\langle a_x|. \quad (13)$$

Therefore, the supports of the operators in (12) and (13) are the same, so that $T_{\text{KL}} = T_{\text{NP}}$.

Now let us compare the merits of Theorem 1, in comparison to the Chernoff bound (11). From now on we always consider $p = \frac{1}{1+\epsilon}$ so that p satisfies the relation $q/p = \epsilon$. The Chernoff bound (11) implies

$$\rho(T_{\text{NP}}) \leq \epsilon \quad \sigma(T_{\text{NP}}) \leq \epsilon^{-s} \langle \Omega_\sigma, \Delta_{\rho|\sigma}^s \Omega_\sigma \rangle \quad (14)$$

whereas Theorem 1 gives

$$\rho(T_{\text{KL}}) \leq \epsilon \quad \sigma(T_{\text{KL}}) \leq \langle \Omega_\sigma, \mathbb{1}_{(\epsilon, +\infty)}(\Delta_{\rho|\sigma})\Omega_\sigma \rangle. \quad (15)$$

If the upper bound for $\sigma(T_{\text{KL}})$ in (15) is bounded by an application of Markov's exponential inequality, then (14) and (15) yield the same pair of bounds. It therefore turns out that the estimates (15) given by Theorem 1 are no better than those (14) given by the Chernoff bound, unless one has an estimate for $\langle \Omega_\sigma, \mathbb{1}_{(\epsilon, +\infty)}(\Delta_{\rho|\sigma})\Omega_\sigma \rangle$ more precise than that given by Markov's exponential inequality. It therefore seems that our result is better suited to situations where specific information on the tails of the distribution of $\Delta_{\rho|\sigma}$ with respect to the state Ω_σ is available. This was the case, in particular, in [3, 9, 12].

These three papers are concerned with the same general question, which we now describe informally. Consider two sequences $(\rho_n)_n$ and $(\sigma_n)_n$ of states on possibly different algebras \mathcal{M}_n , which one would like to discriminate “for large n ”. One will therefore try and construct a sequence $(T_n)_n$ of tests $T_n \in \mathcal{M}_n$ for which $(\alpha(T_n))_n$ and $(\beta(T_n))_n$ are “small” in some sense. The question in [3, 9, 12] is essentially the following: if one only insists on having $\sup_n \alpha(T_n) \leq \epsilon$, what is the best rate of decrease of $\beta(T_n)$ one can hope for? The standard result is called *Stein's lemma* and is well-understood (see [6] and references therein): it gives general assumptions under which, essentially, an exponential decrease of any rate r strictly smaller than a quantity D called the *relative entropy* of $(\rho_n)_n$ and $(\sigma_n)_n$ is possible, but an exponential decrease of a rate strictly larger than D is not possible. The papers [3, 9, 12] ask (in the case where every \mathcal{M}_n is a finite-dimensional algebra) whether a rate *equal*

to D is possible, and what is then the “next order” describing the optimal rate of decrease for $\beta(T_n)$. In particular, these papers identify conditions under which one can construct a sequence (T_n) satisfying both conditions $\sup_n \alpha(T_n) \leq \epsilon$ and

$$-\log \beta(T_n) = nD + \sqrt{n}t + o(\sqrt{n})$$

for any $t < \Phi^{-1}(\epsilon)V$ but not for $t > \Phi^{-1}(\epsilon)V$. Here Φ is the cumulative distribution function of a standard normal distribution, V is a quantity depending on the sequences $(\rho_n)_n$ and $(\sigma_n)_n$.

The article [12] used a submultiplicativity property of the sequences $(\rho_n)_n$ and $(\sigma_n)_n$ to derive concentration inequalities for the distribution of $\Delta_{\rho_n|\sigma_n}$ with respect to the state Ω_{σ_n} ; the nature of the considered sequences $(\rho_n)_n$ and $(\sigma_n)_n$, however, imposes the assumption that every \mathcal{M}_n is finite-dimensional. In that case Theorem 1 follows from Ke Li’s original result and the present general extension is not needed.

One can say in a similar fashion that [9] used a multiplicativity property of the states ρ and σ , but the result in [9] is generalized by Theorem 1 of [3]. The proof of the latter, however, only relies on the Chernoff bound (11) and on Theorem 1. Since the Chernoff bound is proved in the general case in [6], the present proof of Theorem 1 immediately extends Theorem 1 of [3] to the case of general von Neumann algebras.

References

- [1] Konrad Audenaert, Michael Nussbaum, and Arleta Szkoła. Asymptotic error rates in quantum hypothesis testing. *Comm. Math. Phys.*, 279, 2008.
- [2] Ola Bratteli and Derek W. Robinson. *Operator algebras and quantum statistical mechanics. 1*. Texts and Monographs in Physics. Springer-Verlag, New York, second edition, 1987. C^* - and W^* -algebras, symmetry groups, decomposition of states.
- [3] Nilanjana Datta, Yan Pautrat, and Cambyse Rouzé. Second-order asymptotics for quantum hypothesis testing in settings beyond i.i.d.—quantum lattice systems and more. *J. Math. Phys.*, 57(6):062207, 26, 2016.
- [4] Thierry Fack and Hideki Kosaki. Generalized s -numbers of τ -measurable operators. *Pacific J. Math.*, 123(2):269–300, 1986.
- [5] Uffe Haagerup. L^p -spaces associated with an arbitrary von Neumann algebra. In *Algèbres d’opérateurs et leurs applications en physique mathématique (Proc. Colloq., Marseille, 1977)*, volume 274 of *Colloq. Internat. CNRS*, pages 175–184. CNRS, Paris, 1979.
- [6] Vojkan Jakšić, Yoshiko Ogata, Claude-Alain Pillet, and Robert Seiringer. Quantum hypothesis testing and non-equilibrium statistical mechanics. *Rev. Math. Phys.*, 24(6):1230002, 67, 2012.

- [7] Vojkan Jakšić, Yoshiko Ogata, Yan Pautrat, and Claude-Alain Pillet. Entropic fluctuations in quantum statistical mechanics. an introduction. *Quantum Theory from Small to Large Scales, Lecture Notes of the Les Houches Summer School*, 95(978-0-19-965249-5):pp.213–410, 2012.
- [8] Eneet Kaur and Mark M. Wilde. Upper bounds on secret-key agreement over lossy thermal bosonic channels. *Phys. Rev. A*, 96:062318, Dec 2017.
- [9] Ke Li. Second-order asymptotics for quantum hypothesis testing. *Ann. Statist.*, 42(1):171–189, 02 2014.
- [10] Michael Nussbaum and Arleta Szkoła. The Chernoff lower bound for symmetric quantum hypothesis testing. *Ann. Statist.*, 37(2):1040–1057, 2009.
- [11] Samad Khabbazi Oskouei, Stefano Mancini, and Mark M. Wilde. Union bound for quantum information processing. *Proc. R. Soc. A*, 475(2221):20180612, 19, 2019.
- [12] Cambyse Rouzé and Nilanjana Datta. Finite blocklength and moderate deviation analysis of hypothesis testing of correlated quantum states and application to classical-quantum channels with memory. *IEEE Transactions on Information Theory*, 64(1):593–612, 2018.
- [13] Marianne Terp. L^p spaces associated with von neumann algebras. *Notes, Math. Institute, Copenhagen University*, 1981.
- [14] Marco Tomamichel and Vincent Y. F. Tan. Second-order asymptotics for the classical capacity of image-additive quantum channels. *Comm. Math. Phys.*, 338(1):103–137, 2015.
- [15] Mark M. Wilde, Marco Tomamichel, and Mario Berta. Converse bounds for private communication over quantum channels. *IEEE Transactions on Information Theory*, 63(3):1792–1817, 2017.