

Density-dependent nn-potential from subleading chiral three-neutron forces: Long-range terms¹

N. Kaiser

Physik-Department T39, Technische Universität München, D-85747 Garching, Germany

Abstract

The long-range terms of the subleading chiral three-nucleon force [published in Phys. Rev. C77, 064004 (2008)] are specified to the case of three neutrons. From these $3n$ -interactions an effective density-dependent neutron-neutron potential V_{med} in pure neutron matter is derived. Following the division of the pertinent $3n$ -diagrams into two-pion exchange, two-pion-one-pion exchange and ring topology, all self-closings and concatenations of two neutron-lines to an in-medium loop are evaluated. The momentum and k_n -dependent potentials associated with the spin-operators 1 , $\vec{\sigma}_1 \cdot \vec{\sigma}_2$, $\vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q}$, $i(\vec{\sigma}_1 + \vec{\sigma}_2) \cdot (\vec{q} \times \vec{p})$, $(\vec{\sigma}_1 \cdot \vec{p} \vec{\sigma}_2 \cdot \vec{p} + \vec{\sigma}_1 \cdot \vec{p}' \vec{\sigma}_2 \cdot \vec{p}')$ and $\vec{\sigma}_1 \cdot (\vec{q} \times \vec{p}) \vec{\sigma}_2 \cdot (\vec{q} \times \vec{p})$ are expressed in terms of functions, which are either given in closed analytical form or require at most one numerical integration. The subsubleading chiral $3N$ -force is treated in the same way. The obtained results for V_{med} are helpful to implement the long-range chiral three-body forces into advanced neutron matter calculations.

1 Introduction

Three-nucleon forces are an indispensable ingredient in accurate few-nucleon and nuclear structure calculations. Nowadays, chiral effective field theory is the appropriate tool to construct systematically the nuclear interactions in harmony with the symmetries of QCD. Three-nucleon forces appear first at $N^2\text{LO}$, where they consist of a zero-range contact-term ($\sim c_E$), a mid-range 1π -exchange component ($\sim c_D$) and a long-range 2π -exchange component ($\sim c_{1,3,4}$). The complete calculation of the chiral $3N$ -forces to subleading order $N^3\text{LO}$ [1, 2] and even to subsubleading order $N^4\text{LO}$ [3, 4] has been achieved during the past decade by the Bochum-Bonn group. At present the focus lies on constructing $3N$ -forces in chiral effective field theory with explicit $\Delta(1232)$ -isobars, for which the long-range 2π -exchange component has been derived recently in ref. [5] at order $N^3\text{LO}$.

However, for the variety of existing many-body methods, that are commonly employed in calculations of nuclear matter or medium mass and heavy nuclei, it is technically very challenging to include the chiral three-nucleon forces directly. An alternative and approximate approach is to use instead a density-dependent two-nucleon interaction V_{med} that originates from the underlying $3N$ -force. When restricting to on-shell scattering of two nucleons in isospin-symmetric spin-saturated nuclear matter, $N_1(\vec{p}) + N_2(-\vec{p}) \rightarrow N_1(\vec{p}') + N_2(-\vec{p}')$, the resulting in-medium NN-potential V_{med} has the same isospin- and spin-structure as the free NN-potential. The analytical expressions for V_{med} from the leading chiral $3N$ -force at $N^2\text{LO}$ (involving the parameters $c_{1,3,4}$, c_D and c_E) have been presented in ref. [6] and these have found many applications in recent years. But in order to perform nuclear many-body calculations that are consistent with their input at the two-body level, one needs also V_{med} derived from the subleading chiral $3N$ -forces at order $N^3\text{LO}$. In two recent works this task has been completed for the short-range terms and relativistic $1/M$ -corrections in ref. [7], and for the long-range terms in ref. [8]. The continuation of this construction from the intermediate-range terms of the subsubleading chiral $3N$ -force at order $N^3\text{LO}$ has been reported recently in ref. [9]. The normal ordering of chiral $3N$ -forces to density-dependent NN-interactions has also been performed by the

¹This work has been supported in part by DFG and NSFC (CRC110).

Darmstadt group using a decomposition in a Jj -coupled 3N partial-wave momentum basis [10, 11]. An advantage of this purely numerical approach is that the restriction to on-shell kinematics in the center-of-mass frame can be avoided and a regulator function $f_R(p, q) = \exp[-(p^2 + 3q^2/4)/\Lambda_{3N}^8]$ may be included.

In this work the specification to the case of three neutrons is considered and an effective density-dependent neutron-neutron potential V_{med} in pure neutron matter is derived. The 3n-interaction is obtained from a given expression for the 3N-force by the simple substitution of isospin-operators: $\vec{\tau}_1 \cdot \vec{\tau}_2 \rightarrow 1, \vec{\tau}_1 \cdot \vec{\tau}_3 \rightarrow 1, \vec{\tau}_2 \cdot \vec{\tau}_3 \rightarrow 1$ and $\vec{\tau}_1 \cdot (\vec{\tau}_2 \times \vec{\tau}_3) \rightarrow 0$. In the next step three self-closings and six concatenations of two neutron-lines to an in-medium loop need to be evaluated. This introduces a set of loop-functions that depend on the momentum $p = |\vec{p}|$, the momentum-transfer $q = |\vec{p}' - \vec{p}|$, and the neutron Fermi-momentum k_n . Since the calculational procedure follows closely that in previous works [7, 8, 9], it is sufficient to list the results for the nn-potential V_{med} in pure neutron matter without further explanation. The analytical results for the in-medium nn-potential V_{med} derived from the subleading short-range terms and relativistic $1/M$ -corrections have been reported in ref.[12].




Figure 1: 2π -exchange topology, $2\pi 1\pi$ -exchange topology and ring topology which comprise the long- and intermediate-range chiral 3n-forces.

2 Two-pion exchange topology

One starts with the longest-range component of the subleading chiral 3N-interaction. It arises from 2π -exchange with the corresponding symbolic diagram shown on the left of Fig. 1. Using for simplification the relation $q_2^2 = q_1^2 + q_3^2 + 2\vec{q}_1 \cdot \vec{q}_3$, the expression in eq.(2.9) of ref. [1] specified to three neutrons takes the form:

$$2V_{3n} = \frac{g_A^4}{128\pi f_\pi^6} \frac{\vec{\sigma}_1 \cdot \vec{q}_1 \vec{\sigma}_3 \cdot \vec{q}_3}{(m_\pi^2 + q_1^2)(m_\pi^2 + q_3^2)} \left\{ m_\pi(m_\pi^2 + q_1^2 + q_3^2 + 2q_2^2) + (2m_\pi^2 + q_2^2)(3m_\pi^2 + q_1^2 + q_3^2 + 2q_2^2)A(q_2) \right\}, \quad (1)$$

with the pion-loop function $A(s) = (1/2s) \arctan(s/2m_\pi)$.

2.1 Contributions to in-medium nn-potential

The self-closing of neutron-line 2 gives (after relabeling $3 \rightarrow 2$) the contribution

$$V_{\text{med}}^{(0)} = -\frac{g_A^4 m_\pi k_n^3}{2(4\pi f_\pi^2)^3} \frac{\vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q}}{(m_\pi^2 + q^2)^2} \left(q^2 + \frac{5m_\pi^2}{6} \right). \quad (2)$$

From pionic vertex corrections on either neutron-line one obtains the (total) contribution:

$$V_{\text{med}}^{(1)} = \frac{2g_A^4}{(8\pi f_\pi^2)^3} \frac{\vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q}}{m_\pi^2 + q^2} \left\{ m_\pi [2k_n^3 - 8\Gamma_2 - 2m_\pi^2(\Gamma_0 + \Gamma_1) + q^2(\Gamma_0 - \Gamma_1 - 2\Gamma_3)] + S(p) + J(p, q) \right\}, \quad (3)$$

where the functions $\Gamma_\nu(p, k_n)$ are defined in the appendix of ref. [7]. The part linear in m_π comes obviously from the first line in eq.(1) and the decomposition $S(p) + J(p, q)$ is obtained by canceling momentum factors against a pion-propagator. The two functions appearing at the end of eq.(3) read:

$$\begin{aligned} S(p) = & \left[\frac{k_n^5}{15p^3} \left(m_\pi^2 + \frac{k_n^2}{14} \right) + \frac{k_n^3 m_\pi^2}{3p} + \frac{k_n^2}{3} \left(\frac{k_n^2}{4} + 2m_\pi^2 \right) + \frac{k_n^3 p}{6} + \frac{p^2}{5} \left(\frac{k_n^2}{2} - \frac{4m_\pi^2}{3} \right) - \frac{p^4}{84} \right] \arctan \frac{p + k_n}{2m_\pi} \\ & + \left[\frac{k_n^5}{15p^3} \left(m_\pi^2 + \frac{k_n^2}{14} \right) + \frac{k_n^3 m_\pi^2}{3p} - \frac{k_n^2}{3} \left(\frac{k_n^2}{4} + 2m_\pi^2 \right) + \frac{k_n^3 p}{6} + \frac{p^2}{5} \left(\frac{4m_\pi^2}{3} - \frac{k_n^2}{2} \right) + \frac{p^4}{84} \right] \arctan \frac{p - k_n}{2m_\pi} \\ & + m_\pi^3 \left[\frac{7p}{24} + \frac{1}{p} \left(\frac{m_\pi^2}{3} - \frac{k_n^2}{4} \right) + \frac{1}{p^3} \left(\frac{6m_\pi^4}{35} + \frac{k_n^2 m_\pi^2}{15} - \frac{k_n^4}{24} \right) \right] \ln \frac{4m_\pi^2 + (p + k_n)^2}{4m_\pi^2 + (p - k_n)^2} \\ & + \frac{k_n m_\pi}{7} \left[\frac{p^2}{3} - \frac{61m_\pi^2}{30} - \frac{37k_n^2}{15} - \frac{1}{p^2} \left(\frac{2k_n^4}{15} + \frac{k_n^2 m_\pi^2}{6} + \frac{6m_\pi^4}{5} \right) \right], \end{aligned} \quad (4)$$

$$\begin{aligned} J(p, q) = & \frac{1}{4q^2} \int_{p-k_n}^{p+k_n} ds (2m_\pi^2 + s^2)(2m_\pi^2 + q^2 + 2s^2) \arctan \frac{s}{2m_\pi} \\ & \times \left\{ \frac{k_n^2 - (p-s)^2}{p} + \frac{q^2 - m_\pi^2 - s^2}{q} \ln \frac{qX + 2\sqrt{W}}{(2p+q)[m_\pi^2 + (q-s)^2]} \right\}, \end{aligned} \quad (5)$$

with the auxiliary polynomials:

$$\begin{aligned} X &= m_\pi^2 + 2(k_n^2 - p^2) + q^2 - s^2, \\ W &= k_n^2 q^4 + p^2(m_\pi^2 + s^2)^2 + q^2[(k_n^2 - p^2)^2 + m_\pi^2(k_n^2 + p^2) - s^2(k_n^2 + p^2 + m_\pi^2)]. \end{aligned} \quad (6)$$

Finally, the two diagrams related to double exchange lead to the expression:

$$\begin{aligned} V_{\text{med}}^{(3)} = & \frac{g_A^4}{(8\pi f_\pi^2)^3} \left\{ [m_\pi + (2m_\pi^2 + q^2)A(q)] \left[\frac{8k_n^3}{3} + 2q^2(\Gamma_0 - \Gamma_1) \right] - 2m_\pi^2 \Gamma_0 [3m_\pi \right. \\ & + (2m_\pi^2 + q^2)A(q)] + (2m_\pi^2 + q^2)[m_\pi^3 - 2m_\pi q^2 - (2m_\pi^4 + 5m_\pi^2 q^2 + 2q^4)A(q)] G_0 \\ & + i(\vec{\sigma}_1 + \vec{\sigma}_2) \cdot (\vec{q} \times \vec{p}) \left\{ 2[m_\pi + (2m_\pi^2 + q^2)A(q)](\Gamma_0 + \Gamma_1) \right. \\ & \left. \left. + [2m_\pi q^2 - m_\pi^3 + (2m_\pi^4 + 5m_\pi^2 q^2 + 2q^4)A(q)](G_0 + 2G_1) \right\} \right\}. \end{aligned} \quad (7)$$

3 Two-pion-one-pion exchange topology

Next, one considers the $2\pi 1\pi$ -exchange three-neutron interaction represented by the middle diagram in Fig. 1. According to eqs.(2.16)-(2.20) in ref.[1] this chiral 3n-interaction can be written in the form:

$$\begin{aligned} V_{3n} = & \frac{g_A^4}{256\pi f_\pi^6} \frac{\vec{\sigma}_3 \cdot \vec{q}_3}{m_\pi^2 + q_3^2} \left\{ \vec{\sigma}_2 \cdot \vec{q}_1 \vec{q}_1 \cdot \vec{q}_3 f_1(q_1) + \vec{\sigma}_2 \cdot \vec{q}_1 f_2(q_1) \right. \\ & \left. + \vec{\sigma}_2 \cdot \vec{q}_3 f_3(q_1) + \vec{\sigma}_1 \cdot \vec{q}_1 \vec{q}_1 \cdot \vec{q}_3 f_4(q_1) + \vec{\sigma}_1 \cdot \vec{q}_3 f_5(q_1) + \vec{\sigma}_2 \cdot \vec{q}_3 \vec{q}_1 \cdot \vec{q}_3 f_8(q_1) \right\}, \end{aligned} \quad (8)$$

where a common factor $g_A^4/(256\pi f_\pi^6)$ has been pulled out, and thus the reduced functions $f_j(s)$ read:

$$f_1(s) = \frac{m_\pi}{s^2}(1 - 2g_A^2) - \frac{g_A^2 m_\pi}{4m_\pi^2 + s^2} + \left[1 + g_A^2 + \frac{4m_\pi^2}{s^2}(2g_A^2 - 1)\right] A(s), \quad (9)$$

$$f_2(s) = 6m_\pi + 6(2m_\pi^2 + s^2)A(s), \quad (10)$$

$$f_3(s) = (1 - g_A^2)[3m_\pi + (8m_\pi^2 + 3s^2)A(s)], \quad (11)$$

$$f_5(s) = -s^2 f_4(s) = 2g_A^2 s^2 A(s), \quad (12)$$

with $A(s)$ defined after eq.(1). In perspective one notes that at subsubleading order [4] the chiral $2\pi 1\pi$ -exchange $3n$ -interaction has a richer spin- and momentum-dependence and also the last term involving $f_8(q_1)$ is needed to represent all diagrams belonging to this topology at N⁴LO. In the three-neutron case the other relevant functions are combined as: $f_1(s) \equiv f_1(s) + f_6(s)$, $f_2(s) \equiv f_2(s) + f_7(s)$, $f_3(s) \equiv f_3(s) + f_9(s)$ with $f_{1,2,3,6,7,9}(s)$ on the right hand sides given in eqs.(2-12) of ref.[9].

3.1 Contributions to in-medium NN-potential

Again, one lists the four contributions from V_{3n} in eq.(8) to the in-medium nn-potential V_{med} . The self-closing of the neutron lines 1 and 2 lead to nonvanishing contributions, and with the relations $f_3(0) = 5(1 - g_A^2)m_\pi$ and $f_5(q) + q^2 f_4(q) = 0$ one obtains:

$$V_{\text{med}}^{(0)} = \frac{5g_A^4(1 - g_A^2)m_\pi k_n^3}{384\pi^3 f_\pi^6} \frac{\vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q}}{m_\pi^2 + q^2}, \quad (13)$$

which is of the form: $1\pi^0$ -exchange nn-interaction times a factor linear in the neutron density $\rho_n = k_n^3/3\pi^2$. On the other hand the vertex corrections by $1\pi^0$ -exchange, incorporated in eq.(8) through the second factor $\vec{\sigma}_3 \cdot \vec{q}_3/(m_\pi^2 + q_3^2)$, produce the contribution:

$$\begin{aligned} V_{\text{med}}^{(1)} = & \frac{g_A^4}{(8\pi f_\pi^2)^3} \left\{ \left(2m_\pi^2 \Gamma_0 - \frac{4k_n^3}{3}\right) f_3(q) - \left(2\Gamma_2 + \frac{q^2}{2} \tilde{\Gamma}_3\right) q^2 f_1(q) + \tilde{\Gamma}_1 q^2 f_2(q) - 2\vec{\sigma}_1 \cdot \vec{\sigma}_2 \Gamma_2 f_5(q) \right. \\ & - (\vec{\sigma}_1 \cdot \vec{p} \vec{\sigma}_2 \cdot \vec{p} + \vec{\sigma}_1 \cdot \vec{p}' \vec{\sigma}_2 \cdot \vec{p}') \tilde{\Gamma}_3 f_5(q) - \vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q} \left(2\Gamma_2 + \frac{q^2}{2} \tilde{\Gamma}_3\right) f_4(q) \\ & \left. + i(\vec{\sigma}_1 + \vec{\sigma}_2) \cdot (\vec{q} \times \vec{p}) \left[\frac{q^2}{2} \tilde{\Gamma}_3 f_1(q) - \tilde{\Gamma}_1 f_2(q)\right] + \left(\frac{2k_n^3}{3} - m_\pi^2 \tilde{\Gamma}_1\right) q^2 f_8(q) \right\}. \end{aligned} \quad (14)$$

Here, the frequently occuring combinations $\tilde{\Gamma}_1(p) = \Gamma_0(p) + \Gamma_1(p)$ and $\tilde{\Gamma}_3(p) = \Gamma_0(p) + 2\Gamma_1(p) + \Gamma_3(p)$ have been introduced. Moreover, the vertex corrections by 2π -exchange (compiled in the expression in curly brackets of eq.(8)) can be summarized as the $1\pi^0$ -exchange nn-interaction times a (p, q, k_f) -dependent factor:

$$V_{\text{med}}^{(2)} = \frac{g_A^4}{(8\pi f_\pi^2)^3} \frac{\vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q}}{m_\pi^2 + q^2} [S_1(p) + q^2 S_2(p)], \quad (15)$$

with the two functions $S_{1,2}(p, k_n)$ defined by:

$$\begin{aligned} S_1(p) = & \frac{1}{8p^3} \int_{p-k_n}^{p+k_n} ds s [k_n^2 - (p-s)^2] \left\{ [(p+s)^2 - k_n^2] f_2(s) - 8p^2 [f_3(s) + f_5(s)] \right. \\ & \left. + \frac{1}{3} [k_n^2 - (p-s)^2] (k_n^2 - p^2 - 4ps - s^2) [f_1(s) + f_4(s)] \right\}, \end{aligned} \quad (16)$$

$$S_2(p) = \frac{1}{32p^5} \int_{p-k_n}^{p+k_n} ds s [k_n^2 - (p-s)^2] [k_n^2 - (p+s)^2] \left\{ (p^2 + s^2 - k_n^2) [f_1(s) + f_4(s)] - 4p^2 f_8(s) \right\}. \quad (17)$$

Note that the integrals for $S_{1,2}(p)$ with input functions $f_{1,2,3,4,5}(s)$ as in eqs.(9-12) could be solved in terms of the functions $\arctan[(p \pm k_n)/2m_\pi]$ and $\ln[4m_\pi^2 + (p \pm k_n)^2]$.

Finally, the more complicated contribution from double exchange reads:

$$\begin{aligned} V_{\text{med}}^{(3)} = & \frac{g_A^4}{(8\pi f_\pi^2)^3} \left\{ \vec{\sigma}_1 \cdot \vec{\sigma}_2 (2I_{2,2} - 2I_{3,2} - H_{1,2} - \tilde{I}_{1,2} + H_{8,2} + \tilde{I}_{8,2}) + \vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q} \left(\frac{H_{1,1} + \tilde{I}_{1,4} + \tilde{I}_{8,5}}{2} \right. \right. \\ & - I_{2,4} - I_{3,5} + H_{8,0} - H_{8,1} \Big) + (\vec{\sigma}_1 \cdot \vec{p} \vec{\sigma}_2 \cdot \vec{p} + \vec{\sigma}_1 \cdot \vec{p}' \vec{\sigma}_2 \cdot \vec{p}') \left(I_{2,3} - I_{3,3} - \frac{H_{1,3} + \tilde{I}_{1,3}}{2} \right. \\ & + \frac{H_{8,3} + \tilde{I}_{8,3}}{2} \Big) + 2m_\pi^2 I_{5,0} - 2H_{5,0} - 3H_{4,2} - 3\tilde{I}_{4,2} + \frac{q^2}{2} (H_{4,1} + \tilde{I}_{4,4}) - p^2 (H_{4,3} + \tilde{I}_{4,3}) \\ & \left. \left. - \frac{i}{2} (\vec{\sigma}_1 + \vec{\sigma}_2) \cdot (\vec{q} \times \vec{p}) (H_{4,1} + \tilde{I}_{4,1}) \right\}. \quad (18) \end{aligned}$$

The definitions of the double-indexed functions $H_{j,\nu}(p, k_n)$, $I_{j,\nu}(p, q, k_n)$ and $\tilde{I}_{j,\nu}(p, q, k_n)$ can be found in eqs.(25-34) of ref.[8].

One should point to a typing error in the third line of eq.(24) in ref.[8]. There the spin-orbit operator should read $i(\vec{\sigma}_1 + \vec{\sigma}_2) \cdot (\vec{q} \times \vec{p})$ and not $i(\vec{\sigma}_1 + \vec{\sigma}_2) \cdot (\vec{p} \times \vec{q})$. The same typing error has occurred in the second line of eq.(19) in ref.[9]. Further typing errors that need to be corrected in ref.[7] are: $(q^2/4 - 2p^2)$ at the beginning of the last line of eq.(29), $[m_\pi^2(\gamma_0 + \gamma_1) - \Gamma_0 - \Gamma_1]$ at the end of the fourth line of eq.(30), and $(\dots + G_{0*} + 2G_{1*})$ in the fifth line of eq.(45).

4 Ring topology

The three-neutron ring-interaction, represented by the right diagram in Fig.1, possesses a more complicated structure, because any factorization property in the three momentum transfers $\vec{q}_{1,2,3}$ is lost. One starts with the basic expression for V_{3n} in the form of a three-dimensional loop-integral over (static) pion-propagators and momentum-factors [1]:

$$\begin{aligned} V_{3n} = & \frac{g_A^4}{16f_\pi^6} \int \frac{d^3 l_2}{(2\pi)^3} \frac{1}{(m_\pi^2 + l_1^2)(m_\pi^2 + l_2^2)(m_\pi^2 + l_3^2)} \left\{ \vec{l}_1 \cdot \vec{l}_2 \vec{l}_2 \cdot \vec{l}_3 - \vec{\sigma}_1 \cdot (\vec{l}_2 \times \vec{l}_3) \vec{\sigma}_3 \cdot (\vec{l}_1 \times \vec{l}_2) \right. \\ & \left. - \frac{g_A^2}{m_\pi^2 + l_2^2} \left[\vec{l}_1 \cdot \vec{l}_2 \vec{l}_1 \cdot \vec{l}_3 \vec{l}_2 \cdot \vec{l}_3 + 2\vec{\sigma}_2 \cdot (\vec{l}_1 \times \vec{l}_3) \vec{\sigma}_3 \cdot (\vec{l}_1 \times \vec{l}_2) \vec{l}_2 \cdot \vec{l}_3 - \frac{3}{2} \vec{\sigma}_1 \cdot (\vec{l}_2 \times \vec{l}_3) \vec{\sigma}_3 \cdot (\vec{l}_1 \times \vec{l}_2) \vec{l}_1 \cdot \vec{l}_3 \right] \right\}, \quad (19) \end{aligned}$$

where one has to set $\vec{l}_1 = \vec{l}_2 - \vec{q}_3$ and $\vec{l}_3 = \vec{l}_2 + \vec{q}_1$.

4.1 Self-closings of neutron-lines

For the self-closings of neutron-lines the Fermi-sphere integral gives just a factor density $\rho_n = k_n^3/3\pi^2$. Sorted according to powers of g_A^2 , the self-closing contributions to V_{med} from the 3n-ring interaction V_{3n} in eq.(19) read:

$$\begin{aligned} V_{\text{med}}^{(0)} = & \frac{g_A^4 k_n^3}{192\pi^3 f_\pi^6} \left\{ -7m_\pi - \frac{m_\pi^3}{4m_\pi^2 + q^2} - \frac{8m_\pi^2 + 3q^2}{q} \arctan \frac{q}{2m_\pi} \right. \\ & \left. + (\vec{\sigma}_1 \cdot \vec{\sigma}_2 q^2 - \vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q}) \frac{1}{2q} \arctan \frac{q}{2m_\pi} \right\}, \quad (20) \end{aligned}$$

$$V_{\text{med}}^{(0)} = \frac{g_A^6 k_n^3}{192\pi^3 f_\pi^6} \left\{ \frac{19m_\pi}{2} - \frac{5m_\pi^3}{2(4m_\pi^2 + q^2)} + \frac{4m_\pi^5}{(4m_\pi^2 + q^2)^2} + \frac{8m_\pi^2 + 3q^2}{q} \arctan \frac{q}{2m_\pi} \right. \\ \left. + \frac{1}{2q^2} (\vec{\sigma}_1 \cdot \vec{\sigma}_2 q^2 - \vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q}) \left[m_\pi - \frac{5m_\pi^3}{4m_\pi^2 + q^2} + \frac{m_\pi^2 - 7q^2}{2q} \arctan \frac{q}{2m_\pi} \right] \right\}, \quad (21)$$

where the linear divergences in the central parts have been treated by dimensional regularization.

4.2 Concatenations of neutron-lines for ring interaction $\sim g_A^4$

Treating the concatenations of two neutron-lines is somewhat simpler for the g_A^4 -part of the 3n-ring interaction V_{3n} in eq.(19), since for this component the three pion-propagators are on an equal footing. Using the assignments of momenta $\vec{l}_1, \vec{l}_2, \vec{l}_3$ for the six possible concatenations in table 1 of ref.[8], one obtains the following contributions to V_{med} , which are listed individually by specifying first their type.

- Central potential (three pieces added):

$$V_{\text{med}}^{(\text{cc})} = \frac{g_A^4}{64\pi^4 f_\pi^6} \int_0^\infty dl \left\{ l \tilde{\Gamma}_1(l) \left[l \left(\frac{q^2}{2p^2} - 2 \right) + \left(\frac{q^2}{2} + 4m_\pi^2 - 2l^2 + 2p^2 - \frac{q^2}{2p^2} (m_\pi^2 + l^2) \right) \Lambda(l) \right. \right. \\ \left. \left. + (l^2 - m_\pi^2 - p^2)(2m_\pi^2 + q^2) \Omega(l) \right] + l \left[\frac{2k_n^3}{3} - m_\pi^2 \Gamma_0(l) \right] \left[(2m_\pi^2 + q^2) \Omega(l) - 2\Lambda(l) \right] + 4k_n^3 \right\}, \quad (22)$$

with the following auxiliary functions arising from the solid-angle integration:

$$\Lambda(l) = \frac{1}{4p} \ln \frac{m_\pi^2 + (l+p)^2}{m_\pi^2 + (l-p)^2}, \quad (23)$$

$$\Omega(l) = \frac{1}{q\sqrt{B + q^2 l^2}} \ln \frac{ql + \sqrt{B + q^2 l^2}}{\sqrt{B}}, \quad (24)$$

and the abbreviation $B = [m_\pi^2 + (l+p)^2][m_\pi^2 + (l-p)^2]$. In accordance with dimensional regularization the subtraction of the asymptotic constant $-4k_n^3$ ensures the convergence of the radial integral $\int_0^\infty dl$ in eq.(22).

- Spin-spin and tensor potentials (two pieces added):

$$V_{\text{med}}^{(\text{cc})} = \frac{g_A^4}{32\pi^4 f_\pi^6} (\vec{\sigma}_1 \cdot \vec{\sigma}_2 q^2 - \vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q}) \int_0^\infty dl \frac{l \tilde{\Gamma}_1(l)}{4p^2 - q^2} \left[(B + q^2 l^2) \Omega(l) - (m_\pi^2 + l^2 + p^2) \Lambda(l) \right], \quad (25)$$

- Quadratic spin-orbit potential (two pieces added):

$$V_{\text{med}}^{(\text{cc})} = \frac{g_A^4}{16\pi^4 f_\pi^6} \vec{\sigma}_1 \cdot (\vec{q} \times \vec{p}) \vec{\sigma}_2 \cdot (\vec{q} \times \vec{p}) \int_0^\infty dl \frac{l \tilde{\Gamma}_1(l)}{4p^2 - q^2} \left\{ -\frac{l}{2p^2} + \left[\frac{m_\pi^2 + l^2 - p^2}{2p^2} \right. \right. \\ \left. \left. + \frac{4(m_\pi^2 + l^2 + p^2)}{4p^2 - q^2} \right] \Lambda(l) + \left[m_\pi^2 + 3l^2 + p^2 - \frac{4(m_\pi^2 + l^2 + p^2)^2}{4p^2 - q^2} \right] \Omega(l) \right\}, \quad (26)$$

- Spin-orbit term (one piece):

$$V_{\text{med}}^{(\text{cc})} = \frac{g_A^4}{64\pi^4 f_\pi^6} i(\vec{\sigma}_1 + \vec{\sigma}_2) \cdot (\vec{q} \times \vec{p}) \int_0^\infty dl \frac{l}{4p^2 - q^2} \left\{ \Gamma_2(l) \left[(2m_\pi^2 + 2l^2 - 2p^2 + q^2) \Omega(l) - 2\Lambda(l) \right] \right. \\ \left. + \tilde{\Gamma}_3(l) \left[(B + q^2 l^2) \Omega(l) - (m_\pi^2 + l^2 + p^2) \Lambda(l) \right] \right\}. \quad (27)$$

4.3 Concatenations of neutron-lines for ring interaction $\sim g_A^6$

When treating the concatenations of two neutron-lines for the g_A^6 -part of the 3n-ring interaction V_{3n} , one has to distinguish the cases with $\vec{l}_2 = \pm(\vec{l}_4 + \vec{l})$, because this momentum enters a squared pion-propagator in eq.(19). The Fermi-sphere integral over the squared pion-propagator (yielding the functions $\gamma_\nu(l, k_n)$) combined with the solution of the angular integral leads to the following contributions to V_{med} from the concatenations n_1 on n_3 and n_3 on n_1 .

- Central potential:

$$V_{\text{med}}^{(2)} = \frac{g_A^6}{64\pi^4 f_\pi^6} \int_0^\infty dl \left\{ 2l\gamma_2(l) \left[l \left(4 - \frac{q^2}{p^2} \right) + \left((m_\pi^2 + l^2) \frac{q^2}{p^2} - 8m_\pi^2 - 3q^2 \right) \Lambda(l) + (2m_\pi^2 + q^2)^2 \Omega(l) \right] \right. \\ \left. + \frac{l}{8} \tilde{\gamma}_3(l) \left[2l(2m_\pi^2 + 9l^2 + p^2) - \frac{lq^2}{p^2} (m_\pi^2 + 5l^2) + (2m_\pi^2 + q^2) [B + 2l^2(4m_\pi^2 + 3q^2)] \Omega(l) \right] \right. \\ \left. + \left[\frac{q^2}{p^2} (m_\pi^2 + l^2)(m_\pi^2 + 5l^2) - q^2(16l^2 + p^2) - 2B - 4m_\pi^2(m_\pi^2 + 9l^2 + p^2) \right] \Lambda(l) \right] - \frac{4k_n^3}{3} \Big\}, \quad (28)$$

- Spin-orbit potential:

$$V_{\text{med}}^{(2)} = \frac{3g_A^6}{(4\pi)^4 f_\pi^6} i(\vec{\sigma}_1 + \vec{\sigma}_2) \cdot (\vec{q} \times \vec{p}) \int_0^\infty dl \frac{l}{4p^2 - q^2} \left\{ \gamma_2(l) \left[\left[4(p^2 - l^2 - 2m_\pi^2) - 3q^2 + \frac{q^2}{p^2} (m_\pi^2 + l^2) \right] \Lambda(l) \right. \right. \\ \left. \left. + l \left(4 - \frac{q^2}{p^2} \right) + (2m_\pi^2 + q^2)(2m_\pi^2 + 2l^2 - 2p^2 + q^2) \Omega(l) \right] + \tilde{\gamma}_3(l) \left[l \left(1 - \frac{q^2}{4p^2} \right) (m_\pi^2 + l^2 + p^2) \right. \right. \\ \left. \left. + \left[\frac{q^2}{4p^2} (m_\pi^2 + l^2)^2 - B - 2m_\pi^2(m_\pi^2 + l^2 + p^2) - \frac{q^2}{4} (2m_\pi^2 + 6l^2 + 3p^2) \right] \Lambda(l) \right. \right. \\ \left. \left. + (2m_\pi^2 + q^2)(B + q^2 l^2) \Omega(l) \right] \right\}, \quad (29)$$

- Spin-spin and tensor potentials:

$$V_{\text{med}}^{(2)} = \frac{g_A^6}{16\pi^4 f_\pi^6} (\vec{\sigma}_1 \cdot \vec{\sigma}_2 q^2 - \vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q}) \int_0^\infty dl \frac{l}{4p^2 - q^2} \left\{ \gamma_2(l) \left[(m_\pi^2 + l^2 + p^2) \Lambda(l) - (B + q^2 l^2) \Omega(l) \right] \right. \\ \left. + \frac{\tilde{\gamma}_3(l)}{8} \left[l \left(1 - \frac{q^2}{4p^2} \right) (m_\pi^2 + l^2 + p^2) + \left[(l^2 + p^2 - m_\pi^2)^2 + \frac{q^2}{4} (2m_\pi^2 - 2l^2 + p^2) \right. \right. \right. \\ \left. \left. \left. - 4m_\pi^4 - 4m_\pi^2 p^2 - 4p^4 + \frac{q^2}{4p^2} (m_\pi^2 + l^2)^2 \right] \Lambda(l) + 2(m_\pi^2 - l^2 + p^2)(B + q^2 l^2) \Omega(l) \right] \right\}, \quad (30)$$

- Quadratic spin-orbit potential:

$$V_{\text{med}}^{(2)} = \frac{g_A^6}{16\pi^4 f_\pi^6} \vec{\sigma}_1 \cdot (\vec{q} \times \vec{p}) \vec{\sigma}_2 \cdot (\vec{q} \times \vec{p}) \int_0^\infty dl \frac{l}{4p^2 - q^2} \left\{ \gamma_2(l) \left[\frac{l}{p^2} + \left[3 - \frac{m_\pi^2 + l^2}{p^2} \right. \right. \right. \\ \left. \left. \left. - \frac{8(m_\pi^2 + l^2 + p^2)}{4p^2 - q^2} \right] \Lambda(l) + \left[\frac{8(m_\pi^2 + l^2 + p^2)^2}{4p^2 - q^2} - 4m_\pi^2 - 8l^2 - q^2 \right] \Omega(l) \right] \right. \\ \left. + \tilde{\gamma}_3(l) (m_\pi^2 + p^2 - l^2) \left[-\frac{l}{4p^2} + \left[\frac{m_\pi^2 + l^2 - p^2}{4p^2} + \frac{2(m_\pi^2 + l^2 + p^2)}{4p^2 - q^2} \right] \Lambda(l) \right. \right. \\ \left. \left. + \left[\frac{1}{2} (m_\pi^2 + 3l^2 + p^2) - \frac{2(m_\pi^2 + l^2 + p^2)^2}{4p^2 - q^2} \right] \Omega(l) \right] \right\}. \quad (31)$$

For the other four concatenations one has $\vec{l}_{1,3} = \pm(\vec{l}_4 + \vec{l})$ and the Fermi-sphere integral goes over an ordinary pion-propagator (yielding the functions $\gamma_\nu(l, k_n)$). When adding pieces of the same type, one obtains the following additional contributions to V_{med} from concatenations.

- Central potential (two pieces added)

$$\begin{aligned}
V_{\text{med}}^{(\text{cc})} = & \frac{g_A^6}{128\pi^4 f_\pi^6} \int_0^\infty dl \left\{ l \left[\frac{2k_n^3}{3} - m_\pi^2 \Gamma_0(l) \right] \left[\left(8 - \frac{q^2}{p^2} \right) \Lambda(l) \right. \right. \\
& + \frac{l}{B} \left(\frac{q^2}{p^2} (m_\pi^2 + l^2) - 8m_\pi^2 - 3q^2 + \frac{(2m_\pi^2 + q^2)^2}{B + q^2 l^2} (m_\pi^2 + l^2 + p^2) \right) \\
& \left. \left. + (2m_\pi^2 + q^2) \left(\frac{2m_\pi^2 + q^2}{B + q^2 l^2} (m_\pi^2 + l^2 + p^2) - 4 \right) \Omega(l) \right] - \frac{16k_n^3}{3} \right\}, \quad (32)
\end{aligned}$$

- Spin-orbit potential (one piece):

$$\begin{aligned}
V_{\text{med}}^{(\text{cc})} = & \frac{g_A^6}{128\pi^4 f_\pi^6} i(\vec{\sigma}_1 + \vec{\sigma}_2) \cdot (\vec{q} \times \vec{p}) \int_0^\infty dl l \left\{ \Gamma_2(l) \left[\left(\frac{1}{p^2} + \frac{4}{4p^2 - q^2} \right) \Lambda(l) \right. \right. \\
& + \frac{l}{B} \left(1 - \frac{m_\pi^2 + l^2}{p^2} + \frac{2m_\pi^2 + q^2}{B + q^2 l^2} (l^2 - m_\pi^2 - p^2) \right) \\
& + \left[2 - \frac{4(m_\pi^2 + l^2 + p^2)}{4p^2 - q^2} + \frac{2m_\pi^2 + q^2}{B + q^2 l^2} (l^2 - m_\pi^2 - p^2) \right] \Omega(l) \left. \right] \\
& + \tilde{\Gamma}_3(l) \left[-\frac{l}{2p^2} + \left(\frac{m_\pi^2 + l^2 - p^2}{2p^2} + \frac{2(2m_\pi^2 + l^2 + 3p^2)}{4p^2 - q^2} \right) \Lambda(l) \right. \\
& \left. \left. - \left(2B + (m_\pi^2 + p^2)(2m_\pi^2 + q^2) + l^2(2m_\pi^2 + 3q^2) \right) \frac{\Omega(l)}{4p^2 - q^2} \right] \right\}. \quad (33)
\end{aligned}$$

- Spin-spin and tensor potentials (two pieces added):

$$\begin{aligned}
V_{\text{med}}^{(\text{cc})} = & \frac{g_A^6}{64\pi^4 f_\pi^6} (\vec{\sigma}_1 \cdot \vec{\sigma}_2 q^2 - \vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q}) \int_0^\infty dl \frac{l}{4p^2 - q^2} \left\{ 5\Gamma_2(l) \left[(m_\pi^2 + l^2 + p^2) \Omega(l) - \Lambda(l) \right] \right. \\
& + \frac{\tilde{\Gamma}_3(l)}{4} \left[3l \left(\frac{q^2}{2p^2} - 2 \right) + \left[5(3m_\pi^2 + l^2 + 3p^2) - \frac{3q^2}{2p^2} (m_\pi^2 + l^2 + p^2) \right] \Lambda(l) \right. \\
& \left. \left. + \left[4l^2(2p^2 - 2m_\pi^2 - q^2) + l^4 - 9(m_\pi^2 + p^2)^2 \right] \Omega(l) \right] \right\}, \quad (34)
\end{aligned}$$

- Quadratic spin-orbit potential (two pieces added):

$$\begin{aligned}
V_{\text{med}}^{(\text{cc})} = & \frac{g_A^6}{64\pi^4 f_\pi^6} \vec{\sigma}_1 \cdot (\vec{q} \times \vec{p}) \vec{\sigma}_2 \cdot (\vec{q} \times \vec{p}) \int_0^\infty dl \frac{l}{4p^2 - q^2} \left\{ 5\Gamma_2(l) \left[\left(\frac{1}{p^2} + \frac{8}{4p^2 - q^2} \right) \Lambda(l) - \frac{l}{B} \left[\frac{m_\pi^2 + l^2}{p^2} \right. \right. \right. \\
& + \frac{4m_\pi^2 + q^2}{B + q^2 l^2} (m_\pi^2 + l^2 + p^2) - 3 \left. \right] + \left[4 - (m_\pi^2 + l^2 + p^2) \left(\frac{8}{4p^2 - q^2} + \frac{4m_\pi^2 + q^2}{B + q^2 l^2} \right) \right] \Omega(l) \left. \right] \\
& + \tilde{\Gamma}_3(l) \left[2 \left(\frac{l^2 - 9m_\pi^2 - 9p^2}{4p^2 - q^2} - \frac{m_\pi^2}{p^2} \right) \Lambda(l) + \frac{l}{B} \left[\frac{m_\pi^4 - (l^2 - p^2)^2}{q^2} + \frac{2m_\pi^2}{p^2} (m_\pi^2 + l^2) \right. \right. \\
& - \frac{11m_\pi^2}{2} + \frac{5}{2} (l^2 + p^2 - q^2) + \frac{5(4m_\pi^2 + q^2)}{2(B + q^2 l^2)} ((m_\pi^2 + p^2)^2 + l^2(m_\pi^2 - 3p^2 + q^2)) \left. \right] \\
& + \left[\frac{(l^2 - p^2)^2 - m_\pi^4}{q^2} + \frac{2(m_\pi^2 + l^2 + p^2)}{4p^2 - q^2} (9m_\pi^2 + 9p^2 - l^2) - \frac{l^2}{2} \right. \\
& \left. \left. - \frac{5}{2} (5m_\pi^2 + p^2 + q^2) + \frac{5(4m_\pi^2 + q^2)}{2(B + q^2 l^2)} ((m_\pi^2 + p^2)^2 + l^2(m_\pi^2 - 3p^2 + q^2)) \right] \Omega(l) \right] \right\}. \quad (35)
\end{aligned}$$

5 $2\pi^0$ -exchange three-neutron force at N⁴LO

In this section the longest-range $2\pi^0$ -exchange 3n-interaction is treated, following the work of ref.[3]. Modulo terms of shorter range it can be written according to eq.(3.1) in ref.[3] in the general form:

$$2V_{3n} = \frac{g_A^2}{4f_\pi^4} \frac{\vec{\sigma}_1 \cdot \vec{q}_1 \vec{\sigma}_3 \cdot \vec{q}_3}{(m_\pi^2 + q_1^2)(m_\pi^2 + q_3^2)} \tilde{g}_+(q_2), \quad (36)$$

with both sides multiplied by a factor 2 due to the $1 \leftrightarrow 3$ symmetry. The structure function $\tilde{g}_+(q_2)$ is f_π^2 times the isoscalar non-spin-flip πN -scattering amplitude at zero pion-energy $\omega = 0$ and squared momentum-transfer $t = -q_2^2$. The corresponding expression up to N⁴LO is given in eq.(59) of ref.[8].

5.1 Contributions to in-medium nn-potential

The self-closing of neutron-line 2 for V_{3n} in eq.(36) gives (after relabeling $3 \rightarrow 2$) the contribution:

$$V_{\text{med}}^{(0)} = -\frac{g_A^2 k_n^3}{12\pi^2 f_\pi^4} \frac{\vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q}}{(m_\pi^2 + q^2)^2} \tilde{g}_+(0). \quad (37)$$

From pionic vertex corrections on either neutron-line one obtains the (total) contribution:

$$\begin{aligned} V_{\text{med}}^{(1)} = & \frac{g_A^2}{16\pi^2 f_\pi^4} \frac{\vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q}}{(m_\pi^2 + q^2)q^2} \int_{p-k_n}^{p+k_n} ds s \tilde{g}_+(s) \left[\frac{k_n^2 - (p-s)^2}{p} \right. \\ & \left. + \left(q - \frac{m_\pi^2 + s^2}{q} \right) \ln \frac{qX + 2\sqrt{W}}{(2p+q)[m_\pi^2 + (q-s)^2]} \right], \end{aligned} \quad (38)$$

with the auxiliary polynomials X and W defined in eq.(6). Finally, the two diagrams related to double exchange lead to the expression:

$$V_{\text{med}}^{(3)} = \frac{g_A^2 \tilde{g}_+(q)}{16\pi^2 f_\pi^4} \left[2\Gamma_0 - (2m_\pi^2 + q^2)G_0 + i(\vec{\sigma}_1 + \vec{\sigma}_2) \cdot (\vec{q} \times \vec{p})(G_0 + 2G_1) \right], \quad (39)$$

where one reminds that the functions Γ_0 depends on (p, k_n) , and the functions $G_{0,1}$ depend on (p, q, k_n) .

6 Subleading ring diagrams

6.1 3n-ring interaction $\sim g_A^0$

According to eq.(31) in ref.[9] the subleading 3n-ring interaction proportional to g_A^0 reads:

$$V_{3n} = -\frac{1}{f_\pi^6} \int_0^\infty dl_0 \int \frac{d^3 l_2}{(2\pi)^4} \frac{l_0^2}{(\bar{m}^2 + l_1^2)(\bar{m}^2 + l_2^2)(\bar{m}^2 + l_3^2)} \left[2c_1 m_\pi^2 + (c_2 + c_3)l_0^2 + c_3 \vec{l}_2 \cdot \vec{l}_3 \right], \quad (40)$$

with $\bar{m} = \sqrt{m_\pi^2 + l_0^2}$ and one has to set $\vec{l}_1 = \vec{l}_2 - \vec{q}_3$ and $\vec{l}_3 = \vec{l}_2 + \vec{q}_1$. The sum of the three self-closings of a neutron-line gives rise to the central potential:

$$\begin{aligned} V_{\text{med}}^{(0)} = & \frac{k_n^3}{32\pi^4 f_\pi^6} \left\{ \left[\left(2c_1 - \frac{3c_2}{2} - 3c_3 \right) m_\pi^2 - \left(\frac{c_2}{4} + \frac{5c_3}{9} \right) q^2 \right] \ln \frac{m_\pi}{\lambda} + \left(\frac{5c_2}{8} + \frac{37c_3}{27} \right) \frac{q^2}{4} \right. \\ & \left. + \left(\frac{3c_2}{2} - 2c_1 + \frac{23c_3}{9} \right) \frac{m_\pi^2}{4} + \left[\left(2c_1 - c_2 - \frac{17c_3}{9} \right) m_\pi^2 - \left(\frac{c_2}{4} + \frac{5c_3}{9} \right) q^2 \right] L(q) \right\}, \end{aligned} \quad (41)$$

with logarithmic loop-function

$$L(q) = \frac{\sqrt{4m_\pi^2 + q^2}}{q} \ln \frac{q + \sqrt{4m_\pi^2 + q^2}}{2m_\pi}. \quad (42)$$

Note that the four-dimensional euclidean loop-integral in eq.(40) has been regularized with a cutoff λ and the λ^2 -divergence has been dropped in the expression in eq.(41). The total contribution from the six possible concatenations of two neutron-lines is a central potential with the following double-integral representation:

$$V_{\text{med}}^{(\text{cc})} = \frac{1}{4\pi^5 f_\pi^6} \int_0^\lambda dr r \int_0^{\pi/2} d\psi \left\{ l_0^2 l \left\{ \bar{\Gamma}_0(l) \left[c_3 \bar{\Lambda}(l) + \left(6c_1 m_\pi^2 + 3(c_2 + c_3) l_0^2 - \frac{c_3}{2} (2\bar{m}^2 + q^2) \right) \bar{\Omega}(l) \right] \right. \right. \\ \left. \left. + c_3 \bar{\Gamma}_1(l) \left[\bar{\Lambda}(l) + (l^2 - \bar{m}^2 - p^2) \bar{\Omega}(l) \right] \right\} - \frac{k_n^3}{2} (c_3 + c_2 \cos^2 \psi) \sin^2 2\psi \right\}, \quad (43)$$

where one has to set $l_0 = r \cos \psi$ and $l = r \sin \psi$. Note that all barred functions are to be evaluated with $\bar{m} = \sqrt{m_\pi^2 + l_0^2}$ instead of m_π . The behavior of the subtracted double-integral for large λ is:

$$\frac{\pi k_n^3}{16} \left[c_2 \left(3m_\pi^2 + \frac{3k_n^2}{10} + \frac{p^2}{2} + \frac{q^2}{4} \right) - 4c_1 m_\pi^2 + c_3 \left(6m_\pi^2 + \frac{2k_n^2}{3} + \frac{5}{9} (2p^2 + q^2) \right) \right] \ln \frac{m_\pi}{\lambda}. \quad (44)$$

A good check is given by the fact that the λ^2 -divergences behind $V_{\text{med}}^{(0)}$ and $V_{\text{med}}^{(\text{cc})}$ cancel each other. This feature must hold, because there is no 3n-contact coupling that could absorb this divergence.

6.2 3n-ring interaction $\sim g_A^2$

According to eq.(44) in ref.[9] the subleading 3n-ring interaction proportional to g_A^2 reads:

$$V_{3n} = -\frac{g_A^2}{f_\pi^6} \int_0^\infty dl_0 \int \frac{d^3 l_2}{(2\pi)^4} \frac{1}{(\bar{m}^2 + l_1^2)(\bar{m}^2 + l_2^2)(\bar{m}^2 + l_3^2)} \left\{ [2c_1 m_\pi^2 + (c_2 + c_3) l_0^2 + c_3 \vec{l}_2 \cdot \vec{l}_3] \vec{l}_1 \cdot (\vec{l}_2 + \vec{l}_3) \right. \\ \left. + c_4 \left[\bar{m}^2 (\vec{\sigma}_2 \times \vec{l}_3) \cdot (\vec{\sigma}_3 \times \vec{l}_2) + \vec{l}_2 \cdot \vec{l}_3 \vec{\sigma}_2 \cdot \vec{l}_1 \vec{\sigma}_3 \cdot \vec{l}_1 + \vec{l}_1 \cdot \vec{l}_2 \vec{l}_1 \cdot \vec{l}_3 \vec{\sigma}_2 \cdot \vec{\sigma}_3 - 2\vec{l}_1 \cdot \vec{l}_3 \vec{\sigma}_2 \cdot \vec{l}_2 \vec{\sigma}_3 \cdot \vec{l}_1 \right] \right\}. \quad (45)$$

The sum of the three self-closings of a neutron-line gives rise to the following contributions to V_{med} .

- Central potential:

$$V_{\text{med}}^{(0)} = \frac{g_A^2 k_n^3}{96\pi^4 f_\pi^6} \left\{ \left[9m_\pi^2 (4c_1 - c_2 - 6c_3) - \left(\frac{11c_2}{6} + \frac{32c_3}{3} \right) q^2 \right] \ln \frac{m_\pi}{\lambda} \right. \\ \left. + \left(\frac{19c_2}{12} - 5c_1 + \frac{25c_3}{6} \right) m_\pi^2 + \left(\frac{137c_2}{16} + 58c_3 \right) \frac{q^2}{9} \right. \\ \left. + \left[2m_\pi^2 \left(18c_1 - \frac{8c_2}{3} - \frac{43c_3}{3} \right) - \left(\frac{11c_2}{6} + \frac{32c_3}{3} \right) q^2 + \frac{8(c_3 - 2c_1)m_\pi^4}{4m_\pi^2 + q^2} \right] L(q) \right\}, \quad (46)$$

- Spin-spin potential:

$$V_{\text{med}}^{(0)} = \frac{g_A^2 c_4 k_n^3}{96\pi^4 f_\pi^6} \vec{\sigma}_1 \cdot \vec{\sigma}_2 \left\{ - (4m_\pi^2 + q^2) \ln \frac{m_\pi}{\lambda} + \frac{q^2}{4} - m_\pi^2 - q^2 L(q) \right\}, \quad (47)$$

- Tensor potential:

$$V_{\text{med}}^{(0)} = \frac{g_A^2 c_4 k_n^3}{96\pi^4 f_\pi^6} \vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q} \left\{ \ln \frac{m_\pi}{\lambda} - \frac{1}{4} + L(q) \right\}. \quad (48)$$

Next, one evaluates the concatenations n_3 on n_2 and n_2 on n_3 and obtains the following contributions to V_{med} .

- Central potential:

$$\begin{aligned}
V_{\text{med}}^{(1)} = & \frac{g_A^2}{4\pi^5 f_\pi^6} \int_0^\lambda dr r \int_0^{\pi/2} d\psi \left\{ l \tilde{\Gamma}_1(l) \left\{ c_3 l \left(1 - \frac{q^2}{4p^2} \right) + \left[2c_1 m_\pi^2 + c_2 l_0^2 + c_3 \left(r^2 - 2\bar{m}^2 - p^2 \right. \right. \right. \right. \\
& + \left. \left. \left. \frac{q^2}{4p^2} (l^2 + \bar{m}^2 - p^2) \right) \right] \bar{\Lambda}(l) + \left[2c_1 m_\pi^2 + c_2 l_0^2 - c_3 \left(m_\pi^2 + \frac{q^2}{2} \right) \right] (l^2 - \bar{m}^2 - p^2) \bar{\Omega}(l) \right\} \\
& + c_4 l \left\{ [\bar{m}^2 \bar{\Gamma}_0(l) + 2\bar{\Gamma}_2(l)] [2\bar{\Lambda}(l) - (2\bar{m}^2 + q^2) \bar{\Omega}(l)] + \frac{1}{4} \tilde{\Gamma}_3(l) [2(3l^2 - \bar{m}^2 - p^2) \bar{\Lambda}(l) \right. \\
& + l + (\bar{B} - 2l^2(4\bar{m}^2 + q^2)) \bar{\Omega}(l)] \left. \right\} - \frac{4k_n^3}{3} (c_3 + c_4 + c_2 \cos^2 \psi) \sin^4 \psi \left. \right\}, \quad (49)
\end{aligned}$$

where the (subtracted) double-integral has the large- λ behavior:

$$\frac{\pi k_n^3}{12} \left[\frac{c_2}{4} \left(6m_\pi^2 + k_n^2 + \frac{5p^2}{3} + \frac{q^2}{6} \right) - 6c_1 m_\pi^2 + c_3 \left(9m_\pi^2 + k_n^2 + \frac{5p^2}{3} + q^2 \right) + c_4 \left(6m_\pi^2 + \frac{3k_n^2}{5} + p^2 + \frac{7q^2}{6} \right) \right] \ln \frac{m_\pi}{\lambda}. \quad (50)$$

- Spin-orbit potential:

$$\begin{aligned}
V_{\text{med}}^{(1)} = & \frac{c_4 g_A^2}{8\pi^5 f_\pi^6} i(\vec{\sigma}_1 + \vec{\sigma}_2) \cdot (\vec{q} \times \vec{p}) \int_0^\lambda dr r \int_0^{\pi/2} d\psi \frac{l}{4p^2 - q^2} \left\{ [\bar{m}^2 \bar{\Gamma}_0(l) + 2\bar{\Gamma}_2(l)] [2\bar{\Lambda}(l) \right. \\
& + (2p^2 - 2l^2 - 2\bar{m}^2 - q^2) \bar{\Omega}(l)] + \tilde{\Gamma}_3(l) (\bar{m}^2 + p^2 - l^2) [(\bar{m}^2 + l^2 + p^2) \Omega(l) - \Lambda(l)] \left. \right\}, \quad (51)
\end{aligned}$$

with a large- λ behavior of the double-integral: $(\pi k_n^3/24) \ln(m_\pi/\lambda)$.

The other four concatenations (n_3 on n_1 , n_1 on n_3 , n_1 on n_2 , n_2 on n_1) lead to the following contributions to V_{med} .

- Central potential:

$$\begin{aligned}
V_{\text{med}}^{(\text{cc})} = & \frac{g_A^2}{4\pi^5 f_\pi^6} \int_0^\lambda dr r \int_0^{\pi/2} d\psi \left\{ l [2c_1 m_\pi^2 + (c_2 + c_3) l_0^2] \left\{ \bar{\Gamma}_0(l) [2\bar{\Lambda}(l) - (2\bar{m}^2 + q^2) \bar{\Omega}(l)] \right. \right. \\
& + \tilde{\Gamma}_1(l) [\Lambda(l) + (l^2 - p^2 - \bar{m}^2) \Omega(l)] \left. \right\} + c_3 l \left\{ \bar{\Gamma}_2(l) [2\bar{\Lambda}(l) - (2\bar{m}^2 + q^2) \bar{\Omega}(l)] \right. \\
& + \tilde{\Gamma}_3(l) \left[\frac{l}{2} + (l^2 - p^2 - \bar{m}^2) \bar{\Lambda}(l) + \frac{1}{2} (l^2 - p^2 - \bar{m}^2)^2 \bar{\Omega}(l) \right] + \tilde{\Gamma}_1(l) \left[l \left(1 - \frac{q^2}{4p^2} \right) \right. \\
& + \left(l^2 - p^2 - 2\bar{m}^2 - \frac{q^2}{4} + \frac{q^2}{4p^2} (l^2 + \bar{m}^2) \right) \bar{\Lambda}(l) + \left(\bar{m}^2 + \frac{q^2}{2} \right) (\bar{m}^2 + p^2 - l^2) \bar{\Omega}(l) \left. \right] \left. \right\} \\
& - \frac{8k_n^3}{3} (c_3 + c_2 \cos^2 \psi) \sin^4 \psi \left. \right\}, \quad (52)
\end{aligned}$$

with a large- λ behavior of the (subtracted) double-integral:

$$\frac{\pi k_n^3}{4} \left[c_2 \left(m_\pi^2 + \frac{k_n^2}{10} + \frac{p^2}{6} + \frac{5q^2}{36} \right) - 4c_1 m_\pi^2 + c_3 \left(6m_\pi^2 + \frac{11k_n^2}{15} + \frac{11p^2 + 5q^2}{9} \right) \right] \ln \frac{m_\pi}{\lambda}. \quad (53)$$

- Spin-spin potential:

$$\begin{aligned}
V_{\text{med}}^{(\text{cc})} = & \frac{c_4 g_A^2}{16\pi^5 f_\pi^6} \vec{\sigma}_1 \cdot \vec{\sigma}_2 \int_0^\lambda dr r \int_0^{\pi/2} d\psi \left\{ l \tilde{\Gamma}_1(l) \left[\frac{l}{p^2} (3p^2 - l^2 - \bar{m}^2 - q^2) \right. \right. \\
& + \left(2l^2 - 2\bar{m}^2 - 3p^2 + (\bar{m}^2 + l^2 + q^2) \frac{\bar{m}^2 + l^2}{p^2} + \frac{q^2(4\bar{m}^2 + 4l^2 + q^2)}{4p^2 - q^2} \right) \bar{\Lambda}(l) \\
& \left. \left. + \frac{2q^2(\bar{m}^2 + l^2 + p^2)}{4p^2 - q^2} (2p^2 - 2l^2 - 2\bar{m}^2 - q^2) \bar{\Omega}(l) \right] - \frac{32k_n^3}{9} \sin^4 \psi \right\}, \quad (54)
\end{aligned}$$

with a large- λ behavior of the (subtracted) double-integral: $(2\pi k_n^3/9)[6m_\pi^2 + k_n^2 + 2p^2 + 3q^2/4] \ln(m_\pi/\lambda)$.

- Ordinary tensor potential:

$$\begin{aligned}
V_{\text{med}}^{(\text{cc})} = & \frac{c_4 g_A^2}{4\pi^5 f_\pi^6} \vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q} \int_0^\lambda dr r \int_0^{\pi/2} d\psi \frac{l \tilde{\Gamma}_1(l)}{4p^2 - q^2} \left\{ l \left(1 - \frac{q^2}{2p^2} \right) \right. \\
& + \left[\frac{q^2}{4p^2 - q^2} \left(2\bar{m}^2 + 2l^2 + \frac{q^2}{2} \right) + \frac{\bar{m}^2 + l^2}{2p^2} (q^2 - 4p^2) \right] \bar{\Lambda}(l) \\
& \left. + \left[3\bar{B} + 2l^2(5p^2 + q^2) + (\bar{m}^2 + p^2)(q^2 - 2p^2) - \frac{8p^2}{4p^2 - q^2} (\bar{m}^2 + l^2 + p^2)^2 \right] \bar{\Omega}(l) \right\}, \quad (55)
\end{aligned}$$

with a large- λ behavior of the double-integral: $-(\pi k_n^3/24) \ln(m_\pi/\lambda)$.

- Tensor-type potential:

$$\begin{aligned}
V_{\text{med}}^{(\text{cc})} = & \frac{c_4 g_A^2}{4\pi^5 f_\pi^6} (\vec{\sigma}_1 \cdot \vec{p} \vec{\sigma}_2 \cdot \vec{p} + \vec{\sigma}_1 \cdot \vec{p}' \vec{\sigma}_2 \cdot \vec{p}') \int_0^\lambda dr r \int_0^{\pi/2} d\psi \frac{l \tilde{\Gamma}_1(l)}{4p^2 - q^2} \left\{ \frac{l}{2p^2} \left[3\bar{m}^2 + 3l^2 - p^2 + \frac{5q^2}{4} \right. \right. \\
& - \left. \frac{3q^2}{4p^2} (\bar{m}^2 + l^2) \right] + \left[\frac{p^2}{2} + l^2 - \bar{m}^2 - \frac{5q^2}{8} - \frac{6(\bar{m}^2 + l^2)^2 + q^2(\bar{m}^2 + 3l^2)}{4p^2} + \frac{3q^2}{8p^4} (\bar{m}^2 + l^2)^2 \right. \\
& \left. \left. - \frac{q^2(4\bar{m}^2 + 4l^2 + q^2)}{4p^2 - q^2} \right] \bar{\Lambda}(l) + \left[\frac{4(\bar{m}^2 + l^2 + p^2)^2}{4p^2 - q^2} - \bar{m}^2 - 3l^2 - p^2 \right] q^2 \bar{\Omega}(l) \right\}, \quad (56)
\end{aligned}$$

with a large- λ behavior of the double-integral: $-(\pi k_n^3/36) \ln(m_\pi/\lambda)$. The total λ^2 -divergence behind the calculated central and spin-spin potentials adds up to $c_4 g_A^2 k_n^3 \lambda^2 / (96\pi^4 f_\pi^6) [3 + \vec{\sigma}_1 \cdot \vec{\sigma}_2]$. As required by the renormalizability condition this is equivalent to zero, since $\vec{\sigma}_1 \cdot \vec{\sigma}_2$ has the eigenvalue -3 in the 1S_0 -state of two neutrons.

6.3 3n-ring interaction $\sim g_A^4$

In the case of the subleading 3n-ring interaction terms proportional to g_A^4 , as obtained by setting $\vec{\tau}_i \cdot \vec{\tau}_j \rightarrow 1$ and $\vec{\tau}_1 \cdot (\vec{\tau}_2 \times \vec{\tau}_3) \rightarrow 0$ in the very lengthy expression in eq.(58) of ref.[9], only the results for the self-closing contributions are given here.

- Central potential:

$$\begin{aligned}
V_{\text{med}}^{(0)} = & \frac{g_A^4 k_n^3}{96\pi^4 f_\pi^6} \left\{ \left[45m_\pi^2 \left(\frac{11c_3 - 3c_2}{4} - 2c_1 \right) + (177c_3 - 53c_2) \frac{q^2}{8} \right] \ln \frac{m_\pi}{\lambda} - \frac{4m_\pi^4 (2c_1 + 3c_3)}{4m_\pi^2 + q^2} \right. \\
& + (705c_3 - 73c_2 - 528c_1) \frac{m_\pi^2}{32} + \frac{7q^2}{64} \left(\frac{53c_2}{3} - 35c_3 \right) + \left[m_\pi^2 \left(\frac{81c_3 - 35c_2}{2} - 82c_1 \right) \right. \\
& \left. \left. + (177c_3 - 53c_2) \frac{q^2}{8} + \frac{2m_\pi^4}{4m_\pi^2 + q^2} (40c_1 + 3c_2 + 21c_3) - \frac{16m_\pi^6 (2c_1 + 3c_3)}{(4m_\pi^2 + q^2)^2} \right] L(q) \right\}, \quad (57)
\end{aligned}$$

- Spin-spin potential:

$$\begin{aligned}
V_{\text{med}}^{(0)} = & \frac{g_A^4 k_n^3}{96\pi^4 f_\pi^6} \vec{\sigma}_1 \cdot \vec{\sigma}_2 \left\{ \left[6m_\pi^2 \left(6c_1 + c_2 - \frac{13c_3}{2} + 5c_4 \right) + \left(c_2 - \frac{9c_3}{2} + \frac{11c_4}{3} \right) q^2 \right] \ln \frac{m_\pi}{\lambda} \right. \\
& + \left(9c_1 + \frac{c_2}{2} - \frac{33c_3}{4} + \frac{35c_4}{6} \right) m_\pi^2 - \left(\frac{c_2}{6} + \frac{c_4}{9} \right) q^2 + \left[\left(24c_1 + 2c_2 - 11c_3 + \frac{26c_4}{3} \right) m_\pi^2 \right. \\
& \left. \left. + \left(c_2 - \frac{9c_3}{2} + \frac{11c_4}{3} \right) q^2 + \frac{4m_\pi^4}{4m_\pi^2 + q^2} (3c_3 - 6c_1 - 2c_4) \right] L(q) \right\}, \quad (58)
\end{aligned}$$

- Tensor potential:

$$\begin{aligned}
V_{\text{med}}^{(0)} = & \frac{g_A^4 k_n^3}{576\pi^4 f_\pi^6} \vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q} \left\{ \left(\frac{53c_3}{2} - 5c_2 - 21c_4 \right) \ln \frac{m_\pi}{\lambda} + \frac{367c_3}{48} - \frac{131c_2}{120} - \frac{233c_4}{240} \right. \\
& + \frac{4m_\pi^2}{q^2} (18c_1 + 2c_2 - 7c_3 + 6c_4) + \frac{3m_\pi^2}{4m_\pi^2 + q^2} (3c_3 - 6c_1 - 2c_4) \\
& + \left[\frac{53c_3}{2} - 5c_2 - 21c_4 + \frac{4m_\pi^2}{q^2} (7c_3 - 18c_1 - 2c_2 - 6c_4) \right. \\
& \left. \left. + \frac{3m_\pi^2}{4m_\pi^2 + q^2} (2c_2 - 13c_3 + 10c_4) + \frac{12m_\pi^4}{(4m_\pi^2 + q^2)^2} (3c_3 - 6c_1 - 2c_4) \right] L(q) \right\}. \quad (59)
\end{aligned}$$

7 Subleading three-nucleon contact potential

The subleading three-nucleon contact potential (appearing at N⁴LO in the chiral counting) has been reexamined recently in ref.[13]. Its corrected expression, given in eq.(15) of ref.[13], depends quadratically on momenta and it involves 13 parameters, called E_1, \dots, E_{13} . Specifying to the case of three neutrons and closing two neutron-lines in all possible ways to a loop, one obtains the following contribution to the in-medium nn-potential:

$$\begin{aligned}
\frac{V_{\text{med}}}{\rho_n} = & 2(E_1 + E_2) \left(\frac{6k_n^2}{5} + 2p^2 - q^2 \right) + 2(E_3 + E_4) (\vec{\sigma}_1 \cdot \vec{\sigma}_2 + 3) \left(\frac{3k_n^2}{5} + p^2 \right) \\
& + 3(E_5 + E_6) \left[\vec{\sigma}_1 \cdot \vec{p} \vec{\sigma}_2 \cdot \vec{p} + \vec{\sigma}_1 \cdot \vec{p}' \vec{\sigma}_2 \cdot \vec{p}' - \frac{2p^2}{3} \vec{\sigma}_1 \cdot \vec{\sigma}_2 \right] + (E_7 + E_8) i (\vec{\sigma}_1 + \vec{\sigma}_2) \cdot (\vec{q} \times \vec{p}) \\
& + (E_9 + E_{10}) \left[q^2 - 2p^2 - \frac{6k_n^2}{5} - i (\vec{\sigma}_1 + \vec{\sigma}_2) \cdot (\vec{q} \times \vec{p}) \right] \\
& + (E_{11} + E_{12} + E_{13}) \left[q^2 - 2p^2 - \frac{6k_n^2}{5} + i (\vec{\sigma}_1 + \vec{\sigma}_2) \cdot (\vec{q} \times \vec{p}) \right], \quad (60)
\end{aligned}$$

which depends linearly on the neutron density $\rho_n = k_n^3/3\pi^2$ and quadratically on the momenta \vec{p}, \vec{q}, k_n . The above expression for V_{med} produces non-vanishing matrix-elements only in S- and P-wave states:

$$\begin{aligned}
\langle {}^1S_0 | V_{\text{med}} | {}^1S_0 \rangle &= \frac{6\rho_n k_n^2}{5} (2E_1 + 2E_2 - E_9 - E_{10} - E_{11} - E_{12} - E_{13}) = \frac{9\rho_n k_n^2}{5} E_c, \\
\langle {}^3P_0 | V_{\text{med}} | {}^3P_0 \rangle &= \frac{2}{3} \rho_n p^2 (2E_1 + 2E_2 + 2E_7 + 2E_8 - 3E_9 - 3E_{10} + E_{11} + E_{12} + E_{13}) = \rho_n p^2 (E_c - 2E_o), \\
\langle {}^3P_1 | V_{\text{med}} | {}^3P_1 \rangle &= \frac{2}{3} \rho_n p^2 (2E_1 + 2E_2 + E_7 + E_8 - 2E_9 - 2E_{10}) = \rho_n p^2 (E_c - E_o), \\
\langle {}^3P_2 | V_{\text{med}} | {}^3P_2 \rangle &= \frac{2}{3} \rho_n p^2 (2E_1 + 2E_2 - E_7 - E_8 - 2E_{11} - 2E_{12} - 2E_{13}) = \rho_n p^2 (E_c + E_o), \quad (61)
\end{aligned}$$

where the parts $\sim (E_3 + E_4)$ and $\sim (E_5 + E_6)$ in eq.(60) are effectively zero and only a central parameter E_c and a spin-orbit parameter E_o are independent. As a remarkable feature one observes that there is no term proportional to $\rho_n p^2$ in the S -wave matrix element $\langle {}^1S_0 | V_{\text{med}} | {}^1S_0 \rangle$. In order to establish that this restriction is consistent with the renormalizability condition, one exploits $\vec{\sigma}_1 = -\vec{\sigma}_2$ in the spin-singlet state and uses the angular average $\langle q^2 \rangle = 2p^2$ of the squared momentum-transfer. By following in subsections 6.1 and 6.2 all terms that contribute proportional to $c_{2,3} k_n^3 p^2 / (64\pi^4 f_\pi^6) \ln(m_\pi/\lambda)$ or $c_{2,3,4} g_A^2 k_n^3 p^2 / (96\pi^4 f_\pi^6) \ln(m_\pi/\lambda)$ to the S -wave matrix element, one verifies that these indeed sum up to zero. The same exact cancelation occurs also for the terms proportional to $c_{1,2,3} k_n^3 m_\pi^2 / (64\pi^4 f_\pi^6) \ln(m_\pi/\lambda)$ or $c_{1,2,3,4} g_A^2 k_n^3 m_\pi^2 / (96\pi^4 f_\pi^6) \ln(m_\pi/\lambda)$.

From the corrected version of the subleading 3N-contact potential given in eq.(15) of ref.[13] the following in-medium NN-potential in isospin-symmetric nuclear matter of density $\rho = 2k_f^3/3\pi^2$ is derived:

$$\begin{aligned} \frac{V_{\text{med}}}{\rho} = & E_1 \left(\frac{6}{5} k_f^2 + 2p^2 - 3q^2 \right) + E_2 \left[(\vec{\tau}_1 \cdot \vec{\tau}_2 + 3) \left(\frac{3}{5} k_f^2 + p^2 \right) - \vec{\tau}_1 \cdot \vec{\tau}_2 q^2 \right] \\ & + E_3 \left[(\vec{\sigma}_1 \cdot \vec{\sigma}_2 + 3) \left(\frac{3}{5} k_f^2 + p^2 \right) - \vec{\sigma}_1 \cdot \vec{\sigma}_2 q^2 \right] + E_4 \left[(\vec{\tau}_1 \cdot \vec{\tau}_2 \vec{\sigma}_1 \cdot \vec{\sigma}_2 + 9) \left(\frac{3}{5} k_f^2 + p^2 \right) - \vec{\tau}_1 \cdot \vec{\tau}_2 \vec{\sigma}_1 \cdot \vec{\sigma}_2 q^2 \right] \\ & + (E_5 + \vec{\tau}_1 \cdot \vec{\tau}_2 E_6) \left[\vec{\sigma}_1 \cdot \vec{\sigma}_2 (q^2 - p^2) - 3\vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q} + \frac{3}{2} (\vec{\sigma}_1 \cdot \vec{p} \vec{\sigma}_2 \cdot \vec{p} + \vec{\sigma}_1 \cdot \vec{p}' \vec{\sigma}_2 \cdot \vec{p}') \right] \\ & + (7E_7 - 9E_8) \frac{i}{4} (\vec{\sigma}_1 + \vec{\sigma}_2) \cdot (\vec{q} \times \vec{p}) + E_9 \left[\vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q} + \frac{q^2}{2} - p^2 - \frac{3}{5} k_f^2 - \frac{i}{2} (\vec{\sigma}_1 + \vec{\sigma}_2) \cdot (\vec{q} \times \vec{p}) \right] \\ & + E_{10} \left[\vec{\tau}_1 \cdot \vec{\tau}_2 \vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q} + \frac{3}{2} q^2 - 3p^2 - \frac{9}{5} k_f^2 - \frac{3i}{2} (\vec{\sigma}_1 + \vec{\sigma}_2) \cdot (\vec{q} \times \vec{p}) \right] \\ & + E_{11} \left[\vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q} + \frac{q^2}{2} - p^2 - \frac{3}{5} k_f^2 + \frac{i}{2} (\vec{\sigma}_1 + \vec{\sigma}_2) \cdot (\vec{q} \times \vec{p}) \right] \\ & + E_{12} \left[\vec{\tau}_1 \cdot \vec{\tau}_2 \vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q} + \frac{3}{2} q^2 - 3p^2 - \frac{9}{5} k_f^2 + \frac{3i}{2} (\vec{\sigma}_1 + \vec{\sigma}_2) \cdot (\vec{q} \times \vec{p}) \right] \\ & + \frac{E_{13}}{2} \left\{ \vec{\tau}_1 \cdot \vec{\tau}_2 \left[q^2 - 2p^2 - \frac{6}{5} k_f^2 - \vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q} + i(\vec{\sigma}_1 + \vec{\sigma}_2) \cdot (\vec{q} \times \vec{p}) \right] - 3\vec{\sigma}_1 \cdot \vec{q} \vec{\sigma}_2 \cdot \vec{q} \right\}, \end{aligned} \quad (62)$$

which should replace the (incomplete) expression written in eq.(49) of ref.[7].

References

- [1] V. Bernard, E. Epelbaum, H. Krebs, and Ulf-G. Meißner, *Phys. Rev.* **C77**, 064004 (2008).
- [2] V. Bernard, E. Epelbaum, H. Krebs, and Ulf-G. Meißner, *Phys. Rev.* **C84**, 054001 (2011).
- [3] H. Krebs, A. Gasparyan, and E. Epelbaum, *Phys. Rev.* **C85**, 054006 (2012).
- [4] H. Krebs, A. Gasparyan, and E. Epelbaum, *Phys. Rev.* **C87**, 054007 (2013).
- [5] H. Krebs, A.M. Gasparyan, and E. Epelbaum, *Phys. Rev.* **C98**, 014003 (2018).
- [6] J.W. Holt, N. Kaiser, and W. Weise, *Phys. Rev.* **C81**, 024002 (2010).
- [7] N. Kaiser and V. Niessner, *Phys. Rev.* **C98**, 054002 (2018).
- [8] N. Kaiser and B. Singh, *Phys. Rev.* **C100**, 014002 (2019).
- [9] N. Kaiser, *Phys. Rev.* **C101**, 014001 (2020).
- [10] C. Drischler, K. Hebeler, and A. Schwenk, *Phys. Rev.* **C93**, 054314 (2016); *Phys. Rev. Lett.* **122**, 042501 (2019); and references therein.
- [11] C. Drischler, A. Carbone, K. Hebeler, and A. Schwenk, *Phys. Rev.* **C94**, 054307 (2016).
- [12] L. Treuer, Bachelor thesis, TU Munich, nucl-th-2009.11104.
- [13] L. Girlanda, A. Kievsky, and M. Viviani, *Phys. Rev.* **C102**, 019903(E) (2020).