

# Entropy gain in $p$ -adic quantum channels

Evgeny Zelenov

October 3, 2020

## Abstract

The construction of the  $p$ -adic quantum Gaussian channel is suggested. Entropy gain is calculated. The adelic formula for the entropy gain is obtained.

## 1 Introduction

Starting from [1, 2], non-Archimedean analysis has been actively used to build physical models. Thereby, the branch of mathematical physics ( $p$ -adic mathematical physics) has arisen. One can find a bibliography on this subject in the book [3] and reviews [4, 5].

The article is organized as follows. Firstly, we give the necessary facts of the  $p$ -adic analysis. Secondly, we define the  $p$ -adic Gaussian state and the  $p$ -adic Gaussian linear bosonic channel and also give their properties. This information follows [6]; the results are presented without proof. New results are shown below. Thirdly, we prove a formula for the magnitude of the entropy gain in the  $p$ -adic Gaussian channel. Fourthly, we give the adelic formula for the entropy gain and its possible applications.

## 2 $p$ -Adic numbers and symplectic geometry

In this section, some known facts concerning the geometry of lattices in a two-dimensional symplectic space over the field  $\mathbb{Q}_p$  of  $p$ -adic numbers are presented without proof. The necessary information from the  $p$ -adic analysis

is contained, for example, in [7]. Most statements regarding the geometry of lattices can be found in [8].

Let  $F$  be a two-dimensional vector space over the field  $\mathbb{Q}_p$ . A nondegenerate symplectic form  $\Delta : F \times F \rightarrow \mathbb{Q}_p$  is given in the space  $F$ . A free module of rank two over the ring  $\mathbb{Z}_p$  of  $p$ -adic integers considered as a subset of  $F$  will be called a lattice in  $F$ . A lattice is a compact set in the natural topology in the space  $F$ . We introduce the duality relation on the set of lattices. Let  $L$  be a lattice; then the dual lattice  $L^*$  means the following subset of the space  $F$ :  $u \in L^*$  if and only if the condition  $\Delta(u, v) \in \mathbb{Z}_p$  is satisfied for all  $v \in L$ . If  $L$  coincides with  $L^*$ , then the lattice  $L$  will be called a self-dual lattice. For any self-dual lattice  $L$ , there exists a symplectic basis  $(e_1, e_2)$  in the space  $F$  such that the lattice  $L$  has the form

$$L = \mathbb{Z}_p e_1 \oplus \mathbb{Z}_p e_2.$$

By  $Sp(F, \Delta)$  denote the symplectic group (the group of nondegenerate linear transformations of the space  $F$  that preserve the form  $\Delta$ ). The group  $Sp(F, \Delta)$  is isomorphic to the group  $SL_2(\mathbb{Q}_p)$ . In the space  $F$ , there exists a unique translation-invariant measure (Haar measure) up to normalization. We will normalize the measure in such a way that the measure of a self-dual lattice is equal to unity. The action of the symplectic group preserves the measure; therefore, the measure of any self-dual lattice is equal to unity. The measure of the lattice  $L$  will be denoted by  $|L|$ . If  $L$  is a self-dual lattice, then, as noted above,  $|L| = 1$ ; the converse is also true, if  $|L| = 1$ , then the lattice  $L$  is self-dual. It is easy to verify the validity of the relation  $|L||L^*| = 1$ . Let  $S \in Sp(F, \Delta)$ ,  $L \subset F$  be an arbitrary lattice. As already noted, the action of the symplectic group preserves measure, that is,  $|SL| = |L|$ . The converse is also true, if  $L_1, L_2$  are arbitrary lattices in  $F$  with the same measure,  $|L_1| = |L_2|$ , then there exists a symplectic transformation  $S \in Sp(F, \Delta)$ , such that  $SL_1 = L_2$ .

### 3 $p$ -Adic linear bosonic channels

Let  $\mathcal{H}$  be a separable Hilbert space over the field  $\mathbb{C}$  of complex numbers. The scalar product in  $\mathcal{H}$  will be denoted by  $\langle \cdot, \cdot \rangle$ , and we will consider it antilinear in the first argument. The state of the system is described by the density matrix  $\rho$  in the space  $\mathcal{H}$ . Denote by  $\mathfrak{S}$  the set of all states. By  $\mathcal{B}(F)$  we denote the  $\sigma$ -algebra of Borel subsets of  $F$ ; by  $\mathfrak{B}(\mathcal{H})$  we denote the algebra

of bounded operators on space  $\mathcal{H}$ . A quantum observable  $M : \mathcal{B}(F) \rightarrow \mathfrak{B}(\mathcal{H})$  is a projection-valued measure on  $\mathcal{B}(F)$ . Denote by  $\mathfrak{M}$  the set of quantum observables. The Born–von Neumann formula gives the probability distribution of the observable  $M$  in a state  $\rho$

$$\mu_\rho^M = \text{Tr}(\rho M(B)), B \in \mathcal{B}(F).$$

We do not go beyond the framework of the standard statistical model of quantum mechanics ([9]), since the set of real numbers and the set of  $p$ -adic numbers are Borel isomorphic.

Let  $W$  be a mapping from the space  $F$  to the set of unitary operators in the space  $\mathcal{H}$ , satisfying the relation

$$W(z)W(z') = \chi\left(\frac{1}{2}\Delta(z, z')\right)W(z + z')$$

for all  $z, z' \in F$ . Here we use the notation  $\chi(x) = \exp(2\pi i\{x\}_p)$ , where  $\{x\}_p$  denotes the  $p$ -adic fractional part of the number of  $x \in \mathbb{Q}_p$ . Besides, we assume that mapping  $W$  is continuous in the strong operator topology. The mapping  $W$  is called the representation of canonical commutation relations (CCRs) in the Weyl form. As noted above, the state of a quantum system is described by a density matrix  $\rho$  in a Hilbert space  $\mathcal{H}$ . Let an irreducible representation of the CCR  $\{W(z), z \in F\}$  be given in this space. Each density operator can be associated with a function  $\pi_\rho$  in the space  $F$  by the following formula.

$$\pi_\rho(z) = \text{Tr}(\rho W(z)).$$

The function  $\pi_\rho$  is called the characteristic function of the quantum state  $\rho$  and defines this state uniquely. The characteristic function is used to reconstruct the state  $\rho$  according to the relation

$$\rho = \int_F \pi_\rho(z)W(-z)dz.$$

The characteristic function of a quantum state has the property of  $\Delta$ -positive definiteness: for any finite sets  $z_1, z_2, \dots, z_n$  of points in the space  $F$  and complex numbers  $c_1, c_2, \dots, c_n$ , the inequality

$$\sum_{i,j=1}^n c_i \bar{c}_j \pi_\rho(z_i - z_j) \chi\left(-\frac{1}{2}\Delta(z_i, z_j)\right) \geq 0$$

is satisfied. As in the case of the representation of CCR over a real symplectic space, a noncommutative analogue of the Bochner–Khinchin theorem holds for the  $p$ -adic case; this theorem establishes one-to-one correspondence between states and  $\Delta$ -positive definite functions ([10]). We give the following definition.

**Definition 1** *A state  $\rho$  will be called a  $p$ -adic Gaussian state if its characteristic function is the indicator function of some lattice, that is,*

$$\pi_\rho(z) = \text{Tr}(\rho W(z)) = h_L(z) = \begin{cases} 1, & z \in L; \\ 0, & z \notin L. \end{cases}$$

This definition is natural in the following context. Let  $\mathcal{F}$  be the Fourier transform in  $L^2(F)$  defined by the formula

$$\mathcal{F}[f](z) = \int_F \chi(\Delta(z, s)) f(s) ds,$$

$L$  is a lattice in  $F$ . Then, the formula

$$|L|^{-1/2} \mathcal{F}[h_L] = |L^*|^{-1/2} h_{L^*}$$

is satisfied. In other words, the Fourier transform turns the characteristic function of the lattice into the characteristic function of the dual lattice up to a factor. In particular, the characteristic function of the self-dual lattice is invariant under the action of the Fourier transform. In this context, the characteristic function of the lattice is an analogue of the Gaussian function in real analysis.

$p$ -Adic Gaussian states are elementary in structure. Namely, the following statements are true.

**Proposition 1** *The characteristic function  $h_L$  of the lattice  $L$  determines the quantum state, if and only if the condition  $|L| \leq 1$  is satisfied.*

*A Gaussian state  $\rho$  having a characteristic function  $\pi_\rho = h_L$  is  $|L|P_L$ , where  $P_L$  is an orthogonal projector of rank  $1/|L|$ .*

Some obvious properties of Gaussian states are given below.

**Proposition 2** *The following statements are true.*

- *A Gaussian state is pure if and only if the corresponding lattice is self-dual.*
- *The entropy of the Gaussian state defined by the lattice  $L$  is  $-\log |L|$ .*
- *Gaussian states  $\rho_1$  and  $\rho_2$  are unitarily equivalent if and only if the corresponding lattices  $L_1$  and  $L_2$  have the same measure.*
- *The entropy of the Gaussian state determines this state uniquely up to unitary equivalence.*
- *The Gaussian state has the maximum entropy among all states with a fixed rank  $p^m$ ,  $m \in \mathbb{N}$ .*

We will use the notation  $\gamma(L)$  for the density operator of the Gaussian state defined by the lattice  $L$ . We considered only centred Gaussian states. We can similarly view general Gaussian states  $\gamma(L, \alpha)$ , which are defined by a characteristic function of the form

$$\pi_{\gamma(L, \alpha)} = \chi(\Delta(\alpha, z)) h_L(z).$$

It is easy to see that

$$\gamma(L, \alpha) = W(\alpha)\gamma(L)W(-\alpha).$$

Let  $W$  be the irreducible representation of the CCR in the Hilbert space  $\mathcal{H}$ ;  $\mathfrak{S}(\mathcal{H})$  is the set of states.

By analogy with the real case ([11]), a linear bosonic channel (in the Schrodinger representation) is a linear completely positive trace-preserving mapping  $\Phi : \mathfrak{S}(\mathcal{H}) \rightarrow \mathfrak{S}(\mathcal{H})$  such that the characteristic function  $\pi_\rho$  of any state  $\rho \in \mathfrak{S}(\mathcal{H})$  is transformed by the formula

$$\pi_{\Phi[\rho]}(z) = \pi_\rho(Kz)k(z) \tag{1}$$

for some linear transformation  $K$  of the space  $F$  and some complex-valued function  $k$  in  $F$ .

Generally speaking, expression (1) does not always determine the channel; to do this, additional conditions on the transformation  $K$  and the function  $k$  are necessary.

A  $p$ -adic Gaussian channel is a linear bosonic channel for which the function  $k$  is the characteristic function of some lattice  $L \subset F$ , that is,  $k(z) = h_L(z), z \in F$ .

The following proposition holds.

**Proposition 3** *Let  $K$  be a nondegenerate linear transformation of the space  $F$ ,  $L$  be a lattice in the space  $F$ ,  $k(z) = h_L(z)$ . In this case, expression (1) defines a channel if and only if the inequality*

$$|1 - \det K|_p |L| \leq 1 \quad (2)$$

is satisfied. Note that in the case of  $\det K = 1$ , the transformation is symplectic and, therefore, is unitarily representable. Next, we consider the case of  $\det K \neq 1$ .

## 4 Entropy gain

The entropy  $H(\rho)$  of the state  $\rho$  is defined by the following expression

$$H(\rho) = -\text{Tr } \rho \log \rho.$$

According to the definition, the entropy gain is

$$G(\Phi) = \inf_{\rho} \{H(\Phi[\rho]) - H(\rho)\}.$$

**Theorem 1** *The following formula is valid for the entropy gain of the  $p$ -adic Gaussian linear bosonic channel*

$$G(\Phi) = \log |\det K|_p.$$

Let  $K$  be a nondegenerate linear transformation and  $L$  be a lattice. Moreover, we assume that  $K$  and  $L$  define the channel, that is, they satisfy condition (2). By  $L_n$ ,  $n \in \mathbb{N}$  we denote the lattice  $L_n = p^n L$ . Note that  $L_n \subset L$ , and for sufficiently large  $n$ , the condition  $|L_n| < 1$  is satisfied, since  $|L_n| = p^{-n} |L|$ . According to Proposition 1, the Gaussian state  $\gamma(L_n)$  is defined for all such  $n$ . We also note that there exists such  $N \in \mathbb{N}$ , that for all  $n \geq N$ , the conditions  $L_n \subset L$  and  $K^{-1}L_n \subset L$  are simultaneously satisfied. It is not difficult to calculate the entropy gain for state  $\gamma(L_n)$ ,  $n \geq N$ . Indeed, the condition

$$\pi_{\Phi[\gamma(L_n)]}(z) = \pi_{\gamma(L_n)}(Kz)h_L(z) = h_{L_n}(Kz)h_L(z) = h_{K^{-1}p^n L}(z)$$

follows directly from the definition of the Gaussian channel (1). Consequently,

$$\Phi[\gamma(L_n)] = \gamma(K^{-1}L_n) \quad (3)$$

and, in accordance with Proposition 1,

$$H(\Phi[\gamma(L_n)]) - H(\gamma(L_n)) = \log |\det K|_p. \quad (4)$$

The latter formula immediately implies an evaluation for the minimum entropy gain  $G(\Phi) \leq \log |\det K|_p$ . Further, we recall (Proposition 1) that operator  $|L_n|^{-1}\gamma(L_n)$  is an orthogonal projection, and the sequence of these operators strongly converges to the unit operator  $I$  when  $n \rightarrow \infty$ . Therefore, we can correctly determine  $\Phi[I]$  (the channel is regular); moreover, formula (3) and Proposition 1 imply  $\|\Phi[I]\| = |\det K|_p^{-1}$ . Next, we use the evaluation of the minimum entropy gain for a regular channel  $-\log \|\Phi[I]\| \leq G(\Phi)$  obtained in [12]. The theorem is proved. Note that the expression for the entropy gain in the  $p$ -adic linear bosonic Gaussian channel is given by a formula similar to the corresponding expression for the real linear bosonic Gaussian channel obtained in [12].

## 5 Adelic channels and entropy gain

Now let  $F$  be a two-dimensional vector space over the field  $\mathbb{Q}$  of rational numbers,  $\Delta$  be a nondegenerate symplectic form taking values in the field  $\mathbb{Q}$ , and  $K$  be a nondegenerate linear transformation of the space  $F$ . For each prime  $p$ , we construct the corresponding linear bosonic Gaussian channel  $\Phi_p$  and calculate the entropy gain  $G(\Phi_p)$  of such a channel. We can also construct a real linear bosonic Gaussian channel  $\Phi_\infty$  and calculate the entropy gain  $G(\Phi_\infty)$ . Let  $\mathcal{P}$  denotes the set of all prime numbers. The following statement is true.

### Theorem 2

$$G(\Phi_\infty) + \sum_{p \in \mathcal{P}} G(\Phi_p) = 0.$$

Note that  $\det K \in \mathbb{Q}$ . The further follows from a simple adelic formula

$$|\det K| \prod_{p \in \mathcal{P}} |\det K|_p = 1,$$

which is valid for an arbitrary nonzero rational number (for example, see [7]).

Theorem 2 can be interpreted as follows. Consider the adelic linear bosonic Gaussian channel generated by the linear transformation  $K$  of a two-dimensional vector space over the field of rational numbers, that is, the tensor product of the channels  $\Phi_p$  over all primes  $p$  and the real channel  $\Phi_\infty$ . A nontrivial entropy gain  $G(\Phi_p)$  is possible in each component of this adelic channel. However, the total entropy gain in the adelic channel is zero. There is a nontrivial exchange of information between the components of the adelic channel when the total entropy is conserved. If we accept the hypothesis that at the fundamental level, our world ... is adelic ..., then Theorem 2 can be used, for example, to interpret the black hole information paradox.

## References

- [1] *Volovich I.V.* Number theory as the ultimate physical theory // Preprint CERN-TH-4781-87— Cern, Geneva, 1987.
- [2] *Volovich I.V.* p-adic string // Classical and Quantum Gravity— 1987.— V. 4, no. 4— P. L83–L87.
- [3] *Vladimirov V.S., Volovich I.V., Zelenov E.I.* p-Adic analysis and mathematical physics.— World Scientific, Singapore, 1994.
- [4] *Dragovich B, Khrennikov A.Yu., Kozyrev S.V., Volovich I.V.* On p-adic mathematical physics // p-Adic numbers, Ultram. Anal. Appl. — 2009. V. 1, no. 1— P. 1–17.— arXiv:0904.4205 [math-ph].
- [5] *Dragovich B, Khrennikov A.Yu., Kozyrev S.V., Volovich I.V., Zelenov E.I.* p-Adic mathematical physics: the first 30 years // p-Adic numbers, Ultram. Anal. Appl. — 2017. V. 9— P. 87–121.— arXiv:1705.04758 [math-ph].
- [6] *Zelenov E.I.* p-Adic model of quantum mechanics and quantum channels // Proc. Steklov Inst. Math. — 2014. — V. 285 — P. 132–144.
- [7] *Schikhof W.H.* Ultrametric calculus. An introduction to p-adic analysis. — Cambridge University Press, 1984.
- [8] *Serre J.-P.* Trees. — Springer-Verlag, 1980.



- [9] *Holevo A.S.* Statistical structure of quantum theory. — Springer-Verlag, 2001.
- [10] *Zelenov E.I.* Representations of commutations relations for p-adic systems of infinitely many degrees of freedom // Journal of Mathematical Physics — 1992. — V. 33, no. 1. — P. 178-188. —
- [11] *Holevo A.S.* Quantum systems, channels, information. A mathematical introduction. — De Gruyter, Berlin–Boston, 2019.
- [12] *Holevo A.S.* The entropy gain of infinite-dimensional quantum channels — 2010. — arXiv:1003.5765 [math-ph].
- [13] *Manin Yu.I.* Reflections on arithmetical physics. Conformal Invariance and string theory. Poiana Brasov, 1987. — Academic Press, 1989.