

Hidden Wave Function of Twisted Bilayer Graphene: Flat Band as a Landau Level

Fedor K. Popov* and Alexey Milekhin†
(Dated: March 17, 2024)

We study the chirally symmetric continuum model (CS-CM) of the twisted bilayer graphene. The equation on a flat band could be interpreted as a Dirac equation on a torus in the external non-abelian magnetic field. We prove that the existence of the flat band implies that the wave-function has a zero and vice versa. We found a hidden solution in the CS-CM model that has a pole instead of a zero. Our main result is that in the basis of the flat band and hidden wave functions the flat band could be interpreted as Landau level in the external magnetic field. From that interpretation we show the existence of extra flat bands in the magnetic field.

Introduction. Twisted bilayer graphene (TBG) has recently drawn a lot of attention from the physics community due to its interesting properties and applications [1] - [40]. One of the most prominent features is the recent discovery of correlated insulators and superconductivity, that are observed in a narrow range of twist angles near $\theta = 1.05^\circ$, which is usually referred to as the magic angle. At this angle the system develops a nearly flat band near charge neutrality. Recently the flat band was explained in a chiral model of TBG that neglects certain hoppings [43]. In this Letter we continue the exploration of the mathematical structures of the flat band and demonstrate the connection with the vector bundles over the Riemann surfaces of higher genus.

TBG consists of a two graphene sheets placed on top of each other at small angle θ that form a long-period pattern (Moire pattern). One can estimate that the period of the resulting super-lattice is of the order $L(\theta) \sim \frac{a}{\theta} \gg a$, where a is the graphene lattice constant. That allows us to consider a continuum model for the Hamiltonian instead of a lattice one. Such an approach was used by Bistrizer and MacDonald [44, 45] and by Lopes dos Santos [46]. The numerical study of that model confirmed the existence of the flat band at magic angle $\theta_1^* \approx 1.05^\circ$. The modification studied here [43] neglects the coupling between the sublattices of the graphene. The system acquires a chiral symmetry and usually referred to as the *chirally symmetric continuum model* (CS-CM). The Hamiltonian of CS-CM can be written in the following form [?]

$$H = \begin{pmatrix} 0 & \mathcal{D}(\vec{r}) \\ \mathcal{D}^*(-\vec{r}) & 0 \end{pmatrix}, \mathcal{D}(\vec{r}) = \begin{pmatrix} 2i\bar{\partial} & U(\vec{r}) \\ U(-\vec{r}) & 2i\bar{\partial} \end{pmatrix}, \quad (1)$$

where \vec{r} is the vector in 2d graphene sheet, $\bar{\partial} = \partial_x - i\partial_y$ is anti-holomorphic derivative along a sheet and $U(r)$ is the hopping potential between two sheets of the TBG. This Hamiltonian acts on the wave functions Ψ , that are the four-component functions arising because we consider two layers of graphene.

Since we are interested in the existence of a flat band near charge neutrality we should study the following equations

$$H\Psi = H \begin{pmatrix} \phi \\ \psi \end{pmatrix} = 0, \text{ or just } \mathcal{D}(\vec{r})\psi = 2i(\bar{\partial} + \bar{A})\psi = 0, \\ \text{where } \bar{A} = \begin{pmatrix} 0 & U(\vec{r}) \\ U(-\vec{r}) & 0 \end{pmatrix}, \quad (2)$$

where $\psi = (\psi_1, \psi_2)^T$ and $\phi = (\phi_1, \phi_2)^T$ are the components of the wave function Ψ . Due to the enhanced chiral symmetry the equations on ψ and ϕ are decoupled, which allows for a deeper analytical investigation of the properties of the CS-CM. These equations could be interpreted as a Dirac equation in a non-abelian magnetic field $\bar{A} \in \mathfrak{su}(2)$ on a Riemann surface [47]. We will use this interpretation to bolster our intuition and draw conclusions. Bloch wave-functions obey the following boundary conditions

$$\psi_{\vec{k}}(\vec{r} + \vec{a}_{1,2}) = e^{i\vec{k}\vec{a}_{1,2}} U_{\omega} \psi_{\vec{k}}(\vec{r} + \vec{a}_{1,2}), \quad (3)$$

where U_{ω} is a matrix such that $\det U_{\omega} = 1$, $\vec{a}_{1,2}$ are the periods of the Moire super-lattice and the vector \vec{k} defines the location in the Brillouin zone.

If the solution exists for any point k in Brillouin zone then the system has a flat band. In the paper [43] the potential \bar{A} was chosen such that the system of equations (2) and (3) has a solution for any point in the Brillouin zone. But for a general chosen potential \bar{A} , the system of equations (2) and (3) does have a smooth finite solution *only* at some points k in Brillouin zone.

The purpose of this paper is to consider a generic \bar{A} and try to get general properties independent of the concrete form of \bar{A} . Our main result is that once a flat band exists without the magnetic field, there is an additional flat band, that is not physical but appears once the external magnetic field is applied. Therefore, the number of flat bands increases in the presence of the magnetic field.

The paper is organized as follows. First, we will build an integral of motion of the equation (2) that is proportional to the Fermi velocity and prove that the flat band appears if and only if this invariant is equal to zero. We then demonstrate that the system of equations (2) and (3) admits an additional solution, which is singular and

* Princeton University, Physics Department; fpopov@princeton.edu

† UC Santa Barbara, Physics Department; milekhin@ucsb.edu

therefore unphysical. This second solution will allow us to rewrite the system of equations in the form of two Dirac equations on a torus with effective $U(1)$ magnetic fields. That shows the direct connection of flat band to the Landau levels.

Finally, we will show that the second solution becomes non-singular and leads to additional flat bands if one introduces an external magnetic field. The mathematical details are delegated to the Appendix.

Fermi velocity, zeros of wave functions and the flat band. For simplicity, let us consider equation (2) alone without taking into account the boundary conditions (3):

$$D\psi = (\bar{\partial} + \bar{A})\psi = 0, \quad \psi \in \mathbb{C}^2, \quad \bar{A} \in \mathfrak{su}(2). \quad (4)$$

From the mathematical point of view, the external field \bar{A} could be seen as a holomorphic connection on a complex plane.

Therefore equation (4) is just an equation for the meromorphic section of the trivial bundle $E = \mathbb{C} \times \mathbb{C}^2 \rightarrow \mathbb{C}$.

From the physical point of view, a wave function must be finite. So we must assume that ψ is also a *holomorphic* function of vector bundle E [?].

Let us consider two finite solutions of the equation (4), ψ_1, ψ_2 . One can compute the Wronskian of these solutions

$$W(\psi_1, \psi_2) = \det(\psi_1, \psi_2) = W(r), \quad \text{then} \\ \bar{\partial}W(r) = \text{tr } \bar{A} \cdot W(r), \quad \text{tr } \bar{A} = 0 \Rightarrow W(r) = W(z), \quad (5)$$

where we have used a standard argument from the theory of differential equations and the fact that $\bar{A} \in \mathfrak{su}(2)$ is traceless. We come to the conclusion that the Wronskian $W(r)$ must be an analytic function. If $\psi_{1,2}$ are finite on a whole Riemann sphere, $W(z)$ is holomorphic on a compact manifold and therefore must be constant. This property could be generalized to higher genus Riemann surfaces and will provide a necessary and sufficient condition for the existence of flat band in the system. One can actually show [43] that $W(z)$ is proportional to the Fermi velocity of the system.

From flat band to zero Wronskian. Here we prove that we can not have a flat band unless $W(z) = 0$. For the application to TBG we should study the equation (2) on a torus. Namely, we can consider a quotient $\pi : \mathbb{C} \rightarrow \mathbb{C}/\Lambda$, where $\Lambda = ma_1 + na_2, m, n \in \mathbb{Z}, a_{1,2} = a_{1,2}^x + ia_{1,2}^y$. We must impose boundary conditions to glue the wave function as we shift along the lattice Λ (3), which are just the gluing functions of the vector bundle over the torus. Without loss of generality we will set $a_1 = 1$ and $a_2 = \tau$.

The vector bundle defined by boundary conditions (3) with quasi-momentum K we denote as \mathbb{C}_K^2 . Again, equation (4) with the connection \bar{A} defines the meromorphic section of this vector bundle.

For the sake of argument, we assume that there are at least two points K_1, K_2 (not to be confused with the special points K, K') in the Brillouin zone where the solution exists.

Using the projection $\pi : \mathbb{C} \rightarrow \mathbb{C}/\Lambda$ we can lift the section from torus to the complex plane. Hence, the relation (5) should still hold true. If we have two holomorphic solutions $\psi_{K_{1,2}}$ at two different points of the Brillouin zone $K_{1,2}$ we can lift them to the complex plane and compute their Wronskian

$$W_{K_1+K_2}(z) = W(\psi_{K_1}, \psi_{K_2}) = \det(\psi_{K_1}, \psi_{K_2}), \quad (6)$$

but because the $W(z)$ is holomorphic and bounded in the complex plane \mathbb{C} (due to the periodicity conditions (3) and the fact that ψ_{K_1}, ψ_{K_2} are finite) we must conclude that $W(z) = \text{const}$. Moreover, using boundary conditions (3) we have

$$W_{K_1+K_2}(z + a_{1,2}) = W_{K_1+K_2}(z)e^{i(K_1+K_2)a_{1,2}}, \quad (7)$$

But if $K_1 + K_2 \neq 0$ and $W(z)$ is constant, the boundary conditions are satisfied only if $W(z) = 0$. Hence, there are only two possibilities:

1. $W(z) = 0$ and K_1, K_2 are arbitrary.
2. $W(z) \neq 0$ but $K_1 = -K_2$.

We start with the second possibility. We normalize the solutions such that $W(z) = 1$. Then we get immediately that ψ_{K_1}, ψ_{-K_1} are nowhere zero, because otherwise the Wronskian would be equal to zero at points where $\psi_{\pm K_1} = \vec{0}$. Since $W(z)$ is non-zero, the solutions ψ_{K_1} and ψ_{-K_1} are linearly independent at each point of the torus \mathbb{C}/Λ . If we consider now matrix $S = (\psi_{K_1}, \psi_{-K_1})$ it satisfies the following equation

$$(\bar{\partial} + \bar{A})S = 0, \quad \bar{A} = -\bar{\partial}S \cdot S^{-1}, \quad (8)$$

where we have used the fact that if $\det S \neq 0$ the matrix S is invertible. We would like to point out, that the equation (8) does not mean \bar{A} is a flat connection, since $S \in SL(2, \mathbb{C})$ rather than $SU(2)$ and therefore is not a gauge transformations.

Let us consider another solution ψ of the eq. (4). Since $\psi_{\pm K_1}$ are linearly independent we can always represent ψ as a linear combination of these solutions:

$$\psi = v_1(r)\psi_{K_1} + v_2(r)\psi_{-K_1},$$

Applying the operator $D = \bar{\partial} + \bar{A}$ we get

$$D\psi_k = \bar{\partial}v_1\psi_{K_1} + \bar{\partial}v_2\psi_{-K_1} = 0. \quad (9)$$

Since $\psi_{\pm K_1}$ are linearly independent at each point of the torus \mathbb{C}/Λ it follows that the coefficients v_i must be holomorphic $\bar{\partial}v_i = 0$. The functions ψ and $\psi_{\pm K_1}$ are finite and non-zero everywhere, hence we must have that v_i are bounded. From the maximum principle for analytic functions on a complex plane we must have that v_i are just constant. Therefore if we have an arbitrary solution of the equation (4) at the point k of the Brillouin zone we must have

$$\psi_k = v_{k,K_1}\psi_{K_1} + v_{k,-K_1}\psi_{-K_1}, \quad v_{k,\pm K_1} \in \mathbb{C}, \quad (10)$$

but it is easy to see that with any choice of the numbers $v_{k,\pm K_1}$ we are not able to satisfy the boundary conditions (3), unless $k = \pm K_1$. Therefore we can not have a flat band if $W(z) \neq 0$.

From zero Wronskian to a flat band and a wave function with a zero. Let us prove the converse. Namely, if $W(z) = 0$ at some points K_1, K_2 in the Brillouin zone, the system develops a flat band. In other words, the equation (4) has a solution at any point k in the Brillouin zone.

We start the proof by noticing that since $W(z) = 0$ and ψ_{K_1, K_2} satisfy the equation (4), then wave function ψ_{K_1} has a zero. Let us assume the opposite: that $\psi_{K_1}(r) \neq 0$ at any point of the torus \mathbb{C}/Λ . Because torus is compact the minimum $\min_{r \in \mathbb{C}/\Lambda} |\psi_{K_1}(r)| = m > 0$ is reachable. If $W(z) = 0$ the wave functions ψ_{K_1, K_2} are proportional to each other:

$$\psi_{K_2}(r) = \gamma(r)\psi_{K_1}(r), \quad D\psi_{K_2} = \bar{\partial}\gamma(r)\psi_{K_1}(r) = 0,$$

where $\gamma(r)$ is bounded as $|\gamma(r)| < \frac{|\psi_{K_2}(r)|}{m}$ and holomorphic $\bar{\partial}\gamma(r) = 0$. Then the function $\gamma(r) = \gamma(z)$ must be constant by the maximum principle. However, this is impossible since ψ_{K_1, K_2} satisfy different boundary conditions. Hence we must conclude that ψ_K has at least one simple zero [?].

We can now follow the procedure described in [43] and construct solution at any point k of the Brillouin zone.

Hidden wave function. We can draw some additional conclusions the existence of the holomorphic section with a zero at any point K of Brillouin zone. For the sake of argument we will assume that ψ_K has one simple zero, but this could be easily generalized to the case of multiple zeros.

Let us notice that at general K there can be only one holomorphic section ψ_K . Indeed, if there are two holomorphic linearly independent sections $\psi_K^{1,2}$, their Wronskian must be a holomorphic non-zero double periodic function with specific boundary conditions, that must be constant and $2K = 0 \pmod{\Lambda}$. We come to the conclusion if the other solution exists, it must be meromorphic.

The holomorphic section ψ_K (flat-band wave function) of the bundle \mathbb{C}_K^2 forms a subbundle, that we denote as γ . Then one can consider an exact short sequence:

$$0 \rightarrow \gamma \rightarrow \mathbb{C}_K^2 \rightarrow (\gamma)^\perp \rightarrow 0 \quad (11)$$

where $(\gamma)^\perp = \mathbb{C}_K^2/\gamma$. The bundles γ and $(\gamma)^\perp$ are line bundles, that are classified by the first Chern class c_1 [49] (c_1 is just the number of zeros minus the number of poles). From this we have that the first Chern class for the subbundles γ and $(\gamma)^\perp$ must satisfy the following relation

$$c_1((\gamma)^\perp) = -c_1(\gamma), \quad (12)$$

suggesting that if γ has a holomorphic section, $(\gamma)^\perp$ has a section, but instead of a zero it has a pole. This reasoning is not enough for the proof of its existence, because (11) may not split. In other words, an element in

\mathbb{C}_K^2/γ might not have proper boundary conditions: going around a torus cycle will not only produce a phase, but also add a multiple of ψ_K . From physical point of view it means that we can not simply represent this 2d system as a stack of two topological materials with opposite Chern numbers. The theory of vector bundles over the Riemann surfaces was actively studied by Donaldson [51]. Nevertheless, one can show that the short sequence (11) is split over the torus. Here we will present a physical construction. That can be used to find the solution explicitly. In the appendix we will prove the existence using algebro-geometric methods.

Let us start from a holomorphic section ψ_K , that is a solution of the equation (4), satisfies boundary conditions (3), and has a zero at some point z_0 . Let us assume for a moment that we somehow managed to find another solution ϕ_K , that is linearly independent from ψ_K . If such a solution exists, the Wronskian $\widetilde{W}(\psi_K, \phi_K)$ should be a meromorphic function

$$\det(\psi_K, \phi_K) = \widetilde{W}(z), \quad (13)$$

that satisfies double-periodic boundary conditions $\widetilde{W}(z + a_{1,2}) = e^{2i\vec{K}\vec{a}_{1,2}}\widetilde{W}(z)$ (see (3)). Unlike the previous Section, ϕ_K might have poles, so we can not conclude that \widetilde{W} is constant. However, an analytic function with this properties exists and is unique up to a normalization factor [49]. Namely, this function is represented as

$$\widetilde{W}(z) = e^{2i\vec{K}\vec{a}_1 z} \frac{\vartheta(z - z_0; \tau)}{\vartheta(z - z_\infty; \tau)}, \quad z_0 - z_\infty = \vec{K} \frac{\vec{a}_2 - \tau \vec{a}_1}{\pi}$$

where $\vartheta(z; \tau)$ is a Jacobi theta function. Since ψ_K is finite everywhere, ϕ_K must have a pole at the point $z = z_\infty$. From this we get a simple linear equation that ϕ_K should satisfy

$$\phi_K^1 \psi_K^2 - \phi_K^2 \psi_K^1 = \widetilde{W}(z) \quad (14)$$

Having determined \widetilde{W} let us now construct ϕ_K . At any point $z \in \mathbb{C}/\Lambda$ this equation has at least one solution. Since at point $z = z_0$ both sides of the equation (14) has a simple zero we can analytically continue the solution at this point. Let us pick an arbitrary solution to the eq. (14) and denote it as $\zeta_K(r)$. Any other solution of the eq. (14) is

$$\zeta_K^\lambda(r) = \zeta_K(r) + \lambda(r)\psi_K, \quad (15)$$

where $\lambda(r)$ is an arbitrary function.

We can derive a relation for the function $\zeta_K(r)$. Namely, we apply an operator $\bar{\partial}$ to the Wronskian to get

$$\begin{aligned} \bar{\partial}\widetilde{W}(z) &= \bar{\partial}\det(\psi_K, \zeta_K) = \\ &= \det(D\psi_K, \zeta_K) + \det(\psi_K, D\zeta_K) = \\ &= \det(\psi_K, D\zeta_K) = 0 \end{aligned} \quad (16)$$

This means that in general $D\zeta_K$ is proportional to the wave function ψ_K

$$D\zeta_K = \eta(r)\psi_K, \quad (17)$$

for some function $\eta(r)$, that may have singularities. To clarify what we have done, the solution ζ_K is just an arbitrary solution to the equation (14), and does not satisfy the equation (4). In eq. (14) we can arbitrary choose ζ_K^1 . It could have some singularities. To avoid such a problem we set $\zeta_K^1 = 1$ on the torus. Then the singularities of ζ_K^2 come only from the function $\widetilde{W}(z)$.

As we discussed before the function ζ_K is not unique, so we can consider ζ_K^λ from the eq. (15). This freedom allows to set the right hand side of the eq. (17) to zero. Indeed,

$$\zeta_K^\lambda = \zeta_K + \lambda(r)\psi_K \quad (18)$$

$$D\zeta_K^\lambda(r) = D\zeta_K(r) + \bar{\partial}\lambda\psi_K(r) = [\eta(r) + \bar{\partial}\lambda(r)]\psi_K,$$

Therefore we just need to solve the following equation on a torus

$$\bar{\partial}\lambda = -\eta + C\delta^{(2)}(z - z_0), \quad (19)$$

with periodic boundary conditions $\lambda(r + a_{1,2}) = \lambda(r)$. The term proportional to δ function is allowed since ψ_K has a zero at point $z = z_0$. To solve (19) we make a Fourier transform over the torus

$$\bar{k}\lambda(k) = \eta(k) + C\delta_{k,0}, \quad (20)$$

This equation has a solution for any k if $C = -\eta(0)$. The general solution then is

$$\lambda(k) = \frac{k\eta(k)}{|k|^2}, \quad \lambda(0) = 0 \quad (21)$$

Therefore we managed to find a second solution to the equation (4) with boundary conditions (3), that is singular but linearly independent from the holomorphic solution.

One can check that $\lambda(r)\psi_K$ is finite everywhere and therefore the pole of ζ_K . We have checked numerically that if one follows the above procedure we can derive wave function that has a simple pole and satisfies the system of equations (4) and (3).

Hidden Landau Levels. We have two solutions at the Brillouin point K : ψ_K^0 with a zero at a point z_0 and ψ_K^∞ with a pole at a point z_∞ . Then we can introduce wave functions, that are finite defined as

$$\begin{aligned} \hat{\psi}^\infty &= e^{i\vec{K}\vec{a}_1 z - \frac{1}{2}Bz\bar{z}}\vartheta(z - z_\infty; \tau)\psi_K^\infty, \\ \hat{\psi}^0 &= \frac{e^{i\vec{K}\vec{a}_1 z + \frac{1}{2}Bz\bar{z}}}{\vartheta(z - z_0; \tau)}\psi_K^0, \end{aligned} \quad (22)$$

where B is a constant magnetic field corresponding to flux 1 in the Moire lattice.

One can see that

$$\begin{aligned} S &= \begin{pmatrix} \hat{\psi}^0 & \hat{\psi}^\infty \end{pmatrix} \\ \det S &= \det(\hat{\psi}^0, \hat{\psi}^\infty) = 1 \end{aligned} \quad (23)$$

Then since $\det S = 1$ we can invert this matrix at each point of the lattice \mathbb{C}/Λ . This matrix allows to rewrite the Dirac operator as

$$\hat{D} = S^{-1}DS = \begin{pmatrix} \bar{\partial} - \frac{1}{2}Bz & 0 \\ 0 & \bar{\partial} + \frac{1}{2}Bz \end{pmatrix}, \quad (24)$$

This shows that in this basis we have just effectively split TBG into two sheets with some magnetic field B . The magnitude of this field is the same in both sheets, but differs in sign. The boundary conditions and the equations indicate, that the equations correspond to the Landau levels studied on the complex tori (see Appendix for the detailed discussion). In one layer the magnetic field supports the existence of the wave function, while in the other the magnetic field supports a solution, that has a pole instead of a zero and therefore is not physical. Therefore we clearly see that the flat band in the chiral model is a Landau level created by the interlayer interaction. An analogous conclusion was derived from the different arguments [50].

Now we can turn on an external magnetic field. It corresponds to the adding of a component proportional to the identity matrix to the anti-holomorphic connection in the eq. (4). This part is not sensitive to the transformation in the eq. (24). Therefore the equation for the zero mode has the following form

$$\hat{D}_B = \begin{pmatrix} \bar{\partial} - \frac{1}{2}Bz + \bar{A}_{U(1)} & 0 \\ 0 & \bar{\partial} + \frac{1}{2}Bz + \bar{A}_{U(1)} \end{pmatrix}, \quad (25)$$

where $\bar{A}_{U(1)}$ is a gauge potential for the external magnetic field. For simplicity we assume that $\bar{A}_{U(1)}$ has a flux $\Phi \in \mathbb{Z}$ through the Moire super-lattice. The equation (25) shows us that the system decouples into two non-interacting layers with fluxes $\Phi \pm 1$. Then one can see that if $\Phi \leq -1$ there are no flat bands, for $\Phi = 0$ there is exactly 1, for $\Phi \geq 1$ there are 2Φ flat bands.

Note that we have studied only the anti-holomorphic part of the Hamiltonian (1). The holomorphic part (the other chirality) exhibits the same properties but for $\Phi \rightarrow -\Phi$. It means that in total there are $2|\Phi|$ flat bands for $|\Phi| > 0$ and for $\Phi = 0$ there are only 2 flat bands. We can study the case of the rational flux, in this case we would reproduce the Hofstadter's butterfly [52].

Acknowledgments FKP dedicates this paper to the memory of his father, Popov Kalina Fedorovich. We would like to thank Grisha Tarnopolsky and Igor R. Klebanov for the collaboration at the early stages of the work. We would like to thank Vladyslav Kozii, Anastasia Aristova, Alexander Avdoshkin, Nikita Sopenko, Dumitru Calugaru for useful comments and discussions. We would like to thank Alexander Gorsky, Emil Akhmedov

and Peter Czajka for useful comments on the draft of the paper. AM also would like to thank C. King for moral support. The research was supported in part by the US NSF under Grants No. PHY-1620059 and PHY-1914860.

Supplementary Material

Landau Levels on torus. In this section we briefly review the wave functions on torus, we will mostly follow Haldane and Rezayi [53]. We consider that we have a complex torus \mathbb{C}/Λ , $\Lambda = \{n + m\tau; n, m \in \mathbb{Z}\}$. We want to find solutions to the following equation

$$\hat{D}_B \psi = \left(\bar{\partial} + \frac{1}{2} eBz \right) \psi = 0, \quad F = \bar{\partial}A - \partial\bar{A} = B. \quad (1)$$

To establish boundary conditions we consider a shift of z by a lattice vector $a_i = 1, \tau$ to get

$$\left(\bar{\partial} + \frac{1}{2} eBz + \frac{1}{2} eBa_i \right) \psi = 0 \quad (2)$$

To remove the change in the gauge potential, we should make a gauge transformation

$$\psi(z) \rightarrow \psi(z) e^{-\frac{1}{2} eBa_i \bar{z} + \frac{1}{2} eB\bar{a}_i z}, \quad (3)$$

this can be used to define the boundary conditions. Namely,

$$\begin{aligned} T_1 : \psi(z+1) &= \psi(z) e^{-\frac{1}{2} eB\bar{z} + \frac{1}{2} eBz}, \\ T_\tau : \psi(z+\tau) &= \psi(z) e^{-\frac{1}{2} eB\tau\bar{z} + \frac{1}{2} eB\bar{\tau}z} \end{aligned} \quad (4)$$

We should check the consistency of this boundary conditions, that $T_1 T_\tau = T_\tau T_1$. One can check that

$$T_1 T_\tau \psi(z) = e^{eB(\tau - \bar{\tau})} T_\tau T_1 \psi(z) \quad (5)$$

The difference between this phases is $e^{eB(\tau - \bar{\tau})} = 1$. That gives a condition for the consistent boundary conditions (4)

$$\pi\Phi = eB \operatorname{Im} \tau, \quad eB = \frac{\pi\Phi}{\operatorname{Im} \tau}, \quad \Phi \in \mathbb{Z} \quad (6)$$

This conditions give that the integral over a fundamental period is equal to $\Phi = \frac{1}{2\pi} \int F d^2 z = \frac{eB}{\pi} \operatorname{Im} \tau$.

Then if we can consider a general k from Brillouin zone to get

$$\begin{aligned} \left(\bar{\partial} + \frac{1}{2} eBz \right) \psi_k(z) &= 0, \\ \psi_k(z + a_i) &= \psi_k(z) e^{-\frac{1}{2} eBa_i \bar{z} + \frac{1}{2} eB\bar{a}_i z + i(k, a_i)} \end{aligned} \quad (7)$$

Where $(k, z) = k_x x + k_y y = \operatorname{Im} k\bar{z}$, $k = k_x + ik_y$, $z = x + iy$.

For $\Phi = -1$ the equation (7) is easy to solve, we get

$$\psi_{k,-1}(z) = \vartheta \left(z + \frac{ik}{2eB}; \tau \right) e^{\frac{1}{2} eB(z + \frac{ik}{2eB})^2 + i\frac{1}{2} \bar{k}z - \frac{1}{2} eBz\bar{z}}$$

The zero of this function is located at

$$z_0 = \frac{1}{2} + \frac{1}{2} \tau - \frac{ik}{2eB} \quad (8)$$

Whereas for $\Phi = +1$ we get a solution with a pole:

$$\psi_{k,-1}(z) = \frac{1}{\vartheta \left(z + \frac{ik}{2eB}; \tau \right)} e^{\frac{1}{2} eB(z + \frac{ik}{2eB})^2 + i\frac{1}{2} \bar{k}z - \frac{1}{2} eBz\bar{z}}$$

If $\Phi \neq -1$, the solution is just

$$\psi_{K,\Phi}(z) = \prod_{k_1 + \dots + k_\Phi = K} \psi_{k_i,-1}(z) \quad (9)$$

It seems that there is now an infinite number of the wave functions at given K . One can show that there is only a finite number of the linearly independent solutions (9).

To compute this dimension we can use the Riemann–Roch formula [49] for the operator \hat{D}_B in eq. (1). This gives that $\dim \ker \hat{D}_B = \Phi$, so when Φ is negative we don't have any finite solutions for the Landau levels at any point of the Brillouin zone. The case $\Phi = 0$ is special - there is some zero modes but only at special points of the Brillouin zone.

The splitting of short exact sequence and Čech cohomology. In this subsection we would like to clarify the existence of the second solution from the cohomology point of view, that is quite rigorous and could be generalized to the higher Riemann surfaces [54].

We start with the rigorous formulation of the problem. Assume that we have a Riemann surface \mathcal{M} with some holomorphic vector bundle of rank 2, $\pi : E \rightarrow \mathcal{M}$ and some connection \bar{A} . Namely, we have a covering of the Riemann surface with open subsets $\{U_\alpha\}$ where the vector bundle could be trivialized

$$\mathcal{M} = \bigcup_{\alpha} U_{\alpha}, \quad E|_{U_{\alpha}} \approx U_{\alpha} \times \mathbb{C}^2. \quad (10)$$

When we move from one covering U_α to another U_β we need to glue the section with the use of the gluing functions $g_{\alpha\beta}^0$, that are holomorphic $\bar{\partial}g_{\alpha\beta}^0 = 0$. The connection \bar{A}_β transforms as

$$\bar{A}_\beta = \hat{g}_{\alpha\beta}^{-1} \bar{A}_\alpha \hat{g}_{\alpha\beta} + \hat{g}_{\alpha\beta}^{-1} \bar{\partial} \hat{g}_{\alpha\beta} \quad (11)$$

We want to find holomorphic sections of these vector bundles. Namely, we want to find a collection of functions $\{\psi_\alpha\}$, such that the following conditions are satisfied

$$\psi_\alpha^0 = g_{\alpha\beta}^0 \psi_\beta^0, \quad (\bar{\partial} + \bar{A}_\alpha) \psi_\alpha^0 = 0 \quad (12)$$

We can get rid of the connection \bar{A}_α by solving the equation (12) at each covering and performing the gauge transformation.

Then we can generally study the following problem

$$\psi_\alpha = g_{\alpha\beta} \psi_\beta, \quad \bar{\partial} \psi_\alpha = 0. \quad (13)$$

So we just need to find a *meromorphic* sections of the vector bundle E defined by cocycles $g_{\beta\alpha}$ in the assumption

that we have a *holomorphic* sections of the bundle (13) ψ^h . Namely, we have a collection of holomorphic functions $\psi_\alpha^h(z)$ defined at each coverings and satisfy boundary conditions

$$\psi_\alpha^h = g_{\alpha\beta} \psi_\beta^h \quad (14)$$

We want this functions ψ_α^h to be non-zero at any coverings of \mathcal{M} . Whenever we encounter a zero in some covering U_α , $\psi_\alpha^h(z_0^\alpha) = 0$, we redefine holomorphic section and gluing functions $\hat{\psi}_\alpha^h = \frac{1}{z-z_0^\alpha} \psi_\alpha^h$ and $\hat{g}_{\alpha\beta} = \gamma_{\alpha\beta}^1 g_{\alpha\beta}$, $\gamma_{\alpha\beta}^1 = \frac{z-z_0^\alpha}{z-z_0^\beta}$. This new function is nowhere zero and changes as

$$\hat{\psi}_\alpha^h = \gamma_{\alpha\beta}^1 g_{\alpha\beta} \hat{\psi}_\beta^0, \quad \text{where } \gamma_{\alpha\beta}^1 \in \mathbb{C}, \quad \gamma_{\alpha\beta}^1 \gamma_{\beta\gamma}^1 \gamma_{\gamma\alpha}^1 = 1 \quad (15)$$

Since this section is nowhere zero we can find another set of holomorphic functions that is linearly independent from $\hat{\psi}_\alpha$ at each point. We call this set of functions as $\hat{\psi}_\alpha^\infty$ and with analogous procedure introduce $\gamma_{\alpha\beta}^2$ to remove all zeros it can possible have. Because of this at each covering we can change basis to $\hat{\psi}_\alpha$ and $\hat{\psi}_\alpha^\infty$. One can check that gluing functions in this new basis of the vector bundle becomes

$$\hat{g}_{\alpha\beta} = \begin{pmatrix} \gamma_{\alpha\beta}^2 & h_{\alpha\beta} \\ 0 & \gamma_{\alpha\beta}^1 \end{pmatrix}, \quad \text{where} \quad \gamma_{\alpha\beta}^2 \gamma_{\beta\gamma}^2 \gamma_{\gamma\alpha}^2 = 1 \quad \text{and} \quad h_{\alpha\gamma} = \gamma_{\alpha\beta}^2 h_{\beta\gamma} + h_{\alpha\beta} \gamma_{\beta\gamma}^1 \quad (16)$$

If we got that $h_{\alpha\beta} = 0$ then the function $\hat{\psi}_\alpha^\infty$ would change through each other as

$$\hat{\psi}_\alpha^\infty = \gamma_{\alpha\beta}^2 g_{\alpha\beta} \hat{\psi}_\beta^\infty, \quad (17)$$

and define a legitimate section of the vector bundle E . Since the set of function $\gamma_{\alpha\beta}^2$ represents a line bundle, it has a meromorphic section: a set of meromorphic functions $f_\alpha(z)$ with property $\gamma_{\alpha\beta}^2 = \frac{f_\beta}{f_\alpha}$. Then new functions

$$\psi_\alpha^\infty = f_\alpha \hat{\psi}_\alpha^\infty \quad (18)$$

are holomorphic everywhere and transforms as

$$\psi_\alpha^\infty = g_{\alpha\beta} \psi_\beta^\infty, \quad (19)$$

And therefore represents a legitimate section of the original vector bundle E , but contains a pole at some point.

Let us show that we can get rid of $h_{\alpha\beta}$ by a proper redefinition of the arbitrary chosen $\hat{\psi}_\alpha^\infty$. Namely, we notice that the choice of $\hat{\psi}_\alpha^\infty$ is not unique. At each covering we can make a change

$$\hat{\psi}_\alpha^\infty \rightarrow \hat{\psi}_\alpha^\infty + h_\alpha(z) \hat{\psi}_\alpha^0 \quad (20)$$

It changes gluing functions as

$$h_{\alpha\beta} \rightarrow h_{\alpha\beta} + \gamma_{\alpha\beta}^2 h_\beta(z) - h_\alpha(z) \gamma_{\alpha\beta}^1 \quad (21)$$

This gives that $h_{\alpha\beta}$ belongs to $H^1(\mathcal{O}(\gamma_1 \gamma_2^{-1}))$ and by Serre duality are dual to $H^0(\mathcal{O}(\kappa \gamma_1^{-1} \gamma_2))$, where κ is a tangent line bundle. This line bundle does not have any holomorphic section if its Chern class is negative. We get

$$\begin{aligned} c_1(\kappa \gamma_1^{-1} \gamma_2) &= 2g - 2 - c(\gamma_1) + c(\gamma_2) = \\ &= 2g - 2 - 2c(\gamma_1) = -2c(\gamma_1) < 0 \end{aligned} \quad (22)$$

and we have used $g = 1$ (torus) $c_1(\gamma_1) \geq 1$ (because $\hat{\psi}_\alpha$ has at least one simple zero) and $c_1(\gamma_1) + c_1(\gamma_2) = 0$ that is related to the eq. (12).

Since $H^1(\mathcal{O}(\gamma_1 \gamma_2^{-1})) = 0$ the cohomology class represented by $h_{\alpha\beta}$ is trivial. Meaning, that we can always pick h_α such that $h_{\alpha\beta} = 0$ in the eq. (21). Then as we discussed above $\hat{\psi}_\alpha^\infty$ will represent a meromorphic section of the vector bundle E . This procedure could be generalized to the higher genus Riemann surfaces and other vector bundles.

-
- [1] Y. Cao, V. Fatemi, S. Fang, K. Watanabe, T. Taniguchi, E. Kaxiras, and P. Jarillo-Herrero, “Unconventional superconductivity in magic-angle graphene superlattices,” *Nature*, vol. 556, pp. 43–50, mar 2018.
- [2] Y. Cao, V. Fatemi, A. Demir, S. Fang, S. L. Tomarken, J. Y. Luo, J. D. Sanchez-Yamagishi, K. Watanabe, T. Taniguchi, E. Kaxiras, R. C. Ashoori, and P. Jarillo-Herrero, “Correlated Insulator Behaviour at Half-Filling in Magic Angle Graphene Superlattices,” *Nature Publishing Group*, vol. 556, no. 7699, pp. 80–84, 2018.
- [3] H. C. Po, L. Zou, A. Vishwanath, and T. Senthil, “Origin of Mott insulating behavior and superconductivity in twisted bilayer graphene,” pp. 1–32, 2018.
- [4] A. Thomson, S. Chatterjee, S. Sachdev, and M. S. Scheurer, “Triangular antiferromagnetism on the honeycomb lattice of twisted bilayer graphene,” jun 2018.
- [5] L. Zou, H. C. Po, A. Vishwanath, and T. Senthil, “Band Structure of Twisted Bilayer Graphene: Emergent Symmetries, Commensurate Approximants and Wannier Obstructions,” pp. 1–17, 2018.
- [6] F. Guinea and N. R. Walet, “Electrostatic effects and band distortions in twisted graphene bilayers,” *arXiv Mesoscale and Nanoscale Physics*, pp. 1–7, jun 2018.
- [7] F. Guinea and N. R. Walet, “Electrostatic effects and band distortions in twisted graphene bilayers,” pp. 1–9, jun 2018.
- [8] S. Carr, S. Fang, P. Jarillo-Herrero, and E. Kaxiras, “Pressure dependence of the magic twist angle in graphene superlattices,” pp. 1–6, 2018.
- [9] Y. Su and S.-Z. Lin, “Spontaneous vortex-antivortex lattice in superconducting twisted bilayer graphene,” pp. 1–9, 2018.
- [10] J. González and T. Stauber, “Kohn-Luttinger superconductivity in twisted bilayer graphene,” pp. 1–10, 2018.
- [11] F. Wu, T. Lovorn, E. Tutuc, I. Martin, and A. H. MacDonald, “Layer pseudospin skyrmions and topological moiré bands in twisted homobilayers,” pp. 1–8, jul 2018.
- [12] D. K. Efimkin and A. H. MacDonald, “Helical Network Model for Twisted Bilayer Graphene,” pp. 1–6, 2018.
- [13] J. Peng and J.-S. Wang, “Current-Induced Heat Transfer in Double-Layer Graphene,” vol. 12, no. May, 2018.
- [14] N. F. Q. Yuan and L. Fu, “A Model for Metal-Insulator Transition in Graphene Superlattices and Beyond,” 2018.
- [15] C. Xu and L. Balents, “Topological Superconductivity in Twisted Multilayer Graphene,” no. 1, pp. 1–6, 2018.
- [16] E. Riccardi, O. Kashuba, M. Cazayous, A. Sacuto, and Y. Gallais, “Raman Scattering as a Selective Probe of Chiral Electronic Excitations in Bilayer Graphene,” pp. 1–17, may 2018.
- [17] M. Ochi, M. Koshino, and K. Kuroki, “Possible correlated insulating states in magic-angle twisted bilayer graphene under strongly competing interactions,” 2018.
- [18] H. K. Pal, “On magic angles and band flattening in twisted bilayer graphene,” pp. 1–6, may 2018.
- [19] F. Wu, A. H. MacDonald, and I. Martin, “Theory of phonon-mediated superconductivity in twisted bilayer graphene,” vol. 1, no. 3, pp. 1–6, 2018.
- [20] Y.-h. Zhang, D. Mao, Y. Cao, P. Jarillo-Herrero, and T. Senthil, “Moiré Superlattice with Nearly Flat Chern Bands: Platform for (Fractional) Quantum Anomalous Hall Effects and Unconventional Superconductivity,” 2018.
- [21] X.-c. Wu, K. A. Pawlak, C.-m. Jian, and C. Xu, “Emergent Superconductivity in the weak Mott insulator phase of bilayer Graphene Moiré Superlattice,” pp. 1–6, 2018.
- [22] J. Kang and O. Vafek, “Symmetry, maximally localized Wannier states, and low energy model for the twisted bilayer graphene narrow bands,” pp. 1–9, may 2018.
- [23] B. Andrei Bernevig, Zhida Song, Nicolas Regnault, Biao Lian “TBG I: Matrix Elements, Approximations, Perturbation Theory and a $k \cdot p - 2$ -Band Model for Twisted Bilayer Graphene,”
- [24] Zhi-Da Song, Biao Lian, Nicolas Regnault, Andrei B. Bernevig, “TBG II: Stable Symmetry Anomaly in Twisted Bilayer Graphene,”
- [25] Andrei Bernevig, Zhida Song, Nicolas Regnault, Biao Lian “TBG III: Interacting Hamiltonian and Exact Symmetries of Twisted Bilayer Graphene,”
- [26] Biao Lian, Zhi-Da Song, Nicolas Regnault, Dmitri K. Efetov, Ali Yazdani, B. Andrei Bernevig “TBG IV: Exact Insulator Ground States and Phase Diagram of Twisted Bilayer Graphene,”
- [27] B. Andrei Bernevig, Biao Lian, Aditya Cowsik, Fang Xie, Nicolas Regnault, Zhi-Da Song “TBG V: Exact Analytic Many-Body Excitations In Twisted Bilayer Graphene Coulomb Hamiltonians: Charge Gap, Goldstone Modes and Absence of Cooper Pairing,”
- [28] Fang Xie, Aditya Cowsik, Zhida Son, Biao Lian, B. Andrei Bernevig, Nicolas Regnault “TBG VI: An Exact Diagonalization Study of Twisted Bilayer Graphene at Non-Zero Integer Fillings,”
- [29] J. M. Pizarro, M. J. Calderón, and E. Bascones, “The nature of correlations in the insulating states of twisted bilayer graphene,” no. Icm, 2018.
- [30] M. Koshino, N. F. Q. Yuan, T. Koretsune, M. Ochi, K. Kuroki, and L. Fu, “Maximally-localized Wannier orbitals and the extended Hubbard model for the twisted bilayer graphene,” pp. 1–11, may 2018.
- [31] D. M. Kennes, J. Lischner, and C. Karrasch, “Strong Correlations and d+id Superconductivity in Twisted Bilayer Graphene,” pp. 1–5, may 2018.
- [32] H. Isobe, N. F. Q. Yuan, and L. Fu, “Superconductivity and Charge Density Wave in Twisted Bilayer Graphene,” pp. 1–15, may 2018.
- [33] L. Rademaker and P. Mellado, “Charge-transfer insulation in twisted bilayer graphene,” pp. 1–6, 2018.
- [34] J.-B. Qiao and L. He, “Heterostrain engineering on twisted graphene bilayer around the first magic angle,” *arXiv*, pp. 1–23, may 2018.
- [35] M. Zarenia, A. R. Hamilton, F. M. Peeters, and D. Neilson, “Sign Reversal of Electron-Hole Coulomb Drag in Double Bilayer Graphene Heterostructures,” vol. 1, pp. 1–5, jun 2018.
- [36] T.-f. Chung, Y. Xu, and Y. P. Chen, “Transport measurements in twisted bilayer graphene: Studies of electron-phonon coupling and Landau level crossing,” *arXiv Condensed Matter*, pp. 1–35, may 2018.
- [37] M. Fidrysiak, M. Zegrodnik, and J. Spalek, “Unconventional topological superconductivity and phase diagram for a two-orbital model of twisted bilayer graphene,” pp. 1–6, 2018.

- [38] T. J. Peltonen, R. Ojajärvi, and T. T. Heikkilä, “Mean-field theory for superconductivity in twisted bilayer graphene,” pp. 1–7, 2018.
- [39] P. Cazeaux, M. Luskin, and D. Massatt, “Energy minimization of 2D incommensurate heterostructures,” vol. 66049, pp. 1–29, jun 2018.
- [40] K. Wang, T. Hou, Y. Ren, and Z. Qiao, “Enhanced Robustness of Zero-line Modes in Graphene via a Magnetic Field,” vol. 1, pp. 1–5, 2018.
- [41] Vladyslav Kozii, Michael P. Zaletel and Nick Bultinck, “Superconductivity in a doped valley coherent insulator in magic angle graphene: Goldstone-mediated pairing and Kohn-Luttinger mechanism,”
- [42] Vladyslav Kozii, Hiroki Isobe, Jörn W. F. Venderbos, and Liang Fu, “Nematic superconductivity stabilized by density wave fluctuations: Possible application to twisted bilayer graphene,” *Phys. Rev. B* 99, 144507
- [43] Tarnopolsky, Grigory and Kruchkov, Alex Jura and Vishwanath, Ashvin “Origin of Magic Angles in Twisted Bilayer Graphene,” *Physical Review Letters*, vol. 122, no. 10, mar 2019.
- [44] R. Bistritzer and A. H. MacDonald, “Moire bands in twisted double-layer graphene,” *Proceedings of the National Academy of Sciences*, vol. 108, pp. 12233–12237, jul 2011.
- [45] R. Bistritzer and A. H. MacDonald, “Moiré butterflies in twisted bilayer graphene,” *Physical Review B - Condensed Matter and Materials Physics*, vol. 84, no. 3, pp. 1–4, 2011.
- [46] J. M. B. Lopes dos Santos, N. M. R. Peres, and A. H. Castro Neto, “Continuum model of the twisted graphene bilayer,” *Phys. Rev. B*, vol. 86, p. 155449, Oct 2012.
- [47] P. San-Jose, J. González, and F. Guinea, “Non-abelian gauge potentials in graphene bilayers,” *Phys. Rev. Lett.*, vol. 108, p. 216802, May 2012.
- [48] Berline, Nicole and Getzler, Ezra and Vergne, Michele, “Heat kernels and Dirac operators”, Springer Science & Business Media
- [49] , Griffiths, Phillip and Harris, Joseph “Principles of algebraic geometry”,1978, Wiley Online Library
- [50] Jianpeng Liu, Junwei Liu, and Xi Dai. “Pseudo Landau level representation of twisted bilayer graphene: Band topology and implications on the correlated insulating phase”. *Phys. Rev. B* 99, 155415.
- [51] S. K. Donaldson. ”An application of gauge theory to four-dimensional topology”, *J. Differential Geom.* Volume 18, Number 2 (1983), 279-315, no. 2, 279–315.
- [52] Hofstadter, Douglas R. ”Energy levels and wavefunctions of Bloch electrons in rational and irrational magnetic fields”. *Physical Review B.* 14 (6): 2239–2249.
- [53] F. D. M. Haldane and E. H. Rezayi, “Periodic Laughlin-Jastrow wave functions for the fractional quantized Hall effect,” *Phys. Rev. B*, vol. 31, pp. 2529–2531, Feb 1985.
- [54] R.C. Gunning, ”Lectures on Vector Bundles on Riemann surfaces”, Princeton University Press