

On thermal correlators and bosonization duality in Chern-Simons theories with massive fundamental matter

Amiya Mishra

Department of Theoretical Physics,

Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400005, India

E-mail: amiya.mishra@tifr.res.in

ABSTRACT: We consider Chern-Simons theory coupled to massive fundamental matter in three spacetime dimensions at finite temperature, in the large N limit. We compute several thermal correlators in this theory for both fermionic and bosonic matter separately. The results are computed in the large N 't Hooft limit but for arbitrary values of the 't Hooft coupling. Furthermore, we generalize the computations of the four-point function of fundamental scalars in the bosonic theory to finite temperature. As a consistency check, we see that the results obtained here agree with the existing previous results in different limiting cases. Moreover, we check that the results are consistent with the conjectured bosonization duality, providing an additional evidence of it.

Contents

1	Introduction	1
2	A note on holonomy and related conventions	4
3	Thermal correlators in the fermionic theory	6
3.1	Brief review of the theory	6
3.2	Case I: Spin zero	8
3.2.1	Schwinger-Dyson equation for the exact vertex	8
3.2.2	Two-point function :	11
3.3	Case II: Spin one	12
3.3.1	Schwinger-Dyson equation for the exact insertion vertex	13
3.3.2	Two-point function	14
4	Thermal correlators in the bosonic theory	15
4.1	Brief review of the theory	15
4.2	Thermal four-point function of fundamental scalars	17
4.3	Case I: Spin zero	18
4.3.1	Exact insertion vertex	18
4.3.2	Two-point function	19
4.4	Case II: Spin one	20
4.4.1	Exact insertion vertex	20
4.4.2	Two-point function	21
5	Analysis of results	23
5.1	Fermionic result	23
5.1.1	Two-point correlator of spin 0 operator :	24
5.1.2	Two-point correlator of spin 1 current :	25
5.2	Bosonic result	27
5.2.1	Two-point correlator spin 0 operator :	27
5.2.2	Critical theory limit	28
5.2.3	Two-point correlator of spin 1 current :	30
6	Check of duality	31
6.1	Duality map :	32
6.2	For spin 0 :	32
6.2.1	Duality check in the unhiggsed phase of critical bosons:	33
6.2.2	Prediction for the higgsed phase of critical bosons:	33
6.3	For spin 1 :	34
6.3.1	Duality check in the unhiggsed phase of critical bosons :	34
6.3.2	Prediction for the higgsed phase of critical bosons :	34

7	Discussions	34
A	Conventions and useful definitions	37
B	Angular integrals :	38
C	Useful integrals	38
C.1	Integrals in bosonic theory	38
C.2	Integrals in fermionic Theory	40
D	Computation of thermal four-point function of fundamental scalars	41
D.1	Schwinger-Dyson Equation	43
E	Comment on arbitrary spin s current two-point function	47
E.1	Fermionic theory	48
E.2	Bosonic theory	50

Contents

1 Introduction

There is now considerable evidence that bosonic matter coupled to Chern-Simons (CS) gauge theories and fermionic matter coupled to - roughly speaking - ‘level-rank dual’ CS gauge theories are dual to each other in three spacetime dimensions [1–85]¹. More specifically, $SU(N_B)$ CS gauge fields coupled to bosonic matter is level-rank dual to $U(N_F)$ CS gauge fields coupled to the fermionic matter (in the strict large N limit - in which we will be working in - the difference between $SU(N)$ and $U(N)$ effectively disappears)². This is an example of strong-weak coupling duality. One of the exciting and interesting fact about these theories is that, in the ’t Hooft large N limit, exact computations of observables can be performed on both sides of the dual pair of theories and the duality can be explicitly checked in this limit.

Previously there have been numerous computations and checks of this duality in the case of both the massless and massive matter theories at zero temperature. Even the computations have already been performed at finite temperature to compute the thermal

¹There are two pair of conjecturally dual theories in non-supersymmetric matter coupled Chern-Simons theories. One is regular fermion and critical boson which are together called quasi fermionic theories. The other pair is regular boson and critical fermions which are together called quasi bosonic theories. For details about these see e.g. in [3, 4, 10, 11, 17, 27, 62, 67, 68, 83].

²In the dimensional regulation scheme, the Chern-Simons levels are renormalized. In the large N limit, the renormalized levels and ranks of the two gauge groups are related as $\kappa_F = -\kappa_B$ and $N_F = |\kappa_B| - N_B$ (see for details e.g. in [17, 27, 62, 67, 68, 83]). For a precise form of conjectured duality see appendix A of [83].

free energy of these theories and it has already been shown that the thermal free energies in both the fermionic theories and the bosonic theories map to each other under duality [10, 11, 62, 68, 83]. These theories also admit an infinite set of higher spin currents (spin ≥ 1) of single trace operators [1, 3, 4, 14]. These theories have a global $U(1)$ symmetry. The spin one operator is identified as the corresponding $U(1)$ conserved current. The spin two operator is interpreted as the stress tensor of the corresponding theories. Apart from these, these theories also contain a single trace, gauge invariant scalar ‘current’ operator which we refer to as spin-zero operator. Previously, there have been several computations of various correlation functions of these operators in different limiting cases. For example, the correlation functions of single trace operators of various spins were computed in the case of massless fundamental matter (for both fermions and bosons) coupled to Chern-Simons theory at zero temperature (see, e.g. in [7, 9, 65, 79]). In [29, 32], the two-point functions of spin-one current is computed in the massive fermionic matter theory coupled to Chern-Simons gauge fields at zero temperature. At finite temperature, the effects of holonomy of gauge fields - which is basically the zero-mode of the gauge field along the thermal circle - become important and one has to take into account that as well [10, 11]. In the large- N limit, the effect of holonomy can be described by a continuous distribution function. There have been few computations of the current two-point functions in the massless fundamental matter theories at non-zero temperature with a specific holonomy distribution. For example, in [32], the authors first computed the two-point function of $U(1)$ conserved current in the massless fermionic matter theories at finite temperature in the context of studying the Hall conductivity. In their analysis, however, they considered only a particular holonomy distribution of gauge field, i.e., the universal table-top distribution and the final results of that paper was in terms of some integral form. In a recent paper [80] also, the authors have computed two-point functions of several current operators in the massless fermionic and massless bosonic matter theories at non-zero temperature. However, they had chosen a particular (table-top) holonomy distribution which is appropriate for infinite volume limit. In this work, we generalize the computations of various current correlators to the general case of the massive matter at finite temperature considering an arbitrary holonomy distribution for the gauge holonomy, in Chern-Simons coupled to fundamental matter theories³. We consider massive fermionic and bosonic matter theories separately at non-zero temperature. There are two types of these theories: one is the regular matter theories and the other is the critical matter theories. The regular matter theories on one side of the duality is conjectured to be dual to the critical matter on the other side and vice versa. It was discussed in [7, 9] and also pointed out in [80] that the two-point correlators of various spin s operators - for spin $s \geq 1$ - are basically the same whether we consider the regular or critical matter theories, except the difference being implicitly through the exact masses of the fundamental excitations of fermion/boson. The two-point function of spin-zero, single trace scalar operator is different whether one considers the regular matter or critical matter; and it was discussed in [7, 9] that they are related to each other in a particular way, at least

³We have been able to solve the resultant Schwinger-Dyson equations explicitly by ‘effectively performing’ the loop momenta integrals. From this point view, this work can also be thought of as the study of solving an interesting ‘mathematical problem’.

in the massless theory. To be specific, we consider here the massive regular fermionic matter theory coupled to $U(N_F)$ Chern-Simons gauge fields. For convenience of the computation, we also consider the massive regular bosonic matter theory coupled to $SU(N_B)$ Chern-Simons gauge fields. We choose to work in the dimensional regulation scheme in which the Chern-Simons level is given by the renormalized parameter κ . We work in the 't Hooft large- N limit, which is by taking $N, \kappa \rightarrow \infty$ but keeping $\lambda = \frac{N}{\kappa}$ fixed, and the modulus of the 't Hooft coupling λ is less than unity. One of our goals in this paper is to check the conjectured bosonization duality. To do that we take the critical limit of the regular bosonic theory [7, 9, 17] and check the duality. For the computation of bosonic correlators, we take a slightly different route than [80]. Following [7], in the case of massive bosonic theories at finite temperature, we first compute the offshell thermal four-point functions of fundamental scalars; we do so by generalizing the results of [7, 17] to include the finite temperature effects. The offshell four-point function of scalars is then used to compute the various current correlators. In the massive case, even at zero temperature, there are two phases of bosonic matter theories: one is unhiggsed phases of scalars and the other is the higgsed phases of W and Z bosons (for details on the Higgsed phases of bosonic matter coupled to Chern-Simons theories, see e.g. in [62, 68]). In the critical boson theory with a bare mass parameter m_B^{cri} , there are two possible signs. The unhiggsed phase of bosonic scalar corresponds to the case $m_B^{\text{cri}} > 0$ which under bose-fermi duality gets mapped to the regular fermionic matter theory with $\text{sgn}(m_F \lambda_F) = +1$ ⁴, where, m_F is the bare mass parameter of the regular fermionic theory, and $\lambda_F = \frac{N_F}{\kappa_F}$ is 't Hooft coupling. We have explicitly checked the bose-fermi duality between the current correlators in these phases. However, for the fermionic theories, the results are valid for both possible signs of m_F . The other possible case, i.e., the case $\text{sgn}(m_F \lambda_F) = -1$ in the fermionic theory is dual to the higgsed phases of the bosonic theory (which corresponds to the case $m_B^{\text{cri}} < 0$). We use the results of current correlators in the fermionic theory to predict the corresponding results in the Higgsed phases of bosonic matter. We leave the exercise of explicit check of duality between the current correlators by exact computations in this phase for future work.

The organization of this paper is as follows. In section 2, we briefly review and discuss about the effect of holonomy in Chern-Simons matter theories at finite temperature. In section 3, we study the fermionic matter coupled Chern-Simons theory and present the computations of thermal two-point functions of spin-zero and spin-one current operators. In section 4, we study the bosonic matter coupled Chern-Simons theory and present the results of the offshell four-point function of fundamental scalars in the bosonic scalar theory at finite temperature (the details of which is presented in the appendix D) and also present the computations of thermal two-point functions of spin-zero and spin-one current operators. In section 5, we analyze the main results of this paper in various special limits and see that it agrees with the existing results in the limiting cases. In section 6, we explicitly check that

⁴At finite temperature, this condition is replaced by a more general condition $\text{sgn}(h_F \lambda_F) = +1$, where, $\text{sgn}(h_F)$ is defined around (3.10). The regular fermionic theory at nonzero temperature with $\text{sgn}(h_F \lambda_F) = +1$ is dual to the unhiggsed phases of critical boson theory. The other case, i.e., the case $\text{sgn}(h_F \lambda_F) = -1$ is dual to the higgsed phases of critical boson theory. At zero temperature, $\text{sgn}(h_F)$ is replaced by $\text{sgn}(m_F)$. For details about this see e.g., in [62], where $\text{sgn}(h_F)$ is labelled by a different symbol $\text{sgn}(X_F)$.

the results obtained in this paper are consistent with the conjectured bose-fermi duality. In section 7, we draw conclusions about our results, and discuss various outlook and interesting future directions.

2 A note on holonomy and related conventions

In an Euclidean thermal field theory at some temperature T in three dimensions, x^3 direction is put along a circle \mathbf{S}^1 of radius $\beta = T^{-1}$ ⁵. The position space integrals that we encounter here, are given by

$$\int d^3x f(x) = \int d^2\vec{x} \int_0^\beta dx^3 f(\vec{x}, x^3) . \quad (2.1)$$

In Chern-Simons matter theory at finite temperature, holonomy, i.e., the zero mode of the gauge field along the thermal circle, becomes important and crucially effects the physical observables [10, 11]. Holonomy of the gauge field in a $U(N)$ gauge theory is completely specified by its eigenvalues, $e^{i\alpha_j}$ where $j = 1, \dots, N$ and $\alpha_j \in (-\pi, \pi]$ [10, 62, 83]. In the large- N limit, the location of eigenvalues on unit circle can be specified by a continuous distribution function $\rho(\alpha)$ defined by

$$\rho(\alpha) = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{j=1}^N \delta(\alpha - \alpha_j) . \quad (2.2)$$

It follows that the holonomy distribution function is normalized, i.e., $\int_{-\pi}^\pi \rho(\alpha) d\alpha = 1$. We choose the holonomy distribution $\rho(\alpha)$ to be arbitrary, throughout the paper, unless otherwise mentioned⁶. The correlators that we compute in this paper, by summing Feynman diagrams and performing loop integrals, are in momentum space. Technically, the effect of non-trivial holonomy is to shift the loop momenta in the direction of k_3 for a generic loop momentum $k \equiv (k_1, k_2, k_3)$ [10, 32, 62, 80]. Let us briefly explain this point. As x^3 -direction is compactified in the thermal theory, the conjugate momenta along the third direction is quantized. We will not, however, explicitly write every time the quantized version of k_3 but will implicitly assume that is the case. The momentum space integrals are given by

$$\int \frac{\mathcal{D}^3k}{(2\pi)^3} f(k) = \int \frac{d^2\vec{k}}{(2\pi)^2} \int \frac{\mathcal{D}k_3}{2\pi} f(\vec{k}, k_3) , \quad (2.3)$$

where, $d^2\vec{k}$ is the usual integration measure $dk_1 dk_2$ for the spatial momenta, and the momentum integration measure $\mathcal{D}k_3$ is defined by

$$\int \frac{\mathcal{D}k_3}{2\pi} f(\vec{k}, k_3) = \frac{1}{\beta} \int_{-\pi}^\pi \rho(\alpha) d\alpha \sum_{k_3} f\left(\vec{k}, k_3 + \frac{\alpha}{\beta}\right) , \quad (2.4)$$

Here, as mentioned earlier, the effect of holonomy is to shift the k_3 momenta by $\beta^{-1}\alpha$. Exact meaning of ‘integral over k_3 ’ which is really a sum [62], is slightly different for bosons

⁵In other words, the identification $x^3 \sim x^3 + \beta$ is used; here, x^3 is the Euclidean time coordinate.

⁶Throughout, we assume the holonomy distribution $\rho(\alpha)$ to be an even function of α , i.e., $\rho(-\alpha) = \rho(\alpha)$.

and fermions depending upon the periodic/antiperiodic boundary conditions on bosonic and fermionic fields along \mathbf{S}^1 , the precise meaning of which is explained below in (2.6) and (2.8). To see the precise form of the quantization condition on the momenta k_3 , we consider the fourier transform of a function $f(x) \equiv f(\vec{x}, x^3)$ which is defined by [62]

$$f(\vec{x}, x^3) = \int \frac{\mathcal{D}^3 k}{(2\pi)^3} e^{i\vec{k} \cdot \vec{x} + i k_3 x^3} f(\vec{k}, k_3). \quad (2.5)$$

Depending upon the periodicity/antiperiodicity of the boundary conditions on the fields along the thermal circle, there are two possible cases.

Bosonic Theory :

Bosonic fields are periodic along the thermal circle, i.e., $\phi(\vec{x}, x^3 + \beta) = \phi(\vec{x}, x^3)$. This implies $e^{i k_3 \beta} = e^{2 n_k \pi i}$, where $n_k \in \mathbb{Z}$; i.e., k_3 is quantized with the quantization condition $k_3 = \frac{2 n_k \pi}{\beta}$. It follows from (2.3) and (2.4) that the momentum integral with quantized k_3 in the bosonic theory for an arbitrary holonomy distribution $\rho_B(\alpha)$ is given by

$$\int \frac{\mathcal{D}^3 k}{(2\pi)^3} f(\vec{k}, k_3) = \int \frac{d^2 \vec{k}}{(2\pi)^2} \int \frac{\mathcal{D}_B k_3}{2\pi} f(\vec{k}, k_3) = \frac{1}{\beta} \int \frac{d^2 \vec{k}}{(2\pi)^2} \int_{-\pi}^{\pi} \rho_B(\alpha) d\alpha \sum_{n_k \in \mathbb{Z}} f\left(\vec{k}, \frac{2 n_k \pi + \alpha}{\beta}\right). \quad (2.6)$$

For later use, we here define the following function $\chi_B(z)$ of a real variable z

$$\chi_B(z) = \frac{1}{2} \int_{-\pi}^{\pi} \rho_B(\alpha) d\alpha \left[\coth\left(\frac{\beta z + i\alpha}{2}\right) + \coth\left(\frac{\beta z - i\alpha}{2}\right) \right]. \quad (2.7)$$

Following [62], we also define another function $\xi_B(z)$ as $\xi_B(z) = \int^z \chi_B(w) dw$, an (indefinite) integral over $\chi_B(z)$. The explicit form of the function $\xi_B(z)$ is given by (A.1) in the appendix A.

Fermionic Theory :

Fermion fields on the other hand satisfy the anti-periodicity boundary condition along the thermal circle, i.e., $\psi(\vec{x}, x^3 + \beta) = -\psi(\vec{x}, x^3)$, which implies $e^{i k_3 \beta} = e^{i(2 n_k + 1)\pi}$, where $n_k \in \mathbb{Z}$. This means k_3 in the fermionic theory is quantized with the quantization condition $k_3 = \frac{(2 n_k + 1)\pi}{\beta}$. So, the momentum integral with quantized k_3 in the fermionic theory for an arbitrary holonomy distribution $\rho_F(\alpha)$ is given by

$$\int \frac{\mathcal{D}_F^3 k}{(2\pi)^3} f(\vec{k}, k_3) = \int \frac{d^2 \vec{k}}{(2\pi)^2} \int \frac{\mathcal{D}_F k_3}{2\pi} f(\vec{k}, k_3) = \frac{1}{\beta} \int \frac{d^2 \vec{k}}{(2\pi)^2} \int_{-\pi}^{\pi} \rho_F(\alpha) d\alpha \sum_{n_k \in \mathbb{Z}} f\left(\vec{k}, \frac{(2 n_k + 1)\pi + \alpha}{\beta}\right). \quad (2.8)$$

As above in bosonic theory, we define here the following function $\chi_F(z)$

$$\chi_F(z) = \frac{1}{2} \int_{-\pi}^{\pi} \rho_F(\alpha) d\alpha \left[\tanh\left(\frac{\beta z + i\alpha}{2}\right) + \tanh\left(\frac{\beta z - i\alpha}{2}\right) \right], \quad (2.9)$$

which will be used later in the paper. As in the case of bosons, we define another function $\xi_F(z)$ as an integral over $\chi_F(z)$, i.e., as $\xi_F(z) = \int^z \chi_F(w) dw$. The explicit form of $\xi_F(z)$ is given by (A.2). For other conventions and useful definitions see Appendix A.

3 Thermal correlators in the fermionic theory

3.1 Brief review of the theory

As discussed in the introduction, in this paper we study the massive, regular fermionic matter theory with a finite bare mass m_F coupled to $U(N_F)$ Chern-Simons gauge fields at finite temperature in the large- N limit. In the dimensional regulation scheme, the Euclidean action of this theory is given by

$$\mathcal{S}_F[A, \psi] = \frac{i\kappa_F}{4\pi} \int d^3x \epsilon^{\mu\nu\rho} \text{tr} \left(A_\mu \partial_\nu A_\rho - \frac{2i}{3} A_\mu A_\nu A_\rho \right) + \int d^3x (\bar{\psi} \gamma^\mu D_\mu \psi + m_F \bar{\psi} \psi) , \quad (3.1)$$

where, covariant derivatives for fundamental and antifundamental fields are defined by, $D_\mu \psi = \partial_\mu \psi - iA_\mu \psi$ and $D_\mu \bar{\psi} = \partial_\mu \bar{\psi} + i\bar{\psi} A_\mu$, respectively⁷. The gauge field A_μ is in the adjoint representation of $U(N_F)$. The gamma matrices are chosen to be the ordinary 2×2 Pauli matrices $\gamma^\mu = \sigma^\mu$, $\mu = 1, 2, 3$. Following the literature, we work in the gauge $A_- = 0$. With this choice of gauge, the action (3.1) in momentum space takes the following form⁸

$$\begin{aligned} \mathcal{S}_F[A, \psi] = & - \frac{\kappa_F \epsilon^{\mu-\nu}}{4\pi} \int \frac{\mathcal{D}_F^3 k}{(2\pi)^3} A_\mu(-k) k_- A_\nu(k) + \int \frac{\mathcal{D}_F^3 k}{(2\pi)^3} \bar{\psi}(-k) (i\gamma^\mu k_\mu + m_F) \psi(k) \\ & - i \int \frac{\mathcal{D}_F^3 p}{(2\pi)^3} \frac{\mathcal{D}_F^3 k}{(2\pi)^3} \bar{\psi}(p) \gamma^\mu A_\mu(-p-k) \psi(k) . \end{aligned} \quad (3.2)$$

In the momentum space, some of the Feynman rules for this theory are given below.

Propagator for the gauge field :

The propagator for the gauge field is defined as

$$\langle A_\mu(p') A_\nu(p) \rangle = G_{\nu\mu}^F(p) (2\pi)^3 \delta^{(3)}(p + p') , \quad (3.3)$$

where,

$$G_{\nu\mu}^F(p) = \frac{2\pi \epsilon_{\nu-\mu}}{\kappa_F p_-} . \quad (3.4)$$

As discussed in [10], the gauge field propagator is independent of p_3 , so is not effected by the holonomy.

⁷A note on the notations and conventions: we don't explicitly write the color indices, but the color contractions are easily understood from the context. ψ and $\bar{\psi}$ can be thought of as N_F component column and row vectors, respectively and A_μ can be thought of as $N_F \times N_F$ matrices in the color space. We don't show explicitly the spinor indices of ψ and $\bar{\psi}$ which are two-component spinors in three dimensions. Also, we don't explicitly write the adjoint index of the gauge fields $A_\mu \equiv A_\mu^a T^a$, and color trace over the gauge group generators T^a . One can easily restore these indices and explicitly use the normalizations $\text{Tr}(T^a T^b) = C(N_F) \delta^{ab}$. Conventionally, $C(N_F) = \frac{1}{2}$. The convention for the color contraction that is used here, is such that $\bar{\psi} \psi = \bar{\psi}_i \psi^i$; but $\psi \bar{\psi}$ does not necessarily have the contracted color indices. Also, $\bar{\psi} M \psi = \bar{\psi}_i M^i_j \psi^j$. The same convention is used for the bosonic theories where N_F is replaced by N_B .

⁸Normalization for the Levi-Civita tensor used in this paper is such that $\epsilon^{123} = \epsilon_{123} = 1$ and $\epsilon^{+-3} = -\epsilon_{+-3} = -i$.

Exact propagator for the Fermionic field :

The exact fermionic propagator, in the strict large- N limit, to all orders in the 't Hooft coupling parameter λ_F ⁹, is given by

$$\langle \psi(p) \bar{\psi}(p') \rangle = S_F(p) (2\pi)^3 \delta^{(3)}(p + p') , \quad (3.5)$$

where¹⁰,

$$S_F(p) = \frac{1}{i\gamma^\mu p_\mu + m_F \mathbb{1} + \Sigma_F(p)} . \quad (3.7)$$

Here, $\Sigma_F(p)$ is the fermionic self energy. It is useful to expand $\Sigma_F(p)$ in the basis $\{\gamma^\mu, \mathbb{1}\}$ as $\Sigma_F = i\gamma^\mu \Sigma_\mu + \Sigma_{\mathbb{1}} \mathbb{1}$, where, $\mathbb{1}$ is 2×2 identity matrix. At non-zero temperature, the self energy was already computed in the literature (see, e.g. in [1, 10, 11] for details) in the lightcone gauge, and is listed here for the purpose of later use¹¹

$$\tilde{\Sigma}_{\mathbb{1}}(p_s) = m_F + \lambda_F \xi_F(a(p_s)) , \quad \Sigma_+(\vec{p}) = \frac{p_+}{p_s^2} [c_F^2 - \tilde{\Sigma}_{\mathbb{1}}^2(p)] , \quad \Sigma_-(p) = \Sigma_3(p) = 0 , \quad (3.9)$$

where, $\tilde{\Sigma}_{\mathbb{1}} = m_F + \Sigma_{\mathbb{1}}$, $a(p_s) = +\sqrt{p_s^2 + c_F^2}$ and $p^2 = 2p_+p_- + p_3^2 = p_s^2 + p_3^2$. The function $\xi_F(z)$ is defined in (A.2). c_F is the thermal mass of the fermion which is determined by the gap equation

$$c_F = \text{sgn}(h_F)(m_F + \lambda_F \xi_F(c_F)) , \quad (3.10)$$

where, $\text{sgn}(h_F) = \text{sgn}(m_F + \lambda_F \xi_F(c_F))$ (see e.g. in [62]). We use the convention where, c_F (also c_B in the case of bosons) is always positive. There are two possibilities depending upon the two possible signs of $\text{sgn}(h_F \lambda_F)$ [62]¹². At zero temperature, $\text{sgn}(h_F)$ is replaced by $\text{sgn}(m_F)$. Diagrammatically, the gauge field and the exact fermionic propagators are

$$\mu \xrightarrow{p} \nu = G_{\nu\mu}^F(p) \qquad \xrightarrow{p} \bullet = S_F(p)$$

The vertex factor associated with the interaction term $\bar{\psi}(p) A_\mu(-p-k) \psi(k)$ is given by

$$\mathcal{V}^\mu(k, p) = i\gamma^\mu . \quad (3.11)$$

We now compute various correlators of gauge invariant, single trace operators in this theory.

⁹As discussed in the introduction, the 't Hooft coupling parameter for the fermionic theory is defined by $\lambda_F = \frac{N_F}{\kappa_F}$; similarly for the bosonic theory, $\lambda_B = \frac{N_B}{\kappa_B}$. The range of the couplings are $0 < |\lambda| < 1$, with $\lambda = 0$ being the weak coupling limit and $\lambda = 1$ being the strong-coupling limit of the corresponding theories.

¹⁰Alternatively, the exact propagator can be written as

$$S_F(p) = \frac{\tilde{\Sigma}_{\mathbb{1}}(p) \mathbb{1} - i\gamma^\mu (p_\mu + \Sigma_\mu)}{p^2 + c_F^2} \equiv S_\mu^F(p) \gamma^\mu + S_{\mathbb{1}}^F(p) \mathbb{1} . \quad (3.6)$$

¹¹The fermionic self energy is obtained by summing the $1PI$ graphs (see, e.g., Figure 5 in [10]) and is given by [1, 10, 11]

$$\Sigma_F(p) = -N_F \int \frac{\mathcal{D}_F^3 \ell}{(2\pi)^3} \mathcal{V}^\mu(\ell, p) S_F(\ell) \mathcal{V}^\nu(p, \ell) G_{\nu\mu}^F(\ell - p) . \quad (3.8)$$

¹²The choice $\text{sgn}(h_F \lambda_F) = +1$ corresponds to, under duality, the unHiggsed critical bosonic scalar theory, and the choice $\text{sgn}(h_F \lambda_F) = -1$ corresponds to the Higgsed phases of critical bosons under duality.

3.2 Case I: Spin zero

The gauge invariant, single trace, spin 0 operator in the regular fermionic theory is given by $J_0^F(x) = \bar{\psi}(x)\psi(x)$, which in the momentum space takes the form

$$J_0^F(-q) = \int \frac{\mathcal{D}_F^3 k}{(2\pi)^3} \bar{\psi}(-(k+q))\psi(k) \quad (3.12)$$

In order to compute $\langle J_0^F J_0^F \rangle$ two-point correlator, we need to compute the exact J_0^F insertion vertex, which we compute by solving the corresponding Schwinger-Dyson equations shown in Fig.1 in the subsection below.

3.2.1 Schwinger-Dyson equation for the exact vertex

Diagrammatically, the Schwinger-Dyson equation for the exact J_0^F vertex is shown in Fig.1. The exact J_0^F insertion vertex is defined by

$$\langle J_0^F(-q)\psi(k)\bar{\psi}(p) \rangle = V_0^F(k, q) (2\pi)^3 \delta^{(3)}(p+k+q) . \quad (3.13)$$

In the planar limit, the Schwinger-Dyson equation for $V_0^F(k, q)$ is given by




Figure 1. Schwinger-Dyson equations for the exact vertices. The circled cross denote an insertion of an exact vertex and a filled circle denotes the exact fermion propagator. The bare cross denotes the ‘tree level’ insertion.

$$V_0^F(k, q) = \tilde{V}_0^F(k, q) + N_F \int \frac{\mathcal{D}_F^3 \ell}{(2\pi)^3} [\gamma^\mu(\ell+q, k+q) S_F(\ell+q) V_0^F(\ell, q) S_F(\ell) \mathcal{V}^\nu(k, \ell)] G_{\nu\mu}^F(\ell-k) , \quad (3.14)$$

where, the factor of N_F appearing in front of the second term is the contribution from the color factors. \tilde{V}_0^F is the tree-level insertion vertex corresponding to J_0^F . From the definition of J_0^F operator in (3.12), it follows that $\tilde{V}_0^F(k, q) = 1$. We work in the ‘lightcone kinematics’ $q_\pm = 0$, in which case, the calculations simplify considerably¹³. Using (3.11) and (3.4), (3.14) simplifies to

$$V_0^F(k, q) = 1 - 2\pi i \lambda_F \int \frac{\mathcal{D}_F^3 \ell}{(2\pi)^3} [\gamma^{[3]} S_F(\ell+q) V_0^F(\ell, q) S_F(\ell) \gamma^{+] }] \frac{1}{(\ell-k)_-} . \quad (3.15)$$

¹³The difficulty with working in the general case of $q_\pm \neq 0$ lies in the fact that the corresponding integrals over the spatial components ℓ_1 and ℓ_2 of the loop momenta ℓ are difficult to perform exactly.

Here and in the rest of the paper, we use the notation

$$\gamma^{[\rho]} A \gamma^{|\nu]} = \gamma^\rho A \gamma^\nu - \gamma^\nu A \gamma^\rho = 2i\epsilon^{\rho\mu\nu} (A_\mu \mathbb{1} - A_\mathbb{1} \gamma_\mu) , \quad (3.16)$$

for any 2×2 matrix $A = A_\mu \gamma^\mu + A_\mathbb{1} \mathbb{1}$. As $V_0^F(k, q)$ is a 2×2 matrix in the spinor space, it can be expanded in the complete basis $\{\gamma^\mu, \mathbb{1}\}$ of 2×2 matrices as $V_0^F(k, q) = V_{0,\mu}(k, q) \gamma^\mu + V_{0,\mathbb{1}} \mathbb{1}$. Comparing it with the RHS of (3.15), we get $V_{0,-}(k, q) = V_{0,3}(k, q) = 0$ and the following set of two non-trivial coupled integral equations

$$\begin{aligned} V_{0,+}(k, q) &= -4\pi i \lambda_F \int \frac{\mathcal{D}_F^3 \ell}{(2\pi)^3} [S_F(\ell + q) V_0^F(\ell, q) S_F(\ell)]_\mathbb{1} \frac{1}{(\ell - k)_-} , \\ V_{0,\mathbb{1}}(k, q) &= 1 + 4\pi i \lambda_F \int \frac{\mathcal{D}_F^3 \ell}{(2\pi)^3} [S_F(\ell + q) V_0^F(\ell, q) S_F(\ell)]_- \frac{1}{(\ell - k)_-} , \end{aligned} \quad (3.17)$$

where, we have used the notation $[ABC]_a$ to denote the a -th component of the matrix product ABC of three matrices A , B and C , and $a \in (\mu, \mathbb{1})$. The RHS of the above equation (3.17) does not depend upon k_3 . In the ‘lightcone kinematics’ $q_\pm = 0$, the only non-zero component of q is q_3 . So, we use the $SO(2)$ rotational symmetry in the 1-2 plane to write down

$$V_{0,\mathbb{1}}(\vec{\ell}, q_3) \equiv f(\ell_s, q_3) , \quad V_{0,+}(\vec{\ell}, q_3) \equiv \frac{\ell_+}{\ell_s^2} g(\ell_s, q_3) . \quad (3.18)$$

The extra factor of ℓ_s^2 in the denominator of the second expression above is kept for later convenience. Using (3.18) and the explicit components of the exact fermion propagator (3.6), the equations (3.17) can be simplified to give

$$f(k_s, q_3) = 1 - 4\pi i \lambda_F \int \frac{\mathcal{D}_F^3 \ell}{(2\pi)^3} \frac{[-q_3 + 2i\tilde{\Sigma}_\mathbb{1}(\ell_s)]f(\ell_s, q_3) + g(\ell_s, q_3)}{(\ell_3^2 + a^2(\ell_s))((\ell_3 + q_3)^2 + a^2(\ell_s))} \frac{\ell_-}{(\ell - k)_-} , \quad (3.19)$$

and

$$\frac{k_+}{k_s^2} g(k_s, q_3) = 4\pi i \lambda_F \int \frac{\mathcal{D}_F^3 \ell}{(2\pi)^3} \frac{[\ell_3(\ell_3 + q_3) + a^2(\ell_3) - 2\tilde{\Sigma}_\mathbb{1}^2(\ell_s)]f(\ell_s, q_3) + \frac{1}{2}[q_3 + 2i\tilde{\Sigma}_\mathbb{1}(\ell_s)]g(\ell_s, q_3)}{(\ell_3^2 + a^2(\ell_s))((\ell_3 + q_3)^2 + a^2(\ell_s))} \frac{1}{(\ell - k)_-} . \quad (3.20)$$

Due to the $SO(2)$ rotational symmetry in the ℓ_1 - ℓ_2 plane, it is useful to write the integration measure as

$$\mathcal{D}_F^3 \ell \equiv \ell_s d\ell_s d\theta_\ell \mathcal{D}_F \ell_3 . \quad (3.21)$$

where, ℓ_s is the radial momenta in the ℓ_+ - ℓ_+ plane (or equivalently, ℓ_1 - ℓ_2 plane) and is given by $\ell_s^2 = 2\ell_+ \ell_- = \ell_1^2 + \ell_2^2$. θ_ℓ is the angular variable in the lightcone plane. The measure $\mathcal{D}_F \ell_3$ as before is given by (2.8). One can do the angular integration by using the result (B.1) and also perform the integral over the momentum component ℓ_3 by using (C.11) and (C.13); by doing so, we find

$$f(k_s, q_3) = 1 - 2i\lambda_F \int_{k_s}^\infty \frac{\ell_s d\ell_s}{a(\ell_s)} F_F(a(\ell_s), q_3) [[-q_3 + 2i\tilde{\Sigma}_\mathbb{1}(\ell_s)] f(\ell_s, q_3) + g(\ell_s, q_3)] , \quad (3.22)$$

and

$$g(k_s, q_3) = -2i\lambda_F \int_0^{k_s} \frac{\ell_s d\ell_s}{a(\ell_s)} F_F(a(\ell_s), q_3) [4[a^2(\ell_s) - \tilde{\Sigma}_\mathbb{1}^2(\ell_s)] f(\ell_s, q_3) + [q_3 + 2i\tilde{\Sigma}_\mathbb{1}(\ell_s)] g(\ell_s, q_3)] . \quad (3.23)$$

where, the function $F_F(z)$ is defined by $F_F(z) = \frac{\chi_F(z)}{q_3^2 + 4z^2}$, and $\chi_F(z)$ is given by (2.9). In order to solve these two coupled integral equations (3.22) and (3.23), we first introduce a change of variable $a(\ell_s) = +\sqrt{\ell_s^2 + c_F^2} = w$ and $a(k_s) = +\sqrt{k_s^2 + c_F^2} = z$. In terms of these reduced variables (as the functional dependence changes, so, we redefine $f(k_s, q_3) \equiv \tilde{f}(z, q_3)$ and $g(k_s, q_3) \equiv \tilde{g}(z, q_3)$ and $\tilde{\Sigma}_1(\ell_s) \equiv h_F(w)$),

$$\tilde{f}(z, q_3) = 1 - 2i\lambda_F \int_z^\infty dw F_F(w, q_3) [-q_3 + 2ih_F(w)] \tilde{f}(z, q_3) + \tilde{g}(w, q_3) , \quad (3.24)$$

and

$$\tilde{g}(z, q_3) = -2i\lambda_F \int_{c_F}^z dw F_F(w, q_3) [4[w^2 - h_F^2(w)] \tilde{f}(w, q_3) + [q_3 + 2i\tilde{\Sigma}_1(\ell_s)] \tilde{g}(w, q_3)] . \quad (3.25)$$

The above two coupled integral equations (3.24) and (3.25) can be decoupled and solved easily, by converting them first to a set of two differential equations by taking derivatives w.r.t. the free parameter z . The corresponding differential equations take the following form

$$\partial_z \tilde{f}(z, q_3) = 2i\lambda_F [-q_3 + 2ih_F(z)] \tilde{f}(z, q_3) + \tilde{g}(z, q_3) F_F(z, q_3) , \quad (3.26)$$

and

$$\partial_z \tilde{g}(z, q_3) = -2i\lambda_F [4[z^2 - h_F^2(z)] \tilde{f}(z, q_3) + [q_3 + 2ih_F(z)] \tilde{g}(z, q_3)] F_F(z, q_3) . \quad (3.27)$$

There is a particular combination of the two equations (3.26) and (3.27), which can be written as a total derivative w.r.t. the variable z . This is easily done by multiplying (3.26) by $(q_3 + 2ih_F(z))$ and then adding that to (3.27). From the definitions $h_F(z) = m_F + \lambda_F \xi_F(z)$, and $\xi_F(z) = \int^z \chi_F(w) dw$, it follows that $h'_F(z) = \lambda_F \xi'_F(z) = \lambda_F \chi_F(z)$. Constructing the particular combination mentioned in this paragraph, we find that it can be written as a total derivative as given below

$$\partial_z [(q_3 + 2ih_F(z)) \tilde{f}(z, q_3) + \tilde{g}(z, q_3)] = 0 . \quad (3.28)$$

The general solution of (3.28) can be written as

$$(q_3 + 2ih_F(z)) \tilde{f}(z, q_3) + \tilde{g}(z, q_3) = \eta(q_3) , \quad (3.29)$$

where, $\eta(q_3)$ is an unknown function of q_3 to be determined from the boundary conditions. From (3.24) and (3.25), we see that the above differential equations (3.24) and (3.25) must satisfy the following boundary conditions

$$\tilde{f}(z = \Lambda, q_3) = 1, \quad \tilde{g}(z = c_F, q_3) = 0, \quad \text{where, } \Lambda \rightarrow \infty . \quad (3.30)$$

Here and also later in this paper, we have introduced a UV cutoff Λ in the radial momentum integral, in the intermediate steps of calculations to keep track of the divergent terms in the integrals which we have eventually dropped away by regularization. Finally, Λ is taken to infinity. Using (3.29) we substitute $\tilde{g}(z, q_3)$ in (3.24) and get

$$\partial_z \tilde{f}(z, q_3) = 2i\lambda_F [\eta(q_3) - 2q_3 \tilde{f}(z, q_3)] F_F(z, q_3) . \quad (3.31)$$

To solve the above equation (3.31), we choose an ansatz of the form

$$\tilde{f}(z, q_3) = A \left(1 + B \exp \left[4i\lambda_F q_3 \int_z^\Lambda dw F_F(w, q_3) \right] \right). \quad (3.32)$$

Using the definition, $H_F(z, q_3) = \exp \left[4i\lambda_F q_3 \int^z dw F_F(w, q_3) \right]$, (3.32) can be written in the following form

$$\tilde{f}(z, q_3) = A \left(1 + B \frac{H_F(\Lambda, q_3)}{H_F(z, q_3)} \right), \quad (3.33)$$

which is more useful in the intermediate steps of the calculations. There are three unknowns A , B and $\eta(q_3)$ which need to be fixed to completely determine \tilde{f} and \tilde{g} . Using the boundary condition (3.30) on \tilde{f} , we fix A in terms of B . $\eta(q_3)$ is in terms of B by substituting the ansatz (3.32) in (3.31). Finally, we use the boundary condition on \tilde{g} to determine B from equation (3.29). The final result for the exact J_0^F insertion vertex (3.18) is given by

$$\tilde{f}(z, q_3) = \frac{1 + B \frac{H_F(\Lambda, q_3)}{H_F(z, q_3)}}{1 + B}, \quad \tilde{g}(z, q_3) = \frac{2q_3}{1 + B} - (q_3 + 2i(m_F + \lambda_F \xi_F(z))) \tilde{f}(z, q_3), \quad (3.34)$$

where, B is given by

$$B = \frac{(q_3 - 2ih_F(c_F)) H_F(c_F, q_3)}{(q_3 + 2ih_F(c_F)) H_F(\Lambda, q_3)}. \quad (3.35)$$

3.2.2 Two-point function :

The two-point function is computed by computing the Feynman diagram shown in Fig.2. To compute the two-point function $\langle J_0^F(q') J_0^F(q) \rangle$, only a single insertion of an exact vertex $J_0^F(q')$ is required to account for all the perturbative Feynman diagrams without any overcounting. For definiteness, we choose insertion at the left in Fig.2 as the exact vertex.




Figure 2. Diagram contributing to two-point function $\langle J^{(s)}(-q) J^{(s)}(q) \rangle$ for spin zero and spin one. Filled circle denotes the exact fermion propagator.

Insertion on right side in Fig.2 is the ‘free’ insertion vertex ¹⁴ which we define as follows :

$$\langle J_0^F(-q) \psi(k) \bar{\psi}(p) \rangle = U_0^F(k, q) (2\pi)^3 \delta^{(3)}(p + k + q). \quad (3.36)$$

¹⁴We alternately use the terms ‘free’, ‘tree level’ and ‘bare’ insertion vertex for the insertion on the RHS of the schematic diagrams Fig.2 or Fig.5 of two-point correlators. What this exactly means is that it includes only the required diagrams (which may contain loops) to avoid any overcounting [7, 9, 80]. The required diagrams for the ‘bare’ insertion vertex can be got from the definitions of the corresponding current operator.

From the definition of $J_0^F(q)$ operator in (3.12), it follows that $U_0^F(k, q) = 1$. We define the two-point correlation function of J_0^F operator as

$$\langle J_0^F(q') J_0^F(q) \rangle = \mathcal{G}_0^F(q) (2\pi)^3 \delta^{(3)}(q' + q) . \quad (3.37)$$

From Fig.2, the expression for \mathcal{G}_0^F is given by

$$\mathcal{G}_0^F(q) = -N_F \int \frac{\mathcal{D}_F^3 k}{(2\pi)^3} \text{Tr}_F \left[U_0^F(k + q, -q) S_F(k + q) V_0^F(k, q) S_F(k) \right] , \quad (3.38)$$

where, the extra (-1) factor above in (3.38) is because of the integration over fermion loop. Here, Tr_F indicates the trace in spinor space. Inserting the expressions of exact fermion propagator (3.6), the exact insertion vertex V_0^F computed in (3.34), the expression of U_0^F , doing the gamma-matrix algebra and performing the momentum integral over the loop momenta k , the final result for $\langle J_0^F J_0^F \rangle$ that we find is given by

$$\mathcal{G}_0^F(q) = \lim_{\Lambda \rightarrow \infty} \left[\frac{i N_F q_3}{4\pi \lambda_F} \frac{\frac{H_F(\Lambda, q_3)}{H_F(c_F, q_3)} e^{-2i \text{sgn}(h_F) \tan^{-1} \frac{q_3}{2c_F}} + 1}{\frac{H_F(\Lambda, q_3)}{H_F(c_F, q_3)} e^{-2i \text{sgn}(h_F) \tan^{-1} \frac{q_3}{2c_F}} - 1} + \frac{N_F h_F(\Lambda)}{2\pi \lambda_F} \right] , \quad (3.39)$$

where, $H_F(z, q_3) = \exp [4i \lambda_F q_3 \int^z dw F_F(w, q_3)]$, $F_F(z, q_3) = \frac{\chi_F(z)}{q_3^2 + 4z^2}$ and $h_F(z) = m_F + \lambda_F \xi_F(z)$, and $\text{sgn}(h_F)$ is defined in the gap equation (3.10). From the expression of $\xi_F(w)$ in (A.2), it is clear that $\xi_F(\infty)$ diverges linearly. However, we regularize [9, 10] the answer by subtracting the divergent term $\xi_F(\infty)$ and take the $\Lambda \rightarrow \infty$ limit¹⁵. Thus, the renormalized two-point function $\langle J_0^F J_0^F \rangle$ is

$$\mathcal{G}_0^F(q) = \frac{i N_F q_3}{4\pi \lambda_F} \frac{\frac{H_F(\infty, q_3)}{H_F(c_F, q_3)} e^{-2i \text{sgn}(h_F) \tan^{-1} \frac{q_3}{2c_F}} + 1}{\frac{H_F(\infty, q_3)}{H_F(c_F, q_3)} e^{-2i \text{sgn}(h_F) \tan^{-1} \frac{q_3}{2c_F}} - 1} + \frac{N_F m_F}{2\pi \lambda_F} . \quad (3.40)$$

(3.40) is valid at finite temperature and non-zero mass. In various limiting cases, this agrees with the existing results [9, 80, 85].

3.3 Case II: Spin one

In this section we generalize the computations of two-point correlators of spin one operator, done previously [9, 32, 80, 85], to the general case of massive regular fermionic theory at finite temperature with arbitrary holonomy distribution. The regular fermionic theory (3.1) that we study in this paper has a gauge invariant, conserved $U(1)$ current, given by the single trace operator $J_\mu^F(x) = i \bar{\psi}(x) \gamma_\mu \psi(x)$. In momentum space, this operator is given by

$$J_\mu^F(-q) = i \int \frac{\mathcal{D}_F^3 k}{(2\pi)^3} \bar{\psi}(-(k + q)) \gamma_\mu \psi(k) . \quad (3.41)$$

Our goal in this section is to compute the two-point correlator $\langle J_\mu^F J_\nu^F \rangle$, for which as in the case of J_0^F , we need the exact J_μ^F vertex.

¹⁵One can use the dimensional regularization for the radial momenta integrals as discussed in [62] to remove the term $\xi_F(\infty)$. Alternatively, it can be done by adding a mass counterterm for the background source of J_0^F as discussed in [9].

3.3.1 Schwinger-Dyson equation for the exact insertion vertex

Exact J_μ^F insertion vertex is computed by solving the Schwinger-Dyson equation given schematically in Fig.1. The exact J_μ^F vertex is defined as

$$\langle J_\mu^F(-q)\psi(k)\bar{\psi}(p) \rangle = V_{(\mu)}^F(k, q) (2\pi)^3 \delta^{(3)}(p+k+q) . \quad (3.42)$$

The corresponding Schwinger-Dyson equation for $V_\mu^F(k, q)$ is given by

$$V_{(\mu)}^F(k, q) = \tilde{V}_{(\mu)}^F(k, q) + N_F \int \frac{\mathcal{D}_F^3 \ell}{(2\pi)^3} [\mathcal{V}^\nu(\ell+q, k+q) S_F(\ell+q) V_{(\mu)}^F(\ell, q) S_F(\ell) \mathcal{V}^\rho(k, \ell)] G_{\rho\nu}^F(\ell-k) , \quad (3.43)$$

where, $\tilde{V}_{(\mu)}^F(k, q)$ denotes the tree-level insertion of J_μ^F . From (3.41), it follows that $\tilde{V}_{(\mu)}^F(k, q) = i\gamma_\mu$. Using (3.11) and (3.4), the above equation (3.43) can be simplified to

$$V_{(\mu)}^F(k, q) = i\gamma_\mu - 2\pi i \lambda_F \int \frac{\mathcal{D}_F^3 \ell}{(2\pi)^3} [\gamma^{[3]} S_F(\ell+q) V_{(\mu)}^F(\ell, q) S_F(\ell) \gamma^{+]}] \frac{1}{(\ell-k)_-} . \quad (3.44)$$

The only non-zero component of $\langle J_\mu^F(-q) J_\nu^F(q) \rangle$ in the x^1 - x^2 plane, in ‘lightcone kinematics’ $q_\pm = 0$, is the $(\mu\nu) \equiv (-+)$ (or equivalently $(+-)$) component¹⁶. Other components are zero¹⁷. So, from now on, we will be considering only $\langle J_- J_+ \rangle$ (the same argument goes for the bosonic case as well). Also as in the J_0^F case, in order for the computation of two-point functions, only a single exact vertex is required. To compute $\langle J_-^F(-q) J_+^F(q) \rangle$, we will use the exact J_-^F insertion vertex and the ‘tree level’ insertion vertex for J_+^F . So, we need to compute the exact J_-^F insertion vertex. It is clear from (3.44) that $V_{(\mu)}^F(k, q)$ is independent of k_3 . Also, in the lightcone kinematics $q_\pm = 0$, the only non-zero component of the external momenta q is q_3 . So, the momentum dependence of $V_{(\mu)}^F(k, q)$ is basically $V_{(\mu)}^F(\vec{k}, q_3)$. As in the J_0^F case, $V_{(-)}^F(k, q)$ can be expanded as,

$$V_{(-)}^F(\vec{k}, q_3) = V_{(-),\nu}^F(\vec{k}, q_3) \gamma^\nu + V_{(-),\perp}^F(\vec{k}, q_3) \mathbb{1} = g_m(k_s, q_3) \gamma^+ + k_- f_m(k_s, q_3) \mathbb{1} . \quad (3.45)$$

To solve for exact $V_{(-)}^F$, we plug (3.45) in the minus-component equation of (3.44) and get a set of two coupled integral equations involving f_m and g_m . As in the case of J_0^F , these equations for f_m and g_m get simplified a lot once we utilize the $SO(2)$ rotational symmetry in the lightcone plane and decompose the momentum integration measure as (3.21). We perform the angular integration using (B.1) and also perform the integral over momentum component ℓ_3 by using (C.11) and (C.13). Performing a change of variable $a(\ell_s) = +\sqrt{\ell_s^2 + c_F^2} = w$ and $a(k_s) = +\sqrt{k_s^2 + c_F^2} = z$, and relabelling $f_m(k_s, q_3) \equiv \tilde{f}_m(z, q_3)$ and

¹⁶This follows from the $SO(2)$ rotational symmetry in the 1-2 plane.

¹⁷As the $U(1)$ current is classically conserved, it follows from the Ward identity, that in the momentum space, $q^\mu \langle J_\mu(-q) J_\nu(q) \rangle$ should vanish upto (at most) a contact term [80]. With the external momentum choice $q_\pm = 0$, it follows then that the component $\langle J_3 J_3 \rangle$ should vanish upto a contact term. We have also explicitly checked (though not provided here keeping in mind about the length of the paper) in the case of fermionic theory (with the $U(1)$ current given by (3.41)), that this is indeed the case, i.e., $\langle J_- J_- \rangle$, $\langle J_+ J_+ \rangle$, $\langle J_\pm J_3 \rangle$ components vanish in the choice $q_\pm = 0$. Also we have explicitly checked that $\langle J_3 J_3 \rangle$ vanishes exactly in this momentum choice.

$g_m(k_s, q_3) \equiv \tilde{g}_m(z, q_3)$ and $\tilde{\Sigma}_1(\ell_s) \equiv h_F(w)$, the equations for \tilde{f}_m and \tilde{g}_m can be written in a much simpler form, which takes the following form

$$\tilde{f}_m(z, q_3) = 2i\lambda_F \int_z^\infty dw F_F(w, q_3) \left[(q_3 - 2ih_F(w)) \tilde{f}_m(z, q_3) - 2\tilde{g}_m(w, q_3) \right], \quad (3.46)$$

and

$$\tilde{g}_m(z, q_3) = i + 2i\lambda_F \int_z^\infty dw F_F(w, q_3) \left[2(w^2 - h_F^2(w)) \tilde{f}_m(w, q_3) + (q_3 + 2i\tilde{\Sigma}_1(\ell_s)) \tilde{g}_m(w, q_3) \right]. \quad (3.47)$$

To solve the above two equations (3.46) and (3.47), it is best to convert them into a set of differential equations. The boundary conditions that follow from (3.46) and (3.47) are ¹⁸

$$\tilde{f}_m(z = \Lambda, q_3) = 0, \quad \tilde{g}_m(z = \Lambda, q_3) = i, \quad \text{where, } \Lambda \rightarrow \infty. \quad (3.48)$$

Solving the equations (3.46) and (3.47), by first converting them into differential equations with the boundary conditions (3.48), we find the final solution for the exact J_-^F vertex in terms of f_m and g_m as

$$\begin{aligned} \tilde{f}_m(z, q_3) &= \frac{i}{q_3} \left(1 - \exp \left[4i\lambda_F q_3 \int_z^\Lambda dw F_F(w, q_3) \right] \right), \\ \tilde{g}_m(z, q_3) &= i - \frac{1}{2} (q_3 + 2ih_F(z)) \tilde{f}_m(z, q_3). \end{aligned} \quad (3.49)$$

3.3.2 Two-point function

In this subsection, we compute $\langle J_\mu^F(-q) J_\nu^F(q) \rangle$. We define the two-point correlator of spin one operator J_μ^F as

$$\langle J_\mu^F(-q) J_\nu^F(q) \rangle = \mathcal{G}_{\mu\nu}^F(q) (2\pi)^3 \delta^{(3)}(q' + q). \quad (3.50)$$

The corresponding Feynman diagram for this is given in Fig.2. Similar to the J_0^F case, the insertion on left side of the diagram is chosen to be the exact vertex, and the insertion on right side is chosen to be the ‘tree-level’ insertion to avoid any overcounting of the diagrams. We define the ‘tree-level’ insertion J_ν^F vertex as follows

$$\langle J_\nu^F(-q) \psi(k) \bar{\psi}(p) \rangle = U_\nu^F(k, q) (2\pi)^3 \delta^{(3)}(p + k + q). \quad (3.51)$$

From the definition of J_μ^F current in (3.41), it follows that $U_\nu^F(k, q) = i\gamma_\nu$. From Fig.2, we see that the two-point function of the spin one current is given by

$$\mathcal{G}_{\mu\nu}^F(q) = -N_F \int \frac{\mathcal{D}_F^3 k}{(2\pi)^3} \text{Tr}_F \left[S_F(k+q) V_{(\mu)}^F(k, q) S_F(k) U_\nu^F(k+q, -q) \right], \quad (3.52)$$

¹⁸The boundary conditions for the exact vertex $V_{(-)}^F$ is different from the the boundary conditions for that of the exact V_0^F vertex.

where, again the extra (-1) factor above in (3.52) is because of the integration over fermion loop. Using the expression of U_ν^F and performing the gamma matrix algebra¹⁹, (3.52) reduces to

$$\mathcal{G}_{\mu\nu}^F(q) = -2iN_F \int \frac{\mathcal{D}_F^3 k}{(2\pi)^3} [S_F(k+q) V_{(\mu)}^F(k, q) S_F(k)]_\nu. \quad (3.53)$$

Below we consider the $(\mu\nu) \equiv (-+)$ component of (3.53). Using the expressions of S_F and V_-^F in (3.53) and performing the momentum integral, we find

$$\mathcal{G}_{-+}^F(q) = \lim_{\Lambda \rightarrow \infty} \left[\frac{iN_F q_3}{16\pi\lambda_F} \left(1 + \frac{2ih_F(c_F)}{q_3} \right)^2 \left[\frac{H_F(\Lambda, q_3)}{H_F(c_F, q_3)} - 1 \right] - \frac{N_F \xi_F(c_F)}{8\pi} + \frac{N_F \xi_F(\Lambda)}{8\pi} \right] \quad (3.54)$$

As in the case of J_0^F , (3.54) is linearly divergent due to the appearance of $\xi_F(\infty)$. Regularizing the above answer by throwing away the term $\xi_F(\infty)$ ²⁰, we report below the final result for the renormalized current two-point correlator \mathcal{G}_{-+}^F

$$\mathcal{G}_{-+}^F(q) = \frac{iN_F q_3}{16\pi\lambda_F} \left(1 + \frac{2ih_F(c_F)}{q_3} \right)^2 \left[\frac{H_F(\infty, q_3)}{H_F(c_F, q_3)} - 1 \right] - \frac{N_F \xi_F(c_F)}{8\pi} \quad (3.55)$$

Alternatively, one could have considered the exact J_+^F vertex and the ‘tree level’ J_-^F insertion vertex and perform the above exercise and would have got the same result²¹. This follows from the fact that $\mathcal{G}_{-+}^F(q) = \mathcal{G}_{+-}^F(-q)$.

4 Thermal correlators in the bosonic theory

4.1 Brief review of the theory

In this section, we study the mass deformed regular bosonic matter theory coupled to $SU(N_B)$ Chern-Simons gauge fields in the large N_B limit, at finite temperature. The Euclidean action for this theory is given by

$$\begin{aligned} \mathcal{S}_B[A, \phi] = & \frac{i\kappa_B}{4\pi} \int d^3x \epsilon^{\mu\nu\rho} \text{tr} \left(A_\mu \partial_\nu A_\rho - \frac{2i}{3} A_\mu A_\nu A_\rho \right) \\ & + \int d^3x \left((D_\mu \bar{\phi})(D^\mu \phi) + m_B^2 \bar{\phi} \phi + \frac{b_4}{2N_B} (\bar{\phi} \phi)^2 + \frac{b_6}{6N_B^2} (\bar{\phi} \phi)^3 \right). \end{aligned} \quad (4.1)$$

One of our goal is to check the duality between the fermionic and the bosonic theories. We have studied the regular fermionic matter theory in the previous section which is dual to the critical boson theory. We will study critical boson theory by taking critical limit of the regular boson theory defined by the action (4.1), in the next section. We work in the lightcone gauge $A_- = 0$. The Feynman rules in this theory include the following :

¹⁹Use the fact that for a general matrix $M = M_\mu \gamma^\mu + M_\mathbb{1} \mathbb{1}$, $\text{Tr}_F(M \gamma_\nu) = M_\mu \text{Tr}_F(\gamma^\mu \gamma_\nu) = 2M_\nu$.

²⁰This divergence term can be removed by dimensional regularization as discussed in [62]. Alternatively, one can also subtract this by adding the mass counterterm for the source field corresponding to J_μ^F [9].

²¹We also explicitly checked this (not provided here).

Propagator for the gauge field

The gauge boson propagator in the lightcone gauge $A_- = 0$, in Euclidean space, is given by

$$\langle A_\mu(p') A_\nu(p) \rangle = G_{\nu\mu}^B(p) (2\pi)^3 \delta^{(3)}(p + p') , \quad (4.2)$$

where,

$$G_{\nu\mu}^B(p) = \frac{2\pi\epsilon_{\nu-\mu}}{\kappa_B p_-} . \quad (4.3)$$

As also mentioned in the fermionic case, the gauge field propagator (4.3) is independent of p_3 , so is not effected by the holonomy [10]. We label the gauge fields both in the case of fermionic theory and here in the bosonic theory by the same symbol A_μ . However, the distinction should be obvious from the context whether we study fermionic or bosonic theory. Also we denote the gauge field propagator by the same feynman diagram as shown around equation (3.4) but now it is equal to $G_{\nu\mu}^B(p)$.

Propagator for the scalar field

The exact propagator for the scalar fields in the Euclidean space is given by

$$\langle \phi(p) \bar{\phi}(p') \rangle = S_B(p) (2\pi)^3 \delta^{(3)}(p + p') , \quad (4.4)$$

where,

$$S_B(p) = \frac{1}{p^2 + c_B^2} . \quad (4.5)$$

Here, c_B is the thermal mass of the scalar field, which is related to the bare mass-squared m_B^2 as $c_B^2 = m_B^2 + \Sigma_B$, where Σ_B is the bosonic self energy²². The final result for the gap equation of the thermal is given by²³

$$c_B^2 = m_B^2 - \frac{b_4}{4\pi} \xi_B(c_B) + \left(\frac{\lambda_B^2}{4} + \frac{b_6}{32\pi^2} \right) \xi_B^2(c_B) , \quad (4.6)$$

where, the function $\xi_B(x)$ is defined in (A.1). Solving the equation (4.6), one finds the thermal mass c_B of the regular boson theory. In Feynman diagram, the exact scalar propagator is denoted by

$$\text{---} \overset{p}{\bullet} \text{---} = S_B(p)$$

In the lightcone gauge $A_- = 0$, the contribution to the vertex factor corresponding to the term $\bar{\phi} A_3 A^3 \phi$ is $\mathcal{V}_{\bar{\phi} A^2 \phi} = -1$. In momentum space of the action (4.1), the vertex contribution corresponding to the interaction term $\bar{\phi}(p) A_\mu(-(p+k)) \phi(k)$, is given by $\mathcal{V}_{\bar{\phi} A \phi}^\mu(k, p) = (k-p)^\mu$, with explicit momentum conservation at the vertex.

²²As discussed in the literature in great details, one can compute the self energy either by summing the feynman diagrams or by integrating out the matter fields and using the Hubbard-Stratonovich trick.

²³As already mentioned before, we use the convention such that c_B is always positive. In the zero temperature limit, $\xi_B(c_B)$ reduces to c_B . So, in this limit (4.6) reduces to

$$c_B^2 = m_B^2 - \frac{b_4}{4\pi} c_B + \left(\frac{\lambda_B^2}{4} + \frac{b_6}{32\pi^2} \right) c_B^2 .$$

4.2 Thermal four-point function of fundamental scalars

In the zero-temperature theory, the connected scalar four-point function was computed previously in the literature (see, e.g., [7, 17]). In [7], the authors computed the connected scalar four-point function in the massless regular boson theory. On the other hand, in the case of massive regular boson theory as given by the action (4.1), the off-shell connected scalar four-point function was computed in [17], at zero temperature. In this section, we generalize these computations to the finite temperature for the massive theory. We closely follow the procedure given in [17]²⁴ to do the computation of four-point function of scalars. The corresponding Schwinger-Dyson equations that we need to solve are the same as given in [17] (for the relevant Schwinger-Dyson equations see (D.7) and (D.8)²⁵, and for the diagrams see Fig.7,8 and 9²⁶). Here, we generalize these computations to the finite temperature by taking into account the effect of holonomy. In the Appendix D, we present the details of the computation and discuss about the relevant modifications we need to do, because of the holonomy, in the methods of computations given in [17]. In this subsection we present the finite temperature results of the off-shell four-point function of fundamental scalars.

Following [7, 17], we define the exact, connected off-shell scalar four-point function by

$$\langle \phi^i(p+q) \bar{\phi}_j(-(k+q)) \phi^m(k) \bar{\phi}_n(-p) \rangle = \mathcal{A}_{jn}^{im}(p, k, q) (2\pi)^3 \delta^{(3)}(0) . \quad (4.7)$$

Here, we have explicitly shown the color indices of the scalar fields. Without loss of generality, we choose the color contractions to be $\delta_n^i \delta_j^m$ (terms with other possible color contractions are related to this by the permutation of momenta) [7, 17]; so, we consider²⁷

$$\mathcal{A}_{jn}^{im}(p, k, q) = \mathcal{A}(p, k, q) \delta_n^i \delta_j^m . \quad (4.8)$$

Following [17] and including the finite temperature effects as discussed in the Appendix D, we compute $\mathcal{A}(p, k, q)$ in the choice of the overall external momenta $q_{\pm} = 0$, by solving the Schwinger-Dyson equation given in Fig.7. The final result for $\mathcal{A}(p, k, q)$ is (see (D.20) or equivalently (D.29))

$$N_B \mathcal{A}(\vec{p}, \vec{k}, q_3) = \frac{H_B(a(p_s), q_3)}{H_B(a(k_s), q_3)} \left\{ (4\pi i \lambda_B q_3) \frac{(p+k)_-}{(p-k)_-} + j(q_3) \right\} , \quad (4.9)$$

where, $H_B(z, q_3)$ is given by (A.5) and $a(p_s) = +\sqrt{p_s^2 + c_B^2}$. The function $j(q_3)$ is given by

$$\frac{j(q_3)}{4\pi i \lambda_B q_3} = \frac{4\pi i \lambda_B q_3 (H_B(c_B, q_3) - H_B(\infty, q_3)) + \tilde{b}_4 (H_B(c_B, q_3) + H_B(\infty, q_3))}{4\pi i \lambda_B q_3 (H_B(c_B, q_3) + H_B(\infty, q_3)) + \tilde{b}_4 (H_B(c_B, q_3) - H_B(\infty, q_3))} , \quad (4.10)$$

where, \tilde{b}_4 is given by (D.4). An alternative and simplified form of $j(q_3)$ is given by (D.28). In the zero temperature limit, this matches with the existing results given in [17]. We use the result (4.9) to compute the correlation functions of gauge invariant, single trace operators of different spin.

²⁴For details of their method, see section 3.1 and appendix D of [17].

²⁵Or, see e.g. equation 4.6 of [17].

²⁶Or, see figure 5 and figure 4 of [17].

²⁷ $\mathcal{A}(p, k, q)$ here is the same as the quantity $V(p, k, q)$ in [17] but now at finite temperature. We use a different symbol to avoid the notational clash with the rest of the paper.

4.3 Case I: Spin zero

In the regular boson theory that we study here, there is a gauge invariant, single trace, spin 0 operator given by $J_0^B(x) = \bar{\phi}(x)\phi(x)$. In the momentum space, this takes the following form

$$J_0^B(-q) = \int \frac{\mathcal{D}_B^3 k}{(2\pi)^3} \bar{\phi}(-(k+q)) \phi(k) . \quad (4.11)$$

4.3.1 Exact insertion vertex

To compute the correlators involving J_0^B , we need the exact J_0^B insertion vertex which is defined as

$$\langle J_0^B(-q)\phi(k)\bar{\phi}(p) \rangle = V_0^B(k, q) (2\pi)^3 \delta^{(3)}(p+k+q) . \quad (4.12)$$




Figure 3. Exact vertices $\langle J_0^B \phi \bar{\phi} \rangle$. The circled cross denotes an insertion of an exact vertex, the bare cross denotes the insertion vertex in the ‘free’ theory. The filled circle denotes the exact scalar propagator. And the elliptic blob denotes the exact scalar four-point function.

From (4.11), we see that the J_0^B insertion in the ‘free’ theory is $V_{0,\text{free}}^B(k, q) = 1$. We work in the momentum $q_\pm = 0$. The exact J_0^B insertion vertex is shown in Fig.3. In mathematical form, this is given by

$$V_0^B(k, q) = 1 + N_B \int \frac{\mathcal{D}_B^3 p}{(2\pi)^3} \frac{\mathcal{A}(\vec{p}, \vec{k}, q_3)}{(p_3^2 + a^2(p_s))((p_3 + q_3)^2 + a^2(p_s))} , \quad (4.13)$$

where, $a(p_s) = +\sqrt{p_s^2 + c_B^2}$. The factor N_B in the second term on the RHS of (4.13) comes from the color trace in the loop. It is useful to break the momentum integration measure as $\mathcal{D}_B^3 p = p_s dp_s d\theta_p \mathcal{D}_B p_3$ where, p_s is the radial momentum in the lightcone plane and θ_p is the angular direction in that plane. $\mathcal{D}_B p_3$ is the integration measure for the momentum component p_3 taking into account the effect of holonomy as given in (2.6). Doing the integration over the momenta p_3 , we get

$$V_0^B(k, q) = 1 + N_B \int \frac{d^2 \vec{p}}{(2\pi)^2} \frac{\chi_B(a(p_s)) \mathcal{A}(\vec{p}, \vec{k}, q_3)}{a(p_s)(p_3^2 + a^2(p_s))} . \quad (4.14)$$

Using the definition (A.5) and the explicit expression of $\mathcal{A}(\vec{p}, \vec{k}, q_3)$ given by (4.9) and simplifying, we get,

$$V_0^B(k, q) = 1 + \left[\frac{1}{2H(a(k_s), q_3)} \int_0^\infty p_s (\partial_{p_s} H(a(p_s), q_3)) \int_0^{2\pi} \frac{d\theta_p}{2\pi} \left\{ \frac{(p+k)_-}{(p-k)_-} + \frac{j(q_3)}{4\pi i \lambda_B q_3} \right\} \right]. \quad (4.15)$$

Performing the angular integral, we get

$$V_0^B(k, q) = 1 + \left[\frac{1}{2H(a(k_s), q_3)} \int_0^\infty p_s (\partial_{p_s} H(a(p_s), q_3)) \left\{ 2\Theta(p_s - k_s) - 1 + \frac{j(q_3)}{4\pi i \lambda_B q_3} \right\} \right]. \quad (4.16)$$

Finally, performing the radial integral and simplifying further, we find

$$V_0^B(k, q) = \frac{\frac{1}{2} \left(\frac{j(q_3)}{4\pi i \lambda_B q_3} + 1 \right) H(\infty, q_3) - \frac{1}{2} \left(\frac{j(q_3)}{4\pi i \lambda_B q_3} - 1 \right) H(c_B, q_3)}{H(a(k_s), q_3)}. \quad (4.17)$$

For later convenience, we label the numerator on the RHS of (4.17) by the symbol $\tilde{V}(q_3)$, in terms of which (4.17) takes the following form

$$V_0^B(k, q) = \frac{\tilde{V}(q_3)}{H(a(k_s), q_3)}. \quad (4.18)$$

4.3.2 Two-point function

As in the case of fermions described in subsubsection 3.3.2, the two-point function for the spin zero operator J_0^B is computed by evaluating the Feynman diagram shown in Fig.4. To compute the two-point function $\langle J_0^B(q') J_0^B(q) \rangle$, a single exact insertion vertex $J_0^B(q')$ is required to account for all the perturbative Feynman diagrams without any overcounting. We choose the insertion at the left in Fig.4 as the exact vertex. Insertion on right side in




Figure 4. Diagram contributing to two-point function $\langle J_0(-q) J_0(q) \rangle$. The filled circle denotes the exact scalar propagator.

Fig.4 is the ‘free’ insertion vertex which we define as follows :

$$\langle J_0^B(-q) \phi(k) \bar{\phi}(p) \rangle = U_0^B(k, q) (2\pi)^3 \delta^{(3)}(p + k + q). \quad (4.19)$$

From the definition of $J_0^B(q)$ operator in (3.12), it follows that $U_0^B(k, q) = 1$. We define the two-point correlation function of J_0^B operator as

$$\langle J_0^B(q') J_0^B(q) \rangle = \mathcal{G}_0^B(q) (2\pi)^3 \delta^{(3)}(q' + q) . \quad (4.20)$$

From the definition of the J_0^B , it follows that the other ‘free’ insertion vertex is $U_0^B(k, q) = 1$. We now have all the building blocks to compute the $\langle J_0^B J_0^B \rangle$ correlator. As discussed before, we compute the diagram shown in Fig.4, which translates to equation as

$$\mathcal{G}_0^B(q) = N_B \int \frac{\mathcal{D}_B^3 k}{(2\pi)^3} \frac{V_0^B(k, q) U_0^B(k + q, -q)}{(k^2 + c_B^2)((k + q)^2 + c_B^2)} . \quad (4.21)$$

Working in the case $q_\pm = 0$, and inserting the expressions for V_0^B and U_0^B , we find

$$\mathcal{G}_0^B(q) = N_B \tilde{V}(q_3) \int \frac{\mathcal{D}_B^3 k}{(2\pi)^3} \frac{1}{(k_3^2 + a^2(p_s))((k_3 + q_3)^2 + a^2(p_s))H(a(k_s), q_3)} . \quad (4.22)$$

The crucial fact is that the above momentum integral (4.22) can be carried out analytically. As before, we choose to separate the above integral as an integral over the radial momentum $k_s = \sqrt{k_1^2 + k_2^2} = \sqrt{2k_+ k_-}$, and an integral over an angular variable θ_k . It is convenient to perform the k_3 integral first, which can be carried out using the integration result (A.3). The angular integral in (4.22) contributes unity. Once these two integrals are performed, finally, the integration over radial momentum k_s can also be performed. By changing the integration variable k_s to another variable $a(k_s) = +\sqrt{k_s^2 + c_B^2}$, the integrand in the radial integral can be written as a total derivative w.r.t. the variable $a(k_s)$. Thus the remaining integral can be carried out completely and the final result (inserting back the explicit form of $\tilde{V}(q_3)$) for the $\langle J_0^B J_0^B \rangle$ two-point function in the regular boson theory at finite temperature is

$$\mathcal{G}_0^B(q) = -\frac{N_B}{4\pi i \lambda_B q_3 \left(\frac{H_B(c_B, q_3) + H_B(\infty, q_3)}{H_B(c_B, q_3) - H_B(\infty, q_3)} \right) + \tilde{b}_4} . \quad (4.23)$$

4.4 Case II: Spin one

The regular boson theory that we study here, has a global $U(1)$ symmetry and the corresponding conserved spin 1 current is

$$J_\mu^B(x) = i \left[(D_\mu \bar{\phi}) \phi - \bar{\phi} (D_\mu \phi) \right] . \quad (4.24)$$

In the momentum space, the J_μ^B can be written as

$$J_\mu(-q) = \int_k (2k + q)_\mu \bar{\phi}(-(k + q)) \phi(k) - 2 \int_{p, k} \bar{\phi}(p) A_\mu(-(p + k + q)) \phi(k) . \quad (4.25)$$

4.4.1 Exact insertion vertex

In this subsection we compute the exact J_μ^B insertion vertex which is one of the building blocks for computing the corresponding correlators. We define the exact J_μ^B insertion vertex by

$$\langle J_\mu^B(-q) \phi(k) \bar{\phi}(p) \rangle = V_{(\mu)}^B(k, q) (2\pi)^3 \delta^{(3)}(p + k + q) . \quad (4.26)$$

The exact insertion vertex is diagrammatically shown in Fig.3, by a circled cross. The insertion denoted by a bare cross is understood to be the insertion vertex in the ‘free’ theory. In equation, $V_{(\mu)}^B(k, q)$ is given by

$$V_{(\mu)}^B(k, q) = V_{(\mu), \text{free}}^B(k, q) + N_B \int \frac{\mathcal{D}_B^3 p}{(2\pi)^3} \left[S_B(p+k) V_{(\mu), \text{free}}^B(p, q) S_B(p) \mathcal{A}(p, k, q) \right], \quad (4.27)$$

where, $S_B(k)$ is the exact scalar propagator given by (4.5) and $\mathcal{A}(p, k, q)$ is the thermal scalar 4-point function which in the ‘lightcone kinematics’ ($q_{\pm} = 0$), is given by (4.9). Substituting the exact scalar propagator we write this in a more convenient form as

$$V_{(\mu)}^B(k, q) = V_{(\mu), \text{free}}^B(k, q) + N_B \int \frac{\mathcal{D}_B^3 p}{(2\pi)^3} \frac{V_{(\mu), \text{free}}^B(p, q) \mathcal{A}(p, k, q)}{(p^2 + c_B^2)((p+q)^2 + c_B^2)}, \quad (4.28)$$

To perform the integral in (4.28), we need $V_{(\mu), \text{free}}^B(k, q)$, which can be easily read from the definition of J_{μ}^B current given in (4.25).

In this paper the explicit computations are performed in the lightcone gauge $A_- = 0$ and with the external momenta $q_{\pm} = 0$. For the moment, we will be interested in the computation of exact J_-^B vertex. It follows from (4.25) that $V_{(-), \text{free}}^B(k, q) = 2k_-$. From (4.28), it follows that

$$V_{(-)}^B(k, q) = 2k_- + N_B \int \frac{\mathcal{D}_B^3 p}{(2\pi)^3} \frac{(2p_-) \mathcal{A}(\vec{p}, \vec{k}, q_3)}{(p^2 + c_B^2)((p+q)^2 + c_B^2)}. \quad (4.29)$$

To perform the momentum space integral we follow the same procedure as in the case of V_0^B . We perform the integral over p_3 first and then carry out the angular integral by inserting the explicit form of $\mathcal{A}(\vec{p}, \vec{k}, q_3)$ to reduce into an one-dimensional integral over the radial momenta p_s . Interestingly enough, it is noted that the radial momenta integral can also be carried out analytically by writing the corresponding integrand as a total derivative w.r.t. the reduced integration variable $a(p_s) = +\sqrt{p_s^2 + c_B^2}$. Performing the integral, we find the final result for the exact J_-^B insertion vertex as

$$V_{(-)}^B(k, q) = 2k_- \frac{H_B(\infty, q_3)}{H_B(a(k_s), q_3)}. \quad (4.30)$$

4.4.2 Two-point function

In this subsection, we compute the $\langle J_- J_+ \rangle$ two-point correlator. The corresponding feynman diagram is shown in Fig.5. As mentioned before, we choose the insertion at the left of the digram to be the exact vertex which in our present case is the the exact J_-^B vertex. The remaining thing is to compute the contribution from the insertion on the right side. To keep a distinction, we define the insertion on the right to be

$$\langle J_{\nu}^B(-q) \phi(k) \bar{\phi}(p) \rangle = U_{(\nu)}^B(k, q) (2\pi)^3 \delta^{(3)}(p+k+q). \quad (4.31)$$

To compute the two-point function $\langle J_-^B J_+^B \rangle$, we need the ‘free’ J_+^B insertion vertex. From the definition (4.25), it is clear that the ‘bare’ insertion $U_{(+)}^B(k, q)$ has one insertion without




Figure 5. Schematic diagram contributing to two-point function $\langle J^{(s)}(-q)J^{(s)}(q) \rangle$ for spin $s > 0$. The filled circles denote the exact scalar propagator.




Figure 6. Diagrams with nonzero contributions to the ‘bare’ vertex $U_{(+)}^B$ which is the boxed cross on the RHS in Fig.5. The filled circle denotes the exact scalar propagator. The loop momentum in the last two diagrams is ℓ .

the gauge field i.e., the first term and the one involving the gauge field which is the second term; diagrammatically, these are shown in Fig.6. Summing the diagrams in Fig.6, total non-zero contribution to the ‘bare’ insertion vertex $U_{(+)}^B$ is given by

$$U_{(+)}^B(k, q) = 2k_+ - 4\pi\lambda_B\epsilon_{+-\rho}q^\rho \int \frac{\mathcal{D}_B^3\ell}{(2\pi)^3} \frac{S_B(\ell)}{(\ell - k)_-}. \quad (4.32)$$

Here $S_B(\ell)$ is the exact scalar propagator given by (4.5). Performing the angular integral, the integral over ℓ_3 and finally performing the radial integral over ℓ_s (also using the fact that $\epsilon_{+-\rho} = i\delta_{\rho 3}$), we get

$$U_{(+)}^B(k, q) = 2k_+ + \frac{i\lambda_B q_3}{k_-} \left[\xi_B(a(k_s)) - \xi_B(c_B) \right], \quad (4.33)$$

where, the function $\xi_B(z)$ is given by (A.1).

We are now all set to compute the two-point function of the spin-one current operator. We define the two-point function $\langle J_\mu^B(-q)J_\nu^B(q) \rangle$ as

$$\langle J_\mu^B(q')J_\nu^B(q) \rangle = \mathcal{G}_{\mu\nu}^B(q) (2\pi)^3 \delta^{(3)}(q' + q). \quad (4.34)$$

Diagrammatically, this is shown in Fig.5. In equation, it is given by

$$\mathcal{G}_{\mu\nu}^B(q) = N_B \int \frac{\mathcal{D}_B^3 k}{(2\pi)^3} \left[S_B(k+q) V_{(\mu)}^B(k, q) S_B(k) U_{(\nu)}^B(k+q, -q) \right], \quad (4.35)$$

where, $S_B(k)$ is the exact scalar field propagator given by (4.5). Explicitly, we write this as

$$\mathcal{G}_{\mu\nu}^B(q) = N_B \int \frac{\mathcal{D}_B^3 k}{(2\pi)^3} \frac{V_{(\mu)}^B(k, q) U_{(\nu)}^B(k+q, -q)}{(k^2 + c_B^2)((k+q)^2 + c_B^2)}. \quad (4.36)$$

As we are working in the case of external momenta $q_{\pm} = 0$, and in $A_- = 0$ gauge, the non-trivial component in this case is $(\mu\nu) \equiv (-+)$ component. In this case, substituting the expressions (4.30) and (4.33), we find

$$\mathcal{G}_{-+}^B(q) = N_B \int \frac{\mathcal{D}_B^3 k}{(2\pi)^3} (2k_-) \frac{H_B(\infty, q_3)}{H_B(a(k_s), q_3)} \frac{2k_+ - \frac{i\lambda_B q_3}{k_-} [\xi_B(a(k_s)) - \xi_B(c_B)]}{(k_s^2 + a^2(k_s))((k_3 + q_3)^2 + a^2(k_s))}. \quad (4.37)$$

The interesting fact is that the momentum integral here in (4.37) can be carried out completely analytically. We first perform the integration over the momentum component k_3 by using the result (C.4). For the integrations in the lightcone plane, we use the polar coordinates to write the integration effectively as an integration over a radial momentum k_s and an integration over an angular variable θ_k . The angular integral in this case is trivial and contributes unity. Doing these, the integration in (4.37) is reduced to an integration over a single variable k_s . The k_s integral can be performed analytically. The integrand can be written as a total derivative of the reduced variable $a(k_s) = +\sqrt{k_s^2 + c_B^2} \equiv z$, and so carrying out the radial integral, the final result for $\langle J_- J_+ \rangle$ is given by

$$\mathcal{G}_{-+}^B(q) = \lim_{\Lambda \rightarrow \infty} \left[\frac{iN_B q_3}{16\pi\lambda_B} \left(1 + \frac{4c_B^2}{q_3^2} \right) \left[\frac{H_B(\Lambda, q_3)}{H_B(c_B, q_3)} - 1 \right] - \frac{N_B \xi_B(c_B)}{4\pi} + \frac{N_B \xi_B(\Lambda)}{4\pi} \right] \quad (4.38)$$

where, Λ is the UV cut-off in the radial momenta. The term $\xi_B(\Lambda \rightarrow \infty)$ is a pure divergent term. We regularize this answer by dropping this linearly divergent piece²⁸, the renormalized two-point function $\langle J_-^B J_+^B \rangle$, upto the momentum conserving delta function, is given by

$$\mathcal{G}_{-+}^B(q) = \frac{iN_B q_3}{16\pi\lambda_B} \left(1 + \frac{4c_B^2}{q_3^2} \right) \left[\frac{H_B(\infty, q_3)}{H_B(c_B, q_3)} - 1 \right] - \frac{N_B \xi_B(c_B)}{4\pi} \quad (4.39)$$

As discussed in [32] (see e.g. around equation 80 of [32]), the two-point correlator of the $U(1)$ current (4.24) is $\langle J_\mu^B J_\nu^B \rangle - 2\delta_{\mu\nu} \langle \bar{\phi}\phi \rangle$. We report the relevant result by taking the contribution of $\langle \bar{\phi}\phi \rangle$ into account in subsection 5.2.3.

5 Analysis of results

In this section we analyze the results that we have obtained in this paper along with studying various limiting cases.

5.1 Fermionic result

As discussed above in details, we have studied in this paper the massive regular fermions coupled to Chern-Simons theory at finite temperature with arbitrary holonomy distribution. Below we summarize the result and analyze the different relevant structures of the results.

²⁸This can be removed by using dimensional regularization in the integration over k_s [62]. This can also be done by turning on the mass counterterm for the background source which couples to the current J_μ [7].

5.1.1 Two-point correlator of spin 0 operator :

The final result for $\langle J_0^F(-q)J_0^F(q) \rangle$ is given in (3.40) and is rewritten below

$$\mathcal{G}_0^F(q) = \frac{N_F|q_3|}{4\pi\lambda_F} \cot \left[2\lambda_F|q_3| \int_{c_F}^{\infty} \frac{dw}{q_3^2 + 4w^2} \chi_F(w) - \text{sgn}(h_F) \tan^{-1} \frac{|q_3|}{2c_F} \right] + \frac{N_F m_F}{2\pi\lambda_F}, \quad (5.1)$$

where, $\chi_F(w)$ is given by (2.9) and $\text{sgn}(h_F)$ is defined around (3.10) (at zero temperature, $\text{sgn}(h_F)$ effectively reduces to $\text{sgn}(m_F)$). This result is general and is valid in the general case of massive theory at finite temperature. Interestingly, (5.1) is even in q_3 . From the corresponding CFT results [9]²⁹, we know that the two point function of spin zero scalar ‘current’ operator is even in the general momentum q . The spin zero operator is even under parity in the corresponding CFT [9], and it is unlikely that the finite temperature effects will break that structure. Assuming this to be true, we ‘covariantize’ the result (5.1) by replacing $|q_3|$ with corresponding ‘ $SO(3)$ rotation-invariant’ generalization, i.e., with $|q| = +\sqrt{q_1^2 + q_2^2 + q_3^2}$ ³⁰. The corresponding ‘covariant’ generalization is given below

$$\mathcal{G}_0^F(q) = \frac{N_F|q|}{4\pi\lambda_F} \cot \left[2\lambda_F|q| \int_{c_F}^{\infty} \frac{dw}{q^2 + 4w^2} \chi_F(w) - \text{sgn}(h_F) \tan^{-1} \frac{|q|}{2c_F} \right] + \frac{N_F m_F}{2\pi\lambda_F}, \quad (5.2)$$

We now study various limiting cases of this result and check that it matches with the existing previous results.

Zero temperature and zero mass

We start with the simplest possible limiting case, i.e., the case when both temperature and fermion bare mass parameter m_F is zero. In this case, the effect of holonomy vanishes and it is clear from (2.9) that in the zero temperature limit, $\chi_F(w) = 1$. Also from the gap equation (3.10) we see that in this case c_F also vanishes; that is, there is no self energy correction to the pole mass c_F in the corresponding CFT at zero temperature [1]. As bare mass m_F can have both possible signs, we choose to take the mass goes to zero limit from the side in which $\text{sgn}(m_F) = \text{sgn}(\lambda_F)$, which is known to be dual to the unhiggsed phase of bosonic theory. It follows from (5.1) or its ‘covariantized’ form (5.2), that the two-point function of the scalar operator J_0^F in this limiting case takes the form

$$\mathcal{G}_0^F(q) = -\frac{N_F|q|}{4\pi\lambda_F} \tan \left(\frac{\pi\lambda_F}{2} \right) \quad (5.3)$$

As expected, this result matches with the result in [9]. This provides a support to the correctness of the computation performed in this paper.

²⁹By corresponding CFT, we here mean the CFT from which one gets the massive theory that is being studied in this paper by deforming it with relevant deformations.

³⁰Strictly speaking, the external momentum component q_3 is discrete at finite temperature. However, we formally treat this like a continuous variable for the purpose of ‘covariantization’. Also the true rotation symmetry is actually $SO(2)$ in the spatial plane. We however formally write the ‘covariantized’ form by replacing $|q_3|$ with $+\sqrt{q_1^2 + q_2^2 + q_3^2}$ which reduces to $|q_3|$ in the choice $q_{\pm} = 0$. The ‘covariantized’ results are indeed the correct results obtained in this paper in the case $q_{\pm} = 0$ in which case $|q|$ should be thought of as $|q_3|$. It will be nice to have a direct independent computation without setting q_{\pm} to zero to check if this is indeed the case.

Zero temperature and nonzero mass

In this case, as before the effect of holonomy vanishes implying $\chi_F(w) = 1$. The integral in the argument of cotangent in (5.2) can be performed easily in this situation and the final result is given by

$$\mathcal{G}_0^F(q) = \frac{N_F|q|}{4\pi\lambda_F} \cot \left[(\lambda_F - \text{sgn}(m_F)) \tan^{-1} \frac{|q|}{2c_F} \right] + \frac{N_F m_F}{2\pi\lambda_F}. \quad (5.4)$$

Nonzero temperature and zero mass

This is the case, when the effect of holonomy becomes non-trivial. The existing result in this case [80] is only in the case when the holonomy takes the universal table-top distribution form, i.e., $\rho_F(\alpha) = \frac{\Theta(\pi|\lambda_F| - |\alpha|)}{2\pi|\lambda_F|}$. Performing the integral over the holonomy with this particular distribution, we get

$$\chi_F(w) = \frac{i}{\pi|\lambda_F|} \left[\log \cosh \left(\frac{\beta w - i\pi|\lambda_F|}{2} \right) - \log \cosh \left(\frac{\beta w + i\pi|\lambda_F|}{2} \right) \right]. \quad (5.5)$$

Similar to the zero temperature case discussed above, when the bare mass of the fermion is zero, the theory is dual to the bosonic scalar theory for which $\text{sgn}(h_F) = \text{sgn}(\lambda_F)$. The final result of the two-point function of spin zero operator J_0^F in the thermal CFT in the ‘covariantized’ form is given by

$$\mathcal{G}_0^F(q) = \frac{N_F|q|}{4\pi\lambda_F} \cot \left[2\lambda_F|q| \int_{c_F}^{\infty} \frac{dw \chi_F(w)}{q^2 + 4w^2} - \text{sgn}(\lambda_F) \tan^{-1} \frac{|q|}{2c_F} \right]. \quad (5.6)$$

where, c_F is computed from the gap equation (3.10) by using the fact that $m_F = 0$ and using (5.12). $\chi_F(w)$ in this case is given by (5.5). In the special choice of external momenta $q = (0, 0, q_3)$, the result (5.6) agrees with [80].

5.1.2 Two-point correlator of spin 1 current :

The final result for spin 1 current two point function $\langle J_-^F J_+^F \rangle$ is given by (3.55). We rewrite this result below by using the fact that $h_F(c_F) = \text{sgn}(h_F)c_F$ (this follows from the gap equation (3.10)) and the explicit form of $H_F(z, q_3)$. It takes the following form ³¹

$$\mathcal{G}_{-+}^F(q) = \frac{iN_F q_3}{16\pi\lambda_F} \left(1 + \frac{2i\text{sgn}(h_F)c_F}{q_3} \right)^2 \left[\exp \left(4i\lambda_F q_3 \int_{c_F}^{\infty} \frac{dw \chi_F(w)}{q_3^2 + 4w^2} \right) - 1 \right] - \frac{N_F \xi_F(c_F)}{4\pi}. \quad (5.9)$$

³¹This does not have a definite even/odd property under $q_3 \rightarrow -q_3$. However, this can formally be written as a sum of the even and odd parts, i.e., $\mathcal{G}_{-+}^F(q) = \mathcal{G}_{-+}^{F,\text{even}}(q) + \mathcal{G}_{-+}^{F,\text{odd}}(q)$, where, $\mathcal{G}_{-+}^{F,\text{even}}(-q) = \mathcal{G}_{-+}^{F,\text{even}}(q)$ and $\mathcal{G}_{-+}^{F,\text{odd}}(-q) = -\mathcal{G}_{-+}^{F,\text{odd}}(q)$. In this footnote, we propose to formally write the possible ‘covariantized form’ of the correlator by replacing $|q_3|$ with $|q| = +\sqrt{q_1^2 + q_2^2 + q_3^2}$ as (also see footnote 30 for a related clarification)

$$\mathcal{G}_{\mu\nu}^{F,\text{even}}(q) = \left[\frac{N_F}{16\pi\lambda_F} \left(1 - \frac{4c_F^2}{q^2} \right) \sin(4\lambda_F|q|A_F) + \frac{N_F \text{sgn}(h_F)c_F}{4\pi\lambda_F|q|} \left(\cos(4\lambda_F|q|A_F) - 1 \right) + \frac{N_F \xi_F(c_F)}{4\pi|q|} \right] \frac{q_\mu q_\nu - \delta_{\mu\nu} q^2}{|q|}, \quad (5.7)$$

and

$$\mathcal{G}_{\mu\nu}^{F,\text{odd}}(q) = \frac{N_F}{8\pi\lambda_F q^2} \left[(q^2 - 4c_F^2) \sin^2(2\lambda_F|q|A_F) + 2\text{sgn}(h_F)c_F|q| \sin(4\lambda_F|q|A_F) \right] \epsilon_{\mu\nu\rho} q^\rho, \quad (5.8)$$

Below we study limiting cases of this result and verify with the existing results.

Zero temperature and zero mass

As already discussed in the spin zero case, in the zero temperature limit, $\chi_F(w) = 1$ and so, the integral appearing in the exponential in (5.9) can be performed exactly. Also as discussed before, the pole mass c_F vanishes in this case. Using these and the fact that $m_F = 0$ in this case, we get

$$\mathcal{G}_{-+}^F(q) = \frac{iN_F q_3}{16\pi\lambda_F} \left[e^{\pi i \lambda_F \text{sgn}(q_3)} - 1 \right]. \quad (5.10)$$

This exactly matches with the corresponding CFT results at zero temperature reported in [9]³².

Zero temperature and nonzero mass

In this limiting case, the effect of holonomy goes away. As mentioned before, in the zero temperature limit $\chi_F(w)$ equals to unity and the integral appearing in the exponential in (5.9) can be performed exactly. Keeping finite nonzero bare mass m_F , in this case the result (5.9) reduces to

$$\mathcal{G}_{-+}^F(q) = \frac{iN_F q_3}{16\pi\lambda_F} \left(1 + \frac{2ic_F \text{sgn}(m_F)}{q_3} \right)^2 \left[e^{2i\lambda_F \tan^{-1}\left(\frac{q_3}{2c_F}\right)} - 1 \right] - \frac{N_F c_F}{4\pi}, \quad (5.11)$$

where, c_F is the fermionic pole mass determined from the gap equation (3.10) at zero temperature which gives $c_F = \frac{m_F}{\text{sgn}(m_F) - \lambda_F}$. This exactly matches with the results of [32, 85].

Nonzero temperature and zero mass

As discussed before, at finite temperature the effects of holonomy becomes non-trivial. The result in (5.9) is valid for any arbitrary holonomy distribution. In the specific case of table-top holonomy $\rho_F(\alpha) = \frac{\Theta(\pi|\lambda_F| - |\alpha|)}{2\pi|\lambda_F|}$, as used in [32, 80], the final result is given by (5.9), where $\chi_F(w)$ is now given by (5.5). $\xi_F(c_F)$ in this case is given by (A.2) with the table-top holonomy distribution, i.e.,

$$\xi_F(c_F) = \frac{1}{2\pi|\lambda_F|\beta} \int_{-\pi|\lambda_F|}^{\pi|\lambda_F|} d\alpha \left(\ln 2 \cosh\left(\frac{\beta c_F + i\alpha}{2}\right) + \ln 2 \cosh\left(\frac{\beta c_F - i\alpha}{2}\right) \right). \quad (5.12)$$

The results of the holonomy integral above are in terms of the dilogarithm functions as given, e.g., in [10, 32, 80].

where, in the above two expressions $A_F = \int_{c_F}^{\infty} \frac{dw}{q^2 + 4w^2} \chi_F(w)$. It would be nice to have a direct independent check if this is indeed the correct form of $\mathcal{G}_{\mu\nu}^F(q)$ without choosing $q_{\pm} = 0$. Similar arguments apply for the correlator of spin one current in the bosonic case as given in (5.32). The author would like to thank S. Minwalla for asking a question about this.

³²As discussed in [9], this result can be separated into parity even and parity odd parts. And in this case, the covariantized form of the parity even part is given by $\frac{N_F}{16} \frac{\sin(\pi\lambda_F)}{\pi\lambda_F} \frac{q_{\mu}q_{\nu} - \delta_{\mu\nu}q^2}{|q|}$ as reported in [9]. The parity odd part can come from a contact term proportional to $\epsilon_{\mu\nu\rho}q^{\rho}$ [9] (see discussion around equation 35 of [9]).

5.2 Bosonic result

We have studied in this paper the bosonic theory as well. We have considered the massive regular bosonic matter theory coupled to $SU(N_B)$ Chern-Simons theory at finite temperature with arbitrary holonomy distribution. We have computed various correlators which we summarize below considering several limiting cases as well and check that it agrees with the existing results in limiting cases.

5.2.1 Two-point correlator spin 0 operator :

The final result for the renormalized two point function of single trace, spin zero operator J_0^B in the regular boson theory is given by (4.23) which we rewrite below to analyze this result further.

$$\mathcal{G}_0^B(q) = \frac{N_B}{4\pi\lambda_B q_3 \cot\left(2\lambda_B q_3 \int_{c_B}^{\infty} \frac{dw}{q_3^2 + 4w^2} \chi_B(w)\right) - \tilde{b}_4}, \quad (5.13)$$

where, \tilde{b}_4 is given by (D.4) and thermal mass c_B is given by (4.6). First of all, it is easy to see that this result is even under $q_3 \rightarrow -q_3$ (we have also seen the same property in the fermionic result in (5.1)). As described in the J_0^F case (see footnote 30), we ‘covariantize’ this result by replacing $|q_3|$ with $|q| = +\sqrt{q_1^2 + q_2^2 + q_3^2}$. The ‘covariantized’ form of this result is given by

$$\mathcal{G}_0^B(q) = \frac{N_B}{4\pi\lambda_B |q| \cot\left(2\lambda_B |q| \int_{c_B}^{\infty} \frac{dw}{q^2 + 4w^2} \chi_B(w)\right) - \tilde{b}_4}. \quad (5.14)$$

Zero temperature and zero mass

Let us consider the simple case of zero temperature and zero bare mass $m_B = 0$. In the zero temperature limit $\chi_B(w) = 1$ and $\xi_B(w) = w$. In this case, the pole mass of the scalar obtained by solving (4.6). The integration appearing in the argument of (5.14) can be performed explicitly in case of zero temperature. The final result in this case is given by

$$\mathcal{G}_0^B(q) = \frac{N_B}{4\pi\lambda_B |q| \cot\left(\lambda_B \tan^{-1}\left(\frac{|q|}{2c_B}\right)\right) - \tilde{b}_4}. \quad (5.15)$$

Solving (4.6), in this case, one finds the pole mass $c_B = 0$. It follows that with nonzero quartic coupling b_4 , the two-point function of the scalar operator in this case is given by

$$\mathcal{G}_0^B(q) = \frac{N_B}{4\pi\lambda_B |q| \cot\left(\frac{\pi\lambda_B}{2}\right) + b_4}. \quad (5.16)$$

This agrees with the results of [7] (see e.g. equation 66 of [7]). In the special case when b_4 is zero, (5.16) reduces to

$$\mathcal{G}_0^B(q) = \frac{N_B}{4\pi\lambda_B |q|} \tan\left(\frac{\pi\lambda_B}{2}\right). \quad (5.17)$$

This result exactly matches with the one given in [7] (see e.g. equation 35 of [7]).

Zero temperature and nonzero mass

As discussed above, at zero temperature the effect of holonomy disappears. As the bare mass m_B is nonzero, the pole mass c_B in this case is nonzero and is obtained by solving the gap equation (4.6) at zero temperature in which case $\xi_B(c_B) = c_B$. The two-point correlator of J_0^B operator in this case is given by

$$\mathcal{G}_0^B(q) = \frac{N_B}{4\pi\lambda_B|q| \cot\left(\lambda_B \tan^{-1}\left(\frac{|q|}{2c_B}\right)\right) - \tilde{b}_4} . \quad (5.18)$$

Nonzero temperature and zero mass

In this case, the non-trivial effects of holonomy becomes important. The result that we have given in (5.14) is valid for any arbitrary holonomy distribution. In the case of specific table-top holonomy $\rho_B(\alpha) = \frac{\Theta(\pi|\lambda_B| - |\alpha|)}{2\pi|\lambda_B|}$, $\rho_B(\alpha)$, as discussed in [80], the final result is given by (5.14), where $\chi_B(w)$ is now given by

$$\chi_B(w) = \frac{i}{\pi|\lambda_B|} \left[\log \sinh\left(\frac{\beta w - i\pi|\lambda_B|}{2}\right) - \log \sinh\left(\frac{\beta w + i\pi|\lambda_B|}{2}\right) \right] . \quad (5.19)$$

$\xi_B(c_B)$ in this case is given by (A.1) with the table-top holonomy distribution. This can be written as

$$\xi_B(c_B) = \frac{1}{2\pi|\lambda_B|\beta} \int_{-\pi|\lambda_B|}^{\pi|\lambda_B|} d\alpha \left(\ln 2 \sinh\left(\frac{\beta c_B + i\alpha}{2}\right) + \ln 2 \sinh\left(\frac{\beta c_B - i\alpha}{2}\right) \right) , \quad (5.20)$$

and the results of the holonomy integral are in terms of the dilogarithm functions as given, e.g., in [10, 80].

5.2.2 Critical theory limit

One of our goal in this paper is to check the conjectured bose-fermi duality. In this paper, we have studied the regular fermionic matter coupled to Chern-Simons theory which is conjectured to be dual to the critical bosons coupled to Chern-Simons theory. There is a calculational evidence of this conjectured duality from explicit computations of the thermal free energies on both sides of this duality. To check the duality of the two-point functions of gauge invariant operators, i.e., to match with the regular fermionic theory, we need to have the corresponding results in the critical bosonic theory. The ‘critical’ limit of the theory is defined [7, 9, 17] by taking $b_4 \rightarrow \infty$ and $m_B^2 \rightarrow \infty$ with $\frac{4\pi m_B^2}{b_4} = m_B^{\text{cri}}$ and b_6 kept fixed. The action for the critical theory can be obtained from the action (4.1) of the regular boson theory, by introducing a Hubbard-Stratonovich field σ_B , and taking the critical limit (we set $b_6 = 0$) [17]. The action in this case takes the form

$$\begin{aligned} \mathcal{S}_{\text{CB}}[A, \phi, \sigma_B] = & \frac{i\kappa_B}{4\pi} \int d^3x \epsilon^{\mu\nu\rho} \text{tr} \left(A_\mu \partial_\nu A_\rho - \frac{2i}{3} A_\mu A_\nu A_\rho \right) \\ & + \int d^3x \left((D_\mu \bar{\phi})(D^\mu \phi) + \sigma_B \left(\bar{\phi} \phi + \frac{N_B}{4\pi} m_B^{\text{cri}} \right) \right) . \end{aligned} \quad (5.21)$$

We restrict our attention here to the case when $m_B^{\text{cri}} > 0$. In the critical boson theory, the single trace scalar operator J_0^B is simply the Lagrange multiplier field σ_B . The first term in the denominator of (5.14) is finite (here thermal mass c_B is assumed to be finite), however the second term is proportional to b_4 which grows without limit in the critical limit. However, we extract finite result by rescaling the J_0^B operator with b_4 ³³ and define a new operator

$$\tilde{J}_0^B = \frac{b_4}{4\pi\lambda_B} J_0^B. \quad (5.22)$$

Here, we have used the particular normalization to exactly match its two-point function with the dual result in the regular fermionic theory. We define the two-point function of \tilde{J}_0^B operator by $\tilde{\mathcal{G}}_0^B$ upto the momentum conserving delta function, and so, we see that in the critical boson theory,

$$\tilde{\mathcal{G}}_0^B(q) = \lim_{b_4 \rightarrow \infty} \left(\frac{b_4}{4\pi\lambda_B} \right)^2 \mathcal{G}_0^B(q). \quad (5.23)$$

Taking the critical limit, by sending b_4 and m_B^2 to infinity while keeping their ratio fixed, we find the two-point function of the operator \tilde{J}_0^B to be given by (we drop the contact term $\frac{N_B b_4}{(4\pi\lambda_B)^2}$)

$$\tilde{\mathcal{G}}_0^B(q) = -\frac{N_B|q|}{4\pi\lambda_B} \cot \left(2\lambda_B|q| \int_{c_B}^{\infty} \frac{dw \chi_B(w)}{q^2 + 4w^2} \right) \quad (5.24)$$

where, the thermal mass c_B appearing in (5.24) has to be computed from (4.6) in the critical limit which amounts to solving the gap equation $\xi_B(c_B) = m_B^{\text{cri}}$ for a fixed $m_B^{\text{cri}} > 0$.

Zero temperature and zero critical mass : As already mentioned above, in the zero temperature limit as the holonomy becomes trivial, it is easy to see that $\chi_B(w) = 1$ and $\xi_B(w) = w$. Thus, we can explicitly perform the integral appearing in the argument of (5.24). Also in this case, as the critical mass is zero, i.e., $m_B^{\text{cri}} = 0$, so, we see that the pole mass c_B vanishes. In this case, (5.24) reduces to

$$\tilde{\mathcal{G}}_0^B(q) = -\frac{N_B|q|}{4\pi\lambda_B} \cot \left(\frac{\pi\lambda_B}{2} \right) \quad (5.25)$$

This result matches with the existing results [7] in this limiting case.

Zero temperature and nonzero critical mass : This is similar to the above case with only difference being that in this case the pole mass is non zero, and is given by $c_B = m_B^{\text{cri}}$. So, from (5.24) it follows that the final result in this case is

$$\tilde{\mathcal{G}}_0^B(q) = -\frac{N_B|q|}{4\pi\lambda_B} \cot \left[\lambda_B \tan^{-1} \left(\frac{|q|}{2c_B} \right) \right] \quad (5.26)$$

³³The scaling is feasible, because in the corresponding CFT theory, the scaling dimension of the J_0^B operator in the regular bosonic theory is 1 in leading order in N_B whereas, in the critical boson theory, the scaling dimension of the corresponding primary operator is 2.

Nonzero temperature and zero critical mass : Taking into account the effect of holonomy, in this case, $\chi_B(w)$ is not simply unity but is given by (2.7). The result (5.24) is valid for arbitrary holonomy distribution. In the specific case of table-top holonomy, (2.7) reduces to (5.19). So, in this case, the result is (5.24) where, χ_B is now given by (5.19) and the thermal mass c_B is determined from the gap equation $\xi_B(c_B) = 0$, where, the function $\xi_B(c_B)$ is now given by (5.20).

5.2.3 Two-point correlator of spin 1 current :

In (4.39), we have given the result of the two-point function $\langle J_-^B J_+^B \rangle$ of the $U(1)$ current (4.24). However, as discussed in [32] (see e.g. equation 80 of [32]), the gauge-invariant correlator of $U(1)$ current (4.24) is given by

$$\langle J_\mu^B J_\nu^B \rangle - 2\delta_{\mu\nu} \langle \bar{\phi}\phi \rangle \quad (5.27)$$

There is an extra $-2\langle \bar{\phi}\phi \rangle$ term which contributes to the correlator of the $U(1)$ current. The contribution corresponding to this part (apart from the overall momentum conserving delta function) is given by

$$-2N_B \int \frac{\mathcal{D}_B^3 \ell}{(2\pi)^3} \frac{1}{\ell^2 + c_B^2} = \frac{N_B \xi_B(c_B)}{2\pi} \quad (5.28)$$

where, to obtain the RHS of (5.28) we have used the integration result (C.3). We label the correlator (5.27) of the $U(1)$ current by (apart from the overall delta function) $\tilde{\mathcal{G}}_{\mu\nu}^B$, and so we find that

$$\tilde{\mathcal{G}}_{\mu\nu}^B(p) = \mathcal{G}_{\mu\nu}^B(p) + \delta_{\mu\nu} \frac{N_B \xi_B(c_B)}{2\pi} \quad (5.29)$$

where, $\mathcal{G}_{\mu\nu}^B$ is defined by (4.34). The final result of the correlator $\tilde{\mathcal{G}}_{\mu\nu}^B$ of the $U(1)$ current (its $(-+)$ component) is ³⁴

$$\tilde{\mathcal{G}}_{-+}^B(q) = \frac{iN_B q_3}{16\pi\lambda_B} \left(1 + \frac{4c_B^2}{q_3^2}\right) \left[\exp\left(4i\lambda_B q_3 \int_{c_B}^\infty \frac{dw \chi_B(w)}{q_3^2 + 4w^2}\right) - 1 \right] + \frac{N_B \xi_B(c_B)}{4\pi} \quad (5.32)$$

It is worth mentioning at this point that, unlike the spin zero case, there is no explicit appearance of the quartic and sextic coupling parameters b_4 and b_6 in the result (5.32) except implicitly through the thermal mass c_B given by (4.6). So, this result is unchanged also in the critical boson theory except in the critical boson theory the gap equation of the thermal mass c_B is given by $\xi_B(c_B) = m_B^{\text{cri}}$. Below, we analyze various limiting cases of the above result (5.32).

³⁴Following footnotes 30 and 31, a possible ‘covariantized’ form can be written formally as $\tilde{\mathcal{G}}_{\mu\nu}^B(q) = \tilde{\mathcal{G}}_{\mu\nu}^{B,\text{even}}(q) + \tilde{\mathcal{G}}_{\mu\nu}^{B,\text{odd}}(q)$, where,

$$\tilde{\mathcal{G}}_{\mu\nu}^{B,\text{even}}(q) = \left[\frac{N_B}{16\pi\lambda_B} \left(1 + \frac{4c_B^2}{q^2}\right) \sin(4\lambda_B |q| A_B) - \frac{N_B \xi_B(c_B)}{4\pi|q|} \right] \frac{q_\mu q_\nu - \delta_{\mu\nu} q^2}{|q|}, \quad (5.30)$$

and

$$\tilde{\mathcal{G}}_{\mu\nu}^{B,\text{odd}}(q) = \frac{N_B}{8\pi\lambda_B} \left(1 + \frac{4c_B^2}{q^2}\right) \sin^2(2\lambda_B |q| A_B) \epsilon_{\mu\nu\rho} q^\rho, \quad (5.31)$$

where, in the above two expressions, $A_B = \int_{c_B}^\infty \frac{dw \chi_B(w)}{q^2 + 4w^2}$. It would be nice to have a direct independent check if this is indeed the correct form of $\tilde{\mathcal{G}}_{\mu\nu}^B(q)$.

Zero temperature and zero mass

In the zero temperature and massless limit, $m_B = 0$, the pole mass c_B vanishes. So, in this limit, we have

$$\tilde{\mathcal{G}}_{-+}^B(q) = \frac{iN_B q_3}{16\pi\lambda_B} \left[e^{\pi i \lambda_B \text{sgn}(q_3)} - 1 \right] \quad (5.33)$$

This exactly matches with the zero temperature results at zero bare mass reported in [7] (see equation 44 of [7])³⁵.

Zero temperature and nonzero mass

At zero temperature, the contribution of holonomy vanishes which implies $\chi_B(w) = 1$ and $\xi_B(z) = z$. The integral appearing in (5.32) can be performed explicitly and the result is

$$\tilde{\mathcal{G}}_{-+}^B(q) = \frac{iN_B q_3}{16\pi\lambda_B} \left(1 + \frac{4c_B^2}{q_3^2} \right) \left[\exp \left(2i\lambda_B \tan^{-1} \frac{q_3}{2c_B} \right) - 1 \right] - \frac{N_B c_B}{4\pi} \quad (5.34)$$

This exactly agrees with the results [85].

Nonzero temperature and zero mass

At nonzero temperature the holonomy contribution becomes non-trivial. The result (5.32) is valid for an arbitrary holonomy distribution function. In the specific case of tabletop holonomy $\rho_B(\alpha) = \frac{\Theta(\pi|\lambda_B| - |\alpha|)}{2\pi|\lambda_B|}$, the final result for $\tilde{\mathcal{G}}_{-+}^B$ is given by (5.32), where $\chi_B(w)$ is now given by (5.19) and the thermal mass c_B is determined from (4.6) where $\xi_B(c_B)$ is now given by $\xi_B(c_B)$.

6 Check of duality

As already mentioned before, one of our goal in this paper is to check the bose-fermi duality at the level of correlation functions. In this section, we explicitly check the duality between the results that we have obtained in this paper both in the fermionic and in the bosonic theory. We check the dualities between the general results that we have got in the case of massive Chern-Simons matter theory at finite temperature with arbitrary holonomy distribution function. As mentioned in the introduction, the massive regular fermionic theory which is conjecturally dual to the massive critical bosonic theory. Below we list down the parameter map which is required to explicitly show that the results are dual to each other.

³⁵As in the case of fermions, this result can be separated into parity even and parity odd parts as discussed in [7]. The covariantized form of the parity even part in this case is given by $\frac{N_B}{16} \frac{\sin(\pi\lambda_B)}{\pi\lambda_B} \frac{q_\mu q_\nu - \delta_{\mu\nu} q^2}{|q|}$.

6.1 Duality map :

The large N 't Hooft couplings $\lambda_F = \frac{N_F}{\kappa_F}$, $\lambda_B = \frac{N_B}{\kappa_B}$, the holonomies $\rho(\alpha)$ and the exact thermal masses c_F/c_B , under duality, are mapped via

$$\begin{aligned}\lambda_F &= \lambda_B - \text{sgn}(\lambda_B), \quad \kappa_F = -\kappa_B, \quad c_F = c_B, \\ \lambda_F \rho_F(\pi - \alpha) &= \lambda_B \rho_B(\alpha) - \frac{\text{sgn}(\lambda_B)}{2\pi}.\end{aligned}\tag{6.1}$$

In the present case c_F is the thermal mass of the regular boson theory, determined by the gap equation (3.10). The gap equation for the critical bosonic scalar theory is determined from $\xi_B(c_B) = m_B^{\text{cri}}$ ³⁶. The map between the UV parameters of the two theories in this case is given by $m_F = -\lambda_B m_B^{\text{cri}}$. It follows from the definitions of $\chi_B(z)$ and $\chi_F(z)$ as given in (2.7) and (2.9),³⁷

$$\lambda_F \chi_F(z) = \lambda_B \chi_B(z) - \text{sgn}(\lambda_B)\tag{6.2}$$

Using the definition $\xi(z) = \int^z \chi(w)dw$, we can rewrite the above equation in terms of the duality map of $\xi(z)$ as below

$$\lambda_F \xi_F(z) = \lambda_B \xi_B(z) - \text{sgn}(\lambda_B)z\tag{6.3}$$

From the fermionic mass gap equation (3.10), we can write the fermionic mass m_F in terms of the bosonic variables as

$$m_F = (1 - \eta) \text{sgn}(\lambda_B) c_B - \lambda_B \xi_B(c_B)\tag{6.4}$$

where, we have used the definition $\eta = \text{sgn}(h_F \lambda_F)$ ³⁸. The parameter η can have two possible values [62]. In the case of $\eta = +1$, the dual theory in the bosonic side is the scalar theory in the unhiggsed phase, in which case, we get

$$m_F = -\lambda_B \xi_B(c_B)\tag{6.5}$$

However, we will not always write the maps between the bare parameters, instead use the map between the thermal masses, i.e., $c_F = c_B$. As we have considered the bosonic scalar theory (4.1) in the paper, we will use the choice $\eta = +1$ to compare with the results in the bosonic theory. On the other hand, we use $\eta = -1$ to predict the corresponding results for the critical boson theory in the Higgsed phase.

6.2 For spin 0 :

We dualize the result of two-point correlator of spin zero operator J_0^F given by (5.2) in terms of the bosonic variables. Using duality map (6.1) and (6.2), we rewrite (5.2) in terms

³⁶In the critical limit i.e., in the limit $b_4 \rightarrow \infty$, $m_B^2 \rightarrow \infty$ with $\frac{4\pi m_B^2}{b_4} = m_B^{\text{cri}}$ and b_6 fixed, it follows from the gap equation (4.6) that $m_B^{\text{cri}} = \xi_B(c_B)$.

³⁷In the zero temperature limit ($\beta \rightarrow \infty$), this reduces to the familiar expression $\lambda_F = \lambda_B - \text{sgn}(\lambda_B)$.

³⁸Also we use the fact that $\text{sgn}(\lambda_F) = -\text{sgn}(\lambda_B)$.

of bosonic variables as

$$\begin{aligned} \mathcal{G}_0^F(q) = & -\frac{N_B|q|}{4\pi\lambda_B} \cot \left[2\lambda_B|q| \int_{c_B}^{\infty} \frac{dw \chi_B(w)}{q^2 + 4w^2} - (1-\eta)\text{sgn}(\lambda_B) \tan^{-1} \frac{|q|}{2c_B} \right] \\ & + \frac{N_B\xi_B(c_B)}{2\pi} - (1-\eta) \frac{N_B c_B}{2\pi|\lambda_B|} . \end{aligned} \quad (6.6)$$

Depending upon the two possible values $\eta = \pm 1$, there are two cases. The regular fermionic theory is dual to the critical boson theory in the unHiggsed phase when $\eta = +1$. On the other hand, in the case of $\eta = -1$, the regular fermionic theory is dual to the critical bosonic theory in the Higgsed phase (this is the case when $m_B^{\text{cri}} < 0$) (for details about this see [62]).

6.2.1 Duality check in the unhiggsed phase of critical bosons:

As mentioned above, in the unhiggsed phase, $\eta = +1$. This implies that the fermionic result

$$\mathcal{G}_0^{uH}(q) = -\frac{N_B|q|}{4\pi\lambda_B} \cot \left[2\lambda_B|q| \int_{c_B}^{\infty} \frac{dw \chi_B(w)}{q^2 + 4w^2} \right] + \frac{N_B\xi_B(c_B)}{2\pi} . \quad (6.7)$$

This matches with the result of two-point correlator of spin zero operator in the critical boson theory given by (5.24) upto contact terms. Thus, we see that under duality, J_0^F maps to J_0^B (more precisely to \tilde{J}_0^B given in (5.22)) ³⁹.

6.2.2 Prediction for the higgsed phase of critical bosons:

In the higgsed phase, on the other hand, $\eta = -1$. So, the predicted answer of the two-point correlator of single trace, spin zero ‘scalar’ current J_0^H in the higgsed phases of critical bosons is given by ⁴⁰

$$\begin{aligned} \mathcal{G}_0^H(q) = & -\frac{N_B|q|}{4\pi\lambda_B} \cot \left[2\lambda_B|q| \int_{c_B}^{\infty} \frac{dw \chi_B(w)}{q^2 + 4w^2} - 2\text{sgn}(\lambda_B) \tan^{-1} \frac{|q|}{2c_B} \right] \\ & + \frac{N_B\xi_B(c_B)}{2\pi} - \frac{N_B c_B}{\pi|\lambda_B|} . \end{aligned} \quad (6.8)$$

It is an interesting exercise to compute this (exactly in large N) directly from the higgsed phases of bosons, which we leave for future work.

³⁹As mentioned earlier, the single trace spin-zero scalar operator J_0^B in the case of critical boson theory (5.21) is simply the Lagrange multiplier field σ_B appearing in the action (5.21) [7, 9].

⁴⁰The single trace, spin-zero scalar operator in the Higgsed phase of critical bosons which is dual to corresponding spin zero operator $J_0^F = \bar{\psi}\psi$ in the massive regular fermionic theory, is given by $J_0^H = \overline{W}_\mu W^\mu + Z_\mu Z^\mu$ [85].

6.3 For spin 1 :

The final result that we have obtained for spin 1 current correlator in the fermionic theory, is given by (5.9). Under duality maps (6.1), (6.2) and (6.3), $\mathcal{G}_{-+}^F(q)$ can be rewritten as

$$\begin{aligned} \mathcal{G}_{-+}^F(q) = & -\frac{iN_B q_3}{16\pi\lambda_B} \left(1 - \eta \frac{2\text{isgn}(\lambda_B)c_B}{q_3}\right)^2 \left[-\left(\frac{q_3 + 2\text{isgn}(\lambda_B)c_B}{q_3 - 2\text{isgn}(\lambda_B)c_B}\right) \exp\left(4i\lambda_B q_3 \int_{c_B}^{\infty} \frac{dw}{q_3^2 + 4w^2} \chi_B(w)\right) - 1 \right] \\ & + \frac{N_B \xi_B(c_B)}{4\pi} - \frac{N_B c_B}{4\pi|\lambda_B|} . \end{aligned} \quad (6.9)$$

Depending upon the two possible values of η , we consider two phases separately below.

6.3.1 Duality check in the unhiggsed phase of critical bosons :

As mentioned before, in the unhiggsed phase $\eta = +1$. So, it follows from (6.9) that it dualizes to the following result in the unhiggsed phase of bosons,

$$\mathcal{G}_{-+}^{uH}(q) = \frac{iN_B q_3}{16\pi\lambda_B} \left(1 + \frac{4c_B^2}{q_3^2}\right) \left[\exp\left(4i\lambda_B q_3 \int_{c_B}^{\infty} \frac{dw}{q_3^2 + 4w^2} \chi_B(w)\right) - 1 \right] + \frac{N_B \xi_B(c_B)}{4\pi} + \frac{iN_B q_3}{8\pi\lambda_B} . \quad (6.10)$$

This exactly matches with the bosonic results (5.32) upto a contact term $\frac{iN_B q_3}{8\pi\lambda_B}$. Thus, we see that J_μ^F maps to J_μ^B under duality.

6.3.2 Prediction for the higgsed phase of critical bosons :

In the higgsed phase ($\eta = -1$), the predicted result for the two-point correlator of the single trace spin one current is ⁴¹

$$\begin{aligned} \mathcal{G}_{-+}^H(q) = & \frac{iN_B q_3}{16\pi\lambda_B} \left(1 + \frac{2\text{isgn}(\lambda_B)c_B}{q_3}\right)^2 \left[\left(\frac{q_3 + 2\text{isgn}(\lambda_B)c_B}{q_3 - 2\text{isgn}(\lambda_B)c_B}\right) \exp\left(4i\lambda_B q_3 \int_{c_B}^{\infty} \frac{dw}{q_3^2 + 4w^2} \chi_B(w)\right) + 1 \right] \\ & + \frac{N_B \xi_B(c_B)}{4\pi} - \frac{N_B c_B}{4\pi|\lambda_B|} . \end{aligned} \quad (6.11)$$

We leave for the future work, the matching of this prediction with the exact large N computation of $\langle J_\mu^H J_\nu^H \rangle$ directly in the Higgsed phases of critical bosons ⁴².

7 Discussions

In this work we have obtained several two-point momentum space correlators of gauge invariant, single trace operators in Chern-Simons coupled to massive fundamental matter

⁴¹A possible ‘covariant’ form of $\langle J_\mu^H(q') J_\nu^H(q) \rangle = \mathcal{G}_{\mu\nu}^H (2\pi)^3 \delta^{(3)}(q' + q)$, is obtained from (5.7) and (5.8) with the substitution of $\text{sgn}(h_F) = -\text{sgn}(\lambda_F)$ (which is equal to $\text{sgn}(\lambda_B)$) and applying the duality map given in subsection 6.1. It would be nice to have a direct independent check if this is indeed the case.

⁴²In the Higgsed phase of critical bosons in the unitary gauge [62], the $U(1)$ current J_μ^H is proportional to the Z bosons, i.e., $J_\mu^H \propto Z_\mu$ [85]. So, in this case, $\mathcal{G}_{\mu\nu}^H \propto \langle Z_\mu Z_\nu \rangle$.

theories in the large- N 't Hooft limit, at finite temperature and considering arbitrary holonomy distribution for gauge holonomy. One of the challenging aspects of our work was to solve the corresponding Schwinger-Dyson equations for the correlators because we chose to work with an arbitrary holonomy distribution, with massive matter at finite temperature. However, we have been able to overcome this technical difficulty and have solved these equations analytically by explicitly performing the loop momenta integrals. The results for the two-point correlators are indeed very simple in form. We have seen that in different limiting cases, the results obtained in this work agree with the existing previous results [7, 9, 29, 32, 80]. We have also explicitly checked the bose-fermi duality between the results of the regular fermionic theory and the critical bosonic scalar theory. We now discuss about the applications of the results obtained and further possible generalizations of the analysis done in this paper.

We computed the two-point correlators of the spin one conserved current of single trace operators both in the case of massive fermionic theory and in the case of massive bosonic scalar theory. We used the duality map to predict for the two-point function of the spin one current in the case of the higgsed phases of bosons. The analysis done in this paper is in the Euclidean signature and with the external momentum $q_\mu \equiv (0, 0, q_3)$. By analytic continuations, the results in the Lorentzian signature can be obtained by Wick rotating back to Minkowski space $q_3 \rightarrow i\omega$. In [29], it was highlighted that the two-point correlator of the $U(1)$ current can be used to calculate the conductivity tensor by applying the Kubo formula. Their analysis was done at zero temperature in the fermionic theory. Also in [32], the study of conductivity was continued in Chern-Simons fermionic matter theories. However in [32], they considered the massless regular fermionic matter coupled to Chern-Simons gauge fields at finite temperature with table-top holonomy distribution. Moreover the results in [32] were given as an integral expressions. Our results of two-point correlators of spin-one current in the massive fermionic theory at finite temperature is a generalization of [29, 32] and the final result is given in a much simplified form (5.9). Following [29, 32], this can be used to study the conductivity of the massive regular fermionic theory at finite temperature by applying the Kubo formula. In section E, we have commented on the computations of correlators of arbitrary spin s currents and we have seen a general structure of the results; following the same procedure used in the case of spin-zero and spin-one operators, one can get, e.g., the stress tensor two-point correlation function which can be used to study the viscosity of these theories [29].

As already discussed above, in the massive regular fermionic theory the results can be dual to either critical bosonic scalar theory in the unHiggsed phase or to the Higgsed phase of critical bosons depending upon the signs of $\text{sgn}(h_F \lambda_F)$. We have explicitly checked the duality of the regular fermionic theory with the critical bosonic scalar theory in the unHiggsed phase, which is the case with $\text{sgn}(h_F \lambda_F) = +1$. Also, in the case of $\text{sgn}(h_F \lambda_F) = -1$, we have given the predictions for the two-point correlators of the corresponding current operators in the Higgsed phases of critical bosons. One may compute these correlators to all orders in 't Hooft coupling in the large- N limit directly in the Higgsed phases of critical bosons coupled to Chern-Simons theory [62, 68] to explicitly check the duality in the higgsed phase. The technicalities of performing exact computations in the higgsed phases of bosons

are rather involved, and we leave this exercise for future work. It is realized that the spin-one current of the massive fermionic theory is dual to the Z bosons in the Higgsed phases [85]. So, the dualized result of $\langle J_\mu^F J_\nu^F \rangle$ in the higgsed phases is proportional to the Z boson propagator $\langle Z_\mu Z_\nu \rangle$. One can try to explicitly compute the exact Z boson propagator directly in the Higgsed phases of bosonic theory [62], at least in large N limit, and match with the results predicted here.

The pair of theories that we have considered in this paper has global $U(1)$ symmetry. One can consider the analysis presented here in the presence of a chemical potential μ by turning on a constant background gauge field $\mathcal{A}_\nu = i\mu\delta_{\nu 3}$. At least in the case when $|\mu| < c_B$ or $|\mu| < c_F$, the analysis seems to go through exactly the same way and the final result in presence of chemical potential is obtained by making the substitution $\alpha \rightarrow \alpha - i\beta\mu$ in the integrand (except in the holonomy distribution $\rho(\alpha)$) of holonomy integral appearing in the results obtained in this paper at the zero chemical potential. It will be nice to explicitly check if the structure of the results remain the same even in the case when the chemical potential is larger than the thermal mass c_B or c_F [83]. It seems like, even in the case when $|\mu| > c_B$ or $|\mu| > c_F$, the structure of the results of the correlators of this paper will remain the same; the only modifications will be through the functions $\chi_B(w)$ and $\chi_F(w)$ due to the contour deformations of the holonomy integrals away from the unit circle as prescribed in detail in [83]. We leave the careful analysis about this as a future exercise.

As discussed before, as part of the analysis in the bosonic scalar theory, following [17], we have generalized the computations of four-point functions of fundamental scalars to finite temperature. As in [17], this might be useful to study the scattering of fundamental scalars at finite temperature. In footnotes 30, 31, 34 and 41, we have provided a possible ‘covariantized’ form of the two-point correlators of the spin one currents of the fermionic and bosonic theories, valid for arbitrary values of the external momenta q . It would be nice to have an independent check of these by direct computations without assuming $q_\pm = 0$. One can also extend the computations of two-point current correlators to higher point correlators of single trace operators. We have already computed in this paper various exact insertion vertices. And one can use these to explicitly compute the higher-point current correlators by generalizing the existing results (e.g., three-point correlators [7, 9], four-point correlators [65, 79]) to the massive matter theories at finite temperature. Presumably, one may also use the results of two-point correlators of single trace operators obtained in this paper to compute the thermal one-point functions using bootstrap approaches. In this paper we have not paid much attention to the contact terms appearing in the results of the two point-correlators. It would be interesting to understand if the contact terms appearing in this paper have any physical significance. Also, it would be interesting to extend the analysis of this paper by going beyond large- N limit, taking into account the contributions from the non-planar diagrams. we leave all these problems for future work.

Acknowledgements

The author would like to thank S. Chakraborty, S. Jain and N. Prabhakar for helpful conversations. The author would like to thank S. Minwalla for valuable discussions and

comments. The author would also like to acknowledge support from the Infosys Endowment for the study of the Quantum Structure of Spacetime.

A Conventions and useful definitions

We work in the three dimensional spacetime with coordinates (x^1, x^2, x^3) , where x^3 is the Euclidean time coordinate. For a vector $a^\mu \equiv (a^1, a^2, a^3)$, the corresponding lightcone components are defined by $a^\pm \equiv a_\mp = \frac{a^1 \pm i a^2}{\sqrt{2}}$ and the non-zero components of the metric in the lightcone coordinates are given by $\delta_{+-} = \delta_{-+} = \delta_{33} = 1$. The totally antisymmetric Levi-Civita tensor $\epsilon^{\mu\nu\rho}$ that appears in the paper is normalized such that $\epsilon^{123} = \epsilon_{123} = 1$. In the lightcone coordinates, $\epsilon_{+-3} = \epsilon^{-+3} = i$. The momenta squared are given by $p^2 = p_1^2 + p_2^2 + p_3^2 = 2p_+p_- + p_3^2 = p_s^2 + p_3^2$. Here, $p_s = +\sqrt{2p_+p_-}$ is the radial momentum in the spatial momentum plane. Throughout the paper, the overall external momentum flowing through the insertion vertices of the single trace current operators is denoted by q . We work in the momentum choice $q_\pm = 0$. At finite temperature, the component q_3 of the external momentum q entering an insertion vertex is quantized [32]; however, we don't explicitly write the quantized version of it and write it simply as q_3 .

Below, we list down a few definitions of useful quantities that have appeared throughout the paper. Following [62], we use (2.7) to define the function $\xi_B(z)$ as an integral over $\chi_B(z)$, i.e., $\xi_B(z) = \int^z \chi_B(w)dw$. The final expression for $\xi_B(z)$ is given by

$$\xi_B(z) = \frac{1}{\beta} \int_{-\pi}^{\pi} \rho_B(\alpha) d\alpha \left[\ln 2 \sinh \left(\frac{\beta z + i\alpha}{2} \right) + \ln 2 \sinh \left(\frac{\beta z - i\alpha}{2} \right) \right]. \quad (\text{A.1})$$

Following [62], we use the regularization where, $\xi_B(\infty)$ is a pure divergence term and is dropped away (for details, see discussion around equation 2.75 of [62]).

Similarly, for the fermionic theory, using (2.9) we define the function $\xi_F(z) = \int^z \chi_F(w)dw$. As before, we use the regularization where, $\xi_F(\infty)$ is purely a divergent term and is thrown away. Explicitly, $\xi_F(z)$ is given by

$$\xi_F(z) = \frac{1}{\beta} \int_{-\pi}^{\pi} \rho_F(\alpha) d\alpha \left[\ln 2 \cosh \left(\frac{\beta z + i\alpha}{2} \right) + \ln 2 \cosh \left(\frac{\beta z - i\alpha}{2} \right) \right]. \quad (\text{A.2})$$

We define another function which appears in the intermediate steps of the bosonic computations

$$F_B(z, q_3) = \frac{\chi_B(z)}{(q_3^2 + 4z^2)}. \quad (\text{A.3})$$

Similarly, in the case of fermions, we define

$$F_F(z, q_3) = \frac{\chi_F(z)}{(q_3^2 + 4z^2)}. \quad (\text{A.4})$$

Another set of useful definitions are

$$H_B(z, q_3) = \exp \left(4i\lambda_B q_3 \int^z F_B(w, q_3) dw \right) = \exp \left(4i\lambda_B q_3 \int^z \frac{dw \chi_B(w)}{q_3^2 + 4w^2} \right), \quad (\text{A.5})$$

and

$$H_F(z, q_3) = \exp \left(4i\lambda_F q_3 \int^z F_F(w, q_3) dw \right) = \exp \left(4i\lambda_F q_3 \int^z \frac{dw \chi_F(w)}{q_3^2 + 4w^2} \right). \quad (\text{A.6})$$

Before concluding this appendix, we mention here that the multivalued function $\tan^{-1}(x)$ that appears in this paper is defined by choosing the branch $-\frac{\pi}{2} < \tan^{-1}(x) < \frac{\pi}{2}$. With this definition, it follows that $\tan^{-1}(-x) = -\tan^{-1}(x)$. Also, we use the property that $\tan^{-1}(x) + \tan^{-1}(\frac{1}{x}) = \text{sgn}(x)\frac{\pi}{2}$.

B Angular integrals :

The angular integrals that are used in this paper, are given by

$$\int_0^{2\pi} \frac{d\theta_l}{2\pi} \frac{(l_-)^j}{(l-k)_-} = (k_-)^{j-1} \left[-\theta(-j) + \Theta(l_s - k_s) \right]. \quad (\text{B.1})$$

Here, on the right-hand side, θ and Θ both are unit step functions. The small θ - function is used for the power of the momenta l_- appearing in the numerator of the integrand and it is assumed to be defined such that $\theta(j=0) = 1$. This relation is true for $\forall j \in \mathbb{Z}$. The above integration result (B.1) can be easily obtained by choosing $l_- = \frac{\ell_s}{\sqrt{2}} e^{i\theta_\ell}$ and converting the resulting angular integral into a complex contour integral over a unit circle by substituting $z_\ell = e^{i\theta_\ell}$.

C Useful integrals

In this appendix, we list down (non-vanishing) useful integrals which are repeatedly used in the main text.

C.1 Integrals in bosonic theory

We list here a set of useful integrals and we present the integration results in terms of a quantity $a(k_s) = +\sqrt{k_s^2 + c_B^2}$, and the function $\chi_B(z)$ given by (2.7). We don't write the temperature dependence β explicitly in the arguments of the functions that are being defined, but is implicitly assumed. The external momentum component q_3 appearing in the integrals is quantized but the explicit quantized form is not written for simplicity.

Integration B1 :

$$\begin{aligned} \int \frac{\mathcal{D}_B k_3}{2\pi} \frac{1}{k_3^2 + a^2(k_s)} &= \frac{1}{\beta} \int_{-\pi}^{\pi} \rho_B(\alpha) d\alpha \sum_{n_k=-\infty}^{\infty} \frac{1}{\left(\frac{2n_k\pi + \alpha}{\beta}\right)^2 + a^2(k_s)} \\ &= \frac{1}{2a(k_s)} \chi_B(a(k_s)) \end{aligned} \quad (\text{C.1})$$

where, $a(k_s) = +\sqrt{k_s^2 + c_B^2}$ and $\chi_B(z)$ is given by (2.7). In the zero-temperature limit, i.e., in $\beta \rightarrow \infty$ limit, it is clear from (2.7) that $\lim_{\beta \rightarrow \infty} \chi_B(z) = 1$. In the zero temperature

limit, the above integral (C.1) reduces to the usual integral

$$\int_{-\infty}^{\infty} \frac{dk_3}{2\pi} \frac{1}{k_3^2 + a^2(k_s)} = \frac{1}{2a(k_s)} \quad (\text{C.2})$$

We now consider the following integration

$$\begin{aligned} \int \frac{\mathcal{D}_B^3 k}{(2\pi)^3} \frac{1}{k_3^2 + a^2(k_s)} &= \int_0^\infty \frac{k_s dk_s}{2\pi} \frac{\chi_B(a(k_s))}{2a(k_s)} \\ &= \frac{1}{4\pi} \int_{c_B}^\infty da(k_s) \chi_B(a(k_s)) \\ &= \frac{1}{4\pi} \left(\xi_B(\infty) - \xi_B(c_B) \right) \\ &= -\frac{\xi_B(c_B)}{4\pi} \end{aligned} \quad (\text{C.3})$$

where, we have used the definition $\xi'_B(z) = \chi_B(z) \Rightarrow \xi_B(z) = \int^z \chi_B(w) dw$. We use the regularization scheme [62] (see around equation 2.75 of [62]), where, $\xi_B(\infty)$ is a pure divergence term and is dropped away.

Integration B2 :

There is another integration which is mostly used in the main text; this is given by

$$\begin{aligned} &\int \frac{\mathcal{D}_B k_3}{2\pi} \frac{1}{(k_3^2 + a^2(k_s))((k_3 + q_3)^2 + a^2(k_s))} \\ &= \frac{1}{\beta} \int_{-\pi}^{\pi} \rho_B(\alpha) d\alpha \sum_{n_k=-\infty}^{\infty} \frac{1}{\left(\left(\frac{2n_k\pi + \alpha}{\beta} \right)^2 + a^2(k_s) \right) \left(\left(\frac{2n_k\pi + \alpha}{\beta} + q_3 \right)^2 + a^2(k_s) \right)} \\ &= \frac{\chi_B(a(k_s))}{a(k_s)(q_3^2 + 4a^2(k_s))} \end{aligned} \quad (\text{C.4})$$

where, $\chi_B(z)$ is the same function defined in (2.7). As expected, in the zero temperature limit, this reduces to the standard integral

$$\int_{-\infty}^{\infty} \frac{dk_3}{2\pi} \frac{1}{(k_3^2 + a^2(k_s))((k_3 + q_3)^2 + a^2(k_s))} = \frac{1}{a(k_s)(q_3^2 + 4a^2(k_s))} \quad (\text{C.5})$$

Integration B3 :

Another useful integral is given by

$$\begin{aligned}
& \int \frac{\mathcal{D}_B k_3}{2\pi} \frac{k_3(k_3 + q_3)}{(k_3^2 + a^2(k_s))((k_3 + q_3)^2 + a^2(k_s))} \\
&= \frac{1}{\beta} \int_{-\pi}^{\pi} \rho_B(\alpha) d\alpha \sum_{n_k=-\infty}^{\infty} \frac{\left(\frac{2n_k\pi+\alpha}{\beta}\right)\left(\frac{2n_k\pi+\alpha}{\beta} + q_3\right)}{\left(\left(\frac{2n_k\pi+\alpha}{\beta}\right)^2 + a^2(k_s)\right)\left(\left(\frac{2n_k\pi+\alpha}{\beta} + q_3\right)^2 + a^2(k_s)\right)} \\
&= \frac{a(k_s) \chi_B(a(k_s))}{(q_3^2 + 4a^2(k_s))}
\end{aligned} \tag{C.6}$$

where, $\chi_B(z)$ is defined by (2.7). This reduces, in the zero temperature limit, to the usual integral

$$\int_{-\infty}^{\infty} \frac{dk_3}{2\pi} \frac{k_3(k_3 + q_3)}{(k_3^2 + a^2(k_s))((k_3 + q_3)^2 + a^2(k_s))} = \frac{a(k_s)}{(q_3^2 + 4a^2(k_s))} \tag{C.7}$$

C.2 Integrals in fermionic Theory

The integrals that appear in the fermionic theory are of the same structure as that appear in the bosonic theory listed above. Below we provide a set of useful integrals

Integration F1 :

$$\begin{aligned}
& \int \frac{\mathcal{D}_F k_3}{2\pi} \frac{1}{k_3^2 + a^2(k_s)} = \frac{1}{\beta} \int_{-\pi}^{\pi} \rho_F(\alpha) d\alpha \sum_{n_k=-\infty}^{\infty} \frac{1}{\left(\frac{(2n_k+1)\pi+\alpha}{\beta}\right)^2 + a^2(k_s)} \\
&= \frac{1}{2a(k_s)} \chi_F(a(k_s))
\end{aligned} \tag{C.8}$$

where, $a(k_s) = +\sqrt{k_s^2 + c_F^2}$ and $\chi_F(z)$ is given by (2.9). We don't explicitly write the temperature dependence β in the arguments of the functions that are being defined, but is implicitly assumed. In the zero-temperature limit, i.e., in the $\beta \rightarrow \infty$ limit, it follows from (2.9), $\lim_{\beta \rightarrow \infty} \chi_F(a(k_s)) = 1$. Thus this integral reduces, in the zero temperature limit, to the known integral

$$\int_{-\infty}^{\infty} \frac{dk_3}{2\pi} \frac{1}{k_3^2 + a^2(k_s)} = \frac{1}{2a(k_s)} \tag{C.9}$$

As in the case of bosons, we consider here the following integration

$$\begin{aligned}
& \int \frac{\mathcal{D}_F^3 k}{(2\pi)^3} \frac{1}{k_3^2 + a^2(k_s)} = \int_0^{\infty} \frac{k_s dk_s}{2\pi} \frac{\chi_F(a(k_s))}{2a(k_s)} \\
&= \frac{1}{4\pi} \int_{c_F}^{\infty} da(k_s) \chi_F(a(k_s)) \\
&= \frac{1}{4\pi} \left(\xi_F(\infty) - \xi_F(c_F) \right) \\
&= -\frac{\xi_F(c_F)}{4\pi}
\end{aligned} \tag{C.10}$$

where, we have used the definition $\xi'_F(z) = \chi_F(z) \Rightarrow \xi_F(z) = \int^z \chi_F(w)dw$. We use the regularization scheme, where, $\xi_F(\infty)$ is a pure divergent term and is dropped.

Integration F2 :

As in the case of bosons, there is another useful integral

$$\begin{aligned}
& \int \frac{\mathcal{D}_F k_3}{2\pi} \frac{1}{(k_3^2 + a^2(k_s))((k_3 + q_3)^2 + a^2(k_s))} \\
&= \frac{1}{\beta} \int_{-\pi}^{\pi} \rho_F(\alpha) d\alpha \sum_{n_k=-\infty}^{\infty} \frac{1}{\left(\left(\frac{(2n_k+1)\pi+\alpha}{\beta} \right)^2 + a^2(k_s) \right) \left(\left(\frac{(2n_k+1)\pi+\alpha}{\beta} + q_3 \right)^2 + a^2(k_s) \right)} \\
&= \frac{\chi_F(a(k_s))}{a(k_s)(q_3^2 + 4a^2(k_s))}
\end{aligned} \tag{C.11}$$

where, $\chi_F(z)$ is given by (2.9). In the zero temperature limit, this integral reduces to the standard integral

$$\int_{-\infty}^{\infty} \frac{dk_3}{2\pi} \frac{1}{(k_3^2 + a^2(k_s))((k_3 + q_3)^2 + a^2(k_s))} = \frac{1}{a(k_s)(q_3^2 + 4a^2(k_s))} \tag{C.12}$$

Integration F3 :

The following integral is also useful for the computations in this paper

$$\begin{aligned}
& \int \frac{\mathcal{D}_F k_3}{2\pi} \frac{k_3(k_3 + q_3)}{(k_3^2 + a^2(k_s))((k_3 + q_3)^2 + a^2(k_s))} \\
&= \frac{1}{\beta} \int_{-\pi}^{\pi} \rho_B(\alpha) d\alpha \sum_{n_k=-\infty}^{\infty} \frac{\left(\frac{(2n_k+1)\pi+\alpha}{\beta} \right) \left(\frac{(2n_k+1)\pi+\alpha}{\beta} + q_3 \right)}{\left(\left(\frac{(2n_k+1)\pi+\alpha}{\beta} \right)^2 + a^2(k_s) \right) \left(\left(\frac{(2n_k+1)\pi+\alpha}{\beta} + q_3 \right)^2 + a^2(k_s) \right)} \\
&= \frac{a(k_s) \chi_F(a(k_s))}{(q_3^2 + 4a^2(k_s))}
\end{aligned} \tag{C.13}$$

In the zero temperature limit, this integral reduces to the familiar integral

$$\int_{-\infty}^{\infty} \frac{dk_3}{2\pi} \frac{k_3(k_3 + q_3)}{(k_3^2 + a^2(k_s))((k_3 + q_3)^2 + a^2(k_s))} = \frac{a(k_s)}{(q_3^2 + 4a^2(k_s))} \tag{C.14}$$

D Computation of thermal four-point function of fundamental scalars

In this appendix, we generalize the computation of four-point function of fundamental scalars in the regular boson theory presented in [17](see e.g. section 4 and appendix D.3 of [17] for details of computation at zero temperature) to non-zero temperature. The




Figure 7. Schinger-Dyson equation for offshell four-point function of scalars. The elliptic blob denotes the exact scalar four-point function and is denoted by $\mathcal{A}(p, k, q)$. The thick vertical line corresponds to the ‘effective single particle exachange’ four-point amplitude and is denoted by $\mathcal{A}_0(p, k, q)$; this amplitude is given by the sum of the diagrams of Fig.8 and Fig.9. The filled circle denotes the exact scalar propagator.

Schwinger-Dyson equations and the feynman diagrams are all the same as in [17]. We redo the computations here with the relevant modifications due to the finite temperature effect including holonomy of gauge fields. Summing all the diagrams in Fig.8 (or see figure 19 of




Figure 8. Some of the diagrams contributing to one loop four-point amplitude $\mathcal{A}_0(p, k, q)$. The filled circle denotes the exact scalar propagator. The assignments of momenta in the external legs are the same as in Fig.7.

[17] (also including the diagram corresponding to the $(\bar{\phi}\phi)^3$ interaction vertex), the one-loop effective one-particle exchange can be written as

$$\begin{aligned}
 NA_{\text{one-loop}} &= \sum_{i=1}^6 NA_i = -8\pi^2 \lambda_B^2 \int \frac{\mathcal{D}_B^3 \ell}{(2\pi)^3} \frac{1}{\ell^2 + c_B^2} \\
 &= -8\pi^2 \lambda_B^2 \left(-\frac{\xi_B(c_B)}{4\pi} \right) \\
 &= 2\pi \lambda_B^2 \xi_B(c_B)
 \end{aligned} \tag{D.1}$$

where, the function $\xi_B(c_B)$ is given by (A.1).

The contact contribution to the four-point function from the b_4 vertex shown in the Fig.9(a) is given by $-(\frac{b_4}{2N_B}) \cdot 2 = -\frac{b_4}{N_B}$. As given in the action (4.1), contribution from the b_6




Figure 9. Some of the diagrams contributing to the ‘effective single particle exchange’ four-point amplitude $\mathcal{A}_0(p, k, q)$. Diagram (a) corresponds to the four-point interaction vertex. Diagram (b) corresponds to the contribution of the interaction vertex $(\bar{\phi}\phi)^3$ to the four-point amplitude; here, the filled circular dot denotes the exact scalar propagator. Diagram (c) corresponds to the four-point amplitude due to the tree-level gauge boson exchange.

vertex to the four point function is shown in Fig.9(b) and is given by $(-\frac{b_6}{N_B^2})(-\frac{\xi_B(c_B)}{4\pi}N_B) = \frac{\xi_B(c_B)b_6}{4\pi N_B}$. At the tree level, the contribution from the diagram shown in Fig.9(c) is given by

$$\begin{aligned} A_{\text{tree}} &= \mathcal{V}^\mu(p+q, -(k+q))G_{\nu\mu}^B(p-k)\mathcal{V}^\nu(k, -p) = \frac{4\pi}{\kappa_B} \frac{q^\mu(k+p)^\nu \epsilon_{\nu-\mu}}{(p-k)_-} \\ &= -\frac{4\pi i \lambda_B q_3}{N_B} \frac{(k+p)_-}{(k-p)_-} \end{aligned} \quad (\text{D.2})$$

So, the total contribution to the ‘effective one-particle exchange’ four-point amplitude $\mathcal{A}_0(p, k, q)$ is given by

$$\begin{aligned} N_B \mathcal{A}_0(p, k, q_3) &= -4\pi i \lambda_B q_3 \frac{(k+p)_-}{(k-p)_-} + 2\pi \lambda_B^2 \xi_B(c_B) - b_4 + \frac{\xi(c_B)b_6}{4\pi} \\ &= -4\pi i \lambda_B q_3 \frac{(k+p)_-}{(k-p)_-} + \tilde{b}_4 \end{aligned} \quad (\text{D.3})$$

where, in the last line \tilde{b}_4 is defined by

$$\tilde{b}_4 = -b_4 + \frac{\xi_B(c_B)b_6}{4\pi} + 2\pi \lambda_B^2 \xi_B(c_B) \quad (\text{D.4})$$

D.1 Schwinger-Dyson Equation

The exact thermal scalar four-point function that we are interested in is defined by

$$\langle \phi^i(p+q) \bar{\phi}_j(-(k+q)) \phi^m(k) \bar{\phi}_n(-p) \rangle = \mathcal{A}_{jn}^{im}(p, k, q) (2\pi)^3 \delta^3(0) \quad (\text{D.5})$$

For the purpose of calculation of the current correlators, we are interested in the following contraction of the color indices

$$\mathcal{A}_{jn}^{im}(p, k, q) = \mathcal{A}(p, k, q) \delta_n^i \delta_j^m \quad (\text{D.6})$$

As explained in details in [17], in order to calculate the exact four-point amplitude we solve the Schwinger-Dyson equation given in Figure 5 of [17].

In the mathematical form, the Schwinger Dyson equation can be written in the following two alternative but equivalent forms

$$\mathcal{A}(p, k, q_3) = \mathcal{A}_0(p, k, q_3) + N_B \int \frac{\mathcal{D}_B^3 r}{(2\pi)^3} \frac{\mathcal{A}_0(p, r, q_3) \mathcal{A}(r, k, q_3)}{(r^2 + c_B^2)((r + q)^2 + c_B^2)} \quad (\text{D.7})$$

$$\mathcal{A}(p, k, q_3) = \mathcal{A}_0(p, k, q_3) + N_B \int \frac{\mathcal{D}_B^3 r}{(2\pi)^3} \frac{\mathcal{A}(p, r, q_3) \mathcal{A}_0(r, k, q_3)}{(r^2 + c_B^2)((r + q)^2 + c_B^2)} \quad (\text{D.8})$$

These are the two equations that need to be solved for $\mathcal{A}(p, k, q_3)$. We work in the choice $q_{\pm} = 0$, and the expression (D.3) for $\mathcal{A}_0(p, k, q_3)$ is independent of p_3 . So, from the equation (D.7), we can write $\partial_{p_3} \mathcal{A}(p, k, q_3) = 0$. In the same way, we can write from the equation (D.8), $\partial_{k_3} \mathcal{A}(p, k, q_3) = 0$.

This implies that $\mathcal{A}(p, k, q_3)$ is independent of p_3 and k_3 . So, using the integration result (C.4), we integrate r_3 momenta. And so, from (D.7) after integrating over r_3 , and using the definition (A.3) we get

$$\mathcal{A}(\vec{p}, \vec{k}, q_3) = \mathcal{A}_0(\vec{p}, \vec{k}, q_3) + N_B \int \frac{d^2 \vec{r}}{(2\pi)^2} \frac{F_B(a(r_s), q_3)}{a(r_s)} \mathcal{A}_0(\vec{p}, \vec{r}, q_3) \mathcal{A}(\vec{r}, \vec{k}, q_3) \quad (\text{D.9})$$

where, $a(r_s) = +\sqrt{r_s^2 + c_B^2}$. Similarly from (D.8) after integrating over r_3 , we get

$$\mathcal{A}(\vec{p}, \vec{k}, q_3) = \mathcal{A}_0(\vec{p}, \vec{k}, q_3) + N_B \int \frac{d^2 \vec{r}}{(2\pi)^2} \frac{F_B(a(r_s), q_3)}{a(r_s)} \mathcal{A}(\vec{p}, \vec{r}, q_3, \lambda_B) \mathcal{A}_0(\vec{r}, \vec{k}, q_3) \quad (\text{D.10})$$

Treating the lightcone momenta as the complex coordinates, $p_- \rightarrow z$, $p_+ \rightarrow \bar{z}$, and using the standard complex integration formula $\frac{\partial}{\partial \bar{z}} \frac{1}{(z-a)} = 2\pi \delta^{(2)}(\vec{r}-\vec{a})$, from (D.3) we calculate

$$N_B \partial_{p_+} \mathcal{A}_0(\vec{p}, \vec{k}, q_3) = (2\pi)^2 4i \lambda_B q_3 p_- \delta^{(2)}(\vec{p} - \vec{k}) \quad (\text{D.11})$$

From equation (D.9) and the integration result (A.3), one can easily find

$$\partial_{p_+} (\mathcal{A}(\vec{p}, \vec{k}, q_3) - \mathcal{A}_0(\vec{p}, \vec{k}, q_3)) = \frac{4i \lambda_B q_3 p_-}{a(p_s)} F_B(a(k_s), q_3) \mathcal{A}(\vec{p}, \vec{k}, q_3) \quad (\text{D.12})$$

Similarly from (D.10), one gets

$$\partial_{k_+} (\mathcal{A}(\vec{p}, \vec{k}, q_3) - \mathcal{A}_0(\vec{p}, \vec{k}, q_3)) = -\frac{4i \lambda_B q_3 k_-}{a(r_s)} F_B(a(k_s), q_3) \mathcal{A}(\vec{p}, \vec{k}, q_3) \quad (\text{D.13})$$

Equations (D.12) and (D.13) may be regarded as the first order ordinary differential equations in the variables p_+ and k_+ respectively. These equations are easily solved. For convenience, we define the following function

$$H_B(z, q_3) = \exp \left(4i \lambda_B q_3 \int^z F_B(w, q_3) dw \right) \quad (\text{D.14})$$

The inverse of this function is given by $H_B^{-1}(z, q_3) = \frac{1}{H_B(z, q_3)}$. From (D.12), we get

$$\partial_{p_+} \left(H_B^{-1}(a(p_s), q_3) \mathcal{A}(\vec{p}, \vec{k}, q_3) \right) = H_B^{-1}(a(p_s), q_3) \partial_{p_+} \mathcal{A}_0(\vec{p}, \vec{k}, q_3) \quad (\text{D.15})$$

In the same way, the equation (D.13) can be recast into the following form

$$\partial_{k_+} \left(H_B(a(k_s), q_3) \mathcal{A}(\vec{p}, \vec{k}, q_3) \right) = H_B(a(p_s), q_3) \partial_{k_+} \mathcal{A}_0(\vec{p}, \vec{k}, q_3) \quad (\text{D.16})$$

The equations (D.15) and (D.16) can be easily solved by integrations. We have already found in (D.11) that $\partial_{p_+} \mathcal{A}_0(\vec{p}, \vec{k}, q_3) \propto \delta^{(2)}(\vec{p} - \vec{k})$; so, we use the formula

$$f(\vec{r}) \delta^{(2)}(\vec{r} - \vec{a}) = f(\vec{a}) \delta^{(2)}(\vec{r} - \vec{a})$$

Thus, we can replace the prefactors on the RHS of equation (D.15) by replacing \vec{p} by \vec{k} . By integrating both sides of (D.15) and multiplying by N_B and introducing the integration constant $h(\vec{k}, \vec{p}, q_3)$, we find

$$H_B^{-1}(a(p_s), q_3) N_B \mathcal{A}(\vec{p}, \vec{k}, q_3) = H_B^{-1}(a(k_s), q_3) \left\{ (4\pi i \lambda_B q_3) \frac{(p+k)_-}{(p-k)_-} \right\} + h(\vec{k}, \vec{p}, q_3) \quad (\text{D.17})$$

Similarly, integrating (D.16) and introducing another integration constant $\tilde{h}(\vec{k}, \vec{p}, q_3)$, we get

$$H_B(a(k_s), q_3) N_B \mathcal{A}(\vec{p}, \vec{k}, q_3) = H_B(a(p_s), q_3) \left\{ (4\pi i \lambda_B q_3) \frac{(p+k)_-}{(p-k)_-} \right\} + \tilde{h}(k_-, \vec{p}, q_3) \quad (\text{D.18})$$

Comparing (D.17) and (D.18), we see that the k_+ dependence of h and p_+ dependence of \tilde{h} are determined. So, we conclude

$$N_B \mathcal{A}(\vec{p}, \vec{k}, q_3) = \frac{H_B(a(p_s), q_3)}{H_B(a(k_s), q_3)} \left\{ (4\pi i \lambda_B q_3) \frac{(p+k)_-}{(p-k)_-} + j(k_-, p_-, q_3) \right\} \quad (\text{D.19})$$

where, j is now the integration constant to be determined shortly. Following the arguments of [17], we see that in the above equation (D.19), the function $j(k_-, p_-, q_3)$ must be a function of charge zero, and so, must be a function of $\frac{k_-}{p_-}$. It must also be a singularity free, i.e., the derivative w.r.t. both k_+ and p_+ must vanish. This seems impossible unless the function j is a constant w.r.t. k and p . And so, we conclude

$$N_B \mathcal{A}(\vec{p}, \vec{k}, q_3) = \frac{H_B(a(p_s), q_3)}{H_B(a(k_s), q_3)} \left\{ (4\pi i \lambda_B q_3) \frac{(p+k)_-}{(p-k)_-} + j(q_3) \right\} \quad (\text{D.20})$$

In order to evaluate $j(q_3)$, we now replace equation (4.9) back into equations (D.9), and get (A.3) we get

$$N_B (\mathcal{A}(\vec{p}, \vec{k}, q_3) - \mathcal{A}_0(\vec{p}, \vec{k}, q_3)) = \frac{1}{(2\pi) H_B(a(k_s), q_3)} \int_0^\infty \frac{r_s dr_s}{a(r_s)} F_B(a(r_s), q_3) H_B(a(r_s), q_3) I(r_s) \quad (\text{D.21})$$

where,

$$I(r_s) = \int_0^{2\pi} \frac{d\theta_r}{2\pi} \left\{ - (4\pi i \lambda_B q_3) \frac{(r+p)_-}{(r-p)_-} + \tilde{b}_4 \right\} \left\{ (4\pi i \lambda_B q_3) \frac{(r+k)_-}{(r-k)_-} + j(q_3) \right\} \quad (D.22)$$

One can easily perform the angular integrals by using (B.1), and the final result of the integration is

$$\begin{aligned} I(r_s) = & [4\pi i \lambda_B q_3 + \tilde{b}_4] [-4\pi i \lambda_B q_3 + j(q_3)] \\ & - \Theta(r_s - p_s) (8\pi i \lambda_B q_3) \left\{ (4\pi i \lambda_B q_3) \frac{p_- + k_-}{p_- - k_-} + j(q_3) \right\} \\ & + \Theta(r_s - k_s) (8\pi i \lambda_B q_3) \left\{ - (4\pi i \lambda_B q_3) \frac{k_- + p_-}{k_- - p_-} + \tilde{b}_4 \right\} \end{aligned} \quad (D.23)$$

From (D.21), we define

$$\text{LHS} = (8\pi i \lambda_B q_3) H_B(a(k_s), q_3) N_B \left(\mathcal{A}(\vec{p}, \vec{k}, q_3) - \mathcal{A}_0(\vec{p}, \vec{k}, q_3) \right) \quad (D.24)$$

Also, we get from (D.21),

$$\begin{aligned} \text{RHS} = & \int \frac{r_s dr_s}{a(r_s)} (4i \lambda_B q_3) F_B(a(r_s), q_3) H_B(a(r_s), q_3) I(r_s) \\ = & [4\pi i \lambda_B q_3 + \tilde{b}_4] [-4\pi i \lambda_B q_3 + j(q_3)] [H_B(\infty, q_3) - H_B(c_B, q_3)] \\ & - (8\pi i \lambda_B q_3) \left\{ (4\pi i \lambda_B q_3) \frac{(p+k)_-}{(p-k)_-} + j(q_3) \right\} [H_B(\infty, q_3) - H_B(a(p_s), q_3)] \\ & + (8\pi i \lambda_B q_3) \left\{ - (4\pi i \lambda_B q_3) \frac{(k+p)_-}{(k-p)_-} + \tilde{b}_4 \right\} [H_B(\infty, q_3) - H_B(a(p_s), q_3)] \end{aligned} \quad (D.25)$$

From (D.24), by using (D.20) and the expression obtained for $N \mathcal{A}_0(\vec{p}, \vec{k}, q_3)$, we can write

$$\begin{aligned} \text{LHS} = & (8\pi i \lambda_B q_3) \left[H_B(a(p_s), q_3) \left\{ 4\pi i \lambda_B q_3 \frac{(p+k)_-}{(p-k)_-} + j(q_3) \right\} \right. \\ & \left. - H_B(a(k_s), q_3) \left\{ - 4\pi i \lambda_B q_3 \frac{(p+k)_-}{(k-p)_-} + \tilde{b}_4 \right\} \right] \end{aligned} \quad (D.26)$$

Comparing LHS and RHS, we get

$$\frac{j(q_3)}{4\pi i \lambda_B q_3} = \frac{4\pi i \lambda_B q_3 (H_B(c_B, q_3) - H_B(\infty, q_3)) + \tilde{b}_4 (H_B(c_B, q_3) + H_B(\infty, q_3))}{4\pi i \lambda_B q_3 (H_B(c_B, q_3) + H_B(\infty, q_3)) + \tilde{b}_4 (H_B(c_B, q_3) - H_B(\infty, q_3))} \quad (D.27)$$

So, the final result for the thermal four-point function of fundamanetal scalars is given by (D.20) with $j(q_3)$ given by (D.27).

The expression for $j(q_3)$ given in (D.27) can be written as an equivalent but simplified form as below

$$j(q_3) = 4\pi \lambda_B q_3 \tan \left(\tan^{-1} \left(\frac{\tilde{b}_4}{4\pi \lambda_B q_3} \right) + 2\lambda_B q_3 \int_{c_B}^{\infty} \frac{dw}{q_3^2 + 4w^2} \chi_B(w) \right). \quad (D.28)$$

It is clear from (D.28) that $j(q_3)$ is an even function of q_3 , i.e., $j(-q_3) = j(q_3) = j(|q_3|)$. It is also clear from (D.28) that $j(q_3)$ or rather $j(q_3, \lambda_B)$ is also an even function of λ_B separately, i.e., $j(q_3, \lambda_B) = j(q_3, -\lambda_B) = j(q_3, |\lambda_B|)$. The large N exact four-point function (D.20) can be explicitly written as

$$N_B \mathcal{A}(\vec{p}, \vec{k}, q_3) = \left\{ (4\pi i \lambda_B q_3) \frac{(p+k)_-}{(p-k)_-} + j(q_3) \right\} \exp \left(4i \lambda_B q_3 \int_{a(k_s)}^{a(p_s)} \frac{dw}{q_3^2 + 4w^2} \chi_B(w) \right) \quad (\text{D.29})$$

where, $j(q_3)$ is given by (D.27) or equivalently by (D.28).

E Comment on arbitrary spin s current two-point function

Fundamental matter coupled to Chern-Simons theories contain a spectrum of higher spin currents [1, 3, 4, 14]. The relevant modifications in the expressions of the currents $J_{\mu_1 \dots \mu_s}$ in the massive matter theory is discussed in detail in [14]. However, these modifications will not be important to what we are going to discuss below. In this subsection, we outline the generalization of the computations correlators of the previous sections to the case of correlator of arbitrary spin s single trace operator which we schematically label as $J_{(s)}$. The schematic diagram for the two-point correlator $\langle J_{(s)} J_{(s)} \rangle$ is shown in Fig.10. In the




Figure 10. two-point correlator $\langle J_{(s)}(-q) J_{(s)}(q) \rangle$. The circled cross at the left is the exact insertion vertex and the boxed cross at the right is the ‘bare’ insertion vertex. The circular dot denotes the exact propagator.

lightcone gauge $A_- = 0$, the explicit computation of the correlators (specifically, solving the corresponding Schwinger-Dyson equations) becomes simple in the kinematical regime in which the external momentum q is chosen to be such that $q_{\pm} = 0$, i.e., in the explicit component form $q \equiv (0, 0, q_3)$. With these choices, we will specifically talk about the component of the correlator $\langle J_{(s)} J_{(s)} \rangle$ in which one of the $J_{(s)}$ insertion vertex has all the minus signs (s times) and the other $J_{(s)}$ insertion vertex which has all the plus signs (s times). As already mentioned in previous cases, to compute the two-point correlators of current operator $J_{(s)}$, only a single exact insertion vertex is required and the other insertion vertex is the ‘bare’ insertion which is computed from the explicit definition of the corresponding current operator. For definiteness, we choose the exact $J_{(s)}$ insertion vertex to be the one in which the spacetime indices are all minuses, and the ‘bare’ insertion vertex $J_{(s)}$ to be the one in which all spacetime indices are pluses. The definition of the gauge invariant, single-trace, higher spin spin current operator is obtained from the corresponding generating functions (see e.g. [1]). Below, we consider fermionic and bosonic theories separately and discuss about the specific details regarding those theories there.

E.1 Fermionic theory

In the fermionic theory coupled to Chern-Simons theory, there is a spectrum of gauge invariant, single trace, higher spin currents. In the lightcone gauge, and in the case of external momenta $q_{\pm} = 0$, the expression for the current operator - with all component signs being minuses - is given by

$$J_{-...-}^F(-q) \equiv J_{(s_m)}^F(-q) = i\alpha_s \int \frac{\mathcal{D}_F^3 k}{(2\pi)^3} (k_-)^{s-1} \bar{\psi}(-(q+k))\gamma_- \psi(k) , \quad (\text{E.1})$$

for $s \geq 1$. Here, we have kept an explicit factor of i for later convenience (comparing it with the case $s = 1$) and the factor α_s is the normalization factor. Here, a schematic symbol s_m denotes that the spin s current operator carry s minus signs.

Schwinger-Dyson equation for exact vertex

The corresponding Schwinger-Dyson equation for the exact $J_{(s_m)}^F$ vertex is schematically given by the Fig.1 where, the ‘tree level’ insertion in this case is given by $\tilde{V}_{(s_m)}^F = i\alpha_s(k_-)^{s-1}\gamma_-$.

The exact $J_{(s_m)}^F$ insertion vertex is defined as

$$\langle J_{(s_m)}^F(-q)\psi(k)\bar{\psi}(p) \rangle = V_{(s_m)}^F(k, q) (2\pi)^3 \delta^{(3)}(p+k+q) . \quad (\text{E.2})$$

From Fig.1, we find that the corresponding Schwinger-Dyson equation for $V_{(s_m)}^F(k, q)$ is given by

$$V_{(s_m)}^F(k, q) = \tilde{V}_{(s_m)}^F(k, q) + N_F \int \frac{\mathcal{D}_F^3 \ell}{(2\pi)^3} [\mathcal{V}^\nu(\ell+q, k+q) S_F(\ell+q) V_{(s_m)}^F(\ell, q) S_F(\ell) \mathcal{V}^\rho(k, \ell)] G_{\rho\nu}^F(\ell-k) , \quad (\text{E.3})$$

Using (3.11) and (3.4) , the above equation (3.43) can be simplified to

$$V_{(s_m)}^F(k, q) = i\alpha_s(k_-)^{s-1}\gamma_- - 2\pi i\lambda_F \int \frac{\mathcal{D}_F^3 \ell}{(2\pi)^3} [\gamma^{[3]} S_F(\ell+q) V_{(s_m)}^F(\ell, q) S_F(\ell) \gamma^{]+}] \frac{1}{(\ell-k)_-} . \quad (\text{E.4})$$

Following exactly the same procedure that is used to compute the exact J_-^F vertex in the case of spin one two-point correlator, one can solve the above Schwinger-Dyson equation (E.4). Here, briefly outline the steps and present the final result. In ‘lightcone kinematics’ $q_{\pm} = 0$, it is clear from (E.4) that $V_{(s_m)}^F(k, q)$ is independent of k_3 . Also, in the lightcone kinematics $q_{\pm} = 0$, the only non-zero component of the external momenta q is q_3 . So, the momentum dependence of $V_{(s_m)}^F(k, q)$ is basically $V_{(s_m)}^F(\vec{k}, q_3)$. As in the J_-^F case, $V_{(s_m)}^F(k, q)$ can be expanded as,

$$V_{(s_m)}^F(\vec{k}, q_3) = V_{(s_m),\nu}^F(\vec{k}, q_3) \gamma^\nu + V_{(s_m),1}^F(\vec{k}, q_3) \mathbb{1} = (k_-)^{s-1} g_m^{(s)}(k_s, q_3) \gamma^+ + (k_-)^s f_m^{(s)}(k_s, q_3) \mathbb{1} . \quad (\text{E.5})$$

Plugging (3.45) in (E.4), we get a set of two coupled integral equations involving $f_m^{(s)}$ and $g_m^{(s)}$. Simiar to the case of J_-^F , these equations for $f_m^{(s)}$ and $g_m^{(s)}$ simplifies a lot once we utilize the $SO(2)$ rotational symmetry in the lightcone plane and decompose the momentum integration measure as (3.21), perform the angular integration using (B.1) and also perform

the integral over momentum component ℓ_3 by using (C.11) and (C.13). Performing a change of variable $a(\ell_s) = +\sqrt{\ell_s^2 + c_F^2} = w$ and $a(k_s) = +\sqrt{k_s^2 + c_F^2} = z$, and relabelling $f_m^{(s)}(k_s, q_3) \equiv \tilde{f}_m^{(s)}(z, q_3)$ and $g_m(k_s, q_3)^{(s)} \equiv \tilde{g}_m^{(s)}(z, q_3)$ and $\tilde{\Sigma}_1(\ell_s) \equiv h_F(w)$, the equations for $\tilde{f}_m^{(s)}$ and $\tilde{g}_m^{(s)}$ can be written in a much simpler form, which takes the following form

$$\tilde{f}_m^{(s)}(z, q_3) = 2i\lambda_F \int_z^\infty dw F_F(w, q_3) \left[(q_3 - 2ih_F(w)) \tilde{f}_m^{(s)}(z, q_3) - 2\tilde{g}_m^{(s)}(w, q_3) \right], \quad (\text{E.6})$$

and

$$\tilde{g}_m^{(s)}(z, q_3) = i\alpha_s + 2i\lambda_F \int_z^\infty dw F_F(w, q_3) \left[2(w^2 - h_F^2(w)) \tilde{f}_m^{(s)}(w, q_3) + (q_3 + 2i\tilde{\Sigma}_1(\ell_s)) \tilde{g}_m^{(s)}(w, q_3) \right]. \quad (\text{E.7})$$

To solve the above two equations (E.6) and (E.7), it is best to convert them to a set of differential equations. The boundary conditions that follow from (E.6) and (E.7) are

$$\tilde{f}_m^{(s)}(z = \Lambda, q_3) = 0, \quad \tilde{g}_m^{(s)}(z = \Lambda, q_3) = i\alpha_s, \quad \text{where, } \Lambda \rightarrow \infty. \quad (\text{E.8})$$

Solving the equations (E.6) and (E.6), by first converting them into differential equations with the boundary conditions (E.8), we find the final solution for the exact $J_{(s_m)}^F$ vertex in terms of $\tilde{f}_m^{(s)}$ and $\tilde{g}_m^{(s)}$ as

$$\begin{aligned} \tilde{f}_m(z, q_3) &= \frac{i\alpha_s}{q_3} \left(1 - \exp \left[4i\lambda_F q_3 \int_z^\Lambda dw F_F(w, q_3) \right] \right), \\ \tilde{g}_m(z, q_3) &= i\alpha_s - \frac{1}{2} (q_3 + 2ih_F(z)) \tilde{f}_m(z, q_3). \end{aligned} \quad (\text{E.9})$$

Two-point function

Schematic diagram for the two-point function $\langle J_{s_m}^F J_{s_p}^F \rangle$ is shown in Fig.10. One needs the ‘bare’ insertion vertex $J_{s_p}^F$ as shown in the right of Fig.10. Let us define the ‘bare’ insertion $J_{(s_p)}^F(q)$ vertex as

$$\langle J_{(s_p)}^F(-q) \psi(k) \bar{\psi}(p) \rangle = U_{(s_p)}^F(k, q) (2\pi)^3 \delta^{(3)}(p + k + q) \quad (\text{E.10})$$

$U_{(s_p)}^F$ can be computed from the explicit expressions of the corresponding currents and the number of digrams included in this insertion should be such that there is no overcounting. The two-point function $\langle J_{s_m}^F J_{s_p}^F \rangle$ is then given by

$$G_{s_m s_p}^F(q) = -N_F \int \frac{\mathcal{D}_F^3 k}{(2\pi)^3} \text{Tr}_F \left[S_F(k + q) V_{(s_m)}^F(k, q) S_F(k) U_{(s_p)}^F(k + q, -q) \right] \quad (\text{E.11})$$

where, the extra (-1) factor above in (3.38) is because of the integration over fermion loop. For various spins, one needs to compute (E.10) and then perform the integral (E.11).

E.2 Bosonic theory

Similar to the fermionic case, we consider the two point function of spin s current operator, in which one of the operator carries all the minus signs and the other operator carries all the plus signs. We define this two-point function of general spin s current $J_{(s)}^B$ in the bosonic theory as

$$\langle J_{(s_m)}^B(q') J_{(s_p)}^B(q) \rangle = G_{s_m s_p}^B(q) (2\pi)^3 \delta^{(3)}(q' + q), \quad (\text{E.12})$$

where, s_m denotes that the operator $J_{(s_m)}^B(q')$ carries s minus signs and similarly s_p denotes that the operator $J_{(s_p)}^B(q)$ carries s plus signs. To compute this, we first compute the exact $J_{(s_m)}^B(q')$ vertex. The other insertion $J_{(s_p)}^B(q)$ is computed upto the leading non-trivial orders to account for all the diagrams that contribute to (E) without overcounting. We define the exact $J_{(s_m)}^B(q')$ insertion vertex by separating the overall momentum conserving δ -function as follows

$$\langle J_{(s_m)}^B(-q) \phi(k) \bar{\phi}(p) \rangle = V_{(s_m)}^B(k, q) (2\pi)^3 \delta^{(3)}(p + k + q) \quad (\text{E.13})$$

Following the same procedure as spin one current, we compute the exact $J_{(s_m)}^B(q')$ insertion vertex by

$$V_{(s_m)}^B(k, q) = V_{(s_m), \text{free}}^B(k, q) + N_B \int \frac{\mathcal{D}_B^3 p}{(2\pi)^3} \frac{V_{(s_m), \text{free}}^B(k, q) \mathcal{A}(p, k, q)}{(k^2 + c_B^2)((k + q)^2 + c_B^2)} \quad (\text{E.14})$$

In the case of lightcone gauge $A_- = 0$ and in the case $q_\pm = 0$, it follows from the rotational symmetry, that $V_{(s_m), \text{free}}^B(k, q) = \tilde{\alpha}_s(k_-)^s$ for $s \geq 1$, where, $\tilde{\alpha}_s$ depends upon the normalization of the current operator. Following the same procedure as in the case of spin one current, we perform the momentum integral and get

$$V_{(s_m)}^B(k, q) = \tilde{\alpha}_s(k_-)^s \frac{H_B(\infty, q_3)}{H_B(a(k_s), q_3)} \quad (\text{E.15})$$

And we also define the ‘tree level’ insertion $J_{(s_p)}^B(q)$ vertex as

$$\langle J_{(s_p)}^B(-q) \phi(k) \bar{\phi}(p) \rangle = U_{(s_p)}^B(k, q) (2\pi)^3 \delta^{(3)}(p + k + q) \quad (\text{E.16})$$

To compute (E.12) we need (E.16) upto leading non-trivial orders to account for all the diagrams that contribute to (E.12) without overcounting. Once this is computed, it is easy to compute the spin s current two-point function $G_{s_m s_p}^B(q)$ as

$$G_{s_m s_p}^B(q) = N_B \int \frac{\mathcal{D}_B^3 k}{(2\pi)^3} \left[S_B(k + q) V_{(s_m)}^B(k, q) S_B(k) U_{(s_p)}^B(k + q, -q) \right] \quad (\text{E.17})$$

where, $S_B(k)$ is the exact scalar field propagator given by (4.5). Explicitly, this can be written as

$$G_{s_m s_p}^B(q) = N_B H_B(\infty, q_3) \int \frac{\mathcal{D}_B^3 k}{(2\pi)^3} \frac{(k_-)^s U_{s_p}^B(k + q, -q)}{(k^2 + c_B^2)((k + q)^2 + c_B^2)} \quad (\text{E.18})$$

As in the case of fermions, for various spins, one needs to compute (E.16) for various spins and then perform the above integral (E.18).

References

- [1] S. Giombi, S. Minwalla, S. Prakash, S. P. Trivedi, S. R. Wadia and X. Yin, *Chern-Simons Theory with Vector Fermion Matter*, *Eur. Phys. J. C* **72** (2012) 2112 [[1110.4386](#)].
- [2] O. Aharony, G. GurAri and R. Yacoby, *$d=3$ Bosonic Vector Models Coupled to Chern-Simons Gauge Theories*, *JHEP* **1203** (2012) 037 [[1110.4382](#)].
- [3] J. Maldacena and A. Zhiboedov, *Constraining Conformal Field Theories with A Higher Spin Symmetry*, *J.Phys. A* **46** (2013) 214011 [[1112.1016](#)].
- [4] J. Maldacena and A. Zhiboedov, *Constraining conformal field theories with a slightly broken higher spin symmetry*, *Class.Quant.Grav.* **30** (2013) 104003 [[1204.3882](#)].
- [5] C.-M. Chang, S. Minwalla, T. Sharma and X. Yin, *ABJ Triality: from Higher Spin Fields to Strings*, *J. Phys. A* **46** (2013) 214009 [[1207.4485](#)].
- [6] S. Jain, S. P. Trivedi, S. R. Wadia and S. Yokoyama, *Supersymmetric Chern-Simons Theories with Vector Matter*, *JHEP* **1210** (2012) 194 [[1207.4750](#)].
- [7] O. Aharony, G. Gur-Ari and R. Yacoby, *Correlation Functions of Large N Chern-Simons-Matter Theories and Bosonization in Three Dimensions*, *JHEP* **1212** (2012) 028 [[1207.4593](#)].
- [8] S. Yokoyama, *Chern-Simons-Fermion Vector Model with Chemical Potential*, *JHEP* **1301** (2013) 052 [[1210.4109](#)].
- [9] G. Gur-Ari and R. Yacoby, *Correlators of Large N Fermionic Chern-Simons Vector Models*, *JHEP* **1302** (2013) 150 [[1211.1866](#)].
- [10] O. Aharony, S. Giombi, G. Gur-Ari, J. Maldacena and R. Yacoby, *The Thermal Free Energy in Large N Chern-Simons-Matter Theories*, *JHEP* **1303** (2013) 121 [[1211.4843](#)].
- [11] S. Jain, S. Minwalla, T. Sharma, T. Takimi, S. R. Wadia et al., *Phases of large N vector Chern-Simons theories on $S^2 \times S^1$* , *JHEP* **1309** (2013) 009 [[1301.6169](#)].
- [12] T. Takimi, *Duality and higher temperature phases of large N Chern-Simons matter theories on $S^2 \times S^1$* , *JHEP* **1307** (2013) 177 [[1304.3725](#)].
- [13] S. Jain, S. Minwalla and S. Yokoyama, *Chern Simons duality with a fundamental boson and fermion*, *JHEP* **1311** (2013) 037 [[1305.7235](#)].
- [14] Y. Frishman and J. Sonnenschein, *Breaking conformal invariance - Large N Chern-Simons theory coupled to massive fundamental fermions*, *JHEP* **12** (2013) 091 [[1306.6465](#)].
- [15] S. Yokoyama, *A Note on Large N Thermal Free Energy in Supersymmetric Chern-Simons Vector Models*, *JHEP* **1401** (2014) 148 [[1310.0902](#)].
- [16] W. A. Bardeen and M. Moshe, *Spontaneous breaking of scale invariance in a $D=3$ $U(N)$ model with Chern-Simons gauge fields*, *JHEP* **1406** (2014) 113 [[1402.4196](#)].
- [17] S. Jain, M. Mandlik, S. Minwalla, T. Takimi, S. R. Wadia and S. Yokoyama, *Unitarity, Crossing Symmetry and Duality of the S -matrix in large N Chern-Simons theories with fundamental matter*, *JHEP* **04** (2015) 129 [[1404.6373](#)].
- [18] W. A. Bardeen, *The Massive Fermion Phase for the $U(N)$ Chern-Simons Gauge Theory in $D=3$ at Large N* , *JHEP* **1410** (2014) 39 [[1404.7477](#)].
- [19] V. Gurucharan and S. Prakash, *Anomalous dimensions in non-supersymmetric bifundamental Chern-Simons theories*, *JHEP* **1409** (2014) 009 [[1404.7849](#)].

- [20] Y. Dandekar, M. Mandlik and S. Minwalla, *Poles in the S-Matrix of Relativistic Chern-Simons Matter theories from Quantum Mechanics*, *JHEP* **04** (2015) 102 [[1407.1322](#)].
- [21] Y. Frishman and J. Sonnenschein, *Large N Chern-Simons with massive fundamental fermions - A model with no bound states*, *JHEP* **1412** (2014) 165 [[1409.6083](#)].
- [22] M. Moshe and J. Zinn-Justin, *3D Field Theories with Chern-Simons Term for Large N in the Weyl Gauge*, *JHEP* **1501** (2015) 054 [[1410.0558](#)].
- [23] O. Aharony, P. Narayan and T. Sharma, *On monopole operators in supersymmetric Chern-Simons-matter theories*, *JHEP* **05** (2015) 117 [[1502.00945](#)].
- [24] K. Inbasekar, S. Jain, S. Mazumdar, S. Minwalla, V. Umesh and S. Yokoyama, *Unitarity, crossing symmetry and duality in the scattering of $\mathcal{N} = 1$ susy matter Chern-Simons theories*, *JHEP* **10** (2015) 176 [[1505.06571](#)].
- [25] A. Bedhotiya and S. Prakash, *A test of bosonization at the level of four-point functions in Chern-Simons vector models*, *JHEP* **12** (2015) 032 [[1506.05412](#)].
- [26] G. Gur-Ari and R. Yacoby, *Three Dimensional Bosonization From Supersymmetry*, *JHEP* **11** (2015) 013 [[1507.04378](#)].
- [27] S. Minwalla and S. Yokoyama, *Chern Simons Bosonization along RG Flows*, *JHEP* **02** (2016) 103 [[1507.04546](#)].
- [28] D. Radicevic, *Disorder Operators in Chern-Simons-Fermion Theories*, *JHEP* **03** (2016) 131 [[1511.01902](#)].
- [29] M. Geracie, M. Goykhman and D. T. Son, *Dense Chern-Simons Matter with Fermions at Large N*, *JHEP* **04** (2016) 103 [[1511.04772](#)].
- [30] O. Aharony, *Baryons, monopoles and dualities in Chern-Simons-matter theories*, *JHEP* **02** (2016) 093 [[1512.00161](#)].
- [31] S. Yokoyama, *Scattering Amplitude and Bosonization Duality in General Chern-Simons Vector Models*, *JHEP* **09** (2016) 105 [[1604.01897](#)].
- [32] G. Gur-Ari, S. A. Hartnoll and R. Mahajan, *Transport in Chern-Simons-Matter Theories*, *JHEP* **07** (2016) 090 [[1605.01122](#)].
- [33] A. Karch and D. Tong, *Particle-Vortex Duality from 3d Bosonization*, *Phys. Rev.* **X6** (2016) 031043 [[1606.01893](#)].
- [34] J. Murugan and H. Nastase, *Particle-vortex duality in topological insulators and superconductors*, *JHEP* **05** (2017) 159 [[1606.01912](#)].
- [35] N. Seiberg, T. Senthil, C. Wang and E. Witten, *A Duality Web in 2+1 Dimensions and Condensed Matter Physics*, *Annals Phys.* **374** (2016) 395 [[1606.01989](#)].
- [36] S. Giombi, *Higher Spin ? CFT Duality*, in *Proceedings, Theoretical Advanced Study Institute in Elementary Particle Physics: New Frontiers in Fields and Strings (TASI 2015): Boulder, CO, USA, June 1-26, 2015*, pp. 137–214, 2017, DOI [[1607.02967](#)].
- [37] P.-S. Hsin and N. Seiberg, *Level/rank Duality and Chern-Simons-Matter Theories*, *JHEP* **09** (2016) 095 [[1607.07457](#)].
- [38] D. Radicevic, D. Tong and C. Turner, *Non-Abelian 3d Bosonization and Quantum Hall States*, *JHEP* **12** (2016) 067 [[1608.04732](#)].

- [39] A. Karch, B. Robinson and D. Tong, *More Abelian Dualities in 2+1 Dimensions*, *JHEP* **01** (2017) 017 [[1609.04012](#)].
- [40] S. Giombi, V. Gurucharan, V. Kirilin, S. Prakash and E. Skvortsov, *On the Higher-Spin Spectrum in Large N Chern-Simons Vector Models*, *JHEP* **01** (2017) 058 [[1610.08472](#)].
- [41] S. R. Wadia, *Chern-Simons theories with fundamental matter: A brief review of large N results including Fermi-Bose duality and the S -matrix*, *Int. J. Mod. Phys. A* **31** (2016) 1630052.
- [42] O. Aharony, F. Benini, P.-S. Hsin and N. Seiberg, *Chern-Simons-matter dualities with SO and USp gauge groups*, *JHEP* **02** (2017) 072 [[1611.07874](#)].
- [43] S. Giombi, V. Kirilin and E. Skvortsov, *Notes on Spinning Operators in Fermionic CFT*, *JHEP* **05** (2017) 041 [[1701.06997](#)].
- [44] F. Benini, P.-S. Hsin and N. Seiberg, *Comments on global symmetries, anomalies, and duality in $(2 + 1)d$* , *JHEP* **04** (2017) 135 [[1702.07035](#)].
- [45] E. Sezgin, E. D. Skvortsov and Y. Zhu, *Chern-Simons Matter Theories and Higher Spin Gravity*, *JHEP* **07** (2017) 133 [[1705.03197](#)].
- [46] T. Nosaka and S. Yokoyama, *Complete factorization in minimal $\mathcal{N} = 4$ Chern-Simons-matter theory*, *JHEP* **01** (2018) 001 [[1706.07234](#)].
- [47] Z. Komargodski and N. Seiberg, *A symmetry breaking scenario for QCD_3* , *JHEP* **01** (2018) 109 [[1706.08755](#)].
- [48] S. Giombi, *Testing the Boson/Fermion Duality on the Three-Sphere*, [1707.06604](#).
- [49] D. Gaiotto, Z. Komargodski and N. Seiberg, *Time-reversal breaking in QCD_4 , walls, and dualities in $2 + 1$ dimensions*, *JHEP* **01** (2018) 110 [[1708.06806](#)].
- [50] K. Jensen and A. Karch, *Bosonizing three-dimensional quiver gauge theories*, *JHEP* **11** (2017) 018 [[1709.01083](#)].
- [51] K. Jensen and A. Karch, *Embedding three-dimensional bosonization dualities into string theory*, *JHEP* **12** (2017) 031 [[1709.07872](#)].
- [52] J. Gomis, Z. Komargodski and N. Seiberg, *Phases Of Adjoint QCD_3 And Dualities*, *SciPost Phys.* **5** (2018) 007 [[1710.03258](#)].
- [53] K. Inbasekar, S. Jain, P. Nayak and V. Umesh, *All tree level scattering amplitudes in Chern-Simons theories with fundamental matter*, *Phys. Rev. Lett.* **121** (2018) 161601 [[1710.04227](#)].
- [54] K. Inbasekar, S. Jain, S. Majumdar, P. Nayak, T. Neogi, T. Sharma et al., *Dual superconformal symmetry of $\mathcal{N} = 2$ Chern-Simons theory with fundamental matter at large N* , *JHEP* **06** (2019) 016 [[1711.02672](#)].
- [55] C. Cordova, P.-S. Hsin and N. Seiberg, *Global Symmetries, Counterterms, and Duality in Chern-Simons Matter Theories with Orthogonal Gauge Groups*, *SciPost Phys.* **4** (2018) 021 [[1711.10008](#)].
- [56] V. G. Charan and S. Prakash, *On the Higher Spin Spectrum of Chern-Simons Theory coupled to Fermions in the Large Flavour Limit*, *JHEP* **02** (2018) 094 [[1711.11300](#)].
- [57] F. Benini, *Three-dimensional dualities with bosons and fermions*, *JHEP* **02** (2018) 068 [[1712.00020](#)].

- [58] K. Aitken, A. Baumgartner, A. Karch and B. Robinson, *3d Abelian Dualities with Boundaries*, *JHEP* **03** (2018) 053 [[1712.02801](#)].
- [59] K. Jensen, *A master bosonization duality*, *JHEP* **01** (2018) 031 [[1712.04933](#)].
- [60] A. Chattopadhyay, P. Dutta and S. Dutta, *From Phase Space to Integrable Representations and Level-Rank Duality*, *JHEP* **05** (2018) 117 [[1801.07901](#)].
- [61] G. J. Turiaci and A. Zhiboedov, *Veneziano Amplitude of Vasiliev Theory*, *JHEP* **10** (2018) 034 [[1802.04390](#)].
- [62] S. Choudhury, A. Dey, I. Halder, S. Jain, L. Janagal, S. Minwalla et al., *Bose-Fermi Chern-Simons Dualities in the Higgsed Phase*, *JHEP* **11** (2018) 177 [[1804.08635](#)].
- [63] A. Karch, D. Tong and C. Turner, *Mirror Symmetry and Bosonization in 2d and 3d*, *JHEP* **07** (2018) 059 [[1805.00941](#)].
- [64] O. Aharony, L. F. Alday, A. Bissi and R. Yacoby, *The Analytic Bootstrap for Large N Chern-Simons Vector Models*, *JHEP* **08** (2018) 166 [[1805.04377](#)].
- [65] R. Yacoby, *Scalar Correlators in Bosonic Chern-Simons Vector Models*, [1805.11627](#).
- [66] K. Aitken, A. Baumgartner and A. Karch, *Novel 3d bosonic dualities from bosonization and holography*, *JHEP* **09** (2018) 003 [[1807.01321](#)].
- [67] O. Aharony, S. Jain and S. Minwalla, *Flows, Fixed Points and Duality in Chern-Simons-matter theories*, *JHEP* **12** (2018) 058 [[1808.03317](#)].
- [68] A. Dey, I. Halder, S. Jain, L. Janagal, S. Minwalla and N. Prabhakar, *Duality and an exact Landau-Ginzburg potential for quasi-bosonic Chern-Simons-Matter theories*, *JHEP* **11** (2018) 020 [[1808.04415](#)].
- [69] E. Skvortsov, *Light-Front Bootstrap for Chern-Simons Matter Theories*, *JHEP* **06** (2019) 058 [[1811.12333](#)].
- [70] A. Chattopadhyay, D. Suvankar and Neetu, *Chern-Simons Theory on Seifert Manifold and Matrix Model*, *Phys. Rev. D* **100** (2019) 126009 [[1902.07538](#)].
- [71] A. Dey, I. Halder, S. Jain, S. Minwalla and N. Prabhakar, *The large N phase diagram of $\mathcal{N} = 2$ $SU(N)$ Chern-Simons theory with one fundamental chiral multiplet*, *JHEP* **11** (2019) 113 [[1904.07286](#)].
- [72] I. Halder and S. Minwalla, *Matter Chern Simons Theories in a Background Magnetic Field*, *JHEP* **11** (2019) 089 [[1904.07885](#)].
- [73] O. Aharony and A. Sharon, *Large N renormalization group flows in 3d $\mathcal{N} = 1$ Chern-Simons-Matter theories*, *JHEP* **07** (2019) 160 [[1905.07146](#)].
- [74] Z. Li, *Bootstrapping Veneziano Amplitude of Vasiliev Theory and 3D Bosonization*, [1906.05834](#).
- [75] S. Jain, V. Malvimat, A. Mehta, S. Prakash and N. Sudhir, *All order exact result for the anomalous dimension of the scalar primary in Chern-Simons vector models*, *Phys. Rev. D* **101** (2020) 126017 [[1906.06342](#)].
- [76] K. Inbasekar, S. Jain, V. Malvimat, A. Mehta, P. Nayak and T. Sharma, *Correlation functions in $\mathcal{N} = 2$ Supersymmetric vector matter Chern-Simons theory*, *JHEP* **04** (2020) 207 [[1907.11722](#)].

- [77] K. Inbasekar, L. Janagal and A. Shukla, *Mass-deformed $N = 3$ supersymmetric Chern-Simons-matter theory*, *Phys. Rev. D* **100** (2019) 085008 [[1908.08119](#)].
- [78] K. Jensen and P. Patil, *Chern-Simons dualities with multiple flavors at large N* , *JHEP* **12** (2019) 043 [[1910.07484](#)].
- [79] R. R. Kalloor, *Four-point functions in large N Chern-Simons fermionic theories*, [1910.14617](#).
- [80] S. Ghosh and S. Mazumdar, *Thermal Correlators and Bosonization Dualities in Large N Chern Simons Matter Theories*, [1912.06589](#).
- [81] K. Inbasekar, L. Janagal and A. Shukla, *Scattering Amplitudes in $\mathcal{N} = 3$ Supersymmetric $SU(N)$ Chern-Simons-Matter Theory at Large N* , *JHEP* **04** (2020) 101 [[2001.02363](#)].
- [82] S. Jain, R. R. John and V. Malvimat, *Momentum space spinning correlators and higher spin equations in three dimensions*, [2005.07212](#).
- [83] S. Minwalla, A. Mishra and N. Prabhakar, *Fermi seas from Bose condensates in Chern-Simons matter theories and a bosonic exclusion principle*, [2008.00024](#).
- [84] S. Jain, R. R. John and V. Malvimat, *Constraining momentum space correlators using slightly broken higher spin symmetry*, [2008.08610](#).
- [85] I. Halder, S. Jain, S. Minwalla, A. Mishra and N. Prabhakar, *The Z-boson in Chern-Simons Matter Theories*, [To appear](#).