

ERRATA

In the original version of this work, we claimed that the problem DS (find an infinite descending sequence through an ill-founded linear order) and the problem BS (find an infinite bad sequence through a non-well quasi-order) are Weihrauch equivalent (Proposition 4.5). Recently, Takayuki Kihara found a critical gap in our proof. In fact, $DS <_W BS$ [1].

The following is a list of the results that are affected:

- Proposition 4.5;
- Corollary 5.4 and Corollary 5.16: these are one-line relativizations of Proposition 4.5;
- Corollary 5.13: the equivalences $\Pi_k^0\text{-DS} \equiv_W \Delta_{k+1}^0\text{-DS} \equiv_W \Sigma_{k+1}^0\text{-DS}$ are unaffected, but the reductions involving $\Pi_k^0\text{-BS}$ and $\Delta_{k+1}^0\text{-BS}$ were obtained using Corollary 5.4 and transitivity.

At the moment of writing, we are not aware whether all the above-mentioned claims admit a counterexample.

All the other results are not using Proposition 4.5, and either they only deal with DS or they are standalone results about BS which are not affected by the above error. In particular, Theorem 5.3, Theorem 5.14, Proposition 5.15, Theorem 5.23, and Corollary 5.24 are correct to the best of our knowledge.

In order to keep the same structure and numbering of the published version, we only applied minimal changes to the following draft. In particular, we rephrased the parts where the equivalence between DS and BS was stated in words and the proofs of the theorems affected by the error have been replaced with a reference to this erratum. We also updated Figure 2 to reflect our current knowledge on the relation between the coded versions of DS.

REFERENCES

- [1] Jun Le Goh, Arno Pauly, and Manlio Valenti, *The weakness of finding descending sequences in ill-founded linear orders*, Available at <https://arxiv.org/abs/2401.11807>, 2024.

FINDING DESCENDING SEQUENCES THROUGH ILL-FOUNDED LINEAR ORDERS

JUN LE GOH, ARNO PAULY, AND MANLIO VALENTI

ABSTRACT. In this work we investigate the Weihrauch degree of the problem DS of finding an infinite descending sequence through a given ill-founded linear order. We show that DS, despite being hard to solve (it has computable inputs with no hyperarithmetic solution), is rather weak in terms of uniform computational strength. To make the latter precise, we introduce the notion of the deterministic part of a Weihrauch degree. We then generalize DS by considering Γ -presented orders, where Γ is a Borel pointclass or Δ_1^1 , Σ_1^1 , Π_1^1 . We study the obtained DS-hierarchy of problems in comparison with the (effective) Baire hierarchy and show that it does not collapse at any finite level.

CONTENTS

1. Introduction	2
1.1. Summary of our results	3
1.2. Structure of the paper	4
2. Background	5
3. The deterministic part of a problem	9
3.1. Impact of the codomain space	9
3.2. The deterministic part and the first-order part	10
3.3. Interaction with other operations on Weihrauch degrees	10
3.4. Previous appearances in the literature	12
4. Finding descending sequences	12
4.1. The uniform strength of DS	12
4.2. Combinatorial principles on linear orders	17
4.3. Relations with Ramsey theorems	18
5. Presentation of orders	20
5.1. Γ_k^0 -DS and Γ_k^0 -BS	22
5.2. Γ_1^1 -DS and Γ_1^1 -BS	26
6. Conclusions	30
References	31

1. INTRODUCTION

We study the difficulty of the following two computational problems:

- Given an ill-founded countable linear order, find an infinite decreasing sequence in it (DS)
- Given a countable quasi-order which is not well, find a bad sequence in it (BS).

2020 *Mathematics Subject Classification*. Primary: 03D30; Secondary: 03D78, 06A75.

Key words and phrases. Weihrauch reducibility, computable analysis, well-quasiorders, reverse mathematics.

The second author is grateful to Mathieu Hoyrup for answering a question on the impact of the codomain on the deterministic part, and to Linda Westrick for several helpful discussions. The third author thanks Alberto Marcone for many valuable suggestions during the preparation of the draft. He also thanks Damir Dzhafarov, Giovanni Soldà and Vittorio Cipriani for useful discussions on the topics of the paper. His research was partially supported by the Italian PRIN 2017 Grant “Mathematical Logic: models, sets, computability”. The authors made significant progress while all attending the BIRS-CMO workshop “Reverse Mathematics of Combinatorial Principles (19w5111)”.

Motivation for the first stems from the treatment of ordinals in reverse mathematics. When working within submodels of second order arithmetic, the notion of well-order depends on the fixed model. This leads to the so-called pseudo-well-orders, i.e. ill-founded linear orders s.t. no descending sequence exists within the model itself. Such a linear order would appear to be well-founded from the point of view of the model. As a classic example of a pseudo-well-order, consider Kleene’s computable linear order with no hyperarithmetic descending sequence ([38, Lem. III.2.1]). Such a linear order is a well-order when seen within the ω -model HYP consisting exactly of the hyperarithmetic sets. Pseudo-well-orders were first studied in [23] and proved to be a powerful tool in reverse mathematics, especially when working at the level of ATR_0 (see [41, Sec. V.4]). Our first task can essentially be rephrased as being concerned with the difficulty of revealing a pseudo-ordinal as not actually being an ordinal.

Our second task can be seen as a abstraction of the computational content of theorems in well-quasi-order (wqo) theory. There are many famous theorems asserting that wqo’s are closed under certain operations. Examples such as Kruskal’s tree theorem, as well as Extended Kruskal’s theorem and Higman’s theorem, have been well-studied in proof theory via their proof-theoretic ordinals (see [39]). However, in their usual form these results lack computational content. Indeed, these theorems state that a certain quasi-order (Q, \preceq_Q) is a wqo. Phrasing a result of this kind in the classical Π_2^1 -form would yield a statement of the type “given an infinite sequence $(q_n)_{n \in \mathbb{N}}$ in Q , find a pair of indexes $i < j$ s.t. $q_i \preceq_Q q_j$ ”. Such a pair (i, j) would be a witness of the fact that the sequence $(q_n)_{n \in \mathbb{N}}$ is not bad. However, while proving that (Q, \preceq_Q) is a wqo can be “hard” (in particular Extended Kruskal’s theorem is not provable in $\Pi_1^1\text{-CA}_0$ [39]), producing a pair of witnesses for each infinite sequence is a \preceq_Q -computable problem (as it can be solved by an extensive search)!

These theorems are very extreme examples of a well-known difference between reverse mathematics and computable analysis: quoting [19] “the computable analyst is allowed to conduct an unbounded search for an object that is guaranteed to exist by (nonconstructive) mathematical knowledge, whereas the reverse mathematician has the burden of an existence proof with limited means”.

On the other hand, considering the contrapositives of the above theorems can reveal some (otherwise hidden) computational content. For example, to show that a given quasi-order is not a wqo it suffices to produce a bad sequence in it. Extended Kruskal’s theorem or Higman’s theorem can be stated in the form “given a bad sequence for the derived quasi-order, find a bad sequence for the original quasi-order”. Our second problem trivially is an upper bound for all these statements, as we disregard any particular reason for why the given quasi-order is not a wqo, and just start with the promise that it is not. Our results thus lay the groundwork for a future exploration of the computational content of individual theorems from wqo theory.

We use the framework of Weihrauch reducibility for our investigation. This means that we compare the problems under investigation to a scaffolding of benchmark problems by asking whether there is an otherwise computable uniform procedure that solves one problem while invoking a single oracle call to the other problem. We are not constrained to particular weak systems in proving that these procedures are actually correct, but rather use whatever proof techniques of ordinary mathematics are suitable. In particular, we can take aspects like the ill-foundedness of the given linear order as external promises not represented in the coding of the input. We can use the fact freely in reasoning about the correctness of our procedure, but there is no evidence provided as input of the procedure.

1.1. Summary of our results. There are a number of problems whose degrees are milestones in the Weihrauch lattice and are often used as benchmarks to calibrate the uniform strength of the multi-valued function under analysis. Some of them roughly correspond to the so-called *big five* subsystems of second order arithmetic: computable problems correspond to RCA_0 , $\text{C}_{2^{\mathbb{N}}}$ (closed choice on the Cantor space) corresponds to WKL_0 , lim (limit in the Baire space) and its iterations correspond to

ACA_0 , $C_{\mathbb{N}^{\mathbb{N}}}$ (closed choice on the Baire space) and its variants $UC_{\mathbb{N}^{\mathbb{N}}}$ and $TC_{\mathbb{N}^{\mathbb{N}}}$ correspond to ATR_0 , $\Pi_1^1\text{-CA}$ corresponds to $\Pi_1^1\text{-CA}_0$.

We show that DS does not belong to this “explored” part of the lattice. To put it in a nutshell, our results show that it is difficult to solve DS, but that DS is rather weak in solving other problems. For example, DS has computable inputs without any hyperarithmetic solutions. On the other hand, DS uniformly computes only the limit computable functions. We provide a few characterizations that tell us what the greatest Weihrauch degree with representatives of particular types below DS is, and include some general observations on this approach. The diagram in Figure 1 shows the relations between DS and several other Weihrauch degrees. Dashed arrows represent Weihrauch reducibility in the direction of the arrow, solid arrows represent strict Weihrauch reducibility. Next,




FIGURE 1. An overview of some parts of the Weihrauch lattice. The solid frame collects the degrees belonging to the lower cone of DS, while the dashed frame collects principles that are not Weihrauch reducible to DS. The only principle shown which is above DS is $C_{\mathbb{N}^{\mathbb{N}}}$. We do not know whether KL is reducible to DS.

we generalize our results by exploring how different presentations of the same order can affect the uniform strength of the same computational task (finding descending sequences in it). We study the problems $\Gamma\text{-DS}$ and $\Gamma\text{-BS}$, where the name of the input order carries “less accessible information” on the order itself (namely $a \leq_L b$ is assumed to be a Γ -condition relative to the name of the order). We summarize the results in Figure 2.

1.2. Structure of the paper. After a short introduction on the preliminary notions on represented spaces and Weihrauch reducibility (Section 2), we define the deterministic part of a multi-valued function and explore the algebraic properties of the operator $\text{Det}_{\mathbf{X}}(\cdot)$ (Section 3). These results will




FIGURE 2. Diagram presenting the relations between the various generalizations of DS.

be very useful in the study of the problems DS and BS (Section 4) and their generalizations Γ -DS and Γ -BS (Section 5).

2. BACKGROUND

For an introduction to Weihrauch reducibility, we point the reader to [10]; for represented spaces to [36]. Below we briefly introduce the notions we will need, as well as state useful results. Those familiar with Weihrauch reducibility should read Definition 2.2 where we define the first-order part of a problem, recently studied by Dzhafarov, Solomon, Yokoyama [17].

A *represented space* \mathbf{X} is a set X together with a (possibly partial) surjection $\delta_{\mathbf{X}} : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow X$. We can transfer notions of computability from $\mathbb{N}^{\mathbb{N}}$ to \mathbf{X} as follows. For each $x \in X$, we say that p is a $(\delta_{\mathbf{X}})$ -name of x if $\delta_{\mathbf{X}}(p) = x$. We say that $x \in X$ is $(\delta_{\mathbf{X}})$ -computable if it has a computable $(\delta_{\mathbf{X}})$ -name.

We list some relevant examples. Let $\mathbf{LO} = (\mathbf{LO}, \delta_{\mathbf{LO}})$ be the represented space of linear orders with domain contained in \mathbb{N} , where each linear order (L, \leq_L) is represented by the characteristic function of the set $\{(a, b) \in \mathbb{N} : a \leq_L b\}$. Let $\mathbf{WO} = (\mathbf{WO}, \delta_{\mathbf{WO}})$ be the represented space of

well-orders with domain contained in \mathbb{N} , where $\delta_{\mathbf{WO}}$ is the restriction of $\delta_{\mathbf{LO}}$ to codes of well-orders. Similarly, let $\mathbf{QO} = (\mathbf{QO}, \delta_{\mathbf{QO}})$ be the represented space of quasi-orders (represented via the characteristic function of the relation). Let also \mathbf{Tr} be the space of subtrees of $\mathbb{N}^{<\mathbb{N}}$, represented by characteristic functions. For every string $\sigma \in \mathbb{N}^{<\mathbb{N}}$ we denote with $\sigma[n]$ the prefix of length n of σ .

We will formalize the problems under investigation as partial multi-valued functions between represented spaces $f : \subseteq \mathbf{X} \rightrightarrows \mathbf{Y}$. For each $x \in X$, $f(x)$ denotes the set of possible outputs corresponding to the input x . The *domain* $\text{dom}(f)$ is the set of all $x \in X$ such that $f(x)$ is nonempty. We often refer to each $x \in \text{dom}(f)$ as an *f-instance* and each $y \in f(x)$ as an *f-solution to x*. When we define a problem, we will often not specify its domain explicitly, in which case its domain should be taken to be as large as possible. The *codomain* of $f : \subseteq \mathbf{X} \rightrightarrows \mathbf{Y}$ is \mathbf{Y} . If $f : \subseteq \mathbf{X} \rightrightarrows \mathbf{Y}$ is such that $f(x)$ is a singleton for each $x \in \text{dom}(f)$, then we say that f is *single-valued*. We indicate that by writing $f : \subseteq \mathbf{X} \rightarrow \mathbf{Y}$. In this case we will write $f(x) = y$ instead of (the formally correct) $f(x) = \{y\}$. An example of a single-valued problem is the identity function $\text{id} : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$.

We can define the computability or continuity of problems via realizers: we say that a function $F : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ is a *realizer* of a problem $f : \subseteq \mathbf{X} \rightrightarrows \mathbf{Y}$ if whenever p is a name for some $x \in \text{dom}(f)$, $F(p)$ is a name for some $y \in f(x)$. A problem is *computable* (respectively *continuous*) if it has a computable (respectively continuous) realizer.

In order to measure the relative uniform computational strength of problems, we use Weihrauch reducibility. A problem f is *Weihrauch reducible* to a problem g , written $f \leq_{\mathbf{W}} g$, if there are computable maps $\Phi, \Psi : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that if p is a name for some $x \in \text{dom}(f)$, then

- (1) $\Phi(p)$ is a name for some $y \in \text{dom}(g)$;
- (2) if q is a name for some element of $g(y)$, then $\Psi(p, q)$ is a name for some element of $f(x)$.

This means that there is a procedure for solving f which is computable except for a single invocation to an oracle for g . Equivalently, there is a computable procedure which transforms realizers for g into realizers for f . A problem f is *strongly Weihrauch reducible* to a problem g , written $f \leq_{\mathbf{sW}} g$, if there are computable maps Φ and Ψ as above, except that Ψ is not allowed access to p in its computation.

Weihrauch reducibility and strong Weihrauch reducibility are quasi-orders, so they define a degree structure on problems: $f \equiv_{\mathbf{W}} g$ if $f \leq_{\mathbf{W}} g$ and $g \leq_{\mathbf{W}} f$ (likewise for $\leq_{\mathbf{sW}}$). Both the Weihrauch degrees and the strong Weihrauch degrees form lattices (see [10, Thm. 3.9 and Thm. 3.10]). There are several natural operations on problems which also lift to the $\equiv_{\mathbf{W}}$ -degrees and the $\equiv_{\mathbf{sW}}$ -degrees. Below we present the operations that we need in this paper.

The *parallel product* $f \times g$ is defined by $(f \times g)(x, y) = f(x) \times g(y)$. We call f a *cylinder* if $f \equiv_{\mathbf{sW}} f \times \text{id}$. If f is a cylinder, then $g \leq_{\mathbf{W}} f$ if and only if $g \leq_{\mathbf{sW}} f$ ([5, Cor. 3.6]). This is useful for establishing nonreductions because if f is a cylinder, then it suffices to diagonalize against all strong Weihrauch reductions from g to f in order to show that $g \not\leq_{\mathbf{W}} f$. Cylinders will also be useful when working with compositional products (discussed below). Observe that for every problem f , $f \times \text{id}$ is a cylinder which is Weihrauch equivalent to f .

The *parallelization* \hat{f} is defined by $\hat{f}((x_n)_{n \in \mathbb{N}}) = \prod_{n \in \mathbb{N}} f(x_n)$. In other words, given a countable sequence of f -instances, \hat{f} asks for an f -solution for each given f -instance.

The *composition* of $f : \subseteq \mathbf{Y} \rightrightarrows \mathbf{Z}$ and $g : \subseteq \mathbf{X} \rightrightarrows \mathbf{Y}$ is defined by $\text{dom}(f \circ g) = \{x \in \text{dom}(g) : g(x) \subseteq \text{dom}(f)\}$ and $(f \circ g)(x) = \bigcup_{y \in g(x)} f(y)$ for $x \in \text{dom}(f \circ g)$. The composition does *not* respect $\leq_{\mathbf{W}}$ or $\leq_{\mathbf{sW}}$. Instead, for any problems f and g (regardless of domain and codomain), we can consider the *compositional product* $f * g$, which satisfies the following property:

$$f * g \equiv_{\mathbf{W}} \max_{\leq_{\mathbf{W}}} \{f_1 \circ g_1 : f_1 \leq_{\mathbf{W}} f \wedge g_1 \leq_{\mathbf{W}} g\}.$$

This captures what can be achieved by first applying g , possibly followed by some computation, and then applying f . The compositional product was first introduced in [9], and proven to be

well-defined in [12]. A useful tool is the *cylindrical decomposition lemma* ([12, Lem. 3.10]): for all problems f and g , if $F \equiv_W f$ and $G \equiv_W g$ are both cylinders, then there is some computable map Φ such that $f * g \equiv_W F \circ \Phi \circ G$. For each problem f , let $f^{[n]}$ denote the n -fold iteration of the compositional product of f with itself, i.e., $f^{[1]} = f$, $f^{[2]} = f * f$, and so on.

The *jump* of $f : \subseteq \mathbf{X} \rightrightarrows \mathbf{Y}$ is the problem $f' : \subseteq \mathbf{X}' \rightrightarrows \mathbf{Y}$ defined by $f'(x) := f(x)$, where \mathbf{X}' is the represented space $(X, \delta_{\mathbf{X}'})$ and $\delta_{\mathbf{X}'}$ takes in input a convergent sequence $(p_n)_{n \in \mathbb{N}}$ in $\mathbb{N}^{\mathbb{N}}$ and returns $\delta_{\mathbf{X}}(\lim_{n \rightarrow \infty} p_n)$. In other words, f' is the following task: given a sequence which converges to a name of an f -instance, produce an f -solution to that instance. The jump respects \leq_{sW} but does *not* lift to the \equiv_W -degrees. We use $f^{(n)}$ to denote the n -th jump of a problem. If we define $\text{lim} : \subseteq (\mathbb{N}^{\mathbb{N}})^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ by $\text{lim}((p_n)_{n \in \mathbb{N}}) := \lim_{n \rightarrow \infty} p_n$ then it is straightforward from the definition that $f^{(n)} \leq_W f * \text{lim}^{[n]}$. The converse reduction does not hold in general. However if f is a cylinder, then for each n , $f^{(n)}$ is a cylinder and $f^{(n)} \equiv_W f * \text{lim}^{[n]}$ (see [10, Prop. 6.14]). In particular, since lim is a cylinder, $\text{lim}^{(n)} \equiv_W \text{lim}^{[n+1]}$ for each n . We say that a problem is *arithmetic* if it is Weihrauch reducible to $\text{lim}^{(n)}$ for some n .

Next we introduce some problems which are milestones in the Weihrauch lattice. Apart from lim and its jumps, most prominent is the family of problems of (*closed*) *choice* $C_{\mathbf{X}}$ defined below. For a represented space \mathbf{X} , let $\mathcal{A}(X)$ denote the space of closed subsets of \mathbf{X} . These are given by the ability to recognize membership in the complement. We define $C_{\mathbf{X}} : \subseteq \mathcal{A}(X) \rightrightarrows X$ by $C_{\mathbf{X}}(A) := A$. In other words, $C_{\mathbf{X}}$ is the task of producing an element of \mathbf{X} given a way to recognize wrong answers. Define *unique choice* $UC_{\mathbf{X}}$ to be the restriction of $C_{\mathbf{X}}$ to closed sets which are singletons. By definition, $UC_{\mathbf{X}}$ is single-valued.

Of particular interest to us are $C_{\mathbb{N}}$, $C_{\mathbb{N}^{\mathbb{N}}}$ and $UC_{\mathbb{N}^{\mathbb{N}}}$. We can view elements of $\mathcal{A}(\mathbb{N})$ to be given as an enumeration of its complement. Thus, $C_{\mathbb{N}}$ is the task of finding a natural number not occurring in a given list. Given a name for a closed set $A \subseteq \mathbb{N}^{\mathbb{N}}$, we can compute a tree $T \subseteq \mathbb{N}^{<\mathbb{N}}$ such that the set $[T]$ of (infinite) paths on T is A . Conversely, given a tree $T \subseteq \mathbb{N}^{<\mathbb{N}}$, we can compute a name for the closed set $[T]$. Therefore we can view $C_{\mathbb{N}^{\mathbb{N}}}$ as the problem of computing a path on a given ill-founded subtree of $\mathbb{N}^{<\mathbb{N}}$. Similarly, we can view $UC_{\mathbb{N}^{\mathbb{N}}}$ as the problem of computing the unique path on a given subtree of $\mathbb{N}^{<\mathbb{N}}$. Both $C_{\mathbb{N}^{\mathbb{N}}}$ and $UC_{\mathbb{N}^{\mathbb{N}}}$ are closed under compositional product [4, Thm. 7.3]. We have:

Theorem 2.1 ([26, Cor. 3.4]). *If $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbf{X}$ is Weihrauch reducible to $UC_{\mathbb{N}^{\mathbb{N}}}$, then for every $x \in \text{dom}(f)$, $f(x)$ contains some y hyperarithmetical relative to x .*

Another prominent problem is $LPO : \mathbb{N}^{\mathbb{N}} \rightarrow \{0, 1\}$, defined by $LPO(p) := 0$ if $p = 0^{\mathbb{N}}$ and $LPO(p) := 1$ otherwise. Its jump LPO' (and its iterated jumps $LPO^{(k)}$) will play an important technical role. We notice that $\text{lim}^{(n)} \equiv_W \widehat{LPO}^{(n)}$ (see e.g. [10, Thm. 6.7 and Prop. 6.10]).

Next we define the represented space $\Gamma(\mathbf{X})$ of all Γ -definable subsets of a computable metric space X , where $\Gamma \in \{\Sigma_k^0, \Pi_k^0, \Delta_k^0, \Sigma_1^1, \Pi_1^1, \Delta_1^1\}$. This is based on the well-known concept of Borel codes [33]. For a more detailed development in the context of computable analysis we refer to [22]. A more abstract and general treatment is provided in [37]. A $\delta_{\Sigma_1^0}$ -name for a set $B \subseteq X$ is a sequence of indices of rational open balls whose union is B . A $\delta_{\Pi_1^0}$ -name for a set is a $\delta_{\Sigma_k^0}$ -name for its complement. (Note that $\delta_{\Pi_1^0}$ agrees with how we represented closed sets previously.) A $\delta_{\Delta_k^0}$ -name for a set is a pair of $\delta_{\Sigma_k^0}$ -names, one for the set itself and one for its complement. A $\delta_{\Sigma_{k+1}^0}$ -name for a set $B \subseteq X$ is a sequence of names for Π_k^0 sets whose union is B .

A $\delta_{\Sigma_1^1}$ -name for a set $S \subseteq X$ is a $\delta_{\Pi_1^0}$ -name for a set $P \subseteq \mathbb{N}^{\mathbb{N}} \times X$ such that $S = \{x \in X : (\exists g)((g, x) \in P)\}$. We define $\delta_{\Pi_1^1}$ and $\delta_{\Delta_1^1}$ similarly to $\delta_{\Pi_k^0}$ and $\delta_{\Delta_k^0}$. If X is \mathbb{N} , we can think of a $\delta_{\Sigma_1^1}$ -name for $S \subseteq \mathbb{N}$ as a sequence $(T_n)_{n \in \mathbb{N}}$ of subtrees of $\mathbb{N}^{<\mathbb{N}}$ such that $n \in S$ if and only if T_n is ill-founded.

In practice, we rarely construct δ_{Γ} -names explicitly. If we want to construct a δ_{Γ} -name for a set $A \subseteq X$, we typically only check that there is a Γ -formula which defines A . By invoking computable closure properties, one can construct a computable map which takes a Γ -formula ϕ and its parameter

p to a δ_{Γ} -name for the set defined by ϕ . Conversely, one can construct a computable map which takes a δ_{Γ} -name p for a set A to a Γ -formula ϕ with parameter p which defines A .

We define the (single-valued) functions $\Gamma\text{-CA} : \subseteq \Gamma(\mathbb{N}) \rightarrow 2^{\mathbb{N}}$ corresponding to comprehension principles: given a δ_{Γ} -name p for a subset A of \mathbb{N} , produce its characteristic function. Notice that, for each k and each $A \in \Sigma_{k+1}^0$, we can use $\text{LPO}^{(k)}$ to check whether $n \in A$ (intuitively, for every p we can use $\text{LPO}^{(k)}$ to answer a $\Sigma_{k+1}^{0,p}$ question). This shows that, for each k ,

$$\text{lim}^{(k)} \equiv_{\text{W}} \widehat{\text{LPO}^{(k)}} \equiv_{\text{W}} \Sigma_{k+1}^0\text{-CA},$$

somewhat implicitly written in [3]. The problem $\Pi_1^1\text{-CA}$ can be seen as the analogue of $\Pi_1^1\text{-CA}_0$. It is Weihrauch equivalent to the parallelization of $\chi_{\Pi_1^1}$, which is the characteristic function of a Π_1^1 -complete set. It is convenient to think of $\chi_{\Pi_1^1}$ as the function that takes in input a subtree of $\mathbb{N}^{<\mathbb{N}}$ and checks whether it is well-founded.

We can also define Γ -choice $\Gamma\text{-C}_{\mathbf{X}} : \subseteq \Gamma(\mathbf{X}) \rightrightarrows \mathbf{X}$ by $\Gamma\text{-C}_{\mathbf{X}}(A) := A$. In other words, $\Gamma\text{-C}_{\mathbf{X}}$ is the task of producing an element of a nonempty Γ -set A , given a δ_{Γ} -name of A . We can define $\Gamma\text{-UC}_{\mathbf{X}} : \subseteq \Gamma(\mathbf{X}) \rightarrow \mathbf{X}$ analogously. When reducing problems to $\text{C}_{\mathbb{N}^{\mathbb{N}}}$ and $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$, the following facts are helpful: $\Sigma_1^1\text{-C}_{\mathbb{N}^{\mathbb{N}}} \equiv_{\text{W}} \text{C}_{\mathbb{N}^{\mathbb{N}}}$ and $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \equiv_{\text{W}} \Sigma_1^1\text{-UC}_{\mathbb{N}^{\mathbb{N}}}$ (see [26]).

We define the represented spaces $\Gamma(\mathbf{LO})$ and $\Gamma(\mathbf{QO})$ by restricting the codomain of δ_{Γ} to the set of subsets of \mathbb{N} which are characteristic functions of linear orders and quasiorders respectively. This will be used in Section 5.

We will often construct linear orders using the following method. For every tree $T \subseteq \mathbb{N}^{<\mathbb{N}}$, we define the *Kleene-Brouwer order* $\text{KB}(T)$ on T as follows: $\sigma \leq_{\text{KB}} \tau$ if and only if $\tau \sqsubseteq \sigma$ or $\sigma \leq_{\text{lex}} \tau$. The map $T \mapsto \text{KB}(T)$ from \mathbf{Tr} to \mathbf{LO} is computable. It is known that $\text{KB}(T)$ is a well-order if and only if T is well-founded (see e.g. [41, Lem. V.1.3]).

Finally we present a notion recently studied by Dzhafarov, Solomon, Yokoyama [17]:

Definition 2.2. Let \mathcal{F} be the set of *first-order* problems, i.e. the set of problems with codomain \mathbb{N} . For every problem $f : \subseteq \mathbf{Y} \rightrightarrows \mathbf{Z}$, the *first-order part* of f is the multi-valued function ${}^1f : \subseteq \mathbb{N}^{\mathbb{N}} \times \mathbf{Y} \rightrightarrows \mathbb{N}$ defined as follows:

- instances are pairs (p, y) s.t. $y \in \text{dom}(f)$ and for every $z \in f(y)$ and every name t for z , $\Phi_p(t)(0) \downarrow$, where $\Phi_{(\cdot)}$ is a fixed universal Turing functional;
- a solution for (p, y) is any n s.t. there is a name t for a solution $z \in f(y)$ s.t. $\Phi_p(t)(0) \downarrow = n$.

The motivation for this notion comes from the following fact:

Proposition 2.3 ([17]). *For every problem f ,*

$${}^1f \equiv_{\text{W}} \max_{\leq_{\text{W}}} \{g \in \mathcal{F} : g \leq_{\text{W}} f\}.$$

We conclude this section with the following proposition:

Proposition 2.4. ${}^1\text{C}_{\mathbb{N}^{\mathbb{N}}} \equiv_{\text{W}} \Sigma_1^1\text{-C}_{\mathbb{N}}$.

Proof. It is known that $\Sigma_1^1\text{-C}_{\mathbb{N}} <_{\text{W}} \widehat{\Sigma_1^1\text{-C}_{\mathbb{N}}} <_{\text{W}} \text{C}_{\mathbb{N}^{\mathbb{N}}}$ ([1, Thm. 3.34]). On the other hand, if $f : \subseteq \mathbf{X} \rightrightarrows \mathbb{N}$ is s.t. $f \leq_{\text{W}} \text{C}_{\mathbb{N}^{\mathbb{N}}}$ via Φ, Ψ then, for every name p of some $x \in \text{dom}(f)$, $\Phi(p)$ is the name of an ill-founded tree T_p and, for every $t \in [T_p]$ we have $\Psi(t)(0) \in f(x)$. This means that we can compute a solution choosing an element from

$$\{n \in \mathbb{N} : (\exists t \in \mathbb{N}^{\mathbb{N}})(t \in [T_p] \wedge \Psi(t)(0) = n)\},$$

which is a $\Sigma_1^1{}^p$ subset of \mathbb{N} . □

We will characterize the first-order part of DS in Theorem 4.10.

3. THE DETERMINISTIC PART OF A PROBLEM

Definition 3.1. Let \mathbf{X} be a represented space and $f : \subseteq \mathbf{Y} \rightrightarrows \mathbf{Z}$ be a multi-valued function. We define $\text{Det}_{\mathbf{X}}(f) : \subseteq \mathbb{N}^{\mathbb{N}} \times \mathbf{Y} \rightarrow \mathbf{X}$ by

$$\text{Det}_{\mathbf{X}}(f)(p, y) = x : \iff (\forall z \in \delta_{\mathbf{Z}}^{-1}(f(y))) (\delta_{\mathbf{X}}(\Phi_p(z)) = x),$$

where $\Phi_{(\cdot)}$ is a universal Turing functional. The domain of $\text{Det}_{\mathbf{X}}(f)$ is maximal for this to be well-defined. We just write $\text{Det}(f)$ for $\text{Det}_{\mathbb{N}^{\mathbb{N}}}(f)$.

Notice that $\text{Det}(f)$ is always a cylinder. This is not true for all \mathbf{X} (if $\mathbf{X} = \mathbb{N}$ then $\text{Det}_{\mathbf{X}}(f)$ always has computable solutions, and therefore $\text{id} \not\leq_{sW} \text{Det}_{\mathbf{X}}(f)$).

Our interest in the principle $\text{Det}_{\mathbf{X}}(f)$ lies in the fact that it has the maximal Weihrauch degree of all (single-valued!) functions with codomain \mathbf{X} that are Weihrauch below f :

Theorem 3.2. $\text{Det}_{\mathbf{X}}(f) \equiv_W \max_{\leq_W} \{g : \subseteq \mathbf{W} \rightarrow \mathbf{X} : g \leq_W f\}$.

Proof. Clearly, $\text{Det}_{\mathbf{X}}(f)$ is itself present in the set on the right hand side. Assume $g : \subseteq \mathbf{W} \rightarrow \mathbf{X}$ satisfies $g \leq_W f$ with reduction witnesses Φ and Ψ . Given a name q for an input to g , let $y = \delta_{\mathbf{Y}}(\Phi(q))$ be the value f is called on, and let p be a name for the function $\Psi(q, \cdot)$. Then $\text{Det}_{\mathbf{X}}(f)(p, y) = g(\delta_{\mathbf{W}}(q))$. \square

In the same spirit, we can identify several other operators $\Lambda_{\mathcal{Y}}$ of the type $\Lambda_{\mathcal{Y}}(f) := \max_{\leq_W} \{g \in \mathcal{Y} : g \leq_W f\}$. In particular the proof strategy used in Theorem 3.2 can be used to prove that $\Lambda_{\mathcal{U}_N}$ and $\Lambda_{\mathcal{V}_N}$ are total, where \mathcal{U}_N is the set of first-order problems with codomain N , and \mathcal{V}_N is the set of problems in \mathcal{U}_N which are also single-valued. This will come into play in Theorem 4.31 and in Theorem 4.33.

Corollary 3.3. $\text{Det}_{\mathbf{X}}(\cdot)$ is an interior degree-theoretic operator on Weihrauch degrees, i.e.

$$\begin{aligned} \text{Det}_{\mathbf{X}}(\text{Det}_{\mathbf{X}}(f)) &\equiv_W \text{Det}_{\mathbf{X}}(f) \leq_W f; \\ f \leq_W g &\Rightarrow \text{Det}_{\mathbf{X}}(f) \leq_W \text{Det}_{\mathbf{X}}(g). \end{aligned}$$

3.1. Impact of the codomain space. We make some basic observations on how the space \mathbf{X} impacts the degrees $\text{Det}_{\mathbf{X}}(f)$ for arbitrary f . Clearly, whenever \mathbf{Y} computably embeds into \mathbf{X} (i.e. there is a computable injection $\mathbf{Y} \rightarrow \mathbf{X}$ with computable inverse), then $\text{Det}_{\mathbf{Y}}(f) \leq_W \text{Det}_{\mathbf{X}}(f)$. In general, we obtain many different operations. To see this, we consider the point degree spectrum of a represented space as introduced by Kihara and P. [28]. The point degree spectrum of $(X, \delta_{\mathbf{X}})$ is the set of Medvedev degrees of the form $\delta_{\mathbf{X}}^{-1}(x)$ for $x \in X$.

The spectrum of \mathbf{Y} is included in that of \mathbf{X} iff \mathbf{Y} can be decomposed into countably many parts each of which embeds into \mathbf{X} ([28, Lem. 3.6]). If the spectrum of \mathbf{Y} is not included in that of \mathbf{X} , we can consider a constant function y witnessing this. Then $\text{Det}_{\mathbf{X}}(y) <_W \text{Det}_{\mathbf{Y}}(y) \equiv_W y$. We have thus seen that if $\text{Det}_{\mathbf{X}}(f) \equiv_W \text{Det}_{\mathbf{Y}}(f)$ for all f , then \mathbf{X} and \mathbf{Y} must have the same point degree spectrum. Miller [32] has shown that the spectrum of $[0, 1]^{\omega}$ is not contained in the Turing degrees (i.e. the spectrum of $2^{\mathbb{N}}$), which was extended in [28] to the result that the spectrum of a computable Polish space is contained in the Turing degrees relative to some oracle iff that space is countably dimensional. The spectra of further spaces have been explored in [27].

We can extend the separation arguments based on the spectrum by considering sequences rather than just constant functions¹. Whenever we have a sequence $f_0 : \mathbb{N} \rightarrow \mathbf{X}_0$ and a function $g_0 : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbf{X}_1$ with $f_0 \equiv_W g_0$, then there is a sequence $h : \mathbb{N} \rightarrow \mathbf{X}_1$ with $f_0 \equiv_W h$. A Weihrauch reduction $f \leq_W g$ for $f : \mathbb{N} \rightarrow \mathbf{X}$ and $g : \mathbb{N} \rightarrow \mathbf{Y}$ gives rise to a computable partial function $F : \subseteq \mathbf{Y}^{\mathbb{N}} \rightarrow \mathbf{X}^{\mathbb{N}}$ with $F(g) = f$. It follows that it suffices to separate $\mathbf{Y}^{\mathbb{N}}$ and $\mathbf{X}^{\mathbb{N}}$ via their spectrum to conclude that $\text{Det}_{\mathbf{X}}(\cdot)$ and $\text{Det}_{\mathbf{Y}}(\cdot)$ are distinct operators. In particular, Miller's result implies that there is a function with codomain \mathbb{R} that is not equivalent to any function with codomain $\mathbb{N}^{\mathbb{N}}$.

¹The ideas in this paragraph were pointed out to us by Mathieu Hoyrup.

3.2. The deterministic part and the first-order part. Let us now explore the interplay between the deterministic part and the first-order part.

Proposition 3.4. ${}^1\text{Det}(f) \equiv_{\text{W}} \text{Det}_{\mathbb{N}}(f) \leq_{\text{W}} \text{Det}({}^1f)$.

Proof. By considering what the relevant maxima in the characterizations are taken about, it is clear that $\text{Det}_{\mathbb{N}}(f) \leq_{\text{W}} {}^1\text{Det}(f)$ and $\text{Det}_{\mathbb{N}}(f) \leq_{\text{W}} \text{Det}({}^1f)$. To see that ${}^1\text{Det}(f) \leq_{\text{W}} \text{Det}_{\mathbb{N}}(f)$, we consider a function $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ and a multi-valued function $g : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}$ with $g \leq_{\text{W}} f$. But this reduction actually yields some choice function of g , showing that $g \leq_{\text{W}} \text{Det}_{\mathbb{N}}(f)$. \square

Open Question 3.5. Is there some f with $\text{Det}_{\mathbb{N}}(f) <_{\text{W}} \text{Det}({}^1f)$?

The question above asks whether whenever there is a countable cover making a partial function on Baire space piecewise computable, there also is a partition of the same or lower complexity that renders the function piecewise computable. The complexity here is not merely the complexity of the individual pieces, but the Weihrauch degree of the map that assigns the piece to any Baire space element.

Proposition 3.6. $\text{Det}(f) \leq_{\text{W}} \widehat{\text{Det}_{\mathbb{N}}(f)}$.

Proof. A function $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ is reducible to the parallelization of its uncurried form $F : \subseteq \mathbb{N} \times \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}$ where $F(n, p) = f(p)(n)$. \square

Corollary 3.7. $\text{Det}(f) \leq_{\text{W}} \widehat{{}^1f}$.

3.3. Interaction with other operations on Weihrauch degrees. A first straightforward observation is that $\text{Det}(f) \square \text{Det}(g) \leq_{\text{W}} \text{Det}(f \square g)$ whenever \square is a degree-theoretic operator that preserves single-valuedness. We will look at the interaction with the usual well-studied operations on Weihrauch degrees. Besides those introduced in Section 2, we consider \sqcup and \sqcap , the join and meet in the Weihrauch lattice, and the finite parallelization $*$ (which essentially is closure under \times). See [10, Section 3] for definitions. The diamond operator \diamond was introduced in [34], and corresponds to the possibility of using the oracle an arbitrary (but finite) number of times (essentially closure under compositional product).

It is imminent from the definition that $\text{Det}(f) \sqcup \text{Det}(g) \equiv_{\text{W}} \text{Det}(f \sqcup g)$.

Moreover $\text{Det}(f \sqcap g) \leq_{\text{W}} \text{Det}(f)$ and $\text{Det}(f \sqcap g) \leq_{\text{W}} \text{Det}(g)$ by monotonicity, hence $\text{Det}(f \sqcap g) \leq_{\text{W}} \text{Det}(f) \sqcap \text{Det}(g)$, as \sqcap is the meet on Weihrauch degrees ([5, Prop. 3.11]). To see that the inequality can be strict, let $p, q \in 2^{\mathbb{N}}$ be a minimal pair of Turing degrees (which we identify with the constant functions returning these values). Then $\text{Det}(p \sqcap q) \equiv_{\text{W}} \text{id}_{\mathbb{N}^{\mathbb{N}}} <_{\text{W}} \text{Det}(p) \sqcap \text{Det}(q) \equiv_{\text{W}} p \sqcap q$.

Our principle DS (to be defined) already witnesses that the deterministic part does not distribute over \times and $*$, and does not commute with $*$, \diamond and $\widehat{}$: we will prove that $\text{Det}(\text{DS}) \equiv_{\text{W}} \text{lim}$ (Theorem 4.16), while $\text{LPO}' \leq_{\text{W}} \text{DS} \times \text{DS}$ (Theorem 4.18). Here we also give another example with a more computability-theoretic flavour:

Example 3.8. There is a Weihrauch degree f such that:

$$\text{Det}(f) \equiv_{\text{W}} \text{id} <_{\text{W}} f <_{\text{W}} f \times f \equiv_{\text{W}} f^{\diamond} \equiv_{\text{W}} \widehat{f} \equiv_{\text{W}} \text{Det}(f \times f).$$

Indeed, consider the degrees of points in the spaces $\mathbb{R}_{<}$, $\mathbb{R}_{>}$ and \mathbb{R} (see [27] for details). Let $x \in \mathbb{R}$ be neither left-c.e. nor right-c.e.; i.e. it lacks computable names in both $\mathbb{R}_{<}$ and $\mathbb{R}_{>}$. Then $x \in \mathbb{R}_{<}$ and $x \in \mathbb{R}_{>}$ have quasi-minimal degrees, that is do not compute any non-computable elements of Cantor space. We define $f: \mathbf{2} \rightarrow \mathbb{R}_{<} + \mathbb{R}_{>}$ by $f(0) = x \in \mathbb{R}_{<}$ and $f(1) = x \in \mathbb{R}_{>}$. The quasi-minimality implies that $\text{Det}(f) \equiv_{\text{W}} \text{id}$. However, $f \times f$ is equivalent to the constant function returning $x \in \mathbb{R}$, which is also equivalent to the constant function returning the decimal expansion of x . Thus, $f \times f \equiv_{\text{W}} \text{Det}(f \times f)$. Any of f^* , $f * f$, f^{\diamond} and \widehat{f} clearly share the same degree.

Theorem 3.9. For every represented space \mathbf{X} and every problems f, g ,

$$\text{Det}_{\mathbf{X}}(f * g) \leq_{\text{W}} \text{Det}_{\mathbf{X}}(f) * g.$$

Proof. Fix a single-valued h with codomain \mathbf{X} and assume, without loss of generality, that $\text{dom}(h) \subset \mathbb{N}^{\mathbb{N}}$ (if h is single-valued then the map $p \mapsto h \circ \delta(p)$ is single-valued as well, where δ is the representation map for the domain of h). Assume also, for the sake of readability, that f and g are cylinders (if not we can just replace f with $f \times \text{id}$, as $\text{Det}_{\mathbf{X}}(\cdot)$ is a degree-theoretic operation).

By the cylindrical decomposition lemma, there is a computable function Φ_e s.t.

$$h \leq_{sW} f \circ \Phi_e \circ g.$$

Let Φ, Ψ be two maps witnessing this strong reduction. Define ϕ as the restriction of $\delta_{\mathbf{X}} \circ \Psi \circ f \circ \Phi_e$ to $\text{dom}(g \circ \Phi \circ h)$. The choice of the domain of ϕ guarantees that ϕ is single-valued: intuitively ϕ witnesses the “second part” of the reduction $h \leq_{sW} f \circ \Phi_e \circ g$, and the fact that h is single-valued implies that so is ϕ . In particular, $\phi \leq_W \text{Det}_{\mathbf{X}}(f)$ (as $\phi \leq_W f$ trivially). Since $h \leq_W \phi * g$ we have that $h \leq_W \text{Det}_{\mathbf{X}}(f) * g$. \square

Notice that this implies the choice elimination theorem [10, Thm. 7.25], as $\text{Det}(C_{2^{\mathbb{N}}}) \equiv_W \text{id}$ ([5, Cor. 8.8]).

Corollary 3.10. *If g is single-valued with codomain $\mathbb{N}^{\mathbb{N}}$ then $\text{Det}(f * g) \equiv_W \text{Det}(f) * g$.*

Proof. This follows from Theorem 3.9, as $\text{Det}(f) * \text{Det}(g) \leq_W \text{Det}(f * g)$ always holds and $\text{Det}(g) \equiv_W g$ as g is single-valued with codomain $\mathbb{N}^{\mathbb{N}}$. \square

Corollary 3.11. *For every cylinder f and every $k \in \mathbb{N}$*

$$\text{Det}(f)^{(k)} \equiv_W \text{Det}(f^{(k)}).$$

Proof. The left-to-right reduction is straightforward as

$$\text{Det}(f)^{(k)} \leq_W \text{Det}(f) * \lim^{[k]} \leq_W f * \lim^{[k]} \equiv_W f^{(k)},$$

where the last equality follows from the fact that f is a cylinder. Since $\text{Det}(f)^{(k)}$ is single-valued, this implies $\text{Det}(f)^{(k)} \leq_W \text{Det}(f^{(k)})$.

The right-to-left reduction follows from Theorem 3.9 as

$$\text{Det}(f^{(k)}) \equiv_W \text{Det}(f * \lim^{[k]}) \leq_W \text{Det}(f) * \lim^{[k]} \equiv_W \text{Det}(f)^{(k)},$$

where the last equality follows from the fact that $\text{Det}(f)$ is a cylinder. \square

The previous corollary can be generalized in a straightforward way to any represented space \mathbf{X} s.t. $\text{Det}_{\mathbf{X}}(f)$ is a cylinder. Notice that it is false (in general) if f is not a cylinder: take $f = C_2$ and $k = 1$. Since $C'_2 \equiv_W \text{RT}_2^1$ (see e.g. [13, Fact 2.3 and Prop. 3.4]) we have $\text{Det}(C'_2) \leq_W \text{RT}_2^1$, hence in particular $\lim \not\leq_W \text{Det}(C'_2)$. On the other hand $\lim \leq_W \text{Det}(C_2)'$ (as $\text{Det}(C_2)$ is a cylinder).

Definition 3.12. Given some $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ let $?f : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ be defined by $0^\omega \in ?f(0^\omega)$ and $0^n 1 p \in ?f(0^n 1 q)$ iff $p \in f(q)$.

It is easy to see that $?$ defines an operation on Weihrauch degrees, and represents the idea of being able to maybe ask a question to f – but never having to decide to forgo this (which would be the case for $1 \sqcup f$). Many well-studied principles are equivalent to their maybe-variants, this in particular holds for all pointed fractals. We introduce the operation here to be able to express how the deterministic part interacts with the notion of completion $\overline{(\cdot)}$ introduced by Brattka and Gherardi [6, 7].

Proposition 3.13. $\text{Det}(\overline{f}) \equiv_W \text{Det}(?f) \equiv_W ?\text{Det}(f)$.

Proof. To show that $\text{Det}(\overline{f}) \leq_W \text{Det}(?f)$, wlog assume that $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ and consider a function $g : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ with $g \leq_W \overline{f}$ witnessed by Φ, Ψ . Now if for some prefix w the computation of $\Psi(w, \cdot)$ outputs two different things depending on the second part of the input, then in order for g to be a function, we have the guarantee that all extensions of w in the domain of g will be mapped to inputs in the domain of f , i.e. we are actually calling f rather than making use of \overline{f} . On the other

hand, if $\Psi(w, \cdot)$ would output the same thing regardless of the second argument, we can postpone actually calling f (which $?f$ lets us do) and go with that output for the time being. This reasoning establishes that $g \leq_W ?f$.

To see that $\text{Det}(?f) \leq_W \text{Det}(f)$, we just inspect the technical definition of $\text{Det}(\cdot)$.

Finally, for $? \text{Det}(f) \leq_W \text{Det}(\bar{f})$ we observe that $? \text{Det}(f)$ is single-valued with codomain $\mathbb{N}^{\mathbb{N}}$, thus it suffices to show $? \text{Det}(f) \leq_W \bar{f}$. But already $?f \leq_W \bar{f}$ holds: \bar{f} accepts an input that is completely void of information. We provide this as long as our $?f$ instance does not want to use f ; if it ever does, we have the relevant f -instance which we can then feed into \bar{f} . Note that we do not get a strong reduction here, in general. \square

3.4. Previous appearances in the literature. While the deterministic part as such has not been introduced before, and in particular the observation that it is always well-defined is new, there are several results in the literature on Weihrauch degrees that implicitly use it. Already in the first paper introducing the modern definition of Weihrauch reducibility [19], it was shown that $\text{Det}(\mathbb{C}_{2^{\mathbb{N}}}) \equiv_W \text{id}$. It was observed in [29] that the argument actually even establishes that $\text{Det}_{\mathbf{X}}(\mathbb{C}_{2^{\mathbb{N}}}) \equiv_W \text{id}$ for any computably admissible space \mathbf{X} .

In [26] the principle $\text{wList}_{2^{\mathbb{N}}, \leq \omega}$ which produces an enumeration of the elements of a countable closed subset of Cantor space was introduced, and [26, Prop. 6.14] states that $\text{Det}(\text{wList}_{2^{\mathbb{N}}, \leq \omega}) \equiv_W \text{lim}$. The authors also proved the following result, which will be useful in Proposition 5.18:

Theorem 3.14 ([26, Thm. 8.5]). $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \equiv_W \text{Det}(\mathbb{C}_{\mathbb{N}^{\mathbb{N}}}) \equiv_W \text{Det}(\widehat{\mathbb{T}\mathbb{C}_{\mathbb{N}^{\mathbb{N}}})$.

This, in particular, shows that $\text{Det}(\cdot)$ is not useful to separate principles that are between $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$ and $\mathbb{C}_{\mathbb{N}^{\mathbb{N}}}$.

In the context of probabilistic computation [8, 11], the fact that the upper cones of non-trivial enumeration degrees are measure zero is equivalent to the statement that if $f : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ has the property that $f(p)$ has positive measure for every $p \in \text{dom}(f)$ and \mathbf{X} is an effectively countably based space, then there is some g with $\text{Det}_{\mathbf{X}}(f) \leq_W {}^1g$.

4. FINDING DESCENDING SEQUENCES

Let us formally define the problem of finding descending sequences in an ill-founded linear order as a multi-valued function.

Definition 4.1. Let $\text{DS} : \subseteq \mathbf{LO} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ be the multi-valued function defined as

$$\text{DS}(L) := \{x \in \mathbb{N}^{\mathbb{N}} : (\forall i)(x(i+1) <_L x(i))\},$$

with $\text{dom}(\text{DS}) := \mathbf{LO} \setminus \mathbf{WO}$.

4.1. The uniform strength of DS. We can immediately notice the following:

Proposition 4.2. $\text{DS} \leq_W \mathbb{C}_{\mathbb{N}^{\mathbb{N}}}$ but $\text{DS} \not\leq_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$.

Proof. To show that $\text{DS} \leq_W \mathbb{C}_{\mathbb{N}^{\mathbb{N}}}$ it is enough to notice that being a descending sequence in a linear order L is a $\Pi_1^{0,L}$ property. In other words, we can obtain a descending sequence through L by choosing a path through the tree

$$\{\sigma \in \mathbb{N}^{<\mathbb{N}} : (\forall i < |\sigma| - 1)(\sigma(i+1) <_L \sigma(i))\}.$$

To show that $\text{DS} \not\leq_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$, recall that there is a computable linear order with no hyperarithmetic descending sequence (see e.g. [38, Lem. III.2.1]). A reduction $\text{DS} \leq_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ would therefore contradict Theorem 2.1. \square

In particular, this shows that DS is not an arithmetic problem (i.e. $\text{DS} \not\leq_W \text{lim}^{(n)}$, for any n).

Proposition 4.3. $\mathbb{C}_{\mathbb{N}^{\mathbb{N}}} \equiv_W \text{lim} * \text{DS}$.

Proof. The reduction $\text{lim} * \text{DS} \leq_{\text{W}} \text{C}_{\mathbb{N}^{\mathbb{N}}}$ follows from the fact that both lim and DS are reducible to $\text{C}_{\mathbb{N}^{\mathbb{N}}}$ and that $\text{C}_{\mathbb{N}^{\mathbb{N}}}$ is closed under compositional product.

To prove the left-to-right reduction notice that, given a tree T , we can computably build the linear order $\text{KB}(T)$. It is known that $[T] \neq \emptyset$ iff $\text{KB}(T)$ is ill-founded (see e.g. [41, Lem. V.1.3]). Moreover, given a infinite descending sequence $(\sigma_n)_{n \in \mathbb{N}}$ in $\text{KB}(T)$, the sequence $(\sigma_n \hat{\ } 0^\omega)_{n \in \mathbb{N}}$ converges to some $x \in [T]$, and therefore the claim follows. \square

We can generalize the problem DS to the context of quasi-orders. It is easy to see that the problem of finding descending sequences in a quasi-order is Weihrauch equivalent to $\text{C}_{\mathbb{N}^{\mathbb{N}}}$. Indeed, on the one hand, being a descending sequence in a quasi-order P is a $\Pi_1^{0,P}$ property. On the other hand, every tree, ordered by the prefix relation, is a partial order where the descending sequences provide arbitrarily long prefixes of a path.

When working with non-well quasi-orders, it is more natural to ask for bad sequences instead.

Definition 4.4. We define the multi-valued function $\text{BS} : \subseteq \mathbf{QO} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ as

$$\text{BS}(P) := \{x \in \mathbb{N}^{\mathbb{N}} : (\forall i)(\forall j > i)(x(i) \not\leq_P x(j))\},$$

where $\text{dom}(\text{BS})$ is the set of quasi-orders that are not well-quasi-orders.

It follows from the definition that every ill-founded linear order is a non-well quasi-order and that every bad sequence through an ill-founded linear order is indeed a descending sequence.

By expanding a bit on a classical argument we can prove that the two problems are uniformly equivalent.

Proposition 4.5. $\text{DS} \equiv_{\text{W}} \text{BS}$.

Proof. See Errata. \square

We will show that DS is quite weak in terms of uniform computational strength (a fortiori $\text{C}_{\mathbb{N}^{\mathbb{N}}} \not\leq_{\text{W}} \text{DS}$). Let us first underline the following useful proposition.

Proposition 4.6. DS is a cylinder.

Proof. Let $p \in \mathbb{N}^{\mathbb{N}}$ and L be an ill-founded linear order. Define

$$\begin{aligned} M &:= \{(p[n], n) : n \in L\}, \\ (p[n], n) \leq_M (p[m], m) &:\iff n \leq_L m. \end{aligned}$$

It is easy to see that M is computably isomorphic to L , and hence it is a valid input for DS . In particular, letting $((p[n_i], n_i))_{i \in \mathbb{N}} \in \text{DS}(M)$, we have that $(n_i)_{i \in \mathbb{N}}$ is a descending sequence in L and $p = \bigcup_{i \in \mathbb{N}} p[n_i]$. \square

Definition 4.7. Let $\Gamma\text{-Bound} : \subseteq \Gamma(\mathbb{N}) \rightrightarrows \mathbb{N}$ be the first-order problem that takes as input a finite Γ subset of the natural numbers and returns a bound for it. Formally

$$\begin{aligned} \text{dom}(\Gamma\text{-Bound}) &:= \{A \in \Gamma(\mathbb{N}) : (\forall^\infty n)(A(n) = 0)\}, \\ \Gamma\text{-Bound}(A) &:= \{n \in \mathbb{N} : (\forall m \geq n)(A(m) = 0)\}. \end{aligned}$$

The principle $\mathbf{\Pi}_1^1\text{-Bound}$ has been studied in [1] under the name $\Sigma_1^1\text{-C}_{\mathbb{N}}^{\text{cof}}$: notice indeed that the reduction $\Sigma_1^1\text{-C}_{\mathbb{N}}^{\text{cof}} \leq_{\text{sW}} \mathbf{\Pi}_1^1\text{-Bound}$ is trivial. On the other hand, given a finite $\mathbf{\Pi}_1^1$ subset X of \mathbb{N} we can consider the set

$$Y := \{n \in \mathbb{N} : (\exists m \geq n)(m \in X)\}.$$

Clearly Y is a $\mathbf{\Pi}_1^1$ initial segment of \mathbb{N} , and therefore $\mathbb{N} \setminus Y$ is a valid input for $\Sigma_1^1\text{-C}_{\mathbb{N}}^{\text{cof}}$. Moreover a name for Y can be uniformly computed from a name of X and $\Sigma_1^1\text{-C}_{\mathbb{N}}^{\text{cof}}(\mathbb{N} \setminus Y) \subset \mathbf{\Pi}_1^1\text{-Bound}(X)$. This shows that $\mathbf{\Pi}_1^1\text{-Bound} \leq_{\text{sW}} \Sigma_1^1\text{-C}_{\mathbb{N}}^{\text{cof}}$ and hence the two problems are (strongly) Weihrauch equivalent.

In other words, given an instance X of Π_1^1 -Bound we can, without loss of generality, assume that X is an initial segment of \mathbb{N} .

Proposition 4.8. Π_1^1 -Bound $<_W$ DS.

Proof. Let X be a Π_1^1 initial segment of \mathbb{N} . By considering the Kleene-Brouwer ordering, we can think of a name for X as a sequence $(L_n)_{n \in \mathbb{N}}$ of linear orders s.t. $n \in X$ iff L_n is well-founded.

Define the linear order $L := \bigcup_n \{n\} \times L_n$, ordered lexicographically. Notice that L is ill-founded as X is not all of \mathbb{N} . Moreover, for every $<_L$ -descending sequence $((n_i, a_i))_{i \in \mathbb{N}}$, we have that $n_0 \in \Pi_1^1$ -Bound(X). Indeed, for every $n \in X$ and every $a \in L_n$, the pair (n, a) lies in the well-founded part of L .

The fact that the reduction is strict follows from the fact that every solution to Π_1^1 -Bound is computable, whereas there is a computable input for DS with no hyperarithmetic solution. \square

We now show that ${}^1\text{DS} \equiv_W \Pi_1^1$ -Bound. Let us first prove the following lemma, which will also be useful to prove Theorem 4.16.

Lemma 4.9. *Suppose that f is a problem which is Weihrauch reducible to DS via the computable maps Φ, Ψ . For every f -instance X , let \leq^X be the linear order defined by Φ^X . We can uniformly compute a sequence $(F_s)_{s \in \mathbb{N}}$ of finite $<^X$ -descending sequences s.t. (1) for every s , $\Psi^{X \oplus F_s}$ outputs some $j \in \mathbb{N}$; (2) for cofinitely many s , F_s extends to an infinite $<^X$ -descending sequence.*

Proof. Fix an f -instance X and run Φ^X for s steps. This produces a finite linear order \leq_s^X . Define

$$D_s := \{F \subseteq \leq_s^X : F \text{ is a } <_s^X \text{-descending sequence and } |F| \geq 1 \text{ and } \Psi^{X \oplus F} \text{ outputs some } j \in \mathbb{N} \text{ in } s \text{ steps}\}.$$

Note that D_s is finite and $t < s$ implies $D_t \subset D_s$. If $D_s \neq \emptyset$ we define F_s to be the $<_{\mathbb{N}}$ -least element of D_s such that

$$(\forall F \in D_s) (\min_{<^X}(F) \leq_s^X \min_{<^X}(F_s)).$$

This ensures that if any $F \in D_s$ extends to an infinite $<^X$ -descending sequence, then so does F_s . Observe that $(F_s)_s$ is uniformly computable from X . If $D_s = \emptyset$ we define $F_s := F_t$ where t is the first index greater than s s.t. $D_t \neq \emptyset$. (We will show below that such t exists, so we can computably search for it.)

Notice that for cofinitely many s , $D_s \neq \emptyset$. Indeed, let S be an infinite $<^X$ -descending sequence (there must exist one because $<^X$ is a DS-instance). Since $\Psi^{X \oplus S}$ outputs some f -solution j of X , there is some finite nonempty initial segment F of S and some $t \in \mathbb{N}$ such that $\Psi^{X \oplus F}$ outputs j in t steps. Hence for all sufficiently large s , we have that $F \in D_s$. This shows that the sequence $(F_s)_{s \in \mathbb{N}}$ is well-defined. Moreover, as already observed, for every $t \geq s$, F_t extends to an infinite $<^X$ -descending sequence.

The fact that, for every s , $\Psi^{X \oplus F_s}$ outputs some $j \in \mathbb{N}$ follows from the definition of D_s . \square

In particular, if f has codomain \mathbb{N} the above lemma implies that, for cofinitely many s , $\Psi^{X \oplus F_s}$ outputs some f -solution for X .

Theorem 4.10. ${}^1\text{DS} \equiv_W \Pi_1^1$ -Bound.

Proof. If $f \leq_W \Pi_1^1$ -Bound, then $f \leq_W$ DS by Proposition 4.8. Since Π_1^1 -Bound is first order, $f \leq_W {}^1\text{DS}$.

To prove the converse reduction, suppose that $f \leq_W$ DS as witnessed by the maps Φ and Ψ . Given an f -instance X , let $(F_s)_{s \in \mathbb{N}}$ be as in Lemma 4.9. Let \leq^X denote the linear order represented by Φ^X . Define the following Π_1^1, X set:

$$A := \{s \in \mathbb{N} : F_s \notin \text{Ext}\},$$

where Ext denotes the set of finite sequences that extend to an infinite $<^X$ -descending sequence.

Notice that A is finite as, for cofinitely many s , F_s is extendible. In particular A is a valid instance of $\mathbf{\Pi}_1^1\text{-Bound}$ and, for every $b \in \mathbf{\Pi}_1^1\text{-Bound}(A)$, F_b is extendible to an infinite $<^X$ -descending sequence. By construction, $\Psi^{X \oplus F_b}$ commits to some $j \in \mathbb{N}$. The fact that F_b is extendible guarantees that j is a valid f -solution of X . \square

Corollary 4.11. $\text{DS} <_{\text{W}} \mathbf{C}_{\mathbb{N}^{\mathbb{N}}}$.

Proof. If $\mathbf{C}_{\mathbb{N}^{\mathbb{N}}} \leq_{\text{W}} \text{DS}$ then, by Proposition 2.4, $\mathbf{\Sigma}_1^1\text{-C}_{\mathbb{N}} \leq_{\text{W}} \mathbf{\Pi}_1^1\text{-Bound}$. However, this would imply that $\widehat{\mathbf{\Sigma}_1^1\text{-C}_{\mathbb{N}}} \leq_{\text{W}} \widehat{\mathbf{\Pi}_1^1\text{-Bound}}$, contradicting [1, Cor. 3.23]. \square

Definition 4.12. Let $f : \subseteq \mathbf{X} \rightrightarrows \mathbb{N}$ be a multi-valued function. We say that f is *upwards-closed* if whenever $n \in f(x)$, then $m \in f(x)$ for all $m > n$.

It is straightforward from the definition that $\mathbf{\Pi}_1^1\text{-Bound}$ is upwards-closed.

Lemma 4.13. *If f is upwards-closed then $\text{Det}_{\mathbb{N}}(f) \leq_{\text{W}} \mathbf{C}_{\mathbb{N}}$.*

Proof. Let g be a single-valued function with codomain \mathbb{N} and suppose that $g \leq_{\text{W}} f$ as witnessed by Φ, Ψ . Given a name p for a g -instance x , we use $\mathbf{C}_{\mathbb{N}}$ to guess some n, t such that $\Psi(p, n)$ converges to some k in at most t steps, and such that for no $m > n$ it ever happens that $\Psi(p, m)$ converges to anything but k . Since f is upwards-closed and g is single-valued, such n, t must exist. Moreover, the associated k is equal to $g(x)$. \square

Proposition 4.14. $\text{Det}_{\mathbb{N}}(\mathbf{\Pi}_1^1\text{-Bound}) \equiv_{\text{W}} \text{Det}_{\mathbb{N}}(\mathbf{C}_{\mathbb{N}}) \equiv_{\text{W}} \mathbf{C}_{\mathbb{N}}$, and therefore $\text{Det}_{\mathbb{N}}(\text{DS}) \equiv_{\text{W}} \mathbf{C}_{\mathbb{N}}$.

Proof. Let us first notice that $\mathbf{C}_{\mathbb{N}} \equiv_{\text{W}} \text{UC}_{\mathbb{N}}$ ([4, Prop. 6.2]) and therefore $\text{Det}_{\mathbb{N}}(\mathbf{C}_{\mathbb{N}}) \equiv_{\text{W}} \mathbf{C}_{\mathbb{N}}$. The fact that $\text{Det}_{\mathbb{N}}(\mathbf{\Pi}_1^1\text{-Bound}) \leq_{\text{W}} \mathbf{C}_{\mathbb{N}}$ follows from Lemma 4.13. To prove the converse reduction it is enough to show that $\text{UC}_{\mathbb{N}} \leq_{\text{W}} \widehat{\mathbf{\Pi}_1^1\text{-Bound}}$.

Let $(n_i)_{i \in \mathbb{N}}$ be an enumeration of the complement of $\{x\} \subset \mathbb{N}$. Define

$$\begin{aligned} m(s) &:= \min\{j \in \mathbb{N} : (\forall i < s)(n_i \neq j)\}, \\ A &:= \{s \in \mathbb{N} : (\exists t > s)(m(t) \neq m(s))\}. \end{aligned}$$

Clearly $\lim_{s \rightarrow \infty} m(s) = x$, which implies that A is finite. Since m is computable (relative to $(n_i)_{i \in \mathbb{N}}$), A is a valid input for $\mathbf{\Pi}_1^1\text{-Bound}$. Moreover, for every $b \in \mathbf{\Pi}_1^1\text{-Bound}(A)$ we have $m(b) = x$.

This implies that $\mathbf{C}_{\mathbb{N}} \leq_{\text{W}} \text{Det}_{\mathbb{N}}(\text{DS})$. To conclude the proof we notice that, for every single-valued g with codomain \mathbb{N} we have

$$g \leq_{\text{W}} \text{DS} \Rightarrow g \leq_{\text{W}} \mathbf{\Pi}_1^1\text{-Bound} \Rightarrow g \leq_{\text{W}} \text{Det}_{\mathbb{N}}(\mathbf{\Pi}_1^1\text{-Bound}) \equiv_{\text{W}} \mathbf{C}_{\mathbb{N}}. \quad \square$$

Notice that $\mathbf{\Pi}_1^1\text{-Bound} \not\leq_{\text{W}} \mathbf{C}_{\mathbb{N}}$: indeed $\widehat{\mathbf{C}_{\mathbb{N}}} \equiv_{\text{W}} \text{lim}$, while $\text{UC}_{\mathbb{N}^{\mathbb{N}}} <_{\text{W}} \widehat{\mathbf{\Pi}_1^1\text{-Bound}}$ (see Proposition 5.21). This implies that $\text{Det}_{\mathbb{N}}(\mathbf{\Pi}_1^1\text{-Bound}) <_{\text{W}} \widehat{\mathbf{\Pi}_1^1\text{-Bound}}$. In this regard, we observe the following:

Proposition 4.15. *The Weihrauch degree of $\mathbf{C}_{\mathbb{N}}$ is the highest Weihrauch degree containing both of the following:*

- (1) a representative which is single-valued and has codomain \mathbb{N} ;
- (2) a representative which is upwards-closed.

Proof. To prove that $\mathbf{C}_{\mathbb{N}}$ satisfies point 1, consider $\text{UC}_{\mathbb{N}}$, which is Weihrauch equivalent to $\mathbf{C}_{\mathbb{N}}$ ([4, Prop. 6.2]). To prove that $\mathbf{C}_{\mathbb{N}}$ satisfies point 2, consider the problem $\mathbf{\Sigma}_1^0\text{-Bound}$ that produces a bound for a finite $\mathbf{\Sigma}_1^0$ subset of \mathbb{N} . Clearly $\mathbf{\Sigma}_1^0\text{-Bound}$ is upwards closed. The reduction $\mathbf{\Sigma}_1^0\text{-Bound} \leq_{\text{W}} \mathbf{C}_{\mathbb{N}}$ follows from the fact that, for every $A \in \text{dom}(\mathbf{\Sigma}_1^0\text{-Bound})$, the set

$$\{n \in \mathbb{N} : (\forall m \geq n)(m \notin A)\}$$

is a $\mathbf{\Pi}_1^{0,A}$ subset of $\mathbf{\Sigma}_1^0\text{-Bound}(A)$. To prove the converse reduction, let p be a name for some $B \in \text{dom}(\mathbf{C}_{\mathbb{N}})$. Define $m(s)$ to be the least number not enumerated in p by stage s . Clearly

$\lim_{s \rightarrow \infty} m(s) = \min B$. In particular this implies that there are only finitely many stages s s.t. $m(s) \neq \min B$. Using Σ_1^0 -Bound we can obtain a stage b s.t. $m(b) = \min B$, hence solving $\mathbf{C}_{\mathbb{N}}$.

Finally the maximality of $\mathbf{C}_{\mathbb{N}}$ follows from Lemma 4.13: indeed suppose $f : \mathbf{X} \rightarrow \mathbb{N}$ is Weihrauch equivalent to some g which is upwards-closed. By Lemma 4.13, we have $\text{Det}_{\mathbb{N}}(g) \leq_W \mathbf{C}_{\mathbb{N}}$. By definition of $\text{Det}(\cdot)$, we have $f \leq_W \text{Det}_{\mathbb{N}}(g)$, hence $f \leq_W \mathbf{C}_{\mathbb{N}}$. \square

Let us now characterize the deterministic part of DS.

Theorem 4.16. $\text{Det}(\text{DS}) \equiv_W \text{lim}$.

Proof. Let us first prove that $\text{lim} \leq_W \text{DS}$. Let J be the Turing jump operator, i.e. $J(p)(e) = 1$ iff $\varphi_e^p(e)$ halts. It is known that $J \equiv_{sW} \text{lim}$ (see [10, Thm. 6.7]). By relativizing the construction in [31, Lem. 4.2] we have that, for every p , we can p -computably build a linear order L of type $\omega + \omega^*$ s.t. every descending sequence through L computes $J(p)$. This shows that $\text{lim} \equiv_W J \leq_W \text{DS}$.

To prove that $\text{Det}(\text{DS}) \leq_W \text{lim}$, suppose that $f : \subseteq \mathbf{X} \rightarrow \mathbb{N}^{\mathbb{N}}$ is single-valued and $f \leq_W \text{DS}$ as witnessed by the maps Φ, Ψ . For every n , define f_n by $f_n(X) := f(X)(n)$. The maps Φ and Ψ witness that $f_n \leq_W \text{DS}$ as well (modulo a trivial coding). Given an f -instance X , consider the sequences $(F_{s,n})_{s \in \mathbb{N}}$ obtained by applying Lemma 4.9 to each f_n . Define the sequence $(p_s)_{s \in \mathbb{N}}$ in $\mathbb{N}^{\mathbb{N}}$ as $p_s(n) := \Psi^{X \oplus F_{s,n}}(0)$. Notice that, by Lemma 4.9, for every n , $\Psi^{X \oplus F_{s,n}}$ outputs some number, therefore $p_s(n)$ is well-defined and is uniformly computable from X . Moreover, since f_n is single-valued and, for cofinitely many s , $F_{s,n}$ is extendible, the sequence $(\Psi^{X \oplus F_{s,n}}(0))_{s \in \mathbb{N}}$ is eventually constant and equal to $f_n(X)$. In particular this shows that, letting $p := \lim_{s \rightarrow \infty} p_s$, for each n we have $p(n) = f_n(X)$, i.e. $p = f(X)$. \square

This result shows that, despite the fact that DS can have very complicated solutions, it is rather weak from the uniform point of view. In fact, its lower Weihrauch cone misses many arithmetic problems. In particular we have:

Corollary 4.17. $\text{DS} \not\leq_W \text{LPO}'$.

Proof. Since LPO is single-valued, so is LPO'. Since $\text{LPO}' \not\leq_W \text{lim}$ (see [9, Cor. 12.3 and Thm. 12.7]), it follows from Theorem 4.16 that $\text{LPO}' \not\leq_W \text{DS}$. On the other hand, $\text{DS} \leq_W \text{LPO}'$, as LPO' always has computable solutions. \square

Notice that Theorem 4.16 implies also that $\mathbf{C}_{\mathbb{N}^{\mathbb{N}}} \not\leq_W \mathbf{C}_{2^{\mathbb{N}}} * \text{DS}$. Indeed, on the one hand $\text{Det}(\mathbf{C}_{\mathbb{N}^{\mathbb{N}}}) \equiv_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ (Theorem 3.14), while, on the other hand, by Theorem 3.9 if f is single-valued and $f \leq_W \mathbf{C}_{2^{\mathbb{N}}} * \text{DS}$ then $f \leq_W \text{DS}$ (as $\text{Det}(\mathbf{C}_{2^{\mathbb{N}}}) \equiv_W \text{id}$) and hence $\text{Det}(\mathbf{C}_{2^{\mathbb{N}}} * \text{DS}) \equiv_W \text{Det}(\text{DS}) \equiv_W \text{lim}$.

Using Corollary 4.17 we can prove that DS is not closed under (parallel) product:

Theorem 4.18. $\text{LPO}' \leq_W \text{DS} \times \text{lim}$ and therefore DS is not closed under product.

Proof. Let $(p_n)_{n \in \mathbb{N}}$ be a sequence in $\mathbb{N}^{\mathbb{N}}$ converging to an instance p of LPO. For each s define

$$g(s) = \begin{cases} i + 1 & \text{if } i \leq s \wedge p_s(i) \neq 0 \wedge (\forall j < i)(p_s(j) = 0), \\ 0 & \text{otherwise.} \end{cases}$$

Let us define a linear order L inductively: at stage $s = 0$ we put 0 into L . At stage $s + 1$ we do the following:

- (1) if $g(s) = g(s + 1)$ we put $2(s + 1)$ immediately below $2s$;
- (2) if $g(s) \neq g(s + 1)$ and $g(s + 1) = 0$ we put $2(s + 1)$ at the bottom;
- (3) if $g(s) \neq g(s + 1)$ and $g(s + 1) > 0$ we put $2(s + 1)$ at the top and we put $2s + 1$ immediately above 0.

This construction produces a linear order on a computable subset of \mathbb{N} . It is clear that g and L are uniformly computable in $(p_n)_{n \in \mathbb{N}}$. Notice that if $\text{LPO}(p) = 1$ then there is an s s.t. for every $t \geq s$, $g(t) = g(s)$ (this follows by definition of limit in the Baire space). In particular, L has order type $n + \omega^*$. On the other hand, if $\text{LPO}(p) = 0$ we distinguish three cases: if $g(s)$ is

eventually constantly 0 then L has order type ω^* . If there are infinitely many s s.t. $g(s) > 0$ then g is unbounded (because for each i , $\lim_s p_s(i) = p(i) = 0$ so g eventually stays above i). In particular, if there are infinitely many s and infinitely many t s.t. $g(s) = 0$ and $g(t) > 0$ then L has order type $\omega^* + \zeta$, where $\zeta := \omega^* + \omega$ is the order type of the integers. If instead $g(s) > 0$ for all sufficiently large s , then L has order type $n + \zeta$. In all cases, L is ill-founded.

We consider the input $(L, (p_n)_{n \in \mathbb{N}})$ for $\text{DS} \times \text{lim}$. Given an $<_L$ -descending sequence $(q_n)_{n \in \mathbb{N}}$, we compute a solution for $\text{LPO}'((p_n)_{n \in \mathbb{N}}) = \text{LPO}(p)$ as follows: if q_0 is odd or $g(q_0/2) = 0$ then we return 0, otherwise we return $p(i)$ where i is s.t. $g(q_0/2) = i + 1$.

Notice that if $\text{LPO}(p) = 1$ then the ω^* part of \leq_L is the final segment of the even numbers that starts with the first index $2s$ s.t. for every $t \geq s$, $g(t) = i + 1$ and $p(i) = 1$. In particular every $<_L$ -descending sequence starts with some even q_0 s.t. $g(q_0/2) > 0$. On the other hand, if $\text{LPO}(p) = 0$ then, by definition of LPO , we have that $p = 0^{\mathbb{N}}$. In this case, the above procedure must return 0 so it produces the correct solution. This proves that $\text{LPO}' \leq_W \text{DS} \times \text{lim}$.

The fact that DS is not closed under product follows from the fact that $\text{lim} \leq_W \text{DS}$ (Theorem 4.16) and Corollary 4.17. \square

4.2. Combinatorial principles on linear orders. We introduce the following notation to phrase many combinatorial principles from reverse mathematics as multi-valued functions.

Definition 4.19. Let $\text{FindC}_Y^X : \subseteq \mathbf{LO} \rightrightarrows \mathbf{LO}$ be the partial multi-valued function defined as

$$\text{FindC}_Y^X(L) := \{M \in \mathbf{LO} : M \subset L \text{ and } \text{ordtype}(M) \in Y\},$$

with domain being the set of $L \in \mathbf{LO}$ s.t. $\text{ordtype}(L) \in X$ and there is some $M \subset L$ s.t. $\text{ordtype}(M) \in Y$.

Similarly we define $\text{FindS}^X : \subseteq \mathbf{LO} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ to be the partial multi-valued function that takes as input a countable linear order L s.t. $\text{ordtype}(L) \in X$ and produces a string $\langle b, x_0, x_1, \dots \rangle$ s.t. $b \in \{0, 1\}$ and, for all i , if $b = 0$ then $x_i <_L x_{i+1}$ while if $b = 1$ then $x_{i+1} <_L x_i$.

If X or Y is not specified, we assume that it contains every countable order type.

There is an extensive literature that studies the ‘‘ascending/descending sequence principle’’ (ADS) and the ‘‘chain/antichain principle’’ (CAC) (see e.g. [24, 25]). These principles and, several of their variations, have been studied from the point of view of Weihrauch reducibility in [2].

Notice that, in particular, the problem ADS (given a linear order, produce an infinite ascending sequence or infinite descending sequence) corresponds to FindS . Similarly the problem General-SADS (given a stable — i.e. of order type $\omega + n, n + \omega^*$ or $\omega + \omega^*$ — linear order, produce an infinite ascending or descending sequence), corresponds to FindS^X , where $X = \{\omega + n, n + \omega^*, \omega + \omega^*\}$.

Proposition 4.20. $\text{LPO}' \leq_W \text{FindS}^{\{\omega, n + \omega^*\}}$.

Proof. Let $(p_i)_{i \in \mathbb{N}}$ be a sequence in $\mathbb{N}^{\mathbb{N}}$ converging to an instance p of LPO . For every $s \in \mathbb{N}$ we define (as we did in the proof of Theorem 4.18)

$$g(s) = \begin{cases} i + 1 & \text{if } i \leq s \wedge p_s(i) \neq 0 \wedge (\forall j < i)(p_s(j) = 0), \\ 0 & \text{otherwise.} \end{cases}$$

Let us define a linear order \leq_L on \mathbb{N} inductively: for each stage s we define a linear order on $\{0, \dots, s\}$. At stage $s = 0$ there are no decisions to make. At stage $s + 1$ we do the following:

- (1) if $0 = g(s) = g(s + 1)$ we put $s + 1$ immediately above s ;
- (2) if $0 < g(s) = g(s + 1)$ we put $s + 1$ immediately below s ;
- (3) if $g(s) \neq g(s + 1)$ we put $s + 1$ at the top.

It is clear that g and \leq_L are uniformly computable in $(p_n)_{n \in \mathbb{N}}$. Notice that if $\text{LPO}(p) = 1$ then there is an s s.t. for every $t \geq s$, $g(t) = i + 1$, where i is the smallest integer s.t. $p(i) = 1$ (this follows by definition of limit in the Baire space). In particular, \leq_L has order type $n + \omega^*$. On the other hand, if $\text{LPO}(p) = 0$ then g is either eventually constantly 0 or unbounded. In both cases the linear

order \leq_L has order type ω . In other words (\mathbb{N}, \leq_L) has order type ω iff $\text{LPO}'((p_i)_{i \in \mathbb{N}}) = 0$. Since the output of $\text{FindS}^{\{\omega, n+\omega^*\}}((\mathbb{N}, \leq_L))$ comes with an indication of the order type of the solution, this defines a reduction from LPO' to $\text{FindS}^{\{\omega, n+\omega^*\}}$. \square

Corollary 4.21. $\text{FindS}^{\{\omega, n+\omega^*\}} \mid_W \text{DS}$, and hence $\text{General-SADS} \mid_W \text{DS}$.

Proof. The fact that $\text{FindS}^{\{\omega, n+\omega^*\}} \not\leq_W \text{DS}$ follows from Proposition 4.20 and the fact that $\text{LPO}' \not\leq_W \text{DS}$ (Corollary 4.17). Moreover, since $\text{FindS}^{\{\omega, n+\omega^*\}}$ is a restriction of General-SADS , we have $\text{General-SADS} \not\leq_W \text{DS}$.

To show that the converse reduction cannot hold it is enough to notice that General-SADS is an arithmetic problem, while $\text{DS} \not\leq_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ (Proposition 4.2). \square

In particular this implies that ADS , as well as the stable chain/antichain principle SCAC , and the weakly stable chain/antichain principle WSCAC are Weihrauch incomparable with DS (as they are all arithmetic problems, and General-SADS is reducible to all of them, see [2]).

Proposition 4.22. $\text{FindC}_{\{\omega, n+\omega^*\}} \leq_W \text{DS}$.

Proof. Given a linear order (L, \leq_L) we can computably build the linear order $Q := L + L^*$. Formally we define (Q, \leq_Q) as $Q := \{0\} \times L \cup \{1\} \times L$ and

$$(a, p) \leq_Q (b, q) :\iff a < b \vee (a = b = 0 \wedge p \leq_L q) \vee (a = b = 1 \wedge q \leq_L p).$$

Notice that Q is always ill-founded, hence it is a valid input for DS . Given $(q_i)_{i \in \mathbb{N}} \in \text{DS}(Q)$, we computably build the sequence $(x_i)_{i \in \mathbb{N}}$ defined by $x_i := \pi_1 q_i$ where $\pi_i := (a_0, a_1) \mapsto a_i$.

We distinguish 3 cases:

- (1) if $\pi_0 q_i = 0$ for every i then $(x_i)_{i \in \mathbb{N}}$ is an ω^* -sequence in L ;
- (2) if $\pi_0 q_i = 1$ for every i then $(x_i)_{i \in \mathbb{N}}$ is an ω -sequence in L ;
- (3) if there is a k s.t. for all $i < k$ we have $\pi_0 q_i = 1$ and for all $j \geq k$ we have $\pi_0 q_j = 0$ then, by point 1, $(x_j)_{j \geq k}$ is an ω^* -sequence in L , hence $(x_i)_{i \in \mathbb{N}}$ is of type $n + \omega^*$, with $n \leq k$.

In any case the sequence $(x_i)_{i \in \mathbb{N}}$ is a valid solution for $\text{FindC}_{\{\omega, n+\omega^*\}}$. \square

4.3. Relations with Ramsey theorems. We now explore the relations between DS and Ramsey's theorem for n -tuples and k colors. Let us recall the basic definitions.

Definition 4.23. For every $A \subset \mathbb{N}$, let $[A]^n := \{B \subset A : |B| = n\}$ be the set of subsets of A with cardinality n . A map $c : [\mathbb{N}]^n \rightarrow k$ is called a k -coloring of $[\mathbb{N}]^n$, where $k \geq 2$. An infinite set H s.t. $c([H]^n) = \{i\}$ for some $i < k$ is called a *homogeneous solution* for c , or simply *homogeneous*.

The set $\mathcal{C}_{n,k}$ of k -colorings of $[\mathbb{N}]^n$ can be seen as a represented space, where a name for a coloring c is the string $p \in \mathbb{N}^{\mathbb{N}}$ s.t. for each $(i_0, \dots, i_{n-1}) \in [\mathbb{N}]^n$, $p(\langle i_0, \dots, i_{n-1} \rangle) = c(i_0, \dots, i_{n-1})$.

We define $\text{RT}_k^n : \mathcal{C}_{n,k} \rightrightarrows 2^{\mathbb{N}}$ as the total multi-valued function that maps a coloring c to the set of all homogeneous sets for c . Similarly we define $\text{RT}_{\mathbb{N}}^n : \bigcup_{k \geq 1} \mathcal{C}_{n,k} \rightrightarrows 2^{\mathbb{N}}$ as $\text{RT}_{\mathbb{N}}^n(c) := \text{RT}_k^n(c)$, where $k - 1$ is the maximum of the range of c . Note that the input for $\text{RT}_{\mathbb{N}}^n$ does not include information on which color appears in the range of the coloring.

We also define $\text{cRT}_k^n : \mathcal{C}_{n,k} \rightrightarrows k$ as the multi-valued function that produces only the color of a homogeneous solution. We define $\text{cRT}_{\mathbb{N}}^n$ analogously.

Notice that $\text{cRT}_k^n \equiv_W \text{RT}_k^n$ iff $n = 1$. Indeed the output of cRT_k^n is always computable, while for $n > 1$ there are computable k -colorings with no computable homogeneous solutions. Similarly $\text{cRT}_{\mathbb{N}}^n \equiv_W \text{RT}_{\mathbb{N}}^n$ iff $n = 1$. Moreover the equivalence cannot be lifted to a strong Weihrauch equivalence. Indeed RT_k^1 and cRT_k^1 are incomparable from the point of view of strong Weihrauch reducibility. The uniform computational content of Ramsey's theorems is well-studied (see e.g. [13, 14, 16, 35]).

In comparing RT_k^n with DS , we immediately notice that $\text{RT}_2^2 \not\leq_W \text{DS}$. This follows from the fact that $\text{ADS} \leq_W \text{RT}_2^2$ (see e.g. [24]), while $\text{ADS} \not\leq_W \text{DS}$ (see the remarks after Corollary 4.21). Hence $\text{RT}_k^n \not\leq_W \text{DS}$ for all $n, k \geq 2$.

Proposition 4.24. $\text{RT}_{\mathbb{N}}^1 <_{\text{W}} \Pi_1^1\text{-Bound}$, and hence $\text{RT}_{\mathbb{N}}^1 <_{\text{W}} \text{DS}$.

Proof. Given a coloring $c: \mathbb{N} \rightarrow k$, consider the $\Sigma_2^{0,c}$ set

$$X := \{n \in \mathbb{N} : (\forall \infty j)(c(n) \neq c(j))\}.$$

It is easy to see that X is finite, as $\text{ran}(c) \subset k$ and if there is no c -homogeneous set with color i then there are finitely many $j \in \mathbb{N}$ s.t. $c(j) = i$. In particular, given a bound b for X there is a homogeneous solution with color $c(b)$.

The separation follows from the fact that $\Pi_1^1\text{-Bound} \not\leq_{\text{W}} \widehat{\text{UC}}_{\mathbb{N}^{\mathbb{N}}}$ (as $\widehat{\Pi_1^1\text{-Bound}} \not\leq_{\text{W}} \widehat{\text{UC}}_{\mathbb{N}^{\mathbb{N}}}$, see [1, Fact 3.25]), while $\text{RT}_{\mathbb{N}}^1 <_{\text{W}} \widehat{\text{UC}}_{\mathbb{N}^{\mathbb{N}}}$ (in particular $\text{RT}_{\mathbb{N}}^1 <_{\text{W}} \widehat{\text{C}}_{\mathbb{N}}$, see [13, Prop. 7.2 and Cor. 7.6]). The fact that $\text{RT}_{\mathbb{N}}^1 <_{\text{W}} \text{DS}$ follows from $\Pi_1^1\text{-Bound} <_{\text{W}} \text{DS}$ (Proposition 4.8). \square

We now show that $\text{RT}_{\mathbb{N}}^1$ is the strongest problem among those that are reducible to DS and whose instances always have finitely many solutions.

Definition 4.25. Let $f : \subseteq \mathbf{X} \rightrightarrows \mathbb{N}$. We say that f is *pointwise finite* if, for each $x \in \text{dom}(f)$, $|f(x)|$ is finite.

Notice that cRT_k^1 and $\text{cRT}_{\mathbb{N}}^1$ are pointwise finite, as for each k -coloring c we have $|\text{cRT}_k^1(c)| = |\text{cRT}_{\mathbb{N}}^1(c)| \leq k$.

Lemma 4.26. *Let g be upwards-closed and let f be pointwise finite. If $f \leq_{\text{W}} g$ then $f \leq_{\text{W}} \text{RT}_{\mathbb{N}}^1$.*

Proof. Suppose that $f \leq_{\text{W}} g$ as witnessed by Φ, Ψ . Let p be the name for the f -instance x we are given.

We define a coloring c as follows: we dove-tail all computations $\Psi(p, n)$ for $n \in \mathbb{N}$. Whenever some computation converges to some $j \in \mathbb{N}$, we define $c(i) := j$ where i is the first element on which c is not defined yet. Since g is upwards-closed, we know that for all but finitely many n , $\Psi(p, n)$ has to converge to some $j_n \in f(x)$. This implies that $\text{ran}(c)$ contains only finitely many distinct elements. Moreover, any element repeating infinitely often is a correct solution to $f(x)$, therefore we can find a $y \in f(x)$ by applying $\text{RT}_{\mathbb{N}}^1$ to c and returning the color of the solution. \square

Theorem 4.27. *If f is pointwise finite then $f \leq_{\text{W}} \text{DS}$ iff $f \leq_{\text{W}} \text{RT}_{\mathbb{N}}^1$.*

Proof. The right-to-left implication always holds as $\text{RT}_{\mathbb{N}}^1 <_{\text{W}} \text{DS}$ (Proposition 4.24). On the other hand, if f is pointwise finite and $f \leq_{\text{W}} \text{DS}$ then, by Theorem 4.10 we have $f \leq_{\text{W}} \Pi_1^1\text{-Bound}$. Since $\Pi_1^1\text{-Bound}$ is upwards-closed, by Lemma 4.26 we have $f \leq_{\text{W}} \text{RT}_{\mathbb{N}}^1$. \square

By Lemma 4.26 we also have the following:

Proposition 4.28. *The Weihrauch degree of $\text{RT}_{\mathbb{N}}^1$ is the highest Weihrauch degree such that:*

- (1) *it contains a representative which is pointwise finite;*
- (2) *it is Weihrauch reducible to some problem which is upwards-closed.*

Proof. Point 1 holds because $\text{cRT}_{\mathbb{N}}^1$ is pointwise finite and $\text{cRT}_{\mathbb{N}}^1 \equiv_{\text{W}} \text{RT}_{\mathbb{N}}^1$. Point 2 holds because $\text{RT}_{\mathbb{N}}^1 <_{\text{W}} \Pi_1^1\text{-Bound}$ (Proposition 4.24) and $\Pi_1^1\text{-Bound}$ is upwards-closed. Finally, the maximality follows from Lemma 4.26. \square

Lemma 4.29. *If f is upwards-closed and $f \leq_{\text{W}} \text{RT}_{\mathbb{N}}^1$ then $f \leq_{\text{W}} \text{C}_{\mathbb{N}}$.*

Proof. Recall that $\text{RT}_{\mathbb{N}}^1 \equiv_{\text{W}} \text{cRT}_{\mathbb{N}}^1$ and let Φ, Ψ be two computable maps witnessing $f \leq_{\text{W}} \text{cRT}_{\mathbb{N}}^1$. Let p be a name for some $x \in \text{dom}(f)$ and let c be the coloring represented by $\Phi(p)$. We define the following $\Pi_1^{0,p}$ set

$$\begin{aligned} A := \{ \langle n, c_0, \dots, c_k, s \rangle : & (\forall i)(\exists j \leq k)(c(i) = c_j) \text{ and} \\ & (\forall j \leq k)(\exists i < s)(c(i) = c_j) \text{ and} \\ & (\forall j \leq k)(\Psi(p, c_j) \downarrow \rightarrow \Psi(p, c_j) \leq n) \}. \end{aligned}$$

Notice that, if $\langle n, c_0, \dots, c_k, s \rangle \in A$ then, by the first two conditions, there is a $j \leq k$ s.t. c_j is a valid solution for $\text{cRT}_{\mathbb{N}}^1(c)$. In particular $\Psi(p, c_j) \downarrow$ and is a correct solution for $f(x)$ (as Φ and Ψ witness that $f \leq_{\text{W}} \text{cRT}_{\mathbb{N}}^1$). Since f is upwards-closed, every number greater than $\Psi(p, c_j)$ is a valid solution. In particular, the third condition implies that $n \geq \Psi(p, c_j)$ and therefore $n \in f(x)$. \square

Notice that the previous lemma provides an alternative proof for the fact that $\Pi_1^1\text{-Bound} \not\leq_{\text{W}} \text{RT}_{\mathbb{N}}^1$, as $\Pi_1^1\text{-Bound} \not\leq_{\text{W}} \text{C}_{\mathbb{N}}$.

If we consider only bounded pointwise finite functions, we can improve Theorem 4.27 by replacing $\text{RT}_{\mathbb{N}}^1$ with RT_k^1 .

Lemma 4.30. *If f has codomain k , then $f \leq_{\text{W}} \text{RT}_{\mathbb{N}}^1$ iff $f \leq_{\text{W}} \text{RT}_k^1$.*

Proof. The right-to-left implication is trivial, so let us prove the left-to-right one. Since $\text{RT}_{\mathbb{N}}^1 \leq_{\text{W}} \Pi_1^1\text{-Bound}$ and $\Pi_1^1\text{-Bound}$ is upwards-closed, it suffices to show that if g is upwards-closed and $f \leq_{\text{W}} g$, then $f \leq_{\text{W}} \text{RT}_k^1$. The proof closely follows the one of Lemma 4.26. Suppose that $f \leq_{\text{W}} g$ as witnessed by Φ, Ψ . Let p be the name for the f -instance x we are given. We define a k -coloring c as follows: we dove-tail all computations $\Psi(p, n)$ for $n \in \mathbb{N}$. Whenever some computation converges to some $j < k$, we define $c(i) := j$ where i is the first element on which c is not defined yet. Since g is upwards-closed, we know that for all but finitely many n , $\Psi(p, n)$ has to converge to some $j_n < k$ which lies in $f(x)$. Moreover, any element repeating infinitely often is a correct solution to $f(x)$, therefore we can find a $y \in f(x)$ by applying RT_k^1 to c and returning the color of the solution. \square

Theorem 4.31. *If f has codomain k , then $f \leq_{\text{W}} \text{DS}$ iff $f \leq_{\text{W}} \text{RT}_k^1$.*

Proof. The right-to-left implication always holds as $\text{RT}_k^1 \leq_{\text{W}} \text{RT}_{\mathbb{N}}^1$ trivially and $\text{RT}_{\mathbb{N}}^1 <_{\text{W}} \text{DS}$ (Proposition 4.24). The left-to-right implication follows from Theorem 4.27 and Lemma 4.30. \square

To conclude the section we notice how we can improve the results if we restrict our attention to single-valued functions. Define the problem $\text{lim}_k \subseteq k^{\mathbb{N}} \rightarrow k$ as the limit in the discrete space k .

Lemma 4.32. *If f has codomain k and is single-valued, then $f \leq_{\text{W}} \text{lim}_k$ iff $f \leq_{\text{W}} \text{RT}_k^1$.*

Proof. The left-to-right implication is trivial as $\text{lim}_k \leq_{\text{W}} \text{RT}_k^1$. To prove the converse direction recall that $\text{RT}_k^1 \equiv_{\text{W}} \text{cRT}_k^1$ and let the reduction $f \leq_{\text{W}} \text{cRT}_k^1$ be witnessed by the maps Φ, Ψ . Let p be a name for some $x \in \text{dom}(f)$ and let c be the coloring represented by $\Phi(p)$. Notice that, since f is single-valued, for every solution $j \in \text{cRT}_k^1(c)$ we have $\Psi(p, j) = f(x)$. Furthermore, since the range of c is finite, there are only finitely many i such that $c(i)$ is not a solution. If we then define

$$n_i := \begin{cases} \Psi(p, c(i)) & \text{if } \Psi(p, c(i)) \text{ converges in } i \text{ steps and } \Psi(p, c(i)) < k, \\ 0 & \text{otherwise,} \end{cases}$$

we have that the sequence $(n_i)_{i \in \mathbb{N}} \in k^{\mathbb{N}}$ converges to $f(x)$. Therefore we can use lim_k to obtain $f(x)$. \square

Theorem 4.33. *If f has codomain k and is single-valued, then $f \leq_{\text{W}} \text{lim}_k$ iff $f \leq_{\text{W}} \text{DS}$.*

Proof. The left-to-right implication follows from the fact that $\text{lim} <_{\text{W}} \text{DS}$ (Theorem 4.16), while the other direction follows from Theorem 4.31 and Lemma 4.32. \square

5. PRESENTATION OF ORDERS

In this section we study how the presentation of a linear/quasi order can influence the uniform computational strength of the problems DS and BS.

Definition 5.1. For every $\Gamma \in \{\Sigma_k^0, \Pi_k^0, \Delta_k^0, \Sigma_1^1, \Pi_1^1, \Delta_1^1\}$ we define the problem $\Gamma\text{-DS} \subseteq \Gamma(\mathbf{LO}) \rightrightarrows \mathbb{N}^{\mathbb{N}}$ as $\Gamma\text{-DS}(L) := \text{DS}(L)$. Similarly we define $\Gamma\text{-BS} \subseteq \Gamma(\mathbf{QO}) \rightrightarrows \mathbb{N}^{\mathbb{N}}$ as $\Gamma\text{-BS}(P) := \text{BS}(P)$.

It is not the case that $\Gamma\text{-DS} \equiv_{\mathbb{W}} \Gamma\text{-BS}$ in general. In particular, we will show that $\Sigma_k^0\text{-BS} \not\leq_{\mathbb{W}} \Sigma_k^0\text{-DS}$ (Theorem 5.14) and $\Sigma_1^1\text{-BS} \not\leq_{\mathbb{W}} \Sigma_1^1\text{-DS}$ (Corollary 5.24).

Furthermore, we strengthen Corollary 4.11 by showing that $\Sigma_1^1\text{-DS} <_{\mathbb{W}} \mathbb{C}_{\mathbb{N}^{\mathbb{N}}}$ (Theorem 5.22). In other words, even if we are allowed to feed DS a code for a Σ_1^1 linear ordering, we still cannot compute $\mathbb{C}_{\mathbb{N}^{\mathbb{N}}}$. On the other hand, we already showed that if we are allowed to perform a relatively small amount of post-processing (namely lim) on the output of DS, then we can compute $\mathbb{C}_{\mathbb{N}^{\mathbb{N}}}$ (Proposition 4.3). In particular, the use of lim absorbs any difference in uniform strength between DS and $\Sigma_1^1\text{-DS}$ and collapses the whole hierarchy (up to $\Sigma_1^1\text{-DS}$) to $\mathbb{C}_{\mathbb{N}^{\mathbb{N}}}$.

Many of our separations are derived by analyzing the first-order part of the problems in question, or more generally by characterizing the problems satisfying certain properties (such as single-valuedness or having restricted codomain) which lie below the problems in question. On the contrary, we prove Theorem 5.22 using very different techniques due to Anglès d'Auriac and Kihara [1].

Before beginning our analysis, we record some preliminary observations. Note that $\text{DS} = \Delta_1^0\text{-DS}$ and $\text{BS} = \Delta_1^0\text{-BS}$. It is straightforward to see that, for every Γ , $\Gamma\text{-DS} \leq_{\mathbb{W}} \Gamma\text{-BS}$. Moreover, for every Γ, Γ' s.t. $\Gamma(X) \subset \Gamma'(X)$ we have $\Gamma\text{-DS} \leq_{\mathbb{W}} \Gamma'\text{-DS}$ and $\Gamma\text{-BS} \leq_{\mathbb{W}} \Gamma'\text{-BS}$.

Notice also that the set of bad sequences through a Δ_1^1 -quasi-order is Δ_1^1 , hence it is straightforward to see that $\Delta_1^1\text{-BS} \leq_{\mathbb{W}} \Sigma_1^1\text{-C}_{\mathbb{N}^{\mathbb{N}}} \equiv_{\mathbb{W}} \mathbb{C}_{\mathbb{N}^{\mathbb{N}}}$. This shows also that $\Gamma\text{-BS} \leq_{\mathbb{W}} \mathbb{C}_{\mathbb{N}^{\mathbb{N}}}$ for every arithmetic Γ .

Proposition 5.2. *For every $\Gamma \in \{\Sigma_k^0, \Pi_k^0, \Delta_k^0, \Sigma_1^1, \Pi_1^1, \Delta_1^1\}$ the problems $\Gamma\text{-DS}$ and $\Gamma\text{-BS}$ are cylinders.*

Proof. The proof is a straightforward generalization of the proof of Proposition 4.6. \square

Theorem 5.3. *For every $k \in \mathbb{N}$ and every $\Gamma \in \{\Sigma, \Pi, \Delta\}$*

$$\begin{aligned} \Gamma_{k+1}^0\text{-DS} &\equiv_{\mathbb{W}} \Gamma_1^0\text{-DS} * \text{lim}^{[k]} \equiv_{\mathbb{W}} \Gamma_1^0\text{-DS}^{(k)}, \\ \Gamma_{k+1}^0\text{-BS} &\equiv_{\mathbb{W}} \Gamma_1^0\text{-BS} * \text{lim}^{[k]} \equiv_{\mathbb{W}} \Gamma_1^0\text{-BS}^{(k)}. \end{aligned}$$

Proof. Fix k and Γ as above. The reduction $\Gamma_{k+1}^0\text{-DS} \leq_{\mathbb{W}} \Gamma_1^0\text{-DS} * \text{lim}^{[k]}$ follows from the fact that

$$\text{lim}^{[k]} \equiv_{\mathbb{W}} \Sigma_k^0\text{-CA} \equiv_{\mathbb{W}} \Pi_k^0\text{-CA} \equiv_{\mathbb{W}} \Delta_{k+1}^0\text{-CA},$$

hence we can use $\text{lim}^{[k]}$ to compute a Γ_1^0 -name for the input linear order, and then apply $\Gamma_1^0\text{-DS}$ to get a descending sequence.

Let us now prove the converse reduction. Since both $\text{lim}^{[k]}$ and $\Gamma_1^0\text{-DS}$ are cylinders, by the cylindrical decomposition there is an e s.t.

$$\Gamma_1^0\text{-DS} * \text{lim}^{[k]} \equiv_{\mathbb{W}} \Gamma_1^0\text{-DS} \circ \Phi_e \circ \text{lim}^{[k]}.$$

Given any $p \in \text{dom}(\Gamma_1^0\text{-DS} \circ \Phi_e \circ \text{lim}^{[k]})$, the string $q := \Phi_e(\text{lim}^{[k]}(p))$ is a Γ_1^0 -name for a linear order L_p . Since q is $\Delta_{k+1}^{0,p}$, the condition $a \leq_{L_p} b$ is $\Gamma_{k+1}^{0,p}$ for every a, b . This shows that, given an input p we can uniformly compute a Γ_{k+1}^0 -name for the linear order L_p , and hence use $\Gamma_{k+1}^0\text{-DS}$ to compute an $<_{L_p}$ -descending sequence.

The equivalence $\Gamma_1^0\text{-DS} * \text{lim}^{[k]} \equiv_{\mathbb{W}} \Gamma_1^0\text{-DS}^{(k)}$ follows from the fact that $\Gamma_1^0\text{-DS}$ is a cylinder.

The same reasoning works, mutatis mutandis, to show that

$$\Gamma_{k+1}^0\text{-BS} \equiv_{\mathbb{W}} \Gamma_1^0\text{-BS} * \text{lim}^{[k]} \equiv_{\mathbb{W}} \Gamma_1^0\text{-BS}^{(k)}. \quad \square$$

Using this theorem, the relativized version of Proposition 4.5 can be proved explicitly as follows:

Corollary 5.4. *For every $k \geq 1$, $\Delta_k^0\text{-DS} \equiv_{\mathbb{W}} \Delta_k^0\text{-BS}$.*

Proof. See Errata. \square

5.1. Γ_k^0 -DS and Γ_k^0 -BS. We will now show that the hierarchy of Γ -DS problems does not collapse at any finite level. First we study the hierarchy of Δ_k^0 -DS problems by characterizing their first-order parts (Theorem 5.5). Then we prove the analogues of Theorem 4.31 and Theorem 4.33 for Δ_k^0 -DS (Proposition 5.8).

For any sequence of problems $f_s : \subseteq \mathbf{X}_s \rightrightarrows \mathbf{Y}_s$, $s \in \mathbb{N}$, the *countable coproduct* of the sequence is the problem $\bigsqcup_{s \in \mathbb{N}} f_s : \subseteq \bigsqcup_s \mathbf{X}_i \rightrightarrows \bigsqcup_s \mathbf{Y}_i$ defined by $(\bigsqcup_{s \in \mathbb{N}} f_s)(s, x) := \{s\} \times f_s(x)$. The problem $\bigsqcup_{s \in \mathbb{N}} f_s$ allows us access to exactly one f_s of our choice.

Theorem 5.5. *For every $k \geq 1$,*

$${}^1\Delta_k^0\text{-DS} \equiv_{\text{W}} \left(\bigsqcup_{s \in \mathbb{N}} \Delta_k^0\text{-C}_s \right) * \Pi_1^1\text{-Bound}.$$

We split the proof into two lemmas.

Lemma 5.6. *For every $k \geq 1$, if $f : \subseteq \mathbf{X} \rightrightarrows \mathbb{N}$ and $f \leq_{\text{W}} \Delta_k^0$ -DS then*

$$f \leq_{\text{W}} \left(\bigsqcup_{s \in \mathbb{N}} \Delta_k^0\text{-C}_s \right) * \Pi_1^1\text{-Bound}.$$

Proof. Fix Turing functionals Φ and Ψ which witness that $f \leq_{\text{W}} \Delta_k^0$ -DS. Given an f -instance with name x , Φ^x is a $\Delta_k^{0,x}$ -code for the linear order \leq^x . Consider the $\Sigma_k^{0,x}$ set

$$D := \{F \in \mathbb{N} : F \text{ codes a non-empty finite } <^x\text{-descending sequence and } \Psi^{x \oplus F} \text{ outputs some } j \in \mathbb{N}\}.$$

We can uniformly express D as the increasing union over $s \in \mathbb{N}$ of finite sets $D_s \subseteq \{0, \dots, s\}$, which are uniformly $\Pi_{k-1}^{0,x}$.

We now define the set

$$A := \{s \in \mathbb{N} : (\forall F \in D_s)(F \notin \text{Ext}_x)\},$$

where Ext_x is the set of finite sequences that extend to an infinite $<^x$ -descending sequence. It is easy to see that A is $\Pi_1^{1,x}$, as being extendible in a Δ_k^0 -linear order is a Σ_1^1 property.

We show that A is finite. Since \leq^x is a Δ_k^0 -DS-instance, we can fix an infinite $<^x$ -descending sequence S . By definition of Weihrauch reducibility, $\Psi^{x \oplus S}$ outputs some f -solution $j \in \mathbb{N}$. By the continuity of Ψ , there is some finite non-empty initial segment F of S such that $\Psi^{x \oplus F}$ outputs j . Hence for all sufficiently large s , we have $F \in D_s$.

This shows that we can apply Π_1^1 -Bound to A to obtain some $b \in \mathbb{N}$ which bounds A . Note that D_b must be nonempty. We now define the following non-empty subset of D_b :

$$B := \left\{ F \in D_b : (\forall G \in D_b) \left(\min_{<^x}(G) \leq^x \min_{<^x}(F) \right) \right\}.$$

Notice that all the quantifications are bounded. In particular, B is a (non-empty) $\Delta_k^{0,x}$ subset of D_b because D_b is $\Pi_{k-1}^{0,x}$ and \leq^x is $\Delta_k^{0,x}$. Notice also that the definition of B ensures that each of its elements is extendible (as we know that there is some extendible element in D_b). In particular, this shows that, for every $F \in B$, it is enough to run $\Psi^{x \oplus F}$ to compute an f -solution for the original instance. We can find such $F \in B$ by applying $(\bigsqcup_s \Delta_k^0\text{-C}_s)(b, B)$. \square

Notice that $(\bigsqcup_s \Delta_k^0\text{-C}_s)$ is computable, hence in case $k = 1$ we obtain Proposition 4.5.

Lemma 5.7. *For every $k \geq 1$,*

$$\left(\bigsqcup_{s \in \mathbb{N}} \Delta_k^0\text{-C}_s \right) * \Pi_1^1\text{-Bound} \leq_{\text{W}} \Delta_k^0\text{-DS}.$$

Proof. Using the cylindrical decomposition we can write

$$\left(\bigsqcup_{s \in \mathbb{N}} \Delta_k^0\text{-C}_s \right) * \Pi_1^1\text{-Bound} \equiv_W \left(\left(\bigsqcup_{s \in \mathbb{N}} \Delta_k^0\text{-C}_s \right) \times \text{id} \right) \circ \Phi_e \circ (\Pi_1^1\text{-Bound} \times \text{id})$$

for some computable map Φ_e . Let Φ_1, Φ_2 be computable maps s.t. $\Phi_e(p) = \langle \Phi_1(p), \Phi_2(p) \rangle$. Then we have

$$\begin{aligned} & \left(\left(\bigsqcup_{s \in \mathbb{N}} \Delta_k^0\text{-C}_s \right) \times \text{id} \right) \circ \Phi_e \circ (\Pi_1^1\text{-Bound} \times \text{id})(\langle p_1, p_2 \rangle) = \\ & \left\langle \left(\bigsqcup_{s \in \mathbb{N}} \Delta_k^0\text{-C}_s \right) \Phi_1(\Pi_1^1\text{-Bound}(p_1), p_2), \Phi_2(\Pi_1^1\text{-Bound}(p_1), p_2) \right\rangle. \end{aligned}$$

Given an instance $\langle p_1, p_2 \rangle$ of the above composition, we can think of p_1 as coding an input A to $\Pi_1^1\text{-Bound}$ via a tree T s.t. for each $i, i \in A$ iff the subtree $T_i := \{\sigma \in T : \sigma(0) = i\}$ of T is well-founded. For any $b \in \Pi_1^1\text{-Bound}(p_1)$, $\Phi_1(b, p_2)$ must be a name for an instance of $\bigsqcup_{s \in \mathbb{N}} \Delta_k^0\text{-C}_s$. Then $\pi_1 \Phi_1(b, p_2)$ is a number s and $\pi_2 \Phi_1(b, p_2)$ is a Δ_k^0 -name for a non-empty subset A_s of $\{0, \dots, s-1\}$, where $\pi_i(\langle p_1, p_2 \rangle) = p_i$ denotes the projection on the i -th component. Regardless of whether $b \in \Pi_1^1\text{-Bound}(p_1)$, we will interpret $\pi_1 \Phi_1(b, p_2)$ and $\pi_2 \Phi_1(b, p_2)$ as above.

We define a $\Delta_k^{0, \langle p_1, p_2 \rangle}$ linear order as follows. First define

$$\begin{aligned} L := \{ & (\sigma, n) : \sigma \in p_1 \text{ and} \\ & \pi_1 \Phi_1(\sigma(0), p_2) \text{ outputs a number in less than } |\sigma| \text{ steps and} \\ & n \text{ lies in the set named by } \pi_2 \Phi_1(\sigma(0), p_2)\}. \end{aligned}$$

We order the elements of L by

$$(\sigma, n) \leq_L (\tau, m) :\iff \sigma <_{\text{KB}} \tau \vee (\sigma = \tau \wedge n \leq m).$$

It is easy to see that (L, \leq_L) is $\Delta_k^{0, \langle p_1, p_2 \rangle}$. Notice that it is a linear order, as the pairs are ordered lexicographically where the first components are ordered according to the Kleene-Brouwer order on $\mathbb{N}^{<\mathbb{N}}$ and the second components are ordered according to the order on \mathbb{N} .

Let $(q_i)_{i \in \mathbb{N}}$ be an $<_L$ -descending sequence, with $q_i = (\sigma_i, n_i)$. Notice that for each i there is a $j > i$ s.t. $\sigma_j <_{\text{KB}} \sigma_i$. Indeed, if there is an i s.t. for all $j > i$ we have $\sigma_j = \sigma_i$ then, by definition of \leq_L , the sequence $(n_j)_{j > i}$ would be a descending sequence in the natural numbers, which is impossible.

This implies that there is a subsequence $(q_{i_k})_{k \in \mathbb{N}}$ s.t. $(\sigma_{i_k})_{k \in \mathbb{N}}$ is a $<_{\text{KB}}$ -descending sequence. In particular, this implies that $T_{\sigma_0(0)}$ is ill-founded, i.e. $\sigma_0(0) \in \Pi_1^1\text{-Bound}(p_1)$. Moreover, by definition of L , this implies that n_0 lies in the set named by $\pi_2 \Phi_1(\sigma_0(0), p_2)$.

In other words, given an $<_L$ -descending sequence $(q_i)_{i \in \mathbb{N}}$ we have that $(\pi_1 q_0)(0) \in \Pi_1^1\text{-Bound}(p_1)$ and $\pi_2 q_0 \in (\bigsqcup_{s \in \mathbb{N}} \Delta_k^0\text{-C}_s) \Phi_1(\Pi_1^1\text{-Bound}(p_1), p_2)$. From this we can compute $\Phi_2(\pi_1 q_0, p_2)$ as well. This establishes the desired reduction. \square

This completes the proof of Theorem 5.5.

With a small modification of the argument in the proof of Lemma 5.6 we can prove the following:

Proposition 5.8. *Fix $k \geq 1$. For every $f : \subseteq \mathbf{X} \rightrightarrows \mathbb{N}$,*

$$f \leq_W \Delta_k^0\text{-DS} \iff f \leq_W \Pi_1^1\text{-Bound} \times \lim^{[k-1]}.$$

If, in particular, f has codomain N for some $N \geq 1$ then

$$f \leq_W \Delta_k^0\text{-DS} \iff f \leq_W \text{RT}_N^1 * \lim^{[k-1]}.$$

If, additionally, f is single-valued, then

$$f \leq_W \Delta_k^0\text{-DS} \iff f \leq_W \lim_N * \lim^{[k-1]}.$$

Proof. The right-to-left implication follows from Proposition 4.8 and Theorem 5.3:

$$\begin{aligned} \mathbf{\Pi}_1^1\text{-Bound} \times \lim^{[k-1]} &\leq_W \mathbf{\Pi}_1^1\text{-Bound} * \lim^{[k-1]} \\ &\leq_W \text{DS} * \lim^{[k-1]} \equiv_W \mathbf{\Delta}_k^0\text{-DS}. \end{aligned}$$

To prove the left-to-right implication, fix a pair of Turing functionals Φ and Ψ witnessing the reduction $f \leq_W \mathbf{\Delta}_k^0\text{-DS}$. Fix an f -instance with name x and let \leq^x be the $\Delta_k^{0,x}$ linear order defined by Φ^x .

Define D , D_s and A as in the proof of Lemma 5.6. In that proof, we applied $\mathbf{\Pi}_1^1\text{-Bound}$ to A to obtain $b \in \mathbb{N}$. Then we restricted our attention to $B \subseteq D_b$. Here we will still apply $\mathbf{\Pi}_1^1\text{-Bound}$ to A , but we will concurrently consider a subset B_s of each D_s . For each s , define

$$\begin{aligned} B_s &:= \left\{ F \in D_s : (\forall G \in D_s) \left(\min_{<^x}(G) \leq^x \min_{<^x}(F) \right) \right\}, \\ F_s &:= \min B_s, \end{aligned}$$

where F_s is intended to be the empty sequence if B_s (and hence D_s) is empty.

Notice that $B_s \subset \{0, \dots, s-1\}$ is $\mathbf{\Delta}_k^0$ (as each D_s is $\mathbf{\Pi}_{k-1}^0$) and therefore F_s is $\mathbf{\Delta}_k^0$. Since $\lim^{[k-1]} \equiv_W \mathbf{\Delta}_k^0\text{-CA}$, it can determine which B_s is nonempty, and compute F_s if B_s is nonempty. Therefore the sequence $(F_s)_{s \in \mathbb{N}}$ can be computed using $\lim^{[k-1]}$. For every $b \in \mathbf{\Pi}_1^1\text{-Bound}(A)$ we have that F_b is extendible to an infinite $<^x$ -descending sequence and that $\Psi^{x \oplus F_s}$ converges to some f -solution j (see also the proof of Lemma 5.6).

Assume now that f has codomain N for some $N \geq 1$. We can modify the above argument as follows: after computing the sequence $(F_s)_{s \in \mathbb{N}}$, we consider the RT_N^1 -instance c defined as

$$c(s) := \begin{cases} 0 & \text{if } F_s = \langle \rangle, \\ \Psi^{x \oplus F_s}(0) & \text{otherwise.} \end{cases}$$

Since F_s is nonempty and extendible for cofinitely many s , if $c(s) = i$ for infinitely many s (i.e., c has an RT_N^1 -solution of color i), then there is an extendible F_s s.t. $\Psi^{x \oplus F_s}(0) = i$, hence i is an f -solution.

If, additionally, f is single-valued, then there is only one possible i s.t. c has a homogeneous solution with color i . This shows that the sequence $(c(s))_{s \in \mathbb{N}}$ has a limit, and therefore it suffices to use \lim_N to get the solution.

The fact that $\text{RT}_N^1 * \lim^{[k-1]}$ and $\lim_N * \lim^{[k-1]}$ are reducible to $\mathbf{\Delta}_k^0\text{-DS}$ follows from the fact that the compositional product is a degree theoretic operation, as $\text{RT}_N^1 \leq_W \text{DS}$ (Theorem 4.31), $\lim_N \leq_W \text{DS}$ (Theorem 4.16) and $\mathbf{\Delta}_k^0\text{-DS} \equiv_W \text{DS} * \lim^{[k-1]}$ (Theorem 5.3). \square

Notice that $\mathbf{\Pi}_1^1\text{-Bound} \times \lim^{[k-1]}$ is not a first-order problem, so the first statement in Proposition 5.8 is not an alternative characterization of ${}^1\mathbf{\Delta}_k^0\text{-DS}$. It can be rephrased as

$${}^1\mathbf{\Delta}_k^0\text{-DS} \equiv_W {}^1(\mathbf{\Pi}_1^1\text{-Bound} \times \lim^{[k-1]}).$$

This concludes our discussion of the first-order problems that are Weihrauch reducible to $\mathbf{\Delta}_k^0\text{-DS}$. As for the deterministic part of $\mathbf{\Delta}_k^0\text{-DS}$:

Corollary 5.9. *For every $k \geq 1$, $\text{Det}(\mathbf{\Delta}_k^0\text{-DS}) \equiv_W \lim^{[k]}$.*

Proof. This follows from $\text{Det}(\text{DS}) \equiv_W \lim$ (Theorem 4.16) and the fact that, for cylinders, the jump commutes with the deterministic part (Corollary 3.11). \square

Theorem 5.10. *For every $k \geq 1$,*

$$\mathbf{\Delta}_k^0\text{-DS} <_W \mathbf{\Delta}_{k+1}^0\text{-DS}.$$

In particular this shows that the $\mathbf{\Gamma}$ -DS-hierarchy does not collapse at any finite level.

Proof. This follows directly from Proposition 5.8 or, alternatively, from Corollary 5.9. Indeed it suffices to notice that, for every $k \geq 1$, $\text{LPO}^{(k)} \leq_W \lim^{[k+1]}$ but $\text{LPO}^{(k)} \not\leq_W \lim^{[k]}$, as $\text{LPO}^{(k)}$ is the characteristic function of a Σ_{k+1}^0 -complete set while $\lim^{[k]}$ is Σ_{k+1}^0 -measurable. \square

Theorem 5.11. *For every $k \geq 1$, $\Delta_{k+1}^0\text{-DS} \equiv_W \Pi_k^0\text{-DS}$.*

Proof. The right-to-left reduction is trivial. To prove the left-to-right one it suffices to show that $\Delta_1^0\text{-DS}' \equiv_W \Pi_1^0\text{-DS}$ and the proof will follow from Theorem 5.3 as

$$\Delta_{k+1}^0\text{-DS} \equiv_W \Delta_1^0\text{-DS}' * \lim^{[k-1]} \equiv_W \Pi_1^0\text{-DS} * \lim^{[k-1]} \equiv_W \Pi_k^0\text{-DS}.$$

Let $p = (p_n)_{n \in \mathbb{N}}$ be a sequence in $\mathbb{N}^{\mathbb{N}}$ converging to the characteristic function of an ill-founded linear order L . In the following it is convenient to consider also the sequence $q = (q_n)_{n \in \mathbb{N}}$, where $q_n(i) := p_n(\langle i, i \rangle)$. Clearly q converges to the characteristic function of $\text{dom}(L)$ and is uniformly computable from p .

For sake of readability, define the formula

$$\varphi((x_n)_{n \in \mathbb{N}}, \sigma) := (\forall i < |\sigma|)(x_{\sigma(i)}(i) \neq x_{\sigma(i)+1}(i) \wedge (\forall j > \sigma(i))(x_j(i) = x_{j+1}(i))).$$

Intuitively φ says that, for each $i < |\sigma|$, $\sigma(i)$ codes the positions in which the sequence $(x_n)_{n \in \mathbb{N}}$ changes for the last time in the i -th row. Let us also write $x_\sigma := |\sigma| - 1$. We define

$$\begin{aligned} M := \{(\sigma, \tau) \in \mathbb{N}^{<\mathbb{N}} \times \mathbb{N}^{<\mathbb{N}} : \varphi(q, \sigma) \wedge \\ q_{\sigma(x_\sigma+1)}(x_\sigma) = 1 \wedge \\ \varphi(p, \tau) \wedge |\tau| = \langle x_\sigma, x_\sigma \rangle + 1\} \end{aligned}$$

Notice that the first two conditions imply that $x_\sigma \in L$. Intuitively x_σ is the $\leq_{\mathbb{N}}$ -largest element that is witnessed by σ to enter in L . The last line says that τ is exactly as long as needed to witness all the relations between the elements of L that are $\leq_{\mathbb{N}} x_\sigma$.

We order the set M as follows:

$$(\sigma_0, \tau_0) \leq_M (\sigma_1, \tau_1) :\iff x_{\sigma_0} \leq_L x_{\sigma_1}$$

Notice that M is a $\Pi_1^{0,p}$ linear order as M is $\Pi_1^{0,p}$ and the order \leq_M is p -computable: indeed, given two pairs $(\sigma_0, \tau_0), (\sigma_1, \tau_1) \in M$, we can use the longer string between τ_0 and τ_1 to p -compute whether $x_{\sigma_0} \leq_L x_{\sigma_1}$. Notice also that, for each l , there is exactly one string σ of length l witnessing $\varphi(q, \sigma)$ (by minimality). The third line in the definition of M implies that if σ satisfies the first two conditions then there is a unique τ s.t. $(\sigma, \tau) \in M$. The linearity of M follows by the linearity of L .

To conclude the proof it is enough to notice that if $((\sigma_i, \tau_i))_{i \in \mathbb{N}}$ is an $<_M$ -descending sequence then $(x_{\sigma_i})_{i \in \mathbb{N}}$ is an $<_L$ -descending sequence. \square

The following is essentially a classical result (see e.g. [15, Thm. 2.4]). The proof is simple enough that we can briefly sketch it.

Theorem 5.12. *For every $k \geq 1$, $\Sigma_k^0\text{-DS} \equiv_W \Delta_k^0\text{-DS}$.*

Proof. Given a Σ_k^0 linear order L , we can uniformly consider a sequence $((L_s, \leq_s))_{s \in \mathbb{N}}$ of Δ_k^0 linear orders approximating L . We then define

$$\begin{aligned} M := \{(q, s) : q \in L_s \text{ and } (\forall t < s)(q \notin L_t)\}, \\ (p, s) \leq_M (q, t) :\iff p \leq_L q. \end{aligned}$$

Notice that $(p, s) \leq_M (q, t)$ can be written also as $p = q \vee (\forall i)(q \not\leq_i p)$, hence M is $\Delta_k^{0,L}$. Moreover, since for every $q \in L$ there is a unique s s.t. $(q, s) \in M$, it is easy to see that M is computably isomorphic to L . In particular, given an $<_M$ -descending sequence we can obtain an $<_L$ -descending sequence by projection. \square

Corollary 5.13. *For every $k \geq 1$, we have*

$$\Pi_k^0\text{-DS} \equiv_W \Delta_{k+1}^0\text{-DS} \equiv_W \Sigma_{k+1}^0\text{-DS}.$$

Proof. The equivalence $\mathbf{\Pi}_k^0\text{-DS} \equiv_{\text{W}} \mathbf{\Delta}_{k+1}^0\text{-DS}$ was proved in Theorem 5.11. The equivalence $\mathbf{\Delta}_{k+1}^0\text{-DS} \equiv_{\text{W}} \mathbf{\Sigma}_{k+1}^0\text{-DS}$ was proved in Theorem 5.12. \square

Theorem 5.14. *For every $k \geq 1$, $\text{LPO}^{(k)} \leq_{\text{W}} \mathbf{\Sigma}_k^0\text{-BS}$ and therefore $\mathbf{\Sigma}_k^0\text{-BS} \not\leq_{\text{W}} \mathbf{\Sigma}_k^0\text{-DS}$.*

Proof. The second statement follows from the first because $\text{LPO}^{(k)} \not\leq_{\text{W}} \mathbf{\Delta}_k^0\text{-DS}$ (proof of Theorem 5.10) and $\mathbf{\Delta}_k^0\text{-DS} \equiv_{\text{W}} \mathbf{\Sigma}_k^0\text{-DS}$ (Theorem 5.12).

To prove the first statement, it is enough to show that $\text{LPO}' \leq_{\text{W}} \mathbf{\Sigma}_1^0\text{-BS}$, and the claim will follow by Theorem 5.3 as

$$\text{LPO}^{(k)} \leq_{\text{W}} \text{LPO}' * \text{lim}^{[k-1]} \leq_{\text{W}} \mathbf{\Sigma}_1^0\text{-BS} * \text{lim}^{[k-1]} \equiv_{\text{W}} \mathbf{\Sigma}_k^0\text{-BS}.$$

Let $(p_s)_{s \in \mathbb{N}}$ be a sequence in $\mathbb{N}^{\mathbb{N}}$ converging to an instance p of LPO. For every $s \in \mathbb{N}$ we define (as we did in the proofs of Theorem 4.18 and Proposition 4.20)

$$g(s) = \begin{cases} i + 1 & \text{if } i \leq s \wedge p_s(i) \neq 0 \wedge (\forall j < i)(p_s(j) = 0), \\ 0 & \text{otherwise.} \end{cases}$$

Let us define a quasi-order Q inductively: at stage $s = 0$ we add $\langle g(0), 0 \rangle$. At stage $s + 1$ we do the following:

- (1) if $g(s) = g(s + 1)$ we put $\langle g(s), s + 1 \rangle$ immediately below $\langle g(s), s \rangle$;
 - (2) if $g(s) \neq g(s + 1)$ we put $\langle g(s + 1), s + 1 \rangle$ at the top and we put $\langle -1, s + 1 \rangle$ at the bottom.
- Moreover we collapse to a single equivalence class all the elements $\langle g, t \rangle$ with $t \leq s$ and $g \neq -1$.

This construction produces a quasi-order (Q, \preceq_Q) which is computable in $(p_s)_{s \in \mathbb{N}}$.

Notice that if there is an s s.t. for every $t \geq s$, $g(t) = g(s)$ (in particular, this is the case if $\text{LPO}(p) = 1$) then the equivalence classes of \preceq_Q form a linear order of type $n + \omega^*$ and every \preceq_Q -bad sequence is a descending sequence of the form $(\langle g(s), s_n \rangle)_{n \in \mathbb{N}}$ for some strictly increasing sequence $(s_n)_{n \in \mathbb{N}}$. On the other hand, if the sequence $(g(s))_{s \in \mathbb{N}}$ does not stabilize then the equivalence classes of \preceq_Q are linearly ordered as ω^* , where all the elements $\langle g, s \rangle$ with $g \neq -1$ are equivalent and lie in the top equivalence class. This shows that the construction produces a non-well quasi-order.

For every \preceq_Q -bad sequence $(\langle g_n, s_n \rangle)_{n \in \mathbb{N}}$ produced by $\mathbf{\Sigma}_1^0\text{-BS}(Q)$, we compute the solution for $\text{LPO}'((p_s)_{s \in \mathbb{N}}) = \text{LPO}(p)$ by returning 0 if $g_1 \leq 0$ and 1 otherwise. We consider two cases. If the sequence $(g(s))_{s \in \mathbb{N}}$ stabilizes, then the sequence $(g_n)_{n \in \mathbb{N}}$ is constant. Furthermore, its value is 0 if $\text{LPO}(p) = 0$, otherwise its value is positive. On the other hand, if the sequence $(g(s))_{s \in \mathbb{N}}$ does not stabilize, then $\text{LPO}(p) = 0$. Furthermore, for every $n > 0$, we have $g_n = -1 \leq 0$. (The first element $\langle g_0, s_0 \rangle$ may lie in the top equivalence class, in which case g_0 may be positive. Hence we check g_1 instead of g_0 .) \square

5.2. $\mathbf{\Gamma}_1^1\text{-DS}$ and $\mathbf{\Gamma}_1^1\text{-BS}$. We now turn our attention to the analytic classes. Notice first of all that being a descending sequence through a $\mathbf{\Sigma}_1^1$ linear order is a $\mathbf{\Sigma}_1^1$ -property, hence $\mathbf{\Sigma}_1^1\text{-DS} \leq_{\text{W}} \mathbf{\Sigma}_1^1\text{-C}_{\mathbb{N}^{\mathbb{N}}} \equiv_{\text{W}} \text{C}_{\mathbb{N}^{\mathbb{N}}}$. We will show that $\mathbf{\Sigma}_1^1\text{-DS}$ is the strongest DS-principle that is still reducible to $\text{C}_{\mathbb{N}^{\mathbb{N}}}$ (Theorem 5.25).

Proposition 5.15. $\mathbf{\Delta}_1^1\text{-DS} \equiv_{\text{W}} \text{DS} * \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ and $\mathbf{\Delta}_1^1\text{-BS} \equiv_{\text{W}} \text{BS} * \text{UC}_{\mathbb{N}^{\mathbb{N}}}$.

Proof. We will only prove the first statement. The proof of the second statement is similar.

To prove the left-to-right reduction, given a $\mathbf{\Delta}_1^1$ name for L we use $\mathbf{\Delta}_1^1\text{-CA}$ (which is known to be equivalent to $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$, see [26, Thm. 3.11]) to compute a $\mathbf{\Delta}_1^0$ name for L . We can then apply DS to find a descending sequence through L .

To prove the converse reduction, using the cylindrical decomposition we can write

$$\text{DS} * \text{UC}_{\mathbb{N}^{\mathbb{N}}} \equiv_{\text{W}} \text{DS} \circ \Phi_e \circ \text{UC}_{\mathbb{N}^{\mathbb{N}}}$$

for some computable function Φ_e . In particular, given $T \subset \mathbb{N}^{<\mathbb{N}}$ with a unique path x , $\Phi_e(x)$ is the characteristic function of a linear order L . Notice that x is $\Delta_1^{1,T}$ -computable. Indeed,

$$\begin{aligned} x(n) = k &\iff (\exists \sigma \in T)(\sigma \in \text{Ext} \wedge \sigma(n) = k) \\ &\iff (\forall \tau \in T)(\tau \in \text{Ext} \rightarrow \tau(n) = k), \end{aligned}$$

where Ext is the set of finite strings that extend to a path through T ($\sigma \in \text{Ext}$ is a $\Sigma_1^{1,T}$ property). We can therefore obtain a $\Delta_1^{1,T}$ name for L as

$$a \leq_L b \iff \Phi_e(x)(\langle a, b \rangle) = 1,$$

and hence we use Δ_1^1 -DS to find a descending sequence through L . \square

In particular, this implies that Δ_1^1 is the first level at which we can compute $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$. Indeed, for every k , we showed in the proof of Theorem 5.10 that $\text{LPO}^{(k)} \not\leq_W \Delta_k^0$ -DS, while $\lim^{[k]} \leq_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ (see [4, Sec. 6]).

By adapting the proof of Corollary 5.4, we can relativize Proposition 4.5 and obtain the following:

Corollary 5.16. Δ_1^1 -DS \equiv_W Δ_1^1 -BS.

Proof. See Errata. \square

Similarly, the proofs of Theorem 5.5 and of Proposition 5.8 lead to the following equivalences:

Theorem 5.17.

$${}^1\Delta_1^1\text{-DS} \equiv_W {}^1(\Pi_1^1\text{-Bound} \times \text{UC}_{\mathbb{N}^{\mathbb{N}}}) \equiv_W \left(\bigsqcup_{s \in \mathbb{N}} \Delta_1^1\text{-C}_s \right) * \Pi_1^1\text{-Bound}.$$

The deterministic part of Δ_1^1 -DS and Σ_1^1 -DS can be easily characterized using Proposition 5.15, as the following proposition shows.

Proposition 5.18. $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \equiv_W \text{Det}(\Delta_1^1\text{-DS}) \equiv_W \text{Det}(\Sigma_1^1\text{-DS})$.

Proof. The reductions $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \leq_W \text{Det}(\Delta_1^1\text{-DS}) \leq_W \text{Det}(\Sigma_1^1\text{-DS})$ are straightforward from $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \leq_W \Delta_1^1\text{-DS}$ (Proposition 5.15), $\Delta_1^1\text{-DS} \leq_W \Sigma_1^1\text{-DS}$ (trivial) and the fact that $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$ is single-valued. To prove that $\text{Det}(\Sigma_1^1\text{-DS}) \leq_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ it is enough to notice that $\Sigma_1^1\text{-DS} \leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$, and therefore $\text{Det}(\Sigma_1^1\text{-DS}) \leq_W \text{Det}(\text{C}_{\mathbb{N}^{\mathbb{N}}}) \equiv_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ (Theorem 3.14). \square

In particular, the deterministic part does not help us separate Δ_1^1 -DS and Σ_1^1 -DS. Instead, we separate them by considering their first-order parts. We characterized ${}^1\Delta_1^1$ -DS in Theorem 5.17. Notice that our proof (see the proof of Proposition 5.8) cannot be extended to establish the same result for Σ_1^1 -DS, because the definition of the corresponding $(F_s)_{s \in \mathbb{N}}$ would not be Σ_1^1 .

Proposition 5.19. $\widehat{\Sigma_1^1\text{-C}_{\mathbb{N}}} \leq_W \Sigma_1^1\text{-DS}$.

Proof. Let $(A_i)_{i \in \mathbb{N}}$ be a sequence of non-empty Σ_1^1 subsets of \mathbb{N} . We define

$$\begin{aligned} L &:= \{(n, \sigma) \in \mathbb{N} \times \mathbb{N}^{<\mathbb{N}} : |\sigma| = n \wedge (\forall i < n)(\sigma(i) \in A_i)\}, \\ (n, \sigma) \leq_L (m, \tau) &\iff n > m \vee (n = m \wedge \sigma \leq_{lex} \tau). \end{aligned}$$

It is easy to see that L is a Σ_1^1 linear order (the linearity follows from the linearity of \leq and of \leq_{lex}).

Let $((n_i, \sigma_i))_{i \in \mathbb{N}}$ be an $<_L$ -descending sequence. Notice that, since each $A_i \subset \mathbb{N}$, for each n the set $\{\sigma \in \mathbb{N}^{<\mathbb{N}} : (n, \sigma) \in L\}$ is \leq_{lex} -well-founded. Therefore there must be a subsequence $((n_{i_k}, \sigma_{i_k}))_{k \in \mathbb{N}}$ s.t. the sequence $(n_{i_k})_{k \in \mathbb{N}}$ is strictly increasing.

This implies that, for each n , there is some m s.t. $|\sigma_m| \geq n$. In particular, by definition of L , $(\forall i < n)(\sigma_m(i) \in A_i)$ and the claim follows. \square

Proposition 5.19 implies that $\Sigma_1^1\text{-C}_{\mathbb{N}} \leq_W {}^1\Sigma_1^1\text{-DS}$. This, together with ${}^1\text{C}_{\mathbb{N}^{\mathbb{N}}} \equiv_W \Sigma_1^1\text{-C}_{\mathbb{N}}$ (Proposition 2.4) and the observation that $\Sigma_1^1\text{-DS} \leq_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$, immediately yields the following:

Corollary 5.20. ${}^1\text{C}_{\mathbb{N}^{\mathbb{N}}} \equiv_W {}^1\Sigma_1^1\text{-DS} \equiv_W \Sigma_1^1\text{-C}_{\mathbb{N}}$.

As a consequence, $\text{C}_{\mathbb{N}^{\mathbb{N}}}$ and $\Sigma_1^1\text{-DS}$ cannot be separated by means of their first-order part. But $\Delta_1^1\text{-DS}$ and $\Sigma_1^1\text{-DS}$ can, albeit somewhat indirectly:

Proposition 5.21. $\Delta_1^1\text{-DS} <_W \Sigma_1^1\text{-DS}$.

Proof. Notice first of all that $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \leq_W \widehat{\Pi_1^1\text{-Bound}}$. Indeed, given a tree $T \subset \mathbb{N}^{<\mathbb{N}}$ with a unique path, we can consider the following sequence of Π_1^1, T sets:

$$A_n := \{k \in \mathbb{N} : (\forall \sigma \in T)((\exists x \in [T])(\sigma \sqsubset x) \rightarrow \sigma(n) \leq k)\}.$$

Clearly each A_n is bounded by $x(n)$, where x is the unique path through T . Given a solution $f \in \widehat{\Pi_1^1\text{-Bound}}((A_n)_{n \in \mathbb{N}})$, consider the space $X := \{\sigma \in \mathbb{N}^{<\mathbb{N}} : (\forall i < |\sigma|)(\sigma(i) \leq f(i))\}$ and define $T_f := T \cap X$. Notice that $[T_f] = [T]$. In particular, since $[X]$ is f -computably compact, we can uniformly (in f) compute the unique path through $[T_f]$ (see [10, Thm. 7.23 and Cor. 7.26]).

If $\Sigma_1^1\text{-C}_{\mathbb{N}} \leq_W \Delta_1^1\text{-DS}$ then, by Theorem 5.17, $\Sigma_1^1\text{-C}_{\mathbb{N}} \leq_W \text{UC}_{\mathbb{N}^{\mathbb{N}}} \times \widehat{\Pi_1^1\text{-Bound}}$ and therefore

$$\widehat{\Sigma_1^1\text{-C}_{\mathbb{N}}} \leq_W (\text{UC}_{\mathbb{N}^{\mathbb{N}}} \times \widehat{\Pi_1^1\text{-Bound}}) \equiv_W \widehat{\Pi_1^1\text{-Bound}},$$

contradicting $\widehat{\Sigma_1^1\text{-C}_{\mathbb{N}}} \not\leq_W \widehat{\Pi_1^1\text{-Bound}}$ ([1, Cor. 3.23]). \square

To separate $\Sigma_1^1\text{-DS}$ from $\text{C}_{\mathbb{N}^{\mathbb{N}}}$ we generalize a technique based on inseparable Π_1^1 sets, first used in [1] to separate $\widehat{\Sigma_1^1\text{-C}_{\mathbb{N}}}$ from $\text{C}_{\mathbb{N}^{\mathbb{N}}}$. Consider the problem $\text{ATR}_2 : \text{LO} \times 2^{\mathbb{N}} \rightrightarrows \{0, 1\} \times \mathbb{N}^{\mathbb{N}}$ defined in [21, Def. 8.2]. It can be seen as a two-sided version of ATR : it takes in input a pair (L, A) and produces a pair (i, Y) s.t. either $i = 0$ and Y is a $<_L$ -infinite descending sequence or $i = 1$ and Y is a jump (pseudo)hierarchy starting from A . Jun Le Goh proved that $\text{UC}_{\mathbb{N}^{\mathbb{N}}} <_W \text{ATR}_2 <_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$ ([21, Cor. 8.5 and 8.7]).

Before proving the next theorem, we introduce the following notion of reducibility: for every $A, B \subset \mathbb{N}^{\mathbb{N}}$, we say that A is *Muchnik reducible* to B , and write $A \leq_w B$ if, for every $b \in B$ there is a Turing functional Φ_e s.t. $\Phi_e(b) \in A$. Muchnik reducibility is the non-uniform version of Medvedev reducibility. For an extended presentation on these notions of reducibility see e.g. [40].

Theorem 5.22. $\text{ATR}_2 \upharpoonright_W \Sigma_1^1\text{-DS}$, and therefore $\Sigma_1^1\text{-DS} <_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$.

Proof. The fact that $\Sigma_1^1\text{-DS} \not\leq_W \text{ATR}_2$ follows from the fact that $\text{C}_{\mathbb{N}^{\mathbb{N}}} \equiv_W \lim * \Sigma_1^1\text{-DS}$ while $\lim * \text{ATR}_2 <_W \text{C}_{\mathbb{N}^{\mathbb{N}}}$ ([21, Cor. 8.5]).

Let us now prove that $\text{ATR}_2 \not\leq_W \Sigma_1^1\text{-DS}$. Assume towards a contradiction that there is a reduction witnessed by the maps Φ, Ψ . Let $(L_e)_{e \in \mathbb{N}}$ be an enumeration of the computable linear orders. Define the sets

$$\begin{aligned} S_e &:= \Sigma_1^1\text{-DS}(\Phi(L_e)), \\ DS_e &:= \{(x_n)_n \in \mathbb{N}^{\mathbb{N}} : (x_n)_n \text{ is an } <_{L_e}\text{-descending sequence}\}, \\ JH_e &:= \{(y_n)_n \in \mathbb{N}^{\mathbb{N}} : (y_n)_n \text{ is a jump hierarchy on } L_e\}. \end{aligned}$$

Notice that, for each e , S_e is Σ_1^1 (being a descending sequence through a Σ_1^1 linear order is a Σ_1^1 condition) while DS_e and JH_e are arithmetic.

Define now the sets

$$\begin{aligned} B &:= \{e \in \mathbb{N} : DS_e \not\leq_w S_e\}, \\ C &:= \{e \in \mathbb{N} : JH_e \not\leq_w S_e\}, \end{aligned}$$

where \leq_w represents Muchnik reducibility. In particular, if X is (hyper)arithmetic and Y is Σ_1^1 then $X \not\leq_w Y$ is a Σ_1^1 condition, and therefore $B, C \in \Sigma_1^1(\mathbb{N})$.

We now claim that $B \cap C = \emptyset$. Indeed, assume by contradiction that this is not the case and let $e \in B \cap C$. By definition of B and C this means that there are two descending sequences $(q_n)_{n \in \mathbb{N}}$ and $(p_n)_{n \in \mathbb{N}}$ in $\Phi(L_e)$ s.t. $(q_n)_{n \in \mathbb{N}}$ does not compute any $<_{L_e}$ -descending sequence and $(p_n)_{n \in \mathbb{N}}$ does not compute any jump hierarchy on L_e .

In particular, if we run the backward functional Ψ on $(q_n)_{n \in \mathbb{N}}$ and $(p_n)_{n \in \mathbb{N}}$ then, by continuity, there is an n s.t. $\Psi((q_i)_{i < n})$ commits to producing a jump hierarchy on L_e and $\Psi((p_i)_{i < n})$ commits to producing an $<_{L_e}$ -descending sequence. Without loss of generality, assume that $q_n \leq_{\Phi(L_e)} p_n$ (in the opposite case we just swap the roles of $(q_n)_{n \in \mathbb{N}}$ and $(p_n)_{n \in \mathbb{N}}$) and consider the sequence

$$r := \langle p_0, \dots, p_n, q_{n+1}, q_{n+2}, \dots \rangle.$$

Notice that $\Psi(r)$ must produce an $<_{L_e}$ -descending sequence, contradicting the fact that $(q_n)_{n \in \mathbb{N}}$ does not compute any $<_{L_e}$ -descending sequence.

Let $\mathbf{wf}_{\mathbf{LO}}$ be the set of indexes for the computable well-orderings and let \mathbf{hds} be the set of indexes for computable linear orderings with a hyperarithmetic descending sequence. Notice that $\mathbf{wf}_{\mathbf{LO}} \subset B$, because for each e in $\mathbf{wf}_{\mathbf{LO}}$, $DS_e = \emptyset \not\leq_w A$ for every non-empty set A . Likewise, $\mathbf{hds} \subset C$, as any ill-founded linear order which has a hyperarithmetic descending sequence cannot support a jump hierarchy (see² [18, Thm. 4]).

Since B, C are disjoint and Σ_1^1 , by Σ_1^1 -separation there must be a Δ_1^1 set separating them. Such a set would separate $\mathbf{wf}_{\mathbf{LO}}$ and \mathbf{hds} as well. This contradicts the fact that every Σ_1^1 set which separates $\mathbf{wf}_{\mathbf{LO}}$ and \mathbf{hds} must be Σ_1^1 -complete [20]. \square

Finally we turn our attention to Σ_1^1 -BS and Π_1^1 -DS. We show below that these problems are much stronger in uniform computational strength than the problems considered so far. Indeed all the Γ -DS problems, where $\Gamma = \Sigma_1^1$ or below, are s.t.

$$\Gamma\text{-DS} <_W C_{\mathbb{N}^{\mathbb{N}}} \equiv_W \lim * \Gamma\text{-DS}.$$

In other words, Γ -DS is arithmetically Weihrauch equivalent to $C_{\mathbb{N}^{\mathbb{N}}}$, which is prominent among the problems that are considered to be “ATR₀ analogues in the Weihrauch lattice” [26].

On the other hand, a natural analogue of Π_1^1 -CA₀ in the Weihrauch lattice is Π_1^1 -CA, which can be phrased as “given a sequence $(T_n)_{n \in \mathbb{N}}$ of trees in $\mathbb{N}^{<\mathbb{N}}$, produce $x \in 2^{\mathbb{N}}$ s.t., for every n , $x(n) = 1$ iff $[T_n] = \emptyset$ ”.

We can notice that, using [30, Thm. 6.5], Π_1^1 -CA is equivalent to the problem of finding the leftmost path through an ill-founded tree. Using this fact we show that Σ_1^1 -BS and Π_1^1 -DS are in the realm of Π_1^1 -CA₀.

Theorem 5.23. Π_1^1 -CA $\leq_W \Sigma_1^1$ -BS.

Proof. Let $T \subset \mathbb{N}^{<\mathbb{N}}$ be an ill-founded tree. For each $\sigma \in T$, let $T_\sigma := \{\tau \in T : \tau \sqsubseteq \sigma \vee \sigma \sqsubseteq \tau\}$. We define a quasi-order on the extendible strings in T :

$$Q := \{\sigma \in T : [T_\sigma] \neq \emptyset\},$$

$$\sigma \preceq_Q \tau :\iff (\exists \rho \in Q)(\rho <_{lex} \sigma) \vee \tau \sqsubseteq \sigma.$$

It is easy to see that (Q, \preceq_Q) is Σ_1^1 . Moreover, all the σ which are not prefixes of the leftmost path collapse in a bottom equivalence class. This shows that the equivalence classes of Q are linearly ordered as $1 + \omega^*$. To conclude the proof it is enough to notice that any $<_Q$ -descending sequence gives longer and longer prefixes of the leftmost path, hence it computes Π_1^1 -CA. \square

Corollary 5.24. Σ_1^1 -DS $<_W \Sigma_1^1$ -BS.

²Friedman’s result assumes that the linear order is adequate. We do not need this assumption because we choose to define jump hierarchies in a way such that each column (whether limit or successor) uniformly computes earlier columns, such as in [21, Def. 3.1]. This allows us to run Friedman’s proof without assuming adequacy.

Proof. We have $\Sigma_1^1\text{-DS} \leq_W \mathbf{C}_{\mathbb{N}^{\mathbb{N}}} <_W \Pi_1^1\text{-CA} \leq_W \Sigma_1^1\text{-BS}$. \square

Theorem 5.25. $\Pi_1^1\text{-CA} \leq_W \Pi_1^1\text{-DS}$.

Proof. Let $T \subset \mathbb{N}^{<\mathbb{N}}$ be an ill-founded tree. For each $\sigma \in T$, let $T_\sigma := \{\tau \in T : \tau \sqsubseteq \sigma \vee \sigma \sqsubseteq \tau\}$. We define a linear order

$$L := \{\sigma \in T : (\forall \tau \leq_{lex} \sigma)([T_\tau] = \emptyset \vee \tau \sqsubseteq \sigma)\},$$

$$\leq_L := \leq_{KB(T)}.$$

Clearly (L, \leq_L) is a Π_1^1, T linear order. Notice that if $\sigma \in L$ and $[T_\sigma] \neq \emptyset$ then σ must be a prefix of the leftmost path. Moreover if ρ is strictly lexicographically above the leftmost path then $\rho \notin L$. In other words, L is the subset of T that is lexicographically below the leftmost path.

Moreover, every string that is not a prefix of the leftmost path lies in the well-founded part of L (by definition of KB). In particular every $<_L$ -descending sequence computes arbitrarily long prefixes of the leftmost path. \square

6. CONCLUSIONS

In this paper we explored the uniform computational content of the problem DS, and showed how it lies “on the side” w.r.t. the part of the Weihrauch lattice explored so far. We now draw the attention to some of the questions that did not receive an answer.

The problem KL is the multi-valued function corresponding to König’s lemma, and it can be phrased as “find a path through an infinite finitely-branching tree”. It is known that $\text{KL} \equiv_W \mathbf{C}'_{2^{\mathbb{N}}} \equiv_W \widehat{\text{RT}}_2^1$.

Open Question 6.1. $\text{KL} \leq_W \text{DS}$?

We know that, if such a reduction exists, it must be strict (as KL is an arithmetic problem). On the other hand, none of the characterizations we used in Section 4 to describe the lower cone of DS can be used to prove a separation.

In Section 3.4 we introduced the problem $\text{wList}_{2^{\mathbb{N}}, \leq \omega}$. Similarly to DS, this problem does not fit well within the effective Baire hierarchy: $\text{Det}(\text{wList}_{2^{\mathbb{N}}, \leq \omega}) \equiv_W \text{lim}$, but $\text{wList}_{2^{\mathbb{N}}, \leq \omega}^{[3]} \equiv_W \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ ([26, Prop. 6.14 and Cor. 6.16]), hence in particular $\text{wList}_{2^{\mathbb{N}}, \leq \omega}$ is not arithmetic.

Open Question 6.2. $\text{wList}_{2^{\mathbb{N}}, \leq \omega} \leq_W \text{DS}$?

Our results imply that $\text{DS} \not\leq_W \text{wList}_{2^{\mathbb{N}}, \leq \omega}$ (as $\text{DS} * \text{DS} \equiv_W \mathbf{C}_{\mathbb{N}^{\mathbb{N}}}$), and hence a reduction would be strict.

In the context of Γ -DS, there are a few problems that resisted full characterization. In particular:

Open Question 6.3. $\Delta_2^0\text{-DS} \leq_W \Sigma_1^0\text{-BS}$?

We expect that an answer to this question will yield a solution for every k (by relativization).

We notice that, in the statements involving Γ -BS we proved slightly more than what claimed: indeed, in all the reductions, the quasi-order built is a *linear quasi-order*, i.e. a quasi-order whose equivalence classes are linearly ordered. Notice that every bad sequence through a non-well linear quasi-order is actually a descending sequence. If we introduce the problem $\Gamma\text{-DS}_{LQO}$ by restricting Γ -BS to linear quasi-orders, our results imply that

$$\Delta_k^0\text{-DS} <_W \Sigma_{k+1}^0\text{-DS}_{LQO} \leq_W \Sigma_{k+1}^0\text{-BS}.$$

A natural question is therefore

Open Question 6.4. $\Sigma_{k+1}^0\text{-BS} \leq_W \Sigma_{k+1}^0\text{-DS}_{LQO}$?

A negative answer would imply that the possibility of having infinite antichains provides extra uniform strength.

A very important structure that is left out of the picture is the one of partial orders. In the same spirit of the paper we can consider the problems Γ -DS_{PO} and Γ -BS_{PO}. The former is readily seen to be equivalent to $C_{\mathbb{N}^{\mathbb{N}}}$ (see also the comment before Definition 4.4).

Open Question 6.5. What is the relation between Σ_1^0 -BS_{PO} and the problems DS \equiv_W Σ_1^0 -DS, Σ_1^0 -DSLQO and Σ_1^0 -BS?

Answering these questions would yield very interesting insights on how the possibility to have equivalent non-equal elements can enhance the uniform computational strength.

REFERENCES

1. Paul-Elliot Anglès D’Auriac and Takayuki Kihara, *A Comparison Of Various Analytic Choice Principles*, Available at <https://arxiv.org/abs/1907.02769>, 2019.
2. Eric P. Astor, Damir D. Dzhabarov, Reed Solomon, and Jacob Suggs, *The uniform content of partial and linear orders*, *Annals of Pure and Applied Logic* **168** (2017), no. 6, 1153–1171, DOI: 10.1016/j.apal.2016.11.010.
3. Vasco Brattka, *Effective Borel measurability and reducibility of functions*, *Mathematical Logic Quarterly* **51** (2005), no. 1, 19–44, DOI: 10.1002/malq.200310125.
4. Vasco Brattka, Matthew de Brecht, and Arno Pauly, *Closed choice and a Uniform Low Basis Theorem*, *Annals of Pure and Applied Logic* **163** (2012), no. 8, 986–1008, DOI: 10.1016/j.apal.2011.12.020.
5. Vasco Brattka and Guido Gherardi, *Weihrauch degrees, omniscience principles and weak computability*, *The Journal of Logic* **76** (2011), no. 1, 143–176.
6. ———, *Weihrauch goes Brouwerian*, Available at <https://arxiv.org/abs/1809.00380>, 2018.
7. ———, *Completion of Choice*, Available at <https://arxiv.org/abs/1910.13186>, 2019.
8. Vasco Brattka, Guido Gherardi, and Rupert Hözl, *Probabilistic computability and choice*, *Information and Computation* **242** (2015), 249–286, DOI: 10.1016/j.ic.2015.03.005.
9. Vasco Brattka, Guido Gherardi, and Alberto Marcone, *The Bolzano-Weierstrass Theorem is the jump of Weak König’s Lemma*, *Annals of Pure and Applied Logic* **163** (2012), no. 6, 623–655, DOI: 10.1016/j.apal.2011.10.006.
10. Vasco Brattka, Guido Gherardi, and Arno Pauly, *Weihrauch complexity in computable analysis*, To appear. Available at <https://arxiv.org/pdf/1707.03202v1>, 07 2017.
11. Vasco Brattka, Rupert Hözl, and Rutger Kuyper, *Monte Carlo computability*, 34th Symposium on Theoretical Aspects of Computer Science (STACS 2017) (Dagstuhl, Germany) (Heribert Vollmer and Brigitte Vallée, eds.), *Leibniz International Proceedings in Informatics (LIPIcs)*, vol. 66, Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, 2017, DOI: 10.4230/LIPIcs.STACS.2017.17, Available at <http://drops.dagstuhl.de/opus/volltexte/2017/7016>, pp. 17:1–17:14.
12. Vasco Brattka and Arno Pauly, *On the algebraic structure of Weihrauch degrees*, *Logical Methods in Computer Science* **14** (2018), no. 4, 1–36.
13. Vasco Brattka and Tahina Rakotoniaina, *On the uniform computational content of Ramsey’s theorem*, *The Journal of Logic* **82** (2017), no. 4, 1278–1316, DOI: 10.1017/jsl.2017.43.
14. François G. Dorais, Damir D. Dzhabarov, Jeffrey L. Hirst, Joseph R. Mileti, and Paul Shafer, *On uniform relationships between combinatorial problems*, *Transactions of the American Mathematical Society* **368** (2016), 1321–1359, DOI: 10.1090/tran/6465.
15. Rod G. Downey, *Computability theory and linear orderings*, *Handbook of Recursive Mathematics* (Yu. L. Ershov, S.S. Goncharov, A. Nerode, J.B. Remmel, and V.W. Marek, eds.), *Studies in Logic and the Foundations of Mathematics*, vol. 139, Elsevier, 1998, DOI: 10.1016/S0049-237X(98)80047-5, pp. 823–976.

16. Damir D. Dzhafarov, Jun Le Goh, Denis R. Hirschfeldt, Ludovic Patey, and Arno Pauly, *Ramsey's theorem and products in the Weihrauch degrees*, *Computability* **9** (2020), no. 2, 85–110, DOI: 10.3233/COM-180203.
17. Damir D. Dzhafarov, Reed Solomon, and Keita Yokoyama, *On the first-order parts of Weihrauch degrees*, In preparation.
18. Harvey Friedman, *Uniformly Defined Descending Sequences of Degrees*, *The Journal of Logic* **41** (1976), no. 2, 363–367, DOI: 10.2307/2272234.
19. Guido Gherardi and Alberto Marcone, *How Incomputable Is the Separable Hahn-Banach Theorem?*, *Notre Dame Journal of Formal Logic* **50** (2009), no. 4, 393–425.
20. Jun Le Goh, *Inseparable Π_1^1 sets*, In preparation.
21. ———, *Some computability-theoretic reductions between principles around ATR_0* , Available at <https://arxiv.org/abs/1905.06868>, 2019.
22. Vassilios Gregoriades, Tamás Kispéter, and Arno Pauly, *A comparison of concepts from computable analysis and effective descriptive set theory*, *Mathematical Structures in Computer Science* (2016), DOI: 10.1017/S0960129516000128.
23. Joseph Harrison, *Recursive pseudo-well-orderings*, *Transactions of the American Mathematical Society* **131** (1968), no. 2, 526–543, DOI: 10.2307/1994961.
24. Denis R. Hirschfeldt and Richard A. Shore, *Combinatorial principles weaker than Ramsey's Theorem for pairs*, *The Journal of Symbolic Logic* **72** (2007), no. 1, 171–206, DOI: 10.2178/jsl/1174668391.
25. Carl G. Jockusch, Bart Kastermans, Steffen Lempp, Manuel Lerman, and Reed Solomon, *Stability and posets*, *The Journal of Symbolic Logic* **74** (2009), no. 2, 693–711.
26. Takayuki Kihara, Alberto Marcone, and Arno Pauly, *Searching for an analogue of ATR_0 in the Weihrauch lattice*, *Journal of Logic* (To appear), 1–35, Available at <https://arxiv.org/abs/1812.01549>.
27. Takayuki Kihara, Keng Meng Ng, and Arno Pauly, *Enumeration degrees and non-metrizable topology*, Available at <https://arxiv.org/abs/1904.04107>, 2019.
28. Takayuki Kihara and Arno Pauly, *Point degree spectra of represented spaces*, Available at <https://arxiv.org/abs/1405.6866>, 2014.
29. Stéphane Le Roux and Arno Pauly, *Finite choice, convex choice and finding roots*, *Logical Methods in Computer Science* (2015), DOI: 10.2168/LMCS-11(4:6)2015.
30. Alberto Marcone, *On the logical strength of Nash-Williams' theorem on transfinite sequences*, pp. 327–351, Clarendon Press, USA, 1996.
31. Alberto Marcone and Richard A. Shore, *The maximal linear extension theorem in second order arithmetic*, *Archive for Mathematical Logic* **50** (2011), no. 5, 543–564, DOI: 10.1007/s00153-011-0231-1.
32. Joseph S. Miller, *Degrees of unsolvability of continuous functions*, *The Journal of Logic* **69** (2004), no. 2, 555–584.
33. Yiannis N. Moschovakis, *Descriptive Set Theory*, *Mathematical Surveys and Monographs*, vol. 155, AMS, 2009.
34. Eike Neumann and Arno Pauly, *A topological view on algebraic computation models*, *Journal of Complexity* **44** (2018), 1–22, DOI: 10.1016/j.jco.2017.08.003.
35. Ludovic Patey, *The weakness of being cohesive, thin or free in reverse mathematics*, *Israel Journal of Mathematics* **216** (2016), 905–955, DOI: 10.1007/s11856-016-1433-3.
36. Arno Pauly, *On the topological aspects of the theory of represented spaces*, *Computability* **5** (2016), no. 2, 159–180, DOI: 10.3233/COM-150049.
37. Arno Pauly and Matthew de Brecht, *Descriptive Set Theory in the Category of Represented Spaces*, 30th Annual ACM/IEEE Symposium on Logic in Computer Science (LICS), 2015, DOI: 10.1109/LICS.2015.48, pp. 438–449.
38. Gerald E. Sacks, *Higher Recursion Theory*, 1 ed., Springer Verlag, 1990.

39. Stephen G. Simpson, *Nonprovability of Certain Combinatorial Properties of Finite Trees*, Harvey Friedman's Research on the Foundations of Mathematics (L.A. Harrington, M.D. Morley, A. S  dov, and S.G. Simpson, eds.), Studies in Logic and the Foundations of Mathematics, vol. 117, Elsevier, 1985, DOI: 10.1016/S0049-237X(09)70156-9, pp. 87–117.
40. ———, *Mass Problems and Intuitionism*, Notre Dame J. Formal Logic **49** (2008), no. 2, 127–136, DOI: 10.1215/00294527-2008-002.
41. ———, *Subsystems of Second Order Arithmetic*, 2 ed., Cambridge University Press, 2009.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON
Email address: junle.goh@wisc.edu

DEPARTMENT OF COMPUTER SCIENCE, UNIVERSITY OF SWANSEA
Email address: arno.m.pauly@gmail.com

DEPARTMENT OF MATHEMATICS, COMPUTER SCIENCE AND PHYSICS, UNIVERSITY OF UDINE
Email address: manlio.valenti@uniud.it