

UNIQUENESS AND WEAK-BV STABILITY FOR 2×2 CONSERVATION LAWS

GENG CHEN, SAM G. KRUPA, AND ALEXIS F. VASSEUR

ABSTRACT. Let a 1-d system of hyperbolic conservation laws, with two unknowns, be endowed with a convex entropy. We consider the family of small BV functions which are global solutions of this equation. For any small BV initial data, such global solutions are known to exist. Moreover, they are known to be unique among BV solutions verifying either the so-called Tame Oscillation Condition, or the Bounded Variation Condition on space-like curves. In this paper, we show that these solutions are stable in a larger class of weak (and possibly not even BV) solutions of the system. This result extends the classical weak-strong uniqueness results which allow comparison to a smooth solution. Indeed our result extends these results to a weak- BV uniqueness result, where only one of the solutions is supposed to be small BV , and the other solution can come from a large class. As a consequence of our result, the Tame Oscillation Condition, and the Bounded Variation Condition on space-like curves are not necessary for the uniqueness of solutions in the BV theory, in the case of systems with 2 unknowns. The method is L^2 based. It builds up from the theory of a-contraction with shifts, where suitable weight functions a are generated via the front tracking method.

CONTENTS

| | |
|--|----|
| 1. Introduction | 1 |
| 2. Preliminaries | 5 |
| 3. Weighted relative entropy and shifts | 6 |
| 4. Relative entropy for the Riemann problem | 10 |
| 5. Modified front tracking algorithm | 13 |
| 5.1. The Riemann solvers | 13 |
| 5.2. Construction of the approximate solutions | 14 |
| 6. The weight function a | 16 |
| 7. Proof of Proposition 3.2 | 19 |
| Appendix A. Proof of Lemma 2.6 | 23 |
| References | 23 |

1. INTRODUCTION

We consider 1-d system of hyperbolic conservation laws with two unknowns

$$u_t + (f(u))_x = 0 \quad t > 0, x \in \mathbb{R}, \quad (1.1)$$

where $(t, x) \in \mathbb{R}^+ \times \mathbb{R}$ are time and space, and $u = (u_1, u_2) \in \mathcal{V}_0 \subseteq \mathbb{R}^2$ is the unknown. The set of states \mathcal{V}_0 is supposed to be bounded, and we denote \mathcal{V} its interior. Then $f = (f_1, f_2) \in [C(\mathcal{V}_0)]^2 \cap [C^4(\mathcal{V})]^2$ is the flux function, and is assumed to be continuous on \mathcal{V}_0 and C^4 on \mathcal{V} . For any $g \in C^1(\mathcal{V})$, let us denote the vector valued function $g' = Dg$. Then, we denote the

Date: January 18, 2022.

2010 Mathematics Subject Classification. Primary 35L65; Secondary 76N15, 35L45, 35B35.

Key words and phrases. Compressible Euler system, Uniqueness, Stability, Relative entropy, Conservation law.

Acknowledgment. G. Chen is partially supported by NSF with grants DMS-1715012 and DMS-2008504. S. G. Krupa was partially supported by NSF-DMS Grant 1840314. A. F. Vasseur is partially supported by the NSF grant: DMS 1614918.

eigenvalues and associated right eigenvectors of f' on \mathcal{V} as λ_1, r_1 and λ_2, r_2 , corresponding to the 1 and 2 characteristic families respectively. Through the paper, we will make the following general assumptions on the system.

Assumption 1.1. *Assumptions on the system*

- (a) For any $u \in \mathcal{V}$: $\lambda_1(u) < \lambda_2(u)$.
- (b) For any $u \in \mathcal{V}$, and $i = 1, 2$: $\lambda'_i(u) \cdot r_i(u) \neq 0$.
- (c) There exists a strictly convex function $\eta \in C(\mathcal{V}_0) \cap C^3(\mathcal{V})$ and a function $q \in C(\mathcal{V}_0) \cap C^3(\mathcal{V})$ such that

$$q' = \eta' f', \quad \text{on } \mathcal{V}. \quad (1.2)$$

- (d) For any $b \in \mathcal{V}$, and any left eigenvector ℓ of $f'(b)$: the function $u \rightarrow \ell \cdot f(u)$ is either convex or concave on \mathcal{V} .
- (e) There exists $L > 0$ such that for any $u \in \mathcal{V}$ and $i = 1, 2$: $|\lambda_i(u)| \leq L$.
- (f) For $u_L \in \mathcal{V}$, we denote $s \rightarrow S_{u_L}^1(s)$ the 1-shock curve through u_L defined for $s > 0$. We choose the parametrization such that $s = |u_L - S_{u_L}^1(s)|$. Therefore, $(u_L, S_{u_L}^1(s), \sigma_{u_L}^1(s))$ is the 1-shock with left hand state u_L and strength s . Similarly, we define $s \rightarrow S_{u_R}^2$ to be the 2-shock curve such that $(S_{u_R}^2, u_R, \sigma_{u_R}^2(s))$ is the 2-shock with right hand state u_R and strength s . We assume that these curves are defined globally in \mathcal{V} for every $u_L \in \mathcal{V}$ and $u_R \in \mathcal{V}$.
- (g) (for 1-shocks) If (u_L, u_R) is an entropic Rankine-Hugoniot discontinuity with shock speed σ , then $\sigma > \lambda_1(u_R)$.
- (h) (for 1-shocks) If (u_L, u_R) (with $u_L \in B_\epsilon(d)$) is an entropic Rankine-Hugoniot discontinuity with shock speed σ verifying,

$$\sigma \leq \lambda_1(u_L),$$

then u_R is in the image of $S_{u_L}^1$. That is, there exists $s_{u_R} \in [0, s_{u_L})$ such that $S_{u_L}^1(s_{u_R}) = u_R$ (and hence $\sigma = \sigma_{u_L}^1(s_{u_R})$).

- (i) (for 2-shocks) If (u_L, u_R) is an entropic Rankine-Hugoniot discontinuity with shock speed σ , then $\sigma < \lambda_2(u_L)$.
- (j) (for 2-shocks) If (u_L, u_R) (with $u_R \in B_\epsilon(d)$) is an entropic Rankine-Hugoniot discontinuity with shock speed σ verifying,

$$\sigma \geq \lambda_2(u_R),$$

then u_L is in the image of $S_{u_R}^2$. That is, there exists $s_{u_L} \in [0, s_{u_R})$ such that $S_{u_R}^2(s_{u_L}) = u_L$ (and hence $\sigma = \sigma_{u_R}^2(s_{u_L})$).

- (k) For $u_L \in \mathcal{V}$, and for all $s > 0$, $\frac{d}{ds}\eta(u_L|S_{u_L}^1(s)) > 0$ (the shock “strengthens” with s).
Similarly, for $u_R \in \mathcal{V}$, and for all $s > 0$, $\frac{d}{ds}\eta(u_R|S_{u_R}^2(s)) > 0$. Moreover, for each $u_L, u_R \in \mathcal{V}$ and $s > 0$, $\frac{d}{ds}\sigma_{u_L}^1(s) < 0$ and $\frac{d}{ds}\sigma_{u_R}^2(s) > 0$.

These assumptions are fairly general. The first one corresponds to the strict hyperbolicity of the system in \mathcal{V} . The second one means that both characteristics families of the system are genuinely nonlinear in \mathcal{V} in the sense of Lax [31]. The third assumption is related to the second law of thermodynamics. The function η is called an entropy of the system, and q is called the entropy flux associated with η . The next two assumptions are less classical. Assumption (d) ensures a contraction property on rarefaction waves (see Section 4). Assumption (e) provides a global bound on the speeds of propagation. Assumptions (f) to (k) are now standard for the a -contraction theory. It was showed in [33] that they are verified for a large family of systems, including the Full Euler system and the isentropic Euler system.

A typical example of systems verifying the Assumptions 1.1 is the system of isentropic Euler equations for $\gamma > 1$:

$$\begin{aligned} \rho_t + (\rho v)_x &= 0, & t > 0, x \in \mathbb{R}, \\ (\rho v)_t + (\rho v^2 + \rho^\gamma)_x &= 0, & t > 0, x \in \mathbb{R}, \end{aligned} \quad (1.3)$$

endowed with the physical entropy $\eta(u) = \rho v^2/2 + \rho^\gamma/(\gamma - 1)$, where $u = (\rho, \rho v)$. For any fixed constant $C > 0$, we can define the space of states as the invariant region:

$$\mathcal{V}_0 = \{u = (\rho, \rho v) \in \mathbb{R}^+ \times \mathbb{R} : -C < w_1(u) = v - c_1 \rho^{\frac{\gamma-1}{2}} \leq w_2(u) = v + c_1 \rho^{\frac{\gamma-1}{2}} < C\}. \quad (1.4)$$

Note that $\mathcal{V}_0 = \mathcal{V} \cup \{(0, 0)\}$ where $(0, 0)$ is the vacuum state. It justifies the precise distinction of \mathcal{V} and \mathcal{V}_0 (see [33, 45]). The fact that (1.3) verifies the three first assumptions of Assumption 1.1 is well-known (see Serre [38] for instance). We prove in Section 4 that it also verifies assumptions (d) and (e).

We will consider only entropic solutions of (1.1), that is, solutions which verify additionally

$$(\eta(u))_t + (q(u))_x \leq 0, \quad t > 0, x \in \mathbb{R}. \quad (1.5)$$

More precisely, we ask that for all $\phi \in C_0^\infty([0, \infty) \times \mathbb{R})$ verifying $\phi \geq 0$,

$$\int_0^\infty \int_{-\infty}^\infty [\phi_t(t, x)\eta(u(t, x)) + \phi_x(t, x)q(u(t, x))] dx dt + \int_{-\infty}^\infty \phi(0, x)\eta(u^0(x)) dx \geq 0, \quad (1.6)$$

where $u^0 : \mathbb{R} \rightarrow \mathbb{R}$ is the prescribed initial data for the solution u .

We also restrict our study to the solutions verifying the so-called Strong Trace Property.

Definition 1.2 (Strong Trace Property). *Let $u \in L^\infty(\mathbb{R}^+ \times \mathbb{R})$. We say that u verifies the strong trace property if for any Lipschitzian curve $t \rightarrow X(t)$, there exists two bounded functions $u_-, u_+ \in L^\infty(\mathbb{R}^+)$ such that for any $T > 0$*

$$\lim_{n \rightarrow \infty} \int_0^T \sup_{y \in (0, 1/n)} |u(t, X(t) + y) - u_+(t)| dt = \lim_{n \rightarrow \infty} \int_0^T \sup_{y \in (-1/n, 0)} |u(t, X(t) + y) - u_-(t)| dt = 0.$$

For convenience, we will use later the notation $u_+(t) = u(t, X(t)+)$, and $u_-(t) = u(t, X(t)-)$. We can then define the wildest space of solutions that we consider in the paper:

$$\mathcal{S}_{weak} = \{u \in L^\infty(\mathbb{R}^+ \times \mathbb{R} : \mathcal{V}_0) \text{ weak solution to (1.1)(1.5), verifying Definition 1.2}\}. \quad (1.7)$$

Note that this space has no smallness condition.

The aim of this paper is to show the stability of a smaller class of solutions -namely solutions with small BV norms- when perturbations are taken in the wider space \mathcal{S}_{weak} . More precisely, for any domain \mathcal{O} such that $\overline{\mathcal{O}} \subset \mathcal{V}$, consider the following class of solutions:

$$\mathcal{S}_{BV, \varepsilon} = \{u \in L^\infty(\mathbb{R}^+, BV(\mathbb{R} : \mathcal{O})) \text{ solution to (1.1)(1.5), with } \|u(t)\|_{BV(\mathbb{R})} \leq \varepsilon \text{ for } t \geq 0\}. \quad (1.8)$$

Our main result is the following theorem.

Theorem 1.3. *Consider a system (1.1) verifying all the Assumptions 1.1. Then, for any open set \mathcal{O} such that $\overline{\mathcal{O}} \subset \mathcal{V}$, there exists $\varepsilon > 0$ such that the following is true.*

Let $u \in \mathcal{S}_{BV, \varepsilon}$ be a BV solution with initial value u^0 . Assume that $u_n \in \mathcal{S}_{weak}$ is a sequence of wild solutions, uniformly bounded in $L^\infty(\mathbb{R}^+ \times \mathbb{R})$, with initial values $u_n^0 \in L^\infty(\mathbb{R})$. If u_n^0 converges to u^0 in $L^2(\mathbb{R})$, then for every $T > 0, R > 0$, u_n converges to u in $L^\infty(0, T; L^2(-R, R))$. Especially, u is unique in the class \mathcal{S}_{weak} .

In a celebrated paper [20], Glimm showed that for any \mathcal{O} compact subset of \mathcal{V} , any $a \in \mathcal{O}$, and ε small enough, there exists $\varepsilon_{in} > 0$ such that if $\|u^0 - a\|_{L^\infty(\mathbb{R})} \leq \varepsilon_{in}$ and $\|u^0\|_{BV(\mathbb{R})} \leq \varepsilon_{in}$, then there exists a solution $u \in \mathcal{S}_{BV, \varepsilon}$ of (1.1) with $u(0, \cdot) = u^0$. After Glimm's method, currently referred to

as the Glimm scheme or random choice method, there are two other frameworks which can be used to prove the small BV existence for general hyperbolic conservation laws: the front tracking scheme (see [3, 12]) and the vanishing viscosity method [2].

Uniqueness of these solutions was established by Bressan and Goatin [5] under the Tame Oscillation Condition. It improved an earlier theorem by Bressan and LeFloch [6]. Uniqueness was also known to prevail when the Tame Oscillation Condition is replaced by the assumption that the trace of solutions along space-like curves has bounded variation, see Bressan and Lewicka [7]. We will refer to this condition as the Bounded Variation Condition (see Definition 2.2). One can also find these theories in [3] or [12].

These uniqueness theories, which work for general hyperbolic conservation laws with n unknowns, all need some a priori assumption on the solutions, such as Tame Oscillation Condition or Bounded Variation Condition on space-like curves (see [3]).

Note that any BV function verifies the Strong Trace Property (Definition 1.2). Hence, any BV solution to (1.1)(1.5) belongs to \mathcal{S}_{weak} . Therefore a consequence of Theorem 1.3, is that in the case of 2 unknowns, these a priori assumptions are not needed to obtain the uniqueness result. We formulate this result in the following theorem.

Theorem 1.4. *Consider a system (1.1) verifying all the Assumptions 1.1. Then, for any open set \mathcal{O} such that $\overline{\mathcal{O}} \subset \mathcal{V}$, there exists $\varepsilon > 0$ such that the following is true.*

Any solution in $\mathcal{S}_{BV,\varepsilon}$ with initial value u^0 is unique among the functions

$$\{v \in L^\infty([0, T]; BV(\mathbb{R})) \text{ for all } T > 0 \text{ and solution to (1.1)(1.5)}\}$$

with the same initial value.

As another celebrated work for small BV solutions of general systems of hyperbolic conservation laws, the L^1 stability has been established in 1990s [4, 8], or see [3]. In the L^1 stability theory, the perturbations u_n have to stay in the space $\mathcal{S}_{BV,\varepsilon}$. In contrast, this is not required in our L^2 based theory. For this reason, Theorem 1.3 can be seen as a weak-BV stability result, similar to the weak-Strong stability result of Dafermos and DiPerna.

Indeed, since the work of Dafermos and DiPerna [11, 16], it is known that on any span of time $[0, T]$ where a solution of the system is Lipschitz in x , the solution is L^2 stable (for L^2 perturbations on the initial value) among the large class of solutions which are bounded weak entropic solutions to the same system. This implies the well known weak-strong uniqueness principle: as long as a solution is Lipschitz, it is unique among any other bounded weak solution. To be more precise, let us denote the two classes of solutions:

$$\begin{aligned} \mathcal{S}_{reg}^T &= \{u \in L^\infty([0, T] \times \mathbb{R} : \mathcal{V}), \text{ solution to (1.1), with } \|\partial_x u(t)\|_{L^\infty([0, T] \times \mathbb{R})} \leq C, \text{ for } C > 0\} \\ \mathcal{S}_{weak}^T &= \{u \in L^\infty([0, T] \times \mathbb{R} : \mathcal{V}_0) \text{ weak solution to (1.1)(1.5)}\}. \end{aligned}$$

Let \mathcal{O} be a compact subset of \mathcal{V} , and u be a solution in \mathcal{S}_{reg}^T with values in \mathcal{O} , and with initial value u^0 . The result of Dafermos and DiPerna implies that if $(u_n)_{n \in \mathbb{N}}$ is a sequence of solutions in \mathcal{S}_{weak}^T such that their initial values $(u_n^0)_{n \in \mathbb{N}}$ converge in $[L^2(\mathbb{R})]^2$ to u^0 , then $(u_n)_{n \in \mathbb{N}}$ converges in $L^\infty(0, T; L^2(\mathbb{R}))$ to u . Especially, it implies the uniqueness of solutions in \mathcal{S}_{reg}^T among the bigger class of solutions \mathcal{S}_{weak}^T (weak/Strong uniqueness).

Theorem 1.3 extends this result, in the context of 1-d systems with two unknowns, going from the Lipschitz space \mathcal{S}_{reg}^T to the BV space $\mathcal{S}_{BV,\varepsilon}$. Note, however, that the wild solutions of \mathcal{S}_{weak} need to have the extra strong trace property compared to solutions of \mathcal{S}_{weak}^T . Still, they can be valued in \mathcal{V}_0 , including states for which f is not differentiable, like the vacuum for the Euler system (1.3).

Our work is motivated by the construction of bounded solutions via compensated compactness available for 2×2 systems (see Tartar [42], DiPerna [17], Chen [9, 15], Lions-Perthame and etc [34, 35]). Those systems have Riemann invariants (see Smoller's book [41] and Hoff's paper [22]) w_1 and w_2 , such that

$$(w_1)_t + \lambda_1(w_1)_x = 0, \quad (w_2)_t + \lambda_2(w_2)_x = 0.$$

This provides naturally invariant regions as (1.4).

Strong traces properties were first proved for multivariable conservation laws [44], see also ([30, 36]). The technique was later used to get more structural information on the solutions (see [13, 40]). For systems, the question whether bounded weak solutions in \mathcal{S}_{weak}^T verify the Strong Trace Property is mostly open.

The Euler system (1.3) with $\gamma = 3$ is an interesting case. Indeed, it was proved in [43] that for any initial values in \mathcal{V}_0 , one can construct global solutions with values in \mathcal{V}_0 verifying a similar strong property in time. It would be interesting to investigate whether this property can be extended to the property of Definition 1.2 in this context.

In a parallel program, it has been shown that, considering inviscid limit of Navier-Stokes equation, instead of weak solution to the inviscid conservation laws, one can avoid the need of the strong traces property. The case of the inviscid limit of the barotropic Navier-Stokes equation in the Lagrangian variables is considered in [23, 24]. It is shown that single shocks are stable (and so unique) in the class of inviscid weak limit of energy bounded solutions to Navier-Stokes equations. Neither Boundedness of the function, nor the strong traces property are needed in this context. This result is a first milestone in the program of the authors to show the convergence from Navier-Stokes to Euler for initial values small in BV, a major open problem in the field [3]. Theorem 1.3 is a second major milestone in this direction. It provides several tools needed in the program to leap from the study of a single wave solution to general Cauchy data.

The paper is structured as follows. We begin in Section 2 with preliminaries linked to the Bounded Variation Condition along space-like curve needed for the L^1 uniqueness theory. The proof of our result is based on the weighted entropy method with shifts, and the front tracking method. The main ideas of our proof are presented in Section 3. This section proves Theorem 1.3 from Proposition 3.2. The rest of the paper is dedicated to the proof of Proposition 3.2. Section 4 is dedicated to the L^2 study of single waves. The most important one concerns the study of a single shock. The exact needed version, Proposition 4.1, is proved in a companion paper [21]. The modified front tracking algorithm is introduced in Section 5. The construction of the weight functions are performed in Section 6. Finally, Section 7 is dedicated to the proof of Proposition 3.2.

2. PRELIMINARIES

This section gather tools from the L^1 theory that will be useful later. Every result and notion for this section comes from [3]. Our proof uses the L^1 uniqueness result of [7]. Let us write precisely the statement here. Following [3], we introduce first the notion of space-like curve.

Definition 2.1 (Space-like curves). *Let $\hat{\lambda}$ be a fixed constant. Then we define a space-like curve to be a curve of the form $\{t = \gamma(x) : x \in (a, b)\}$, with*

$$|\gamma(x_2) - \gamma(x_1)| < \frac{x_2 - x_1}{\hat{\lambda}} \quad \text{for all } a < x_1 < x_2 < b. \quad (2.1)$$

In this paper, the value of $\hat{\lambda}$ will be determined by Proposition 4.1. Still following [3], we now introduce the extra condition needed for the classical L^1 uniqueness theorem.

Definition 2.2 (Bounded Variation Condition). *We say that a function $u \in L^\infty(\mathbb{R}^+ \times \mathbb{R})$ verifies the Bounded Variation Condition if there exists $\delta > 0$ such that, for every bounded space-like curve $\{t = \gamma(x) : x \in [a', b']\}$ with*

$$|\gamma(x_1) - \gamma(x_2)| \leq \delta |x_1 - x_2| \quad \text{for all } x_1, x_2 \in [a', b'], \quad (2.2)$$

the function $x \mapsto u(\gamma(x), x) := u_\gamma(x)$ is well defined and has bounded variation.

Note that taking constant functions γ shows that these functions u are BV in x . Let us now state a uniqueness result of [3, 7], rephrased in our context.

Theorem 2.3 (From [3, 7]). *For any $d \in \mathcal{V}$, there exists $\varepsilon > 0$ such that for any u^0 initial value with $\|u^0\|_{BV(\mathbb{R})} \leq \varepsilon$ and $\|u^0 - d\|_{L^\infty(\mathbb{R})} \leq \varepsilon$, we have the following uniqueness result.*

There exists only one solution u of (1.1)–(1.5) with initial value u^0 and verifying the Bounded Variation Condition of Definition 2.2.

Note that Theorem 1.4 replaces the condition of Definition 2.2, by only $u \in L^\infty(\mathbb{R}^+; BV(\mathbb{R}))$, and Theorem 1.3 by $u \in \mathcal{S}_{weak}$.

We will need to prove that a certain limit of solutions to a modified front tracking algorithm inherits the Bounded Variation Condition. Still following [3], we introduce the following domination principle.

Definition 2.4 (Domination). *Given two space-like curves $\gamma : (a, b) \rightarrow \mathbb{R}$ and $\gamma' : (a', b') \rightarrow \mathbb{R}$, we say that γ dominates γ' if $a \leq a' < b' \leq b$ and, moreover,*

$$\gamma(x) \leq \gamma'(x) \leq \min \left\{ \gamma(a) + \frac{x-a}{\hat{\lambda}}, \gamma(b) + \frac{b-x}{\hat{\lambda}} \right\} \quad \text{for all } x \in (a', b'). \quad (2.3)$$

This property implies that γ' is entirely contained in a domain of determinacy for the curve γ . We introduce now the following property.

Condition 2.5. *Let $C > 0$. Let a function $\psi \in L^\infty(\mathbb{R}^+; BV(\mathbb{R}))$ be piecewise constant. We say that it verifies the Condition 2.5 with constant C , if it verifies the following.*

Let γ and γ' be any two space-like curves with γ dominating γ' (Definition 2.4). Then,

$$\text{Tot.Var.}\{\psi; \gamma'\} \leq C \text{Tot.Var.}\{\psi; \gamma\}. \quad (2.4)$$

We will use the following lemma.

Lemma 2.6. *Let $\{\psi_n\}_{n \in \mathbb{N}}$ be a family of piecewise constant functions uniformly bounded in $L^\infty(\mathbb{R}^+, BV(\mathbb{R}))$. Assume that there exists $C > 0$ such that for every $n \in \mathbb{N}$, ψ_n verifies Condition 2.5 for this constant C , and*

$$\|\psi_n(t, \cdot) - \psi_n(s, \cdot)\|_{L^1} \leq C|t - s|, \quad 0 < s < t < T. \quad (2.5)$$

Then, there exists $\psi \in L^\infty(\mathbb{R}^+ \times \mathbb{R})$ verifying the Bounded Variation Condition 2.2 such that, up to a subsequence, ψ_n converges to ψ when $n \rightarrow \infty$ in $C^0(0, T; L^2(-R, R))$ for every $T > 0$, $R > 0$, and almost everywhere in $\mathbb{R}^+ \times \mathbb{R}$.

This lemma is very similar to [3, Lemma 7.3], where the same result is stated for classical piecewise constant approximate solutions constructed by the front tracking algorithm (without shifts). For the sake of completeness, we provide a proof of it in the appendix.

3. WEIGHTED RELATIVE ENTROPY AND SHIFTS

The proof of our result is based on the relative entropy method first introduced by Dafermos [11] and DiPerna [16]. From the assumption of the existence of a convex entropy η , we define an associated pseudo-distance defined for any $a, b \in \mathcal{V}_0 \times \mathcal{V}$:

$$\eta(a|b) = \eta(a) - \eta(b) - \nabla \eta(b)(a - b). \quad (3.1)$$

The quantity $\eta(a|b)$ is called the relative entropy of a with respect to b , and is equivalent to $|a - b|^2$. We also define the relative entropy-flux: For $a, b \in \mathbb{R}^2$,

$$q(a; b) = q(a) - q(b) - \nabla \eta(b)(f(a) - f(b)). \quad (3.2)$$

The strength of this notion is that if u is a weak solution of (1.1), (1.5), then u verifies also the full family of entropy inequalities for any $b \in \mathcal{V}$ constant:

$$(\eta(u|b))_t + (q(u; b))_x \leq 0. \quad (3.3)$$

Similar to the Kruzkov theory for scalar conservation laws, (3.3) provides a full family of entropies measuring the distance of the solution to any fixed values b in \mathcal{V} . The main difference is that the distance is equivalent to the square of the L^2 norm rather than the L^1 norm. Same as for the Kruzkov theory, (3.3) provides directly the stability of constant solutions (by integrating in x the inequality). Modulating the inequality with a smooth function $t, x \rightarrow b(t, x)$ provides the well-known weak-strong uniqueness result. Precisely, the relative entropy is an L^2 theory in the following sense:

Lemma 3.1. *For any fixed compact set $V \subset \mathcal{V}$, there exists $c^*, c^{**} > 0$ such that for all $(u, v) \in \mathcal{V}_0 \times V$,*

$$c^*|u - v|^2 \leq \eta(u|v) \leq c^{**}|u - v|^2. \quad (3.4)$$

The constants c^, c^{**} depend on bounds on the second derivative of η in V , and on the continuity of η on \mathcal{V}_0 .*

This elementary lemma follows directly from Taylor's theorem (see [33, 45]).

For the family of Euler systems, it is well known that the relative entropy provides a contraction property for rarefaction function $t, x \rightarrow b(t, x)$, even in multi-D [19]. This is because it verifies Assumption 1.1 (d) (see Section 4).

However, when modulating the inequality with discontinuous functions b with shocks, the situation diverges significantly from the Kruzkov situation. This is due to the fact that the L^2 norm is not as well suited as the L^1 norm for the study of stability of shocks. However, the method was used by DiPerna [16] to show the uniqueness of single shocks (see also Chen Frid [10] for the Riemann problem of the Euler equation). In [45], it was proposed to use the method to obtain stability of discontinuous solutions. The main idea was that the L^2 norm can capture very well the stability of the profile of the shock (up to a shift), even if the shift itself is more sensitive [33]. Leger in [32] showed that in the scalar settings, the shock profiles (modulo shifts) have a contraction property in L^2 , reminiscent to the L^1 contraction of the Kruzkov theory. It was shown in [39] that the contraction property is usually false for systems. However, it can be recovered by weighting the relative entropy [25]. More precisely, consider a fixed shock (u_l, u_r, s) . It was shown that there exists $0 < a_1 < a_2$ such that, for any wild solutions $u \in \mathcal{S}_{weak}$, we can construct a Lipschitz shift function $h : \mathbb{R}^+ \rightarrow \mathbb{R}$ with

$$\frac{d}{dt} \left\{ a_1 \int_{-\infty}^{h(t)} \eta(u(t, x)|u_L) dx + a_2 \int_{h(t)}^{\infty} \eta(u(t, x)|u_R) dx \right\} \leq 0. \quad (3.5)$$

Note that this formula for $a_1 = a_2$, and $h(t) = st$ would imply the contraction property of the shock for the relative entropy. But the result, to be valid, needs the weights a_i , and the shifts h , giving the name to the method: a-contraction with shifts.

Let us emphasize that the L^2 based a-contraction is not true without the notion of shifts. This is a major obstruction to consider solutions with several waves. Conservation laws have finite speeds of propagation. Therefore, usually, considering a finite amount of waves is equivalent to studying a single one, at least, as long as they do not interact. Because of the shifts, it is not obvious anymore in this theory. The general idea, is that one shift by singularity is needed. Those shifts depend

crucially on the perturbation. It is therefore needed to prevent that this artificial shifts do not force a 1-shock to stick and holds to a 2-shock, making the whole process to collapse. This problem was solved in [27], allowing the treatment of the Riemann problem. The main idea is that the shifts can be constructed based on perturbed characteristic curves associated to the wild solution.

This article is making the leap going from the stability of the Riemann problem, to the stability of small BV solutions. Because of the generation of infinitely many shifts, the estimate (3.5) is significantly weakened in this case. Our main proposition is the following.

Proposition 3.2. *Consider a system (1.1) verifying all the Assumptions 1.1. Let $d \in \mathcal{V}$. Then there exist $C, v, \varepsilon > 0$ such that the following is true.*

For any $m > 0$, $R, T > 0$. $u^0 \in BV(\mathbb{R})$ such that $\|u^0\|_{BV(\mathbb{R})} \leq \varepsilon$ and $\|u^0 - d\|_{L^\infty(\mathbb{R})} \leq \varepsilon$, and any wild solution $u \in \mathcal{S}_{weak}$, there exists $\psi: \mathbb{R}^+ \times \mathbb{R} \rightarrow \mathcal{V}$ such that for almost every $0 < s < t < T$:

$$\begin{aligned} \|\psi(t, \cdot)\|_{BV(\mathbb{R})} &\leq C\|u^0\|_{BV(\mathbb{R})}, \\ \|\psi(t, \cdot) - \psi(s, \cdot)\|_{L^1} &\leq C|t - s|, \\ \|\psi(t, \cdot) - u(t, \cdot)\|_{L^2((-R+vt, R-vt))} &\leq C \left(\|u^0 - u(0, \cdot)\|_{L^2(-R, R)} + \frac{1}{m} \right), \end{aligned}$$

The function ψ verifies the Condition 2.5 with constant C .

It would be natural to try to take for the function ψ , the unique BV solution with initial value u^0 of Theorem 2.3. However, functions ψ which verify the proposition are not solutions to (1.1). Instead, the proposition shows that if the initial value $u(0, \cdot)$ is L^2 close to a set of small BV functions, then $u(t, \cdot)$ stays L^2 close, for every time $t > 0$, to a slightly bigger set of small BV functions.

Despite the finite speed propagation of the equation, there are two major difficulties to obtain this result: one is the shifts, the other is the weights. Let us give an example of the difficulties the shifts introduce. Consider a piecewise-constant solution $\bar{u} \in \mathcal{S}_{BV, \varepsilon}$ to (1.1) (1.5). Until the first time that there is an interaction between the shocks in \bar{u} , we can represent the function \bar{u} as

$$\bar{u}(t, x) = u_i, \quad \text{for } x_i + s_i t < x < x_{i+1} + s_{i+1} t,$$

where (u_{i-1}, u_i, s_i) are admissible shocks. For any weak solution $u \in \mathcal{S}_{weak}$, the general theory of weighted relative entropy with shifts ensure the existence of shifts $t \rightarrow h_i(t)$ and a piecewise weighted function

$$a(t, x) = a_i, \quad \text{for } h_{i-1}(t) \leq x \leq h_i(t),$$

such that, as long as the functions $h_i(t)$ do not cross:

$$\frac{d}{dt} \int_{\mathbb{R}} a(t, x) \eta(u(t, x) | \psi(t, x)) dx \leq 0, \quad (3.6)$$

where

$$\psi(t, x) = u_i \quad \text{for } h_{i-1}(t) \leq x_{i+1} \leq h_i(t).$$

Let $t^* > 0$ be such that there are no collisions between any of the wavefronts in either \bar{u} or ψ for $t \in [0, t^*]$. One might then hope to control $\|\bar{u}(t, \cdot) - u(t, \cdot)\|_{L^2}$ by using both (3.6) and by controlling $\|\bar{u}(t, \cdot) - \psi(t, \cdot)\|_{L^2}$. However, it is difficult to control $\|\bar{u}(t, \cdot) - \psi(t, \cdot)\|_{L^2}$. For time $t \in [0, t^*]$, the function ψ can be reconstructed from the function \bar{u} via a change of variables. However, if for example after time t^* a collision between two waves occurs in \bar{u} , but the corresponding waves do not collide in ψ , then after the local Riemann problem in \bar{u} is solved and the clock restarted, the functions \bar{u} and ψ cannot be related through a change of variables. Furthermore, as in the scalar case, the best control we have on the shifts gets worse and worse as the strength of the shock (being controlled by the shift) decreases (see [26, Theorem 1.1] and [29, Theorem 1.2]). This is problematic, because we want the initial data of the function ψ to approach the initial data for the wild solution u , and in general the sizes of the shocks in ψ will get arbitrarily small.

The idea then is to give up on trying to control $\bar{u} - \psi$ or the shifts. Instead, we construct an artificial function ψ which stays L^2 close to u while sharing the structural property of \bar{u} (the smallness in BV). If we now consider a sequence of such solutions u_n such that the initial values converge to the initial value of \bar{u} , we can transfer, at the limit, the structural property from the ψ_n functions to the limit u . This implies that u belongs to $\mathcal{S}_{BV,\varepsilon}$ and still verifies the Bounded Variation Condition, and so is equal to \bar{u} by the uniqueness theorem (Theorem 2.3).

Note that this strategy was first introduced in [28] in the scalar case with convex fluxes. The paper [28] gives a new proof for the uniqueness of solutions verifying a single entropy condition. Previous proofs of this result were obtained by Panov [37] and De Lellis, Otto, and Westdickenberg [14]. Their proofs were based on the link between conservation laws and Hamilton-Jacobi equations, and it seems difficult to extend them to the system case where no such relation exists.

The strategy is now to construct the function ψ of Proposition 3.2 which stays L^2 close to u , while enjoying the small BV property of \bar{u} . We construct it via the front tracking method, from the initial value of \bar{u}^0 , but with the propagation of discontinuities following the shifts $\{h_i\}$ (which depend on the weak solution u). A key point is that the BV estimates obtained from the front tracking method do not depend on the propagation of these fronts. We can then control the BV norm of ψ . Note that ψ is not a solution to the equation (1.1) since the Rankin-Hugoniot conditions are not verified anymore. It cannot be easily compared to \bar{u} either since the waves can interact in a different order. We remark that although we limit ourselves in this paper to 2×2 systems with genuinely nonlinear wave families, the front tracking algorithm we use works for general $n \times n$ systems with either genuinely nonlinear or linearly degenerate wave families. In particular, in this paper, we have incorporated non-physical waves (also known as pseudoshocks) into our modified front tracking algorithm. The use of non-physical waves is not required for the 2×2 case, but it is necessary for applying the front tracking method to general $n \times n$ systems.

The last difficulty is due to the weight function a . In order to obtain the contraction property (3.6), we have constraints on the variations of the weights $a_i - a_{i-1}$ which depend both on the size and the family of the shock (u_{i-1}, u_i, s_i) . This means that the weight function has to be reconstructed at each collision time between two waves. The variations of the weight function has to be controlled precisely to ensure that it stays bounded away from 0. The construction of the weight is closely related to the front tracking method, and the control of its BV norm mirrors the BV control on the function ψ itself.

For this procedure, a key refinement of the a-contraction for a single shock is provided by the companion paper [21]. It shows that the size of variation of the weight $|a_i - a_{i-1}|$ can be chosen proportionally to the strength of the associated shock wave $|\sigma_i| \approx |u_{i-1} - u_i|$. This property was first showed in the class of inviscid limit of Navier-Stokes [23, 24]. Surprisingly, the proof based directly on the inviscid model is very different, and quite delicate.

We finish this section showing how Proposition 3.2 implies Theorem 1.3.

Proof of Main Theorem (Theorem 1.3). For each $d \in \overline{\mathcal{O}}$, consider $\varepsilon_d > 0$ such that both Proposition 3.2 and Theorem 2.3 are valid. The union (over d) of the balls $B_{\varepsilon_d/2}(d)$ cover the compact $\overline{\mathcal{O}}$, so there exists a finite subcover. Denote $\varepsilon > 0$ the smallest of the $\varepsilon_{d_i}/2$ for this finite subcover.

By passing to a subsequence if necessary, we assume that $\|u_m^0 - u^0\|_{L^2} \leq \frac{1}{m}$. From Proposition 3.2 we have a sequence of functions ψ_m (for all $m \in \mathbb{N}$), uniformly bounded in $L^\infty(\mathbb{R}^+, BV(\mathbb{R}))$. Moreover ψ_m verify Condition 2.5 and (2.5) uniformly, and they verify for all time $t > 0$:

$$\|\psi_m(t, \cdot) - u_m(t, \cdot)\|_{L^2(\mathbb{R})} \leq \frac{2}{m}. \quad (3.7)$$

From Lemma 2.6, there exists $\psi \in L^\infty(\mathbb{R}^+ \times \mathbb{R})$ verifying the Bounded Variation Condition (Definition 2.2) such that for every $T > 0, R > 0$, ψ_m converges in $C^0(0, T : L^2(-R, R))$ to ψ . Together with (3.7), u_m converges in $L^\infty(0, T : L^2(-R, R))$ to ψ . Since the convergence is strong and u_m verifies (1.1) (1.5), the limit ψ is also solution to (1.1) (1.5), with initial value u^0 . From Theorem 2.3, it is the unique solution verifying Definition 2.2.

Applying the result to the constant sequence $\bar{u}_n = u$, the fixed BV function with initial value u^0 from the hypotheses of the theorem, shows that u is also this unique solution. Therefore $\psi = u$. This ends the proof of Theorem 1.3. The rest of the paper is dedicated to the proof of Proposition 3.2.

4. RELATIVE ENTROPY FOR THE RIEMANN PROBLEM

We first state the refined a-contraction property of shocks for the weighted relative entropy with shifts. This result is proved in [21]. Note that the constant L is defined in Assumption 1.1 (e).

Proposition 4.1. *Consider a system (1.1) verifying all the Assumptions 1.1. Let $d \in \mathcal{V}$. Then there exist constants $\alpha_1, \alpha_2, \hat{\lambda}$ and $C, \varepsilon > 0$, with $\alpha_1 < \alpha_2$ and $\hat{\lambda} \geq 2L$, such that the following is true.*

Consider any shock (u_L, u_R) with $|u_L - d| + |u_R - d| \leq \varepsilon$, any $u \in \mathcal{S}_{weak}$, any $\bar{t} \in [0, \infty)$, and any $x_0 \in \mathbb{R}$. Let σ be the strength of the shock $\sigma = |u_L - u_R|$. Then for any $a_1 > 0, a_2 > 0$ verifying

$$1 - 2C\sigma \leq \frac{a_2}{a_1} \leq 1 - \frac{C\sigma}{2} \quad \text{if } (u_L, u_R) \text{ is a 1-shock}$$

$$1 + \frac{C\sigma}{2} \leq \frac{a_2}{a_1} \leq 1 + 2C\sigma \quad \text{if } (u_L, u_R) \text{ is a 2-shock,}$$

there exists a Lipschitz shift function $h : [\bar{t}, \infty) \rightarrow \mathbb{R}$, with $h(\bar{t}) = x_0$, such that the following dissipation functional verifies

$$\begin{aligned} & a_1 \left[q(u(t, h(t)+); u_R) - \dot{h}(t) \eta(u(t, h(t)+)|u_R) \right] \\ & - a_2 \left[q(u(t, h(t)-); u_L) - \dot{h}(t) \eta(u(t, h(t)-)|u_L) \right] \leq 0 \end{aligned} \quad (4.1)$$

for almost all $t \in [\bar{t}, \infty)$.

Moreover, if (u_L, u_R) is a 1-shock, then for almost all $t \in [\bar{t}, \infty)$:

$$-\frac{\hat{\lambda}}{2} \leq \dot{h}(t) \leq \alpha_1 < \inf_{v \in B_{2\varepsilon}(d)} \lambda_2(v).$$

Similarly, if (u_L, u_R) is a 2-shock, then for almost all $t \in [\bar{t}, \infty)$:

$$\sup_{v \in B_{2\varepsilon}(d)} \lambda_1(v) < \alpha_2 \leq \dot{h}(t) \leq \frac{\hat{\lambda}}{2}.$$

Integrating (3.3) with $b = u_L$ for $x \in (-\infty, h(t))$, integrating (3.3) with $b = u_R$ for $x \in (h(t), \infty)$, summing the results, and using (4.1) together with the strong traces property Definition 1.2 provides the contraction property (3.6) in the case of a single shock as long as,

$$a_2/a_1 \text{ is between } 1 + \frac{C}{2}(-1)^i\sigma \text{ and } 1 + 2C(-1)^i\sigma,$$

when (u_L, u_R) is a i -shock. It shows that the variation of the a function has to be negative for a 1-shock, positive for a 2-shock, and can be chosen with strength of the same order as the size of the shock. The estimates on \dot{h} show that we keep a finite speed of propagation, and that a shift of a 1-shock cannot overtake the shift of a 2-shock if it started on its left. This is important because when we introduce shifts into the solution to a Riemann problem with two shocks, both shock speeds move with artificial velocities. We need to ensure that the positions of the shocks do not touch at some

time after the initial time to preserve the property of classical solutions to the Riemann problem, where shocks born from a solution to a Riemann problem will never touch.

We need a similar control for approximations of rarefactions via the front tracking method. We begin to show that, under the Assumption 1.1 (b)(d), the real rarefaction has a contraction property without the need of shift.

Lemma 4.2. *Consider a system (1.1) verifying all the Assumptions 1.1. Let $\bar{u}(y)$ $v_L \leq y \leq v_R$ be a rarefaction wave for (1.1). Then for any $u \in \mathcal{S}_{weak}$ and every $t > 0$ we have*

$$\begin{aligned} \frac{d}{dt} \int_{v_L t}^{v_R t} \eta(u(t, x) | \bar{u}(x/t)) dx &\leq q(u(t, v_L t+); \bar{u}(v_L)) - q(u(t, v_R t-); \bar{u}(v_R)) \\ &\quad - v_L \eta(u(t, v_L t+); \bar{u}(v_L)) + v_R \eta(u(t, v_R t-); \bar{u}(v_R)). \end{aligned}$$

Remark 4.3. *We generalize the result and proof known for Euler. For one possible reference for this, see [45].*

Proof. Following [11] or [45] we have that for any $u \in \mathcal{S}_{weak}$ and any $v \in \mathcal{S}_{reg}^T$:

$$\partial_t \eta(u|v) + \partial_x q(u; v) + \partial_x \{\eta'(v)\} f(u|v) \leq 0.$$

The relative flux is defined analogously to the relative entropy. For $a, b \in \mathcal{V}_0 \times \mathcal{V}$, we define it as:

$$f(a|b) = f(a) - f(b) - \nabla f(b)(a - b).$$

Since \bar{u} is a rarefaction, for all y , $\bar{u}'(y)$ is a right eigenvector of $f'(\bar{u}(y))$, and so from Assumption 1.1 (b): $\bar{u}'(y) \cdot \lambda'(\bar{u}(y)) > 0$. For $v(t, x) = \bar{u}(x/t)$, $\partial_x v = \frac{1}{t} \bar{u}'(x/t) = r$ is also a right eigenvector of $f'(\bar{u}(y)) = f'(v)$ with the same property from Assumption 1.1 (b):

$$r \cdot \lambda'(v) > 0. \tag{4.2}$$

Denote $\ell = \partial_x \{\eta'(v)\} = \eta''(v) \partial_x v = \eta''(v) r$. For v fixed, since η'' is positive

$$\ell \cdot r = r^T \eta''(v) r > 0. \tag{4.3}$$

Since $\eta''(v) f'(v)$ is symmetric

$$\ell^T f'(v) = r^T \eta''(v) f'(v) = \eta''(v) f'(v) r = \lambda(v) \eta''(v) r = \lambda(v) \ell.$$

Hence ℓ is a left eigenvector of $f'(v)$. From Assumption 1.1 (d), we have that $\ell \cdot f$ is either convex or concave. Let $r(u)$ be a right eigenvector of $f'(u)$ for the same family. Using that ℓ is a left eigenvector of $f'(v)$, we get

$$\ell^T (f'(u) - f'(v)) r(u) = \ell \cdot r(u) (\lambda(u) - \lambda(v)).$$

passing to the limit u goes to v , and taking the value along $r = r(v)$ we find:

$$\ell f''(v)(r, r) = (\lambda'(v) \cdot r)(\ell \cdot r) > 0$$

thanks to (4.2) and (4.3). Hence, $\ell \cdot f$ is convex, and

$$u \rightarrow \partial_x \{\eta'(v)\} f(u|v)$$

is non-negative for all $u \in \mathcal{V}$. Finally

$$\partial_t \eta(u|v) + \partial_x q(u; v) \leq 0.$$

Integrating in x between $v_L t$ and $v_R t$, and using the Strong Trace Property (Definition 1.2) gives the result. \square

We can now give the control needed for the error due to the approximation of the rarefaction via the front tracking method.

Proposition 4.4. *There exists a constant $C > 0$ such that the following is true. For any $\bar{u}(y)$ $v_L \leq y \leq v_R$ rarefaction wave for (1.1), denote*

$$\delta = |v_L - v_R| + \sup_{y \in [v_L, v_R]} |u_L - \bar{u}(y)|, \quad \bar{u}(v_L) = u_L, \quad \bar{u}(v_R) = u_R.$$

Then for any $u \in \mathcal{S}_{weak}$, any $v_L \leq v \leq v_R$, and any $t > 0$ we have:

$$\int_0^t \left\{ q(u(t, tv+); u_R) - q(u(t, tv-); u_L) - v (\eta(u(t, tv+)|u_R) - \eta(u(t, tv-)|u_L)) \right\} dt \leq C\delta |u_L - u_R|t.$$

Proof. Consider the quantity

$$D = \frac{d}{dt} \left\{ \int_{v_L t}^{v t} \eta(u(t, x)|u_L) dx + \int_{v t}^{v_R t} \eta(u(t, x)|u_R) dx - \int_{v_L t}^{v_R t} \eta(u(t, x)|\bar{u}(x/t)) dx \right\}.$$

The dissipation due to the shocks of u cancels out. Therefore, using twice (3.3) with equality, (once with $b = u_L$, and once with $b = u_R$), and the inequality of Lemma 4.2, we find:

$$D \geq q(u(t, tv+); u_R) - q(u(t, tv-); u_L) - v (\eta(u(t, tv+)|u_R) - \eta(u(t, tv-)|u_L)).$$

Integrating in time gives that

$$\begin{aligned} & \int_0^t \left\{ q(u(t, tv+); u_R) - q(u(t, tv-); u_L) - v (\eta(u(t, tv+)|u_R) - \eta(u(t, tv-)|u_L)) \right\} dt \\ & \leq \int_0^t D dt \\ & \leq \int_{v_L t}^{v t} \eta(u(t, x)|u_L) dx + \int_{v t}^{v_R t} \eta(u(t, x)|u_R) dx - \int_{v_L t}^{v_R t} \eta(u(t, x)|\bar{u}(x/t)) dx \\ & \leq \int_{v_L t}^{v t} (\eta(\bar{u}(x/t)) - \eta(u_L) + \eta'(\bar{u}(x/t))(u - \bar{u}(x/t)) - \eta'(u_L)(u - u_L)) dx \\ & \quad + \int_{v t}^{v_R t} (\eta(\bar{u}(x/t)) - \eta(u_R) + \eta'(\bar{u}(x/t))(u - \bar{u}(x/t)) - \eta'(u_R)(u - u_R)) dx \\ & \leq C\delta |u_L - u_R|t. \end{aligned}$$

□

For the sake of completeness, we conclude this section by proving that Assumptions 1.1 are always verified for the isentropic Euler system 1.3.

Lemma 4.5. *Consider the system (1.3), with state set (1.4). Then Assumptions 1.1 are verified.*

Proof. The properties 1.1 (a), (b), (c) are well known (see [38] for instance). Denote the flux function $f(\rho, \rho v) = (\rho v, \rho v^2 + \rho^\gamma)$, and the conservative quantities $(u_1, u_2) = (\rho, \rho v)$. Then

$$f_1(u_1, u_2) = u_2, \quad f_2(u_1, u_2) = \frac{u_2^2}{u_1} + u_1^\gamma.$$

Note that f_1 is linear in u , and f_2 is convex. Hence, for any vector ℓ (left eigenvector of $f'(v)$ or not):

$$\ell \cdot f = \ell_1 f_1 + \ell_2 f_2$$

is convex if $\ell_2 \geq 0$, and concave if $\ell_2 \leq 0$. This proves 1.1 (d).

The eigenvalues of f' are given by the formula $\lambda_\pm = v \pm \sqrt{\gamma \rho^{\gamma-1}}$ (see [38]). But from (1.4), $|v| \leq C$, and $c_1 \rho^{(\gamma-1)/2} \leq 2C$. This shows 1.1 (e).

□

5. MODIFIED FRONT TRACKING ALGORITHM

In the proof of Proposition 3.2, the function ψ will be defined through a modification of the front tracking algorithm. For an excellent introduction to the front tracking algorithm, we refer the reader to Chapter 14 of Dafermos's book [12] and also the succinct paper of Baiti-Jenssen [1]. These two references, together, make an excellent introduction.

For completeness, we include here a brief description of the front tracking algorithm as we use it. In this paper, we do not make use of any of the convergence properties of the front tracking algorithm or related analysis. We use instead the fact that the algorithm gives a sequence of functions with uniformly bounded total variation.

For the construction of the ψ we are about to give, the modification to the front tracking algorithm (as presented in Baiti-Jenssen [1]) consists in changing the velocity of the shocks. The shocks move with an artificial velocity dictated by the shift functions of Proposition 4.1, instead of moving with the Rankine-Hugoniot speed. Thus, performing analysis on our version of the front tracking algorithm is nearly identical to performing analysis on the front tracking algorithm when shocks move with Rankine-Hugoniot speed.

We now give the details of the construction of ψ , following Baiti-Jenssen [1]. The main idea is to take a piecewise-constant approximation of the initial data, solve (approximately) all of the local Riemann problems within the class of piecewise-constant functions, until a time when two of the Riemann problems interact. Then, the procedure is repeated: the local Riemann problems are again solved, etc.

The key point is to show that the number of wave-fronts (i.e., curves of discontinuity in time-space) remains finite, so this inductive process does not terminate in finite time. This is done by using two different Riemann solvers: an accurate Riemann solver is used to continue the solution in time after the interaction of two wavefronts when the product of the two strengths of the wavefronts is large. With the accurate solver, the number of wave fronts in the solution might increase. On other hand, when the product of the two strengths of the colliding wavefronts is small, a simplified Riemann solver is used which will prevent an explosion in the number of wavefronts. The key is that the accurate Riemann solver will only need to be used a small number of times, keeping the number of wavefronts in our solution finite.

Recall that given a Riemann problem with two constant states u_- and u_+ sufficiently close, a solution with at most three constant states, connected by either shocks or rarefaction fans, can always be found. More precisely, there exist C^2 curves $\sigma \mapsto T_i(\sigma)(u_-)$, $i = 1, 2$, parametrized by arclength, such that

$$u_+ = T_2(\sigma_2) \circ T_1(\sigma_1)(u_-), \quad (5.1)$$

for some σ_1 and σ_2 . We define $u_0 := u_-$ and

$$u_1 := T_1(\sigma_1)(u_0), \quad (5.2)$$

$$u_2 := T_2(\sigma_2) \circ T_1(\sigma_1)(u_0). \quad (5.3)$$

We use the convention that, when σ_i is positive (negative) the states u_{i-1} and u_i are separated by an i-shock (i-rarefaction) wave. Further, the strength of the i-wave is defined as $|\sigma_i|$.

For given initial data u^0 , let ψ_ν^0 be a sequence of piecewise-constant functions approximating u^0 in L^2 on $(-R, R)$. (We will choose ν later such as to give us the required $\psi = \psi_\nu$.) Let N_ν be the number of discontinuities in the function ψ_ν and choose a parameter δ_ν controlling the maximum strength of the (approximate) rarefaction fronts.

We now introduce the two Riemann solvers. One will be used when the product of the strengths of the colliding waves is large, the other will be used when the product of the strengths is small or one of the incoming waves is non-physical (also known as a pseudoshock).

5.1. The Riemann solvers. The Riemann solvers will use *non-physical waves* (also known as *pseudoshocks*). These are waves connecting two states (let's call them u_- and u_+), and traveling

with a fixed velocity $\hat{\lambda} > 0$ defined in Proposition 4.1. Therefore, it is greater than all characteristic speeds on \mathcal{V} and greater than the speed of the shifts (which have a uniform bound on their speeds). We define this non-physical wave to have strength $|\sigma| := |u_- - u_+|$ and we say it belongs to the third wave family. Remark that since all non-physical waves travel with the same speed $\hat{\lambda}$, they cannot interact with each other.

Assume that at a positive time \bar{t} , there is an interaction at the point \bar{x} between two waves of families i_α, i_β and strengths $\sigma'_\alpha, \sigma'_\beta$, respectively, with $1 \leq i_\alpha, i_\beta \leq 3$. Let σ'_α denote the left incoming wave. Let u_-, u_+ be the Riemann problem generated by the interaction, and let σ_1, σ_2 and u_0, u_1, u_2 be defined as in (5.1). Finally, we can now define the accurate and simplified Riemann solvers.

(A) *Accurate solver*: If $\sigma_i < 0$ then we let

$$p_i := \lceil \sigma_i / \delta_\nu \rceil, \quad (5.4)$$

where $\lceil s \rceil$ denotes the smallest integer number greater than s . For $l = 1, \dots, p_i$ we define

$$u_{i,l} := T_i(l\sigma_i/p_i)(u_{i-1}), \quad x_{i,l}(t) := \bar{x} + (t - \bar{t})\lambda_i(u_{i,l}). \quad (5.5)$$

On the other hand, if $\sigma_i > 0$, we define $p_i := 1$ and

$$u_{i,l} := u_i, \quad x_{i,l}(t) := h_i(t). \quad (5.6)$$

Here, h_i is the shift function coming from Proposition 4.1. Within the context of Proposition 4.1, we take $u_L = u_{i-1}$ and $u_R = u_i$.

Then, we define the approximate solution to the Riemann problem as follows:

$$v_a(t, x) := \begin{cases} u_-, & \text{if } x < x_{1,1}(t), \\ u_+, & \text{if } x > x_{2,p_2}(t), \\ u_i, & \text{if } x_{i,p_i}(t) < x < x_{i+1,1}(t), \\ u_{i,l}, & \text{if } x_{i,l}(t) < x < x_{i,l+1}(t) \end{cases} \quad (l = 1, \dots, p_i - 1). \quad (5.7)$$

Note that thanks to the two last properties of Proposition 4.1, we have : $x_{i,p_i}(t) < x_{i+1,1}(t)$ for all $t > 0$, so the function is well defined.

(B) *Simplified solver*: for each $i = 1, 2$ let σ''_i be the sum of the strengths of the strengths of all incoming i -waves. Define

$$u' := T_2(\sigma''_2) \circ T_1(\sigma''_1)(u_-). \quad (5.8)$$

Let $v_a(t, x)$ be the approximate solution of the Riemann problem (u_-, u') given by (5.7). Remark that in general $u' \neq u_+$ and thus we are introducing a non-physical front between these states. Hence, we define the simplified solution as follows:

$$v_s(t, x) := \begin{cases} v_a(t, x), & \text{if } x - \bar{x} < \hat{\lambda}(t - \bar{t}), \\ u_+, & \text{if } x - \bar{x} > \hat{\lambda}(t - \bar{t}). \end{cases} \quad (5.9)$$

Notice that by construction, the simplified solution to the Riemann problem contains at most two physical waves and an additional non-physical wave. Thus, by strategically employing the simplified solver for small collisions, we can prevent an explosion in the number of wavefronts.

5.2. Construction of the approximate solutions. Given ν we construct the approximate solution $\psi_\nu(t, x)$ as follows. At time $t = 0$ all of the Riemann problems in ψ_ν^0 are solved accurately as in (A) (the accurate solver). By slightly perturbing the speed of a wave if necessary, we can ensure that at each time we have at most one collision, which will involve only two wavefronts. Suppose that at some time $t > 0$ there is a collision between two waves from the i_α th and i_β th families. Denote the strengths of the two waves by σ_α and σ_β , respectively. The Riemann problem generated by this interaction is solved as follows. Let ϵ_ν be a fixed small parameter which will be chosen later.

- if $|\sigma_\alpha \sigma_\beta| > \epsilon_\nu$ and the two waves are physical, then we use the accurate solver (A);

- if $|\sigma_\alpha \sigma_\beta| < \epsilon_\nu$ and the two waves are physical, or one wave is non-physical, then we use the simplified solver (B).

By the following Lemma, for any ϵ_ν this algorithm will yield an approximate solution defined for all times $t > 0$.

Lemma 5.1 (from [1, Lemma 2.1]). *The number of wavefronts in $\psi_\nu(t, x)$ is finite. Hence, the approximate solutions ψ_ν are defined for all $t > 0$.*

This Lemma is stated and proved in [1, Lemma 2.1] for piecewise constant front tracking solutions where shocks move according to Rankine-Hugoniot. We do not repeat the proof here, because using shifts in the front tracking algorithm (instead of Rankine-Hugoniot speeds) does not impact the proof. The proof is identical.

We introduce the total variation of ψ_ν as

$$L(t) = \sum |\sigma_i| = \text{TV}(\psi_\nu)(t),$$

namely the sum of the strengths of all jump discontinuities that cross the t -time line. Clearly, $L(t)$ stays constant along time intervals between consecutive collisions of fronts and changes only across points of wave interaction.

A j -wave and an i -wave, with the former crossing the t -time line to the left of the latter, are called *approaching* when either $i < j$, or $i = j$ and at least one of these waves is a shock. We recall then the definition of the potential for wave interactions

$$Q(t) = \sum_{i,j:\text{approaching waves}} |\sigma_i| |\sigma_j|,$$

where the summation runs over all paires of approaching waves, with strengths $|\sigma_i|$ and $|\sigma_j|$, which cross the t -line. Let us summarize some well known fact of the front tracking method which are still valid in our situation.

Proposition 5.2. *There exists $\kappa > 0$ such that for any ε small enough, the following is true.*

The functional $L(t) + \kappa Q(t)$ is decreasing in time. Moreover, for any time t where waves with strength $|\sigma_i|$ and $|\sigma_j|$ interact the jump of Q at this time verifies

$$\Delta L(t) + \kappa \Delta Q(t) \leq -\frac{\kappa}{2} |\sigma_i| |\sigma_j|.$$

Especially, there exists a constant $C > 0$, such that for every $\nu > 0$, $T > 0$:

$$\begin{aligned} \|\psi_\nu\|_{L^\infty(0,T,BV(\mathbb{R}))} &\leq 2\varepsilon, \\ \|\psi_\nu(t, \cdot) - \psi_\nu(s, \cdot)\|_{L^1} &\leq C|t - s|, \quad 0 < s < t < T, \\ \text{The function } \psi_\nu &\text{ verifies the Condition 2.5 with constant } C. \end{aligned}$$

Proof. The definitions of the functional Q and L do not depend on the propagation speed of the waves, as long as one verifies the rules that only approaching waves can interact in the future, and (after interaction) interacting waves will not be approaching anymore in the future. These two rules are still valid, thanks to the separation of wave speeds by families in Proposition 4.1. Therefore, the evolution rule of $L(t)$ and $Q(t)$ after each collision depends only on the Riemann solvers which are identical to the real front tracking algorithm. So we recover the estimates involving $\Delta Q(t)$ and $\Delta L(t)$ in the exact same way as the original front tracking method (see [3]).

This estimate shows that $L(t) + \kappa Q(t)$ is a non increasing function in time. We have also that $Q(t) \leq \varepsilon L(t)$. So for every time $t > 0$

$$L(t) \leq L(t) + \kappa Q(t) \leq L(0) + \kappa Q(0) \leq (1 + \kappa \varepsilon) \varepsilon,$$

which provides the uniform BV bound for ε small enough.

As in [3, (7.79)], we have that

$$\begin{aligned} \|\psi_\nu(t, \cdot) - \psi_\nu(s, \cdot)\|_{L^1} &\leq \mathcal{O}(1)|t - s| \left(\sup_{\tau \in \mathbf{R}^+} L(\tau) \right) [\text{maximum speed}] \\ &\leq C|t - s|, \end{aligned}$$

since the maximum speed is uniformly bounded by $\hat{\lambda}$ defined in Proposition 4.1.

The proof of the last statement is identical to [3, Lemma 7.3], since it depends only on the interaction rules, and on the finite speed of propagation. \square

For every time $r > 0$, we denote by $\mathcal{P}(r)$ the set of i corresponding to non-physical waves. For the same reasons, the following lemma is unchanged from [1, Lemma 3.1].

Lemma 5.3 (from [1, Lemma 3.1]). *If*

$$\lim_{\nu \rightarrow \infty} \epsilon_\nu \left(N_\nu + \frac{1}{\delta_\nu} \right)^k = 0, \quad (5.10)$$

for every positive integer k , then the total strength of non-physical waves in ψ_ν goes to zero uniformly in t as $\nu \rightarrow \infty$:

$$\sup_{r \in [0, T]} \sum_{i \in \mathcal{P}(r)} |\sigma_i| \rightarrow 0, \text{ when } \nu \rightarrow \infty.$$

Lemma 5.3 is based on [1, Lemma 3.1], where the result is given for piecewise-constant front tracking solutions with shocks moving with Rankine-Hugoniot speed. A proof is also provided in [1]. The proof of our Lemma 5.3, where shocks move according to shift functions, is identical.

6. THE WEIGHT FUNCTION a

For any pairwise interaction between two small (shock or rarefaction) waves, one has the following estimates (see [3, 20, 41]). See Figure 1.

Proposition 6.1. *Call σ', σ'' the strengths of two interacting wave-fronts, and let σ_1, σ_2 be the strengths of the outgoing waves of the first and second family. σ takes positive sign on a shock and negative value on a rarefaction front.*

Then there exists a constant C_0 (uniformly valid for $u \in B_\varepsilon(d)$) such that

- *If both σ' and σ'' belong to the first family, then*

$$|\sigma_1 - (\sigma' + \sigma'')| + |\sigma_2| \leq C_0 |\sigma' \sigma''| (|\sigma'| + |\sigma''|). \quad (6.1)$$

- *If σ'' is a 1-wave and σ' is a 2-wave, then*

$$|\sigma_1 - \sigma''| + |\sigma_2 - \sigma'| \leq C_0 |\sigma' \sigma''| (|\sigma'| + |\sigma''|). \quad (6.2)$$

- *If both σ' and σ'' belong to the second family, then*

$$|\sigma_1| + |\sigma_2 - (\sigma' + \sigma'')| \leq C_0 |\sigma' \sigma''| (|\sigma'| + |\sigma''|). \quad (6.3)$$

Remark that $B_\varepsilon(d)$ is in the context of Theorem 2.3.

Here we can always choose ε small enough, especially smaller than the ε of Proposition 4.1, and such that

$$C_0 \varepsilon \leq 1. \quad (6.4)$$




FIGURE 1. Pairwise interactions. Each line segment represent a shock or a rarefaction jump. To distinguish from each other, we represent 1-waves by line segment with negative speeds, and 2-waves with positive speeds. The middle picture is for the head-on interaction. The other two pictures are for overtaking interactions.

We now define the following measure $\mu(t, \cdot)$ as a sum of Dirac measures in x :

$$\begin{aligned} \mu(t, x) &= - \sum_{i:1\text{-shock}} \sigma_i \delta_{\{x_i(t)\}} + \sum_{i:2\text{-shock}} \sigma_i \delta_{\{x_i(t)\}} \\ &= - \sum_{i:1\text{-wave}} (\sigma_i)_+ \delta_{\{x_i(t)\}} + \sum_{i:2\text{-wave}} (\sigma_i)_+ \delta_{\{x_i(t)\}}. \end{aligned}$$

The weight function is then defined as

$$a(t, x) = 1 + C \left(L(t) + \kappa Q(t) + \int_{-\infty}^x \mu(t, z) dz \right), \quad (6.5)$$

where the constant C is defined in Proposition 4.1.

Note that the function a is piecewise constant, with discontinuities only along shock curves. In particular it is constant across rarefaction curves and pseudoshock curves. We show that the function a has the following properties.

Proposition 6.2. *There exists $C_0 > 0$, such that for every $\varepsilon > 0$ small enough,*

$$|a(t, x) - 1| \leq C_0 \varepsilon.$$

For every time without wave interaction, and for every x such that a 1-shock σ_i is located at $x = x_i(t)$:

$$1 - 2C|\sigma_i| \leq \frac{a(t, x_i(t)+)}{a(t, x_i(t)-)} \leq 1 - \frac{C}{2}|\sigma_i|, \quad (6.6)$$

For every time without wave interaction, and for every x such that a 2-shock σ_i is located at $x = x_i(t)$:

$$1 + \frac{1}{2}C|\sigma_i| \leq \frac{a(t, x_i(t)+)}{a(t, x_i(t)-)} \leq 1 + 2C|\sigma_i|, \quad (6.7)$$

For every time t with a wave interaction, and almost every x :

$$a(t+, x) \leq a(t-, x). \quad (6.8)$$

Proof. Note that $\|\mu(t)\|_{\mathcal{M}} \leq L(t)$, and

$$L(0) + \kappa Q(0) = O(\varepsilon).$$

Since $L(t) + \kappa Q(t)$ is decreasing, it means that

$$|a(t, x) - 1| = O(\varepsilon).$$

So for ε small enough, $1/2 < 1/a(t, x(t)-) < 2$. Now:

$$\frac{a(t, x_i(t)+)}{a(t, x(t)-)} - 1 = \frac{1}{a(t, x(t)-)} (a(t, x_i(t)+) - a(t, x(t)-)) = \frac{1}{a(t, x(t)-)} C |\sigma_i| \alpha,$$

with $\alpha = 1$ if the shock σ_i is a 2-shock, and $\alpha = -1$ if it is a 1-shock. This shows (6.6) and (6.7).

Consider a time t with a wave interaction. From the definition of the a function,

$$\sup_{\mathbb{R}} (a(t+, x) - a(t-, x)) \leq C \left(\int_{\mathbb{R}} |\mu(t+) - \mu(t-)| dx + (\Delta L(t) + \kappa \Delta Q(t)) \right). \quad (6.9)$$

Assume that the waves interacts at $x = x_0$. The interacting wave fronts are σ' σ'' leading to outgoing waves σ_1, σ_2 . We study $\mu(t+) - \mu(t-)$ by considering separately all the possible kind of interactions.

If the interaction involves a pseudoshock, then the strengths of the shocks are not affected, so $\mu(t+) - \mu(t-) = 0$, and since $\Delta L + \kappa \Delta Q \leq 0$, we have also $a(t+, x) - a(t-, x) \leq 0$ for every $x \in \mathbb{R}$.

If the simplified solver is used, then

$$\mu(t+) - \mu(t-) = \delta_{\{x_0\}} [(\sigma' + \sigma'')_+ - (\sigma')_+ - (\sigma'')_+] \leq 0.$$

We still have $a(t+, x) - a(t-, x) \leq 0$ in this case.

It remains to consider the cases involving the accurate solver. They correspond to the three cases of Proposition 6.1.

(i). If σ'' is a 1-wave and σ' is a 2-wave. Using the definition of μ to justify the first equality below, the fact that $y \rightarrow (y)_+$ is Lipschitz with constant 1 for the second inequality, (6.2) for the third inequality, and Proposition 5.2 for ε small enough to get the last inequality, we have:

$$\begin{aligned} |\mu(t+) - \mu(t-)| &= \delta_{\{x_0\}} |(\sigma_2)_+ - (\sigma_1)_+ - ((\sigma')_+ - (\sigma'')_+)| \\ &\leq \delta_{\{x_0\}} |(\sigma_2)_+ - (\sigma')_+| + |(\sigma_1)_+ - (\sigma'')_+| \\ &\leq \delta_{\{x_0\}} (|\sigma_2 - \sigma'| + |\sigma_1 - \sigma''|) \\ &\leq \delta_{\{x_0\}} C_0 |\sigma' \sigma''| (|\sigma'| + |\sigma''|) \\ &\leq -(\Delta L(t) + \kappa \Delta Q(t)) \delta_{\{x_0\}}. \end{aligned}$$

Remark also that we used (6.4). But using (6.9) and the fact that $\Delta L(t) + \kappa \Delta Q(t) < 0$, gives

$$\sup_{\mathbb{R}} (a(t+, x) - a(t-, x)) \leq 0.$$

(ii). The cases (6.1) and (6.3) are similar. Let us do (6.1) in detail. Both σ' and σ'' belong to the first family. In this case,

$$\begin{aligned} |\mu(t+) - \mu(t-)| &= \delta_{\{x_0\}} |(\sigma_2)_+ - (\sigma_1)_+ - (-(\sigma')_+ - (\sigma'')_+)| \\ &\leq \delta_{\{x_0\}} (|(\sigma_2)_+| + |(\sigma_1)_+ - (\sigma')_+ - (\sigma'')_+|) \\ &\leq \delta_{\{x_0\}} (|(\sigma_2)| + |(\sigma_1)_+ - (\sigma')_+ - (\sigma'')_+|) \end{aligned}$$

We need to separate cases depending on the nature of the incoming waves, and the kind of Riemann solver used.

(ii)-1. If σ' and σ'' are rarefactions, then σ_1 is also a rarefaction, and

$$|(\sigma_1)_+ - (\sigma')_+ - (\sigma'')_+| = 0.$$

(ii)-2. If σ' and σ'' are shocks, then σ_1 is also a shock, and

$$|(\sigma_1)_+ - (\sigma')_+ - (\sigma'')_+| = |\sigma_1 - \sigma' - \sigma''|.$$

(ii)-3. Finally, if one of σ' and σ'' is a shock, let say σ' , and the other a rarefaction, let say σ'' . Then

$$\begin{aligned} |(\sigma_1)_+ - (\sigma')_+ - (\sigma'')_+| &= |(\sigma_1)_+ - \sigma'| \leq |\sigma'' - (\sigma_1)_-| + |\sigma_1 - \sigma' - \sigma''| \\ &\leq -(\Delta L(t) + \kappa \Delta Q(t)), \end{aligned}$$

since $|\sigma'' - (\sigma_1)_-| \leq -\Delta L(t) + C_0 \varepsilon |\sigma'| |\sigma''|$.

Gathering all the cases, we obtain:

$$\sup_{\mathbb{R}} (a(t+, x) - a(t-, x)) \leq 0.$$

□

7. PROOF OF PROPOSITION 3.2

This section is devoted to the proof of Proposition 3.2. In our front-tracking procedure, we are stopping and restarting the clock every time there is a collision between waves (when the waves initiated from distinct Riemann problems). Weak solutions u to (1.1) naturally lies in $C^0(\mathbb{R}^+; W^{-1,\infty}(\mathbb{R}))$. Note that the formulation of the entropy inequality (1.6) holds with a boundary term for $t = 0$, and this classically implies that u is continuous in time at $t = 0$ with values in $L^1_{\text{loc}}(\mathbb{R})$. Because $L^1_{\text{loc}}(\mathbb{R})$ is a strong topology, it implies that $\eta(u)$ is also continuous at $t = 0$ in the same topology in x . However, because $\eta(u)$ verifies only Inequality (1.5), $\eta(u)$ does not share this regularity in time for $t > 0$. Therefore $\eta(u)$ is well defined only almost everywhere in time. However, this technicality of stopping and restarting the clock at any time t is not a real issue, and its resolution can be formalized with the use of approximate limits as follows. For a reference on approximate limits, see [18, p. 55-57].

Lemma 7.1 (A technical lemma on stopping and restarting the clock (from [28, Lemma 2.5])).

Let $u \in L^\infty(\mathbb{R}^+ \times \mathbb{R})$ be a weak solution to (1.1) with initial data u^0 . Further, assume that u is entropic for the entropy η , i.e. verifies (1.5) in the sense of distribution. Assume also that u verifies the strong trace property (Definition 1.2). Then for all $v \in \mathcal{V}_0$, and for all $c, d \in \mathbb{R}$ with $c < d$, the following approximate right- and left-hand limits

$$\text{ap} \lim_{t \rightarrow t_0^\pm} \int_c^d \eta(u(t, x)|v) dx \quad (7.1)$$

exist for all $t_0 \in (0, \infty)$ and verify

$$\text{ap} \lim_{t \rightarrow t_0^-} \int_c^d \eta(u(t, x)|v) dx \geq \text{ap} \lim_{t \rightarrow t_0^+} \int_c^d \eta(u(t, x)|v) dx. \quad (7.2)$$

Furthermore, the approximate right-hand limit exists at $t_0 = 0$ and verifies

$$\int_c^d \eta(u^0(x)|v) dx \geq \text{ap} \lim_{t \rightarrow t_0^+} \int_c^d \eta(u(t, x)|v) dx. \quad (7.3)$$

The proof of Lemma 7.1 follows exactly the proof of [28, Lemma 2.5]. For this reason, we do not include a proof here.

We gather in the following lemma useful simple properties of the relative quantities.

Lemma 7.2. For any \mathcal{O} open subset of \mathcal{V} with $\overline{\mathcal{O}} \subset \mathcal{V}$, there exists a constant $C > 0$ such that

$$\begin{aligned} |q(a; b)| &\leq C \eta(a|b), \quad \forall (a, b) \in \mathcal{V}_0 \times \overline{\mathcal{O}}, \\ |q(a; b_1) - q(a; b_2)| &\leq C |b_1 - b_2|, \quad \forall (b_1, b_2) \in \overline{\mathcal{O}}^2, \quad a \in \mathcal{V}, \\ |\eta(a|b_1) - \eta(a|b_2)| &\leq C |b_1 - b_2|, \quad \forall (b_1, b_2) \in \overline{\mathcal{O}}^2, \quad a \in \mathcal{V}. \end{aligned}$$

Proof. Consider an open set \mathcal{O}' such that $\overline{\mathcal{O}} \subset \mathcal{O}'$ and $\overline{\mathcal{O}'} \subset \mathcal{V}$. Since both $f, q \in C^0(\mathcal{V}_0)$ and $f' \in C^0(\overline{\mathcal{O}})$, $q(\cdot; \cdot)$ is uniformly bounded on $\mathcal{V}_0 \times \overline{\mathcal{O}}$. Moreover, from Lemma 3.1, $\eta(\cdot, \cdot)$ is bounded above and below uniformly on $(\mathcal{V}_0 \setminus \mathcal{O}') \times \overline{\mathcal{O}}$. Therefore there exists a constant such that the first equality holds for those values. But from the definition in (3.2), $q(b; b) = \partial_1 q(b, b) = 0$ for all $b \in \mathcal{V}$. So using Lemma 3.1, and the fact that $q'' \in C^0(\overline{\mathcal{O}'})$ we have that there exists a constant C such that

$$|q(a; b)| \leq C|a - b|^2 \leq \frac{C}{c^*} \eta(a|b), \quad \text{for } (a, b) \in \overline{\mathcal{O}'} \times \overline{\mathcal{O}}.$$

This proves the first inequality of the lemma.

From the definition of $q(\cdot; \cdot)$ in (3.2), denoting $h = q - \eta'f \in C^1(\overline{\mathcal{O}})$, we have for every $(a, b_1, b_2) \in \mathcal{V}_0 \times \overline{\mathcal{O}^2}$:

$$|q(a; b_1) - q(a; b_2)| = |h(b_2) - h(b_1) + [\eta'(b_2) - \eta'(b_1)]f(a)| \leq \sup_{\overline{\mathcal{O}}}(|h'| + |\eta''|)(1 + \sup_{\mathcal{V}_0}|f|)|b_2 - b_1|.$$

The proof of the last statement is similar. \square

We now prove Proposition 3.2. First we fix the value v to be bigger than both $\hat{\lambda}$ and the constant C of Lemma 7.2. Take $0 < \varepsilon < 1/2$ small enough such that Theorem 2.3, Proposition 4.1, and Proposition 5.2 hold true. For any initial value u^0 , and wild solution $u \in \mathcal{S}_{weak}$, we consider the family of solutions ψ_ν of the modified front tracking method. We want now to choose a particular one. Fix $T, R > 0$, and $p \in \mathbb{N}$. First we insure that the initial value verifies

$$\|u^0 - \psi_\nu(0, \cdot)\|_{L^2(-R, R)} \leq \frac{1}{p}.$$

This fixes N_ν . Then we fix $\delta_\nu = 1/(pT)$. Thanks to Lemma 5.3, we can choose ε_ν such that

$$\sup_{r \in [0, T]} \sum_{i \in \mathcal{P}(r)} |\sigma_i| \leq \frac{1}{pT}.$$

We denote by ψ the associated solution to the modified front tracking method ψ_ν . Especially, it verifies

$$\|u(0, \cdot) - \psi(0, \cdot)\|_{L^2(-R, R)} \leq \|u^0 - u(0, \cdot)\|_{L^2(-R, R)} + \frac{1}{p}, \quad (7.4)$$

$$T\delta_\nu \sup_{t \in [0, T]} L(t) \leq \frac{1}{p}, \quad (7.5)$$

$$T \sup_{r \in [0, T]} \sum_{i \in \mathcal{P}(r)} |\sigma_i| \leq \frac{1}{p}. \quad (7.6)$$

Proposition 5.2 provides three of the four properties of Proposition 3.2. It remains only to show the control in L^2 of $\psi(t, \cdot) - u(t, \cdot)$. Recall that as in Section 5, for every time $r > 0$, we denote by $\mathcal{P}(r)$ the set of i corresponding to non-physical waves.

Consider two successive interaction times $t_j < t_{j+1}$ of the front tracking solution ψ . Let the curves of discontinuity between the two times $t_j < t_{j+1}$ be h_1, \dots, h_N for $N \in \mathbb{N}$ such that

$$h_1(t) < \dots < h_N(t), \quad (7.7)$$

for all $t \in (t_j, t_{j+1})$. We only work on the cone of information, so we define for all times t

$$h_0(t) = -R + vt, \quad (7.8)$$

$$h_{N+1} = R - vt. \quad (7.9)$$

Note that there are no interactions between wave fronts in ψ and the cone of information (coming from h_0 and h_{N+1}). For any $t \in [t_j, t_{j+1}]$, note that on $Q = \{(r, x) : t_j < r < t, h_i(r) < x < h_{i+1}(r)\}$,

the function $\psi(r, x) = b$ is constant. Moreover, by construction, the weight function $a(r, x)$ is also constant on this set. Therefore, integrating (3.3) on Q , and using the strong trace property of Definition 1.2, we find

$$\begin{aligned} & \text{ap} \lim_{s \rightarrow t^-} \int_{h_i(t)}^{h_{i+1}(t)} a(t-, x) \eta(u(s, x)) |\psi(t, x)| dx \\ & \leq \text{ap} \lim_{s \rightarrow t_j^+} \int_{h_i(t_j)}^{h_{i+1}(t_j)} a(t_j+, x) \eta(u(s, x)) |\psi(x, t_j)| dx + \int_{t_j}^t (F_i^+(r) - F_{i+1}^-(r)) dr, \end{aligned}$$

where

$$\begin{aligned} F_i^+(r) &= a(r, h_i(r)+) \left[q(u(r, h_i(r)+); \psi(r, h_i(r)+)) - \dot{h}_i(r) \eta(u(r, h_i(r)+)) |\psi(r, h_i(r)+)| \right], \\ F_i^-(r) &= a(r, h_i(r)-) \left[q(u(r, h_i(r)-); \psi(r, h_i(r)-)) - \dot{h}_i(r) \eta(u(r, h_i(r)-)) |\psi(r, h_i(r)-)| \right]. \end{aligned}$$

We sum in i , and combine the terms corresponding to i into one sum, and the terms corresponding to $i+1$ into another sum, to find

$$\begin{aligned} & \text{ap} \lim_{s \rightarrow t^-} \int_{-R+vt}^{R-vt} a(t-, x) \eta(u(s, x)) |\psi(t, x)| dx \\ & \leq \text{ap} \lim_{s \rightarrow t_j^+} \int_{-R+vt_j}^{R-vt_j} a(t_j+, x) \eta(u(s, x)) |\psi(t_j, x)| dx + \sum_{i=1}^N \int_{t_j}^t (F_i^+(r) - F_i^-(r)) dr, \end{aligned}$$

where we have used that $F_0^+ \leq 0$ and $F_{N+1}^- \geq 0$ thanks to the first statement of Lemma 7.2, the definition of v , and the fact that $\dot{h}_0 = -v = -\dot{h}_{N+1}$.

We decompose the sum into three sums, one corresponding to the shock fronts, one for the rarefaction fronts, and one for the pseudoshocks. Thanks to Proposition 4.1 and Proposition 6.2, for any i corresponding to a shock front:

$$F_i^+(r) - F_i^-(r) \leq 0, \quad \text{for almost every } t_j < r < t.$$

Denote \mathcal{R} the set of i corresponding to approximated rarefaction fronts. Then for any $i \in \mathcal{R}$ by construction, $a(h_i(r)+, r) = a(h_i(r)-, r)$. And from Proposition 4.4, and (7.5):

$$\sum_{i \in \mathcal{R}} \int_{t_j}^t (F_i^+(r) - F_i^-(r)) dr \leq C \delta_\nu (t - t_j) \sum_{i \in \mathcal{R}} |\sigma_i| \leq C \delta_\nu (t - t_j) L(t) \leq \frac{C}{pT} (t - t_j).$$

Consider now the case when $i \in \mathcal{P}(r)$. Recall that pseudoshocks travel with supersonic (greater-than-characteristic) speed $\hat{\lambda}$. Thus, we must have that for almost every time r : $u(r, h_i(r)+) = u(r, h_i(r)-)$. This is because if $u(r, h_i(r)+) \neq u(r, h_i(r)-)$, then the shock $(u(r, h_i(r)+), u(r, h_i(r)-), \hat{\lambda})$ would be traveling with speed greater than any of the eigenvalues of Df , a contradiction. By construction of the a function, we know that a does not have a jump across pseudoshocks, so we have also $a(r, h_i(r)+) = a(r, h_i(r)-)$. Therefore, thanks to the second and third estimates of Lemma 7.2,

$$F_i^+(r) - F_i^-(r) \leq C |\psi(r, h_i(r)+) - \psi(r, h_i(r)-)| = C |\sigma_i|.$$

Then, from (7.6) we receive

$$\sum_{i \in \mathcal{P}(r)} \int_{t_j}^t (F_i^+(r) - F_i^-(r)) dr \leq C (t - t_j) \sum_{i \in \mathcal{P}(t)} |\sigma_i| \leq \frac{C(t - t_j)}{pT}.$$

Gathering all the families of waves, we find:

$$\begin{aligned} & \text{ap} \lim_{s \rightarrow t^-} \int_{-R+vt}^{R-vt} a(t-, x) \eta(u(s, x) | \psi(t, x)) dx \\ & \leq \text{ap} \lim_{s \rightarrow t_j^+} \int_{-R+vt_j}^{R-vt_j} a(t_j+, x) \eta(u(s, x) | \psi(t_j, x)) dx + \frac{C(t - t_j)}{pT}. \end{aligned}$$

Consider now any $0 < t < T$, and denote $0 < t_1 < \dots < t_J$ the times of wave interactions before t , $t_0 = 0$, and $t_{J+1} = t$. Using the convexity of η , Lemma 7.1, and (6.8) we find:

$$\begin{aligned} & \int_{-R+vt}^{R-vt} a(t, x) \eta(u(t, x) | \psi(t, x)) dx - \int_{-R}^R a(0, x) \eta(u(0, x) | \psi(0, x)) dx \\ & \leq \text{ap} \lim_{s \rightarrow t^+} \int_{-R+vt}^{R-vt} a(t, x) \eta(u(s, x) | \psi(t, x)) dx - \int_{-R}^R a(0, x) \eta(u(0, x) | \psi(0, x)) dx \\ & \leq \sum_{j=1}^{J+1} \left(\text{ap} \lim_{s \rightarrow t_j^+} \int_{-R+vt_j}^{R-vt_j} a(t_j-, x) \eta(u(s, x) | \psi(t, x)) dx \right. \\ & \quad \left. - \text{ap} \lim_{s \rightarrow t_{j-1}^+} \int_{-R+vt_{j-1}}^{R-vt_{j-1}} a(t_{j-1}-, x) \eta(u(s, x) | \psi(t, x)) dx \right) \\ & \leq \sum_{j=1}^{J+1} \left(\text{ap} \lim_{s \rightarrow t_j^-} \int_{-R+vt_j}^{R-vt_j} a(t_j-, x) \eta(u(s, x) | \psi(t, x)) dx \right. \\ & \quad \left. - \text{ap} \lim_{s \rightarrow t_{j-1}^+} \int_{-R+vt_{j-1}}^{R-vt_{j-1}} a(t_{j-1}-, x) \eta(u(s, x) | \psi(t, x)) dx \right) \\ & \leq \sum_{j=1}^{J+1} \left(\text{ap} \lim_{s \rightarrow t_j^-} \int_{-R+vt_j}^{R-vt_j} a(t_j-, x) \eta(u(s, x) | \psi(t, x)) dx \right. \\ & \quad \left. - \text{ap} \lim_{s \rightarrow t_{j-1}^+} \int_{-R+vt_{j-1}}^{R-vt_{j-1}} a(t_{j-1}+, x) \eta(u(s, x) | \psi(t, x)) dx \right) \\ & \leq \sum_{j=1}^{J+1} C \frac{(t_j - t_{j-1})}{Tp} \leq C \frac{t}{Tp} \leq \frac{C}{p}. \end{aligned}$$

Using that $|a - 1| < 1/2$ and (7.4), we get that for every $0 < t < T$

$$\int_{-R+vt}^{R-vt} \eta(u(t, x) | \psi(t, x)) dx \leq 2 \|u^0 - u(0, \cdot)\|_{L^2(-R, R)}^2 + \frac{C}{p}.$$

Choosing p big enough such that $C/p < 1/m$ gives the result.

APPENDIX A. PROOF OF LEMMA 2.6

For any $R > 0$, Consider $B_{-1} = L^1(-R, R)$, $B_0 = L^2(-R, R)$, $B_1 = BV(-R, R)$. These Banach Spaces are nested into each other, and B_1 is injected in B_0 compactly. Moreover, the ψ_n are uniformly bounded in $L^\infty(0, R; B_1)$ and, from (2.5), $\{\partial_t \psi_n\}_{n \in \mathbb{N}}$ is uniformly bounded in $L^\infty(0, R; B_{-1})$. Therefore, thanks to the Aubin Lions lemma, $\{\psi_n\}_{n \in \mathbb{N}}$ is precompact in $C^0(0, R; L^2(-R, R))$. Therefore there exists a subsequence, still denoted $\{\psi_n\}$, and $\psi \in L^\infty(\mathbb{R}^+ \times \mathbb{R})$ such that ψ_n converges to ψ in $C^0(0, R; L^2(-R, R))$ for all $R > 0$. Obviously, the convergence holds also in $L^2((0, R) \times (-R, R))$. Then, by a diagonal argument, we can re-extract a subsequence such that the convergence holds for almost every $(t, x) \in \mathbb{R}^+ \times \mathbb{R}$. The function ψ still verifies $\|\psi\|_{L^\infty(\mathbb{R}^+; BV(\mathbb{R}))} \leq C$. Thus, we can choose a representative of ψ that is right-continuous (in x).

It remains to show that ψ verifies the Bounded Variation Condition of Definition 2.2 for $\delta := \frac{1}{\lambda}$. Let γ , and a', b' be a space-like curve as in Definition 2.1. We will now show that the function $x \mapsto \psi(\gamma(x), x)$ has bounded variation. Let any points $P_i = (\gamma(x_i), x_i)$ be given, with $x_0 < x_1 < \dots < x_m$. At each time $t_i = \gamma(x_i)$, we know that $\psi_n(t_i, \cdot)$ will converge to $\psi(t_i, \cdot)$ almost everywhere. Due to $\psi(t_i, \cdot)$ being right-continuous for each i , for any $\epsilon > 0$ we can find $x'_i \in [x_i, x_i + \epsilon]$ such that

$$|\psi(t_i, x'_i) - \psi(t_i, x_i)| < \epsilon, \quad \lim_{n \rightarrow \infty} \psi_n(t_i, x'_i) = \psi(t_i, x'_i), \quad (\text{A.1})$$

$$\hat{\lambda}|t_i - t_{i-1}| < x'_i - x'_{i-1}, \quad \hat{\lambda}t_m < b - x_m, \quad \hat{\lambda}t_0 < x_0 - a, \quad (\text{A.2})$$

for constants a and b to be chosen momentarily.

Define γ' to be the polygonal line with vertices at the points $P'_i := (t_i, x'_i)$ and also define $\bar{\gamma}$ to be the constant map $\bar{\gamma}(x) = 0$ for all $x \in (a, b)$, where a and b are from (A.2).

Then, by choosing a and b such that $a' - a > 0$ is suitably large and $b - b' > 0$ is suitably large, the properties (A.1) and (A.2) imply that $\bar{\gamma}$ dominates γ' (as in Definition 2.4).

Since ψ_n verifies uniformly the property of Condition 2.5, from (2.4), we have

$$\sum_{i=1}^m |\psi(t_i, x_i) - \psi(t_{i-1}, x_{i-1})| \leq 2m\epsilon + \sum_{i=1}^m |\psi(t_i, x'_i) - \psi(t_{i-1}, x'_{i-1})| \quad (\text{A.3})$$

$$\leq 2m\epsilon + \limsup_{n \rightarrow \infty} (\text{Tot. Var.} \{\psi_n; \gamma'\}) \quad (\text{A.4})$$

$$\leq 2m\epsilon + C \limsup_{n \rightarrow \infty} (\text{Tot. Var.} \{\psi_n(0, \cdot); (a, b)\}). \quad (\text{A.5})$$

Recall then that the total variation of $\psi_n(0, \cdot)$ on the interval (a, b) is uniformly bounded in n . Then, since $\epsilon > 0$ and the points P_i were arbitrary, we conclude that ψ verifies the Bounded Variation Condition (Definition 2.2).

REFERENCES

- [1] Paolo Baiti and Helge Kristian Jenssen, *On the front-tracking algorithm*, J. Math. Anal. Appl. **217** (1998), no. 2, 395–404. MR1492096
- [2] Stefano Bianchini and Alberto Bressan, *Vanishing viscosity solutions of nonlinear hyperbolic systems*, Ann. of Math. (2) **161** (2005), no. 1, 223–342. MR2150387
- [3] Alberto Bressan, *Hyperbolic systems of conservation laws*, Oxford Lecture Series in Mathematics and its Applications, vol. 20, Oxford University Press, Oxford, 2000. The one-dimensional Cauchy problem. MR1816648
- [4] Alberto Bressan, Graziano Crasta, and Benedetto Piccoli, *Well-posedness of the Cauchy problem for $n \times n$ systems of conservation laws*, Mem. Amer. Math. Soc. **146** (2000), no. 694, viii+134. MR1686652
- [5] Alberto Bressan and Paola Goatin, *Oleinik type estimates and uniqueness for $n \times n$ conservation laws*, J. Differential Equations **156** (1999), no. 1, 26–49. MR1701818
- [6] Alberto Bressan and Philippe LeFloch, *Uniqueness of weak solutions to systems of conservation laws*, Arch. Rational Mech. Anal. **140** (1997), no. 4, 301–317. MR1489317
- [7] Alberto Bressan and Marta Lewicka, *A uniqueness condition for hyperbolic systems of conservation laws*, Discrete Contin. Dynam. Systems **6** (2000), no. 3, 673–682. MR1757395
- [8] Alberto Bressan, Tai-Ping Liu, and Tong Yang, *L^1 stability estimates for $n \times n$ conservation laws*, Arch. Ration. Mech. Anal. **149** (1999), no. 1, 1–22. MR1723032

- [9] Gui Qiang Chen, *Convergence of the Lax-Friedrichs scheme for isentropic gas dynamics. III*, Acta Math. Sci. (English Ed.) **6** (1986), no. 1, 75–120. MR924671
- [10] Gui-Qiang Chen, Hermano Frid, and Yachun Li, *Uniqueness and stability of Riemann solutions with large oscillation in gas dynamics*, Comm. Math. Phys. **228** (2002), no. 2, 201–217. MR1911734
- [11] Constantine M. Dafermos, *The second law of thermodynamics and stability*, Arch. Rational Mech. Anal. **70** (1979), no. 2, 167–179. MR546634
- [12] ———, *Hyperbolic conservation laws in continuum physics*, Fourth, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 325, Springer-Verlag, Berlin, 2016. MR3468916
- [13] Camillo De Lellis, Felix Otto, and Michael Westdickenberg, *Structure of entropy solutions for multi-dimensional scalar conservation laws*, Arch. Ration. Mech. Anal. **170** (2003), no. 2, 137–184. MR2017887
- [14] ———, *Minimal entropy conditions for Burgers equation*, Quart. Appl. Math. **62** (2004), no. 4, 687–700. MR2104269
- [15] Xia Xi Ding, Gui Qiang Chen, and Pei Zhu Luo, *Convergence of the Lax-Friedrichs scheme for isentropic gas dynamics. I, II*, Acta Math. Sci. (English Ed.) **5** (1985), no. 4, 415–432, 433–472. MR922139
- [16] Ronald J. DiPerna, *Uniqueness of solutions to hyperbolic conservation laws*, Indiana Univ. Math. J. **28** (1979), no. 1, 137–188. MR523630
- [17] ———, *Convergence of the viscosity method for isentropic gas dynamics*, Comm. Math. Phys. **91** (1983), no. 1, 1–30. MR719807
- [18] Lawrence C. Evans and Ronald F. Gariepy, *Measure theory and fine properties of functions*, Revised, Textbooks in Mathematics, CRC Press, Boca Raton, FL, 2015. MR3409135
- [19] Eduard Feireisl, Ondřej Kreml, and Alexis F. Vasseur, *Stability of the isentropic Riemann solutions of the full multidimensional Euler system*, SIAM J. Math. Anal. **47** (2015), no. 3, 2416–2425. MR3357629
- [20] James Glimm, *Solutions in the large for nonlinear hyperbolic systems of equations*, Comm. Pure Appl. Math. **18** (1965), 697–715.
- [21] William M. Golding, Sam G. Krupa, and Alexis F. Vasseur, *Sharp α -contraction estimates for small extremal shocks*, arXiv e-prints (December 2020), arXiv:2012.09261, available at 2012.09261.
- [22] David Hoff, *Invariant regions for systems of conservation laws*, Trans. Amer. Math. Soc. **289** (1985), no. 2, 591–610. MR784005
- [23] Moon-Jin Kang and Alexis Vasseur, *Contraction property for large perturbations of shocks of the barotropic Navier-Stokes system*, arXiv e-prints (2017Dec), available at 1712.07348. To appear in Journal of the European Mathematical Society.
- [24] ———, *Uniqueness and stability of entropy shocks to the isentropic Euler system in a class of inviscid limits from a large family of Navier-Stokes systems*, arXiv:1902.01792, To appear in Inventiones Mathematicae (2020).
- [25] Moon-Jin Kang and Alexis F. Vasseur, *Criteria on contractions for entropic discontinuities of systems of conservation laws*, Arch. Ration. Mech. Anal. **222** (2016), no. 1, 343–391. MR3519973
- [26] Sam G. Krupa, *Criteria for the α -contraction and stability for the piecewise-smooth solutions to hyperbolic balance laws*, Commun. Math. Sci. **18** (2020), no. 6, 1493–1537.
- [27] Sam G. Krupa, *Finite time stability for the Riemann problem with extremal shocks for a large class of hyperbolic systems*, Journal of Differential Equations **273** (2021), 122–171.
- [28] Sam G. Krupa and Alexis F. Vasseur, *On uniqueness of solutions to conservation laws verifying a single entropy condition*, J. Hyperbolic Differ. Equ. **16** (2019), no. 1, 157–191. MR3954680
- [29] ———, *Stability and uniqueness for piecewise smooth solutions to a nonlocal scalar conservation law with applications to Burgers-Hilbert equation*, SIAM J. Math. Anal. **52** (2020), no. 3, 2491–2530. MR4101366
- [30] Young-Sam Kwon and Alexis Vasseur, *Strong traces for solutions to scalar conservation laws with general flux*, Arch. Ration. Mech. Anal. **185** (2007), no. 3, 495–513. MR2322819
- [31] Peter David Lax, *Hyperbolic systems of conservation laws. II*, Comm. Pure Appl. Math. **10** (1957), 537–566. MR0093653
- [32] Nicholas Leger, *L^2 stability estimates for shock solutions of scalar conservation laws using the relative entropy method*, Archive for Rational Mechanics and Analysis **199** (2011), no. 3, 761–778.
- [33] Nicholas Leger and Alexis F. Vasseur, *Relative entropy and the stability of shocks and contact discontinuities for systems of conservation laws with non-BV perturbations*, Archive for Rational Mechanics and Analysis **201** (2011), no. 1, 271–302.
- [34] P.-L. Lions, B. Perthame, and E. Tadmor, *Kinetic formulation of the isentropic gas dynamics and p -systems*, Comm. Math. Phys. **163** (1994), no. 2, 415–431. MR1284790
- [35] Pierre-Louis Lions, Benoît Perthame, and Panagiotis E. Souganidis, *Existence and stability of entropy solutions for the hyperbolic systems of isentropic gas dynamics in Eulerian and Lagrangian coordinates*, Comm. Pure Appl. Math. **49** (1996), no. 6, 599–638. MR1383202
- [36] E. Yu. Panov, *Existence of strong traces for quasi-solutions of multidimensional conservation laws*, J. Hyperbolic Differ. Equ. **4** (2007), no. 4, 729–770. MR2374223

- [37] Evgueni Yu. Panov, *Uniqueness of the solution of the cauchy problem for a first order quasilinear equation with one admissible strictly convex entropy*, Mathematical Notes **55** (1994May), no. 5, 517–525.
- [38] Denis Serre, *Systems of conservation laws. 1*, Cambridge University Press, Cambridge, 1999. Hyperbolicity, entropies, shock waves. Translated from the 1996 French original by I. N. Sneddon. MR1707279
- [39] Denis Serre and Alexis F. Vasseur, *L^2 -type contraction for systems of conservation laws*, J. Éc. polytech. Math. **1** (2014), 1–28. MR3322780
- [40] Luis Silvestre, *Oscillation properties of scalar conservation laws*, Comm. Pure Appl. Math. **72** (2019), no. 6, 1321–1348. MR3948559
- [41] Joel Smoller, *Shock waves and reaction-diffusion equations*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Science], vol. 258, Springer-Verlag, New York-Berlin, 1983.
- [42] L. Tartar, *Compensated compactness and applications to partial differential equations*, Nonlinear analysis and mechanics: Heriot-Watt Symposium, Vol. IV, 1979, pp. 136–212. MR584398
- [43] Alexis F. Vasseur, *Time regularity for the system of isentropic gas dynamics with $\gamma = 3$* , Comm. Partial Differential Equations **24** (1999), no. 11-12, 1987–1997. MR1720782
- [44] ———, *Strong traces for solutions of multidimensional scalar conservation laws*, Arch. Ration. Mech. Anal. **160** (2001), no. 3, 181–193. MR1869441
- [45] ———, *Recent results on hydrodynamic limits*, Handbook of Differential Equations: Evolutionary Equations **4** (2008), 323–376.

(Geng Chen)

DEPARTMENT OF MATHEMATICS,
UNIVERSITY OF KANSAS, LAWRENCE, KS, 66049, USA
Email address: gengchen@ku.edu

(Sam G. Krupa)

MAX PLANCK INSTITUTE FOR MATHEMATICS IN THE SCIENCES,
INSELSTR. 22, 04103 LEIPZIG, GERMANY
Email address: Sam.Krupa@mis.mpg.de

(Alexis F. Vasseur)

DEPARTMENT OF MATHEMATICS,
THE UNIVERSITY OF TEXAS AT AUSTIN, AUSTIN, TX 78712, USA
Email address: vasseur@math.utexas.edu