Perturbations in a Chaplygin gas Cosmology

Shambel Sahlu^{1,2,3}, Heba Sami³, Anna-Mia Swart³, Thato Tsabone³, Maye Elmardi⁴ and Amare Abebe³

¹ Astronomy and Astrophysics Department, Entoto Observatory and Research Center,

Ethiopian Space Science and Technology Institute, Addis Ababa, Ethiopia.

² Department of Physics, College of Natural and Computational Science, Wolkite University, Wolkite, Ethiopia.

³ Center for Space Research, North-West University, Mahikeng 2735, South Africa.

⁴ Center for Space Research, North-West University, Potchefstroom 2520, South Africa.

E-mail: hebasami.abdulrahman@gmail.com

Abstract. In this paper we study the perturbations of a cosmic multi-fluid medium consisting of radiation, dust and a Chaplygin gas. To do so, we follow the 1 + 3 covariant formalism and derive the evolution equations of the fluctuations in the energy density for each species of fluid in the multi-fluid system. The solutions to these coupled systems of equations are then computed in both short-wavelength and long-wavelength modes. Our preliminary results suggest that unlike most dark energy models that discourage large-scale structure formation due to the rapid cosmological expansion (which gives little time for fluctuations to coalesce), the Chaplygin-gas model supports the formation of cosmic structures. This is manifested in the solutions of the perturbation equations which show the growth of density fluctuations over time.

1. Introduction

In the last two decades, it has become apparent that the standard cosmological model (ACDM) fails to explain the accelerating expansion of the Universe [1], the existence of dark matter in galaxy clusters [2], the formation of the large-scale structure [3], the inherent inhomogeneity and anisotropy of the Universe on the small scales [4] and so forth. To solve the first two problems, scholars have proposed an exotic fluid, the so-called Chaplygin gas (CG), which acts as a cosmological fluid with an equation of state (EoS) of the form $p_{ch} = -A/\rho_{ch}^{\alpha}$, where p_{ch} and ρ_{ch} are the pressure and energy density of the CG, and A and α are constants such that A > 0 and $0 < \alpha \leq 1$ [5]. From this EoS we can conclude that the EoS parameter for CG can be written as $w_{ch} = -A/\rho_{ch}^{1+\alpha} = -A/(A+B(1+z)^{3(1+\alpha)})$, where z is the redshift. The purpose of this CG fluid is to substitute the dark matter and dark-energy components of the Universe. This model acts as dark-matter in the early universe and dark-energy in the late times of the cosmos. In a universe in which we assume the energy density to consist of matter and CG fluids, the conservation equation reads as

$$\dot{\rho}_t + 3\frac{\dot{a}}{a}(\rho_t + p_t) = 0, \tag{1}$$

where ρ_t is the energy density and p_t is the pressure for the total fluid¹, *a* is the cosmological scale factor and the subscript *t* stands for total (matter+CG) fluids. From this equation, the energy

¹ When referring to matter, we mean either radiation or dust, and total fluid implies matter and Chaplygin gas.

density becomes $\rho_t(z) = [A + B(1+z)^{3(1+\alpha)}]^{\frac{1}{1+\alpha}} + \rho_m(1+z)^{3(1+w_m)}$, where $B = e^{C(1+\alpha)}$ with Ca constant of integration.² Whereas much has been said of the Chaplygin gas as an alternative to a unified dark matter and dark energy model that mimics the cosmic expansion history of a Friedmann-Lemaître-Robertson-Walker (FLRW) background, to our knowledge there is no work in the literature that considered the cosmological perturbations of this fluid model in the 1+3 covariant formalism [6–8]. Henceforth we derive the density perturbation equations and present the solutions both analytically and numerically. From the results, we analyse cosmological implications as far as large-scale structure formation is concerned on both sub- and superhorizon scales [8]. The outline of the manuscript is as follows: in the next section we review the basic spatial gradient variables and we also derive the linear evolution equations. We apply scalar and harmonic decomposition techniques and obtain the wave-number dependent energy density fluctuations of density perturbations by considering radiation-CG, dust-CG and CG like fluids for both wave-length ranges. We then devote Sec. 4 to discussions of our results and the conclusions.

2. Perturbations

For a perfect-fluid system, the following fluid equations hold: $\dot{\rho}_t = -\Theta(\rho_t + p_t)$, and $(\rho_t + p_t)\dot{u}_a + \tilde{\nabla}_a p_t = 0$, from which one can conclude that the 4-acceleration is $\dot{u}_a = \alpha w_{ch} \rho_{ch} D_a^{ch} / (a(\rho_t + p_t)) - w_m \rho_m D_a^m / (a(\rho_t + p_t))$. Another key equation for a general fluid is the so-called Raychaudhuri equation which can be expressed as $\dot{\Theta} = -\Theta^2/3 - (\rho_t + 3p_t)/2 - \tilde{\nabla}^a \dot{u}_a$, where Θ is the cosmological expansion scalar. After finding the comoving gradients of the cosmological expansion and the comoving fractional density gradient of the matter components in a covariant and gauge-invariant way, and taking the harmonically decomposed scalar parts, it can be shown that the scalar perturbations for the matter and CG energy densities, respectively, evolve according to

$$\ddot{\Delta}_{m}^{k} = \Theta(w_{m} - \frac{2}{3})\dot{\Delta}_{m} + \left\{ (1 + w_{m}) \left[\frac{1}{2} (1 + 3w_{m}) + \frac{\Theta^{2} w_{m}}{3\rho_{t}(1 + w_{t})} + \frac{w_{m}(1 + 3w_{t})}{2(1 + w_{t})} + \frac{k^{2} w_{m}}{a^{2}\rho_{t}} \right] \rho_{m} + \left[\frac{1}{3} w_{m} \Theta^{2} (1 + w_{t}) \rho_{t} \right] \Delta_{m}^{k} + (1 + w_{m}) \left[\frac{1}{2} - \frac{3w_{ch}}{2} - \frac{\Theta^{2} w_{ch}}{3\rho_{t}(1 + w_{t})} - \frac{w_{ch}(1 + 3w_{t})}{2(1 + w_{t})} + \frac{k^{2} w_{ch}}{a^{2}\rho_{t}} \right] \rho_{ch} \Delta_{ch}^{k} , \quad (2)$$

$$\begin{split} \ddot{\Delta}_{ch}^{k} &= -\Theta\left(w_{ch} + \frac{2}{3}\right)\dot{\Delta}_{ch}^{k} + (1+w_{ch})\left[\frac{1}{2}(1+3w_{m}) + \frac{\Theta^{2}w_{m}}{3\rho_{t}(1+w_{t})} + \frac{w_{m}(1+3w_{t})}{2(1+w_{t})} + \frac{k^{2}w_{m}}{a^{2}\rho_{t}}\right]\rho_{m}\Delta_{m}^{k} \\ &+ \left\{(1+w_{ch})\left[\frac{1}{2} - \frac{3w_{ch}}{2} - \frac{\Theta^{2}w_{ch}}{3\rho_{t}(1+w_{t})} - \frac{w_{ch}(1+3w_{t})}{2(1+w_{t})} + \frac{k^{2}w_{ch}}{a^{2}\rho_{t}}\right]\rho_{ch} - w_{ch}\left(\frac{1}{3}\Theta^{2} + \frac{1}{2}(1+3w_{t})\rho_{t}\right) \\ &+ \Theta^{2}w_{ch}\left[2(1+2w_{ch})\rho_{ch} + \frac{2}{3}\right]\right\}\Delta_{ch}^{k}, \quad (3)$$

where $k = 2\pi a/\lambda$, k being the co-moving wave-number and λ the wavelength of the perturbations. In GR, with normal forms of matter, one would obtain a closed second-order equation of the density fluctuations and the equations are generally easier to solve. But here we get a coupled system of second-order equations for the density fluctuations of both matter and CG, the solutions to which are more complicated to compute analytically. As such, we consider short-wavelength $(k/aH \gg 1)$ and long-wavelength $(k/aH \ll 1)$ limits of the perturbations [8] and analyse large-scale structure implications of the resulting solutions.

² w_m is the equation of state parameter for the matter fluid which takes the value of 1/3 for the radiation component (w_r) and 0 for the dust component (w_d) , and $w_t = p_t/\rho_t = p_t/(\rho_{ch} + \rho_m)$.

In redshift space, the corresponding equations can respectively be found to be 3

$$\Delta_m'' = \frac{1}{1+z} \left(\frac{1}{2} - 4w_m\right) \Delta_m' + \left\{\frac{(1+w_m)}{(1+z)^2} \left[\frac{1}{2}(1+3w_m) + \frac{w_m}{(\Omega_{ch} + \Omega_m)(1+w_t)} + \frac{w_m(1+3w_t)}{2(1+w_t)} + \frac{9\pi^2(1+w_m)^2w_m}{3\lambda^2(1+z)^{3(1+w_m)}(\Omega_{ch} + \Omega_m)}\right] 3\Omega_m + \left[3w_m - \frac{3}{2}w_m(\Omega_{ch} + \Omega_m)(1+w_t)\right] \Delta_m^k + \frac{(1+w_m)}{(1+z)^2} \left[\frac{1}{2} - \frac{3\alpha w_{ch}}{2} - \frac{\alpha w_{ch}}{(\Omega_{ch} + \Omega_m)(1+w_t)} - \frac{\alpha w_{ch}(1+3w_t)}{2(1+w_t)} + \frac{9\pi^2(1+w_m)^2\alpha w_{ch}}{3\lambda^2(1+z)^{3(1+w_m)}(\Omega_{ch} + \Omega_m)}\right] 3\Omega_{ch}\Delta_{ch}^k , \quad (4)$$

$$\Delta_{ch}^{\prime\prime} = \frac{3\sqrt{4} - 6}{\sqrt{4}(1+z)} \left(w_{ch}(1+\alpha) - w_{ch} + \frac{2}{3} \right) \Delta_{ch}^{\prime} + \frac{(1+w_{ch})}{(1+z)^2} \left[\frac{1}{2} (1+3w_m) + \frac{w_m}{3(\Omega_m + \Omega_{ch})(1+w_t)} + \frac{w_m(1+3w_t)}{3\lambda^2(1+z)^{3(1+w_m)}(\Omega_{ch} + \Omega_m)} \right] 3\Omega_m \Delta_m^k + \frac{1}{(1+z)^2} \left\{ (1+w_{ch}) \left[\frac{1}{2} - \frac{3\alpha w_{ch}}{2} - \frac{\alpha w_{ch}}{(\Omega_m + \Omega_{ch})(1+w_t)} - \frac{\alpha w_{ch}(1+3w_t)}{2(1+w_t)} + \frac{9\pi^2(1+w_m)^2 \alpha w_{ch}}{\lambda^2(A+B(1+z)^{3(1+\alpha)})^{\frac{1}{1+\alpha}}(\Omega_{ch} + \Omega_m)} \right] 3\Omega_{ch} - w_{ch} \left(3 + \frac{3}{2} (1+3w_t)(\Omega_m + \Omega_{ch}) \right) + 9w_{ch} \left[(1+\alpha)(1+2w_{ch})3\Omega_{ch} + \frac{2}{3} \right] \right\} \Delta_{ch}^k.$$
(5)

3. Results and Discussion

For further discussion of the growth of the energy density fluctuations with redshift, we assume non-interacting fluids, namely radiation-CG, dust-CG, and only CG fluids in the following subsections.⁴

3.1. GR and ΛCDM approaches

We first consider the well-known results of GR and ACDM approaches. For the case where the entire universe has been filled with dust and radiation fluids, the evolution equation becomes

$$\frac{d^2 \Delta_m(z)}{dz^2} - \frac{1}{2(1+z)} \frac{d\Delta_m(z)}{dz} - \frac{1}{(1+z)^2} \left[\frac{3}{2} \Omega_m \left(1+3w_m \right) (1-w_m) + 6w_m \Omega_\Lambda - w_m k^2 (1+z)^2 \right] \Delta_m(z) = 0$$
(6)

By eliminating the cosmological constant term, Λ , the Λ CDM model reduces to GR as shown in [9]. For illustrative purpose, we redefined the normalized energy density for fluid as $\delta(z) = \Delta(z)$ (as was done in [9]), and $\delta_{in} = \Delta(z_{in})$ is the initial value of the density contrast, $\Delta(z)$, at $z_{in} = 1100$.

3.2. Radiation-CG dominated Universe

The EoS parameter of the total fluid consisting of radiation (implying that $w_m = w_r$) and CG is given as

$$w_t = \frac{\rho_{0,r}(1+z)^4 - 3\frac{A}{\sqrt{A+B(1+z)^6}}}{3\rho_{0,r}(1+z)^4 + 3\sqrt{A+B(1+z)^6}},$$
(7)

where $\rho_{0,r}$ is the initial energy density of radiation at the time of the Big Bang. In the early universe, the redshift is large and CG acts as dark matter, whereas, in the late universe (where redshift is small and negative) the CG behaves like dark energy. The EoS parameter, w_t ,

³ A single dot () and double dots () represent the first- and second-order derivatives, respectively, with respect to time, t, and a single prime (') and double prime (") represent the first- and second-order derivatives, respectively, with respect to redshift, z.

⁴ For all analysis, we consider the original CG model, $\alpha = 1$.



Figure 1. $\delta(z)$ versus z for dust for GR.



Figure 3. $\delta(z)$ versus z for long wavelengths for Λ CDM.



Figure 2. $\delta(z)$ versus z for short wavelengths for Λ CDM with different wavelength.

The numerical results for a dust dominated universe, and a radiation dominated universe are presented in Fig. 1, and Fig. 2 and 3, for short and long-wavelength modes, respectively. From the plots, we observe the growth of matter density fluctuations with cosmic-time. For shortwavelength, the amplitude has an oscillatory behavior. The variation of CMB temperature detected observationally is of the order of 10^{-3} [10]. Therefore, we set the initial conditions at $\delta_{in} = \Delta_i(z_{in} \simeq 1100) = 10^{-3}$ and $\dot{\delta}_{in} = \dot{\Delta}_i(z_{in} = 1100) = 0$. The normalized energy densities are defined as $\Omega_m + \Omega_{\Lambda} = 1$.

approaches -1 when $z \ll 1$ (which means that the dominant component in late times is dark energy, corresponding to our Λ CDM model predictions). Now, taking large redshift, $z \gg 1$, then $w_t \approx 1/3 = w_r$ (corresponding to a dominant radiation component in early times). If we assume a dominant radiation fluid, and background CG fluid, we have $\Delta_r \gg \Delta_{ch}$, i.e., Δ_{ch} is very small. Similar analysis is done in [8] for a dust-radiation system. Then the solutions of Eq. (4), for the short-wavelength range, are found to be

$$\Delta(z) = C_1(1+z)^{\frac{1}{12}} \text{BesselJ}\left(\frac{\Sigma}{24(\Omega_{ch}+\Omega_r)}, \frac{4\pi\sqrt{\Omega_r}}{3\lambda\sqrt{\Omega_{ch}+\Omega_r}(1+z)^2}\right) + C_2(1+z)^{\frac{1}{12}} \text{BesselY}\left(\frac{\Sigma}{24(\Omega_{ch}+\Omega_r)}, \frac{4\pi\sqrt{\Omega_r}}{3\lambda\sqrt{\Omega_{ch}+\Omega_r}(1+z)^2}\right),$$
(8)

where $\Sigma = \sqrt{624\Omega_r^2 - 96\Omega_{ch}^2 + 528\Omega_{ch}\Omega_r + 145\Omega_{ch} + 289\Omega_r(\Omega_{ch} + \Omega_r)}$, C_1 and C_2 are integration constants and BesselJ and BesselY are the first- and second-order Bessel functions. The solution of Eq. (4) for the long-wavelength range is given as

$$\Delta(z) = C_3 (1+z)^{\Omega_{ch} + \Omega_r + \frac{\psi}{12(\Omega_{ch} + \Omega_r)}} + C_4 (1+z)^{-(\Omega_{ch} + \Omega_r - \frac{\psi}{12(\Omega_{ch} + \Omega_r)})}, \qquad (9)$$

where $\psi = \sqrt{432\Omega_{ch}^2\Omega_r + 1152\Omega_{ch}\Omega_r^2 - 96\Omega_{ch}^3 + 624\Omega_r^3 + 145\Omega_{ch}^2 + 289\Omega_r^2 + 434\Omega_{ch}\Omega_r}$. We assume initial conditions given as $\Delta_{in} \equiv \Delta^k(z_{in} = 1100) = 10^{-3}$ and $\Delta'_{in} \equiv \Delta'(z_{in} = 1100) = 0$ for every mode, k, to deal with the growth of matter fluctuations [8]. Therefore, we determine the integration constant C_i (i = 1, 2, 3, 4, ...) by imposing these initial conditions. The numerical result of Eq. (8) is presented in Fig. 4. In this figure we observe the oscillatory motions of the density perturbations in the short-wavelength modes. The solutions for for the long-wavelength modes (given by Eq. (9)) are shown in Fig. 5. In this figure, we clearly observe the growth of the density fluctuations. We used $\Omega_r = 4.48 \times 10^{-5}$ [11].



Figure 4. $\delta(z)$ versus z for short wavelengths for radiation-CG fluids.



Figure 5. $\delta(z)$ versus z for long wave-lengths for radiation-CG fluids.

3.3. Dust-CG dominated Universe

If we consider a total fluid which consists of dust (implying that $w_m = w_d$) and CG, then the EoS parameter of the total fluid becomes

$$w_t = \frac{-\frac{A}{\sqrt{A+B(1+z)^6}}}{\rho_{0,d}(1+z)^3 + \sqrt{A+B(1+z)^6}}.$$
(10)

For large values of redshift, $z \gg 1$ (i.e. in the early universe), $w_t \approx 0$ (corresponding to a dominant dust component in early times). When $z \ll 1$, the EoS parameter becomes $w_t \approx -1$, which again corresponds to dark energy being the dominant component at late times. The solutions of our evolution equations for the density perturbations, which are given by Eq. (4), in a dust-CG system is presented graphically in Fig. 6 and reads

$$\Delta(z) = C_5 (1+z)^{\frac{3}{4} + \frac{1}{4}\sqrt{9 + 24\Omega_d}} + C_6 (1+z)^{\frac{3}{4} - \frac{1}{4}\sqrt{9 + 24\Omega_d}} .$$
⁽¹¹⁾



Figure 6. $\delta(z)$ versus z for dust-CG fluids and for $\Omega_d = 1 - \Omega_{ch}$. We used $\Omega_d = 0.32$ [12].



Figure 7. $\delta(z)$ versus z for CG alone, using $1 - \Omega_r - \Omega_d = \Omega_{ch}$.

3.4. CG-dominated Universe

When the energy density of the CG fluid is dominant over that of the matter fluid, we can let $\Delta_{ch} \gg \Delta_m$, causing Δ_m to be very small. The solution of Eq. (5) then reads as

$$\Delta(z) = C_7 \log(1+z) \sin(\Omega_r + \Omega_d + \Omega_{ch} + 5) + C_8 \log(1+z) \cos(\Omega_r + \Omega_d + \Omega_{ch} + 5) .$$
(12)

The numerical results are presented in Fig. 7 and depicts the growth of the CG density fluctuations in terms of redshift, and the role of CG fluid for the formation of large-scale

structure. In this paper we observe that CG acts as a dark energy and it is an alternative approach to explaining the current phase of universe. From all numerical results, we observe that the CG component becomes dominant over time and the growth of matter density contrast is fast growing with cosmic time compared to the matter-CG fluid system, since the Universe becomes less structured in the accelerating universe. We think our current results show that even at the level of the perturbations, the CG fluid offers an alternative to the narrative of large-scale structure formation. This is because, contrary to what one would expect in a darkenergy-dominated universe where there would be less chance of large-scale structure formation due to the rapid cosmological expansion, we see the growth of the density perturbations with time.

4. Conclusions

In this work, we explored the solutions of cosmological perturbations in a multi-fluid cosmic medium where one of the fluids is a Chaplygin gas. We applied the 1 + 3 covariant and gauge-invariant formalism to define the spatial gradient variables and applied scalar and harmonic decomposition methods to analyse the scalar perturbations of the different energy densities involved. We considered different systems such as radiation-CG, dust-CG and CG fluids in both short- and long-wavelength modes to present the numerical and analytical solutions to the perturbation equations. Our results show that, at least in the simplest CG model, the formation of large-scale structures is enhanced, rather than discouraged (as one would expect from dark energy fluid models), since all our preliminary calculations show the growth of density fluctuations with time.

Acknowledgments

SS gratefully acknowledges financial support from WKU, EORC and ESSTI, as well as the hospitality of the Physics Department of North-West University (NWU) during the preparation of part of this manuscript. HS acknowledges the financial support from the Mwalimu Nyerere African Union scholarship and the National Research Foundation (NRF) free-standing scholarship. TT has been funded through the National Research Foundation (NRF) free-standing scholarship. AS acknowledges funding through the National Astrophysics and Space Science Program (NASSP) scholarship. ME acknowledges that this work is supported by an NWU/NRF postdoctoral fellowship. AA acknowledges that this work is based on the research supported in part by the NRF of South Africa with grant numbers 109257 and 112131.

- [1] Riess A G et al. 1998 The Astronomical Journal **116** 1009
- [2] Fall S M and Efstathiou G 1980 Monthly Notices of the Royal Astronomical Society 193 189–206
- [3] Blumenthal G R, Faber S, Primack J R and Rees M J 1984 Nature 311 517
- [4] Rees M J 1980 Physica Scripta 21 614
- [5] Bento M, Bertolami O and Sen A A 2002 Physical Review D 66 043507
- [6] Ellis G F, Hwang J and Bruni M 1989 Physical Review D 40 1819
- [7] Dunsby P K 1991 Class. Quant. Grav. 8 1785
- [8] Abebe A, Abdelwahab M, De la Cruz-Dombriz Á and Dunsby P K 2012 Class. Quant. Grav. 29 135011
- [9] Sahlu S et al. 2020 European Physical Journal C 80 1-19
- [10] Smoot G F, Bennett C L, Kogut A, Wright E, Aymon J, Boggess N, Cheng E, De Amici G, Gulkis S, Hauser M et al. 1992 The Astrophysical Journal 396 L1–L5
- [11] Chavanis P H 2015 Physical Review D 92 103004
- [12] Ade P A, Aghanim N, Arnaud M, Ashdown M, Aumont J, Baccigalupi C, Banday A, Barreiro R, Bartlett J, Bartolo N et al. 2016 Astronomy & Astrophysics 594 A13