

# CRITICAL 2D ZAKHAROV-KUZNETSOV EQUATION POSED ON A HALF-STRIP

N. A. LARKIN

DEPARTAMENTO DE MATEMÁTICA,  
UNIVERSIDADE ESTADUAL DE MARINGÁ  
87020-900, MARINGÁ, PARANA, BRAZIL

**ABSTRACT.** An initial-boundary value problem for the critical generalized 2D Zakharov-Kuznetsov equation posed on the right half-strip is considered. Existence, uniqueness and exponential decay rate of global regular solutions for small initial data are established.

## 1. INTRODUCTION

We are concerned with an initial-boundary value problem (IBVP) for the critical Zakharov-Kuznetsov (ZK) equation posed on the right half-strip

$$u_t + u^2 u_x + u_{xxx} + u_{xyy} = 0 \quad (1.1)$$

which is a two-dimensional analog of the generalized Korteweg-de Vries (KdV) equation

$$u_t + u^k u_x + u_{xxx} = 0 \quad (1.2)$$

with plasma physics applications [38] that has been intensively studied last years [4, 7, 12, 13].

Equations (1.1) and (1.2) are typical examples of so-called dispersive equations attracting considerable attention of both pure and applied mathematicians. The KdV equation is more studied in this context. The theory of the initial-value problem (IVP henceforth) for (1.2) is considerably advanced today [3, 14, 15, 34].

Although dispersive equations were deduced for the whole real line, necessity to calculate numerically the Cauchy problem approximating the real line by finite intervals implies to study initial-boundary value problems posed on bounded and unbounded intervals [4, 5, 7, 23, 24, 36, 37]. What concerns (1.2) with  $k > 1$ ,  $l = 1$ , called generalized KdV equations, the Cauchy problem was studied in [31, 32] and later in [8, 10, 11, 15], where it has been established that for  $k = 4$  (the critical case) the problem is well-posed for small initial data, whereas

---

2020 *Mathematics Subject Classification.* 35M20, 35Q72.

*Key words and phrases.* ZK equation, stabilization.

email:nlarkine@uem.br.

for arbitrary initial data solutions may blow-up in a finite time. The generalized Korteweg-de Vries equation was studied for understanding the interaction between the dispersive term and the nonlinearity in the context of the theory of nonlinear dispersive evolution equations [12, 13, 15]. In [29], the initial-boundary value problem for the generalized KdV equation with an internal damping posed on a bounded interval was studied in the critical case; exponential decay of weak solutions for small initial data has been established. In [2], decay of weak solutions in the case  $l = 2$ ,  $k = 2$  has been established.

Recently, due to physics and numerics needs, publications on initial-boundary value problems in both bounded and unbounded domains for dispersive equations have been appeared [23, 24, 27, 28, 33, 34]. In particular, it has been discovered that the KdV equation posed on a bounded interval possesses an implicit internal dissipation. This allowed to prove the exponential decay rate of small solutions for (1.2) with  $k = 1$  posed on bounded intervals without adding any artificial damping term [4]. Similar results were proved for a wide class of dispersive equations of any odd order with one space variable [7, 23, 24].

The interest on dispersive equations became to extend their study for multi-dimensional models such as Kadomtsev-Petviashvili (KP) and ZK equations. We call (1.1) a critical ZK equation by analogy with the critical KdV equation (1.2) for  $k = 4$ . It means that we could not prove the existence and uniqueness of global regular solutions without smallness restrictions for initial data similarly to the critical case for the KdV equation [8, 10, 11, 24, 31, 32]. As far as the ZK equation is concerned, the results on both IVP and IBVP can be found in [5, 6, 8, 27, 28, 30]. Our work has been inspired by [29] where critical KdV equation with internal damping posed on a bounded interval was considered and exponential decay of weak solutions has been established. We must note that solvability of initial-boundary value problems in classes of global regular solutions for the regular case of the 2D ZK equation ( $uu_x$ ) has been established in [4, 6, 18, 19, 22, 26, 30, 36, 37] for arbitrary smooth initial data. On the other hand, for the 3D ZK equation, the convective term  $uu_x$ , which is regular for the 2D ZK equation, corresponds to a critical case. It means that to prove the existence and uniqueness of global regular solutions one must put restrictions of small initial data [20, 21, 25].

The main goal of our work is to prove for small initial data the existence and uniqueness of global-in-time regular solutions for (1.1) posed on bounded rectangles and the exponential decay rate of these solutions.

The paper is outlined as follows: Section I is the Introduction. Section 2 contains formulation of the problem and auxiliaries. In Section 3, Galerkin's approximations are used to prove the existence and exponential decay of strong solutions. In Section 4, regularity of strong solutions their uniqueness and decay are established.

## 2. PROBLEM AND PRELIMINARIES

Let  $(x, y) \equiv (x_1, x_2) \in \Omega$  and  $\Omega$  be a domain in  $\mathbb{R}^2$ . We use the usual notations of Sobolev spaces  $W^{k,p}$ ,  $L^p$  and  $H^k$  and the following notations for the norms [1]:

$$\|f\|_{L^p(\Omega)}^p = \int_{\Omega} |f|^p d\Omega, \quad \|f\|_{W^{k,p}(\Omega)} = \sum_{0 \leq |\alpha| \leq k} \|D^\alpha f\|_{L^p(\Omega)}, \quad p \in (1, +\infty).$$

$$\|f\|_{L^\infty(\Omega)} = \text{ess sup}_\Omega |f(x, y)|.$$

Let

$$\mathcal{D}(\Omega) = \{f \in C^\infty(\Omega); \text{supp } f \text{ is a compact set of } \Omega, \}$$

$$\mathbb{R}^+ = \{t \in \mathbb{R}, \quad t > 0.\}$$

The closure of  $\mathcal{D}(\Omega)$  in  $W^{k,p}(\Omega)$  is denoted by  $W_0^{k,p}(\Omega)$ ,  $H_0^k(\Omega)$  when  $p = 2$ .

Let  $L, B$  be finite positive numbers. Define

$$D = \{(x, y) \in \mathbb{R}^2 : x \in (0, L), y \in (0, B)\}, \quad Q = D \times \mathbb{R}^+;$$

$$\gamma = \partial D \text{ is a boundary of } D.$$

Consider the following IBVP:

$$Au \equiv u_t + u^2 u_x + u_{xxx} + u_{xyy} = 0, \quad \text{in } Q; \quad (2.1)$$

$$u_{\gamma \times t} = 0, \quad t > 0; \quad (2.2)$$

$$u_x(L, y, t) = 0, \quad y \in (0, B), \quad t > 0; \quad (2.3)$$

$$u(x, y, 0) = u_0(x, y), \quad (x, y) \in D, \quad (2.4)$$

where  $u_0 : D \rightarrow \mathbb{R}$  is a given function.

Hereafter subscripts  $u_x$ ,  $u_{xy}$ , etc. denote the partial derivatives, as well as  $\partial_x$  or  $\partial_{xy}^2$  when it is convenient. Operators  $\nabla$  and  $\Delta$  are the gradient and Laplacian acting over  $D$ . By  $(\cdot, \cdot)$  and  $\|\cdot\|$  we denote the inner product and the norm in  $L^2(D)$ , and  $\|\cdot\|_{H^k(D)}$  stands for the norm in  $L^2$ -based Sobolev spaces.

We will need the following result [16, 17].

**Lemma 2.1.** *Let  $u \in H^1(D)$  and  $\gamma$  be the boundary of  $D$ .*

*If  $u|_\gamma = 0$ , then*

$$\|u\|_{L^q(D)} \leq \beta \|\nabla u\|^\theta \|u\|^{1-\theta}. \quad (2.5)$$

*We will use frequently the following inequalities:*

$$\|u\|_{L^4(D)} \leq 2^{1/2} \|\nabla u\|^{1/2} \|u\|^{1/2}, \quad \|u\|_{L^8(D)} \leq 4^{3/4} \|\nabla u\|^{3/4} \|u\|^{1/4}.$$

*If  $u|_\gamma \neq 0$ , then*

$$\|u\|_{L^q(D)} \leq C_D \|u\|_{H^1(D)}^\theta \|u\|^{1-\theta}, \quad (2.6)$$

*where  $\theta = 2(\frac{1}{2} - \frac{1}{q})$ .*

**Lemma 2.2.** *Let  $v \in H_0^1(0, L)$ . Then*

$$\|v_x\|^2 \geq \frac{\pi^2}{L^2} \|v\|^2. \quad (2.7)$$

*Proof.* The proof is based on the Steklov inequality [35]: let  $v(t) \in H_0^1(0, \pi)$ , then by the Fourier series  $\int_0^\pi v_t^2(t) dt \geq \int_0^\pi v^2(t) dt$ . Inequality (2.7) follows by a simple scaling.  $\square$

**Proposition 2.1.** *Let for a.e. fixed  $t$   $u(x, y, t) \in H_0^1(D)$  and  $u_{xy}(x, y, t) \in L^2(D)$ . Then*

$$\begin{aligned} \sup_{(x,y) \in D} u^2(x, y, t) &\leq 2 \left[ \|u\|_{H_0^1(D)}^2(t) + \|u_{xy}\|_{L^2(D)}^2(t) \right] \\ &\leq 2 \|u\|^2(t)_{H^2(D) \cap H_0^1(D)}. \end{aligned} \quad (2.8)$$

*Proof.* For a fixed  $x \in (0, L)$  and for any  $y \in (0, B)$ , it holds

$$\begin{aligned} u^2(x, y, t) &= \int_0^y \partial_s u^2(x, s, t) ds \leq \int_0^B u^2(x, y, t) dy + \int_0^B u_y^2(x, y, t) dy \\ &\equiv \rho^2(x, t). \end{aligned}$$

On the other hand,

$$\begin{aligned} \sup_{(x,y) \in \mathcal{D}} u^2 &\leq \sup_{x \in (0, L)} \rho^2(x) = \sup_{x \in (0, L)} \left| \int_0^x \partial_s \rho^2(s) ds \right| \\ &\leq 2 \int_0^L \int_0^B (u^2 + u_x^2 + u_y^2 + u_{xy}^2) dx dy \leq 2 \|u\|_{H^2(D) \cap H_0^1(D)}^2. \end{aligned}$$

The proof of Proposition 2.1 is complete.  $\square$

## 3. EXISTENCE THEOREM

Define the space  $W(D)$  with the norm

$$\|u\|_{W(D)} = \|u\|_{H^2(D) \cap H_0^1(D)} + \|\Delta u_x\|.$$

**Theorem 3.1.** *Given  $u_0 \in W(D)$  and  $D$  such that  $u_0|_\gamma = u_{0x}|_{x=L} = 0$  and*

$$\|u_0\| < \min\left(\frac{1}{2}, m\right), \quad (3.1)$$

where

$$\begin{aligned} m &< \left( \frac{\pi^2}{4(1+L)^2} \left( \frac{1}{L^2} + \frac{1}{B^2} \right) \left[ 5 \times 2^7 (1+L) \|u_t(0)\| (1 \right. \right. \\ &\quad \left. \left. + 5^2 \times 2^{15} (1+L)^6 \|u_t(0)\|) \right]^{-1} \right)^{1/3}, \\ \|u_t(0)\| &\leq \|\Delta u_{0x}\| + \|u_0^2 u_{0x}\| \leq C(\|u_0\|_{H^2(D) \cap H_0^1(D)}). \end{aligned}$$

Then for all finite positive  $B, L$  there exists a unique regular solution to (2.1)-(2.4) such that

$$\begin{aligned} u &\in L^\infty(\mathbb{R}^+; H^2(D)) \cap L^2(\mathbb{R}^+; H^3(D)); \\ \Delta u_x &\in L^\infty(\mathbb{R}^+; L^2(D)) \cap L^2(\mathbb{R}^+; H^1(D)); \\ u_t &\in L^\infty(\mathbb{R}^+; L^2(D)) \cap L^2(\mathbb{R}^+; H^1(D)) \end{aligned}$$

and

$$\begin{aligned} \|u\|_{H^2(D)}^2(t) + \|\Delta u_x\|^2(t) + \|u_t\|^2(t) &\leq C(\|u_0\|_{W(D)}) e^{(-\chi t)}, \quad t > 0, \\ \text{where } \chi &= \frac{\pi^2}{2(1+L)} \left[ \frac{5}{L^2} + \frac{1}{B^2} \right]; \end{aligned} \quad (3.2)$$

$$\begin{aligned} \int_{\mathbb{R}^+} \left\{ \|u\|_{H^3(D)}^2(t) + \|\Delta u_x\|_{H^1(D)}^2(t) + \|u_x(0, y, t)\|_{H^2(0, B)}^2 \right\} dt \\ \leq C(\|u_0\|_{W(D)}, L, B). \end{aligned} \quad (3.3)$$

To prove this theorem, we will use the Faedo-Galerkin approximations. Let  $w_j(y)$  be orthonormal in  $L^2(D)$  eigenfunctions to the following Dirichlet Problem:

$$w_{jyyy} + \lambda_j w_j = 0, \quad y \in (0, B); \quad w_j(0) = w_j(L) = 0; \quad j \in \mathbb{N}. \quad (3.4)$$

Define approximate solutions of (2.1)-(2.4) in the form:

$$u^N(x, y, t) = \sum_{j=1}^N g_j^N(x, t) w_j(y) \quad (3.5)$$

and  $g_j^N(x, t)$  are solutions to the following Korteweg-de Vries system:

$$g_{jt}^N + g_{jxxx}^N - \lambda_j g_{jx}^N + \int_0^B |u^N|^2 u_x^N w_j(y) dy = 0, \quad (3.6)$$

$$g_j^N(0, t) = g_j^N(L, t) = g_{jx}^N(L, t) = 0; \quad t > 0, \quad (3.7)$$

$$g_j^N(x, 0) = (u_{0N}, w_j), \quad x \in (0, L), \quad (3.8)$$

where  $u_{0N} = \sum_{i=1}^N \alpha_{iN} w_i \rightarrow u_0$  in  $W(D) \cap H_0^1(D)$ ,  $j = 1, \dots, N$ .

Since each regularized KdV equation from (3.6) is not critical, it is known [23, 24, 36, 37] that there exists a unique regular solution of (3.5)-(3.8) at least locally in  $t$ .

Our goal is to obtain global in  $t$  a priori estimates for the  $u^N$  independent of  $t$  and  $N$ , then to pass the limit as  $N$  tends to  $\infty$  getting a solution to (2.1)-(2.4).

**Lemma 3.1.** *Under the conditions of Theorem 3.1, the following independent of  $N$  and  $t$  estimates hold:*

$$u^N \text{ is bounded in } L^\infty(\mathbb{R}^+; L^2(D)) \cap L^2(\mathbb{R}^+; H^1(D)) \quad (3.9)$$

and

$$\begin{aligned} \|u^N\|^2(t) &\leq ((1+x)^{1/2}, u^N)^2(t) \leq ((1+x), u_0^2) e^{(-\chi t)} \\ &\leq \frac{1+L}{4} e^{(-\chi t)}, \text{ where } \chi = \frac{\pi^2}{2(1+L)} \left[ \frac{5}{L^2} + \frac{1}{B^2} \right]. \end{aligned} \quad (3.10)$$

*Proof. Estimate I.* Multiply (3.6) by  $g_j^N$ , sum up over  $j = 1, \dots, N$  and integrate over  $\Omega \times (0, t)$  to obtain

$$\begin{aligned} \|u^N\|^2(t) + \int_0^t \int_0^B (u_x^N)^2(0, y, \tau) dy d\tau \\ = \|u_0^N\|^2 \leq \|u_0\|^2, \quad t > 0. \end{aligned} \quad (3.11)$$

**Estimate II.** Write the inner product

$$2(Au^N, (1+x)u^N)(t) = 0,$$

dropping the index  $N$ , in the form:

$$\begin{aligned} \frac{d}{dt} ((1+x), u^2)(t) + \int_0^B u_x^2(0, y, t) dy + 3\|u_x\|^2(t) + \|u_y\|^2(t) \\ = \frac{1}{2} \int_{\mathcal{D}} u^4 dx dy. \end{aligned}$$

Taking into account (2.5) and (3.11), we obtain

$$\frac{1}{2} \int_{\mathcal{D}} u^4 dx dy \leq \frac{1}{2} \|u\|_{L^4(\mathcal{D})}^4(t) \leq 2 \|\nabla u\|^2(t) \|u_0\|^2(t).$$

This implies

$$\begin{aligned} \frac{d}{dt} ((1+x), u^2)(t) + \frac{1}{2} \|\nabla u\|^2(t) + \left(\frac{1}{2} - 2\|u_0\|^2\right) \|\nabla u\|^2(t) \\ + 2\|u_x\|^2(t) + \int_0^B u_x^2(0, y, t) dy \leq 0. \end{aligned} \quad (3.12)$$

Making use of (3.1) and Lemma 2.2, we get

$$\frac{d}{dt} ((1+x), u^2)(t) + \frac{\pi^2}{2(1+L)} \left[ \frac{5}{L^2} + \frac{1}{B^2} \right] ((1+x), u^2)(t) \leq 0.$$

This gives

$$\begin{aligned} \|u^N\|^2(t) &\leq ((1+x)^{1/2}, u^N)^2(t) \leq ((1+x), u_0^2) e^{(-\chi t)} \\ &\leq \frac{1+L}{4} e^{(-\chi t)}. \end{aligned} \quad (3.13)$$

Returning to (3.12), we obtain

$$\begin{aligned} ((1+x), |u^N|^2)(t) + \int_0^t \int_0^B |u_x^N|^2(0, y, \tau) dy d\tau + \frac{1}{2} \int_0^t \|\nabla u^N\|^2(\tau) d\tau \\ \leq ((1+x), u_0^2), \quad t > 0. \end{aligned} \quad (3.14)$$

Moreover, we can rewrite (3.12) as

$$\begin{aligned} 4\|u_x^N\|^2(t) + \|\nabla u^N\|^2(t) + \int_0^B |u_x^N|^2(0, y, t) dy \leq 4|((1+x)u, u_t^N)(t)| \\ \leq 4\|(1+x)^{1/2}u^N\|(t)\|(1+x)^{1/2}u_t^N\|(t). \end{aligned} \quad (3.15)$$

The proof of Lemma 3.1 is complete.  $\square$

**Lemma 3.2.** *Under the conditions of Theorem 3.1, the following inequalities are true:*

$$((1+x), |u_t^N|^2)(t) \leq C_0 e^{(-\chi t)}, \quad (3.16)$$

$$\begin{aligned} ((1+x), |u_t^N|^2)(t) + \int_0^t \int_0^B (\partial_{x\tau}^2 u^N)^2(0, y, \tau) dy d\tau \\ + \frac{1}{2} \int_0^t \|\nabla \partial_\tau u^N\|^2(\tau) d\tau \leq C_0, \quad t > 0. \end{aligned} \quad (3.17)$$

It follows from (2.1) that

$$\begin{aligned} C_0 &= (1+x, |u_t^N|^2)(0) \leq (1+L)\|u_t^N(0)\|^2 \\ &\leq (1+L)\left[\|\Delta u_{0x}\| + \|u_0^2 u_{0x}\|\right]^2. \end{aligned} \quad (3.18)$$

*Proof.* Making use of (2.8) for  $t = 0$ , we get

$$\sup_D u_0^2(x, y) = \sup_D u^2(x, y, 0) \leq 2(\|u_0\|_{H_0^1(D)}^2 + \|u_{0xy}\|^2) \equiv C_s. \quad (3.19)$$

Substituting (3.19) into (3.18), we find

$$(1+x, |u_t^N|^2)(0) \leq C_0(L, C_s, \|u_0\|_{W(D)}). \quad (3.20)$$

### Estimate III

Dropping the index  $N$ , write the inner product

$$2\left((1+x)u_t^N, \partial_t(Au^N)\right)(t) = 0$$

as

$$\begin{aligned} &\frac{d}{dt} \left((1+x), u_t^2\right)(t) + \int_0^B u_{xt}^2(0, y, t) dy + 3\|u_{xt}\|^2(t) + \|u_{yt}\|^2(t) \\ &= 2\left((1+x)u^2 u_t, u_{xt}\right)(t) + 2(u^2, u_t^2)(t). \end{aligned} \quad (3.21)$$

Making use of Lemmas 2.1, 2.2, (3.15) and taking into account the first inequality of (3.1), we estimate

$$\begin{aligned} I_1 &= 2\left((1+x)u^2 u_t, u_{xt}\right)(t) \leq 2(1+L)\|u_{xt}\|(t)\|u\|^2(t)_{L^8(D)}\|u_t\|(t)_{L^4(D)} \\ &\leq 2^{9/2}(1+L)\|u_{xt}\|(t)\|u_0\|^{1/2}\|\nabla u\|^{3/2}(t)\|u_t\|^{1/2}(t)\|\nabla u_t\|^{1/2}(t) \\ &\leq 2^{9/2}(1+L)\|\nabla u_t\|^{3/2}(t)\|u_0\|^{1/2}\|\nabla u\|^{3/2}(t)\|u_t\|^{1/2}(t) \\ &\leq \frac{3\delta}{4}\|\nabla u_t\|^2(t) + \frac{2^{16}(1+L)^4}{\delta^3}\|u\|^2(t)\|\nabla u\|^6(t)\|u_t\|^2(t) \\ &\leq \frac{3\delta}{4}\|\nabla u_t\|^2(t) + \frac{2^{22}(1+L)^{11/2}}{\delta^3}\|u_0\|^5\left((1+x), u_t^2\right)^{3/2}(t)\|u_t\|^2(t). \end{aligned}$$

Here and henceforth  $\delta$  is an arbitrary positive number. Similarly,

$$\begin{aligned} I_2 &= 2(u^2, u_t^2)(t) \leq 2\|u\|^2(t)_{L^4(D)}\|u_t\|^2(t)_{L^4(D)} \\ &\leq 2^3\|u\|(t)\|\nabla u\|(t)\|u_t\|(t)\|\nabla u_t\|(t) \\ &\leq \frac{\delta}{2}\|\nabla u_t\|^2(t) + \frac{2^5}{\delta}\|u\|^2(t)\|\nabla u\|^2(t)\|u_t\|^2(t) \\ &\leq \frac{\delta}{2}\|\nabla u_t\|^2(t) + \frac{2^7}{\delta}(1+L)^{1/2}\|u_0\|^3\|(1+x)^{1/2}u_t\|(t)\|(1+x), u_t^2)(t). \end{aligned}$$



Substituting  $I_1, I_2$  into (3.21), making use of Lemma 2.2, taking into account the first inequality of (3.1) and setting  $\delta = \frac{1}{5}$ , we come to the inequality

$$\begin{aligned} & \frac{d}{dt}((1+x), u_t^2)(t) + \int_0^B u_{xt}^2(0, y, t) dy + 2\|u_{xt}\|^2(t) + \frac{1}{2}\|\nabla u_t\|^2(t) \\ & + \left[ \frac{1}{4(1+L)}\pi^2\left(\frac{1}{L^2} + \frac{1}{B^2}\right) - 5 \times 2^7(1+L)^{1/2}\|u_0\|^3((1+x), u_t^2)^{1/2}(t) \right. \\ & \left. + 5^2 \times 2^{15}(1+L)^5\|u_0\|^2((1+x), u_t^2)(t) \right]((1+x), u_t^2)(t) \leq 0. \end{aligned} \quad (3.22)$$

Using (3.1), Lemma 2.2 and standard arguments, [7, 25], we obtain that

$$\begin{aligned} & \frac{1}{4(1+L)}\pi^2\left(\frac{1}{L^2} + \frac{1}{B^2}\right) - 5 \times 2^7(1+L)^{1/2}\|u_0\|^3((1+x), u_t^2)^{1/2}(t) \{1 \\ & + 5^2 \times 2^{15}(1+L)^5\|u_0\|^2((1+x), u_t^2)(t)\} > 0, \quad t > 0. \end{aligned}$$

Returning to (3.21) and using the Steklov inequalities (2.7), we can rewrite it as

$$\begin{aligned} & \frac{d}{dt}((1+x), u_t^2)(t) + \int_0^B u_{xt}^2(0, y, t) dy \\ & + \frac{\pi^2}{2(1+L)}\left[\frac{5}{L^2} + \frac{1}{B^2}\right]((1+x), u_t^2)(t) \leq 0. \end{aligned} \quad (3.23)$$

This implies

$$((1+x), u_t^2)(t) \leq ((1+x), u_t^2)(0)e^{-\chi t} \leq C_0 e^{(-\chi t)}. \quad (3.24)$$

Since (3.23) can be rewritten as

$$\frac{d}{dt}((1+x), u_t^2)(t) + \int_0^B u_{xt}^2(0, y, t) dy + \frac{1}{2}\|\nabla u_t\|^2(t) \leq 0,$$

integrating it we get

$$\begin{aligned} & ((1+x), u_t^2)(t) + \int_0^t \int_0^B (\partial_{x\tau}^2 u)^2(0, y, \tau) dy d\tau + \frac{1}{2} \int_0^t \|\nabla \partial_\tau u\|^2(\tau) d\tau \\ & \leq ((1+x), u_t^2)(0) = C_0. \end{aligned} \quad (3.25)$$

Inequalities (3.24), (3.25) complete the proof of Lemma 3.2.  $\square$

Returning to (3.15), we find

$$\|\nabla u^N\|^2(t) \leq C e^{(-\chi t)}, \quad (3.26)$$

where the constant  $C$  does not depend on  $N, t > 0$ .

**Lemma 3.3.** *Under the conditions of Theorem 3.1, the following inequality holds:*

$$\begin{aligned} & \|\nabla u_y^N\|^2(t) + \int_0^B |u_{xy}^N|^2(0, y, t) dy \\ & \leq C(L, \|u_0\|, \|u_t(0)\|) e^{(-\chi t)}. \end{aligned} \quad (3.27)$$

*Proof.* Multiplying  $j$ -th equation of (3.6) by  $\lambda_j$ , and summing up the results over  $J = 1, \dots, N$ , dropping the index  $N$ , we transform the inner product

$$-2 \left( (1+x) \partial_y^2 u, Au \right) (t) = 0$$

into the inequality

$$\begin{aligned} & 3\|u_{xy}\|^2(t) + \|u_{yy}\|^2(t) + \int_0^B u_{xy}^2(0, y, t) dy + \\ & = \frac{2}{3} \left( (1+x)(u^3)_x, u_{yy} \right) (t) + 2((1+x)u_{yy}, u_t)(t) \\ & = -\frac{2}{3} \left( (1+x)(u^3)_{yx}, u_y \right) (t) + 2((1+x)u_{yy}, u_t)(t) \\ & \leq \frac{2}{3} |(u^3)_y, u_y + (1+x)u_{xy}| + 2(1+L)\|u_t\|(t)\|u_{yy}\|(t) \\ & \leq \delta \|\nabla u_y\|^2(t) + \frac{(1+L)^2}{\delta} \|u_t\|^2(t) \\ & + 2(u^2, u_y^2)(t) + 2(1+x)u^2 u_y, u_{xy}(t). \end{aligned} \quad (3.28)$$

Making use of Lemma 2.1, we estimate

$$\begin{aligned} I_1 & \equiv 2(u^2, u_y^2)(t) \leq 2\|u\|_{L^4(D)}^2(t)\|u_y\|_{L^4(D)}^2(t) \\ & \leq 4\|u\|(t)\|\nabla u\|(t)C_D^2\|u_y\|(t)\|\nabla u_y\|(t) \\ & \leq \delta\|\nabla u_y\|^2(t) + \frac{4C_D^4}{\delta}\|u\|^2(t)\|\nabla u\|^4(t), \end{aligned}$$

$$\begin{aligned} I_2 & \equiv 2(1+x)u^2 u_y, u_{xy}(t) \leq 2(1+L)\|u_{xy}\|(t)\|u^2\|_{L^4(D)}(t)\|u_y\|_{L^4(D)} \\ & \leq 2^{3/2}C_D^{1/2}(1+L)\|u_{xy}\|(t)\|u_y\|^{1/2}(t)\|\nabla u_y\|^{1/2}(t)\|u\|_{L^8(D)}^2 \\ & \leq 2^{9/2}C_D^{1/2}(1+L)\|\nabla u_y\|^{3/2}(t)\|u_0\|^{1/2}\|\nabla u\|^{3/2}(t) \\ & \leq \frac{3\delta}{4}\|\nabla u_y\|^2(t) + \frac{2^{16}(1+L)^4C_D^2}{\delta^3}\|u_0\|^2\|\nabla u\|^6(t). \end{aligned}$$

Substituting  $I_1, I_2$  into (3.28), we transform it into the following inequality:

$$\begin{aligned} & \frac{1}{2} \|\nabla u_y\|^2(t) + \left(\frac{1}{2} - \frac{11\delta}{4}\right) \|\nabla u_y\|^2(t) + \int_0^B u_{xy}^2(0, y, t) dy \\ & \leq \frac{C}{\delta} \left[ (1+L)^2 \|u_t\|^2(t) + \|u_0\|^2 \|\nabla u\|^4(t) \right. \\ & \quad \left. + \frac{2^{16}}{\delta^2} (1+L)^4 \|u_0\|^2 \|\nabla u\|^6(t) \right]. \end{aligned} \quad (3.29)$$

Taking  $11\delta = 2$  and making use of (3.14), (3.15), we come to (3.27). The proof of Lemma 3.3 is complete.  $\square$

**Lemma 3.4.** *Under the conditions of Theorem 3.1, the following inequality holds:*

$$\begin{aligned} & \int_0^t \left[ \|\nabla u_{yy}^N\|^2(\tau) + \int_0^B |u_{xyy}^N(0, y, \tau)|^2 dy \right] d\tau \\ & \leq C(D, \|u_0\|, \|u_t\|(0), \|u_{yy}\|(0)), \quad t > 0. \end{aligned} \quad (3.30)$$

*Proof. Estimate IV*

Multiply each of the  $j$ -th equation of (3.6) by  $\lambda_j^2$ , sum up over  $j = 1, \dots, N$  and, dropping the index  $N$ , write the scalar product

$$2 \left( (1+x) \partial_y^4 u, Au \right) (t) = 0$$

in the form

$$\begin{aligned} & \frac{d}{dt} \left( (1+x), u_{yy}^2 \right) (t) + 3 \|\partial_y^2 u_x\|^2(t) + \|\partial_y^3 u\|^2(t) + \int_0^B u_{xyy}^2(0, y, t) dy \\ & = -\frac{2}{3} \left( (1+x) u_{yy}, (u^3)_{yyx} \right) (t). \end{aligned} \quad (3.31)$$

Denote

$$\begin{aligned} I &= -\frac{2}{3} \left( (1+x) u_{yy}, (u^3)_{yyx} \right) (t) = \frac{2}{3} \left( u_{yy}, (u^3)_{yy} \right) (t) \\ & \quad + \frac{2}{3} \left( (1+x) u_{xyy}, (u^3)_{yy} \right) (t) \equiv I_1 + I_2, \end{aligned}$$

where

$$I_1 = \frac{2}{3} \left( u_{yy}, (u^3)_{yy} \right) (t) = 4(u u_y^2, u_{yy})(t) + 2(u^2, u_{yy}^2)(t) = I_{11} + I_{12}.$$

By Proposition 2.1 and (3.13), (3.18), (3.26), (3.27),

$$\begin{aligned} \sup_{(x,y) \in D, t > 0} u^2(x, y, t) &\leq 2 \left[ \|u\|_{H_0^1(D)}^2(t) + \|u_{xy}\|_{L^2(D)}^2(t) \right] \\ &\leq C e^{(-\chi t)}, \quad t > 0. \end{aligned} \quad (3.32)$$

Then

$$\begin{aligned} I_{11} &= 2(u^2, u_{yy}^2)(t) \leq 2 \sup_{(x,y) \in D} |u(x, y, t)|^2 \|u_{yy}\|^2(t) \\ &\leq C \left( (1+x), u_{yy}^2 \right)(t) \leq C e^{(-\chi t)} \end{aligned}$$

and

$$\begin{aligned} I_{12} &= 2(u_{yy}, u u_y^2)(t) \leq 2 \sup_D |u(x, y, t)| \|u_{yy}\|(t) \|u_y\|_{L^4(D)}^2(t) \\ &\leq 2C_D^2 \sup_D |u(x, y, t)| \|u_{yy}\|(t) \|u_y\|(t) \|u_y\|_{H^1(D)}(t) \\ &\leq C e^{(-\chi t)}. \end{aligned}$$

Similarly,

$$\begin{aligned} I_2 &= \frac{2}{3}((1+x)u_{xyy}, (u^3)_{yy})(t) = 2((1+x)u_{xyy}, (u^2 u_y)_y)(t) \\ &\leq 4|((1+x)u_{xyy}, u u_y^2)(t)| + 2|((1+x)u_{xyy}, u^2 u_{yy})(t)| \\ &\leq 4|((1+x)u_{xyy}, u u_y^2)(t)| + 2 \sup_D u^2(x, y, t) |(1+x)u_{xyy}, u_{yy}(t)| \\ &\leq 4(1+L) \sup_D |u(x, y, t)| \|u_{xyy}\| \|u_y\|^2(t)_{L^4(D)} \\ &\quad + 2(1+L) \sup_D u^2(x, y, t) \|u_{yy}\|(t) \|u_{xyy}\|(t) \\ &\leq \delta \|u_{xyy}\|^2(t) + \frac{C(D)}{\delta} \left[ \sup u^2(x, y, t) \|u_y\|^2(t) \|\nabla u_y\|^2(t) \right. \\ &\quad \left. + \sup_D u^4(x, y, t) \|u_{yy}\|^2(t) \right]. \end{aligned}$$

Making use of Proposition 2.1, (3.31) and Lemmas 3.2, 3.3, we get

$$I_2 \leq \delta \|u_{xyy}\|^2(t) + \frac{C(L, \|u_0\|, \|u_t(0)\|)}{\delta} e^{(-\chi t)}.$$

Taking  $\delta = \frac{1}{2}$  and substituting  $I_1, I_2$  into (3.25), we find that

$$\begin{aligned} \frac{d}{dt} \left( (1+x), u_{yy}^2 \right)(t) &+ \frac{1}{2} \|\nabla u_{yy}\|^2(t) + \int_0^B u_{xyy}^2(0, y, t) dy \\ &\leq C(D, \|u_0\|, \|u_t(0)\|) e^{(-\chi t)}. \end{aligned}$$

Simple integration completes the proof of Lemma 3.4.  $\square$

Taking into account (3.3),(3.4), write (3.6) in the form

$$\int_0^B (u_{xxx}^N - \lambda_j u_x^N) w_j dy = - \int_0^B [u_t^N + |u^N|^2 u_x^N] w_j dy.$$

Multiplying it by  $g_{jxx} - \lambda_j g_{jx}$ , summing over  $j = 1, \dots, N$  and integrating with respect to  $x$  over  $(0, L)$ , we obtain

$$\|\Delta u_x^N\|^2(t) \leq \|u_t^N\|(t) \|\Delta u_x^N\|(t) + \| |u^N|^2 u_x^N \|(t) \|\Delta u_x^N\|(t)$$

or

$$\|\Delta u_x^N\|(t) \leq \|u_t^N\|(t) + \| |u^N|^2 u_x^N \|(t).$$

Making use of (2.8), (3.24), we get

$$\|\Delta u_x^N\|(t) \leq C e^{(-\frac{\kappa t}{2})} \quad (3.33)$$

where the constant  $C$  does not depend on  $N, t > 0$ .

**3.1. Passage to the limit as  $N \rightarrow \infty$ .** Since the constants in Lemmas 3.1-3.4 and (3.33) do not depend on  $N, t > 0$ , then making use of the standard arguments, see [16, 37], one may pass to the limit as  $N \rightarrow \infty$  in (3.6) to obtain for all  $\psi(x, y) \in L^2(D)$  :

$$\int_D [u_t + u^2 u_x + \Delta u_x] \psi dx dy = 0. \quad (3.34)$$

Taking into account Lemmas 3.1-3.4, we establish the following result:

**Lemma 3.5.** *Let all the conditions of Theorem 3.1 hold. Then there exists a strong solution  $u(x, y, t)$  to (2.1)-(2.4) such that*

$$\begin{aligned} & \|u\|_{H_0^1(D)}^2(t) + \|\nabla u_y\|^2(t) + \|u_t\|^2(t) + \|\Delta u_x\|^2(t) \\ & + \|u_x(0, y, t)\|_{H_0^1(0, B)}^2 \leq C e^{(-\kappa t)}, \end{aligned} \quad (3.35)$$

$$\begin{aligned} & \int_0^t \left\{ \|\nabla u_{yy}\|^2(\tau) + \|\Delta u_x\|_{L^2(D)}^2(\tau) + \|u_t\|_{H_0^1(D)}^2(\tau) \right. \\ & \left. + \|u_x(0, y, \tau)\|_{H^2(0, B)}^2 d\tau \right\} \\ & \leq C(D, \|u_0\|, \|u_t\|(0), \|u_{yy}\|(0)), \quad t > 0. \end{aligned} \quad (3.36)$$

#### 4. MORE REGULARITY

In order to complete the proof of the existence part of Theorem 3.1, it suffices to show that

$$\begin{aligned} u & \in L^\infty(\mathbb{R}^+; H^2(D)) \cap u \in L^2(\mathbb{R}^+; H^3(D)), \\ u_t & \in L^\infty(\mathbb{R}^+; L^2(D)) \cap u \in L^2(\mathbb{R}^+; H^1(D)). \end{aligned}$$

These inclusions will be proved in the following lemmas.

**Lemma 4.1.** *A strong solution from Lemma 3.5 satisfies the following inequality:*

$$\begin{aligned} & \int_{\mathbb{R}^+} \left\{ \|u\|_{H^3(D)}^2(t) + \|\Delta u_x\|_{H^1(D)}^2(t) \right\} dt \\ & \leq C(D, \|u_0\|, \|u_t\|(0), \|u_{yy}\|(0)) e^{(-\chi t)}. \end{aligned} \quad (4.1)$$

*Proof.* Taking into account (3.35), (3.36) and Proposition 2.1, we write (3.34) in the form

$$\begin{aligned} \Delta u_x &= -u_t - u^2 u_x \equiv f(x, y, t) \in L^\infty(\mathbb{R}^+; L^2(D)), \\ u_x(0, y, t) &\equiv \phi(y, t) \in L^2(\mathbb{R}^+; H^2(0, B)) \cap L^\infty(\mathbb{R}^+; H_0^1(0, B)), \\ u_x(x, 0, t) &= u_x(x, B, t) = u_x(L, y, t) = 0. \end{aligned}$$

Denote  $\Phi(x, y, t) = \phi(y, t)(1 - x/L)$ . Obviously,

$$\Phi \in L^2(\mathbb{R}^+; H^2(D)).$$

Then the function

$$v = u_x - \Phi(x, y, t)$$

solves in  $D$  the elliptic problem

$$\Delta v = f(x, y, t) - \Phi_{yy}(x, y, t) \in L^2(R^+; L^2(0, D)), \quad v|_\gamma = 0$$

which admits a unique solution  $v \in L^2(\mathbb{R}^+; H^2(D))$ , see [16]. Consequently,  $u_x \in L^2(\mathbb{R}^+; H^2(D))$ . Therefore (3.36) implies (4.1). This completes the proof of Lemma 4.1.  $\square$

**Lemma 4.2.** *A strong solution given by Lemma 3.5 satisfies the following inequality:*

$$\begin{aligned} & \|u\|_{H^2(D)}^2(t) + \|\Delta u_x\|^2(t) \\ & \leq C(D, \|u_0\|, \|u_t\|(0), \|u_{yy}\|(0)) e^{(-\chi t)}, \quad t > 0. \end{aligned} \quad (4.2)$$

*Proof.* Making use of (3.35) and acting as by the proof of Lemma 4.1, we get

$$\begin{aligned} \Delta u_x &= -u_t - u^2 u_x \equiv f(x, y, t) \in L^\infty(\mathbb{R}^+; L^2(D)), \\ u_x(0, y, t) &\equiv \varphi(y, t) \in L^\infty(\mathbb{R}^+; H^1(0, B)), \\ u_x(x, 0, t) &= u_x(x, B, t) = u_x(L, y, t) = 0. \end{aligned}$$

Denote  $\Phi(x, y, t) = \varphi(y, t)(1 - x/L)$ . Obviously,

$$\Phi \in L^\infty(\mathbb{R}^+; H_0^1(D)).$$

Then the function

$$v = u_x - \Phi(x, y, t)$$

solves in  $D$  the elliptic problem

$$\Delta v = f(x, y, t) - \Phi_{yy}(x, y, t) \in L^\infty(\mathbb{R}^+; H^{-1}(D)), \quad v|_\gamma = 0. \quad (4.3)$$

By the elliptic equations theory [16], there exists a unique weak solution to (4.3),

$$v \in L^\infty(\mathbb{R}^+; H_0^1(D)).$$

Consider the scalar product

$$-(v, \Delta v)(t) = -(v, f - \Phi_{yy})(t) = -(v, f)(t) - (v_y, \Phi_y)(t)$$

that can be rewritten in the form

$$\|v_x\|^2(t) + \|v_y\|^2(t) \leq \frac{1}{2} [\|f\|^2(t) + \|v\|^2(t) + \|\Phi_y\|^2(t) + \|v_y\|^2(t)].$$

This implies

$$\|v\|_{H^1(D)}^2(t) \leq C[\|f\|^2(t) + \|\Phi_y\|^2(t)] \leq Ce^{(-\chi t)}. \quad (4.4)$$

Hence,  $u_x = v + \Phi \in L^\infty(\mathbb{R}^+; H^1(D))$  and since  $u, u_y \in L^\infty(\mathbb{R}^+; H^1(D))$ , then

$$u \in L^\infty(\mathbb{R}^+; H^2(D) \cap H_0^1(D)).$$

This completes the proof of Lemma 4.2.  $\square$

**Lemma 4.3.** *The strong solution from Lemmas 3.5-3.7 is uniquely defined.*

*Proof.* Let  $u_1$  and  $u_2$  be two distinct solutions to (2.1)-(2.4). Then  $z = u_1 - u_2$  solves the following IBVP:

$$Az \equiv z_t + z_x + \frac{1}{2}(u_1^3 - u_2^3)_x + \Delta z_x = 0 \quad \text{in } Q \quad t > 0, \quad (4.5)$$

$$\begin{aligned} z(0, y, t) &= z(L, y, t) = z_x(L, y, t) = z(x, 0, t) \\ &= z(x, B, t) = 0, \end{aligned} \quad (4.6)$$

$$z(x, y, 0) = 0, \quad (x, y) \in D. \quad (4.7)$$

From the scalar product

$$2(Az, (1+x)z)(t) = 0,$$

we infer

$$\begin{aligned} &\frac{d}{dt}((1+x), z^2)(t) + 3\|z_x\|^2(t) + \|z_y\|^2(t) + \int_0^B z_x^2(0, y, t) dy \\ &= -2((1+x)(u_1^3 - u_2^3)_x, z)(t) = 2((u_1^2 + u_1 u_2 + u_2^2)z, z \\ &+ (1+x)z_x)(t) \leq 2\|z_x\|^2(t) + C(M)((1+x), z^2)(t), \end{aligned} \quad (4.8)$$

where  $M = \sup_D |u_1^2 + u_1 u_2 + u_2^2|(x, y, t)$ . Due to Proposition 2.1 and (4.2),  $M$  does not depend on  $t > 0$ . Hence (4.8) becomes

$$\frac{d}{dt}((1+x), z^2)(t) \leq C(M)((1+x), z^2)(t).$$

Since  $z(x, y, 0) \equiv 0$ , by the Gronwall lemma,

$$\|z\|^2(t) \leq ((1+x), z^2)(t) \equiv 0, \quad t > 0.$$

The proof of Lemma 4.3 is complete.  $\square$

**Conclusions.** An initial-boundary value problem for the 2D critical generalized Zakharov-Kuznetsov equation posed on rectangles has been considered. Assuming small initial data, the existence of a regular global solution, uniqueness and exponential decay of  $\|u\|(t)_{H^2(D)}$  have been established.

**Acknowledgements.** This research has been supported by Fundação Araucaria, Parana, Brazil; convenio No 307/2015, Protocolo No 45.703. The author appreciates profound and concrete comments of the reviewer.

**Data Availability Statement.** This research does not contain any data necessary to confirm.

## REFERENCES

- [1] R. Adams, J. Fournier, Sobolev Spaces, Second Edition (2003), Elsevier Science Ltd.
- [2] Araruna F. D., Capistrano-Filho R. A., Doronin G. G.: Energy decay for the modified Kawahara equation posed in a bounded domain. J. Math. Anal. Appl. 385, 743–756 (2012).
- [3] J. Colliander, M. Keel, G. Staffilani, H. Takaoka and T. Tao, Sharp global well-posedness results for periodic and non-periodic KdV and modified KdV on  $\mathbb{R}$  and  $\mathbb{T}$ , J. Amer. Math. Soc. 16 (2003), 705–749.
- [4] G. G. Doronin and N. A. Larkin, Stabilization of regular solutions for the Zakharov-Kuznetsov equation posed on bounded rectangles and on a strip, Proc. of the Edinburgh Math. Soc. 58 (2015), 661–682.
- [5] A. V. Faminskii, Well-posed initial-boundary value problems for the Zakharov-Kuznetsov equation, Electronic Journal of Differential equations 127 (2008), 1–23.
- [6] A. V. Faminskii, Initial-boundary value problems in a rectangle for the two-dimensional Zakharov-Kuznetsov equation, JMAA 463 (2018), 760–793.
- [7] Faminskii A. V., Larkin N. A.: Initial-boundary value problems for quasilinear dispersive equations posed on a bounded interval, Electron. J. Differ. Equations. 1–20 (2010).
- [8] Farah L. G., Linares F., Pastor A.: The supercritical generalized KDV equation: global well-posedness in the energy space and below, Math. Res. Lett. 18, no. 02, 357–377 (2011).



- [9] L. G. Farah, F. Linares and A. Pastor, A note on the 2D generalized Zakharov-Kuznetsov equation: Local, global, and scattering results, *J. Differential Equations* 253 (2012), 2558–2571.
- [10] G. Fonseca, F. Linares, G. Ponce, Global well-posedness for the modified Korteweg-de Vries equation, *Comm. Part. Diff. Equats*, 24 (3,4) (1999), 683–705.
- [11] G. Fonseca, F. Linares, G. Ponce, Global existence for the critical generalized KDV equation, *Proc. of the AMS*, vol. 131, Number 6 (2002), 1847–1855.
- [12] Jeffrey A. and Kakutani T.: Weak nonlinear dispersive waves: a discussion centered around the Korteweg-de Vries equation, *SIAM Review*, vol 14 no 4, pp. 582–643 (1972).
- [13] Kakutani T. and Ono H.: Weak non linear hydromagnetic waves in a cold collision free plasma, *J. Phys. Soc. Japan*, 26, 1305–1318 (1969).
- [14] T. Kato, On the Cauchy problem for the (generalized) Korteweg-de Vries equations, *Advances in Mathematics Supplementary Studies, Stud. Appl. Math.* 8 (1983), 93–128.
- [15] C. E. Kenig, G. Ponce and L. Vega, Well-posedness and scattering results for the generalized Korteweg-de Vries equation and the contraction principle, *Commun. Pure Appl. Math.* 46 (1993), 527–620.
- [16] O. A. Ladyzhenskaya, *The Boundary Value Problems of Mathematical Physics. Applied Math. Sci.* 49, Springer-Verlag, New York, 1985.
- [17] O. A. Ladyzhenskaya, V. A. Solonnikov and N. N. Uraltseva, *Linear and Quasilinear Equations of Parabolic Type*. American Mathematical Society, Providence, Rhode Island, 1968.
- [18] N. A. Larkin; Correct initial boundary value problems for dispersive equations, *J. Math. Anal. Appl.* 344 (2008) 1079–1092.
- [19] N. A. Larkin; Exponential decay of the  $H^1$ -norm for the 2D Zakharov-Kuznetsov equation on a half-strip, *J. Math. Anal. Appl.* 405 (2013) 326–335.
- [20] N. A. Larkin; Global regular solutions for the 3D Zakharov-Kuznetsov equation posed on a bounded domain, *Differential and Integral equations*, vol.29, N 7-8, (2016), 775–790.
- [21] N. A. Larkin; Global regular solutions for the 3D Zakharov-Kuznetsov equation posed on unbounded domains, *J. of Mathematical Physics*, 56, 091508 (2015); doi.org 10.1063/1.4928924.
- [22] N. A. Larkin; The 2D Zakharov-Kuznetsov-Burgers equation on a strip, *Bol. Soc. Paran. Mat.*, (3s), v.34 (2016), 151–172.
- [23] N. A. Larkin and J. Luchesi, Generalized dispersive equations of higher orders posed on bounded intervals: local theory, arXiv:1812.04146 v 1 [math. AP] 10 Dec 2018.
- [24] N. A. Larkin and J. Luchesi, Initial-boundary value problems for generalized dispersive equations of higher orders posed on bounded intervals, *Appl. Math and Optmization*, doi.org/10.1007/s00245-019-09579-w.
- [25] N. A. Larkin and M. Padilha, Global regular solutions to one problem of Saut-Temam for the 3D Zakharov-Kuznetsov equation, *Appl. Mat. and Optimization*, (2018) 77:253–274; doi.org/10.1007/sw00245-016-9373-1
- [26] N. A. Larkin and E. Tronco, Regular solutions of the 2D Zakharov-Kuznetsov equation on a half-strip, *J. Differential Equations* 254 (2013), 81–101.

- [27] F. Linares and A. Pastor, Local and global well-posedness for the 2D generalized Zakharov-Kuznetsov equation, *J. Funct. Anal.* 260 (2011), 1060–1085.
- [28] F. Linares, A. Pastor and J.-C. Saut, Well-posedness for the ZK equation in a cylinder and on the background of a KdV Soliton, *Comm. Part. Diff. Equations* 35 (2010), 1674–1689.
- [29] F. Linares and A. Pazoto; *On the exponential decay of the critical generalized Korteweg-de Vries equation with localized damping.* *Proc. Amer. Math. Soc.*, 135, 1 (2007) 1515-1522.
- [30] M. Panthee, M. Scialom, Asymptotic behaviour for a class of solutions to the critical modified Zakharov-Kuznetsov equation, *Stud. Appl. Math.* 124 (2010) 229–245.
- [31] Y. Martel and F. Merle; Instability of solutions for the critical generalized Korteweg-de Vries equation, *Geometrical and Funct. Analysis*, 11 (2001) 74-123.
- [32] F. Merle; Existence of blow up solutions in the energy space for the critical generalized KdV equation, *J. Amer. Math. Soc.*, 14 (2001) 555-578.
- [33] Pilod D.: *On the Cauchy problem for higher-order nonlinear dispersive equations.* *Journal of Differential Equations.* 245, 2055–2077 (2008).
- [34] J. C. Saut, Sur quelques généralisations de l'équation de Korteweg-de Vries (French), *J. Math. Pures Appl.* 58 (1979), 21–61.
- [35] A. V. Steklov, The problem of cooling of an heterogeneous rigid rod, *Communs. Kharkov Math. Soc.*, Ser. 2, 5 (1896) 136-181 (Russian).
- [36] J.-C. Saut and R. Temam, An initial boundary-value problem for the Zakharov-Kuznetsov equation, *Advances in Differential Equations* 15 (2010), 1001–1031.
- [37] J.-C. Saut, R. Temam and C. Wang, An initial and boundary-value problem for the Zakharov-Kuznetsov equation in a bounded domain, *J. Math. Phys.* 53 115612(2012).
- [38] V. E. Zakharov and E. A. Kuznetsov, On three-dimensional solitons, *Sov. Phys. JETP* 39 (1974), 285–286.