

Quantum State Dependence of Thermodynamic Phase Transition in 4D AdS Gauss-Bonnet Quantum Black Holes Surrounded With Cloud of Strings

H. Ghaffarnejad ^{a,1}, E. Yaraie ^{a,b,2} and M. Farsam ^{a,b,3}

^a *Faculty of Physics, Semnan University, P.C. 35131-19111, Semnan, Iran*

^b *Instituut-Lorentz for Theoretical Physics, ITP, Leiden University, Niels Bohrweg 2, Leiden 2333 CA, The Netherlands*

Abstract

According to the Lovelock theorem where the model [27] could not applicable for Einstein Gauss Bonnet (EGB) gravity in all 4D curved spacetimes, authors of the reference [31] presented an effective model by applying break of diffeomorphism property. Hence we use the latter model instead of former for study of thermodynamic behavior of a 4D AdS EGB spherically symmetric static black hole which surrounded with a cloud of string. In short our work is extension of the works given by [33, 34] but not by using [27] but by applying [31]. Our metric solutions are obtained versus the Hermite polynomials (quantum harmonic Oscillator) for which eigen values come from single scale defined by multiplication of the coupling constants of the model: Namely the regularized GB parameter, AdS radius, the black hole ADM mass and the string tension. Hence we claim the obtained metric solution is in fact behavior of quantized black hole. Because the GB term is originated from renormalization of quantum matter fields. Also we should pointed that this kind of quantization is different with the canonical quantization (Wheeler De Witt). Our study shows that all phase transitions of this quantum black hole are dependent to the Hermite quantum numbers.

1 Introduction

In absence of a pure quantum gravity theory which should be valid for the Planck scale of the spacetime [1], its semi classical approach is applicable

¹E-mail address: hghafarnejad@semnan.ac.ir

²E-mail address: eyaraie@semnan.ac.ir

³E-mail address: mhdarsam@semnan.ac.ir

still. It is known as quantum field theories propagating on curved spacetimes [2, 3]. In this sense renormalization theory for expectation value of quantum matter fields stress energy tensor operator, suggests nonsingular expectation value for it but with an anomaly trace. It is well known that this conformal anomaly is given by the geometrical Ricci or Kretschmann scalars or Gauss Bonnet topological invariant when quantum matter fields [2, 3] are massless. By regarding these anomaly terms as geometric perspective of quantum matter field corrections on the Einstein metric equation which is called as the backreaction equation (see for instance [4] and references therein), one can infer that this metric backreaction equation or corresponding action functional can be consider as quantum gravity model in absence of pure quantum gravity theory which does not still presented. Form this point of view the EGB gravity which we will use in this work can be considered as alternative gravity which originates from quantum matter effects. Other approaches such as generalization of number of dimensions of the curved spacetimes to larger than four $D > 4$ via string theories [5, 6] is also generate such these anomalies [7, 8, 9]. Among the various higher order derivative gravitational models which are given in the literature the Lovelock gravity [10] is quite special, because it is free of ghost [11, 12, 13, 14, 15, 16, 17]. In fact, there are presented many higher order derivative metric theories which exhibit with Ostrogradsky instability (see [18, 19] for a good review). In this sense the actions which contains higher order curvature terms introduce equations of motions with fourth order or higher metric derivatives where linear perturbations disclose that the graviton should be a ghost. Fortunately the Lovelock model is free of ghost term which means that this model have field equations involving not more than second order derivatives of the metric. Action functional of the Lovelock gravity, is given by combinations of various terms as follows. The first term is the cosmological constant Λ , the second term is the Ricci scalar $R = R^\mu_\mu$, and the third and fourth terms are the second order Gauss-Bonnet [20] and third order Lovelock terms (see Eq. 22 in ref. [21]), respectively. Without the latter term the Lovelock gravity reduces to a simplest form called as the Einstein-Gauss-Bonnet (EGB) theory in which the Einstein-Hilbert action is supplemented with the quadratic curvature GB term. Importance of this form of the gravity model is appeared more when we see that it is generated from effective Lagrangian of low energy string theory [22, 23, 24, 25, 26]. In fact for curved spacetimes with $D > 4$ dimensions the Gauss Bonnet coupling parameter which is calculated by dimensional regularization method have some regular values but not have for $D = 4$. To

resolve this problem the author Glavan and his collaborator presented a proposal [27] but we know now that their initial proposal does not lead to every well-defined theory because regularization is guaranteed just some large class of metrics but not for all metrics. In this sense one can see [28, 29] where its authors explained several inconsistencies of the original 4DEGB paper by Glavan and Lin. Particularly, besides pointing out possible problems in defining the limit or finding an action for the theory, their work also adds new results to the discussion concerning the ill-definiteness of second order perturbations even around a Minkowskian background or the geodesic incompleteness of the spherically-symmetric black hole geometry presented by Glavan and Lin (See also [30]). Thus there must be presented other proposals that can be cover all metric theories. Recently a well defined theory is presented [31] by breaking the diffeomorphism property of the curved space-time. Instead of the former work [27] the latter model is in concordance with the Lovelock theorem and thereby, seems more to be physical and applicable. For instance FRW cosmology of the latter model is studied in [32] and showed success of the model versus [27].

Despite that the gravity model [27] is sick in 4 dimensional curved space time but idea of authors of [33] is still useful in which by adding action of string clouds with the model [27] they obtained metric solution of a charged 4DEGB black hole. Then they are studied thermodynamic properties of the obtained black hole metric solution. By looking at their work one can infer that the string tension changes all thermodynamic variables of the black hole except its entropy. By studying the heat capacity behavior they obtained that the smaller black holes are locally stable. Due to the surrounding cloud of strings, they obtained a black hole phase transition where a large unstable black holes transit to a globally thermodynamically small stable black holes with negative free energy. Their results demonstrate that the Hawking's evaporation leads to a thermodynamically stable remnant with no temperature. We should point that authors of the work [33] did not considered the cosmological constant effect as vacuum de Sitter or Anti de Sitter space.

On the other side we should point to idea which is used in [34] by applying [27] as follows. Authors of the work [34] studied thermodynamics of the 4DAdSEGB black hole in the extended phase space where the cosmological constant behaves as pressure of the AdS vacuum space. They obtained that the black hole exhibits with a phase transition similar to that of van der Waals system. They also showed that this black hole exhibited with the Joule-Thomson expansion. They do not used effects of surrounding clouds

of string tension in their work.

As an extension of the works [33, 34] we like to have study effects of string clouds on thermodynamic behavior of 4DAdSEGB chargeless black hole by applying the gravity model [31] not the sick model [27].

There are many published works in the literature where the higher order derivative gravities such as Lovelock or its reduced forms are applied to study black hole thermodynamics (see for instance [35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62]). The Lovelock gravity or its reduced models are studied in cosmological approaches too. In this sense we like to point to some published works such as follows. For instance authors of the work [63] have assumed that the graviton is massless or nearly massless motivated by results of *GW170817*. This indicated that speed of the gravitational waves is equal to that of light's and thus the fundamental carrier of the gravitational force is nearly massless. The gravitational effects is not short range interaction. Specifically by using an EGB extension of Horndeski theories they showed that some classes of the latter theories may be revived and become viable in view of the *GW170817* result. In ref. [64] they showed how the normal EGB theories can be viable in view of *GW170817*. As an extension of non-minimally coupled scalar field to Ricci scalar theory in the context of EGB model one can see [65] and seek what is the swampland criteria in the context of a massless graviton EGB theory? To do so [66, 67] should be followed. Also authors of the work [68] checked the validity of the weak cosmic censorship conjecture for the 4D charged EGB black hole and considered the effect of the GB coupling constant on the validity of the weak cosmic censorship conjecture. Hawking studied in a new attempt in a fascinating way, AdS black holes phase transition at a first time which has been called as extended phase space where the negative cosmological constant behaves as pressure of AdS vacuum space acts on the local black hole horizon . [69]. The extended phase space represents a phase space in which the traditional first law of black hole thermodynamics is corrected by an additional VdP work so that the cosmological constant is regarded as thermodynamic pressure of AdS space which affects on the black hole. In this sense its conjugate variable is thermodynamic volume. A large number of investigations have been done on this concept during last years and one can follow them for instance at these works: [70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93]. Particularly for Van der Waals behavior of AdS black holes which can be can followed some published works such as follows [95, 96, 97, 98, 99]. As a different model one can see

[100] where the authors obtained a small to large black hole phase transition by generalizing the well known Einstein Maxwell Dilaton gravity with an additional vector field supporting spherical or hyperbolic horizons. In this view the strings fluids as gravitational matter source which has received less attention in solving the gravitational equations and we like consider in this work. As an important role of the string fluids at cosmology it is enough to say that the rapid expansion of the universe during inflation is thought to be related to the expansion of cosmic strings. [94, 102, 103, 104, 105, 106]. In this paper we would like to investigate extended thermodynamics phase transition of 4DAdSGB black hole metric solution obtained from [31] in presence of strings fluid. We will show important and critical role of the used string tension which how can present quantization condition on the event horizon of the obtained black hole metric solution. Setup of this work is as follows. In section 2, we review 4DAdSGB gravity [31] in presence of strings fluid. Then in sections 3 and 4 we solve metric solution of a spherically symmetric static curved space time with a black hole topology. Our obtained metric solutions are described by the well known Hermite polynomials. In section 5 we investigate various aspect of the extended thermodynamics of 4DAdSGB black hole in presence of strings fluid for ground state and first excited state. To do so we study $P - V$ criticality and probe global stability of the black hole system by plotting diagram of the Gibbs free energy versus the temperature where there is seen a swallowtail behavior (coexistence of phases) for pressures higher than the critical pressure (see figures 3 and 8). Then we investigate possibility of small to large black hole phase transition and also Hawking-Page phase transition. Last section denotes to concluding remark and outlook.

2 4D AdS GB gravity with strings fluid

According to the work [31] we know a constraint EGB gravity in $D \rightarrow 4$ limit which is given by the first term in the following action functional. I_{matter} Second matter term in the following action is assumed to be clouds of strings fluid.

$$I = \frac{1}{16\pi G} \int dt d^3x N \sqrt{\gamma} \mathcal{L}_{EGB}^{4D} + I_{\text{matter}}, \quad (2.1)$$

with

$$\mathcal{L}_{\text{EGB}}^{4D} = 2R - 2\Lambda - \mathcal{M} + \frac{\tilde{\alpha}}{2} [8R^2 - 4R\mathcal{M} - \mathcal{M}^2 - \frac{8}{3}(8R_{ij}R^{ij} - 4R_{ij}\mathcal{M}^{ij} - \mathcal{M}_{ij}\mathcal{M}^{ij})], \quad (2.2)$$

where G is the Newton's gravitational coupling constant, R and R_{ij} are the Ricci scalar and the Ricci tensor of the spatial metric γ_{ij} respectively and

$$\mathcal{M}_{ij} = R_{ij} + \mathcal{K}_k^k \mathcal{K}_{ij} - \mathcal{K}_{ik} \mathcal{K}_j^k, \quad \mathcal{M} = \mathcal{M}_i^i \quad (2.3)$$

with

$$\mathcal{K}_{ij} = \frac{1}{2N} (\dot{\gamma}_{ij} - 2D_i N_j - 2D_j N_i - \gamma_{ij} D_k D^k \lambda_{GF}). \quad (2.4)$$

Here a dot denotes time derivative t and all the effects of the constraint stemming from the gauge-fixing (GF) are now encoded in Lagrange multiplier λ_{GF} . D_i is spatial covariant derivative and re-scaled regular EGB coupling constant $\tilde{\alpha}$ is defined versus the irregular GB coupling constant itself α_{GB} in $D \rightarrow 4$ dimensions limit as $\tilde{\alpha} = (D - 4)\alpha_{GB}$. The above EGB gravity action satisfies the following gauge condition for all spherically symmetric and cosmological backgrounds (see [31] and [32]).

$$\sqrt{\gamma} D_k D^k (\pi^{ij} \gamma_{ij} / \sqrt{\gamma}) \approx 0 \quad (2.5)$$

In fact the above EGB action is generated from ADM decomposition of the 4D background metric as 1 + 3 dimension as follows.

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = -N^2 dt^2 + \gamma_{ij} (dx^i + N^i dt)(dx^j + N^j dt) \quad (2.6)$$

where N , N_i , γ_{ij} are the lapse function, the shift vector, and the spatial metric respectively. γ factor in the action (2.1) is absolute value of determinant of the spatial metric γ_{ij} . This ADM decomposition is done on the background metric to remove divergent boundary term of the higher order metric derivative in the GB term of the action functional (2.1) in general 4D form [31]. First term in the theory defined by (2.1) has the time re-parametrization symmetry $t \rightarrow t = t(t')$. We now set the matter source I_{matter} to be the Nambu-Goto action [102] (see also page 100 in ref. [5]) which explains the dynamics of relativistic strings as follows.

$$I_{\text{NG}} = \int_{\Sigma} \rho \sqrt{\mathbf{g}} d\sigma^0 d\sigma^1 \quad (2.7)$$

where ρ is tension (mass per unit length or linear mass density) in the string and worldsheet of strings can be parameterized by local coordinates (σ^0, σ^1) . \mathfrak{g} is absolute value of determinant of induced metric \mathfrak{g}_{ab} and the bivector $\Sigma^{\mu\nu}$ related to strings worldsheet are given respectively by

$$\mathfrak{g}_{ab} = g_{\mu\nu} \frac{\partial x^\mu}{\partial \sigma^a} \frac{\partial x^\nu}{\partial \sigma^b} \quad (2.8)$$

and

$$\Sigma^{\mu\nu} = \epsilon^{ab} \frac{\partial x^\mu}{\partial \sigma^a} \frac{\partial x^\nu}{\partial \sigma^b}, \quad (2.9)$$

where ϵ^{ab} is two dimensional Levi-Civita tensor density $\epsilon^{01} = -\epsilon^{10} = 1$. Strings fluid is described by the energy momentum tensor

$$T^{\mu\nu} = \mathfrak{g}^{-\frac{1}{2}} \rho \Sigma^{\mu\delta} \Sigma_\delta^\nu = \frac{2\partial \mathfrak{L}_{NG}}{\partial g^{\mu\nu}}. \quad (2.10)$$

We now investigate to solve the metric equation of the above model for a spherically symmetric static black hole space time.

3 4D AdS GBBH surrounded by strings cloud

By comparing the metric line element (2.6) with general form of a spherically symmetric state 4D metric line element

$$ds^2 = f(r) dt^2 - f(r)^{-1} dr^2 - r^2 d^2\Omega \quad (3.1)$$

we infer that the lapse function, and the shift vector, the spatial metric components and lagrange multiplier should be r dependent so that we can write

$$N = f(r) = \frac{1}{\gamma_{rr}}, \quad N_i = 0, \quad \gamma_{\theta\theta} = r^2, \quad \gamma_{\varphi\varphi} = r^2 \sin^2 \theta, \quad \lambda_{GF} = \lambda_{GF}(r) \quad (3.2)$$

where

$$\begin{aligned} \mathcal{L}_{EGB}^{4D} = & -2\Lambda - \frac{3q^2}{2r^4 f} - \frac{2(rf' + f - 1)}{r^2} \\ & + \tilde{\alpha} \left\{ + \frac{4(f-1)f'}{r^3} - \frac{2(f-1)^2}{r^4} + \frac{q'^2(rf' + f - 1)}{r^6 f} - \frac{q^4}{8r^8 f^2} \right\} \end{aligned} \quad (3.3)$$

with

$$q = r^2 \sqrt{f} \lambda'_{GF} \quad (3.4)$$

in which \prime denotes to derivative with respect to r . To calculate explicit form of the string action functional (2.7) for the spherically symmetric static background metric (3.1) we should use a static gauge [5] where the commoving time σ^0 in worldsheet $x^\mu(\sigma^0, \sigma^1)$ is equal to the time t as $\sigma^0 = t = t_0 = \text{constant}$ and we should assume the static string is along to the radial direction r of worldsheet $x^\mu(\sigma^0, \sigma^1)$ such that

$$r = r(\sigma^0, \sigma^1) = F(\sigma^1), \quad t = \sigma^0, \quad \theta(\sigma^0, \sigma^1) = \varphi(\sigma^0, \sigma^1) = 0. \quad (3.5)$$

Here we choose an open string which one edge of the world sheet to be the curve $\sigma^1 = 0$ and the other edge to be the curve $\sigma^1 = a$ such that $\sigma^1 \in [0, a]$ for an open string with arbitrary shape $F(\sigma^1)$. In this case non-vanishing components of the induced metric (2.8) reads

$$\mathfrak{g}_{tt} = f(r), \quad \mathfrak{g}_{rr} = \frac{-1}{f(r)} \left(\frac{dF(\sigma^1)}{d\sigma^1} \right)^2, \quad \sqrt{\mathfrak{g}} = \sqrt{|\det(\mathfrak{g})|} = \frac{dF(\sigma^1)}{d\sigma^1}. \quad (3.6)$$

Spherically symmetric property of the background metric (3.1) causes that the string tension ρ and bivector $\Sigma^{\mu\nu}$ to be dependent to r alone. Covariant conservation of the string stress tensor (2.10) leads to the condition $\nabla_\mu(\rho \Sigma^{\mu\nu}) = 0$ (see ref. [102]) which for the metric equation (3.1) we will have stress energy tensor components of the Nambu Goto (NG) relativistic string as follows.

$$T_t^t = T_r^r = -\rho \Sigma^{tr} = \frac{C}{r^2} \quad (3.7)$$

where dimensionless parameter C is a integral constant. By regarding the stress energy tensor definition $T^{\mu\nu} = \frac{2\partial \mathfrak{L}_{NG}}{\partial g^{\mu\nu}}$ given by (2.10) and by substituting (3.7) and metric components (3.1) we can obtain explicit form of the NG string Lagrangian density as follows.

$$\mathfrak{L}_{NG} = \sqrt{\mathfrak{g}} \rho = \frac{C}{4r^2 f^2}. \quad (3.8)$$

By substituting (3.3) and by integrating first term of the action functional (2.1) on the 2-sphere $0 \leq \theta \leq \pi$, $0 \leq \varphi \leq 2\pi$ we obtain $I_{EGB} = \int dt dr \mathfrak{L}$ for the first term of the action functional (2.1) where \mathfrak{L} is defined by

$$\mathfrak{L} = \frac{r^2 f}{4G} \mathcal{L}_{EGB}^{AD}. \quad (3.9)$$

Now we are in position to obtain total lagrangian density of the system by adding (3.8) and (3.9) as follows.

$$\begin{aligned} \mathfrak{L}_{total} = \frac{r^2 f}{4G} \left\{ \frac{6}{\ell^2} - \frac{3q'^2}{2r^4 f} - \frac{2(rf' + f - 1)}{r^2} + \tilde{\alpha} \left\{ \frac{4(f-1)f'}{r^3} - \frac{2(f-1)^2}{r^4} \right. \right. \\ \left. \left. + \frac{q'^2(rf' + f - 1)}{r^6 f} - \frac{q'^4}{8r^8 f^2} \right\} \right\} + \frac{C}{4r^2 f^2} \end{aligned} \quad (3.10)$$

where we substituted identity between the cosmological constant and radius of 4D AdS space as $\Lambda = -\frac{3}{\ell^2}$. By substituting

$$q'^2 = Q'^2 + \xi \quad (3.11)$$

in which

$$\xi = 4r^2 f(rf' + f - 1) - \frac{6r^4 f}{\tilde{\alpha}} \quad (3.12)$$

the total Lagrangian density (3.10) reads

$$\begin{aligned} \mathfrak{L}_{total} = \frac{r^2 f}{4G} \left\{ \frac{6}{\ell^2} - \frac{2(rf' + f - 1)}{r^2} + \tilde{\alpha} \left\{ \frac{4(f-1)f'}{r^3} - \frac{2(f-1)^2}{r^4} \right\} \right\} \\ \frac{r^2 f}{4G} \left\{ \xi \left[\frac{\tilde{\alpha}(rf' + f - 1)}{r^6 f} - \frac{3}{2r^4 f} \right] - \frac{\tilde{\alpha}(Q'^4 + \xi^2)}{8r^8 f^2} \right\} + \frac{C}{4r^2 f^2}. \end{aligned} \quad (3.13)$$

It is easy to check that Euler Lagrange equation for the field Q reduces to the following conservation equation.

$$\frac{\partial \mathfrak{L}_{total}}{\partial Q'} = -\frac{\tilde{\alpha} Q'^3}{8G f r^6} = constant \equiv D^3 \quad (3.14)$$

for which we have

$$Q' = -2Dr^2 \left(\frac{Gf}{\tilde{\alpha}} \right)^{\frac{1}{3}}. \quad (3.15)$$

This conservation equation helps us to obtain explicit form of the undetermined gauge fixing Lagrange multiplier as follows.

$$\lambda_{GF}(r) = \int^r \frac{dr_1}{r_1^2 \sqrt{f(r_1)}} \times \quad (3.16)$$

$$\left[\int^{r_1} dr_2 \sqrt{\left(\frac{8GD^3 r_2^6}{\tilde{\alpha} f(r_2)} \right)^{\frac{2}{3}} + r_2^2 f(r_2) f'(r_2) + r_2^2 f^2(r_2) - r_2^2 f(r_2) - \frac{6r_2^4 f(r_2)}{\tilde{\alpha}}} \right]$$

where we substituted (3.11), (3.12) and (3.15) into the relation (3.4). By substituting (3.12) and (3.15), the total Lagrangian density (3.13) reduces to the following form.

$$\begin{aligned} \mathfrak{L}_{total} = & \frac{\tilde{\alpha} f f'^2}{2G} + \frac{2}{G} \left(\frac{\tilde{\alpha}(f-1)}{r} - r \right) f f' - \frac{2f(f-1)}{G} \\ & + \frac{3f r^2}{2G} \left(\frac{1}{\ell^2} + \frac{3}{4\tilde{\alpha}} \right) - \frac{D^4 r^2}{2} \left(\frac{Gf}{\tilde{\alpha}} \right)^{\frac{1}{3}} + \frac{C}{4r^2 f^2} \end{aligned} \quad (3.17)$$

for which one can obtain Euler Lagrange equation for the metric potential $f(r)$ as follows.

$$\begin{aligned} 2f^4 f'' + f^3 f'^2 + \frac{4f^3(f-1)}{\tilde{\alpha}} - \frac{4f^4(f-1)}{r^2} \\ - \frac{3r^2 f^3}{2\tilde{\alpha}} \left(\frac{1}{\ell^2} + \frac{3}{4\tilde{\alpha}} \right) + \frac{D^4 r^2}{3} \left(\frac{G}{\tilde{\alpha}} \right)^{\frac{4}{3}} f^{\frac{7}{3}} + \frac{GC}{\tilde{\alpha} r^2} = 0. \end{aligned} \quad (3.18)$$

This is a nonlinear second order ordinary differential equation and so has not an analytic closed solution. To solve it we must use numerical method or perturbation series expansion method. The latter method is used to solve the metric potential equation (3.18) in the subsequent section.

4 Metric solutions and Discretization

As a physical boundary condition to solve the equation (3.18) we choose

$$\lim_{r \rightarrow \infty} f(r) \rightarrow 1 \quad (4.1)$$

which by substituting the equation (3.18) reads to the following condition.

$$D^4 \left(\frac{G}{\tilde{\alpha}} \right)^{\frac{4}{3}} = \frac{9}{2\tilde{\alpha}} \left(\frac{1}{\ell^2} + \frac{3}{4\tilde{\alpha}} \right). \quad (4.2)$$

To solve the equation (3.18) by regarding the above condition it is useful we define

$$\frac{1}{d^2} = \frac{1}{\ell^2} + \frac{3}{4\tilde{\alpha}}, \quad x = \frac{r}{d}, \quad \epsilon = \frac{GC}{\tilde{\alpha}}, \quad \beta^4 = \frac{d^2}{\tilde{\alpha}}. \quad (4.3)$$

By substituting (4.2) and (4.3) into the equation (3.18) we obtain

$$2f^4\ddot{f} + f^3\dot{f}^2 - \frac{4f^4(f-1)}{x^2} + 4\beta^4 f^3(f-1) + \frac{3}{2}\beta^4 x^2(f^{\frac{7}{3}} - f^3) + \frac{\epsilon}{x^2} = 0 \quad (4.4)$$

where dot denotes to derivative with respect to x . Now we choose ϵ to be order parameter in the following perturbation series function.

$$f(x) = 1 + \epsilon g(x) + \epsilon^2 h(x) + \mathcal{O}(\epsilon^3). \quad (4.5)$$

By substituting (4.5) into (4.4) and solving order by order of the metric equation (4.4) we obtain $g(x)$ and $h(x)$. First and second order of the equation (4.5) against ϵ order parameter are obtained respectively as follows.

$$\ddot{g}(x) + \left(2\beta^4 - \frac{\beta^4 x^2}{2} - \frac{2}{x^2}\right)g(x) + \frac{1}{2x^2} = 0 \quad (4.6)$$

and

$$\begin{aligned} & \ddot{h}(x) + \left(2\beta^4 - \frac{\beta^4 x^2}{2} - \frac{2}{x^2}\right)h(x) \\ &= -\frac{1}{2}\dot{g}^2(x) + \frac{2}{x^2}g(x) + \left(2\beta^4 - \frac{11x^2\beta^4}{12}\right)g^2(x). \end{aligned} \quad (4.7)$$

We do not bring here higher order terms of series form of the equation (4.4) because if we solve just the zero order equation (4.6) then higher order solutions can be obtained easily step by step after where by substituting zero order solution $g(x)$ into the right hand side of the equation (4.7) and obtain their particular solutions so on. Our strategy to solve (4.6) is as follows: At first step we obtain asymptotically behavior of the function $g(x)$ for large distances $x \gg 1$ for which (4.6) reads

$$\ddot{g} + (2\beta^4 - \beta^4 x^2/2)g(x) \approx 0 \quad (4.8)$$

which has the following solutions in terms of the Hermite polynomials.

$$g_n^\infty(y) \approx \frac{e^{-y^2/2} H_n(y)}{\Gamma_n} \quad (4.9)$$

by setting

$$y = \beta x/2^{\frac{1}{4}}, \quad \beta^2 = \frac{(2n+1)}{2\sqrt{2}} \quad (4.10)$$

where Γ_n is normalization coefficient so that

$$\Gamma_n = 2^{\frac{n}{2}} \pi^{\frac{1}{4}} (n!)^{\frac{1}{2}} \quad (4.11)$$

and $n = 0, 1, 2, 3, 4, \dots$ are order of the Hermite polynomials $H_n(y)$. At second step we substitute (4.9), (4.10) and definition

$$g(y) = g_n^\infty(y) F(y) \quad (4.12)$$

into the equation (4.6) to obtain differential equation for unknown $F(y)$ as follows.

$$\frac{d^2 F(y)}{dy^2} + 2 \left(-y + \frac{2n H_{n-1}(y)}{H_n(y)} \right) \frac{dF(y)}{dy} - \frac{2F(y)}{y^2} + \frac{1}{2g_n^\infty(y)y^2} = 0 \quad (4.13)$$

in which we used the identity $2n H_{n-1}(y) = \frac{dH_n(y)}{dy}$ for the Hermite polynomials and we supposed $F(y)$ gives behavior of the metric function $g(y)$ at $y \rightarrow 0$ limits. It is easy to check that in $y \rightarrow 0$ limits we have

$$\lim_{y \rightarrow 0} \frac{2(2n)H_{2n-1}(y)}{H_{2n}(y)} = 0, \quad \lim_{y \rightarrow 0} g_{2n}^\infty(y) = \frac{H_{2n}(0)}{\Gamma_{2n}} = \frac{(-1)^n (2n)!}{n! \Gamma_{2n}} \equiv \frac{1}{e_{2n}} \quad (4.14)$$

and

$$\lim_{y \rightarrow 0} \frac{2(2n+1)H_{2n}(y)}{H_{2n+1}(y)} = \frac{1}{y}, \quad \lim_{y \rightarrow 0} g_{2n+1}^\infty(y) \approx \frac{2(2n+1)H_{2n}(0)}{\Gamma_{2n+1}} y \equiv \frac{y}{o_{2n+1}} \quad (4.15)$$

respectively. By substituting these limits into the equation (4.13) we obtain

$$\frac{d^2 F_e(y)}{dy^2} - \frac{2F_e(y)}{y^2} + \frac{e_{2n}}{2y^2} = 0 \quad (4.16)$$

and

$$\frac{d^2 F_o(y)}{dy^2} + \frac{2}{y} \frac{dF_o(y)}{dy} - \frac{2F_o(y)}{y^2} + \frac{o_{2n+1}}{2y^3} = 0 \quad (4.17)$$

in $y \rightarrow 0$ limits respectively. One can obtain solutions of the equations (4.16) and (4.17) respectively as follows.

$$F_e(y) = \frac{e_{2n}}{2} + \frac{\sigma_1}{y} + \sigma_2 y^2 \quad (4.18)$$

and

$$F_o(y) = \frac{o_{2n+1}}{2y} + \frac{\eta_1}{y^2} + \eta_2 y \quad (4.19)$$

where $\sigma_{1,2}$ and $\eta_{1,2}$ are integral constants and they should be fixed by physical characteristics of local gravitational objects for instance mass and charge of black holes. Now we are in position to write explicit form of the metric potential (4.5) which up to second order terms become

$$f_{2n}(y) = 1 + \frac{\epsilon e^{-y^2/2}}{\Gamma_{2n}} H_{2n}(y) \left(\frac{e_{2n}}{2} + \frac{\sigma_1}{y} + \sigma_2 y^2 \right) \quad (4.20)$$

and

$$f_{2n+1}(y) = 1 + \frac{\epsilon e^{-y^2/2}}{\Gamma_{2n+1}} H_{2n+1}(y) \left(\frac{o_{2n+1}}{2y} + \frac{\eta_1}{y^2} + \eta_2 y \right) \quad (4.21)$$

respectively. We know from (4.14) and (4.15)

$$H_{2n}(0) = (-1)^n \frac{(2n)!}{n!}, \quad \lim_{y \rightarrow 0} H_{2n+1}(y) \sim \frac{2(-1)^n (2n+1)!}{n!} y. \quad (4.22)$$

Now we apply to fix integral constants $\sigma_{1,2}$ and $\eta_{1,2}$ as follows. By substituting (4.3), (4.10) and (4.22), the metric potentials (4.20) and (4.21) read

$$\lim_{r < < d} f_{2n}(r) \sim 1 + \frac{\epsilon}{2} - \frac{2M}{r} + \frac{r^2}{\ell^2} \quad (4.23)$$

and

$$\lim_{r < < d} f_{2n+1}(r) \sim 1 + \frac{\epsilon}{2} - \frac{2M}{r} + \frac{r^2}{\ell^2} \quad (4.24)$$

respectively where we set

$$\sigma_1 = -\frac{2^{\frac{3}{4}} \beta n! \Gamma_{2n}}{(-1)^n (2n)! \epsilon} \frac{M}{d} = -\frac{2^{n-1} n!}{(-1)^n} \sqrt{\frac{(2n+1)\sqrt{\pi}}{(2n)!}} \left(\frac{M\tilde{\alpha}}{CG} \right) \sqrt{\frac{4}{\ell^2} + \frac{3}{\tilde{\alpha}}}, \quad (4.25)$$

$$\sigma_2 = \frac{\sqrt{2} n! \Gamma_{2n}}{\beta^2 \epsilon (-1)^n (2n)!} \frac{d^2}{\ell^2} = \frac{16n! 2^n}{(2n+1)(-1)^n} \sqrt{\frac{\sqrt{\pi}}{(2n)!}} \frac{\tilde{\alpha}^2}{CG(4\tilde{\alpha} + 3\ell^2)} \quad (4.26)$$

and

$$\eta_1 = -\frac{\beta n! \Gamma_{2n+1}}{\epsilon (-1)^n (2n+1)! 2^{\frac{1}{4}}} \frac{M}{d} = -\frac{2^{n-1} n!}{(-1)^n} \sqrt{\frac{\sqrt{\pi}}{2(2n)!}} \left(\frac{M\tilde{\alpha}}{CG} \right) \sqrt{\frac{4}{\ell^2} + \frac{3}{\tilde{\alpha}}} \quad (4.27)$$

,

$$\eta_2 = \frac{n! \Gamma_{2n+1}}{\sqrt{2} \epsilon \beta^2 (-1)^n (2n+1)!} \frac{d^2}{\ell^2} = \frac{n! 2^{n+3}}{(2n+1)(-1)^n} \sqrt{\frac{2\sqrt{\pi}}{(2n+1)!}} \frac{\tilde{\alpha}^2}{CG(4\tilde{\alpha} + 3\ell^2)}. \quad (4.28)$$

By comparing the above identities one can infer that

$$\frac{\sigma_1}{\eta_1} = \frac{\sigma_2}{\eta_2} = \sqrt{2(2n+1)}. \quad (4.29)$$

It is easy to see that for fixed values of parameters of the model called as $\{M, \ell, \tilde{\alpha}, C, G\}$ one can call dimensionless eigenvalues for mass and AdS radiuses of the discretized (quantum) 4DAdSGB black hole metric surrounded by NG cloud of string as follows.

$$\sigma_1 = -2m_{2n}, \quad \sigma_2 = \frac{1}{\ell_{2n}^2}, \quad \eta_1 = -2m_{2n+1}, \quad \eta_2 = \frac{1}{\ell_{2n+1}^2}. \quad (4.30)$$

Looking at the relations (4.3), (4.10) and metric solutions (4.23) and (4.24), one can infer that the quantized distance

$$d_n = \sqrt{\frac{\tilde{\alpha}}{2}} \left(n + \frac{1}{2} \right) \quad (4.31)$$

is a particular region for a fixed regularized GB parameter $\tilde{\alpha}$ where the quantum behavior of this black hole appears but not for distances less than it. In fact conditions given by (4.25), (4.26), (4.27) and (4.28) are quantization conditions on the black hole solution. They make the obtained black hole metric solution be discretized. To end of this section we substitute (4.30) into the metric solutions (4.20) and (4.21) to obtain exact form of the discretized metric for the 4DAdSGB black hole surrounded with string cloud. In this sense we obtain discretized metric solutions for even and odd Hermite eigenvalues as follows.

$$f_{2n}(y) = 1 + \frac{n!}{(-1)^n(2n)!} e^{-y^2/2} H_{2n}(y) \left(1 + \frac{\epsilon}{2} - \frac{2m_{2n}}{y} + \frac{y^2}{\ell_{2n}^2} \right) \quad (4.32)$$

and

$$f_{2n+1}(y) = 1 + \frac{n!}{(-1)^n(2n+1)!} e^{-y^2/2} H_{2n+1}(y) \left(1 + \frac{\epsilon}{2} - \frac{2m_{2n+1}}{y} + \frac{y^2}{\ell_{2n+1}^2} \right). \quad (4.33)$$

We now study thermodynamics behavior of the quantum black hole for different quantum states $n = 0, 1, 2, 3, \dots$ as follows.

5 Thermodynamics of eigen states

Using the eigen (discretized) metric potentials (4.32) and (4.33) one can obtain eigen states for event horizons by solving $f_n(y_+) = 0$ for *even* and *odd* states as follows.

$$m_{2n} = \frac{y_+}{2} \left(1 + \frac{\epsilon}{2} + \frac{8\pi}{3} p_{2n} y_+^2 + \frac{(-1)^n (2n)! e^{y_+^2/2}}{n! H_{2n}(y_+)} \right) \quad (5.1)$$

and

$$m_{2n+1} = \frac{y_+}{2} \left(1 + \frac{\epsilon}{2} + \frac{8\pi}{3} p_{2n+1} y_+^2 + \frac{(-1)^n (2n+1)! e^{y_+^2/2}}{n! H_{2n+1}(y_+)} \right) \quad (5.2)$$

where dimensionless ADM quantum mass of the black hole m_{2n} and m_{2n+1} are called as eigen enthalpy of the black hole in the extended thermodynamics of the black holes and they are described versus the black hole horizon y_+ for *even* and *odd* states respectively. In context of the extended phase of black hole thermodynamics the eigen radiuses of the AdS space treat as thermodynamic eigen pressure as

$$p_{2n} = \frac{3}{8\pi \ell_{2n}^2}, \quad p_{2n+1} = \frac{3}{8\pi \ell_{2n+1}^2} \quad (5.3)$$

and corresponding eigen thermodynamics volume is obtained from the equation $v_n = \frac{\partial m_n}{\partial p_n}$ which by substituting (5.1) and (5.2) reads

$$v_n = \frac{4\pi}{3} y_+^3. \quad (5.4)$$

It is easy to see that the above quantized thermodynamics eigen volume is equal to geometrical volume of the discretized black hole and corresponding event horizon radius y_+ should be substituted from $f_n(y_+) = 0$ for *even* and *odd* eigen states respectively. Applying the definition of the black hole Hawking temperature $T = \frac{1}{4\pi} \frac{df(y)}{dy} \Big|_{y=y_+}$ we obtain eigen temperatures for *even* and *odd* states as follows.

$$T_{2n} = \frac{1}{4\pi y_+} + \frac{y_+}{4\pi} - \frac{n H_{2n-1}(y_+)}{\pi H_{2n}(y_+)} + \frac{n! e^{-y_+^2/2} H_{2n}(y_+)}{4\pi (-1)^n (2n)!} \left(\frac{1 + \frac{\epsilon}{2} + 8\pi p_{2n} y_+^2}{y_+} \right) \quad (5.5)$$

and

$$T_{2n+1} = \frac{1}{4\pi y_+} + \frac{y_+}{4\pi} \quad (5.6)$$

$$- \frac{(2n+1)H_{2n}(y_+)}{2\pi H_{2n+1}(y_+)} + \frac{n!e^{-y_+^2/2}H_{2n+1}(y_+)}{4\pi(-1)^n(2n+1)!} \left(\frac{1 + \frac{\epsilon}{2} + 8\pi p_{2n+1}y_+^2}{y_+} \right).$$

Entropy of this quantum black hole is obtained for *even* and *odd* eigen states respectively as follows.

$$s_{2n} = \int_0^{y_+} \frac{1}{T_{2n}} \left(\frac{\partial m_{2n}}{\partial y_+} \right) dy_+ = \int \Sigma_{p_{2n}}^\epsilon(y_+) dy_+ \quad (5.7)$$

and

$$s_{2n+1} = \int_0^{y_+} \frac{1}{T_{2n+1}} \left(\frac{\partial m_{2n+1}}{\partial y_+} \right) dy_+ = \int \Sigma_{p_{2n+1}}^\epsilon(y_+) dy_+ \quad (5.8)$$

in which we defined

$$\Sigma_{p_{2n}}^\epsilon(y_+) = \frac{\frac{1}{2} + \frac{\epsilon}{4} + 4\pi p_{2n}y_+^2 + \frac{(-1)^n(2n)!e^{y_+^2/2}}{2n!} \left(\frac{(1+y_+^2)H_{2n}(y_+) - 4nH_{2n-1}(y_+)}{H_{2n}^2(y_+)} \right)}{\frac{1}{4\pi y_+} + \frac{y_+}{4\pi} - \frac{nH_{2n-1}(y_+)}{\pi H_{2n}(y_+)} + \frac{n!e^{-y_+^2/2}H_{2n}(y_+)}{4\pi(-1)^n(2n)!} \left(\frac{1 + \frac{\epsilon}{2} + 8\pi p_{2n}y_+^2}{y_+} \right)} \quad (5.9)$$

and

$$\Sigma_{p_{2n+1}}^\epsilon(y_+) = \frac{\frac{1}{2} + \frac{\epsilon}{4} + 4\pi p_{2n+1}y_+^2 + \frac{(-1)^n(2n+1)!(1+y_+^2)e^{y_+^2/2}}{2n!H_{2n+1}(y_+)} - \frac{(-1)^n(2n+1)(2n+1)!y_+e^{y_+^2/2}H_{2n}(y_+)}{n!H_{2n+1}^2(y_+)}}{\frac{1}{4\pi y_+} + \frac{y_+}{4\pi} - \frac{(2n+1)H_{2n}(y_+)}{2\pi H_{2n+1}(y_+)} + \frac{n!e^{-y_+^2/2}H_{2n+1}(y_+)}{4\pi(-1)^n(2n+1)!} \left(\frac{1 + \frac{\epsilon}{2} + 8\pi p_{2n+1}y_+^2}{y_+} \right)} \quad (5.10)$$

Applying the above formulas one can obtain the Gibbs free energy of this quantum black hole for *even* and *odd* states as

$$G_n(y_+) = m_n(y_+) - T_n(y_+)s_n(y_+) \quad (5.11)$$

which its form is long and so does not shown here. As an example we continue our thermodynamics study for ground state $n = 0$ and first excited state $n = 1$ in the next subsection.

5.1 Thermodynamics phase transition in ground state

$$n = 0$$

By substituting $n = 0$ into the obtained general form of equation of states in the previous section we obtain enthalpy $m_0(y_+)$, the Hawking temperature $T_0(y_+)$ and the entropy $s_0(y_+)$ respectively as follows.

$$m_0(y_+) = \frac{y_+}{2} \left(1 + \frac{\epsilon}{2} + \frac{8\pi y_+^2}{3} p_0 + e^{y_+^2/2} \right) \quad (5.12)$$

$$T_0(y_+) = \frac{1}{4\pi y_+} + \frac{y_+}{4\pi} + \frac{e^{-y_+^2/2}}{4\pi} \left(\frac{1 + \frac{\epsilon}{2} + 8\pi p_0 y_+^2}{y_+} \right) \quad (5.13)$$

$$s_0(y_+) = 2\pi(e^{y_+^2/2} - 1). \quad (5.14)$$

Looking at the temperature equation (5.13) and comparing with the ideal gas equation of state one can infer that the corresponding specific volume in ground state become

$$\nu_0 = 2y_+ e^{-y_+^2/2}. \quad (5.15)$$

By eliminating y_+ between this specific volume equation and the Hawking temperature (5.13) we can obtain PVT imperfect equation of state of the quantum black hole in ground state as follows.

$$p_0(\nu_0) = -\frac{(1 + \frac{\epsilon}{2})e^{L(\nu_0)}}{4\pi\nu_0} + e^{\frac{\nu_0^2 e^{-2L(\nu_0)}}{8}} \left(\frac{T_0 e^{L(\nu_0)}}{\nu_0} - \frac{e^{2L(\nu_0)}}{2\pi\nu_0^2} - \frac{1}{8\pi} \right) \quad (5.16)$$

where we defined

$$L(\nu_0) = \frac{1}{2} \text{LambertW} \left(-\frac{\nu_0^2}{2} \right) \quad (5.17)$$

One can obtain critical point by solving the equations $\frac{\partial p_0}{\partial \nu_0} = 0$ and $\frac{\partial^2 p_0}{\partial \nu_0^2} = 0$ at constant temperature. In fact these equations are equivalent with the equations $\frac{\partial p_0}{\partial y_+} = 0$ and $\frac{\partial^2 p_0}{\partial y_+^2} = 0$ at constant temperature which lead to the following parametric critical points if we substitute the pressure $p_0(y_+)$ given by (5.13) into them.

$$T_{0c} = \frac{y_{+c}^3(5 + y_{+c}^2)}{4\pi(y_{+c}^4 + 2y_{+c}^2 - 1)}, \quad p_{0c} = \frac{(3 + y_{+c}^4)e^{y_{+c}^2/2}}{16\pi(y_{+c}^4 + 2y_{+c}^2 - 1)}, \quad \nu_{0c} = 2y_{+c}e^{-y_{+c}^2/2} \quad (5.18)$$

in which the critical radius of the event horizon in ground state y_{+c} is determined versus the string tension ϵ by solving the following equation

$$\epsilon = \frac{2 - 4y_{+c}^2 - 2y_{+c}^4 + e^{y_{+c}^2/2}(2 - 5y_{+c}^2 + 4y_{+c}^4 - y_{+c}^6)}{y_{+c}^4 + 2y_{+c}^2 - 1}. \quad (5.19)$$

We plotted diagrams of the above equation $\epsilon(y_{+c})$ and all possible numerical values for critical points given by (5.18) in figure 1. To continue study of thermodynamics of this black hole we should choose sample numerical value from diagrams of the figure 1 for instance $\epsilon = -1$. In this case we obtain

$$\begin{aligned} y_{+c} &= 0.7394220670, & \nu_{0c} &= 1.125118423, \\ p_{0c} &= 0.2198245577, & T_{0c} &= 0.4547307702 \end{aligned} \quad (5.20)$$

in which critical enthalpy of this black hole is obtained as

$$m_{0c} = 1.043056198 \quad (5.21)$$

which is useful to study Joule-Thomson expansion by plotting isenthalpic $T - P$ diagrams. $p - v$ diagram in figure 2 shows a small to large black hole phase transition which is happened at temperature higher than the critical temperature $T_0 > T_{0c}$ at ground state $n = 0$. Other diagrams namely $p - y_+$ or $T - v$ predict similar situations. We know that a thermodynamic system is stable when its Gibbs free energy takes negative values. For this black hole it is seen at figure 3 for $G - v$ or $G - y_+$ diagrams. In these diagrams minimum point of the Gibbs free energy with negative value is at bigger size black holes. This means that the black hole participates at small to large black hole phase transition. This is happened at pressures higher than the critical pressure $p_0 > p_{0c}$. One can see from $G - v$ or $G - y_+$ diagrams that the enormous black holes have not negative Gibbs energy which means that this quantum black holes at ground state can not to participate in the Hawking-Page phase transition where a black hole evaporates and its final state reaches to a vacuum AdS space. We will see this is possible in the first excited states by plotting $G - v$ or $G - y_+$ curves of this black hole (see figure 7 in the next section).

5.2 Thermodynamics phase transition in first excited state $n = 1$

By substituting $n = 1$ into the obtained general form of equation of states in the previous section we obtain enthalpy $m_1(y_+)$, the Hawking temperature

$T_1(y_+)$ and the entropy $s_1(y_+)$ respectively as follows.

$$m_1(y_+) = \frac{y_+}{2} \left(1 + \frac{\epsilon}{2} + \frac{8\pi y_+^2}{3} p_1 + \frac{e^{y_+^2/2}}{2y_+} \right) \quad (5.22)$$

$$T_1(y_+) = \frac{y_+}{4\pi} + \frac{e^{-y_+^2/2}}{2\pi} \left(1 + \frac{\epsilon}{2} + 8\pi p_1 y_+^2 \right) \quad (5.23)$$

$$s_1(y_+) = -i\pi \sqrt{\frac{\pi}{2}} \operatorname{erf} \left(\frac{iy_+}{\sqrt{2}} \right). \quad (5.24)$$

By looking at the temperature equation (5.23) and by comparing with the ideal gas equation of state one can infer that the corresponding specific volume in first excited state become

$$\nu_1 = 4y_+^2 e^{-y_+^2/2}, \quad y_+^{(\pm)} = \pm \sqrt{2L(\nu_1)} \quad (5.25)$$

where we defined

$$L(\nu_1) = -\operatorname{LambertW} \left(-\frac{\nu_1}{96} \right). \quad (5.26)$$

By eliminating y_+ between this specific volume equation (5.25) and the Hawking temperature (5.23) we can obtain PVT equation of state of the quantum black hole in first excited state as follows.

$$p_1^{(\pm)}(\nu_1) = \frac{192\pi L(\nu_1) T_1 \mp 48\sqrt{2} L(\nu_1)^{\frac{3}{2}} - (1 + \frac{\epsilon}{2})\nu_1}{16\pi\nu_1 L(\nu_1)}. \quad (5.27)$$

One can obtain critical point by solving the equations $\frac{\partial p_1}{\partial \nu_1} = 0$ and $\frac{\partial^2 p_1}{\partial \nu_1^2} = 0$. We checked these equations to obtain some real positive values for the critical specific volume ν_{1c} and saw that there are complex (analytic continuation) values for the critical event horizons y_{+c} which still give out real values for the specific volume ν_{1c} . To exclude these analytic continuation horizon radiuses we solve the equivalent critical points equations $\frac{\partial p_1}{\partial y_+} = 0$ and $\frac{\partial^2 p_1}{\partial y_+^2} = 0$ to obtain real positive values for the critical radius of the horizons and the corresponding specific volume of the black hole horizons as follows. After solving the equations $\frac{\partial p_1}{\partial y_+} = 0$ and $\frac{\partial^2 p_1}{\partial y_+^2} = 0$ and some simple mathematical derivations we obtain general form of the critical points against y_{+c} as follows.

$$T_{1c} = \frac{y_{+c}^4 + 2y_{+c}^2 - 1}{4\pi y_{+c}^3}, \quad p_{1c} = \frac{(y_{+c}^2 - 1)e^{-y_{+c}^2/2}}{32\pi y_{+c}^3}, \quad \nu_{1c} = 4y_{+c}^2 e^{-y_{+c}^2/2} \quad (5.28)$$

where the critical radius of the event horizon y_{+c} is obtained from largest positive real root of the following equation.

$$\epsilon = -2 - \frac{e^{y_{+c}^2/2}}{2} \left(y_{+c} - \frac{5}{y_{+c}} + \frac{2}{y_{+c}^3} \right). \quad (5.29)$$

This root will be correspond to exterior horizon of the black hole which is visible for observers located at outside of the black hole. We show all possible real roots of the above equation for a given string tension ϵ in figure 4. We continue our study as numerically by choosing a particular value from the figure 4 as follows.

$$\nu_{1c} = 2.198851498, \quad p_{1c} = 0.0005220136425, \quad T_{1c} = 0.2279369128, \quad (5.30)$$

for

$$\epsilon = -1, \quad y_{+c} = 1.984472401 \quad (5.31)$$

where critical black hole enthalpy at $n = 1$ state is obtained as follows

$$m_{1c} = 2.304201063 \quad (5.32)$$

which is applicable to study isenthalpic $T - p$ curves and Joule-Thomson expansion of the system. All possible critical points given by the equations (5.28) are plotted versus the string tension ϵ at first excited state $n = 1$ in figure 4. By looking at the isobaric $T - v$ and $T - y_+$ curves in figure 5 one can understand that the temperature has local maximum (minimum) value for smaller (larger) black holes at pressures higher than the critical pressure. This means that the black hole participates at the small to large scale black hole phase transition for pressures higher than the critical pressure. Isobaric $T - v$ curves show that the black hole have maximum temperature at largest scale at $p > p_c$. This diagram shows that the black hole can not to participate at the Hawking-Page phase transition apparently where the black hole evaporates and finally reaches to the Ads vacuum space. But by looking at $G - v$ or $G - y_+$ isobaric curves one can infer that negativity sign of the Gibbs free energy is happened at large black holes and huge sized black holes (see figures 7 and 8) in which the first small regions with $G < 0$ is related to the small to large phase transition while the larger regions with $G < 0$ is corresponds to the Hawking-Page phase transition [69]. One can obtain similar statements by looking at the isothermal $p - v$ curves given by figure 6. In context of black holes thermodynamics the small to large scale black hole phase transition is

confirmed that is similar to solid-fluid-gas phase transition in imperfect Van der Waals fluid. In fact the diagrams show different behavior for the black hole in the below and the above of the critical points of the phase space. If we want to explain accurately, in fact the small to large black hole phase transition is happened by connecting an unstable medium size black hole near the maximum point of $p - v$ diagram in the ground state (see figure 2) and the first excited state (see figure 7). In any thermodynamic system always exist a phase with minimum value for its Gibbs free energy. When two branches of Gibbs free energy at minimum state cross each other then the phase transition is happened, while if these two branches have the same value of Gibbs free energy then this two phases are called coexist. This swallowtail behavior indicates coexistence of two different phases which is happened in ground state $n = 0$ (see G-T curve at constant pressure $p_0 > p_{0c}$ in figure 3) and at first excited state $n = 1$ (see G-T curve at constant pressure $p_1 > p_{1c}$ in figure 8). At end of this section we can understand by looking the G-y or G-v curves given in the figures 3 and 8 where an unstable small black hole with $G > 0$ reaches to an stable large black hole with $G < 0$. While an unstable huge black hole with $G > 0$ evaporate completely and reduces to a vacuum AdS space with $G < 0$ by participating in the Hawking-Page phase transition.

6 Conclusion

According to the 4DAdS EGB gravity model [31] we obtained discretized metric potentials for spherically symmetric static quantum black holes surrounded with cloud of string which is described by the Hermite polynomials (spherical quantum harmonic oscillator). Thus we called our obtained solutions as quantized (quantumized) black hole. We should point that our used ‘quantization’ adjective for this black hole metric solution is different with the well known canonical quantization. In fact we used this adjective because of two reasons as follows: (a) The metric solution is obtained by the Hermite polynomials with ‘discrete’ eigenvalues. (b) The used EGB gravity as an higher order derivative of metric theory originates from renormalization of quantum fields propagating in curved space times. We used the extended thermodynamic formalism to show that 4D AdS EGB black holes surrounded by strings fluid mimics Van der Waals fluid behavior. In extended thermodynamic paradigm, pressure of AdS background spacetime which is affect on a

local black hole metric is described by negative cosmological constant and so its conjugate variable plays the role of thermodynamic volume of the black hole. To examine the analogy between 4D AdS EGB black holes surrounded by cloud of strings with ordinary liquid-gas thermodynamic system we derived analytical solutions of the critical points in different eigenstates which are fixed by the string tension. With in detail we studied thermodynamics of this black hole at the ground state and first excited state respectively and obtained dependence of the black hole thermodynamic phase transition to the black hole quantum states. In this sense the black hole in ground state participates just at the phase transition of small to large scale black holes if it has small Gibbs energy while it participates at the Hawking-page phase transition if it has large Gibbs free energy. On the other side for a black hole with small Gibbs free energy when is at the first excited state there is possibility of participation of the black hole in both of the small to large and also the Hawking-Page phase transitions. Also we checked diagrams of the isenthalpic P-T curves and obtained that this black hole does not participates in the Joule-Thomson expansion phenomena in ground state and first excited state. As a future work we like to study canonical quantization of this black hole by solving its Wheeler De Witt wave equation and obtain its relation with the thermodynamic phase transition by comparing with our obtained results in this work.

References

- [1] B. Fauser, J. Tolksdorf and E. Zeidler, '*Quantum Gravity, Mathematical models and Experimental Bounds*', (Birkhauser verlag, P.O.Box133, CH-4010 Basel, Switzerland 2007).
- [2] Birrell and Davies, '*Quantum fields in curved space*', (Cambridge University press, England, 1982).
- [3] L. Parker and D. Toms, '*Quantum Field Theory in Curved Spacetime, Quantized fields and Gravity*', (Cambridge University press, 2009).
- [4] H. Ghaffarnejad, 'Quantum field backreaction corrections and remnant stable evaporating Schwarzschild de Sitter dynamical black hole', Phys. Rev. D75, 084009, (2007).

- [5] B. Zwiebach, '*A first course in string theory*', (Cambridge University press 2004).
- [6] M. Gasperini '*Elements of String Cosmology*' (Cambridge University press 2007).
- [7] F. Bastianelli and P. V. Nieuwenhuizen, '*Path integrals and anomalies in curved space*', (Cambridge University press 2006).
- [8] A. Hatzinikitas, K. Schalm, P.v. Nieuwenhuizen, '*Trace and chiral anomalies in string and ordinary field theory from Feynman diagrams for nonlinear sigma models*', Nucl.Phys. **B518**, 424 (1998); arXiv:hep-th/9711088.
- [9] M. Henningson and K. Skenderis, '*The Holographic Weyl Anomaly*', JHEP **9807**, 023 (1998).
- [10] D. Lovelock, '*The Einstein Tensor and Its Generalizations*', J. Math. Phys. (N.Y.)**12**, 498 (1971).
- [11] B. Zwiebach, '*Curvature squared terms and string theories*', Phys. Lett. B **156**, 315 (1985).
- [12] B. Zumino, '*Gravity theories in more than four dimensions*', Phys. Rept.**137**, 109 (1986).
- [13] N. Deruelle and L. Farina-Busto, '*Lovelock gravitational field equations in cosmology*', Phys. Rev. D**41**, 3696 (1990).
- [14] R. G. Cai and N. Ohta, '*Black Holes in Pure Lovelock Gravities*', Phys. Rev. D**74**, 064001 (2006).
- [15] D.G. Boulware, S. Deser, '*String-Generated Gravity Models*', Phys. Rev. Lett.**55**, 2656 (1985).
- [16] J.T. Wheeler, Nucl. '*Symmetric solutions to the Gauss-Bonnet extended Einstein equations*', Phys. **B268**, 737 (1986).
- [17] R.C.Myers, J.Z. Simon, '*Black-hole thermodynamics in Lovelock gravity*', Phys. Rev. D**38**, 2434 (1988).

- [18] R. P. Woodard, '*The Theorem of Ostrogradsky*' arXiv:1506.02210 [hep-th].
- [19] H. Motohashi and T. Suyama, '*Third order equations of motion and the Ostrogradsky instability*', Phys. Rev. D **91**, 085009 (2015).
- [20] C. Lanczos, '*A Remarkable property of the Riemann-Christoffel tensor in four dimensions*', Annals Math**39**, 842 (1938).
- [21] S. G. Ghosh and S. D. Maharaj, '*Cloud of strings for radiating black holes in Lovelock gravity*', Phys. Rev. D **89**, 084027 (2014); arXiv:1409.7874 [gr-qc].
- [22] D. J. Gross and E. Witten, '*Superstring modifications of Einstein's equations*', Nucl. Phys.B**277**, 1 (1986).
- [23] D. J. Gross and J. H. Sloan, '*The quartic effective action for the heterotic string*', Nucl. Phys.B**291**, 41 (1987).
- [24] R.R. Metsaev and A. A. Tseytlin, '*Two-loop β -function for the generalized bosonic sigma model*', Phys. Lett. B**191**, 354 (1987).
- [25] B. Zwiebach, '*Curvature squared terms and string theories*', Phys. Lett. B**156**, 315 (1985).
- [26] R. R. Metsaev and A. A. Tseytlin, '*Order α (two-loop) equivalence of the string equations of motion and the σ -model Weyl invariance conditions: Dependence on the dilaton and the antisymmetric tensor*', Nucl. Phys.B**293**, 385 (1987).
- [27] D. Glavan and C. Lin, '*Einstein-Gauss-Bonnet Gravity in 4D Spacetime*', Phys. Rev. Lett. **124**, 081301 (2020), [arXiv:1905.03601 [gr-qc]].
- [28] J. Arrechea, A. Delhom and A. J. Cano, '*Inconsistencies in four-dimensional Einstein-Gauss-Bonnet gravity*', arXiv:2004.12998 [gr-qc].
- [29] J. Arrechea, A. Delhom and A. J. Cano, '*Comment on Einstein-Gauss-Bonnet Gravity in Four-Dimensional Spacetime*', Phys. Rev. Lett. **125**, 149002; arXiv:2009.10715 [gr-qc].

- [30] M. Gurses, T. C. Sisman and B. Tekin, '*Comment on Einstein-Gauss-Bonnet Gravity in Four-Dimensional Spacetime*', Phys. Rev. Lett. **125**, 149001 (2020).
- [31] K. Aoki, M. A. Gorji and S. Mukohyama, '*A consistent theory of $D=4$ Einstein-Gauss-Bonnet gravity*', Phys. Lett. **B810**, 135843 (2020); [arXiv:2005.03859[gr-qc]].
- [32] K. Aoki, M. A. Gorji and S. Mukohyama, '*Cosmology and gravitational waves in consistent $D \rightarrow 4$ Einstein-Gauss-Bonnet gravity*', JCAP**2020**, 014 (2020); [arXiv:2005.08428 [gr-qc]].
- [33] D. V. Singh, S. G. Ghosh and S.D. Maharaj, '*Clouds of strings in the novel 4D Einstein-Gauss-Bonnet black holes*', Phys. Dark Univ. **30** 100730 (2020); [arXiv:2003.14136 [gr-qc]].
- [34] K. Hegde, A. N. Kumara, C. L. A. Rizwan, Ajith K. M, and M. S. Ali, '*Thermodynamics, Phase Transition and Joule Thomson Expansion of novel 4-D Gauss Bonnet AdS Black Hole*', [arXiv:2003.08778 [gr-qc]].
- [35] R Cai, L. Cao, and R. Yang,, '*P-V criticality in the extended phase space of Gauss-Bonnet black holes in AdS space*', JHEP, **1309**, 005 (2013).
- [36] J. Mo, G. Li, and X. Xu, '*Combined effects of $f(R)$ gravity and conformally invariant Maxwell field on the extended phase space thermodynamics of higher-dimensional black holes*', Eur. Phys. J. C**76** 545 (2016).
- [37] H. Ghaffarnejad and E. Yaraie, '*Effects of a cloud of strings on the extended phase space of Einstein-Gauss-Bonnet AdS black holes*', Phys. Lett.B, **785**, 105 (2018).
- [38] E. Herscovich and M. G. Richarte, '*Black holes in Einstein Gauss-Bonnet gravity with a string cloud background*', Phys. Lett.B **689**, 192, (2020).
- [39] S. G. Ghosh and S. D. Maharaj, '*Cloud of strings for radiating black holes in Lovelock gravity*', Phys. Rev. D **89**, 084027 (2014).
- [40] S. G. Ghosh, U. Papnoi and S. D. Maharaj, '*Cloud of strings in third order Lovelock gravity*', Phys. Rev. D **90**, 044068 (2014).

- [41] T. H. Lee, D. Baboolal and S. G. Ghosh, '*Lovelock black holes in a string cloud background*', Eur. Phys. J. C **75**, 297 (2015).
- [42] R. A. Konoplya and A. F. Zinhailo, '*Quasinormal modes, stability and shadows of a black hole in the novel 4D Einstein Gauss-Bonnet gravity*', Eur. Phys. J. C **80**, 1049 (2020); [arXiv:2003.01188 [gr-qc]].
- [43] M. Guo and P. C. Li, '*The innermost stable circular orbit and shadow in the novel 4D Einstein-Gauss-Bonnet gravity*', Eur. Phys. J. C **80**, 588 (2020) ; [arXiv:2003.02523 [gr-qc]].
- [44] P. G. S. Fernandes, '*Charged Black Holes in AdS Spaces in 4D Einstein Gauss-Bonnet Gravity*', Phys. Lett. B **805**, 135468 (2020); [arXiv:2003.05491[gr-qc]].
- [45] Y.P. Zhang, S.W. Wei, and Y.X. Liu, '*Spinning test particle in four-dimensional Einstein-Gauss-Bonnet Black Hole*', [arXiv:2003.10960 [gr-qc]].
- [46] H. Lu and Y. Pang, '*Horndeski Gravity as $D \rightarrow 4$ Limit of Gauss-Bonnet*', Phys. Lett. B **809** , 135717 (2020); [arXiv:2003.11552 [gr-qc]].
- [47] D. V. Singh and S. Siwach, '*Thermodynamics and P - v criticality of Bardeen-AdS Black Hole in 4-D Einstein-Gauss-Bonnet Gravity*', Phys. Lett. B **808**, 135658 (2020); [arXiv:2003.11754 [gr-qc]].
- [48] R. A. Konoplya, A. Zhidenko, '*BTZ black holes with higher curvature corrections in the 3D Einstein-Lovelock theory*', Phys. Rev. D **102**, 064004 (2020); [arXiv:2003.12171 [gr-qc]].
- [49] S. G. Ghosh and R. Kumar, '*Generating black holes in the novel 4D Einstein-Gauss-Bonnet gravity*', [arXiv:2003.12291 [gr-qc]].
- [50] R. A. Konoplya and A. Zhidenko, '*(In)stability of black holes in the 4D Einstein-Gauss-Bonnet and Einstein-Lovelock gravities*', Phys. Dark Universe **30**, 100697 (2020); [arXiv:2003.12492 [gr-qc]].
- [51] T. Kobayashi, '*Effective scalar-tensor description of regularized Lovelock gravity in four dimensions*', JCAP **07**, 013 (2020); [arXiv:2003.12771 [gr-qc]].

- [52] A. Kumar and R. Kumar, '*Bardeen black holes in the novel 4D Einstein-Gauss-Bonnet gravity*', [arXiv:2003.13104 [gr-qc]].
- [53] C. Y. Zhang, P. C. Li, and M. Guo, '*Greybody factor and power spectra of the Hawking radiation in the novel 4D Einstein-Gauss-Bonnet de-Sitter gravity*', Eur. Phys. J. C **80**, 874 (2020) [arXiv:2003.13068 [hep-th]].
- [54] S. A. H. Mansoori, '*Thermodynamic geometry of novel 4D Gauss Bonnet AdS Black Hole*', [arXiv:2003.13382 [gr-qc]].
- [55] S.W. Wei and Y.X. Liu, '*Testing the nature of Gauss-Bonnet gravity by four-dimensional rotating black hole shadow*', [arXiv:2003.07769 [gr-qc]].
- [56] R. Kumar and S. G. Ghosh, '*Rotating black holes in the novel 4D Einstein Gauss Bonnet gravity*', JCAP **2020**, 053 (2020); [arXiv:2003.08927[gr-qc]].
- [57] Jia Xi Feng, Bao Min Gu, Fu Wen Shu, '*Theoretical and observational constraints on regularized 4D Gauss-Bonnet Gravity*', arxiv: 2006.16751[gr-qc]
- [58] Fu Wen Shu '*Vacua in novel 4D Einstein Gauss Bonnet Gravity: pathology and instability?*', arXiv:2004.09339
- [59] K.Yang, B.M.Gu, S.W.Wei and Y.X.Liu '*Born-Infeld Black Holes in novel 4D Einstein-Gauss-Bonnet gravity*', arXiv:2004.14468
- [60] A. Aragon, R. Becar, P. A. Gonzalez, Y. Vasquez, '*Perturbative and nonperturbative quasinormal modes of 4D Einstein Gauss Bonnet black holes*', Eur.Phys.J. C **80** 773 (2020); arXiv:2004.05632 [gr-qc].
- [61] R. A. Konoplya and A. F. Zinhailo, '*Quasinormal modes, stability and shadows of a black hole in the novel 4D Einstein-Gauss-Bonnet gravity*', Eur. Phys. J. C **80**, 1049 (2020); arXiv:2003.01188 [gr-qc].
- [62] R.A. Konoplya and A. F. Zinhailo, '*Grey body factors and Hawking radiation of black holes in 4D Einstein-Gauss-Bonnet gravity*', Phys. Lett. B **810**, 135793 (2020); [arXiv:2004.02248 [gr-qc]].
- [63] V. Oikonomou and F. Fronimos, '*Reviving non-Minimal Horndeski-like Theories after GW170817: Kinetic Coupling Corrected Einstein-Gauss-Bonnet Inflation*', [arXiv:2006.05512 [gr-qc]].

- [64] S. Odintsov, V. Oikonomou and F. Fronimos, ‘*Rectifying Einstein Gauss Bonnet Inflation in View of GW170817*’, [arXiv:2003.13724 [gr-qc]].
- [65] S. D. Odintsov, V. K.Oikonomou and F. P.Fronimos, ‘*Non-Minimally Coupled Einstein Gauss Bonnet Inflation Phenomenology in View of GW170817*’,[arXiv:2007.02309 [gr-qc]].
- [66] V. K. Oikonomou and F. P. Fronimos, ‘*A Nearly Massless Graviton in Einstein Gauss Bonnet Inflation with Linear Coupling Implies Constant-roll for the Scalar Field,*, [arXiv:2007.11915 [gr-qc]].
- [67] S.Odintsov and V.Oikonomou, ‘*Swampland Implications of GW170817 compatible Einstein Gauss Bonnet Gravity*’, Phys. Lett. B **805** 135437 (2020); arXiv:2004.00479 [gr-qc].
- [68] S. J. Yang, J. J. Wan, J. Chen, J. Yang and Y. Q. Wang, ‘*Weak cosmic censorship conjecture for the novel 4D charged Einstein-Gauss-Bonnet black hole with test scalar field and particle*’, Eur. Phys. J. C**80**, 937 (2020); [arXiv:2004.07934 [gr-qc]].
- [69] S. W. Hawking, D.N. Page, ‘*Thermodynamics of black holes in anti-de Sitter space*’, Comm. Math. Phys. **87** 577 (1983).
- [70] D. Kastor, S. Ray, and J. Traschen, ‘*Enthalpy and mechanics of AdS black holes*’, Class. Quant. Grav. **26** 195011 (2009).
- [71] B. P. Dolan, ‘*The cosmological constant and black hole thermodynamic potential*’, Class. Quant. Grav. **28** 125020 (2011).
- [72] D. Kubiznak and R. B. Mann, ‘*P-V criticality of charged AdS black holes*’, JHEP **1207**, 033, (2012).
- [73] S. Dutta, A. Jain and R. Soni, ‘*Dyonic Black Hole and Holography*’, JHEP **2013**, 60 (2013) [arXiv:1310.1748 [hep-th]].
- [74] Gu-Qiang Li, ‘*Effects of dark energy on P-V criticality of charged AdS black holes*’, Phys. Lett. B **735**, 256 (2014).
- [75] M. Zhang and W Liu,l, ‘*Coexistent physics of massive black holes in the phase transitions*’, [arXiv:1610.03648 [gr-qc]].

- [76] R. G. Cai, Y.P. Hu, Q.Y. Pan, and Y.L. Zhang, '*Thermodynamics of black holes in massive gravity*', Phys. Rev. D **91**, 024032; (2015); arXiv:1409.2369 [hep-th].
- [77] R. Zhao, H. H. Zhao, M. S. Ma and L. C. Zhang, '*On the critical phenomena and thermodynamics of charged topological dilaton AdS black holes*', Eur. Phys. J. C **73**, 2645(2013).
- [78] J. X. Mo, X. X. Zeng, G. Q. Li, X. Jiang, W. B. Liu, '*A unified phase transition picture of the charged topological black hole in Horava-Lifshitz gravity*', JHEP **1310**, 056(2013).
- [79] N. Altamirano, D. Kubiznak, R. Mann and Z. Sherkatghanad, '*Kerr AdS analogue of tricritical point and solid-liquid gas phase transition*', Class. Quantum Grav. **31**, 042001(2014); arXiv:1308.2672 [hep-th].
- [80] J. X. Mo and W. B. Liu, '*Ehrenfest scheme for P-V criticality in the extended phase space of black holes*', Phys. Lett. B **727**, 336 (2013).
- [81] N. Altamirano, D. Kubiznak and R. Mann, '*Reentrant phase transitions in rotating AdS black holes*', Phys. Rev. D **88**, 101502 (2013).
- [82] D. C. Zou, S. J. Zhang and B. Wang, '*Critical behavior of Born-Infeld AdS black holes in the extended phase space thermodynamics*', Phys. Rev. D **89**, 044002 (2014).
- [83] R. Zhao, H. H. Zhao, M. S. Ma and L. C. Zhang, '*On the critical phenomena and thermodynamics of charged topological dilaton AdS black holes*', Eur. Phys. J. C **73**, 2645(2013).
- [84] H. Ghaffarnejad, E. Yaraie and M. Farsam '*Quintessence Reissner Nordström anti de Sitter black holes and Joule Thomson effect*', Int. J. Theor. Phys. **57**,1671, (2018).
- [85] Ö. Ökcü, and E. Aydiner, '*Joule-Thomson expansion of the charged AdS black holes*', Eur. Phys. J. C **77**, 24 (2017).
- [86] D. Hansen, D. Kubiznak and R. B. Mann, '*Universality of P-V Criticality in Horizon Thermodynamics*', arXiv:1603.05689 [gr-qc].

- [87] H. Ghaffarnejad and M. Farsam. '*The last lost charge and phase transition in Schwarzschild AdS minimally coupled to a cloud of strings*', Eur. Phys. J. Plus **134**, 110 (2019).
- [88] H. Ghaffarnejad, E. Yaraie, M. Farsam, and K. Bamba, '*Hairy black holes and holographic heat engine*', Nucl. Phys.B **952**, 114941 (2020).
- [89] H. Ghaffarnejad, E. Yaraie and M. Farsam, '*Thermodynamic phase transition for quintessence dyonic anti-de Sitter black holes*', Eur. Phys. J. Plus **135**, 179 (2020).
- [90] J. X. Feng, B. M. Gu and F. W. Shu, '*Theoretical and observational constraints on regularized 4D Gauss Bonnet Gravity*', arxiv: 2006.16751 [gr-qc].
- [91] F. W. Shu, '*Vacua in novel 4D Einstein Gauss Bonnet Gravity: pathology and instability?*', Phys. Lett.B**811**, 135907, (2020); arXiv:2004.09339[gr-qc].
- [92] K. Yang, B.M.Gu, S.W.Wei and Y.X.Liu, '*Born Infeld Black Holes in novel 4D Einstein Gauss Bonnet gravity*', Eur. Phys. J. C**80**, 662 (2020); arXiv:2004.14468 [gr-qc].
- [93] A. Aragon, R. Becar, P. A. Gonzalez, Y. Vasquez, '*Perturbative and nonperturbative quasinormal modes of 4D Einstein Gauss Bonnet black holes*', Eur. Phys.J.C**80** 773 (2020); arXiv:2004.05632 [gr-qc].
- [94] A. Strominger and C. Vafa, '*Microscopic origin of the Bekenstein Hawking entropy*', Phys. Lett.B **379**, 99 (1996).
- [95] B. R. Majhi and S. Samanta, '*P-V criticality of AdS black holes in a general framework*', Phys.Lett. B**773** 203 (2017), arXiv:1609.06224 [gr-qc].
- [96] K. Bhattacharya and B. R. Majhi, '*Thermogeometric description of the van der Waals like phase transition in AdS black holes*', Phys. Rev. D **95**, 104024 (2017); arXiv:1702.07174 [gr-qc].
- [97] K. Bhattacharya, B. R. Majhi and S. Samanta, '*Van der Waals criticality in AdS black holes: a phenomenological study*', Phys. Rev. D **96**, 084037 (2017); arXiv:1709.02650 [gr-qc].

- [98] K. Bhattacharya and B. R. Majhi, '*Thermogeometric study of van der Waals like phase transition in black holes: an alternative approach*', Phys. Lett. B **802** 135224, (2020), arXiv:1903.10370 [gr-qc].
- [99] E. Caceres, P. H. Nguyen and J. F. Pedraza, '*Holographic entanglement entropy and the extended phase structure of STU black holes*', JHEP **1509** 184(2015); arXiv:1507.06069 [hep-th].
- [100] J. F. Pedraza, W. Sybesma and M. R. Visser, '*Hyperscaling violating black holes with spherical and hyperbolic horizons*', Class. Quantum Grav. **36**, 054002 (2019); arXiv:1807.09770 [hep-th].
- [101] S. W. Wei and Y. X. Liu, '*Extended thermodynamics and microstructures of four-dimensional charged Gauss Bonnet black hole in AdS space*', Phys. Rev. D **101**, 104018 (2020); arXiv:2003.14275 [gr-qc].
- [102] P. S. Letelier, '*Clouds Of Strings In General Relativity*', Phys. Rev. D **20**, 1294 (1979).
- [103] M. G. Richarte and C. Simeone, '*Traversable wormholes in a string cloud*', Int. J. Mod. Phys. D **17**, 1179 (2008).
- [104] A. K. Yadav, V. K. Yadav and L. Yadav, '*Cylindrically symmetric inhomogeneous universes with a cloud of strings*', Int. J. Theor. Phys. **48**, 568 (2009).
- [105] A. Ganguly, S. G. Ghosh and S. D. Maharaj, '*Accretion onto a black hole in a string cloud background*', Phys. Rev. D **90**, 064037 (2014).
- [106] K. A. Bronnikov, S. W. Kim and M. V. Skvortsova,, '*The Birkhoff theorem and string clouds*', Class. Quant. Grav. **33**, 195006 (2016).
- [107] N. Goldenfeld, '*Lectures on Phase Transitions and the Renormalization Group*', CRC Press, ISBN-10:9780429973123, (2018).




Figure 1: Numerical values of all possible critical points versus the string tension ϵ at ground state of the AdS 4DGB quantum black hole $n = 0$




Figure 2: Diagrams for $p(v)$, $p(y_+)$ and $T(v)$ at ground state $n=0$




Figure 3: Diagrams for Gibbs free energy at ground state $n=0$.




Figure 4: Critical points versus the string tension at first excited state $n = 1$.




Figure 5: Diagram of the temperature is plotted versus the specific volume and the event horizon at $n = 1$ state.




Figure 6: Diagram of the pressure is plotted versus the specific volume and the event horizon at $n = 1$ state.




Figure 7: Diagram of the Gibbs free energy is plotted versus the specific volume and the event horizon at $n = 1$ state.




Figure 8: $G - T$ curve is plotted in $n = 1$ state for different values of the constant pressure.