

On the equivalence of Batyrev and BHK Mirror symmetry constructions

Alexander Belavin,^{a,c,d} Boris Eremin^{a,b,c,d}

^a*Landau Institute for Theoretical Physics, 142432 Chernogolovka, Russia*

^b*Skolkovo Institute of Science and Technology, 143026 Moscow, Russia*

^c*Moscow Institute of Physics and Technology, 141700 Dolgoprudny, Russia*

^d*Kharkevich Institute for Information Transmission Problems, 127994 Moscow, Russia*

E-mail: Belavin@itp.ac.ru, Eremin.ba@phystech.edu

ABSTRACT: We consider the connection between two constructions of the mirror partner for the Calabi-Yau orbifold. This orbifold is defined as a quotient by some suitable subgroup G of the phase symmetries of the hypersurface X_M in the weighted projective space, cut out by a quasi-homogeneous polynomial W_M . The first, Berglund-Hübsch-Krawitz (BHK) construction, uses another weighted projective space and the quotient of a new hypersurface X_{M^T} inside it by some dual group G^T . In the second, Batyrev construction, the mirror partner is constructed as a hypersurface in the toric variety defined by the reflexive polytope dual to the polytope associated with the original Calabi-Yau orbifold. We give a simple evidence of the equivalence of these two constructions.

KEYWORDS: Mirror Symmetry, Calabi-Yau manifolds, Compactification

Contents

1	Introduction	1
2	Berglund–Hübsch–Krawitz mirror construction	4
3	Batyrev construction and verification of equivalence	7

1 Introduction

Mirror symmetry provides a geometrical connection between a pair of algebraic manifolds and has been studied for decades. According to mirror conjecture for a pair of Calabi-Yau threefolds (X, Y) there is an isomorphism of the cohomologies

$$H^{p,q}(X, \mathbb{C}) = H^{3-p,q}(Y, \mathbb{C}). \quad (1.1)$$

Calabi-Yau manifolds X and Y , being complex and Kähler manifolds, allow deformations of complex and Kähler structures. Thus, moduli space of complex $M_C(X)$ and Kähler $M_K(X)$ structure deformations arise [1]. The mirror symmetry can be interpreted as matching the Special geometries [2], on the moduli spaces

$$M_C(X) \simeq M_K(Y), \quad M_K(X) \simeq M_C(Y), \quad (1.2)$$

and the equality of their dimensions (or the Hodge numbers)

$$h_{21}(X) = h_{11}(Y), \quad h_{21}(Y) = h_{11}(X). \quad (1.3)$$

In this paper we consider Calabi-Yau manifolds cut out by quasi-homogeneous polynomial

$$W_M(x) = \sum_{i=1}^5 \prod_{j=1}^5 x_j^{M_{ij}} \quad (1.4)$$

in a weighted projective space

$$\mathbb{P}_{(k_1, k_2, k_3, k_4, k_5)}^4 = \left\{ (x_1, \dots, x_5) \in \mathbb{C}^5 \setminus \{0\} \mid x_i \sim \lambda^{k_i} x_i \right\}. \quad (1.5)$$

Such polynomials are also called *invertible*, meaning the number of monomials is equal to the number of variables, also the polynomial satisfy the following conditions:

- (i) the matrix M is integer, and invertible;

- (ii) the polynomial W_M is quasi-homogeneous, i.e., there exist positive integers k_i (weights of the projective space), and d (degree), such that

$$\sum_{j=1}^5 M_{ij} k_j = d = \sum_{i=1}^5 k_i, \quad \forall i; \quad (1.6)$$

- (iii) the polynomial W_M is a non-degenerate potential away from the origin. It follows that the W_M must be a sum of invertable potentials of one of three atomic types: Fermat, loop, or chain [3].

These facts provide that the W_M defines a Calabi-Yau manifold. Note some useful properties of the matrix M , and their inverse matrix $B = M^{-1}$. There also exist positive integer numbers $\bar{k}_j$, and $\bar{d}$ such that

$$\sum_{i=1}^5 \bar{k}_i M_{ij} = \bar{d} = \sum_{l=1}^5 \bar{k}_l, \quad \forall j. \quad (1.7)$$

Using the quasi-homogeneity condition (1.6),(1.7) and the definition $\sum_l B_{il} M_{lj} = \delta_{ij}$ one can obtain the relations

$$\sum_{j=1}^5 B_{ij} = \frac{k_i}{d}, \quad (1.8)$$

and

$$\sum_{i=1}^5 B_{ij} = \frac{\bar{k}_j}{\bar{d}}. \quad (1.9)$$

Calabi–Yau X_M allows the deformation of complex structure. In terms of the polynomials, it is realized as a deformation of the W_M . Thus the full family of X_M is given by zero locus of

$$W(x, \varphi) = \sum_{i=1}^5 \prod_{j=1}^5 x_j^{M_{ij}} + \sum_{s=1}^h \varphi_s e_s(x), \quad (1.10)$$

where e_s are also quasi-homogeneous and form the basis in the space of deformations of complex structure. The φ_s are moduli of complex structure, and h is the Hodge number of the family X_M .

In this paper we present a simple verification of the equivalence between Berglund – Hübsch – Krawitz and Batyrev constructions for mirror symmetry. Berglund and Hübsch proposed [4] that the mirror partner for the hypersurface X_M is related to the hypersurface X_{M^T} cut out by

$$W_{M^T}(z) = \sum_{i=1}^5 \prod_{j=1}^5 x_j^{(M^T)_{ij}} \quad (1.11)$$

in another weighted projective space $\mathbb{P}_{(\bar{k}_1, \bar{k}_2, \bar{k}_3, \bar{k}_4, \bar{k}_5)}^4$. They suggested that the mirror of X_M

is realized as a quotient of X_{M^T} by some subgroup of the phase symmetries of the W_{M^T} . This approach has been generalized by Krawitz [5]. The construction starts with the polynomial W_M and the hypersurface X_M . Let $SL(M)$ be the group of phase symmetries preserving $H^{3,0}(X_M)$, and $J_M \subseteq SL(M)$ – its subgroup that consists of the phase symmetries induced by $\mathbb{C}^*$ action on $\mathbb{P}^4_{(k_1, k_2, k_3, k_4, k_5)}$. Choose a group G_0 to be some subgroup of $SL(M)$ containing J_M . It was shown that the quotient space $Z(M, G) := X_M/G$ is a Calabi-Yau orbifold, where $G = G_0/J_M$. A similar procedure can be applied to the transposed polynomial W_{M^T} to obtain Calabi-Yau orbifold $Z(M^T, G^T) := X_{M^T}/G^T$. Krawitz has shown how to choose the dual group G^T such that $Z(M, G)$ and $Z(M^T, G^T)$ form a mirror pair on the level of cohomologies [5], see also [6–8]. We describe this construction in more detail in the next section.

On the other hand, following Batyrev [9], we can build a mirror of the original orbifold as follows. The superpotential W_M after adding to it admissible, that is invariant with respect to the group G , quasi-homogeneous monomials, that correspond to deformation of the conformal structure, defines a set of vectors $\vec{V}_a \in \mathbb{Z}^5$. These vectors are the exponents of the monomials in W_M and its admissible deformations. The $\vec{V}_a$, after subtracting the vector $\vec{v}_0$ with components $(1, 1, 1, 1, 1)$ from each of them, begin to belong to the four-dimensional sublattice, determined by its orthogonality to the vector $\vec{k} := (k_1, k_2, k_3, k_4, k_5)$. Moreover, more precisely, the vectors $\vec{V}'_a := \vec{V}_a - \vec{v}_0$ lie in a sublattice of the L , since they satisfy additional constraints, namely, they correspond only to the G -invariant monomials. The lattice points $\vec{V}'_a$ consist the Batyrev reflexive polyhedron $\Delta(M, G)$ for the orbifold $Z(M, G)$. On the other hand, the vectors $\vec{V}'_a$ correspond to the edges of the *fan* for the dual reflexive polyhedron. Having the data of the *fan*, we build the mirror for the $Z(M, G)$, let us call it Y , in two steps. First, we construct the toric variety T [9, 10]. On the second step, we find the homogeneous polynomial W^Y whose critical locus cut out a hypersurface which is nothing but the mirror manifold Y . After that, by reducing the toric variety T to a weighted projective space $\mathbb{P}^4_{(\bar{k}_1, \bar{k}_2, \bar{k}_3, \bar{k}_4, \bar{k}_5)}$ which appears in the BHK construction, we demonstrate that the results of the two constructions coincide. The ”strong parallel between the Berglund-Hübsch-Krawitz and Batyrev mirror symmetry”, based on Landau-Ginzburg/Calabi-Yau correspondence, was stated by Clader and Ruan in [8]. In this paper we confirm exactly the equivalence of these two constructions by the explicit computation for the CY orbifolds connected with the invertible singularities.

In Section 2, we fix notations and give a short review of the BHK mirror construction. In Section 3, we formulate a version of the Batyrev approach for constructing mirror orbifolds and show its equivalence to the BHK construction.

2 Berglund–Hübsch–Krawitz mirror construction

Let Calabi-Yau hypersurface X_M be defined in $\mathbb{P}_{(k_1, k_2, k_3, k_4, k_5)}^4$ by zero locus of

$$W_M(x) = \sum_{i=1}^5 \prod_{j=1}^5 x_j^{M_{ij}}. \quad (2.1)$$

Taking into account the quasi-homogeneity of W_M we obtain that it is invariant under the action of the group J_M generated by the following action

$$x_i \mapsto \omega^{k_i} x_i, \quad \omega^d = 1. \quad (2.2)$$

Moreover, the polynomial W_M has a larger group of diagonal automorphisms

$$\text{Aut}(M) := \{(\lambda_1, \dots, \lambda_5) \in (\mathbb{C}^*)^5 \mid W_M(\lambda_1 x_1, \dots, \lambda_5 x_5) = W_M(x_1, \dots, x_5), \forall x_i\}, \quad (2.3)$$

of order $|\text{Aut}(M)| = d$ and $J_M \subseteq \text{Aut}(M)$. The group $\text{Aut}(M)$ is generated by $q_i(M)$ which act on coordinates as

$$q_i(M) : x_j \mapsto e^{2\pi i B_{ji}} x_j, \quad (2.4)$$

where the matrix $B = M^{-1}$. Indeed, the generators of $\text{Aut}(M)$ act on each term in (1.4) as

$$q_l(M) \cdot \prod_{j=1}^5 x_j^{M_{ij}} = e^{2\pi i B_{ij} M_{jl}} \prod_{l=1}^5 x_l^{M_{il}} = e^{2\pi i \delta_{il}} \prod_{l=1}^5 x_l^{M_{il}} = \prod_{l=1}^5 x_l^{M_{il}}. \quad (2.5)$$

In these terms, the generator of the group J_M is

$$\prod_{i=1}^5 q_i(M). \quad (2.6)$$

Calabi-Yau threefold X_M admits the existence of holomorphic, nowhere vanishing 3-form Ω . Subgroups preserving this form Ω , or, equivalently, preserving the product $\prod_i x_i$ are called allowable. Let $SL(M)$ be the maximal allowable group with generators $p_s(M)$,

$$SL(M) := \left\{ p_s(M) \in \text{Aut}(M) \mid p_s(M) \cdot \prod_{j=1}^5 x_j = \prod_{j=1}^5 x_j \right\}. \quad (2.7)$$

The obvious fact is that $J_M \subseteq SL(M)$. Consider an allowable subgroup G_0 such that $J_M \subseteq G_0 \subseteq SL(M)$. Define the quotient group

$$G := G_0 / J_M. \quad (2.8)$$

Then we obtain Calabi-Yau orbifold X as

$$X := Z(M, G) = X_M/G. \quad (2.9)$$

For the first time, such orbifolds and their mirrors were considered for the case of Calabi-Yau threefold in [11]. In the general case, the Berglund–Hübsch–Krawitz construction starts from the quasi-homogeneous, invertible polynomial W_M and the group G_0

$$J_M \subseteq G_0 \subseteq SL(M) \subseteq \text{Aut}(M). \quad (2.10)$$

The full family of the orbifold X cut out by the equation $\{W^X = 0\} \subseteq \mathbb{P}_{(k_1, k_2, k_3, k_4, k_5)}^4/G$. Here the polynomial is

$$W^X(x, \varphi) = \sum_{i=1}^5 \prod_{j=1}^5 x_j^{M_{ij}} + \sum_{l=1}^{h_X} \varphi_l \prod_{j=1}^5 x_j^{S_{lj}}, \quad (2.11)$$

where φ_l are moduli of complex structure deformations, and $h_X := h_{21}(X)$ is Hodge number. The monomials

$$e_l := \prod_{j=1}^5 x_j^{S_{lj}} \quad (2.12)$$

in (2.11) are quasi-homogeneous since $\sum_j S_{lj} k_j = d$ and invariant under the group G . They belong to G -invariant subring of the Milnor ring $\mathbb{C}[x_1, \dots, x_5]/\langle \frac{\partial W_M}{\partial x_j} \rangle$ [12]. We denote the monomial with $S_{h_X, i} = 1$, which plays a distinguished role, e_{h_X} ,

$$e_{h_X} = \prod_{i=1}^5 x_i. \quad (2.13)$$

Actually, the monomials e_l in (2.11) are the subset in the basis of deformation of complex structure e_s , $s = 1, \dots, h$ (see (1.10)) of the original CY family X_M . If we denote by ρ_s the generator of the group G_0 , then we will see that

$$\rho_s \cdot \prod_{j=1}^5 x_j^{S_{lj}} = \prod_{j=1}^5 x_j^{S_{lj}}, \quad l = 1, \dots, h_X. \quad (2.14)$$

We can define a similar (2.10) set of groups for the transposed matrix: $J_{M^T} \subseteq G_0^T \subseteq SL(M^T) \subseteq \text{Aut}(M^T)$. Recall, that the polynomial

$$W_{M^T}(z) = \sum_{i=1}^5 \prod_{j=1}^5 z_j^{M_{ji}} \quad (2.15)$$

of the degree $\bar{d} = \sum_j \bar{k}_j M_{ji}$ defines the Calabi-Yau hypersurface X_{M^T} in another projective

space $\mathbb{P}^4_{(\bar{k}_1, \bar{k}_2, \bar{k}_3, \bar{k}_4, \bar{k}_5)}$. The weights $\bar{k}_i$ satisfy the Calabi–Yau condition $\sum_i \bar{k}_i = \bar{d}$.

Taking the quotient $G^T := G_0^T / J_{M^T}$, we define the Calabi-Yau orbifold as

$$Z(M^T, G^T) := X_{M^T} / G^T. \quad (2.16)$$

The fact is that groups G and G^T can be chosen in different ways. The question arises: is it possible to choose a group G^T for a given group G , and if so, how to do this so that the manifolds $Z(M, G)$ and $Z(M^T, G^T)$ form a mirror pair?

The answer was given by Krawitz [5]. His construction allows one to define the generators of the group G_0^T . Namely, they are constructed using the exponents S_{li} of the invariant monomials as (2.12) as follows

$$\rho_l^T := \prod_{i=1}^5 q_i(M^T)^{S_{li}}, \quad (2.17)$$

where $q_i(M^T)$ are generators of the $\text{Aut}(M^T)$ acting on each coordinate z_j in $\mathbb{P}^4_{(\bar{k}_1, \bar{k}_2, \bar{k}_3, \bar{k}_4, \bar{k}_5)}$ as

$$q_i(M^T) : z_j \mapsto e^{2\pi i B_{ij}} z_j. \quad (2.18)$$

It follows that the group G_0^T acts on the coordinates z_j as

$$\rho_l^T : z_j \mapsto e^{2\pi i \sum_i S_{li} B_{ij}} z_j. \quad (2.19)$$

The full family of the mirror Calabi-Yau orbifold $Z(M^T, G^T)$ is given by zero locus of

$$W^{Z(M^T, G^T)}(z, \psi) = \sum_{i=1}^5 \prod_{j=1}^5 z_j^{M_{ji}} + \sum_{m=1}^{h_Y} \psi_m \prod_{j=1}^5 z_j^{R_{mj}}, \quad (2.20)$$

where ψ_m are moduli of complex structure of the family $Z(M^T, G^T)$. The monomials $\bar{e}_m := \prod_{j=1}^5 z_j^{R_{mj}}$ are invariant under the G_0^T action (2.19):

$$\rho_l^T \cdot \bar{e}_m = e^{2\pi i \sum_{ij} B_{ij} S_{li} R_{mj}} \prod_{j=1}^5 z_j^{R_{mj}} = \prod_{j=1}^5 z_j^{R_{mj}}. \quad (2.21)$$

The exponents S_{li} , and R_{mj} can be interpreted as a five-component integer vectors $(\vec{S}_l)_j = S_{lj}$, and $(\vec{R}_m)_i = R_{mi}$. The invariance condition (2.21) can be rewritten in terms of the pairing of these vectors, defined with matrix B as

$$(\vec{S}_l, \vec{R}_m) = \sum_{i,j=1}^5 B_{ij} S_{li} R_{mj} \in \mathbb{Z}. \quad (2.22)$$

This relation is a strong restriction because although the matrices S and R are integers,

at the same time the entries in B are rational. Taking into account also the condition of quasi-homogeneity

$$\sum_{i=1}^5 R_{mi} \bar{k}_i = \bar{d}, \quad (2.23)$$

we conclude that the equations (2.22) have finite non-negative number of solutions. Denote this number as h_Y .

The Chiodo-Ruan theorem [13] states that orbifolds $Z(M, G)$ and $Z(M^T, G^T)$ form a mirror pair on the level of cohomology

$$H^{p,q}(Z(M, G), \mathbb{C}) = H^{3-p,q}(Z(M^T, G^T), \mathbb{C}). \quad (2.24)$$

Thus the Berglund–Hübsch–Krawitz construction allows one to determine the polynomial $W^{Z(M^T, G^T)}$ which defines the full family of $Z(M^T, G^T)$.

3 Batyrev construction and verification of equivalence

In this section, following the Batyrev’s approach [9], briefly described in the section 1, we will build the mirror Y of the orbifold $Z(M, G)$ as a hypersurface in toric variety. We find the mirror polynomial W^Y as a function of toric coordinates. We also show that the mirror hypersurface Y is equivalent to the Calabi-Yau orbifold $Z(M^T, G^T)$ (2.20) in a weighted projective space obtained above using the BHK construction. We do this in two steps.

The first step is to construct the toric variety. To do this, we begin with the invertible quasi-homogeneous polynomial W_M (2.11) with the deformations e_l . Rewrite it in a form

$$W^X(x, \varphi) = \sum_{i=1}^5 \prod_{j=1}^5 x_j^{M_{ij}} + \sum_{l=1}^{h_X} \varphi_l \prod_{j=1}^5 x_j^{S_{lj}} = \sum_{a=1}^{h_X+5} C_a \prod_{j=1}^5 x_j^{V_{aj}}. \quad (3.1)$$

The exponents V_{aj} are components of integer vectors, namely $V_{aj} = (\vec{V}_a)_j$. They are equal:

$$V_{aj} = \begin{cases} M_{aj}, & 1 \leq a \leq 5 \\ S_{a-5,j}, & 6 \leq a \leq h \end{cases} \quad (3.2)$$

and $\vec{V}_{h_X+5}$ has components $(1, 1, 1, 1, 1)$. From the condition

$$\sum_{i=1}^5 V_{ai} k_i = d \quad (3.3)$$

it follows, that the vectors $\vec{V}'_a := \vec{V}_a - \vec{V}_{h_X+5}$ lie in the four-dimensional lattice $L \subset \mathbb{Z}^5$. More precisely, they belong to the sublattice of L defined by the group G .

These vectors $\vec{V}'_a$ correspond to the lattice points of the reflexive polytope $\Delta(M, G)$. They also correspond to the edges of the fan of the reflexive polyhedron dual to the original one [9, 10]. The fan defines the toric variety T . Let us construct this toric variety. The vectors $\vec{V}'_a$, $a = 1, \dots, h_X + 4$, being four-dimensional, satisfy to h_X linear relations

$$\sum_{a=1}^{h_X+4} Q_{la} \vec{V}'_a = 0, \quad l = 1, \dots, h_X, \quad (3.4)$$

where the coefficients Q_{la} are integers. The solution of the equations (3.4) is [14]

$$Q_{la} = \begin{cases} S_{lj} B_{ja}, & 1 \leq a \leq 5 \\ -\delta_{l, a-5}, & a > 5. \end{cases} \quad (3.5)$$

The numbers Q_{la} are the weights of this toric variety T . Knowing them allows one to define T as [10]

$$T = \frac{\mathbb{C}^{h_X+4} - Z}{(\mathbb{C}^*)^{h_X}}, \quad (3.6)$$

where Z is the $(\mathbb{C}^*)^{h_X}$ -invariant subset [10]. The abelian group $(\mathbb{C}^*)^{h_X}$ acts on the coordinates in T as

$$y_a \mapsto \lambda^{Q_{la}} y_a, \quad l = 1, \dots, h_X; \quad a = 1, \dots, h_X + 4. \quad (3.7)$$

The mirror Calabi-Yau manifold Y is realized as a hypersurface in the toric variety T , cut out by an invariant under the action of the abelian group $(\mathbb{C}^*)^{h_X}$ homogeneous polynomial $W^Y(y)$ [14, 15]

$$W^Y(\lambda^{Q_{l1}} y_1, \dots, \lambda^{Q_{lh_X+5}} y_{h_X+4}) = W^Y(y_1, \dots, y_{h_X+4}), \quad l = 1, \dots, h_X - 1, \quad (3.8)$$

$$W^Y(\lambda^{Q_{h_X,1}} y_1, \dots, \lambda^{Q_{h_X, h_X+5}} y_{h_X+4}) = \lambda W^Y(y_1, \dots, y_{h_X+4}), \quad (3.9)$$

We are looking for the W^Y as a sum quasi-homogeneous monomials

$$W^Y(y) = \sum_{b=1}^{h_Y+5} \tilde{C}_b \prod_{a=1}^{h_X+4} y_a^{N_{ba}}. \quad (3.10)$$

The exponents N_{ba} are non-negative integer numbers. We know 5 invariant monomials of the form

$$P_i = \prod_{a=1}^{h_X+4} y_a^{V_{ai}} = \prod_{j=1}^5 y_j^{M_{ji}} \times \prod_{l=1}^{h_X-1} y_{l+5}^{S_{li}} = \prod_{j=1}^5 y_j^{(M^T)_{ij}} \times \prod_{l=1}^{h_X-1} y_{l+5}^{S_{li}}, \quad (3.11)$$

where $i = 1, \dots, 5$. We can check that monomials P_i are invariant under the $(\mathbb{C}^*)^{h_X}$ action,

because $\sum_a Q_{la} V_{ai} = 0$. Also, we have one more invariant monomial of the form

$$\bar{e}_{h_Y} = \prod_{a=1}^{h_X+4} y_a. \quad (3.12)$$

Other invariant monomials we seek in the form

$$\bar{e}_m = \prod_{a=1}^{h_X+4} y_a^{N_{ma}} = \prod_{j=1}^5 y_j^{N_{mj}} \times \prod_{l=1}^{h_X-1} y_{l+5}^{N_{m,l+5}}. \quad (3.13)$$

From the invariance of the $\bar{e}_m$ under $(\mathbb{C}^*)^{h_X}$ action one can obtain that the non-negative integers N_{mi} need to satisfy the relations

$$B_{ji} S_{lj} N_{mi} - N_{m,l+5} = \delta_{l,h_X}. \quad (3.14)$$

Since the numbers in the r.h.s. $N_{m,l+5}$ are assumed to be integers, we get that the integer numbers N_{mi} satisfy the relation

$$B_{ji} S_{lj} N_{mi} \in \mathbb{Z}, \quad (3.15)$$

which exactly coincides with the relation for the exponents R_{mi} in (2.22).

Using the fact that $S_{h_X,j} = 1$ and the relation (1.9) for the matrix B we get from (3.14) for $l = h_X$

$$N_{mj} \bar{k}_j = \bar{d}. \quad (3.16)$$

Therefore, as it follows from the (3.15) and (3.16) that these equations have the same solutions as the equations (2.22) in the BHK approach, so that

$$N_{mi} = R_{mi}. \quad (3.17)$$

Now we can find the remaining exponents. Using the (3.14) we obtain that

$$N_{m,l+5} = S_{lj} B_{ji} R_{mi}. \quad (3.18)$$

Thus, we finally obtain that the polynomial W^Y has the form

$$W^Y(y_1, \dots, y_{h_X+4}) = \sum_{i=1}^5 \prod_{j=1}^5 y_j^{M_{ji}} \prod_{l=1}^{h_X-1} y_{l+5}^{S_{li}} + \sum_{m=1}^{h_Y} \psi_m \prod_{j=1}^5 y_j^{R_{mj}} \prod_{l=1}^{h_X-1} y_{l+5}^{(S_{lj} B_{ji} R_{mi})}. \quad (3.19)$$

Then, the mirror Calabi-Yau manifold Y is given by zero locus of W^Y (3.19) in T , similar as it was in [16]. To see the correspondence of the just described construction to

the previous BHK construction, we introduce the new coordinates

$$(y_1, \dots, y_{h_X+4}) \mapsto (z_1, z_2, z_3, z_4, z_5), \quad (3.20)$$

where

$$z_j = y_j \prod_{l=1}^{h_X-1} y_{l+5}^{S_{li}B_{ij}}, \quad 1 \leq z_j \leq 5. \quad (3.21)$$

The change of the coordinates (3.20) is mapping of the toric variety T into the quotient of the weighted projective space $\mathbb{P}_{(\bar{k}_1, \bar{k}_2, \bar{k}_3, \bar{k}_4, \bar{k}_5)}^4 / G^T$. Indeed, the abelian group $(\mathbb{C}^*)^{h_X}$ acts on z_j as

$$z_j \mapsto \lambda^{Q_{sj}+Q_{s,l+5}S_{li}B_{ij}} z_j = \lambda^{S_{sm}B_{mj}-\delta_{sl}S_{li}B_{ij}} z_j = z_j, \quad s = 1, \dots, h_X - 1. \quad (3.22)$$

And, in view of the fact that $S_{h_X j} = 1$ and the property of the matrix B (1.9),

$$z_j \mapsto \lambda^{S_{h_X m}B_{mj}} z_j = \lambda^{\sum_m B_{mj}} z_j = \lambda^{\nu_j} z_j, \quad (3.23)$$

where $\nu_j := \bar{k}_j / \bar{d}$.

It is easy to check, that the polynomial W^Y can be rewritten in these coordinates as

$$W^Y(z) = \sum_{i=1}^5 \prod_{j=1}^5 z_j^{(M^T)_{ij}} + \sum_{m=1}^{h_Y} \psi_m \prod_{j=1}^5 z_j^{R_{mj}} \quad (3.24)$$

and coincides with the result of Berglund–Hübsch–Krawitz construction (2.20).

Thus, in this paper, we have provided a simple evidence of equivalence of two mirror partners obtained by Berglund–Hübsch–Krawitz, and Batyrev constructions. Namely started with an invertible polynomial W_M , suitable subgroup G and the weighted projective space $\mathbb{P}_{(k_1, k_2, k_3, k_4, k_5)}^4$ we have obtained the mirror in another projective space $\mathbb{P}_{(\bar{k}_1, \bar{k}_2, \bar{k}_3, \bar{k}_4, \bar{k}_5)}^4$. Following to Batyrev approach, we have constructed the toric variety and defined the mirror polynomial W_M . Finally, we showed the connection between the toric variety and the weighted projective space. The results of these two mirror constructions coincide. Earlier, we checked a similar statement for the particular case of quintic orbifolds in [17].

Acknowledgments

We are grateful to V. Belavin, M. Bershtein and G. Koshevoy for useful discussions. This work was performed in Landau Institute for Theoretical Physics. The work of A. Belavin was done within the framework of the state assignment. The work of B. Eremin has been supported by the Russian Science Foundation under the grant 18-12-00439.

References

- [1] P. Candelas and X. de la Ossa, “Moduli Space of Calabi-Yau Manifolds,” Nucl. Phys. B **355**, 455-481 (1991).
- [2] A. Strominger, “Special Geometry,” Commun. Math. Phys. **133**, 163-180 (1990).
- [3] M. Kreuzer and H. Skarke, “On the classification of quasi-homogeneous functions,” Commun. Math. Phys. **150**, 137 (1992), [arXiv:hep-th/9202039](#).
- [4] P. Berglund and T. Hubsch, “A Generalized construction of mirror manifolds,” AMS/IP Stud. Adv. Math. **9**, 327-346 (1998), [arXiv:hep-th/9201014](#).
- [5] M. Krawitz, “FJRW rings and Landau-Ginzburg Mirror Symmetry,” (2009), [arXiv:0906.0796 \[math.AG\]](#).
- [6] T. L. Kelly, “Berglund–Hübsch–Krawitz mirrors via Shioda maps,” Adv. Theor. Math. Phys. **17**, no.6, 1425-1449 (2013), [arXiv:1304.3417 \[math.AG\]](#).
- [7] M. Shoemaker, “Birationality of Berglund-Huebsch-Krawitz Mirrors,” Commun. Math. Phys. **331**, no.2, 417-429 (2014), [arXiv:1209.5016 \[math.AG\]](#).
- [8] E. Clader and Y. Ruan, “Mirror Symmetry Constructions,” [arXiv:1412.1268 \[math.AG\]](#).
- [9] V. V. Batyrev, “Dual polyhedra and mirror symmetry for Calabi-Yau hypersurfaces in toric varieties,” J. Alg. Geom. **3**, 493-545 (1994), [arXiv:alg-geom/9310003](#).
- [10] K. Hori, S. Katz, A. Klemm, R. Pandharipande, R. Thomas, C. Vafa, R. Vakil and E. Zaslow, “Mirror symmetry,” pages 101-142, AMS, Clay Mathematical Institute.
- [11] B. R. Greene and M. R. Plesser, “Duality in Calabi-Yau Moduli Space,” Nucl. Phys. B **338**, 15-37 (1990).
- [12] W. Lerche, C. Vafa and N. P. Warner, “Chiral Rings in N=2 Superconformal Theories,” Nucl. Phys. B **324**, 427-474 (1989).
- [13] A. Chiodo and Y. Ruan “LG/CY correspondence: the state space isomorphism,” (2009), [arXiv:0908.0908 \[math.AG\]](#)
- [14] K. Aleshkin, A. Belavin and A. Litvinov, “JKLMR conjecture and Batyrev construction,” J. of Stat. Mech.: Theory and Experiment, **2019**, March (2019) [arXiv:1812.00478 \[hep-th\]](#).
- [15] A. Belavin and B. Eremin, “Partition Functions of $\mathcal{N} = (2, 2)$ Supersymmetric Sigma Models and Special Geometry on the Moduli Spaces of Calabi-Yau Manifolds,” Theor. Math. Phys. **201**, no.2, 1606-1613 (2019), [arXiv:1907.11102 \[hep-th\]](#).
- [16] E. Witten, “Phases of N=2 theories in two-dimensions,” AMS/IP Stud. Adv. Math. **1**, 143-211 (1996), [arXiv:hep-th/9301042](#).
- [17] A. Belavin and B. Eremin, “Mirror pairs of Quintic orbifolds,” JETP Letters. **112**, no.6, 388-393 (2020).