

Consistency of archetypal analysis*

Braxton Osting[†], Dong Wang[‡], Yiming Xu[§], and Dominique Zosso[¶]

Abstract. Archetypal analysis is an unsupervised learning method that uses a convex polytope to summarize multivariate data. For fixed k , the method finds a convex polytope with k vertices, called *archetype points*, such that the polytope is contained in the convex hull of the data and the mean squared distance between the data and the polytope is minimal. In this paper, we prove a consistency result that shows if the data is independently sampled from a probability measure with bounded support, then the archetype points converge to a solution of the continuum version of the problem, of which we identify and establish several properties. We also obtain the convergence rate of the optimal objective values under appropriate assumptions on the distribution. If the data is independently sampled from a distribution with unbounded support, we also prove a consistency result for a modified method that penalizes the dispersion of the archetype points. Our analysis is supported by detailed computational experiments of the archetype points for data sampled from the uniform distribution in a disk, the normal distribution, an annular distribution, and a Gaussian mixture model.

Key words. Archetypal analysis; principal convex hull; consistency; multivariate data summarization; unsupervised learning

AMS subject classifications. 62H12, 62H30, 68T10, 65D18

1. Introduction. Fix $k \in \mathbb{N}$ and write $[k] = \{1, 2, \dots, k\}$. For given data $X_N = \{x_i\}_{i \in [N]} \subset \mathbb{R}^d$, the *archetypal analysis* problem is to find a cardinality k pointset $A = \{a_\ell\}_{\ell \in [k]} \subset \mathbb{R}^d$ that solves

$$(1.1a) \quad \min_{A \subset \mathbb{R}^d} F(A)$$

$$(1.1b) \quad A \subset \text{co}(X_N),$$

where

$$F(A) = \left(\frac{1}{N} \sum_{i=1}^N d^2(x_i, \text{co}(A)) \right)^{1/2} = \left(\int_{\mathbb{R}^d} d^2(x, \text{co}(A)) d\mu_N(x) \right)^{1/2}.$$

Here, $\text{co}(\cdot)$ denotes the convex hull, $d^2(\cdot, \cdot)$ denotes the squared Euclidean distance, and $\mu_N(x) = \frac{1}{N} \sum_{i \in [N]} \delta_{x_i}(x)$ is the empirical measure associated with the data X_N . The constraint in (1.1) imposes that the convex hull of the dataset should contain the pointset A , and hence $\text{co}(A)$. The objective function, F , in (1.1) is the root mean squared distance from the

*Braxton Osting acknowledges partial support from NSF DMS 16-19755 and 17-52202. Dong Wang acknowledges support by the University Development Fund from The Chinese University of Hong Kong, Shenzhen (UDF01001803). Dominique Zosso acknowledges support by a Simons collaboration grant for mathematicians (#586942).

[†]Department of Mathematics, University of Utah, Salt Lake City, UT (osting@math.utah.edu).

[‡]School of Science and Engineering, The Chinese University of Hong Kong, Shenzhen, Guangdong 518172, China (wangdong@cuhk.edu.cn).

[§]Department of Mathematics, University of Utah, Salt Lake City, UT (yxu@math.utah.edu).

[¶]Department of Mathematical Sciences, Montana State University, Bozeman, MT (dominique.zosso@montana.edu).

data to the convex hull of A . By pointset, we mean the unordered collection of k points in $\mathbb{R}^d$, which we denote by $\{\mathbb{R}^d\}^k$. One may think of $\{\mathbb{R}^d\}^k$ as the product space $(\mathbb{R}^d)^k$ modulo permutation of its components, *i.e.*,

$$\{a_\ell\}_{\ell \in [k]} = \left\{ (b_\ell)_{\ell \in [k]} \in (\mathbb{R}^d)^k : a_\ell = b_{\sigma(\ell)} \text{ for some } \sigma \in S[k] \right\},$$

where $S[k]$ denotes the set of permutations on $[k]$. We refer to a minimizing pointset $A = \{a_\ell\}_{\ell \in [k]} \in \{\mathbb{R}^d\}^k$ solving (1.1) as an *archetype pointset*, the points $a_i \in A$ as *archetype points*, and the convex hull, $\text{co}(A)$ as the *archetype polytope*.

Archetypal analysis was introduced in 1994 by A. Cutler and L. Breiman as an unsupervised method to summarize multivariate data [9]. They proved the following results: (i) If $k = 1$, then the archetype point is the mean of the data, X_N . (ii) For $1 < k < N$, there exists an archetype pointset, $A = \{a_\ell\}_{\ell \in [k]}$, *i.e.*, there exists a solution to (1.1) and furthermore, there exists an archetype pointset on the boundary of $\text{co}(X_N)$. (iii) Finally for $k \geq N$, the archetype pointset is given by $A = X_N$, which attains the value $F(A) = 0$. They demonstrated that archetypal analysis can be reformulated as a nonlinear least squares problem and efficiently solved using an alternating minimization algorithm. In their concluding remarks, they note that “Because the archetypes are located on the boundary of the convex hull of the data, the procedure can be sensitive to outliers. Robust versions could be developed using convex hull peeling or the outlyingness idea of Donoho and Gasko (1992)”. Subsequently, there has been a lot of work to extend archetypal analysis to improve robustness and sensitivity to outliers [8, 34, 22] and to develop improved computational methods for computing archetypes [1]. M. Mørup and L. K. Hansen carefully compared archetypal analysis to other matrix factorization and clustering methods such as SVD/PCA, NMF, soft k -means, k -means, and k -medoids, and showed how archetypal analysis can be effectively used as an unsupervised machine learning tool for a variety of data analysis problems [23]. Interestingly, nonlinear versions of archetypal analysis based on neural networks have recently been proposed [31, 20, 19]. Finally, archetypal analysis has been applied to a variety of real-world applications, including finding archetype soccer players based on performance data [27], query-focused multi-document summarization [6], classification of galaxy spectra [7], spatio-temporal dynamics [29], *etc.*. We comment that archetypal analysis is also sometimes referred to as *principal convex hull analysis* as it approximates the “best” convex hull, although we do not use this language here.

In this work, we consider the following consistency problem. Suppose that $x_1, x_2, \dots$ are independently sampled from the probability measure μ and denote the first N points by $X_N = \{x_i\}_{i \in [N]}$. Let A_N denote the optimal solution to (1.1) for each N . *Is there a set A (depending on μ), such that $A_N \rightarrow A$ as $N \rightarrow \infty$ in some sense?*

Consistency results are fundamental in statistics and have important consequences in applications. Namely, an estimate (*e.g.*, archetype polytope) obtained using a consistent method will asymptotically stabilize, and so the collection of more data will yield diminishing returns. Generally, a consistency result for a problem posed on sampled data requires two ingredients: (i) a model from which the samples are drawn from; the original problem is viewed as a finite sample size problem for this model and (ii) a notion of convergence for the estimated quantity. We give a consistency result for the archetypal analysis problem where the data is

sampled from a measure with compact support and a consistency result for a modification of the archetypal analysis problem where the data is sampled from a measure with non-compact support. In both cases, we will use a Euclidean notion of distance; see [Definition 1.4](#).

1.1. Consistency for measures with compact support. In the following, let ν be a general probability measure on $\mathbb{R}^d$ and μ be the sampling measure for data. For convenience, we assume that μ has a bounded density. Note that (1.1) is a special case of the following minimization problem when we take $\nu = \mu_N$:

$$(1.2a) \quad \min_{A \in \{\mathbb{R}^d\}^k} F_\nu(A)$$

$$(1.2b) \quad \text{s.t. } A \subset \text{co}(\text{supp}(\nu)).$$

where $F_\nu(A) = \left(\int_{\mathbb{R}^d} d^2(x, \text{co}(A)) d\nu(x) \right)^{1/2}$. Since $\mu_N \rightarrow \mu$ as $N \rightarrow \infty$, a natural candidate limiting problem for (1.1) is thus given by

$$(1.3a) \quad \min_{A \in \{\mathbb{R}^d\}^k} F_\mu(A)$$

$$(1.3b) \quad \text{s.t. } A \subset \text{co}(\text{supp}(\mu)).$$

In [section 2](#), we prove the following elementary existence result.

Theorem 1.1. *Suppose that ν is a probability measure on $\mathbb{R}^d$ with compact support. Then the minimization problem (1.2) admits at least one minimizer with optimal value $F_\nu^* \leq \left(\int_{\mathbb{R}^d} \|x - \bar{x}\|_2^2 d\nu(x) \right)^{1/2}$, where $\bar{x} = \int_{\mathbb{R}^d} x d\nu(x)$ is the mean.*

Taking $\nu = \mu_N$ and μ in [Theorem 1.1](#) deduces the existence of minimizers for both problems (1.1) and (1.3). However, it should be noted that solutions to (1.3) are not unique in general; non-unique examples can easily be constructed when the support of μ has symmetries.

As proven in [\[9\]](#) for the discrete problem, the following theorem states that if ν has bounded support, then there exists a minimizer for problem (1.2) on the boundary of the convex hull of the support of ν .

Theorem 1.2. *Let $k \geq 2$. Suppose that ν is supported on a bounded set in $\mathbb{R}^d$. Then there exists a minimizing pointset $\{a_\ell\}_{\ell \in [k]}$ of problem (1.2) such that $\{a_\ell\}_{\ell \in [k]} \subset \partial(\text{co}(\text{supp}(\nu)))$.*

As the proof of [Theorem 1.2](#) is similar to that in [\[9\]](#), we omit the proof.

Remark 1.3. *We remark that [Theorem 1.2](#) does not imply that every minimizing pointset of problem (1.2) is on the boundary of $\text{co}(\text{supp}(\nu))$. For example, consider the case when $k = 3$ and ν is the empirical measure of the magenta data points X_N , as illustrated in [Figure 1](#). It is clear from our construction that all red and blue triangles are archetype triangles (optimal solutions). The topmost vertex of the blue triangle is in the interior of the $\text{co}(X_N)$. Note that due to the two clusters of magenta data points, the gray triangle is not optimal.*

We first consider the consistency problem under the assumption that μ has compact support. To describe the convergence of archetype points, we will use the $d_{2,\infty}$ metric on $\{\mathbb{R}^d\}^k$ which is defined as follows:




Figure 1. A illustration of an example where the archetype pointset is attained on the interior of $\text{co}(X_N)$.

Definition 1.4. For fixed integers d and k , the metric $d_{2,\infty}(\cdot, \cdot)$ on $\{\mathbb{R}^d\}^k$ is defined by

$$d_{2,\infty}(\{a_\ell\}_{\ell \in [k]}, \{b_\ell\}_{\ell \in [k]}) := \min_{\sigma \in S[k]} \max_{\ell \in [k]} \|a_{\sigma(\ell)} - b_\ell\|_2, \quad \{a_\ell\}_{\ell \in [k]}, \{b_\ell\}_{\ell \in [k]} \in \{\mathbb{R}^d\}^k,$$

where $S[k]$ is the set of permutations on $[k]$.

Properties of $d_{2,\infty}$ will be given in [section 2](#). The following Theorem, which is proven in [section 2](#), gives a consistency result under the assumption that μ has compact support.

Theorem 1.5. Fix $k \in \mathbb{N}$. Let μ be a probability measure on $\mathbb{R}^d$ with compact support. Let $x_1, x_2, \dots$ be iid samples from μ and denote by $X_N = \{x_i\}_{i \in [N]}$ the first N samples. Let A_N be an archetype pointset solving (1.1) for each N . Then, A_N has a convergent subsequence A_{N_m} (in the $d_{2,\infty}$ sense) whose limit lies in the solution set of (1.3) μ -a.s.. If (1.3) has a unique solution $A_\star$, then A_N converge to $A_\star$ in $d_{2,\infty}$ μ -a.s..

Although the consistency result stated in [Theorem 1.5](#) only holds for a subsequence in general, the optimal objective values of the discrete problems are, in fact, convergent. Moreover, under appropriate assumptions on μ , we can further establish the convergence rate using a result from random geometry [[11](#), [4](#)] and Dudley's inequality [[32](#)]. Before stating the convergence rate in [Theorem 1.8](#), we recall the α -cap condition on a probability measure which is introduced in [[4](#), Assumption 1].

Definition 1.6 (α -cap condition). Let $\alpha > 0$. A probability measure ν defined on $\mathbb{R}^d$ is said to satisfy the α -cap assumption with parameters L and η if it has bounded support and

$$(1.4) \quad \min_{u \in \mathbb{S}^{d-1}} \nu \left(\left\{ y \in \text{co}(\text{supp}(\nu)) : \langle y, u \rangle \geq \max_{z \in \text{co}(\text{supp}(\nu))} \langle z, u \rangle - \varepsilon \right\} \right) \geq L\varepsilon^\alpha, \quad \forall \varepsilon \in (0, \eta].$$

Remark 1.7. The α -cap condition can be interpreted as that for any $x \in \partial(\text{co}(\text{supp}(\nu)))$, the slab between the supporting hyperplane at x and its ε -perturbation captures $\mathcal{O}(\varepsilon^\alpha)$ mass under ν . One can check that μ satisfies the α -cap assumption with $\alpha = d$ when μ is supported on a convex domain with density bounded from below. Moreover, μ satisfies the α -cap assumption with $\alpha \geq d$ whenever μ has a density, since Lebesgue measure of the slab defined in (1.4) scales as $\mathcal{O}(\varepsilon^d)$ for $\varepsilon \rightarrow 0$.

Theorem 1.8. Under the same conditions as [Theorem 1.5](#), $F_\mu(A_N) \rightarrow F_\mu(A_\star)$ as $N \rightarrow \infty$. Moreover, if μ satisfies the α -cap condition with parameters L and η , then with probability at least $1 - 4N^{-2}$,

$$(1.5) \quad F_\mu(A_N) - F_\mu(A_\star) \lesssim \left(\frac{\log N}{N} \right)^{1/\alpha} \quad N > N(\eta),$$

where $N(\eta)$ is a non-increasing function in η and the implicit constant in (1.5) depends only on μ . In particular, if μ is supported on a convex set with positive density, $F_\mu(A_N) - F_\mu(A_\star) = \mathcal{O}((\log N/N)^{1/d})$ μ -a.s..

The proof of Theorem 1.8 is given in section 2. The asymptotic rate in (1.5) mainly depends on the α in the α -cap definition, which, roughly speaking, depends on the geometry of the support and the density near the boundary of the support of μ . In Remark 2.4, we discuss the rate of convergence for densities with very large variance.

In section 3, we demonstrate the consistency of the archetype analysis problem for the example when $d = 2$ and μ is a uniform measure on the Euclidean unit disk. In this case, we prove that archetype points are the vertices of regular k -polygons inscribed in the unit disk. Furthermore, in section 5, we perform numerical experiments for the discrete problem (1.1) with $k = 3$ to demonstrate the convergence, as $N \rightarrow \infty$, of the archetype triangle to a regular triangle inscribed in a unit disk.

1.2. Consistency for measures with non-compact support. We now consider the consistency problem when the probability measure, μ , has non-compact support. Here we have that $\text{co}(X_1) \subseteq \text{co}(X_2) \subseteq \dots$ and $\text{co}(X_N)$ is a.s. unbounded as $N \rightarrow \infty$. In this case, it is clear that there can be no limiting problem for the archetype pointset A_N as $N \rightarrow \infty$; the problem, as stated, is inconsistent. Consequently, we must modify (1.1) or (1.3) to obtain a consistency result.

When the probability measure ν has non-compact support, we add a ‘penalty term’ to the energy that prevents the archetype points from tending to infinity and consider the objective function

$$(1.6) \quad F_{\nu,\alpha}(A) = (F_\nu^2(A) + \alpha V(A))^{1/2} = \left(\int_{\mathbb{R}^d} d^2(x, \text{co}(A)) \, d\nu(x) + \frac{\alpha}{k} \sum_{\ell \in [k]} \|a_\ell - \bar{a}\|_2^2 \right)^{1/2}.$$

Here, $\alpha > 0$ is a fixed parameter, $V(A) = \frac{1}{k} \sum_{\ell \in [k]} \|a_\ell - \bar{a}\|_2^2$ is the variance of the archetype points, and $\bar{a} = \frac{1}{k} \sum_{\ell \in [k]} a_\ell$ is the mean of the archetype points. We think of $F_{\nu,\alpha}(A)$ as a trade-off between the fidelity to the data and the size of the archetype polytope.

Remark 1.9. *In practice, datasets are always finite so that one can directly apply the archetypal analysis to them. Adding a regularization term can potentially make the algorithm more robust to unexpected outliers. However, the regularization parameter should be small, otherwise the solution will be close to the mean of the dataset (see Theorem 1.14), contradicting the original goal of summarizing datasets by their extreme patterns.*

Remark 1.10. *Other regularization terms like the k -means type of penalties considered in [34] could be discussed similarly under our methodology. Another penalty term one might consider would be $V(A) = \text{vol}(\text{co}(A))$. However, in this case, it is possible for $\text{co}(A)$ to be a degenerate (lower dimensional) set to avoid the penalization.*

We consider the variance-regularized archetype problem,

$$(1.7a) \quad \min_{A \in \{\mathbb{R}^d\}^k} F_{\nu, \alpha}(A)$$

$$(1.7b) \quad \text{s.t. } A \subset \text{co}(\text{supp}(\nu)).$$

For fixed N and square-integrable ν , the following result establishes the existence of minimizers of (1.7).

Theorem 1.11. *Let $\alpha > 0$ be a fixed number. Suppose that ν is square-integrable, i.e., $\int_{\mathbb{R}^d} \|x\|_2^2 d\nu(x) < \infty$. Then the minimization problem (1.7) admits at least one minimizer with optimal value $F_{\nu, \alpha}^* \leq (\int_{\mathbb{R}^d} \|x - \bar{x}\|_2^2 d\nu(x))^{1/2}$, where $\bar{x} = \int_{\mathbb{R}^d} x d\nu(x)$ is the mean.*

Taking $\nu = \mu_N$ and μ in Theorem 1.11 deduces the existence of minimizers for both the discrete and continuous archetype problems with the variance regularization.

We next state the consistency result for the variance-regularized archetype problem. Since the support of the distribution may be unbounded, a technical condition is needed on the distribution and discussed further below.

Theorem 1.12. *Fix $k \in \mathbb{N}$. Let μ be a probability measure on $\mathbb{R}^d$ which is square-integrable. Let $x_1, x_2, \dots$ be iid samples from μ and denote by $X_N = \{x_i\}_{i \in [N]}$ the first N samples. Let A_N be the archetype pointset solving (1.7) with $\nu = \mu_N$ each N . Suppose that μ -a.s., for any $r > 0$, there exists some $N(r) \in \mathbb{N}$ such that $B(r) \subset \text{co}(X_{N(r)})$. Then, A_N has a convergent subsequence A_{N_m} (in the $d_{2, \infty}$ sense) whose limit lies in the solution set of (1.3) μ -a.s.. If (1.7) has a unique solution A_* , then A_N converge to A_* in $d_{2, \infty}$ μ -a.s..*

Remark 1.13. *The technical assumption on μ states that μ -a.s., for any $r > 0$, there exists some $N(r) \in \mathbb{N}$ such that $B(r) \subset \text{co}(X_{N(r)})$. The same proof applies if all the minimizers of the continuous problem are in the interior of $\text{co}(\text{supp}(\mu))$. In particular, the theorem applies to any μ defined on $\mathbb{R}^d$ with strictly positive density.*

Note that as $\alpha \rightarrow \infty$, problem (1.7) reduces to the following problem

$$(1.8a) \quad \min_{a \in \mathbb{R}^d} \left(\int_{\mathbb{R}^d} d^2(x, a) d\nu(x) \right)^{1/2}$$

$$(1.8b) \quad \text{s.t. } a \in \text{co}(\text{supp}(\nu)),$$

which has a unique minimizer $\bar{x} = \int_{\mathbb{R}^d} x d\nu(x)$. Therefore, one should expect that solutions of (1.7), $A_{\star, \alpha}$, shrink to $A_{\star, \infty} = \{\bar{x}\}^k$ as α tends to infinity. The following theorem quantifies this observation by giving an upper bound on the convergence rate.

Theorem 1.14. *Fix $k \in \mathbb{N}$. Let ν be a probability measure on $\mathbb{R}^d$ which is square-integrable. Let $A_{\star, \alpha}$ be a solution to the problem (1.7). Let $\bar{x} = \int_{\mathbb{R}^d} x d\nu(x)$ and define $A_{\star, \infty} = \{\bar{x}\}^k$. For sufficiently large α ,*

$$d_{2, \infty}(A_{\star, \alpha}, A_{\star, \infty}) \leq 8k^{1/2} \alpha^{-1/4} \left(\int_{\mathbb{R}^d} \|x - \bar{x}\|_2^2 d\nu(x) \right)^{1/2}.$$

In section 5, we perform several numerical experiments for data sampled from the normal distribution, an annular distribution, and a Gaussian mixture distribution to study the

solution to problem (1.7). Furthermore, in the experiments, we study how the area of $\text{co}(A)$ depends on α and observe that $\text{co}(A)$ always contains $\bar{x}$ for all initializations and samples of the data.

1.3. A brief review of consistency results for other unsupervised learning methods. In Theorem 1.5 and Theorem 1.12, we state consistency results for archetypal analysis. Here, we briefly review related work proving consistency for other unsupervised methods.

A related and well-known unsupervised method for data summary is k -means clustering, which finds the k points in the space that minimize the within class variance of the dataset. In the seminal work, Pollard established a strong consistency property for the theoretic k -means solutions by assuming the data points are independently sampled from a fixed distribution [25]. An asymptotic normality result on the convergence rate was later proven in [26] using a functional central limit theorem. The rich theory behind the k -means clustering boils down to the convenient form of the objective function as well as the fact that the admissible set is data-independent, and allows generalization to other related problems. For instance, Sun *et al.* extended the consistency result in [25] to the regularized k -means clustering, where a group LASSO/adaptive LASSO term is considered to balance the trade-off between model fitting and sparsity [30]. Consistency results concerning other unsupervised clustering methods are also worth noting. For example, Hartigan provided a weak notion of consistency for Linkage algorithms, proving that the algorithm can obtain a couple of high-density regions [14]. Finally, there are many consistency and convergence results for the graph Laplacian on geometric graphs [3, 21, 2, 16, 28, 15, 5], as well as graph-based methods for data analysis based on the graph Laplacian, including spectral clustering [33, 12], Cheeger and ratio graph cuts [13], Dirichlet partitions [24], and the PageRank algorithm [35].

1.4. Outline. In section 2, we discuss the consistency problem for distributions with compact support; in particular, we prove Theorem 1.1, Theorem 1.5, and Theorem 1.8. In section 3, we consider finding the archetype points for the uniform distribution on the unit disk. In section 4, we consider distributions with unbounded support and prove Theorem 1.11, Theorem 1.12 and Theorem 1.14. In section 5, we present a variety of numerical examples that strongly support the analysis. We conclude in section 6 with a discussion.

2. Proof of Theorems 1.1, 1.5, and 1.8 for probability measures with compact support.

Before proving the main results on the consistency of archetype problems for probability measures with bounded support, we discuss a few properties of the $d_{2,\infty}$ distance, defined in Definition 1.4, which will be used throughout the convergence analysis. To see that the $d_{2,\infty}$ distance is well-defined, it suffices to check the triangle inequality holds, as the non-negativity, identity and symmetry are obvious. For any $\{a_\ell\}_{\ell \in [k]}, \{b_\ell\}_{\ell \in [k]}, \{c_\ell\}_{\ell \in [k]} \in \{\mathbb{R}^d\}^k$, there exist some $\sigma_1, \sigma_2 \in S[k]$ such that

$$\begin{aligned} d_{2,\infty}(\{a_\ell\}_{\ell \in [k]}, \{c_\ell\}_{\ell \in [k]}) &= \max_{\ell \in [k]} \|a_{\sigma_1(\ell)} - c_\ell\|_2 \\ d_{2,\infty}(\{b_\ell\}_{\ell \in [k]}, \{c_\ell\}_{\ell \in [k]}) &= \max_{\ell \in [k]} \|b_{\sigma_2(\ell)} - c_\ell\|_2. \end{aligned}$$

By definition,

$$\begin{aligned}
 d_{2,\infty}(\{a_\ell\}_{\ell \in [k]}, \{b_\ell\}_{\ell \in [k]}) &\leq \max_{\ell \in [k]} \|a_{\sigma_1(\ell)} - b_{\sigma_2(\ell)}\|_2 \\
 &\leq \max_{\ell \in [k]} \|a_{\sigma_1(\ell)} - c_\ell\|_2 + \max_{\ell \in [k]} \|b_{\sigma_2(\ell)} - c_\ell\|_2 \\
 &= d_{2,\infty}(\{a_\ell\}_{\ell \in [k]}, \{c_\ell\}_{\ell \in [k]}) + d_{2,\infty}(\{b_\ell\}_{\ell \in [k]}, \{c_\ell\}_{\ell \in [k]}).
 \end{aligned}$$

It is clear that any bounded closed subset of $\{\mathbb{R}^d\}^k$ is compact in the topology induced by $d_{2,\infty}$.

Another commonly used metric to measure the distance between two closed sets in a metric space $(\mathcal{M}, d)$ is the Hausdorff distance. For any two closed sets $X, Y \subset \mathcal{M}$, the Hausdorff distance between them is given by

$$(2.1) \quad d_H(X, Y) := \max \left\{ \sup_{x \in X} \inf_{y \in Y} d(x, y), \sup_{y \in Y} \inf_{x \in X} d(x, y) \right\}.$$

It is easy to check that $d_{2,\infty}$ is stronger than the Hausdorff distance for sets consisting of k points in $\mathbb{R}^d$. In particular, for $\{a_\ell\}_{\ell \in [k]}, \{b_\ell\}_{\ell \in [k]} \in \{\mathbb{R}^d\}^k$,

$$(2.2) \quad d_H(\{a_\ell\}_{\ell \in [k]}, \{b_\ell\}_{\ell \in [k]}) \leq d_{2,\infty}(\{a_\ell\}_{\ell \in [k]}, \{b_\ell\}_{\ell \in [k]}).$$

In general, $d_{2,\infty}$ and d_H are not exactly the same. For example, consider $A = \{(0, 0), (1, 0), (2, 0)\}$ and $B = \{(0.5, 0), (2, 0.1), (2, -0.1)\}$ in $\{\mathbb{R}^2\}^3$. It is easy to check that $d_H(A, B) = 0.5 < \sqrt{1.01} = d_{2,\infty}(A, B)$.

We will first establish the existence of solutions to (1.2) as stated in Theorem 1.1. We use the following lemma about the objective function, $F_\nu: \{\mathbb{R}^d\}^k \rightarrow \mathbb{R}$, defined in (1.2).

Lemma 2.1. *For any probability measure ν on $\mathbb{R}^d$ which is square-integrable, the objective function defined by*

$$F_\nu(\{a_\ell\}_{\ell \in [k]}) = \left(\int_{\mathbb{R}^d} d^2(x, \text{co}(\{a_\ell\}_{\ell \in [k]})) d\nu(x) \right)^{1/2}$$

is a 1-Lipschitz continuous function from the metric space $(\{\mathbb{R}^d\}^k, d_{2,\infty})$ to $\mathbb{R}$, where $d_{2,\infty}$ is defined in Definition 1.4.

Proof. Square-integrability guarantees that F_ν is finite for all $\{a_\ell\}_{\ell \in [k]} \in \{\mathbb{R}^d\}^k$. The rest of the proof follows from direct computation. For any $\{a_\ell\}_{\ell \in [k]}, \{b_\ell\}_{\ell \in [k]} \in \{\mathbb{R}^d\}^k$, we compute

$$\begin{aligned}
 |F_\nu(\{a_\ell\}_{\ell \in [k]}) - F_\nu(\{b_\ell\}_{\ell \in [k]})| &\leq \left(\int_{\mathbb{R}^d} |d(x, \text{co}(\{a_j\}_{j \in [k]})) - d(x, \text{co}(\{b_j\}_{j \in [k]}))|^2 d\nu(x) \right)^{1/2} \\
 &\leq \left(\int_{\mathbb{R}^d} |d_H(\text{co}(\{a_j\}_{j \in [k]}), \text{co}(\{b_j\}_{j \in [k]}))|^2 d\nu(x) \right)^{1/2} \\
 &= d_H(\text{co}(\{a_j\}_{j \in [k]}), \text{co}(\{b_j\}_{j \in [k]})) \\
 &\leq d_H(\{a_j\}_{j \in [k]}, \{b_j\}_{j \in [k]}) \\
 &\stackrel{(2.2)}{\leq} d_{2,\infty}(\{a_j\}_{j \in [k]}, \{b_j\}_{j \in [k]}). \quad \blacksquare
 \end{aligned}$$

Proof of Theorem 1.1. Note that the admissible set $\{\text{supp}(\mu)\}^k \subset \{\mathbb{R}^d\}^k$ is closed in $d_{2,\infty}$ and has bounded diameter, therefore is compact. Theorem 1.1 is an immediate consequence of Lemma 2.1 together with the fact that a continuous function achieves its infimum on a compact set. This proves the existence of minimizers. The second part follows by noting that $\bar{x} \in \text{co}(\text{supp}(\nu))$. ■

We will next prove the consistency result stated in Theorem 1.5.

Proof of Theorem 1.5. Let $\mathcal{A}$ be the solution set of (1.3). It is clear that $\mathcal{A}$ is non-empty by Theorem 1.1. Our goal is to show that $\{A_N\}_{N \in \mathbb{N}}$ has a convergent subsequence whose limit lies in $\mathcal{A}$. Since $\text{supp}(\mu)$ is compact, we can apply the Bolzano–Weierstrass theorem to the k elements of $\{A_N\}_{N \in \mathbb{N}}$ with a diagonal argument to find a convergent subsequence $\{A_{N_m}\}_{m \in \mathbb{N}}$ of $\{A_N\}_{N \in \mathbb{N}}$ such that $A_{N_m} \rightarrow A_\star$ in $d_{2,\infty}$ for some $A_\star \subset \text{co}(\text{supp}(\mu))$, i.e., $d_H(A_{N_m}, A_\star) \rightarrow 0$ as $m \rightarrow \infty$. We claim $A_\star \in \mathcal{A}$ and this will be shown via a triangle-inequality type of argument.

Compared to (1.3), the discrete problem (1.1) has not only discrete objective function but also discrete constraints. To bridge the gap, we introduce an intermediate discrete minimization problem with the objective function in (1.1) but the constraint in (1.3):

$$(2.3) \quad \begin{aligned} \min_{\{a_\ell\}_{\ell \in [k]}} \quad & F_{\mu_N}(A) \\ \text{s.t.} \quad & a_\ell \in \text{co}(\text{supp}(\mu)), \quad \ell \in [k] \end{aligned}$$

and denote a solution to (2.3) by A'_N for each N . Similar to as in Theorem 1.2, we can take A'_N to be located on the boundary of $\text{co}(\text{supp}(\mu))$. Below, it is also convenient to define the set of points consisting of the convex projection of the points in A'_N onto $\text{co}(\text{supp}(\mu_N))$,

$$(2.4) \quad A''_N = \text{Proj}_{\text{co}(\text{supp}(\mu_N))}(A'_N).$$

Note that since $A'_N \subset \overline{\mathbb{R}^d \setminus \text{co}(\text{supp}(\mu_N))}$, we have that $A''_N \subset \partial(\text{co}(\text{supp}(\mu_N)))$.

The rest of the proof can be summarized as two steps: (i) first show that $d_H(A'_{N_m}, A''_{N_m}) \rightarrow 0$ as $m \rightarrow \infty$ μ -a.s. and (ii) then use it as a bridge to obtain that $A_\star$ is an optimal solution for (1.3).

To show (i), observe that $\text{co}(\{x_j\}_{j \in [N]}) \subseteq \text{co}(\{x_j\}_{j \in [N+1]}) \subseteq \text{co}(\text{supp}(\mu))$. Since μ has compact support, $K := \lim_{N \rightarrow \infty} \text{co}(\{x_j\}_{j \in [N]})$ exists and is a convex set contained in $\text{co}(\text{supp}(\mu))$. It follows from a contradiction argument that $d_H(K, \text{co}(\text{supp}(\mu))) = 0$ μ -a.s.. If not, there would exist a point $x \in \text{co}(\text{supp}(\mu))$ such that $d(x, K) > 0$. Since x can be written as a convex combination of finitely many points in $\text{supp}(\mu)$, say $x^{(1)}, \dots, x^{(r)}$ for some $r > 0$, at least one of them has positive distance to K . Without loss of generality we assume $d(x^{(1)}, K) > 0$. This implies that the sequence $\{x_j\}$ does not intersect with a ball centered at $x^{(1)}$ with radius $\frac{1}{2}d(x^{(1)}, K)$, i.e., $\{x_j\} \cap B(x^{(1)}, \frac{1}{2}d(x^{(1)}, K)) = \emptyset$. But this happens on a μ -null set:

$$\mu \left(\{x_j\} \cap B(x^{(1)}, \frac{1}{2}d(x^{(1)}, K)) = \emptyset \right) \leq \lim_{N \rightarrow \infty} (1 - \mu(B(x^{(1)}, \frac{1}{2}d(x^{(1)}, K))))^N = 0,$$

where $\mu(B(x^{(1)}, \frac{1}{2}d(x^{(1)}, K))) > 0$ because $x^{(1)} \in \text{supp}(\mu)$. Therefore, $d_H(K, \text{co}(\text{supp}(\mu))) = 0$. The desired result follows by observing that

$$(2.5) \quad d_H(A'_{N_m}, A''_{N_m}) \leq d_H(\text{co}\{x_j\}_{j \in [N]}, \text{co}(\text{supp}(\mu))) \rightarrow 0.$$

For step (ii), we note that (2.5) together with $A_{N_m} \rightarrow A_\star$ and $\mu_{N_m} \rightharpoonup \mu$ μ -a.s., implies that for any $A \in \mathcal{A}$ and $\varepsilon > 0$, there exists a sufficiently large M such that for $m > M$,

$$(2.6) \quad d_H(A_{N_m}, A_\star) \leq \varepsilon$$

$$(2.7) \quad d_H(A'_{N_m}, A''_{N_m}) < \varepsilon$$

$$(2.8) \quad \left(\int_{\mathbb{R}^d} d^2(x, A_\star) d\mu(x) \right)^{1/2} \leq \left(\int_{\mathbb{R}^d} d^2(x, A_\star) d\mu_{N_m}(x) \right)^{1/2} + \varepsilon$$

$$(2.9) \quad \left(\int_{\mathbb{R}^d} d^2(x, A) d\mu_{N_m}(x) \right)^{1/2} \leq \left(\int_{\mathbb{R}^d} d^2(x, A) d\mu(x) \right)^{1/2} + \varepsilon.$$

Therefore, for $m > M$,

$$\begin{aligned} & \left(\int_{\mathbb{R}^d} d^2(x, A_\star) d\mu(x) \right)^{1/2} \\ & \stackrel{(2.8)}{\leq} \left(\int_{\mathbb{R}^d} d^2(x, A_\star) d\mu_{N_m}(x) \right)^{1/2} + \varepsilon \\ & \stackrel{\text{pf. Lem 2.1}}{\leq} \left(\int_{\mathbb{R}^d} d^2(x, A_{N_m}) d\mu_{N_m}(x) \right)^{1/2} + \left(\int_{\mathbb{R}^d} d_H^2(A_\star, A_{N_m}) d\mu_{N_m}(x) \right)^{1/2} + \varepsilon \\ & \stackrel{\text{def. } A_{N_m}, (2.6)}{\leq} \left(\int_{\mathbb{R}^d} d^2(x, A''_{N_m}) d\mu_{N_m}(x) \right)^{1/2} + 2\varepsilon \\ & \stackrel{\text{pf. Lem 2.1}}{\leq} \left(\int_{\mathbb{R}^d} d^2(x, A'_{N_m}) d\mu_{N_m}(x) \right)^{1/2} + \left(\int_{\mathbb{R}^d} d_H^2(A''_{N_m}, A'_{N_m}) d\mu_{N_m}(x) \right)^{1/2} + 2\varepsilon \\ & \stackrel{\text{def. } A'_{N_m}, (2.7)}{\leq} \left(\int_{\mathbb{R}^d} d^2(x, A) d\mu_{N_m}(x) \right)^{1/2} + 3\varepsilon \\ & \stackrel{(2.9)}{\leq} \left(\int_{\mathbb{R}^d} d^2(x, A) d\mu(x) \right)^{1/2} + 4\varepsilon. \end{aligned}$$

Setting $\varepsilon \rightarrow 0$ yields

$$\int_{\mathbb{R}^d} d^2(x, A_\star) d\mu(x) \leq \int_{\mathbb{R}^d} d^2(x, A) d\mu(x).$$

This finishes the proof of the first part of the theorem. The second part of the theorem follows by noting that whenever $\mathcal{A}$ contains only one element, $A_\star$ does not depend on the choice of the subsequence $\{A_{N_m}\}$. A moment thought on the compactness of $\text{supp}(\mu)$ reveals that the result in the first part holds for the whole sequence. $\blacksquare$

For the proof of Theorem 1.8, we will need a uniform bound on the convergence rate on the empirical measure of μ in a class of test functions, which is given by the following lemmas.

Lemma 2.2 (Dudley's inequality). *Let $(X_t)_{t \in T}$ be a random process on a metric space (T, d) with sub-gaussian increments:*

$$\|X_t - X_s\|_{\Psi_2} \leq Kd(t, s), \quad \forall s, t \in T,$$

where $\|\cdot\|_{\Psi_2}$ is the sub-gaussian norm and $K > 0$ is a constant. Then, for every $u \geq 0$, the event

$$\sup_{t,s \in T} |X_t - X_s| \leq CK \left(\int_0^\infty \sqrt{\log \mathcal{N}(T, d, \varepsilon)} d\varepsilon + u \cdot \text{diam}(T) \right)$$

holds with probability at least $1 - 2\exp(-u^2)$, where $\mathcal{N}(T, d, \varepsilon)$ is the ε -covering number of T under metric d .

The proof of [Lemma 2.2](#) can be found in [\[32, Theorem 8.16\]](#).

[Lemma 2.3](#). Under the same conditions as [Theorem 1.5](#),

$$\sup_{A \in \{\text{co}(\text{supp}(\mu))\}^k} \left| \frac{1}{N} \sum_{i \in [N]} d^2(x_i, \text{co}(A)) - \int_{\mathbb{R}^d} d^2(x, A) d\mu \right| \lesssim \left(\frac{\log N}{N} \right)^{1/2} \quad \mu\text{-a.s.}$$

Proof. For fixed N , consider the random process

$$Y_A = N^{-1} \sum_{i \in [N]} d^2(x_i, \text{co}(A)) - \int_{\mathbb{R}^d} d^2(x, A) d\mu,$$

where the index set is equipped with the metric $d_{2,\infty}$. One can verify that $\{d^2(x_i, \text{co}(A)) - d^2(x_i, \text{co}(B))\}_{i \in [N]}$ are i.i.d. random variables which are bounded by $2Dd_{2,\infty}(A, B)$ for $A, B \in \{\text{co}(\text{supp}(\mu))\}^k$, where D is the diameter of $\text{co}(\text{supp}(\mu))$. It follows from a similar computation as in [\[32, Section 8.2.2\]](#) that

$$\|Y_A - Y_B\|_{\Psi_2} \lesssim \frac{D}{\sqrt{N}} d_{2,\infty}(A, B).$$

Therefore, by [Lemma 2.2](#), with probability at least $1 - 2\exp(-u^2)$,

$$(2.10) \quad \sup_{A \in \{\text{co}(\text{supp}(\mu))\}^k} |Y_A| \lesssim \frac{D}{\sqrt{N}} \left(\int_0^\infty \sqrt{\log \mathcal{N}(\{\text{co}(\text{supp}(\mu))\}^k, d_{2,\infty}, \varepsilon)} d\varepsilon + kDu \right) + |Y_B|,$$

where $B \in \{\text{co}(\text{supp}(\mu))\}^k$ is some fixed element. Note that

$$(2.11) \quad \begin{aligned} \mathcal{N}(\{\text{co}(\text{supp}(\mu))\}^k, d_{2,\infty}, \varepsilon) &\leq (\mathcal{N}(\text{co}(\text{supp}(\mu)), \|\cdot\|_2, \varepsilon))^k \\ &\leq (\mathcal{N}(B_z(D), \|\cdot\|_2, \varepsilon/2))^k \leq \left(\frac{6D}{\varepsilon} \right)^{dk}, \end{aligned}$$

where $z \in \text{co}(\text{supp}(\mu))$, $B_z(D) = \{x: \|x - z\|_2 \leq D\}$, and the last inequality follows from estimates on the Euclidean covering number of ℓ_2 balls, see [\[32\]](#). Plugging (2.11) into (2.10) and setting $u = \sqrt{2\log N}$ yields that with probability at least $1 - 2N^{-2}$,

$$(2.12) \quad \sup_{A \in \{\text{co}(\text{supp}(\mu))\}^k} |Y_A| \lesssim \left(\frac{\log N}{N} \right)^{1/2} + |Y_B|.$$

On the other hand, Y_B is the sum of i.i.d. centered bounded random variables, by Hoeffding's inequality, $|Y_B| \lesssim (\log N/N)^{1/2}$ with probability at least $1 - N^{-2}$. An application of the Borel–Cantelli lemma finishes the proof. ■

Proof of Theorem 1.8. Fix $N > 0$. Adopting the same notation as in the previous proof, we let $A_\star$ and A_N denote two archetype pointsets for the continuum and discrete problems, respectively. Moreover, by Theorem 1.2, we may assume that they are on the boundary of the convex hull of their respective support. Similar to (2.4), define $A'_\star$ as the projection of $A_\star$ to $\text{co}(\text{supp}(\mu_N))$. It is easy to check that

$$(2.13) \quad d_H(A_\star, A'_\star) \leq d_H(\text{co}(\text{supp}(\mu)), \text{co}(\text{supp}(\mu_N))).$$

Without loss of generality, we assume that $F_{\mu_N}(A_N) \geq F_\mu(A_\star)$. Repeating the same trick in the proof of Theorem 1.5,

$$(2.14) \quad \begin{aligned} F_\mu(A_N) - F_\mu(A_\star) &\leq |F_\mu(A_N) - F_{\mu_N}(A_N)| + F_{\mu_N}(A_N) - F_\mu(A_\star) \\ &\leq |F_\mu(A_N) - F_{\mu_N}(A_N)| + F_{\mu_N}(A'_\star) - F_\mu(A_\star) \\ &\leq |F_\mu(A_N) - F_{\mu_N}(A_N)| + |F_{\mu_N}(A_\star) - F_\mu(A_\star)| + |F_{\mu_N}(A'_\star) - F_{\mu_N}(A_\star)| \\ &\leq \underbrace{|F_\mu(A_N) - F_{\mu_N}(A_N)| + |F_{\mu_N}(A_\star) - F_\mu(A_\star)|}_{(i)} + \underbrace{|F_{\mu_N}(A'_\star) - F_{\mu_N}(A_\star)|}_{(ii)} \rightarrow 0 \end{aligned}$$

as $N \rightarrow \infty$, where (i) $\rightarrow 0$ due to Lemma 2.3 and (ii) $\rightarrow 0$ due to the proof of Theorem 1.5. This finishes the first part of the proof.

To obtain the convergence rate, note that (2.12) combined with the triangle inequality implies that (i) $\lesssim (\log N/N)^{1/2}$ with probability at least $1 - 3N^{-2}$. Bounding (ii) for an arbitrary μ is intractable in general. However, under the α -cap condition (with parameters L and η), it has been established in [4, Theorem 1] that there exists a sufficiently large integer $N(\eta)$ (non-increasing in η), such that for $N > N(\eta)$, with μ at least $1 - \delta$,

$$(2.15) \quad d_H(\text{co}(\text{supp}(\mu)), \text{co}(\text{supp}(\mu_N))) \lesssim \left(\frac{d \log(1/\delta)}{NL} \right)^{1/\alpha},$$

Setting $\delta = N^{-2}$ and noting $\alpha \geq d \geq 2$ when μ has a density completes the proof. $\blacksquare$

Remark 2.4. A very large variance¹ may accelerate the convergence in (1.5). To see this, for fixed set $D \subset \mathbb{R}^d$, let

$$\mathcal{M} = \{\nu : \nu \text{ is a probability measure on } \mathbb{R}^d \text{ such that } \text{supp}(\nu) \subset D\}.$$

Consider the elements in $\mathcal{M}$ that maximize the variance:

$$(2.16) \quad \mathcal{V} = \arg \max_{\nu \in \mathcal{M}} \mathbb{E}_\nu [\|x - \mathbb{E}_\nu[x]\|_2^2].$$

We claim that every $\mu \in \mathcal{V}$ is supported on at most $d + 1$ affinely independent points in D ; for a proof, see Appendix A. Therefore, any element in $\mathcal{M}$ with variance approximating the variance of μ should concentrate on some finite set of affinely independent points.

¹Note that variance is defined for real-valued random variables. For a random vector $X \in \mathbb{R}^d$, we consider $\mathbb{E}[\|X - \mathbb{E}[X]\|_2^2]$ as its (total) variance, which is the trace of its variance-covariance matrix.

3. Example: uniform probability measure on the unit disk. In this section, we consider the continuous archetypal analysis problem (1.3) for the uniform probability measure on the unit disk.

Proposition 3.1. *Let $k \geq 3$, $\mathbb{D} \subset \mathbb{R}^2$ be the Euclidean unit disk, and μ the uniform probability measure on $\mathbb{D}$. The minimizers to (1.3) are extremal points for regular k -polygons inscribed in $\mathbb{D}$ and have squared objective value*

$$F_\mu^2 = kI\left(\frac{2\pi}{k}\right),$$

where $I: [0, \pi] \rightarrow \mathbb{R}$ is defined in (3.2). The minimizers are unique modulo rotation.

Proof. Let R_k denote a regular k -polygon inscribed in the unit disc and A_k its vertices. Using symmetry, the squared objective value for A_k can be simplified,

$$(3.1) \quad F_\mu^2(A_k) = \int_{\mathbb{D}} d^2(x, R_k) d\mu(x) = \frac{1}{\pi} \int_{\mathbb{D}} d^2(x, R_k) dx = \frac{k}{\pi} \int_{\vartriangle_{\frac{2\pi}{k}}} d^2(x, R_k) dx.$$

Here, $\vartriangle_{\frac{2\pi}{k}}$ denotes a sector of the disk with angle $\frac{2\pi}{k}$.

For future reference, it is useful to record a slightly more general calculation. Let

$$I(\alpha) = \frac{1}{\pi} \int_{\vartriangle_\alpha} d^2(x, \triangle_\alpha) dx, \quad \alpha \in [0, \pi].$$

Here, denoting $A = (1, 0)$, $O = (0, 0)$, and $B = (\cos \alpha, \sin \alpha)$, we have used the notation that $\vartriangle_\alpha$ is the sector of the unit disc $\vartriangle AOB$ and $\triangle_\alpha$ is the triangle with vertices A , O , and B . The chord $\overline{AB}$ lies on the line

$$\left\{ (x, y) \in \mathbb{R}^2 : \begin{pmatrix} \cos \frac{\alpha}{2} \\ \sin \frac{\alpha}{2} \end{pmatrix} \cdot \begin{pmatrix} x \\ y \end{pmatrix} = \cos \frac{\alpha}{2} \right\},$$

so $\overline{AB}$ can be explicitly parameterized by

$$\overline{AB} = \left\{ (r \cos \theta, r \sin \theta) : r = \frac{\cos \frac{\alpha}{2}}{\cos(\theta - \frac{\alpha}{2})}, \theta \in [0, \alpha] \right\}.$$

For a point, $(r \cos \theta, r \sin \theta) \in \vartriangle_\alpha \setminus \triangle_\alpha$, we compute

$$\begin{aligned} d^2((r \cos \theta, r \sin \theta), \triangle_\alpha) &= d^2((r \cos \theta, r \sin \theta), \overline{AB}) \\ &= \left(\begin{pmatrix} \cos \frac{\alpha}{2} \\ \sin \frac{\alpha}{2} \end{pmatrix} \cdot \begin{pmatrix} r \cos \theta \\ r \sin \theta \end{pmatrix} - \cos \frac{\alpha}{2} \right)^2 \\ &= \left(r \cos \left(\theta - \frac{\alpha}{2} \right) - \cos \frac{\alpha}{2} \right)^2. \end{aligned}$$




Figure 2. The function $I: [0, \pi] \rightarrow \mathbb{R}$ defined in (3.2).

For $\alpha \in [0, \pi]$, we compute

$$(3.2a) \quad I(\alpha) = \frac{1}{\pi} \int_{\triangle_\alpha} d^2(x, \triangle_\alpha) dx$$

$$(3.2b) \quad = \frac{2}{\pi} \int_0^{\frac{\alpha}{2}} \int_{\frac{\cos \frac{\alpha}{2}}{\cos(\frac{\alpha}{2}-\theta)}}^1 \left(r \cos\left(\theta - \frac{\alpha}{2}\right) - \cos \frac{\alpha}{2} \right)^2 r dr d\theta$$

$$(3.2c) \quad = \frac{1}{2\pi} \left(\frac{\alpha}{4} - \frac{13}{12} \sin \alpha + \alpha \cos^2 \frac{\alpha}{2} - \frac{1}{3} \sin \frac{\alpha}{2} \cos^3 \frac{\alpha}{2} \right),$$

It can easily be checked that

- $I(0) = 0$ and $I(\pi) = \frac{1}{8}$,
- $I(\alpha)$ is an increasing positive function on $[0, \pi]$, and
- $I(\alpha)$ is strictly convex on $[0, \pi]$.

From (3.1), we obtain that $F_\mu^2(A_k) = kI\left(\frac{2\pi}{k}\right)$, which is a decreasing function in k .

We now consider the optimization problem (1.3). We first note that all the boundary points of $\mathbb{D}$ are extreme points, which has infinite cardinality. In this particular case, it is easy to check that for each $k \in \mathbb{N}$, the minimizers to (1.3) must lie on $\partial\mathbb{D}$, so their convex hull must be a cyclic polygon inscribed in the unit disc. Let P_α be a cyclic polygon inscribed in the unit disc with central angles $\alpha_1, \dots, \alpha_k$ where $\alpha_j \in [0, 2\pi]$, $j \in [k]$ and $\sum \alpha_j = 2\pi$. We first argue that if P_α were to be optimal, then it could not be contained in a half disc, H , and therefore, we may assume that $\alpha_j \in [0, \pi]$, $j \in [k]$. This follows from the monotonicity property (if $P_\alpha \subset H$, then $F_\mu(P_\alpha) > F_\mu(H)$) and the fact that $F_\mu(H) > F_\mu(A_3)$. To see this last inequality, we compute

$$F_\mu^2(H) = \int_{\mathbb{D}} d^2(x, H) d\mu(x) = \frac{1}{\pi} \int_{\mathbb{D}} \int_{\mathbb{D}} d^2(x, H) dx = I(\pi) = \frac{1}{8}$$

and $F_\mu^2(A_3) = 3I\left(\frac{2\pi}{3}\right) \approx 0.035$.

The optimization problem in (1.3) then reduces to

$$\begin{aligned} \min \quad & \sum_{j=1}^k I(\alpha_j) \\ \text{s.t.} \quad & \alpha_j \in [0, \pi] \\ & \sum_{j=1}^k \alpha_j = 2\pi. \end{aligned}$$

This is a convex optimization problem with convex constraints. By Jensen's inequality, the unique solution is given by $\alpha_1 = \dots = \alpha_k = \frac{2\pi}{k}$, as desired. $\blacksquare$

4. Proof of Theorems 1.11, 1.12, and 1.14 for probability measures with unbounded support. We will establish existence of solutions to (1.7) as stated in Theorem 1.11. We use the following lemma about the objective function, $F_{\nu, \alpha}: \{\mathbb{R}^d\}^k \rightarrow \mathbb{R}$, defined in (1.7).

Lemma 4.1. *For any square-integrable ν , $F_{\nu, \alpha}$ is a $(1+2\sqrt{\alpha})$ -Lipschitz continuous function from $(\{\mathbb{R}^d\}^k, d_{2, \infty})$ to $\mathbb{R}$.*

Proof. For any $A_1 = \{a_\ell^{(1)}\}_{\ell \in [k]}, A_2 = \{a_\ell^{(2)}\}_{\ell \in [k]} \in \{\mathbb{R}^d\}^k$, let σ be the permutation of $[k]$ such that

$$d_{2, \infty}(A_1, A_2) = \max_{\ell \in [k]} \|a_\ell^{(1)} - a_{\sigma(\ell)}^{(2)}\|_2.$$

First applying the triangle inequality and then the elementary inequality $\sqrt{a+b} \leq \sqrt{a} + \sqrt{b}$,

$$\begin{aligned} & |F_{\nu, \alpha}(A_1) - F_{\nu, \alpha}(A_2)| \\ & \leq |F_\nu(A_1) - F_\nu(A_2)| + \sqrt{\alpha} \left| \left(\frac{1}{k} \sum_{\ell \in [k]} \|a_\ell^{(1)} - \bar{a}^{(1)}\|_2^2 \right)^{1/2} - \left(\frac{1}{k} \sum_{\ell \in [k]} \|a_\ell^{(2)} - \bar{a}^{(2)}\|_2^2 \right)^{1/2} \right|. \end{aligned}$$

By Lemma 2.1, the first term on the right-hand side is bounded by $d_{2, \infty}(A_1, A_2)$. The second term is further bounded using the triangle inequality as follows:

$$\begin{aligned} & \sqrt{\alpha} \left| \left(\frac{1}{k} \sum_{\ell \in [k]} \|a_\ell^{(1)} - \bar{a}^{(1)}\|_2^2 \right)^{1/2} - \left(\frac{1}{k} \sum_{\ell \in [k]} \|a_\ell^{(2)} - \bar{a}^{(2)}\|_2^2 \right)^{1/2} \right| \\ & \leq \sqrt{\alpha} \left(\left(\frac{1}{k} \sum_{\ell \in [k]} \|a_\ell^{(1)} - a_{\sigma(\ell)}^{(2)}\|_2^2 \right)^{1/2} + \left(\frac{1}{k} \sum_{\ell \in [k]} \|\bar{a}^{(1)} - \bar{a}^{(2)}\|_2^2 \right)^{1/2} \right) \\ & \leq 2\sqrt{\alpha} d_{2, \infty}(A_1, A_2). \end{aligned}$$

Putting the estimates together completes the proof. $\blacksquare$

Proof of Theorem 1.11. Denote by $F_{\nu,\alpha}^* \geq 0$ as the infimum objective value. Square-integrability of ν ensures that $F_{\nu,\alpha}^* < \infty$. Let $A_m = \{a_\ell^{(m)}\}_{\ell \in [k]}$ be a non-increasing minimizing sequence such that for $m \geq 1$,

$$2F_{\nu,\alpha}^* \geq F_{\nu,\alpha}(A_1) \geq F_{\nu,\alpha}(A_m) \geq F_{\nu,\alpha}(A_{m+1}).$$

We first observe that the diameter of $\{A_m\}_{m \in \mathbb{N}}$ is uniformly bounded. Let $r_1 = 4\sqrt{\frac{k}{\alpha}}F_{\nu,\alpha}^*$. For any set $A = \{a_\ell\}_{\ell \in [k]}$ with $\text{diam}(A) > r_1$,

$$F_{\nu,\alpha}(A) \geq \sqrt{\frac{\alpha}{k}} \sum_{\ell \in [k]} \|a_\ell - \bar{a}\|_2^2 \geq \sqrt{\frac{\alpha}{k}} \frac{\text{diam}(A)}{2} > 2F_{\nu,\alpha}^* \geq F_{\nu,\alpha}(A_1).$$

This implies that $\sup_m \text{diam}(A_m) \leq r_1$. We next show that $\{A_m\}_{m \in \mathbb{N}}$ cannot escape to infinity as m increases, which is equivalent to showing that $\{a_1^{(m)}\}_{m \in \mathbb{N}}$ is uniformly bounded. To see this, note that most mass of ν is concentrated in a ball centered at the origin. In particular, we can find a ball $B(r_2)$ with radius $r_2 > 0$ such that $\nu(B(r_2)) \geq 1/2$. We claim that $\{a_1^{(m)}\}_{m \in \mathbb{N}}$ is contained in a ball of radius $r_3 := r_1 + r_2 + 2\sqrt{2}F_{\nu,\alpha}^*$. Suppose $\|a_1^{(m)}\|_2 > r_3$ for some m . Then the distance between any $x \in B(r_2)$ and A_m is greater than $r_3 - r_1 - r_2 = 2\sqrt{2}F_{\nu,\alpha}^*$. Therefore,

$$F_{\nu,\alpha}(A_m) \geq \left(\int_{B(r_2)} d^2(x, A_m) d\nu(x) \right)^{1/2} > \left(\int_{B(r_2)} 8(F_{\nu,\alpha}^*)^2 d\nu(x) \right)^{1/2} \geq 2F_{\nu,\alpha}^* \geq F_{\nu,\alpha}(A_1),$$

contradicting the hypothesis that $\{A_m\}_{m \in \mathbb{N}}$ is a decreasing minimizing sequence. Hence we have proved that $\{A_m\}_{m \in \mathbb{N}}$ is contained in $B(r_1 + r_3)$, i.e., $\{A_m\}_{m \in \mathbb{N}}$ is uniformly bounded. Applying the Bolzano–Weierstrass theorem componentwise yields that $\{A_m\}_{m \in \mathbb{N}}$ admits a convergent subsequence in the $d_{2,\infty}$ metric, say $A_{m_n} \rightarrow A_\star$ as $n \rightarrow \infty$. Since $\text{co}(\text{supp}(\nu))$ is closed and $A_m \subset \text{co}(\text{supp}(\nu))$ for every m , $A_\star \subset \text{co}(\text{supp}(\nu))$. It follows from the fact that $F_{\nu,\alpha}$ is continuous with respect to $d_{2,\infty}$ -metric that

$$F_{\nu,\alpha}(A_\star) = \lim_{n \rightarrow \infty} F_{\nu,\alpha}(A_{m_n}) = F_{\nu,\alpha}^*.$$

Hence, $A_\star$ is a minimizer of problem (1.7). The inequality $F_{\nu,\alpha}(A_\star) \leq (\int_{\mathbb{R}^d} \|x - \bar{x}\|_2^2 d\nu(x))^{1/2}$ follows from evaluating $F_{\nu,\alpha}$ at $A = \{\bar{x}\}^k$. ■

We will next prove the consistency result stated in Theorem 1.12.

Proof of Theorem 1.12.. Let $\mathcal{A}$ be the solution set of (1.7) with $\nu = \mu$. Let A_N be a minimizer to problem (1.7) with $\nu = \mu_N$ and $A \in \mathcal{A}$. Since $\mu_N \rightarrow \mu$ μ -a.s., it is possible to find a centered ball of sufficiently large radius r_2 such that

$$\min \left\{ \sup_{N \in \mathbb{N}} \mu_N(B(r_2)), \mu(B(r_2)) \right\} \geq \frac{1}{2}.$$

It follows from the proof of [Theorem 1.11](#) that

$$\begin{aligned} A_N &\subset B\left((8k^{1/2}\alpha^{-1/2} + 2\sqrt{2})F_{\mu_N, \alpha}^* + r_2\right) \\ A &\subset B\left((8k^{1/2}\alpha^{-1/2} + 2\sqrt{2})F_{\mu, \alpha}^* + r_2\right). \end{aligned}$$

To show that $\{A_N\}_{N \in \mathbb{N}}$ admits a convergent subsequence in $d_{2, \infty}$, it suffices to show that $F_{\mu_N, \alpha}(A_N)$ is uniformly bounded in N . To see this, note that for every N ,

$$F_{\mu_N, \alpha}(A_N) \leq F_{\mu_N, \alpha}(\{x_1\}^k) = \frac{1}{N} \sum_{i \in [N]} (x_i - x_1)^2 \rightarrow \int_{\mathbb{R}^d} \|x - x_1\|_2^2 d\mu(x) < \infty.$$

Hence, for

$$R := \left(8k^{1/2}\alpha^{-1/2} + 2\sqrt{2}\right) \max \left\{ \sup_{N \in \mathbb{N}} F_{\mu_N, \alpha}^*, F_{\mu, \alpha}^* \right\} + r_2 < \infty,$$

we have

$$(\cup_{N \in \mathbb{N}} A_N) \cup A \subset B(R) \cap \text{co}(\text{supp}(\mu)).$$

Since $B(R) \cap \text{co}(\text{supp}(\mu))$ is compact, by the Bolzano–Weierstrass theorem, $\{A_N\}_{N \in \mathbb{N}}$ has a convergent subsequence $\{A_{N_m}\}_{m \in \mathbb{N}}$ such that $A_{N_m} \rightarrow A_\star$ in $d_{2, \infty}$ and in the Hausdorff metric. It is clear that $A_\star \subset \text{co}(\text{supp}(\mu))$ thus is a natural candidate for the minimizer of the continuous problem. To show that it is, we use the fact that for any $\varepsilon > 0$, there exists a sufficiently large M such that the following statements hold true whenever $m > M$:

$$(4.1) \quad A \subset B(R) \subset \text{co}(X_{N_m})$$

$$(4.2) \quad d_H(A_{N_m}, A_\star) \leq \varepsilon$$

$$(4.3) \quad \int_{\mathbb{R}^d} d^2(x, A_\star) d\mu(x) \leq \int_{\mathbb{R}^d} d^2(x, A_\star) d\mu_{N_m}(x) + \varepsilon$$

$$(4.4) \quad \int_{\mathbb{R}^d} d^2(x, A) d\mu_{N_m}(x) \leq \int_{\mathbb{R}^d} d^2(x, A) d\mu(x) + \varepsilon.$$

Therefore, for $m > M$,

$$\begin{aligned} F_{\mu, \alpha}(A_\star) &= \left(\int_{\mathbb{R}^d} d^2(x, A_\star) d\mu(x) + \alpha V(A_\star) \right)^{1/2} \\ &\stackrel{(4.3)}{\leq} \left(\int_{\mathbb{R}^d} d^2(x, A_\star) d\mu_{N_m}(x) + \alpha V(A_\star) + \varepsilon \right)^{1/2} \\ &\leq \left(\int_{\mathbb{R}^d} d^2(x, A_\star) d\mu_{N_m}(x) + \alpha V(A_\star) \right)^{1/2} + \varepsilon^{1/2} \\ &\stackrel{(4.2), \text{ pf. Lem 4.1}}{\leq} \left(\int_{\mathbb{R}^d} d^2(x, A_{N_m}) d\mu_{N_m}(x) + \alpha V(A_{N_m}) \right)^{1/2} + (1 + 2\sqrt{\alpha})\varepsilon + \varepsilon^{1/2} \\ &\stackrel{\text{def. } A_{N_m}, (4.1)}{\leq} \left(\int_{\mathbb{R}^d} d^2(x, A) d\mu_{N_m}(x) + \alpha V(A) \right)^{1/2} + (1 + 2\sqrt{\alpha})\varepsilon + \varepsilon^{1/2} \\ &\stackrel{(4.4)}{\leq} F_{\mu, \alpha}(A) + (1 + 2\sqrt{\alpha})\varepsilon + 2\varepsilon^{1/2}. \end{aligned}$$

Taking $\varepsilon \rightarrow 0$ yields $F_{\mu,\alpha}(A_\star) \leq F_{\mu,\alpha}(A)$ which completes the first part of the theorem. The second part of the theorem follows by noting that whenever $\mathcal{A}$ contains only one element, $A_\star$ does not depend on the choice of the subsequence $\{A_{N_m}\}$, hence the whole sequence converges to $A_\star$. $\blacksquare$

Proof of Theorem 1.14.. Let $A_{\star,\alpha}$ be a minimizer of (1.7) and $\bar{a}_\alpha = \int_{A_{\star,\alpha}} x \, d\nu(x)$ be the center of $A_{\star,\alpha}$. It follows from the proof of Theorem 1.11 that

$$\text{diam}(A_{\star,\alpha}) \leq 4k^{1/2}\alpha^{-1/2}F_{\nu,\alpha}^\star \leq 4k^{1/2}\alpha^{-1/2} \left(\int_{\mathbb{R}^d} \|x - \bar{x}\|_2^2 \, d\nu(x) \right)^{1/2}.$$

where the second inequality follows by evaluating $F_{\nu,\alpha}$ at $A = A_{\star,\alpha}$. Therefore,

$$\begin{aligned} F_{\nu,\alpha}^\star &= F_{\nu,\alpha}(A_{\star,\alpha}) \\ &\geq \left(\int_{\mathbb{R}^d} d^2(x, A_{\star,\alpha}) \, d\nu(x) \right)^{1/2} \\ &\geq \left(\int_{\mathbb{R}^d} \|x - \bar{a}_\alpha\|_2^2 \, d\nu(x) \right)^{1/2} - \text{diam}(A_{\star,\alpha}) \\ (4.5) \quad &\geq \left(\int_{\mathbb{R}^d} \|x - \bar{x}\|_2^2 \, d\nu(x) + \|\bar{a}_\alpha - \bar{x}\|_2^2 \right)^{1/2} - 4k^{1/2}\alpha^{-1/2} \left(\int_{\mathbb{R}^d} \|x - \bar{x}\|_2^2 \, d\nu(x) \right)^{1/2}. \end{aligned}$$

Since $F_{\nu,\alpha}^\star \leq \left(\int_{\mathbb{R}^d} \|x - \bar{x}\|_2^2 \, d\nu(x) \right)^{1/2}$, $\|\bar{a}_\alpha - \bar{x}\|_2 \rightarrow 0$ as $\alpha \rightarrow \infty$. Therefore, there exists a sufficiently large T (which only depends on ν and k) such that

$$(4.6) \quad \|\bar{a}_\alpha - \bar{x}\|_2^2 \leq 3 \int_{\mathbb{R}^d} \|x - \bar{x}\|_2^2 \, d\nu(x) \quad \alpha > T.$$

Inserting (4.6) this into (4.5) and then applying the elementary inequality $\sqrt{x+y} \geq \sqrt{x} + \frac{y}{2\sqrt{x+y}}$ ($x, y > 0$) yields

$$\begin{aligned} F_{\nu,\alpha}^\star &\geq \left(\int_{\mathbb{R}^d} \|x - \bar{x}\|_2^2 \, d\nu(x) \right)^{1/2} + \frac{1}{2} \left(\int_{\mathbb{R}^d} \|x - \bar{x}\|_2^2 \, d\nu(x) + \|\bar{a}_\alpha - \bar{x}\|_2^2 \right)^{-1/2} \|\bar{a}_\alpha - \bar{x}\|_2^2 \\ &\quad - 4k^{1/2}\alpha^{-1/2} \left(\int_{\mathbb{R}^d} \|x - \bar{x}\|_2^2 \, d\nu(x) \right)^{1/2} \\ &\geq \left(\int_{\mathbb{R}^d} \|x - \bar{x}\|_2^2 \, d\nu(x) \right)^{1/2} + \frac{1}{4} \left(\int_{\mathbb{R}^d} \|x - \bar{x}\|_2^2 \, d\nu(x) \right)^{-1/2} \|\bar{a}_\alpha - \bar{x}\|_2^2 \\ &\quad - 4k^{1/2}\alpha^{-1/2} \left(\int_{\mathbb{R}^d} \|x - \bar{x}\|_2^2 \, d\nu(x) \right)^{1/2} \\ &\geq F_{\nu,\alpha}^\star + \frac{1}{4} \left(\int_{\mathbb{R}^d} \|x - \bar{x}\|_2^2 \, d\nu(x) \right)^{-1/2} \|\bar{a}_\alpha - \bar{x}\|_2^2 - 4k^{1/2}\alpha^{-1/2} \left(\int_{\mathbb{R}^d} \|x - \bar{x}\|_2^2 \, d\nu(x) \right)^{1/2}, \end{aligned}$$

from which one deduces that

$$\|\bar{a}_\alpha - \bar{x}\|_2 \leq 4k^{1/4}\alpha^{-1/4} \left(\int_{\mathbb{R}^d} \|x - \bar{x}\|_2^2 \, d\nu(x) \right)^{1/2}.$$

Consequently,

$$d_{2,\infty}(A_{\star,\alpha}, A_{\star,\infty}) \leq \|\bar{a}_\alpha - \bar{x}\|_2 + \text{diam}(A_{\star,\alpha}) \leq 8k^{1/2}\alpha^{-1/4} \left(\int_{\mathbb{R}^d} \|x - \bar{x}\|_2^2 d\nu(x) \right)^{1/2},$$

as desired. ■

5. Numerical implementation and experiments. In this section, we first discuss an algorithm and its numerical implementation for the archetypal analysis problem for a distribution with bounded support (1.1) and the modified method for measures with non-compact support (1.6). We then present the results of various numerical experiments that support and complement our analysis.

5.1. Numerical methods and their implementation.

5.1.1. Numerical method and implementation for (1.1). The archetypal analysis problem (1.1) can be equivalently rewritten as

$$\begin{aligned} (5.1a) \quad & \min_{\mathcal{A} \in \mathbb{R}^{N \times k}, \mathcal{B} \in \mathbb{R}^{k \times N}} \frac{1}{N} \|X - X\mathcal{A}\mathcal{B}\|_F^2 \\ (5.1b) \quad & \text{s.t., } \mathcal{A}_{ij}, \mathcal{B}_{ij} \geq 0 \\ (5.1c) \quad & \mathcal{A}^t \mathbf{1} = 1, \quad \mathcal{B}^t \mathbf{1} = 1 \end{aligned}$$

where $X = [x_1 \mid \cdots \mid x_N] \in \mathbb{R}^{d \times N}$, and $\mathbf{1}$ denotes vector of the all ones of compatible size. Furthermore, we denote $\mathcal{Z} := X\mathcal{A} = [a_1 \mid \cdots \mid a_k] \in \mathbb{R}^{d \times k}$.

Motivated by [9, 23], we use an alternating algorithm to update $\mathcal{Z}$ (i.e., update $\mathcal{A}$) and $\mathcal{B}$, respectively. That is, starting from some initial $\mathcal{Z}^{(0)}$, we find a sequence of solutions,

$$\mathcal{B}^{(0)}, \mathcal{Z}^{(1)}, \mathcal{B}^{(1)}, \dots, \mathcal{B}^{(p)}, \mathcal{Z}^{(p+1)}, \dots,$$

where

$$\begin{aligned} (5.2) \quad & \mathcal{B}^{(p)} = \arg \min_{\mathcal{B}} \frac{1}{N} \|X - \mathcal{Z}^{(p)}\mathcal{B}\|_F^2 \\ & \text{s.t., } \mathcal{B}_{ij} \geq 0 \text{ and } \mathcal{B}^t \mathbf{1} = 1 \end{aligned}$$

and

$$\begin{aligned} (5.3) \quad & \mathcal{Z}^{(p+1)} = \arg \min_{\mathcal{Z}} \frac{1}{N} \|X - \mathcal{Z}\mathcal{B}^{(p)}\|_F^2 \\ & \text{s.t., } \mathcal{A}_{ij} \geq 0, \quad \mathcal{A}^t \mathbf{1} = 1, \text{ and } \mathcal{Z} = X\mathcal{A}. \end{aligned}$$

We first use the Gauss–Seidel strategy to solve (5.3) by updating the columns of $\mathcal{Z}$ (i.e., a_i , $i \in [k]$) sequentially. For $\ell \in [k]$, the ℓ -th column of $\mathcal{Z}^{(p)}$ is updated by

$$\begin{aligned} (5.4) \quad & a_\ell^{(p+1)} = \arg \min_{a_\ell} \frac{1}{N} \left\| \frac{\sum_{i=1}^N \mathcal{B}_{\ell i}^{(p)} \xi_i}{\sum_{i=1}^N \mathcal{B}_{\ell i}^{(p)}} - a_\ell \right\|_2^2 \\ & \text{s.t., } \mathcal{A}_{ij} \geq 0, \quad \mathcal{A}^t \mathbf{1} = 1, \text{ and } a_\ell = \{X\mathcal{A}\}_\ell \end{aligned}$$

Algorithm 5.1 The projected gradient descent method to approximately solve (5.6).

Input: Let ω^0 be the initialization, $\tau > 0$

Output: ω^n that approximately solves (5.6).

Set $s = 1$

While *not converged* **do**

1. Gradient Descent. Solve the initial value problem until time τ with initial value given by ω^{s-1} :

$$(5.7) \quad \begin{cases} \frac{d\omega}{dt} = -C^T C \omega + C^T u \\ \omega(0) = \omega^{s-1}. \end{cases}$$

Let $\tilde{\omega} = \omega(\tau)$.

2. Projection Step. Project $\tilde{\omega}$ to $\mathcal{S}$ to obtain ω^s .

Set $s = s + 1$

where $\{\cdot\}_\ell$ denotes the ℓ -th column, and

$$(5.5) \quad \xi_i = x_i - \sum_{s \neq \ell}^k \mathcal{B}_{si}^{(p)} a_s \quad \forall i \in [N].$$

The details of the derivation of (5.4) is given in [Appendix B](#). It's easy to see that (5.2) and (5.4) are summarized by the following general convex least squares problem:

$$(5.6) \quad \min_{\omega \in \mathcal{S}} \|u - C\omega\|_2^2$$

where

$$\mathcal{S} = \{\omega = (\omega_1, \omega_2, \dots, \omega_q) : \omega_i \geq 0 \text{ and } \sum_{i=1}^q \omega_i = 1\}$$

is the unit simplex and $u \in \mathbb{R}^n$ and $C = [t_1 \mid t_2 \mid \dots \mid t_q] \in \mathbb{R}^{n \times q}$ are given. (5.6) is a standard convex programming problem which can be effectively solved by the projected gradient descent method, as summarized in [Algorithm 5.1](#).

When n and q are relatively small, the exact solution of step 1 in [Algorithm 5.1](#) can be efficiently computed as follows. Assume that $C^T C$ has eigenvalue decomposition $V \Sigma V^T$ with eigenvalues σ_i , $i \in [q]$. The solution to (5.7) is

$$(5.8) \quad \omega(t) = V P$$

where $P \in \mathbb{R}^{q \times 1}$ with entries

$$(5.9) \quad P_i = \begin{cases} e^{-\sigma_i t} P_i^0 - \frac{1}{\sigma_i} e^{-\sigma_i t} (V^T C^T u)_i + \frac{1}{\sigma_i} (V^T C^T u)_i, & \text{if } \sigma_i \neq 0, \\ P_i^0 + t (V^T C^T u)_i, & \text{if } \sigma_i = 0 \end{cases}$$

and $P^0 = V^T \omega(0)$. For the high-dimensional and large data set, one can simply use the forward Euler method to approximate the solution [23].

Algorithm 5.2 Algorithm for the projection onto the simplex [10].

Input: A vector $\tilde{\omega} \in \mathbb{R}^n$

Output: ω solves (5.10).

Step 1. Sort $\tilde{\omega}$ into μ : $\mu_1 \geq \mu_2 \geq \dots \geq \mu_n$;

Step 2. Find

$$\rho = \max \left\{ j \in [n] : \mu_j - \frac{1}{j} \left(\sum_{r=1}^j \mu_r - 1 \right) > 0 \right\};$$

Step 3. Define

$$\theta = \frac{1}{\rho} \left(\sum_{i=1}^{\rho} \mu_i - 1 \right);$$

Step 4. Set

$$\omega_i = \max\{\tilde{\omega}_i - \theta, 0\}.$$

Step 2 of Algorithm 5.1 can be efficiently solved by writing it as the optimization problem

$$(5.10a) \quad \min_{\omega} \frac{1}{2} \|\omega - \tilde{\omega}\|_2^2$$

$$(5.10b) \quad \text{s.t. } \sum_{i=1}^n w_i = 1, \quad w_i \geq 0$$

and using the projection algorithm proposed in [10], which we summarize in Algorithm 5.2.

5.1.2. Numerical method and implementation for (1.6). Similarly to subsection 5.1.1, we can equivalently write the variance-regularized archetypal analysis problem (1.6) in the following form:

$$(5.11a) \quad \min_{\mathcal{A} \in \mathbb{R}^{N \times k}, \mathcal{B} \in \mathbb{R}^{k \times N}} \frac{1}{N} \|X - X\mathcal{A}\mathcal{B}\|_F^2 + \frac{\alpha}{k} \sum_{i=1}^k \left\| a_i - \frac{1}{k} \sum_{j=1}^k a_j \right\|_2^2$$

$$(5.11b) \quad \text{s.t., } \mathcal{A}_{ij}, \mathcal{B}_{ij} \geq 0$$

$$(5.11c) \quad \mathcal{A}^t \mathbf{1} = \mathbf{1}, \quad \mathcal{B}^t \mathbf{1} = \mathbf{1}$$

where $\alpha > 0$ is a parameter. Again, $X = [x_1 \mid \dots \mid x_k] \in \mathbb{R}^{d \times N}$, $\mathbf{1}$ denotes the all 1 vector of compatible size, and $\mathcal{Z} := X\mathcal{A} = [a_1 \mid \dots \mid a_k] \in \mathbb{R}^{d \times k}$.

Again, we use an alternating algorithm to update $\mathcal{Z}$ (i.e., update $\mathcal{A}$) and $\mathcal{B}$. That is, starting from some initial $\mathcal{Z}^{(0)}$, we find a sequence of solutions, $\mathcal{B}^{(0)}, \mathcal{Z}^{(1)}, \mathcal{B}^{(1)}, \dots, \mathcal{B}^{(p)}$,

$\mathcal{Z}^{(p+1)}, \dots$, where

$$(5.12) \quad \begin{aligned} \mathcal{B}^{(p)} = \arg \min_{\mathcal{B}} \quad & \frac{1}{N} \|X - \mathcal{Z}^{(p)} \mathcal{B}\|_F^2 + \frac{\alpha}{k} \sum_{i=1}^k \left\| a_i^{(p)} - \frac{1}{k} \sum_{j=1}^k a_j^{(p)} \right\|_2^2 \\ \text{s.t., } & \mathcal{B}_{ij} \geq 0 \text{ and } \mathcal{B}^t \mathbf{1} = 1 \end{aligned}$$

and

$$(5.13) \quad \begin{aligned} \mathcal{Z}^{(p+1)} = \arg \min_{\mathcal{Z}} \quad & \frac{1}{N} \|X - \mathcal{Z} \mathcal{B}^{(p)}\|_F^2 + \frac{\alpha}{k} \sum_{i=1}^k \left\| a_i^{(p)} - \frac{1}{k} \sum_{j=1}^k a_j^{(p)} \right\|_2^2 \\ \text{s.t., } & \mathcal{A}_{ij} \geq 0, \mathcal{A}^t \mathbf{1} = 1, \text{ and } \mathcal{Z} = X \mathcal{A}. \end{aligned}$$

We note that (5.12) is exactly same as (5.2) when $\mathcal{Z}^{(p)}$ is fixed because the additional term is independent of $\mathcal{B}$. As for (5.13), we continue to use the Gauss–Seidel strategy. For $\ell \in [k]$, the ℓ -th column of $\mathcal{Z}^{(p)}$ is updated by

$$(5.14) \quad \begin{aligned} a_{\ell}^{(p+1)} = \arg \min_{a_{\ell}} \quad & \left\| \frac{\frac{1}{N} \sum_{i=1}^N \mathcal{B}_{\ell i}^{(p)} \xi_i + \frac{\alpha}{k^2} \sum_{s \neq \ell}^k a_s}{\frac{1}{N} \sum_{i=1}^N \mathcal{B}_{\ell i}^{(p)^2} + \frac{\alpha(k-1)}{k^2}} - a_{\ell} \right\|_2^2 \\ \text{s.t., } & \mathcal{A}_{ij} \geq 0, \mathcal{A}^t \mathbf{1} = 1, \text{ and } a_{\ell} = \{X \mathcal{A}\}_{\ell} \end{aligned}$$

where $\{\cdot\}_{\ell}$ denotes the ℓ -th column and ξ_i is the same as (5.5). The details of this derivation are given in [Appendix B](#). Again, both problems (*i.e.*, (5.12) and (5.13)) can be reduced to solving (5.6) which can be efficiently solved via [Algorithms 5.1](#) and [5.2](#).

Remark 5.1. In (5.8), as we mentioned, the exact solution is only possible when n and q are relatively small. It would be very expensive for high-dimensional and large data. However, if $q \gg n$ when n is relatively small, to accelerate this procedure, we choose a convenient representation based on Carathéodory’s theorem, which states that every point in a compact convex set in $\mathbb{R}^n$ can be written as a convex combination of $n+1$ extreme points. This suffices to set X in (5.4) as extreme points of $\partial(\text{co}(X_N))$, and the resulting q in the computation is dramatically reduced.

5.2. Numerical experiments. In this section, we illustrate the performance of the algorithm and verify the above consistency results via several computational experiments. We implemented the algorithms in MATLAB. All reported results were obtained on a laptop with a 2.7GHz Intel Core i5 processor and 8GB of RAM. In all experiments, we set $\tau = 0.5$. We choose the stopping criteria for the iteration to be $\|\mathcal{Z}^{(p+1)} - \mathcal{Z}^{(p)}\|_F^2 < \text{tol}$ and, in all experiments, set $\text{tol} = 1 \times 10^{-6}$.

5.2.1. Example 1: Uniform distribution in a disk. We first consider the case where the distribution is the uniform distribution in a unit disc. That is, each data point (x, y) is randomly generated by

$$\begin{cases} x = \sqrt{r} \cos(2\pi\theta) \\ y = \sqrt{r} \sin(2\pi\theta) \end{cases}$$

where r and θ are both random numbers generated from the uniform distribution on $[0, 1]$ (i.e., $U([0, 1])$).

In this experiment, we study, when k is fixed, how the convex hull of archetype points, $\text{co}(\{a_\ell\}_{\ell \in [k]})$, changes as the number of data points, N , increases. From the consistency result in [Theorem 1.5](#) and the analytical result from [section 3](#) for this problem, we know that the optimal polytope should converge to the regular one with unit-length vertices. To measure this convergence, we report the maximum and minimum values of the angles at extreme points. The reported results were obtained by repeating the same experiment 100 times and taking their mean for each value of N . [Figure 3](#) shows the change of angles as N increases from $N = 4$ to 13 with $\Delta N = 1$, from $N = 13$ to 333 with $\Delta N = 10$, and from $N = 333$ to 30,033 with $\Delta N = 300$, where ΔN is the step size. We observe that, as N increases, the maximum angle and the minimum angle converge to the same value, indicating that $\text{co}(\{a_\ell\}_{\ell \in [k]})$ converges to an equilateral triangle. The right figure in [Figure 3](#) is the plot of a minimizer for 30,033 random data points. The red curve is the boundary of the convex hull of the archetype points and the magenta curve is the boundary of the convex hull of random data points.




Figure 3. (left) The maximum and minimum angles of the convex polytope as N increases from 4 to 30,033. This result is obtained by averaging over 100 numerical experiments. (right) A minimizer for 30,033 random data points. The red curve is the boundary of the convex hull of the archetype points and the magenta curve is the boundary of the convex hull of random data points. See [subsection 5.2.1](#).

5.2.2. Example 2: Normal distribution. In this example, we consider the case where the support of the distribution is non-compact. In this case, to obtain a solution with bounded support in the limit as $N \rightarrow \infty$, a penalty term is added to the objective functional as in [\(1.6\)](#).

In this experiment, for a fixed $k = 3$, we study the behavior of the solution as α increases. We first generate $N = 30,000$ random points from a standard normal distribution $\mathcal{N}(0, 10 \cdot I)$ where I is the identity matrix. In [Figure 4](#), we plot the initialization and the solution to the archetypal analysis problem for $\alpha = 0$. We observe that all archetype points are located on the boundary of the convex hull of the data set, as expected by [Theorem 1.2](#). Here, we note that, when $\alpha = 0$, as $N \rightarrow \infty$, the convex polytope should grow to infinity in all directions. However, for a fixed N , we have a bounded data set and thus observe similar behavior to that in [subsection 5.2.1](#). In this case with 30,000 random data points, the area of the final $\text{co}(A)$ is

233. When α is slightly increased to 0.1, the area of $\text{co}(A)$ decreases to about 39. We further study the dependence of the area of $\text{co}(A)$ on α in the next experiment.

In this experiment, we gradually increase the value of α and find the corresponding archetype points, A , by solving (1.6). In Figure 5, we plot the area of $\text{co}(A)$ for various values of α . Here, we let α vary from 0.1 to 5 with step size 0.1. We also plot the convex polytope, $\text{co}(A)$, for some values of α . We observe that $\text{co}(A)$ shrinks as α increases, which is consistent with intuition. Also, for this isotropic distribution, we observe that the solution is always close to a regular triangle. In each subfigure in Figure 5, we indicate the mean of the distribution $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$ with a black point for comparison with Theorem 1.14. In all following numerical experiments, *i.e.*, initializations and data samples X_N , we observe that the mean of the distribution is contained in $\text{co}(A)$.

We expect the cost function of a minimizer $A_\star$ to be balanced in the sense that $F_\nu^2(A_\star)$ and $\alpha V(A_\star)$ are on the same order, *i.e.*,

$$F_\nu^2(A_\star) \sim \alpha V(A_\star) \sim (F_{\nu,\alpha}^\star)^2.$$

Since we also have that $V(A_\star) \sim \text{diam}(A_\star)^2$, where $\text{diam}(A)$ denotes the diameter of $\text{co}(A)$, we obtain

$$\text{diam}(A_\star) \sim \frac{F_\nu^2(A_\star)}{\alpha}.$$

The scaling in Figure 5 is consistent with this argument.




Figure 4. (left) $N = 30,000$ random points sampled from a normal distribution. The red curve represents the initialization of the archetype polygon. (right) The optimal solution of A when $\alpha = 0$. See subsection 5.2.2.

5.2.3. Example 3: Annular distribution. In this experiment, we consider the case when the distribution is an annular distribution where N random points (x_i, y_i) , $i \in [N]$ are generated by

$$(5.15) \quad \begin{cases} x_i = \sqrt{|r_i|} \cos(2\pi\theta_i) \\ y_i = \sqrt{|r_i|} \sin(2\pi\theta_i). \end{cases}$$




Figure 5. The area of $\text{co}(A)$ as we vary α with $N = 30,000$ random data points generated from a normal distribution. See subsection 5.2.2.

Here θ_i is a random number generated from the uniform distribution in $[0, 1]$ (i.e., $U([0, 1])$) and r_i is a random number generated from the normal distribution $\mathcal{N}(25, 50)$ (See Figure 6 for an example with $N = 30,000$). Note that the random points are concentrated around the circle with radius 5.




Figure 6. (left) $N = 30,000$ random points generated from an annular distribution as in (5.15). The boundary of the convex hull is indicated. (right) The optimal solution of (1.7) with $\alpha = 0$. See subsection 5.2.3.

In Figure 7, we plot the area of $\text{co}(A)$ as we vary α . We observe that even for random points generated from this annular distribution, as $\alpha \rightarrow 0$, $\text{co}(A)$ shrinks to a regular triangle containing the mean of the distribution.

To study how the algorithm depends on the initialization, we fix $\alpha = 2$ and choose several different initializations for $\{a_\ell\}_{\ell \in [3]}$. Figure 8 displays selected snapshots of the iterations for different initializations. The final $\text{co}(A)$ is more or less independent of the initialization, which indicates the robustness of the algorithm.




Figure 7. The area of $\text{co}(A)$ as we vary α for $N = 30,000$ random points generated from an annular distribution. See [subsection 5.2.3](#).




Figure 8. Snapshots of the iterates for different initializations. Last column is the final iterate. See [subsection 5.2.3](#).

5.2.4. Example 4: A Gaussian mixture model. In this experiment, we consider an asymmetric distribution where $N = 30,000$ random data points are generated from the mixed Gaussian model with two normal distributions $\mathcal{N}(\mu_1, \Sigma_1)$ and $\mathcal{N}(\mu_2, \Sigma_2)$ with $\mu_1 = \begin{pmatrix} 8 \\ 8 \end{pmatrix}$, $\mu_2 = \begin{pmatrix} -8 \\ -8 \end{pmatrix}$, $\Sigma_1 = 9 \cdot I$, and varying Σ_2 . We note that the mean of this bimodal distribution is $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$. In this experiment, $\alpha = 2$. In the first two rows of [Figure 9](#), we display snapshots of iterations for the data randomly generated from the Gaussian mixture model with $\Sigma_2 = 4 \cdot I$ for two different initializations. We observe that the iterates converge to a triangle containing the data mean in just a few iterations. In the last two rows of [Figure 9](#), we change Σ_2 to

$1 \cdot I$ and $0.25 \cdot I$ to increase the “anisotropy”. After about 20 iterations, the solution also converges to a triangle containing the mean $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$. In this experiment, we note that, because the “anisotropy” of the random data, $\text{co}(A)$ is not a regular triangle, but contains the mean. This numerically supports the claim [Theorem 1.14](#) that penalization makes the archetype points tend towards the mean.




Figure 9. Snapshots of the iterations for different initializations and different sampled data from a Gaussian mixture model. The last column is the final iterate. See [subsection 5.2.4](#).

6. Discussion. In this paper, we studied the archetypal analysis problem (1.1). We proved the consistency result that shows if the data are independently sampled from a probability measure with bounded support, then the archetype points converge to the solution of the continuum version of the problem, (1.2); see [Theorem 1.5](#). If the points are independently sampled from a distribution with unbounded support, we also prove a consistency result for a modified method (1.7) that penalizes the dispersion of the archetype points; see [Theorem 1.12](#). Our analysis is supported by detailed computational experiments of the archetype points for data sampled from the uniform distribution in a disk, the normal distribution, an annular distribution, and a bimodal distribution.

To handle distributions with unbounded support, we introduced a modified method (1.7) that penalizes the dispersion of the archetype points. There are several other potential modified methods that we could have used. (i) One choice would be the boundary of the convex set of the archetype points, $\partial(\text{co}(A))$ rather than $\text{co}(A)$ in the definition of $F_\nu(A)$, but this is a more difficult cost to compute. (ii) Interpreting extremal data as outliers, we could have followed the suggestion of [9] and used convex peeling to discard a small fraction of the data. This is equivalent to sampling from a conditional distribution. Finally, we could have changed

the measure of distance in (1.2) to, for example, use the 2-Wasserstein metric W_2 and consider the problem of minimizing $W_2(\mu, \xi)$, where $\xi = \frac{1}{\text{vol}(\text{co}(A))} \chi_{\text{co}(A)}$. We intend to pursue this direction in subsequent work.

In this work, we used a projected gradient descent method to approximately solve (5.6); see Algorithm 5.1. For a more scalable implementation, Algorithm 5.1 could be combined with sampling techniques to accelerate the convergence of the iterations. A subsampling approach would be to apply Algorithm 5.1 to a relatively small subsample of the dataset to obtain an approximate solution and then iteratively use this as an initial guess for increasingly larger (nested) subsamples of the dataset. An adaptive sampling technique for distributions with compact support would involve ‘ignoring’ samples of the dataset that lie far from $\partial(\text{co}(\text{supp}(\mu)))$ and ‘focusing’ on samples that lie near $\partial(\text{co}(\text{supp}(\mu)))$.

Appendix A. Proof of the claim in Remark 2.4. We prove that for every $\mu \in \mathcal{V}$ (which is defined in (2.16)), $\text{supp}(\mu)$ is affinely independent. First note that existence of μ is guaranteed by Prokhorov’s theorem. For $a \in \text{co}(D)$, define $\mathcal{M}_a = \{\nu \in \mathcal{M} : \mathbb{E}_\nu[x] = a\}$, so that

$$\mu \in \arg \max_{\nu \in \mathcal{M}_{\mathbb{E}_\mu[x]}} \mathbb{E}_\nu[\|x - \mathbb{E}_\mu[x]\|_2^2] = \arg \max_{\nu \in \mathcal{M}_{\mathbb{E}_\mu[x]}} \mathbb{E}_\nu[\|x\|_2^2].$$

Note that $\mathcal{M}_{\mathbb{E}_\mu[x]}$ is a convex compact set, and $l(\nu) := \mathbb{E}_\nu[\|x\|_2^2]$ is a non-zero bounded linear functional on $\mathcal{M}_{\mathbb{E}_\mu[x]}$. The maximum of l must be obtained on the extreme set of $\mathcal{M}_{\mathbb{E}_\mu[x]}$. A result in [18, Theorem 2.1] implies that the extreme set of $\mathcal{M}_{\mathbb{E}_\mu[x]}$ consists of probability measures in $\mathcal{M}$ whose support is affinely independent, particularly, $\text{supp}(\mu) \leq d + 1$.

Appendix B. Derivation of problem (5.4) and (5.14). In the follows, we include the derivation of problem (5.14). (5.4) can be reduced from (5.14) by setting $\alpha = 0$. From (5.13), assume we are updating the ℓ -th column of $\mathcal{Z}$ (i.e., a_ℓ) with other columns being fixed, direct calculation yields

$$\begin{aligned} & \frac{1}{N} \|X - \mathcal{Z}\mathcal{B}\|_F^2 + \frac{\alpha}{k} \sum_{q=1}^k \left\| a_q - \frac{1}{k} \sum_{s=1}^k a_s \right\|_2^2 \\ &= \frac{1}{N} \sum_{j=1}^d \sum_{i=1}^N \left(x_{ji}^2 - 2x_{ji} \sum_{s=1}^k \mathcal{Z}_{js} \mathcal{B}_{si} + \left(\sum_{s=1}^k \mathcal{Z}_{js} \mathcal{B}_{si} \right)^2 \right) + \frac{\alpha}{k} \sum_{q=1}^k \left\| a_q - \frac{1}{k} \sum_{s=1}^k a_s \right\|_2^2 \\ &= \frac{1}{N} \sum_{j=1}^d \sum_{i=1}^N \left(-2x_{ji} \mathcal{Z}_{j\ell} \mathcal{B}_{\ell i} + (\mathcal{Z}_{j\ell} \mathcal{B}_{\ell i})^2 + 2\mathcal{Z}_{j\ell} \mathcal{B}_{\ell i} \sum_{s \neq \ell}^k \mathcal{Z}_{js} \mathcal{B}_{si} \right) + \frac{\alpha(k-1)}{k^2} \|a_\ell\|_2^2 \\ &\quad - 2\frac{\alpha}{k} \left\langle a_\ell, \frac{k-1}{k^2} \sum_{s \neq \ell}^k a_s + \frac{1}{k} \sum_{q \neq \ell}^k \left(a_q - \frac{1}{k} \sum_{s \neq \ell}^k a_s \right) \right\rangle + \mathcal{O} \\ &= \left(\frac{1}{N} \sum_{i=1}^N \mathcal{B}_{\ell i}^2 + \frac{\alpha(k-1)}{k^2} \right) \|a_\ell\|_2^2 - 2 \left\langle a_\ell, \frac{1}{N} \sum_{i=1}^N \left[\mathcal{B}_{\ell i} \left(x_i - \sum_{s \neq \ell}^k a_s \mathcal{B}_{si} \right) \right] + \frac{\alpha}{k^2} \sum_{s \neq \ell}^k a_s \right\rangle + \mathcal{O} \end{aligned}$$

where $\mathcal{O}$ contains all terms independent of a_ℓ . Completing the squares for a_ℓ gives the desired formula.

Acknowledgments. We would like to thank the anonymous referees for their very helpful comments and especially for pointing us towards [4], which we used in the proof of [Theorem 1.8](#).

REFERENCES

- [1] C. BAUCKHAGE AND C. THURAU, *Making archetypal analysis practical*, in Lecture Notes in Computer Science, Springer Berlin Heidelberg, 2009, pp. 272–281, https://doi.org/10.1007/978-3-642-03798-6_28.
- [2] M. BELKIN AND P. NIYOGI, *Towards a theoretical foundation for Laplacian-based manifold methods*, in International Conference on Computational Learning Theory, Springer, 2005, pp. 486–500, https://doi.org/10.1007/11503415_33.
- [3] O. BOUSQUET, O. CHAPPELLE, AND M. HEIN, *Measure based regularization*, in Advances in Neural Information Processing Systems, 2004, pp. 1221–1228.
- [4] V.-E. BRUNEL, *Uniform behaviors of random polytopes under the Hausdorff metric*, Bernoulli, 25 (2019), pp. 1770–1793, <https://doi.org/10.3150/18-bej1035>.
- [5] J. CALDER AND N. GARCÍA TRILLOS, *Improved spectral convergence rates for graph Laplacians on ε -graphs and k -NN graphs*, arXiv:1910.13476, (2019).
- [6] E. CANHASI AND I. KONONENKO, *Weighted archetypal analysis of the multi-element graph for query-focused multi-document summarization*, Expert Systems with Applications, 41 (2014), pp. 535–543, <https://doi.org/10.1016/j.eswa.2013.07.079>.
- [7] B. H. P. CHAN, D. A. MITCHELL, AND L. E. CRAM, *Archetypal analysis of galaxy spectra*, Monthly Notices of the Royal Astronomical Society, 338 (2003), pp. 790–795, <https://doi.org/10.1046/j.1365-8711.2003.06099.x>.
- [8] Y. CHEN, J. MAIRAL, AND Z. HARCHAoui, *Fast and robust archetypal analysis for representation learning*, in 2014 IEEE Conference on Computer Vision and Pattern Recognition, IEEE, jun 2014, <https://doi.org/10.1109/cvpr.2014.192>.
- [9] A. CUTLER AND L. BREIMAN, *Archetypal analysis*, Technometrics, 36 (1994), p. 338, <https://doi.org/10.2307/1269949>.
- [10] J. DUCHI, S. SHALEV-SHWARTZ, Y. SINGER, AND T. CHANDRA, *Efficient projections onto the ℓ_1 -ball for learning in high dimensions*, in Proceedings of the 25th international conference on Machine learning - ICML 08, ACM Press, 2008, <https://doi.org/10.1145/1390156.1390191>.
- [11] L. DÜMBGEN AND G. WALTHER, *Rates of convergence for random approximations of convex sets*, Advances in Applied Probability, 28 (1996), pp. 384–393, <https://doi.org/10.2307/1428063>, <https://doi.org/10.2307/2F1428063>.
- [12] N. GARCÍA TRILLOS AND D. SLEPČEV, *A variational approach to the consistency of spectral clustering*, Applied and Computational Harmonic Analysis, 45 (2018), pp. 239–281, <https://doi.org/10.1016/j.acha.2016.09.003>.
- [13] N. GARCÍA TRILLOS, D. SLEPČEV, J. VON BRECHT, T. LAURENT, AND X. BRESSON, *Consistency of Cheeger and ratio graph cuts*, The Journal of Machine Learning Research, 17 (2016), pp. 6268–6313.
- [14] J. A. HARTIGAN, *Consistency of single linkage for high-density clusters*, Journal of the American Statistical Association, 76 (1981), pp. 388–394, <https://doi.org/10.1080/01621459.1981.10477658>.
- [15] M. HEIN, J.-Y. AUDIBERT, AND U. V. LUXBURG, *Graph Laplacians and their convergence on random neighborhood graphs*, Journal of Machine Learning Research, 8 (2007), pp. 1325–1368.
- [16] M. HEIN, J.-Y. AUDIBERT, AND U. VON LUXBURG, *From graphs to manifolds—weak and strong pointwise consistency of graph Laplacians*, in International Conference on Computational Learning Theory, Springer, 2005, pp. 470–485, https://doi.org/10.1007/11503415_32.
- [17] W. Hoeffding, *Probability inequalities for sums of bounded random variables*, in The Collected Works of Wassily Hoeffding, Springer, 1994, pp. 409–426.
- [18] A. F. KARR, *Extreme points of certain sets of probability measures, with applications*, Mathematics of Operations Research, 8 (1983), pp. 74–85, <https://doi.org/10.1287/moor.8.1.74>, <https://doi.org/10.1287/moor.8.1.74>.

- 1287%2Fmoor.8.1.74.
- [19] S. M. KELLER, M. SAMARIN, F. A. TORRES, M. WIESER, AND V. ROTH, *Learning extremal representations with deep archetypal analysis*, arXiv preprint arXiv:2002.00815, (2020).
 - [20] S. M. KELLER, M. SAMARIN, M. WIESER, AND V. ROTH, *Deep archetypal analysis*, in *Lecture Notes in Computer Science*, Springer International Publishing, 2019, pp. 171–185, https://doi.org/10.1007/978-3-030-33676-9_12.
 - [21] S. S. LAFON, *Diffusion maps and geometric harmonics*, PhD thesis, Yale University PhD dissertation, 2004.
 - [22] S. MAIR AND U. BREFELD, *Coresets for archetypal analysis*, in *Advances in Neural Information Processing Systems*, 2019, pp. 7245–7253.
 - [23] M. MØRUP AND L. K. HANSEN, *Archetypal analysis for machine learning and data mining*, *Neurocomputing*, 80 (2012), pp. 54–63, <https://doi.org/10.1016/j.neucom.2011.06.033>.
 - [24] B. OSTING AND T. H. REEB, *Consistency of Dirichlet partitions*, *SIAM Journal on Mathematical Analysis*, 49 (2017), pp. 4251–4274, <https://doi.org/10.1137/16M1098309>.
 - [25] D. POLLARD, *Strong consistency of k-means clustering*, *The Annals of Statistics*, 9 (1981), pp. 135–140, <https://doi.org/10.1214/aos/1176345339>.
 - [26] D. POLLARD ET AL., *A central limit theorem for k-means clustering*, *The Annals of Probability*, 10 (1982), pp. 919–926, <https://doi.org/10.1214/aop/1176993713>.
 - [27] S. SETH AND M. J. A. EUGSTER, *Probabilistic archetypal analysis*, *Machine Learning*, 102 (2015), pp. 85–113, <https://doi.org/10.1007/s10994-015-5498-8>.
 - [28] A. SINGER, *From graph to manifold Laplacian: The convergence rate*, *Applied and Computational Harmonic Analysis*, 21 (2006), pp. 128–134, <https://doi.org/10.1016/j.acha.2006.03.004>.
 - [29] E. STONE AND A. CUTLER, *Archetypal analysis of spatio-temporal dynamics*, *Physica D: Nonlinear Phenomena*, 90 (1996), pp. 209–224, [https://doi.org/10.1016/0167-2789\(95\)00244-8](https://doi.org/10.1016/0167-2789(95)00244-8).
 - [30] W. SUN, J. WANG, AND Y. FANG, *Regularized k-means clustering of high-dimensional data and its asymptotic consistency*, *Electronic Journal of Statistics*, 6 (2012), pp. 148–167, <https://doi.org/10.1214/12-ejs668>.
 - [31] D. VAN DIJK, D. B. BURKHARDT, M. AMODIO, A. TONG, G. WOLF, AND S. KRISHNASWAMY, *Finding archetypal spaces for data using neural networks*, *CoRR*, (2019).
 - [32] R. VERSHYNIN, *High-Dimensional Probability*, Cambridge University Press, sep 2018, <https://doi.org/10.1017/9781108231596>, <https://doi.org/10.1017%2F9781108231596>.
 - [33] U. VON LUXBURG, M. BELKIN, AND O. BOUSQUET, *Consistency of spectral clustering*, *The Annals of Statistics*, 36 (2008), pp. 555–586, <https://doi.org/10.1214/009053607000000640>.
 - [34] C. WU AND E. G. TABAK, *Prototypal analysis and prototypal regression*, arXiv preprint arXiv:1701.08916, (2017).
 - [35] A. YUAN, J. CALDER, AND B. OSTING, *A continuum limit for the pagerank algorithm*, arXiv:2001.08973, (2020).