

THE QUARTIC INTEGRABILITY AND LONG TIME EXISTENCE OF STEEP WATER WAVES IN 2D

SIJUE WU

ABSTRACT. It is known since the work of Dyachenko & Zakharov [54] that for the weakly nonlinear 2d infinite depth water waves, there are no 3-wave interactions and all of the 4-wave interaction coefficients vanish on the non-trivial resonant manifold. In this paper we study this partial integrability from a different point of view. We construct, directly in the physical space, a sequence of energy functionals $\mathfrak{E}_j(t)$ which are explicit in the Riemann mapping variable and involve material derivatives of order j of the solutions for the 2d water wave equation, so that $\frac{d}{dt}\mathfrak{E}_j(t)$ is quintic or higher order. We show that if some scaling invariant norm, and a norm involving one spatial derivative above the scaling of the initial data are of size no more than ε , then the lifespan of the solution for the 2d water wave equation is at least of order $O(\varepsilon^{-3})$, and the solution remains as regular as the initial data during this time. If only the scaling invariant norm of the data is of size ε , then the lifespan of the solution is at least of order $O(\varepsilon^{-5/2})$. Our long time existence results do not impose size restrictions on the slope of the initial interface and the magnitude of the initial velocity, they allow the interface to have arbitrary large steepnesses and initial velocities to have arbitrary large magnitudes.

1. INTRODUCTION

A class of water wave problems concerns the motion of the interface separating an inviscid, incompressible, irrotational fluid, under the influence of gravity, from a region of zero density (i.e. air) in n -dimensional space. It is assumed that the fluid region is below the air region. Assume that the density of the fluid is 1, the gravitational field is $-\mathbf{k}$, where \mathbf{k} is the unit vector pointing in the upward vertical direction, and at time $t \geq 0$, the free interface is $\partial\Omega(t)$, and the fluid occupies region $\Omega(t)$. When surface tension is zero, the motion of the fluid is described by

$$(1.1) \quad \begin{cases} \mathbf{v}_t + (\mathbf{v} \cdot \nabla)\mathbf{v} = -\mathbf{k} - \nabla P & \text{on } \Omega(t), t \geq 0, \\ \operatorname{div} \mathbf{v} = 0, \quad \operatorname{curl} \mathbf{v} = 0, & \text{on } \Omega(t), t \geq 0, \\ P = 0, & \text{on } \partial\Omega(t) \\ (1, \mathbf{v}) \text{ is tangent to the free surface } (t, \partial\Omega(t)), \end{cases}$$

where \mathbf{v} is the fluid velocity, P is the fluid pressure. There is an important condition for these problems:

$$(1.2) \quad -\frac{\partial P}{\partial \mathbf{n}} \geq 0$$

pointwise on the interface, where \mathbf{n} is the outward unit normal to the fluid interface $\partial\Omega(t)$ [43]; it is well known that when surface tension is neglected and the Taylor sign condition (1.2) fails, the water wave motion can be subject to the Taylor instability [43, 14, 10, 25].

The study of water waves dates back centuries to Newton [36], Stokes [42], Levi-Civita [33], and G.I. Taylor [43]. Nalimov [35], Yosihara [52] and Craig [21] gave early local in time existence and uniqueness results for the 2d water wave equation (1.1) for small and smooth initial data. In [46, 47], Wu showed that

Financial support in part by NSF grants DMS-1764112.

in dimensions $n \geq 2$, the strong Taylor sign condition

$$(1.3) \quad -\frac{\partial P}{\partial \mathbf{n}} \geq c_0 > 0$$

always holds for the infinite depth water wave problem (1.1),¹ and the initial value problem of equation (1.1) is locally well-posed in Sobolev spaces H^s , $s \geq 4$ for arbitrary given data. Since then, local wellposedness for water waves with additional effects such as the surface tension, bottom and non-zero vorticity, under the assumption (1.3),² were obtained, c.f. [9, 16, 17, 29, 32, 34, 37, 39, 57]; local wellposedness of (1.1) in low regularity Sobolev spaces, cf. [6, 7, 27, 1], and in a regime allowing for non- C^1 interfaces, cf. [31, 51, 3, 4, 5] were proved.

There are also numerous studies on the long time existence and regularity of solutions for the water wave equation (1.1) over the last ten years. Wu [48, 49], Germain, Masmoudi & Shatah [26], Ionescu & Pusateri [30] and Alazard & Delort [8] proved almost global and global existence of solutions for two and three dimensional water waves equation (1.1) for small, smooth and sufficiently localized data, further developments were given in [27, 28, 24, 44, 45, 13, 11, 40, 58, 1, 2]. There are two main ideas involved in these works. The first is to use a normal form transformation to remove all the quadratic nonlinearities in the equation (1.1)³; and the second is to use the dispersive decay properties of the spatially localized solutions for the equation (1.1) to obtain extended life spans of the solutions.

Despite these advances, many important questions remain open. For instance, in all the aforementioned works on the long time existence of water waves, norms involving derivatives both above and below the scaling are assumed small at the initial time; such smallness cannot be preserved after a rescaling. However physical phenomena do not disappear with a rescaling, a valid description of a physical phenomenon should only involve scaling invariant quantities and norms; also in all of these works, it is assumed that the steepness of the initial wave is sufficiently small. The questions are whether the smallness assumption on the steepness can be removed? and whether a long time existence result holds assuming smallness only on some scaling invariant quantities?

The study of the 2d water wave equation (1.1) in the Hamiltonian point of view began in [53], where Zakharov discovered that the 2d equation (1.1) can be written as a Hamiltonian system. In [54] Dyachenko & Zakharov showed that (formally) there are no three-wave resonant interactions in the Hamiltonian 2d periodic water wave equation (1.1) and all of the four-wave interaction coefficients vanish on the non-trivial resonant manifold.⁴ Dyachenko & Zakharov [54] and Craig & Wolfolk [22] derived a formal⁵ symplectic transformation that maps the Hamiltonian system of the 2d periodic water waves to its Birkhoff normal form of order 4; and mapping properties of the transformation were studied in [23]. Building on [54, 22, 23], Berti, Feola & Pusateri [12] gave a rigorous construction of a bounded and invertible (non-symplectic)

¹More precisely, the strong Taylor sign condition (1.3) holds for $C^{1+\gamma}$ interfaces, where $\gamma > 0$. For less regular interfaces, such as interfaces with angled crests, only the weak Taylor sign condition (1.2) holds, with degeneracies at the singularities, cf. [31, 51].

²When there is surface tension, or bottom, or vorticity, (1.3) does not always hold, it needs to be assumed. See [32], [40], [41] for some interesting computations of $-\frac{\partial P}{\partial \mathbf{n}}$ in the cases of finite depth and point vortices.

³For water waves with surface tension [24] or non-zero vorticity [40], only part of the quadratic nonlinearities can be removed.

⁴While there are no 3-wave resonant interactions, there are two types of 4-wave resonances: the trivial ones and the Benjamin & Feir resonances. What [54] proved was that all the coefficients of the 4th order terms in the power expansion of the Hamiltonian vanish on the Benjamin & Feir resonance manifold, cf. [12] for further explanations. In the whole line case, there are in addition near resonances.

⁵That is, it is unbounded and non-invertible.

transformation in a neighborhood of the origin in phase space, mapping the 2d periodic water wave equation (1.1) to its Birkhoff normal form up to order 4, and showed that for sufficiently small and smooth initial data of size ε ,⁶ the 2d periodic water wave equation (1.1) is solvable for time of order $O(\varepsilon^{-3})$. [12] is the first and the only work putting the formal computations in [54, 22] in a rigorous framework. The transformation in [12] is constructed via composing several paradifferential flow conjugations. The final resonant Poincaré-Birkhoff normal form system constructed in [12] is not a priori explicit; an important step in [12] is a normal form uniqueness argument that allows the authors to identify the Poincaré-Birkhoff normal form system of [12] with the Birkhoff normal form system constructed in [54, 22], up to degree 4 of homogeneity.

The computations in [54, 22, 23, 12] were all carried out in the Fourier space using tools such as the Birkhoff normal forms from the Hamiltonian system. As the water wave equation (1.1) is explicitly formulated in the physical space, it is natural to ask then how do the structural results on the 3-wave and 4-wave interactions in [54] manifest, explicitly, in the physical space? and what are the specific features in equation (1.1) that lead to these structural properties? The motivation is clear: by better understanding the structure of equation (1.1) we hope to obtain better information on the long time behavior of the water waves. Moreover, we hope to gain insight on the non-linear structure of a wider class of equations.

In this paper we construct, directly in the physical space, an infinite sequence of energy functionals $\mathfrak{E}_j(t)$ that involve material derivatives of order j of the solutions of the 2d water wave equation (1.1), so that $\frac{d}{dt}\mathfrak{E}_j(t)$ is quintic.⁷ Our energy functionals $\mathfrak{E}_j(t)$ and their time derivatives $\frac{d}{dt}\mathfrak{E}_j(t)$ are explicit in the Riemann mapping variable, and the process of constructing the sequence of our energy functionals $\mathfrak{E}_j(t)$, as well as $\mathfrak{E}_j(t)$, $\frac{d}{dt}\mathfrak{E}_j(t)$ themselves, reveal remarkable novel symmetric structures of the 2d water wave equation (1.1). Moreover for $j \geq 2$, $\mathfrak{E}_j(t)$ are energy functionals in terms of the derivatives of the velocity and the derivatives of the steepness, water waves with large velocity and large steepness can be small in $\mathfrak{E}_j(t)$; similarly, $\frac{d}{dt}\mathfrak{E}_j(t)$ are quintic in terms of only the sizes of the derivatives of the velocity and the derivatives of the steepness of the interface. As a consequence we show that if some scaling invariant norm and a norm involving one order spatial derivative above the scaling of the initial data are of size no more than ε , then the lifespan of the solution for the 2d water wave equation (1.1) is at least of order $O(\varepsilon^{-3})$; if only the scaling invariant norm of the initial data is of size ε , then the solution will exist up to time of order $O(\varepsilon^{-5/2})$.⁸ Our long time existence results do not impose size restrictions on the slope of the initial interface and the magnitude of the initial velocity, the initial interface can have arbitrary large steepness, and the magnitude of the initial velocity can be arbitrarily large. To the best of our knowledge, this is the first long time existence result assuming smallness only on a scaling invariant quantity at the initial time, and it is the first allowing steep initial interfaces.

We will give rigorous statements of our main results, in the whole line setting, in §2 and §3. The same construction and derivation as in this paper work and analogous results hold for the 2d periodic water wave equation (1.1). The only difference is that in the periodic case, there are some additional (spatially) constant terms arising from the means of the quantities involved, c.f. Proposition 1 and equation (290) of

⁶In [12], the steepness, and quantities both above and below the scaling are assumed small at the initial time.

⁷If not specified, the "quintic" in §1 means it is a finite sum of terms homogeneous of degree 5 or higher of the unknown functions and their derivatives; similar for "quartic".

⁸No dispersive properties of equation (1.1) are used.

[31]. However these constants are harmless, and results analogous to Theorem 2.8 and Theorem 3.1, as well as the identities in §2 leading to Theorem 2.8 hold for the periodic case.⁹

Just as this work continues the investigation initiated in [48, 49, 13, 51], we think that the results in §2 will themselves have further consequences. The next subsection gives a more detailed explanation of the main results, motivation for the method and relations to earlier work.

1.1. Further explanations on the motivation, the ideas and the main results in this paper. In this paper we regard \mathbb{R}^2 as a complex plane, and identify a point $(x, y) \in \mathbb{R}^2$ with its complex form $x + iy \in \mathbb{C}$. $\operatorname{Re} z, \operatorname{Im} z$ are the real and imaginary parts of z , $\bar{z} = \operatorname{Re} z - i \operatorname{Im} z$.

Let $z = x + iy = z(\alpha, t)$, $\alpha \in \mathbb{R}$ be the interface $\partial\Omega(t)$ in Lagrangian coordinate α , so $z_t(\alpha, t) = \mathbf{v}(z(\alpha, t), t)$ is the velocity of the fluid particle on the interface, and $z_{tt}(\alpha, t) = (\mathbf{v}_t + \mathbf{v} \cdot \nabla \mathbf{v})(z(\alpha, t), t)$ is the acceleration. We know (cf. [48]) the interface equation for the 2d water waves (1.1) is given by

$$(1.4) \quad \begin{cases} z_{tt} + i = i\mathbf{a}z_\alpha, \\ \bar{z}_t = \mathfrak{H}\bar{z}_t, \end{cases}$$

where $\mathbf{a} = -\frac{\partial P}{\partial \mathbf{n}} \frac{1}{|z_\alpha|}$, and \mathfrak{H} is the Hilbert transform associated to the fluid domain $\Omega(t)$; and the quasilinear equation, obtained by applying ∂_t to the first equation in (1.4), is

$$(1.5) \quad (\partial_t^2 + i\mathbf{a}\partial_\alpha)\bar{z}_t = \frac{\mathbf{a}_t}{\mathbf{a}}(\bar{z}_{tt} - i).$$

Let $\Phi : \Omega(t) \rightarrow \mathcal{P}_-$ be the Riemann mapping from the fluid domain $\Omega(t)$ to the lower half plane \mathcal{P}_- , satisfying $\lim_{z \rightarrow \infty} \Phi(z) = 1$ and $\Phi(z(0, t), t) = 0$; let $\mathbf{h}(\alpha, t) := \Phi(z(\alpha, t), t)$, $b(\alpha', t) := \mathbf{h}_t \circ \mathbf{h}^{-1}(\alpha', t)$. And let $Z(\alpha', t) = X(\alpha', t) + iY(\alpha', t) := z(\mathbf{h}^{-1}(\alpha', t), t)$, $Z_t(\alpha', t) := z_t(\mathbf{h}^{-1}(\alpha', t), t)$ and $Z_{tt}(\alpha', t) := z_{tt}(\mathbf{h}^{-1}(\alpha', t), t)$ be the position, velocity and acceleration of the interface in the Riemann mapping variable α' ; let $D_t := \partial_t + b\partial_{\alpha'}$ be the material derivative, i.e. $Z_t = D_t Z$ and $Z_{tt} = D_t Z_t$. Let

$$(1.6) \quad \mathfrak{P} := D_t^2 + i \frac{A_1}{|Z_{,\alpha'}|^2} \partial_{\alpha'},$$

and \mathbb{H} be the Hilbert transform associated to the lower half plane \mathcal{P}_- :

$$(1.7) \quad \mathbb{H}f(\alpha') = \frac{1}{\pi i} p.v. \int \frac{1}{\alpha' - \beta'} f(\beta') d\beta',$$

$\mathbb{P}_H = \frac{1}{2}(I + \mathbb{H})$, $\mathbb{P}_A = \frac{1}{2}(I - \mathbb{H})$ be the projections onto the space of holomorphic, and respectively, anti-holomorphic functions in \mathcal{P}_- . The quasilinear equation for the 2d water waves in the Riemann mapping variable α' is (cf [46, 51])

$$(1.8) \quad \mathfrak{P}\bar{Z}_t = \frac{\mathbf{a}_t}{\mathbf{a}} \circ \mathbf{h}^{-1}(\bar{Z}_{tt} - i),$$

where A_1 , b and $\frac{\mathbf{a}_t}{\mathbf{a}} \circ \mathbf{h}^{-1}$ are as given in (B.2) and (B.5). The main operator in (1.5) is $\partial_t^2 + i\mathbf{a}\partial_\alpha$ and in (1.8), \mathfrak{P} . It is clear that the quasilinear equation (1.5) or (1.8) contains quadratic nonlinear terms.

In [48] we found a pair of transformations which can be used to completely transform away the quadratic nonlinearities in (1.5). Let φ be the velocity potential, $\psi := \varphi(z(\alpha, t), t)$ be the trace of the velocity potential on the interface. Let

$$(1.9) \quad \Lambda = (I - \mathfrak{H})\psi, \quad \Pi = (I - \mathfrak{H})(z - \bar{z}); \quad \text{and} \quad \kappa = 2 \operatorname{Re} z - \mathbf{h}.$$

⁹In the periodic case, there are some extra (spatially) constant terms $c(t)$ in the corresponding quantities $\mathbb{P}_H G^{(j)}$, cf. §2 for the definition, however they do not affect the energy functional sequence, since they show up in the energy functionals as $\int_I \partial_{\alpha'} \Theta c(t) d\alpha'$. Since $\int_I \partial_{\alpha'} \Theta c(t) d\alpha' = 0$ for any periodic function Θ on I , these constants do not affect the outcome.

In [48] we showed that the quantities Λ and Π , after the change of coordinates κ , satisfy equations of the type

$$(1.10) \quad (\partial_t^2 - i\partial_{\tilde{\alpha}})(\Lambda \circ \kappa^{-1}) = G_1, \quad (\partial_t^2 - i\partial_{\tilde{\alpha}})(\Pi \circ \kappa^{-1}) = G_2,$$

where G_1 and G_2 consist of only cubic and higher order terms, $\tilde{\alpha} = \kappa(\alpha, t)$, c.f. Propositions 2.3 and 2.4 of [48]. Using equations (1.10) and applying the basic energy functional¹⁰

$$(1.12) \quad \tilde{E}_0(t) = \operatorname{Re} \int \left(\frac{1}{a} |\partial_t \theta|^2 + i\theta \partial_{\alpha} \bar{\theta} \right) d\alpha$$

on Λ , Π and their derivatives, we constructed an infinite sequence of energy functionals $\tilde{E}_j(t)$, so that

$$(1.13) \quad \frac{d}{dt} \tilde{E}_j(t) = \text{quartic};^{11}$$

we then proved our almost global existence result by employing the quartic energy identities (1.13)¹² and the method of vector fields, cf. [48]. Quartic energy estimates¹³ similar to (1.13) also played key roles in the subsequent works [30, 8, 27, 28]. While [30] directly improved upon the quartic energy estimates in [48], [8] obtained theirs using the Zakharov-Craig-Sulem formulation of the water wave equation and a normal form transformation via resonant analysis of the 3-wave interactions in the Fourier space. In [27, 28] the authors used

$$Q := (I + \mathbb{H})(\psi \circ h^{-1}), \quad \text{and} \quad W := \frac{1}{2}(I + \mathbb{H})(Z - \bar{Z}) = Z - \alpha',$$

which are the counter-parts¹⁴ of the quantities Λ and Π in (1.9) in the Riemann mapping coordinate, and a basic energy functional similar to (1.12);¹⁵ they first applied their basic energy functional to (W, Q) and their derivatives, and then directly modified the resulting energy functionals to obtain their quartic energy estimates. In all of [30, 8, 27, 28] dispersive properties of the water wave equation and modified scatterings were additionally used to obtain their global existence results.¹⁶

In what follows we call an energy functional $\tilde{E}_j(t)$ satisfying (1.13) a quartic energy functional. Similarly if $\frac{d}{dt} \mathfrak{E}_j(t)$ is quintic, we call $\mathfrak{E}_j(t)$ a quintic energy functional.

¹⁰cf. Lemma 4.1 of [48]. Upon changing to the Riemann mapping coordinate yields

$$(1.11) \quad \tilde{E}_0(t) = \operatorname{Re} \int \left(\frac{1}{a} |\partial_t \theta|^2 + i\theta \partial_{\alpha} \bar{\theta} \right) d\alpha = \operatorname{Re} \int \left(\frac{1}{A_1} |D_t \Theta|^2 + i\Theta \partial_{\alpha'} \bar{\Theta} \right) d\alpha',$$

where $\Theta = \theta \circ h^{-1}$. As was explained in [48], it is advantageous to replace the Dirichlet-to-Neumann operator ∇_n by $i\partial_{\alpha}$ when dealing with the cancellations in the quadratic nonlinearities of the 2d water wave equation, since ∇_n is nonlinear, while $i\partial_{\alpha}$ is linear.

¹¹The "quartic" in this paragraph means it is a finite sum of terms homogeneous of degree 4 or higher of the amplitude, the steepness and the velocity of the interface, as well as their derivatives.

¹²The quartic energy identities obtained by directly applying $E_0(t)$ to the derivatives of the basic quantities Λ and Π did not quite work, some further modifications were introduced to produce desired estimates, cf. [48].

¹³The basic mechanism behind these quartic energy estimates is the absence of three-wave interactions.

¹⁴Applying the basic energy functional (1.11) to $D_t^j Q$ yields

$$\tilde{E}_j(t) = \operatorname{Re} \int \left(\frac{1}{A_1} \left| D_t^{j+1} Q \right|^2 + i(D_t^j \bar{Q}) \partial_{\alpha'} (D_t^j Q) \right) d\alpha',$$

which satisfies the quartic energy identity (1.13) due to (1.10), and analogous result holds for $D_t^j W$. However in the Riemann mapping coordinate, not all derivatives of Q or W will lead to energy functionals satisfying (1.13). This is solely because of the change of the coordinates.

¹⁵In the basic energy functional of [27, 28] the first term is slightly different from the first term in (1.11) while the second term is the same as that in (1.11).

¹⁶The work [48, 30, 8, 27, 28] discussed here are all about small, smooth and sufficiently localized solutions, the smallness assumption is not scaling invariant, and the steepness of the interface is assumed small.

However none of the existing quartic energy functionals $\tilde{E}_j(t)$, including those in [48, 30, 8, 27], can be easily modified to turn them into quintic, since the remainders $\frac{d}{dt}\tilde{E}_j(t)$ are too complex to be used to identify further cancellations. Our strategy in constructing our quintic energy functional sequence $\mathfrak{E}_j(t)$ is to first re-understand and re-construct appropriate quartic energy functionals; we choose the Riemann mapping coordinate since it is advantageous in understanding the structure of the 2d water wave equation, cf. [46, 31, 50, 51]. We begin with the quantity Q (the counter-part of Λ in the Riemann mapping coordinates), and look for useful symmetries and structures hidden in the equations of the quantities involved and in the energy functionals. The construction of our new quartic energy functionals $E_j(t)$ involve three new interrelated ingredients: the sequence of quantities $\Theta^{(j)} := (\mathbb{P}_H D_t)^j Q$, their associated equations

$$(1.14) \quad D_t \mathbb{P}_H D_t \Theta^{(j)} + i \frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \Theta^{(j)} := G^{(j)},$$

cf. (2.7), (2.8), and the symmetric basic energy functional, cf. (2.14),

$$(1.15) \quad E(t) = \operatorname{Re} \left(\int i \partial_{\alpha'} \Theta_2 \overline{D_t \Theta_1} d\alpha' - \int i \partial_{\alpha'} \Theta_1 \overline{D_t \Theta_2} d\alpha' \right).$$

By Proposition 2.2, $\mathbb{P}_H G^{(j)}$ is cubic and possesses remarkable symmetries. Applying the basic energy $E(t)$ to $\Theta_1 = \Theta^{(j)}$ and $\Theta_2 = \Theta^{(j+1)}$, and using (2.15) we arrive at the new quartic energy identity (2.19)-(2.20)-(2.21).¹⁷ What is important is that (2.20)-(2.21) possesses desirable symmetric structure that allows us, with further use of the new identities (2.28)-(2.29), to systematically construct the quartic correcting functionals $C_{1,j}$ and $C_{2,j}$ in (2.33), (2.34) and obtain our quintic energy functionals $\mathfrak{E}_j(t)$, cf. (2.38). The main result of §2 is stated in Theorem 2.8.

The main difficulty in proving our second main theorem, Theorem 3.1, is that the quantity $\left\| \frac{1}{Z_{,\alpha'}} - 1 \right\|_{L^\infty}$ is not small, and no norms below the scaling are assumed small. This requires us to explore further cancellations in the quintic energy identities in Theorem 2.8 and employ stronger inequalities, including those in Propositions C.17 and C.14. The details of the proof for Theorem 3.1 can be found in §4.

Finally we mention that the transformations in [48], cf. (1.9), are used in [40, 15] to prove the recent important results on the long time behavior of point vortices and the nonlinear modulation instability of the Stokes waves in 2d water waves. Analogous transformations as those in (1.9) were also constructed to remove the quadratic nonlinearities in the 3d water wave equation [49] for perturbations near the flat equilibrium and in the free boundary problem of a self-gravitating incompressible fluid [13] for perturbations near the circular equilibrium.

1.2. Outline of the paper. We present our results on the algebraic aspects of the water wave equation, namely the quartic integrability, and their proofs in §2. In §3 we state our long time existence theorem, its proof is given in §4. A complete list of notations and conventions can be found in Appendix A. Some of the basic equations and formulas derived in our earlier works [46, 48, 51], as well as some additional identities that are used in the derivations in this paper are collected in Appendix B. Appendix C contains the inequalities that are used for our proofs. In Appendix D we summarize the estimates obtained in §4.3.1 and §4.3.2 for easy referencing.

¹⁷The "quartic" here is in terms of only the derivatives of the steepness of the interface and the derivatives of the velocity; similar for the "quintic" energies $\mathfrak{E}_j(t)$. Observe that the construction of $E_j(t)$ and $\mathfrak{E}_j(t)$ is based on a sequence of equations, instead of repeated derivatives to a single transformed equation.

1.3. Conventions. We consider solutions of 2d the water wave equation (1.1) in the setting where the fluid domain $\Omega(t)$ is simply connected in \mathbb{R}^2 , with the free interface $\partial\Omega(t) := \partial\Omega(t)$ being a non-self-intersecting curve,

$$\mathbf{v}(z, t) \rightarrow 0, \quad \text{as } |z| \rightarrow \infty$$

and the interface $\partial\Omega(t)$ tending to horizontal lines at infinity. We will primarily use the Riemann mapping variable in this paper.

1.4. Acknowledgement. The author would like to thank Jeffrey Rauch for carefully going through the first draft of the paper and for his helpful suggestions.

2. THE MAIN RESULTS ON THE STRUCTURE OF THE WATER WAVE EQUATION

In this section we construct a sequence of energy functionals $\mathfrak{E}_j(t)$ which involve material derivatives of order j of the solutions, such that $\frac{d}{dt}\mathfrak{E}_j(t)$ is quintic.¹⁸ The construction is based on a series of observations, given in Lemma 2.1 through Proposition 2.6, and equations (2.20)-(2.21), (2.31)-(2.32).

We continue with the notations introduced in §1, further notations can be found in Appendix A.

As derived in earlier work, cf. [46, 50] or §2.2 of [51], we have

$$(2.1) \quad b = \operatorname{Re}(I - \mathbb{H}) \left(\frac{Z_t}{Z_{,\alpha'}} \right) = \mathbb{P}_H \left(\frac{\overline{Z}_t}{\overline{Z}_{,\alpha'}} \right) + \mathbb{P}_A \left(\frac{Z_t}{Z_{,\alpha'}} \right).$$

Since the fluid is incompressible and irrotational, so $\mathbf{v} = \nabla\varphi$, where the velocity potential φ satisfies the Bernoulli equation in the fluid domain $\Omega(t)$:

$$(2.2) \quad \begin{cases} \varphi_t + \frac{1}{2}|\nabla\varphi|^2 + P + y = 0, & \Delta\varphi = 0, & \text{in } \Omega(t), \\ P = 0, & & \text{on } \partial\Omega(t). \end{cases}$$

Let $\psi(\alpha, t) := \varphi(z(\alpha, t), t)$ be the trace of the velocity potential on the interface, and

$$Q := (I + \mathbb{H})(\psi \circ \mathbf{h}^{-1}) = 2\mathbb{P}_H(\psi \circ \mathbf{h}^{-1}).$$

Lemma 2.1. *We have¹⁹*

$$(2.3) \quad D_t Q = i(Z - \alpha') + \mathbb{P}_A(|Z_t|^2),$$

$$(2.4) \quad D_t \mathbb{P}_H D_t Q + i \frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} Q = i \mathbb{P}_A \left(Z_t \left(1 - \frac{1}{Z_{,\alpha'}} \right) + \overline{Z}_t \left(\frac{1}{\overline{Z}_{,\alpha'}} - 1 \right) \right).$$

Proof. By chain rule, (2.2) and the fact that $\nabla\varphi(z, t) = z_t$ on the interface z , we have

$$(2.5) \quad \begin{cases} \psi_t = \varphi_t + \nabla\varphi \cdot z_t = -y + \frac{1}{2}|z_t|^2, \\ \psi_\alpha = \nabla\varphi \cdot z_\alpha = z_t \cdot z_\alpha. \end{cases}$$

Changing to the Riemann mapping variable α' in the second equation yields $\partial_{\alpha'}(\psi \circ \mathbf{h}^{-1}) = Z_t \cdot Z_{,\alpha'} = \operatorname{Re}(\overline{Z}_t Z_{,\alpha'})$, which in turn gives

$$(2.6) \quad \partial_{\alpha'} Q = \overline{Z}_t Z_{,\alpha'},$$

¹⁸The "quintic" in §2 in general means it is a finite sum of terms homogeneous of degree 5 or higher of the unknown functions Z_t , $\frac{1}{Z_{,\alpha'}} - 1$ and their derivatives. However at the end what we construct in this section are the energy functionals $\mathfrak{E}_j(t)$, cf.

Theorem 2.8, which are quintic in terms of only the spatial derivatives of $\left(Z_t, \frac{1}{Z_{,\alpha'}} - 1 \right)$.

¹⁹Applying $D_{\alpha'}$ to both sides of (2.3) gives the first equation in the interface system (B.1).

because the holomorphic quantities $\partial_{\alpha'} Q$ and $\overline{Z}_t Z_{,\alpha'}$ have the same real parts. A change of variables in the first equation in (2.5) yields $D_t(\psi \circ h^{-1}) = -Y + \frac{1}{2}|Z_t|^2$.

We compute, by first commuting D_t with \mathbb{H} , then applying (2.1), Proposition B.1, (A.3) and (2.6), that

$$\begin{aligned} D_t Q &= (I + \mathbb{H})D_t(\psi \circ h^{-1}) + [D_t, \mathbb{H}](\psi \circ h^{-1}) \\ &= (I + \mathbb{H})(-Y + \frac{1}{2}|Z_t|^2) + \left[\frac{Z_t}{Z_{,\alpha'}}, \mathbb{H} \right] \partial_{\alpha'} \mathbb{P}_H(\psi \circ h^{-1}) + \left[\frac{\overline{Z}_t}{\overline{Z}_{,\alpha'}}, \mathbb{H} \right] \partial_{\alpha'} \mathbb{P}_A(\psi \circ h^{-1}) \\ &= (I + \mathbb{H})(-Y + \frac{1}{2}|Z_t|^2) + \mathbb{P}_A \left(\frac{Z_t}{Z_{,\alpha'}} \partial_{\alpha'} Q \right) - \mathbb{P}_H \left(\frac{\overline{Z}_t}{\overline{Z}_{,\alpha'}} \partial_{\alpha'} \overline{Q} \right) \\ &= i(Z - \alpha') + \mathbb{P}_A(|Z_t|^2), \end{aligned}$$

where $(I + \mathbb{H})(-Y) = i(Z - \alpha')$ because both of the holomorphic quantities $(I + \mathbb{H})(-Y)$ and $i(Z - \alpha')$ have the same real parts. This gives (2.3).²⁰

Observe that $\mathbb{P}_H D_t Q = i(Z - \alpha')$ by (2.3). We further compute, by (2.1) and the fact that $\mathbb{P}_A Z_t = Z_t$ and $\mathbb{P}_A \overline{Z}_t = 0$, that

$$D_t \{i(Z - \alpha')\} = i(Z_t - b) = -i \frac{\overline{Z}_t}{\overline{Z}_{,\alpha'}} + i \mathbb{P}_A \left(Z_t \left(1 - \frac{1}{Z_{,\alpha'}} \right) + \overline{Z}_t \left(\frac{1}{\overline{Z}_{,\alpha'}} - 1 \right) \right).$$

Observe that $i \frac{\overline{Z}_t}{\overline{Z}_{,\alpha'}} = i \frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} Q$ by (2.6). This gives (2.4). \square

Let

$$(2.7) \quad \Theta^{(0)} := Q, \quad \Theta^{(j)} := (\mathbb{P}_H D_t)^j Q, \quad \text{and}$$

$$(2.8) \quad D_t \mathbb{P}_H D_t \Theta^{(j)} + i \frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \Theta^{(j)} =: G^{(j)}.$$

We know $\mathbb{P}_H G^{(0)} = 0$ by (2.4). We would like to find a formula for $\mathbb{P}_H G^{(j)}$. To this end we derive a recursive relation, which in turn gives the formula for $\mathbb{P}_H G^{(j)}$.

Proposition 2.2. *1. Let Θ be holomorphic, i.e. $\mathbb{P}_A \Theta = 0$, and $\Theta_1 = \mathbb{P}_H D_t \Theta$. Assume that*

$$(2.9) \quad D_t \mathbb{P}_H D_t \Theta + i \frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \Theta = G, \quad \text{and} \quad D_t \mathbb{P}_H D_t \Theta_1 + i \frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \Theta_1 = G_1.$$

Then

$$(2.10) \quad \mathbb{P}_H G_1 - \mathbb{P}_H D_t \mathbb{P}_H G = \frac{1}{2} \mathbb{P}_H \left\{ \frac{1}{\overline{Z}_{,\alpha'}} \left(\langle \overline{Z}_t, i \frac{1}{\overline{Z}_{,\alpha'}} D_{\alpha'} \Theta \rangle + \langle -i \frac{1}{Z_{,\alpha'}}, Z_t, D_{\alpha'} \Theta \rangle \right) \right\},$$

where the cubic-form $\langle \cdot, \cdot, \cdot \rangle$ is defined by

$$\langle f, g, h \rangle := \frac{1}{\pi i} \int \frac{(f(\alpha') - f(\beta'))(g(\alpha') - g(\beta'))(h(\alpha') - h(\beta'))}{(\alpha' - \beta')^2} d\beta'.$$

2. We have

$$(2.11) \quad \mathbb{P}_H(G^{(j)}) = \sum_{l=0}^{j-1} (\mathbb{P}_H D_t)^l \left(\mathbb{P}_H(G^{(j-l)}) - \mathbb{P}_H D_t \mathbb{P}_H(G^{(j-l-1)}) \right),$$

where $\mathbb{P}_H(G^{(j-l)}) - \mathbb{P}_H D_t \mathbb{P}_H(G^{(j-l-1)})$ is given by (2.10) with $\Theta = \Theta^{(j-l-1)}$.

²⁰Equation (2.3) was also derived in [27] using a slightly different approach.

Proof. Let $F = \mathbb{P}_A D_t \Theta$ and $F_1 = \mathbb{P}_A D_t \Theta_1$. We know by (2.1) and the identity (A.4) that

$$(2.12) \quad F = \mathbb{P}_A D_t \Theta = \mathbb{P}_A \left(\frac{Z_t}{|Z, \alpha'|^2} \partial_{\alpha'} \Theta \right).$$

It is easy to check, by the definitions and the decomposition identity (A.3) that

$$D_t \mathbb{P}_H G - G_1 = -D_t \mathbb{P}_A G + D_t F_1 + i \frac{1}{|Z, \alpha'|^2} \partial_{\alpha'} F + \left[D_t, i \frac{1}{|Z, \alpha'|^2} \partial_{\alpha'} \right] \Theta;$$

using the definitions again yields

$$\mathbb{P}_A G = F_1 + \mathbb{P}_A \left(i \frac{1}{|Z, \alpha'|^2} \partial_{\alpha'} \Theta \right),$$

so

$$D_t \mathbb{P}_H G - G_1 = -D_t \mathbb{P}_A \left(i \frac{1}{|Z, \alpha'|^2} \partial_{\alpha'} \Theta \right) + i \frac{1}{|Z, \alpha'|^2} \partial_{\alpha'} F + \left[D_t, i \frac{1}{|Z, \alpha'|^2} \partial_{\alpha'} \right] \Theta;$$

this gives, by (2.1), (A.4), (2.12) and (B.20), that

$$(2.13) \quad \begin{aligned} \mathbb{P}_H (D_t \mathbb{P}_H G - G_1) &= -\mathbb{P}_H \left(\frac{\overline{Z}_t}{\overline{Z}, \alpha'} \partial_{\alpha'} \mathbb{P}_A \left(i \frac{1}{|Z, \alpha'|^2} \partial_{\alpha'} \Theta \right) \right) \\ &\quad + \mathbb{P}_H \left(i \frac{1}{|Z, \alpha'|^2} \partial_{\alpha'} \mathbb{P}_A \left(\frac{Z_t}{Z, \alpha'} \partial_{\alpha'} \Theta \right) \right) + i \mathbb{P}_H \left(\frac{b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t}{|Z, \alpha'|^2} \partial_{\alpha'} \Theta \right). \end{aligned}$$

Now by identity (A.4) and the definition of $D_{\alpha'} \Theta$ we can rewrite the first two terms in (2.13) as

$$\begin{aligned} -\mathbb{P}_H \left(\frac{\overline{Z}_t}{\overline{Z}, \alpha'} \partial_{\alpha'} \mathbb{P}_A \left(i \frac{1}{|Z, \alpha'|^2} \partial_{\alpha'} \Theta \right) \right) &= -\mathbb{P}_H \left(\frac{1}{\overline{Z}, \alpha'} \mathbb{P}_H \left(\overline{Z}_t \partial_{\alpha'} \mathbb{P}_A \left(i \frac{1}{\overline{Z}, \alpha'} D_{\alpha'} \Theta \right) \right) \right), \\ \mathbb{P}_H \left(i \frac{1}{|Z, \alpha'|^2} \partial_{\alpha'} \mathbb{P}_A \left(\frac{Z_t}{Z, \alpha'} \partial_{\alpha'} \Theta \right) \right) &= \mathbb{P}_H \left(\frac{1}{\overline{Z}, \alpha'} \mathbb{P}_H \left(i \frac{1}{Z, \alpha'} \partial_{\alpha'} \mathbb{P}_A (Z_t D_{\alpha'} \Theta) \right) \right). \end{aligned}$$

We use (B.11) and the fact that \overline{Z}_t and $\frac{1}{Z, \alpha'}$ are holomorphic to further rewrite

$$-\mathbb{P}_H \left(\overline{Z}_t \partial_{\alpha'} \mathbb{P}_A \left(i \frac{1}{\overline{Z}, \alpha'} D_{\alpha'} \Theta \right) \right) = \frac{1}{2} \mathbb{P}_H \left[\overline{Z}_t, i \frac{1}{\overline{Z}, \alpha'}; D_{\alpha'} \Theta \right]$$

and

$$\mathbb{P}_H \left(i \frac{1}{Z, \alpha'} \partial_{\alpha'} \mathbb{P}_A (Z_t D_{\alpha'} \Theta) \right) = \frac{1}{2} \mathbb{P}_H \left[-i \frac{1}{Z, \alpha'}, Z_t; D_{\alpha'} \Theta \right].$$

Replacing $b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t$ in (2.13) by (B.7) and using part 2. of Proposition B.1 gives (2.10).

It is easy to check (2.11) using the fact that $\mathbb{P}_H G^{(0)} = 0$. \square

We next present a new energy identity. The basic energy form, such as $\int |\partial_t \theta|^2 + \mathfrak{a} \nabla_n \theta \bar{\theta} d\alpha$, is commonly used in the proof of the local well-posedness of the water wave equations. In [48] we found that the energy form

$$\int \frac{1}{\mathfrak{a}} |\partial_t \theta|^2 + i \partial_{\alpha} \theta \bar{\theta} d\alpha,$$

with the coefficient \mathfrak{a} moved to the first term and the Dirichlet-Neumann operator ∇_n replaced by $i \partial_{\alpha}$ was advantageous in the study of the quadratic and cubic cancellations in the energy functionals and proving long time existence, see Lemma 4.1 of [48]; however it is not adequate for the purpose of this paper. In the following Proposition we introduce a new, symmetric, basic energy form. We will see that this new energy form makes it easier for us to find the quartic correcting functionals to cancel out the quartic terms in the time derivatives of the basic energies.

Proposition 2.3. *Let Θ_1, Θ_2 be holomorphic, i.e. $\mathbb{P}_A \Theta_1 = \mathbb{P}_A \Theta_2 = 0$, smooth and decay fast at infinity. Define*

$$(2.14) \quad E(t) = \operatorname{Re} \left(\int i \partial_{\alpha'} \Theta_2 \overline{D_t \Theta_1} d\alpha' - \int i \partial_{\alpha'} \Theta_1 \overline{D_t \Theta_2} d\alpha' \right).$$

Then

$$(2.15) \quad \frac{d}{dt} E(t) = \operatorname{Re} \left(\int i \partial_{\alpha'} \Theta_2 \overline{(\mathbb{P}_H G_1)} d\alpha' - \int i \partial_{\alpha'} \Theta_1 \overline{(\mathbb{P}_H G_2)} d\alpha' \right).$$

where $G_k := D_t \mathbb{P}_H D_t \Theta_k + i \frac{1}{|Z_{\alpha'}|^2} \partial_{\alpha'} \Theta_k$, for $k = 1, 2$.

Proof. We use (B.12) to compute $E'(t)$. We have, after applying (B.18) and cancelling out equal terms,²¹

$$(2.16) \quad \begin{aligned} \frac{d}{dt} E(t) &= \operatorname{Re} \left(\int i \partial_{\alpha'} \Theta_2 \overline{D_t^2 \Theta_1} d\alpha' - \int i \partial_{\alpha'} \Theta_1 \overline{D_t^2 \Theta_2} d\alpha' \right) \\ &= \operatorname{Re} \left(\int i \partial_{\alpha'} \Theta_2 \overline{(D_t^2 \Theta_1 + i \frac{1}{|Z_{\alpha'}|^2} \partial_{\alpha'} \Theta_1)} d\alpha' - \int i \partial_{\alpha'} \Theta_1 \overline{(D_t^2 \Theta_2 + i \frac{1}{|Z_{\alpha'}|^2} \partial_{\alpha'} \Theta_2)} d\alpha' \right); \end{aligned}$$

it is clear that the terms inserted in the second step above sum up to zero. We now want to show

$$(2.17) \quad \operatorname{Re} \int i \partial_{\alpha'} \Theta_2 \overline{D_t \mathbb{P}_A D_t \Theta_1} d\alpha' = \operatorname{Re} \int i \partial_{\alpha'} \Theta_1 \overline{D_t \mathbb{P}_A D_t \Theta_2} d\alpha'.$$

We begin with the term on the left hand side. Using Cauchy integral formula to insert a \mathbb{P}_H gives

$$\int i \partial_{\alpha'} \Theta_2 \overline{D_t \mathbb{P}_A D_t \Theta_1} d\alpha' = \int i \partial_{\alpha'} \Theta_2 \overline{\mathbb{P}_H D_t \mathbb{P}_A D_t \Theta_1} d\alpha'.$$

By (2.1) and (A.3), $\mathbb{P}_H D_t \mathbb{P}_A D_t \Theta_1 = \mathbb{P}_H (\frac{\overline{Z_t}}{\overline{Z_{\alpha'}}} \partial_{\alpha'} \mathbb{P}_A D_t \Theta_1)$, and using Cauchy integral formula again to remove the \mathbb{P}_H and insert a \mathbb{P}_A , and then use (2.12), we get

$$(2.18) \quad \int i \partial_{\alpha'} \Theta_2 \overline{\mathbb{P}_H D_t \mathbb{P}_A D_t \Theta_1} d\alpha' = \int i \partial_{\alpha'} \Theta_2 \overline{\frac{\overline{Z_t}}{\overline{Z_{\alpha'}}} \partial_{\alpha'} \mathbb{P}_A D_t \Theta_1} d\alpha' = i \int \mathbb{P}_A D_t \Theta_2 \partial_{\alpha'} \overline{\mathbb{P}_A D_t \Theta_1} d\alpha'.$$

Exchanging roles between Θ_1 and Θ_2 in (2.18) gives us a similar identity for the right hand term in (2.17), a further integration by parts then yields (2.17). From (2.16) and (2.17), and using Cauchy integral formula again to insert a \mathbb{P}_H gives (2.15). \square

Let $\Theta_1 = \Theta^{(j)}$, $\Theta_2 = \Theta^{(j+1)}$, and define

$$(2.19) \quad E_j(t) = \operatorname{Re} \left(\int i \partial_{\alpha'} \Theta^{(j+1)} \overline{\Theta^{(j+1)}} d\alpha' - \int i \partial_{\alpha'} \Theta^{(j)} \overline{\Theta^{(j+2)}} d\alpha' \right).$$

From (2.15) we get,

$$\frac{d}{dt} E_j(t) = \operatorname{Re} \left(\int i \partial_{\alpha'} \Theta^{(j+1)} \overline{\mathbb{P}_H G^{(j)}} d\alpha' - \int i \partial_{\alpha'} \Theta^{(j)} \overline{\mathbb{P}_H G^{(j+1)}} d\alpha' \right);$$

compute further by using (B.12), (2.11), (2.4) and Proposition B.4, we obtain

$$(2.20) \quad \begin{aligned} \frac{d}{dt} E_j(t) &= \frac{d}{dt} \operatorname{Re} \int i \partial_{\alpha'} \Theta^{(j)} \overline{\mathbb{P}_H G^{(j)}} d\alpha' \\ &\quad + 2 \operatorname{Re} \sum_{l=0}^{j-1} \left\{ \int i \partial_{\alpha'} \overline{\Theta^{(j)}} (\mathbb{P}_H D_t)^{l+1} \mathbb{P}_H \left(G^{(j-l)} - D_t \mathbb{P}_H G^{(j-1-l)} \right) d\alpha' \right\} \\ &\quad + \operatorname{Re} \left\{ \int i \partial_{\alpha'} \overline{\Theta^{(j)}} \mathbb{P}_H \left(G^{(j+1)} - D_t \mathbb{P}_H G^{(j)} \right) d\alpha' \right\}. \end{aligned}$$

²¹Observe that (2.16) holds without the assumption that Θ_1, Θ_2 are holomorphic.

where

$$(2.21) \quad \begin{aligned} & \mathbb{P}_H \left(G^{(j-l)} - D_t \mathbb{P}_H G^{(j-1-l)} \right) \\ &= \frac{1}{2} \mathbb{P}_H \left\{ \frac{1}{\overline{Z}_{,\alpha'}} \left(\langle \overline{Z}_t, i \frac{1}{\overline{Z}_{,\alpha'}} \rangle, D_{\alpha'} \Theta^{(j-l-1)} \rangle + \langle -i \frac{1}{\overline{Z}_{,\alpha'}}, Z_t, D_{\alpha'} \Theta^{(j-l-1)} \rangle \right) \right\}, \end{aligned}$$

for $l = -1, 0, \dots, j-1$, by Proposition 2.2.

Observe that $\frac{d}{dt} E_j(t)$ is quartic with a few desirable features: 1. the integrands depend only on the spatial derivatives of the quantities \overline{Z}_t , $\frac{1}{\overline{Z}_{,\alpha'}}$ and $\Theta^{(j)}$; 2. it has remarkable symmetries, with the order of derivatives evenly distributed among the factors: we note that $D_{\alpha'} \Theta^{(0)} = \overline{Z}_t$. This should allow us to readily derive very low regularity estimates. However our main interest here is to find a quartic correcting functional for $E_j(t)$, so that the time derivative of the corrected energy functional is quintic in terms of the spatial derivatives of the quantities \overline{Z}_t , $\frac{1}{\overline{Z}_{,\alpha'}}$ and $\Theta^{(j)}$ and prove the new long time existence result. The feature of $\frac{d}{dt} E_j(t)$ allows us to do so, thanks to a few additional observations on the structure of the water wave equation.

We note that by (B.1), $-i \frac{1}{\overline{Z}_{,\alpha'}} + i = \overline{Z}_{tt} + \text{quadratic}$, and by (2.6), $D_{\alpha'} \Theta^{(k)} = D_t^k \overline{Z}_t + \text{quadratic}$. Let²²

$$(2.22) \quad \mathfrak{D}_t := \partial_t + b(\alpha', t) \partial_{\alpha'} + b(\beta', t) \partial_{\beta'},$$

$$(2.23) \quad \mathcal{P} := \mathfrak{D}_t^2 + i \frac{A_1(\alpha', t)}{|\overline{Z}_{,\alpha'}|^2} \partial_{\alpha'} + i \frac{A_1(\beta', t)}{|\overline{Z}_{,\beta'}|^2} \partial_{\beta'}.$$

And let

$$(2.24) \quad \theta := \overline{Z}_t(\alpha', t) - \overline{Z}_t(\beta', t),$$

observe that

$$(2.25) \quad \mathfrak{D}_t^j \theta = D_t^j \overline{Z}_t(\alpha', t) - D_t^j \overline{Z}_t(\beta', t), \quad \text{for } j \geq 0.$$

So the quartic terms in

$$2 \operatorname{Re} \sum_{l=0}^{j-1} \left\{ \int i \partial_{\alpha'} \overline{\Theta^{(j)}} (\mathbb{P}_H D_t)^{l+1} \mathbb{P}_H \left(G^{(j-l)} - D_t \mathbb{P}_H G^{(j-1-l)} \right) d\alpha' \right\}$$

consists of

$$(2.26) \quad I_{1,j} := \frac{1}{\pi} \operatorname{Re} \sum_{l=0}^{j-1} \iint D_t^j Z_t(\alpha', t) \frac{\mathfrak{D}_t^{l+1} \left\{ \mathfrak{D}_t \left(\theta(\alpha', \beta', t) \overline{\theta(\alpha', \beta', t)} \right) \mathfrak{D}_t^{j-l-1} \theta(\alpha', \beta', t) \right\}}{(\alpha' - \beta')^2} d\beta' d\alpha'.$$

Similarly the quartic terms in $\operatorname{Re} \int i \partial_{\alpha'} \overline{\Theta^{(j)}} \mathbb{P}_H (G^{(j+1)} - D_t \mathbb{P}_H G^{(j)}) d\alpha'$ are

$$(2.27) \quad I_{2,j} := \frac{1}{2\pi} \operatorname{Re} \iint D_t^j Z_t(\alpha', t) \frac{\mathfrak{D}_t \left(\theta(\alpha', \beta', t) \overline{\theta(\alpha', \beta', t)} \right) \mathfrak{D}_t^j \theta(\alpha', \beta', t)}{(\alpha' - \beta')^2} d\beta' d\alpha'.$$

We know it is possible to find quartic functionals $C_{i,j}$, $i = 1, 2$, so that $\frac{d}{dt} C_{i,j}(t) = I_{i,j} + \text{quintic}$, based on the following observations.

²²Observe that for any function $f = f(\alpha', t)$, $\mathfrak{D}_t f = D_t f$ and $\mathcal{P} f = (D_t^2 + i \frac{A_1}{|\overline{Z}_{,\alpha'}|^2} \partial_{\alpha'}) f =: \mathfrak{P} f$, where \mathfrak{P} is defined in (B.3).

Lemma 2.4. Assume that f and \mathcal{G} are smooth and decay fast at infinity. We have

$$\begin{aligned}
 (2.28) \quad & \frac{d}{dt} \iint \frac{\bar{f} \mathfrak{D}_t \mathcal{G} - \mathfrak{D}_t \bar{f} \mathcal{G}}{(\alpha' - \beta')^2} d\alpha' d\beta' = \iint \frac{\bar{f} \mathcal{P} \mathcal{G}}{(\alpha' - \beta')^2} d\alpha' d\beta' \\
 & + \iint \left(-\frac{\overline{\mathcal{P} f \mathcal{G}}}{(\alpha' - \beta')^2} + \left(b_{\alpha'} + b_{\beta'} - 2 \frac{b(\alpha') - b(\beta')}{\alpha' - \beta'} \right) \frac{\bar{f} \mathfrak{D}_t \mathcal{G} - \mathfrak{D}_t \bar{f} \mathcal{G}}{(\alpha' - \beta')^2} \right) d\alpha' d\beta' \\
 & + i \iint \left(\partial_{\alpha'} \frac{A_1(\alpha')}{|Z_{,\alpha'}|^2} + \partial_{\beta'} \frac{A_1(\beta')}{|Z_{,\beta'}|^2} - 2 \frac{\frac{A_1(\alpha')}{|Z_{,\alpha'}|^2} - \frac{A_1(\beta')}{|Z_{,\beta'}|^2}}{\alpha' - \beta'} \right) \frac{\bar{f} \mathcal{G}}{(\alpha' - \beta')^2} d\alpha' d\beta'.
 \end{aligned}$$

(2.28) follows easily from (B.13) and integration by parts. We omit the details.

Lemma 2.5. Let $\mathcal{G} = g \bar{h} q$. Then

$$(2.29) \quad \mathcal{P} \mathcal{G} = 2 \mathfrak{D}_t (g \bar{h}) \mathfrak{D}_t q + 2 \mathfrak{D}_t (g \mathfrak{D}_t \bar{h}) q + \bar{h} q \mathcal{P} g + g \bar{h} \mathcal{P} q - g q \overline{\mathcal{P} h}.$$

The verification of (2.29) is straightforward. We omit the details.

To find the quartic functionals $C_{i,j}$, we will use Lemma 2.4 with $\mathcal{G} = g \bar{h} q$. From (2.28), (2.29), we have

$$(2.30) \quad \frac{d}{dt} \iint \frac{\bar{f} \mathfrak{D}_t (g \bar{h} q) - (\mathfrak{D}_t \bar{f}) g \bar{h} q}{(\alpha' - \beta')^2} d\alpha' d\beta' = 2 \iint \frac{\bar{f} \mathfrak{D}_t (g \bar{h}) \mathfrak{D}_t q + \bar{f} \mathfrak{D}_t (g \mathfrak{D}_t \bar{h}) q}{(\alpha' - \beta')^2} d\alpha' d\beta' + \text{quintic},$$

provided $\mathcal{P} f$, $\mathcal{P} g$, $\mathcal{P} h$ and $\mathcal{P} q$ are quadratic. Observe that on the right hand side of (2.30) the operator \mathfrak{D}_t acting on q in the first term is moved to the second term, acting on \bar{h} . And we know $\mathcal{P} \mathfrak{D}_t^j \theta = \text{quadratic}$ by (B.4). Instead of going through the straightforward but tedious process of finding the correcting functionals $C_{i,j}$, we will directly give the results. Before doing so, we use Lemmas 2.4 and 2.5 to write the following equation. We have, for $j, k, i, l, m \geq 0$, and $D_t^j Z_t = D_t^j Z_t(\alpha', t)$,

$$\begin{aligned}
 (2.31) \quad & \frac{d}{dt} \iint \frac{\left(D_t^j Z_t \mathfrak{D}_t - \mathfrak{D}_t (D_t^j Z_t) \right) \mathfrak{D}_t^m (\mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta)}{(\alpha' - \beta')^2} d\alpha' d\beta' \\
 & = 2 \iint \frac{D_t^j Z_t \mathfrak{D}_t^m \left\{ \mathfrak{D}_t (\mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta}) \mathfrak{D}_t^{k+1} \theta + \mathfrak{D}_t (\mathfrak{D}_t^l \theta \mathfrak{D}_t^{i+1} \bar{\theta}) \mathfrak{D}_t^k \theta \right\}}{(\alpha' - \beta')^2} d\alpha' d\beta' + R_{j;l,\bar{i},k}^{(m)},
 \end{aligned}$$

where

$$\begin{aligned}
 (2.32) \quad & R_{j;l,\bar{i},k}^{(m)} = \iint \frac{D_t^j Z_t [\mathcal{P}, \mathfrak{D}_t^m] (\mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta)}{(\alpha' - \beta')^2} d\alpha' d\beta' \\
 & + \iint \frac{D_t^j Z_t \mathfrak{D}_t^m \left\{ (\mathcal{P} \mathfrak{D}_t^l \theta) \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta - \mathfrak{D}_t^l \theta (\overline{\mathcal{P} \mathfrak{D}_t^i \bar{\theta}}) \mathfrak{D}_t^k \theta + \mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} (\mathcal{P} \mathfrak{D}_t^k \theta) \right\}}{(\alpha' - \beta')^2} d\alpha' d\beta' \\
 & - \iint \frac{\overline{(\mathcal{P} D_t^j \bar{Z}_t)} \mathfrak{D}_t^m (\mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta)}{(\alpha' - \beta')^2} d\alpha' d\beta' \\
 & + \iint \left(b_{\alpha'} + b_{\beta'} - 2 \frac{b(\alpha') - b(\beta')}{\alpha' - \beta'} \right) \frac{\left(D_t^j Z_t \mathfrak{D}_t - \mathfrak{D}_t (D_t^j Z_t) \right) \mathfrak{D}_t^m (\mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta)}{(\alpha' - \beta')^2} d\alpha' d\beta' \\
 & + i \iint \left(\partial_{\alpha'} \frac{A_1(\alpha')}{|Z_{,\alpha'}|^2} + \partial_{\beta'} \frac{A_1(\beta')}{|Z_{,\beta'}|^2} - 2 \frac{\frac{A_1(\alpha')}{|Z_{,\alpha'}|^2} - \frac{A_1(\beta')}{|Z_{,\beta'}|^2}}{\alpha' - \beta'} \right) \frac{D_t^j Z_t \mathfrak{D}_t^m (\mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta)}{(\alpha' - \beta')^2} d\alpha' d\beta'.
 \end{aligned}$$

It is clear that $R_{j;l,\bar{i},k}^{(m)}$ is quintic.

We now give the correcting functionals. Let²³

$$\begin{aligned}
(2.33) \quad C_{1,j} = & \frac{1}{2\pi} \sum_{l=0}^{j-1} \sum_{k=0}^l \iint \frac{(D_t^j Z_t \mathfrak{D}_t - D_t^{j+1} Z_t) \mathfrak{D}_t^{l-k} (\mathfrak{D}_t^k \theta \bar{\theta} \mathfrak{D}_t^{j-l-1} \theta)}{(\alpha' - \beta')^2} d\beta' d\alpha' \\
& + \frac{1}{4\pi} \sum_{l=0}^{j-2} \sum_{k=0}^{j-l-2} (-1)^k \iint \frac{(D_t^j Z_t \mathfrak{D}_t - D_t^{j+1} Z_t) \mathfrak{D}_t^{1+l} \theta \mathfrak{D}_t^k \bar{\theta} \mathfrak{D}_t^{j-l-2-k} \theta}{(\alpha' - \beta')^2} d\beta' d\alpha' \\
& - \frac{1}{8\pi} \sum_{l=0}^{j-2} \sum_{k=0}^{j-l-2} (-1)^k \iint \frac{(\theta \mathfrak{D}_t - \mathfrak{D}_t \theta) \mathfrak{D}_t^{j-l-1} \bar{\theta} \mathfrak{D}_t^{j-k-1} \theta \mathfrak{D}_t^{k+l+1} \bar{\theta}}{(\alpha' - \beta')^2} d\beta' d\alpha' \\
& + \frac{1}{2\pi} \sum_{l=0}^{j-1} \iint D_t^j Z_t \frac{\mathfrak{D}_t^{j-l-1} \theta \bar{\theta} \mathfrak{D}_t^{1+l} \theta}{(\alpha' - \beta')^2} d\beta' d\alpha',
\end{aligned}$$

and

$$\begin{aligned}
(2.34) \quad C_{2,j} = & \frac{1}{4\pi} \sum_{k=0}^{j-1} (-1)^k \iint \frac{(D_t^j Z_t \mathfrak{D}_t - D_t^{j+1} Z_t) \theta \mathfrak{D}_t^k \bar{\theta} \mathfrak{D}_t^{j-k-1} \theta}{(\alpha' - \beta')^2} d\beta' d\alpha' \\
& + \frac{1}{4\pi} (-1)^j \iint D_t^j Z_t \frac{\theta \mathfrak{D}_t^j \bar{\theta} \theta}{(\alpha' - \beta')^2} d\beta' d\alpha'.
\end{aligned}$$

Proposition 2.6. *Let $j \geq 0$. We have*

$$(2.35) \quad I_{1,j} + I_{2,j} - \frac{d}{dt} \operatorname{Re}(C_{1,j} + C_{2,j}) = R_{IC,j}$$

where

$$\begin{aligned}
(2.36) \quad R_{IC,j} = & -\frac{1}{2\pi} \sum_{l=0}^{j-1} \operatorname{Re} \iint \left(b_{\alpha'} + b_{\beta'} - 2 \frac{b(\alpha') - b(\beta')}{\alpha' - \beta'} \right) \frac{D_t^j Z_t \mathfrak{D}_t^{j-l-1} \theta \bar{\theta} \mathfrak{D}_t^{l+1} \theta}{(\alpha' - \beta')^2} d\alpha' d\beta' \\
& - \frac{1}{4\pi} \operatorname{Re} \left(2 \sum_{l=0}^{j-1} \sum_{k=0}^l R_{j;k,0,j-l-1}^{(l-k)} + \sum_{l=0}^{j-2} \sum_{k=0}^{j-l-2} (-1)^k \left(R_{j;1+l,\bar{k},j-l-2-k}^{(0)} - \overline{R_{0;j-l-1,j-k-1,l+k+1}^{(0)}} \right) \right) \\
& - (-1)^j \frac{1}{4\pi} \operatorname{Re} \iint \left(b_{\alpha'} + b_{\beta'} - 2 \frac{b(\alpha') - b(\beta')}{\alpha' - \beta'} \right) \frac{D_t^j Z_t \theta \mathfrak{D}_t^j \bar{\theta} \theta}{(\alpha' - \beta')^2} d\alpha' d\beta' \\
& - \frac{1}{4\pi} \operatorname{Re} \sum_{k=0}^{j-1} (-1)^k R_{j;0,\bar{k},j-k-1}^{(0)}.
\end{aligned}$$

Observe that $\mathfrak{D}_t D_t^j Z_t = D_t^{j+1} Z_t$. It is easy to prove Proposition 2.6 by (2.28)-(2.29) or (2.31)-(2.32),²⁴ (B.13) and the symmetry,²⁵ we omit the details.

²³We define $\sum_{l=0}^{-1} = \sum_{l=0}^{-2} = 0$. $D_t^j Z_t = D_t^j Z_t(\alpha', t)$, $D_t^{j+1} Z_t = D_t^{j+1} Z_t(\alpha', t)$ in (2.33) and (2.34).

²⁴When computing $\frac{d}{dt} C_{1,j}$, use the identity

$$\mathfrak{D}_t^{l-k} \left(\mathfrak{D}_t (\mathfrak{D}_t^k \theta \bar{\theta}) \mathfrak{D}_t^{j-l-1} \theta + \mathfrak{D}_t (\mathfrak{D}_t^k \theta \mathfrak{D}_t \bar{\theta}) \mathfrak{D}_t^{j-l-1} \theta \right) = \mathfrak{D}_t^{l-k+1} \left(\mathfrak{D}_t (\mathfrak{D}_t^k \theta \bar{\theta}) \mathfrak{D}_t^{j-l-1} \theta \right) - \mathfrak{D}_t^{l-k} \left(\mathfrak{D}_t (\mathfrak{D}_t^{k+1} \theta \bar{\theta}) \mathfrak{D}_t^{j-l-1} \theta \right).$$

Also, when using (2.28)-(2.29) or (2.31)-(2.32) to compute, apply \mathfrak{D}_t to the factors in the exact order given in the right hand side of (2.29) or (2.31) to facilitate cancellations.

²⁵By interchanging α' with β' and use symmetry we have

$$\begin{aligned}
(2.37) \quad & \iint \frac{f(\alpha')(g(\alpha') - g(\beta'))(h(\alpha') - h(\beta'))(q(\alpha') - q(\beta'))}{(\alpha' - \beta')^2} d\alpha' d\beta' \\
& = \frac{1}{2} \iint \frac{(f(\alpha') - f(\beta'))(g(\alpha') - g(\beta'))(h(\alpha') - h(\beta'))(q(\alpha') - q(\beta'))}{(\alpha' - \beta')^2} d\alpha' d\beta'.
\end{aligned}$$

Remark 2.7. It is clear from (2.36) that the remainder $R_{IC,j}$ is quintic.

We now sum up (2.20)-(2.21)-(2.26)-(2.27) and Proposition 2.6, and present it in the following. Let

$$(2.38) \quad \mathfrak{E}_j(t) = E_j(t) - \operatorname{Re} \left(\int i \partial_{\alpha'} \Theta^{(j)} \overline{\mathbb{P}_H G^{(j)}} d\alpha' + C_{1,j}(t) + C_{2,j}(t) \right).$$

Theorem 2.8. *We have*

$$(2.39) \quad \frac{d}{dt} \mathfrak{E}_j(t) = \mathfrak{R}_j(t),$$

where

$$(2.40) \quad \begin{aligned} \mathfrak{R}_j = & \left(2 \operatorname{Re} \sum_{l=0}^{j-1} \int i \partial_{\alpha'} \overline{\Theta^{(j)}} (\mathbb{P}_H D_t)^{l+1} \mathbb{P}_H \left(G^{(j-l)} - D_t \mathbb{P}_H G^{(j-1-l)} \right) d\alpha' - I_{1,j} \right) \\ & + \left(\operatorname{Re} \int i \partial_{\alpha'} \overline{\Theta^{(j)}} \mathbb{P}_H \left(G^{(j+1)} - D_t \mathbb{P}_H G^{(j)} \right) d\alpha' - I_{2,j} \right) + R_{IC,j} \end{aligned}$$

is quintic.

In particular for $j = 0$, equations (2.38)-(2.40) gives

$$(2.41) \quad \mathfrak{E}_0(t) = \int \left(i \partial_{\alpha'} (Z - \alpha') \overline{(Z - \alpha')} + |Z_t|^2 \right) d\alpha' - \frac{1}{8\pi} \iint \frac{|Z_t(\alpha') - Z_t(\beta')|^4}{(\alpha' - \beta')^2} d\alpha' d\beta'$$

and

$$(2.42) \quad \begin{aligned} \frac{d}{dt} \mathfrak{E}_0(t) = & \frac{1}{2} \int i Z_t \left(\langle \overline{Z}_t, i \frac{1-A_1}{\overline{Z}_{\alpha'}} \overline{Z}_t \rangle + \langle -i \frac{1-A_1}{Z_{\alpha'}}, Z_t, \overline{Z}_t \rangle \right) d\alpha' \\ & - \frac{1}{8\pi} \iint \left(b_{\alpha'} + b_{\beta'} - 2 \frac{b(\alpha') - b(\beta')}{\alpha' - \beta'} \right) \frac{|Z_t(\alpha') - Z_t(\beta')|^4}{(\alpha' - \beta')^2} d\alpha' d\beta'. \end{aligned}$$

In our proof for the long time existence result, Theorem 3.1, we will only use (2.39)-(2.40) for $1 \leq j \leq 4$. Similar to (2.20)-(2.21), equations (2.39)-(2.40) can be readily used to derive very low regularity results. However our focus in this paper is on the new long time existence result for solutions to the water wave equation.

2.1. Scaling. The solutions for the 2d water wave equation (B.1)-(B.2) obey the following scaling law: If (\overline{Z}_t, Z) is a solution of (B.1)-(B.2), then

$$(2.43) \quad (\overline{Z}_t^\lambda, Z^\lambda) := (\lambda^{-1/2} \overline{Z}_t(\lambda\alpha', \lambda^{1/2}t), \lambda^{-1} Z(\lambda\alpha', \lambda^{1/2}t))$$

is also a solution of the equation (B.1)-(B.2). The following norms are scaling invariant for the water wave equation (B.1)-(B.2): $\left\| \frac{1}{Z_{\alpha'}} \right\|_{\dot{H}^{1/2}(\mathbb{R})}$, $\|\overline{Z}_{t,\alpha'}\|_{L^2(\mathbb{R})}$, and $\left\| \frac{1}{Z_{\alpha'}} - 1 \right\|_{L^\infty(\mathbb{R})}$.

3. THE LONG TIME EXISTENCE OF STEEP WATER WAVES

We are now ready to present our new long time existence result, Theorem 3.1.

Let

$$(3.1) \quad L(t) = \left\| \frac{1}{Z_{\alpha'}}(t) \right\|_{\dot{H}^{1/2}(\mathbb{R})} + \|\overline{Z}_{t,\alpha'}(t)\|_{L^2(\mathbb{R})} + \left\| \partial_{\alpha'} \frac{1}{Z_{\alpha'}}(t) \right\|_{\dot{H}^{1/2}(\mathbb{R})} + \|\partial_{\alpha'}^2 \overline{Z}_t(t)\|_{L^2(\mathbb{R})}.$$

Theorem 3.1. *1. Let $J \geq 2$. Assume that the initial data $(\overline{Z}_t(0), \frac{1}{Z_{\alpha'}}(0) - 1) \in \cap_{\frac{1}{2} \leq s \leq J} \dot{H}^s(\mathbb{R}) \times \dot{H}^{s-\frac{1}{2}}(\mathbb{R})$. Then there are constants $m_0 > 0$, and $0 < \varepsilon_0 \leq 1$, such that for all $0 < \varepsilon \leq \varepsilon_0$, if the data satisfies*

$$(3.2) \quad L(0) \leq \varepsilon, \quad \left\| \frac{1}{Z_{\alpha'}}(0) - 1 \right\|_{L^\infty} < 1, \quad \text{and} \quad E_1(0)E_3(0) \leq m_0^2,$$

there is a constant $\mathcal{T}_0 > 0$, depending only on m_0 , so that the initial value problem for the water wave equation (B.1)-(B.2) or equivalently (1.1) has a unique classical solution for the time period $[0, \frac{\mathcal{T}_0}{\varepsilon^3}]$. During this time, the solution is as regular as the initial data, $L(t) \lesssim \varepsilon$, $\left\| \frac{1}{Z_{,\alpha'}}(t) - 1 \right\|_{L^\infty} < 1$ and $E_1(t)E_3(t) \lesssim m_0^2$.

2. If instead of (3.2) the data satisfies

$$(3.3) \quad \left\| \frac{1}{Z_{,\alpha'}}(0) \right\|_{\dot{H}^{1/2}(\mathbb{R})} + \left\| \overline{Z}_{t,\alpha'}(0) \right\|_{L^2(\mathbb{R})} \leq \varepsilon, \quad \left\| \frac{1}{Z_{,\alpha'}}(0) - 1 \right\|_{L^\infty} < 1, \quad \text{and} \quad E_1(0)E_3(0) \leq m_0^2,$$

then there is a constant $\mathcal{T}_1 > 0$, depending on m_0 , $\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}}(0) \right\|_{\dot{H}^{1/2}}$ and $\left\| \partial_{\alpha'}^2 \overline{Z}_t(0) \right\|_{L^2}$, so that the initial value problem for the water wave equation (B.1)-(B.2) or equivalently (1.1) has a unique classical solution for the time period $[0, \frac{\mathcal{T}_1}{\varepsilon^{5/2}}]$. During this time, the solution is as regular as the initial data.

Remark 3.2. Observe that $E_1(t)E_3(t)$ is scaling invariant. The sole reason for the assumption on $E_1(0)E_3(0)$ is to control the evolution of the norm $\left\| \frac{1}{Z_{,\alpha'}}(t) - 1 \right\|_{L^\infty}$. We will show that for ε small enough, $E_1(t)$ controls $\left\| \frac{1}{Z_{,\alpha'}}(t) - 1 \right\|_{L^2}^2 + \left\| \Theta^{(2)}(t) \right\|_{\dot{H}^{1/2}}^2$, and $E_3(t)$ controls $\frac{1}{4} \left\| D_{\alpha'} \frac{1}{Z_{,\alpha'}^2}(t) \right\|_{L^2}^2 + \left\| \Theta^{(4)}(t) \right\|_{\dot{H}^{1/2}}^2$, see Proposition 4.4; and the quantities $\Theta^{(2)}(t)$, $\Theta^{(4)}(t)$ are mainly related to the velocity Z_t , see Step 1, Step 3 in §4.3.2. We know by Lemma 4.2 that, assuming $\left\| 1 - \frac{1}{Z_{,\alpha'}} \right\|_{L^\infty} \leq 1$, there is a constant $c > 0$, such that

$$(3.4) \quad \left\| \frac{1}{Z_{,\alpha'}} - 1 \right\|_{L^\infty(\mathbb{R})}^2 \leq c \left\| \frac{1}{Z_{,\alpha'}} - 1 \right\|_{L^2(\mathbb{R})} \left\| D_{\alpha'} \frac{1}{Z_{,\alpha'}^2} \right\|_{L^2(\mathbb{R})},$$

so

$$(3.5) \quad \left\| \frac{1}{Z_{,\alpha'}}(t) - 1 \right\|_{L^\infty}^4 \leq 4c^2 E_1(t)E_3(t).$$

We will show that the growth of $E_1(t)E_3(t)$ can be controlled for time of order $O(\varepsilon^{-3})$, which in turn gives control of $\left\| \frac{1}{Z_{,\alpha'}}(t) - 1 \right\|_{L^\infty}$ for the same time period. The constants $m_0 > 0$ is chosen so that $\left\| \frac{1}{Z_{,\alpha'}}(0) - 1 \right\|_{L^\infty} < 1$.²⁶ Since $\left\| \frac{1}{Z_{,\alpha'}}(0) - 1 \right\|_{L^\infty}$ can be arbitrarily close to 1, the slope of the (initial) interface can be arbitrary large. Observe also that no assumption is imposed on the magnitude of Z_t . So the magnitude of the (initial) velocity can be arbitrary large.

Remark 3.3. Part 2 of Theorem 3.1 is a direct consequence of part 1. Observe that the rescaled data $(\overline{Z}_t^\varepsilon(0), Z^\varepsilon(0))$ satisfies

$$\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}^\varepsilon}(0) \right\|_{\dot{H}^{1/2}} + \left\| \partial_{\alpha'}^2 \overline{Z}_t^\varepsilon(0) \right\|_{L^2} = \varepsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}}(0) \right\|_{\dot{H}^{1/2}} + \varepsilon \left\| \partial_{\alpha'}^2 \overline{Z}_t(0) \right\|_{L^2}.$$

So by part 1, the assumption (3.3) implies that the lifespan of the rescaled solution $(\overline{Z}_t^\varepsilon, Z^\varepsilon)$ is of order $O(\varepsilon^{-3})$, therefore the solution (\overline{Z}_t, Z) has a lifespan of order $O(\varepsilon^{-5/2})$. In fact, the same scaling argument shows that for data satisfying (3.3) and $(Z_t(0), \frac{1}{Z_{,\alpha'}}(0) - 1) \in \cap_{\frac{1}{2} \leq s \leq J} \dot{H}^s \times \dot{H}^{s-1/2}$, the life span of the solution is at least of order $O(\varepsilon^{-3+\frac{1}{2J-2}})$, and the solution remains as smooth as the initial data during this time.

By Remark 3.3, it suffices to prove part 1 of Theorem 3.1.

²⁶Assuming c is the optimal constant so that (3.4) holds, we choose m_0 such that $m_0^2 < \frac{1}{4c^2}$.

Remark 3.4. In part 1 of Theorem 3.1, the smallness assumption is only imposed on the scaling invariant quantity $\left\| \frac{1}{Z_{,\alpha'}}(0) \right\|_{\dot{H}^{1/2}(\mathbb{R})} + \left\| \overline{Z}_{t,\alpha'}(0) \right\|_{L^2(\mathbb{R})}$ and the quantity $\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}}(0) \right\|_{\dot{H}^{1/2}(\mathbb{R})} + \left\| \partial_{\alpha'}^2 \overline{Z}_t(0) \right\|_{L^2(\mathbb{R})}$ which involves one higher order derivative. In this paper we do not try to lower this higher than scaling derivative to avoid technicality. We note that in all earlier works on long time existence [48, 49, 26, 8, 30, 24, 27, 28, 44, 1, 12], smallness is assumed on norms involving derivatives both above and below the scaling, so the smallness of the quantities can not be preserved after rescaling. And in all these earlier results, smallness is assumed on the slope of the initial interface.

Remark 3.5. In the regime where $\left\| \frac{1}{Z_{,\alpha'}}(t) - 1 \right\|_{L^\infty} < 1$, the interface $Z = Z(\alpha', t)$, $\alpha' \in \mathbb{R}$ is a graph.

4. THE PROOF OF THEOREM 3.1

4.1. Some additional quintic correcting functionals. To prove Theorem 3.1, we need some additional quintic correcting functionals to the energies $\mathfrak{E}_j(t)$, for $j \geq 2$. The goal is to have no loss of derivatives when controlling the higher order energy growth, to not involve $\left\| \frac{1}{Z_{,\alpha'}} - 1 \right\|_{L^\infty(\mathbb{R})}$ and any norms of order lower than scaling when bounding the cubic growth rates for the lower order energies, and to have proper control on the growth of $E_1(t)E_3(t)$.

Let $j \geq 2$. Observe that the factors $D_t^{j+1} Z_t$ and $\overline{\mathcal{P} D_t^j \overline{Z}_t}$ in $R_{j;l,\tilde{l},k}^{(m)}$ of (2.32) have more derivatives than are controlled by $\mathfrak{E}_j(t)$; we perform an "integration by parts" in the time variable, moving one D_t from the factors $D_t^{j+1} Z_t$ and $\overline{\mathcal{P} D_t^j \overline{Z}_t}$ onto others, resulting in the following quintic correcting functional.

Let

$$(4.1) \quad \begin{aligned} F_{j;l,\tilde{l},k}^{(m)}(t) = & \iint \frac{\overline{\mathcal{P} D_t^{j-1} \overline{Z}_t} \mathfrak{D}_t^m(\mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta)}{(\alpha' - \beta')^2} d\alpha' d\beta' \\ & + \iint \left(b_{\alpha'} + b_{\beta'} - 2 \frac{b(\alpha') - b(\beta')}{\alpha' - \beta'} \right) \frac{D_t^j Z_t \mathfrak{D}_t^m(\mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta)}{(\alpha' - \beta')^2} d\alpha' d\beta' \end{aligned}$$

Computing using (B.13) and (B.18) yields

$$(4.2) \quad \begin{aligned} \frac{d}{dt} F_{j;l,\tilde{l},k}^{(m)}(t) = & \iint \frac{\left(\overline{\mathcal{P} D_t^j \overline{Z}_t} + [D_t, \mathcal{P}] D_t^{j-1} \overline{Z}_t \right) \mathfrak{D}_t^m(\mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta)}{(\alpha' - \beta')^2} d\alpha' d\beta' \\ & + \iint \frac{\overline{\mathcal{P} D_t^{j-1} \overline{Z}_t} \mathfrak{D}_t^{m+1}(\mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta)}{(\alpha' - \beta')^2} d\alpha' d\beta' \\ & + \iint \left(b_{\alpha'} + b_{\beta'} - 2 \frac{b(\alpha') - b(\beta')}{\alpha' - \beta'} \right) \frac{\overline{\mathcal{P} D_t^{j-1} \overline{Z}_t} \mathfrak{D}_t^m(\mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta)}{(\alpha' - \beta')^2} d\alpha' d\beta' \\ & + \iint \left(b_{\alpha'} + b_{\beta'} - 2 \frac{b(\alpha') - b(\beta')}{\alpha' - \beta'} \right) \frac{D_t^{j+1} Z_t \mathfrak{D}_t^m(\mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta)}{(\alpha' - \beta')^2} d\alpha' d\beta' \\ & + \iint \left(b_{\alpha'} + b_{\beta'} - 2 \frac{b(\alpha') - b(\beta')}{\alpha' - \beta'} \right) \frac{D_t^j Z_t \mathfrak{D}_t^{m+1}(\mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta)}{(\alpha' - \beta')^2} d\alpha' d\beta' \\ & + \iint \left(\partial_{\alpha'} D_t b + \partial_{\beta'} D_t b - 2 \frac{D_t b(\alpha') - D_t b(\beta')}{\alpha' - \beta'} \right) \frac{D_t^j Z_t \mathfrak{D}_t^m(\mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta)}{(\alpha' - \beta')^2} d\alpha' d\beta' \\ & + \iint \left(6 \frac{(b(\alpha') - b(\beta'))^2}{(\alpha' - \beta')^2} - 4(b_{\alpha'} + b_{\beta'}) \frac{(b(\alpha') - b(\beta'))}{(\alpha' - \beta')} + 2b_{\alpha'} b_{\beta'} \right) \frac{D_t^j Z_t \mathfrak{D}_t^m(\mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta)}{(\alpha' - \beta')^2} d\alpha' d\beta'; \end{aligned}$$

in the sum $R_{j;l,\tilde{l},k}^{(m)}(t) + \frac{d}{dt} F_{j;l,\tilde{l},k}^{(m)}(t)$ those terms containing the factors $D_t^{j+1} Z_t$ and $\overline{\mathcal{P} D_t^j \overline{Z}_t}$ are all cancelled out.

We also need the following correcting term for $\frac{1}{4\pi} \overline{R_{0;1,\bar{1},1}^{(0)}}$ in $R_{IC,2}$, cf. (2.36)-(2.32). Define

$$(4.3) \quad D_2(t) = \frac{1}{4\pi} \iint (\mathbb{H}b_{\alpha'}) \frac{\bar{\theta} \mathfrak{D}_t \theta \mathfrak{D}_t \bar{\theta} \mathfrak{D}_t \theta}{(\alpha' - \beta')^2} d\alpha' d\beta', \quad D_3(t) = D_4(t) = 0;$$

we have by (B.13), (B.18),

$$(4.4) \quad \begin{aligned} \frac{d}{dt} D_2(t) &= \frac{1}{4\pi} \iint \partial_{\alpha'} D_t \mathbb{H}b \frac{\bar{\theta} \mathfrak{D}_t \theta \mathfrak{D}_t \bar{\theta} \mathfrak{D}_t \theta}{(\alpha' - \beta')^2} d\alpha' d\beta' + \frac{1}{4\pi} \iint \mathbb{H}b_{\alpha'} \frac{\mathfrak{D}_t (\bar{\theta} \mathfrak{D}_t \theta \mathfrak{D}_t \bar{\theta} \mathfrak{D}_t \theta)}{(\alpha' - \beta')^2} d\alpha' d\beta' \\ &+ \frac{1}{4\pi} \iint \left(b_{\beta'} - 2 \frac{b(\alpha') - b(\beta')}{\alpha' - \beta'} \right) \mathbb{H}b_{\alpha'} \frac{\bar{\theta} \mathfrak{D}_t \theta \mathfrak{D}_t \bar{\theta} \mathfrak{D}_t \theta}{(\alpha' - \beta')^2} d\alpha' d\beta'. \end{aligned}$$

Now define

$$(4.5) \quad \lambda^j := D_{\alpha'} \Theta^{(j)}(\alpha', t) - D_{\beta'} \Theta^{(j)}(\beta', t).$$

We next introduce

$$(4.6) \quad H_j(t) := \frac{1}{4\pi} \operatorname{Re} \left(\iint \frac{\bar{\lambda}^j \lambda^j \theta \bar{\theta} + \bar{\lambda}^j \bar{\lambda}^j \theta \theta}{(\alpha' - \beta')^2} d\alpha' d\beta' - \iint \frac{\mathfrak{D}_t^j \bar{\theta} \mathfrak{D}_t^j \theta \theta \bar{\theta} + \mathfrak{D}_t^j \bar{\theta} \mathfrak{D}_t^j \bar{\theta} \theta \theta}{(\alpha' - \beta')^2} d\alpha' d\beta' \right)$$

to deal with the derivative loss in the term

$$2 \operatorname{Re} \sum_{l=0}^{j-1} \int i \partial_{\alpha'} \overline{\Theta^{(j)}} (\mathbb{P}_H D_t)^{l+1} \mathbb{P}_H \left(G^{(j-l)} - D_t \mathbb{P}_H G^{(j-1-l)} \right) d\alpha' - I_{1,j},$$

and to ensure the control of $E_1(t)E_3(t)$. We compute by (B.13) and using the symmetry (2.37) to get

$$(4.7) \quad \begin{aligned} \frac{d}{dt} H_j &= \frac{1}{\pi} \operatorname{Re} \left(\iint \frac{\overline{D_{\alpha'} \Theta^{(j)}} \mathfrak{D}_t \lambda^j \theta \bar{\theta} + \mathfrak{D}_t \bar{\lambda}^j \theta \theta}{(\alpha' - \beta')^2} d\alpha' d\beta' - \iint D_t^j Z_t \frac{\mathfrak{D}_t^{j+1} \theta \theta \bar{\theta} + \mathfrak{D}_t^{j+1} \bar{\theta} \theta \theta}{(\alpha' - \beta')^2} d\alpha' d\beta' \right) \\ &+ R_{H,j} \end{aligned}$$

where

$$(4.8) \quad \begin{aligned} R_{H,j} &= \frac{1}{2\pi} \operatorname{Re} \iint \frac{(|\lambda^j|^2 - |\mathfrak{D}_t^j \theta|^2) \theta \mathfrak{D}_t \bar{\theta} + ((\bar{\lambda}^j)^2 - (\mathfrak{D}_t^j \bar{\theta})^2) \theta \mathfrak{D}_t \theta}{(\alpha' - \beta')^2} d\alpha' d\beta' \\ &+ \frac{1}{4\pi} \operatorname{Re} \iint \left(b_{\alpha'} + b_{\beta'} - 2 \frac{b(\alpha') - b(\beta')}{\alpha' - \beta'} \right) \frac{(|\lambda^j|^2 - |\mathfrak{D}_t^j \theta|^2) |\theta|^2 + ((\bar{\lambda}^j)^2 - (\mathfrak{D}_t^j \bar{\theta})^2) \theta^2}{(\alpha' - \beta')^2} d\alpha' d\beta'. \end{aligned}$$

Observe that in the sum $I_{1,j} + \frac{d}{dt} H_j$, the term

$$\frac{1}{\pi} \operatorname{Re} \iint D_t^j Z_t \frac{\mathfrak{D}_t^{j+1} \theta \theta \bar{\theta} + \mathfrak{D}_t^{j+1} \bar{\theta} \theta \theta}{(\alpha' - \beta')^2} d\alpha' d\beta'$$

in $I_{1,j}$ is replaced with

$$\frac{1}{\pi} \operatorname{Re} \iint \frac{\overline{D_{\alpha'} \Theta^{(j)}} \mathfrak{D}_t \lambda^j \theta \bar{\theta} + \mathfrak{D}_t \bar{\lambda}^j \theta \theta}{(\alpha' - \beta')^2} d\alpha' d\beta',$$

which matches better with the corresponding term in $2 \operatorname{Re} \int i \partial_{\alpha'} \overline{\Theta^{(j)}} (\mathbb{P}_H D_t)^j \mathbb{P}_H (G^{(1)}) d\alpha'$, and the harmless terms in $R_{H,j}$.

Let

$$(4.9) \quad F_j(t) := \frac{1}{2\pi} \sum_{l=0}^{j-1} \sum_{k=0}^l F_{j;k,\bar{0},j-l-1}^{(l-k)} + \frac{1}{4\pi} \sum_{l=0}^{j-2} \sum_{k=0}^{j-l-2} (-1)^k F_{j;1+l,\bar{k},j-l-2-k}^{(0)} + \frac{1}{4\pi} \sum_{k=0}^{j-1} (-1)^k F_{j;0,\bar{k},j-k-1}^{(0)},$$

$$(4.10) \quad \mathcal{R}_{j;l,\bar{i},k}^{(m)}(t) := R_{j;l,\bar{i},k}^{(m)}(t) + \frac{d}{dt} F_{j;l,\bar{i},k}^{(m)}(t);$$

and define

$$(4.11) \quad \begin{aligned} \mathcal{E}_j(t) &:= \mathfrak{E}_j(t) - \operatorname{Re} F_j(t) + \operatorname{Re} D_j(t) - H_j(t) \\ &= E_j(t) - \operatorname{Re} \left(\int i \partial_{\alpha'} \Theta^{(j)} \overline{\mathbb{P}_H G^{(j)}} d\alpha' + C_{1,j}(t) + C_{2,j}(t) + F_j(t) - D_j(t) + H_j(t) \right). \end{aligned}$$

By Theorem 2.8 and (4.10), we have

$$(4.12) \quad \frac{d}{dt} \mathcal{E}_j(t) = \mathcal{R}_j(t)$$

where

$$(4.13) \quad \begin{aligned} \mathcal{R}_j(t) &= \left(2 \operatorname{Re} \sum_{l=0}^{j-1} \int i \partial_{\alpha'} \overline{\Theta^{(j)}} (\mathbb{P}_H D_t)^{l+1} \mathbb{P}_H \left(G^{(j-l)} - D_t \mathbb{P}_H G^{(j-1-l)} \right) d\alpha' - I_{1,j} - \frac{d}{dt} H_j \right) \\ &\quad + \left(\operatorname{Re} \int i \partial_{\alpha'} \overline{\Theta^{(j)}} \mathbb{P}_H \left(G^{(j+1)} - D_t \mathbb{P}_H G^{(j)} \right) d\alpha' - I_{2,j} \right) + \mathcal{R}_{IC,j} + \operatorname{Re} \frac{d}{dt} D_j(t), \end{aligned}$$

and

$$(4.14) \quad \mathcal{R}_{IC,j} := R_{IC,j} - \frac{d}{dt} \operatorname{Re} F_j(t)$$

is as given in (2.36), except that the quantities $R_{j;k,\bar{0},j-l-1}^{(l-k)}$, $R_{j;1+l,\bar{k},j-l-2-k}^{(0)}$ and $R_{j;0,\bar{k},j-k-1}^{(0)}$ are replaced by $\mathcal{R}_{j;k,\bar{0},j-l-1}^{(l-k)}$, $\mathcal{R}_{j;1+l,\bar{k},j-l-2-k}^{(0)}$ and $\mathcal{R}_{j;0,\bar{k},j-k-1}^{(0)}$ respectively.

4.2. The main components in the proof of Theorem 3.1. We are now ready to prove Theorem 3.1. We begin with the following lemmas.

Lemma 4.1. *Let $0 < \delta < 1$. Assume that*

$$(4.15) \quad \left\| \frac{1}{Z_{,\alpha'}} - 1 \right\|_{L^\infty(\mathbb{R})} \leq 1 - \delta.$$

Then for any holomorphic function f , i.e. $\mathbb{P}_H f = f$, we have

$$(4.16) \quad \delta \|f\|_{\dot{H}^{1/2}(\mathbb{R})} \leq \left\| \mathbb{P}_H \left(\frac{f}{\overline{Z}_{,\alpha'}} \right) \right\|_{\dot{H}^{1/2}(\mathbb{R})}.$$

Proof. We know

$$f = \mathbb{P}_H f = \mathbb{P}_H \left(\frac{f}{\overline{Z}_{,\alpha'}} \right) + \mathbb{P}_H \left\{ f \left(1 - \frac{1}{\overline{Z}_{,\alpha'}} \right) \right\}$$

By (C.38),

$$\left\| \mathbb{P}_H \left\{ f \left(1 - \frac{1}{\overline{Z}_{,\alpha'}} \right) \right\} \right\|_{\dot{H}^{1/2}}^2 = \frac{1}{2\pi} \iint \frac{|f(\alpha') - f(\beta')|^2}{(\alpha' - \beta')^2} \left| 1 - \frac{1}{\overline{Z}_{,\alpha'}} \right|^2 d\alpha' d\beta' \leq (1 - \delta)^2 \|f\|_{\dot{H}^{1/2}}^2.$$

This gives (4.16). □

Lemma 4.2 (Sobolev). *Assume that $\left\| 1 - \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{L^\infty} \leq 1$, and $\frac{1}{\overline{Z}_{,\alpha'}}$ is sufficiently smooth. Then*

$$(4.17) \quad \left\| \frac{1}{\overline{Z}_{,\alpha'}} - 1 \right\|_{L^\infty(\mathbb{R})}^2 \leq 18 \left\| \frac{1}{\overline{Z}_{,\alpha'}} - 1 \right\|_{L^2(\mathbb{R})} \left\| D_{\alpha'} \frac{1}{\overline{Z}_{,\alpha'}^2} \right\|_{L^2(\mathbb{R})}.$$

Proof. (4.17) is essentially a Sobolev inequality. By Fundamental Theorem of Calculus and Cauchy-Schwarz inequality, we know for any smooth complex valued function f tending to 1 at infinity,

$$\left| \frac{1}{4} (f^4 - 1) - \frac{1}{3} (f^3 - 1) \right| \leq \int |(f^3 - f^2) \partial_{\alpha'} f| d\alpha' \leq \|f - 1\|_{L^2} \|f^2 \partial_{\alpha'} f\|_{L^2}.$$

On the other hand,

$$\frac{1}{4}(f^4 - 1) - \frac{1}{3}(f^3 - 1) = (f - 1)^2 \left(\frac{1}{4}f^2 + \frac{1}{6}f + \frac{1}{12} \right).$$

It is easy to check that if $|1 - f| \leq 1$, then $\operatorname{Re} \left(\frac{1}{4}f^2 + \frac{1}{6}f + \frac{1}{12} \right) \geq \frac{1}{36}$, which implies

$$\left| \frac{1}{4}(f^4 - 1) - \frac{1}{3}(f^3 - 1) \right| \geq \frac{1}{36} |f - 1|^2.$$

Replacing f by $\frac{1}{Z_{,\alpha'}}$ yields (4.17). \square

Part 1 of Theorem 3.1 follows from the following Propositions.

Proposition 4.3. *Let $0 < \delta < 1$, and $T_0 > 0$. Assume that*

$$(4.18) \quad \sup_{t \in [0, T_0]} \left\| \frac{1}{Z_{,\alpha'}}(t) - 1 \right\|_{L^\infty} \leq 1 - \delta, \quad \sup_{t \in [0, T_0]} L(t) \leq 2\epsilon.$$

There is a constant $0 < \epsilon_0(\delta) \leq 1$, such that for all $0 < \epsilon \leq \epsilon_0(\delta)$, we have 1.

$$(4.19) \quad c_2(\delta)L(t)^2 \leq \sum_{j=2}^4 \mathcal{E}_j(t) \leq c_1L(t)^2, \quad \forall t \in [0, T_0]$$

where c_1 and $c_2(\delta)$ are some positive constants. 2.

$$(4.20) \quad \sum_{j=2}^4 \mathcal{E}_j(t) \leq \sum_{j=2}^4 \mathcal{E}_j(0) + \epsilon^5 t, \quad \forall t \in [0, T_0]$$

Proposition 4.4. *Assume that the assumption of Proposition 4.3 holds. There is a constant $0 < \epsilon_0(\delta) \leq 1$, such that for all $0 < \epsilon \leq \epsilon_0(\delta)$, we have 1.*

$$(4.21) \quad (1 + \delta)^{-1} E_1(t) \leq \left\| \frac{1}{Z_{,\alpha'}}(t) - 1 \right\|_{L^2}^2 + \left\| \Theta^{(2)}(t) \right\|_{\dot{H}^{1/2}}^2 \leq (1 + \delta) E_1(t), \quad \forall t \in [0, T_0];$$

$$(4.22) \quad (1 + \delta)^{-1} E_3(t) \leq \frac{1}{4} \left\| D_{\alpha'} \frac{1}{Z_{,\alpha'}^2}(t) \right\|_{L^2}^2 + \left\| \Theta^{(4)}(t) \right\|_{\dot{H}^{1/2}}^2 \leq (1 + \delta) E_3(t), \quad \forall t \in [0, T_0];$$

$$(4.23) \quad (1 + \delta)^{-1} E_1(t) E_3(t) \leq \mathfrak{E}_1(t) \mathcal{E}_3(t) \leq (1 + \delta) E_1(t) E_3(t), \quad \forall t \in [0, T_0].$$

2. There is a constant $c_3(\delta^{-1}) > 0$, such that

$$(4.24) \quad \mathfrak{E}_1(t) \mathcal{E}_3(t) \leq \mathfrak{E}_1(0) \mathcal{E}_3(0) e^{c_3(\delta^{-1})\epsilon^3 t}, \quad \forall t \in [0, T_0].$$

Let $J \geq 2$ and assume that the initial data satisfies $\left(\bar{Z}_t(0), \frac{1}{Z_{,\alpha'}}(0) - 1 \right) \in \cap_{\frac{1}{2} \leq s \leq J} \dot{H}^s(\mathbb{R}) \times \dot{H}^{s-\frac{1}{2}}(\mathbb{R})$. By the local existence result in [4], we know there is a unique classical solution for the Cauchy problem of the water wave equation (B.1)-(B.2) for some positive time period $[0, T]$, and the solution exists and is as regular as the initial data for as long as $L(t)$ and $\|Z_{,\alpha'}(t)\|_{L^\infty}$ remain finite.

Now assume that the initial data satisfies (3.2). Let $\delta > 0$ be fixed, such that $\left\| 1 - \frac{1}{Z_{,\alpha'}}(0) \right\|_{L^\infty} \leq 1 - 2\delta$, $\epsilon := \max \left\{ 1, \sqrt{\frac{c_1}{c_2(\delta)}} \right\} \epsilon$, so $L(0) \leq \min \left\{ 1, \sqrt{\frac{c_2(\delta)}{c_1}} \right\} \epsilon$. Let $T_0 > 0$ be the maximum time such that the solution satisfies

$$(4.25) \quad \sup_{t \in [0, T_0]} \left\| \frac{1}{Z_{,\alpha'}}(t) - 1 \right\|_{L^\infty} \leq 1 - \delta, \quad \sup_{t \in [0, T_0]} L(t) \leq 2\epsilon.$$

By Proposition 4.4, there is a $\epsilon_0 = \epsilon_0(\delta) > 0$, such that for all $\epsilon \leq \epsilon_0$,

$$(4.26) \quad E_1(t) E_3(t) \leq (1 + \delta)^2 E_1(0) E_3(0) e^{c_3(\delta^{-1})\epsilon^3 t}, \quad \forall 0 \leq t \leq T_0$$

so

$$(4.27) \quad E_1(t)E_3(t) \leq (1+\delta)^2 m_0^2 (1+\delta), \quad \forall \ 0 \leq t \leq T_1 := \min\{T_0, \frac{\delta}{2c_3(\delta-1)\epsilon^3}\}.$$

By Lemma 4.2 and (4.21), (4.22), there is a constant c_0 such that

$$(4.28) \quad \left\| \frac{1}{Z_{\alpha'}}(t) - 1 \right\|_{L^\infty(\mathbb{R})}^4 \leq c_0(1+\delta)^2 E_1(t)E_3(t), \quad \forall \ t \in [0, T_0].$$

Let $m_0^2 = \frac{(1-2\delta)^4}{c_0(1+\delta)}$. Then we have

$$(4.29) \quad \left\| \frac{1}{Z_{\alpha'}}(t) - 1 \right\|_{L^\infty(\mathbb{R})}^4 \leq c_0(1+\delta)^5 m_0^2 = (1+\delta)^4 (1-2\delta)^4 = (1-\delta-2\delta^2)^4, \quad \forall \ t \in [0, T_1].$$

By Proposition 4.3, there is $\epsilon_0 = \epsilon_0(\delta)$, such that for all $\epsilon \leq \epsilon_0$,

$$(4.30) \quad L(t)^2 \leq \frac{1}{c_2(\delta)} \left(\sum_{j=2}^4 \mathcal{E}_j(0) + \epsilon^5 t \right) \leq \epsilon^2 + \frac{1}{c_2(\delta)} \epsilon^5 t \leq 2\epsilon^2, \quad \text{for all } t \in [0, T_2],$$

where $T_2 = \min\{T_0, \frac{c_2(\delta)}{\epsilon^3}\}$. By the maximality of T_0 we must have $\min\{T_1, T_2\} < T_0$, so there is $\mathcal{T}_0 = \mathcal{T}_0(\delta) > 0$, such that $T_0 \geq \frac{\mathcal{T}_0}{\epsilon^3}$. This proves part 1 of Theorem 3.1.

In the remainder of this paper we prove Propositions 4.3 and 4.4.²⁷ We assume that (4.18) holds, namely for some $0 < \delta < 1$, $0 < \epsilon \leq 1$ and $T_0 > 0$,

$$(4.31) \quad \sup_{t \in [0, T_0]} \left\| \frac{1}{Z_{\alpha'}}(t) - 1 \right\|_{L^\infty} \leq 1 - \delta, \quad \sup_{t \in [0, T_0]} L(t) \leq 2\epsilon.$$

We assume $t \in [0, T_0]$ in what follows.

4.3. The proof of Propositions 4.3 and 4.4. From definition (2.7), (2.8), and equations (2.6), (2.3) we know

$$(4.32) \quad \Theta^{(0)} = Q, \quad \Theta^{(1)} = i(Z - \alpha'), \quad \Theta^{(2)} = -i\mathbb{P}_H b = -i\mathbb{P}_H \frac{\overline{Z}_t}{\overline{Z}_{\alpha'}},$$

$$(4.33) \quad D_{\alpha'} \Theta^{(0)} = \overline{Z}_t, \quad D_{\alpha'} \Theta^{(1)} = i \left(1 - \frac{1}{Z_{\alpha'}} \right); \quad \text{and}$$

$$(4.34) \quad \Theta^{(j+2)} = -i\mathbb{P}_H \left(\frac{1}{|Z_{\alpha'}|^2} \partial_{\alpha'} \Theta^{(j)} \right) + \mathbb{P}_H(G^{(j)}), \quad \text{for } j \geq 1.$$

Therefore

$$(4.35) \quad \begin{aligned} E_j(t) &= \operatorname{Re} \left(\int i \partial_{\alpha'} \Theta^{(j+1)} \overline{\Theta^{(j+1)}} d\alpha' - \int i \partial_{\alpha'} \Theta^{(j)} \overline{\Theta^{(j+2)}} d\alpha' \right) \\ &= \operatorname{Re} \left(\int i \partial_{\alpha'} \Theta^{(j+1)} \overline{\Theta^{(j+1)}} d\alpha' + \int |D_{\alpha'} \Theta^{(j)}|^2 d\alpha' + \int i \partial_{\alpha'} \overline{\Theta^{(j)}} \mathbb{P}_H(G^{(j)}) d\alpha' \right). \end{aligned}$$

²⁷We will be brief on straightforward but tedious details for the sake of a concise presentation. We suggest the reader get familiar with the basic tools in Appendices B, C before continuing.

4.3.1. *Quantities controlled by $L(t)$, $\left\|1 - \frac{1}{Z_{,\alpha'}}\right\|_{L^2}$ and $\|Z_t\|_{\dot{H}^{1/2}}$, and $\|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} + \left\|\partial_{\alpha'} \frac{1}{Z_{,\alpha'}}\right\|_{L^2}$.* We begin with deriving some basic estimates for the quantities involved in the proof of Propositions 4.3, 4.4. First note that by assumption (4.31),

$$(4.36) \quad \sup_{t \in [0, T_0]} \left\| \frac{1}{Z_{,\alpha'}}(t) \right\|_{L^\infty} \leq 2, \quad \sup_{t \in [0, T_0]} \|Z_{t,\alpha'}(t)\|_{L^\infty} \leq \delta^{-1};$$

and by interpolation and Sobolev embedding (C.6),

$$(4.37) \quad \|Z_{t,\alpha'}\|_{L^\infty} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \lesssim L(t) \leq \epsilon.$$

We estimate, by (B.1),

$$\bar{Z}_{tt} = -i \frac{A_1 - 1}{Z_{,\alpha'}} + i \left(1 - \frac{1}{Z_{,\alpha'}} \right),$$

(B.2) and (C.17), that

$$(4.38) \quad \|A_1 - 1\|_{L^2} \lesssim \|Z_t\|_{\dot{H}^{1/2}} \|Z_{t,\alpha'}\|_{L^2} \lesssim \epsilon \|Z_t\|_{\dot{H}^{1/2}},$$

$$(4.39) \quad \|Z_{tt}\|_{L^2} \lesssim \left\| 1 - \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|A_1 - 1\|_{L^2} \lesssim \left\| 1 - \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \epsilon \|Z_t\|_{\dot{H}^{1/2}};$$

by (C.18), (C.40), (B.2) that

$$(4.40) \quad A_1 \geq 1, \quad \|A_1 - 1\|_{L^\infty} + \|A_1\|_{\dot{H}^{1/2}} \lesssim \|Z_{t,\alpha'}\|_{L^2}^2 \lesssim \epsilon^2;$$

by (C.3), (B.1) and (4.40),

$$(4.41) \quad \|Z_{tt}\|_{\dot{H}^{1/2}} \lesssim \|Z_{t,\alpha'}\|_{L^2}^2 + \left\| \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \lesssim \epsilon, \quad \|Z_{tt}\|_{L^\infty} \lesssim 1;$$

by (B.6), (B.7), (C.24) and (C.23) that

$$(4.42) \quad \|D_t A_1\|_{L^2} \lesssim \|Z_{t,\alpha'}\|_{L^2} \|Z_{tt}\|_{\dot{H}^{1/2}} + \|Z_{t,\alpha'}\|_{L^2}^2 \|b_{\alpha'}\|_{L^2},$$

$$(4.43) \quad \|b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t\|_{L^2} \lesssim \|Z_{t,\alpha'}\|_{L^2} \left\| \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \lesssim \epsilon^2;$$

this implies that

$$(4.44) \quad \|b_{\alpha'}\|_{L^2} \lesssim \|Z_{t,\alpha'}\|_{L^2} \left\| \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \|Z_{t,\alpha'}\|_{L^2} \lesssim \epsilon, \quad \text{and} \quad \|D_t A_1\|_{L^2} \lesssim \epsilon^2;$$

and by (4.32) and (B.4) that

$$(4.45) \quad \left\| \partial_{\alpha'} \Theta^{(2)} \right\|_{L^2} \lesssim \epsilon, \quad \left\| \mathcal{P} \bar{Z}_t \right\|_{L^2} \lesssim \epsilon^2, \quad \|D_t^2 Z_t\|_{L^2} \lesssim \|Z_{t,\alpha'}\|_{L^2} + \epsilon^2 \lesssim \epsilon.$$

In what follows we pay particular attention to the bounds $\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}$ and $\|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}$ in the estimates for the sake of proving Proposition 4.4. We estimate, by (B.7), and (C.40), (C.18),

$$(4.46) \quad \|b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t\|_{\dot{H}^{1/2}} + \|b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t\|_{L^\infty} \lesssim \|Z_{t,\alpha'}\|_{L^2} \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \lesssim \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2},$$

so by (C.4), (B.8),

$$(4.47) \quad \|b_{\alpha'}\|_{\dot{H}^{1/2}} + \left\| D_t \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \lesssim \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}};$$

this gives, by using (4.32) and (C.4), that

$$(4.48) \quad \left\| \partial_{\alpha'} \Theta^{(2)} \right\|_{\dot{H}^{1/2}} + \left\| D_{\alpha'} \Theta^{(2)} \right\|_{\dot{H}^{1/2}} \lesssim \|b_{\alpha'}\|_{\dot{H}^{1/2}} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \|b_{\alpha'}\|_{L^2} \lesssim \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}};$$

and as a consequence of (4.46) and (4.37), we also have

$$(4.49) \quad \|b_{\alpha'}\|_{L^\infty} \lesssim \|Z_{t,\alpha'}\|_{L^2} \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{L^\infty} \lesssim \epsilon;$$

and we record, by (B.8),

$$(4.50) \quad \left\| D_t \frac{1}{Z_{,\alpha'}} \right\|_{L^\infty} \lesssim \epsilon.$$

Now by (B.2), (C.17), (C.19),

$$(4.51) \quad \|\partial_{\alpha'} A_1\|_{L^2} \lesssim \|Z_{t,\alpha'}\|_{L^2} \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \lesssim \epsilon \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}};$$

and by (B.1),

$$(4.52) \quad \|Z_{tt,\alpha'}\|_{L^2} \lesssim \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{L^2} \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \lesssim \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \epsilon \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}.$$

We write

$$(4.53) \quad [Z_t, b; \overline{Z}_{t,\alpha'}] = - \langle Z_t, b, \overline{Z}_{t,\alpha'} \rangle + \overline{Z}_{t,\alpha'} [Z_t, b; 1],$$

estimating $\|\langle Z_t, b, \overline{Z}_{t,\alpha'} \rangle\|_{\dot{H}^{1/2}}$ by (C.31), and the $\dot{H}^{1/2}$ norm of the second term by (C.4) and (C.23), (C.24) yield

$$(4.54) \quad \|[Z_t, b; \overline{Z}_{t,\alpha'}]\|_{\dot{H}^{1/2}} \lesssim \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \|b_{\alpha'}\|_{L^2} \|Z_{t,\alpha'}\|_{L^2} + \|Z_{t,\alpha'}\|_{L^2} \|Z_{t,\alpha'}\|_{L^2} \|b_{\alpha'}\|_{\dot{H}^{1/2}};$$

from (B.6), further using (C.40) gives

$$(4.55) \quad \|D_t A_1\|_{\dot{H}^{1/2}} \lesssim \|Z_{t,\alpha'}\|_{L^2} \|Z_{tt,\alpha'}\|_{L^2} + \|[Z_t, b; \overline{Z}_{t,\alpha'}]\|_{\dot{H}^{1/2}} \lesssim \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \epsilon \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}};$$

by (B.4) and (C.3), (C.4), this gives

$$(4.56) \quad \|Z_{ttt}\|_{\dot{H}^{1/2}} \lesssim \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}.$$

We also record, by (B.4), (4.46), (4.55), (4.51) and other relevant earlier estimates that

$$(4.57) \quad \|\mathcal{P}\overline{Z}_t\|_{\dot{H}^{1/2}} \lesssim \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \epsilon \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}.$$

By (B.6), (C.18), (C.27), (C.42),

$$(4.58) \quad \|D_t A_1\|_{L^\infty} \lesssim \|Z_{t,\alpha'}\|_{L^2} \|Z_{tt,\alpha'}\|_{L^2} + \|Z_{t,\alpha'}\|_{L^4}^2 \|b_{\alpha'}\|_{L^2} \lesssim \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \epsilon \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}},$$

therefore by (B.4), we also have

$$(4.59) \quad \|\mathcal{P}\overline{Z}_t\|_{L^\infty} \lesssim \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \epsilon \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}},$$

and

$$(4.60) \quad \|Z_{ttt}\|_{L^\infty} \lesssim \|Z_{t,\alpha'}\|_{L^\infty} + \epsilon^2 \lesssim \epsilon.$$

Now by (B.7), (B.34), and (C.17), (C.19), (C.23), (4.47),

$$(4.61) \quad \begin{aligned} & \|D_t(b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t)\|_{L^2} \lesssim \|Z_{tt}\|_{\dot{H}^{1/2}} \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{L^2} \left\| D_t \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \\ & + \|Z_{t,\alpha'}\|_{L^2} \|b_{\alpha'}\|_{L^2} \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \lesssim \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \epsilon \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}, \end{aligned}$$

this gives, by (4.52), (B.17), (B.18), (C.42) and earlier relevant estimates in this section that

$$(4.62) \quad \|\partial_{\alpha'} D_t b\|_{L^2} \lesssim \|D_t(b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t)\|_{L^2} + \|D_t D_{\alpha'} Z_t\|_{L^2} + \|b_{\alpha'}\|_{L^4}^2 \lesssim \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \epsilon \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}.$$

We estimate similarly, by (B.6), (B.34), (B.31), and (C.17), (C.19), (C.23), (C.27), (C.42) and the estimates above in this section to obtain

$$(4.63) \quad \|D_t^2 A_1\|_{L^2} \lesssim \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \epsilon \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}},$$

this in turn gives, by (B.4), (B.21), because $\mathcal{P}\bar{Z}_{tt} = D_t \mathcal{P}\bar{Z}_t + [\mathcal{P}, D_t] \bar{Z}_t$, that

$$(4.64) \quad \|\mathcal{P}\bar{Z}_{tt}\|_{L^2} + \|[\mathcal{P}, D_t] \bar{Z}_t\|_{L^2} \lesssim \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \epsilon \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}},$$

$$(4.65) \quad \|D_t^3 Z_t\|_{L^2} \lesssim \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \epsilon \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}.$$

We compute $D_t^2 \frac{1}{Z_{,\alpha'}}$ starting from (B.8) and use (B.20), (C.42) and the earlier estimates in this section and get

$$(4.66) \quad \left\| D_t^2 \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \lesssim \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}.$$

From (4.109) we estimate by (2.10), (C.29), (C.32) to get

$$(4.67) \quad \left\| \partial_{\alpha'} \Theta^{(3)} \right\|_{L^2} \lesssim \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \|Z_{t,\alpha'}\|_{L^2}^2 \lesssim \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2};$$

we compute by (B.17) and the definition (2.7):

$$(4.68) \quad D_t D_{\alpha'} \Theta^{(2)} = D_{\alpha'} \Theta^{(3)} + D_{\alpha'} [\mathbb{P}_A, b] \partial_{\alpha'} \Theta^{(2)} - D_{\alpha'} Z_t D_{\alpha'} \Theta^{(2)},$$

and using (C.17), (C.20), (C.42) and the earlier estimates we obtain

$$(4.69) \quad \begin{aligned} \|D_t D_{\alpha'} \Theta^{(2)}\|_{L^2} &\lesssim \left\| \partial_{\alpha'} \Theta^{(3)} \right\|_{L^2} + \|b_{\alpha'}\|_{\dot{H}^{1/2}} \left\| \partial_{\alpha'} \Theta^{(2)} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{L^4} \left\| \partial_{\alpha'} \Theta^{(2)} \right\|_{L^4} \\ &\lesssim \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \epsilon \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}. \end{aligned}$$

Now by (B.7), (C.19), (C.20),

$$(4.70) \quad \|\partial_{\alpha'}(b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t)\|_{L^2} \lesssim (\|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} + \|Z_{t,\alpha'}\|_{L^\infty}) \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \lesssim \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2};$$

therefore

$$(4.71) \quad \|\partial_{\alpha'} b_{\alpha'}\|_{L^2} \lesssim \|\partial_{\alpha'}(b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t)\|_{L^2} + \|Z_{t,\alpha'\alpha'}\|_{L^2} + \|Z_{t,\alpha'}\|_{L^\infty} \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2};$$

and we compute $\partial_{\alpha'} D_t \frac{1}{Z_{,\alpha'}}$ by (B.8), and use (4.70), (4.46), (4.37) to obtain

$$(4.72) \quad \left\| \partial_{\alpha'} D_t \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \lesssim \|Z_{t,\alpha'\alpha'}\|_{L^2} + \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}.$$

Now by (B.2), and (C.40),

$$(4.73) \quad \|\partial_{\alpha'} A_1\|_{\dot{H}^{1/2}} \lesssim \|Z_{t,\alpha'}\|_{L^2} \|Z_{t,\alpha'\alpha'}\|_{L^2} \lesssim \epsilon \|Z_{t,\alpha'\alpha'}\|_{L^2};$$

therefore by (B.1), (C.4),

$$\begin{aligned}
(4.74) \quad \|Z_{tt,\alpha'}\|_{\dot{H}^{1/2}} &\lesssim \|\partial_{\alpha'} A_1\|_{\dot{H}^{1/2}} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \|\partial_{\alpha'} A_1\|_{L^2} + \|A_1\|_{L^\infty} \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \\
&\lesssim \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \epsilon \|Z_{t,\alpha'\alpha'}\|_{L^2} + \epsilon^2 \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}.
\end{aligned}$$

Starting from (B.7), we use (B.34) to expand, and use the Sobolev inequality (C.6) to estimate the $\dot{H}^{1/2}$ norm of $\left[\frac{1}{Z_{,\alpha'}}, b; Z_{t,\alpha'} \right]$ and $\left[Z_t, b; \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right]$, we have, by (C.40), (C.23), (C.24), (C.25), (C.21) and (C.6),

$$\begin{aligned}
(4.75) \quad \|D_t(b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t)\|_{\dot{H}^{1/2}} &\lesssim \left\| \partial_{\alpha'} D_t \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \|Z_{t,\alpha'}\|_{L^2} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \|Z_{tt,\alpha'}\|_{L^2} \\
&\quad + \epsilon^2 \left(\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \|Z_{t,\alpha'\alpha'}\|_{L^2} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \right) \\
&\lesssim \epsilon \left(\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \|Z_{t,\alpha'\alpha'}\|_{L^2} \right) + \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2};
\end{aligned}$$

therefore by (B.17), (B.18), (C.3), (C.4),

$$\begin{aligned}
(4.76) \quad \|\partial_{\alpha'} D_t b\|_{\dot{H}^{1/2}} &\lesssim \|D_t(b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t)\|_{\dot{H}^{1/2}} + \|(b_{\alpha'})^2\|_{\dot{H}^{1/2}} + \|D_{\alpha'} Z_{tt}\|_{\dot{H}^{1/2}} + \|(D_{\alpha'} Z_t)^2\|_{\dot{H}^{1/2}} \\
&\lesssim \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \epsilon \|Z_{t,\alpha'\alpha'}\|_{L^2} + \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \epsilon \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}};
\end{aligned}$$

we also have by (C.18), (C.26) that

$$\begin{aligned}
(4.77) \quad \|D_t(b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t)\|_{L^\infty} &\lesssim \left\| \partial_{\alpha'} D_t \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \|Z_{t,\alpha'}\|_{L^2} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \|Z_{tt,\alpha'}\|_{L^2} \\
&\quad + \|b_{\alpha'}\|_{L^\infty} \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \|Z_{t,\alpha'}\|_{L^2} \lesssim \epsilon^2.
\end{aligned}$$

Compute $D_t^2 \frac{1}{Z_{,\alpha'}}$ starting from (B.8), and use (B.20), (C.3), (C.4), (4.46), (4.47), (4.52), (4.61), (4.74) and (4.75), and Sobolev embedding (C.6) to find

$$(4.78) \quad \left\| D_t^2 \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \lesssim \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \epsilon \|Z_{t,\alpha'\alpha'}\|_{L^2} + \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}.$$

Now from (B.6), using (C.19), (C.20), and (C.24), (C.21), (C.25), we have

$$\begin{aligned}
(4.79) \quad \|\partial_{\alpha'} D_t A_1\|_{L^2} &\lesssim \|Z_{tt,\alpha'}\|_{L^2} (\|Z_{t,\alpha'}\|_{L^\infty} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}) \\
&\quad + \|b_{\alpha'}\|_{L^\infty} \|Z_{t,\alpha'}\|_{L^2} \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} + \|b_{\alpha'}\|_{\dot{H}^{1/2}} \|Z_{t,\alpha'}\|_{L^2} \|Z_{t,\alpha'}\|_{L^\infty} \\
&\lesssim \epsilon \left(\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right);
\end{aligned}$$

therefore by (B.4),

$$(4.80) \quad \|Z_{ttt,\alpha'}\|_{L^2} \lesssim \|Z_{t,\alpha'\alpha'}\|_{L^2} + \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \epsilon \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}.$$

We begin with (B.7), and use (B.32) to expand. By (C.17), (C.19), (C.20), (C.23), (C.27), (C.42), (C.25), and (C.21), we have

$$\begin{aligned}
(4.81) \quad & \|D_t^2(b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t)\|_{L^2} \lesssim \left\| D_t^2 \frac{1}{Z_{,\alpha'}} \right\|_{L^2} (\|Z_{t,\alpha'}\|_{L^\infty} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}) \\
& + \left\| D_t \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \|Z_{t,\alpha'}\|_{L^2} \|b_{\alpha'}\|_{L^\infty} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \|Z_{t,\alpha'}\|_{L^2} \|\partial_{\alpha'} D_t b\|_{L^2} \\
& + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \|Z_{t,\alpha'}\|_{L^2} \|b_{\alpha'}\|_{L^4}^2 + \left\| D_t \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \|Z_{tt,\alpha'}\|_{L^2} \\
& + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \|Z_{tt,\alpha'}\|_{L^2} \|b_{\alpha'}\|_{L^2} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \|Z_{ttt}\|_{\dot{H}^{1/2}} \\
& \lesssim \epsilon \left(\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \|Z_{t,\alpha'\alpha'}\|_{L^2} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right);
\end{aligned}$$

therefore by (B.18), (B.17),

$$(4.82) \quad \|\partial_{\alpha'} D_t^2 b\|_{L^2} \lesssim \|Z_{t,\alpha'\alpha'}\|_{L^2} + \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \epsilon \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}.$$

Now begin with (B.8), and use (B.17), we get

$$\begin{aligned}
(4.83) \quad & \left\| D_t^3 \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \lesssim \|b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t\|_{L^\infty} \left\| D_t^2 \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|D_t^2(b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t)\|_{L^2} \\
& + \|D_t(b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t)\|_{L^2} \left\| D_t \frac{1}{Z_{,\alpha'}} \right\|_{L^\infty} + \|D_{\alpha'} Z_t\|_{L^\infty} \left\| D_t^2 \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|D_t^2 D_{\alpha'} Z_t\|_{L^2} \\
& + \|D_t D_{\alpha'} Z_t\|_{L^2} \left\| D_t \frac{1}{Z_{,\alpha'}} \right\|_{L^\infty} \lesssim \|Z_{t,\alpha'\alpha'}\|_{L^2} + \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \epsilon \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}.
\end{aligned}$$

We compute $D_t^2 A_1$ by (B.6), (B.34), (B.31), estimating the $\dot{H}^{1/2}$ norms of $[Z_{tt}, b; \overline{Z}_{t,\alpha'}]$, $[Z_t, b; \overline{Z}_{tt,\alpha'}]$ and $D_t [Z_t, b; \overline{Z}_{t,\alpha'}]$ by the Sobolev inequality (C.6), and estimating the $\dot{H}^{1/2}$ norms of the remaining commutators in the expansion of $D_t^2 A_1$ by (C.40). We get, by further using (C.23), (C.24), (C.25), (C.21), (C.27),

$$(4.84) \quad \|D_t^2 A_1\|_{\dot{H}^{1/2}} \lesssim \epsilon \left(\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \|Z_{t,\alpha'\alpha'}\|_{L^2} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right);$$

this gives, by (B.1) and the estimates above in this section,

$$(4.85) \quad \|D_t^2 Z_{tt}\|_{\dot{H}^{1/2}} \lesssim \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \epsilon \left(\|Z_{t,\alpha'\alpha'}\|_{L^2} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right).$$

We also have, by (C.18) and (C.27) that

$$(4.86) \quad \|D_t^2 A_1\|_{L^\infty} \lesssim \epsilon^2.$$

We compute $D_t^3 A_1$ by (B.6), (B.32), (B.31), and obtain, by using the inequalities in Appendix C,

$$(4.87) \quad \|D_t^3 A_1\|_{L^2} \lesssim \epsilon \left(\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \|Z_{t,\alpha'\alpha'}\|_{L^2} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right);$$

therefore by (B.4),

$$(4.88) \quad \|D_t^3 Z_{tt}\|_{L^2} \lesssim \|Z_{t,\alpha'\alpha'}\|_{L^2} + \epsilon \left(\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right).$$

We note that (4.81) also gives

$$(4.89) \quad \|D_t^2(b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t)\|_{L^2} \lesssim \epsilon \left(\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right);$$

(4.89) is needed for the sake of estimating $\frac{d}{dt} \mathcal{E}_3(t)$ and proving (4.24). We claim that a similar estimate also holds for $\|D_t^3 A_1\|_{L^2}$, that is

$$(4.90) \quad \|D_t^3 A_1\|_{L^2} \lesssim \epsilon \left(\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right);$$

expanding $D_t^3 A_1$ via (B.6), (B.32), (B.31), (4.90) can be checked straightforwardly using the inequalities in Appendix C, the only term in the expansion that needs some extra explanation is the term $[Z_t, D_t^2 b; \overline{Z}_{t,\alpha'}]$. Observe that by (B.17), (B.18),

$$(4.91) \quad \begin{aligned} \partial_{\alpha'} D_t^2 b &= D_t^2 (b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t) + 3b_{\alpha'} \partial_{\alpha'} D_t b - 2(b_{\alpha'})^3 \\ &\quad - 2 \operatorname{Re} (3D_{\alpha'} Z_t D_{\alpha'} Z_{tt} - 2(D_{\alpha'} Z_t)^3) - 2 \operatorname{Re} \left(Z_{ttt} \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right) + 2 \operatorname{Re} \partial_{\alpha'} \frac{Z_{ttt}}{Z_{,\alpha'}}, \end{aligned}$$

so we have, by (4.89), (4.62), (4.49), (4.47), (4.44), (C.42), (4.37), (4.52), (4.60),

$$(4.92) \quad \left\| \partial_{\alpha'} \left(D_t^2 b - 2 \operatorname{Re} \frac{Z_{ttt}}{Z_{,\alpha'}} \right) \right\|_{L^2} \lesssim \epsilon \left(\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right).$$

We write

$$(4.93) \quad [Z_t, D_t^2 b; \overline{Z}_{t,\alpha'}] = \left[Z_t, D_t^2 b - 2 \operatorname{Re} \frac{Z_{ttt}}{Z_{,\alpha'}}; \overline{Z}_{t,\alpha'} \right] + \left[Z_t, 2 \operatorname{Re} \frac{Z_{ttt}}{Z_{,\alpha'}}; \overline{Z}_{t,\alpha'} \right]$$

and apply (C.23) to the first term and (C.24) to the second term, we get, by further using (C.4),

$$(4.94) \quad \|[Z_t, D_t^2 b; \overline{Z}_{t,\alpha'}]\|_{L^2} \lesssim \epsilon^2 \left(\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right).$$

This proves (4.90). By (B.4), (B.21), (B.24), (4.89), (4.90) and the earlier estimates in this section, we conclude

$$(4.95) \quad \|\mathcal{P} D_t^2 \overline{Z}_t\|_{L^2} + \|[D_t, \mathcal{P}] \overline{Z}_{tt}\|_{L^2} + \|[D_t^2, \mathcal{P}] \overline{Z}_t\|_{L^2} \lesssim \epsilon \left(\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right).$$

Finally we consider $\mathcal{P} D_t^3 \overline{Z}_t$. We write it as

$$\mathcal{P} D_t^3 \overline{Z}_t = D_t^3 \mathcal{P} \overline{Z}_t - [D_t, \mathcal{P}] D_t^2 \overline{Z}_t - D_t [D_t, \mathcal{P}] D_t \overline{Z}_t - D_t^2 [D_t, \mathcal{P}] \overline{Z}_t$$

and use (B.4), (B.21) to compute. By the earlier estimates in this section, we have

$$(4.96) \quad \|[D_t, \mathcal{P}] \overline{Z}_{ttt}\|_{L^2} + \|D_t [D_t, \mathcal{P}] \overline{Z}_{tt}\|_{L^2} + \|D_t^2 [D_t, \mathcal{P}] \overline{Z}_t\|_{L^2} \lesssim \epsilon^2;$$

we compute $D_t^3(b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t)$ from (B.7), expanding by (B.32), (B.31). We have, by (C.17), (C.19), (C.20), (C.23), (C.24), (C.22), (C.21), (C.27),

$$(4.97) \quad \|D_t^3(b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t)\|_{L^2} \lesssim \epsilon^2.$$

we compute by (B.18),

$$(4.98) \quad \partial_{\alpha'} D_t^3 b = D_t \partial_{\alpha'} D_t^2 b + b_{\alpha'} \partial_{\alpha'} D_t^2 b$$

and use (B.17), (B.18), and (4.91) to expand, by (C.42) and earlier estimates in this section, we have

$$(4.99) \quad \left\| \partial_{\alpha'} \left(D_t^3 b - 2 \operatorname{Re} \frac{D_t^3 Z_t}{Z_{,\alpha'}} \right) \right\|_{L^2} \lesssim \epsilon^2;$$

now we compute $D_t^4 A_1$ from (B.6), expanding by (B.32), (B.31). Using a similar argument as that of (4.93) for the term $[Z_t, D_t^3 b; \overline{Z}_{t,\alpha'}]$ and use the inequalities in Appendix C to estimate the remaining terms in $D_t^4 A_1$ we get

$$(4.100) \quad \|D_t^4 A_1\|_{L^2} \lesssim \epsilon^2.$$

This gives that

$$(4.101) \quad \|\mathcal{P} D_t^3 \overline{Z}_t\|_{L^2} \lesssim \epsilon^2.$$

4.3.2. *The estimates for $E_1(t)$, $\mathfrak{E}_1(t)$ and $E_j(t)$, $\mathcal{E}_j(t)$, $j \geq 2$.* In this section we prove the inequalities (4.19) and (4.21), (4.22), (4.23) of Propositions 4.3, 4.4.

Step 1. We begin with $E_1(t)$. From (4.35), (4.33),

$$E_1(t) = \|\Theta^{(2)}\|_{\dot{H}^{1/2}}^2 + \left\|1 - \frac{1}{Z_{,\alpha'}}\right\|_{L^2}^2 + \int i \partial_{\alpha'} \overline{\Theta^{(1)}} \mathbb{P}_H(G^{(1)}) d\alpha',$$

and by Proposition 2.2, and (C.29),

$$(4.102) \quad \left| \int i \partial_{\alpha'} \overline{\Theta^{(1)}} \mathbb{P}_H(G^{(1)}) d\alpha' \right| \lesssim \left\|1 - \frac{1}{Z_{,\alpha'}}\right\|_{L^2}^2 \|Z_{t,\alpha'}\|_{L^2}^2 \lesssim \epsilon^2 \left\|1 - \frac{1}{Z_{,\alpha'}}\right\|_{L^2}^2,$$

so there is $\epsilon_0 = \epsilon_0(\delta) > 0$, such that for all $0 < \epsilon \leq \epsilon_0$, (4.21) holds. We note that by Lemma 4.1,

$$(4.103) \quad \delta \|Z_t\|_{\dot{H}^{1/2}} \leq \|\Theta^{(2)}\|_{\dot{H}^{1/2}}.$$

Now we consider $\mathfrak{E}_1(t)$. We estimate, by (C.29), (C.30) and (4.39), (4.45), (4.103) that

$$\begin{aligned} |C_{1,1}(t)| + |C_{2,1}(t)| &\lesssim \|Z_{tt}\|_{L^2}^2 \|Z_{t,\alpha'}\|_{L^2}^2 + \|Z_{ttt}\|_{L^2} \|Z_{t,\alpha'}\|_{L^2} \|Z_t\|_{\dot{H}^{1/2}}^2 \\ &\lesssim \frac{\epsilon^2}{\delta^2} \left(\|\Theta^{(2)}\|_{\dot{H}^{1/2}}^2 + \left\|1 - \frac{1}{Z_{,\alpha'}}\right\|_{L^2}^2 \right), \end{aligned}$$

so there is $\epsilon_0 = \epsilon_0(\delta) > 0$, such that for all $0 < \epsilon \leq \epsilon_0$,

$$(4.104) \quad (1 + \delta)^{-1/2} E_1(t) \leq \mathfrak{E}_1(t) \leq (1 + \delta)^{1/2} E_1(t).$$

Step 2. We next consider $E_2(t)$ and $\mathcal{E}_2(t)$. By (4.35),

$$(4.105) \quad E_2(t) = \|\Theta^{(3)}\|_{\dot{H}^{1/2}}^2 + \|D_{\alpha'} \Theta^{(2)}\|_{L^2}^2 + \int i \partial_{\alpha'} \overline{\Theta^{(2)}} \mathbb{P}_H(G^{(2)}) d\alpha'.$$

First we have by Lemma 4.1, and (C.1), (C.3)

$$(4.106) \quad \delta \left\|1 - \frac{1}{Z_{,\alpha'}}\right\|_{\dot{H}^{1/2}} \leq \left\| \mathbb{P}_H \left(\frac{1}{\overline{Z}_{,\alpha'}} \left(1 - \frac{1}{Z_{,\alpha'}}\right) \right) \right\|_{\dot{H}^{1/2}} \lesssim \left\|1 - \frac{1}{Z_{,\alpha'}}\right\|_{\dot{H}^{1/2}};$$

and from the identity

$$\frac{\overline{Z}_{t,\alpha'}}{\overline{Z}_{,\alpha'}} = \partial_{\alpha'} \mathbb{P}_H \frac{\overline{Z}_t}{\overline{Z}_{,\alpha'}} - [\mathbb{P}_H, \overline{Z}_t] \partial_{\alpha'} \left(\frac{1}{\overline{Z}_{,\alpha'}} \right) + \left[\mathbb{P}_A, \frac{1}{\overline{Z}_{,\alpha'}} \right] \overline{Z}_{t,\alpha'},$$

and (4.32) (C.17), (C.19), that

$$(4.107) \quad \left\| \frac{\overline{Z}_{t,\alpha'}}{\overline{Z}_{,\alpha'}} - i \partial_{\alpha'} \Theta^{(2)} \right\|_{L^2} \lesssim \left\| \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \|\overline{Z}_{t,\alpha'}\|_{L^2} \lesssim \epsilon \|\overline{Z}_{t,\alpha'}\|_{L^2},$$

so by (4.36), there is $\epsilon_0 = \epsilon_0(\delta) > 0$, such that for $0 < \epsilon \leq \epsilon_0$,

$$(4.108) \quad \delta \|\overline{Z}_{t,\alpha'}\|_{L^2} \lesssim \|\partial_{\alpha'} \Theta^{(2)}\|_{L^2} \lesssim \|\overline{Z}_{t,\alpha'}\|_{L^2}, \quad \delta^2 \|\overline{Z}_{t,\alpha'}\|_{L^2} \lesssim \|D_{\alpha'} \Theta^{(2)}\|_{L^2} \lesssim \|\overline{Z}_{t,\alpha'}\|_{L^2}.$$

Now by (4.35), (4.32),

$$(4.109) \quad \Theta^{(3)} = \mathbb{P}_H \left(\frac{1}{\overline{Z}_{,\alpha'}} \left(1 - \frac{1}{Z_{,\alpha'}} \right) \right) + \mathbb{P}_H(G^{(1)});$$

we estimate, by (2.11), (4.36), (4.106) and (C.3), (C.31), (C.29) to get

$$(4.110) \quad \left\| \mathbb{P}_H(G^{(1)}) \right\|_{\dot{H}^{1/2}} \lesssim \left\| \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \|Z_{t,\alpha'}\|_{L^2}^2 \lesssim \frac{\epsilon^2}{\delta} \left\| \mathbb{P}_H \left(\frac{1}{\overline{Z}_{,\alpha'}} \left(1 - \frac{1}{Z_{,\alpha'}} \right) \right) \right\|_{\dot{H}^{1/2}},$$

and by (2.11)-(2.21), (B.25), (C.15), (B.28), (C.29), (C.30), (C.27), (4.36) to yield

$$(4.111) \quad \begin{aligned} \left\| \mathbb{P}_H(G^{(2)}) \right\|_{L^2} &\lesssim \|b_{\alpha'}\|_{L^2} \|Z_{t,\alpha'}\|_{L^2}^2 + \|Z_{t,\alpha'}\|_{L^2} \left\| \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \left(\|Z_{tt}\|_{\dot{H}^{1/2}} + \left\| \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \right) \\ &+ \|Z_{t,\alpha'}\|_{L^2}^2 \left\| D_t \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \lesssim \frac{\epsilon^2}{\delta^2} \|D_{\alpha'} \Theta^{(2)}\|_{L^2}, \end{aligned}$$

so there is $\epsilon_0 = \epsilon_0(\delta) > 0$, such that for $0 < \epsilon \leq \epsilon_0$,

$$(4.112) \quad (1 + \delta)^{-1} E_2(t) \leq \left\| \mathbb{P}_H \left(\frac{1}{\overline{Z}_{,\alpha'}} \left(1 - \frac{1}{Z_{,\alpha'}} \right) \right) \right\|_{\dot{H}^{1/2}}^2 + \|D_{\alpha'} \Theta^{(2)}\|_{L^2}^2 \leq (1 + \delta) E_2(t).$$

Now we consider the terms $C_{1,2} + C_{2,2} + F_2 + H_2$ in $\mathcal{E}_2(t)$. Observe that those terms with the factor $D_t^3 Z_t$ in $C_{1,2}$, $C_{2,2}$ can be combined with those in F_2 with the factor $\overline{\mathcal{P} D_t \overline{Z}_t}$, and we know

$$(4.113) \quad -D_t^3 Z_t + \overline{\mathcal{P} D_t \overline{Z}_t} = -i \frac{A_1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} Z_{tt}.$$

So by (C.29), (C.30), (C.31), (C.3), (C.27), (C.37), we have

$$(4.114) \quad \begin{aligned} |C_{1,2} + C_{2,2} + F_2 + H_2| &\lesssim \|Z_{ttt}\|_{L^2}^2 \|Z_{t,\alpha'}\|_{L^2}^2 + \|Z_{ttt}\|_{L^2} \|Z_{t,\alpha'}\|_{L^2} \|Z_{tt}\|_{\dot{H}^{1/2}}^2 \\ &+ \|Z_{t,\alpha'}\|_{L^2}^2 \|Z_{tt}\|_{\dot{H}^{1/2}} \left(\|Z_{tt}\|_{\dot{H}^{1/2}} + \left\| \frac{A_1}{|Z_{,\alpha'}|^2} \right\|_{\dot{H}^{1/2}} \right) + \|b_{\alpha'}\|_{L^2} \|Z_{ttt}\|_{L^2} \|Z_{t,\alpha'}\|_{L^2}^2 \\ &+ \|Z_{tt}\|_{\dot{H}^{1/2}}^4 + \|D_{\alpha'} \Theta^{(2)}\|_{L^2}^2 \|Z_{t,\alpha'}\|_{L^2}^2 \\ &\lesssim \frac{\epsilon^2}{\delta^4} \left(\|D_{\alpha'} \Theta^{(2)}\|_{L^2}^2 + \left\| \mathbb{P}_H \left(\frac{1}{\overline{Z}_{,\alpha'}} \left(1 - \frac{1}{Z_{,\alpha'}} \right) \right) \right\|_{\dot{H}^{1/2}}^2 \right); \end{aligned}$$

we also have by (C.36),

$$(4.115) \quad |D_2(t)| \lesssim \|b_{\alpha'}\|_{L^2} \|Z_{t,\alpha'}\|_{L^2} \|Z_{tt}\|_{\dot{H}^{1/2}}^2 \lesssim \frac{\epsilon^2}{\delta^4} \left(\|D_{\alpha'} \Theta^{(2)}\|_{L^2}^2 + \left\| \mathbb{P}_H \left(\frac{1}{\overline{Z}_{,\alpha'}} \left(1 - \frac{1}{Z_{,\alpha'}} \right) \right) \right\|_{\dot{H}^{1/2}}^2 \right).$$

By (4.106), (4.108), there is $\epsilon_0 = \epsilon_0(\delta) > 0$, such that for $0 < \epsilon \leq \epsilon_0$,

$$(4.116) \quad c_2(\delta) \left(\left\| 1 - \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}}^2 + \|Z_{t,\alpha'}\|_{L^2}^2 \right) \leq \mathcal{E}_2(t) \leq c_1 \left(\left\| 1 - \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}}^2 + \|Z_{t,\alpha'}\|_{L^2}^2 \right).$$

for some constants $c_1 > 0$ and $c_2(\delta) > 0$.

Step 3. We now study $E_3(t)$ and $\mathcal{E}_3(t)$. By (4.32), (4.34), (4.35), we know

$$(4.117) \quad E_3(t) = \left\| \Theta^{(4)} \right\|_{\dot{H}^{1/2}}^2 + \|D_{\alpha'} \Theta^{(3)}\|_{L^2}^2 + \int i \partial_{\alpha'} \overline{\Theta}^{(3)} \mathbb{P}_H(G^{(3)}) d\alpha',$$

where

$$\Theta^{(4)} = -\mathbb{P}_H \left(\frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \mathbb{P}_H \frac{\overline{Z}_t}{\overline{Z}_{,\alpha'}} \right) + \mathbb{P}_H(G^{(2)}), \quad D_{\alpha'} \Theta^{(3)} = -D_{\alpha'} \mathbb{P}_H \left(\frac{1}{|Z_{,\alpha'}|^2} \right) + D_{\alpha'} \mathbb{P}_H(G^{(1)}).$$

We compute

$$(4.118) \quad \begin{aligned} D_{\alpha'} \mathbb{P}_H \left(\frac{1}{|Z_{,\alpha'}|^2} \right) - \frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} &= \\ &= \left[\frac{1}{Z_{,\alpha'}}, \mathbb{P}_H \right] \partial_{\alpha'} \frac{1}{|Z_{,\alpha'}|^2} + \left[\mathbb{P}_H, \frac{1}{Z_{,\alpha'}^2} \right] \partial_{\alpha'} \frac{1}{\overline{Z}_{,\alpha'}} - \left[\mathbb{P}_A, \frac{1}{|Z_{,\alpha'}|^2} \right] \partial_{\alpha'} \frac{1}{Z_{,\alpha'}}, \end{aligned}$$

so by (C.17), (C.3), and (4.36),

$$(4.119) \quad \left\| D_{\alpha'} \mathbb{P}_H \left(\frac{1}{|Z_{,\alpha'}|^2} \right) - \frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \lesssim \left\| \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \lesssim \frac{\epsilon}{\delta^2} \left\| \frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}.$$

We estimate by (2.10), (C.32) and (C.29), and get

$$(4.120) \quad \left\| \partial_{\alpha'} \mathbb{P}_H(G^{(1)}) \right\|_{L^2} \lesssim \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \|Z_{t,\alpha'}\|_{L^2}^2 \lesssim \frac{\epsilon^2}{\delta^2} \left\| \frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}.$$

Now

$$(4.121) \quad \mathbb{P}_H \frac{\overline{Z}_{t,\alpha'}}{\overline{Z}_{,\alpha'}} = \partial_{\alpha'} \mathbb{P}_H \frac{\overline{Z}_t}{\overline{Z}_{,\alpha'}} - [\mathbb{P}_H, \overline{Z}_t] \partial_{\alpha'} \frac{1}{\overline{Z}_{,\alpha'}},$$

and by Lemma 4.1, (C.40) and (C.5),

$$(4.122) \quad \begin{aligned} \delta \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} &\leq \left\| \mathbb{P}_H \frac{\overline{Z}_{t,\alpha'}}{\overline{Z}_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \lesssim \left\| \partial_{\alpha'} \mathbb{P}_H \frac{\overline{Z}_t}{\overline{Z}_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \|Z_{t,\alpha'}\|_{L^2} \left\| \partial_{\alpha'} \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{L^2} \\ &\lesssim \frac{1}{\delta} \left(\left\| D_{\alpha'} \mathbb{P}_H \frac{\overline{Z}_t}{\overline{Z}_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \left\| \partial_{\alpha'} \mathbb{P}_H \frac{\overline{Z}_t}{\overline{Z}_{,\alpha'}} \right\|_{L^2} \right) + \|Z_{t,\alpha'}\|_{L^2} \left\| \partial_{\alpha'} \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{L^2}; \end{aligned}$$

using Lemma 4.1 again gives

$$(4.123) \quad \delta \left\| D_{\alpha'} \mathbb{P}_H \frac{\overline{Z}_t}{\overline{Z}_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \leq \left\| \mathbb{P}_H \left(\frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \mathbb{P}_H \frac{\overline{Z}_t}{\overline{Z}_{,\alpha'}} \right) \right\|_{\dot{H}^{1/2}},$$

therefore

$$(4.124) \quad \delta^3 \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \lesssim \left\| \mathbb{P}_H \left(\frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \mathbb{P}_H \frac{\overline{Z}_t}{\overline{Z}_{,\alpha'}} \right) \right\|_{\dot{H}^{1/2}} + \epsilon \delta \left\| \partial_{\alpha'} \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{L^2};$$

on the other hand, by (C.4), (4.45), (4.48) we also have

$$(4.125) \quad \left\| \mathbb{P}_H \left(\frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \mathbb{P}_H \frac{\overline{Z}_t}{\overline{Z}_{,\alpha'}} \right) \right\|_{\dot{H}^{1/2}} \lesssim \epsilon \left\| \partial_{\alpha'} \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}.$$

We next estimate $\|\mathbb{P}_H(G^{(2)})\|_{\dot{H}^{1/2}}$. By (2.11)-(2.21), (C.3), (C.31), and (C.29),

$$(4.126) \quad \left\| \mathbb{P}_H \left(G^{(2)} - D_t \mathbb{P}_H(G^{(1)}) \right) \right\|_{\dot{H}^{1/2}} \lesssim \|Z_{t,\alpha'}\|_{L^2} \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \left\| \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \lesssim \epsilon^2 \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2},$$

by (B.25), (B.28), (C.40), (C.32), (C.29), (C.4), (C.30), (C.27),

$$(4.127) \quad \begin{aligned} &\left\| \mathbb{P}_H D_t \mathbb{P}_H(G^{(1)}) \right\|_{\dot{H}^{1/2}} \lesssim \|b_{\alpha'}\|_{L^2} \|Z_{t,\alpha'}\|_{L^2}^2 \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \left\| D_t \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \|Z_{t,\alpha'}\|_{L^2}^2 \\ &+ \left\| D_t \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \|Z_{t,\alpha'}\|_{L^2}^2 \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{L^2} \|Z_{tt}\|_{\dot{H}^{1/2}} \left\| \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \\ &+ \left\| D_t \langle \overline{Z}_t, i \frac{1}{\overline{Z}_{,\alpha'}} \rangle \right\|_{\dot{H}^{1/2}} + \left\| D_t \langle Z_t, -i \frac{1}{Z_{,\alpha'}} \rangle \right\|_{\dot{H}^{1/2}}. \end{aligned}$$

To estimate $\left\| D_t < \overline{Z}_t, i \frac{1}{\overline{Z}_{,\alpha'}}, \overline{Z}_t > \right\|_{\dot{H}^{1/2}}$ and $\left\| D_t < Z_t, -i \frac{1}{\overline{Z}_{,\alpha'}}, \overline{Z}_t > \right\|_{\dot{H}^{1/2}}$, we use (B.28) to expand. We use Sobolev inequality (C.6) to estimate the following $\dot{H}^{1/2}$ norm. By (C.27), (C.42), we have

$$(4.128) \quad \begin{aligned} & \left\| \int \partial_{\beta'} \mathfrak{D}_t \left(\frac{1}{\alpha' - \beta'} \right) \theta \left(\frac{1}{\overline{Z}_{,\alpha'}} - \frac{1}{\overline{Z}_{,\beta'}} \right) d\beta' \right\|_{\dot{H}^{1/2}}^2 \\ & \lesssim \|b_{\alpha'}\|_{L^2} \|Z_{t,\alpha'}\|_{L^2}^2 \|b_{\alpha'}\|_{L^2} \|Z_{t,\alpha'}\|_{L^4}^2 \left\| \partial_{\alpha'} \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{L^2} \lesssim \epsilon^5 \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \left\| \partial_{\alpha'} \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{L^2} \end{aligned}$$

For the $\dot{H}^{1/2}$ norm of the remaining term in $D_t < \overline{Z}_t, i \frac{1}{\overline{Z}_{,\alpha'}}, \overline{Z}_t >$, we use (C.31). We get

$$(4.129) \quad \begin{aligned} \left\| D_t < \overline{Z}_t, i \frac{1}{\overline{Z}_{,\alpha'}}, \overline{Z}_t > \right\|_{\dot{H}^{1/2}} & \lesssim \epsilon^2 \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}^{1/2} \left\| \partial_{\alpha'} \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{L^2}^{1/2} \\ & + \|Z_{t,\alpha'}\|_{L^2} \|Z_{tt}\|_{\dot{H}^{1/2}} \left\| \partial_{\alpha'} \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{L^2}^2 \left\| D_t \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{\dot{H}^{1/2}}. \end{aligned}$$

Therefore

$$(4.130) \quad \left\| \mathbb{P}_H \left(G^{(2)} \right) \right\|_{\dot{H}^{1/2}} \lesssim \epsilon^2 \left\| \partial_{\alpha'} \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{L^2} + \epsilon^2 \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}.$$

We now consider the term $\int i \partial_{\alpha'} \overline{\Theta}^{(3)} \mathbb{P}_H(G^{(3)}) d\alpha'$ in (4.117). The estimate for $\|\mathbb{P}_H(G^{(3)})\|_{L^2}$ is as usual, by using the formula (2.11)-(2.21), the expansions (B.25) and (B.29), (B.34), and the inequalities (C.29), (C.30), (C.19), (C.20), (C.27) and (C.42); observe that by (C.19), the estimates (4.126)-(4.129) for (4.130) can be used to treat the second term in the expansion by (B.25). We have

$$(4.131) \quad \begin{aligned} \left\| \mathbb{P}_H(G^{(3)}) \right\|_{L^2} & \lesssim \epsilon^2 \left\| \partial_{\alpha'} \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{L^2} + \epsilon^2 \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} + \epsilon^2 \|Z_{tt,\alpha'}\|_{L^2} + \epsilon^2 \left\| D_t \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \\ & + \epsilon^2 \left\| D_t^2 \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{L^2} + \epsilon^2 \|b_{\alpha'}\|_{\dot{H}^{1/2}} + \epsilon^2 \|\partial_{\alpha'} D_t b\|_{L^2} + \epsilon^2 \|Z_{ttt}\|_{\dot{H}^{1/2}} \\ & \lesssim \epsilon^2 \left\| \partial_{\alpha'} \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{L^2} + \epsilon^2 \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}; \end{aligned}$$

so there is a $\epsilon_0 = \epsilon(\delta) > 0$, such that for all $0 < \epsilon \leq \epsilon_0$, the inequality (4.22) in Proposition 4.4 holds. Finally we can estimate the correcting terms $C_{1,3} + C_{2,3} + F_3 + H_3$ as in **Step 2** by combining the terms in $C_{1,3}$ and $C_{2,3}$ with the factor $D_t^4 \overline{Z}_t$ with the terms in F_3 with the factor $\overline{\mathcal{P} D_t^2 \overline{Z}_t}$ and use (C.29), (C.30), (C.27), (C.31), (C.4), and (C.42) to obtain

$$(4.132) \quad \begin{aligned} |C_{1,3} + C_{2,3} + F_3 + H_3| & \lesssim \|D_t^2 Z_t\|_{L^2} \|Z_{t,\alpha'}\|_{L^2} \|Z_{ttt}\|_{\dot{H}^{1/2}}^2 + \|D_t^2 Z_t\|_{L^2}^2 \|Z_{tt,\alpha'}\|_{L^2}^2 \\ & + \|D_t^3 Z_t\|_{L^2} \left(\|Z_{t,\alpha'}\|_{L^2}^2 \|D_t^2 Z_t\|_{L^2} + \|Z_{t,\alpha'}\|_{L^2} \|Z_{tt,\alpha'}\|_{L^2} \|D_t^2 Z_t\|_{L^2} + \|Z_{tt,\alpha'}\|_{L^2} \|Z_{tt}\|_{\dot{H}^{1/2}}^2 \right) \\ & + \|D_t^3 Z_t\|_{L^2} \left(\|b_{\alpha'}\|_{L^4} \|D_t^2 Z_t\|_{L^4} \|Z_{t,\alpha'}\|_{L^2}^2 + \|b_{\alpha'}\|_{L^2} \|Z_{t,\alpha'}\|_{L^2} \|Z_{tt,\alpha'}\|_{L^2} \|Z_{tt}\|_{L^\infty} \right) \\ & + \|Z_{t,\alpha'}\|_{L^2}^2 \|D_{\alpha'} \Theta^{(3)}\|_{L^2}^2 + \|Z_{ttt}\|_{\dot{H}^{1/2}} \left(\|Z_{t,\alpha'}\|_{L^2}^2 \|Z_{ttt}\|_{\dot{H}^{1/2}} + \|Z_{t,\alpha'}\|_{L^2} \|Z_{tt,\alpha'}\|_{L^2} \|Z_{tt}\|_{\dot{H}^{1/2}} \right) \\ & + \|Z_{ttt}\|_{\dot{H}^{1/2}} \left\| \partial_{\alpha'} \frac{A_1}{|\overline{Z}_{,\alpha'}|^2} \right\|_{L^2} \left(\|Z_{t,\alpha'}\|_{L^2}^2 \|Z_{ttt}\|_{L^2} + \|Z_{t,\alpha'}\|_{L^2} \|Z_{tt}\|_{\dot{H}^{1/2}}^2 \right) \\ & \lesssim \epsilon^2 \left(\left\| \partial_{\alpha'} \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{L^2}^2 + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}^2 + \|D_{\alpha'} \Theta^{(3)}\|_{L^2}^2 \right). \end{aligned}$$

So there is $\epsilon_0 = \epsilon(\delta) > 0$, such that for all $0 < \epsilon \leq \epsilon_0$,

$$(4.133) \quad (1 + \delta)^{-1/2} E_3(t) \leq \mathcal{E}_3(t) \leq (1 + \delta)^{1/2} E_3(t).$$

This together with (4.104) proves the inequality (4.23) in Proposition 4.4. Moreover, there are constants $c_1 > 0$ and $c_2(\delta) > 0$, such that for all $0 < \epsilon \leq \epsilon_0(\delta)$,

$$(4.134) \quad c_2(\delta) \left(\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}^2 \right) \leq \mathcal{E}_3(t) \leq c_1 \left(\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}^2 \right).$$

We also note that by (4.125), (4.130),

$$(4.135) \quad \left\| \Theta^{(4)} \right\|_{\dot{H}^{1/2}} \lesssim \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \lesssim \epsilon.$$

Because

$$(4.136) \quad D_t \Theta^{(3)} = \Theta^{(4)} + \mathbb{P}_A D_t \Theta^{(3)} = \Theta^{(4)} + [\mathbb{P}_A, b] \partial_{\alpha'} \Theta^{(3)},$$

$$(4.137) \quad D_t^2 \Theta^{(2)} = \Theta^{(4)} + [\mathbb{P}_A, b] \partial_{\alpha'} \Theta^{(3)} + D_t [\mathbb{P}_A, b] \partial_{\alpha'} \Theta^{(2)},$$

using the identity

$$(4.138) \quad [b, b; \partial_{\alpha'} \Theta^{(2)}] = - \langle b, b, \partial_{\alpha'} \Theta^{(2)} \rangle + \partial_{\alpha'} \Theta^{(2)} [b, b; 1],$$

we have, by (C.40), (B.34), (C.31), (C.4), (C.24), (C.23),

$$(4.139) \quad \left\| D_t \Theta^{(3)} \right\|_{\dot{H}^{1/2}} + \left\| D_t^2 \Theta^{(2)} \right\|_{\dot{H}^{1/2}} \lesssim \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \lesssim \epsilon.$$

Step 4. We now consider $E_4(t)$ and $\mathcal{E}_4(t)$. By (4.35),

$$(4.140) \quad E_4(t) = \left\| \Theta^{(5)} \right\|_{\dot{H}^{1/2}}^2 + \left\| D_{\alpha'} \Theta^{(4)} \right\|_{L^2}^2 + \int i \partial_{\alpha'} \bar{\Theta}^{(4)} \mathbb{P}_H(G^{(4)}) d\alpha',$$

and by (4.34), (4.109),

$$(4.141) \quad \begin{aligned} \Theta^{(5)} &= -i \mathbb{P}_H \left(\frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \Theta^{(3)} \right) + \mathbb{P}_H(G^{(3)}) \\ &= i \mathbb{P}_H \left(\frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \mathbb{P}_H \frac{1}{|Z_{,\alpha'}|^2} \right) - i \mathbb{P}_H \left(\frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \mathbb{P}_H(G^{(1)}) \right) + \mathbb{P}_H(G^{(3)}), \end{aligned}$$

$$(4.142) \quad D_{\alpha'} \Theta^{(4)} = -D_{\alpha'} \mathbb{P}_H \left(\frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \mathbb{P}_H \frac{\bar{Z}_t}{\bar{Z}_{,\alpha'}} \right) + D_{\alpha'} \mathbb{P}_H(G^{(2)}).$$

Since

$$(4.143) \quad \begin{aligned} &\mathbb{P}_H \left(\frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \mathbb{P}_H \frac{1}{|Z_{,\alpha'}|^2} \right) - \frac{1}{|Z_{,\alpha'}|^2 \bar{Z}_{,\alpha'}} \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \\ &= - \left[\mathbb{P}_H, \frac{1}{|Z_{,\alpha'}|^2} \right] \partial_{\alpha'} \mathbb{P}_A \frac{1}{|Z_{,\alpha'}|^2} + \left[\mathbb{P}_H, \frac{1}{|Z_{,\alpha'}|^2 Z_{,\alpha'}} \right] \partial_{\alpha'} \frac{1}{\bar{Z}_{,\alpha'}} - \left[\mathbb{P}_A, \frac{1}{|Z_{,\alpha'}|^2 \bar{Z}_{,\alpha'}} \right] \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} =: I, \end{aligned}$$

and by (C.40),

$$(4.144) \quad \|I\|_{\dot{H}^{1/2}} \lesssim \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 \lesssim \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2};$$

by (C.4), (C.5),

$$(4.145) \quad \left\| \frac{1}{|Z_{,\alpha'}|^2 \bar{Z}_{,\alpha'}} \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \lesssim \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2,$$

$$(4.146) \quad \delta^3 \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \lesssim \left\| \frac{1}{|Z_{,\alpha'}|^2 \bar{Z}_{,\alpha'}} \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2;$$

therefore

$$(4.147) \quad \begin{aligned} \delta^3 \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} - \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} &\lesssim \left\| \mathbb{P}_H \left(\frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \mathbb{P}_H \frac{1}{|Z_{,\alpha'}|^2} \right) \right\|_{\dot{H}^{1/2}} \\ &\lesssim \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}. \end{aligned}$$

Also, we know $\mathbb{P}_H b = \mathbb{P}_H \frac{\bar{Z}_t}{Z_{,\alpha'}} = i\Theta^{(2)}$, and

$$(4.148) \quad \begin{aligned} &\partial_{\alpha'} \mathbb{P}_H \left(\frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \mathbb{P}_H \frac{\bar{Z}_t}{Z_{,\alpha'}} \right) - \frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'}^2 \mathbb{P}_H \frac{\bar{Z}_t}{Z_{,\alpha'}} \\ &= \mathbb{P}_H \left(\partial_{\alpha'} \frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \mathbb{P}_H \frac{\bar{Z}_t}{Z_{,\alpha'}} \right) - \left[\mathbb{P}_A, \frac{1}{|Z_{,\alpha'}|^2} \right] \partial_{\alpha'}^2 \mathbb{P}_H \frac{\bar{Z}_t}{Z_{,\alpha'}}, \end{aligned}$$

applying (C.14) on the second term, we get

$$(4.149) \quad \left\| \partial_{\alpha'} \mathbb{P}_H \left(\frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \mathbb{P}_H \frac{\bar{Z}_t}{Z_{,\alpha'}} \right) - \frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'}^2 \mathbb{P}_H \frac{\bar{Z}_t}{Z_{,\alpha'}} \right\|_{L^2} \lesssim \|b_{\alpha'}\|_{L^\infty} \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \lesssim \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2};$$

now since

$$(4.150) \quad \partial_{\alpha'}^2 \mathbb{P}_H \frac{\bar{Z}_t}{Z_{,\alpha'}} - \frac{\bar{Z}_{t,\alpha'\alpha'}}{\bar{Z}_{,\alpha'}} = - \left[\mathbb{P}_A, \frac{1}{\bar{Z}_{,\alpha'}} \right] \partial_{\alpha'} \bar{Z}_{t,\alpha'} + 2 [\mathbb{P}_H, \bar{Z}_{t,\alpha'}] \partial_{\alpha'} \frac{1}{\bar{Z}_{,\alpha'}} + [\mathbb{P}_H, \bar{Z}_t] \partial_{\alpha'}^2 \frac{1}{\bar{Z}_{,\alpha'}},$$

applying (C.17), (C.19), (C.20) yields

$$(4.151) \quad \left\| \partial_{\alpha'}^2 \mathbb{P}_H \frac{\bar{Z}_t}{Z_{,\alpha'}} - \frac{\bar{Z}_{t,\alpha'\alpha'}}{\bar{Z}_{,\alpha'}} \right\|_{L^2} \lesssim \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \lesssim \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2};$$

therefore

$$(4.152) \quad \begin{aligned} \delta^4 \|Z_{t,\alpha'\alpha'}\|_{L^2} - \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} &\lesssim \left\| D_{\alpha'} \mathbb{P}_H \left(\frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \mathbb{P}_H \frac{\bar{Z}_t}{Z_{,\alpha'}} \right) \right\|_{L^2} \\ &\lesssim \|Z_{t,\alpha'\alpha'}\|_{L^2} + \epsilon \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}. \end{aligned}$$

We next consider $\left\| \mathbb{P}_H \left(\frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \mathbb{P}_H (G^{(1)}) \right) \right\|_{\dot{H}^{1/2}}$. By (2.10), and applying (C.4), (4.120), (2.11), (C.27), (C.32), (C.31) gives

$$(4.153) \quad \begin{aligned} &\left\| \mathbb{P}_H \left(\frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \mathbb{P}_H (G^{(1)}) \right) \right\|_{\dot{H}^{1/2}} \lesssim \left\| \partial_{\alpha'} \mathbb{P}_H (G^{(1)}) \right\|_{\dot{H}^{1/2}} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 \|Z_{t,\alpha'}\|_{L^2}^2 \\ &\lesssim \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \|Z_{t,\alpha'}\|_{L^2}^2 + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 \|Z_{t,\alpha'}\|_{L^2}^2 + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \|Z_{t,\alpha'}\|_{L^2} \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \\ &\lesssim \epsilon^2 \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \epsilon^2 \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}. \end{aligned}$$

We now estimate $\|\partial_{\alpha'} \mathbb{P}_H(G^{(2)})\|_{L^2}$. By (2.11)-(2.21), using (B.25), (B.28) to expand, then applying (C.29), (C.32), (C.17), (C.20), (C.27), we get

$$\begin{aligned}
(4.154) \quad & \left\| \partial_{\alpha'} \mathbb{P}_H(G^{(2)}) \right\|_{L^2} \lesssim \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 \|Z_{t,\alpha'}\|_{L^2} + \|b_{\alpha'}\|_{\dot{H}^{1/2}} \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \|Z_{t,\alpha'}\|_{L^2}^2 \\
& + \left\| \partial_{\alpha'} D_t \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \|Z_{t,\alpha'}\|_{L^2}^2 + \left\| D_t \frac{1}{Z_{,\alpha'}} \right\|_{L^\infty} \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \|Z_{t,\alpha'}\|_{L^2}^2 \\
& + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \|Z_{t,\alpha'}\|_{L^2} \|Z_{tt,\alpha'}\|_{L^2} + \|b_{\alpha'}\|_{L^\infty} \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \|Z_{t,\alpha'}\|_{L^2}^2 \\
& \lesssim \epsilon^2 \|Z_{t,\alpha'\alpha'}\|_{L^2} + \epsilon^2 \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \epsilon^2 \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}.
\end{aligned}$$

And we estimate $\|\mathbb{P}_H(G^{(3)})\|_{\dot{H}^{1/2}}$ by (2.11)-(2.21), expanding with (B.25), (B.29) and (B.34). Again the estimate is straightforward but tedious. We use (C.31) to estimate $\|\mathbb{P}_H(G^{(3)} - D_t \mathbb{P}_H G^{(2)})\|_{\dot{H}^{1/2}}$ and the $\dot{H}^{1/2}$ norm of the term

$$\frac{1}{\pi} \int \frac{\mathfrak{D}_t^2 \left((\overline{Z}_t(\alpha') - \overline{Z}_t(\beta'))^2 \left(\frac{1}{\overline{Z}_{,\alpha'}} - \frac{1}{\overline{Z}_{,\beta'}} \right) \right)}{(\alpha' - \beta')^2} d\beta'$$

in the expansion of $D_t^2 < \overline{Z}_t, i \frac{1}{\overline{Z}_{,\alpha'}}, \overline{Z}_t >$; we use Sobolev inequality (C.6) to estimate the $\dot{H}^{1/2}$ norms of the remaining terms in the expansion of $D_t^2 < \overline{Z}_t, i \frac{1}{\overline{Z}_{,\alpha'}}, \overline{Z}_t >$, as well as $\left\| D_t < \overline{Z}_t, i \frac{1}{\overline{Z}_{,\alpha'}}, i \frac{1}{\overline{Z}_{,\alpha'}} > \right\|_{\dot{H}^{1/2}}$ and $\left\| \left[b, b, \partial_{\alpha'} \left\{ \frac{1}{\overline{Z}_{,\alpha'}} < \overline{Z}_t, i \frac{1}{\overline{Z}_{,\alpha'}}, \overline{Z}_t > \right\} \right] \right\|_{\dot{H}^{1/2}}$. Observe that we have (4.129), and the estimate in (4.154) is useful for the estimate of the second term in the expansion of $\mathbb{P}_H D_t \mathbb{P}_H G^{(2)}$ by (B.25). We have

$$\begin{aligned}
(4.155) \quad & \left\| \mathbb{P}_H(G^{(3)}) \right\|_{\dot{H}^{1/2}} \lesssim \epsilon^2 \|Z_{t,\alpha'\alpha'}\|_{L^2} + \epsilon^2 \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \epsilon^2 \left\| \partial_{\alpha'} D_t \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \epsilon^2 \left\| D_t^2 \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \\
& + \epsilon^2 \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \epsilon^2 \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \\
& \lesssim \epsilon^2 \|Z_{t,\alpha'\alpha'}\|_{L^2} + \epsilon^2 \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \epsilon^2 \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \epsilon^2 \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}.
\end{aligned}$$

We estimate $\|\mathbb{P}_H(G^{(4)})\|_{L^2}$ by (2.11)-(2.21), expanding with (B.25)-(B.26), (B.29), (B.32). The estimate is routine, using the inequalities in Appendix C and the estimates in §4.3.1. Going through the terms carefully, we get

$$(4.156) \quad \left\| \mathbb{P}_H(G^{(4)}) \right\|_{L^2} \lesssim \epsilon^2 \|Z_{t,\alpha'\alpha'}\|_{L^2} + \epsilon^2 \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \epsilon^2 \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \epsilon^2 \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}.$$

Finally the correcting terms $C_{1,4} + C_{2,4} + F_4 + H_4$ can be estimated similarly as in Step 2 by combining the terms in $C_{i,4}$ with the factor $D_t^5 Z_t$ with those terms in F_4 with the factor $\overline{\mathcal{P} D_t^3 Z_t}$ and use the equation

$$(4.157) \quad -D_t^5 Z_t + \overline{\mathcal{P} D_t^3 Z_t} = -i \frac{A_1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} D_t^3 Z_t$$

and we have, by applying the inequalities in Appendix C,

$$\begin{aligned}
(4.158) \quad & |C_{1,4} + C_{2,4} + F_4 + H_4| \\
& \lesssim \epsilon^2 \left(\|Z_{t,\alpha'\alpha'}\|_{L^2} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} + \|D_{\alpha'} \Theta^{(4)}\|_{L^2} \right)^2.
\end{aligned}$$

This together with (4.134) shows that there is a $\epsilon_0 = \epsilon_0(\delta) > 0$, such that for all of $0 < \epsilon \leq \epsilon_0$,

$$(4.159) \quad \mathcal{E}_3(t) + \mathcal{E}_4(t) \lesssim \|Z_{t,\alpha'\alpha'}\|_{L^2}^2 + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}}^2 + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}^2 \leq c_2(\delta)^{-1} (\mathcal{E}_3(t) + \mathcal{E}_4(t)),$$

for some constant $c_2(\delta) > 0$. Combining with (4.37), (4.116) proves the inequality (4.19) in Proposition 4.3. We also note that by (4.142), (4.149), (4.151), (4.154), (4.141), (4.147), (4.153), (4.155), (4.109),

$$(4.160) \quad \left\| \partial_{\alpha'} \Theta^{(4)} \right\|_{L^2} + \left\| \Theta^{(5)} \right\|_{\dot{H}^{1/2}} + \left\| \partial_{\alpha'} \Theta^{(3)} \right\|_{\dot{H}^{1/2}} \lesssim \epsilon.$$

Because

$$(4.161) \quad \begin{aligned} \partial_{\alpha'} D_t \Theta^{(3)} &= \partial_{\alpha'} \Theta^{(4)} + \partial_{\alpha'} [\mathbb{P}_A, b] \partial_{\alpha'} \Theta^{(3)}, \\ D_t^2 \Theta^{(3)} &= \Theta^{(5)} + [\mathbb{P}_A, b] \partial_{\alpha'} \Theta^{(4)} + D_t [\mathbb{P}_A, b] \partial_{\alpha'} \Theta^{(3)}, \\ D_t D_{\alpha'} \Theta^{(3)} &= D_{\alpha'} D_t \Theta^{(3)} - D_{\alpha'} Z_t D_{\alpha'} \Theta^{(3)}, \end{aligned}$$

using the identity

$$(4.162) \quad [b, b; \partial_{\alpha'} \Theta^{(3)}] = - \langle b, b, \partial_{\alpha'} \Theta^{(3)} \rangle + \partial_{\alpha'} \Theta^{(3)} [b, b; 1],$$

we have by (C.17), (C.20), (C.40), (B.34), (C.31), (C.4), (C.24), (C.23),

$$(4.163) \quad \left\| \partial_{\alpha'} D_t \Theta^{(3)} \right\|_{L^2} + \left\| D_t^2 \Theta^{(3)} \right\|_{\dot{H}^{1/2}} + \left\| D_t D_{\alpha'} \Theta^{(3)} \right\|_{L^2} \lesssim \epsilon.$$

4.3.3. The estimates for $\frac{d}{dt} \mathfrak{E}_1(t)$, $\frac{d}{dt} \mathcal{E}_j(t)$, $j \geq 2$. In this section we prove the inequalities (4.20) and (4.24). This requires us to estimate $\frac{d}{dt} \mathfrak{E}_1(t)$, $\frac{d}{dt} \mathcal{E}_j(t)$, $2 \leq j \leq 4$. We will use Theorem 2.8 for $\frac{d}{dt} \mathfrak{E}_1(t)$, and (4.12)-(4.13)-(4.14) for $\frac{d}{dt} \mathcal{E}_j(t)$, $2 \leq j \leq 4$. We refer the reader to Appendix D for a list of the quantities controlled by ϵ .

Step 5. In this step we want to use (2.39)-(2.40) to show that for $0 < \epsilon \leq \epsilon_0(\delta)$, where $\epsilon_0(\delta)$ is as in Steps 1-4,

$$(4.164) \quad \frac{d}{dt} \mathfrak{E}_1(t) \lesssim \frac{\epsilon^3}{\delta^3} \mathfrak{E}_1(t), \quad \text{for } t \in [0, T_0].$$

The prove for (4.164) is quite straightforward. Observe that $D_{\alpha'} \Theta^{(1)} = i \left(1 - \frac{1}{Z_{,\alpha'}} \right)$ by (4.33), and by (B.1),

$$(4.165) \quad \overline{Z}_{tt} = i \left(1 - \frac{1}{Z_{,\alpha'}} \right) - i \frac{A_1 - 1}{Z_{,\alpha'}} = D_{\alpha'} \Theta^{(1)} - i \frac{A_1 - 1}{Z_{,\alpha'}};$$

recall the notation (4.5), (2.24), (2.27), we write

$$(4.166) \quad \begin{aligned} & \operatorname{Re} \int i \partial_{\alpha'} \overline{\Theta^{(1)}} \mathbb{P}_H \left(G^{(2)} - D_t \mathbb{P}_H G^{(1)} \right) d\alpha' - I_{2,1} \\ &= \frac{1}{2\pi} \operatorname{Re} \iint \overline{(D_{\alpha'} \Theta^{(1)})(\alpha', t) - Z_{tt}(\alpha', t)} \frac{\mathfrak{D}_t(\theta \bar{\theta}) \mathfrak{D}_t \theta}{(\alpha' - \beta')^2} d\beta' d\alpha' \\ &+ \frac{1}{2\pi} \operatorname{Re} \iint \overline{D_{\alpha'} \Theta^{(1)}(\alpha', t)} \frac{\mathfrak{D}_t(\theta \bar{\theta})(\lambda^1 - \mathfrak{D}_t \theta)}{(\alpha' - \beta')^2} d\beta' d\alpha' \\ &+ \frac{1}{2} \operatorname{Re} \int i \overline{D_{\alpha'} \Theta^{(1)}(\alpha', t)} \left(\langle \overline{Z}_t, i \frac{1 - A_1}{\overline{Z}_{,\alpha'}}, -i \frac{1}{Z_{,\alpha'}} \rangle + \langle i \frac{A_1 - 1}{Z_{,\alpha'}}, Z_t, -i \frac{1}{Z_{,\alpha'}} \rangle \right) d\alpha', \end{aligned}$$

so by (C.29), and the estimates in §4.3.1,

$$\begin{aligned}
(4.167) \quad & \left| \operatorname{Re} \int i \partial_{\alpha'} \overline{\Theta^{(1)}} \mathbb{P}_H \left(G^{(2)} - D_t \mathbb{P}_H G^{(1)} \right) d\alpha' - I_{2,1} \right| \lesssim \|A_1 - 1\|_{L^2} \|Z_{t,\alpha'}\|_{L^2} \|Z_{tt,\alpha'}\|_{L^2} \|Z_{tt}\|_{L^2} \\
& + \|A_1 - 1\|_{L^2} \|Z_{t,\alpha'}\|_{L^2} \left\| 1 - \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \left(\|Z_{tt,\alpha'}\|_{L^2} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \right) \\
& \lesssim \epsilon^3 \left(\|Z_t\|_{\dot{H}^{1/2}} + \left\| 1 - \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \right)^2;
\end{aligned}$$

observe that $D_{\alpha'} \Theta^{(0)} = \overline{Z}_t$, we write

$$\begin{aligned}
(4.168) \quad & 2 \operatorname{Re} \int i \partial_{\alpha'} \overline{\Theta^{(1)}} \mathbb{P}_H D_t \mathbb{P}_H \left(G^{(1)} - D_t \mathbb{P}_H G^{(0)} \right) d\alpha' - I_{1,1} \\
& = \frac{1}{\pi} \operatorname{Re} \int Z_{tt} \left(D_t \int \frac{\mathfrak{D}_t(\theta \bar{\theta}) \theta}{(\alpha' - \beta')^2} d\beta' - \int \frac{\mathfrak{D}_t(\theta \bar{\theta}) \theta}{(\alpha' - \beta')^2} d\beta' \right) d\alpha' \\
& + \frac{1}{\pi} \operatorname{Re} \int \partial_{\alpha'} \overline{\Theta^{(1)}} \left(\mathbb{P}_H D_t \mathbb{P}_H \frac{1}{\overline{Z}_{,\alpha'}} - \frac{1}{\overline{Z}_{,\alpha'}} D_t \right) \int \frac{\mathfrak{D}_t(\theta \bar{\theta}) \theta}{(\alpha' - \beta')^2} d\beta' d\alpha' \\
& + \frac{1}{\pi} \operatorname{Re} \int \left(\overline{D_{\alpha'} \Theta^{(1)}} - Z_{tt} \right) D_t \int \frac{\mathfrak{D}_t(\theta \bar{\theta}) \theta}{(\alpha' - \beta')^2} d\beta' d\alpha' \\
& + \operatorname{Re} \int i \partial_{\alpha'} \overline{\Theta^{(1)}} \mathbb{P}_H D_t \mathbb{P}_H \left\{ \frac{1}{\overline{Z}_{,\alpha'}} \left(\langle \overline{Z}_t, i \frac{1 - A_1}{\overline{Z}_{,\alpha'}}, \overline{Z}_t \rangle + \langle i \frac{A_1 - 1}{\overline{Z}_{,\alpha'}}, Z_t, \overline{Z}_t \rangle \right) \right\} d\alpha';
\end{aligned}$$

now

$$(4.169) \quad \mathbb{P}_H D_t \mathbb{P}_H \left(\frac{1}{\overline{Z}_{,\alpha'}} f \right) - \mathbb{P}_H \left(\frac{1}{\overline{Z}_{,\alpha'}} D_t f \right) = \mathbb{P}_H \left\{ D_t \left(\frac{1}{\overline{Z}_{,\alpha'}} \right) f \right\} - \mathbb{P}_H [b, \mathbb{P}_A] \partial_{\alpha'} \left(\frac{1}{\overline{Z}_{,\alpha'}} f \right),$$

so by (B.28), (B.25), (C.27), (C.29), (C.30), (C.3),

$$(4.170) \quad \left| 2 \operatorname{Re} \int i \partial_{\alpha'} \overline{\Theta^{(1)}} \mathbb{P}_H D_t \mathbb{P}_H \left(G^{(1)} - D_t \mathbb{P}_H G^{(0)} \right) d\alpha' - I_{1,1} \right| \lesssim \frac{\epsilon^3}{\delta} \left(\|Z_t\|_{\dot{H}^{1/2}}^2 + \left\| 1 - \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 \right).$$

Now the estimates for all the terms in $R_{IC,1}$ are straightforward, using (C.27), (C.29), (C.30), (C.35), (C.36), (C.33), (C.46), (C.48) and the estimates in §4.3.1, we have

$$(4.171) \quad |R_{IC,1}| \lesssim \epsilon^3 \left(\|Z_t\|_{\dot{H}^{1/2}} + \left\| 1 - \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \right)^2.$$

This together with (4.167), (4.170), (4.103), (4.104), (4.21) proves (4.164).

In the remainder of this paper we will show that for $0 < \epsilon \leq \epsilon_0(\delta)$, where $\epsilon_0(\delta)$ is as in Steps 2-4,

$$(4.172) \quad \frac{d}{dt} \mathcal{E}_2(t) \lesssim \epsilon^5, \quad \frac{d}{dt} \mathcal{E}_3(t) \lesssim \epsilon^3 \left(\|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}^2 + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 \right), \quad \frac{d}{dt} \mathcal{E}_4(t) \lesssim \epsilon^5.$$

this together with (4.164), (4.134) proves the inequalities (4.20), (4.24) and finishes the proof for Propositions 4.3, 4.4 as well as Theorem 3.1.

Step 6. We begin with those terms in the remainder $\mathcal{R}_{IC,j} + \operatorname{Re} \frac{d}{dt} D_j(t)$, $j \geq 2$, (cf. (4.13)-(4.14)) for which the desired estimates can be directly derived from the inequalities in Appendix C, keep in mind that lower order norms such as $\left\| 1 - \frac{1}{Z_{,\alpha'}} \right\|_{L^2}$ or $\|Z_t\|_{\dot{H}^{1/2}}$ are NOT allowed, as they are not controlled by $L(t)$; and since $\left\| \frac{1}{\overline{Z}_{,\alpha'}} - 1 \right\|_{L^\infty} \leq 1$, $\|Z_{tt}\|_{L^\infty} \lesssim 1$, we can involve $\left\| \frac{1}{\overline{Z}_{,\alpha'}} - 1 \right\|_{L^\infty}$ and $\|Z_{tt}\|_{L^\infty}$ only in sextic or higher order terms, at most once.

We first consider the following type of terms

$$(4.173) \quad M_{1,j} := \iint \left(b_{\alpha'} + b_{\beta'} - 2 \frac{b(\alpha') - b(\beta')}{\alpha' - \beta'} \right) \frac{D_t^j Z_t \mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta}{(\alpha' - \beta')^2} d\alpha' d\beta'$$

where $l + i + k = j$. This type of terms appears in (4.2), (2.36) and (2.32), we use (C.35), (C.36) and the estimates in §4.3.1 to handle $M_{1,j}$.

For $j = 2$, $(l, i, k) = (0, 0, 2), (0, 1, 1)$ and their permutations, so

$$(4.174) \quad |M_{1,2}| \lesssim \|D_t^2 Z_t\|_{L^2} \|b_{\alpha'}\|_{L^\infty} \left(\|Z_{t,\alpha'}\|_{L^2}^2 \|D_t^2 Z_t\|_{L^2} + \|Z_{t,\alpha'}\|_{L^2} \|Z_{tt}\|_{\dot{H}^{1/2}}^2 \right) \lesssim \epsilon^5.$$

For $j = 3$, $(l, i, k) = (0, 0, 3), (0, 1, 2), (1, 1, 1)$ and their permutations, so

$$(4.175) \quad \begin{aligned} |M_{1,3}| &\lesssim \|D_t^3 Z_t\|_{L^2} \|b_{\alpha'}\|_{L^\infty} \left(\|Z_{t,\alpha'}\|_{L^2}^2 \|D_t^3 Z_t\|_{L^2} + \|Z_{t,\alpha'}\|_{L^2} \|Z_{tt,\alpha'}\|_{L^2} \|D_t^2 Z_t\|_{L^2} \right) \\ &\quad + \|D_t^3 Z_t\|_{L^2} \|b_{\alpha'}\|_{L^\infty} \|Z_{tt,\alpha'}\|_{L^2} \|Z_{tt}\|_{\dot{H}^{1/2}}^2 \lesssim \epsilon^3 \left(\|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}^2 + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 \right). \end{aligned}$$

For $j = 4$, $(l, i, k) = (0, 0, 4), (0, 1, 3), (0, 2, 2), (1, 1, 2)$ and their permutations, so a similar argument also gives

$$(4.176) \quad |M_{1,4}| \lesssim \epsilon^5.$$

We next consider terms of the type

$$(4.177) \quad M_{2,j} := \iint \frac{\overline{\mathcal{P} D_t^m \bar{Z}_t} \mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta}{(\alpha' - \beta')^2} d\alpha' d\beta'$$

where $m + l + i + k = 2j - 1$. We discuss the following cases:

1. $m = j - 1$, $l + i + k = j$, as appeared in (4.2);
2. $i = j$, $m + l + k = j - 1$, as appeared in (2.32)-(2.36);
3. $i = 0$, $m + l + k = 2j - 1$, $0 \leq m, l, k \leq j - 1$, as appeared in (2.32)-(2.36);
4. $m = 0$, $l + i + k = 2j - 1$, $0 \leq l, i, k \leq j - 1$, as appeared in (2.32)-(2.36).

Observe that we can use the symmetry (2.37) to rewrite the corresponding terms in (2.32) as (4.177), and vice versa. We use (C.29), (C.30) and the estimates in §4.3.1 to obtain the inequalities (4.178), (4.179), and (4.180) below.

For $j = 2$, we have the following cases: $m = 1$, $(l, i, k) = (0, 0, 2), (0, 1, 1)$ and permutations; $i = 2$, $(m, l, k) = (0, 0, 1)$ and permutations; $i = 0$, $(m, l, k) = (1, 1, 1)$; $m = 0$, $(l, i, k) = (1, 1, 1)$. It is clear that by (C.29), (C.30) and the estimates in §4.3.1 we have

$$(4.178) \quad |M_{2,2}| \lesssim \epsilon^5.$$

For $j = 3$, we have the following cases: $m = 2$, $(l, i, k) = (0, 0, 3), (0, 1, 2), (1, 1, 1)$ and permutations; $i = 3$, $(m, l, k) = (0, 0, 2), (0, 1, 1)$ and permutations; $i = 0$, $(m, l, k) = (1, 2, 2)$ and permutations; $m = 0$, $(l, i, k) = (1, 2, 2)$ and permutations. Observe that for the case where $i = 3$, we need to use the symmetry (2.37) to rewrite $M_{2,3}$ as in (2.32). We have by (C.29), (C.30) and the estimates in §4.3.1,

$$(4.179) \quad |M_{2,3}| \lesssim \epsilon^3 \left(\|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}^2 + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 \right).$$

For $j = 4$, we have the following cases: $m = 3$, $(l, i, k) = (0, 0, 4), (0, 1, 3), (0, 2, 2), (1, 1, 2)$ and permutations; $i = 4$, $(m, l, k) = (0, 0, 3), (0, 1, 2), (1, 1, 1)$ and permutations; $i = 0$, $(m, l, k) = (3, 3, 1), (3, 2, 2)$ and

permutations; $m = 0$, $(l, i, k) = (3, 3, 1), (3, 2, 2)$ and permutations. Again we use the form in (2.32) for the case where $i = 4$. We have quite straightforwardly that

$$(4.180) \quad |M_{2,4}| \lesssim \epsilon^5.$$

Observe that those terms

$$\iint \frac{D_t^j Z_t \mathfrak{D}_t^m \left\{ (\mathcal{P} \mathfrak{D}_t^l \theta) \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta - \mathfrak{D}_t^l \theta \overline{(\mathcal{P} \mathfrak{D}_t^i \theta)} \mathfrak{D}_t^k \theta + \mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} (\mathcal{P} \mathfrak{D}_t^k \theta) \right\}}{(\alpha' - \beta')^2} d\alpha' d\beta'$$

in (2.32), with $m + l + i + k = j - 1$ can be treated exactly in the same way as $M_{2,j}$, we do not specifically go over these terms.

Now we consider the terms $\iint \frac{\overline{[D_t, \mathcal{P}] D_t^{j-1} \bar{Z}_t} \mathfrak{D}_t^m (\mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta)}{(\alpha' - \beta')^2} d\alpha' d\beta'$ in (4.2). We use product rules and complex conjugate to reduce it to the following form

$$(4.181) \quad M_{3,j} := \iint \frac{[D_t, \mathcal{P}] D_t^{j-1} \bar{Z}_t \mathfrak{D}_t^l \bar{\theta} \mathfrak{D}_t^i \theta \mathfrak{D}_t^k \bar{\theta}}{(\alpha' - \beta')^2} d\alpha' d\beta',$$

where $l + i + k = j - 1$. We know by (B.21),

$$(4.182) \quad [D_t, \mathcal{P}] D_t^{j-1} \bar{Z}_t = i \left(\frac{D_t A_1}{A_1} + b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t \right) \frac{A_1}{|Z_{\alpha'}|^2} \partial_{\alpha'} D_t^{j-1} \bar{Z}_t,$$

so $M_{3,j}$ is sextic; and we have, by (C.3) and the estimates in §4.3.1,

$$(4.183) \quad \begin{aligned} |M_{3,j}| &\lesssim \left\| D_t^{j-1} Z_t \right\|_{\dot{H}^{1/2}} \left\| \left(\frac{D_t A_1}{A_1} + b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t \right) \frac{A_1}{|Z_{\alpha'}|^2} \langle D_t^l Z_t, D_t^i \bar{Z}_t, D_t^k Z_t \rangle \right\|_{\dot{H}^{1/2}} \\ &\lesssim \left\| D_t^{j-1} Z_t \right\|_{\dot{H}^{1/2}} (\|D_t A_1\|_{L^\infty} + \|b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t\|_{L^\infty}) \|\langle D_t^l Z_t, D_t^i \bar{Z}_t, D_t^k Z_t \rangle\|_{\dot{H}^{1/2}} \\ &\quad + \left\| D_t^{j-1} Z_t \right\|_{\dot{H}^{1/2}} \left(\left\| \frac{D_t A_1}{|Z_{\alpha'}|^2} \right\|_{\dot{H}^{1/2}} + \left\| \frac{(b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t) A_1}{|Z_{\alpha'}|^2} \right\|_{\dot{H}^{1/2}} \right) \|\langle D_t^l Z_t, D_t^i \bar{Z}_t, D_t^k Z_t \rangle\|_{L^\infty}; \end{aligned}$$

we can also use (C.4) to get

$$(4.184) \quad \begin{aligned} |M_{3,j}| &\lesssim \left\| D_t^{j-1} Z_t \right\|_{\dot{H}^{1/2}} (\|D_t A_1\|_{L^\infty} + \|b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t\|_{L^\infty}) \|\langle D_t^l Z_t, D_t^i \bar{Z}_t, D_t^k Z_t \rangle\|_{\dot{H}^{1/2}} \\ &\quad + \left\| D_t^{j-1} Z_t \right\|_{\dot{H}^{1/2}} \left(\left\| \frac{D_t A_1}{|Z_{\alpha'}|^2} \right\|_{\dot{H}^1} + \left\| \frac{(b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t) A_1}{|Z_{\alpha'}|^2} \right\|_{\dot{H}^1} \right) \|\langle D_t^l Z_t, D_t^i \bar{Z}_t, D_t^k Z_t \rangle\|_{L^2}. \end{aligned}$$

For $j = 2$, we have $(l, i, k) = (0, 0, 1)$ and permutations; for $j = 3$, we have $(l, i, k) = (0, 0, 2), (0, 1, 1)$ and permutations; for $j = 4$, we have $(l, i, k) = (0, 0, 3), (0, 1, 2), (1, 1, 1)$ and permutations. We use (4.183) for $j = 2, 3$ and (4.184) for $j = 4$. By (C.3), (C.31), (C.29), (C.30) and the estimates in §4.3.1 we get

$$(4.185) \quad |M_{3,2}| \lesssim \epsilon^5, \quad |M_{3,3}| \lesssim \epsilon^3 \left(\|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}^2 + \left\| \partial_{\alpha'} \frac{1}{Z_{\alpha'}} \right\|_{L^2}^2 \right), \quad |M_{3,4}| \lesssim \epsilon^6.$$

Next we look at the terms of type

$$(4.186) \quad M_{4,j} := \iint \left(b_{\alpha'} + b_{\beta'} - 2 \frac{b(\alpha') - b(\beta')}{\alpha' - \beta'} \right) \frac{\overline{\mathcal{P} D_t^{j-1} \bar{Z}_t} \mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta}{(\alpha' - \beta')^2} d\alpha' d\beta'$$

and

$$(4.187) \quad M_{5,j} := \iint \left(6 \frac{(b(\alpha') - b(\beta'))^2}{(\alpha' - \beta')^2} - 4(b_{\alpha'} + b_{\beta'}) \frac{(b(\alpha') - b(\beta'))}{(\alpha' - \beta')} + 2b_{\alpha'} b_{\beta'} \right) \frac{D_t^j Z_t \mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta}{(\alpha' - \beta')^2} d\alpha' d\beta'$$

in (4.2), where $l + i + k = j - 1$. Observe that both terms are sextic, and we used product rules to reduce the terms in (4.2) to the forms of (4.186), (4.187). We take advantage of the fact that these terms are sextic and use (C.27), (C.42) to deduce for $m = 4, 5$,

$$(4.188) \quad |M_{m,2}| \lesssim \epsilon^5, \quad |M_{m,3}| \lesssim \epsilon^3 \left(\|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}^2 + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 \right), \quad |M_{m,4}| \lesssim \epsilon^5.$$

Now we consider the term

$$(4.189) \quad M_{6,j} := \iint \frac{D_t^j Z_t [\mathcal{P}, \mathfrak{D}_t^m] (\mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta)}{(\alpha' - \beta')^2} d\alpha' d\beta' = \iint \frac{[\mathfrak{P}, D_t^m] (\mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta) \mathfrak{D}_t^j \bar{\theta}}{(\alpha' - \beta')^2} d\alpha' d\beta'$$

in (2.32), where $m + l + i + k = j - 1$. Here we used the symmetry (2.37) to get the second equality above. We expand $[\mathfrak{P}, D_t^m]$ by (B.21), (B.24), and use (C.29) and the estimates in §4.3.1 to obtain

$$(4.190) \quad |M_{6,2}| \lesssim \epsilon^6, \quad |M_{6,3}| \lesssim \epsilon^4 \left(\|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}^2 + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 \right), \quad |M_{6,4}| \lesssim \epsilon^6.$$

Now we treat the terms

$$(4.191) \quad M_{7,j} := i \iint \left(\partial_{\beta'} \frac{A_1(\beta')}{|Z_{,\beta'}|^2} - 2 \frac{\frac{A_1(\alpha')}{|Z_{,\alpha'}|^2} - \frac{A_1(\beta')}{|Z_{,\beta'}|^2}}{\alpha' - \beta'} \right) \frac{D_t^j Z_t(\alpha') \mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta}{(\alpha' - \beta')^2} d\alpha' d\beta'$$

and

$$(4.192) \quad M_{8,j} := \iint \left(\partial_{\beta'} D_t b(\beta') - 2 \frac{D_t b(\alpha') - D_t b(\beta')}{\alpha' - \beta'} \right) \frac{D_t^j Z_t(\alpha') \mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta}{(\alpha' - \beta')^2} d\alpha' d\beta'$$

where $l + i + k = j - 1$ in (2.32)-(2.36) and (4.2). Here we used the product rules to convert the terms in (2.32) and (4.2) to the forms in (4.191), (4.192). We use (C.46) and (C.48) and the estimates in §4.3.1 to obtain for $m = 7, 8$,

$$(4.193) \quad |M_{m,2}| \lesssim \epsilon^5, \quad |M_{m,3}| \lesssim \epsilon^3 \left(\|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}^2 + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 \right), \quad |M_{m,4}| \lesssim \epsilon^5.$$

We also have the following terms from $R_{0;l,\bar{i},k}^{(0)}$, where $l + i + k = 2j - 1$, $0 \leq l, i, k \leq j - 1$, in (2.32)-(2.36),

$$(4.194) \quad \iint \left(b_{\alpha'} + b_{\beta'} - 2 \frac{b(\alpha') - b(\beta')}{\alpha' - \beta'} \right) \frac{(Z_t \mathfrak{D}_t - Z_{tt}) (\mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta)}{(\alpha' - \beta')^2} d\alpha' d\beta';$$

we use the symmetry (2.37) to rewrite it as

$$(4.195) \quad M_{9,j} := \frac{1}{2} \iint \left(b_{\alpha'} + b_{\beta'} - 2 \frac{b(\alpha') - b(\beta')}{\alpha' - \beta'} \right) \frac{(\bar{\theta} \mathfrak{D}_t - \mathfrak{D}_t \bar{\theta}) (\mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta)}{(\alpha' - \beta')^2} d\alpha' d\beta';$$

and use product rules to expand. By (C.35), (C.36) and the estimates in §4.3.1, we have

$$(4.196) \quad \begin{aligned} |M_{9,3}| &\lesssim \|b_{\alpha'}\|_{L^\infty} \|Z_{t,\alpha'}\|_{L^2} \left(\|D_t^3 Z_t\|_{L^2} \|Z_{tt,\alpha'}\|_{L^2} \|Z_{ttt}\|_{L^2} + \|Z_{ttt}\|_{\dot{H}^{1/2}}^2 \|Z_{ttt}\|_{L^2} \right) \\ &\quad + \|b_{\alpha'}\|_{L^\infty} \|Z_{tt,\alpha'}\|_{L^2}^2 \|Z_{ttt}\|_{L^2}^2 \lesssim \epsilon^3 \left(\|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}^2 + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 \right), \end{aligned}$$

and

$$(4.197) \quad |M_{9,4}| \lesssim \epsilon^5.$$

For $j = 2$, $(l, i, k) = (1, 1, 1)$, we have, by (C.36),

$$(4.198) \quad \left| \iint \left(b_{\alpha'} + b_{\beta'} - 2 \frac{b(\alpha') - b(\beta')}{\alpha' - \beta'} \right) \frac{\bar{\theta} \mathfrak{D}_t (\mathfrak{D}_t \theta \mathfrak{D}_t \bar{\theta} \mathfrak{D}_t \theta)}{(\alpha' - \beta')^2} d\alpha' d\beta' \right| \\ \lesssim \|Z_{ttt}\|_{L^2} \|Z_{t,\alpha'}\|_{L^2} \|Z_{tt}\|_{\dot{H}^{1/2}}^2 \|b_{\alpha'}\|_{L^\infty} \lesssim \epsilon^5;$$

and by (C.37),

$$(4.199) \quad \left| \iint \left(b_{\alpha'} + b_{\beta'} - 2 \frac{b(\alpha') - b(\beta')}{\alpha' - \beta'} \right) \frac{\mathfrak{D}_t \bar{\theta} \mathfrak{D}_t \theta \mathfrak{D}_t \bar{\theta} \mathfrak{D}_t \theta}{(\alpha' - \beta')^2} d\alpha' d\beta' \right| \lesssim \|b_{\alpha'}\|_{L^\infty} \|Z_{tt}\|_{\dot{H}^{1/2}}^4 \lesssim \epsilon^5;$$

so

$$(4.200) \quad |M_{9,2}| \lesssim \epsilon^5.$$

We can estimate the second and third term on the right hand side of (4.4) similarly and obtain

$$(4.201) \quad \left| \iint \mathbb{H} b_{\alpha'} \frac{\mathfrak{D}_t (\bar{\theta} \mathfrak{D}_t \theta \mathfrak{D}_t \bar{\theta} \mathfrak{D}_t \theta)}{(\alpha' - \beta')^2} d\alpha' d\beta' \right| \lesssim \|\mathbb{H} b_{\alpha'}\|_{L^\infty} \|Z_{tt}\|_{\dot{H}^{1/2}}^4 \\ + \|\mathbb{H} b_{\alpha'}\|_{L^2} \|Z_{ttt}\|_{L^\infty} \|Z_{tt}\|_{\dot{H}^{1/2}}^2 \|Z_{t,\alpha'}\|_{L^2} \lesssim \epsilon^5;$$

$$(4.202) \quad \left| \iint \left(b_{\beta'} - 2 \frac{b(\alpha') - b(\beta')}{\alpha' - \beta'} \right) \mathbb{H} b_{\alpha'} \frac{\bar{\theta} \mathfrak{D}_t \theta \mathfrak{D}_t \bar{\theta} \mathfrak{D}_t \theta}{(\alpha' - \beta')^2} d\alpha' d\beta' \right| \\ \lesssim \|\mathbb{H} b_{\alpha'}\|_{L^2} \|b_{\alpha'}\|_{L^\infty} \|Z_{tt}\|_{\dot{H}^{1/2}}^2 \|Z_{t,\alpha'}\|_{L^2} \|Z_{tt}\|_{L^\infty} \lesssim \epsilon^5.$$

Observe that the first two (non-zero) terms in $R_{0;l,\bar{l},k}^{(0)}$, $l+i+k=2j-1$, $0 \leq l, i, k \leq j-1$, cf. (2.32)-(2.36), have been covered in $M_{2,j}$, there is one more term left, which is

$$(4.203) \quad i \iint \left(\partial_{\alpha'} \frac{A_1(\alpha')}{|Z_{,\alpha'}|^2} + \partial_{\beta'} \frac{A_1(\beta')}{|Z_{,\beta'}|^2} - 2 \frac{\frac{A_1(\alpha')}{|Z_{,\alpha'}|^2} - \frac{A_1(\beta')}{|Z_{,\beta'}|^2}}{\alpha' - \beta'} \right) \frac{Z_t \mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta}{(\alpha' - \beta')^2} d\alpha' d\beta'.$$

We use symmetry (2.37) to rewrite it as

$$(4.204) \quad M_{10,j} := \frac{1}{2} i \iint \left(\partial_{\alpha'} \frac{A_1(\alpha')}{|Z_{,\alpha'}|^2} + \partial_{\beta'} \frac{A_1(\beta')}{|Z_{,\beta'}|^2} - 2 \frac{\frac{A_1(\alpha')}{|Z_{,\alpha'}|^2} - \frac{A_1(\beta')}{|Z_{,\beta'}|^2}}{\alpha' - \beta'} \right) \frac{\bar{\theta} \mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta}{(\alpha' - \beta')^2} d\alpha' d\beta'.$$

By (C.27), (C.42) and the estimates in §4.3.1, we have

$$(4.205) \quad |M_{10,3}| \lesssim \epsilon^3 \left(\|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}^2 + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 \right), \quad |M_{10,4}| \lesssim \epsilon^5.$$

For $j = 2$ and $(l, i, k) = (1, 1, 1)$, we have by (C.37) and Hölder's inequality,

$$(4.206) \quad \left| \iint \left(\frac{\frac{A_1(\alpha')}{|Z_{,\alpha'}|^2} - \frac{A_1(\beta')}{|Z_{,\beta'}|^2}}{\alpha' - \beta'} \right) \frac{\bar{\theta} \mathfrak{D}_t \theta \mathfrak{D}_t \bar{\theta} \mathfrak{D}_t \theta}{(\alpha' - \beta')^2} d\alpha' d\beta' \right| \lesssim \|Z_{tt}\|_{\dot{H}^{1/2}}^3 \left\| \frac{A_1}{|Z_{,\alpha'}|^2} \right\|_{\dot{H}^{1/2}} \|Z_{t,\alpha'}\|_{L^\infty} \lesssim \epsilon^5.$$

Step 7. We have treated all the terms in $\mathcal{R}_{IC,j} + \text{Re} \frac{d}{dt} D_j(t)$ but the following two: the first is

$$(4.207) \quad M_{11,j} := \iint \partial_{\alpha'} \left(i \frac{A_1(\alpha')}{|Z_{,\alpha'}|^2} + D_t b(\alpha') \right) \frac{D_t^j Z_t(\alpha') \mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta}{(\alpha' - \beta')^2} d\alpha' d\beta'$$

where $l+i+k=j-1$, $2 \leq j \leq 4$; the second is for $j = 2$ only, from $\text{Re} \left(\frac{1}{4\pi} R_{0;1,\bar{1},1}^{(0)} + \frac{d}{dt} D_2(t) \right)$:

$$(4.208) \quad M_{12,2} := \iint \partial_{\alpha'} \left(i \frac{A_1(\alpha')}{|Z_{,\alpha'}|^2} + D_t \mathbb{H} b(\alpha') \right) \frac{\bar{\theta} \mathfrak{D}_t \theta \mathfrak{D}_t \bar{\theta} \mathfrak{D}_t \theta}{(\alpha' - \beta')^2} d\alpha' d\beta',$$

here we used the symmetry (2.37) to rewrite the term from (2.32), and used product rules to arrive at the term in (4.207).

By (B.39),

$$(4.209) \quad \mathbb{P}_H \left(D_t b + i \frac{A_1}{|Z_{,\alpha'}|^2} - i \right) = i \mathbb{P}_H \left(\frac{A_1 - 1}{|Z_{,\alpha'}|^2} \right) + i \mathbb{P}_H(G^{(1)}) + i [b, \mathbb{P}_H] \partial_{\alpha'} \overline{\Theta^{(2)}};$$

using (B.22), (B.38) gives

$$(4.210) \quad \begin{aligned} D_t \mathbb{H} b + i \frac{A_1}{|Z_{,\alpha'}|^2} - i &= [D_t, \mathbb{H}] b + 2i \operatorname{Im} \mathbb{P}_H \left(D_t b + i \frac{A_1}{|Z_{,\alpha'}|^2} - i \right) \\ &= i \frac{A_1 - 1}{|Z_{,\alpha'}|^2} + 2i \operatorname{Re} \mathbb{P}_H(G^{(1)}) + \frac{1}{2} [b, \mathbb{H}] \partial_{\alpha'} b; \end{aligned}$$

and we know for $j \geq 2$

$$(4.211) \quad \mathbb{P}_A D_t^j \overline{Z}_t = -i \left[\mathbb{P}_A, \frac{A_1}{|Z_{,\alpha'}|^2} \right] \partial_{\alpha'} D_t^{j-2} \overline{Z}_t + \mathbb{P}_A \mathcal{P} D_t^{j-2} \overline{Z}_t;$$

so by (C.17), (C.19), (C.20), (4.120) and the estimates in §4.3.1, we have for $2 \leq j \leq 4$,

$$(4.212) \quad \left\| \partial_{\alpha'} \left(D_t \mathbb{H} b + i \frac{A_1}{|Z_{,\alpha'}|^2} \right) \right\|_{L^2} + \left\| \partial_{\alpha'} \mathbb{P}_H \left(D_t b + i \frac{A_1}{|Z_{,\alpha'}|^2} \right) \right\|_{L^2} + \left\| \mathbb{P}_A D_t^j \overline{Z}_t \right\|_{L^2} \lesssim \epsilon^2,$$

in particular for $j = 3$,

$$(4.213) \quad \left\| \partial_{\alpha'} \mathbb{P}_H \left(D_t b + i \frac{A_1}{|Z_{,\alpha'}|^2} \right) \right\|_{L^2} + \left\| \mathbb{P}_A D_t^3 \overline{Z}_t \right\|_{L^2} \lesssim \epsilon \left(\|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} \right);$$

by (C.40), (C.4), (4.153), and the estimates in §4.3.1, the following also holds,

$$(4.214) \quad \left\| \partial_{\alpha'} \mathbb{P}_H \left(D_t b + i \frac{A_1}{|Z_{,\alpha'}|^2} \right) \right\|_{\dot{H}^{1/2}} \lesssim \epsilon^2.$$

Observe that $M_{12,2}$ is sextic. Applying (C.36) gives

$$(4.215) \quad |M_{12,2}| \lesssim \left\| \partial_{\alpha'} \left(D_t \mathbb{H} b + i \frac{A_1}{|Z_{,\alpha'}|^2} \right) \right\|_{L^2} \|Z_{t,\alpha'}\|_{L^2} \|Z_{tt}\|_{\dot{H}^{1/2}}^2 \|Z_{tt}\|_{L^\infty} \lesssim \epsilon^5.$$

We rewrite $M_{11,j}$ as

$$(4.216) \quad \begin{aligned} M_{11,j} &= \iint \partial_{\alpha'} \mathbb{P}_H \left(i \frac{A_1(\alpha')}{|Z_{,\alpha'}|^2} + D_t b(\alpha') \right) \frac{D_t^j Z_t(\alpha') \mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta}{(\alpha' - \beta')^2} d\alpha' d\beta' \\ &+ \iint \partial_{\alpha'} \mathbb{P}_A \left(i \frac{A_1(\alpha')}{|Z_{,\alpha'}|^2} + D_t b(\alpha') \right) \frac{\mathbb{P}_H(D_t^j Z_t)(\alpha') \mathfrak{D}_t^l \theta \mathfrak{D}_t^i \bar{\theta} \mathfrak{D}_t^k \theta}{(\alpha' - \beta')^2} d\alpha' d\beta' \\ &+ \int \partial_{\alpha'} \mathbb{P}_A \left(i \frac{A_1(\alpha')}{|Z_{,\alpha'}|^2} + D_t b(\alpha') \right) [\mathbb{P}_H, < D_t^l \overline{Z}_t, D_t^i Z_t, D_t^k \overline{Z}_t >] \mathbb{P}_A(D_t^j Z_t)(\alpha') d\alpha' \\ &= I_j + II_j + III_j, \end{aligned}$$

where in the last term we used the Cauchy integral formula to rewrite it as a commutator. We have for $j = 2$, $(l, i, k) = (0, 0, 1)$ and permutations; $j = 3$, $(l, i, k) = (0, 0, 2), (0, 1, 1)$ and permutations; $j = 4$, $(l, i, k) = (0, 0, 3), (0, 1, 2), (1, 1, 1)$ and permutations. Observe that the first two terms in (4.216), I_j , II_j , are sextic; we apply (C.27), (C.42) to obtain

$$(4.217) \quad |I_2| + |II_2| + |I_4| + |II_4| \lesssim \epsilon^5, \quad |I_3| + |II_3| \lesssim \epsilon^3 \left(\|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}^2 + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 \right);$$

applying (C.17), (C.31) on III_j yields

$$(4.218) \quad |III_2| + |III_4| \lesssim \epsilon^5, \quad |III_3| \lesssim \epsilon^3 \left(\|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}^2 + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 \right),$$

therefore

$$|M_{11,2}| + |M_{11,4}| \lesssim \epsilon^5, \quad |M_{11,3}| \lesssim \epsilon^3 \left(\|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}^2 + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 \right).$$

Sum up the estimates in **Step 6** and **Step 7** we have

$$(4.219) \quad \left| \mathcal{R}_{IC,2} + \operatorname{Re} \frac{d}{dt} D_2(t) \right| + |\mathcal{R}_{IC,4}| \lesssim \epsilon^5, \quad |\mathcal{R}_{IC,3}| \lesssim \epsilon^3 \left(\|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}^2 + \left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2}^2 \right).$$

Step 8. In this step we treat the remaining terms in (4.13), namely

$$(4.220) \quad 2 \operatorname{Re} \sum_{l=0}^{j-1} \int i \partial_{\alpha'} \overline{\Theta^{(j)}} (\mathbb{P}_H D_t)^{l+1} \mathbb{P}_H \left(G^{(j-l)} - D_t \mathbb{P}_H G^{(j-1-l)} \right) d\alpha' - I_{1,j} - \frac{d}{dt} H_j,$$

$$(4.221) \quad \operatorname{Re} \int i \partial_{\alpha'} \overline{\Theta^{(j)}} \mathbb{P}_H \left(G^{(j+1)} - D_t \mathbb{P}_H G^{(j)} \right) d\alpha' - I_{2,j}.$$

Let

$$(4.222) \quad J_{l,j} = \frac{1}{2\pi} \iint D_t^j Z_t(\alpha', t) \frac{\mathfrak{D}_t^{l+1} \left\{ \mathfrak{D}_t \left(\theta(\alpha', \beta', t) \overline{\theta(\alpha', \beta', t)} \right) \mathfrak{D}_t^{j-l-1} \theta(\alpha', \beta', t) \right\}}{(\alpha' - \beta')^2} d\beta' d\alpha',$$

By (2.26), (2.27),

$$I_{1,j} = 2 \operatorname{Re} \sum_{l=0}^{j-1} J_{l,j}, \quad I_{2,j} = \operatorname{Re} J_{-1,j}.$$

Because

$$D_{\alpha'} \Theta^{(1)} = i \left(1 - \frac{1}{Z_{,\alpha'}} \right)$$

by (2.10), (4.33) and the notations (2.24) and (4.5)²⁸ we can write

$$(4.223) \quad \mathbb{P}_H \left(G^{(j-l)} - D_t \mathbb{P}_H G^{(j-1-l)} \right) = \frac{1}{2\pi i} \mathbb{P}_H \left(\frac{1}{\overline{Z}_{,\alpha'}} \int \frac{(\theta \overline{\lambda^1} + \overline{\theta} \lambda^1) \lambda^{j-l-1}}{(\alpha' - \beta')^2} d\beta' \right),$$

²⁸Observe that $\lambda^0 = \theta$.

and

$$\begin{aligned}
(4.224) \quad N_{l,j} &:= 2\pi \left(\int i \partial_{\alpha'} \overline{\Theta^{(j)}} (\mathbb{P}_H D_t)^{l+1} \mathbb{P}_H \left(G^{(j-l)} - D_t \mathbb{P}_H G^{(j-1-l)} \right) d\alpha' - J_{l,j} \right) \\
&= \int \partial_{\alpha'} \overline{\Theta^{(j)}} \left\{ (\mathbb{P}_H D_t)^{l+1} \mathbb{P}_H \frac{1}{\overline{Z}_{\alpha'}} - \frac{1}{\overline{Z}_{\alpha'}} D_t^{l+1} \right\} \int \frac{(\theta \overline{\lambda^1} + \overline{\theta} \lambda^1) \lambda^{j-l-1}}{(\alpha' - \beta')^2} d\beta' d\alpha' \\
&\quad + \int \overline{D_{\alpha'} \Theta^{(j)}} \left\{ D_t^{l+1} \int \frac{(\theta \overline{\lambda^1} + \overline{\theta} \lambda^1) \lambda^{j-l-1}}{(\alpha' - \beta')^2} d\beta' - \int \frac{\mathfrak{D}_t^{l+1} \left((\theta \overline{\lambda^1} + \overline{\theta} \lambda^1) \lambda^{j-l-1} \right)}{(\alpha' - \beta')^2} d\beta' \right\} d\alpha' \\
&\quad + \int \overline{D_{\alpha'} \Theta^{(j)}} \int \frac{\mathfrak{D}_t^{l+1} \left(\theta (\overline{\lambda^1} - \mathfrak{D}_t \overline{\theta}) \lambda^{j-l-1} + \overline{\theta} (\lambda^1 - \mathfrak{D}_t \theta) \lambda^{j-l-1} \right)}{(\alpha' - \beta')^2} d\beta' d\alpha' \\
&\quad + \int \overline{D_{\alpha'} \Theta^{(j)}} \int \frac{\mathfrak{D}_t^{l+1} \left(\mathfrak{D}_t (\theta \overline{\theta}) (\lambda^{j-l-1} - \mathfrak{D}_t^{j-l-1} \theta) \right)}{(\alpha' - \beta')^2} d\beta' d\alpha' \\
&\quad + \int \left(\overline{D_{\alpha'} \Theta^{(j)}} - D_t^j Z_t \right) \int \frac{\mathfrak{D}_t^{l+1} \left(\mathfrak{D}_t (\theta \overline{\theta}) \mathfrak{D}_t^{j-l-1} \theta \right)}{(\alpha' - \beta')^2} d\beta' d\alpha' \\
&= N_{l,j,1} + N_{l,j,2} + N_{l,j,3} + N_{l,j,4} + N_{l,j,5}.
\end{aligned}$$

Observe that there is a derivative loss in $N_{j-1,j,3}$ and $N_{j-1,j,5}$.²⁹ Because $\lambda^0 = \theta$, so $N_{j-1,j,4} = 0$. We combine $N_{j-1,j,3} + N_{j-1,j,5}$ with $-\pi \frac{d}{dt} H_j$, cf. (4.7), and write

$$\begin{aligned}
(4.225) \quad &\text{Re}(N_{j-1,j,3} + N_{j-1,j,5}) - \pi \frac{d}{dt} H_j = \tilde{N}_{j,3} + \tilde{N}_{j,4} + \tilde{N}_{j,5} - \pi R_{H,j} \\
&:= \text{Re} \int \overline{D_{\alpha'} \Theta^{(j)}} \int \frac{\overline{\theta} \theta \left(\mathfrak{D}_t^j \lambda^1 - \mathfrak{D}_t \lambda^j \right) + \theta \theta \left(\mathfrak{D}_t^j \overline{\lambda^1} - \mathfrak{D}_t \overline{\lambda^j} \right)}{(\alpha' - \beta')^2} d\beta' d\alpha' \\
&\quad + \text{Re} \iint D_t^j Z_t \frac{\mathfrak{D}_t^j \left(\theta (\overline{\lambda^1} - \mathfrak{D}_t \overline{\theta}) \theta + \overline{\theta} (\lambda^1 - \mathfrak{D}_t \theta) \theta \right) - \overline{\theta} \theta \mathfrak{D}_t^j (\lambda^1 - \mathfrak{D}_t \theta) - \theta \theta \mathfrak{D}_t^j (\overline{\lambda^1} - \mathfrak{D}_t \overline{\theta})}{(\alpha' - \beta')^2} d\beta' d\alpha' \\
&\quad + \text{Re} \int \left(\overline{D_{\alpha'} \Theta^{(j)}} - D_t^j Z_t \right) \int \frac{\mathfrak{D}_t^j \left((\theta \overline{\lambda^1} + \overline{\theta} \lambda^1) \theta \right) - \overline{\theta} \theta \mathfrak{D}_t^j \lambda^1 - \theta \theta \mathfrak{D}_t^j \overline{\lambda^1}}{(\alpha' - \beta')^2} d\beta' d\alpha' - \pi R_{H,j};
\end{aligned}$$

this cancels out the derivative lossing terms for $j = 4$, and enables us to get the desired estimates for $j = 3$.

We sum up the above decomposition:

$$\begin{aligned}
(4.226) \quad &\pi \left(2 \text{Re} \sum_{l=0}^{j-1} \int i \partial_{\alpha'} \overline{\Theta^{(j)}} (\mathbb{P}_H D_t)^{l+1} \mathbb{P}_H \left(G^{(j-l)} - D_t \mathbb{P}_H G^{(j-1-l)} \right) d\alpha' - I_{1,j} - \frac{d}{dt} H_j(t) \right) \\
&= \text{Re} \left(\sum_{l=0}^{j-2} \sum_{k=1}^5 N_{l,j,k} + \sum_{k=1}^2 N_{j-1,j,k} \right) + \sum_{k=3}^5 \tilde{N}_{j,k} - \pi R_{H,j},
\end{aligned}$$

and

$$(4.227) \quad 2\pi \left(\text{Re} \int i \partial_{\alpha'} \overline{\Theta^{(j)}} \mathbb{P}_H \left(G^{(j+1)} - D_t \mathbb{P}_H G^{(j)} \right) d\alpha' - I_{2,j} \right) = \text{Re} \sum_{k=1}^5 N_{-1,j,k}.$$

Observe that all the terms in (4.224), (4.225) are quintic.

²⁹Namely $N_{j-1,j,3}$ and $N_{j-1,j,5}$ contain factors that cannot be controlled by $\mathcal{E}_j(t)$.

Now we use (B.26) to write

$$(4.228) \quad \begin{aligned} \mathbb{P}_H \left((\mathbb{P}_H D_t)^{l+1} \mathbb{P}_H \frac{1}{\overline{Z}_{,\alpha'}} - \frac{1}{\overline{Z}_{,\alpha'}} D_t^{l+1} \right) &= ((\mathbb{P}_H D_t)^{l+1} \mathbb{P}_H - \mathbb{P}_H D_t^{l+1}) \frac{1}{\overline{Z}_{,\alpha'}} + \mathbb{P}_H \left[D_t^{l+1}, \frac{1}{\overline{Z}_{,\alpha'}} \right] \\ &= - \sum_{k=0}^l (\mathbb{P}_H D_t)^k \mathbb{P}_H [b, \mathbb{P}_A] \partial_{\alpha'} D_t^{l-k} \frac{1}{\overline{Z}_{,\alpha'}} + \mathbb{P}_H \sum_{k=0}^l \binom{l+1}{k+1} \left(D_t^{k+1} \frac{1}{\overline{Z}_{,\alpha'}} \right) D_t^{l-k}, \end{aligned}$$

because

$$(4.229) \quad D_{\alpha'} \Theta^{(1)} - \overline{Z}_{tt} = i \frac{A_1 - 1}{\overline{Z}_{,\alpha'}},$$

by (B.41) we have

$$(4.230) \quad \begin{aligned} D_{\alpha'} \Theta^{(l+1)} - D_t^{l+1} \overline{Z}_t &= D_{\alpha'} \Theta^{(l+1)} - D_t^l D_{\alpha'} \Theta^{(1)} + D_t^l (D_{\alpha'} \Theta^{(1)} - \overline{Z}_{tt}) \\ &= \sum_{k=0}^{l-1} (\mathbb{P}_H D_t)^k \left[\frac{1}{\overline{Z}_{,\alpha'}}, \mathbb{P}_H \right] \partial_{\alpha'} D_t \Theta^{(l-k)} + \sum_{k=0}^{l-1} (\mathbb{P}_H D_t)^k \left[\mathbb{P}_H, \frac{1}{\overline{Z}_{,\alpha'}} D_{\alpha'} \Theta^{(l-k)} \right] Z_{t,\alpha'} \\ &\quad + i \sum_{k=0}^{l-1} D_t^k [\mathbb{P}_A, b] \partial_{\alpha'} (\mathbb{P}_H D_t)^{l-1-k} \frac{1}{\overline{Z}_{,\alpha'}} + i D_t^l \left(\frac{A_1 - 1}{\overline{Z}_{,\alpha'}} \right), \end{aligned}$$

and

$$(4.231) \quad \begin{aligned} D_t D_{\alpha'} \Theta^{(j)} - D_t^j D_{\alpha'} \Theta^{(1)} &= \sum_{k=0}^{j-2} D_t (\mathbb{P}_H D_t)^k \left[\frac{1}{\overline{Z}_{,\alpha'}}, \mathbb{P}_H \right] \partial_{\alpha'} D_t \Theta^{(j-1-k)} \\ &\quad + \sum_{k=0}^{j-2} D_t (\mathbb{P}_H D_t)^k \left[\mathbb{P}_H, \frac{1}{\overline{Z}_{,\alpha'}} D_{\alpha'} \Theta^{(j-1-k)} \right] Z_{t,\alpha'} + i \sum_{k=0}^{j-2} D_t^{k+1} [\mathbb{P}_A, b] \partial_{\alpha'} (\mathbb{P}_H D_t)^{j-2-k} \frac{1}{\overline{Z}_{,\alpha'}}. \end{aligned}$$

We are now ready to do the estimates. We begin with $N_{l,j,1}$. Observe that by the Cauchy integral formula, we can insert a \mathbb{P}_H to write it as

$$(4.232) \quad N_{l,j,1} = \int \partial_{\alpha'} \overline{\Theta^{(j)}} \mathbb{P}_H \left\{ (\mathbb{P}_H D_t)^{l+1} \mathbb{P}_H \frac{1}{\overline{Z}_{,\alpha'}} - \frac{1}{\overline{Z}_{,\alpha'}} D_t^{l+1} \right\} \int \frac{(\theta \overline{\lambda^1} + \overline{\theta} \lambda^1) \lambda^{j-l-1}}{(\alpha' - \beta')^2} d\beta' d\alpha'$$

and it is clear that $N_{-1,j,1} = 0$. We estimate $|N_{l,j,1}|$ for $0 \leq l \leq j-1$, $2 \leq j \leq 4$. By (4.228) we need to estimate, for $0 \leq k \leq l$,

$$(4.233) \quad A_{k,l,j} := \left\| (\mathbb{P}_H D_t)^k \mathbb{P}_H [b, \mathbb{P}_A] \partial_{\alpha'} D_t^{l-k} \left(\frac{1}{\overline{Z}_{,\alpha'}} \int \frac{(\theta \overline{\lambda^1} + \overline{\theta} \lambda^1) \lambda^{j-l-1}}{(\alpha' - \beta')^2} d\beta' \right) \right\|_{L^2},$$

$$(4.234) \quad B_{k,l,j} := \left\| \mathbb{P}_H \left\{ \left(D_t^{k+1} \frac{1}{\overline{Z}_{,\alpha'}} \right) D_t^{l-k} \int \frac{(\theta \overline{\lambda^1} + \overline{\theta} \lambda^1) \lambda^{j-l-1}}{(\alpha' - \beta')^2} d\beta' \right\} \right\|_{L^2}.$$

By (B.29), (C.29), (C.30), (C.27), we have, for $k < l$ or $l < j-1$ and $2 \leq j \leq 4$,

$$(4.235) \quad \left\| D_t^{l-k} \int \frac{(\theta \overline{\lambda^1} + \overline{\theta} \lambda^1) \lambda^{j-l-1}}{(\alpha' - \beta')^2} d\beta' \right\|_{L^2} \lesssim \epsilon^3,$$

and by (4.129) and (C.31), (C.32),

$$(4.236) \quad \left\| \int \frac{(\theta \overline{\lambda^1} + \overline{\theta} \lambda^1) \lambda^1}{(\alpha' - \beta')^2} d\beta' \right\|_{\dot{H}^{1/2}} + \left\| D_t \int \frac{(\theta \overline{\lambda^1} + \overline{\theta} \lambda^1) \lambda^0}{(\alpha' - \beta')^2} d\beta' \right\|_{\dot{H}^{1/2}} \lesssim \epsilon^2 \left(\left\| \partial_{\alpha'} \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right),$$

$$(4.237) \quad \left\| \int \frac{(\theta \overline{\lambda^1} + \overline{\theta} \lambda^1) \lambda^0}{(\alpha' - \beta')^2} d\beta' \right\|_{\dot{H}^{1/2}} \lesssim \epsilon^3, \quad \left\| \partial_{\alpha'} \int \frac{(\theta \overline{\lambda^1} + \overline{\theta} \lambda^1) \lambda^0}{(\alpha' - \beta')^2} d\beta' \right\|_{L^2} \lesssim \epsilon^2 \left\| \partial_{\alpha'} \frac{1}{\overline{Z}_{,\alpha'}} \right\|_{L^2},$$

therefore by (B.25), (B.32), (C.19), (C.20), (C.23), (C.21), (C.27), we have, except for the cases where $l = k = j - 1$ for $j = 3, 4$,

$$(4.238) \quad A_{k,l,2} \lesssim \epsilon^4, \quad A_{k,l,3} \lesssim \epsilon^3 \left(\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right), \quad A_{k,l,4} \lesssim \epsilon^4.$$

We also have, by (B.29), (C.29), (C.27), (C.33), (C.42), that except for the cases where $l = k = j - 1$ for $2 \leq j \leq 4$,

$$(4.239) \quad B_{k,l,2} \lesssim \epsilon^4, \quad B_{k,l,3} \lesssim \epsilon^3 \left(\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right), \quad B_{k,l,4} \lesssim \epsilon^4.$$

Now all the terms in $A_{j-1,j-1,j}$, $j = 3, 4$, after expanding by (B.25), (B.32), can be handled similarly, except for one, namely

$$(4.240) \quad \mathbb{P}_H \left[D_t^{j-1} b, \mathbb{P}_A \right] \partial_{\alpha'} \left(\frac{1}{\overline{Z}_{,\alpha'}} \int \frac{(\theta \overline{\lambda}^1 + \overline{\theta} \lambda^1) \lambda^0}{(\alpha' - \beta')^2} d\beta' \right);$$

for this term, we write

$$(4.241) \quad D_t^{j-1} b = D_t^{j-1} b - 2 \operatorname{Re} \frac{D_t^{j-1} Z_t}{Z_{,\alpha'}} + 2 \operatorname{Re} \frac{D_t^{j-1} Z_t}{Z_{,\alpha'}}$$

and use (4.92) and (4.99) and (C.19) to estimate the first term and (C.17) to estimate the second term,

$$(4.242) \quad \begin{aligned} & \left\| \mathbb{P}_H \left[D_t^{j-1} b, \mathbb{P}_A \right] \partial_{\alpha'} \left(\frac{1}{\overline{Z}_{,\alpha'}} \int \frac{(\theta \overline{\lambda}^1 + \overline{\theta} \lambda^1) \lambda^0}{(\alpha' - \beta')^2} d\beta' \right) \right\|_{L^2} \\ & \lesssim \left\| \partial_{\alpha'} \left(D_t^{j-1} b - 2 \operatorname{Re} \frac{D_t^{j-1} Z_t}{Z_{,\alpha'}} \right) \right\|_{L^2} \left\| \frac{1}{\overline{Z}_{,\alpha'}} \int \frac{(\theta \overline{\lambda}^1 + \overline{\theta} \lambda^1) \lambda^0}{(\alpha' - \beta')^2} d\beta' \right\|_{\dot{H}^{1/2}} \\ & + \left\| \frac{D_t^{j-1} Z_t}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}} \left\| \partial_{\alpha'} \left(\frac{1}{\overline{Z}_{,\alpha'}} \int \frac{(\theta \overline{\lambda}^1 + \overline{\theta} \lambda^1) \lambda^0}{(\alpha' - \beta')^2} d\beta' \right) \right\|_{L^2}, \end{aligned}$$

and we get

$$(4.243) \quad A_{2,2,3} \lesssim \epsilon^3 \left(\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right), \quad A_{3,3,4} \lesssim \epsilon^4.$$

To estimate $B_{j-1,j-1,j}$ for $2 \leq j \leq 4$, we first compute using (B.8),

$$(4.244) \quad \begin{aligned} D_t^j \frac{1}{\overline{Z}_{,\alpha'}} &= D_t^{j-1} \left(\frac{1}{\overline{Z}_{,\alpha'}} (b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t) \right) + \left[D_t^{j-1}, \frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \right] Z_t \\ &+ \frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \mathbb{P}_H D_t^{j-1} Z_t + \frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \mathbb{P}_A D_t^{j-1} Z_t, \end{aligned}$$

and by (B.20), we write

$$(4.245) \quad \left[D_t^{j-1}, \frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \right] Z_t = \sum_{k=0}^{j-2} D_t^k \left(\frac{b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t}{|Z_{,\alpha'}|^2} \partial_{\alpha'} D_t^{j-2-k} Z_t \right),$$

and using the fact $\mathbb{P}_H Z_t = 0$ we write

$$(4.246) \quad \mathbb{P}_H D_t^{j-1} Z_t = \sum_{k=0}^{j-2} D_t^k [\mathbb{P}_H, D_t] D_t^{j-2-k} Z_t = \sum_{k=0}^{j-2} D_t^k [\mathbb{P}_H, b] \partial_{\alpha'} D_t^{j-2-k} Z_t;$$

using the expansions in (4.244), (4.245) and (4.246) and the estimates in §4.3.1, we have,

$$(4.247) \quad \left\| D_t^j \frac{1}{\overline{Z}_{,\alpha'}} - \frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \mathbb{P}_A D_t^{j-1} Z_t \right\|_{L^2} \lesssim \epsilon^2, \quad j = 2, 4,$$

$$(4.248) \quad \left\| D_t^3 \frac{1}{\bar{Z}_{,\alpha'}} - \frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \mathbb{P}_A D_t^2 Z_t \right\|_{L^2} \lesssim \epsilon \left(\left\| \partial_{\alpha'} \frac{1}{\bar{Z}_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right);$$

now we rewrite

$$(4.249) \quad \begin{aligned} & \mathbb{P}_H \left\{ \left(D_t^j \frac{1}{\bar{Z}_{,\alpha'}} \right) \int \frac{(\theta \bar{\lambda}^1 + \bar{\theta} \lambda^1) \lambda^0}{(\alpha' - \beta')^2} d\beta' \right\} \\ &= \mathbb{P}_H \left\{ \left(D_t^j \frac{1}{\bar{Z}_{,\alpha'}} - \frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \mathbb{P}_A D_t^{j-1} Z_t \right) \int \frac{(\theta \bar{\lambda}^1 + \bar{\theta} \lambda^1) \lambda^0}{(\alpha' - \beta')^2} d\beta' \right\} \\ &+ \left[\mathbb{P}_H, \frac{1}{|Z_{,\alpha'}|^2} \int \frac{(\theta \bar{\lambda}^1 + \bar{\theta} \lambda^1) \lambda^0}{(\alpha' - \beta')^2} d\beta' \right] \partial_{\alpha'} \mathbb{P}_A D_t^{j-1} Z_t, \end{aligned}$$

by (C.29), (C.32),

$$(4.250) \quad \left\| \int \frac{(\theta \bar{\lambda}^1 + \bar{\theta} \lambda^1) \lambda^0}{(\alpha' - \beta')^2} d\beta' \right\|_{L^\infty} \lesssim \|Z_{t,\alpha'}\|_{L^2}^2,$$

$$(4.251) \quad \left\| \int \frac{(\theta \bar{\lambda}^1 + \bar{\theta} \lambda^1) \lambda^0}{(\alpha' - \beta')^2} d\beta' \right\|_{\dot{H}^1} \lesssim \|Z_{t,\alpha'}\|_{L^2}^2 \left\| \partial_{\alpha'} \frac{1}{\bar{Z}_{,\alpha'}} \right\|_{L^2};$$

therefore by (C.19), (4.247), (4.248), we have

$$(4.252) \quad B_{2,2,3} \lesssim \epsilon^3 \left(\left\| \partial_{\alpha'} \frac{1}{\bar{Z}_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right), \quad B_{1,1,2} + B_{3,3,4} \lesssim \epsilon^4.$$

Sum up (4.232)-(4.252) we conclude

$$(4.253) \quad \begin{aligned} & |N_{l,j,1}| \lesssim \epsilon^5, \quad -1 \leq l \leq j-1, \quad j=2,4; \\ & |N_{l,3,1}| \lesssim \epsilon^3 \left(\left\| \partial_{\alpha'} \frac{1}{\bar{Z}_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right)^2, \quad -1 \leq l \leq 2. \end{aligned}$$

Now we handle $N_{l,j,2}$. Observe that $N_{-1,j,2} = 0$, so we work on the cases where $0 \leq l \leq j-1$. We use (B.29) to expand. The estimates are routine for all the terms after expansion, using (C.46), (C.48), (C.27), (C.42), (C.35), (C.49), except that when $l = j-1$ for $j = 3, 4$, we need to again decompose $D_t^{j-1} b$ by (4.241), and treat all the terms as usual, except

$$(4.254) \quad \int \partial_{\beta'} \operatorname{Re} \frac{D_t^{j-1} Z_t(\beta')}{Z_{,\beta'}} \frac{(\theta \bar{\lambda}^1 + \bar{\theta} \lambda^1) \lambda^0}{(\alpha' - \beta')^2} d\beta',$$

for which we first perform integration by parts, then apply (C.46), (C.48) for $\mathbf{b} = \operatorname{Re} \frac{D_t^{j-1} Z_t}{Z_{,\alpha'}}$. We have

$$(4.255) \quad \begin{aligned} & |N_{l,j,2}| \lesssim \epsilon^5, \quad -1 \leq l \leq j-1, \quad j=2,4; \\ & |N_{l,3,2}| \lesssim \epsilon^3 \left(\left\| \partial_{\alpha'} \frac{1}{\bar{Z}_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right)^2, \quad -1 \leq l \leq 2. \end{aligned}$$

We consider the remaining terms in (4.226)-(4.227). Using (4.230) and the estimates in §4.3.1, §4.3.2, (C.17), (C.20), (C.19), (C.23), (C.27) we have

$$(4.256) \quad \left\| D_{\alpha'} \Theta^{(1)} - D_t \bar{Z}_t \right\|_{\dot{H}^{1/2}} \lesssim \epsilon^2,$$

$$(4.257) \quad \left\| D_{\alpha'} \Theta^{(1)} - D_t \bar{Z}_t \right\|_{\dot{H}^1} \lesssim \epsilon \left(\left\| \partial_{\alpha'} \frac{1}{\bar{Z}_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right),$$

$$(4.258) \quad \left\| D_t^l \left(D_{\alpha'} \Theta^{(j-l)} - D_t^{j-l} \bar{Z}_t \right) \right\|_{L^2} \lesssim \epsilon^2, \quad j=2,4, \quad 0 \leq l \leq j,$$

$$(4.259) \quad \left\| D_t^l \left(D_{\alpha'} \Theta^{(3-l)} - D_t^{3-l} \bar{Z}_t \right) \right\|_{L^2} \lesssim \epsilon \left(\left\| \partial_{\alpha'} \frac{1}{Z_{\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right), \quad 0 \leq l \leq 3;$$

and using (4.231) we have³⁰

$$(4.260) \quad \begin{aligned} \left\| D_t D_{\alpha'} \Theta^{(j)} - D_t^j D_{\alpha'} \Theta^{(1)} \right\|_{L^2} &\lesssim \epsilon \left(\left\| \partial_{\alpha'} \frac{1}{Z_{\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right), \quad \text{for } j = 2, 3, \\ \left\| D_t D_{\alpha'} \Theta^{(4)} - D_t^4 D_{\alpha'} \Theta^{(1)} \right\|_{L^2} &\lesssim \epsilon^2; \end{aligned}$$

this gives, by (C.29), (C.30),

$$(4.261) \quad \begin{aligned} \sum_{l=-1}^{j-2} \sum_{k=3}^5 |N_{l,j,k}| + \sum_{k=3}^5 |\tilde{N}_{j,k}| + |R_{H,j}| &\lesssim \epsilon^5, \quad \text{for } j = 2, 4; \\ \sum_{l=-1}^{j-2} \sum_{k=3}^5 |N_{l,j,k}| + \sum_{k=3}^5 |\tilde{N}_{j,k}| + |R_{H,j}| &\lesssim \epsilon^3 \left(\left\| \partial_{\alpha'} \frac{1}{Z_{\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right)^2, \quad \text{for } j = 3. \end{aligned}$$

Sum up the results in Steps 6-8, we get

$$(4.262) \quad \frac{d}{dt} (\mathcal{E}_2(t) + \mathcal{E}_3(t) + \mathcal{E}_4(t)) \lesssim \epsilon^5, \quad \frac{d}{dt} \mathcal{E}_3(t) \lesssim \epsilon^3 \left(\left\| \partial_{\alpha'} \frac{1}{Z_{\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right)^2.$$

This together with (4.164), (4.134) gives the inequalities (4.20) and (4.24), and finishes the proof for Propositions 4.3, 4.4 and Theorem 3.1.

APPENDIX A. NOTATIONS AND CONVENTIONS

We use the following notations and conventions throughout the paper: compositions are always in terms of the spatial variables and we write for $f = f(\cdot, t)$, $g = g(\cdot, t)$, $f(g(\cdot, t), t) := f \circ g(\cdot, t) := U_g f(\cdot, t)$. We identify (x, y) with the complex number $x + iy$; $\operatorname{Re} z$, $\operatorname{Im} z$ are the real and imaginary parts of z ; $\bar{z} = \operatorname{Re} z - i \operatorname{Im} z$ is the complex conjugate of z . $\bar{\Omega}$ is the closure of the domain Ω , $\partial\Omega$ is the boundary of Ω , $\mathcal{P}_- := \{z \in \mathbb{C} : \operatorname{Im} z < 0\}$ is the lower half plane. $[A, B] := AB - BA$ is the commutator of operators A and B .

We use $z = x + iy = z(\alpha, t)$, $z_t = z_t(\alpha, t)$ and $z_{tt}(\alpha, t)$ to denote the position, velocity and acceleration of the interface in Lagrangian coordinate α ; $Z = X + iY = Z(\alpha', t)$, $Z_t = Z_t(\alpha', t)$ and $Z_{tt}(\alpha', t)$ denote the position, velocity and acceleration of the interface in the Riemann mapping variable α' ; $\mathbf{h}(\alpha, t) = \alpha'$ is the coordinate change from the Lagrangian variable α to the Riemann mapping variable α' ,³¹ $b = \mathbf{h}_t \circ \mathbf{h}^{-1}$, and the material derivative is $D_t = \partial_t + b \partial_{\alpha'}$. We write

$$Z_{,\alpha'} = \partial_{\alpha'} Z(\alpha', t), \quad Z_{t,\alpha'} = \partial_{\alpha'} Z_t, \quad Z_{tt,\alpha'} = \partial_{\alpha'} Z_{tt}, \quad \text{etc.}$$

Let \mathbb{H} be the Hilbert transform associated with the lower half plane \mathcal{P}_- :

$$(A.1) \quad \mathbb{H}f(\alpha') = \frac{1}{\pi i} \operatorname{pv.} \int \frac{1}{\alpha' - \beta'} f(\beta') d\beta'.$$

We know $\mathbb{H}^2 = I$, and a function $f \in L^p(\mathbb{R})$, $1 \leq p < \infty$, is the boundary value of a holomorphic function in \mathcal{P}_- if and only if $f = \mathbb{H}f$. We define the projections to the space of holomorphic, and respectively, anti-holomorphic functions in the lower half plane by

$$(A.2) \quad \mathbb{P}_H := \frac{1}{2}(I + \mathbb{H}), \quad \text{and} \quad \mathbb{P}_A := \frac{1}{2}(I - \mathbb{H}).$$

³⁰Again we use the decomposition (4.241) to treat the term $[\mathbb{P}_A, D_t^{j-1} b] \partial_{\alpha'} \frac{1}{Z_{\alpha'}}$.

³¹ $\mathbf{h}(\alpha, t) = \Phi(z(\alpha, t); t)$, where $\Phi(\cdot, t) : \Omega \rightarrow \mathcal{P}_-$ is the Riemann mapping satisfying $\Phi(z(0, t); t) = 0$ and $\lim_{z \rightarrow \infty} \Phi_z(z, t) = 1$.

It is clear that the decomposition identity

$$(A.3) \quad \mathbb{P}_H + \mathbb{P}_A = I$$

and the projection identity

$$(A.4) \quad \mathbb{P}_H \mathbb{P}_A = \mathbb{P}_A \mathbb{P}_H = 0$$

hold. We will often call a function $f \in L^p$, $1 \leq p < \infty$, that is the boundary value of a holomorphic function in \mathcal{S}_- simply by "holomorphic".

We define

$$(A.5) \quad D_{\alpha'} = \frac{1}{Z_{\alpha'}} \partial_{\alpha'},$$

$$(A.6) \quad [f, g; h] := \frac{1}{\pi i} \int \frac{(f(x) - f(y))(g(x) - g(y))}{(x - y)^2} h(y) dy,$$

and

$$(A.7) \quad \langle f, g, h \rangle := \frac{1}{\pi i} \int \frac{(f(x) - f(y))(g(x) - g(y))(h(x) - h(y))}{(x - y)^2} dy.$$

We use the following notations for functional spaces: $H^s = H^s(\mathbb{R})$ is the Sobolev space with norm $\|f\|_{H^s} := (\int (1 + |\xi|^2)^s |\hat{f}(\xi)|^2 d\xi)^{1/2}$, $\dot{H}^s = \dot{H}^s(\mathbb{R})$ is the homogeneous Sobolev space with norm $\|f\|_{\dot{H}^s} = c(\int |\xi|^{2s} |\hat{f}(\xi)|^2 d\xi)^{1/2}$, and we define

$$(A.8) \quad \|f\|_{\dot{H}^{1/2}}^2 = \|f\|_{\dot{H}^{1/2}(\mathbb{R})}^2 := \int i\mathbb{H} \partial_x f(x) \bar{f}(x) dx = \frac{1}{2\pi} \iint \frac{|f(x) - f(y)|^2}{(x - y)^2} dx dy.$$

$L^p = L^p(\mathbb{R})$ is the L^p space with $\|f\|_{L^p} := (\int |f(x)|^p dx)^{1/p}$ for $1 \leq p < \infty$, and $f \in L^\infty$ if $\|f\|_{L^\infty} := \text{ess sup } |f(x)| < \infty$. When not specified, all the norms $\|f\|_{H^s}$, $\|f\|_{\dot{H}^s}$, $\|f\|_{L^p}$, $1 \leq p \leq \infty$ are in terms of the spatial variable only, and $\|f\|_{H^s(\mathbb{R})}$, $\|f\|_{\dot{H}^s(\mathbb{R})}$, $\|f\|_{L^p(\mathbb{R})}$, $1 \leq p \leq \infty$ are in terms of the spatial variable. $C^j(X)$ is the space of j -times continuously differentiable functions on the set X ; $C_0^j(\mathbb{R})$ is the space of j -times continuously differentiable functions that decays at the infinity.

We use c , C to denote universal constants. $c(a_1, \dots)$, $C(a_1, \dots)$, $M(a_1, \dots)$ are constants depending on a_1, \dots ; constants appearing in different contexts need not be the same. We write $f \lesssim g$ if there is a universal constant c , such that $f \leq cg$.

The following are some additional notations used in this paper:

$Q := (I + \mathbb{H})\psi \circ \mathbf{h}^{-1}$, where $\psi \circ \mathbf{h}^{-1}$ is the trace of the velocity potential on the interface; $\Theta^{(0)} := Q$, $\Theta^{(j)} := (\mathbb{P}_H D_t)^j Q$, and $G^{(j)} := D_t \mathbb{P}_H D_t \Theta^{(j)} + i \frac{1}{|Z_{\alpha'}|^2} \partial_{\alpha'} \Theta^{(j)}$.

We define $\theta := \bar{Z}_t(\alpha', t) - \bar{Z}_t(\beta', t)$; $\lambda^j = D_{\alpha'} \Theta^{(j)}(\alpha', t) - D_{\beta'} \Theta^{(j)}(\beta', t)$; $\mathfrak{D}_t := \partial_t + b(\alpha', t) \partial_{\alpha'} + b(\beta', t) \partial_{\beta'}$, and $\mathcal{P} := \mathfrak{D}_t^2 + i \frac{A_1(\alpha', t)}{|Z_{\alpha'}|^2} \partial_{\alpha'} + i \frac{A_1(\beta', t)}{|Z_{\beta'}|^2} \partial_{\beta'}$. Observe that when acting on a function independent of β' , i.e. $f = f(\alpha', t)$, $\mathcal{P}f = (D_t^2 + i \frac{A_1}{|Z_{\alpha'}|^2} \partial_{\alpha'})f$.

We denote by $M(f)$ the Hardy-Littlewood maximum function of f .

APPENDIX B. EQUATIONS AND IDENTITIES

Here we give some basic equations and identities that will be used in this paper. First we recall some of the equations and formulas derived in our earlier work, see [46, 50] or §2.2, §2.3 and §2.7 of [51].

B.1. Interface equations. We know that the interface equations for the 2d water waves is given by

$$(B.1) \quad \begin{cases} \overline{Z}_{tt} - i = -i \frac{A_1}{Z_{,\alpha'}} \\ \overline{Z}_t = \mathbb{H} \overline{Z}_t, \quad \frac{1}{Z_{,\alpha'}} - 1 = \mathbb{H} \left(\frac{1}{Z_{,\alpha'}} - 1 \right), \end{cases}$$

where the quantities A_1 and b satisfy

$$(B.2) \quad A_1 = 1 - \operatorname{Im} [Z_t, \mathbb{H}] \overline{Z}_{t,\alpha'} = 1 - \frac{1}{2} \operatorname{Im} [Z_t, \overline{Z}_t; 1], \quad b = \operatorname{Re} (I - \mathbb{H}) \left(\frac{Z_t}{Z_{,\alpha'}} \right).$$

Let

$$(B.3) \quad \mathfrak{P} := D_t^2 + i \frac{A_1}{|Z_{,\alpha'}|^2} \partial_{\alpha'}.$$

The quasi-linear equation for the water waves is

$$(B.4) \quad \mathcal{P} \overline{Z}_t = \mathfrak{P} \overline{Z}_t = \frac{\mathfrak{a}_t}{\mathfrak{a}} \circ \mathfrak{h}^{-1} (\overline{Z}_{tt} - i)$$

where

$$(B.5) \quad \frac{\mathfrak{a}_t}{\mathfrak{a}} \circ \mathfrak{h}^{-1} = \frac{D_t A_1}{A_1} + b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t,$$

with

$$(B.6) \quad D_t A_1 = -\operatorname{Im} ([Z_{tt}, \mathbb{H}] \overline{Z}_{t,\alpha'} + [Z_t, \mathbb{H}] \partial_{\alpha'} \overline{Z}_{tt} - [Z_t, b; \overline{Z}_{t,\alpha'}]) = -\operatorname{Im} ([\overline{Z}_t, Z_{tt}; 1] - [Z_t, b; \overline{Z}_{t,\alpha'}]),$$

and³²

$$(B.7) \quad b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t = \operatorname{Re} \left(\left[\frac{1}{Z_{,\alpha'}}, \mathbb{H} \right] Z_{t,\alpha'} + [Z_t, \mathbb{H}] \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right) = \frac{1}{2} \left[\frac{1}{Z_{,\alpha'}}, Z_t; 1 \right] - \frac{1}{2} \left[\overline{Z}_t, \frac{1}{\overline{Z}_{,\alpha'}}; 1 \right].$$

We also have

$$(B.8) \quad D_t \left(\frac{1}{Z_{,\alpha'}} \right) = \frac{1}{Z_{,\alpha'}} (b_{\alpha'} - D_{\alpha'} Z_t) = \frac{1}{Z_{,\alpha'}} (b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t) + \frac{\overline{Z}_{t,\alpha'}}{|Z_{,\alpha'}|^2}.$$

B.2. Basic identities. We give some basic identities that will be used in our calculations. We begin with Proposition B.1, which is a consequence of the fact that the product of holomorphic functions is holomorphic and (A.4).

Proposition B.1. *Assume that $f, g \in L^2(\mathbb{R})$.*

1. Assume either both f, g are holomorphic: $f = \mathbb{H}f, g = \mathbb{H}g$, or both are anti-holomorphic: $f = -\mathbb{H}f, g = -\mathbb{H}g$. Then

$$(B.9) \quad [f, \mathbb{H}]g = 0.$$

2. If $\mathbb{P}_A f = \mathbb{P}_A g = 0$, then $\mathbb{P}_A(fg) = 0$; and if $\mathbb{P}_H f = \mathbb{P}_H g = 0$, then $\mathbb{P}_H(fg) = 0$.

Proposition B.2. *1. We have*

$$(B.10) \quad [f, g; h] = [f, \mathbb{H}] \partial_{\alpha'} (gh) + [g, \mathbb{H}] \partial_{\alpha'} (fh) - [fg, \mathbb{H}] \partial_{\alpha'} h.$$

2. If h is holomorphic, i.e. $\mathbb{P}_A h = 0$, then

$$(B.11) \quad \mathbb{P}_H [f, g; h] = -2 \mathbb{P}_H (f \partial_{\alpha'} \mathbb{P}_A (gh)) - 2 \mathbb{P}_H (g \partial_{\alpha'} \mathbb{P}_A (fh)).$$

³²The second equality in (B.6) and (B.7) are obtained by integration by parts.

(B.10) is obtained by integration by parts. (B.11) follows from (B.10), (A.4) and (A.3) .

We often use the following equalities to calculate the time and spatial derivatives to our energy functionals, and the following version of the Cauchy integral formula to derive our equations and formulas.

Proposition B.3. *For f, g smooth and decay fast at infinity,*

$$(B.12) \quad \frac{d}{dt} \int f(\alpha', t) d\alpha' = \int (D_t + b_{\alpha'}) f(\alpha', t) d\alpha';$$

$$(B.13) \quad \frac{d}{dt} \iint g(\alpha', \beta', t) d\alpha' d\beta' = \iint (\mathfrak{D}_t + b_{\alpha'} + b_{\beta'}) g(\alpha', \beta', t) d\alpha' d\beta';$$

$$(B.14) \quad D_t \int g(\alpha', \beta', t) d\beta' = \int (\mathfrak{D}_t + b_{\beta'}) g(\alpha', \beta', t) d\beta';$$

$$(B.15) \quad \partial_{\alpha'} \int g(\alpha', \beta', t) d\beta' = \int (\partial_{\alpha'} + \partial_{\beta'}) g(\alpha', \beta', t) d\beta'.$$

(B.12), (B.13), (B.14), (B.15) follow from the simple fact that $\int (b\partial_{\alpha'} + b_{\alpha'}) f d\alpha' = \int \partial_{\alpha'} (bf) d\alpha' = 0$, $\int (b\partial_{\beta'} + b_{\beta'}) g d\beta' = \int \partial_{\beta'} (bg) d\beta' = 0$, and $\int \partial_{\beta'} g d\beta' = 0$.

Proposition B.4 (Cauchy integral formula). *For any $\Theta \in L^1(\mathbb{R})$, satisfying $\mathbb{P}_A \Theta = 0$ or $\mathbb{P}_H \Theta = 0$,*

$$(B.16) \quad \int \Theta(\alpha') d\alpha' = 0.$$

B.3. Commutator identities. We include here various commutator identities that are necessary for our proofs. Some have already appeared in Appendix B.5 of [31] and Appendix B of [51]. We have

$$(B.17) \quad [D_t, D_{\alpha'}] = -(D_{\alpha'} Z_t) D_{\alpha'},$$

$$(B.18) \quad [D_t, \partial_{\alpha'}] = -b_{\alpha'} \partial_{\alpha'}.$$

By product rules,

$$(B.19) \quad [D_t, \frac{1}{Z_{,\alpha'}}] f = D_t \left(\frac{1}{Z_{,\alpha'}} \right) f.$$

From (B.8) and (B.18) it implies that

$$(B.20) \quad \left[D_t, \frac{1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \right] f = \frac{b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t}{|Z_{,\alpha'}|^2} \partial_{\alpha'} f,$$

$$(B.21) \quad [D_t, \mathfrak{P}] f = \left[D_t, i \frac{A_1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} \right] f = i \left(\frac{D_t A_1}{A_1} + b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t \right) \frac{A_1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} f.$$

We also have

$$(B.22) \quad [D_t, \mathbb{H}] = 2[D_t, \mathbb{P}_H] = [b, \mathbb{H}] \partial_{\alpha'} = \left[\frac{Z_t}{Z_{,\alpha'}}, \mathbb{H} \right] \partial_{\alpha'} \mathbb{P}_H + \left[\frac{\overline{Z}_t}{\overline{Z}_{,\alpha'}}, \mathbb{H} \right] \partial_{\alpha'} \mathbb{P}_A,$$

where the last equality in (B.22) is a consequence of (2.1), (A.3) and (B.9). In general, for operators A, B and C ,

$$(B.23) \quad [A, BC] = [A, B]C + B[A, C].$$

We use the following identities in our computations. We have

$$(B.24) \quad [\mathcal{P}, \mathfrak{D}_t^m] = \sum_{k=0}^{m-1} \mathfrak{D}_t^k [\mathcal{P}, \mathfrak{D}_t] \mathfrak{D}_t^{m-k-1},$$

$$(B.25) \quad \mathbb{P}_H D_t \mathbb{P}_H = \mathbb{P}_H D_t - \mathbb{P}_H D_t \mathbb{P}_A = \mathbb{P}_H D_t - \mathbb{P}_H [b, \mathbb{P}_A] \partial_{\alpha'},$$

and we compute

$$\begin{aligned} (\mathbb{P}_H D_t)^{l+1} \mathbb{P}_H &= \mathbb{P}_H D_t^{l+1} + \sum_{k=0}^l ((\mathbb{P}_H D_t)^{k+1} \mathbb{P}_H D_t^{l-k} - (\mathbb{P}_H D_t)^k \mathbb{P}_H D_t^{l+1-k}) \\ &= \mathbb{P}_H D_t^{l+1} - \sum_{k=0}^l (\mathbb{P}_H D_t)^k \mathbb{P}_H D_t \mathbb{P}_A D_t^{l-k} = \mathbb{P}_H D_t^{l+1} - \sum_{k=0}^l (\mathbb{P}_H D_t)^k \mathbb{P}_H [b, \mathbb{P}_A] \partial_{\alpha'} D_t^{l-k}, \end{aligned}$$

so

$$(B.26) \quad (\mathbb{P}_H D_t)^{l+1} \mathbb{P}_H = \mathbb{P}_H D_t^{l+1} - \sum_{k=0}^l (\mathbb{P}_H D_t)^k \mathbb{P}_H [b, \mathbb{P}_A] \partial_{\alpha'} D_t^{l-k}.$$

A similar computation also gives

$$(B.27) \quad (\mathbb{P}_H D_t)^{l+1} \mathbb{P}_H = D_t^{l+1} \mathbb{P}_H - \sum_{k=0}^l D_t^k [\mathbb{P}_A, b] \partial_{\alpha'} (\mathbb{P}_H D_t)^{l-k} \mathbb{P}_H.$$

Let $\mathfrak{f} := f(\alpha') - f(\beta')$, $\mathfrak{g} := g(\alpha') - g(\beta')$ and $\mathfrak{h} := h(\alpha') - h(\beta')$. We use (B.14) and (B.18) to get

$$(B.28) \quad D_t < f, g, h > = \frac{1}{\pi i} \int \frac{\mathfrak{D}_t(\mathfrak{f} \mathfrak{g} \mathfrak{h})}{(\alpha' - \beta')^2} d\beta' + \frac{1}{\pi i} \int \partial_{\beta'} \mathfrak{D}_t \left(\frac{1}{\alpha' - \beta'} \right) \mathfrak{f} \mathfrak{g} \mathfrak{h} d\beta';$$

and by induction,

$$(B.29) \quad D_t^n < f, g, h > = \frac{1}{\pi i} \sum_{k=0}^n \binom{n}{k} \int \partial_{\beta'} \mathfrak{D}_t^k \left(\frac{1}{\alpha' - \beta'} \right) \mathfrak{D}_t^{n-k}(\mathfrak{f} \mathfrak{g} \mathfrak{h}) d\beta',$$

Similarly,

$$(B.30) \quad D_t^n[f, g; h] = \frac{1}{\pi i} \sum_{k=0}^n \binom{n}{k} \int \partial_{\beta'} \mathfrak{D}_t^k \left(\frac{1}{\alpha' - \beta'} \right) \mathfrak{D}_t^{n-k}((f(\alpha') - f(\beta'))(g(\alpha') - g(\beta'))h(\beta')) d\beta';$$

$$(B.31) \quad D_t^n[f, g; \partial_{\alpha'} h] = \frac{1}{\pi i} \sum_{k=0}^n \binom{n}{k} \int \mathfrak{D}_t^k \left(\frac{(f(\alpha') - f(\beta'))(g(\alpha') - g(\beta'))}{(\alpha' - \beta')^2} \right) \partial_{\beta'} D_t^{n-k} h(\beta') d\beta';$$

$$(B.32) \quad D_t^n[f, \mathbb{H}] \partial_{\alpha'} g = \frac{1}{\pi i} \sum_{k=0}^n \binom{n}{k} \int \mathfrak{D}_t^k \left(\frac{f(\alpha') - f(\beta')}{\alpha' - \beta'} \right) \partial_{\beta'} D_t^{n-k} g(\beta') d\beta';$$

where

$$(B.33) \quad \mathfrak{D}_t \left(\frac{1}{\alpha' - \beta'} \right) = -\frac{b(\alpha') - b(\beta')}{(\alpha' - \beta')^2}, \quad \mathfrak{D}_t^2 \left(\frac{1}{\alpha' - \beta'} \right) = -\frac{D_t b(\alpha') - D_t b(\beta')}{(\alpha' - \beta')^2} + 2 \frac{(b(\alpha') - b(\beta'))^2}{(\alpha' - \beta')^3}, \text{ etc};$$

in particular,

$$(B.34) \quad D_t[f, \mathbb{H}] \partial_{\alpha'} g = [D_t f, \mathbb{H}] \partial_{\alpha'} g + [f, \mathbb{H}] \partial_{\alpha'} D_t g - [f, b; \partial_{\alpha'} g].$$

B.4. Some additional equations. We know by definition (2.7), (2.8) and equation (2.3) that

$$(B.35) \quad \Theta^{(0)} = Q, \quad \Theta^{(1)} = i(Z - \alpha'), \quad \Theta^{(2)} = -i\mathbb{P}_H b = -i\mathbb{P}_H \frac{\overline{Z}_t}{\overline{Z}_{, \alpha'}},$$

$$(B.36) \quad \Theta^{(j+2)} = -i\mathbb{P}_H \left(\frac{1}{|\overline{Z}_{, \alpha'}|^2} \partial_{\alpha'} \Theta^{(j)} \right) + \mathbb{P}_H(G^{(j)}), \quad j \geq 1.$$

This implies

$$(B.37) \quad D_{\alpha'} \Theta^{(0)} = \overline{Z}_t, \quad D_{\alpha'} \Theta^{(1)} = i \left(1 - \frac{1}{\overline{Z}_{, \alpha'}} \right);$$

$$(B.38) \quad b = i \Theta^{(2)} - i \overline{\Theta^{(2)}}, \quad D_t b = i D_t \Theta^{(2)} - i \overline{D_t \Theta^{(2)}}, \quad \text{and}$$

$$(B.39) \quad \mathbb{P}_H \left(D_t b + i \frac{A_1}{|Z_{,\alpha'}|^2} - i \right) = i \mathbb{P}_H \left(\frac{A_1 - 1}{|Z_{,\alpha'}|^2} \right) + i \mathbb{P}_H(G^{(1)}) + i [b, \mathbb{P}_H] \partial_{\alpha'} \overline{\Theta^{(2)}}.$$

We compute, by (B.24), (B.21) that

$$(B.40) \quad \begin{aligned} [\mathcal{P}, D_t^l] \overline{Z}_t &= \sum_{k=0}^{l-1} D_t^{l-1-k} [\mathcal{P}, D_t] D_t^k \overline{Z}_t \\ &= -i \sum_{k=0}^{l-1} D_t^{l-1-k} \left\{ \left(\frac{D_t A_1}{A_1} + b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t \right) \frac{A_1}{|Z_{,\alpha'}|^2} \partial_{\alpha'} D_t^k \overline{Z}_t \right\}. \end{aligned}$$

and we have, by (B.23), (B.17) and (B.27), that

$$(B.41) \quad \begin{aligned} D_{\alpha'} \Theta^{(l+1)} - D_t^l D_{\alpha'} \Theta^{(1)} &= \sum_{k=0}^{l-1} (\mathbb{P}_H D_t)^k \left[\frac{1}{Z_{,\alpha'}}, \mathbb{P}_H \right] \partial_{\alpha'} D_t \Theta^{(l-k)} \\ &+ \sum_{k=0}^{l-1} (\mathbb{P}_H D_t)^k \mathbb{P}_H (D_{\alpha'} Z_t D_{\alpha'} \Theta^{(l-k)}) - \sum_{k=0}^{l-1} D_t^k [\mathbb{P}_A, b] \partial_{\alpha'} (\mathbb{P}_H D_t)^{l-1-k} D_{\alpha'} \Theta^{(1)}. \end{aligned}$$

APPENDIX C. BASIC INEQUALITIES

We will use the following equalities or inequalities in this paper. The first set: Lemma C.1 through Proposition C.14 are either classical results, or simple consequences of definitions and classical results, and some have already appeared in Appendix A of [51]. Lemma C.15 and Proposition C.16 are from Section 5.1 of [51].

Lemma C.1. *For any function $f \in \dot{H}^{1/2}(\mathbb{R})$,*

$$(C.1) \quad \|f\|_{\dot{H}^{1/2}}^2 = \|\mathbb{P}_H f\|_{\dot{H}^{1/2}}^2 + \|\mathbb{P}_A f\|_{\dot{H}^{1/2}}^2;$$

$$(C.2) \quad \int i \partial_{\alpha'} f \overline{f} d\alpha' = \|\mathbb{P}_H f\|_{\dot{H}^{1/2}}^2 - \|\mathbb{P}_A f\|_{\dot{H}^{1/2}}^2.$$

Proposition C.2. *Let $f, g \in C^1(\mathbb{R})$. Then*

$$(C.3) \quad \|fg\|_{\dot{H}^{1/2}} \lesssim \|f\|_{L^\infty} \|g\|_{\dot{H}^{1/2}} + \|g\|_{L^\infty} \|f\|_{\dot{H}^{1/2}};$$

$$(C.4) \quad \|fg\|_{\dot{H}^{1/2}} \lesssim \|f\|_{L^\infty} \|g\|_{\dot{H}^{1/2}} + \|f'\|_{L^2} \|g\|_{L^2};$$

$$(C.5) \quad \|g\|_{\dot{H}^{1/2}} \lesssim \|f^{-1}\|_{L^\infty} (\|fg\|_{\dot{H}^{1/2}} + \|f'\|_{L^2} \|g\|_{L^2}).$$

(C.4) is straightforward from the definition of $\dot{H}^{1/2}$ and Hardy's inequality. The remaining two are from Appendix A of [51].

Proposition C.3 (Sobolev inequality). *Let $f \in C_0^1(\mathbb{R})$. Then*

$$(C.6) \quad \|f\|_{L^\infty}^2 \leq 2 \|f\|_{L^2} \|f'\|_{L^2}, \quad \|f\|_{\dot{H}^{1/2}}^2 \leq \|f\|_{L^2} \|f'\|_{L^2}.$$

Proposition C.4 (Maximum inequality). *Let $1 < p \leq \infty$, Then for all $f \in L^p$,*

$$(C.7) \quad \|M(f)\|_{L^p} \lesssim \|f\|_{L^p}.$$

Proposition C.5 (Hardy's inequalities). *Let $1 < p < \infty$, $f \in C^1(\mathbb{R})$, with $f' \in L^p(\mathbb{R})$. Then*

$$(C.8) \quad \sup_{x \in \mathbb{R}} \int \frac{|f(x) - f(y)|^p}{|x - y|^p} dy \lesssim \|f'\|_{L^p}^p;$$

and

$$(C.9) \quad \iint \frac{|f(x) - f(y)|^{2p}}{|x - y|^{2p}} dx dy \lesssim \|f'\|_{L^p}^{2p}.$$

Let $H \in C^1(\mathbb{R}; \mathbb{R}^d)$, $A_i \in C^1(\mathbb{R})$, $i = 1, \dots, m$, and $F \in C^\infty(\mathbb{R})$. Define

$$(C.10) \quad C_1(A_1, \dots, A_m, f)(x) = \text{pv.} \int F \left(\frac{H(x) - H(y)}{x - y} \right) \frac{\prod_{i=1}^m (A_i(x) - A_i(y))}{(x - y)^{m+1}} f(y) dy.$$

Proposition C.6. *There exist constants $c_1 = c_1(F, \|H'\|_{L^\infty})$, $c_2 = c_2(F, \|H'\|_{L^\infty})$, such that*

1. *For any $f \in L^2$, $A'_i \in L^\infty$, $1 \leq i \leq m$,*

$$(C.11) \quad \|C_1(A_1, \dots, A_m, f)\|_{L^2} \leq c_1 \|A'_1\|_{L^\infty} \dots \|A'_m\|_{L^\infty} \|f\|_{L^2}.$$

2. *For any $f \in L^\infty$, $A'_i \in L^\infty$, $2 \leq i \leq m$, $A'_1 \in L^2$,*

$$(C.12) \quad \|C_1(A_1, \dots, A_m, f)\|_{L^2} \leq c_2 \|A'_1\|_{L^2} \|A'_2\|_{L^\infty} \dots \|A'_m\|_{L^\infty} \|f\|_{L^\infty}.$$

(C.11) is a result of Coifman, McIntosh and Meyer [20]. (C.12) is a consequence of the Tb Theorem, a proof is given in [48].

Let H, A_i, F satisfy the same assumptions as in (C.10). Define

$$(C.13) \quad C_2(A, f)(x) = \int F \left(\frac{H(x) - H(y)}{x - y} \right) \frac{\prod_{i=1}^m (A_i(x) - A_i(y))}{(x - y)^m} \partial_y f(y) dy.$$

The following are consequences of Proposition C.6 and integration by parts.

Proposition C.7. *There exist constants c_3, c_4 and c_5 , depending on F and $\|H'\|_{L^\infty}$, such that*

1. *For any $f \in L^2$, $A'_i \in L^\infty$, $1 \leq i \leq m$,*

$$(C.14) \quad \|C_2(A, f)\|_{L^2} \leq c_3 \|A'_1\|_{L^\infty} \dots \|A'_m\|_{L^\infty} \|f\|_{L^2}.$$

2. *For any $f \in L^\infty$, $A'_i \in L^\infty$, $2 \leq i \leq m$, $A'_1 \in L^2$,*

$$(C.15) \quad \|C_2(A, f)\|_{L^2} \leq c_4 \|A'_1\|_{L^2} \|A'_2\|_{L^\infty} \dots \|A'_m\|_{L^\infty} \|f\|_{L^\infty}.$$

3. *For any $f' \in L^2$, $A_1 \in L^\infty$, $A'_i \in L^\infty$, $2 \leq i \leq m$,*

$$(C.16) \quad \|C_2(A, f)\|_{L^2} \leq c_5 \|A_1\|_{L^\infty} \|A'_2\|_{L^\infty} \dots \|A'_m\|_{L^\infty} \|f'\|_{L^2}.$$

Proposition C.8. *Assume that f, g, h are smooth and decay fast at infinity. Then*

$$(C.17) \quad \|[f, \mathbb{H}]g\|_{L^2} \lesssim \|f\|_{\dot{H}^{1/2}} \|g\|_{L^2};$$

$$(C.18) \quad \|[f, \mathbb{H}]g\|_{L^\infty} \lesssim \|f'\|_{L^2} \|g\|_{L^2};$$

$$(C.19) \quad \|[f, \mathbb{H}]\partial_{\alpha'} g\|_{L^2} \lesssim \|f'\|_{L^2} \|g\|_{\dot{H}^{1/2}};$$

$$(C.20) \quad \|[f, \mathbb{H}]\partial_{\alpha'} g\|_{L^2} \lesssim \|f'\|_{\dot{H}^{1/2}} \|g\|_{L^2};$$

$$(C.21) \quad \|[f, h; \partial_{\alpha'} g]\|_{L^2} \lesssim \|f'\|_{L^2} \|h'\|_{L^\infty} \|g\|_{\dot{H}^{1/2}},$$

$$(C.22) \quad \|[f, h; \partial_{\alpha'} g]\|_{L^2} \lesssim \|f'\|_{L^2} \|h'\|_{\dot{H}^{1/2}} \|g\|_{L^\infty} + \|f'\|_{\dot{H}^{1/2}} \|h'\|_{L^2} \|g\|_{L^\infty}.$$

(C.20) follows from integration by parts, Cauchy-Schwarz inequality and the definition (A.8), and (C.22) follows from integration by parts, Cauchy-Schwarz inequality, Hardy's inequality and the definition (A.8). The remaining inequalities are from Appendix A of [51].

Proposition C.9. *For any f, g, h smooth and decay fast at spatial infinity, we have*

$$(C.23) \quad \|[f, g; h]\|_{L^p} \lesssim \|f'\|_{L^2} \|g'\|_{L^2} \|h\|_{L^p}, \quad 1 \leq p \leq \infty;$$

$$(C.24) \quad \|[f, g; h]\|_{L^2} \lesssim \|f'\|_{L^2} \|g\|_{\dot{H}^{1/2}} \|h\|_{L^\infty};$$

$$(C.25) \quad \|[f, g; h]\|_{L^2} \lesssim \|f\|_{\dot{H}^{1/2}} \|g'\|_{L^\infty} \|h\|_{L^2};$$

$$(C.26) \quad \|[f, g; h]\|_{L^\infty} \lesssim \|f'\|_{L^2} \|g'\|_{L^\infty} \|h\|_{L^2}.$$

(C.23) directly follows from Hölder's inequality and Hardy's inequality. (C.24) and (C.25) are direct consequences of Cauchy-Schwarz inequality and the definition (A.8). (C.26) is from Appendix A of [51].

Proposition C.10. *Assume that $f_i \in C^1(\mathbb{R})$, with $f'_i \in L^{p_i}$, $g \in L^q$, where*

$$\sum_{i=1}^n \frac{1}{p_i} + \frac{1}{q} = \frac{1}{p} + 1, \quad 1 < p_i \leq \infty, \quad 1 \leq q \leq \infty, \quad p > 0.$$

Then

$$(C.27) \quad \left\| \int \frac{\prod_{i=1}^n (f_i(x) - f_i(y))}{(x-y)^n} g(y) dy \right\|_{L^p} \lesssim \prod_{i=1}^n \|f'_i\|_{L^{p_i}} \|g\|_{L^q}.$$

Observe that $\left| \frac{f(x)-f(y)}{x-y} \right| \leq \min\{M(f')(x), M(f')(y)\}$. (C.27) is a direct consequence of the Maximum inequality (C.7) and Hölder's inequality.

We also have the following inequalities for the cubic form $\langle \cdot, \cdot, \cdot \rangle$.

Proposition C.11. *For any f, g, h smooth and decay fast at spatial infinity, we have*

$$(C.28) \quad \|\langle f, g, h \rangle\|_{L^2} \lesssim \|f'\|_{L^2} \|g\|_{L^\infty} \|h\|_{\dot{H}^{1/2}};$$

$$(C.29) \quad \|\langle f, g, h \rangle\|_{L^p} \lesssim \|f'\|_{L^2} \|g'\|_{L^2} \|h\|_{L^p}, \quad 1 \leq p \leq \infty;$$

$$(C.30) \quad \|\langle f, g, h \rangle\|_{L^2} \lesssim \|f'\|_{L^2} \|g\|_{\dot{H}^{1/2}} \|h\|_{\dot{H}^{1/2}};$$

$$(C.31) \quad \|\langle f, g, h \rangle\|_{\dot{H}^{1/2}} \lesssim \|f'\|_{L^2} \|g'\|_{L^2} \|h\|_{\dot{H}^{1/2}};$$

$$(C.32) \quad \|\langle f, g, h \rangle\|_{\dot{H}^1} \lesssim \|f'\|_{L^2} \|g'\|_{L^2} \|h'\|_{L^2};$$

$$(C.33) \quad \|\langle f, g, h \rangle\|_{L^\infty} \lesssim \|f'\|_{L^2} (\|g'\|_{L^2} \|h\|_{\dot{H}^{1/2}} + \|h'\|_{L^2} \|g\|_{\dot{H}^{1/2}}).$$

(C.28), (C.29), and (C.32) are easy consequences of Hölder's inequality, Hardy's inequality, and the definition (A.8). (C.30) and (C.31) follow from interpolation; (C.33) follows from the inequality

$$\|\partial_{\alpha'} \langle f, g, h \rangle\|_{L^1} \leq \|f'\|_{L^2} (\|g'\|_{L^2} \|h\|_{\dot{H}^{1/2}} + \|h'\|_{L^2} \|g\|_{\dot{H}^{1/2}}),$$

which in turn follows from Cauchy-Schwarz inequality and Hardy's inequality.

Proposition C.12. *Assume that f, g, h are holomorphic, i.e. $f = \mathbb{P}_H f$, $g = \mathbb{P}_H g$ and $h = \mathbb{P}_H h$. Then*

$$(C.34) \quad \|\langle f, g, h \rangle\|_{L^\infty} \lesssim \|f'\|_{L^2} \|g'\|_{L^2} \|h\|_{\dot{H}^{1/2}}.$$

Observe that for f, g holomorphic, $[f, g; 1] = 0$. The same argument for (C.33) gives (C.34).

Proposition C.13. *Assume that f, g, h are smooth and decay fast at infinity, $b = b(x, y)$ is bounded.*

Then

$$(C.35) \quad \left\| \int b(x, y) \frac{(f(x) - f(y))(g(x) - g(y))(h(x) - h(y))}{(x-y)^2} dy \right\|_{L^2} \lesssim \|b\|_{L^\infty(\mathbb{R}^2)} \|f'\|_{L^2} \|g'\|_{L^2} \|h\|_{L^2};$$

$$(C.36) \quad \left\| \int b(x, y) \frac{(f(x) - f(y))(g(x) - g(y))(h(x) - h(y))}{(x-y)^2} dy \right\|_{L^2} \lesssim \|b\|_{L^\infty(\mathbb{R}^2)} \|f'\|_{L^2} \|g\|_{\dot{H}^{1/2}} \|h\|_{\dot{H}^{1/2}}.$$

(C.35) follows from Cauchy-Schwarz inequality and Hardy's inequality; (C.36) follows from interpolation.

Proposition C.14. *Assume that $f_k \in \dot{H}^{1/2}(\mathbb{R})$, $k = 1, 2, 3, 4$. Then*

$$(C.37) \quad \left| \int \frac{\Pi_{k=1}^4(f_k(x) - f_k(y))}{(x-y)^2} dx dy \right| \lesssim \Pi_{k=1}^4 \|f_k\|_{\dot{H}^{1/2}}.$$

Proof. Observe that by symmetry, we can write

$$\int \frac{\Pi_{k=1}^4(f_k(x) - f_k(y))}{(x-y)^2} dx dy = 2 \int f_1(x) \int \frac{\Pi_{k=2}^4(f_k(x) - f_k(y))}{(x-y)^2} dy dx.$$

By (C.30), we have, for $f_k \in H^1(\mathbb{R})$, $k = 1, 2$,

$$\left| \int \frac{\Pi_{k=1}^4(f_k(x) - f_k(y))}{(x-y)^2} dx dy \right| \lesssim \|f_1\|_{L^2} \|f_2'\|_{L^2} \|f_3\|_{\dot{H}^{1/2}} \|f_4\|_{\dot{H}^{1/2}};$$

similarly we also have

$$\left| \int \frac{\Pi_{k=1}^4(f_k(x) - f_k(y))}{(x-y)^2} dx dy \right| \lesssim \|f_1'\|_{L^2} \|f_2\|_{L^2} \|f_3\|_{\dot{H}^{1/2}} \|f_4\|_{\dot{H}^{1/2}}.$$

(C.37) then follows by interpolation. \square

The equality and inequalities in Lemma C.15 and Proposition C.16 are from Section 5.1 and Lemma 6.3 of [51].

Lemma C.15 (cf. Lemma 5.4, [51]). *Assume that $f, g, f_1, g_1 \in H^1(\mathbb{R})$ are the boundary values of some holomorphic functions on \mathcal{P}_- . Then*

$$(C.38) \quad \int \partial_{\alpha'} \mathbb{P}_A(\bar{f}g)(\alpha') f_1(\alpha') \bar{g}_1(\alpha') d\alpha' = -\frac{1}{2\pi i} \iint \frac{(\bar{f}(\alpha') - \bar{f}(\beta'))(f_1(\alpha') - f_1(\beta'))}{(\alpha' - \beta')^2} g(\beta') \bar{g}_1(\alpha') d\alpha' d\beta'.$$

Proposition C.16 (cf. Proposition 5.6, Lemma 6.3, [51]). *Assume that $f, g \in H^1(\mathbb{R})$. We have*

$$(C.39) \quad \|[f, \mathbb{H}]g\|_{\dot{H}^{1/2}} \lesssim \|f\|_{\dot{H}^{1/2}} (\|g\|_{L^\infty} + \|\mathbb{H}g\|_{L^\infty});$$

$$(C.40) \quad \|[f, \mathbb{H}]g\|_{\dot{H}^{1/2}} \lesssim \|\partial_{\alpha'} f\|_{L^2} \|g\|_{L^2};$$

$$(C.41) \quad \|[f, \mathbb{H}]\partial_{\alpha'} g\|_{\dot{H}^{1/2}} \lesssim \|g\|_{\dot{H}^{1/2}} (\|\partial_{\alpha'} f\|_{L^\infty} + \|\partial_{\alpha'} \mathbb{H}f\|_{L^\infty});$$

$$(C.42) \quad \|f\|_{L^4}^2 \lesssim \|f\|_{L^2} \|f\|_{\dot{H}^{1/2}}.$$

Finally, we need the following inequalities for our proof. Let $f_x^y \mathbf{b} = \frac{f_y^x \mathbf{b}(\alpha) d\alpha}{x-y}$. We have

Proposition C.17. *Let $\mathbf{b} \in BMO(\mathbb{R})$, $f_1, \dots, f_n \in C^1(\mathbb{R})$ such that $f_1', \dots, f_n' \in L^2(\mathbb{R})$, and $g \in L^2(\mathbb{R})$. Then*

$$(C.43) \quad \left\| \int \left(\mathbf{b}(y) - \int_x^y \mathbf{b} \right) \frac{\Pi_{i=1}^2(f_i(x) - f_i(y))}{(x-y)^2} dy \right\|_{L^\infty} \lesssim \|\mathbf{b}\|_{BMO} \|f_1'\|_{L^2} \|f_2'\|_{L^2};$$

$$(C.44) \quad \left\| \int \left(\mathbf{b}(y) - \int_x^y \mathbf{b} \right)^2 \frac{\Pi_{i=1}^2(f_i(x) - f_i(y))}{(x-y)^2} dy \right\|_{L^\infty} \lesssim \|\mathbf{b}\|_{BMO}^2 \|f_1'\|_{L^2} \|f_2'\|_{L^2};$$

$$(C.45) \quad \left\| \int \left(\mathbf{b}(x) + \mathbf{b}(y) - 2 \int_x^y \mathbf{b} \right) \frac{\Pi_{i=1}^2(f_i(x) - f_i(y))}{(x-y)^2} g(y) dy \right\|_{L^2} \lesssim \|\mathbf{b}\|_{BMO} \|f_1'\|_{L^2} \|f_2'\|_{L^2} \|g\|_{L^2};$$

$$(C.46) \quad \left\| \int \frac{(\mathbf{b}(x) - \mathbf{b}(y)) \Pi_{i=1}^2(f_i(x) - f_i(y))}{(x-y)^2} g(y) dy \right\|_{L^2} \lesssim \|\mathbf{b}\|_{BMO} \|f_1'\|_{L^2} \|f_2'\|_{L^2} \|g\|_{L^2};$$

$$(C.47) \quad \left\| \int \left(\mathbf{b}(x) + \mathbf{b}(y) - 2 \int_x^y \mathbf{b} \right) \frac{\prod_{i=1}^3 (f_i(x) - f_i(y))}{(x-y)^3} dy \right\|_{L^2} \lesssim \|\mathbf{b}\|_{BMO} \|f'_1\|_{L^2} \|f'_2\|_{L^2} \|f'_3\|_{L^2};$$

$$(C.48) \quad \left\| \int \frac{(\mathbf{b}(x) - \mathbf{b}(y)) \prod_{i=1}^3 (f_i(x) - f_i(y))}{(x-y)^3} dy \right\|_{L^2} \lesssim \|\mathbf{b}\|_{BMO} \|f'_1\|_{L^2} \|f'_2\|_{L^2} \|f'_3\|_{L^2};$$

and for $n \geq 3$,

$$(C.49) \quad \left\| \int \frac{(\mathbf{b}(x) - \mathbf{b}(y)) \prod_{i=1}^n (f_i(x) - f_i(y))}{(x-y)^n} g(y) dy \right\|_{L^2} \lesssim \|\mathbf{b}\|_{BMO} \|f'_1\|_{L^2} \|f'_2\|_{L^2} \|g \prod_{i=3}^n M(f'_i)\|_{L^2},$$

where $M(f)$ is the Hardy-Littlewood maximum function of f .

Remark C.18. Observe that $\dot{H}^{1/2}(\mathbb{R}) \subset BMO(\mathbb{R})$, and $\|\mathbf{b}\|_{BMO} \lesssim \|\mathbf{b}\|_{\dot{H}^{1/2}}$ for all $\mathbf{b} \in \dot{H}^{1/2}$. We will primarily use Proposition C.17 for $\mathbf{b} \in \dot{H}^{1/2}(\mathbb{R})$. We sometimes also use Proposition C.17 for $\mathbf{b} \in L^\infty(\mathbb{R})$, with the inequality $\|\mathbf{b}\|_{BMO} \lesssim \|\mathbf{b}\|_{L^\infty}$.

Proof. It suffices to prove for the case that $f_1 = f_2$. We begin with the following lemmas.

Lemma C.19 (Schur test). *Let*

$$Tg(x) = \int e^{\mathbf{b}(x)} K(x, y) e^{\mathbf{b}(y)} g(y) dy.$$

where \mathbf{b} is a real valued measurable function, K is measurable on \mathbb{R}^2 . Assume that

$$\max\left\{ \sup_x \int e^{2\mathbf{b}(y)} |K(x, y)| dy, \sup_y \int e^{2\mathbf{b}(x)} |K(x, y)| dx \right\} := M < \infty.$$

Then for any $g \in L^2$,

$$\|Tg\|_{L^2} \leq M \|g\|_{L^2}.$$

Let $\mathbf{b} \in BMO(\mathbb{R})$ be real valued, and $h(x) = \int_0^x e^{\mathbf{b}(y)} dy$. We know there is a constant $\gamma_0 > 0$, such that for any $\mathbf{b} \in BMO(\mathbb{R})$ satisfying $\|\mathbf{b}\|_{BMO} \leq \gamma_0$, $e^{\mathbf{b}}$ is a A_2 weight and $U_{h^{-1}}$ is a bounded map from $BMO(\mathbb{R})$ to $BMO(\mathbb{R})$, with $\|f \circ h^{-1}\|_{BMO} \leq 2 \|f\|_{BMO}$ for all $f \in BMO(\mathbb{R})$.³³ And for any real valued BMO function \mathbf{b} with $\|\mathbf{b}\|_{BMO} \leq \gamma < 1$, $z(x) = \int_0^x e^{i\mathbf{b}(\alpha)} d\alpha$ defines a chord-arc curve, cf. [18, 19], with

$$(C.50) \quad (1 - \gamma)|x - y| \leq |z(x) - z(y)| \leq |x - y|, \quad \text{for all } x, y \in \mathbb{R}.$$

We have

Lemma C.20. *Let $\mathbf{b} \in BMO(\mathbb{R})$ be real valued with $\|\mathbf{b}\|_{BMO} \leq \gamma_0/2$, and $h'(x) = e^{\mathbf{b}(x)}$. For any f such that $f' \in L^2(\mathbb{R})$, there is a constant $c(\gamma_0) > 0$, such that*

$$(C.51) \quad \sup_x \int e^{2\mathbf{b}(y)} \left| \frac{f(x) - f(y)}{h(x) - h(y)} \right|^2 dy \leq c(\gamma_0) \|f'\|_{L^2}^2.$$

Proof. We know

$$f(x) - f(y) = \int_y^x f'(\alpha) d\alpha = \int_{h(y)}^{h(x)} f' \circ h^{-1}(\beta) (h^{-1})'(\beta) d\beta,$$

hence

$$\sup_x \left| \frac{f(x) - f(y)}{h(x) - h(y)} \right| \leq M (f' \circ h^{-1} (h^{-1})') (h(y))$$

³³These are consequences of John-Nirenberg's inequality and the theory of A_p weights, cf. [18, 19].

where $M(g)$ is the Hardy-Littlewood Maximal function of g . Because $e^{\mathbf{b} \circ h^{-1}}$ is a A_2 weight, we have

$$(C.52) \quad \begin{aligned} \sup_x \int e^{2\mathbf{b}(y)} \left| \frac{f(x) - f(y)}{h(x) - h(y)} \right|^2 dy &\leq \int e^{2\mathbf{b}(y)} M^2(f' \circ h^{-1}(h^{-1})')(h(y)) dy \\ &= \int e^{\mathbf{b} \circ h^{-1}(y)} M^2(f' \circ h^{-1}(h^{-1})')(y) dy \lesssim \int e^{\mathbf{b} \circ h^{-1}(y)} |f' \circ h^{-1}(h^{-1})'|^2(y) dy = \|f'\|_{L^2}^2. \end{aligned}$$

This proves Lemma C.20. \square

Now let \mathbf{b} be a BMO function satisfying $\|\mathbf{b}\|_{BMO} \leq 1$, and z be a complex number in $D := \{|z| < \min\{\frac{1}{4}, \frac{\gamma_0}{2}\}\}$. Let $h_z(x) = \int_0^x e^{z\mathbf{b}(\alpha)} d\alpha$ and

$$(C.53) \quad T_1(f, \mathbf{b}; z)(x) := \int e^{2z\mathbf{b}(y)} \left(\frac{f(x) - f(y)}{h_z(x) - h_z(y)} \right)^2 dy,$$

$$(C.54) \quad T_2(f, g, \mathbf{b}; z)(x) := \int e^{z\mathbf{b}(x)} e^{z\mathbf{b}(y)} \left(\frac{f(x) - f(y)}{h_z(x) - h_z(y)} \right)^2 g(y) dy.$$

By Lemma C.20, and (C.50), we have for all f satisfying $f' \in L^2$,

$$(C.55) \quad \|T_1(f, \mathbf{b}; z)\|_{L^\infty} \leq c(\gamma_0) \|f'\|_{L^2}^2,$$

and by Lemmas C.19, C.20, and (C.50), we have for all f, g , satisfying $f', g \in L^2$,

$$(C.56) \quad \|T(f, g, \mathbf{b}; z)\|_{L^2(\mathbb{R})} \leq c(\delta_o) \|f'\|_{L^2}^2 \|g\|_{L^2},$$

where $c(\gamma_0) > 0$ is a constant depending only on γ_0 . Let $q_1 \in L^1(\mathbb{R})$, $q_2 \in L^2(\mathbb{R})$, and

$$(C.57) \quad F_1(z) := \int q_1(x) T_1(f, \mathbf{b}; z)(x) dx,$$

$$(C.58) \quad F_2(z) := \int q_2(x) T_2(f, g, \mathbf{b}; z)(x) dx.$$

Then F_1, F_2 are holomorphic functions in the domain D , satisfying

$$(C.59) \quad |F_1(z)| \leq c(\gamma_0) \|q_1\|_{L^1} \|f'\|_{L^2}^2, \quad |F_2(z)| \leq c(\gamma_0) \|q_2\|_{L^2} \|f'\|_{L^2}^2 \|g\|_{L^2}, \quad \text{for all } z \in D.$$

And by Cauchy integral theorem, we have

$$(C.60) \quad |F_1'(0)| \lesssim c(\gamma_0) \|q_1\|_{L^1} \|f'\|_{L^2}^2, \quad |F_1''(0)| \lesssim c(\gamma_0) \|q_1\|_{L^1} \|f'\|_{L^2}^2,$$

$$(C.61) \quad \text{and} \quad |F_2'(0)| \lesssim c(\gamma_0) \|q_2\|_{L^2} \|f'\|_{L^2}^2 \|g\|_{L^2}.$$

We compute $F_1'(0), F_1''(0)$, and obtain

$$(C.62) \quad \begin{aligned} F_1'(0) &= 2 \iint q_1(x) \left(\mathbf{b}(y) - \int_x^y \mathbf{b} \right) \frac{(f(x) - f(y))^2}{(x - y)^2} dy dx \\ F_1''(0) &= \iint \frac{q_1(x) (f(x) - f(y))^2}{(x - y)^2} \left(4 \left(\mathbf{b}(y) - \int_x^y \mathbf{b} \right)^2 - 2 \int_x^y (\mathbf{b}(\alpha) - \int_x^y \mathbf{b})^2 d\alpha \right) dx dy. \end{aligned}$$

Because $\sup_{x,y} \int_x^y |\mathbf{b}(\alpha) - \int_x^y \mathbf{b}|^2 d\alpha \lesssim \|\mathbf{b}\|_{BMO}^2 \leq 1$ and by Hardy's inequality,

$$\iint |q_1(x)| \left| \frac{(f(x) - f(y))^2}{(x - y)^2} \right| dx dy \lesssim \|q_1\|_{L^1} \|f'\|_{L^2}^2,$$

so

$$\left| \iint \frac{q_1(x) (f(x) - f(y))^2}{(x - y)^2} \left(\int_x^y (\mathbf{b}(\alpha) - \int_x^y \mathbf{b})^2 d\alpha \right) dx dy \right| \lesssim \|q_1\|_{L^1} \|f'\|_{L^2}^2,$$

(C.43) and (C.44) then follow from (C.60) and (C.62). We compute

$$(C.63) \quad F'_2(0) = \iint q_2(x) \left(\mathbf{b}(x) + \mathbf{b}(y) - 2 \int_x^y \mathbf{b} \right) \frac{(f(x) - f(y))^2}{(x - y)^2} g(y) dy dx.$$

By (C.61) this gives us (C.45).

Now the Schur test, (C.44) and Hardy's inequality yields

$$(C.64) \quad \left\| \int \left(\mathbf{b}(y) - \int_x^y \mathbf{b} \right) \frac{(f(x) - f(y))^2}{(x - y)^2} g(y) dy \right\|_{L^2} \lesssim \|\mathbf{b}\|_{BMO} \|f'\|_{L^2}^2 \|g\|_{L^2}.$$

This together with (C.45) gives (C.46).

(C.47), (C.48), and (C.49) can be proved similarly by considering

$$(C.65) \quad T_3(f, f_3, \mathbf{b}; z) := \int e^{z\mathbf{b}(x)} e^{z\mathbf{b}(y)} \left(\frac{f(x) - f(y)}{h_z(x) - h_z(y)} \right)^2 \frac{f_3(x) - f_3(y)}{x - y} dy,$$

$$(C.66) \quad T_4(f, f_3, \dots, f_n, g, \mathbf{b}; z) := \int e^{z\mathbf{b}(x)} e^{z\mathbf{b}(y)} \left(\frac{f(x) - f(y)}{h_z(x) - h_z(y)} \right)^2 \frac{\prod_{i=3}^n (f_i(x) - f_i(y))}{(x - y)^{n-2}} g(y) dy,$$

and using the fact that

$$\left| \frac{f_i(x) - f_i(y)}{x - y} \right| \leq M(f'_i)(y), \quad 3 \leq i \leq n.$$

We omit the details. □

APPENDIX D. MAIN QUANTITIES CONTROLLED BY $L(t)$ AND $\left\| \partial_{\alpha'} \frac{1}{Z_{\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}$

Assume that

$$L(t) = \|Z_{t,\alpha'}\|_{L^2} + \left\| \frac{1}{Z_{\alpha'}} \right\|_{\dot{H}^{1/2}} + \|Z_{t,\alpha'\alpha'}\|_{L^2} + \left\| \partial_{\alpha'} \frac{1}{Z_{\alpha'}} \right\|_{\dot{H}^{1/2}} \leq 2\epsilon, \quad \left\| \frac{1}{Z_{\alpha'}} - 1 \right\|_{L^\infty} \leq 1 - \delta < 1.$$

We have shown in §4.3.1 and §4.3.2 that the following quantities are controlled by ϵ :

$$(D.1) \quad \begin{aligned} & \|Z_{t,\alpha'}\|_{L^\infty}, \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}, \left\| \partial_{\alpha'} \frac{1}{Z_{\alpha'}} \right\|_{L^2}, \|Z_{tt}\|_{\dot{H}^{1/2}}, \|b_{\alpha'}\|_{L^2}, \left\| \partial_{\alpha'} \Theta^{(2)} \right\|_{L^2}, \|D_t^2 Z_t\|_{L^2}, \|b_{\alpha'}\|_{L^\infty}, \\ & \left\| D_t \frac{1}{Z_{\alpha'}} \right\|_{L^\infty}, \|Z_{ttt}\|_{L^\infty}, \|\partial_{\alpha'} b_{\alpha'}\|_{L^2}, \left\| \partial_{\alpha'} D_t \frac{1}{Z_{\alpha'}} \right\|_{L^2}, \|Z_{tt,\alpha'}\|_{\dot{H}^{1/2}}, \|\partial_{\alpha'} D_t b\|_{\dot{H}^{1/2}}, \\ & \left\| D_t^2 \frac{1}{Z_{\alpha'}} \right\|_{\dot{H}^{1/2}}, \|Z_{ttt,\alpha'}\|_{L^2}, \|\partial_{\alpha'} D_t^2 b\|_{L^2}, \left\| D_t^3 \frac{1}{Z_{\alpha'}} \right\|_{L^2}, \|D_t^2 Z_{tt}\|_{\dot{H}^{1/2}}, \|D_t^3 Z_{tt}\|_{L^2}, \\ & \left\| \partial_{\alpha'} \Theta^{(4)} \right\|_{L^2}, \left\| \Theta^{(5)} \right\|_{\dot{H}^{1/2}}, \left\| \partial_{\alpha'} \Theta^{(3)} \right\|_{\dot{H}^{1/2}}, \left\| \partial_{\alpha'} D_t \Theta^{(3)} \right\|_{L^2}, \|D_t^2 \Theta^{(3)}\|_{\dot{H}^{1/2}}, \|D_t D_{\alpha'} \Theta^{(3)}\|_{L^2}; \end{aligned}$$

the following quantities are controlled by ϵ^2 :

$$(D.2) \quad \begin{aligned} & \|A_1 - 1\|_{L^\infty}, \|A_1\|_{\dot{H}^{1/2}}, \|D_t A_1\|_{L^2}, \|b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t\|_{L^2}, \|\mathcal{P} \bar{Z}_t\|_{L^2}, \\ & \|\partial_{\alpha'} A_1\|_{\dot{H}^{1/2}}, \|D_t(b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t)\|_{\dot{H}^{1/2}}, \|D_t(b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t)\|_{L^\infty}, \\ & \|D_t^2 A_1\|_{\dot{H}^{1/2}}, \|D_t^2 A_1\|_{L^\infty}, \|[D_t, \mathcal{P}] \bar{Z}_{ttt}\|_{L^2}, \|D_t [D_t, \mathcal{P}] \bar{Z}_{tt}\|_{L^2}, \|D_t^2 [D_t, \mathcal{P}] \bar{Z}_t\|_{L^2}, \\ & \|D_t^3(b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t)\|_{L^2}, \left\| \partial_{\alpha'} \left(D_t^3 b - 2 \operatorname{Re} \frac{D_t^3 Z_t}{Z_{\alpha'}} \right) \right\|_{L^2}, \|D_t^4 A_1\|_{L^2}, \|\mathcal{P} D_t^3 \bar{Z}_t\|_{L^2}; \end{aligned}$$

the following quantities are controlled by $\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}}$:

$$(D.3) \quad \begin{aligned} & \|b_{\alpha'}\|_{\dot{H}^{1/2}}, \left\| D_t \frac{1}{Z_{,\alpha'}} \right\|_{\dot{H}^{1/2}}, \left\| \partial_{\alpha'} \Theta^{(2)} \right\|_{\dot{H}^{1/2}}, \left\| D_{\alpha'} \Theta^{(2)} \right\|_{\dot{H}^{1/2}}, \|Z_{tt,\alpha'}\|_{L^2}, \|Z_{ttt}\|_{\dot{H}^{1/2}}, \\ & \left\| \partial_{\alpha'} D_t b \right\|_{L^2}, \|D_t D_{\alpha'} Z_t\|_{L^2}, \|D_t^3 Z_t\|_{L^2}, \left\| D_t^2 \frac{1}{Z_{,\alpha'}} \right\|_{L^2}, \\ & \left\| \partial_{\alpha'} \Theta^{(3)} \right\|_{L^2}, \|D_t D_{\alpha'} \Theta^{(2)}\|_{L^2}, \left\| \Theta^{(4)} \right\|_{\dot{H}^{1/2}}, \|D_t \Theta^{(3)}\|_{\dot{H}^{1/2}}, \|D_t^2 \Theta^{(2)}\|_{\dot{H}^{1/2}}; \end{aligned}$$

the following quantities are controlled by $\epsilon \left(\left\| \partial_{\alpha'} \frac{1}{Z_{,\alpha'}} \right\|_{L^2} + \|Z_{t,\alpha'}\|_{\dot{H}^{1/2}} \right)$:

$$(D.4) \quad \begin{aligned} & \|b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t\|_{\dot{H}^{1/2}}, \|b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t\|_{L^\infty}, \|\partial_{\alpha'} A_1\|_{L^2}, \|D_t A_1\|_{\dot{H}^{1/2}}, \|\mathcal{P} \bar{Z}_t\|_{\dot{H}^{1/2}}, \\ & \|D_t A_1\|_{L^\infty}, \|\mathcal{P} \bar{Z}_t\|_{L^\infty}, \|D_t(b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t)\|_{L^2}, \|D_t^2 A_1\|_{L^2}, \|\mathcal{P} \bar{Z}_{tt}\|_{L^2}, \|[\mathcal{P}, D_t] \bar{Z}_t\|_{L^2}, \\ & \|D_t^2(b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t)\|_{L^2}, \|D_t^3 A_1\|_{L^2}, \left\| \partial_{\alpha'} \left(D_t^2 b - 2 \operatorname{Re} \frac{Z_{ttt}}{Z_{,\alpha'}} \right) \right\|_{L^2}, \|\mathcal{P} D_t^2 \bar{Z}_t\|_{L^2}, \\ & \|[D_t, \mathcal{P}] \bar{Z}_{tt}\|_{L^2}, \|[D_t^2, \mathcal{P}] \bar{Z}_t\|_{L^2}, \|\partial_{\alpha'}(b_{\alpha'} - 2 \operatorname{Re} D_{\alpha'} Z_t)\|_{L^2}, \|\partial_{\alpha'} D_t A_1\|_{L^2}. \end{aligned}$$

REFERENCES

- [1] A. Ai, M. Ifrim & D. Tataru *Two dimensional gravity waves at low regularity I: energy estimates* arXiv:1910.05323 (Cited on 2, 16.)
- [2] A. Ai, M. Ifrim & D. Tataru *Two dimensional gravity waves at low regularity II: global solutions* arXiv:2009.11513 (Cited on 2.)
- [3] S. Agrawal *Rigidity of singularities of 2d gravity water waves* J. Diff. Equ. 268 (2020) no.3, 1220-1249 (Cited on 2.)
- [4] S. Agrawal *Angled crested type water waves with surface tension: Wellposedness of the problem* To appear in Comm. Math. Phys. (Cited on 2, 19.)
- [5] S. Agrawal *Angled crested type water waves with surface tension II: Zero surface tension limit* arXiv:2009.13469 (Cited on 2.)
- [6] T. Alazard, N. Burq & C. Zuily *On the Cauchy problem for gravity water waves*. Invent. Math. Vol.198 (2014) pp.71-163 (Cited on 2.)
- [7] T. Alazard, N. Burq & C. Zuily *Strichartz estimates and the Cauchy problem for the gravity water waves equations*. Mem. Amer. Math. Soc. 256 (2018), no. 1229 (Cited on 2.)
- [8] T. Alazard & J.-M. Delort *Global solutions and asymptotic behavior for two dimensional gravity water waves* Ann. Sci. Éc. Norm. Supér. (4) 48 (2015), no. 5, 1149-1238. (Cited on 2, 5, 6, 16.)
- [9] D. Ambrose, N. Masmoudi *The zero surface tension limit of two-dimensional water waves*. Comm. Pure Appl. Math. 58 (2005), no. 10, 1287-1315 (Cited on 2.)
- [10] T. Beale, T. Hou & J. Lowengrub *Growth rates for the linearized motion of fluid interfaces away from equilibrium* Comm. Pure Appl. Math. 46 (1993), no.9, 1269-1301. (Cited on 1.)
- [11] M. Berti & J.-M. Delort *Almost global existence of solutions for capillary-gravity water wave equations with periodic spatial boundary conditions* arXiv:1702.04674 (Cited on 2.)
- [12] M. Berti, R. Feola & F. Pusateri *Birkhoff normal form and long time existence for periodic gravity water waves* arXiv:1810.11549 (Cited on 2, 3, 16.)
- [13] L. Bieri, S. Miao, S. Shahshahani, & S. Wu *On the Motion of a Self-Gravitating Incompressible Fluid with Free Boundary* Comm. Math. Phys. Vol. 355, no. 1, (2017), 161-243. (Cited on 2, 4, 6.)
- [14] G. Birkhoff *Helmholtz and Taylor instability* Proc. Symp. in Appl. Math. Vol. XIII, pp.55-76. (Cited on 1.)
- [15] G. Chen, Q. Su *Nonlinear modulational instability of the Stokes waves in 2d full water waves* arXiv:2012.15071, (2020) (Cited on 6.)
- [16] D. Christodoulou, H. Lindblad *On the motion of the free surface of a liquid* Comm. Pure Appl. Math. 53 (2000) no. 12, 1536-1602 (Cited on 2.)
- [17] D. Coutand, S. Shkoller *Wellposedness of the free-surface incompressible Euler equations with or without surface tension* J. AMS. 20 (2007), no. 3, 829-930. (Cited on 2.)
- [18] R.R. Coifman, Y. Meyer *Nonlinear harmonic analysis* Beijing Lectures in Harmonic Analysis, ed. by E. M. Stein, 1986. (Cited on 55.)
- [19] R.R. Coifman, Y. Meyer *Laurentiev's curves and conformal mappings* Rep.5, 1983, Mittag-Leffler Inst. Sweden. (Cited on 55.)
- [20] R. Coifman, A. McIntosh and Y. Meyer *L'intégrale de Cauchy définit un opérateur borne sur L^2 pour les courbes lipschitziennes* Annals of Math, 116 (1982), 361-387. (Cited on 52.)

- [21] W. Craig *An existence theory for water waves and the Boussinesq and Korteweg-devries scaling limits* Comm. in P. D. E. 10(8), 1985 pp.787-1003 (Cited on 1.)
- [22] W. Craig & P. Worfolk *An integrable normal form for water waves in infinite depth* Physica D 84 (1995) 513-531 (Cited on 2, 3.)
- [23] W. Craig & C. Sulem *Mapping properties of normal forms transformations for water waves* Boll. Unione Mat. Ital. 9, no. 2, 289-318, 2016. (Cited on 2, 3.)
- [24] Y. Deng, A. D. Ionescu, B. Pausader, & F. Pusateri *Global solutions of the gravity-capillary water wave system in 3 dimensions* Acta Math. 219 (2017), no. 2, 213-402. (Cited on 2, 16.)
- [25] D. G. Ebin. *The equations of motion of a perfect fluid with free boundary are not well posed* Comm. P. D. E. 12(10), 1987, pp.1175-1201. (Cited on 1.)
- [26] P. Germain, N. Masmoudi, & J. Shatah *Global solutions of the gravity water wave equation in dimension 3* Ann. of Math (2) 175 (2012), no.2, 691-754. (Cited on 2, 16.)
- [27] J. Hunter, M. Ifrim & D. Tataru *Two dimensional water waves in holomorphic coordinates* Comm. Math. Phys. 346 (2016), no. 2, 483-552. (Cited on 2, 5, 6, 8, 16.)
- [28] M. Ifrim & D. Tataru *Two dimensional water waves in holomorphic coordinates II: global solutions* Bull. Soc. Math. France 144 (2016), no. 2, 369-394. (Cited on 2, 5, 16.)
- [29] T. Iguchi *Well-posedness of the initial value problem for capillary-gravity waves* Funkcial. Ekvac. 44 (2001) no. 2, 219-241. (Cited on 2.)
- [30] A. Ionescu & F. Pusateri. *Global solutions for the gravity water waves system in 2d*, Invent. Math. 199 (2015), no. 3, 653-804. (Cited on 2, 5, 6, 16.)
- [31] R. Kinsey & S. Wu *A priori estimates for two-dimensional water waves with angled crests* Cambridge J. Math. vol.6, no.2, (2018), pp.93-181 (Cited on 2, 4, 6, 49.)
- [32] D. Lannes *Well-posedness of the water-wave equations* J. Amer. Math. Soc. 18 (2005), 605-654 (Cited on 2.)
- [33] T. Levi-Civita. *Détermination rigoureuse des ondes permanentes d'ampleur finie*. Math. Ann., 93(1), 1925. pp.264-314 (Cited on 1.)
- [34] H. Lindblad *Well-posedness for the motion of an incompressible liquid with free surface boundary* Ann. of Math. 162 (2005), no. 1, 109-194. (Cited on 2.)
- [35] V. I. Nalimov *The Cauchy-Poisson Problem* (in Russian), Dynamika Splosh. Sredy 18, 1974, pp. 104-210. (Cited on 1.)
- [36] Sir Issac Newton *Philosophi Naturalis Principia Mathematica* 1726 (Cited on 1.)
- [37] M. Ogawa, A. Tani *Free boundary problem for an incompressible ideal fluid with surface tension* Math. Models Methods Appl. Sci. 12, (2002), no.12, 1725-1740. (Cited on 2.)
- [38] E. M. Stein *Singular integrals and differentiability properties of functions* Princeton University Press, 1970. (Cited on .)
- [39] J. Shatah, C. Zeng *Geometry and a priori estimates for free boundary problems of the Euler's equation* Comm. Pure Appl. Math. V. 61. no.5 (2008) pp.698-744 (Cited on 2.)
- [40] Q. Su *Long time behavior of 2d water waves with point vortices*, To appear in Comm. Math. Phys. (Cited on 2, 6.)
- [41] Q. Su *On the transition of the Rayleigh-Taylor instability in 2d water waves* arXiv:2007.13849 (Cited on 2.)
- [42] G. G. Stokes. *On the theory of oscillatory waves*. Trans. Cambridge Philos. Soc., 8: 1847, pp.441- 455. (Cited on 1.)
- [43] G. I. Taylor *The instability of liquid surfaces when accelerated in a direction perpendicular to their planes I*. Proc. Roy. Soc. London A 201, 1950, 192-196 (Cited on 1.)
- [44] X. Wang *Global infinite energy solutions for the 2D gravity water waves system* Comm. Pure Appl. Math. 71 (2018), no. 1, 90-162. (Cited on 2, 16.)
- [45] X. Wang *Global solution for the 3D gravity water waves system above a flat bottom* Adv. Math. 346 (2019), 805-886. (Cited on 2.)
- [46] S. Wu *Well-posedness in Sobolev spaces of the full water wave problem in 2-D* Inventiones Mathematicae 130, 1997, pp. 39-72 (Cited on 1, 4, 6, 7, 47.)
- [47] S. Wu *Well-posedness in Sobolev spaces of the full water wave problem in 3-D* Journal of the AMS. 12. no.2 (1999), pp. 445-495. (Cited on 1.)
- [48] S. Wu *Almost global wellposedness of the 2-D full water wave problem* Invent. Math, 177, (2009), no.1, pp. 45-135. (Cited on 2, 4, 5, 6, 9, 16, 52.)
- [49] S. Wu *Global wellposedness of the 3-D full water wave problem* Invent. Math. 184 (2011), no.1, pp.125-220. (Cited on 2, 4, 6, 16.)
- [50] S. Wu *Wellposedness and singularities of the water wave equations* Notes of the lectures given at the Newton Institute, Cambridge, UK, Aug. 2014. (Cited on 6, 7, 47.)
- [51] S. Wu *Wellposedness of the 2d full water wave equation in a regime that allows for non- C^1 interfaces* Invent. Math. 217 (2019), no.2, pp.241-375 (Cited on 2, 4, 6, 7, 47, 49, 51, 52, 53, 54.)
- [52] H. Yoshida *Gravity waves on the free surface of an incompressible perfect fluid of finite depth*, RIMS Kyoto 18, 1982, pp. 49-96 (Cited on 1.)
- [53] V. E. Zakharov, *Stability of periodic waves of finite amplitude on the surface of a deep fluid* J. Appl. Mech. Tech. Phys. Vol. 9, 1968, 190-194. (Cited on 2.)
- [54] A. I. Dyachenko & V.E. Zakharov *Is free-surface hydrodynamics an integrable system?* Physics Letters A 190 144-148, 1994. (Cited on 1, 2, 3.)

- [55] A. I. Dyachenko, Y. V. Lvov & Zakharov *Five-wave interaction on the surface of deep fluid* Physica D 87, 233-261, 1995 (Cited on .)
- [56] V. E. Zakharov, A. I. Dyachenko & O. A. Vasilyev *New method for numerical simulation of a nonstationary potential flow of incompressible fluid with a free surface* European Journal of Mechanics B/Fluids, Vol. 21, (2002), 283-291 (Cited on .)
- [57] P. Zhang, Z. Zhang *On the free boundary problem of 3-D incompressible Euler equations*. Comm. Pure. Appl. Math. V. 61. no.7 (2008), pp. 877-940 (Cited on 2.)
- [58] F. Zheng *Long-term regularity of 3D gravity water waves* arXiv:1910.01912 (Cited on 2.)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI