FUKAYA CATEGORY FOR LANDAU-GINZBURG ORBIFOLDS

CHEOL-HYUN CHO, DONGWOOK CHOA, AND WONBO JEONG

ABSTRACT. For a weighted homogeneous polynomial and a choice of a diagonal symmetry group, we define a new Fukaya category for a Landau-Ginzburg orbifold (of Fano or Calabi-Yau type). The construction is based on the wrapped Fukaya category of its Milnor fiber together with the monodromy of the singularity, and it is analogous to the variation operator in singularity theory. The new A_{∞} -structure is constructed using popsicle maps with interior insertions of the monodromy orbit. This requires new compactifications of popsicle moduli spaces where conformal structures of some of the spheres and discs are aligned due to the popsicle structures. In particular, codimension one popsicle sphere bubbles might exist and become obstructions to define the A_{∞} -structure. For log Fano and Calabi-Yau cases, we show that the sphere bubbles do not arise from action and degree estimates, together with the computation of indices of twisted Reeb orbits for Milnor fiber quotients.

CONTENTS

1.	Introduction	1
2.	New compactification of popsicles with interior marked points: examples	10
3.	New compactification of popsicles with interior marked points: formal definitions	18
4.	Quantum cap action and J-holomorphic popsicles with Γ -insertions	22
5.	The new A_{∞} -category \mathscr{C}_{Γ}	27
6.	Mirror counterpart: Hypersurface restriction	37
7.	A_∞ -category for a weighted homogeneous polynomial with a symmetry group	41
8.	Twisted Reeb orbits in the Milnor fiber quotients and their indices	47
9.	Monodromy orbit Γ_W and vanishing of spheres with Γ_W	53
Ap	pendix A. Moduli space of pseudo-holomorphic curves and perturbations	56
Ap	pendix B. Popsicle moduli space and its charts	61
Re	References	

1. INTRODUCTION

Singularity theory is a fascinating branch of mathematics with a long history and has deep relations to many branches of mathematics, such as algebraic, complex and symplectic geometry, Lie groups and algebras, commutative algebra and mathematical physics. Classifications as well as the topology and geometry of singularities have been well-established (see [AGZV85], [AGZV12]). In commutative algebras, Cohen-Macaulay rings as coordinate rings of singularities have been investigated in 80's and its indecomposable maximal Cohen-Macaulay modules has been classified for ADE singularities (see [Knö87],[Yos90]). Eisenbud has shown that maximal Cohen-Macaulay modules are equivalent to matrix factorizations of a singularity [Eis80]. In mathematical physics, singularities are often called Landau-Ginzburg (LG for short) models, and together with a finite group *G* preserving the singularity, they are called LG orbifolds.

There has been much attention to the mirror symmetry of singularities, which revealed quite unexpected connections to different branches of mathematics. Symplectic study of Picard-Lefschetz theory by Seidel is mirror to the corresponding algebraic geometry of coherent sheaves [Sei08], and quantum singularity theory developed by Fan, Jarvis and Ruan on Witten equation is mirror to the corresponding integrable hierarchies [FJR13] just to name a few.

Recently, there has been a lot of research on mirror symmetry between LG orbifolds, called Berglund-Hübsch mirror symmetry [BH93]. A polynomial

$$W(x_1,\ldots,x_n) = \sum_{i=1}^n \prod_{j=1}^n x_j^{a_{ij}}$$

is called *invertible* if the matrix of exponents $A = (a_{ij})$ is an $n \times n$ invertible matrix. Its *Berglund*-*Hübsch dual* W^T is an invertible polynomial whose exponent matrix is the transpose A^T . The group of diagonal symmetries

$$G_W = \left\{ (\lambda_1, \dots, \lambda_n) \in (\mathbb{C}^*)^n \mid W(\lambda_1 z_1, \dots, \lambda_n z_n) = W(z_1, \dots, z_n) \right\}.$$

also plays an important role. For a subgroup $G < G_W$, define its *dual group* G^T following [BH95] (see also [Kra09]) as

$$G^T := \operatorname{Hom}(G_W/G, \mathbb{C}^*) \subset G_{W^T}.$$

Berglund-Hübsch mirror symmetry is a duality between two Landau-Ginzburg orbifolds

$$(W,G) \stackrel{\text{mirror}}{\Longleftrightarrow} (W^T, G^T)$$

For closed string mirror symmetry, Fan-Jarvis-Ruan defined quantum singularity theory (FJRW invariants) of (W, G). It should be mirror to Saito-Givental theory of (W^T, G^T) (see [FJR13], [Kra09] for example).

Kontsevich's homological mirror symmetry conjecture [Kon95] for Berglund-Hübsch pairs predicts a derived equivalence between the following two categories;

Fukaya category of
$$(W, G) \stackrel{\text{mirror}}{\longleftrightarrow}$$
 Matrix factorization category of (W^T, G^T)

The left hand side for a nontrivial subgroup G of G_W has **not** been defined. For the case of trivial subgroup $G = \{0\}$, the left hand side is defined as a *Fukaya-Seidel category* of W, $\overrightarrow{FS}(W)$, which is a directed \mathbb{Z} -graded A_{∞} -category of vanishing cycles [Sei08]. The right hand side is the dg-category of G^T -equivariant matrix factorizations $\mathcal{MF}(W^T, G^T)$.

This BH HMS conjecture has been studied extensively for the case of the trivial subgroup $G = \{0\}$ of G_W . In this case $G^T = G_{W^T}$ and the RHS (together with \mathbb{Z} -grading) is known as the category of maximally graded matrix factorization of W^T . For this case of trivial G, HMS conjecture of this form was proposed by Takahashi [Tak], and there have been many interesting works in this direction. See Seidel [Sei01], Kajiura-Saito-Takahashi [KST07], Auroux-Katzarkov-Orlov

[AKO08], Futaki-Ueda [FU11], [FU13], Lekili-Ueda [LU18], [LU20], Harbermann-Smith [HS19] and the references therein. We also refer readers to a nice survey by Ebeling [Ebe].

The candidate for the Fukaya category of the pair (W, G) for a nontrivial subgroup G would be an orbifold version of Fukaya-Seidel category, but the latter is currently out of reach (see [FU09], Problem 3). The main difficulty is that one needs to perturb W to Morse function W_{ϵ} to obtain a legitimate collection of Lagrangian vanishing cycles but this procedure destroys the original symmetry G_W .

The purpose of this paper is to define the Fukaya category of the pair (W, G) when W is a weighted homogeneous polynomial of log Fano and Calabi-Yau type and G is any subgroup of the diagonal symmetry group G_W .

For this purpose, we propose a new approach which does not require Morsification of W. Namely, our approach uses wrapped Fukaya category of Milnor fiber, maximal symmetry group G_W and monodromy of the singularity. Our approach is orthogonal to that of Fukaya-Seidel category in the sense that we will mainly work with non-compact Lagrangians (K in Figure 1) whereas Fukaya-Seidel category uses vanishing cycles (L in Figure 1).




FIGURE 1. Milnor fiber, vanishing cycle L and noncompact Lagrangian K

A topological precursor of our construction is a variation operator

(1.1)
$$\operatorname{var}: H_{n-1}\left(\overline{M}_W, \partial \overline{M}_W\right) \to H_{n-1}\left(\overline{M}_W\right)$$

where $M_W = W^{-1}(1)$ is a Milnor fiber of W. It is defined as a difference of a cycle itself and its monodromy image (fixing the boundary), and it provides an alternative way to describe vanishing cycles.

Recall that a polynomial W is called weighted homogeneous if

$$W(\lambda^{w_1}z_1,\ldots,\lambda^{w_n}z_n)=\lambda^h W(z_1,\ldots,z_n)$$

for $w_1, ..., w_n, h \in \mathbb{N}$ with $gcd(w_1, ..., w_n, h) = 1$. We say *W* has weight $(w_1, ..., w_n; h)$. *W* is assumed to be a weighted homogeneous polynomial for the rest of the paper.

Monodromy homeomorphism (not fixing the boundary) for *W* is known to be an action given by its weights (Milnor [Mil68], see (7.1)).

We define what we call, the **monodromy orbit** Γ_W as follows. We first observe that monodromy homeomorphism is an action of an element $J \in G_W$, hence becomes trivial for the quotient orbifold $[M_W/G_W]$. But on the link of the singularity, monodromy (not fixing the boundary) may be taken as time one Reeb flow of this contact manifold. Therefore on the quotient orbifold $[M_W/G_W]$, they form principal orbits on its contact boundary. We define a distinguished Hamiltonian orbit Γ_W of $[M_W/G_W]$ representing fundamental class of these principal orbits.




FIGURE 2. Two ways of representing vanishing cycle

Let us explain this in the case of $W = x^2 + y^2$. Monodromy ϕ of W which fixes the boundary of the Milnor fiber, is a right-handed Dehn twist as illustrated in Figure 2 (A). Monodromy ψ given by the action of weights is nothing but $(x, y) \rightarrow (-x, -y)$, and at the boundary p of the Milnor fiber, there is a Reeb flow from p to $\psi(p)$ as illustrated in Figure 2 (B). The quotient of the Milnor fiber by the G_W -action is an orbifold sphere $\mathbb{P}^1_{2,2,\infty}$ with a puncture and two $\mathbb{Z}/2$ -orbifold points. The boundary Reeb flow from monodromy becomes a Reeb orbit winding around the puncture once. The monodromy orbit Γ_W is the fundamental class of these principal Reeb orbits.

We remark that monodromy (fixing the boundary) on symplectic Lefschetz fibration has played important roles also in the study of Fukaya-Seidel category as well. For example, Seidel has shown that monodromy provides a natural transformation from identity to monodromy functor [Sei08]. From this point of view, triviality of the monodromy for $[M_W/G_W]$ should provide a natural transformation from identity functor to itself which is an element of Hochschild cohomology of the wrapped Fukaya category, or a symplectic cohomology class [Sei06]. This should be (conjecturally) the geometric orbit Γ_W that we described above.

Now, let us explain our construction of the new Fukaya category using the orbit Γ_W . First, classical variation operator may be viewed as the coequalizer sequence for monodromy ϕ ,

$$H_{n-1}\left(\overline{M}_W,\partial\overline{M}_W\right) \xrightarrow{\phi} H_{n-1}\left(\overline{M}_W,\partial\overline{M}_W\right) \xrightarrow{var} H_{n-1}\left(\overline{M}_W\right) \longrightarrow 0$$

To construct a symplectic analogue, we consider the quantum cap action of Γ_W on the wrapped Fukaya category of $[M_W/G_W]$ given by a version of closed-open map. It gives an A_∞ -bimodule map $\cap \Gamma_W : \mathcal{WF}([M_W/G_W]) \to \mathcal{WF}([M_W/G_W])$. As an analogue of coequalizer, we consider the cone of $\cap \Gamma_W$, which is again an A_∞ -bimodule over $\mathcal{WF}([M_W/G_W])$. We show that it also carries an A_∞ -category structure by constructing higher A_∞ -operations using *J*-holomorphic maps from popsicles with Γ_W -insertions (see Section 7).

Theorem 1.1 (Theorem 7.9). Let W be a non-degenerate weighted homogeneous polynomial $W(z_1, \dots, z_n)$ and G_W be its group of maximal diagonal symmetry group. Suppose that the weight

 $(w_1, \cdots, w_n; h)$ of W satisfies

(1.2)
$$(\sum_{i=1}^{n} w_i) - h \ge 0.$$

Then, there is an A_{∞} -category $\mathscr{F}(W, G_W)$ which fits to a distinguished diagram of bimodules;

$$\mathscr{WF}([M_W/G_W]) \xrightarrow{\cap 1_W} \mathscr{WF}([M_W/G_W]) \longrightarrow \mathscr{F}(W,G_W) \longrightarrow$$

(For a precise definition of Γ_W and its action $\cap \Gamma_W$, see Section 7).

Cohomology groups of the A_{∞} -category $\mathscr{F}(W, G_W)$ can be computed as a cone of $\cap \Gamma_W$ -action, but the rest of the A_{∞} -structure are not constructed algebraically but rather geometrically from *J*-holomorphic maps from popsicles. This requires an important step of popsicle compactifications and we will explain more about it below. Also (1.2) is equivalent to Γ_W having a non-negative Robbin-Salamon index as we show in Corollary 9.6 that Robbin-Salamon index of the monodromy orbit is given by the weights

$$\mu_{RS}(\Gamma_W) = \frac{2((\sum_i w_i) - h)}{h}$$

More explanation will be given in the Proposition 1.3 below.

The constructed A_{∞} -category $\mathscr{F}(W, G_W)$ is the new Fukaya category for the maximal symmetry group G_W . For any subgroup $G < G_W$, we can define the associated Fukaya category using semi-direct product (following Seidel [Sei15]) $\mathscr{F}(W, G) := \mathscr{F}(W, G_W) \rtimes G^T$. This gives the desired Fukaya category for any pair (W, G).

Note that for the trivial subgroup case, the constructed Fukaya category $\mathscr{F}(W, \{0\})$ is $\mathscr{F}(W, G_W) \rtimes G_{W^T}$, and it is natural to conjecture the following.

Conjecture 1.2. The Fukaya category $\mathscr{F}(W, G_W) \rtimes G_{W^T}$ is derived equivalent to the Fukaya-Seidel category of W.

This conjecture is a categorical version of the classical theorem that variation operator (1.1) is an isomorphism and Figure 2 (B) illustrates how $\cap \Gamma_W$ -action recovers vanishing cycles.

It may be surprising that Hom spaces of wrapped Fukaya category is infinite dimensional whereas we expect that $\mathscr{F}(W, G_W)$ have finite dimensional Hom spaces (We will show this in the case of curve singularities in the sequel). This is because most of (but not all) wrapped generators are killed by the quantum cap action of Γ_W .

The conjectural relationship would be quite interesting. Recall that for $\overrightarrow{FS}(W)$, one chooses a Morsification of W and define the vanishing cycles of this Morsification as objects. Other vanishing cycles from a different choice of vanishing paths or a Morsification are included only as twisted complexes of initial vanishing cycles. On the other hand, $\mathscr{F}(W,G)$ includes all (non-compact) Lagrangians as objects, and under the categorical variation operator, some of them would correspond to the vanishing cycles. Furthermore, directness of Fukaya-Seidel category is imposed as a definition, but should be an intrinsic property of $\mathscr{F}(W,G)$. We will discuss further on the conjecture elsewhere in the future.

Now let us explain about the main ingredients for the proof of the main theorem.

CHO, CHOA, AND JEONG

A crucial step in the construction of A_{∞} -category $\mathscr{F}(W, G_W)$ is to define a new compactification of popsicles with interior marked points. Recall that popsicle structure of a disc was introduced and developed by Abouzaid-Seidel [AS10] and Seidel [Sei18] in a different geometric context. Roughly speaking, given a disc D^2 with boundary marked points $\{z_0, \dots, z_n\}$, one considers (hyperbolic) geodesic lines connecting one of z_i and z_0 in D^2 , and interior marked points of a popsicle disc are only allowed to live on these lines. In the construction of Abouzaid and Seidel, interior marked points are allowed to overlap as they were used as a place to put the support of certain one forms.

Unlike [AS10] or [Sei18], we will use interior marked points to place the monodromy Hamiltonian orbit Γ_W to define a new A_∞ -category $\mathscr{F}(W, G_W)$. Therefore, interior marked points should not overlap with each other and when interior marked points collide in a sequence, they will create sphere bubbles in accordance with the conventional Floer theory. Thus, it is necessary to compactify popsicles differently from [AS10] and [Sei18].

In fact, the convergence becomes much more delicate compared to the cases of Abouzaid and Seidel (we thank Paul Seidel for this observation). Let us explain it in more detail.

When the interior marked points collide and create sphere bubbles, the popsicle geodesic lines also collide and induces a popsicle structure on the sphere bubble as well. But depending on the relative positions of marked points, the resulting popsicle structure on the sphere bubble may or may not be trivial. Namely, geodesic lines may remain distinct in the limit bubble or may become a single line in the limit bubble. If popsicle lines remain distinct in the bubbled sphere, the conformal structures of this sphere bubble and a certain disc component (with the same set of non-trivial popsicle lines) should be related. Thus the standard compactification using stable maps is not enough to describe the compactifications.

To overcome these difficulties, we define the concept of *alignment data* which captures such additional relations between conformal structures of discs and spheres. We define a new compactification of popsicle discs using stable popsicles with alignment data. In this new compactification, dimension of a stratum also becomes subtle because alignment might give restrictions to the possible conformal structures. We find that stratum with sphere bubbles can be of codimension one when sphere bubbles are aligned to a popsicle disc. For example, the stratum in Figure 3 is codimension one!




FIGURE 3. A possible codimension one stratum in the popsicle compactification

It is interesting to note that similar phenomena were already observed in the beginning of 90's in the study of modular operads. Fox-Neuwirth [FN62] introduced a cellular decomposition of $S^{2n} = \mathbb{R}^{2n} \cup \{\infty\}$, from the configuration of *n*-points on \mathbb{R}^{2n} . Getzler-Jones in [GJ94] developed the

relationship between operads and homotopy algebras, and studied modular operad to prove a version of Deligne's conjecture. The above phenomenon of sphere bubbles of codimension one also appears in this context and provide obstructions to the cellular structure of the relevant moduli spaces (see Voronov [Vor00] where Getzler-Jone's approach was completed to prove the conjecture).

We can analogously define the compactification of the moduli space of popsicle spheres, and we expect to obtain new homotopy algebraic structures on symplectic cohomology, such as (shifted) L_{∞} - structure on symplectic cohomology. We will explore this elsewhere.

Coming back to our construction of the new Fukaya category, the desired A_{∞} -equation holds if these sphere bubbles of codimension one do not contribute. Let us first work on the case of a symplectic cohomology class Γ in a Liouville manifold M (satisfying Assumption 5.1). If we consider popsicle discs with Γ insertions, the following proposition turns out to be crucial.

Proposition 1.3 (Prop. 5.6, Prop. 9.8). *If Robbin-Salamon index of a principle Reeb orbit component is non-negative, then sphere bubbles of codimension one (with* Γ *-insertions) do not arise.*

We will prove the proposition using estimates of degrees and action values. This proposition enables us to exclude codimension one sphere bubbles and obtain the A_{∞} -relations as in [AS10] and [Sei18]. This defines an A_{∞} -category $\mathscr{C}_{\Gamma}(M)$ (see Section 5).

We run into more challenges with the construction in the case of the Milnor fiber quotient orbifold $[M_W/G_W]$ (to define $\mathscr{F}(W, G_W)$).

The first one is to define the monodromy orbit Γ_W representing principal Reeb orbits at the boundary. This requires a choice of a Morse-Smale function on the contact boundary. As G_W acts diagonally, orbifold strata are somewhat special in the sense that they are always given by setting some of the coordinates to be zero. This can be used to define a desired Morse-Smale function (Proposition 9.1) whereas it is not always possible for a general orbifold.

The next difficulty is that a general definition of orbifold symplectic cohomology is not known. Our case is somewhat simpler in the sense that we only use Γ_W as insertions, but the output of a possible sphere bubble could be an arbitrary orbifold symplectic cochain. Even though we will not define the orbifold symplectic cochain complex of $[M_W/G_W]$, we will identify their generators and compute their Robbin-Salamon indices.

Namely, we will consider the space of G_W -twisted Reeb orbits, i.e., $\gamma : [0, l] \rightarrow M_W$ satisfying $g \cdot \gamma(1) = \gamma(0)$ for some $g \in G_W$, and γ being a Reeb flow. The problem can be quite complicated as the period *l* could be a fractional number. Their Robbin-Salamon indices are explicitly computed in Proposition 8.15. We obtain an inequality relating the weights of *W* and the Robbin-Salamon indices of twisted Reeb orbits in Proposition 8.19. Hence the log Fano or Calai-Yau assumption (1.2) combined with this inequality implies that the associated Robbin-Salamon index is non-negative. This enables us to prove vanishing of sphere bubbles with Γ_W insertions (see Proposition 9.8). This finishes the construction of the new A_{∞} -category $\mathscr{F}(W, G)$.

Our construction heavily depends on the fact that W is weighted homogeneous, especially in the definition of the monodromy orbit Γ_W . But as variational operator exists for general W, it is natural to expect that its categorification exists in general as well. In the case that W is a polynomial of two variables, which are not necessarily weighted homogeneous, we can indeed define such a category (in a joint work in progress with Hanwool Bae).

CHO, CHOA, AND JEONG

In the sequel, we will apply the construction in this paper to find a geometric understanding of Berglund-Hübsch Mirror symmetry. Given an invertible polynomial W of two variables, we can find the transpose polynomial W^T entirely in a geometric way. The Milnor fiber quotient $[M_W/G_W]$ has a Landau-Ginzburg mirror $W_{\mathbb{L}}$ given as a Lagrangian Floer potential function of a Seidel Lagrangian \mathbb{L} . Then, the closed open map of the monodromy orbit Γ_W can be used to define a polynomial g via Kodaira-Spencer map ([FOOO16], [ACHL20]). We can show that quantum cap action $\cap \Gamma_W$ corresponds to the restriction of the potential $W_{\mathbb{L}}$ to the hypersurface g = 0 (see also Section 6). Then $W_{\mathbb{L}}$ restricted to the hypersurface g = 0 turns out to be the transpose polynomial W^T !

To exclude the sphere bubbles in our main construction, we need to require log Fano or Calabi-Yau condition. Unfortunately, only $x^2 + y^2$ satisfies this condition and all the other invertible curve singularities are of log general type. But we can still use low dimensionality to exclude certain sphere bubbles, and more detailed discussion will be given in the sequel.

Let us finish the introduction by working out the example of $x^2 + y^2$.

1.1. **Example:** $x^2 + y^2$. To illustrate our construction, let us explain the case of $W = x^2 + y^2$. Milnor fiber $M_W = \{x^2 + y^2 = 1\}$ is T^*S^1 or a cylinder. Its zero section is a vanishing cycle of W and its cotangent fiber K generates its wrapped Fukaya category (see Figure 1). The maximal diagonal symmetry group G_W is $\mathbb{Z}/2 \times \mathbb{Z}/2$ and the quotient $[M_W/G_W]$ is an orbifold sphere $\mathbb{P}^1_{2,2,\infty}$ with two $\mathbb{Z}/2$ -orbifold points, say A, B and a single puncture C as in Figure 4.




FIGURE 4. Quotient of Milnor fiber

We claim that

$$H^{\bullet}(\operatorname{Hom}_{\mathscr{F}(W,G_W)}(K,K)) \simeq \operatorname{Cliff}_2(\mathbb{C}).$$

A category $\mathscr{F}(W, G_W)$ is essentially the wrapped Fukaya category of $[M_W/G_W] = \mathbb{P}^1_{2,2,\infty}$ with additional relation " $\Gamma_W = 0$ ". We realize it by considering quantum cap action of Γ_W on wrapped Fukaya category of $\mathbb{P}^1_{2,2,\infty}$. As a complex, we have

$$\left(\operatorname{Hom}_{\mathscr{F}(W,G_W)}(K,K),d\right) \simeq \left((CW^{\bullet}(K,K),m_1) \xrightarrow{\cap \Gamma_W} (CW^{\bullet}(K,K),m_1) \right)$$

We view *K* as an object of the A_{∞} -category $\mathcal{WF}([M_W/G_W])$ by considering G_W -orbit of *K* consisting of $K, \psi(K)$ and their disjoint $\mathbb{Z}/2$ -images. Let us take two minimal wrapped generators, name them as $\alpha, \beta \in CW(K, K)$. The computation shows

$$(CW^{\bullet}(K,K),m_1) \simeq (\mathbb{C}\langle \alpha,\beta\rangle/\{\alpha^2=\beta^2=1\},m_1=0)$$

The element Γ_W is an S^1 family of orbits (quotient of a Reeb flow drawn in Figure 2 (B)). The same flow determines an element $\alpha\beta$ and $\beta\alpha$ in $CW^{\bullet}(K, K)$. The map $\cap \Gamma_W$ turns out to be the

multiplication by $\alpha\beta + \beta\alpha$. Therefore,

$$H^{\bullet}(\operatorname{Hom}_{\mathscr{F}(W,G_W)}(K,K)) \simeq \mathbb{C}\langle \alpha,\beta\rangle/\{\alpha^2 = \beta^2 = 1, \alpha\beta = -\beta\alpha\},$$

as is claimed.

Intuitively cone of a morphism corresponds to the Lagrangian surgery as illustrated in Figure 2 (B). But we should note that the new category $\mathscr{C}_{\Gamma_W}([M_W/G_W]) = \mathscr{F}(W, G_W)$ is different from the category of cones (where cones are considered as twisted complexes), and hence this surgery interpretation should be taken only as an intuition. The actual A_∞ -category is defined by using popsicle maps with Γ_W -insertions.

We would like to notice that this computation matches to the expectation from Berglund-Hübsch version of homological mirror symmetry. A mirror for $\mathbb{P}^1_{2,2,\infty}$ is a polynomial $x^2 + y^2 + xyz$. To see this, take Seidel's Lagrangian \mathbb{L} ([Sei11]), which is an immersed Lagrangian with three odd immersed generators X, Y, Z. It is weakly unobstructed with bounding cochain $\boldsymbol{b} = xX + yY + zZ$ and its Lagrangian potential function is exactly $W_{\mathbb{L}} = x^2 + y^2 + xyz$ (see [CHL17]). This can be seen by taking lifts of \mathbb{L} to the cylinder, which gives four circles as in Figure 5. Pick a generic point, and count all rigid polygons passing through it. The reader can find two bigons with corners labeled by X, X and Y, Y together with a minimal triangle XYZ.




FIGURE 5. Lifts of Seidel Lagrangian and K

If we set z = 0, $W_{\mathbb{L}}$ becomes a desired dual polynomial $W^T = x^2 + y^2$. This is related to the monodromy of W as follows. Monodromy of W (fixing the boundary) is given by a Dehn twist ϕ as in Figure 2 (A). We will consider a version of monodromy ψ which does not fix the boundary as in Figure 2 (B). It turns out that the monodromy ψ on the Milnor fiber M_W is is the same as the action by the element (-1, -1) of a maximal symmetry group G_W . Also, consider the Reeb flows on the boundary $\partial \overline{M}_W$ describing the monodromy ψ , which give a Reeb orbit Γ_W in the quotient $\mathbb{P}^1_{2,2,\infty}$. In this case, Γ_W is the orbit that winds the puncture of the quotient once. Closed-open map takes this orbit to $z \cdot 1_{\mathbb{L}}$.

We have A_{∞} -functor from $\mathscr{F}(W, G_W) \to \mathscr{MF}(W^T)$ which is induced by localized mirror functor [CHL17] (and setting z = 0). It sends K to Floer complex $\left(CW^{\bullet}(K, \mathbb{L}), -m_1^{0, \boldsymbol{b}}\right)\Big|_{z=0}$, which is a

matrix factorization of W^T . We can calculate it in Figure 5 and it is given by

$$-\begin{pmatrix} y & x \\ x & -y \end{pmatrix} \cdot - \begin{pmatrix} y & x \\ x & -y \end{pmatrix}$$

This factorization is the compact generator of $\mathcal{MF}(W^T)$ considered in [Dyc11], and it's endomorphism ring is the Clifford algebra in two variables.

1.2. Structure of the paper. In Section 2, we explain the new phenomenon that arises in popsicle compactifications with interior insertions, and present the concept of alignment data using a number of instructive examples. In Section 3, we give a formal definition of the compactified moduli space of popsicles with interior insertions. In Section 4, we recall the definition of quantum cap action and define *J*-holomorphic popsicles with insertions of a fixed symplectic cohomology cycle Γ in a Liouville manifold *M*. In Section 5, we construct an A_{∞} -category \mathscr{C}_{Γ} from wrapped Fukaya category of M on which the quantum cap action of Γ vanishes using *J*-holomorphic popsicles with Γ -insertions. This construction will be generalized to orbifold quotients of Milnor fibers of weighted homogeneous polynomials in Section 7. A parallel construction in algebraic geometry is described in Section 6, which is a restriction to a hypersurface. In Section 7, we explain our choice of distinguished Hamiltonian orbit Γ_W for a Milnor fiber of a given weighted homogeneous polynomial W which encodes the monodromy information. Then we define a new A_{∞} -category $\mathscr{F}(W,G)$ by generalizing the construction of \mathscr{C}_{Γ} to this particular class of orbifolds. In order to show its well-definedness, we classify Reeb orbits of a Milnor fiber and computes their Conley-Zehnder indices in Section 8. Finally, we show that $\mathscr{F}(W,G)$ is well-defined for a log Fano/Calabi-Yau type polynomial in Section 9.

In Appendix A, we briefly describe the moduli spaces and perturbation scheme we use throughout the paper. Appendix B explains a compactification of popsicle moduli spaces.

1.3. **Acknowledgement.** We would like to thank Paul Seidel for informing us a crucial error in the previous version of our compactification of popsicles and the encouragements. We would like to thank Otto van Koert, Hanwool Bae for the discussion on symplectic cohomology theory on orbifolds, and popsicle compactifications. We would like to thank Atsushi Takahashi, Philsang Yoo, Kaoru Ono for helpful discussions. The second author was partially supported by the T.J.Park science fellowship grant.

2. New compactification of popsicles with interior marked points: examples

Let us first informally explain the new compactification of popsicles through several examples of increasing complexity, and give a formal treatment later in the section.

Let us first recall the definition of a popsicle by Abouzaid and Seidel [AS10] (we follow Seidel [Sei18] to use rational ends).

Definition 2.1 ([AS10],[Sei18]). A popsicle is a disc D^2 with following decorations;

- (1) **boundary marked points**: denoted by $z_0, z_1, ..., z_n$ according to their cyclic order.
- (2) **popsicle lines**: regarding the interior of D^2 as a hyperbolic disc, the geodesic connecting z_i and z_0 (at infinity), denoted as Q_i .

(3) *flavor*: a set of flavors $F = \{1, ..., l\}$ and a non-decreasing map

$$\phi: F \to \{1, \ldots, n\}.$$

(4) *sprinkles (interior marked points): l* distinct interior marked points $z_1^+, ..., z_l^+$, such that z_j^+ lies on the geodesic $Q_{\phi(j)}$ for j = 1, ..., l.

We say that the above popsicle has type (n, ϕ) (see Figure 6). We call it stable if $n + l \ge 2$. We denote a moduli space of popsicles of type (n, ϕ) by $P_{n, F, \phi}$.




FIGURE 6. Example of a popsicle such that $\phi(1) = \phi(2) = 1$ and $\phi(l) = k$.

Abouzaid and Seidel defined compactifications of popsicle moduli spaces while allowing the interior marked points to coincide. For example, z_1^+ and z_2^+ in Figure 6 may coincide in their limit. But we will not allow our interior marked points to coincide, as they will be used to place inputs.

For example, let us consider a popsicle of type $(2, \phi)$, which has two sprinkles with the identity map $\phi : \{1, 2\} \rightarrow \{1, 2\}$. Our compactification of the moduli space of type $(2, \phi)$ is illustrated in the Figure 7. To obtain the Abouzaid-Seidel's compactification, one has to contract the bottom edge to a point (see Figure 3 of [AS10]), as sphere bubble does not arise in the limit.

Note that at the two vertices of the bottom edge in the Figure 7 two popsicle lines coincide in the limiting sphere bubble. On the interior of the bottom edge, sphere bubbles inherit two popsicle lines from discs, and popsicle lines of the sphere and disc are aligned.

2.1. Aligning conformal structures on a disc and a sphere. We introduce the notion of a popsicle sphere, and explain the notion of alignment between a popsicle disc and a popsicle sphere.

Definition 2.2. A popsicle sphere is the following decoration on a rational sphere \mathbb{P}^1 ;

- (1) fixed outgoing marked point point: denoted by $w_0^- \in \mathbb{P}^1$,
- (2) incoming marked points: denoted by $\{w_i^+ | 1 \le i \le n\}$,
- (3) **popsicle lines**: regarding $\mathbb{P}^1 \setminus w_0^-$ as a complex plane, a set of *m* distinct oriented lines on a plane with the same direction. We can arrange them using rotation of a complex plane so that they are all oriented upward. We denote them by $Q_1, \ldots, Q_m \subset \mathbb{C}$ from the left to the right after this arrangement.
- (4) $a \operatorname{map} \phi : \{w_1^+, \dots, w_n^+\} \to \{1, \dots, m\}$ such that w_i^+ lies on $Q_{\phi(w_i^+)}$.

If $n \ge 2$, then we call it stable. We write such a moduli space of popsicle structure by $\mathbb{P}_{m,n,\phi}$.




FIGURE 7. Compactification of popsicles when sprinkles cannot overlap with each other




FIGURE 8. Example of a popsicle sphere and its planar model

Popsicle structure on a sphere and a disc can be compared in a natural way. First, take out z_0 (resp. w_0^-) from D^2 (resp. \mathbb{P}^1) and fix an identification with the upper half plane H (resp. \mathbb{C} via stereographic projection).

In the case of a disc, boundary marked points z_1, \dots, z_n are determined by ordered real numbers x_1, \dots, x_n , and a popsicle line between z_0 and z_i on D^2 correspond to the vertical (upward) lay at x_i . Thus the position of sprinkle z_i^+ is determined by its *y*-coordinate.

$$z_i \mapsto x_i \in \mathbb{R}, \quad Q_i \mapsto \{\operatorname{Re} z = x_i\}, \quad z_j^+ \mapsto \left(x_{\phi(j)} + \sqrt{-1}y_j\right) \in Q_{\phi(j)}.$$

The forgetful map $f : P_{n, E, \phi} \to \mathcal{M}_{0, n+1}$ is well-defined.

In the case of a sphere, we may further assume that w_1^+ maps to $\mathbb{R} \in \mathbb{C}$ and each and every Q_i head strictly upward. Then popsicle lines are determined by ordered real numbers x_i and

sprinkles are determined by real numbers y_i ;

$$Q_i \mapsto \{\operatorname{Re} z = x_i\}, \quad w_1^+ \mapsto x_{\phi(w_1^+)}, \quad w_j^+ \mapsto \left(x_{\phi(w_j^+)} + \sqrt{-1}y_j\right) \in Q_{\phi(j)}.$$

We get a well-defined map $f : \mathbb{P}_{m,n,\phi} \to \mathcal{M}_{0,m+1}$, just remembering *x*-coordinates of popsicle lines.

Definition 2.3. We say a popsicle disc Σ and a popsicle sphere Σ^+ are **aligned** if their forgetful map images coincide in $\mathcal{M}_{0,m+1}$. Namely, we have

$$(\Sigma, \Sigma^+) \in P_{m, F, \phi} \times_{\mathcal{M}_{0, m+1}} \mathbb{P}_{m, n, \phi'}.$$

When Σ_1 *and* Σ^+ *are aligned, we will write*

$$\Sigma \bowtie \Sigma^+$$
 or $\Sigma^+ \bowtie \Sigma$

Intuitively, if popsicle lines when seen in the upper half plane H and \mathbb{C} agree with each other, then they are aligned. Note also that a disc and a sphere with exactly two popsicle lines each, are always aligned with each other.

A popsicle which consists of discs and sphere bubbles is called stable if it is stable in the usual sense. Namely, each sphere component has at least 3 special points (marked or nodal), and each disc component has either at least three special points (marked or nodal) or one boundary and one interior nodal point. Hence, stable popsicle do not contain a sphere component with several popsicle lines but with just one or zero sprinkle. But in order to describe a neighborhood of a stable popsicle, sometimes we need to consider such unstable components for the gluing. Such unstable components will be uniquely determined from a given stable popsicle and thus they will be called auxiliary data.

Remark 2.4. Let us remark on the alignment for the unstable case of only one popsicle line. The space of pairs (Σ, Σ^+) have extra one dimension because $\mathcal{M}_{0,1+1} \simeq [\bullet/\mathbb{R}]$ is a stacky point. Explicitly, suppose two different sphere component Σ_1^+ and Σ_2^+ are isomorphic by a homothety automorphism. Even though Σ_i^+ represent isomorphic element in $\mathbb{P}_{1,F',\phi'}$, a pair (Σ, Σ_1^+) and (Σ, Σ_2^+) represents different element in the fiber product. This ambiguity provides extra one more dimension.

Let us discuss examples of compactifications and the alignment therein.

2.2. Examples of new compactifications. Let us start with the simplest example, which appears at the bottom stratum of Figure 7.

Example 2.5. Consider (D^2, z_0, z_1, z_2) with two sprinkles z_1^+, z_2^+ , whose type is $(2, \phi)$ where ϕ : $\{1,2\} \rightarrow \{1,2\}$ is an identity map. The moduli space of popsicles of type $(2, \phi)$ has dimension two.

The new codimension one stratum is illustrated in Figure 9 (A). Denote by Σ^+ , Σ_0 , Σ_1 , the unique sphere component, the middle disc Σ^+ and the popsicle disc as in the figure, and by Σ the whole stable popsicle. Here, we will require that $\Sigma^+ \bowtie \Sigma_1$ (this condition is automatically satisified in this case of n = 2, but will be non-trivial for the general case n > 2).

This limit can appear if z_1^+, z_2^+ approach to each other (producing a sphere bubble) and at the same time they approach z_0 (producing a disc bubble). This stratum at first might appear to be of codimension two or higher, but it is of codimension one, hence is one dimensional.




FIGURE 9. Popsicle with sphere bubble and gluing using planar model 1

Note that Σ_0, Σ_1 has the unique conformal structure up to automorphism. Since $\Sigma_0 \bowtie \Sigma^+$, there is one dimensional automorphisms of Σ^+ which preserve p and two popsicle lines and we use it to fix the position of z_1^+ . Thus there only remains the freedom to choose z_2^+ on the popsicle line, which gives one dimensional family as suggested.

It is instructive to describe the explicit neighborhood of Σ using the gluing parameter $r (0 < r \ll 1)$ and the parameter y for the position of z_2^+ . Here y is the vertical coordinate of z_2^+ in $\mathbb{C} \cong \Sigma^+ \setminus \{p\}$ when z_1^+ is at the origin of \mathbb{C} . Given (r, y), we can describe the planar model (see Figure 9 (B)) of the glued disc: upper half plane is Σ_0 and Σ^+ is glued at (0, 1) and Σ_1 is glued at (0, 0). Note that we use the same gluing parameter r for the aligned disc and sphere so that popsicle lines are aligned and extend to all of H. We can place z_2^+ on the second vertical ray whose height is $r^2y + 1$. This describes regular popsicles (with r > 0) near Σ .

We can generalize this example to the case of *n* sprinkles. Namely, consider Σ_1 with (n + 1) marked points and Σ^+ with *n* popsicle lines. This configuration will be again of codimension one, with the similar planar model with several popsicle lines.

Example 2.6. Consider the new codimension two stratum is illustrated in Figure 10 (which appear as a vertex at the bottom of Figure 7). Denote the components of Figure 10(A) by Σ^+ , Σ_0 , Σ_1 as before. In this case, two popsicle lines on Σ^+ coincide and two sprinkles are placed on it. Note that an automorphism preserving the popsicle lines and fixing two sprinkles z_1^+ , z_2^+ should be trivial and this stratum is 0 dimensional.

Let us explain how this stratum may arise as a limit along two different codimension one strata.

One of the limit is the usual one: consider stable popsicle with two disk components Σ_0, Σ_1 with Σ_1 as before, but Σ_0 has the horizontal popsicle line with z_1^+, z_2^+ on it. If z_1^+, z_2^+ collide with each other, we obtain the sphere bubble with an induced popsicle line, and the corresponding limit is 10(A).

The other one is as a limit of Example 2.5 when z_2^+ approaches p_0 . It might appear that this will create another sphere bubble, but this will not happen as the resulting configuration would not be




FIGURE 10. Popsicle with sphere bubble and gluing using planar model 2

stable. It is better seen from the planar gluing model. In Figure 9 (B), suppose that y-coordinate of z_2^+ is 1 + cr for a small constant c, whereas the popsicle line is at $x = r^2$. Note that the order of popsicle lines approaching to each other and that of z_1^+ and z_2^+ are r^2 and r respectively, and the limit $r \to \infty$ is the desired stratum. Note also that in the limit the alignment between Σ_1 and Σ^+ no longer exists.

Another way to see the second limit is as follows. Identify Σ^+ with S^2 , with south pole p and north pole z_1^+ . Fix a popsicle line through p and z_1^+ . Use the remaining automorphism of Σ^+ to place z_2^+ at the equator, which fixes the popsicle line through z_2^+ . Then z_2^+ has a freedom to move along the half of the equator and as z_2^+ approaches the first popsicle line, we obtain two limits corresponding to two vertices of the bottom edge of Figure 7.

Let us explain the alignment data which we assign for this stratum. As explained above, Σ_1 and Σ^+ has different order of Gromov-convergence, and we record it using the following alignment data. We first introduce the unstable popsicle disc Σ_{un} (that is D^2 with z_0, z_1 and popsicle line(s) between them) and associate it to the nodal point p_1 as an auxiliary data. Then, we align this unstable disc with Σ^+ . As an alignment data, it will be denoted as

 $p_1 \bowtie \Sigma^+$.

By aligning each sphere bubble with a disc or a nodal point, we will keep track of the order of convergence, relate conformal structures and also use it to find the glued model.

It is helpful to describe the glued model Figure 10(B) in the neighborhood of this codimension two stratum. Since it is of codimension two, there are two gluing parameters, say $r_1, r_2 \ll 1$. Here r_1 is the gluing parameter between Σ^+ and Σ_0 . Given another parameter r_2 , a standard gluing of Σ_0 and Σ_1 with parameter r_2 is not the correct one. Instead, we make use of the unstable disc Σ_{un} at p_1 that is aligned with Σ^+ , and glue Σ_0, Σ_{un} and Σ_1 . This unstable disc (with two striplike ends removed) corresponds to the half of annulus (centered at $0 \in H$ between two circles of radius r_1r_2 and r_1) after gluing in the Figure 10(B). Alternatively one simply forget Σ_{un} and

CHO, CHOA, AND JEONG

glue Σ_0 and Σ_1 with gluing parameter r_1r_2 to get the same glued model. This is the role of the alignment $p_1 \bowtie \Sigma^+$.

Another feature of Figure 10(B) is that the neighborhood of two marked points z_1^+ , z_2^+ in Σ^+ are replaced by two discs of radius r_1r_2 which is the same as that of Σ_1 . In this way, popsicle lines extends consistently to the entire glued model. We may say that $z_1^+ \bowtie \Sigma_1$ and $z_2^+ \bowtie \Sigma_1$. But since it is clear how to align each sprinkle, we will not put these alignments explicitly in the alignment data.

In the above two examples, we have illustrated that how the alignment data arises from compactifications and how we can use it to describe neighborhoods of stable popsicles.

Now, let us consider another example, which illustrates that stable popsicle alone is not enough to describe the limit in the compactification of popsicles. In the previous two examples, the alignment data is uniquely determined from each stable popsicle. But a general stable popsicle might allow several compatible alignment data. For example the order of convergence may not be uniquely determined from the given stable popsicle. Therefore, alignment data should be specified to describe the limit correctly.

Example 2.7. The following stable popsicle has three string of disks $\Sigma_0, \Sigma_1, \Sigma_2$ and a sphere Σ^+ attached to Σ_0 at p_0 . Let us denote by p_i the intersection of Σ_{i-1} and Σ_i for i = 1, 2. Σ_2 has marked points z_1, z_2, z_3 with three popsicle lines to z_0, Σ_1 has z_3^+ on the common popsicle line and on Σ^+, z_1^+, z_2^+ lie on the common popsicle line (for all colors). Here we put a (distinct) color to each popsicle line for convenience.




FIGURE 11

Let us discuss where to align Σ^+ . There are three choices, Σ_1 , p_1 , p_2 , and each choice corresponds to a different limit. The planar models for each case are illustrated in Figure 12. The codimension of each stratum are 2,3,3 respectively. Since popsicle lines become distinct in Σ_2 , so Σ^+ cannot be aligned to Σ_2 .

(1) The case of $\Sigma_1 \bowtie \Sigma^+$: This is when the order of convergence for Σ_1 and Σ^+ are the same. Note that conformal structure of Σ_1 and Σ^+ are unique up to biholomorphic map, but as they are aligned, we need to use the same biholomorphic map for Σ_1 and Σ^+ . As a result, we get one dimensional freedom. More precisely, let us fix the position of p_1, p_2 for Σ_1 and fix p_0, z_1^+ and the popsicle line for Σ^+ . Each of Σ_1 and Σ^+ has the translation symmetry, which we use it to fix z_3^+ in Σ_1 . Therefore, the position of z_2^+ on the unique popsicle line in Σ^+ is free. As Σ_2 has one dimensional moduli, the stratum is two dimensional, hence is of codimension two.




FIGURE 12. Glued models for a stable popsicle with 3 different alignment data

(2) The case of p₁ ⋈ Σ⁺: Conformal structure of Σ⁺, Σ₀, Σ₁ is unique up to automorphism and hence the stratum is one dimensional (due to Σ₂) and of codimension 3.
 In the glued model Figure 12(P) Σ is identified with H and Σ⁺ is glued at (0, 1) with

In the glued model Figure 12(B), Σ_0 is identified with H, and Σ^+ is glued at (0,1) with radius r_1 . From the alignment $p_1 \bowtie \Sigma^+$, we consider a unstable disc Σ_{un} , which is glued at $0 \in H$ with the same gluing parameter r_1 , which gives the half annulus between radius r_1r_2 and r_1 in H. Then Σ_1 is glued to Σ_{un} to give another half annulus between radius $r_1r_2r_3$ and r_1r_2 in H. Lastly Σ_2 is glued to Σ_1 to give the half disc of radius $r_1r_2r_3$, and popsicle lines are extended from Σ_2 to all of H.

 (3) The case of p₂ ⋈ Σ⁺: This is similar to the previous case but we glue the unstable disc (aligned with Σ⁺) at p₂.

Note that to properly describe the glued model of a stable popsicle, we need to keep track of unstable components called auxiliary data. One key feature is that such an auxiliary data can be uniquely determined from the stable components and the alignment data. For example, an aligned sphere with several distinct popsicle lines through a nodal point but with only one sprinkle is regarded as an unstable sphere, although it has the unique conformal structure. This is in fact the auxiliary sphere that we glue in at each sprinkle in the above example. One might choose to include such auxiliary data as a part of the definition of a stable popsicle, but for simplicity we will omit them from the definition.

Let us summarize informally the rules of alignment before we give a formal treatment in the next section. Given a general stable popsicle with discs and sphere bubbles, each sphere bubble is equipped with an induced popsicle lines on it. We need to specify an alignment for each sphere component. A sphere component may be aligned to a disc component if the configuration of popsicle lines of the disc and sphere is the same. If there are more than 2 distinct popsicle lines on the sphere, then it should be aligned to a disc (not to a vertex) and there is a unique disc, whose double has the desired configuration of popsicle lines of the sphere. So we align the sphere with such a disc component and this will align their conformal structures as well.

If a sphere component has a unique circle which is the common popsicle line, it is more delicate to describe the rule, but roughly, it should be aligned to either a nodal point between disc components where exactly the same set of popsicle lines pass through, or to a disc component with the same popsicle lines. In this case, the nodal point or the disc component that is aligned should be further away from the root than the the disc component that the sphere sits on.

When several sphere components are aligned to the same disc nodal point, then we need to also describe their respective orders of convergence, and this will be a part of the alignment data as well.

3. New compactification of popsicles with interior marked points: formal Definitions

In this section, we define a new compactification of popsicles with interior marked points. We will first give a combinatorial description and then define a stable popsicles of a given combinatorial type.

Let us recall the notion of a rooted tree. A **rooted tree** with *n* leaves has n + 1 semi-infinite edges with a single preferred one called the root, and the rest are called leaves. All rooted trees are oriented from the root toward the leaves.

Definition 3.1. This orientation induces the following **partial order** on vertices and edges: For any $a, b \in Vertex(T) \sqcup Edge(T)$, we declare a < b when a is contained in the unique path from the root to b.

For each edge e of T, we denote by T_e a subtree consisting of e itself and all the vertices and edges that are greater than e.

Definition 3.2. A rooted ribbon tree is a rooted tree T with cyclic orders on adjacent edges for each vertex v of T. Each vertex v of a rooted ribbon tree has a single distinguished edge coming from the roots, denoted as $e_{v,0}$. The rests are arranged according to the cyclic structure as $\{e_{v,1}, \ldots, e_{v,val(v)-1}\}$. The leaves are arranged in a similar way, denoted as $\{l_1, \ldots, l_n\}$.

Using a rooted ribbon tree, we can describe limits of popsicles without sphere bubbles following [AS10].

Definition 3.3. (Broken popsicles without sphere bubbles) Let *T* be a rooted ribbon tree with *n* leaves and let ϕ : $F \rightarrow \{1, ..., n\}$ be a non-decreasing flavor function. A ϕ -flavored stable broken popsicle (without sphere bubbles) modelled on *T* consists of the following data;

- (1) *decomposition of* F: a decomposition $F = \bigsqcup_{v} F_{v}$ where v runs through vertex(T).
- (2) *induced decomposition of* ϕ : maps $\phi_v : F_v \to \{1, \dots, val(v) 1\}$ for each vertices v so that the edge $e_{\phi_v(f)}$ lies on the unique path from the root to $l_{\phi(f)}$ of T.
- (3) **popsicles**: for each vertex v, a popsicle of type $(val(v), F_v, \phi_v)$

We call a broken popsicle (without sphere bubble) is stable if popsicles assigned to each vertices are stable.

Let us define a combinatorial notion called a layer. Recall that *n* interior marked points of a popsicle sphere in $\mathbb{P}_{m,n,\phi}$ are lying on *m* popsicles lines, which partitions *n* interior marked

points into *m* disjoint subsets. Also, the natural ordering of popsicle lines (as vertical lines in \mathbb{C}) induces the ordering on these partitioned subsets.

Definition 3.4. A rooted layered tree is a rooted tree T' together with ordered partitions of outgoing edges of from every vertices. Namely, for each vertex w with outgoing edges $\{e_{w,1}, \dots, e_{w,val(w)-1}\}$, we take its ordered partition

$$E_w := \{E_{w,1}, \dots E_{w,m}\}$$

with the following ordering : we declare $E_{w,i} < E_{w,j}$ for i < j. We call $E_{w,i}$ an i-th layers at w. It is called stable if it is stable as a rooted tree.

We remark that a rooted tree modeling sphere bubbles does not carry a ribbon structure.

Recall from Definition 2.1 that $F = \{1, \dots, l\}$ labels interior marked points, and flavor function ϕ describes a popsicles line for each interior marking to sit on. Thus, the induced marked points in the sphere bubble (which corresponds to leaves of a rooted tree) should carry this data as well.

Definition 3.5. An *F*-label of a rooted tree *T* is an injective map ψ : Leaf(*T*) \rightarrow *F*.

As we explained in the previous section, a sphere bubble of popsicles are not random and is aligned to a disc component or a nodal point. To describe the alignment data, we introduce the notion of a coloring.

When *C* is a set of some colors, a map $f : \text{Leaf}(T) \to C$ associates a color to each leaf of *T*.

Definition 3.6. Let *T* be a rooted tree, equipped with a map f: Leaf(*T*) \rightarrow *C*. For an edge *e* of *T*, denote by Leaf_e(*T*) \subset Leaf(*T*) the leaves of *T* that are contained in the subtree *T*_e (see Definition 3.1).

The image of an induced map $f : \text{Leaf}_{e}(T) \to C$ describes a **color set** of e with respect to f

$$C^{f}(e) := \left\{ f(l) \mid l \in \operatorname{Leaf}_{e}(T) \right\} \subset C.$$

We call this assignment a C-coloring of a rooted tree with respect to f, denoted as C^{f} .

The intuition behind the definition is simple. We "color" all the edges along the unique path from the root to each leaf l by f(l). Each edge $e \in \text{Edge}(T)$ may carry multiple colors, and the set of those colors are denoted as $C^{f}(e)$. There are two different colorings we would like to consider (corresponding to a disc and a sphere):

- Let *T* be a rooted ribbon tree with *n* leaves. A cyclic order on leaves determine the unique order preserving map *f*: Leaf(*T*) → *C* = {1,..., *n*} given by *f*(*l_i*) = *i*. We call it a **canonical coloring** of a rooted ribbon tree. We denote the color set of edge *e* with respect to a canonical coloring simply as *C*(*e*), omitting *f*.
- Let T' be a rooted tree with an F-label ψ: Leaf(T') → F = {1,..., l}. Further suppose that we are given a nondecreasing flavor function F → C = {1,..., n} as in 3.3. The composition φ∘ψ: Leaf(T') → {1,..., n} determines a C-coloring of T'. We call this coloring an **induced coloring** on an F-labeled rooted tree, denoted as C^(φ,ψ)

In particular, if in addition T' is layered (Definition 3.4), then we can define the color set of a single layer $E_{w,k}$ from the coloring $C^{(\phi,\psi)}$:

$$C^{(\phi,\psi)}(E_{w,k}) := \bigcup_{e \in E_{w,k}} C^{(\phi,\psi)}(e) \subset C.$$

The notion of coloring will help us to describe the alignment data, which we will explain from now.

Given one popsicle sphere or trees of popsicle spheres, we should be able to tell whether they may arise in the limit of popsicle discs. If they arise from such limits, there are alignments of conformal structures. We will first describe when it is possible to align a popsicle sphere component to a popsicle disc component or a nodal point. Next, we will define an appropriate compatibility between the choices of alignment data for every sphere components.

Let us define the notion of alignability.

Definition 3.7. Let *T* be a rooted ribbon tree with *n*-leaves, and let ϕ : $F = \{1, ..., l\} \rightarrow C = \{1, ..., n\}$ be a (non-decreasing) flavor function. Let *T'* be a rooted layered tree with an *F*-label ψ : Leaf(*T'*) \rightarrow *F*. Then *T* carries a canonical coloring *C* and *T'* carries an induced coloring $C^{(\phi,\psi)}$ as above.

(1) We say $w \in Vertex(T')$ and $v \in Vertex(T)$ are alignable if for each $k \ge 1$

$$C^{(\phi,\psi)}\left(E_{w,k}\right) \subset C\left(e_{v,i_k}\right)$$

for some $i_k \ge 1$, and $i_{k+1} > i_k$ for $\forall k \ge 1$. In fact, such i_k is unique if it exists. (2) We say $w \in \text{Vertex}(T')$ and $e \in \text{Edge}(T)$ are alignable if w has only one layer E_w and

$$C^{(\phi,\psi)}(E_w) \subset C(e).$$

The first case is when a sphere component is aligned to a disc component. Each layer $E_{w,k}$ in the sphere bubble is from a vertical line given by the (possibly overlapping) popsicle line(s) of color(s) $C^{(\phi,\psi)}(E_{w,k})$. The first alignability is to guarantee that the alignable disc should at least carry such popsicle lines as well. More precisely, the disc have a nodal (or marked) point that contains such colors. This condition should hold for every layer of w. Later, this will allow us to relate the conformal structures of a disc and a sphere.

If a popsicle sphere is aligned to a vertex, we will match the conformal structure of the sphere with that of the unstable disc (with an input and an output that are connected by a single common popsicle line), and hence the second alignability condition is needed.

Example 3.8. Let us give an example of the first alignability condition using the Figure 13. The tree on the left describes a neighborhood of a vertex in a rooted ribbon tree with colors containing $\{1,2,\ldots,8\}$. This vertex corresponds to a disc with 4 inputs and one output marked point. Each input is connected to an output via a geodesic in the disc, each of which are colored by $\{\{1,2\},\{3\},\{4\},\{5,6,7\}\}$. The layered tree on the right describes a neighborhood of a vertex in a rooted layered tree with colors containing $\{1,2,3,5,7\}$. This vertex corresponds to a sphere with 5 inputs and one output, whose popsicle lines as drawn on the right. We can check that the first alignability condition holds for each layer.

Let us make a few remarks. In this example, the colors {4,6,8} does not appear on the right. This is because we only keep track of a popsicle line whose sprinkle lies on the particular tree of sphere bubbles. The sprinkles for {4,6,8} should be in other trees of sphere bubbles or in some disc components.

Even though we do not record it, the popsicle lines of the color {4,6,8} do exist, as one can see the dotted line on the right for the color 4. Later, if we align the conformal structures of the disc and sphere component, the vertical lines on the left and on the right will be aligned as in the Figure 12.




FIGURE 13. Example of an alignable pairs and associated planar models

Now, let us explain the tree model for a general stable popsicle with sphere bubbles. For each disc D_v^2 for the vertex v, there may be several tree of sphere bubbles that are attached to D_v^2 . We use i, j to label these trees $\{T_{v,i,j}\}$, where i labels the order of distinct popsicle lines on D_v^2 , and j labels the order of trees on each popsicle line.

Definition 3.9. (*Tree model for general popsicles*) *Let us fix a non-decreasing flavor function* ϕ : $\tilde{F} = \{1, ..., l\} \rightarrow \{1, ..., n\}$. A tree model for popsicles with n-boundary input is a tuple $(\tilde{T}, \tilde{F}, \tilde{\psi}, \Phi, \Psi)$ consists of

- (1) *a decomposition of* \tilde{T} : *a decomposition* $\tilde{T} = T \bigsqcup_{v \in Vertex(T)} \{T_{v,i,j}\}$ where T is a rooted ribbon tree with n-leaves, and $\{T_{v,i,j}\}$ are rooted layered trees.
- (2) *a decomposition of* \widetilde{F} : *a decomposition* $\widetilde{F} = \bigsqcup_{v \in Vertex(T)} \widetilde{F}_v$ and $\widetilde{F}_v = F_v \sqcup (\sqcup_{i,j} F_{v,i,j})$ such that

$$\phi(F_{\nu}) \subset C_{e_{\nu,0}}, \quad \phi(F_{\nu,i,j}) \subset C_{e_{\nu,i}},$$

(3) *an induced decomposition of* ϕ : *a collection of induced map*

$$\phi_{\nu}: F_{\nu} \sqcup (\sqcup_{i,j} F_{\nu,i,j}) \to \{1, \dots, \operatorname{val}(\nu) - 1\}$$

such that $\phi(v)$, $v \in F_v$ is defined as in Definition 3.3, and $\phi(F_{v,i,j}) := i$

- (4) **F-labels**: a collection of bijective $F_{v,i,j}$ -labels $\tilde{\psi} = \{\psi_{v,i,j} : \text{Leaf}(T_{v,i,j}) \leftrightarrow F_{v,i,j}\},\$
- (5) alignment data I: an order-preserving function

 Φ : Vertex($\sqcup T_{\nu,i,j}$) \rightarrow Vertex(T) \sqcup Edge(T)

satisfying the following compatibility condition;

- w and $\Phi(w)$ are alignable,
- $\Phi|_{\operatorname{Vertex}(T_{v,i,j})} \subset \operatorname{Vertex}(T)_{>v} \cup \operatorname{Edge}(T)_{>v}$,
- Φ is strictly order-preserving on $\Phi^{-1}(\text{Vertex}(T))$.
- (6) alignment data II: a collection of strictly order-preserving surjection

 $\Psi = \left\{ \Psi_e : \Phi^{-1}(e) \to \{1, \dots, m_e\}, \ m_e \in \mathbb{N} \right\}$

for each e whose $\Phi^{-1}(e)$ is nonempty.

Here, order-preserving means with respect to the partial order on a rooted tree given by its orientation. We call a pair (Φ, Ψ) an alignment data for the model.

The alignment data I describes the alignment for each sphere component. Note that the order preserving condition naturally arises from the compactification process. The alignment data II concerns the case when there are several sphere components that are aligned to an edge *e*. In

this case, we need to specify their relative speed of formation of bubbles. The strictly order preserving condition is enforced within each tree of sphere bubbles.

Now, it is not difficult to associate an actual broken popsicle for the given tree model.

Definition 3.10. For a given tree model $(\tilde{T}, \tilde{F}, \tilde{\psi}, \Phi, \Psi)$ for popsicles, we obtain a general broken popsicle of type $(\tilde{T}, \tilde{F}, \tilde{\psi}, \Phi, \Psi)$ in the following way;

- (1) (popsicle discs on a ribbon tree) For each vertex v of a rooted ribbon tree T, assign a popsicles Σ_v of type $(val(v), \phi_v)$. Our popsicle disc carries not only F_v -many interior marked points, but also # {{ $F_{v,i,j}$ }-many nodal points over which sphere bubbles are attached.
- (2) (aligned with vertex) For a vertex w of $T_{v,i,j}$ such that $\Phi(w)$ is a vertex, we assign a popsicle sphere of type $(val(\Phi(w)) 1, val(w) 1, \phi_w)$, where ϕ_w is constructed as follows. A color of a layer satisfies $C^{(\phi,\psi)}(E_{w,k}) \subset C(e_{v,i_k})$ for the unique i_k . If $e_j \in E_{w,k}$, then we define $\phi_w(w_j^+) := i_k$. Finally, we require that a popsicle sphere Σ_w^+ and a popsicle disc $\Sigma_{\Phi(w)}$ are aligned to each other.
- (3) (aligned with edges) for a vertex w of $T_{v,i,j}$ such that $\Phi(w)$ is an edge, we assign a popsicle sphere of type $(1, val(w), \phi_w)$, where ϕ_w is just a constant map with value 1. It means that all incoming special points are on the single line.

Here is a simple example of a tree model and its associated popsicle.




FIGURE 14. Examples of combinatorial model for a general popsicle

We call the model is stable if it is stable as stable maps (after forgetting the popsicle structure and alignments). We write a moduli space of general stable popsicles of type $\star = (\tilde{T}, \tilde{F}, \tilde{\psi}, \Phi, \Psi)$ with *n*-boundary inputs by $P_{n,\tilde{F},\phi,\star}$. It is not hard to see that for a fixed *n* and $\phi : F \to \{1, ..., n\}$, only finitely many combinatorial types of stable trees are possible. The compactified moduli space we consider is given by

$$\overline{P}_{n,F,\phi} := \bigsqcup_{\star} P_{n,\widetilde{F},\phi,\star}.$$

4. Quantum CAP action and J-holomorphic popsicles with Γ -insertions

Let us recall quantum cap action and introduce *J*-holomorphic popsicles with Γ -insertions.

4.1. **Quantum cap action.** A quantum cap action of quantum cohomology can be defined on Fukaya category of a compact symplectic manifold [Aur07], and analogously that of symplectic cohomology can be defined on wrapped Fukaya category of a Liouville manifold [Gan13]. Let us briefly recall the definition. We refer readers to Appendix A for a brief account of symplectic cohomology $SH^{\bullet}(M)$ and the wrapped Fukaya category $\mathcal{WF}(M)$.

Let $P_{n,\{i\}}$ $(1 \le i \le n)$ be a moduli space of discs with one interior marked point z_1^+ and boundary marked points $z_0, ..., z_n$ such that z_0, z_1^+, z_i lie on a geodesic of D^2 . By applying an automorphism of the disc, we may say that z_1^+, z_0, z_i are precisely 0, 1, -1 of the disc. Consider a moduli space

$$\mathscr{P}_{n,\{i\}}(\Gamma; a_1, \ldots, a_{i-1}, b, a_{i+1}, \ldots, a_n, a_0)$$

of pseudo-holomorphic maps from $S \in P_{n,\{i\}}$ with an interior insertion $\Gamma \in SH^{\bullet}(M)$ at the puncture z_1^+ and with boundary insertions $(a_1, \dots, a_{i-1}, b, a_{i+1}, \dots, a_n)$. Here, we use a different symbol b to emphasize that it is a bimodule input.

Definition 4.1. A cochain level quantum cap action of Γ is an A_{∞} -bimodule map is defined by

$$\cap \Gamma : T(\mathcal{WF}) \otimes \underline{\mathcal{WF}} \otimes T(\mathcal{WF}) \to \mathcal{WF}$$

$$(a_1, \dots, a_{i-1}, \underline{b}, a_{i+1}, \dots, a_n) \mapsto (-1)^{\star_{n,\{i\}} - \deg\Gamma} \mathbf{P}_{n,\{i\}}(\Gamma; a_1, \dots, a_{i-1}, b, a_{i+1}, \dots, a_n),$$

$$\star_{n,\{i\}} = \sum_{j < i} (j-1) \cdot \deg a_i + i \cdot \deg b + \sum_{i < j} j \deg a_j$$

where $\mathbf{P}_{n,\{i\}}(\Gamma; a_1, \dots, a_{i-1}, b, a_{i+1}, \dots, a_n)$ is a sum of orientation operators associated to the moduli spaces for all possible a_0 (see Appendix A).

It is easy to check that $\cap \Gamma$ is a A_{∞} -bimodule map from diagonal A_{∞} -bimodule $\mathscr{WF}(M)_{\Delta}$ to itself:

$$\cap \Gamma: \mathscr{WF}(M)_{\Delta} \to \mathscr{WF}(M)_{\Delta}.$$

Also, this action is closely related to the closed-open map (see Appendix A) in the following way.

Proposition 4.2. Up to homotopy, we have

$$\cap \Gamma(b) = m_2(b, \operatorname{CO}_{L_2}(\Gamma))$$

where $CO_{L_2}: SH^{\bullet}(M) \to CW^{\bullet}(L_2, L_2)$ is a word-length zero component of the closed-open map.

Proof. Consider a 1-parameter family of moduli space of holomorphic discs with

- one outgoing boundary marking z_0 at 1
- one incoming boundary marking z_1 at -1
- one moving interior marking z_1^+ at -it, $t \in [0, 1]$

At t = 0, we get $P_{1,\{1\}}$. At t = 1, we get a moduli space of discs with disc bubble containing interior marked point. See Figure 15. It corresponds to a disc moduli space governing $m_2(b, CO_{L_2}(\Gamma))$.

We will construct a new A_{∞} -category \mathscr{C}_{Γ} associated to the quantum cap action $\cap \Gamma$. We first define its Hom space and the differential M_1 using the following A_{∞} -bimodule structure.

Definition 4.3. We first define \mathscr{C}_{Γ} as an A_{∞} -bimodule over $\mathscr{WF}(M)$ from the following distinguished triangle:

$$\mathscr{WF}(M) \xrightarrow{\cap \Gamma} \mathscr{WF}(M) \longrightarrow \mathscr{C}_{\Gamma} \longrightarrow$$

In particular, cohomologically we have

$$\operatorname{Hom}_{\mathscr{C}_{\Gamma}}^{\bullet}(L_{1},L_{2}) \simeq \operatorname{Cone} \left(CW^{\bullet}(L_{1},L_{2}) \xrightarrow{\cap \Gamma} CW^{\bullet}(L_{1},L_{2}) \right)$$




FIGURE 15. Homotopy between $\cap \Gamma$ and CO(Γ)

Instead of the A_{∞} -bimodule structure on \mathscr{C}_{Γ} , we will construct an A_{∞} -category structure on \mathscr{C}_{Γ} in the next section. The action $\cap \Gamma$ vanishes up to homotopy in \mathscr{C}_{Γ} .

Intuitively,

• objects of \mathscr{C}_{Γ} are twisted complexes

$$\left(\begin{array}{c} L \xrightarrow{\operatorname{CO}(\Gamma)} L \end{array} \right), \ L \in \mathcal{WF}(M).$$

In many cases, they can be realized as a geometric surgery of *L* with itself along $CO(\Gamma)$.

• the space of morphisms is a "half" of the morphisms between twist complexes. It consists of

$$a + \epsilon b'' = \begin{pmatrix} a & 0 \\ b & a \end{pmatrix} \in CW^{\bullet} \left(L_1 \xrightarrow{\operatorname{CO}(\Gamma)} L_1 , L_2 \xrightarrow{\operatorname{CO}(\Gamma)} L_2 \right)$$

This intuition works well in differential graded world (see Section 6), but fails in all possible ways in the A_{∞} -world. For example, A_{∞} -compositions do not preserve the above space of morphisms. We will consider a new type of *J*-holomorphic curves to construct a precise A_{∞} -structure: Namely, the new A_{∞} -structure that we will put on \mathscr{C}_{Γ} is not an algebraic one, but rather a geometric one, which are *J*-holomorphic maps from popsicles. It is natural in a sense that the cap action uses one hyperbolic geodesic line, and for the A_{∞} -structure, we work with arbitrarily many geodesic lines on a disc.

4.2. **Popsicle maps with** Γ **-insertions.** Let us introduce pseudo-holomorphic maps from popsicles with Γ -insertions. Consider Lagrangians L_0, \dots, L_n in the wrapped Fukaya category of M.

Definition 4.4. Let $a_i \in CF^{\bullet}(L_{i-1}, L_i)$ and $\Gamma \in SH^{\bullet}(M)$. Define

$$\mathscr{P}_{n,F,\phi}(\Gamma; a_1, \ldots, a_n, a_0)$$

be a moduli space of pseudo-holomorphic maps

$$\left\{ u: S \to M \mid S \in P_{n,F,\phi} \right\}$$

that satisfies

- a boundary segment from z_i to z_{i+1} goes to L_i ,
- a boundary marking z_i goes to a_i ,

• all interior markings are asymptotic to Γ .

Such a map u is called a regular pseudo-holomorphic popsicle.

Using the compactification $\overline{P}_{n,F,\phi}$ of the domain popsicles in the last section, the compactification for regular pseudo-holomorphic popsicles can be carried out in a standard way. The new feature of alignment data that was used in the compactification of the domain, concerns restriction of conformal structures on various disc and sphere components in the limiting stable popsicles. And we do **not** need to align Floer data for the aligned domains as the alignment is a restriction only on the domain.

As we have shown that $\overline{P}_{n,F,\phi}$ is a manifold with corners, we can choose Floer data continuous and consistently by extending it from the lowest dimensional strata in an inductive manner as usual. We put no other restriction on the choice of domain-dependent Hamiltonians H_S , F_S and almost complex structures J_S . Transversality can be achieved in a standard way as we do not require compatibility of almost complex structures assigned at aligned domains.

Remark 4.5. We also want this choice to be mildly equivariant following [AS10], [Sei18]. Suppose several sprinkles are on the same popsicle line for a broken popsicle of type \star . Then a symmetry group acts on the corresponding strata $P_{n,\tilde{F},\star}$ by permuting their positions. We want our choice of Floer data to be invariant with respect to this action. It does not harm transversality argument because $P_{n,\tilde{F},\star}$ does not have any fixed point of the action and hence does not pose any problem on surjectivity of linearized Fredholm problem.

Fix a collection of Hamiltonian chords $a_i \in \chi(L_{i-1}, L_i)$, i = 1, ..., n. Let us describe possible limits in the compactification

$$\overline{\mathscr{P}}_{n,F,\phi}(\Gamma;a_1,\ldots,a_n,a_0).$$

As a first step, we allow our a tree model \tilde{T} of a broken popsicle may have semistable vertex. More precisely, a rooted ribbon tree $T \subset \tilde{T}$ may have a bivalent vertices v with $\tilde{F}_v = \emptyset$, and a rooted layered tree $T_{v,i,j} \in \tilde{T}$ may contains bivalent vertices. We need to specify an alignment data for the new bivalent vertices on a tree of sphere bubbles so that it is also compatible with the original alignment data. This can be done as in the last section and we omit the details. In addition, we need to add cylinders (semi-stable spheres) splitting off at the interior marked points of a popsicle disc (say v). We need to add the corresponding new trees to the family $\{T_{v,i,j}\}$ but bubbling is local and do not carry alignment data.

Next, choose a Hamiltonian chord a_e for each finite edge $e \in \text{Edge}(T)$, and choose a Hamiltonian orbit $\gamma_{e'}$ for each edge $e' \in \text{Edge}(T_{v,i,j})$. Let $v \in \text{Vertex}(T)$ be a vertex and let $\Phi^{-1}(v) = \{w_i\}$. We assign the following moduli space of maps

(4.1)
$$\mathscr{P}_{\nu} := \left\{ (u_{\nu}, \{u_{w_i}\}) | u_{\nu} : \Sigma_{\nu} \to M, \ u_{w_i} : \Sigma_{w_i}^+ \to M, \ (\Sigma_{\nu}, \Sigma_{w_i}^+) \text{ are aligned} \right\}$$

such that

- each u_v and u_{w_i} satisfy pseudo-holomorphic equation (A.1).
- u_v satisfies Lagrangian boundary condition induced on Σ_v ,
- u_v and u_{w_i} satisfy asymptotic condition on their strip-like/cylindrical ends.
- If $\Sigma_v \simeq Z$ (resp. $\Sigma_{w_i} \simeq S^1 \times \mathbb{R}$) is semistable, then u_v (resp. u_{w_i}) is non-constant.

An aligned pair $(\Sigma_v, \{\Sigma_{w_i}\})$ carries a translation \mathbb{R} symmetry if Σ_v and $\sigma_{w_i}^+$ are all semi-stable. In that case, we define a similar moduli space as a quotient with respect to that symmetry group

(4.2)
$$\mathscr{P}_{\nu} := \left\{ (u_{\nu}, \{u_{w_i}\}) | u_{\nu} : \Sigma_{\nu} \to M, \ u_{w_i} : \Sigma_{w_i}^+ \to M, \ (\Sigma_{\nu}, \Sigma_{w_i}^+) \text{ are aligned} \right\} / \mathbb{R}.$$

Similarly, when $\Psi_e^{-1}(k) = \{w_j\}$ for $e \in \text{Edge}(T)$, we can define (following the case of (4.1))

(4.3)
$$\mathscr{P}_{e,k} := \left\{ \{u_{w_j}\} | u_{w_j} : \Sigma_{w_j}^+ \to M, \ (\Sigma_{w_j}^+)_j \text{ are aligned} \right\},$$

Define

$$\mathscr{P}_{n,\widetilde{F},\phi,\star}(\Gamma;a_1,\ldots,a_n,a_0) := \bigcup_{a_e,\gamma_{e'}} \left(\bigcup_{v \in \operatorname{Vertex}(T)} \mathscr{P}_v \right) \cup \left(\bigcup_{e \in \operatorname{Edge}(T)} \bigcup_{1 \le k \le m_e} \mathscr{P}_{e,k} \right)$$

and finally we set

$$\overline{\mathscr{P}}_{n,F,\phi}(\Gamma;a_1,\ldots,a_n,a_0) := \bigcup_{\star} \mathscr{P}_{n,\widetilde{F},\phi,\star}(\Gamma;a_1,\ldots,a_n,a_0)$$

Lemma 4.6. For a generic choice of universal and consistent Floer data, we have the following.

- (1) Moduli spaces $\overline{\mathscr{P}}_{n,F,\phi}(\Gamma; a_1, \ldots, a_n, a_0)$ are smooth manifolds with corners and compact.
- (2) For a given input Γ and a_i , (i = 1, ..., n), there are only finitely many a_0 for which

$$\overline{\mathscr{P}}_{n,F,\phi}(\Gamma;a_1,\ldots,a_n,a_0)$$

is non-empty . (3) *It is a manifold of dimension*

$$|F|(1 - \deg \Gamma) + n - 2 + \deg a_0 - \sum_{i=1}^n \deg a_i$$

Proof. Recall that the domain broken popsicles are shown to be smooth manifolds with corners in Theorem B.1. The desired transversality and compactness for pseudo-holomorphic maps can be proved as well using the standard methods and the above theorem. A standard index formula tells us that

$$\dim_{\mathbb{R}}\overline{\mathscr{P}}_{n,F,\phi}(\Gamma;a_1,\ldots,a_n,a_0) = \dim_{\mathbb{R}}\overline{\mathscr{M}}_{|F|;n,1}(\Gamma,\ldots,\Gamma;a_1,\ldots,a_n,a_0) - |F|$$
$$= (2|F|+n-2) + \deg a_0 - \sum_{i=1}^n \deg a_i - |F| \cdot \deg \Gamma - |F|$$
$$= |F|(1-\deg\Gamma) + n - 2 + \deg a_0 - \sum_{i=1}^n \deg a_i.$$

Let $\mathbf{P}_{n,E\phi}^{\Gamma}$ be a sum of orientation operators associated to the zero-dimensional component of

$$\overline{\mathscr{P}}_{n,F,\phi}(\Gamma;a_1,\ldots,a_n,a_0)$$

for all possible a_0 . A degree of this operator is $2 - n - |F|(1 - \deg \Gamma)$. Also, $\mathbf{P}_{n,F,\phi}^{\Gamma}$ describes A_{∞} -structure $\{m_n\}$ if F is empty, and describes quantum cap action $\cap \Gamma$ if |F| = 1.

FUKAYA CATEGORY FOR LANDAU-GINZBURG ORBIFOLDS

5. The New A_{∞} -category \mathscr{C}_{Γ}

In this section, we will define a new A_{∞} -category \mathscr{C}_{Γ} of a Liouville manifold M using popsicles with Γ -insertions, where Γ is a certain symplectic cohomology class of M. A key feature of the new A_{∞} -category is that the image of quantum cap actions by Γ vanishes in cohomology of \mathscr{C}_{Γ} .

To control the new sphere bubbling phenomenon of the popsicle moduli space, we restrict ourselves to the following setting.

Assumption 5.1. We will assume that a Liouville manifold M satisfies the following in this section.

- (1) $c_1(M) = 0$.
- (2) Reeb flow \mathscr{R}_t on ∂M defines a free S^1 -action.
- (3) *linearized Reeb flow is complex linear.*
- (4) Robbin-Salamon index of a principal Reeb orbit component is non-negative.
- (5) Symplectic cochain Γ_M that represents the fundamental class of principal orbits is closed.

More details on these assumptions will be given in Section 5.2. The Milnor fiber quotients that we will investigate later in this paper do not satisfy the first two hypotheses, and we will explain later how to generalize the construction in those cases.

Under the above assumptions, we prove the following theorem in this section.

Theorem 5.2. Given a Liouville manifold M and the element $\Gamma_M \in SH^{\bullet}(M)$ satisfying the above assumption, we define a new \mathbb{Z} -graded A_{∞} -category \mathscr{C}_{Γ} such that

- (1) \mathscr{C}_{Γ} has the same set of objects as the wrapped Fukaya category $\mathscr{WF}(M)$,
- (2) for two objects L_1, L_2 , its morphisms are given by

$$\operatorname{Hom}_{\mathscr{C}_{\Gamma}}(L_1, L_2) = CW(L_1, L_2) \oplus CW(L_1, L_2)\epsilon$$

where $CW(L_1, L_2)$ is the morphism space for $\mathcal{WF}(M)$ and $\deg \epsilon = -1 + \deg \Gamma_M$,

- (3) a natural inclusion $\Psi : \mathcal{WF}(M) \to \mathcal{C}_{\Gamma}$ is an A_{∞} -functor,
- (4) regarding the A_{∞} -category \mathscr{C}_{Γ} as an A_{∞} -bimodule over $\mathscr{WF}(M)$ (using Ψ), we have a distinguished triangle of A_{∞} -bimodules

$$\mathscr{WF}(M)_{\Delta} \xrightarrow{\cap \Gamma} \mathscr{WF}(M)_{\Delta} \longrightarrow \mathscr{C}_{\Gamma} \longrightarrow$$

Roughly speaking, A_{∞} -operations are defined by *popsicles with* Γ -*insertions at sprinkles*. One of the difficulty is to rule out any codimension one (aligned) sphere bubbles with Γ -insertions, and we achieve it using Conley-Zehnder indices and action values under the Assumption 5.1. The rest of codimension one strata are analogous to those of Abouzaid-Seidel [AS10], and Seidel [Sei18] and we obtain the desired A_{∞} -category.

Remark 5.3. *Popsicle structures were introduced by Abouzaid-Seidel* [AS10] *and Seidel* [Sei18]. *It is a good moment to point out differences and similarities.*

In [AS10],[Sei18], sprinkles were allowed to coincide resulting a different compactification of popsicles. In [AS10], popsicles were used to mark the places to put sub-closed one forms for continuation maps between linear Hamiltonians. They localize the continuation map and obtain a big chain complex involving countable family of Hamiltonians $\{nH\}_{n\in\mathbb{N}}$. This is a definition of

wrapped Fukaya category in a linear Hamiltonian setting. Seidel [Sei18] considered the continuation map for Lefschetz fibration, but he also considered its cone

$$CF^{\bullet}(L_0, L_1; -H) \rightarrow CF^{\bullet}(L_0, L_1; 0) \rightarrow CF^{\bullet}(L_0, L_1; \text{conti}),$$

not only its localization. Popsicle maps were used to construct an A_{∞} -algebra structure on $CF^{\bullet}(L_0, L_1; \text{conti})$. This brings out the effect of removing contributions from compact part of Lefschetz fibration, and recovers the Floer cohomology of the fiber.

Our geometric setting is different from the references because we regard sprinkles as genuine inputs for the symplectic cohomology class Γ . But the algebraic properties we desire are similar to those of [Sei18]. In our case, we want to remove the effect of the quantum cap action $\cap\Gamma$ from the wrapped Fukaya category. Namely, wrapped generator in the image of quantum cap action will be killed.

5.1. **Codimension one analysis.** Let us discuss the codimension one strata of the compactification

$$\overline{\mathscr{P}}_{n,F,\phi}(\Gamma;a_1,\ldots,a_n,a_0)$$

The obvious ones are given by splitting off a non-trivial cylinder (at sprinkles) or a strip (at boundary markings). These will correspond to the differential of Γ or a_i 's. Beyond these ones, the remaining codimension one strata arise when the domain regular popsicle degenerates. Recall from (B.1), that we can explicitly determine the codimension of the broken popsicle of type $\star = (\tilde{T}, \tilde{F}, \tilde{\psi}, \Phi, \Psi)$ to be

$$|\operatorname{Vertex}(T)| - 1 + \sum_{e \in \operatorname{Edge}(T)} m_e.$$

Since $m_e \ge 0$, codimension can may arise in the following cases:

- (1) |Vertex(T)| = 1 and $m_e = 1$ for a single *e* and $m_{e'} = 0$ for other *e'*.
- (2) |Vertex(T)| = 2 and $m_e = 0$.

The first case do not arise, since there is only one disc component, there is no possible edge *e*. The second case is quite interesting. Label two disc components as $Vertex(T) = \{v_1, v_2\}$ with an edge *e* from v_1 to v_2 .

If there are no sphere components, then any such stratum is given by a broken popsicle consisting of two popsicles without sphere bubbles, and these will describe the A_{∞} -relations (as in Abouzaid-Seidel [AS10] and Seidel [Sei18]).

Let us consider the case that there is at least one sphere component. As the alignment data m_e counts sphere bubbles that are aligned to the edge e, the condition $m_e = 0$ implies that all sphere components Σ_w^+ (if there is) should be aligned to the disc component v_2 . Note that there is no restriction on the number of sphere components. From the restriction on the alignment data, all sphere components should be attached to the disc v_1 (if they are stacked on each other, then it violates the strictly order preserving condition). We may have several sphere bubbles in codimension one! (see Figure 3 for an example).

Therefore, sphere components in the compactification provide genuine obstructions of defining A_{∞} -structure from $\overline{\mathscr{P}}_{n,F,\phi}(\Gamma; a_1, \ldots, a_n, a_0)$. Fortunately, we can show these obstructions vanish in certain favorable situations. The following degree estimate that we establish will be crucial for this purpose. Let us first set up the notations for such codimension one strata with sphere bubbles. Consider the following pseudo-holomorphic broken popsicle with appropriate boundary conditions:

$$u := (u_{\nu_1}, u_{\nu_2}, \{u_{w_i}\}) : \Sigma = \Sigma_{\nu_1} \cup \Sigma_{\nu_2} \cup \Sigma_{w_1}^+ \cup \cdots \cup \Sigma_{w_m}^+ \to M,$$

We assume that all spheres $\Sigma_{w_i}^+$ are aligned to the stable disc Σ_{v_2} . We assume that *u* is Fredholm-regular and rigid. We make the following notations.

- Σ_{ν_2} has *n* boundary insertions,
- $k_{v_i} = F_{v_i}$ denotes a number of sprinkles on Σ_{v_i} for i = 1, 2,
- $k_{w_i} = \text{val}(w_i) 1$ denotes a number of incoming insertion of $\Sigma_{w_i}^+$ (where Γ 's are placed) and $k_{w_i} \ge 2$,
- $\gamma_{w_i} \in \mathcal{O}$ denotes an output of u_{w_i} ,
- \mathcal{M} denotes a moduli space of underlying aligned pairs of popsicles $\Sigma = \left(\Sigma_{\nu_1}, \Sigma_{\nu_2}, \{\Sigma_{w_i}^+\}\right)$.

Observe that we can decompose its tangent space as

(5.1)
$$T_{\Sigma}\mathcal{M} \simeq \left(\mathbb{R}^{k_{\nu_{1}}} \times \mathbb{R}^{m-1}\right) \times \left(T\mathcal{M}_{0,n+1} \times \mathbb{R}^{k_{\nu_{2}}}\right) \times \mathbb{R}^{k_{w_{1}}-1} \times \cdots \times \mathbb{R}^{k_{w_{m}}-1}.$$

Here, the first two components handle the deformation of marked points on the discs Σ_{v_1} , Σ_{v_2} respectively and the component $\mathbb{R}^{k_{w_i}-1}$ for that of the aligned sphere Σ_{w_i} for $i = 1, \dots, m$.

Proposition 5.4. In this situation, the following inequality holds;

(5.2)
$$k_{w_i} \deg \Gamma - k_{w_i} - n + 3 \le \deg \gamma_{w_i} \le k_{w_i} \deg \Gamma - k_{w_i} + 1$$

Proof. Consider a linearized Fredholm problem associated to u. It can be written (using the decomposition (5.1))

$$(5.3) D_u: T_{\Sigma}\mathcal{M} \times \left(V_{\nu_1} \times V_{\nu_2} \times V_{w_1} \times \dots \times V_{w_k} \right) \to \left(W_{\nu_1} \times W_{\nu_2} \times W_{w_1} \times \dots \times W_{w_k} \right)$$

Let us write *D* in the matrix form.

(5.4)
$$D_{u} = \begin{pmatrix} D_{u_{v_{1}}} & 0 & 0 & 0 & 0 & \cdots & 0 \\ 0 & D_{0v_{2}} & D_{v_{2}v_{2}} & 0 & 0 & \cdots & 0 \\ 0 & D_{0,w_{1}} & 0 & D_{w_{1}w_{1}} & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & & \\ 0 & D_{0w_{k}} & 0 & 0 & 0 & \cdots & D_{w_{k}w_{k}} \end{pmatrix}.$$

Let us explain each components. For v_1 and v_2 , we have

$$D_{u_{\nu_1}}: (\mathbb{R}^{k_{\nu_1}} \times \mathbb{R}^{m-1}) \times V_{\nu_1} \to W_{\nu_1}$$

$$D_{u_{v_2}} = (D_{0v_2}, D_{v_2v_2}) : T\mathcal{M}_{0,n+1} \times \left(\mathbb{R}^{k_{v_2}} \times V_{v_2}\right) \to W_{v_2}$$

is a linearized Fredholm operator associated to u_{v_2} and

$$D_{u_{w_i}} = (D_{0w_i}, D_{w_iw_i}) : T\mathcal{M}_{0,n+1} \times \left(\mathbb{R}^{k_{w_i}-1} \times V_{w_i}\right) \to W_{w_i}$$

are linearized Fredholm operators associated to u_{w_i} respectively. In this way, we can separate the contribution of $T\mathcal{M}_{0,n+1}$, which affects the disc v_2 as well as the spheres $\{w_i\}$ due to the alignment.

Since *u* is Fredholm regular and rigid, D_u is surjective and $\text{Ker}(D_u) = \{0\}$. And each D_{u_v} and $D_{u_{w_i}}$ is surjective because D_u is surjective. It means that u_v and u_{w_i} are Fredholm regular in their own rights.

Next, observe that $\operatorname{Ker} D_{u_{w_i}} \cap \{0\} \times \left(\mathbb{R}^{k_{w_i}-1} \times V_v\right) = \{0\}$. If not, we can extend a non-zero vector $(0, \alpha, \beta) \in \operatorname{Ker} D_{u_{w_i}} \cap \{0\} \times \left(\mathbb{R}^{k_{w_i}-1} \times V_v\right)$ to

 $(0,\ldots,0,\alpha,\beta,0,\ldots,0) \in \operatorname{Ker} D_u,$

which contradicts to Ker $D_u = \{0\}$. Therefore, Ker $D_{u_{w_i}}$ must projects isomorphically into $T\mathcal{M}_{0,n+1}$ as a vector space, which again implies dimKer $D_{u_{w_i}} \leq \dim T\mathcal{M}_{0,n+1}$. Combine these two observations with Fredholm index theorem, we get

$$0 \leq \operatorname{dim}\operatorname{Ker} D_{u_{w_i}} = \operatorname{Index} D_{u_{w_i}} = \operatorname{dim} T \mathcal{M}_{0,n+1} + (k_{w_i} - 1) + \operatorname{deg} \gamma_{w_i} - k_{w_i} \operatorname{deg} \Gamma \leq \operatorname{dim} T \mathcal{M}_{0,n+1}.$$

The proposition follows from these inequalities.

Corollary 5.5. In this situation, there is at least one component u_{w_i} such that

$$\deg \gamma_{w_i} \le k_{w_i} \deg \Gamma - \frac{k_{w_i}}{2}.$$

Proof. Since $k_{w_i} \ge 2$ for all *i*, we know that the number of sphere component (denoted by *m*) is less then the half of the number of interior marked points. Therefore,

(5.5)
$$\sum \deg \gamma_{w_i} \le (\sum k_{w_i}) \deg \Gamma - (\sum k_{w_i}) + m \le (\sum k_{w_i}) \deg \Gamma - (\sum k_{w_i})/2.$$
The corollary follows by the pigeon hole principle.

5.2. **Vanishing of sphere bubbles.** We prove the following key proposition, which allows us to define the desired A_{∞} -structure.

Proposition 5.6. Under the assumption 5.1, sphere bubbles of codimension one do **not** appear in the compactification $\overline{\mathcal{P}}_{n,F,\phi}(\Gamma; a_1, ..., a_n, a_0)$.

Let us first explain the assumptions in more detail. The first assumption $c_1(M) = 0$ is to use degree arguments for the vanishing. The second assumption on the existence of free S^1 action on ∂M is given for two purposes. The first is that we will have such a S^1 -action (although it is not free) in our main application, and the second reason is that S^1 -action make the symplectic cohomology setting to be Morse-Bott and we can use associated spectral sequences. We recall the third assumption from [KvK16] which makes the local system associated to the Morse-Bott components to be trivial:

Definition 5.7. A linearized Reeb flow is complex linear if there exist a compatible complex structure *J* for the contact structure ξ (with $d\lambda$) such that for every periodic Reeb orbit, its linearized Reeb flow is complex linear with respect to some unitary trivialization of $(\xi, J, d\lambda)$.

Under the assumption of the S^1 -action, Reeb orbits are degenerate and it is more convenient to work with the Robbin-Salamon index. We will make a perturbation in the Morse-Bott setting to define the Hamiltonian orbit Γ_M that corresponds to the (fundamental class of) the set of principal orbits. We suppose that Γ_M is a closed element in the symplectic cochain complex. Perturbation only guarantees that it is closed in the neighborhood of the Morse-Bott component,

 \square

but pseudo-holomorphic cylinders might escape such a neighborhood, and closedness has to be checked using the geometry of *M*.

Now, let us explain the proof of the proposition in two steps.

5.2.1. Step I: spectral sequence associated to an energy filtration.

Lemma 5.8. There is a spectral sequence converging to $SH^*(M)$

$$E_1^{pq} = \begin{cases} H^q(M) & (p=0) \\ H^{p+q-p \cdot \mu_{RS}(\partial M_1)}(\partial M) & (p<0) \\ 0 & (p>0) \end{cases}$$

Proof. We only sketch the construction (see [Sei06] and [KvK16] for more details). Choose a Hamiltonian function H which is C^2 -small in the interior, and also a C^2 -small time dependent Hamiltonian F_t which is constructed from Morse function f on ∂M so that $H_t = H + F_t$. Non-trivial orbits of H_t appear as a small perturbation of possible family of (degenerate) orbits. Since the action of \Re_t is free, there are integer many such families, classified by its period, all diffeomorphic to ∂M . An action value of orbits of period n is given by

$$\begin{aligned} \mathscr{A}_{H_{S^{1}}}(\gamma) &:= -\int_{S^{1}} \gamma^{*} \lambda + \int_{0}^{1} H_{S^{1}}(\gamma(t)) dt \\ &= -2 \int_{0}^{1} r^{2} dt + \int_{0}^{1} H(\gamma(t)) dt + \int_{0}^{1} F(\gamma(t)) dt, \ (H_{S^{1}} = H(r) + F(r, t)) \\ &= -\int_{0}^{1} r^{2} + \operatorname{error} \\ &\in (-r^{2} - \epsilon, -r^{2} + \epsilon), \ \epsilon = \max f - \min f. \end{aligned}$$

Here, we choose small enough $\epsilon \ll 1$ so that $[-(r+1)^2 - 2\epsilon, -(r+1)^2 + 2\epsilon]$ is disjoint from $[-r^2 - 2\epsilon, -r^2 + 2\epsilon]$ for all $r \in \mathbb{Z}_{\geq 0}$. We define an action filtration on $CH^*(M; H_{S^1})$ as

$$F^{-n} := \left\{ \gamma \in \mathcal{O}_{H_{S^1}} | \mathcal{A}_{H_{S^1}}(\gamma) > -n^2 - \epsilon \right\},\$$

the spectral sequence associated to F^* converges to $SH^*(M)$, and its first page is

$$E_1^{pq} = \begin{cases} H^q(M) & (p=0) \\ H^{p+q+\star_p}(\partial M) & (p<0) \end{cases}$$

A degree shifting number \star_p (for p < 0) is given as $\mu_{RS}(\partial M_p)$, where ∂M_p denotes a component corresponds to the period -p component. Notice that, we have $\mu_{RS}(\partial M_p) = p \cdot \mu_{RS}(\partial M_1) = -p \cdot \deg(\Gamma_M)$ because Γ_M is chosen to be $1_{\partial M} \in H^0(\partial M_1)$.

5.2.2. Step II: vanishing.

Proposition 5.9. If Robbin-Salamon index of a principle component (that is, $\mu_{RS}(\partial M_1)$) is nonnegative, then popsicle spheres with two or more Γ_M inputs vanish.

Proof. For simplicity, we write $\mu := \mu_{RS}(\partial M_1)$.

Let ξ be an output of a sphere component satisfying Corollary 5.5 with *N*-many Γ_M inputs. Since deg(Γ_W) = $-\mu$, we obtain the following inequality by Corollary 5.5

(5.6)
$$\deg \xi \leq N \cdot \deg \Gamma_M - \frac{N}{2} = -N \cdot \mu - \frac{N}{2}.$$

Suppose $\mu = 0$. Then Lemma 5.8 implies SH^* is concentrated in positive degrees, while 5.6 implies the degree of ξ is less then -N/2. This implies the proposition.

For the general case $\mu \ge 0$, we proceed as follows. Suppose $\xi \in F^{-l}$. Since μ is non-negative, we must have deg $\xi \ge -l \cdot \mu$ from Lemma 5.8. Combining with the inequality from Corollary 5.5, we have

(5.7)
$$-l \cdot \mu \le \deg \xi \le -N \cdot \mu - \frac{N}{2} \Longrightarrow l \ge N \left(1 + \frac{1}{2\mu} \right)$$

On the other hand, from the action inequality, we have

(5.8)
$$\frac{-l^2 + \epsilon}{N} \ge \frac{\mathscr{A}_{H_{S^1}}(\xi)}{N} \ge \sum_{i=1}^N \mathscr{A}_{H_{S^1}}(\Gamma_M) \ge N \cdot (-1 - \epsilon)$$

Here, we choose our weights at each cylindrical ends as $w_0^- = N$ and $w_i^+ = 1$. Thus we obtain

$$(5.9) label{eq:loss} l^2 \le N^2(1-\epsilon) - \epsilon$$

Combine 5.7 and 5.9, we should have the following inequality due to the sphere component.

$$N^2 \left(1 + \frac{1}{\mu} \right)^2 \le l^2 \le N^2 (1 - \epsilon) - \epsilon$$

If we choose ϵ small enough, this inequality does not hold for any integer $N \ge 2$. Hence such a sphere bubble does not arise.

5.3. A_{∞} -category \mathscr{C}_{Γ} . We construct a new \mathbb{Z} -graded A_{∞} -category \mathscr{C}_{Γ} under the assumption 5.1.

We start with the following cancellation results from Abouzaid-Seidel.

Proposition 5.10. ([AS10]) If $\phi: F \to \{1, ..., n\}$ is not injective, then $\mathbf{P}_{n, E, \phi}^{\Gamma}$ vanishes.

Proof. The assumption means that at least one popsicle stick carries more than two interior markings. Then Sym^{ϕ} contains a nontrivial transposition. Since we put the same class Γ for all interior markings, the transposition extends to $\overline{\mathscr{P}}_{n,F,\phi}(\Gamma; a_1, \ldots, a_n)$ also. It induces an orientation-reversal automorphism on $\overline{\mathscr{P}}_{n,F,\phi}$. Therefore the contribution of this moduli space should vanish.

Now we can focus on the case when $\phi: F \to \{1, ..., n\}$ is injective. Then *F* can be considered as a subset $\{i_1, ..., i_k\} \subset \{1, ..., n\}$. In this case, we omit a notation ϕ and simply write $\overline{\mathscr{P}}_{n,F}$ and $\mathbf{P}_{n,F}^{\Gamma}$.

Definition 5.11. An admissible cut of F consists of

- (1) $n_1, n_2 \ge 1$ such that $n_1 + n_2 = n + 1$,
- (2) a number $i \in \{1, ..., n\}$,
- (3) $F_1 \subset \{1, ..., n_1\}$ and $F_2 \subset \{1, ..., n_2\}$ such that $|F_1| + |F_2| = |F|$,

satisfying the following properties;

• $F \supset \{k | k \in F_1, k < i\}$ and $F \supset \{k + n_2 - 1 | k \in F_1, k > i\}$,

• $F \supset \{k + i - 1 | k \in F_2\}.$

If $i \notin F_1$, then this completely recovers F. Otherwise, F has one more element among $\{i, i+1, ..., (i+n_2-1)\}$.

An admissible cut describes a stratum $P_{n_1,F_1} \times P_{n_2,F_2}$ of a moduli space $\overline{P}_{n,F}$. They describe precisely codimension 1 strata whose associated flavour $\phi_j : F_j \rightarrow \{1, ..., n_j\}$ (j = 1, 2) is still injective (when sphere bubbles of codimension one do not appear). Combined with 5.10, we get a quadratic relation

$$\sum_{\text{\forall admissible cuts}} (-1)^{\clubsuit} \mathbf{P}_{n_1,F_1}^{\Gamma}(a_1,\ldots,a_{i-1},\mathbf{P}_{n_2,F_2}^{\Gamma}(a_i,\ldots,a_{i+|F_2|-1}), a_{i+|F_2|},\ldots,a_n) = 0,$$

$$\clubsuit = n_2 i + i + 1 + n_1 |F_2| + (n_2 + |F_2|) \left(\sum_{j \le i+|F_2|} \deg a_j\right) + \left| \left\{ (f_1, f_2) \in F_1 \times F_2 | f_1 < f_2 \text{ inside } F \right\} \right|$$

Now we are ready to define a new A_{∞} -category.

Definition 5.12. The A_{∞} -category \mathscr{C}_{Γ} for $\Gamma_M \in SH^{\bullet}(M)$ from Assumption 5.1, consists of

- (1) a set of objects $Ob(\mathscr{C}_{\Gamma}) = Ob(\mathscr{WF}(M))$.
- (2) morphisms between two objects

$$\operatorname{Hom}_{\mathscr{C}_{\Gamma}}(L_1, L_2) = CW(L_1, L_2) \oplus CW(L_1, L_2)\epsilon$$

Here deg $\epsilon = -1 + \mu_{RS}(\Gamma)$, and we denotes the element of this complex by $c := a + \epsilon b$. (3) An A_{∞} -structure $\{M_n\}_{n=1}^{\infty}$ is given as follows. We may write

$$M_n(c_1,...,c_n) = M_n^a(c_1,...,c_n) + \epsilon M_n^b(c_1,...,c_n)$$

(a) Suppose $c_i = a_i$ for all *i* (all the inputs do not have ϵ components), then we set

$$M_n(a_1,\ldots,a_n)=m_n(a_1,\ldots,a_n)$$

where $\{m_n\}$ is the A_∞ -operation for $\mathcal{WF}(M)$. (b) Suppose $c_i = \epsilon b_i$ for $i \in \{i_1, \dots, i_k\}$, and $c_i = a_i$ for $i \notin \{i_1, \dots, i_k\}$. Then we set

$$F = \{i_1, \dots, i_k\}, \ \widehat{F}^j = \{i_1, \dots, \widehat{i_j}, \dots, i_k\},$$

and define

$$M_n^a(c_1,...,c_n) = (-1)^{\star_{n,F}^a} \mathbf{P}_{n,F}^{\Gamma}(a_1,...,b_{i_1},...,b_{i_j},...,b_{i_k},...,a_n)$$

$$M_n^b(c_1,...,c_n) = \sum_{j=1}^k (-1)^{\star_{j-1} + \star_{n,\widehat{F}^j}^b} \mathbf{P}_{n,\widehat{F}^j}^{\Gamma}(a_1,...,b_{i_1},...,b_{i_j},...,b_{i_k},...,a_n)$$

If we use the common notion x_i to denote a_i and b_i . Then

$$\star_{n,F}^{a} = \sum_{j} j \deg x_{j} + \sum_{f \in F, l > f} (\deg x_{l} - 1)$$

$$\star_{n,\widehat{F}^{j}}^{b} = \sum_{j} j \deg x_{j} + \sum_{f \in \widehat{F}^{j}, l > j} (\deg x_{l} - 1)$$

$$\star_{j} = \sum_{l=1}^{j} (\deg x_{l} - 1)$$

As a sanity check, let us check the degree of M_n . Recall that the degree of $\mathbf{P}_{n,F}^{\Gamma}$ is $2 - n - |F|(1 - \mu_{RS}(\Gamma_M))$, and the corresponding operation in terms of M_n has degree 2 - n because the degree of ϵ is $-1 + \mu_{RS}(\Gamma_M)$ and we have |F| many ϵ -inputs.

We follow the sign analysis of Abouzaid-Seidel [AS10], to which we refer readers for full details. Using the orientation operators, they have separated the signs from the popsicle domain and the popsicle maps. They used the standard gluing theorem identify the sign factors from the popsicle maps, and the signs from degeneration of popsicle domains and the rearrangements of factors determined the sign convention. As we have a fixed interior input Γ_M , the contribution from the popsicle maps can be still glued using the standard gluing theorem, whereas the remaining part of the sign analysis is the same as the reference. Thus, the same sign convention can be used in our cases as well.

5.4. **Proof of** A_{∞} **-identity.**

Proposition 5.13. \mathscr{C}_{Γ} is an A_{∞} -category. Namely, for any composable (c_1, \ldots, c_n) , we have

$$\sum_{n_1+n_2=n+1} (-1)^{\sum_{j=1}^{l-1} |c_j|'} M_{n_1}(c_1,\ldots,c_{i-1},M_{n_2}(c_i,\ldots,c_{i+n_2-1}),\ldots,c_n) = 0.$$

Proof. We check the identity on each component of the output. We first show that

$$\sum \left(M_{n_1}^a(\dots, M_{n_2}^a(\dots), \dots) \pm M_{n_1}^a(\dots, M_{n_2}^b(\dots), \dots) \right) = 0.$$

This identify follows from the compactification of popsicle moduli spaces. Namely, a codimension one stratum of the popsicle moduli space corresponds to a term in the above equation. In Figure 16, we illustrated corresponding broken popsicles for the case |F| = 4. Even for broken popsicles, we can consider a sequence of hyperbolic geodesics connecting z_0 and z_{i_j} . In the figure, dotted lines are such geodesics that do not contain a sprinkle.




FIGURE 16. A_{∞} -identity with *a*-output

Next we show that

$$\sum \left(M_{n_1}^b(\ldots, M_{n_2}^a(\ldots), \ldots) \pm M_{n_1}^b(\ldots, M_{n_2}^b(\ldots), \ldots) \right) = 0.$$

This identify follows from the compactification of popsicle moduli spaces for \hat{F}^{j} for all j. In Figure 17, we illustrated corresponding broken popsicles for the case |F| = 4 and j = 1. We also expressed the forgotten geodesic for M^{b} -operation as dotted lines.




FIGURE 17. A_{∞} -identity with *b*-output

5.5. **Example:** M_2 -operation. Let us examine the following Leibniz rule for the input $(a, \epsilon b)$.

(5.10)
$$M_1(M_2(a,\epsilon b)) + M_2(M_1(a),\epsilon b) + (-1)^{|a|'} M_2(a,M_1(\epsilon b)) = 0.$$

For simplicity, we will omit the signs from the formulas. From the definition

$$M_2(a,\epsilon b) = \mathbf{P}_{2,\{2\}}^{\Gamma}(a,b) + \epsilon m_2(a,b)$$
$$M_1(\epsilon b) = \mathbf{P}_{1,\{1\}}^{\Gamma}(b) + \epsilon m_1(b)$$

We have

$$M_{1}(M_{2}(a,\epsilon b)) = M_{1}(\mathbf{P}_{2,\{2\}}^{\Gamma}(a,b) + \epsilon m_{2}(a,b)) = m_{1}(\mathbf{P}_{2,\{2\}}^{\Gamma}(a,b)) + (\mathbf{P}_{1,\{1\}}^{\Gamma}(m_{2}(a,b)) + \epsilon m_{1}(m_{2}(a,b)))$$

$$M_{2}(M_{1}(a),\epsilon b) = M_{2}(m_{1}(a),\epsilon b) = \mathbf{P}_{2,\{2\}}^{\Gamma}(m_{1}(a),b) + \epsilon m_{2}(m_{1}(a),b)$$

$$M_{2}(a,M_{1}(\epsilon b)) = M_{2}(a,\mathbf{P}_{1,\{1\}}^{\Gamma}(b) + \epsilon m_{1}(b)) = m_{2}(a,\mathbf{P}_{1,\{1\}}^{\Gamma}(b)) + \mathbf{P}_{2,\{2\}}^{\Gamma}(a,m_{1}(b)) + \epsilon m_{2}(a,m_{1}(b))$$

If we collect the terms with ϵ in (5.10), we obtain the original A_{∞} -identity

$$\epsilon (m_1(m_2(a,b)) + m_2(m_1(a),b) + (-1)^{|a|'}m_2(a,m_1(b))) = 0$$

Collecting the terms without ϵ in (5.10), we get the following(up to sign)

$$\mathbf{P}_{2,\{2\}}^{\Gamma}(m_1(a),b) + \mathbf{P}_{2,\{2\}}^{\Gamma}(a,m_1(b)) + m_2(a,\mathbf{P}_{1,\{1\}}^{\Gamma}(b)) + \mathbf{P}_{1,\{1\}}^{\Gamma}(m_2(a,b)) + m_1\mathbf{P}_{2,\{2\}}^{\Gamma}(a,b) = 0.$$

These terms correspond to the codimension one degenerations (given by disc bubblings) in Figure 18. Here dotted lines just indicate paths to the 0-th vertex, and do not give any restriction to the domain. Hence one may remove dotted lines to find the corresponding A_{∞} -operations.

....

Let us examine Leibniz rule for the input $(\epsilon b_1, \epsilon b_2)$. Namely, we want to verify

(5.11)
$$M_1(M_2(\epsilon b_1, \epsilon b_2)) + M_2(M_1(\epsilon b_1), \epsilon b_2) + (-1)^{|b_1|} M_2(\epsilon b_1, M_1(\epsilon b_2)) = 0.$$




FIGURE 18. Leibniz rule for the inputs $(a, \epsilon b)$

We have

$$\begin{split} M_{1}(M_{2}(\epsilon b_{1},\epsilon b_{2})) &= M_{1}(\mathbf{P}_{2,\{1,2\}}^{\Gamma}(b_{1},b_{2}) + \epsilon \mathbf{P}_{2,\{2\}}^{\Gamma}(b_{1},b_{2}) + \epsilon \mathbf{P}_{2,\{1\}}^{\Gamma}(b_{1},b_{2})) \\ &= m_{1}(\mathbf{P}_{2,\{1,2\}}^{\Gamma}(b_{1},b_{2})) + \mathbf{P}_{1,\{1\}}^{\Gamma}(\mathbf{P}_{2,\{2\}}^{\Gamma}(b_{1},b_{2}) + \mathbf{P}_{2,\{1\}}^{\Gamma}(b_{1},b_{2})) + \epsilon m_{1}(\mathbf{P}_{2,\{2\}}^{\Gamma}(b_{1},b_{2}) + \mathbf{P}_{2,\{1\}}^{\Gamma}(b_{1},b_{2})) \\ M_{2}(M_{1}(\epsilon b_{1}),\epsilon b_{2}) &= M_{2}(\mathbf{P}_{1,\{1\}}^{\Gamma}(b_{1}) + \epsilon m_{1}(b_{1}),\epsilon b_{2}) \\ &= \mathbf{P}_{2,\{2\}}^{\Gamma}(\mathbf{P}_{1,\{1\}}^{\Gamma}(b_{1}),b_{2}) + \epsilon m_{2}(\mathbf{P}_{1,\{1\}}^{\Gamma}(b_{1}),b_{2}) \\ &+ \mathbf{P}_{2,\{1,2\}}^{\Gamma}(m_{1}(b_{1}),b_{2}) + \epsilon \mathbf{P}_{2,\{2\}}^{\Gamma}(m_{1}(b_{1}),b_{2}) + \epsilon \mathbf{P}_{2,\{1\}}^{\Gamma}(m_{1}(b_{1}),b_{2}) \\ M_{2}(\epsilon b_{1},M_{1}(\epsilon b_{2})) &= M_{2}(\epsilon b_{1},\mathbf{P}_{1,\{1\}}^{\Gamma}(b_{2}) + \epsilon m_{1}(b_{2})) \\ &= \mathbf{P}_{2,\{1,2\}}^{\Gamma}(b_{1},\mathbf{P}_{1,\{1\}}^{\Gamma}(b_{2})) + \epsilon m_{2}(b_{1},\mathbf{P}_{1,\{1\}}^{\Gamma}(b_{2})) \\ &+ \mathbf{P}_{2,\{1,2\}}^{\Gamma}(b_{1},m_{1}(b_{2})) + \epsilon \mathbf{P}_{2,\{2\}}^{\Gamma}(b_{1},m_{1}(b_{2})) + \epsilon \mathbf{P}_{2,\{1\}}^{\Gamma}(b_{1},m_{1}(b_{2})) \\ &+ \mathbf{P}_{2,\{1,2\}}^{\Gamma}(b_{1},m_{1}(b_{2})) + \epsilon \mathbf{P}_{2,\{2\}}^{\Gamma}(b_{1},m_{1}(b_{2})) + \epsilon \mathbf{P}_{2,\{1\}}^{\Gamma}(b_{1},m_{1}(b_{2})) \\ &+ \mathbf{P}_{2,\{1,2\}}^{\Gamma}(b_{1},m_{1}(b_{2})) + \epsilon \mathbf{P}_{2,\{2\}}^{\Gamma}(b_{1},m_{1}(b_{2})) + \epsilon \mathbf{P}_{2,\{1\}}^{\Gamma}(b_{1},m_{1}(b_{2})) \\ &+ \mathbf{P}_{2,\{1,2\}}^{\Gamma}(b_{1},m_{1}(b_{2})) + \epsilon \mathbf{P}_{2,\{2\}}^{\Gamma}(b_{1},m_{1}(b_{2})) + \epsilon \mathbf{P}_{2,\{2\}}^{\Gamma}(b_{1},m_{1}(b_{2})) \\ &+ \mathbf{P}_{2,\{1,2\}}^{\Gamma}(b_{1},m_{1}(b_{2})) + \epsilon \mathbf{P}_{2,\{2\}}^{\Gamma}(b_{1},m_{1}(b_{2})) \\ &+ \mathbf{P}_{2,\{1,2\}}^{\Gamma}(b_{1},m_{1}(b_{2})) + \epsilon \mathbf{P}_{2,\{2\}}^{\Gamma}(b_{1},m_{1}(b_{2})) \\ &+ \mathbf{P}_{2,\{1,2\}}^{\Gamma}(b_{1},m_{1}(b_{2})) + \epsilon \mathbf{P}_{2,\{2\}}^{\Gamma}(b_{1},m_{1}(b_{2})) \\ &+ \mathbf{P}_{2,\{2\}}^{\Gamma}(b_{1},m_{1}(b_{2})) + \epsilon \mathbf{P}_{2,\{2\}}^{\Gamma}(b_{1},m_{1}(b_{2})) \\ &+ \mathbf{P}_{2,\{2\}}^{\Gamma}(b_{1}$$

The following figure 19 describes the terms without ϵ in the above (in the same order). It is not




FIGURE 19. Leibniz rule for the inputs ($\epsilon b_1, \epsilon b_2$)

hard to see that these arise from codimension one boundary strata of $\overline{P}_{2,\{1,2\}}$. The terms with ϵ are similar.

6. MIRROR COUNTERPART: HYPERSURFACE RESTRICTION

A mirror construction of last section for algebraic geometry is just a categorical reformulation of the restriction to a hypersurface.

6.1. Restriction to a hypersurface in D^bCoh . Let *S* be an algebra. Choose an element

$$g \in Z(S) \cong HH^0(S,S)$$

The DG-bimodule $S \xrightarrow{g} S$ is quasi-isomorphic to an ideal quotient S/(g) which carries a natural algebra structure. One can directly construct DG algebra structure on the bimodule itself:

Definition 6.1. Define a DG algebra

$$\mathscr{B} := S[\epsilon] / \begin{pmatrix} \epsilon^2 = 0 \\ d\epsilon = g \end{pmatrix},$$

Here deg $\epsilon = -1$ *and the differential d on S is set to be zero.*

Further assume that *S* is commutative. Consider an affine variety X = Spec(S) and a hypersurface Y = V(g) with an inclusion $i : Y \hookrightarrow X$. We have the following elementary lemma whose proof is omitted.

Lemma 6.2. We have an isomorphism $\mathscr{B} \simeq i_* \mathscr{O}_Y$. Moreover, we have the following.

- (1) A sheaf \mathscr{F} on a hypersurface Y corresponds to an \mathscr{B} -module object. It is a pair $(i_*\mathscr{F}, h_{\mathscr{F}})$ where $i_*\mathscr{F}$ is a pushforward of \mathscr{F} equipped with a homotopy $h_{\mathscr{F}}$ between the zero map and a multiplication of g. It is an action of $\epsilon \in \mathscr{B}$.
- (2) Moreover,

$$\operatorname{Hom}_{Y}(\mathscr{F}_{1},\mathscr{F}_{2}) \simeq \operatorname{Hom}_{\mathscr{B}}((i_{*}\mathscr{F}_{1},h_{\mathscr{F}_{1}}),(i_{*}\mathscr{F}_{2},h_{\mathscr{F}_{2}}))$$

For the sheaf \mathcal{O}_Y on *Y*, its pushfoward $i_*\mathcal{O}_Y$ has a simple free resolution.

$$0 \longrightarrow \mathscr{O}_X \stackrel{g}{\longrightarrow} \mathscr{O}_X \longrightarrow i_* \mathscr{O}_Y \longrightarrow 0$$

An action of degree -1 element ϵ , or a homotopy *h*, is given as follows.

A category of coherent sheaves on *Y* is described as \mathscr{B} -modules of *X*.

Theorem 6.3. *Let* $Y \subset X$ *as before. Then*

$$DCoh(Y) \simeq \mathscr{B} - mod(DCoh(X))$$

Proof. A concise categorical proof using Lurie's Barr-Beck theorem can be found in Corollary 3.3.1 in [Pre11]. We present an elementary proof to illustrate the idea. Since everything is affine, it is enough to consider a structure sheaf $\mathcal{O}_Y \in DCoh(Y)$. Computation shows that the morphism complex is

Therefore, $\operatorname{Hom}_{\mathscr{B}}^{\bullet}((i_*\mathscr{O}_Y, h), (i_*\mathscr{O}_Y, h))$ is isomorphic to

$$H^{\bullet}(\operatorname{Hom}_{X}(\mathcal{O}_{X},\mathcal{O}_{X}) \xrightarrow{g} \operatorname{Hom}_{X}(\mathcal{O}_{X},\mathcal{O}_{X})) \simeq \operatorname{Hom}_{Y}^{\bullet}(\mathcal{O}_{Y},\mathcal{O}_{Y}).$$

Intuitively, objects of $\mathscr{B} - mod(DCoh(X))$ are cones

$$\left(\mathscr{F}[1] \xrightarrow{g} \mathscr{F} \right), \mathscr{F} \in DCoh(X).$$

It is quasi-isomorphic to a quotient $\mathcal{F}/(g)$ and the space of morphisms is a half of the original one. It consists of

$$\begin{pmatrix} a & 0 \\ b & a \end{pmatrix} \in \operatorname{Hom}_{X}^{\bullet} \left(\ \mathscr{F}_{1}[1] \xrightarrow{g} \mathscr{F}_{1} \ , \ \mathscr{F}_{2}[1] \xrightarrow{g} \mathscr{F}_{2} \ \right).$$

This is closed under DG operations whereas A_{∞} -analogue is not (see the paragraph after the Definition 4.3).

6.2. **Restriction to a graph hypersurface in Matrix factorizations.** Let us explain an analogous construction for matrix factorizations. First, let us recall its definition for readers' convenience.

Let *R* be a polynomial algebra over the algebraically closed field *k* of characteristic 0.

Definition 6.4. For $W \in R$, a DG category of matrix factorization of W, denoted by MF(W) consists of the following:

- (1) its objects are matrix factorization (P, δ_P) of W. P is $\mathbb{Z}/2$ -graded free R-module and δ_P is an odd degree endomorphism such that $\delta_P^2 = W \cdot id$.
- (2) $\operatorname{Hom}^{\bullet}_{\operatorname{MF}(W)}((P, \delta_P), (P', \delta_{P'})) := (\operatorname{Hom}^{\bullet}_R(P, \dot{P'}), d)$ with usual composition \circ . A differential d on morphisms is defined as

$$d(\phi) = \delta_{P'} \circ \phi - (-1)^{\deg(\phi)} \phi \circ \delta_P.$$

We are interested in the following situation. Consider a polynomial of the form $U = U_1(x_1, ..., x_{n-1}) + x_n \cdot U_2(x_1, ..., x_{n-1})$. We consider a graph of some polynomial *f*

$$\mathbb{A}^{n-1} \to \mathbb{A}^n$$
$$(x_1, \dots, x_{n-1}) \mapsto (x_1, \dots, x_{n-1}, f)$$

and a pull-back

$$V(x_1, \dots, x_{n-1}) = U(x_1, \dots, x_{n-1}, f) = U_1 + f \cdot U_2$$

along the graph. We explain how to obtain a similar relation between MF(U) and MF(V). We start by collecting functorial properties between two matrix factorization categories, which we refer to [Orl09] and [EP15].

Let $X = \{U = 0\} \subset \mathbb{C}^n$ and $Y = \{V = 0\} \subset \mathbb{C}^{n-1}$. We view *Y* as a hypersurface $\{x_n = f\} \subset X$. A closed embedding $Y \hookrightarrow X$ is proper and has a finite tor-dimension. A usual adjoint pair of functors (i^*, i_*) extends to categories of singularities.

$$i^*: D^b_{sg}(X) \longleftrightarrow D^b_{sg}(Y): i_*$$

On the other hand, there are Orlov's equivalences

$$\operatorname{MF}(U) \simeq \overline{D_{sg}^b(X)}, \ \operatorname{MF}(V) \simeq \overline{D_{sg}^b(Y)}$$

Here, $\overline{\mathscr{C}}$ denotes Karoubi completion of a category \mathscr{C} . This functor sends

$$M = (M^{odd} \xrightarrow{\phi_{10}} M^{even}) \mapsto \operatorname{coker}(\phi_{10})$$

We have an induced pair

$$i^*: \mathrm{MF}(U) \longleftrightarrow \mathrm{MF}(V): i_*$$

Proposition 6.5. Let

$$\begin{split} M &= (M^{odd} \xrightarrow{\phi_{10}} M^{even}) \in \mathrm{MF}(U), \\ N &= (N^{odd} \xrightarrow{\psi_{10}} N^{even}) \in \mathrm{MF}(V). \end{split}$$

Then

(1)
$$(i^*, i_*)$$
 is an adjoint pair.
(2) $i^*M \simeq M|_{x_n=f} \in MF(V)$.
(3) $i_*N \simeq N \otimes \begin{pmatrix} & & \\ & & & \\ & & \\ & & \\ & & & \\ & & \\ &$

Proof. The first proposition is proven in more general setup. See [EP15] Section 2.1. Second proposition follows from the fact that cokernel commutes with tensor product.

$$\operatorname{coker}(\phi_{10}) \otimes_{\mathbb{C}[x_1,\ldots,x_n]} \mathbb{C}[x_1,\ldots,x_{n-1}] \simeq \operatorname{coker}(\phi_{10}|_{x_n=f}).$$

To prove a third proposition, we should specify Fourier-Mukai kernel of a push-forward functor. Write

$$V(x_1,...,x_{n-1}) - V(y_1,...,y_{n-1}) = \sum_{i=1}^{n-1} (x_i - y_i) \cdot V_i$$

Define a Koszul-type matrix factorization Γ of $V(\vec{x}) - U(\vec{y}) = V(\vec{x}) - (V(\vec{y}) + (y_n - f)U_2(\vec{y}))$ as

$$\Gamma := \left(\Lambda^{\bullet} \langle e_1, \dots, e_n \rangle, \left(\sum_{i=1}^{n-1} (x_i - y_i) i_{e_i} + (y_n - f) i_{e_n} + \sum_{i=1}^{n-1} V_i (\cdot \wedge e_i) + U_2 (\cdot \wedge V)\right)\right).$$

Under Orlov's equivalence Γ corresponds to a stabilization of a graph $\Gamma_{Y \to X}$. Therefore a Fourier-Mukai functor associated to Γ is a pushforward functor. Notice that

$$-\otimes\Gamma\simeq-\otimes\Delta_V\otimes\left(\begin{array}{c}x_n-f\\ \mathbb{C}[x_1,\ldots,x_n]\\\mathbb{C}[x_1,\ldots,x_n]\\\mathbb{V}_2\end{array}\right)$$

where Δ_V is a stabilized diagonal of *V*. This proves the third proposition.

For the fourth proposition, observe that $i_* \circ i^* M$ goes to

$$\operatorname{coker}\left(\phi_{10}|_{x_n=f}: M^{odd}|_{x_n=f} \to M^{even}|_{x_n=f}\right)$$

under Orlov's equivalence. It is easy to see that the periodic tail of the following double complex realizes the matrix factorization associated to that module.

$$\cdots \xrightarrow{\phi_{10}} M^{even} \xrightarrow{\phi_{01}} M^{odd} \xrightarrow{\phi_{10}} M^{even} \\ \downarrow x_n - f \qquad \downarrow x_n - f \qquad \downarrow x_n - f \\ \cdots \xrightarrow{\phi_{10}} M^{even} \xrightarrow{\phi_{01}} M^{odd} \xrightarrow{\phi_{10}} M^{even}$$

This is equal to $\text{Cone}((x_n - f) : M[1] \rightarrow M)$.

The fourth of Proposition 6.5 means

$$(i_* \circ i^*)M \simeq \left(\begin{array}{c} M[1] \xrightarrow{x_n - f} M \end{array} \right) = M[\epsilon] / \left(\begin{array}{c} \epsilon^2 = 0 \\ d\epsilon = x_n - f \end{array} \right).$$

This is a perfect analogy of \mathcal{B} -module objects we considered in the last subsection.

Definition 6.6. Define a DG category $MF(U)|_{x_n-f}$ as follows:

- *its objects consist of the matrix factorizations* $(i_* \circ i^*)M$ *for* $M \in MF(U)$,
- *its morphisms* Hom_{MF(U)|xn-f} ($(i_* \circ i^*)M_1, (i_* \circ i^*)M_2$) consist of

 $a + \epsilon b \in \operatorname{Hom}_{\operatorname{MF}(U)}\left((i_* \circ i^*)M_1, (i_* \circ i^*)M_2\right) = \operatorname{Hom}(M_1[\epsilon], M_2[\epsilon])$

Differentials and compositions are induced from MF(U).

The next proposition explains the relation between MF of the restriction $V = U|_{x_n-f=0}$ and the DG category MF(*U*)|_{*x_n-f*.}

Corollary 6.7. Let

$$i_* \circ i^* : \mathrm{MF}(U) \to \mathrm{MF}(U)|_{x_n - f}$$

be a natural inclusion and

$$c: \mathrm{MF}(U)|_{x_n-f} \to \mathrm{MF}(V)$$

be a DG functor sending $(i_* \circ i^*)M$ to the matrix factorization $M|_{x_n-f=0}$.

(1) The following diagram commutes.

$$\begin{array}{cccc}
\operatorname{MF}(U) & \xrightarrow{i_* \circ i^*} \operatorname{MF}(U)|_{x_n - f} \\
& & \downarrow^c \\
& & & \downarrow^c \\
& & & \operatorname{MF}(V)
\end{array}$$

(2) A category $MF(U)|_{x_n-f}$ fits into a diagram of distinguished triangle of DG bimodules:

$$MF(U) \xrightarrow{x_n-f} MF(U) \xrightarrow{i_* \circ i^*} MF(U)|_{x_n-f} \longrightarrow$$

(3) *c* is fully faithful. It is a quasi-equivalence whenever $i^* : MF(U) \to MF(V)$ is essentially surjective.

Proof. The first and the second proposition follows directly from the definition. For the third one, observe

$$i_* \circ i^* M \simeq M[\epsilon] \Big/ \left(\begin{array}{c} \epsilon^2 = 0 \\ d\epsilon = x_n - f \end{array} \right).$$

Therefore, we have

$$\operatorname{Hom}_{\mathrm{MF}(U)|_{x_n-f}}\left((i_* \circ i^*)M_1, (i_* \circ i^*)M_2\right) \simeq \operatorname{Hom}_{\mathbb{C}[x_1, \dots, x_n, \epsilon]/(d\epsilon = x_n - f)}(M_1[\epsilon], M_2[\epsilon])$$
$$\simeq \operatorname{Hom}_{\mathrm{MF}(U)}(M_1, M_2[\epsilon]) \Big/ \left(\begin{array}{c} \epsilon^2 = 0\\ d\epsilon = x_n - f \end{array}\right)$$
$$\simeq \operatorname{Hom}_{\mathrm{MF}(U)}(M_1, (i_* \circ i^*)M_2)$$
$$\simeq \operatorname{Hom}_{\mathrm{MF}(V)}(i^*M_1, i^*M_2).$$

If i^* is essentially surjective, we have $i_*N \simeq \left(\begin{array}{c} M[1] \xrightarrow{x_n-f} M \end{array}\right)$ for some $M \in MF(U)$. This implies *c* is also essentially surjective.

In this way, we can regard $MF(U)|_{x_n-f}$ as another DG-model representing MF(V).

7. A_{∞} -category for a weighted homogeneous polynomial with a symmetry group

In this section, we will generalize the construction of Section 5 and define a new A_{∞} -category $\mathscr{F}(W, G_W)$ for a weighted homogeneous polynomial W paired with its group G_W of diagonal symmetries. From the monodromy transformation of W and its flow induced on the boundary of Milnor fiber M_W , we define a Reeb orbit Γ_W in the quotient orbifold $[M_W/G_W]$ and use J-holomorphic popsicles with Γ_W -insertions to define such an A_{∞} -category.

7.1. Weighted homogeneous polynomials and variation operator.

Definition 7.1. We say a weighted homogeneous polynomial W is

- (1) $\log Fano \ if \sum_{k=1}^{n} w_i h > 0$ (2) $\log Calabi-Yau \ if \sum_{k=1}^{n} w_i h = 0$ (3) $\log general \ if \sum_{k=1}^{n} w_i h < 0$

Assume that *W* has an isolated singularity at the origin. Set $V_t = V_t(W) = \{z \in \mathbb{C}^n \mid W(z) = t\}$. V_0 is an hypersurface of isolated singularity at 0 and V_t ($t \neq 0$) is non-singular. Milnor fiber M_W is defined to be $V_1(W)$. For the well-known Milnor fibration

$$\frac{W}{|W|}: S_{\epsilon}^{2n-1} \setminus K \to S^{1}$$

with $K = (S_{\varepsilon}^{2n-1} \cap V_0)$, its fiber is diffeomorphic to M_W . Geometric monodromy $\Phi_W : \mathbb{C}^n \to \mathbb{C}^n$ is defined by

(7.1)
$$\Phi_W(x_1, \dots, x_m) = (e^{2\pi i w_1/h} x_1, \dots, e^{2\pi i w_m/h} x_m)$$

which restricts to M_W . It is known that $S_{\epsilon}^{2n-1} \setminus K$ is diffeomorphic to the manifold obtained by identifying two ends of $M_W \times [0, 1]$ by Φ_W (see [Mil68] Lemma 9.4).

It is another famous theorem of Milnor that the homotopy type of M_W is a bouquet of (n-1)spheres. One may define the notion of vanishing skeleton, but it is usually very singular (for Brieskorn-Pham singularities, it is called *Pham's spine*). If we perturb W to a complex Morse function W', then we may obtain vanishing spheres in the Milnor fiber of W', and Fukaya-Seidel directed A_{∞} -category of W is defined on these objects. But since G_W -symmetry is broken after the perturbation, it is not known how to work equivariantly within Fukaya-Seidel category framework.

On the other hand, it is well-known that non-compact Lagrangians are related to vanishing cycles via the variation operator. Consider monodromy homomorphism (from a parallel transport fixing the boundary)

(7.2)
$$h_*: H_*(\overline{M}_W) \to H_*(\overline{M}_W), \ h_*: H_*(\overline{M}_W, \partial \overline{M}_W) \to H_*(\overline{M}_W, \partial \overline{M}_W).$$

Definition 7.2. A variation operator (around the origin in \mathbb{C})

(7.3)
$$\operatorname{var}: H_{n-1}(\overline{M}_W, \partial \overline{M}_W) \to H_*(\overline{M}_W).$$

is defined by sending $[c] \mapsto (h_* - id)([c])$.

This map is known to be an isomorphism. We will find a symplectic categorical analogue of this variation operator for weighted homogenous polynomials in the next section. We define a distinguished Reeb orbit Γ_W from the geometric monodromy (7.1). The analogue of monodromy homomorphism (7.2) will be given by the quantum cap action by Γ_W

$$\cap \Gamma_W : \mathscr{WF}(L,L) \to \mathscr{WF}(L,L).$$

Then, the analogue of the variation operator (7.3) will be the new Fukaya category \mathscr{C}_{Γ_W} , with a distinguished triangle of A_{∞} -bimodules

$$\mathscr{WF}(M) \xrightarrow{\cap \Gamma_{W}} \mathscr{WF}(M) \longrightarrow \mathscr{C}_{\Gamma} \longrightarrow$$

7.2. **Diagonal symmetry group of weighted homogeneous polynomials.** We collect some facts about weighted homogeneous polynomials and their diagonal symmetry group from [FJR13] and a reference therein.

Let *W* be a weighted homogeneous polynomial of weight $(w_1, \ldots, w_n; h)$.

Definition 7.3. The maximal diagonal symmetry group G_W of W is defined as

$$G_W := \left\{ \vec{\lambda} \in (\mathbb{C}^*)^n \mid W(\lambda_1 \cdot x_1, \dots, \lambda_n \cdot x_n) = W(x_1, \dots, x_n) \right\}.$$

Note that geometric monodromy (7.1) is an element of G_W . A class of polynomials that we are interested in are those with finite diagonal symmetries.

Definition 7.4 ([FJR13] Definition 2.1.5). *A weighted homogeneous polynomial W is called nondegenerate if*

- (1) W contains no monomial of the form $x_i x_j$ for some $i \neq j$;
- (2) W is an isolated singularity at the origin.

If *W* is non-degenerate, the number of monomials of *W* must be greater than or equal to the number of variables. The first finiteness condition that we need is the following.

Proposition 7.5 ([HK12], [FJR13]). *Let W be a non-degenerate weighted homogeneous polynomial. Then*

- (1) The weight $(w_1, ..., w_n; h)$ of W is bounded by $\frac{w_i}{h} \le \frac{1}{2}$ and it is unique.
- (2) The maximal diagonal symmetry group G_W is a finite abelian group.

For a subgroup $K \subset G_W$, let $\mathbb{C}^{N_K} := (\mathbb{C}^n)^K$ be the set of fixed points of K. Notice that they are always a coordinate plane. Let W_K be a restriction of W on \mathbb{C}^{N_K} . The second finiteness condition that we will use is the following.

Proposition 7.6 ([FJR13] Lemma 2.10). *Let W be a non-degenerate weighted homogeneous polynomial. Then*

- (1) W_K is a non-degenerate weighted homogeneous polynomial on \mathbb{C}^{N_K} for $\forall K < G_W$.
- (2) G_{W_K} is canonically embedded inside G_W .

An orbifold strata $\{Fix(K)\}_{K < G_W}$ on \mathbb{C}^n is compatible with restrictions. This will enable us to do inductive arguments on the number of variables later on. A pair (W, G) of non-degenerate homogeneous polynomial W and a subgroup $G < G_W$ of diagonal symmetries is called a *orbifold Landau-Ginzburg model*.

7.3. **Orbifold wrapped Fukaya category.** The starting point of our construction is an orbifold wrapped Fukaya category of $[M_W/G_W]$. We follow [Sei15] closely.

We consider the collection \mathcal{W} of Lagrangians $L \in [M_W/G_W]$ with the following properties.

- It is given by the family of embedded Lagrangians $\{L_i\}_{i \in I}$ and an action of G_W on the index set *I* such that $g(L_i) = L_{g \cdot i}$.
- *L* lies away from the singular locus.
- It is either compact or conical at the end.

- It is equipped with an orientation and a spin structure (compatible with G_W -action).
- G_W -orbits intersect transversally to each other without triple intersections.

Definition 7.7. $A\mathbb{Z}/2$ -graded orbifold wrapped Fukaya category $\mathcal{WF}([M_W/G_W])$ consists of;

- (1) a set of objects W;
- (2) For two such Lagrangians $L_0, L_1 \in W$, its morphism space is defined by

$$CW^{\bullet}(L_0, L_1) := \left(\bigoplus_{g, h \in G_W} CW^{\bullet}\left(g \cdot \widetilde{L_0}, h \cdot \widetilde{L_1}\right)\right)^{G_W}$$

where \tilde{L}_i is a lift of L_i and we take G_W -invariant part. (3) A_∞ -operations are induced from the m_k -operation of $\mathcal{WF}(M_W)$.

Following Seidel [Sei15], we can make the A_{∞} -structure G_W -equivariant. We omit the details and refer readers to [Sei15]. In addition to the usual $\mathbb{Z}/2$ -grading, we can also define $\frac{1}{|G_W|}\mathbb{Z}$ graded category. Consider a canonical holomorphic volume form $\Omega = \operatorname{Res}_W dz_1 \wedge \cdots \wedge dz_n$ on a fiber of $W : \mathbb{C}^n \to C$. It is not difficult to show that we can define the Milnor fiber M_W so that Ω induces a non-vanishing *n*-form on Ω_{M_W} on M_W . Regarding a finite group action of G_W on M_W , Ω_{M_W} does not descend to $[M_W/G_W]$ because $G_W \not\subset SL_n$ (hence \mathbb{Z} -grading does not exist). But its tensor power $\Omega_{M_W}^{\otimes |G_W|}$ does descend. This allows us to define $\frac{1}{|G_W|}\mathbb{Z}$ -graded version of wrapped Fukaya category following [Sei11], [AAE⁺13]. We can also forget the fractional grading and simply work with $\mathbb{Z}/2$ -grading as well.

A holomorphic disc $u: S \to M_W$ defining A_∞ -structure can be considered as a smooth holomorphic disc $\overline{u}: S \to [M_W/G_W]$ ramified at orbifold points accordingly. Conversely, because S has a vanishing orbifold fundamental group, all smooth holomorphic discs lifts to M_W in a G_W -equivariant manner. Therefore, Fukaya category of $[M_W/G_W]$ is the same as G_W -equivariant Fukaya category of M_W .

7.4. **Monodromy flow and orbits.** Fix a non-degenerate weighted homogeneous polynomial *W* of weight $(w_1, \ldots, w_n; h)$ once and for all. Choose a slightly rescaled symplectic form ω and a Liouville form λ on \mathbb{C}^n (see [KvK16] for the relation to the standard one) as

$$\omega = \sum_{k} \frac{1}{2\pi i w_{k}} dz_{k} \wedge d\overline{z}_{k}, \quad \lambda = \sum_{k} \frac{i}{4\pi w_{k}} (z_{k} d\overline{z}_{k} - \overline{z}_{k} dz_{k}).$$

Definition 7.8. The monodromy flow is the Hamiltonian flow

$$\Phi_W(s)(x_1,\ldots,x_n):=(e^{\frac{2\pi iw_1s}{h}}x_1,\ldots,e^{\frac{2\pi iw_ns}{h}}x_n)$$

of a quadratic Hamiltonian $H := \frac{1}{2} \sum_{i=1}^{n} |x_i|^2$. The monodromy transformation $\Phi_W = \Phi_W(1)$ is a time-1 flow

$$x_i \mapsto e^{\frac{2\pi i w_i}{h}} x_i.$$

Geometrically, a Hamiltonian action of *H* is a lifting of a rotation action on the base of a fibration $W : \mathbb{C}^n \to \mathbb{C}$. We have $W \circ \Phi_W(s) = e^{2\pi i s} W$, which means that the flow of *H* acts as a circle action of an S^1 family of Milnor fiber $W = e^{2\pi i s}$. Set s = 1 then we get a desired automorphism.

The monodromy flow restricts to a singular fiber $W^{-1}(0)$ and a *link* of W

$$L_{W,\delta} := W^{-1}(0) \cap S_{\delta}^{2n-1}$$

for small $\delta > 0$. The monodromy flow $\Phi_W(s)$ becomes a Reeb flow \mathscr{R} on $L_{W,\delta}$, where the contact one form is given by a restriction of λ . Starting from any point $x \in L_{W,\delta}$, we get a Reeb chords

$$\gamma: [0,1] \to L_{W,\delta}, \ \gamma(0) = x, \ \gamma(1) = \Phi_W(x)$$

Notice that Φ_W always gives an element *J* of a maximal symmetry group G_W . Reeb chords become **orbits** of a quotient $(L_{W,\delta}/G_W)$. A space of time-1 Reeb orbits of the quotient is a total space $L_{W,\delta}/G_W$.

The link $L_{W,\delta}$ can be symplectically identified with $W^{-1}(1) \cap S_{\delta}^{2n-1}$ in the following way (see [Sei00]). Choose a cutoff function ψ with $\psi(t^2) = 1$ for $t^2 \leq \frac{\delta}{3}$ and $\psi(t^2) = 0$ for $t^2 \geq \frac{2\delta}{3}$. Define

$$F := \{ x \in B_{\delta}^{2n} \mid W(x) = \psi(|x|^2) \}.$$

It was shown in [Sei00] that *F* is symplectic manifold with $\partial F = L_{W,\delta}$. Moreover *F* is diffeomorphic to $W^{-1}(1) \cap B^{2n}$ by a smooth cobordism

$$Bor_s := \{ x \in B_{\delta}^{2n} \mid W(x) = s\psi(|x|^2) + (1-s) \}.$$

Shrinking δ if it is necessary, we see that F and M_W contain $W^{-1}(1) \cap B^{2n}_{\delta/3}$ as a Liouville submanifold and Liouville isotopy compresses both F and M_W to $Int(W^{-1}(1) \cap B^{2n}_{\delta/3})$. Since ψ only depends on $|x|^2$, the whole construction is compatible with the action of G_W . Therefore $L_{W,\delta}/G_W$ serves as a model for the contact type boundary of M_W/G_W .

7.5. **Main theorem.** The reader would realize that we are in a similar situation to the case of Assumption 5.1. Reeb flow \mathscr{R}_t is an S^1 -action on L_W/G_W , and therefore L_W/G_W provides a Morse-Bott component of Hamiltonian orbits. The orbit that corresponds to its fundamental class will denote by Γ_W . We state our main theorem.

Theorem 7.9. Let W be a nondegenerate weighted homogeneous polynomial $\sum_{i=1}^{n} w_i - h \ge 0$. With the monodromy orbit Γ_W of $[M_W/G_W]$, we define a new $\frac{\mathbb{Z}}{|G_W|}$ -graded A_∞ -category \mathscr{C}_{Γ_W} such that

- (1) \mathscr{C}_{Γ_W} has the same set of objects as the $\frac{\mathbb{Z}}{|G_W|}$ -graded wrapped Fukaya category $\mathscr{W}\mathscr{F}([M_W/G_W])$,
- (2) for two objects L_1, L_2 , its morphisms are given by

$$\operatorname{Hom}_{\mathscr{C}_{\Gamma}}(L_1, L_2) = CW(L_1, L_2) \oplus CW(L_1, L_2)\epsilon$$

where $CW(L_1, L_2)$ is the morphism space for $\mathcal{WF}([M_W/G_W])$ and $\deg \epsilon = -1 + \mu_{RS}(\Gamma_W)$,

- (3) a natural inclusion $\Psi : \mathscr{WF}([M_W/G_W]) \to \mathscr{C}_{\Gamma_W}$ is an A_{∞} -functor,
- (4) regarding the A_{∞} -category \mathscr{C}_{Γ_W} as an A_{∞} -bimodule over $\mathscr{WF}([M_W/G_W])$ (using Ψ), we have a distinguished triangle of A_{∞} -bimodules

$$\mathscr{WF}([M_W/G_W])_{\Delta} \xrightarrow{\cap \Gamma_W} \mathscr{WF}([M_W/G_W])_{\Delta} \longrightarrow \mathscr{C}_{\Gamma} \longrightarrow$$

It turns out that we still need a substantial amount of work to generalize Theorem 5.2 to the current setting, and we will give the proof in the next section.

Let us just mention about the fractional grading here. One can assign factional gradings to Hamiltonian orbits as in the case of wrapped Fukaya category following [Sei11], [AAE⁺13]. First, let us explain the underlying $\mathbb{Z}/2$ -grading. Recall that $\mathbb{Z}/2$ -grading for symplectic cohomology complex is easy to define. Choose a trivialization of the tangent bundle along a hamiltonian orbit, and the parity of the Conley-Zehnder index of the linearized flow gives the desired grading. Note that the change of trivialization does not change the parity of the index. Robbin-Salamon

index of a symplectic loop under any trivialization is even. Thus one find that $\mathbb{Z}/2$ -grading for Γ_W is still even, since under the trivialization of the tangent bundle over a principal orbit, we still have a symplectic loop.

Now, additional fractional grading can be assigned as usual. Recall that we have non-vanishing form $\Omega_{M_W}^{\otimes |G_W|}$ on $[M_W/G_W]$, which can be used to define fractional Conley-Zehnder or Robbin-Salamon indices (see [McL16]). In particular, for Γ_W , Robbin-Salamon index of the orbit with respect to the form $\Omega_{M_W}^{\otimes |G_W|}$ is denoted as $\mu_{RS}^{\Omega^{\otimes |G_W|}}$, and we set

$$\mu_{RS}(\Gamma_W) = \frac{1}{|G_W|} \mu_{RS}^{\Omega^{\otimes |G_W|}}$$

7.6. Fukaya category $\mathscr{F}(W, G)$ for any subgroup $G < G_W$.

Definition 7.10. The $\mathbb{Z}/2$ -graded Fukaya category $\mathscr{F}(W, G_W)$ of a Landau-Ginzburg orbifold (W, G_W) is defined to be the underlying $\mathbb{Z}/2$ -graded A_{∞} -category of \mathscr{C}_{Γ_W} .

For any subgroup $G < G_W$, we define the Fukaya category $\mathscr{F}(W, G)$ using the dual group G_W^T -action on $\mathscr{F}(W, G_W)$.

Let us first explain the algebraic setting. Let \mathscr{C} be an A_{∞} -category with an action by finite abelian group H. By taking a H-invariant part, we may obtain an A_{∞} -category \mathscr{C}^{H} whose object is a H-family of objects $\{h \cdot \tilde{K}\}_{h \in H}$, denoted as K, and its morphisms between K and K' are given by

$$\bigoplus_{h \in H} \operatorname{Hom}_{\mathscr{C}}(K, hK') \cong \left(\bigoplus_{h_1, h_2 \in H} \operatorname{Hom}_{\mathscr{C}}(h_1 \tilde{K}, h_2 \tilde{K}') \right)^H$$

Then the character group $H^* = \text{Hom}(H, \mathbb{C}^*)$ naturally acts on the A_{∞} -category \mathscr{C}^H : we define $\chi \in H^*$ action for an element $X \in \text{Hom}_{\mathscr{C}}(K, hK')$ (resp. its dual formal variable *x*) as

$$\chi \cdot X = \chi(h^{-1})X$$
 (resp. $\chi \cdot x = \chi(h)x$).

In [Sei15], Seidel observed that one can recover \mathscr{C} from \mathscr{C}^H using semi-direct product. By taking a tensor product of $\operatorname{Hom}_{\mathscr{C}^H}(K, K')$ with the group ring $\mathbb{C}[H^*]$, and extending the A_{∞} -operation suitably, we obtain a semi-direct product A_{∞} -category $\mathscr{C}^H \rtimes H^*$, which can be canonically identified with \mathscr{C} . More precisely, the component $\operatorname{Hom}_{\mathscr{C}^H}(K, K') \otimes \chi$ in the morphism space can be identified with the χ -eigenspace of H-action on \mathscr{C} (see [CL20] for an explicit identification). Geometrically, it means that A_{∞} -operations in \mathscr{C}^H actually come from \mathscr{C} . For the case of Fukaya category, this corresponds to the fact that a smooth holomorphic discs in the quotient can be always lifted.

Let us discuss the case of the A_{∞} -category $\mathscr{F}(W, G_W)$. For a *J*-holomorphic disc without Γ_W -insertion,

$$m_k(\chi \cdot w_1, \dots, \chi \cdot w_k) = \chi \cdot m_k(w_1, \dots, w_k)$$

holds because disc has a lift to upstairs and still is a disc. When we walk along the boundary Lagrangians, we come back to the original branch that we start with.

We have additional operations of popsicles which have Γ_W interior insertions. These *J*-holomorphic discs do not have lifts to upstairs, so we may proceed as follows. Denote by $g_W \in G_W$ be the group element for the monodromy of *W* in (7.1). For each sprinkle with Γ_W insertion, we can make a

branch cut in the domain D^2 to the boundary arc between two marked points z_0, z_1 . We can make the cuts disjoint from each other. Then, the resulting disc lifts to the cover, but if we restrict to the arc z_0z_1 , it has discontinuous lift. Namely, z_0z_1 boundary arc has as many g_W jumps as the number of sprinkles. This can be used to show that

$$m_{k,F}^{\Gamma_W}(\chi \cdot w_1, \dots, \chi \cdot w_k) = \chi(g_W^{-|F|}) m_{k,F}^{\Gamma_W}(w_1, \dots, w_k)$$

To accommodate this new factor, we introduce G_W^* -action on the formal parameter ϵ in the Definition 5.12. It is not hard to check that this gives the desired G_W^* -action on \mathscr{C}_{Γ_W} .

Proposition 7.11. We set $\chi \cdot \epsilon = \chi(g_W^{-1})\epsilon$. Then A_{∞} -category \mathscr{C}_{Γ_W} admits G_W^* -action.

For any subgroup $G < G_W$, we define its dual group $G^T = \text{Hom}(G_W/G, \mathbb{C}^*)$ following Berglund-Henningson [BH95]. We are ready to define a Fukaya category of the Landau-Ginzburg orbifold (W, G).

Definition 7.12. We define a Fukaya A_{∞} -category for (W, G) by

$$\mathscr{F}(W,G) := \mathscr{C}_{\Gamma_W} \rtimes G^T.$$

Here, the quotient map $G_W \to G_W/G$ induces the map $G^T \to G_W^*$ and hence a G^T -action on \mathscr{C}_{Γ_W} .

8. Twisted Reeb orbits in the Milnor Fiber quotients and their indices

We give a proof of Theorem 7.9 in this section and the next, generalizing the construction of Section 5. The main scheme is the same, but our orbifold $[M_W/G_W]$ does not satisfy Assumption 5.1, and hence we have the following issues.

- (1) $[M_W/G_W]$ have a nontrivial c_1 .
- (2) The action of Reeb flow \mathcal{R}_t is not free. We need a classification of Morse-Bott components.
- (3) Inequality relating a period and CZ index is required.
- (4) We need to define Γ_W and show that it is closed in a suitable sense.

We will address each of these items, and then Theorem 7.9 should follow as in construction of Section 5.

The first item can be overcome by introducing the fractional grading as we explained. Let us discuss the remaining issues one by one.

8.1. Classifying twisted Reeb orbits. The second issue is that there can be several Morse-Bott components of Hamiltonian orbits in $[M_W/G_W]$. We investigate such orbits and their fractional gradings in this subsection.

In the construction of \mathscr{C}_{Γ_W} , we only use the principal orbit Γ_W . Even though we do not use other Reeb orbits as interior insertions, they might appear in the compactification of popsicles. It is well-known that a perturbed *J*-holomorphic cylinder with bounded energy in the smooth locus of the orbifold can limit to a periodic orbit in the orbifold locus. And the purpose of the study of these non-principal orbits is to rule out their appearances in the compactification of our popsicles.

A loop in the orbifold $[M_W/G_W]$ is given by a G_W -twisted loop in M_W .

Definition 8.1. For an element $g \in G_W$, $a \neq g$ -twisted loop (g, γ) of period l is a path $\gamma : [0, l] \rightarrow M_W$ such that $g \cdot \gamma(l) = \gamma(0)$. We have a G_W -action on the space of twisted loops given by

$$k \cdot (g, \gamma) = (kgk^{-1}, k\gamma), \ k \in G_W$$

Now, let us restrict ourselves to twisted Reeb orbits on the link of the singularity *W*. Let us first introduce some notations.

Definition 8.2. For each element $g \in G_W$, we set

$$\hat{\theta}_g = \text{diag}(\theta_{g,1}, \dots, \theta_{g,n}), \ \theta_{g,i} \in [0,1) \ \forall i.$$

so that we have

$$g = \exp(2\pi i \vec{\theta}_g) := \left(\exp(2\pi i \theta_{g,1}), \dots, \exp(2\pi i \theta_{g,n}) \right).$$

In particular, for a given system of weights $(w_1, ..., w_n; h)$ of W, we have

$$\vec{\theta}_I = (w_1/h, \dots, w_n/h), \quad J = \exp(2\pi i \vec{\theta}_I) \in G_W.$$

Notice that Reeb flow \mathcal{R}_t is given by $\exp(2\pi i t \vec{\theta}_J)$ and J is just \mathcal{R}_1 .

Definition 8.3. We denote by $(\mathbb{C}^n)^{g,l}$ the space of g-twisted Reeb orbits of period l in \mathbb{C}^n . We also denote by $\Sigma_{g,l}$ the space of g-twisted Reeb orbits in the link of W:

$$\Sigma_{g,l} = (\mathbb{C}^n)^{g,l} \cap L_W.$$

Notice that $p \in (\mathbb{C}^n)^{g,l}$ precisely when $g \circ \mathcal{R}_l(p) = p$, and hence $(\mathbb{C}^n)^{g,l}$ is a fixed locus of $g \circ \mathcal{R}_l$ -action on \mathbb{C}^n . Equivalently, we have

(8.1)
$$(\mathbb{C}^n)^{g,l} := \left\{ (z_1, \dots, z_n) \in \mathbb{C}^n : \text{ if } z_i \neq 0, \text{ then } \theta_{g,i} + l\theta_{J,i} = 0 \text{ mod}\mathbb{Z} \right\}$$

A set $\{\Sigma_{g,l}\}_{g \in G_W, l \in \mathbb{R}_{>0}}$ classifies all possible Morse-Bott components of twisted Reeb orbits.

Lemma 8.4. We have

$$\Sigma_{g,l} \simeq \Sigma_{gJ^{-1},l+1}, \ \Sigma_{J^{-1},1} \simeq L_W.$$

The lemma follows immediately from $\Re_1 = J$. One can also easily check that there are finitely many diffeomorphism types of $\Sigma_{g,l}$.

Example 8.5. (Principal component) Since $\Re_1 = J$, every point in L_W arises J^{-1} -twisted Reeb orbit of period 1, and hence, $\Sigma_{J^{-1},1} \simeq L_W$. This gives the principal component of the J^{-1} -twisted Reeb orbits in $[M_W/G_W]$ of period 1. The orbit Γ_W , a fundamental class of of this component, is called principal orbits. We have J^{-l} -twisted Reeb orbits of period l in $[M_W/G_W]$ for any $l \in \mathbb{N}$ in a similar manner.

We may ask whether the period of a twisted Reeb orbit is always an integer as we saw in the previous section. The answer is no in general, and this complicates the story quite a bit. Namely, we can have Reeb orbits of shorter periods (starting at *p*) if some of the coordinates (of *p*) vanish. Here is one example of a fractional period of $l = \frac{1}{3}$.

Example 8.6. (A component with a fractional period) Consider $W = x^2 y + y^4$. Its weight is (3,2;8), and $\vec{\theta}_J = (\frac{3}{8}, \frac{1}{4})$. For $(g, l) = (J^{-3}, \frac{1}{3})$, we have $(\mathbb{C}^2)^{g,l} = \mathbb{C}_x \times \{0\}$. Indeed,

$$J^{-3} \circ \mathscr{R}_{\frac{1}{3}} = (e^{\frac{7\pi i}{4}}, e^{\frac{\pi i}{2}}) \circ (e^{\frac{\pi i}{4}}, e^{\frac{\pi i}{6}}) = (1, e^{\frac{2\pi i}{3}}).$$

Note that $\mathbb{C} \times \{0\}$ *does not arise as a fixed locus for any* $g \in G_W$ *, and moreover* $W|_{\mathbb{C} \times \{0\}} = 0$ *.*

Let us comment on the relationship between twisted sectors and twisted Reeb orbits, although we do not use it in this paper. As G_W is abelian, a twisted sector of the quotient orbifold $[M_W/G_W]$ for an element $k \in G_W$ is given by $[(M_W)^k/G_W]$ where $(M_W)^k$ is the *k*-fixed locus. On the other hand, as *k* acts linearly on \mathbb{C}^n we can consider the restriction of *W* on the *k*-fixed subspace $\operatorname{Fix}(k) \subset \mathbb{C}^n$ to obtain W^k : $\operatorname{Fix}(k) \to \mathbb{C}$. One can easily check that $(M_W)^k$ can be identified with M_{W^k} .

As \mathscr{R}_t commutes with G_W -action, the canonical Reeb flow on the link L_{W^k} is the restriction of that of L_W . In particular, if a point of the Reeb flow γ is fixed by k, then the whole trajectory is fixed by k. A twisted Reeb orbit (g, γ) gives an element of the chain complex $SC^{\bullet}([M_W/G_W])$, but if (a point of) it is fixed by some k, then it also gives rise to an element of $SC^{\bullet}([M_{W^k}/G_W])$ as well.

Remark 8.7. Recall that W raises a canonical exact sequence of symmetry groups

 $0 \longrightarrow G_W \longrightarrow \widetilde{G}_W \xrightarrow{\chi} \mathbb{C}_R^* \longrightarrow 0$

and \widetilde{G}_W acts naturally on $W^{-1}(0)$. Notice that Morse-Bott components $\{\Sigma_{g,l}\}$ corresponds to twisted sectors of $[(W^{-1}(0) \setminus \{0\})/\widetilde{G}_W]$.

8.1.1. Conley-Zehnder indices of twisted orbits. We have identified generators of the chain complex $SC^*([M_W/G_W])$. We compute their Robbin-Salamon indices.

Remark 8.8. We do not know how to define the differential and products in our setting of perturbation scheme (which follows Abouzaid-Seidel), but we do not need them in this paper (except m_1 of Γ_W).

We start with the following elementary observations.

Lemma 8.9. Consider a weighted homogeneous polynomial W, and its maximal diagonal symmetry group G_W . For each $g \in G_W$, we define $F_g \subset \{1, ..., n\}$ so that

 $i \in F_g$ if and only if $g \cdot x_i = x_i$.

We write

$$W = W_{F_g} + W_{move,g}$$

where W_{F_g} is the sum of monomials of W that consists of variables whoses indices are in F_g , and $W_{move,g}$ is the sum of remaining monomials. Let **m** be the ideal generated by x_i with $i \notin F_g$. Then we have

$$W_{move,g} \in \mathbf{m}^2$$

Proof. Note that *g*-action leaves W_{F_g} to be invariant. Since *g*-action leaves *W* to be invariant, the same holds for each monomial of $W_{\text{move},g}$. Note that $W_{\text{move},g} \in \mathbf{m}$. If a monomial of it does not lie in \mathbf{m}^2 , then it cannot be invariant under *g*-action.

Lemma 8.10. We have the following identities of partial derivatives.

(1) For $g \in G_W$, we have

$$\partial_i W(x) = \exp(2\pi i \theta_{g,i}) \partial_i W(gx)$$

(2) *We have*

(8.2)
$$W(R_t(x)) = \exp(2\pi i t) W(x)$$

(8.3)
$$\partial_i W(x) = \exp(2\pi i l(-1 + \theta_{J,i})) \partial_i W(R_l(x))$$

To compute the Robbin-Salamon index of a twisted Reeb orbit (g, γ) , we first find a good frame for the tangent bundle of the Milnor fiber M_W with respect to *g*-action.

Definition 8.11. We define another index set $I_{g,l} \subset \{1, ..., n\}$

$$I_{g,l} = \left\{ i : \partial_i W |_{(\mathbb{C})^{g,l}} = 0 \right\}$$

We will simply write it as I sometimes.

Proposition 8.12. Let γ be a g-twisted loop of period l. The following frame (for a fixed $j_0 \in I^c$) trivializes $T(M_W)|_{\gamma}$:

$$\left\{e^{-2\pi i\theta_{g,i}t/l}e_{i}\colon i\in I\right\}\cup\left\{\frac{e_{j}}{\partial_{j}W}-\frac{e_{j_{0}}}{\partial_{j_{0}}W}\colon j\in I^{c}\right\}.$$

Proof. From the definition of *I*, we have

$$dW|_{(\mathbb{C}^n)^{g,l}} = \sum_{i\in I^c} \partial_i W dz_i.$$

Thus it is clear that the above vectors lie in the Kernel of dW. It remains to check that it is compatible with *g*-twisting $g \cdot \gamma(l) = \gamma(0)$. It is clear for the basis vectors of the first type.

For the second type, from the above lemma, we have

$$\left(\partial_{j}W(\gamma(l))\right) = e^{2\pi i\theta_{g,j}} \left(\partial_{j}W(g\gamma(l))\right) = e^{2\pi i\theta_{g,j}} \left(\partial_{j}W(\gamma(0))\right)$$

Therefore, we obtain the claim as follows.

$$e^{2\pi i\theta_{g,j}} \cdot \frac{e_j}{\partial_j W(\gamma(l))} = \frac{e^{2\pi i\theta_{g,j}}e_j}{e^{2\pi i\theta_{g,j}}\partial_j W(\gamma(0))} = \frac{e_j}{\partial_j W(\gamma(0))}$$

Now, we measure how much the frame rotates with respect to the canonical volume form Ω_{M_W} .

Lemma 8.13. The rotation number (along γ) of the frame in Proposition 8.12 w.r.t. Ω_{M_W} is

$$\left(-\sum_{i\in I}\theta_{g,i}+\sum_{j\in I^c}(l\theta_{J,j}-l)\right)$$

Proof. Given the frame, $\partial_{z_{j_0}} W \neq 0$ for the chosen j_0 in I^c . (If I^c is empty, then choose any j_0 with such property). Hence we have

(8.4)
$$\Omega_{\mathbb{C}^n}^n/dW = \frac{dz_1\cdots dz_{j_0}\cdots dz_n}{\partial_{z_{j_0}}W}$$

By evaluating the basis vectors of the first type (for $i \in I$), we obtain the rotation number $-\sum_{i \in I} \theta_{g,i}$. For the basis vectors of the second type (for $i \in I^c$), note that if we evaluate such vectors to the form Ω_{M_W} , we obtain the factor

$$\prod_{j\in I^c}\frac{1}{\partial_{z_j}W}$$

where the contribution of $j_0 \in I^c$ come from the coefficient of (8.4). From the lemma 8.10, the rotation number of such a factor is

$$\sum_{j \in I^c} l(\theta_{J,j} - 1)$$

To compute the Maslov index of the linearized Reeb flow, we write the linearized Reeb flow in terms of this frame: for $t \in [0, l]$,

$$\left\{e^{2\pi i(\theta_{J,i}+\theta_{g,i}/l)t}\left(e^{-2\pi i\theta_{g,i}t/l}e_{i}\right):i\in I\right\}\cup\left\{e^{2\pi it}\left(\frac{e_{j}}{\partial_{j}W}-\frac{e_{j_{0}}}{\partial_{j_{0}}W}\right):j,j_{0}\in I^{c}\right\}$$

Lemma 8.14. *Robbin-Salamon index of the linearized Reeb flow w.r.t. the frame in Proposition* 8.12 is

(8.5)
$$\sum_{i \in I} Q(l\theta_{J,i} + \theta_{g,i}) + (|I^c| - 1)Q(l),$$

(8.6)
$$Q(s) = \begin{cases} 2s & s \in \mathbb{Z} \\ 2\lfloor s \rfloor + 1 & s \notin \mathbb{Z} \end{cases}$$

Combining Lemma 8.13, 8.14, we obtain the following

Proposition 8.15. Robbin-Salamon index of the linearized Reeb flow in $\Sigma_{g,l}$ is

$$\mu_{\mathrm{RS}}(\Sigma_{g,l}) = 2\left(-\sum_{i\in I}\theta_{g,i} + \sum_{j\in I^c}(l\theta_{J,j}-l)\right) + \sum_{i\in I}Q(l\theta_{J,i}+\theta_{g,i}) + (|I^c|-1)Q(l)$$

Remark 8.16. $\mu_{\text{RS}}(\Sigma_{g,l})$ always lies in $\frac{1}{|G_W|}\mathbb{Z}$.

Now, we rewrite the above expression of $\mu_{\text{RS}}(\Sigma_{g,l})$ as

(8.7)
$$\mu_{\text{RS}}(\Sigma_{g,l}) = 2l(\sum_{i=1}^{n} \theta_{J,i} - 1) + \sum_{i \in I} \left(Q(l\theta_{J,i} + \theta_{g,i}) - 2(l\theta_{J,i} + \theta_{g,i}) \right) + (|I^c| - 1)(Q(l) - 2l)$$

Let us illustrate our formula in some examples.

Example 8.17. (1) For $g = J^{-l}$, we have $(\mathbb{C}^n)^{g,l} = \mathbb{C}^n$ and $I = \emptyset$.

$$\mu_{\rm RS}(\Sigma_{J^{-l},l}) = 2l(\sum_{j=1}^{l} \theta_{J,i} - 1) = \frac{2l(\sum_{k=1}^{l} w_i - h)}{h}.$$

(2) For the example 8.6 of $W = x^2 y + y^4$ with $(g, l) = (J^{-3}, \frac{1}{3})$. We have

$$\vec{\theta}_g = \left(\frac{7}{8}, \frac{1}{4}\right), \vec{\theta}_J = \left(\frac{3}{8}, \frac{1}{4}\right).$$

$$On \mathbb{C} \times \{0\}, \, \partial_1 W = 2xy = 0. \text{ Thus } I = \{1\}, I^c = \{2\}, l = \frac{1}{3}.$$
$$\mu_{\text{RS}}(\Sigma_{g,l}) = 2\left(-\frac{7}{8} + \frac{1}{3}\left(\frac{1}{4} - 1\right)\right) + Q\left(\frac{1}{3} \cdot \frac{3}{8} + \frac{7}{8}\right) + 0 = -2 + 2 = 0.$$

8.2. **Degree shifting number and its non-negativity.** The following number $\star_{g,l} \in \mathbb{Q}$ will be used for degree shifting.

Definition 8.18. We define $\star_{g,l} \in \mathbb{Q}$ as

$$(n-1) - \mu_{\mathrm{RS}}(\Sigma_{g,l}) - \frac{1}{2}(\dim_{\mathbb{R}}\Sigma_{g,l} + 1)$$

It is easy to see that $\star_{g,l} = n - \dim_{\mathbb{C}}(\mathbb{C}^n)^{g,l} - \mu_{\mathrm{RS}}(\Sigma_{g,l})$

Proposition 8.19. We have the following inequality.

$$\star_{g,l} \ge -\frac{2l(\sum_{k=1}^{n} w_i - h)}{h}$$

In particular, $\star_{g,l} \ge 0$ for log Fano and log Calabi-Yau cases.

Proof. (1) Suppose $W|_{(\mathbb{C}^n)^{g,l}} \neq 0$.

There is a j_0 such that $j_0 \in I^c$. Since $\partial_{j_0} W \neq 0$ on $(\mathbb{C}^n)^{g,l}$, from the Lemma 8.10, we obtain the condition

$$l\theta_{J, i_0} + \theta_{g, i_0} \equiv l \mod \mathbb{Z}.$$

Also, as z_{i_0} is fixed under the action of $g \circ \mathcal{R}_l$, we obtain another condition

$$l\theta_{J,i_0} + \theta_{g,i_0} \equiv 0 \mod \mathbb{Z}.$$

Therefore, $l \in \mathbb{N}$ and $g \circ \mathscr{R}_l = g \circ J^l \in G_W$. Hence $(\mathbb{C}^n)^{g,l}$ is a fixed locus of gJ^l -action. Recall from Lemma 8.9 that

$$W = W_{F_{\alpha I}l} + W_{move,gJ^l}$$

and $W_{move,gJ^l} \in \mathbf{m}^2$. Hence, if $j \notin F_{gJ^l}$, $\partial_j W_{move,gJ^l}$ still contains at least one moving variable hence vanishes on $(\mathbb{C}^n)^{g,l}$. Therefore,

$$F_{gJ^l}^c \subset I_{g,l}.$$

Since *W* is a non-degenerate polynomial, we also have $F_{gJ^l} \subset I_{g,l}^c$, which implies that $I_{g,l}^c = F_{gJ^l}$.

Therefore, using the identity 8.7, we have

(8.8)
$$\star_{g,l} = n - \dim_{\mathbb{C}}(\mathbb{C}^n)^{g,l} - \mu_{\mathrm{RS}}(\Sigma_{g,l}) = |I_{g,l}| - \mu_{\mathrm{RS}}(\Sigma_{g,l})$$

(8.9)
$$= -2l(\sum_{i=1}^{n} \theta_{J,i} - 1) + \sum_{i \in I_{g,l}} \left(1 - (Q(l\theta_{J,i} + \theta_{g,i}) - 2(l\theta_{J,i} + \theta_{g,i})) \right) - (|I_{g,l}^{c}| - 1)(Q(l) - 2l)$$

(8.10)
$$\geq -2l(\sum_{i=1}^{n} \theta_{J,i} - 1)$$

Here, we use the fact that Q(s) - 2s < 1 for all *t* and Q(l) - 2l = 0 as $l \in \mathbb{Z}$.

(2) Suppose $W|_{(\mathbb{C}^n)^{g,l}} = 0$. If $i \in I_{g,l}^c$, then $\partial_i W|_{(\mathbb{C}^n)^{g,l}} \neq 0$, but the latter vanishes from the assumption. Hence, $I_{g,l}^c = \emptyset$ and $I_{g,l} = \{1, \dots, n\}$. Consider the action of $g \circ R^l$ on (z_1, \dots, z_n)

and say $i \in K \subset \{1, \dots, n\}$ if and only if z_i is fixed under this action. Then $\dim_{\mathbb{C}}(\mathbb{C}^n)^{g,l} = |K|$. Also, we have $l\theta_{J,i} + \theta_{g,i} \in \mathbb{Z}$ for $i \in K$. Therefore, we have

(8.11)
$$\star_{g,l} = n - \dim_{\mathbb{C}}(\mathbb{C}^n)^{g,l} - \mu_{\mathrm{RS}}(\Sigma_{g,l})$$

(8.12)
$$= n - |K| - 2l(\sum_{i=1}^{n} \theta_{J,i} - 1) - \sum_{i=1}^{n} \left(Q(l\theta_{J,i} + \theta_{g,i}) - 2(l\theta_{J,i} + \theta_{g,i}) \right) + (Q(l) - 2l)$$

3)
$$= -2l(\sum_{i=1}^{n} \theta_{J,i} - 1) + \sum_{i \notin K} \left(1 - (Q(l\theta_{J,i} + \theta_{g,i}) - 2(l\theta_{J,i} + \theta_{g,i})) \right) + (Q(l) - 2l)$$

(8.14)
$$\geq -2l(\sum_{i=1}^{n} \theta_{J,i} - 1)$$

This completes the proof.

9. MONODROMY ORBIT Γ_W and vanishing of spheres with Γ_W

9.1. Morse-Smale function for perturbation. Let us first define a Morse-Smale function on $(L_{W,\delta}/G_W)$ that is needed for perturbations of Morse-Bott components. Although Morse-Smale function might not exist for an orbifold, our situation is rather special so that we can construct one as follows.

Lemma 9.1. There is an G_W -equivariant Morse function h on $L_{W,\delta}$ such that

- *if a critical point* $p \in crit(h)$ *lies in* Fix(K) *for some* $K < G_W$ *, then the unstable manifold* $W^{-}(p)$ of p is also contained in Fix(K).
- its Morse-Smale-Witten complex is well-defined and computes the cohomology of $(L_{W,\delta}/G_W)$.

Proof. Since orbifold strata $\{Fix(K)\}_{K < G_W}$ of \mathbb{C}^n intersect transversally to the link, it induces orbifold strata for $L_{W,\delta}$. We construct a Morse function h' in an inductive way. The action of a diagonal symmetry group is effective on the link. Therefore the deepest strata does not have a nontrivial fixed locus. Choose a Morse function on it and extend it strata by strata in a way that it depends on the radial direction of the normal bundle of the strata inside the next one. We may choose our extension so that the negative Morse flow line of our extension flows into the lower strata. This is possible because our orbifold strata are induced from coordinate planes. When several stratum intersect along a lower dimensional one, their normal directions are compatible.

Morse flow of h' respects orbifold strata. If $p \in \operatorname{crit}(h') \cap \operatorname{Fix}(K)$, then its unstable manifold $W^{-}(p)$ is contained in Fix(K). If $q \in \operatorname{crit}(h') \cap \operatorname{Fix}(K')$ for some $K \leq K'$, then its stable manifold $W^+(q)$ intersect $W^-(p)$ at smooth points of Fix(K) only. Therefore, the moduli space of gradient flow trajectories can be defined as in the smooth case. They intersect transversally after a small perturbation of h' in the smooth part of Fix(K). We do it strata by strata to obtain h.

Because of this first property, Morse-Witten complex of h makes sense. i.e, its differential does square to zero. Moreover, the unstable manifolds of h provide a cell decomposition of $(L_{W,\delta}/G_W)$. Hence Morse-Witten complex computes the singular cohomology of $(L_{W,\delta}/G_W)$ (for a similar argument for $G = S^1$ in the context of contact homology, see [Bou03]).

9.2. A symplectic cohomology cocycle Γ_W . Finally, let us define Γ_W .

We adapt the idea of [CFHW96] and [KvK16]. Choose a small normal neighborhood $v(\Sigma_{J^{-1},1}/G_W) \simeq L_{W,\delta}/G_W \times (-\epsilon,\epsilon)$ inside M_W/G_W . Choose a sufficiently small, nonnegative bump function ρ supported inside $(-\epsilon,\epsilon)$. Use our Morse function $h = h_{J^{-1},1}$ in 9.1 to define an equivariant time-dependent perturbation term by

$$\overline{h}: \nu \left(L_{W,\delta}/G_W \right) \times S^1 \to \mathbb{R}$$
$$(p, v, t) \mapsto h(Fl_t^{\mathscr{R}}(p))\rho(v)$$

Definition 9.2. A local Floer cochain complex

$$CF_{loc}^{\bullet}(L_{W,\delta}/G_W, H_{S^1}) = \bigoplus_{\gamma \in \mathcal{O}(H_{S^1})} o_{\gamma}$$

is defined to be a free module generated by Hamiltonian orbits of time-dependent Hamiltonian $H_{S^1} := H + \overline{h}$. Its differential is defined similarly as a counting of pseudo-holomorphic cylinders, but only those inside $v(L_{W,\delta}/G_W)$. Its cohomology, called local Floer cohomology, is denoted by $HF_{loc}^{\bullet}(L_{W,\delta}/G_W, H_{S^1})$.

Theorem 9.3. As a $\mathbb{Z}/2$ -graded vector space,

$$HF_{loc}^{\bullet}(L_{W,\delta}/G_W, H_{S^1}) \simeq H_{Morse}^{\bullet}(L_{W,\delta}/G_W, h; \mathbb{Z})$$

Proof. See [KvK16], Proposition 8.4. One can check that $L_{W,\delta}/G_W$ satisfies all the conditions in the proposition except the first Chern class condition, which makes the isomorphism only $\mathbb{Z}/2$ -graded. We can also check that flows and Hamiltonians in the proof are equivariant.

Definition 9.4. The Hamiltonian orbit $\Gamma_W \in CF^{\bullet}_{loc}(L_{W,\delta}/G_W, H)$ is defined to be a cochain which corresponds to a fundamental class

$$\Gamma_W \leftrightarrow \left[L_{W,\delta} / G_W \right] \in H^{\bullet} \left(L_{W,\delta} / G_W; \mathbb{Z} \right).$$

It will be shown that Γ_W is indeed a cocycle. We call Γ_W the monodromy orbit.

In Lemma 9.7, we prove that there are no rigid Floer cylinder emanating from Γ_W other then the ones in local Floer complex of $\Sigma_{f^{-1},1}/G_W$. Although we do not define full-fledged orbifold symplectic cohomology complex, it is legitimate to say Γ_W is closed is a Floer theoretic sense.

We also apply perturbations of $\Sigma_{g,l}$ using Morse functions $h_{g,l}$ on each Morse-Bott components. Such perturbation is well-defined since $h_{g,l}$ are G_W -equivariant. Also, there are only finitely many diffeomorphism types of $\Sigma_{g,l}$. Therefore, we can assume that there is an $\epsilon > 0$ such that

- (1) $\max h_{g,l} \min h_{g,l} \le \epsilon$ for $\forall g \in G_W, l \in \mathbb{R}_{\ge 0}$,
- (2) if γ arises as a critical point of $h_{g,l}$, then $\mathscr{A}_{H_{S^1}}(\gamma) \in [-l^2 \epsilon, -l^2 + \epsilon]$,
- (3) if Σ_{g_1,l_1} and Σ_{g_2,l_2} are nonempty and $l_1 \neq l_2$, then $[-l_1^2 \epsilon, -l_1^2 + \epsilon] \cap [-l_2^2 \epsilon, -l_2^2 + \epsilon] = \emptyset$.

Define a filtration $\mathscr{F}^{-l} := \{ \gamma : \mathscr{A}_{H_{S^1}}(\gamma) \ge -l^2 - \epsilon \}$. Then similar to 5.8, we get

Lemma 9.5. An associated $\mathbb{Q} \times \mathbb{Q}$ -graded of the filtration $\mathscr{F}^{-\bullet}$ is

$$E_0^{pq} = \begin{cases} \bigoplus_{g \in G_W} C^{q-2age(g)} \left((M_W)^g / G_W \right) & (p=0) \\ \bigoplus_{g \in G_W} C^{p+q-\star_{g,-p}} (\Sigma_{g,-p} / G_W) & (p<0) \\ 0 & (p>0) \end{cases}$$

Proof. The cases of p = 0 correspond to the constant orbits. In Example 3.7 of [GZ21], Gironella and Zhou explained that the relation between the Conley-Zehnder indices of constant orbits and age of the corresponding Chen-Ruan cohomology class, which explains the degree shifting for the case of p = 0. For p < 0, \star was defined in Definition 8.18 to denote the degree shifting number, and hence the lemma follows.

Corollary 9.6. deg(Γ_W) = $-\mu_{\text{RS}}(\Gamma_W) = -\frac{2((\sum_{i=1}^n w_i) - h)}{h}$

Proof. Since Γ_W corresponds to a fundamental class of a Morse-Bott family $\Sigma_{J^{-1},1}$, we get the second inequality from Proposition 8.15 as computed in Example 8.17. It can be used to compute the degree of Γ_W as follows. Note that the degree shifting number $\star_{J^{-1},1} \in \mathbb{Q}$ is given by

$$(n-1) - \mu_{\rm RS}(\Sigma_{J^{-1},l}) - \frac{1}{2}(\dim_{\mathbb{R}}\Sigma_{J^{-1},1} + 1) = -\mu_{\rm RS}(\Sigma_{J^{-1},l})$$

As Γ_W is from the fundamental class of the principal orbits, deg $(\Gamma_W) - \star_{J^{-1},1} = 0$, which implies the claim.

The construction of the full orbifold symplectic cohomology class will face a transversality problem in the current perturbation scheme of pseudo-holomorphic curves, since some generators are contained in singular locus of the orbifold. Fortunately, the element Γ_W will comes from $\Sigma_{J^{-1},1}/G_W \simeq L_W/G_W$, which is inside a smooth locus of $[M_W/G_W]$. Pseudo-holomorphic curves with such input do not suffer transversality issue since it is not contained in a singular locus of $[M_W/G_W]$.

Lemma 9.7. Suppose H_{S^1} is *G*-equivariant, $H_{S^1} > 0$, and C^2 -small Morse perturbation of H inside a compact region. Then there is no pseudo-holomorphic cylinder of finite energy satisfying

$$u: S^1 \times \mathbb{R} \to [M_W/G_W] \quad \lim_{s \to \infty} u(t, s) = \Gamma_W(t).$$

whose output

$$\gamma_{-}(t) := \lim_{s \to -\infty} u(s, t)$$

does not lie in $v(\Sigma_{J^{-1},1}/G_W) \times \{1\}$.

Proof. At first, we can rule out the case when the output is on outside of a compact region using the idea of [Sei06] using action values: for a non-trivial orbit $\gamma \in \mathcal{O}(H_{S^1})$ at the end, its action is given by

$$\begin{split} A_{H_{S^1}}(\gamma) &:= -\int_{S^1} \gamma^* \lambda + \int_0^1 H_{S^1}(\gamma(t)) dt \\ &= -2 \int_0^1 r^2 dt + \int_0^1 H(\gamma(t)) dt + \int_0^1 F(\gamma(t)) dt, \ (H_{S^1} = H(r) + F(r, t)) \\ &= -\int_0^1 r^2 + \epsilon \ (\epsilon <<1) \end{split}$$

Nontrivial Hamiltonian orbits are appears as a small perturbation of $\Sigma_{g,l}$. An action value of such orbit is dominated by $-l^2$. Therefore, the output γ_- cannot be an orbit of level l > 1 by the positivity of the topological energy.

Suppose γ_{-} is an orbit of period $0 < l \le 1$. Since *u* provides a homotopy between the loop Γ_W and the orbifold loop (γ_{-}, g) , the (conjugacy class of) *g* must be equal to J^{-1} . This means that

 $\gamma_{-}(0)$ is a fixed point of \mathscr{R}_{1-l} . Because $w_i/h \leq \frac{1}{2}$ for $\forall i$, Reeb flow \mathscr{R}_{1-l} has a fixed point only at the origin (which is not in M_W). This gives the contradiction.

Similarly, if γ_{-} is Morse critical point in a compact region, we must have $\Phi(\gamma_{-}(0)) = \gamma_{-}(1)$. It must be a fixed point of a monodromy Φ_{W} . It is impossible because the only fixed point of Φ_{W} is the origin, which is not contained in M_{W} .

Also notice that an orbifold nodal degeneration does not affect a standard analysis of codimension 1 boundary strata of moduli spaces because they are of codimension two. It follows from the fact that a local model of such degeneration is given by a family $z_1 z_2 : [\mathbb{C}^2/(\mathbb{Z}/n)] \to \mathbb{C}$. Here, a group \mathbb{Z}/n acts on \mathbb{C}^2 by $(z_1, z_2) \to (\xi \cdot z_1, \xi^{-1} \cdot z_2)$ with ξ an *n*-th root of unity.

Therefore operations such as a closed-open map $CO(\Gamma_W)$, quantum cap action $\cap \Gamma_W$ and popsicle operations $\mathbf{P}_{n,F,\phi}^{\Gamma_W}$ on $\mathcal{WF}([M_W/G_W])$ make sense, and we use them to define the A_{∞} -operation $\{M_k\}$.

The final important piece, the vanishing of sphere operations can be shown as follows.

Proposition 9.8. If $\sum_{k=1}^{n} w_i - h \ge 0$, then popsicle spheres with two or more Γ_M inputs vanish.

Proof. Proof is similar to that of Proposition 5.9.

Suppose such a sphere bubble from popsicle moduli space exist. As Corollary 5.5 still holds, we have a sphere component with N insertions of Γ_W with an output ξ , satisfying the same inequality (5.6).

From Lemma 9.5, the degree of generators of symplectic cochains are either non-negative (in the constant orbit cases) or satisfies the inequality $p + q - \star_{g,-p} \ge 0$ or $p + q \ge \star_{g,-p}$ (for other cases). Let us denote

$$\mu := \mu_{\rm RS}(\Sigma_{J^{-1},1}) = \frac{2(\sum_{k=1}^{n} w_i - h)}{h}$$

for simplicity. From Proposition 8.19, we have

$$\star_{g,l} \ge -\frac{2l(\sum_{k=1}^{n} w_i - h)}{h} = -l\mu$$

Combining these, we have

if
$$\mu$$
 is non-negative and $\xi \in F^{-l}$, then deg $\xi \ge -l \cdot \mu$.

Now, rest of the proof is exactly the same as that of Proposition 5.9.

Remark 9.9. Although we only prove log-Fano/log Calabi-Yau cases, the readers can easily check that the same proof works for when $\mu_{\text{RS}}(\Sigma_{J^{-1},1}) = \frac{2(\sum_{i=1}^{n} w_i - h)}{h} > -\frac{1}{2}$.

APPENDIX A. MODULI SPACE OF PSEUDO-HOLOMORPHIC CURVES AND PERTURBATIONS

We briefly describe a perturbation scheme for moduli spaces of pseudo-holomorphic curves that we use. We refer [Sei08], [AS10], [Abo12], and especially [Abo10] and [Sei18] from which most of the material has been borrowed.

A.1. **Setup.** Let $(M^{2n}, \omega = d\lambda)$ be a Liouville manifold with cylindrical ends. By definition, *M* can be decomposed into a union of a compact part and cylindrical ends

$$M = M_{cpt} \bigcup_{\partial M_{cpt}} \partial M_{cpt} \times [1, \infty),$$

and Liouville flow *Z* is of the form $Z = r \frac{\partial}{\partial r}$ at the cylindrical end where *r* is a coordinate of $[1, \infty)$. We also assume $c_1(TM) = 0$ to put \mathbb{Z} -grading everywhere.

- We will work with a function $H \in C^{\infty}(M,\mathbb{R})$ such that H > 0, C^2 -small on M_{cpt} and qua*dratic at infinity* $(H(x, r) = r^2)$, and denote the class of such functions by $\mathcal{H}(M)$.
- Whenever we consider a time dependent perturbation $H_{S^1} = H + F : S^1 \times M \to \mathbb{R}$, we assume $H_{S^1} > 0$, C^2 -small on M_{cpt} so that the time-1 periodic orbits of H_{S^1} are nondegenerate. This is true for a generic perturbation.
- a ω -compatible almost complex structure *J* is called *contact type* if $\lambda \circ J = dr$ at the end. We denote a class of such almost complex structure by $\mathscr{J}(M)$.

 \mathcal{W} is a collection of exact properly embedded Lagrangian submanifolds in M, such that if non-compact, $L \cap \partial M_{cpt}$ is a Legendrian submanifold, and L is **conical at the end**. Furthermore, all such L is required to have vanishing relative first Chern class $2c_1(M, L)$. We attach a spin structure and a grading function on each L.

Fix a small, time dependent perturbation $H_{S^1}: S^1 \times M \to \mathbb{R}$ of H. Let $\mathcal{O} := \mathcal{O}(M, H_{S^1})$ be a set of time-1 orbits γ of H_{S^1} , and $\chi(L_0, L_1; H)$ to be the set of time-1 Hamiltonian chords *a* of *H* from L_0 to L_1 for two Lagrangians $L_0, L_1 \in \mathcal{W}$. We use a notation o_{γ} (resp. o_a) to denote its orientation operator. Its degree deg o_{γ} (resp. deg o_a) is given by its cohomological Conley-Zehnder index (resp. Maslov index).

A.2. Moduli spaces. The moduli space of the Riemann sphere with m + 1 marked points (m for positive punctures and 1 for the negative puncture) is denoted as $S_{m,1;0}$. The moduli space of a disc with *m* interior marked points $\{z_1^+, \ldots, z_m^+\}$ (all for positive punctures) and with n + 1cyclically ordered boundary marked points $(\{z_1, \ldots, z_n\}$ for positive punctures and $\{z_0\}$ for the negative puncture) is denoted as $S_{m;n,1}$. An oriented real blow-up of \overline{S} at marked points is a surface |S such that boundary puncture z_k (resp. interior puncture z_l^+) is replaced by $[0, 1] \times \{z_k\}$ (resp. $S^1 \times \{z_1^+\}$). Let us denote (with coordinate (s, t))

$$Z^+ = [0,\infty) \times [0,1], Z^- = (-\infty,0] \times [0,1], C^+ = [0,\infty) \times S^1, C^- = (-\infty,0] \times S^1.$$

Definition A.1. A set of ends \mathfrak{S} for $S \in S_{m;n,1}$ is a choice of

- strip-like ends $\epsilon_k^{\pm} : Z_{\pm} \to S$, $\lim_{s \to \pm \infty} \epsilon_k^{\pm}(s, t) = z_k$, $(\epsilon_k^{\pm})^{-1}(\partial S) = \{t = 0, 1\}$ cylindrical ends $\delta_l^{\pm} : C_{\pm} \to S$, $\lim_{s \to \pm \infty} \delta_l^{\pm}(s, t) = z_l^{\pm}$

Such collection is said to be weighted if each strip and cylinder is endowed with a positive real number

- w[±]_{S,k} for each strip-like end ε^k_±
 v[±]_{S,l} for each cylindrical end δ^l_±

such that $\sum w_{Sk}^{+} + \sum v_{Sl}^{+} = \sum w_{Sk}^{-} + \sum v_{Sl}^{-}$.

The choice of ends (without weights) is a choice complex coordinate *z* of *S* near punctures. It also provides an analytic coordinate (σ , *t*) of |*S* near puncture in the following way.

$$\sigma = e^{\mp \pi s} = |z|, \quad t = \mp \arg(z)/\pi$$

We denote Deligne-Mumford compactification of a family of surfaces |S by $|S_{m;n,1}$.

Definition A.2. Let (S, \mathfrak{S}) denote a holomorphic disc S with ends $\{\kappa\}$ with weight $\{v_{\kappa}\}$.

(1) A basic, asymptotically compatible 1-form α_S is a closed 1-form on S whose restriction on ∂S vanishes, extends smoothly to |S| and $\alpha_S = v_{\kappa}dt$ at the interval / circle at infinity. It implies

$$\kappa^* \alpha_S = \nu_{\kappa} dt + d(e^{\mp \pi s} g_{\kappa}(e^{\mp \pi s}, t)), \quad \forall \kappa$$

for sufficiently large |s| and for some smooth function g_{κ} that vanishes for t = 0, 1.

(2) A secondary, asymptotically compatible 1-form β_S is a sub-closed 1-form whose restriction on ∂S vanishes, extends smoothly to $|S, \beta_S = v_{\kappa} dt$ at the circle at infinity, and vanishes at the intervals at infinity. It implies

$$\kappa^* \beta_S = \begin{cases} \nu_{\kappa} dt + d(e^{\mp \pi s} h_{\kappa}(e^{\mp \pi s}, t)) & \forall cylindrical end \kappa \\ d(e^{\mp \pi s} h_{\kappa}(e^{\mp \pi s}, t)) & \forall strip-like end \kappa \end{cases}$$

for sufficiently large |s| and for some smooth function h_{κ} that vanishes for t = 0, 1 whenever κ is a strip-like ends.

(3) An \mathfrak{S} -adapted time-shifting map is a function $a_S: \partial \overline{S} \to [1,\infty)$ such that

$$\kappa^* a_S = v_{\kappa}, \quad \forall strip-like \ end \ \kappa.$$

Here, κ^* should be thought of the restriction along a small neighborhood of x_k .

(4) For a fixed Hamiltonian $H \in \mathcal{H}(M)$, an S-dependent Hamiltonian H_S is said to be compatible with (S, \mathfrak{S}, H) if

$$\kappa^* H_S = \frac{H \circ \psi^{\nu_\kappa}}{\nu_\kappa^2}, \quad \forall \kappa$$

(5) For a fixed absolutely bounded S^1 -dependent Hamiltonian perturbation F, an S-dependent Hamiltonian F_S is said to be compatible with (S, \mathfrak{S}, F) if it is also absolutely bounded and

$$\kappa^* F_S = \begin{cases} \frac{F \circ \psi^{\nu_{\kappa}}}{v_{\kappa}^2} & \forall cylindrical end \kappa \\ 0 & \forall strip-like end \kappa \end{cases}$$

(6) For a fixed S¹ or [0,1]-dependent almost complex structure J_t, an S-dependent almost complex structure J_S is called asymptotically compatible to (S, S, a_S, J_t) if it extends smoothly to |S and

$$J_p \in \mathscr{J}(M), \quad \forall p \in S$$

It implies

$$\kappa^* J_S = \left(\phi^{\nu_{\kappa}}\right)^* J_t + e^{\mp \pi s} O_{\kappa,(s,t)}, \quad \forall \kappa$$

for some error term $O_{\kappa,(s,t)}$.

Remark A.3. In [Abo10], (1), (2), and (6) is required to be strictly compatible to \mathfrak{S} which means

$$\kappa^* \alpha_S = v_\kappa dt$$

and similar for β_S and J_S . In particular, it depends on the choice \mathfrak{S} . An idea of asymptotic compatibility is due to [Sei18]. It is more flexible because it only depends smoothly on |S, not on

the choice of the set of ends $\{\kappa_S\}$. It allows us to choose a special kind of ends to deal with popsicle structures later on.

Finally, we define

Definition A.4. For a fixed surface S, Hamiltonian $H \in \mathcal{H}(M)$ and absolutely bounded timedependent Hamiltonian F, a Floer data Floer_S consists of

- (1) A collection of weighted strip and cylinder data \mathfrak{S} ;
- (2) A basic 1-form α_S and secondary 1-form β_S asymptotically compatible to (S, \mathfrak{S}) ;
- (3) An (S, \mathfrak{S}) -adapted time-shifting map a_S ;
- (4) An S-dependent, (S, \mathfrak{S}, H, F) -compatible Hamiltonian H_S and F_S ;
- (5) An S-dependent, asymptotically compatible almost complex structure J_S .

Also, we say Floer¹_S and Floer²_S are conformally equivalent if Floer²_S is a rescaling by Liouville flow of Floer¹_S, up to constant ambiguity in the Hamiltonian terms.

In the simplest case of a strip $S \in S_{0;1,1}$ or a cylinder $S \in S_{1,1;0}$, we choose a canonical striplike/cylindrical end with weights 1 for all ends. A form dt is a compatible sub-closed one form. A universal and consistent choice of Floer data is a choice of Floer data Floer_S for all $S \in S_{m;n,1}$ which varies smoothly over the moduli space. Since the space of Floer data is contractible, we can extend it to $\overline{|S_{m;n,1}|}$.

Definition A.5. Let $\gamma_i \in \mathcal{O}$ be a time-1 Hamiltonian orbits and $a_j \in \chi(L_{j-1}, L_j)$, j = 1, ..., n and $a_0 \in \chi(L_n, L_0)$ be Hamiltonian chords. Define the moduli space $\overline{\mathcal{M}}_{m;n,1}(\gamma_1, ..., \gamma_m; a_1, ..., a_n, a_0)$ of maps

$$\left\{u:S\to M\middle|S\in\overline{S}_{m;n,1}\right\}$$

satisfying the inhomogeneous Cauchy-Riemann equation with respect to J_S

(A.1) $(du - X_{H_S} \otimes \alpha_S - X_{F_S} \otimes \beta_S)^{0,1} = 0$

and the following asymptotic boundary conditions;

$$\lim_{s \to \infty} u \circ \epsilon_{+}^{k}(s, \cdot) = a_{k}$$
$$\lim_{s \to -\infty} u \circ \epsilon_{-}^{0}(s, \cdot) = a_{0}$$
$$\lim_{s \to \infty} u \circ \delta_{+}^{l}(s, \cdot) = \gamma_{l}$$

 $u(z) \in \psi^{a_S(z)}L_i, z \in \partial_i S, an i-th boundary component of S.$

Additional perturbation terms α_S , β_S , H_S , F_S come from the universal and consistent choice of *Floer data* Floer_S.

The following compactness and transversality result is standard.

Lemma A.6. For a generic choice of universal and consistent Floer data,

- (1) The moduli spaces $\mathcal{M}(\gamma_1, \ldots, \gamma_m; a_1, \ldots, a_n, a_0)$ are compact.
- (2) For a given input γ_i , i = 1, ..., m and a_j , j = 1, ..., n, there are only finitely many a_0 for which $\overline{\mathcal{M}}_{m;n,1}(\gamma_1, ..., \gamma_m; a_1, ..., a_n, a_0)$ is non-empty.

(3) It is a manifold of dimension

$$\dim_{\mathbb{R}} \overline{\mathcal{M}}_{m;n,1}(\gamma_1,...,\gamma_m;a_1,...,a_n,a_0) = (2m+n-2) + \deg o_{a_0} - \sum_{i=1}^m \deg o_{\gamma_i} - \sum_{j=1}^n \deg o_{a_j}$$

Proof. See [Abo10]. For a compactness result, one need to assure that the energy of pseudoholomorphic curves are a priori bounded in M. This estimate is carefully done therein. Transversality result is a standard application of Sard-Smale argument. The dimension formula is also a standard application of Atiyah-Singer index theorem on a linearized Fredholm operator.

When $\mathcal{M}_{m;n,1}(\gamma_1, \dots, \gamma_m; a_1, \dots, a_n, a_0)$ has dimension zero so that it is rigid, then a map $u : S \to M$ in that moduli space is isolated. An orientation of the moduli space provides a canonical isomorphism

$$Q_u: \bigotimes_{i=1}^m o_{\gamma_i} \otimes \bigotimes_{j=1}^n o_{a_j} \to o_{a_0}$$

We sum up Q_u for all $u \in \mathcal{M}(\gamma_1, \dots, \gamma_m; a_1, \dots, a_n, a_0)$ and all a_0 and define

$$\mathbf{F}_{m;n,1}(\gamma_1,\ldots,\gamma_m;a_1,\ldots,a_n) = \sum_{\dim_{\mathbb{R}}\mathcal{M}(\gamma_1,\ldots,\gamma_m;a_1,\ldots,a_n,a_0)=0} \sum_{u \in \mathcal{M}(\gamma_1,\ldots,\gamma_m;a_1,\ldots,a_n,a_0)} Q_u\left(\bigotimes_{i=1}^m o_{\gamma_i} \otimes \bigotimes_{j=1}^n o_{a_j}\right)$$

(m

)

n

We define $\mathcal{M}(\gamma_1, \ldots, \gamma_m, \gamma_0)$ and $\mathbf{F}_{m,1;0}$ in a similar way.

Let us first recall the our setup and explain the detailed construction.

A.3. Wrapped Fukaya category and symplectic cohomology. Symplectic cohomology $SH^{\bullet}(M)$ is a version of Hamiltonian Floer cohomology for Liouville domain introduced by Cieliebak, Floer and Hofer [CFH95] and Viterbo [Vit99]. For Lagrangian submanifolds in M that are either compact or cylindrical at infinity, a wrapped Fukaya A_{∞} -category $\mathcal{WF}(M)$ was defined by Abouzaid-Seidel [AS10]. We use the version with quadratic Hamiltonian of Abouzaid [Abo12] which we recall briefly here.

For two Lagrangian submanifolds $L_0, L_1 \in W$, a *wrapped Floer cochain complex* is a vector space

$$CW^{\bullet}(L_0, L_1; H) = \bigoplus_{a \in \chi(L_0, L_1; H)} o_a$$

It is graded by the degree deg o_a . We will use the notation a instead of o_a for generators, and $CW^{\bullet}(L_0, L_1)$ instead of $CW^{\bullet}(L_0, L_1; H)$ if it cause no confusion.

Definition A.7. A wrapped Fukaya category $\mathcal{WF}(M)$ consists of a set of objects \mathcal{W} with the space of morphisms $CW^{\bullet}(L_0, L_1)$ for $L_i \in \mathcal{W}$ equipped with an A_{∞} structure

$$m_k : CW^{\bullet}(L_0, L_1) \otimes \cdots \otimes CW^{\bullet}(L_{k-1}, L_k) \to CW^{\bullet}(L_0, L_k)$$
$$m_k(a_1, \dots, a_k) = (-1)^{\diamond_k} F_{0;k,1}(a_1, \dots, a_k);$$
$$\diamond_k = \sum_{i=1}^k i \cdot \deg a_i$$

The Liouville flow ψ^{ρ} for time log(ρ) defines a canonical isomorphism

$$CW^{\bullet}(L_0, L_1; H, J_t) \simeq CW^{\bullet}\left(\psi^{\rho} L_0, \psi^{\rho} L_1; \frac{H \circ \psi^{\rho}}{\rho}, (\psi^{\rho})^* J_t\right)$$

60

Also, $H \circ \psi^{\rho} = \rho^2 r^2$ at the cylindrical end. Therefore $\frac{H \circ \psi^{\rho}}{\rho^2} \in \mathcal{H}(M)$. The proof of A_{∞} relation follows from the degeneration patterns of pseudo-holomorphic discs which correspond to a codimension 1 boundary strata of Gromov bordification $\overline{\mathcal{M}}_{0;k,1}(a_1,\ldots,a_k,a_0)$. In particular, we have $m_1^2 = 0$. Its m_1 -cohomology $HW^{\bullet}(L_0, L_1)$ is called the wrapped Floer cohomology.

We are also interested in the symplectic cohomology of Liouville manifold.

Definition A.8. A symplectic cochain complex is a \mathbb{Z} -graded cochain complex

$$CH^{\bullet}(M;H_{S^1}) = \bigoplus_{\gamma \in \mathcal{O}(M;H_{S^1})} o_{\gamma}$$

graded by the degree deg o_{γ} . We will use the notation γ instead of o_{γ} for generators if it cause no confusion. A differential of this complex is

$$d_{CH}(o_{\gamma_1}) = (-1)^{\deg o_{\gamma_1}} F_{1,1;0}(\gamma_1).$$

Recall that $\mathbf{F}_{1,1;0}(\gamma_1)$ is given by a counting of a zero-dimensional component of a moduli space of pseudo-holomorphic cylinders $\mathcal{M}_{1,1;0}(\gamma_1,\gamma_0)$. Its cohomology is denoted as $SH^{\bullet}(M)$.

A ring structure on its cohomology is induced from

$$CH^{\bullet}(M)^{\otimes 2} \to CH^{\bullet}(M) : (\gamma_1, \gamma_2) \mapsto (-1)^{\deg \gamma_1} \mathbf{F}_{2,1;0}(\gamma_1, \gamma_2)$$

Let us recall the definition of a closed-open map to Hochschild cochain complex of wrapped Fukaya category.

Definition A.9. A closed-open map is a map

$$CO = \{CO_l\}_{l \ge 0} : CH^{\bullet}(M) \to CC^{\bullet}(\mathscr{WF}(M), \mathscr{WF}(M))$$
$$CO_l(\gamma)(a_1, \dots, a_l) := (-1)^{\diamond_l} F_{1;l,1}(\gamma; a_1, \dots, a_l, a_0)$$

A degeneration pattern of a moduli space $\overline{\mathcal{M}}_{1;n,1}(\gamma; a_1, \dots, a_n, a_0)$ proves that CO is a cochain map.

APPENDIX B. POPSICLE MODULI SPACE AND ITS CHARTS

In this section, we prove the following theorem.

Theorem B.1. The compactified moduli space of popsicles $\overline{P}_{n,\tilde{F},\phi}$ is a manifold with corners.

For the compactification, we need to consider a mixture of two disjoint degenerations; one is when an underlying disc component breaks into several pieces, and the other is when several sprinkles collide. The first part can be covered by the result of [AS10], and we will focus on the second type of degeneration. To prove the theorem, we will describe a coordinate chart of a given stable popsicle with an alignment data (Φ, Ψ). Let us first recall the rational strip-like/cylindrical ends from [Sei18].

Definition B.2. A choice of strip-like ends $(\epsilon_0^-; \epsilon_1^+, \dots, \epsilon_k^+)$ is called rational if

(1) ϵ_0^- extends to a map $\epsilon_0^-: Z \to \overline{\Sigma}$ such that $\epsilon_0^-(-\infty) = z_0$.

(2) ϵ_k^+ extends to a map $\epsilon_k^+: Z \to \overline{\Sigma}$ such that $\epsilon_k^+(-\infty) = z_0$ and $\epsilon_k^+(+\infty) = z_k$.

CHO, CHOA, AND JEONG

Likewise, a choice of cylindrical ends $(\eta_0^-; \eta_1^+, \dots, \eta_k^+)$ *is called rational if*

- (1) η_0^- extends to a map $\eta_0^-: \hat{\mathbb{C}} \to \overline{\Sigma}^+$ such that $\eta_0^-(0) = w_0$. (2) η_k^+ extends to a map $\eta_k^+: \hat{\mathbb{C}} \to \overline{\Sigma}^+$ such that $\eta_k^+(0) = w_0$ and $\varepsilon_k^+(\infty) = w_k$.

If $\Sigma \simeq Z$ is semi-stable with two boundary marking, two choices of strip-like ends are equivalent if they are differ by a translation of Z. In a similar way, if $\Sigma^+ \simeq \mathbb{C}^*$ is semi-stable with two interior markings, two choices of cylindrical ends are equivalent if they are differ by a dialation of \mathbb{C}^* .

A rational end at the negative end can be viewed as a choice of a particular planar models of popsicles: ε_0^- determines an isomorphism $\overline{\Sigma} \simeq \mathbb{H}$ which sends z_0 to ∞ , and the image of ε_0^- is a complement of a unit half disc $\{|z| \ge 1\}$. The rest of its boundary markings can be identified with points $x_k \in \mathbb{R}$, and their rational ends ϵ_k^+ determine semicircle centered at x_k of radius ρ_k . Note that equivalent strip-like ends have the same image of ϵ -maps, hence radius is well-defined.

Similarly, a rational end η_0^- determines an isomorphism $\overline{\Sigma}^+ \simeq \hat{\mathbb{C}}$ which sends w_0 to ∞ , and the image of η_0^- is a complement of a unit disc $\{|w| \ge 1\}$. Then, the rest of the markings w_k^+ can be identified with a complex number $\{z_k\}$ and the rational end η_k^+ is determined by a disc with radius ξ_k centered at each z_k .

Next, we discuss the alignment of ends.

Definition B.3. Suppose a popsicle disc Σ and a popsicle sphere Σ^+ are aligned to each other. A choice of strip-like/cylindrical ends for Σ and Σ^+ is said to be **aligned** if the following condition holds:

- Strip-like ends of Σ and cylindrical ends of Σ^+ are rational.
- If you compare two planar models of Σ and Σ^+ associated to ϵ_0 and η_0 , the x-coordinates for corresponding popsicle lines are exactly the same.
- If $z^+ \in \Sigma$ be a sprinkle on a disc which lies on a popsicle line coming out of $z_k \in \mathbb{R}$, then the image of a cylindrical end for z^+ is a disc of radius ρ_k . Here ρ_k is a radius of a strip-like end for z_k .
- If $w \in \Sigma^+$ is a sprinkle on a sphere which lies on a line corresponds to $z_k \in \mathbb{R}$, then the image of a rational cylindrical end for w is disc of radius ρ_k . Here ρ_k is a radius of a strip-like end for z_k .

If every ends on a broken popsicle satisfy these conditions, we say that choices of ends are **aligned**. See Figure 20.

One can choose the ends of a popsicle smoothly and consistently along the moduli space while the choices are all aligned. From now on, we assume that all choice of ends are always aligned.

Let us describe a corner chart for a point in $P_{n,\tilde{F},\phi,\star} \subset \overline{P}_{n,\tilde{F},\phi}$. As usual, a corner chart is described in terms of gluing parameters, but alignment of conformal structure forces us to align the gluing parameters as well. We will describe how to align gluing parameters according to the alignment data.

For a broken popsicle, we will select a collection of gluing parameters of outgoing marked points, one for each aligned disc/spheres pair.

FUKAYA CATEGORY FOR LANDAU-GINZBURG ORBIFOLDS




FIGURE 20. Example of aligned pairs with the choice of aligned ends

Describing a gluing procedure requires a little more work. We will first define, what we call, an **auxiliary model**, by adding suitable semi-stable discs and spheres in the following way.

- (1) (adding semi-stable discs) For a given data $\Psi_e : \Phi^{-1}(e) \to \{1, ..., m\}$, we add a string of *m* many semi-stable discs at the nodal point corresponding to the edge *e*. If $\Psi_e(w) = k$, the corresponding sphere component of *w* is aligned with *k*-th semi-stable disc component.
- (2) (adding semi-stable spheres) Let $w \in \text{Vertex}(T_{v,i,j})$ be a vertex connected to $v \in \text{Vertex}(T)$. If $\Phi(w)$ is adjacent to v (even after adding semi-stable discs), then we do nothing. If not, consider a unique path from v to $\Phi(w)$. Then we add semi-stable popsicle spheres (at the output of the component w) which are aligned to disc components along this path. Here, each added semi-stable popsicle sphere have one input and one output, and aligned to the corresponding disc component.

Similarly, let $w_1 < w_2$ are two adjacent vertices of $T_{v,i,j}$. If $\Phi(w_1)$ and $\Phi(w_2)$ are not adjacent, take a unique path from $\Phi(w_1)$ and $\Phi(w_2)$, and we add corresponding semistable popsicle spheres for disc components along this path. Here, each sphere has one input and one output, and aligned to the corresponding disc component.

We apply the same procedure to leaves. Let l be a leaf of $T_{v,i,j}$ attached to a vertex w with induced color m. Then we add a semi-stable sphere for each disc component along a unique path from $\Phi(w)$ to the leaf l_m . Again, each sphere component is aligned to a corresponding disc component.

(3) (choose aligned ends) We extend the choice of ends to added discs and spheres so that the overall choice is still aligned. Since we only add semi-stable components, we can copy-and-paste original choice of ends without modification.

Remark B.4. We record a minimal amount of information in the alignment data. Auxiliary model recovers the full alignment from the prescribed ones by adding suitable semi-stable discs and spheres. They are just tracking devices for aligned components and compatible choices of ends so that the gluing process is well-defined. For example, in the gluing model of the broken popsicle described in Figure 10, two disc neighborhoods of interior marked points as well as the upper-half of the annulus (between radius r_1r_2 and r_2) corresponds to the added semi-stable spheres and discs. The broken popsicle in Figure 9 does not require additional semi-stable components.

We call the resulting surface with decoration the auxiliary model for a broken popsicle of type \star . Every sphere component in the auxiliary model is aligned to a disc component. Moreover, if several spheres are aligned to a single disc, at least one of the disc or spheres are stable. Therefore, we can move back and forth between stable broken popsicles and its auxiliary model by simply forgetting (and recovering) semi-stable components.

The gluing process takes place in two stages. At first, we replace a broken popsicle to its auxiliary model. Second, we turn on gluing parameters and obtain a new auxiliary model. Finally, we contract semi-stable components to get a stable broken popsicle.

The process is well-defined in the following sense. In the auxiliary model, suppose that we are gluing Σ_1 and Σ_2 with a gluing parameter ϵ . If Σ'_1 is aligned to Σ_1 and Σ'_2 is aligned to Σ_2 , then our procedure glues two pairs (Σ_1, Σ_2) and (Σ'_1, Σ'_2) simultaneously. Since our choice of ends are compatible with respect to the alignment, they are still aligned after gluing.

Rational ends help us to put a canonical popsicle structure on a gluing of two. Let S_{γ} be a gluing of $S_1 \in P_{n_1,F_1}$ and $S_2 \in P_{n_2,F_2}$ along $z_{0,1} \in S_1$ and $z_{i,2} \in S_2$. Then positive boundary markings of S_{γ} are $(n_1 + n_2 - 1)$ points on $\partial \mathbb{H}$:

$$z_k \leftrightarrow x_{k,\gamma} = \begin{cases} x_{k,1} & k < i, \\ x_{i,1} + (\rho_{1,i} \times \gamma) x_{k-i+1,2} & i \le k \le i + n_2 - 1, \\ x_{k-n_2+1,1} & i + n_2 \le k. \end{cases}$$

Moreover, sprinkles are given as:

$$z_{l}^{+} \leftrightarrow x_{\phi(f),\gamma} + y_{f,\gamma} = \begin{cases} x_{\phi_{1}(f),1} + iy_{f} & f \in F_{1}, \ \phi_{1}(f) \neq i, \\ (x_{i,1} + (\rho_{1,i} \times \gamma)x_{\phi_{1}(f) - i + 1,2}) + iy_{f} & f \in F_{1} \ \phi_{1}(f) = i, \\ (x_{i,1} + (\rho_{1,i} \times \gamma)x_{\phi_{2}(f),2}) + iy_{\phi_{2}(f)} & f \in F_{2}. \end{cases}$$

Let us describe more details of the gluing, and how a combinatorial type of general popsicle changes afterward. Every finite edge $e \in \text{Edge}(\tilde{T})$ there is the unique vertex v such that $e = e_{v,0}$. We denote a gluing parameter associated to e by ϵ_v . Recall that if v_1 and v_2 are aligned to each other, then ϵ_{v_1} and ϵ_{v_2} must be turned on simultaneously. Therefore, we get two types of gluing parameters.

- (1) for each $v \in \text{Vertex}(T)$, we get a single parameter from a tuple $(\epsilon_v, \epsilon_{w_1}, \dots, \epsilon_{w_{n_v}})$ where $\{w_1, \dots, w_{n_v}\} = \Phi^{-1}(v)$. We simply denote it by ϵ_v .
- (2) for each $e \in \text{Edge}(T)$ such that $\Phi^{-1}(e) \neq \emptyset$, we get m_e -many parameters, one for each $1 \le k \le m_e$, from each tuple $(\epsilon_{w_1}, \dots, \epsilon_{w_{n_{e,k}}})$ where $\{w_1, \dots, w_{n_{e,k}}\} = \Psi_e^{-1}(k)$. We simply denote them by $\epsilon_{e,k}$.

Now let us see what happens on a combinatorial type when we turn on a single gluing parameter ϵ .

- (Case 1) : $\epsilon = \epsilon_v$ for some $v \in \text{Vertex}(T)$ and $\Phi^{-1}(e_{v,0}) = \emptyset$. Let us say the edge $e_{v,0}$ goes from u to v. Because of the second condition, popsicle discs Σ_u and Σ_v are still adjacent in the auxillary model. They are glued together if we turn on ϵ_v . It means that the edge $e_{v,0}$ is contracted and two vertices u and v become a single vertex v' after gluing. Now we have to see what happen to the vertices which are aligned to v.
 - (a) Suppose you see two vertices $w_1 < w_2 \in \text{Vertex}(T \setminus T)$ connected by an edge e' such that $\Phi(w_1) = u$ and $\Phi(w_2) = v$. Two components $\Sigma_{w_1}^+$ and $\Sigma_{w_2}^+$ will be glued together

with the same gluing parameter ϵ_v . For the trees, the edge e' is also contracted and two vertices w_i are replaced with a single vertex w'. Now, we align w' to v' for an obvious reason.

- (b) Suppose $\Phi(w) = v$, but there is no adjacent w' such that $\Phi(w') = u$. There will be no change in combinatorial type except $\Phi(w)$ becomes v' after gluing.
- (c) Suppose $\Phi(w) = u$, but there is no adjacent w' such that $\Phi(w') = v$. There will be no change in combinatorial type except $\Phi(w)$ becomes v' after gluing, similar as above.
- (d) Suppose you see $w \in \text{Vertex}(T_{u,i,j})$ such that $\Phi(w) = v$ and the incoming edge $e_{0,w}$ is actually the root of $T_{u,i,j}$ connected to u. It means Σ_w^+ and Σ_v are both adjacent to Σ_u , and they will be glued together to a single disc. Therefore the vertices u, v, w and the edge e and $e_{0,w}$ are contracted to a single vertex v'. Also, a tree $T_{u,i,j}$ becomes $T_{u,i,j} \setminus \{w, e_{0,w}\}$, which is a disjoint union of (val(w)-1)-many trees. A decomposition of $F_{u,i,j}$ and its F-label changes accordingly.
- (Case 2) : $\epsilon = \epsilon_v$ for some $v \in \text{Vertex}(T)$ and $\Phi^{-1}(e_{v,0}) \neq \emptyset$. Again, let us say that $e_{v,0}$ goes from u to v. The second condition implies that there will be a string of m_e -many semi-stable components in between Σ_u and Σ_v . Let w_1 be a stable sphere component that is aligned to the node $e_{v,0}$ and $\Psi_{e_{v,0}}(w_1) = m_e$. In the auxiliary model, we should have another sphere component (either stable or semi-stable) that is aligned with v, and we denote it by w_2 .

Turning on ϵ_v will glue Σ_v and the m_e -th semi-stable disc components (call the resulting component by v'), and the same parameter will also glue two sphere components $\Sigma_{w_1}^+$ and $\Sigma_{w_2}^+$ as well (call the resulting component by w'_1). For the resulting broken popsicle, the tree *T* remains the same, and now w'_1 should be aligned to v'. Note that m_e decrease by one.

(Case 3) : $\epsilon = \epsilon_{e,k}$ for some $e \in \text{Edge}(T)$ and $1 \le k \le m_e$. $\epsilon_{e,k}$ is a gluing parameter for the *k*-th and (k + 1)-th semi-stable disc components (say u_1 and u_2) attached at the node *e*. We can identify it with the corresponding gluing parameters for the aligned sphere components: Consider any two (semi-stable or stable) sphere components in the auxiliary model, say w_1 and w_2 that are aligned to u_1 and u_2 respectively. Turning on $\epsilon_{e,k}$, we glue u_1 and u_2 (call the resulting component u') and with the same parameter, we glue w_1 and w_2 (call the resulting component w'). Then, w' is now aligned to u'. This process will reduce m_e by one, and we need to reassign the function Ψ_e accordingly: for all the vertices w'' such that $\Phi(w'') = e$ and $\Psi_e(w'') \ge k + 1$, we lower the value $\Psi_e(w'')$ by one. This is because the function Ψ_e measures the relative order of the sphere bubbles. In this way, the rooted ribbon tree *T* for the resulting glued popsicle remains the same and only the associated alignment data is modified.

In any cases, a gluing procedure decreases either the number Vertex(T), or decreases a certain m_e by one. In fact, the following lemma can be obtained from the above description.

Lemma B.5. $\overline{P}_{n,\widetilde{F},\phi}$ is a manifold with corners of dimension $n-2+|\widetilde{F}|$, and

(B.1)
$$\operatorname{codim}(P_{n,\widetilde{F},\star}) = |\operatorname{Vertex}(T)| - 1 + \sum_{\rho} m_{e}$$

CHO, CHOA, AND JEONG

REFERENCES

- [AAE⁺13] Mohammed Abouzaid, Denis Auroux, Alexander Efimov, Ludmil Katzarkov, and Dmitri Orlov, *Homolog-ical mirror symmetry for punctured spheres*, Journal of the American Mathematical Society 26 (2013), no. 4, 1051–1083.
- [Abo10] Mohammed Abouzaid, *A geometric criterion for generating the Fukaya category*, Publications Mathématiques de l'IHÉS **112** (2010), 191–240.
- [Abo12] _____, *On the wrapped Fukaya category and based loops*, Journal of Symplectic Geometry **10** (2012), no. 1, 27–79.
- [ACHL20] Lino Amorim, Cheol-Hyun Cho, Hansol Hong, and Siu-Cheong Lau, *Big quantum cohomology of orbifold spheres*, arXiv preprint arXiv:2002.11180 (2020).
- [AGZV85] Vladimir Igorevich Arnold, Sabir Medgidovich Gusein-Zade, and Aleksandr Nikolaevich Varchenko, *Sin*gularities of differentiable maps. Vol. I, Monographs in Mathematics **82** (1985).
- [AGZV12] _____, Singularities of differentiable maps: Volume II monodromy and asymptotic integrals, vol. 83, Springer Science & Business Media, 2012.
- [AKO08] Denis Auroux, Ludmil Katzarkov, and Dmitri Orlov, *Mirror symmetry for weighted projective planes and their noncommutative deformations*, Ann. of Math. (2) **167** (2008), no. 3, 867–943. MR 2415388 (2009f:53142)
- [AS10] Mohammed Abouzaid and Paul Seidel, *An open string analogue of Viterbo functoriality*, Geometry & Topology **14** (2010), no. 2, 627–718.
- [Aur07] Denis Auroux, *Mirror symmetry and T-duality in the complement of an anticanonical divisor*, J. Gökova Geom. Topol. GGT 1 (2007), 51–91.
- [BH93] Per Berglund and Tristan Hübsch, *A generalized construction of mirror manifolds*, Nuclear Physics B **393** (1993), no. 1-2, 377–391.
- [BH95] Per Berglund and Måns Henningson, *Landau-Ginzburg orbifolds, mirror symmetry and the elliptic genus*, Nuclear Physics B **433** (1995), no. 2, 311–332.
- [Bou03] Frédéric Bourgeois, *A Morse-Bott approach to contact homology*, Symplectic and contact topology: interactions and perspectives (Toronto, ON/Montreal, QC, 2001) **35** (2003), 55–77.
- [CFH95] Kai Cieliebak, Andreas Floer, and Helmut Hofer, *Symplectic homology II*, Mathematische Zeitschrift **218** (1995), no. 1, 103–122.
- [CFHW96] Kai Cieliebak, Andreas Floer, Helmut Hofer, and Kris Wysocki, *Applications of symplectic homology II: Stability of the action spectrum*, Mathematische Zeitschrift **223** (1996), no. 1, 27–45.
- [CHL17] Cheol-Hyun Cho, Hansol Hong, and Siu-Cheong Lau, *Localized mirror functor for Lagrangian immersions, and homological mirror symmetry for* $\mathbb{P}^1_{a,b,c}$, J. Differential Geom. **106** (2017), no. 1, 45–126. MR 3640007
- [CL20] Cheol-Hyun Cho and Sangwook Lee, *Kodaira-Spencer map, Lagrangian Floer theory and orbifold Jacobian algebras*, arXiv preprint arXiv:2007.11732 (2020).
- [Dyc11] Tobias Dyckerhoff, *Compact generators in categories of matrix factorizations*, Duke Math. J. **159** (2011), no. 2, 223–274.
- [Ebe] Wolfgang Ebeling, *Homological mirror symmetry for singularities*, Representation Theory Current Trends and Perspectives, European Mathematical Society Publishing House, pp. 75–107.
- [Eis80] David Eisenbud, *Homological algebra on a complete intersection, with an application to group representations,* Transactions of the American Mathematical Society **260** (1980), no. 1, 35–64.
- [EP15] Alexander I Efimov and Leonid Positselski, *Coherent analogues of matrix factorizations and relative singularity categories*, Algebra & Number Theory **9** (2015), no. 5, 1159–1292.
- [FJR13] Huijun Fan, Tyler Jarvis, and Yongbin Ruan, *The Witten equation, mirror symmetry, and quantum singularity theory*, Annals of Mathematics (2013), 1–106.
- [FN62] Ralph Fox and Lee Neuwirth, *The braid groups*, Mathematica Scandinavica **10** (1962), 119–126.
- [FOOO16] Kenji Fukaya, Yong-Geun Oh, Hiroshi Ohta, and Kaoru Ono, *Lagrangian Floer theory and mirror sym*metry on compact toric manifolds, Astérisque (2016), no. 376, vi+340. MR 3460884
- [FU09] Masahiro Futaki and Kazushi Ueda, *Homological mirror symmetry for Brieskorn-Pham singularities*, Proceeding of the 56th Japan Geometry Symposium (2009), 98–107.

[FU11]	, Homological mirror symmetry for Brieskorn-Pham singularities, Selecta Mathematica 17 (2011),
[FU13]	no. 2, 435–452. , <i>Homological mirror symmetry for singularities of type d</i> , Mathematische Zeitschrift 273 (2013),
	no. 3-4, 633–652.
[Gan13]	Sheel Ganatra, <i>Symplectic cohomology and duality for the wrapped Fukaya category</i> , arXiv preprint arXiv:1304.7312 (2013).
[GJ94]	Ezra Getzler and John DS Jones, <i>Operads, homotopy algebra and iterated integrals for double loop spaces,</i> arXiv preprint hep-th/9403055 (1994).
[GZ21]	Fabio Gironella and Zhengyi Zhou, Exact orbifold fillings of contact manifolds, arXiv preprint
[[][[]]]	arXiv:2108.12247 (2021).
[HK12]	Claus Hertling and Ralf Kurbel, <i>On the classification of quasihomogeneous singularities</i> , Journal of Singularities 4 (2012), 131–153.
[HS19]	Matthew Habermann and Jack Smith, Homological Berglund-Hübsch mirror symmetry for curve singu-
[11010]	<i>larities</i> , arXiv preprint arXiv:1903.01351 (2019).
[Knö87]	Horst Knörrer, <i>Cohen-Macaulay modules on hypersurface singularities I</i> , Inventiones mathematicae 88 (1987), no. 1, 153–164.
[Kon95]	Maxim Kontsevich, <i>Homological algebra of mirror symmetry</i> , Proceedings of the International Congress
[101100]	of Mathematicians, Vol. 1, 2 (Zürich, 1994) (Basel), Birkhäuser, 1995, pp. 120–139.
[Kra09]	Marc Krawitz, FJRW rings and Landau-Ginzburg mirror symmetry, arXiv preprint arXiv:0906.0796
	(2009).
[KST07]	Hiroshige Kajiura, Kyoji Saito, and Atsushi Takahashi, Matrix factorizations and representations of quiv-
	ers II: type ADE case, Advances in Mathematics 211 (2007), no. 1, 327–362.
[KvK16]	Myeonggi Kwon and Otto van Koert, Brieskorn manifolds in contact topology, Bulletin of the London
	Mathematical Society 48 (2016), no. 2, 173–241.
[LU18]	Yanki Lekili and Kazushi Ueda, <i>Homological mirror symmetry for K3 surfaces via moduli of</i> A_{∞} - <i>structures</i> , arXiv preprint arXiv:1806.04345 (2018).
[LU20]	, Homological mirror symmetry for Milnor fibers of simple singularities, arXiv preprint arXiv:2004.07374 (2020).
[McL16]	Mark McLean, Reeb orbits and the minimal discrepancy of an isolated singularity, Inventiones mathe-
	maticae 204 (2016), no. 2, 505–594.
[Mil68]	John Milnor, Singular points of complex hypersurfaces, no. 61, Princeton University Press, 1968.
[Orl09]	Dmitri Orlov, <i>Derived categories of coherent sheaves and triangulated categories of singularities</i> , Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II, Progr. Math., vol. 270, Birkhäuser Boston Inc., Boston, MA, 2009, pp. 503–531.
[Pre11]	Anatoly Preygel, Thom-Sebastiani & duality for matrix factorizations, arXiv preprint arXiv:1101.5834
	(2011).
[Sei00]	Paul Seidel, <i>Graded Lagrangian submanifolds</i> , Bulletin de la Société Mathématique de France 128 (2000), no. 1, 103–149.
[Sei01]	, <i>More about vanishing cycles and mutation</i> , Symplectic geometry and mirror symmetry (Seoul, 2000) (2001), 429–465.
[Sei06]	, <i>A biased view of symplectic cohomology</i> , Current developments in mathematics 2006 (2006), no. 1, 211–254.
[Sei08]	, Fukaya categories and Picard-Lefschetz theory, Zurich Lectures in Advanced Mathematics, Eu-
[Sei11]	ropean Mathematical Society (EMS), Zürich, 2008. MR 2441780 , <i>Homological mirror symmetry for the genus two curve</i> , J. Algebraic Geom. 20 (2011), no. 4, 727–
(0, 1) = 1	769. Here is the interview of the second sec
[Sei15]	, Homological mirror symmetry for the quartic surface, Mem. Amer. Math. Soc. 236 (2015), no. 1116, vi+129. MR 3364859
[Sei18]	, Fukaya A_{∞} -structures associated to Lefschetz fibrations. VI, arXiv preprint arXiv:1810.07119 (2018).
[Tak]	Atsushi. Takahashi, Weighted projective lines associated to regular systems of weights of dual type, Adv. Stud. Pure Math 59 .

CHO, CHOA, AND JEONG

- [Vit99] Claude Viterbo, *Functors and computations in Floer homology with applications, I*, Geometric & Functional Analysis GAFA **9** (1999), no. 5, 985–1033.
- [Vor00] Alexander A Voronov, *Homotopy gerstenhaber algebras*, Conférence Moshé Flato 1999, Springer, 2000, pp. 307–331.
- [Yos90] Yuji Yoshino, *Maximal Cohen-Macaulay modules over Cohen-Macaulay rings*, vol. 146, Cambridge University Press, 1990.

DEPARTMENT OF MATHEMATICAL SCIENCES, RESEARCH INSTITUTE IN MATHEMATICS, SEOUL NATIONAL UNIVER-SITY, SEOUL, SOUTH KOREA

Email address: chocheol@snu.ac.kr

KOREA INSTITUTE OF ADVANCED STUDIES, DONGDAEMOON-GU, SEOUL, SOUTH KOREA

Email address: dwchoa@kias.re.kr

DEPARTMENT OF MATHEMATICAL SCIENCES, RESEARCH INSTITUTE IN MATHEMATICS, SEOUL NATIONAL UNIVER-SITY, GWANAK-GU, SEOUL, SOUTH KOREA

Email address: wonbo.jeong@gmail.com

68