

Optimal building block of multipartite quantum battery

Tian-Ran Yang,¹ Yu-Yu Zhang,^{1,2,*} Hui Dong,² Libin Fu,² and Xiaoguang Wang^{3,†}

¹*Department of Physics, Chongqing University, Chongqing 401330, China*

²*Graduate School, China Academy of Engineering Physics, Beijing 100193, China*

³*Zhejiang Institute of Modern Physics, Department of Physics, Zhejiang University, Hangzhou 310027, China*

(Dated: October 27, 2021)

To take quantum advantage of collective effects in many-body system, we design an elementary block for building multipartite quantum battery, which enables charging an atomic ensemble with optimal numbers in a common thermal bath. One achieves maximum free energy as the stored energy in the steady state, which is prior to each atom parallel charging independently. It ascribes to quantum collective effects in the ensemble of atoms induced by the competition between the coherent driving and decoherent dissipation. The corresponding thermodynamic efficiency of the energy storage is analyzed. The existence of the optimal elementary units of multipartite quantum battery provide a guideline for designing a realizable charging scheme.

I. INTRODUCTION

Recently, diligent efforts are devoted to explore the possibility of taking advantage of quantum resources to achieve superior performances in the energy conversion and storage with the control achievement on multipartite quantum system [1–3]. Quantum battery (QB) is a quantum system for storing energy supplied by an external source. The battery exploits quantum effects for efficient charging in comparison to its classical counterpart [4–7]. A renewed effort is devoted to enhance charging of multipartite batteries in a closed system as a consequence of quantum correlations in many-body systems, which is known as collective effects [8–11].

When a multipartite battery is subjected to a common thermal bath, it gives rise to interesting phenomenons such as an increasing entropy, which establishes a link to quantum thermodynamics [12–15]. An emergence of collective effects in quantum thermodynamics like the free energy as the stored energy and work extraction are attractive quantum phenomenon in open systems, while the influence of thermodynamics on such quantum effects are overlooked. Many efforts have been devoted to investigate the QB for the energy storage in the thermal environment using different charging protocols [16–18]. More recently, a dissipative charging process of a battery was suggested [16], in which an efficient thermodynamic equilibrium state is approached by work extraction under cyclic unitary operations with help of auxiliary systems. This engineering needs to find a unitary evolution for the battery and the auxiliary system with a globally conserved quantity, which is controlled by a post selected driving agent.

A harmonic driving as an external source has been proposed as an alternative powerful charging field due to the tunable driving frequency for maximal stored energy [9]. Inspired by the advantage of the collective effects and the

harmonic driving, a multipartite QB in a common thermal bath, collectively coupled to a harmonic driving field, is an attractive battery model for optimal energy storage. In such many-body system, the interplay between the coherent driving and decoherent charging induced by the thermal dissipation plays an essential role in the cooperative many-body effects for the charging. The question is that whether such collective effects in many-body system be harnessed to improve thermodynamically meaningful features in the driven-dissipative charging protocols.

In this paper, we present the irreversible thermodynamic charging of an ensemble of N two-level atoms charged by a harmonic driving field in a common thermal bath. It is different from previous entropy-preserving or energy-conserving charging process under unitary operations. With the increment of atoms, the quantum collective effects lead to a non-monotonic behavior of free energy and a decreasing entropy per atom, which are induced by the competition between the coherent driving force and the decoherent dissipation of the common thermal bath. We find the optimal number of atoms as an elementary unit of the QB interacting with a common thermal bath, which results in maximizing the stored energy per atom. Engineering such optimal-atom battery as a building block one achieves more free energy by compared to parallel charging for N independent atoms. Meanwhile, thermodynamic efficiency of the energy storage in terms of the work done by the external charging field is analyzed.

II. A PROTOCOL FOR A MULTIPARTITE QB CHARGING

We consider an open charging system of a multipartite battery, which consists of two-level atoms coupled to an external driving as a charger to transfer energy. Fig. 1 (a) shows normal parallel charging strategy with independent thermal bath. Our charging protocol focus on the elementary building block (blue box), illustrated in Fig. 1 (b), with finite number of atoms with shared thermal bath. Atoms in each units are collectively charged by

*Electronic address: yuyuzh@cqu.edu.cn

†Electronic address: xgwang1208@zju.edu.cn




FIG. 1: (a) Charging protocol of N two-level atoms in parallel. Each atom is charged by a harmonic driving and couples to a thermal environment. (b) The elementary building block (blue box) consists of a few atoms with a common thermal bath. During the charging time, the QB interacts with a harmonic driving field $A \cos(\omega t)$.

a harmonic field. The total Hamiltonian consists of the QB-system part and the interacting part as $H = H_s + H_I$ with

$$H_s = \omega_0 J_z, H_I = A \cos(\omega t) J_x, \quad (1)$$

where $J_\alpha = \sum_i^N \sigma_i^\alpha / 2$ ($\alpha = x, y, z$) is the collective operator of N two-level atoms with the energy level splitting ω_0 . A and ω are the driving amplitude and the modulated frequency.

In the practical application, the QB are coupled also to the environment, which is modeled as the thermodynamics dissipation. The temporal evolution of the density matrix $\rho(t)$ for the QB is described by the master equation,

$$\begin{aligned} \frac{d\rho(t)}{dt} = & -i[\omega_0 J_z + A \cos(\omega t) J_x, \rho] \\ & + \gamma[n(T) + 1](2J_- \rho J_+ - \{J_+ J_-, \rho\}) \\ & + \gamma n(T)(2J_+ \rho J_- - \{J_- J_+, \rho\}), \end{aligned} \quad (2)$$

where $n(T) = [e^{\omega/(k_B T)} - 1]^{-1}$ is the mean occupation number of bath mode with the frequency ω at the bath temperature T . The Boltzmann's constant is set by $k_B = 1$ for later discussion. The first term on the right side describes the normal-parallel charging for N independent atoms. The basis set for representing the QB system of N atoms is the Dicke states $|J, l - N/2\rangle$ ($l = 0, 1, \dots, N$), which are eigenstates of J^2 and J_z with the total pseudospin $J = N/2$. Both external driving and the dissipation induce the state transitions among energy levels of the Dicke states $|N/2, l - N/2\rangle$ with $l = 0, 1, \dots, N$.

III. AN OPTIMAL ELEMENTARY UNIT FOR CHARGING

The importance is to define the usable stored energy. In the normal charging without thermal environment, the

internal energy change of QB can be utilized in the later retraction. Yet, only part of the internal energy can be extracted in the charging process within the thermal environment. For such evaluation, we have considered the entropy $S(\rho_s) = -\text{Tr}(\rho_s \ln \rho_s)$ of the QB system. The useful energy stored in the QB is measured by the free energy

$$F(\rho_s) = E(\rho_s) - k_B T S(\rho_s), \quad (3)$$

where $E(\rho_s) = \text{Tr}(H_s \rho_s)$ is the internal energy of the QB system. The difference in free energy $\Delta F = \Delta E - k_B T \Delta S$ measures the useful energy stored in the QB. At zero temperature $T = 0$, the free energy change is equivalent to the mean energy, $\Delta F(\rho_s) = \Delta E(\rho_s)$. Here we only consider the situation with one thermal bath during the charging and later retraction process with the same temperature. The similar definition of the useful work is well discussed in the quantum thermodynamic resource discussions [19].

Initially, N atoms decouple with the charging field, and the QB system is prepared in the Gibbs thermal state of N atoms, $\rho_0 = e^{-H_s/(k_B T)}/Z$ with the partition function $Z = \text{Tr}[e^{-H_s/(k_B T)}]$. For the single-atom QB, the temporal mean energy $E(\rho_s) = \omega_0 \langle \sigma_z \rangle / 2$ is obtained analytically as (see the appendix)

$$\begin{aligned} E(\rho_s)/\omega_0 = & -\frac{\gamma^2 \chi}{\gamma^2 \chi^2 + A^2/2} \left\{ 1 - \frac{e^{-3t\gamma\chi/2}}{2\chi\gamma^2} \right. \\ & \left\{ [(2\chi\gamma^2(1 + \alpha\chi) + \alpha A^2) \cos(\Omega t) \right. \\ & \left. \left. - \frac{\gamma}{2\Omega} [2\gamma^2 \chi^2(1 + \alpha\chi) + A^2(4 + \alpha\chi)] \sin(\Omega t) \right] \right\}, \end{aligned} \quad (4)$$

where the oscillation Rabi frequency is $\Omega = \sqrt{A^2 - \gamma^2 \chi^2 / 4}$ with $\chi = [1 + 2n(T)]$. The mean energy E of the QB becomes larger as the driving amplitude A increases or the dissipative rate γ decreases. At the zero temperature with $n(T) = 0$, when the driving strength A becomes larger than the dissipative rate γ , E tends to be zero in the steady state and the corresponding stored energy ΔE equals to $\omega_0/2$. (See Appendix A)

For the large value of N , we use the Holshtein-Primakoff transformation in terms of auxiliary bosonic operators b^\dagger and b : $J_z = b^\dagger b - N/2$, $J_+ = b^\dagger \sqrt{N}$ and $J_- = b \sqrt{N}$. It gives $d\rho(t)/dt = -i[\omega_0(b^\dagger b - N/2) + A\sqrt{N}\cos(\omega t)(b^\dagger + b), \rho] + \gamma N[n(T) + 1](2b\rho b^\dagger - \{b^\dagger b, \rho\}) + \gamma N n(T)(2b^\dagger \rho b - \{bb^\dagger, \rho\})$. The driving strength is proportional to $A\sqrt{N}$, while the decay rate is proportional to γN . Due to the competition between the external driving force and the thermal dissipation, one would expect the existence of the efficient elementary unit of the QB with optimal number of atoms for maximizing the stored energy.

Due to the difficulty of finding analytical results for the multipartite batteries, we present dynamic charging process via the free energy and the entropy by numerical calculations. Fig. 2(a) shows oscillations of the scaled free energy change $\Delta F/N$, which exhibits a stable value at the steady state. The charging period for




FIG. 2: (a) Variance of the scaled free energy $\Delta F/(N\omega_0)$ and (b) variance of the entropy $\Delta S/N$ as a function of the charging time t for different $N = 1, 10$ and 28 with $A = 0.5\omega$ and $n(T) = 0.2$. (c) $\Delta F/\omega_0$ in the steady state as a function of N for different dissipation rate $\gamma = 0, 0.03\omega, 0.05\omega$ and 0.1ω with $A = 0.5\omega$.

the steady state becomes shorter as the atom number N increases, because the system relaxes to the steady state rapidly due to the dominated dissipation. Interesting, the stored free energy in the steady state increases firstly with the incremental of N , then decreases for $N = 28$. It demonstrates the existence of the optimal number N_{op} of atoms to achieve maximum free energy. Meanwhile, it emerges inevitably an increasing of entropy in the dissipative charging process, which is different from the preserved entropy in a closed system. The corresponding change of the scaled entropy $\Delta S/N$ decrease as N increases in Fig. 2(b). It means that the heat flow induced by the entropy $Q_s = T\Delta S$ becomes lower than that obtained by N atoms charging independently in parallel in Fig. 1(a). It exhibits an improvement over the parallel charging of N atoms.

One expects a steady state with larger population in high-level states to achieve maximal stored energy. N atoms with a strong driving amplitude A would result in the occupation of the higher-energy states $|N/2, l - N/2\rangle$ with a larger value of l , yet the occupation of the lower-energy states with the increasing of dissipation γ . As a consequence of such competition between the coherent external driving and dissipation, an efficient elementary unit of the QB with the optimal atoms is predicted to achieve maximum free energy per atom. Fig. 2(c) shows that the non-monotonic dependence of the free energy ΔF on the number of atoms N in the steady state, exhibiting an maximum value at an optimal number N_{op} of atoms. Especially, in the absence of the dissipation $\gamma = 0$, the free energy is proportional to N , $\Delta F \propto N$,




FIG. 3: Work done W , free energy ΔF , variance of the internal energy $\Delta\langle H\rangle$, and heat Q as a function of time t for $N = 1$ quantum battery for different dissipation rate $\gamma = 0$ (a) and $\gamma = 0.03\omega$ (b). (c) The efficiency η as a function of time t for different number of atoms $N = 1, 17$ and 25 with $\gamma = 0.03\omega$. The parameters are $A = 0.5\omega$ and $n(T) = 0.2$.

which is consistent with previous results [9, 10]. For different dissipation rate γ and the driving strength A , the efficient elementary unit of the QB consists of the optimal number N_{op} of atoms in a common thermal bath in Fig. 1(b), which has maximum stored energy by comparing to results of atoms charging independently in parallel.

IV. CHARGING EFFICIENCY

In the charging process, the stored free energy is an important quantity to evaluate the performance. However, we have shown the energy from the agent dissipates into the thermal bath. The extent, to which the work done by the agent is stored, is also important and typically evaluated as the efficiency. We quantify the efficiency of the charging process by the ratio

$$\eta = \frac{\Delta F}{W}, \quad (5)$$

where the amount of the work done on the QB by the external driving field is given by

$$W = \text{Tr} \left(\int_0^\tau \frac{dH}{dt} \rho dt \right) = -A\omega \int_0^\tau \sin(\omega t) \text{Tr}(J_x \rho) dt. \quad (6)$$

Due to the thermal dissipation of the irreversible process, some amount of work W is transferred into heat Q flowing into the thermal bath, which is measured by $Q = \text{Tr}(\int_0^\tau \dot{\rho} H dt)$. The intrinsic energy of the charging system including the interactions driving part H_I is measured as $\langle H \rangle = \text{Tr}[H(\rho(t))] = E + \langle H_I \rangle$, which is different from the mean energy E of the QB system.

Fig. 3 shows behaviors of the work W done by the external driving force, the heat flowing to the common bath Q . For a closed system with $\gamma = 0$ in Fig. 3(a), the work W is fully transferred into the variance of the intrinsic energy $\Delta\langle H \rangle$ periodically without heat dissipation. One interesting finding is that the variance of the intrinsic energy $\Delta\langle H \rangle$ sometimes is lower than the variance of free energy $\Delta F = \Delta E$ due to the negative value of the interacting energy $\Delta\langle H_I \rangle < 0$, resulting in the efficiency $\eta > 1$. Such efficiency is caused by the energy exchange between the coherent driving field and the QB. For the dissipation case with $\gamma = 0.03\omega$ in Fig. 3(b), the laws of thermodynamics, $W = Q + \Delta H$, holds in the open charging system as well as the closed system. The corresponding efficiency η decreases as the number N of atoms for an elementary unit of the QB increases in Fig. 3(c). The efficiency for the efficient elementary unit with the optimal number of atoms $N_{op} = 17$ is worse than that of single-battery case, which ascribes to heat dissipation.

V. CONCLUSION

We have shown the driven-dissipative charging of the multipartite battery comprising an ensemble of two-level atoms, which interact with an external harmonic driving and a common thermal bath. The stored energy in the battery is measured by the free energy due to increasing of the entropy in the open charging system. Due to the competition between the coherent driving and the decoherent dissipation in atoms system, the free energy per atom in the steady state behaves non-monotonically dependent on the number of the atoms. We find the optimal

elementary unit of the QB with finite atoms in a common thermal bath, which provides maximal average stored energy. To character the thermodynamic efficiency of the charging process, we elucidate the work done by the external field as well as the intrinsic energy of the QB system. We prove the first law of thermodynamics holds in the open charging system. The quantum thermodynamics in the open charging system involves quantum collective effects induced by quantum coherence in many-body system in comparison to classical thermodynamics, resulting in optimal building block of multipartite quantum battery. Our results is a fundamental attributions to the powerful energy storage with optimal atoms for physically realizable charging schemes.

Acknowledgments

This work was supported by the NSFC under Grants No.12075040, the Chongqing Research Program of Basic Research and Frontier Technology under Grants No.cstc2020jcyj-msxmX0890, and by the Fundamental Research Funds for the Central Universities under Grant No. 2020CDJ-LHZZ-012. H. Dong acknowledges the support from the NSFC (Grant No.11875049) and NSAF (Grant No.U1930403). L. B. Fu acknowledges the support from the NSFC(Grant No.11725417) and NSAF (Grant No.U1940403). X. G. Wang acknowledges supports from the NSFC (Grant No.11875231) and the National Key Research and Development Program of China (Contracts No. 2017YFA0304202 and No. 2017YFA0205700).

-
- [1] P. Strasberg, G. Schaller, T. Brandes, and M. Esposito, *Phys. Rev. X* **7**, 021003 (2017).
 - [2] G. Watanabe, B. P. Venkatesh, P. Talkner, and A. del Campo, *Phys. Rev. Lett.* **118**, 050601 (2017).
 - [3] M. Campisi, P. Hänggi, and P. Talkner, *Rev. Mod. Phys.* **83**, 1653 (2011).
 - [4] K. V. Hovhannisyan, M. Perarnau-Llobet, and A. Acín, *Phys. Rev. Lett.* **111**, 240401 (2013).
 - [5] G. M. Andolina, M. Keck, A. Mari, M. Campisi, V. Giovannetti, and M. Polini, *Phys. Rev. Lett.* **122**, 047702 (2019).
 - [6] N. Friis, M. Huber, and M. Perarnau-Llobet, *Phys. Rev. E* **93**, 042135 (2016).
 - [7] N. Friis, and M. Huber, *Quantum* **2**, 61 (2018).
 - [8] F. Campaioli, F. A. Pollock, F. C. Binder, L. Cleri, J. Goold, S. Vinjanampathy, and K. Modi, *Phys. Rev. Lett.* **118**, 150601 (2017).
 - [9] Y. Y. Zhang, T. R. Yang, L. B. Fu, and X. G. Wang, *Phys. Rev. E* **99**, 052106 (2019).
 - [10] D. Ferraro, M. Campisi, G. M. Andolina, V. Pellegrini, and M. Polini, *Phys. Rev. Lett.* **120**, 117702 (2018).
 - [11] F. Binder, S. Vinjanampathy, K. Modi, and J. Goold, *New. J. Phys.* **17**, 075015 (2015).
 - [12] P. Faist, F. Dupuis, J. Oppenheim, and R. Renner, *Nat. Commun.* **6**, 7669 (2015).
 - [13] P. Skrzypack, A. J. Short, and S. Popescu, *Nat. Commun.* **5**, 4185 (2014).
 - [14] M. N. Bera, A. Riera, M. Lewenstein, and A. Winter, *Nat. Commun.* **8**, 2180 (2017).
 - [15] Y. H. Ma, R. X. Zhai, C. P. Sun, and H. Dong, *arXiv*: 1910.13434.
 - [16] F. Barra, *Phys. Rev. Lett.* **122**, 210601 (2019).
 - [17] D. Farina, G. M. Andolina, A. Mari, M. Polini, and V. Giovannetti, *Phys. Rev. B* **99**, 035421 (2019).
 - [18] F. Pirmoradian, and K. Mølmer, *Phys. Rev. A* **100**, 043833 (2019).
 - [19] R. Alicki, *J. Phys. A Math. Gen.* **12**, L103-L107 (1979).




FIG. A.1: Scaled mean energy E/ω_0 obtained by the analytical solution (dashed black line) as a function of τ for one-atom quantum battery with different driving strength (a)(c) $A/\omega_0 = 0.05$, and (b)(d) $A/\omega_0 = 0.5$ at temperature $n(T) = 0$ and 1. The driving frequency is chosen as $\omega = \omega_0 = 2$, and the dissipation rate is $\gamma = 0.01\omega_0$. The numerical results are listed for comparison (solid red line).

Appendix A: Analytical solutions for single-atom battery

We analyze the thermodynamics of a single-atom battery charging process with the master equation. The two-level system evolution involves energy transfer from the external driving field and the dissipation of the energy into the thermal environment.

The master equation of the single-atom battery system becomes

$$\begin{aligned} d\rho_s/dt = & -i\frac{\omega_0}{2}[\sigma_z, \rho_s] - i\frac{A}{4}(e^{i\omega t} + e^{-i\omega t})[\sigma_x, \rho_s] \\ & + \gamma[n(T) + 1](2\sigma_- \rho_s \sigma_+ - \{\sigma_+ \sigma_-, \rho_s\}) \\ & + \gamma n(T)(2\sigma_+ \rho_s \sigma_- - \{\sigma_- \sigma_+, \rho_s\}). \end{aligned} \quad (\text{A1})$$

We perform a rotating-frame transformation using $U = \exp(i\omega t \sigma_z/2)$ to give

$$\begin{aligned} d\rho'_s/dt = & -i\frac{\omega_0 - \omega}{2}[\sigma_z, \rho'_s] \\ & - i\frac{A}{4}(e^{i\omega t} + e^{-i\omega t})[e^{i\omega t} \sigma_+ + e^{-i\omega t} \sigma_-, \rho'_s] \\ & + \gamma[n(T) + 1](2\sigma_- \rho'_s \sigma_+ - \{\sigma_+ \sigma_-, \rho'_s\}) \\ & + \gamma n(T)(2\sigma_+ \rho'_s \sigma_- - \{\sigma_- \sigma_+, \rho'_s\}), \end{aligned} \quad (\text{A2})$$

where $\rho'_s = U \rho_s U^\dagger$. When the driving strength A is much smaller than the two-level energy ω_0 on resonance with the QB system $\omega_0 = \omega$, it is reasonable to making a rotating-wave approximation (RWA) by ignoring fast

oscillating terms. The Bloch equations are derived as

$$\langle \dot{\sigma}_z \rangle_\tau = i\frac{A}{2}(\langle \sigma_- \rangle_\tau - \langle \sigma_+ \rangle_\tau) - 2\gamma[\langle \sigma_z \rangle_\tau(2n(T) + 1) + 1], \quad (\text{A3})$$

$$\langle \dot{\sigma}_+ \rangle_\tau = i\frac{A}{4}\langle \sigma_z \rangle_\tau - \gamma\langle \sigma_- \rangle_\tau(2n(T) + 1), \quad (\text{A4})$$

$$\langle \dot{\sigma}_- \rangle_\tau = -i\frac{A}{4}\langle \sigma_z \rangle_\tau - \gamma\langle \sigma_+ \rangle_\tau(2n(T) + 1). \quad (\text{A5})$$

For the initial Gibbs state with $\alpha = Tr[\sigma_z \rho(0)]$, it gives analytically

$$\begin{aligned} \langle \sigma_z \rangle_\tau = & -\frac{2\gamma^2\eta}{2\gamma^2\eta^2 + A^2} \left\{ 1 - e^{-3\tau\gamma\eta/2} \frac{1}{2\eta\gamma^2} \right. \\ & \left. \{ [(2\eta\gamma^2(1 + \alpha\eta) + \alpha A^2) \cos(\Omega\tau) \right. \\ & \left. - \frac{\gamma[2\gamma^2\eta^2(1 + \alpha\eta) + A^2(4 + \alpha\eta)]}{2\Omega} \sin(\Omega\tau)] \right\} \end{aligned} \quad (\text{A6})$$

where $\eta = [1 + 2n(T)]$, and the oscillation Rabi frequency is

$$\Omega = \sqrt{A^2 - \frac{\gamma^2\eta^2}{4}}. \quad (\text{A7})$$

Especially, at zero temperature with $n(T) = 0$ for $\langle \sigma_z \rangle = -1$, $\langle \sigma_- \rangle = 0$, $\langle \sigma_+ \rangle = 0$, the corresponding temporal energy $E(\rho_s) = \omega_0 \langle \sigma_z \rangle_\tau / 2$ is obtained

$$E(\rho_s) = \frac{-4\omega_0\gamma^2}{8\gamma^2 + A^2} \left[1 + \frac{A^2}{8\gamma^2} e^{-3\gamma\tau/2} (\cos \Omega\tau + \frac{3\gamma}{2\Omega} \sin \Omega\tau) \right], \quad (\text{A8})$$

with $\Omega = \sqrt{A^2 - \gamma^2/4}$. When the driving strength A becomes larger than the dissipative rate γ , $\langle \sigma_z \rangle_\infty$ tends to be zero and the stored energy ΔE equals to $\omega_0/2$.

We calculate $E(\rho_s)$ according to Eq.(A1) numerically without the RWA for the dissipative charging process in Fig. A.1. The analytical solutions in Eq.(A8) agree well with numerical ones for $A/\omega_0 = 0.05$ and 0.5. And the oscillation frequency of the mean energy are correctly captured by the analytical one Ω dependent on the driving strength A and dissipative rate γ . As A increases to be much larger than γ , $A = 0.5\omega_0$, $E(\rho_s)$ increases to be 0 in the steady state in Fig. A.1 (b), which is consistent with the analytical ones from Eq.(A8). For a thermal bath with $n(T) = 1$, excellent agreement between the analytical results and the numerical ones is observed in Fig. A.1 (c) and (d). As the temperature increases, the oscillation Rabi frequency Ω becomes smaller in Eq.(A7) due to the stronger thermal dissipation. It demonstrates that the charging time to reach the steady state becomes shorter at finite temperature, and the oscillation Rabi frequency of the QB energy E can be modulated by the driving strength A , the loss rate γ and the temperature T .

Fig. A.2 shows the thermodynamics charging controlled by the temperature. Since the decay rate of the




FIG. A.2: (a) Scaled free energy $\Delta F/\omega_0$ and (b) the Von Neumann entropy ΔS of one-atom quantum battery for different temperature $n(T) = 0.1, 1$ with dissipation rate $\gamma = 0.05\omega$. (c) Scaled free energy $\Delta F/\omega_0$ and (d) the Von Neumann entropy for different dissipative rate $\gamma = 0$ and 0.05ω with $n(T) = 0.2$. The driving frequency is chosen as $\omega = \omega_0 = 2$, and the driving strength $A/\omega_0 = 0.5$.

two-level atom is proportional to the rate $\gamma[n(T)+1]$ and $\gamma n(T)$ in Eq.(A1). As $n(T)$ increases, the mean energy of the QB system and the free energy decay rapidly as in Fig. A.2(a). The corresponding variance of the entropy ΔS decreases in Fig. A.2(b). The increasing of the temperature induces more energy flowing to the environment, while it reduces the information flow. In the entropy non-preserving process, the heat flow to the thermal bath decreases of the energy of the QB system due to the entropy production. In the absence of the dissipation $\gamma = 0$, it exhibits periodic oscillation of the free energy ΔF and the entropy is conserved in Fig. A.2(c) (d).