

A conjecture of Cameron and Kiyota on sharp characters with prescribed values

A. Abdollahi^{a,c}, J. Bagherian^{a1}, M. Khatami^a, Z. Shahbazi^a, R. Sobhani^b

^a Department of Pure Mathematics, Faculty of Mathematics and Statistics, University of Isfahan, Isfahan 81746-73441, Iran

^b Department of Applied Mathematics and Computer Science, Faculty of Mathematics and Statistics, University of Isfahan, Isfahan 81746-73441, Iran

^c School of Mathematics, Institute for Research in Fundamental Sciences (IPM), 19395-5746 Tehran, Iran

Emails: a.abdollahi@sci.ui.ac.ir, bagherian@sci.ui.ac.ir, m.khatami@sci.ui.ac.ir, z.shahbazi@sci.ui.ac.ir, r.sobhani@sci.ui.ac.ir

Abstract

Let χ be a virtual (generalized) character of a finite group G and $L = L(\chi)$ be the image of χ on $G - \{1\}$. The pair (G, χ) is said to be sharp of type L if $|G| = \prod_{l \in L} (\chi(1) - l)$. If the principal character of G is not an irreducible constituent of χ , the pair (G, χ) is called normalized. In this paper, we first provide some counterexamples to a conjecture that was proposed by Cameron and Kiyota in 1988. This conjecture states that if (G, χ) is sharp and $|L| \geq 2$, then the inner product $(\chi, \chi)_G$ is uniquely determined by L . We then prove that this conjecture is true in the case that (G, χ) is normalized, χ is a character of G , and L contains at least an irrational value.

Keywords: Character value; Finite group; Sharp character; Sharp pair.

MSC(2010): 20C15.

1 Introduction

Let G be a finite group and χ be a virtual (generalized) character of G . We define

$$L(\chi) := \{\chi(g) \mid 1 \neq g \in G\}$$

and

$$Sh(\chi) = \prod_{l \in L(\chi)} (\chi(1) - l).$$

¹Corresponding author

It is known that $Sh(\chi)$ is a multiple of $|G|$; see [2]. The pair (G, χ) (or briefly, the virtual character χ) is called sharp of type L or L -sharp if $L = L(\chi)$ and $Sh(\chi) = |G|$. Clearly, χ is faithful whenever (G, χ) is L -sharp. The pair (G, χ) is said to be normalized if $(\chi, 1_G)_G = 0$, where 1_G is the principal character of G .

The notion of sharpness was first introduced for permutation characters by Ito and Kiyota as a generalization of sharply t -transitive permutation representations [4]. Then, Cameron and Kiyota extended the definition of sharp pairs to what is given above and posed the following conjecture [3].

Conjecture 1: If (G, χ) is L -sharp with $|L| \geq 2$, then $(\chi, \chi)_G$ is uniquely determined by L .

As a main result in this paper, we provide some counterexamples to the above conjecture. In Section 2, we give a group G and two characters χ and θ of G such that the pairs (G, χ) and (G, θ) are L -sharp, but $(\chi, \chi)_G \neq (\theta, \theta)_G$ (see Example 2.1). Moreover, we show that if (G, χ) in Conjecture 1 is assumed to be normalized, then there are some counterexamples which show $(\chi, \chi)_G$ is not uniquely determined by L (see Example 2.2). It should be emphasized that we can not find any counterexample to Conjecture 1 in the case that (G, χ) is normalized and χ is a character. It seems that this conjecture is true whenever (G, χ) is normalized and χ is a character. Therefore, we give an improvement of Conjecture 1 as follows.

Conjecture 2: If (G, χ) is L -sharp and normalized where χ is a character and $|L| \geq 2$, then $(\chi, \chi)_G$ is uniquely determined by L .

In Section 3, we show that Conjecture 2 is true whenever L contains at least an irrational value. To do this, we use a characterization of L -sharp pairs, given in [1], when L contains at least an irrational value.

2 L -Sharp characters with different numbers of irreducible constituents

In this section, we first give a group G with two characters χ and θ such that (G, χ) and (G, θ) are L -sharp and $(\chi, \chi)_G \neq (\theta, \theta)_G$ (see Example 2.1). This shows that

Conjecture 1 is not true, in general.

Example 2.1. Let G be the group `SmallGroup(32,6)` in GAP [5]. Put $\chi = \chi_1 + 2\chi_2 + \chi_5$. Then as we can see in Table 1, χ is sharp of type $L = \{-1, 3\}$ and $(\chi, \chi)_G = 6$. On the other hand, set $\theta = \chi_2 + \chi_3 + \chi_4 + \chi_5$. Then θ is also sharp of type $L = \{-1, 3\}$ such that $(\theta, \theta)_G = 4$. But we have $(\chi, \chi)_G \neq (\theta, \theta)_G$.

$ Cl_G(g) $	1	4	4	2	2	1	4	4	4	2	4
χ_1	1	1	1	1	1	1	1	1	1	1	1
χ_2	1	-1	1	1	1	1	-1	-1	1	1	-1
χ_3	1	1	-1	1	1	1	-1	1	-1	1	-1
χ_4	1	-1	-1	1	1	1	1	-1	-1	1	1
χ_5	4	0	0	0	0	-4	0	0	0	0	0

Table 1: Sharp characters $\chi = \chi_1 + 2\chi_2 + \chi_5$ and $\theta = \chi_2 + \chi_3 + \chi_4 + \chi_5$

The example below shows that Conjecture 1 is not true if we further assume that (G, χ) is normalized.

Example 2.2. Let G be the group `SmallGroup(192,1494)` in GAP [5]. Put $\chi = \chi_2 + \chi_5$. Then χ is sharp of type $L = \{-2, 0, 2\}$ and $(\chi, \chi)_G = 2$; see Table 2. On the other hand, set $\theta = \chi_3 + \chi_4 - \chi_1$. Then θ is also sharp of type $L = \{-2, 0, 2\}$ such that $(\theta, \theta)_G = 3$. Clearly, $(\chi, \chi)_G \neq (\theta, \theta)_G$.

$ Cl_G(g) $	1	24	32	6	6	1	12	24	32	12	6	24	12
χ_1	1	-1	1	1	1	1	-1	-1	1	1	1	-1	-1
χ_2	2	0	-1	2	2	2	0	0	-1	2	2	0	0
χ_3	3	1	0	-1	-1	3	-1	-1	0	-1	3	1	-1
χ_4	4	0	1	0	0	-4	-2	0	-1	0	0	0	2
χ_5	4	0	1	0	0	-4	2	0	-1	0	0	0	-2

Table 2: Sharp characters $\chi = \chi_2 + \chi_5$ and $\theta = \chi_3 + \chi_4 - \chi_1$

In the following, we give some remarks related to Conjecture 2.

Remark 2.3. *We can express a strong version of Conjecture 2 as follows:*

If (G, χ) and (H, θ) are L -sharp and normalized such that χ and θ are characters and $|L| \geq 2$, then $(\chi, \chi)_G = (\theta, \theta)_H$.

By Example 2.4 below, we show that the above conjecture is not true, in general. In fact, there exist non-isomorphic groups G and H with L -sharp characters χ and θ with $|L| \geq 2$ such that $(\chi, \chi)_G \neq (\theta, \theta)_H$. Moreover, if we further assume that $|G| = |H|$, then there exists a counterexample to the above conjecture; see Example 2.5.

Example 2.4. Let G be the alternating group A_7 of degree 7. Using the character table of G , one can check that G has an irreducible sharp character χ of type $L = \{-1, 0, 2\}$; see Table 3. On the other hand, let H be the dihedral group D_{12} of order 12. Then H has the sharp character $\theta = \chi_2 + \chi_6$ of type $L = \{-1, 0, 2\}$; see Table 4. Clearly, in this case $(\chi, \chi)_G \neq (\theta, \theta)_H$.

$ Cl_G(g) $	1	105	70	210	280	630	504	360	360
χ	14	2	2	2	-1	0	-1	0	0

Table 3: Sharp character χ

$ Cl_G(g) $	1	1	3	3	2	2
χ_1	1	1	1	1	1	1
χ_2	1	1	-1	-1	1	1
χ_3	1	-1	-1	1	1	-1
χ_4	1	-1	1	-1	1	-1
χ_5	2	2	0	0	-1	-1
χ_6	2	-2	0	0	-1	1

Table 4: Character table of D_{12}

Example 2.5. Let G be the group `SmallGroup(192,955)` in GAP [5]. Then as we can see in Table 5, G has an irreducible sharp character χ of type $L = \{-2, 0, 2\}$. On the other hand, let H be the group `SmallGroup(192,1494)` in GAP [5]. Using

$ Cl_G(g) $	1	12	4	32	6	3	12	24	12	32	12	6	24	12
χ	6	-2	0	0	2	-2	0	0	2	0	0	-2	0	0

Table 5: Sharp character χ

the character table of H , $\theta = \theta_1 + \theta_2$ is a sharp character of type $L = \{-2, 0, 2\}$ (see Table 6). Clearly, in this case $(\chi, \chi)_G \neq (\theta, \theta)_H$.

$ Cl_H(h) $	1	24	32	6	6	1	12	24	32	12	6	24	12
θ_1	2	0	-1	2	2	2	0	0	-1	2	2	0	0
θ_2	4	0	1	0	0	-4	2	0	-1	0	0	0	-2

Table 6: Sharp character $\theta = \theta_1 + \theta_2$

3 Sharp characters with at least an irrational value

In this section, we show that if (G, χ) is L -sharp and normalized such that χ is a character and L contains at least an irrational value, then Conjecture 2 holds. Our key tool is the following result given by Alvis and Nozawa in [1].

For the remainder of this paper, whenever (G, χ) is sharp, we assume that χ is a character of G .

Theorem 3.1. (*[1, Theorem 1.3]*) *Suppose (G, χ) is L -sharp and normalized such that L contains at least an irrational value. Then one of the following holds.*

- (i) *G is cyclic of order m , and either $m \geq 3$ and χ is linear, or else $m \geq 5$ and χ is the sum of two complex conjugate linear characters of G .*
- (ii) *G is dihedral of order $2m$, where $m \geq 5$ is odd, and χ is irreducible of degree 2.*
- (iii) *G is dihedral or generalized quaternion of order $2m$, where $m \geq 8$ is even, and $\chi = \psi$ or $\chi = \psi + \varepsilon$, where ψ is irreducible of degree 2 and ε is linear with cyclic kernel of order m .*
- (iv) *G is isomorphic to the binary octahedral group and χ is irreducible of degree 2.*
- (v) *G is isomorphic to $SL(2, 5)$ and χ is irreducible of degree 2.*
- (vi) *G is isomorphic to A_5 and χ is irreducible of degree 3.*

The next lemmas give some results about the sharp characters of cyclic groups, dihedral groups, and generalized quaternion groups which we need in the sequel.

Lemma 3.2. ([3, Proposition 1.8]) Let G be a cyclic group of order m and λ be a faithful linear character of G . Then (G, λ) is sharp of type

$$L = \{\omega^r \mid 1 \leq r \leq m-1\}$$

where $\omega = e^{2\pi i/m}$.

Lemma 3.3. Let G be a cyclic group of order m , where $m \geq 5$ is odd, and let λ be a faithful linear character of G . Then $(G, \lambda + \bar{\lambda})$ is a sharp pair of type $L = \{\omega^r + \omega^{-r} \mid 1 \leq r \leq (m-1)/2\}$, where $\omega = e^{2\pi i/m}$.

Proof. Set $\chi = \lambda + \bar{\lambda}$. First we show that χ is faithful. Let $L = \{\chi(g) \mid 1 \neq g \in G\}$. Then $L = \{2 \cos(2\pi r/m) \mid 1 \leq r \leq (m-1)/2\}$. If there exists $1 \leq r \leq (m-1)/2$ such that $2 \cos(2\pi r/m) = 2$, then $2\pi r/m = 2k\pi$ for some $k \in \mathbb{Z}$, a contradiction. Hence χ is faithful. Moreover,

$$\begin{aligned} \prod_{r=1}^{(m-1)/2} \left(2 - 2 \cos\left(\frac{2\pi r}{m}\right)\right) &= \prod_{r=1}^{(m-1)/2} 4 \sin^2\left(\frac{\pi r}{m}\right) \\ &= \prod_{r=1}^{(m-1)/2} 2 \sin\left(\frac{\pi r}{m}\right) \prod_{r=1}^{(m-1)/2} 2 \sin\left(\frac{\pi r}{m}\right) \\ &= \prod_{r=1}^{(m-1)/2} 2 \sin\left(\frac{\pi r}{m}\right) \prod_{r=1}^{(m-1)/2} 2 \sin\left(\frac{(m-r)\pi}{m}\right) \\ &= \prod_{r=1}^{(m-1)/2} 2 \sin\left(\frac{\pi r}{m}\right) \prod_{j=(m+1)/2}^{m-1} 2 \sin\left(\frac{\pi j}{m}\right) \\ &= \prod_{r=1}^{m-1} 2 \sin\left(\frac{\pi r}{m}\right) = m, \end{aligned}$$

(See [7, Lemma 3.1]). This shows that χ is a sharp character, as desired. ■

Lemma 3.4. Let D_{2m} be the dihedral group of order $2m$, where m is odd, and let χ be a faithful irreducible character of D_{2m} of degree 2. Then (D_{2m}, χ) is a sharp pair of type $L = \{0, \omega^r + \omega^{-r} \mid 1 \leq r \leq (m-1)/2\}$, where $\omega = e^{2\pi i/m}$.

Proof. See the proof of Theorem 3.2 in [7]. ■

Lemma 3.5. Let G be a dihedral or generalized quaternion group of order $2m$ where m is even and let $\omega = e^{2\pi i/m}$. Then

(1) if $m/4 \in \mathbb{Z}$ and ψ is a faithful irreducible character of degree 2, then (G, ψ) is a sharp pair of type $L = \{-2, 0, \omega^r + \omega^{-r} \mid 1 \leq r \leq m/2 - 1\}$;

(2) if ψ is a faithful irreducible character of degree 2 and ε is linear with cyclic kernel of order m , then $(G, \psi + \varepsilon)$ is a sharp pair of type

$$L = \{-1, 1 + \omega^r + \omega^{-r} \mid 1 \leq r \leq m/2 - 1\}.$$

Proof. Assume that $m = 2t$ for some integer t . If $G = D_{2m}$, then

$$G = \langle a, b : a^m = b^2 = 1, b^{-1}ab = a^{-1} \rangle$$

and it is known that Table 7 is the character table of G where

$$\{1\}, \{a^t\}, \{a^r, a^{-r}\} \text{ for } 1 \leq r \leq t-1, \{a^s b \mid s \text{ even}\}, \{a^s b \mid s \text{ odd}\}$$

are the conjugacy classes of G . Similarly, if $G = Q_{2m}$ is the generalized quaternion

$Cl_G(g)$	1	a^t	$a^r (1 \leq r \leq t-1)$	b	ab
$ Cl_G(g) $	1	1	2	$m/2$	$m/2$
χ_1	1	1	1	1	1
χ_2	1	1	1	-1	-1
χ_3	1	$(-1)^t$	$(-1)^r$	1	-1
χ_4	1	$(-1)^t$	$(-1)^r$	-1	1
ψ_j	2	$2(-1)^j$	$\omega^{jr} + \omega^{-jr}$	0	0
$(1 \leq j \leq t-1)$					

Table 7: Character table of D_{2m} (m even)

group of order $2m$, then

$$G = \langle a, b : a^{2t} = 1, a^t = b^2, b^{-1}ab = a^{-1} \rangle.$$

In this case

$$\{1\}, \{a^t\}, \{a^r, a^{-r}\}$$

for $1 \leq r \leq t-1$, and

$$\{a^{2r}b \mid 0 \leq r \leq t-1\}, \{a^{2r+1}b \mid 0 \leq r \leq t-1\}$$

are the conjugacy classes of G and the character tables of the groups Q_{4t} (t odd) and Q_{4t} (t even) are given in Table 8 and Table 9, respectively; (see [6]). Now suppose that G is one of the two groups D_{2m} or Q_{2m} , (m even). We show that the irreducible character ψ_1 is a sharp character of G of type

$$L = \{-2, 0, \omega^r + \omega^{-r} \mid 1 \leq r \leq m/2 - 1\}.$$

It is easy to see that ψ_1 is faithful and we have

$$\begin{aligned} \prod_{r=1}^{m/2-1} \left(2 - 2 \cos\left(\frac{2\pi r}{m}\right)\right) &= \prod_{r=1}^{m/2-1} 4 \sin^2\left(\frac{\pi r}{m}\right) \\ &= \prod_{r=1}^{m/2-1} 2 \sin\left(\frac{\pi r}{m}\right) \prod_{r=1}^{m/2-1} 2 \sin\left(\frac{\pi r}{m}\right) \\ &= \prod_{r=1}^{m/2-1} 2 \sin\left(\frac{\pi r}{m}\right) \prod_{r=1}^{m/2-1} 2 \sin\left(\frac{(m-r)\pi}{m}\right) \\ &= \prod_{r=1}^{m/2-1} 2 \sin\left(\frac{\pi r}{m}\right) \prod_{j=m/2+1}^{m-1} 2 \sin\left(\frac{\pi j}{m}\right) \\ &= \frac{\prod_{r=1}^{m-1} 2 \sin\left(\frac{\pi r}{m}\right)}{2 \sin\left(\frac{\pi}{2}\right)} = \frac{m}{2}, \end{aligned}$$

(See [7, Lemma 3.1]). Since for $r = m/4$, we have $2 \cos(\frac{2\pi r}{m}) = 0$ it follows that $\prod_{l \in L(\psi_1)} (2 - l) = 2m$ and hence ψ_1 is a sharp character of G of type

$$L = \{-2, 0, \omega^r + \omega^{-r} \mid 1 \leq r \leq m/2 - 1\}.$$

$Cl_G(g)$	1	a^t	$a^r (1 \leq r \leq t-1)$	b	ab
$ Cl_G(g) $	1	1	2	$m/2$	$m/2$
χ_1	1	1	1	1	1
χ_2	1	1	1	-1	-1
χ_3	1	-1	$(-1)^r$	i	$-i$
χ_4	1	-1	$(-1)^r$	$-i$	i
ψ_j	2	$2(-1)^j$	$\omega^{jr} + \omega^{-jr}$	0	0
$(1 \leq j \leq t-1)$					

Table 8: Character table of Q_{4t} (t odd)

$Cl_G(g)$	1	a^t	$a^r (1 \leq r \leq t-1)$	b	ab
$ Cl_G(g) $	1	1	2	$m/2$	$m/2$
χ_1	1	1	1	1	1
χ_2	1	1	1	-1	-1
χ_3	1	1	$(-1)^r$	1	-1
χ_4	1	1	$(-1)^r$	-1	1
ψ_j	2	$2(-1)^j$	$\omega^{jr} + \omega^{-jr}$	0	0
$(1 \leq j \leq t-1)$					

Table 9: Character table of Q_{4t} (t even)

Now we consider irreducible character ψ_j where $2 \leq j \leq t-1$. Then one can see that ψ_j is faithful if and only if j is odd and $(j, m) = 1$. In this case we have $L(\psi_1) = L(\psi_j)$ and the argument above shows that ψ_j is a sharp character for G of type $L = \{-2, 0, \omega^r + \omega^{-r} \mid 1 \leq r \leq m/2 - 1\}$.

Similarly, we can show that $\chi_2 + \psi_j$, where $1 \leq j \leq t-1$ is odd and $(j, m) = 1$, is a sharp character of type $L = \{-1, 1 + \omega^r + \omega^{-r} \mid 1 \leq r \leq m/2 - 1\}$ for G . ■

For the remainder of this paper we put $\omega = e^{2\pi i/m}$ and we assume that

- (a) $L_1 = \{\omega^r \mid 1 \leq r \leq m-1\}$ such that $m \geq 3$;
- (b) $L_2 = \{\omega^r + \omega^{-r} \mid 1 \leq r \leq (m-1)/2\}$ such that $m \geq 5$ is odd;
- (c) $L_3 = \{0, \omega^r + \omega^{-r} \mid 1 \leq r \leq (m-1)/2\}$ such that $m \geq 5$ is odd;
- (d) $L_4 = \{-2, 0, \omega^r + \omega^{-r} \mid 1 \leq r \leq m/2 - 1\}$ such that $m \geq 8$ is even;
- (e) $L_5 = \{-1, 1 + \omega^r + \omega^{-r} \mid 1 \leq r \leq m/2 - 1\}$ such that $m \geq 8$ is even;
- (f) $L_6 = \{-2, -1, 0, 1, \sqrt{2}, -\sqrt{2}\}$;
- (g) $L_7 = \{-2, -1, 0, 1, \frac{1+\sqrt{5}}{2}, \frac{1-\sqrt{5}}{2}, \frac{-1+\sqrt{5}}{2}, \frac{-1-\sqrt{5}}{2}\}$;

$$(h) \ L_8 = \{-1, 0, \frac{1+\sqrt{5}}{2}, \frac{1-\sqrt{5}}{2}\}.$$

Then we can see that $L_i \cap \mathbb{Z} \subseteq \{-2, -1, 0, 1, 2\}$ for $1 \leq i \leq 5$. In the next lemmas for cyclic groups, dihedral groups, and generalized quaternion groups we determine $L_i \cap \mathbb{Z}$ whenever (G, χ) is L_i -sharp.

Lemma 3.6. *Let G be a cyclic group of order m and (G, χ) be L -sharp with $L = L_1$ or $L = L_2$, then $L \cap \mathbb{Z} \subseteq \{-1\}$.*

Proof. First let $L = L_1$. Since L_1 is the set of m th roots of unity (distinct from 1), it is clear that $L_1 \cap \mathbb{Z} \subseteq \{-1\}$.

Now suppose that $L = L_2$. Since m is odd, it is easy to see that $-2, 0, 1, 2 \notin L_2$. Moreover, $\omega^r + \omega^{-r} = -1$ if and only if $m/3 \in \mathbb{Z}$. So $L_2 \cap \mathbb{Z} \subseteq \{-1\}$. ■

Lemma 3.7. *Let G be a dihedral group of order $2m$, ($m \geq 5$ odd) and (G, χ) be L -sharp with $L = L_3$, then $\{0\} \subseteq L \cap \mathbb{Z} \subseteq \{-1, 0\}$.*

Proof. Since $0 \in L_3 \cap \mathbb{Z}$, it suffices to show that $-2, 1, 2 \notin L_3$. If $-2, 1 \in L_3$, then we obtain a contradiction under assumption $m \geq 5$ is odd. Moreover, $r \leq (m-1)/2$ shows that $2 \notin L_3$. Note that $\omega^r + \omega^{-r} = -1$ if and only if $m/3$ is an integer. Therefore if $m/3$ is integer, then $L \cap \mathbb{Z} = \{-1, 0\}$. ■

Lemma 3.8. *Let G be a dihedral or generalized quaternion group of order $2m$, ($m \geq 8$ even) and (G, χ) be L -sharp with $L = L_4$, then $L \cap \mathbb{Z} = \{-2, 0\}$ or $L \cap \mathbb{Z} = \{-2, -1, 0, 1\}$.*

Proof. Obviously, $-2, 0 \in L_4$. Since $r \leq m/2 - 1$, we have $2 \notin L_4$. It is easy to see that $-1, 1 \in L_4$ if and only if $m/3$ is an integer. Hence $L_4 \cap \mathbb{Z} = \{-2, 0\}$ or $L_4 \cap \mathbb{Z} = \{-2, -1, 0, 1\}$. ■

Lemma 3.9. *Let G be a dihedral or generalized quaternion group of order $2m$, ($m \geq 8$ even) and (G, χ) be L -sharp with $L = L_5$. Then $L \cap \mathbb{Z} \in \{\{-1\}, \{-1, 1\}, \{-1, 0, 2\}, \{-1, 0, 1, 2\}\}$.*

Proof. It follows from the assumption that $-1 \in L_5$ and $-2 \notin L_5$. Since m is even, we have $0, 2 \in L_5$ if and only if $m/3$ is an integer. Moreover, $1 \in L_5$ if and only if $r = m/4$. Therefore $L \cap \mathbb{Z}$ is equal to $\{-1\}$, $\{-1, 1\}$, $\{-1, 0, 2\}$ or $\{-1, 0, 1, 2\}$. ■

Theorem 3.10. *Suppose (G, χ) is L -sharp and normalized such that L contains at least an irrational value. If $L \subseteq \mathbb{R}$ and*

$$L \cap \mathbb{Z} \in \{\emptyset, \{-1\}, \{-1, 1\}, \{-1, 0, 2\}, \{-1, 0, 1, 2\}\},$$

then $(\chi, \chi)_G = 2$. Otherwise $(\chi, \chi)_G = 1$.

Proof. First suppose that $L \subseteq \mathbb{R}$ and

$$L \cap \mathbb{Z} \in \{\emptyset, \{-1\}, \{-1, 1\}, \{-1, 0, 2\}, \{-1, 0, 1, 2\}\}.$$

Then it follows from Theorem 3.1 and Lemmas 3.6, 3.7, 3.8, and 3.9 that

- (1) either G is a cyclic group of odd order and χ is the sum of two complex conjugate linear characters of G ;
- (2) or G is a dihedral or generalized quaternion group of order $2m$ ($m \geq 8$ even) and χ is the sum of an irreducible character of degree 2 and a linear character with cyclic kernel of order m .

So in each of these two cases we have $(\chi, \chi)_G = 2$.

Now we consider the other cases as follows.

- (i) If $L \not\subseteq \mathbb{R}$, then by Theorem 3.1 and Lemmas 3.2, 3.3, 3.4, and 3.5, G is a cyclic group and χ is linear.
- (ii) If $L \subseteq \mathbb{R}$ and $L \cap \mathbb{Z} \notin \{\emptyset, \{-1\}, \{-1, 1\}, \{-1, 0, 2\}, \{-1, 0, 1, 2\}\}$, then by Theorem 3.1 and Lemmas 3.6, 3.7, 3.8, and 3.9, we have one of the following cases:
 - (a) G is dihedral of order $2m$, where $m \geq 5$ is odd, and χ is irreducible of degree 2.
 - (b) G is dihedral or generalized quaternion of order $2m$, where $m \geq 8$ is even, and χ is irreducible of degree 2.
 - (c) G is isomorphic to the binary octahedral group and χ is irreducible sharp character of degree 2 and type $L = L_6$.
 - (d) G is isomorphic to $SL(2, 5)$ and χ is irreducible sharp character of degree 2 and type $L = L_7$.

- (e) G is isomorphic to A_5 and χ is irreducible sharp character of degree 3 and type $L = L_8$.

As we see in the cases (i) and (ii) above, the sharp character χ is irreducible and so we have $(\chi, \chi)_G = 1$. ■

Now we are ready to state and prove the main result of this section.

Theorem 3.11. *If (G, χ) is L -sharp and normalized with $|L| \geq 2$ and L contains at least an irrational value, then $(\chi, \chi)_G$ is uniquely determined by L .*

Proof. It follows from Theorem 3.10 that if (G, χ) is L -sharp with $|L| \geq 2$ and L contains at least an irrational value, then $(\chi, \chi)_G \in \{1, 2\}$ and it can be uniquely determined by L . ■

Remark 3.12. *The example below shows that the pair (G, χ) in Theorem 3.11 must be normalized as a necessary condition.*

Example 3.13. Let G be the dihedral group D_{16} . Then $\chi = \chi_3 + 2\chi_1$ is a sharp character of type $\{2, 0, \sqrt{2} + 2, -\sqrt{2} + 2\}$; see Table 10. On the other hand, $\theta = \chi_2 + \chi_3 + \chi_1$ is also a sharp character of type $\{2, 0, \sqrt{2} + 2, -\sqrt{2} + 2\}$. But we have $(\chi, \chi)_G \neq (\theta, \theta)_G$.

$ Cl_G(g) $	1	4	2	2	1	4	2
χ_1	1	1	1	1	1	1	1
χ_2	1	-1	1	1	1	-1	1
χ_3	2	0	$\sqrt{2}$	0	-2	0	$-\sqrt{2}$

Table 10: Sharp characters $\chi = \chi_3 + 2\chi_1$ and $\theta = \chi_2 + \chi_3 + \chi_1$

References

- [1] D. Alvis and S. Nozawa, Sharp characters with irrational values, J. Math. Soc. Japan, 48(3) (1996) 567–591.
- [2] H. F. Blichfeldt, A theorem concerning the invariants of linear homogeneous groups, with some applications to substitution-groups, Trans. Amer. Math. Soc. 5 (1904) 461–466.

- [3] P. J. Cameron and M. Kiyota, Sharp characters of finite groups, *J. Algebra*, 115(1) (1988) 125–143.
- [4] T. Ito and M. Kiyota, Sharp permutation groups, *J. Math. Soc. Japan* 33 (1981) 435–444.
- [5] The GAP Group, GAP – Groups, Algorithms, and Programming, Version 4.8.8; 2017. (<http://www.gap-system.org>)
- [6] G. James and M. Liebeck, *Representations and Characters of Groups*, Cambridge Mathematical Textbooks, Cambridge, 1995.
- [7] E. Poimenidou and A. Cottrell, Total characters of dihedral groups and sharpness, *Missouri J. Math Sci.* 12(1) (2000) 12–25.