Complemented subspaces of polynomial ideals

Sergio A. Pérez *

Universidad Pedagógica y Tecnológica de Colombia Email:Sergio.2060@hotmail.com

January 28, 2022

Abstract

Given the polynomial ideal $\mathcal{J} \circ \mathcal{P}({}^{n}E; F)$, we prove that if $\mathcal{J} \circ \mathcal{P}({}^{n}E; F)$ contains an isomorphic copy of c_0 , then $\mathcal{J} \circ \mathcal{P}({}^{n}E; F)$ is not complemented in $\mathcal{P}({}^{n}E; F)$ for every closed operator ideal $\mathcal{J} \subset \mathcal{L}_K$ and every $n \in \mathbb{N}$. Likewise we show that if $(\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}({}^{n}E; F)$ contains an isomorphic copy of c_0 , then $(\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}({}^{n}E; F)$ is not complemented in $\mathcal{P}({}^{n}E; F)$ for every closed operator ideal $\mathcal{J} \subset \mathcal{L}_K$ and every n > 1. When $\mathcal{J} = \mathcal{L}_K$, these results generalizes results of several authors [2],[13],[16],[18],[22], among others.

1 Introduction

The problem of establishing sufficient conditions for the complementation of the subspace of compact linear operators $\mathcal{L}_K(E; F)$ in the space $\mathcal{L}(E; F)$ of all continuous linear operators, has been widely studied by many authors. For example, see Bator and Lewis [2], Emmanuelle [13], John [15], Kalton [16] and Ghenciu [18], among others.

Emmanuele [13] and John [15] showed that if c_0 embeds in $\mathcal{L}_K(E; F)$ then $\mathcal{L}_K(E; F)$ is not complemented in $\mathcal{L}(E; F)$ for every E and F infinite dimensional Banach spaces.

John [15] proved that if E and F are arbitrary Banach spaces and $T : E \to F$ is a non compact operator which admits a factorization $T = A \circ B$ through a Banach space G with an unconditional basis, then the subspace $\mathcal{L}_K(E; F)$ of compact operators contains an isomorphic copy of c_0 and thus $\mathcal{L}_K(E; F)$ is not complemented in $\mathcal{L}(E; F)$. John [15] also proved that if E and F are infinite dimensional Banach spaces, such that each non compact operator $T \in \mathcal{L}(E; F)$ factors through a Banach space G with an unconditional basis, then the following conditions are equivalent:

- 1. $\mathcal{L}_K(E;F) = \mathcal{L}(E;F).$
- 2. $\mathcal{L}(E;F)$ contains no copy of ℓ_{∞} .
- 3. $\mathcal{L}_K(E; F)$ contains no copy of c_0 .
- 4. $\mathcal{L}_K(E; F)$ is complemented in $\mathcal{L}(E; F)$.

Ghenciu [18] obtained the following result:

Theorem 1.1. ([18, Theorem 1]) Let E and F be Banach spaces, and let G be a Banach space with an unconditional basis (g_n) and coordinate functionals (g'_n) .

(a) If there exist operators $R \in \mathcal{L}(G; F)$ and $S \in \mathcal{L}(E; G)$ such that $(R(g_n))$ is a semi-normalized basic sequence in F and $(S'(g'_n))$ is not relatively compact in E', then $\mathcal{L}_K(E; F)$ is not complemented in $\mathcal{L}(E; F)$.

^{*}S.Pérez was supported by Minciencias, Colombia. (corresponding author)

(b) If there exist operators $R \in \mathcal{L}(G; F)$ and $S \in \mathcal{L}(E; G)$ such that $(R(g_n))$ is a semi-normalized basic sequence in F and $(S'(g'_n))$ is not relatively weakly compact in E', then the space of all weakly compact linear operators $\mathcal{L}_{wK}(E; F)$ is not complemented in $\mathcal{L}(E; F)$.

This result generalizes results of several authors [12], [2], [14]. Clearly Theorem 1.1 (*a*) follows from results of Emmanuele [13] and John [15] previously aforementioned.

We consider the space $P_w({}^{n}E;F)$ of all n-homogeneous polynomials from E into F which are weakly continuous on bounded sets. When n = 1 we have that $P_w({}^{n}E;F) = \mathcal{L}_K(E;F)$.

Pérez [22] proved the polynomial versions of the previously results, among the most prominent theorems we find the following:

Theorem 1.2. Let E and F be Banach spaces, and let G be a Banach space with an unconditional basis (g_n) and coordinate functionals (g'_n) . If there exist operators $R \in \mathcal{L}(G; F)$ and $S \in \mathcal{L}(E; G)$ such that $(R(g_n))$ is a semi-normalized basic sequence in F and $(S'(g'_n))$ is not relatively compact in E', then $\mathcal{P}_w(^nE; F)$ is not complemented in $\mathcal{P}(^nE; F)$ for every $n \in \mathbb{N}$.

Theorem 1.3. Let E be an infinite dimensional Banach space and n > 1. If $\mathcal{P}_w(^nE;F)$ contains a copy of c_0 , then $\mathcal{P}_w(^nE;F)$ is not complemented in $\mathcal{P}(^nE;F)$.

Theorem 1.4. Let E and F be Banach spaces, with E infinite dimensional, and let n > 1. If each $P \in \mathcal{P}(^{n}E; F)$ such that $P \notin \mathcal{P}_{w}(^{n}E; F)$ admits a factorization $P = Q \circ T$, where $T \in \mathcal{L}(E; G)$, $Q \in \mathcal{P}(^{n}G; F)$ and G is a Banach space with an unconditional finite dimensional expansion of the identity, then the following conditions are equivalent:

- (1) $\mathcal{P}_w(^{n}E;F)$ contains a copy of c_0 .
- (1') $\mathcal{P}_K(^{n}E;F)$ contains a copy of c_0 .
- (2) $\mathcal{P}_w(^{n}E;F)$ is not complemented in $\mathcal{P}(^{n}E;F)$.
- (2') $\mathcal{P}_K(^{n}E;F)$ is not complemented in $\mathcal{P}(^{n}E;F)$.
- (3) $\mathcal{P}_w(^nE;F) \neq \mathcal{P}(^nE;F).$
- (3') $\mathcal{P}_K(^{n}E;F) \neq \mathcal{P}(^{n}E;F).$
- (4) $\mathcal{P}(^{n}E;F)$ contains a copy of c_{0} .
- (5) $\mathcal{P}(^{n}E;F)$ contains a copy of ℓ_{∞} .

In this paper, we obtain polynomial ideal versions of the preceding results.

2 Preliminaries

Let E and F denote Banach spaces over \mathbb{K} , where \mathbb{K} is \mathbb{R} or \mathbb{C} . Let E' denote the dual of E. Let B_E denote the unit ball of E. By J_E we mean the canonical embedding from E to E''. We denote by (e_n) the canonical basis of c_0 . Denote by $\mathcal{L}(E;F)$, $\mathcal{L}_K(E;F)$, $\mathcal{F}(E;F)$, $\mathcal{K}_p(E;F)$, $\mathcal{QN}_p(E;F)$ and $\mathcal{L}_{wK}(E;F)$, respectively, the spaces of all bounded, all compact, all finite range, all p-compact, all quasi p-nuclear and all weakly compact linear operators from E into F. Let $\mathcal{P}(^nE;F)$ denote the Banach space of all continuous n-homogeneous polynomials from E into F. We omit F when $F = \mathbb{K}$. Given a continuous n-homogeneous polynomial $P : E \to F$, by \check{P} we mean the (unique) symmetric n- linear operator such that $\check{P}(x, \ldots, x) = P(x)$ for every $x \in E$. Let $\mathcal{P}_w(^nE;F)$ denote the subspace of all $P \in \mathcal{P}(^nE;F)$ which are weakly continuous on bounded sets, that is the restriction

 $P|_B: B \to F$ is continuous for each bounded set $B \subset E$, when B and F are endowed with the weak topology and the norm topology, respectively. Let $\mathcal{P}_K(^nE;F)$ denote the subspace of all $P \in \mathcal{P}(^nE;F)$ which map bounded sets onto relatively compact sets. Let $\mathcal{P}_{wK}(^{n}E;F)$ denote the subspace of all $P \in \mathcal{P}(^{n}E;F)$ which map bounded sets onto relatively weakly compact sets.

We always have the inclusions

$$\mathcal{P}_w(^{n}E;F) \subset \mathcal{P}_K(^{n}E;F) \subset \mathcal{P}_{wK}(^{n}E;F) \subset \mathcal{P}(^{n}E;F),$$

as can be derived from results in [1]. We refer to [21] for background information on the theory of polynomials on Banach spaces.

By $E_1 \hat{\otimes}_{\pi} \dots \hat{\otimes}_{\pi} E_n$ we denote the completed projective tensor product of E_1, \dots, E_n . If $E_1 = \dots = E_n = E$ we write $\hat{\otimes}_{n,\pi} E$. By $\hat{\otimes}_{n,s,\pi} E$ we denote the *n*-fold completed symmetric tensor product of *E*.

E is isomorphic to a complemented subspace of F if and only if there are $A \in \mathcal{L}(E;F)$ and $B \in \mathcal{L}(F;E)$ such that $B \circ A = I$. E is said to have an unconditional finite dimensional expansion of the identity if there is a sequence of bounded linear operators $A_n: E \to E$ of finite rank, such that for $x \in E$

$$\sum_{n=1}^{\infty} A_n(x) = x$$

unconditionally.

We will say that the series $\sum_{n=1}^{\infty} x_n$ of elements of E is weakly unconditionally Cauchy if $\sum_{n=1}^{\infty} |x'(x_n)| < \infty$ for $x' \in E'$ or convict whether if all $x' \in E'$ or, equivalently if

$$\sup\left\{\left\|\sum_{n\in F}x_n\right\|; F\subset\mathbb{N}, Ffinite\right\}<\infty.$$

A sequence $(x_n) \subset E$ is a semi-normalized basic sequence if (x_n) is a Schauder basis for the closed subspace $M = [x_n : n \in \mathbb{N}]$, and moreover there are constant a and b such that $0 < a < ||x_n|| < b$ for all $n \in \mathbb{N}$. We denote by (e_n) the canonical basis of c_0 . If Σ is an algebra of subsets of a set Ω , then a finitely additive vector measure $\mu: \Sigma \to E$ is said to be strongly additive if the series $\sum_{n=1}^{\infty} \mu(A_n)$ converges in norm for each sequence (A_n) of pairwise disjoint members of Σ . The Diestel-Faires Theorem (see [11, p. 20, Theorem 2]) asserts that if Σ is a σ algebra and there is a measure $\mu: \Sigma \to E$ which is not strongly additive, then E contains an isomorphic copy of ℓ_{∞} .

Remark 2.1. The proof of the Diestel-Faires Theorem [11, p. 20, Theorem 2] does not ensure the existence of a topological isomorphism, indeed, only can be proved that there exists a copy of c_0 and ℓ_{∞} in E, respectively.

In [6] is defined an ideal of *multilinear mappings* (or *multi-ideal*) \mathcal{M} as a subclass of the class of all continuous multilinear mappings between Banach spaces such that for all $n \in \mathbb{N}$ and Banach spaces E_1, \ldots, E_n and F the components $\mathcal{M}(E_1,\ldots,E_n;F) = \mathcal{L}(E_1,\ldots,E_n;F) \cap \mathcal{M}$ satisfy:

- (i) $\mathcal{M}(E_1,\ldots,E_n;F)$ is a linear subspace of $\mathcal{L}(E_1,\ldots,E_n;F)$ which contains the *n*-linear mappings of finite type.
- (ii) The ideal property: if $A \in \mathcal{M}(E_1, \ldots, E_n; F)$, $u_j \in \mathcal{L}(G_j, E_j)$ for $j = 1, \ldots, n$ and $t \in \mathcal{L}(F; H)$, then $t \circ A \circ (u_1, \ldots, u_n) \in \mathcal{M}(G_1, \ldots, G_n; H).$

If $\|\|_{\mathcal{M}} : \mathcal{M} \to \mathbb{R}^+$ satisfies

(*i'*) $(\mathcal{M}(E_1,\ldots,E_n;F), \|\cdot\|_{\mathcal{M}})$ is a normed (Banach) space for all Banach spaces E_1,\ldots,E_n , F and all $n \in \mathbb{N}$.

- $(ii') ||A^n : \mathbb{K}^n \to \mathbb{K} : A^n(x_1, \dots, x_n) = x_1 \dots x_n ||_{\mathcal{M}} = 1 \text{ for all } n \in \mathbb{N}.$
- (*iii'*) Si $A \in \mathcal{M}(E_1, \ldots, E_n; F)$, $u_j \in \mathcal{L}(G_j, E_j)$ for $j = 1, \ldots, n$ and $t \in \mathcal{L}(F; H)$, then $||t \circ A \circ (u_1, \ldots, u_n)||_{\mathcal{M}} \le ||t|| ||A||_{\mathcal{M}} ||u_1|| \ldots ||u_n||$, then $(\mathcal{M}, ||\cdot||_{\mathcal{M}})$ is called a normed (Banach) multi-ideal.

The multi-ideal \mathcal{M} is said to be closed if each $\mathcal{M}(E_1, \ldots, E_n; F)$ is a closed subspace of $\mathcal{L}(E_1, \ldots, E_n; F)$.

An ideal of *homogeneous polynomials* (or *polynomial ideal*) Q is a subclass of the class of all continuous homogeneous polynomials between Banach spaces such that for all $n \in \mathbb{N}$ and Banach spaces E and F, the components $Q(^{n}E; F) = \mathcal{P}(^{n}E; F) \cap Q$ satisfy:

- (i) $\mathcal{Q}(^{n}E;F)$ is a linear subspace of $\mathcal{P}(^{n}E;F)$ which contains the *n*-homogeneous polynomials of finite type.
- (ii) The ideal property: if $u \in \mathcal{L}(G; E)$, $P \in \mathcal{Q}(^{n}E; F)$ and $t \in \mathcal{L}(F, H)$, then the composition $t \circ P \circ u$ is in $\mathcal{Q}(^{n}G; H)$.

When $n = 1 \mathcal{Q}$ is called an **operator ideal**.

Definición 2.2. Let \mathcal{I} be an operator ideal.

- (a) An application n- linear $A \in \mathcal{L}(E_1, E_2, \dots, E_n; F)$ belongs to $\mathcal{I} \circ \mathcal{L}$, in this case we write $A \in \mathcal{I} \circ \mathcal{L}(E_1, E_2, \dots, E_n; F)$, if there are a Banach space G, an n- linear mapping $B \in \mathcal{L}(E_1, E_2, \dots, E_n; G)$ and an operator $u \in \mathcal{I}(G; F)$, such that $A = u \circ B$.
- (b) An *n*-homogeneous polynomial $P \in \mathcal{P}(^{n}E; F)$ belongs to $\mathcal{I} \circ \mathcal{P}$ in this case we write $P \in \mathcal{I} \circ \mathcal{P}(^{n}E; F)$ if there are a Banach space G, an *n*-homogeneous polynomial $Q \in \mathcal{P}(^{n}E; G)$ and an operator $u \in \mathcal{I}(G; F)$ such that $P = u \circ Q$.

Example 2.3. Let \mathcal{K} and \mathcal{W} be the closed operator ideals formed by all compact and weakly compact linear operators, respectively. By $\mathcal{P}_{\mathcal{K}}$ and $\mathcal{P}_{\mathcal{W}}$ we mean the classes of all compact and weakly compact polynomials, respectively. The equalities $\mathcal{P}_{\mathcal{K}} = \mathcal{K} \circ \mathcal{P}$ and $\mathcal{P}_{\mathcal{W}} = \mathcal{W} \circ \mathcal{P}$ were proved by *R*. Ryan [23].

Definición 2.4. *Given an operator ideal* \mathcal{I} *and Banach spaces* E *and* F*, we define* $\mathcal{I}^{dual}(E;F) = \{T \in \mathcal{L}(E;F) : T' \in \mathcal{I}(F';E')\}.$

Definición 2.5. Given a Banach space E and an operator ideal \mathcal{I} . The collection of all \mathcal{I} bounded sets of E is defined by $C_I(E) = \{A \subset E : \exists T \in \mathcal{I}(Z; E) \text{ for some Banach space } Z \text{ such that } A \subset T(B_Z)\}.$

If $\mathcal{I} = \mathcal{K}$ or $\mathcal{I} = \mathcal{W}$ the compact and weakly compact operator ideals, respectively, then $C_I(E)$ are the relatively compact and relatively weakly compact sets of E, respectively.

In [17], given a multi-ideal \mathcal{M} , is constructed an ideal \mathcal{M}^{fac} such that every n-linear operator A in \mathcal{M}^{fac} is strongly \mathcal{M} - factorable in the sense that it factors through multilinear operator in \mathcal{M} with respect to any partition of the set $\{1, \ldots, n\}$. For $\mathcal{L}_K = \mathcal{K} \circ \mathcal{L}$ (compact multilinear operators) we have that $\mathcal{L}_K^{fac} = \mathcal{L}_{wb}$ (weakly continuous on bounded sets multilinear operators).

3 The main results

The next result is a generalization of theorem 1.1. The proof of the next theorem is based in ideas of [19, Theorem 6(i)].

Theorem 3.1. Let *E* and *F* be Banach spaces, and let *G* be a Banach space with an unconditional basis (g_n) and coordinate functionals (g'_n) .

- (a) If there exist operators $R \in \mathcal{L}(G; F)$ and $S \in \mathcal{L}(E; G)$ such that $(R(g_n))$ is a semi-normalized basic sequence in F and $(S'(g'_{n_k})) \notin C_{\mathcal{I}}(E')$ for all subsequence $(n_k) \subset \mathbb{N}$, where \mathcal{I} and \mathcal{J} be closed operator ideals such that $\mathcal{J} \subset \mathcal{I}^{dual}$. Then $\mathcal{J} \circ \mathcal{L}(E; F)$ is not complemented in $\mathcal{L}(E; F)$.
- (b) If E is a reflexive Banach space and there exist operators $R \in \mathcal{L}(G; F)$ and $S \in \mathcal{L}(E; G)$ such that $(R(g_n))$ is a semi-normalized basic sequence in F and $(S'(g'_{n_k})) \notin \mathcal{C}_{\mathcal{I}}(E')$ for all subsequence $(n_k) \subset \mathbb{N}$, where \mathcal{I} and \mathcal{J} be closed operator ideals such that $\mathcal{J}^{dual} \subset \mathcal{I}$. Then $\mathcal{J} \circ \mathcal{L}(E; F)$ is not complemented in $\mathcal{L}(E; F)$.

Proof. (a) Let (P_A) be the family of projections associated with (g_n) , R and S as in the hypothesis, and set $\mu(A) = R \circ P_A \circ S$, $A \subseteq \mathbb{N}$. Let E_0 be a separable subspace of E such that $||x'|| = ||x'|_{E_0}||$ for all $x' \in [S'(g'_n): n \ge 1]$. Let (y'_n) be the sequence of biorthogonal coefficients corresponding to $(R(g_n))$, and (f'_n) be the sequence of Hahn-Banach extension to F'. Now suppose that there exist a projection $\pi : \mathcal{L}(E; F) \to \mathcal{J} \circ \mathcal{L}(E; F)$. Let $J: F \to \ell_\infty$ be an operator that is an isometry on $[R(g_n): n \ge 1]$. We define $v: \mathcal{P}(\mathbb{N}) \to \mathcal{L}(E_0; \ell_\infty)$ by $v(A) = (J \circ \pi \circ \mu(A) - J \circ \mu(A))|_{E_0}$, $A \subseteq \mathbb{N}$. Since $\mu(\{n\})$ is an operator of finite range, then $v(\{n\}) = 0$ for every $n \in \mathbb{N}$. Apply [19, Lemma 4] to obtain an infinite subset M of \mathbb{N} so that $J \circ \pi \circ \mu(M) = J \circ \mu(M)$ on E_0 . Therefore $(J \circ \mu(M))|_{E_0} \in \mathcal{J}(E_0; \ell_\infty)$. Since J is an isometry on $[R(g_n): n \ge 1]$ and $\mu(M)(E_0) \subset [R(g_n): n \ge 1]$, we have that $\mu(M)|_{E_0} \in \mathcal{J}(E_0; F)$. As $\mathcal{J} \subset \mathcal{I}^{dual}$ it implies that $(\mu(M)|_{E_0})' \in \mathcal{I}(F'; E'_0)$. On the other hand, there exist a constant c > 0 such that $||f'_n|| \le c$ for all $n \in \mathbb{N}$. Is easy to see that $c(\mu(M)|_{E_0})'(\frac{f'_n}{c}) = (S'(g'_n))|_{E_0}$ for all $n \in M$. Consider the operator $\psi : [S'(g'_n)|_{E_0}: n \in M] \to [S'(g'_n): n \in M]$, defined by $\psi(S'(g'_n)|_{E_0}) = S'(g'_n)$. Since $||x'|| = ||x'|_{E_0}||$ for all $x' \in [S'(g'_n): n \ge 1]$, then ψ is an operator linear and continuo, thus $\psi \circ c(\mu(M)|_{E_0})') \in \mathcal{I}(F'; E')$ and $(S'(g'_n))_{n \in M} \subset \psi \circ c(\mu(M)|_{E_0})'(B_{F'})$. Thus $(S'(g'_n))_{n \in M} \in C_{\mathcal{I}}(E')$, but it is a contradiction with the hypothese. Therefore $\mathcal{J} \circ \mathcal{L}(E; F)$ is not complemented in $\mathcal{L}(E; F)$.

(b) If we assume that exist a projetion $\pi : \mathcal{L}(E; F) \to \mathcal{J} \circ \mathcal{L}(E; F)$, then analogously as in (i), we have that $\mu(M)|_{E_0} \in \mathcal{J}(E_0; F)$ for any infinite subset M of \mathbb{N} . Is easy to see that $(\mu(M)|_{E_0})'' \circ J_{E_0} = J_F \circ \mu(M)|_{E_0} \in \mathcal{J}(E_0; F'')$. Since E is reflexive then E_0 is also reflexive, therefore $(\mu(M)|_{E_0})'' = (\mu(M)|_{E_0})'' \circ J_{E_0} \circ J_{E_0}^{-1} \in \mathcal{J}(E_0''; F'')$. By hypothesis $\mathcal{J}^{dual} \subset \mathcal{I}$, thus $(\mu(M)|_{E_0})' \in \mathcal{I}(F'; E_0')$. As in the proof of (i) it implies that $(S'(g'_n))_{n \in M} \in \mathcal{C}_{\mathcal{I}}(E')$, but it is absurd. Finally we obtain the desired result.

Remark 3.2. By [8, Corollary 3.4] and [8, Proposition 3.8], $QN_p = \mathcal{K}_p^{dual}$ and $\mathcal{K}_p = Q\mathcal{N}_p^{dual}$, respectively. Thus, if we take $\mathcal{J} = Q\mathcal{N}_p$, $\mathcal{I} = \mathcal{K}_p$ and $\mathcal{J} = \mathcal{K}_p$, $\mathcal{I} = Q\mathcal{N}_p$, then the previously theorem can be applied.

In [3, Example 4.4] we can to see more examples of known ideals of operators that satisfy the theorem previously mentioned.

Theorem 3.3. Let *E* and *F* be Banach spaces, and let *G* be a Banach space with an unconditional basis (g_n) and coordinate functionals (g'_n) .

- (a) If there exist operators $R \in \mathcal{L}(G; F)$ and $S \in \mathcal{L}(E; G)$ such that $(R(g_n))$ is a semi-normalized basic sequence in F and $(S'(g'_{n_k})) \notin \mathcal{C}_{\mathcal{I}}(E')$ for all subsequence $(n_k) \subset \mathbb{N}$, where \mathcal{I} and \mathcal{J} be closed operator ideals such that $\mathcal{J} \subset \mathcal{I}^{dual}$. Then $\mathcal{J} \circ \mathcal{P}(^nE; F)$ is not complemented in $\mathcal{P}(^nE; F)$ for every $n \in \mathbb{N}$.
- (b) If E is a reflexive Banach space and there exist operators $R \in \mathcal{L}(G; F)$ and $S \in \mathcal{L}(E; G)$ such that $(R(g_n))$ is a semi-normalized basic sequence in F and $(S'(g'_{n_k})) \notin \mathcal{C}_{\mathcal{I}}(E')$ for all subsequence $(n_k) \subset \mathbb{N}$, where \mathcal{I} and \mathcal{J} be closed operator ideals such that $\mathcal{J}^{dual} \subset \mathcal{I}$. Then $\mathcal{J} \circ \mathcal{P}(^nE; F)$ is not complemented in $\mathcal{P}(^nE; F)$ for every $n \in \mathbb{N}$.

Proof. (a) The case n = 1 follows from Theorem 3.1 (a). Suppose that exists a projection $\pi : \mathcal{P}(^{n}E;F) \to \mathcal{J} \circ \mathcal{P}(^{n}E;F)$ for some n > 1. By a result of Ryan [23] there exists an isomorphism

$$P \in \mathcal{P}(^{n}E;F) \to \check{P} \in \mathcal{L}(\hat{\otimes}_{n,s,\pi}E;F)$$

Using [6, Proposition 3.2] we have that $\mathcal{J} \circ \mathcal{P}({}^{n}E;F)$ is isomorphic to $\mathcal{J}(\hat{\otimes}_{n,s,\pi}E;F)$, then there exists a projection $\rho : \mathcal{L}(\hat{\otimes}_{n,s,\pi}E;F) \to \mathcal{J}(\hat{\otimes}_{n,s,\pi}E;F)$. By a result of Blasco [5, Theorem 3] E is isomorphic to a complemented subspace of $\hat{\otimes}_{n,s,\pi}E$. Hence there exist operators $A \in \mathcal{L}(E; \hat{\otimes}_{n,s,\pi}E)$ and $B \in \mathcal{L}(\hat{\otimes}_{n,s,\pi}E;E)$ such that $B \circ A = I$. Consider the operator

$$\varphi: T \in \mathcal{L}(E; F) \to \rho(T \circ B) \circ A \in \mathcal{J}(E; F).$$

If $T \in \mathcal{J}(E; F)$, then $T \circ B \in \mathcal{J}(\hat{\otimes}_{n,s,\pi}E; F)$ and therefore $\rho(T \circ B) \circ A = T \circ B \circ A = T$. Thus $\varphi: T \in \mathcal{L}(E; F) \to \rho(T \circ B) \circ A \in \mathcal{J}(E; F)$ is a projection, contradicting the case n = 1. (b) The proof of (b) is almost identical to the proof of (a).

When $\mathcal{J} = \mathcal{I} = \mathcal{L}_K$ Theorem 3.3 correspond to part (*a*) of [22, Theorem 3.1], and part (b) of [22, Theorem 3.1] is obtained when $\mathcal{J} = \mathcal{I} = \mathcal{L}_{wK}$.

Proposition 3.4. Let $T \in (\mathcal{I} \circ \mathcal{L})^{fac}(^{n}E; F)$ and $\varphi \in E'$. Then $\varphi T \in (\mathcal{I} \circ \mathcal{L})^{fac}(^{n+1}E; F)$, where

 $\varphi T(x_1,\ldots,x_{n+1}) = \varphi(x_{n+1})T(x_1,\ldots,x_n).$

Proof. Let $\pi = \{j_1^1, \dots, j_{k_1}^1\} \cup \dots \cup \{j_1^m, \dots, j_{k_m}^m\}$ be a partition of $\{1, \dots, n+1\}$.

Case 1: Suppose that $\{n+1\} \in \pi$. Let $\hat{\pi} = \pi - \{n+1\}$ be a partition of $\{1, \ldots, n\}$. We can write

$$T \stackrel{\pi}{=} C \circ (B_1, \ldots, B_{m-1}),$$

where $B_i \in \mathcal{I} \circ \mathcal{L}(\underbrace{E \times \ldots \times E}_{k_i \text{ times}}; G_i), i = 1, \ldots, m-1$, and $C \in \mathcal{L}(G_1 \times \ldots \times G_{m-1}; F)$. We consider

 $G_m = \mathbb{K}$ and $B_m = \varphi$. Define $\widehat{C} \in \mathcal{L}(G_1 \times \ldots \times G_m; F)$ by

$$C(g_1,\ldots,g_m)=C(g_1,\ldots,g_{m-1})g_m,$$

for all $g_i \in G_i$, i = 1, ..., m. We can see easily that

$$\varphi T(x_1, \dots, x_{n+1}) = \widehat{C}(B_1(x_{j_1}^1, \dots, x_{j_{k_1}}^1), \dots, B_{m-1}(x_{j_1}^{m-1}, \dots, x_{j_{k_{m-1}}}^{m-1}), B_m(x_{n+1})).$$

Therefore $\varphi T \in (\mathcal{I} \circ \mathcal{L})^{fac}(^{n+1}E; F).$

Case 2: Suppose that $n + 1 \in \{j_1^l, \dots, j_{k_l}^l\}$ for some $1 \le l \le m$, where $k_l > 1$. Let $\hat{\pi} = \pi - \{n + 1\}$ be a partition of $\{1, \dots, n\}$. Then

$$T \stackrel{\pi}{=} C \circ (B_1, \ldots, B_m),$$

where $B_l \in \mathcal{I} \circ \mathcal{L}(\underbrace{E \times \ldots \times E}_{k_l-1 \ times}; G_l), B_i \in \mathcal{I} \circ \mathcal{L}(\underbrace{E \times \ldots \times E}_{k_i \ times}; G_i), i = 1, \ldots l - 1, l + 1, \ldots, m, \text{ and } C \in \mathcal{L}(G_1 \times \ldots \times G_m; F).$ We define $\widehat{B_l} \in \mathcal{I} \circ \mathcal{L}(\underbrace{E \times \ldots \times E}_{k_l \ times}; G_l), \underset{k_l \ times}{}$

by

$$\widehat{B}_l(x_1,\ldots,x_{k_l})=\varphi(x_{k_l})B_l(x_1,\ldots,x_{k_l-1}).$$

Finally, we have that

$$\varphi T(x_1, \dots, x_{n+1}) = \varphi(x_{n+1})T(x_1, \dots, x_n) = C(B_1(x_{j_1}^1, \dots, x_{j_{k_1}}^1), \dots, \widehat{B_l}(x_{j_1}^l, \dots, x_{n+1}), \dots, B_m(x_{j_1}^m, \dots, x_{j_{k_m}}^m))$$

This complete the proof.

Proposition 3.5. Let $T \in (\mathcal{I} \circ \mathcal{L})^{fac}(^{n}E; F)$ and $e \in E$. Then $\overline{T_j} \in (\mathcal{I} \circ \mathcal{L})^{fac}(^{n-j}E; F)$, where $\overline{T_j}(x_1, \ldots, x_{n-j}) = T(x_1, \ldots, x_{n-j}, \underbrace{e, \ldots, e}_{j \text{ times}})$, for every j = 1, 2, ..., n-1.

Proof. To prove the proposition by induction on j it suffices to prove for the case j = 1. Let $\pi = \{j_1^1, \ldots, j_{k_1}^1\} \cup \ldots \cup \{j_1^m, \ldots, j_{k_m}^m\}$ be a partition of $\{1, \ldots, n-1\}$. Consider the partition $\hat{\pi} = \pi \cup \{n\}$ of $\{1, \ldots, n\}$. By hipothesis $T \in (\mathcal{I} \circ \mathcal{L})^{fac}({}^nE; F)$, then we can write

$$T \stackrel{\pi}{=} C \circ (B_1, \ldots, B_{m+1}),$$

where $B_i \in \mathcal{I} \circ \mathcal{L}(\underbrace{E \times \ldots \times E}_{k_i \text{ times}}; G_i), i = 1, \ldots, m + 1, \text{ and } C \in \mathcal{L}(G_1 \times \ldots \times G_{m+1}; F)$. Let $\widehat{C} \in \mathcal{C}(G_1 \times \ldots \times G_{m+1}; F)$ define here

 $\mathcal{L}(G_1 \times \ldots \times G_m; F)$ define by

$$\widehat{C}(g_1,\ldots,g_m)=C(g_1,\ldots,g_m,B_{m+1}(e)),$$

then

$$\overline{T_1}(x_1, \dots, x_{n-1}) = T(x_1, \dots, x_{n-1}, e) = C(B_1(x_{j_1}^1, \dots, x_{j_{k_1}}^1), \dots, B_m(x_{j_1}^m, \dots, x_{j_{k_m}}^m), B_{m+1}(e))$$
$$= \widehat{C}(B_1(x_{j_1}^1, \dots, x_{j_{k_1}}^1), \dots, B_m(x_{j_1}^m, \dots, x_{j_{k_m}}^m))$$

Thus $\overline{T_1} \in (\mathcal{I} \circ \mathcal{L})^{fac}(^{n-1}E; F)$. This complete the proof.

Theorem 3.6. Let E and F be Banach spaces, and let G be a Banach space with an unconditional basis (g_n) and coordinate functionals (g'_n) . If there exist operators $R \in \mathcal{L}(G; F)$ and $S \in \mathcal{L}(E; G)$ such that $(R(g_n))$ is a semi-normalized basic sequence in F and $(S'(g'_{n_k})) \notin \mathcal{C}_{\mathcal{I}}(E')$ for all subsequence $(n_k) \subset \mathbb{N}$, where \mathcal{I} and \mathcal{J} be closed operator ideals such that $\mathcal{J} \subset \mathcal{I}^{dual}$. Then $(\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}({}^{n}E; F)$ is not complemented in $\mathcal{P}({}^{n}E; F)$ for every $n \in \mathbb{N}$.

Proof. When n = 1 we have that $(\mathcal{J} \circ \mathcal{L})^{fac}({}^{n}E;F) = \mathcal{J} \circ \mathcal{L}(E;F)$ and $\mathcal{P}({}^{n}E;F) = \mathcal{L}(E;F)$. Thus the case n = 1 follows from Theorem 3.1 (a). To prove the theorem by induction on n it suffices to prove that if $(\mathcal{J} \circ \mathcal{L})^{fac}({}^{n+1}E;F)$ is complemented in $\mathcal{P}({}^{n+1}E;F)$, then $(\mathcal{J} \circ \mathcal{L})^{fac}({}^{n}E;F)$ is complemented in $\mathcal{P}({}^{n}E;F)$. Aron and Schottenloher [1, Proposition 5.3] proved that $\mathcal{P}({}^{n}E;F)$ is isomorphic to a complemented subspace of $\mathcal{P}({}^{n+1}E;F)$ when F is the scalar field, but their proof works equally well when F is an arbitrary Banach space (see [5, Proposition 5]). Thus there exist operators $A \in \mathcal{L}(\mathcal{P}({}^{n}E;F);\mathcal{P}({}^{n+1}E;F))$ and $B \in \mathcal{L}(\mathcal{P}({}^{n+1}E;F);\mathcal{P}({}^{n}E;F))$ such that $B \circ A = I$. The operator A is of the form

$$A(P)(x) = \varphi_0(x)P(x)$$

for every $P \in \mathcal{P}({}^{n}E;F)$ and $x \in E$, where $\varphi_{0} \in E'$ verifies that $\|\varphi_{0}\| = 1 = \varphi_{0}(x_{0})$, where $x_{0} \in E$ and $\|x_{0}\| = 1$. Using Proposition 3.4 we have that if $P \in (\mathcal{J} \circ \mathcal{L})^{fac}({}^{n}E;F)$ then $A(P) \in (\mathcal{J} \circ \mathcal{L})^{fac}({}^{n+1}E;F)$. On the other hand, the operator B is of the form $B = A^{-1} \circ D$, where $D : \mathcal{P}({}^{n+1}E;F) \to \mathcal{P}({}^{n+1}E;F)$ is defined by $D(P)(x) = P(x) - P(x - \varphi_{0}(x)x_{0})$ for every $P \in \mathcal{P}({}^{n+1}E;F)$ and $x \in E$. More exactly B(P) is given by

$$B(P)(x) = \sum_{j=1}^{n+1} \binom{n+1}{j} (-1)^{j+1} \varphi_0^{j-1}(x) \check{P}(\underbrace{x, \dots, x}_{n+1-j \text{ times}}, \underbrace{e, \dots, e}_{j \text{ times}})$$

for all $x \in E$, (see [7, p. 597]). Applying simultaneously Propositions 3.4 and 3.5 we can to see that if $P \in (\mathcal{J} \circ \mathcal{L})^{fac}(^{n+1}E; F)$ then $B(P) \in (\mathcal{J} \circ \mathcal{L})^{fac}(^{n}E; F)$.

Let us assume that $(\mathcal{J} \circ \mathcal{L})^{fac}(^{n+1}E; F)$ is complemented in $\mathcal{P}(^{n+1}E; F)$, and let $\pi : \mathcal{P}(^{n+1}E; F) \to (\mathcal{J} \circ \mathcal{L})^{fac}(^{n+1}E; F)$ be a projection. Consider the operator

$$\rho = B \circ \pi \circ A : \mathcal{P}(^{n}E;F) \to (\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}(^{n}E;F).$$

If $P \in (\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}({}^{n}E; F)$, then $A(P) \in (\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}({}^{n+1}E; F)$, and therefore

$$\rho(P) = B \circ \pi \circ A(P) = B \circ A(P) = P.$$

Thus $\rho : \mathcal{P}({}^{n}E;F) \to (\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}({}^{n}E;F)$ is a projection, and therefore $(\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}({}^{n}E;F)$ is complemented in $\mathcal{P}({}^{n}E;F)$. This completes the proof.

Ghenciu [18] derived as corollaries of Theorem 1.1 results of several authors [2],[12], [14], [16] and [15]. We now apply Theorems 3.3 and 3.6 to obtain versions about ideals of polynomials of those corollaries.

Corollary 3.7. If $c_0 \to F$ and E' contains a weak star-null sequence (x'_n) such that $(x'_{n_k}) \notin C_{\mathcal{I}}(E')$ for all subsequence $(n_k) \subset \mathbb{N}$, where \mathcal{I} and \mathcal{J} be closed operator ideal such that $\mathcal{J} \subset \mathcal{I}^{dual}$. Then $\mathcal{J} \circ \mathcal{P}(^nE;F)$ is not complemented in $\mathcal{P}(^nE;F)$ for every $n \in \mathbb{N}$.

Corollary 3.8. If $c_0 \hookrightarrow F$ and E contains a complemented copy of c_0 . Then $\mathcal{J} \circ \mathcal{P}(^nE; F)$ is not complemented in $\mathcal{P}(^nE; F)$ for every $n \in \mathbb{N}$, where $\mathcal{J} \subset \mathcal{L}_{wK}$ is a closed operator ideal.

Corollary 3.9. If F contains a copy of ℓ_1 and $\mathcal{L}(E; \ell_1) \neq \mathcal{L}_K(E; \ell_1)$. Then $\mathcal{J} \circ \mathcal{P}(^nE; F)$ is not complemented in $\mathcal{P}(^nE; F)$ for every $n \in \mathbb{N}$, where $\mathcal{J} \subset \mathcal{L}_{wK}$ is a closed operator ideal.

When n = 1 and $\mathcal{J} = \mathcal{L}_{wK}$, Corollaries 3.7, 3.8 and 3.9 correspond to [18, Corollaries 2,3 and 5]. Ghenciu derived those corollaries by observing that E and F satisfy the hypothesis of Theorem 1.1 (b). Since the hypothesis of Theorem 1.1 (b) implies the hypothesis of Theorem 3.3 (a), we see that Corollaries 3.7, 3.8 and 3.9 follow from Theorem 3.3 (a).

Corollary 3.10. If F contains a copy of c_0 and E is infinite dimensional, then for all closed operator ideal $\mathcal{J} \subset \mathcal{L}_K$ we have that:

- (a) $\mathcal{J} \circ \mathcal{P}(^{n}E; F)$ is not complemented in $\mathcal{P}(^{n}E; F)$ for every $n \in \mathbb{N}$.
- (b) $(\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}(^{n}E; F)$ is not complemented in $\mathcal{P}(^{n}E; F)$ for every $n \in \mathbb{N}$.

Corollary 3.11. If *E* contains a complemented copy of ℓ_1 and *F* is infinite dimensional, then for all closed operator ideal $\mathcal{J} \subset \mathcal{L}_K$ we have that:

- (a) $\mathcal{J} \circ \mathcal{P}(^{n}E; F)$ is not complemented in $\mathcal{P}(^{n}E; F)$ for every $n \in \mathbb{N}$.
- (b) $(\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}(^{n}E; F)$ is not complemented in $\mathcal{P}(^{n}E; F)$ for every $n \in \mathbb{N}$.

When n = 1 and $\mathcal{J} = \mathcal{L}_K$, Corollaries 3.10 and 3.11 correspond to [18, Corollaries 4 and 6]. Ghenciu derived those corollaries by observing that E and F satisfy the hypothesis of Theorem 1.1 (*a*). Since the hypothesis of Theorem 1.1 (*a*) implies the hypothesis of Theorems 3.3 (*a*) and 3.6, we see that Corollaries 3.10 and 3.11 follow from Theorems 3.3 (*a*) and 3.6.

Corollary 3.12. If E contains a copy of ℓ_1 and F contains a copy of ℓ_p , with $2 \le p < \infty$, then for all closed operator ideal $\mathcal{J} \subset \mathcal{L}_K$ we have that:

(a) $\mathcal{J} \circ \mathcal{P}(^{n}E; F)$ is not complemented in $\mathcal{P}(^{n}E; F)$ for every $n \in \mathbb{N}$.

(b) $(\mathcal{J} \circ \mathcal{L})^{fac}(^{n}E; F)$ is not complemented in $\mathcal{P}(^{n}E; F)$ for every $n \in \mathbb{N}$.

When $\mathcal{J} = \mathcal{L}_K$, Corollary 3.12 correspond to [22, Corollary 3.8]. Since the hypothesis of [22, Corollary 3.8] implies the hypothesis of Theorems 3.3 (*a*) and 3.6, we see that Corollary 3.12 follow from Theorems 3.3 (*a*) and 3.6.

The next Corollary is a generalization of [13, Theorem 2] and [15, Theorem 2]. The proof is based in ideas of [19, Corollary 11].

Corollary 3.13. Let E and F be infinite dimensional Banach spaces. If $c_0 \hookrightarrow \mathcal{J}(E;F)$, then $\mathcal{J}(E;F)$ is not complemented in $\mathcal{L}(E;F)$ for all closed operator ideal $\mathcal{J} \subset \mathcal{L}_K$.

Proof. By Corollaries 3.10 and 3.11 we may suppose without loss of generality that F contains no copy of c_0 and E contains no complemented copy of ℓ_1 . Thus, by [16, Theorem 4] $\mathcal{J}(E;F)$ contains no copy of ℓ_{∞} . If T_n is a

copy of the unit vector basis of c_0 in $\mathcal{J}(E; F)$. Then $\sum_{n=1}^{\infty} T_n(x)$ converges unconditionally for each $x \in E$. Define

 $\mu: \wp(\mathbb{N}) \to \mathcal{J}(E;F)$ by $\mu(A)(x) = \sum_{n \in A} T_n(x), x \in E, A \subset \mathbb{N}$. Suppose there is a projection $\pi: \mathcal{L}(E;F) \to \mathbb{C}$

 $\mathcal{J}(E; F)$. Then $\pi(T_n) = T_n$ for every $n \in \mathbb{N}$. If $(||T_n||)$ does not converge to zero, we can apply the Diestel-Faires Theorem [11, p. 20, Theorem 2] to the measure $\pi \circ \mu$ and obtain $\ell_{\infty} \hookrightarrow \mathcal{J}(E; F)$. Therefore $||T_n|| \to 0$, but this is absurd too, because (T_n) is a copy of unit vectors. This complete the proof.

Proposition 3.14. Let E and F be infinite dimensional Banach spaces. If $c_0 \hookrightarrow \mathcal{J} \circ \mathcal{P}(^nE; F)$, then $\mathcal{J} \circ \mathcal{P}(^nE; F)$ is not complemented in $\mathcal{P}(^nE; F)$ for every $n \in \mathbb{N}$, and every closed ideal $\mathcal{J} \subset \mathcal{L}_K$.

Proof. By [6, Proposition 3.2] we have that $\mathcal{J} \circ \mathcal{P}(^{n}E; F)$ is isomorphic to $\mathcal{J}(\hat{\otimes}_{n,s,\pi}E; F)$. Thus the result follows from Corollary 3.13.

The proof of [16, Lemma 2] can be applied to get the following lemma.

Lemma 3.15. Assume E is separable, $(\mathcal{J} \circ \mathcal{L})^{fac}({}^{k}E; F)$ is complemented in $\mathcal{P}({}^{k}E; F)$, and an linear bounded operator $\phi : \ell_{\infty} \to \mathcal{P}({}^{k}E; F)$ is given with the following properties:

- (a) $\phi(e_n) \in (\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}({}^kE; F)$ for all $n \in \mathbb{N}$.
- (b) $\{\phi(\xi)(x): \xi \in \ell_{\infty}, x \in E\} \subset F$ is separable.

Then, for every infinite subset $M \subset \mathbb{N}$, there exists an infinite subset $M_0 \subset M$ with $\phi(\xi) \in (\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}({}^kE;F)$ for all $\xi \in \ell_{\infty}(M_0)$.

Lemma 3.16. Suppose E contains a complemented copy of ℓ_1 . Then $(\mathcal{J} \circ \mathcal{L})^{fac}(^nE; F)$ is not complemented in $\mathcal{P}(^nE; F)$ for every F and n > 1, where \mathcal{J} is a closed operator ideal $\mathcal{J} \subset \mathcal{L}_K$.

Proof. As in [20, Lemma 5], we can reduce the problem to the case $E = \ell_1$.

Fix $v \in B_F$ and define the operator $\phi : \ell_{\infty} \to \mathcal{P}(^k\ell_1; F)$ by $\phi(\xi)(x) = \sum_{i=1}^{\infty} \xi_i x_i^k v$ for $\xi = (\xi_i) \in \ell_{\infty}$,

 $x = (x_i) \in \ell_1$ and k > 1. Note that $\phi(e_n)$ are polynomials of finite type, so $\phi(e_n) \in (\mathcal{J} \circ \mathcal{L})^{fac}({}^k\ell_1; F)$ for all $n \in \mathbb{N}$. Suppose there exists a projection $\pi : \mathcal{P}({}^k\ell_1; F) \to (\mathcal{J} \circ \mathcal{L})^{fac}({}^k\ell_1; F)$ for any k > 1. Using Lemma 3.15, there is an infinite subset $M \subset \mathbb{N}$ such that $\phi(\xi) \in (\mathcal{J} \circ \mathcal{L})^{fac}({}^k\ell_1; F)$ for all $\xi \in \ell_{\infty}(M)$. But it is absurd because $\phi(\xi) \in \mathcal{P}_{wb}({}^k\ell_1; F)$ if and only if $\xi \in c_0$ (to see [20, Lemma 5]). Therefore $(\mathcal{J} \circ \mathcal{L})^{fac}({}^nE; F)$ is not complemented in $\mathcal{P}({}^nE; F)$ for every F and n > 1.

Proposition 3.17. Let *E* be an infinite dimensional Banach space and n > 1. If $c_0 \hookrightarrow (\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}(^nE; F)$, then $(\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}(^nE; F)$ is not complemented in $\mathcal{P}(^nE; F)$ for every closed operator ideal $\mathcal{J} \subset \mathcal{L}_K$.

Proof. By Corollary 3.10 and Lemma 3.16 we way suppose that F contains no copy of c_0 and E contains no complemented copy of ℓ_1 . By [20, Theorem 3] $(\mathcal{J} \circ \mathcal{L})^{fac}({}^nE; F)$ contains no copy of ℓ_{∞} . Let (P_i) be a copy of the unit vector basis (e_i) of c_0 in $(\mathcal{J} \circ \mathcal{L})^{fac}({}^nE; F)$. Then

$$\sup\left\{ \bigg\| \sum_{i \in F} e_i \bigg\|; F \subset \mathbb{N}, Ffinite \right\} = 1.$$

By a result of Bessaga and Pelczynski [4] (see also [9, p. 44, Theorem 6]) the series $\sum_{i=1}^{1} e_i$ is weakly unconditionally

Cauchy in c_0 . This implies that the series $\sum_{i=1}^{\infty} P_i$ is weakly unconditionally Cauchy in $(\mathcal{J} \circ \mathcal{L})^{fac}({}^{n}E; F)$. For every $\varphi \in F'$ and $x \in E$ we consider the continuous linear functional

$$\psi: P \in (\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}(^{n}E; F) \to \varphi(P(x)) \in \mathbb{C}.$$

Since the series $\sum_{i=1}^{\infty} P_i$ is weakly unconditionally Cauchy in $(\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}(^nE; F)$, $\sum_{i=1}^{\infty} |\psi(P_i)| = \sum_{i=1}^{\infty} |\varphi(P_i(x))| < \infty$ for every $\varphi \in F'$ and $x \in E$. This shows that $\sum_{i=1}^{\infty} P_i(x)$ is weakly unconditionally Cauchy in F for each

 $x \in E$. Finally since F contains no copy of c_0 , an application of [9, p. 45, Theorem 8] shows that $\sum P_i(x)$ converges unconditionally in F for each $x \in E$. Let $\mu : \wp(\mathbb{N}) \to \mathcal{P}(^{n}E;F)$ be the finitely additive vector measure defined by $\mu(A)(x) = \sum_{i \in A} P_i(x)$ for each $x \in E$ and $A \subset \mathbb{N}$. Suppose there is a projection $\pi : \mathcal{P}(^nE;F) \to \mathbb{P}(^nE;F)$

 $(\mathcal{J} \circ \mathcal{L})^{fac}({}^{n}E; F)$ for any $n \in \mathbb{N}$. Then $\pi(P_i) = P_i$ for each $i \in \mathbb{N}$. If the sequence $(||P_i||)$ does not converge to zero, then there is $\epsilon > 0$ and a subsequence (i_k) of \mathbb{N} , such that $||P_{i_k}|| > \epsilon$ for each $k \in \mathbb{N}$. But this implies that the measure $\pi \circ \mu : \wp(\mathbb{N}) \to (\mathcal{J} \circ \mathcal{L})^{fac}({}^{n}E; F)$ is not strongly additive. Then the Diestel-Faires Theorem [11, p. 20, Theorem 2] would imply that $(\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}({}^{n}E;F)$ contains a copy of ℓ_{∞} . Therefore $||P_i|| \to 0$, but this is absurd too, because (P_i) is a copy of (e_i) . This complete the proof.

When $\mathcal{J} = \mathcal{L}_K$ Proposition 3.17 gives [22, Proposition 3.10].

Theorem 3.18. Let E and F be Banach spaces and $P \in \mathcal{P}(^{n}E;F)$ such that $P \notin (\mathcal{J} \circ \mathcal{L})^{fac}(^{n}E;F)$, where $\mathcal{J} \subset \mathcal{L}_K$ is a closed operator ideal. Suppose that P admits a factorization $P = Q \circ T$ through a Banach space G with an unconditional finite dimensional expansion of the identity, where $T \in \mathcal{L}(E;G)$ and $Q \in \mathcal{P}(^{n}G;F)$. Then $(\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}({}^{n}E;F)$ contains a copy of c_0 and thus $(\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}({}^{n}E;F)$ is not complemented in $\mathcal{P}({}^{n}E;F)$.

Proof. There exists a sequence $(A_i) \in \mathcal{F}(G;G)$ such that, for every $g \in G$, we have $g = \sum_{i=1}^{\infty} A_i(g)$ unconditionally. Then,

$$Q\left(\sum_{i=1}^{k} A_{i}(g)\right) = \sum_{i_{1},\dots,i_{n}=1}^{k} \check{Q}(A_{i_{1}}(g),\dots,A_{i_{n}}(g))$$
$$= \sum_{m=1}^{k} \left(\sum_{\max\{i_{1},\dots,i_{n}\}=m} \check{Q}(A_{i_{1}}(g),\dots,A_{i_{n}}(g))\right)$$
$$= \sum_{m=1}^{k} Q_{m}(g).$$

Since Q is continuous, we have that $Q(g) = \sum_{m=1}^{\infty} Q_m(g)$ for all $g \in G$.

To prove that $Q_m \in (\mathcal{J} \circ \mathcal{L})^{fac}({}^nG; F)$ for every $m \in \mathbb{N}$, it suffices to prove that the multilinear map $\varphi \in \mathcal{L}({}^nG; F)$ defined by,

 $\mathcal{L}(^{n}G;F) \text{ defined by,} \\ \varphi(g_{1},\ldots,g_{n}) = \check{Q}(C_{1}(g_{1}),\ldots,C_{n}(g_{n})) \text{ is contained in } (\mathcal{J} \circ \mathcal{L})^{fac}(^{n}G;F), \text{ for any } C_{1},\ldots,C_{n} \in \mathcal{F}(G;G). \\ \text{Let } \pi = \{j_{1}^{1},\ldots,j_{k_{1}}^{1}\} \cup \ldots \cup \{j_{1}^{m},\ldots,j_{k_{m}}^{m}\} \text{ be a partition of } \{1,\ldots,n\}. \text{ We consider } B_{i} : \underbrace{G \times \ldots \times G}_{k_{i} \text{ times}} \rightarrow \underbrace{G \times \ldots \times G}_{k_{i} \text{ times}}$

 $\hat{\otimes}_{k_i,s,\pi}G, i=1,\ldots,m$, defined by

$$B_{i}(g_{j_{1}^{i}},\ldots,g_{j_{k_{i}}^{i}})=C_{j_{1}^{i}}(g_{j_{1}^{i}})\widehat{\otimes}_{s,\pi}\ldots\widehat{\otimes}_{s,\pi}C_{j_{k_{i}}^{i}}(g_{j_{k_{i}}^{i}}),$$

for all $g_{j_1^1}, \ldots, g_{j_{k_m}^m} \in G$.

 $B_i = \widehat{B}_i \circ \psi, \text{ where } \psi : \underbrace{G \times \ldots \times G}_{k_i \text{ times}} \to \hat{\otimes}_{k_i, s, \pi} G \text{ is the canonical isomorphism, and } \widehat{B}_i : \hat{\otimes}_{k_i, s, \pi} G \to \hat{\otimes}_{k_i, s, \pi} G$

is defined by

 $\hat{B}_{i}(g_{j_{1}^{i}} \overset{\circ}{\otimes} \pi \dots \overset{\circ}{\otimes} \pi g_{j_{k_{i}}^{i}}) = C_{j_{1}^{i}}(g_{j_{1}^{i}}) \overset{\circ}{\otimes} \pi \dots \overset{\circ}{\otimes} \pi C_{j_{k_{i}}^{i}}(g_{j_{k_{i}}^{i}}).$ Like $C_{j_{1}^{i}}(G), \dots, C_{j_{k_{i}}^{i}}(G)$ are spaces finite dimensional, then $\hat{B}_{i} \in \mathcal{J}(\overset{\circ}{\otimes}_{k_{i},\pi}G; \overset{\circ}{\otimes}_{k_{i},\pi}G)$ for every $i = 1, \dots, m$. Therefore $B_{i} \in \mathcal{J} \circ \mathcal{L}(\underbrace{G \times \dots \times G}_{k_{i}}; \overset{\circ}{\otimes}_{k_{i},\pi}G), \underbrace{K_{i} \in \mathcal{K}}_{k_{i},\pi}G)$

 $i=1,\ldots,m.$ Let $C\in\mathcal{L}(\hat{\otimes}_{k_1,\pi}G,\ldots,\hat{\otimes}_{k_m,\pi}G;F)$ given by

$$C(g_{j_1^1} \hat{\otimes}_{\pi} \dots \hat{\otimes}_{\pi} g_{j_{k_1^1}}, \dots, g_{j_1^m} \hat{\otimes}_{\pi} \dots \hat{\otimes}_{\pi} g_{j_{k_m}^m}) = \check{Q}(g_{j_1^1}, \dots, g_{j_{k_1^1}}, \dots, g_{j_1^m}, \dots, g_{j_{k_m}^m})$$

We have that

 $\varphi(g_1, \dots, g_n) = \check{Q}(C_1(g_1), \dots, C_n(g_n)) = \check{Q}(C_{j_1^1}(g_{j_1^1}), \dots, C_{j_{k_m}^m}(g_{j_{k_m}^m})) = C(B_1(g_1^1, \dots, g_{k_1}^1), \dots, B_m(g_{j_1^m}, \dots, g_{j_{k_m}^m}))$ for all g_1 , $g_1 \in C$.

for all $g_1, \ldots, g_n \in G$. Therefore, $Q_m \in (\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}({}^nG; F)$ for every $m \in \mathbb{N}$, and so, $Q_m \circ T \in (\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}({}^nE; F)$ for every $m \in \mathbb{N}$.

Choosing finite subsets I_1, \ldots, I_k of integers, we have

$$\left\|\sum_{i_{1}\in I_{1},\dots,i_{k}\in I_{k}}\check{Q}(A_{i_{1}}(g),\dots,A_{i_{k}}(g))\right\| = \left\|\check{Q}\left(\sum_{i_{1}\in I_{1}}A_{i_{1}}(g),\dots,\sum_{i_{k}\in I_{k}}A_{i_{k}}(g)\right)\right\| \le \|\check{Q}\|\left\|\sum_{i_{1}\in I_{1}}A_{i_{1}}(g)\right\|\dots\left\|\sum_{i_{k}\in I_{k}}A_{i_{k}}(g)\right\|$$

Hence, the series

$$\sum_{i_1\in I_1,\ldots,i_k\in I_k}^{\infty}\check{Q}(A_{i_1}(g),\ldots,A_{i_k}(g))$$

is unconditionally convergent for all $g \in G$ [10, Theorem 1.9]. Therefore, $P(x) = \sum_{m=1}^{\infty} Q_m(T(x))$ unconditionally. Moreover, by the uniform boundedness principle [21, Theorem 2.6], we have

$$\sup\left\{\left\|\sum_{m\in F}Q_m\circ T\right\|; F\subset \mathbb{N}, \ F \ finite\right\}<\infty.$$

So $\sum_{m=1}^{\infty} Q_m \circ T$ is weakly unconditionally Cauchy in $(\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}({}^nE;F)$. Since $P \notin (\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}({}^nE;F)$, an

application of [9, p.45, Theorem 8] shows that $(\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}({}^{n}E; F)$ contains a copy of c_0 , and therefore by Corollary 3.13 and Proposition 3.17 $(\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}({}^{n}E; F)$ is not complemented in $\mathcal{P}({}^{n}E; F)$.

Corollary 3.19. Let E and F be Banach spaces, with E infinite dimensional, and let n > 1. If each $P \in \mathcal{P}(^{n}E; F)$ such that $P \notin (\mathcal{J} \circ \mathcal{L})^{fac}(^{n}E; F)$ admits a factorization $P = Q \circ T$, where $T \in \mathcal{L}(E; G)$, $Q \in \mathcal{P}(^{n}G; F)$ and G is a Banach space with an unconditional finite dimensional expansion of the identity, then the following conditions are equivalent for any closed operator ideal $\mathcal{J} \subset \mathcal{L}_{K}$.

- (1) $(\mathcal{J} \circ \mathcal{L})^{fac}(^{n}E; F)$ contains a copy of c_0 .
- (2) $(\widehat{\mathcal{J} \circ \mathcal{L}})^{fac}(^{n}E;F)$ is not complemented in $\mathcal{P}(^{n}E;F)$.
- (3) $(\mathcal{J} \circ \mathcal{L})^{fac}(^{n}E; F) \neq \mathcal{P}(^{n}E; F).$
- (4) $\mathcal{P}(^{n}E;F)$ contains a copy of c_{0} .
- (5) $\mathcal{P}(^{n}E;F)$ contains a copy of ℓ_{∞} .

Proof. $(1) \Rightarrow (2)$ by Proposition 3.17.

 $(2) \Rightarrow (3)$ is obvious.

- $(3) \Rightarrow (1)$ by Theorem 3.18.
- $(1) \Rightarrow (4)$ is obvious.

 $(4) \Rightarrow (3)$ suppose (4) holds and (3) does not hold. Then $(\mathcal{J} \circ \mathcal{L})^{fac}({}^{n}E; F) = \mathcal{P}({}^{n}E; F) \supset c_{0}$. Thus (1) holds, and therefore (3) holds, a contradiction.

 $(5) \Rightarrow (4)$ is obvious.

(4) \Rightarrow (5) Since (4) \Rightarrow (1) $(\mathcal{J} \circ \mathcal{L})^{fac}({}^{n}E;F) \subset \mathcal{P}_{K}({}^{n}E;F)$ contains a copy of c_{0} . By a result of Ryan [23] $\mathcal{P}({}^{n}E;F)$ and $\mathcal{P}_{K}({}^{n}E;F)$ are isometrically isomorphic to $\mathcal{L}(\widehat{\otimes}_{n,s,\pi}E;F)$ and $\mathcal{L}_{K}(\widehat{\otimes}_{n,s,\pi}E;F)$, respectively. Thus $\mathcal{L}_{K}(\widehat{\otimes}_{n,s,\pi}E;F)$ contains a copy of c_{0} . Since E is infinite dimensional, $\widehat{\otimes}_{n,s,\pi}E$ is also infinite dimensional. Then by combining the proofs of [16, Theorem 6, $(iii) \Rightarrow (ii)$] and [15, Remark 3 e) 2 \Rightarrow 3] we can conclude that $\mathcal{L}(\widehat{\otimes}_{n,s,\pi}E;F)$ contains a copy of ℓ_{∞} and the result follows.

Thus (1), (2), (3), (4) and (5) are equivalent.

In particular if $\mathcal{J} = \mathcal{L}_K$ and E has an unconditional finite dimensional expansion of the identity we obtain [20, Theorem 7]. The assumptions of this corollary apply also if F is a complemented subspace of a space with an unconditional basis.

References

- [1] R.M. Aron, M. Schottenloher, *Compact holomorphic mappings on Banach spaces and the approximation property*. J. Funct. Anal. 21 (1976), 7–30.
- [2] E. Bator and P. Lewis, Complemented spaces of operators, Bull. Pol. Acad. Sci. Math. 50 (4) (2002), 413-416.
- [3] S. Berrio and G. Botelho, *Approximation properties determined by operator ideals*, Studia. Mathematica. 208 (2012), 97-116.
- [4] C. Bessaga, A. Pelczynski, On bases and unconditional convergence of series in Banach spaces, Studia Math. 17 (1958), 151-164.
- [5] F. Blasco, Complementation of symmetric tensor products and polynomials, Studia Math. 123 (1997), 165-173.
- [6] G. Botelho, D. Pellegrino and P. Rueda, On Composition Ideals of Multilinear Mappings and Homogeneous Polynomials, Publ. RIMS, Kyoto Univ. 43 (2007), 1139–1155.

- [7] E. Çaliskan and P. Rueda, On distinguished polynomials and their projections, Ann. Acad. Sci. Fenn. Math. 37(2012), 595–603.
- [8] J. M. Delgado, C. Piñeiro and E. Serrano, Operators whose adjoints are quasi p-nuclear, Studia Math. 197 (2010), 291–304.
- [9] J. Diestel, Sequences and Series in Banach Spaces, Springer, New York, 1984.
- [10] J. Diestel, H. Jarchow and A. Tonge, *Absolutely Summing Operators*, Cambridge Stud. Adv. Math. 43, Cambridge Univ. Press, Cambridge 1995.
- [11] J. Diestel, J. Uhl, *Vector measures*, Mathematical Surveys Number 15, American Mathematical Society, Providence, Rhode Island, 1977.
- [12] G. Emmanuele, Remarks on the uncomplemented subspace $\mathcal{W}(E; F)$, J. Funct. Anal. 99 (1991), 125-130.
- [13] G. Emmanuele, A remark on the containment of c_0 in spaces of compact operators, Math. Proc. Cambridge Phil. Soc., 111 (1992), 331–335.
- [14] M. Feder, On the non-existance of a projection onto the space of compact Operators, Canad. Math. Bull. 25 (1982), 78-81.
- [15] K. John, On the uncomplemented subspace $\mathcal{K}(X;Y)$, Czechoslovak Math. J. 42 (1992), 167–173.
- [16] N. Kalton, Spaces of compact operators, Math. Ann. 208 (1974), 267-278.
- [17] B. Geraldo and E. Torres, Strongly factorable multilinear operators on Banach spaces, Colloquium Mathematicum 154 (2018), 15-30.
- [18] I. Ghenciu, Complemented spaces of operators, Proc. Amer. Math. Soc. 133 (9) (2005), 2621–2623.
- [19] I. Ghenciu and P. Lewis, Unconditional convergence in the strong operator topology and ℓ_{∞} , Glasg. Math. J. 53 (2011), 583–598.
- [20] González, M.G., Gutiérrez, J.M, The polynomial property (V), Arch. Math. (Basel) 75(4)(2000), 299–306.
- [21] J. Mujica, Complex Analysis in Banach Spaces, Math. Studies 120, North-Holland, Amsterdam 1986.
- [22] S. Pérez, Complemented subspaces of homogeneous polynomials, Rev. Mat. Complut., 31 (2018), 153-161.
- [23] R. Ryan, *Applications of topological tensor products to infinite dimensional holomorphy*, Ph. D. Thesis, Trinity College, Dublin, 1980.