

ON ASYMPTOTIC EXPANSIONS OF OSCILLATORY INTEGRALS WITH PHASE FUNCTIONS EXPRESSED BY A PRODUCT OF POSITIVE REAL POWER FUNCTION AND REAL ANALYTIC FUNCTION IN ONE VARIABLE

TOSHIO NAGANO

ABSTRACT. In this paper, by using asymptotic expansions of oscillatory integrals with positive real power phase functions in one variable, we obtain asymptotic expansions of oscillatory integrals with phase functions expressed by a product of positive real power function and real analytic function in one variable. Moreover we show an example which we can compute all coefficients of terms in asymptotic expansions concretely.

1. INTRODUCTION

We study for **degenerate** phase functions $\phi(x)$ and suitable amplitude functions $a(x)$ in one variable, asymptotic expansions of oscillatory integrals:

$$Os\text{-}\int_{-\infty}^{\infty} e^{i\lambda\phi(x)} a(x) dx := \lim_{\varepsilon \rightarrow +0} \int_{-\infty}^{\infty} e^{i\lambda\phi(x)} a(x) \chi(\varepsilon x) dx,$$

as $\lambda \rightarrow \infty$, where χ is a rapidly decreasing function of class C^∞ on \mathbb{R} with $\chi(0) = 1$ and $0 < \varepsilon < 1$.

In the previous paper [17], we studied the case of $\phi(x) = x^p$ for any positive real number $p > 0$ and $a(x) \in \mathcal{A}_\delta^\tau(\mathbb{R})$, where for any $\tau \in \mathbb{R}$ and $\delta \in [-1, p-1)$, $\mathcal{A}_\delta^\tau(\mathbb{R})$ is the set of all $a \in C^\infty(\mathbb{R})$ such that for any $l \in \mathbb{Z}_{\geq 0}$,

$$|a|_l^{(\tau)} := \max_{k \leq l} \sup_{x \in \mathbb{R}} (1 + |x|^2)^{-(\tau + \delta k)/2} |a^{(k)}(x)| < \infty.$$

Then we obtained the following results (Theorem 3.3 (i) and Theorem 5.2 in [17]):

Theorem 1.1. *Assume that $\lambda > 0$ and $p > 0$. Let $\varphi \in C_0^\infty(\mathbb{R})$. Then there exist the following oscillatory integrals, and the following holds:*

$$Os\text{-}\int_0^\infty e^{\pm i\lambda x^p} \varphi(x) dx = \int_0^\infty e^{\pm i\lambda x^p} \varphi(x) dx,$$

where double signs \pm are in the same order.

Theorem 1.2. *Assume that $\lambda > 0$ and $a \in \mathcal{A}_\delta^\tau(\mathbb{R})$. Then the following hold:*

Date: October 22, 2020.

2010 Mathematics Subject Classification. Primary 42B20 ; Secondary 41A60, 33B20.

Key words and phrases. oscillatory integral, stationary phase method, asymptotic expansion.

The author was supported by Tokyo University of Science Graduate School doctoral program scholarship and an exemption of the cost of equipment from 2016 to 2018 and would like to thank to Professor Minoru Ito for giving me an opportunity of studies and preparing the environment.

(i) If $p > 0$, then for any $N \in \mathbb{N}$ such that $N \geq p + 1$, as $\lambda \rightarrow \infty$,

$$Os- \int_0^\infty e^{\pm i \lambda x^p} a(x) dx = \sum_{k=0}^{N-[p]-1} \tilde{I}_{p,k+1}^\pm \frac{a^{(k)}(0)}{k!} \lambda^{-\frac{k+1}{p}} + O\left(\lambda^{-\frac{N-p+1}{p}}\right),$$

where $[p]$ is the Gauss' symbol, that is, $[p] \in \mathbb{Z}$ such that $p - 1 < [p] \leq p$,

$$(1.1) \quad \tilde{I}_{p,k+1}^\pm = p^{-1} e^{\pm i \frac{\pi}{2} \frac{k+1}{p}} \Gamma\left(\frac{k+1}{p}\right),$$

and double signs \pm are in the same order.

(ii) If $p = m \in \mathbb{N}$, then for any $N \in \mathbb{N}$ such that $N > m$, as $\lambda \rightarrow \infty$,

$$Os- \int_0^\infty e^{\pm i \lambda (-y)^m} a(-y) dy = \sum_{k=0}^{N-m-1} (-1)^k \tilde{I}_{m,k+1}^{\pm \pm m} \frac{a^{(k)}(0)}{k!} \lambda^{-\frac{k+1}{m}} + O\left(\lambda^{-\frac{N-m+1}{m}}\right),$$

where

$$(1.2) \quad \tilde{I}_{m,k+1}^{\pm \pm m} = m^{-1} e^{\pm (-1)^m i \frac{\pi}{2} \frac{k+1}{m}} \Gamma\left(\frac{k+1}{m}\right),$$

and double signs \pm are in the same order.

In this paper, by using Theorem 1.1, as an application of Theorem 1.2, we obtain the following results:

Theorem 1.3. Assume that $\lambda > 0$ and $p > 0$. Let $\{a_j\}_{j \in \mathbb{N}}$ be a sequence of real numbers such that $l_0 := \sup_{j \in \mathbb{N}} |a_j|^{1/j} < \infty$, and let

$$\phi(x) := x^p \left(1 + \sum_{j=1}^{\infty} a_j x^j \right)$$

for $|x| < R_0 := (2l_0)^{-1}$ where if $l_0 = 0$, then $R_0 = \infty$. Then there exists a neighborhood U of the origin in \mathbb{R} such that $U \subset (-R_0, R_0)$, and a diffeomorphism $x = \Phi(y)$ for $x, y \in U$ such that $x(1 + \sum_{j=1}^{\infty} a_j x^j)^{1/p} = y$, $\Phi(U \cap [0, \infty)) = U \cap [0, \infty)$ and $\Phi(U \cap (-\infty, 0]) = U \cap (-\infty, 0]$, for any $a \in C_0^\infty(\mathbb{R})$ such that $\text{supp } a \subset U$, the following hold:

(i) For any $N \in \mathbb{N}$ such that $N \geq p + 1$, as $\lambda \rightarrow \infty$,

$$\begin{aligned} & \int_0^\infty e^{\pm i \lambda x^p (1 + \sum_{j=1}^{\infty} a_j x^j)} a(x) dx \\ &= \sum_{k=0}^{N-[p]-1} \frac{\tilde{I}_{p,k+1}^\pm}{k!} \left(\frac{d}{dy} \right)^k \Big|_{y=0} \left\{ a(\Phi(y)) \frac{d\Phi}{dy}(y) \right\} \lambda^{-\frac{k+1}{p}} + O\left(\lambda^{-\frac{N-p+1}{p}}\right), \end{aligned}$$

where $\tilde{I}_{p,k+1}^\pm$ is given by (1.1) and double signs \pm are in the same order.

(ii) If $p = m \in \mathbb{N}$, then for any $N \in \mathbb{N}$ such that $N > m$, as $\lambda \rightarrow \infty$,

$$\begin{aligned} & \int_{-\infty}^\infty e^{\pm i \lambda x^m (1 + \sum_{j=1}^{\infty} a_j x^j)} a(x) dx \\ &= \sum_{k=0}^{N-m-1} \frac{c_{m,k}^\pm}{k!} \left(\frac{d}{dy} \right)^k \Big|_{y=0} \left\{ a(\Phi(y)) \frac{d\Phi}{dy}(y) \right\} \lambda^{-\frac{k+1}{m}} + O\left(\lambda^{-\frac{N-m+1}{m}}\right), \end{aligned}$$

where

$$c_{m,k}^{\pm} = m^{-1} \left\{ e^{\pm i \frac{\pi}{2} \frac{k+1}{m}} + (-1)^{k+1} e^{\pm (-1)^m i \frac{\pi}{2} \frac{k+1}{m}} \right\} \Gamma \left(\frac{k+1}{m} \right),$$

and double signs \pm are in the same order.

In particular, when $a_j = 1/j!$, by using differential coefficients of all orders, of Lambert W-function $X = W(Y)$ (cf.[1]) at $Y = 0$, which is an inverse function of $Y = Xe^X$ defined on a certain neighborhood of the origin in \mathbb{R} , then we can compute all coefficients of terms in asymptotic expansions concretely as follows:

Corollary 1.4. (i)

$$\begin{aligned} & \int_0^{\infty} e^{\pm i \lambda x^p} e^x a(x) dx \\ &= \sum_{k=0}^{N-[p]-1} \frac{\tilde{I}_{p,k+1}^{\pm}}{k!} \left(\frac{d}{dy} \right)^k \Big|_{y=0} \left\{ a(\Phi(y)) \frac{dW}{dY} \left(\frac{y}{p} \right) \right\} \lambda^{-\frac{k+1}{p}} + O \left(\lambda^{-\frac{N-p+1}{p}} \right), \end{aligned}$$

where $\tilde{I}_{p,k+1}^{\pm}$ is given by (1.1) and double signs \pm are in the same order.

(ii) If $p = m \in \mathbb{N}$, then for any $N \in \mathbb{N}$ such that $N > m$, as $\lambda \rightarrow \infty$,

$$\begin{aligned} & \int_{-\infty}^{\infty} e^{\pm i \lambda x^m} e^x a(x) dx \\ &= \sum_{k=0}^{N-m-1} \frac{c_{m,k}^{\pm}}{k!} \left(\frac{d}{dy} \right)^k \Big|_{y=0} \left\{ a(\Phi(y)) \frac{dW}{dY} \left(\frac{y}{m} \right) \right\} \lambda^{-\frac{k+1}{m}} + O \left(\lambda^{-\frac{N-m+1}{m}} \right), \end{aligned}$$

where

$$c_{m,k}^{\pm} = m^{-1} \left\{ e^{\pm i \frac{\pi}{2} \frac{k+1}{m}} + (-1)^{k+1} e^{\pm (-1)^m i \frac{\pi}{2} \frac{k+1}{m}} \right\} \Gamma \left(\frac{k+1}{m} \right),$$

and double signs \pm are in the same order.

To the end of §1, we note notation used in this paper:

$C^{\infty}(\mathbb{R}^n)$ is the set of complex-valued functions of class C^{∞} on \mathbb{R}^n . $C_0^{\infty}(\mathbb{R}^n)$ is the set of all $f \in C^{\infty}(\mathbb{R}^n)$ with compact support.

O means the Landau's symbol, that is, $f(x) = O(g(x))$ ($x \rightarrow a$) if $|f(x)/g(x)|$ is bounded as $x \rightarrow a$ for functions f and g , where $a \in \mathbb{R} \cup \{\pm\infty\}$.

2. PROOFS OF MAIN RESULTS

Proof of Theorem 1.3. (i) Since $l := \limsup_{j \rightarrow \infty} |a_j|^{1/j} \leq \sup_{j \in \mathbb{N}} |a_j|^{1/j} =: l_0 < \infty$, by Cauchy-Hadamard's formula, the radius of convergence of the power series $\sum_{j=1}^{\infty} a_j x^j$ is $R := 1/l > 0$. Moreover if $|x| < R_0 := (2l_0)^{-1} \leq (2l)^{-1} < R$, then

$$\left| \sum_{j=1}^{\infty} a_j x^j \right| \leq \sum_{j=1}^{\infty} |a_j| |x|^j < \sum_{j=1}^{\infty} |a_j| \left\{ \left(2 \sup_{j \in \mathbb{N}} |a_j|^{1/j} \right)^{-1} \right\}^j < \sum_{j=1}^{\infty} 2^{-j} = 1.$$

Hence if $|x| < R_0$, since $1 + \sum_{j=1}^{\infty} a_j x^j > 0$, then we can define

$$f(x) := x \left(1 + \sum_{j=1}^{\infty} a_j x^j \right)^{1/p}$$

for $x \in (-R_0, R_0)$. Since f is a function of class C^∞ on $(-R_0, R_0)$ with $f(0) = 0$ and $f'(0) = 1$, by inverse function theorem, there exists a neighborhood U of the origin in \mathbb{R} such that $U \subset (-R_0, R_0)$, and a diffeomorphism $x = \Phi(y)$ for $x, y \in U$ such that $f(x) = y$, $\Phi(U \cap [0, \infty)) = U \cap [0, \infty)$ and $\Phi(U \cap (-\infty, 0]) = U \cap (-\infty, 0]$. Then for any $a \in C_0^\infty(\mathbb{R})$ such that $\text{supp } a \subset U$, the following improper integrals are absolutely convergent:

$$\int_0^\infty e^{\pm i\lambda x^p(1+\sum_{j=1}^\infty a_j x^j)} a(x) dx \text{ and } \int_{-\infty}^0 e^{\pm i\lambda x^m(1+\sum_{j=1}^\infty a_j x^j)} a(x) dx \text{ for } m \in \mathbb{N}.$$

By change of variable $x = \Phi(y)$ on $U \cap [0, \infty)$, since $x^p(1+\sum_{j=1}^\infty a_j x^j) = (f(x))^p = y^p$,

$$\int_0^\infty e^{\pm i\lambda x^p(1+\sum_{j=1}^\infty a_j x^j)} a(x) dx = \int_0^\infty e^{\pm i\lambda y^p} a(\Phi(y)) \frac{d\Phi}{dy}(y) dy.$$

Here since $a(\Phi(y)) \frac{d\Phi}{dy}(y) \in C_0^\infty(\mathbb{R})$, by Theorem 1.1 and Theorem 1.2 (i), for any $N \in \mathbb{N}$ such that $N \geq p+1$, as $\lambda \rightarrow \infty$,

$$\begin{aligned} \int_0^\infty e^{\pm i\lambda x^p(1+\sum_{j=1}^\infty a_j x^j)} a(x) dx &= Os- \int_0^\infty e^{\pm i\lambda y^p} a(\Phi(y)) \frac{d\Phi}{dy}(y) dy \\ &= \sum_{k=0}^{N-[p]-1} \frac{\tilde{I}_{p,k+1}^\pm}{k!} \left(\frac{d}{dy} \right)^k \Big|_{y=0} \left\{ a(\Phi(y)) \frac{d\Phi}{dy}(y) \right\} \lambda^{-\frac{k+1}{p}} + O\left(\lambda^{-\frac{N-p+1}{p}} \right). \end{aligned}$$

(ii) If $p = m \in \mathbb{N}$, then we take $N \in \mathbb{N}$ such that $N > m$, since $N \geq m+1$ and $[m] = m$, by (i), as $\lambda \rightarrow \infty$,

$$\begin{aligned} \int_0^\infty e^{\pm i\lambda x^m(1+\sum_{j=1}^\infty a_j x^j)} a(x) dx \\ (2.1) \quad = \sum_{k=0}^{N-m-1} \frac{\tilde{I}_{m,k+1}^\pm}{k!} \left(\frac{d}{dy} \right)^k \Big|_{y=0} \left\{ a(\Phi(y)) \frac{d\Phi}{dy}(y) \right\} \lambda^{-\frac{k+1}{m}} + O\left(\lambda^{-\frac{N-m+1}{m}} \right), \end{aligned}$$

where $\tilde{I}_{m,k+1}^\pm$ is given by (1.1) and double signs \pm are in the same order.

On the other hand, by (i), the following improper integrals is absolutely convergent, and by change of variable $x = \Phi(y)$ on $U \cap (-\infty, 0]$ and $y = -u$, since $x^m(1+\sum_{j=1}^\infty a_j x^j) = (f(x))^m = y^m = (-u)^m$, the following holds:

$$\begin{aligned} \int_{-\infty}^0 e^{\pm i\lambda x^m(1+\sum_{j=1}^\infty a_j x^j)} a(x) dx &= \int_{-\infty}^0 e^{\pm i\lambda y^m} a(\Phi(y)) \frac{d\Phi}{dy}(y) dy \\ &= - \int_0^\infty e^{\pm i\lambda (-u)^m} a(\Phi(-u)) \frac{d\Phi}{du}(-u) du. \end{aligned}$$

Here since $a(\Phi(-u)) \frac{d\Phi}{du}(-u) \in C_0^\infty(\mathbb{R})$, by Theorem 1.1 and Theorem 1.2 (ii), for any $N \in \mathbb{N}$ such that $N > m$, as $\lambda \rightarrow \infty$,

$$\begin{aligned} \int_{-\infty}^0 e^{\pm i\lambda x^m(1+\sum_{j=1}^\infty a_j x^j)} a(x) dx &= -Os- \int_0^\infty e^{\pm i\lambda (-u)^m} a(\Phi(-u)) \frac{d\Phi}{du}(-u) du \\ (2.2) \quad &= - \sum_{k=0}^{N-m-1} \frac{\tilde{I}_{m,k+1}^{\pm\pm m}}{k!} \left(\frac{d}{du} \right)^k \Big|_{u=0} \left\{ a(\Phi(-u)) \frac{d\Phi}{du}(-u) \right\} \lambda^{-\frac{k+1}{m}} + O\left(\lambda^{-\frac{N-m+1}{m}} \right), \end{aligned}$$

where $\tilde{I}_{m,k+1}^{\pm\pm m}$ is given by (1.2) and double signs \pm are in the same order.

Therefore by (2.1) and (2.2), for any $N \in \mathbb{N}$ such that $N > m$, as $\lambda \rightarrow \infty$,

$$\begin{aligned} & \int_{-\infty}^{\infty} e^{\pm i\lambda x^m (1 + \sum_{j=1}^{\infty} a_j x^j)} a(x) dx \\ &= \sum_{k=0}^{N-m-1} \frac{c_{m,k}^{\pm}}{k!} \left(\frac{d}{dy} \right)^k \Big|_{y=0} \left\{ a(\Phi(y)) \frac{d\Phi}{dy}(y) \right\} \lambda^{-\frac{k+1}{m}} + O\left(\lambda^{-\frac{N-m+1}{m}}\right), \end{aligned}$$

where

$$\begin{aligned} c_{m,k}^{\pm} &= \tilde{I}_{m,k+1}^{\pm} + (-1)^{k+1} \tilde{I}_{m,k+1}^{\pm \pm m} \\ &= m^{-1} \left\{ e^{\pm i \frac{\pi}{2} \frac{k+1}{m}} + (-1)^{k+1} e^{\pm (-1)^m i \frac{\pi}{2} \frac{k+1}{m}} \right\} \Gamma\left(\frac{k+1}{m}\right), \end{aligned}$$

and double signs \pm are in the same order. \square

Proof of Corollary 1.4. (i) In Theorem 1.3 (i), if $a_j = 1/j!$, since

$$\sup_{j \in \mathbb{N}} |1/j!|^{1/j} = \sup_{j \in \mathbb{N}} \prod_{k=1}^j (1/k)^{1/j} \leq 1,$$

then

$$f(x) = x \left(1 + \sum_{j=1}^{\infty} \frac{x^j}{j!} \right)^{1/p} = x e^{x/p}.$$

Here let $y = f(x)$, since $y/p = (x/p)e^{x/p}$, then

$$\frac{x}{p} = W\left(\frac{y}{p}\right),$$

where $X = W(Y)$ is a Lambert W-function on $(1/p)U \cap [0, \infty)$. And then since $\Phi(y) = x = pW(y/p)$,

$$\frac{d\Phi}{dy}(y) = \frac{d}{dy} \left\{ pW\left(\frac{y}{p}\right) \right\} = \frac{dW}{dY}\left(\frac{y}{p}\right).$$

(ii) When $p = m \in \mathbb{N}$, if $x < 0$, then we take a Lambert W-function $X = W(Y)$ on $(1/m)U \cap (\infty, 0]$, we can prove (ii) in the same manner to (i). \square

REFERENCES

- [1] R. CORLESS, G. GONNET, D. HARE, D. JEFFREY AND D. KNUTH, *On the Lambert W function*, Advances in Computational Mathematics, **5** (1996), 329–359.
- [2] J.J. DUISTERMAAT, *Fourier integral operators*, Birkhäuser, Boston. 1996.
- [3] D. FUJIWARA, *The stationary phase method with an estimate of the remainder term on a space of large dimension*, Nagoya Math. J., **124** (1991), 61–97.
- [4] D. FUJIWARA, *The Stationary phase method with estimate of remainder term over a space of large dimension*, Advanced studies in pure Mathematics **23** Spectral and Scattering Theory and applications (1994), 57–67, Mathematical Society of Japan.
- [5] D. FUJIWARA, *Mathematical method for Feynman path integrals*, Springer, Tokyo, 1999 (in Japanese).
- [6] D. FUJIWARA, *Rigorous time slicing approach to Feynman path integrals*, Springer, 2017.
- [7] D. FUJIWARA, *Asymptotic method in linear partial differential equations*, Iwanami Shoten, 2019 (in Japanese).
- [8] A. GRIGIS AND J. SJÖSTRAND, *Microlocal analysis for differential operators, an introduction*, London Mathematical Society Lecture Note Series **196**, Cambridge University Press, Cambridge, 1994.

- [9] V. GUILLMIN AND S. STERNBERG, *Geometric asymptotics*, Math. Survey **14**, American Mathematical Society, 1977.
- [10] V. GUILLMIN AND S. STERNBERG, *Symplectic techniques in physics*, Cambridge University Press, 1990.
- [11] L. HÖRMANDER, *Fourier integral operators I*, Acta Math., **127** (1971), 79–183.
- [12] L. HÖRMANDER, *The analysis of linear partial differential operators I*, Springer, 1983.
- [13] H. KUMANO-GO, *Pseudo-differential operators*, MIT Press, 1981.
- [14] H. KUMANO-GO, *Theory of Fourier integral operators and pseudo-differential operators and fundamental solution for hyperbolic equations*, Department of Mathematics, Faculty of Science, Osaka University, 1983 (in Japanese).
- [15] T. NAGANO, *An extension of phase functions on the singular points in the stationary phase method*, Master's Thesis in Tokyo University of Science, 1994 (in Japanese).
- [16] T. NAGANO AND N. MIYAZAKI, *On singular points and oscillatory integrals*, arXiv:1906.01438v1 [math.CA], 2 Jun 2019.
- [17] T. NAGANO AND N. MIYAZAKI, *Generalized Fresnel integrals as oscillatory integrals with positive real power phase functions and applications to asymptotic expansions*, arXiv:2005.12754v1 [math.CA], 24 May 2020.
- [18] T. NAGANO AND N. MIYAZAKI, *Asymptotic expansions of oscillatory integrals for certain phase functions with a degenerate singular point including type A_k , E_6 , E_8 -function germs*, arXiv:2009.09620v1 [math.CA], 21 September 2020.
- [19] H. OMORI, *Infinite dimensional Lie groups*, A.M.S. Translation Monograph, **158**, American Mathematical Society, 1997.
- [20] H. OMORI, *Algebra of calculus from the operational viewpoint I,II*, Gendai Suugaku Sya, I, 2018, II, 2019 (in Japanese).
- [21] H. OMORI, Y. MAEDA, Y. MIYAZAKI AND A. YOSHIOKA, *Deformation expressions for elements on an algebra*, Symplectic, Poisson and Noncommutative Geometry MSRI Publications, vol.**62** (2014), 171–209, Cambridge University Press.
- [22] M. SUGIURA, *Introduction to analysis I,II*, Univ. of Tokyo Press, I, 1980, II, 1985 (in Japanese).

DEPARTMENT OF LIBERAL ARTS, FACULTY OF SCIENCE AND TECHNOLOGY, TOKYO UNIVERSITY OF SCIENCE, 2641, YAMAZAKI, NODA, CHIBA 278-8510, JAPAN
 Email address: `tonagan@rs.tus.ac.jp`