

# A STATISTICAL THEORY OF HEAVY ATOMS: ENERGY AND EXCESS CHARGE

HONGSHUO CHEN, RUPERT L. FRANK, AND HEINZ SIEDENTOP

**ABSTRACT.** The purpose of this note is to give an elementary derivation of a lower bound on the relativistic Thomas-Fermi-Weizsäcker-Dirac functional of Thomas-Fermi type and to apply it to get an upper bound on the excess charge of this model.

## 1. INTRODUCTION

The description of heavy atoms suffered for a long time from the fact that the naive adaptation of Thomas-Fermi to the relativistic setting leads to a functional that is unbounded from below (see Gombas [6, §14] and [7, Chapter III, Section 16.] for reviews). As late as 1987 Engel and Dreizler [3] solved this problem deriving a relativistic Thomas-Fermi-Weizsäcker-Dirac functional  $\mathcal{E}_Z^{\text{TFWD}}$  from quantum electrodynamics. For atoms of atomic number  $Z$  and electron density  $\rho$  and velocity of light  $c$  the functional, written in Hartree units, is

$$(1) \quad \mathcal{E}_Z^{\text{TFWD}}(\rho) := \mathcal{T}^{\text{W}}(\rho) + \mathcal{T}^{\text{TF}}(\rho) - \mathcal{X}(\rho) + \mathcal{V}(\rho).$$

The first summand on the right is an inhomogeneity correction of the kinetic energy generalizing the Weizsäcker correction. Using the abbreviation  $p(x) := (3\pi^2\rho(x))^{1/3}$ ,

$$(2) \quad \mathcal{T}^{\text{W}}(\rho) := \int_{\mathbb{R}^3} dx \frac{3\lambda}{8\pi^2} (\nabla p(x))^2 c f(p(x)/c)^2$$

with  $f(t)^2 := t(t^2+1)^{-\frac{1}{2}} + 2t^2(t^2+1)^{-1} \mathfrak{A} \text{rsin}(t)$  where  $\mathfrak{A} \text{rsin}$  is the inverse function of the hyperbolic sine and  $\lambda \in \mathbb{R}_+$  is given by the gradient expansion as  $1/9$  but in the non-relativistic analogue sometimes taken as an adjustable parameter (Weizsäcker [10], Yonei and Tomishima [11], Lieb [9, 8]). The second summand is the relativistic generalization of the Thomas-Fermi kinetic energy. It is

$$(3) \quad \mathcal{T}^{\text{TF}}(\rho) := \int_{\mathbb{R}^3} dx \frac{c^5}{8\pi^2} T^{\text{TF}}\left(\frac{p(x)}{c}\right)$$

with  $T^{\text{TF}}(t) := t(t^2+1)^{3/2} + t^3(t^2+1)^{1/2} - \mathfrak{A} \text{rsin}(t) - \frac{8}{3}t^3$ . The third summand is a relativistic generalization of the exchange energy. It is

$$(4) \quad \mathcal{X}(\rho) := \int_{\mathbb{R}^3} dx \frac{c^4}{8\pi^3} X\left(\frac{p(x)}{c}\right)$$

with  $X(t) := 2t^4 - 3[t(t^2+1)^{\frac{1}{2}} - \mathfrak{A} \text{rsin}(t)]^2$ , and, eventually, the last summand is the potential energy, namely the sum of the electron-nucleus energy and the electron-electron energy. It is

$$(5) \quad \mathcal{V}(\rho) := -Z \int_{\mathbb{R}^3} dx \rho(x) |x|^{-1} + \underbrace{\frac{1}{2} \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} dy \rho(x) \rho(y) |x-y|^{-1}}_{=: \mathcal{D}[\rho]}.$$

We note that, as  $c \rightarrow \infty$ , all integrands of  $\mathcal{E}_Z^{\text{TFWD}}$  tend pointwise to the corresponding part of the non-relativistic Thomas-Fermi-Weizsäcker-Dirac functional

$$\mathcal{E}_Z^{\text{nr}}(\rho) = \int_{\mathbb{R}^3} dx \left( \frac{\lambda}{2} |\nabla \sqrt{\rho}(x)|^2 + \frac{3}{10} \gamma_{\text{TF}} \rho(x)^{\frac{5}{3}} - \frac{3}{4} \left( \frac{3}{\pi} \right)^{\frac{1}{3}} \rho(x)^{\frac{4}{3}} - \frac{Z}{|x|} \rho(x) \right) + \mathcal{D}[\rho]$$

with  $\gamma_{\text{TF}} := (3\pi^2)^{\frac{2}{3}}$  suggesting that we might expect a lower bound of Thomas-Fermi type when  $c$  is large. We will prove in Section 2 that this is indeed true. The bound will allow us to implement the method of [4] in the present context and give an improved bound on atomic excess charges. This is carried through in Section 3.

## 2. BOUND ON THE ENERGY

**2.1. The Domain of  $\mathcal{E}_Z^{\text{TFWD}}$ .** First we discuss the domain of the functional. To this end, we write  $F(t) := \int_0^t f(s) ds$  for the antiderivative of  $f$ . Then

$$(6) \quad \mathcal{T}^{\text{W}}(\rho) = \frac{3\lambda c^3}{8\pi^2} \int_{\mathbb{R}^3} dx |\nabla(F \circ (p/c)(x))|^2.$$

This allows to define  $\mathcal{E}_Z^{\text{TFWD}}$  on

$$(7) \quad P := \{\rho \in L^{\frac{4}{3}}(\mathbb{R}^3) | \rho \geq 0, \mathcal{D}[\rho] < \infty, F \circ p \in D^1(\mathbb{R}^3)\}.$$

**2.2. Lower Bound.** We turn to the lower bound itself and address the parts separately.

**2.2.1. The Weizsäcker Energy.** Since  $F(t) \geq t\sqrt{\mathfrak{A}\text{rsin}(t)}/2$  (see [2, Formula (90)]), Hardy's inequality gives the lower bound

$$(8) \quad \mathcal{T}^{\text{W}}(\rho) \geq \frac{3\lambda c}{2^7 \pi^2} \int_{\mathbb{R}^3} dx \frac{p(x)^2 \mathfrak{A}\text{rsin}(p(x)/c)}{|x|^2} = \frac{3^{\frac{5}{3}} \lambda c}{2^7 \pi^{\frac{2}{3}}} \underbrace{\int_{\mathbb{R}^3} dx \frac{\rho(x)^{\frac{2}{3}} \mathfrak{A}\text{rsin}(\frac{p(x)}{c})}{|x|^2}}_{=: \mathcal{H}(\rho)}.$$

**2.2.2. The Potential Energy.** Pick a density  $\sigma \in P$  of finite mass and set  $\varphi_{\sigma} := Z|\cdot|^{-1} - \sigma * |\cdot|^{-1}$ . Since  $\sigma$  is nonnegative, we have  $\varphi_{\sigma}(x) \leq Z/|x|$ . Then

$$(9) \quad \mathcal{V}(\rho) = - \int_{\mathbb{R}^3} dx \varphi_{\sigma}(x) \rho(x) - 2\mathcal{D}(\sigma, \rho) + \mathcal{D}[\rho] \geq - \int_{\mathbb{R}^3} dx \varphi_{\sigma}(x) \rho(x) - \mathcal{D}[\sigma].$$

Splitting the integrals at  $s$ , using (9), and Schwarz's inequality yields

$$(10) \quad \begin{aligned} \mathcal{V}(\rho) &\geq - \int_{p(x)/c < s} dx \varphi_{\sigma}(x) \rho(x) \\ &\quad - Z \int_{p(x)/c \geq s} dx \frac{\rho(x)^{\frac{1}{3}}}{|x|} \mathfrak{A}\text{rsin}(\frac{p(x)}{c})^{\frac{1}{2}} \frac{\rho(x)^{\frac{2}{3}}}{\mathfrak{A}\text{rsin}(\frac{p(x)}{c})^{\frac{1}{2}}} - \mathcal{D}[\sigma] \\ &\geq - \frac{Z}{\mathfrak{A}\text{rsin}(s)^{\frac{1}{2}}} \mathcal{H}(\rho)^{\frac{1}{2}} \mathcal{T}_{>}(\rho)^{\frac{1}{2}} - \int_{p(x)/c < s} dx \varphi_{\sigma}(x) \rho(x) - \mathcal{D}[\sigma] \end{aligned}$$

with  $\mathcal{T}_{>}(\rho) := \int_{p(x)/c > s} dx \rho(x)^{\frac{4}{3}}$ .

**2.2.3. The Thomas-Fermi Term.** First, we note that

$$(11) \quad \mathbb{R}_+ \rightarrow \mathbb{R}_+, \quad t \mapsto T^{\text{TF}}(t)/t^5,$$

is strictly monotone decreasing from  $4/5$  to  $0$  and

$$(12) \quad \mathbb{R}_+ \rightarrow \mathbb{R}_+, \quad t \mapsto T^{\text{TF}}(t)/t^4,$$

is strictly increasing from 0 to 2. Thus

$$\begin{aligned}
 \mathcal{T}^{\text{TF}}(\rho) &= \int_{p(x)/c < s} dx \frac{c^5}{8\pi^2} T^{\text{TF}}\left(\frac{p(x)}{c}\right) + \int_{p(x)/c \geq s} dx \frac{c^5}{8\pi^2} T^{\text{TF}}\left(\frac{p(x)}{c}\right) \\
 (13) \quad &\geq \int_{p(x)/c < s} dx \frac{3}{10} \frac{5T^{\text{TF}}(s)}{4s^5} \gamma_{\text{TF}} \rho(x)^{\frac{5}{3}} + \int_{p(x)/c \geq s} dx \frac{T^{\text{TF}}(s)}{s^4} \frac{3}{8} (3\pi^2)^{\frac{1}{3}} c \rho(x)^{\frac{4}{3}} \\
 &= \frac{3}{10} \frac{5T^{\text{TF}}(s)}{4s^5} \gamma_{\text{TF}} \int_{p(x)/c < s} dx \rho(x)^{\frac{5}{3}} + \frac{3}{8} \frac{T^{\text{TF}}(s)}{s^4} \gamma_{\text{TF}}^{\frac{1}{2}} c \mathcal{T}_{>}(\rho).
 \end{aligned}$$

**2.2.4. Exchange Energy.** Since  $X$  is bounded from above and  $X(t) = O(t^4)$  at  $t = 0$ , we have that for every  $\alpha \in [0, 4]$  there is a  $\xi_0$  such that  $X(t) \leq \xi_0 t^\alpha$ . We pick  $\alpha = 3$  in which case  $\xi_0 \approx 1.15$ . Thus

$$(14) \quad \mathcal{X}(\rho) \leq \frac{c\xi_0}{4\pi} N = \xi cN.$$

with  $\xi := \xi_0/(4\pi) \approx 0.0914$ .

**2.2.5. The Total Energy.** Adding everything up yields

$$\begin{aligned}
 \mathcal{E}_Z^{\text{TFWD}}(\rho) &\geq \frac{3^{\frac{5}{3}} \lambda c}{2^7 \pi^{\frac{2}{3}}} \mathcal{H}(\rho) + \frac{3}{8} \frac{T^{\text{TF}}(s)}{s^4} \gamma_{\text{TF}}^{\frac{1}{2}} c \mathcal{T}_{>}(\rho) - \frac{Z}{2\mathfrak{A}\text{rsin}(s)^{\frac{1}{2}}} \mathcal{H}(\rho)^{\frac{1}{2}} \mathcal{T}_{>}(\rho)^{\frac{1}{2}} \\
 (15) \quad &+ \int_{\frac{p(x)}{c} < s} dx \left( \underbrace{\frac{3}{10} \frac{5T^{\text{TF}}(s)}{4s^5} \gamma_{\text{TF}} \rho(x)^{\frac{5}{3}}}_{=: \gamma_e(s)} - \varphi_\sigma(x) \rho(x) \right) - \mathcal{D}[\sigma] - \xi cN.
 \end{aligned}$$

We pick  $s \in \mathbb{R}_+$  such that the sum of the first three summands of (15) is a complete square, i.e., fulfilling

$$(16) \quad \sqrt{\frac{3^{\frac{5}{3}}}{2^7 \pi^{\frac{2}{3}}} \frac{3T^{\text{TF}}(s)(3\pi^2)^{\frac{1}{3}}}{8s^4}} = \frac{Z}{c\sqrt{\lambda}} \frac{1}{2\mathfrak{A}\text{rsin}(s)^{\frac{1}{2}}}.$$

The solution is uniquely determined, since  $T^{\text{TF}}(s)/s^4$  is strictly monotone increasing from 0 to 2 and  $\mathfrak{A}\text{rsin}(s)$  is also monotone increasing from 0 to  $\infty$ . Call the corresponding  $s$   $s_0$ . Obviously,  $s_0$  depends only on  $\kappa := Z/(c\sqrt{\lambda})$  and is strictly monotone increasing from 0 to  $\infty$ .

Eventually we pick  $\sigma(x) := \rho(x)\theta(sc - p(x))$ . Summing the first three terms of the second line of (15) yields the Thomas-Fermi functional with Thomas-Fermi constant  $\gamma_e(s_0)$  evaluated at  $\sigma$ . Minimizing this functional and scaling in  $\gamma$  yields

$$(17) \quad \mathcal{E}_Z^{\text{TFWD}}(\rho) \geq -\frac{4s_0^5}{5T^{\text{TF}}(s_0)} e^{\text{TF}} Z^{\frac{7}{3}} - \xi cN$$

where  $-e^{\text{TF}}$  is the Thomas-Fermi energy of hydrogen (with the physical value of the Thomas-Fermi constant, namely  $\gamma_{\text{TF}}$ ).

The function  $s_0$  tends exponentially to  $\infty$  as  $\kappa \rightarrow \infty$ . Thus (17) is merely an exponential lower bound for large  $Z$  and fixed  $\lambda$  and  $c$ . However, if we fix  $\kappa \in \mathbb{R}_+$ , then we have a Thomas-Fermi type lower bound with a correction term linear in  $cN$ . In conclusion we have

**Theorem 1.** For given  $c, \lambda, Z \in \mathbb{R}_+$ , set  $\kappa := Z/(c\sqrt{\lambda})$ . Define  $s_0 : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  by (16), set  $\xi := \max\{X(t)/t^3 | t \in \mathbb{R}_+\}/(4\pi)$ , and write  $-e^{\text{TF}}$  for the Thomas-Fermi energy of hydrogen. Then, for all  $\rho \in P$  with  $\int_{\mathbb{R}^3} dx \rho(x) = N$ ,

$$(18) \quad \mathcal{E}_Z^{\text{TFWD}}(\rho) \geq -\frac{4s_0(\kappa)^5}{5T^{\text{TF}}(s_0(\kappa))} e^{\text{TF}} Z^{\frac{7}{3}} - \xi cN.$$

Moreover,  $s_0$  is strictly monotone increasing with  $s_0(0) = 0$  and  $s_0(\kappa) \rightarrow \infty$  as  $\kappa \rightarrow \infty$ .

## 3. APPLICATION ON THE EXCESS CHARGE PROBLEM

In this section we will show that the bound (17) allows for an adaptation of the ideas of [4, 5] and show a bound on the excess charge of the relativistic Thomas-Fermi-Weizsäcker-Dirac atom which complements the bound obtained in [2] in the absence of the exchange term.

We define a monotone increasing function  $\alpha : \mathbb{R} \rightarrow [0, \pi/2]$  by

$$(19) \quad \alpha(s) := \begin{cases} 0 & s \leq 0 \\ \frac{\pi}{2}s & s \in (0, 1) \\ \frac{\pi}{2} & s \geq 1 \end{cases}.$$

We introduce two localization functions

$$(20) \quad R := \sin \circ \alpha \text{ and } L := \cos \circ \alpha,$$

and corresponding localization functions  $U$  and  $O$  on  $\mathbb{R}^3$  defined by

$$(21) \quad U(x) := L\left(\frac{\omega \cdot x - l}{s}\right), \quad O(x) := R\left(\frac{\omega \cdot x - l}{s}\right)$$

with the parameters  $\omega \in \mathbb{S}^2$ ,  $l \in \mathbb{R}_+$ , and  $s \in (0, \infty)$ . For later use, we write  $A := \text{supp}(\nabla U)$  for the support of the gradient of  $U$  and  $O$ .

Assume  $\rho_N$ , with associated  $p_N := (3\pi^2\rho_N)^{\frac{1}{3}}$ , is a minimizer of  $\mathcal{E}_Z^{\text{TFWD}}$  under the constraint

$$(22) \quad \int_{\mathbb{R}^3} \rho(x) dx = N.$$

In abuse of notation, we sometimes write the occurring energy functionals instead of depending on  $\rho$  as depending on  $p$ , i.e.,  $p$  instead of  $p^3/(3\pi^2)$ .

Our starting point is the binding condition following directly from the variational principle by pushing the  $O$ -part away from the  $U$ -part

$$(23) \quad \mathcal{E}_Z^{\text{TFWD}}(Up_N) + \mathcal{E}_0^{\text{TFWD}}(Op_N) - \mathcal{E}_Z^{\text{TFWD}}(p_N) \geq 0$$

which is true, since

$$(24) \quad \begin{aligned} & \frac{1}{3\pi^2} \int_{\mathbb{R}^3} (U(x)^3 + O(x)^3) p_N(x)^3 dx \leq \frac{1}{3\pi^2} \int_{\mathbb{R}^3} (U(x)^2 + O(x)^2) p_N(x)^3 dx \\ & = \int_{\mathbb{R}^3} \rho_N(x) dx = N \end{aligned}$$

and the infima under the constraint (22) and the constraint

$$(25) \quad 3\pi^2 \int_{\mathbb{R}^3} p(x)^3 dx = \int_{\mathbb{R}^3} \rho(x) dx \leq N$$

agree by [1, Section 3.5]. The corresponding argument, namely pushing the charge difference between  $N$  and the charge of the minimizer to infinity, is a standard argument and works also when the Dirac term is included.

We also have by the product rule

$$(26) \quad \begin{aligned} & \int_{\mathbb{R}^3} dx [|\nabla(Up)(x)|^2 f(Up(x)/c)^2 + |\nabla(Op)(x)|^2 f(Op(x)/c)^2] \\ & \leq \int_{\mathbb{R}^3} dx [|\nabla(Up)(x)|^2 + |\nabla(Op)(x)|^2] f(p(x)/c)^2 \\ & = \int_{\mathbb{R}^3} dx |\nabla p(x)|^2 f(p(x)/c)^2 + \int_{\mathbb{R}^3} dx p(x)^2 [|\nabla U(x)|^2 + |\nabla O(x)|^2] f(p(x)/c)^2 \\ & = \int_{\mathbb{R}^3} dx |\nabla p(x)|^2 f(p(x)/c)^2 + s^{-2} \int_{\mathbb{R}^3} dx \alpha'((\omega \cdot x - l)/s)^2 p(x)^2 f(p(x)/c)^2, \end{aligned}$$

since  $f$  is monotone increasing.

An elementary calculation shows that there is a constant  $\mu$  such for all  $t \in \mathbb{R}_+$

$$(27) \quad f(t)^2 \leq \mu t.$$

The optimal constant, namely  $\max\{f(t)^2/t | t > 0\}$ , is  $\mu \approx 1.66$  achieved at  $t \approx 1.45$ .

For  $\alpha \in \mathbb{R}_+$  and  $x \in \mathbb{R}^3$  we claim

$$(28) \quad \int_{\mathbb{S}^2} \frac{d\omega}{4\pi} (\omega \cdot x - \alpha)_+ = \frac{|x|}{4} \left[ \left(1 - \frac{\alpha}{|x|}\right)_+ \right]^2$$

(see [4] for a related formula). Since the left side is independent of the direction of  $x$  and equals  $|x| \int_{\mathbb{S}^2} d\omega (4\pi)^{-1} (\omega \cdot x/|x| - \alpha/|x|)$ , it suffices to show (28) for  $x = \mathbf{e}_3$ :

$$\begin{aligned} \int_{\mathbb{S}^2} \frac{d\omega}{4\pi} (\omega \cdot \mathbf{e}_3 - \alpha)_+ &= \frac{1}{2} \int_0^\pi d\vartheta \sin \vartheta (\cos \vartheta - \alpha)_+ = \frac{1}{2} \int_{\min\{1, \alpha\}}^1 du (u - \alpha) \\ &= \frac{1}{4} [(1 - \alpha)_+]^2. \end{aligned}$$

We estimate the various parts of (23) separately. We begin with the Weizsäcker terms and get using (26) and (27)

$$\begin{aligned} &\mathcal{T}^W(U p_N) + \mathcal{T}^W(Op_N) - \mathcal{T}^W(p_N) \\ &\leq \frac{3\lambda}{8\pi^2 s^2} \int_{0 < \omega \cdot x - l < s} dx \alpha' ((\omega \cdot x - l)/s)^2 p_N(x)^2 c f(p_N(x)/c)^2 \\ (29) \quad &= \frac{3\lambda}{32s^2} \int_{0 < \omega \cdot x - l < s} dx p_N(x)^2 c f(p_N(x)/c)^2 \leq \frac{3\lambda\mu}{32s^2} \int_{0 < \omega \cdot x - l < s} dx p_N(x)^3 \\ &= \frac{9\pi^2 \lambda \mu}{32s^2} \int_{0 < \omega \cdot x - l < s} dx \rho_N(x). \end{aligned}$$

Integration over  $l \in \mathbb{R}_+$  and  $\omega \in \mathbb{S}^2$  and using (28) yields

$$\begin{aligned} &\int_{\mathbb{S}^2} \frac{d\omega}{4\pi} \int_{\mathbb{R}_+} dl (\mathcal{T}^W(U p_N) + \mathcal{T}^W(Op_N) - \mathcal{T}^W(p_N)) \\ &\leq \frac{9\pi^2 \lambda \mu}{128s^2} \int_{\mathbb{S}^2} \frac{d\omega}{\pi} \int_{\mathbb{R}_+} dx (\omega \cdot x - (\omega \cdot x - s)_+)_+ \rho_N(x) \\ (30) \quad &= \frac{9\pi^2 \lambda \mu}{128s^2} \int_{\mathbb{S}^2} \frac{d\omega}{\pi} \int_{\mathbb{R}_+} dx [(\omega \cdot x)_+ - (\omega \cdot x - s)_+] \rho_N(x) \\ &= \frac{9\pi^2 \lambda \mu}{128s^2} \int_{\mathbb{R}_+} dx |x| \left[ 1 - \left(1 - \frac{s}{|x|}\right)_+ \right]^2 \rho_N(x) \\ &= \frac{9\pi^2 \lambda \mu}{128s^2} \left( \int_{s < |x|} dx \left( 2s - \frac{s^2}{|x|} \right) \rho_N(x) + \int_{s > |x|} dx |x| \rho_N(x) \right) \leq \frac{3^2 \pi^2 \lambda \mu}{2^6} \frac{N}{s}. \end{aligned}$$

Next we estimate the combined Thomas-Fermi-Exchange part of (23). To this end we introduce the functions  $a$  and  $b$  on  $\mathbb{R}_+$  defined by

$$(31) \quad a(t) := \frac{c^5}{8\pi^2} T^{\text{TF}}(t) + \frac{c^4}{8\pi^3} 3[t(t^2 + 1)^{\frac{1}{2}} - \mathfrak{A} \text{rsin}(t)]^2.$$

$$(32) \quad b(t) := \frac{c^4}{8\pi^3} 2t^4.$$

Pick now  $f_1, \dots, f_n \in \mathbb{R}_+$  with  $f_1^2 + \dots + f_n^2 = 1$ . Since  $a, \dots, a^{(iv)}$  are all positive, we have

$$(33) \quad a'''(f_i t) \leq a'''(t)$$

because  $f_i \leq 1$ . Since also  $a(0) = a'(0) = a''(0) = a'''(0)$ , integration of (33) yields successively  $a''(f_\nu t) \leq f_\nu a''(t)$ ,  $a'(f_\nu t) \leq f_\nu^2 a'(t)$ , and  $a(f_\nu t) \leq f_\nu^3 a(t)$ . (See [1, Formula 3.135] for a similar argument for  $T^{\text{TF}}(t)$ .)

Thus, we get

$$\begin{aligned}
& \mathcal{T}^{\text{TF}}(Up_N) - \mathcal{X}(Up_N) + \mathcal{T}^{\text{TF}}(Op_N) - \mathcal{X}(Op_N) - (\mathcal{T}^{\text{TF}}(p_N) - \mathcal{X}(p_N)) \\
&= \int_{\mathbb{R}^3} dx [a(U(x)\frac{p_N(x)}{c}) + a(O(x)\frac{p_N(x)}{c}) - a(\frac{p_N(x)}{c}) \\
&\quad + b(\frac{p_N(x)}{c}) - b(U(x)\frac{p_N(x)}{c}) - b(O(x)\frac{p_N(x)}{c})] \\
&\leq \int_{\mathbb{R}^3} dx [(U(x)^3 + O(x)^3 - 1)a(\frac{p_N(x)}{c}) - (U(x)^4 + O(x)^4 - 1)b(\frac{p_N(x)}{c})] \\
(34) \quad &\leq \frac{1}{4} \max \left\{ \underbrace{\frac{(1 - \cos(t)^4 - \sin(t)^4)^2}{1 - \cos(t)^3 - \sin(t)^3}}_{=(2+\sqrt{2})/4} \middle| t \in [0, \pi/2] \right\} \int_A dx \frac{b(\frac{p_N(x)}{c})^2}{a(\frac{p_N(x)}{c})} \\
&\leq \frac{2 + \sqrt{2}}{2^5} \frac{c^3}{\pi^4} \int_A dx \frac{(\frac{p_N(x)}{c})^8}{T^{\text{TF}}(\frac{p_N(x)}{c})}.
\end{aligned}$$

Using (11) and (12) we get for any  $S \in \mathbb{R}_+$

$$\begin{aligned}
& \mathcal{T}^{\text{TF}}(Up_N) - \mathcal{X}(Up_N) + \mathcal{T}^{\text{TF}}(Op_N) - \mathcal{X}(Op_N) \\
&\quad - (\mathcal{T}^{\text{TF}}(p_N) - \mathcal{X}(p_N)) \\
(35) \quad &\leq \frac{2 + \sqrt{2}}{2^5 \pi^4} \int_{A, p_N(x)/c < S} dx \frac{S^5}{T^{\text{TF}}(S)} p_N(x)^3 \\
&\quad + \frac{2 + \sqrt{2}}{2^5 \pi^4} \int_{A, p_N(x)/c \geq S} dx \frac{S^4}{c T^{\text{TF}}(S)} p_N(x)^4 \\
&\leq \frac{(2 + \sqrt{2})3}{2^5 \pi^2} \frac{S^5}{T^{\text{TF}}(S)} N + \frac{(2 + \sqrt{2})3^{\frac{4}{3}}}{2^5 \pi^{\frac{4}{3}} c} \frac{S^4}{T^{\text{TF}}(S)} \int_{\mathbb{R}^3} dx \rho_N(x)^{\frac{4}{3}}.
\end{aligned}$$

The external potential part yields

$$(36) \quad -Z \int_{\mathbb{R}^3} dx \frac{(U(x)^3 - 1)\rho_N(x)}{|x|} \leq Z \int_{\omega \cdot x - l > 0} dx \frac{\rho_N(x)}{|x|}.$$

Integration over  $l$  and averaging over the sphere yields

$$\begin{aligned}
& -Z \int_{\mathbb{S}^2} \frac{d\omega}{4\pi} \int_{\mathbb{R}_+} dl \int_{\mathbb{R}^3} dx \frac{(U(x)^3 - 1)\rho_N(x)}{|x|} \\
(37) \quad &\leq Z \int_{\mathbb{S}^2} \frac{d\omega}{4\pi} \int_{\mathbb{R}_+} dl \int_{\omega \cdot x - l > 0} dx \frac{\rho_N(x)}{|x|} \\
&= Z \int_{\mathbb{S}^2} \frac{d\omega}{4\pi} \int_{\mathbb{R}^3} dx \frac{(\omega \cdot x) + \rho_N(x)}{|x|} = \frac{Z}{4} \int_{\mathbb{R}^3} dx \rho_N(x) = \frac{Z}{4} N.
\end{aligned}$$

Finally, we address the electron-electron repulsion in (23). We have

$$\begin{aligned}
(38) \quad W(l, \omega) &:= D(U^3 \rho_N, U^3 \rho_N) + D(O^3 \rho_N, O^3 \rho_N) \\
&\quad - D((U^2 + O^2)\rho_N, (U^2 + O^2)\rho_N) \\
&\leq -2D(U^2 \rho_N, O^2 \rho_N) \leq - \int_{\omega \cdot x - l < 0} dx \int_{\omega \cdot y - l > s} \frac{\rho_N(x)\rho_N(y)}{|x - y|}.
\end{aligned}$$

Integration in  $l$  and  $\omega$  and using (28) yields

$$\begin{aligned}
(39) \quad & \int_{\mathbb{S}^2} \frac{d\omega}{4\pi} \int_{\mathbb{R}_+} dl W(l, \omega) \leq - \int_{\mathbb{S}^2} \frac{d\omega}{4\pi} \int_{\mathbb{R}_+} dl \int_{\omega \cdot x - l < 0} dx \int_{\omega \cdot y - l > s} dy \frac{\rho_N(x) \rho_N(y)}{|x - y|} \\
& = - \int_{\mathbb{S}^2} \frac{d\omega}{8\pi} \int_{\mathbb{R}_+} dl \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} dy \\
& \quad [\theta(l - \omega \cdot x) \theta(\omega \cdot y - s - l) + \theta(l - (-\omega \cdot y)) \theta(-\omega \cdot x - s - l)] \frac{\rho_N(x) \rho_N(y)}{|x - y|} \\
& = - \int_{\mathbb{S}^2} \frac{d\omega}{8\pi} \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} dy \frac{(\omega \cdot (y - x) - s)_+ \rho_N(x) \rho_N(y)}{|x - y|}.
\end{aligned}$$

Thus, with (28),

$$\begin{aligned}
(40) \quad & \int_{\mathbb{S}^2} \frac{d\omega}{4\pi} \int_{\mathbb{R}_+} dl W(l, \omega) \leq - \frac{1}{8} \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} dy \rho_N(x) \rho_N(y) \left[ \left( 1 - \frac{s}{|x - y|} \right)_+ \right]^2 \\
& = - \frac{N^2}{8} + \frac{1}{8} \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} dy \rho_N(x) \rho_N(y) \left\{ 1 - \left[ \left( 1 - \frac{s}{|x - y|} \right)_+ \right]^2 \right\} \\
& = - \frac{N^2}{8} + \frac{1}{8} \int_{\mathbb{R}^3} dx \int_{\mathbb{R}^3} dy \rho_N(x) \rho_N(y) \times \begin{cases} 1 & s \geq |x - y| \\ \frac{2s}{|x - y|} - \left( \frac{s}{|x - y|} \right)^2 & s < |x - y| \end{cases} \\
& \leq - \frac{N^2}{8} + \frac{s}{2} \mathcal{D}[\rho_N].
\end{aligned}$$

Inserting these estimates in (23) gives

$$(41) \quad \frac{3^2 \pi^2 \lambda \mu}{2^6} \frac{N}{s} + \frac{ZN}{4} - \frac{N^2}{8} + c_1(S)N + \frac{c_2(S)}{c} \int_{\mathbb{R}^3} dx \rho_N(x)^{\frac{4}{3}} + \frac{s}{2} \mathcal{D}[\rho_N] \geq 0$$

or

$$(42) \quad N \leq 2Z + \frac{3^2 \pi^2 \lambda \mu}{2^3 s} + 2^2 s \frac{\mathcal{D}[\rho_N]}{N} + 8c_1(S) + \frac{8c_2(S)}{cN} \int_{\mathbb{R}^3} dx \rho_N(x)^{\frac{4}{3}}$$

and after optimization in  $s$

$$(43) \quad N \leq 2Z + 3\pi\sqrt{2} \sqrt{\frac{\lambda\mu\mathcal{D}[\rho_N]}{N}} + 8c_1(S) + \frac{8c_2(S)}{cN} \int_{\mathbb{R}^3} dx \rho_N(x)^{\frac{4}{3}}.$$

Now, we assume  $\kappa = Z/(c\sqrt{\lambda})$  fixed and apply Theorem 1 with a factor 2 in front of the exchange term and  $Z$  replaced by  $2Z$ . This gives (18) but with the corresponding replacements, namely  $Z$  by  $2Z$  and a factor 2 in front of  $\xi cN$ . Therefore we get

$$\begin{aligned}
(44) \quad & 0 \geq \mathcal{E}_Z^{\text{TFWD}}(\rho_N) = \frac{1}{2} \mathcal{T}^W(\rho_N) + \frac{1}{2} \mathcal{T}^{\text{TF}}(\rho_N) + \frac{1}{2} \mathcal{D}[\rho_N] \\
& \quad + \frac{1}{2} \left( \mathcal{T}^W(\rho_N) + \mathcal{T}^{\text{TF}}(\rho_N) + \mathcal{D}[\rho_N] - \int_{\mathbb{R}^3} dx \frac{Z \rho_N(x)}{2|x|} - 2\mathcal{X}(\rho_N) \right) \\
& \geq \frac{1}{2} \mathcal{T}^W(\rho_N) + \frac{1}{2} \mathcal{T}^{\text{TF}}(\rho_N) + \frac{1}{2} \mathcal{D}[\rho_N] - C_\kappa Z^{7/3} - \xi cN.
\end{aligned}$$

Thus, all three terms,  $\mathcal{T}^W(\rho_N)$ ,  $\mathcal{T}^{\text{TF}}(\rho_N)$ , and  $\mathcal{D}(\rho_N)$  are bounded by a constant times  $Z^{7/3} + cN$ .

Now,  $T^{\text{TF}}(t) \geq 2t^4 - (8/3)t^3$ . Thus

$$(45) \quad \begin{aligned} \int_{\mathbb{R}^3} \rho_N(x)^{\frac{4}{3}} &= \frac{c^4}{(2\pi^2)^{\frac{4}{3}}} \int_{\mathbb{R}^3} dx \left( \frac{p_N(x)}{c} \right)^4 \\ &\leq \left( \frac{2}{\pi} \right)^{\frac{2}{3}} (c^{-1} \mathcal{T}^{\text{TF}}(\rho_N) + cN) \leq D_\kappa (Z^{\frac{4}{3}} \lambda^{\frac{1}{2}} + N + cN) \end{aligned}$$

with a  $\kappa$ -dependent constant  $D_\kappa$ . Thus, (43) yields the following bound on the excess charge.

**Theorem 2.** *Assume that  $\rho \in P$  with  $\mathcal{E}_Z^{\text{TFWD}}(\rho) = \inf \mathcal{E}_Z^{\text{TFWD}}(P)$ , set  $N := \int_{\mathbb{R}^3} \rho(x) dx$ , and assume  $\kappa$  and  $\lambda$  positive and fixed. Then, for large  $Z$ ,*

$$(46) \quad N \leq 2Z + O(Z^{\frac{2}{3}}).$$

This should be compared to the bound  $N \leq 2.56Z$  of [2, Formula (18)] for the relativistic Thomas-Fermi-Weizsäcker functional without exchange, i.e., even with exchange term included we are lead to an improved leading order. Note, however, it comes at a price, namely the ratio  $Z/c$  and  $\lambda$  is now fixed.

#### ACKNOWLEDGMENTS

Partial support by the U.S. National Science Foundation through grants DMS-1363432 and DMS-1954995 (R.L.F.) and by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) through Germany's Excellence Strategy EXC - 2111 - 390814868 (R.L.F., H.S.) is gratefully acknowledged. H.C. and H.S. thank the Institute of Mathematical Sciences at the National University of Singapore for support through the program *Density Functionals for Many-Particle Systems: Mathematical Theory and Physical Applications of Effective Equations* which inspired this work.

#### REFERENCES

- [1] Hongshuo Chen. *On the Excess Charge Problem in Relativistic Quantum Mechanics*. PhD thesis, Ludwig-Maximilians-Universität München, July 2019.
- [2] Hongshuo Chen and Heinz Siedentop. On the excess charge of a relativistic statistical model of molecules with an inhomogeneity correction. *Journal of Physics A: Mathematical and Theoretical*, 53(39):395201, September 2020.
- [3] E. Engel and R. M. Dreizler. Field-theoretical approach to a relativistic Thomas-Fermi-Dirac-Weizsäcker model. *Phys. Rev. A*, 35:3607–3618, May 1987.
- [4] Rupert L Frank, Phan Thành Nam, and Hanne Van Den Bosch. The ionization conjecture in Thomas-Fermi-Dirac-von Weizsäcker theory. *Communications on Pure and Applied Mathematics*, 71(3):577–614, 2018.
- [5] Rupert L. Frank, Phan Thành Nam, and Hanne Van Den Bosch. The maximal excess charge in Müller density-matrix-functional theory. *Ann. Henri Poincaré*, 19(9):2839–2867, 2018.
- [6] P. Gombás. *Die statistische Theorie des Atoms und ihre Anwendungen*. Springer-Verlag, Wien, 1 edition, 1949.
- [7] P. Gombás. Statistische Behandlung des Atoms. In S. Flügge, editor, *Handbuch der Physik. Atome II*, volume 36, pages 109–231. Springer-Verlag, Berlin, 1956.
- [8] Elliott H. Lieb. Analysis of the Thomas-Fermi-von Weizsäcker equation for an infinite atom without electron repulsion. *Comm. Math. Phys.*, 85(1):15–25, 1982.
- [9] Elliott H. Lieb and David A. Liberman. Numerical calculation of the Thomas-Fermi-von Weizsäcker function for an infinite atom without electron repulsion. Technical Report LA-9186-MS, Los Alamos National Laboratory, Los Alamos, New Mexico, April 1982.
- [10] C. F. v. Weizsäcker. Zur Theorie der Kernmassen. *Z. Phys.*, 96:431–458, 1935.
- [11] Katsumi Yonei and Yasuo Tomishima. On the Weizsäcker correction to the Thomas-Fermi theory of the atom. *Journal of the Physical Society of Japan*, 20(6):1051–1057, 1965.



COLLEGE OF MATHEMATICS AND STATISTICS, CHONGQING UNIVERSITY, CHONGQING 401331,  
CHINA

*Email address:* hongshuo.chen@gmail.com

MATHEMATISCHES INSTITUT, LUDWIG-MAXIMILANS UNIVERSITÄT MÜNCHEN, THERESIENSTR.  
39, 80333 MÜNCHEN, GERMANY, AND MUNICH CENTER FOR QUANTUM SCIENCE AND TECHNOLOGY  
(MCQST), SCHELLINGSTR. 4, 80799 MÜNCHEN, GERMANY, AND MATHEMATICS 253-37, CALTECH,  
PASADENA, CA 91125, USA

*Email address:* rlfrank@caltech.edu

MATHEMATISCHES INSTITUT, LUDWIG-MAXIMILANS UNIVERSITÄT MÜNCHEN, THERESIENSTRASSE  
39, 80333 MÜNCHEN, GERMANY, AND MUNICH CENTER FOR QUANTUM SCIENCE AND TECHNOLOGY  
(MCQST), SCHELLINGSTR. 4, 80799 MÜNCHEN, GERMANY

*Email address:* h.s@lmu.de