A possible connection between the reflection symmetry and existence of equatorial circular orbit

Sayak Datta * and Sajal Mukherjee[†]

Inter-University Centre for Astronomy and Astrophysics, Post Bag 4, Pune-411007, India

October 26, 2020

Abstract

We study a viable connection between the circular-equatorial orbits and reflection symmetry across the equatorial plane of a stationary axis-symmetric metric. Behavior of the circular equatorial orbits in the direction perpendicular to the equatorial plane are studied, and different outcomes in the presence and in the absence of the reflection symmetry are discussed. We conclude that in the absence of the equatorial reflection symmetry, neither stable nor unstable circular orbit can exist on the equatorial plane. Moreover, to address the observational aspects, we provide two possible examples relating gravitational wave (GW) astronomy and structure formation in galaxy which can put constrain on the symmetry breaking parameters.

1 Introduction

The Kerr metric uniquely describes a stationary, axissymmetric and asymptotically flat black hole (BH) solution of vacuum Einstein's field equations [1]. Beside the stationary and axis-symmetry properties, Kerr spacetime is also endowed with an additional feature of reflection symmetry across the equatorial plane. Even if the former characteristics are likely to be associated with astrophysical objects with rotation, both asymptotic flatness and equatorial reflection symmetry can be relaxed in order to probe larger domain of compact objects other than BH. Various possible distinction between BHs and other exotic compact objects (ECOs) based on tidal deformability [2-5], tidal heating [4, 6-9], multipole moments [6, 10], echoes in postmerger [11–16] has been proposed in literature. Similarly, distinguishing them on the basis of equatorial reflection symmetry can be useful to detect them or rule out as viable astrophysical bodies. Not only these studies may provide a fresh outlook to model astrophysical objects, but also assign an observational impact to it. In the present article, we aim to elaborate on the equatorial reflection symmetry in a generic spacetime, and outline its possible theoretical and observational implications in depth.

In order to study any particular effect appearing from spacetime geometry, the ideal approach is to begin with orbital dynamics. Based on the fact how orbits behave in a

given spacetime, more involved astrophysical searches are constructed. In Kerr, the orbital properties are well studied, and extensively explored in literature [17–19]. Similar exploration is carried out for Kerr-NUT spacetime [20-22], which violates the equatorial reflection symmetry, and describes an asymptotically non-flat geometry [23, 24]. While in Kerr, we know stable/unstable equatorial circular orbits exist, the same is not true in the presence of NUT charge. In particular, neither stable nor unstable equatorial circular orbits can exist for massive or massless particles in Kerr-NUT spacetime [20, 21]. In fact, this stark contrast between Kerr and Kerr-NUT is the primary source of our motivation to study further, and investigate whether equatorial reflection symmetry and existence of the equatorial circular orbits can be generically connected. We introduce a perturbative approach for confronting equatorial circular geodesics in geometries where equatorial reflection symmetry is absent. By assuming that the orbits reside on the equatorial plane initially, we study the growth of perturbation in time, and aim to realize which parameters engineer any possible deviation from the equatorial plane. Assuming the perpendicular to $\theta = \pi/2$ is along z-axis, we consider the z-perturbation in our work. Due to the absence of the equatorial reflection symmetry, it is likely that the potential will not be an even function of z. As a result, there will be an intrinsic force in the z-direction that will lift the orbits from the equatorial plane. To study this in the context of a general stationary

^{*}skdatta@iucaa.in

[†]sajal@iucaa.in

axis-symmetric metric we will consider Ernst potential and write metric components in terms of it.

The existence of the planner circular orbits is crucial to discuss any physical effect associated with a binary. The orbits in extreme mass ratio inspirals (EMRI), that will be observed with LISA [25], are most likely to be generic [26–28]. Binaries with stellar masses, as already detected by gravitational wave (GW) detectors LIGO and VIRGO, can have precession [29]. This means that the understandings found in the current paper will not only have a theoretical ground but also it will have an observational impact, which will be discussed later.

The rest of the manuscript is organized as follows. In Section (2), we start with the Ernst potential, and in Section (3), we will briefly discuss the geodesic equations in terms of the metric components. The primary findings of the paper are given in Section (4) and Section (5) respectively. In Section (6), we will study observational impacts of current findings, and finally, we will conclude in Section (7).

2 Metric components and the Ernst potential

We start with a general stationary and axis-symmetric spacetime written in cylindrical coordinates (t, ρ, z, ϕ) as follows [30, 31]:

$$ds^{2} = -F(dt - \omega d\phi)^{2} + \frac{1}{F} \left[e^{2\gamma} (d\rho^{2} + dz^{2}) + \rho^{2} d\phi^{2} \right], \quad (1)$$

where F, ω , and γ are functions of ρ and z. Substituting the metric in the Einstein equation, it is possible to find governing equations for these entities. In passing, we should note that the above metric does not guarantee to be asymptotically flat, and remains general otherwise.

The above metric components can be written in a more compact form by using the complex Ernst potential which is a combination of both *norm* (λ) and *twist* (ω). In particular, $\lambda = -g_{tt} = F$, and $\omega_{\mu} = \sqrt{-g} \epsilon_{\mu\nu\gamma\delta} \xi^{\nu} \nabla^{\gamma} \xi^{\delta}$, where ξ^{μ} is the timelike killing vector [31]. Given that the spacetime is stationary and axis-symmetric, both norm and twist are expected to be nonzero. Finally, the Ernst potential takes the following form:

$$\mathcal{E} = F + i\psi = \frac{(\rho^2 + z^2)^{1/2} - \tilde{\xi}}{(\rho^2 + z^2)^{1/2} + \tilde{\xi}},$$
(2)

where $\tilde{\xi}$ can be written as [32]:

$$\tilde{\xi} = \sum_{j,k=0}^{\infty} a_{jk} \frac{\rho^j z^k}{(\rho^2 + z^2)^{j+k}}.$$
(3)

The reasons to choose Ernst potential formalism as a tool to express metric components are two-folded. Firstly, the metric components can be written directly in terms of \mathcal{E} . As a result, from the behaviour of \mathcal{E} under the absence of the symmetries, the nature of the orbits can easily be extracted. And secondly, reflection symmetry manifests itself through the values of a_{ik} , making it easier to impose the presence or the absence of equatorial reflection symmetry. The a_{jk} is nonzero only for non-negative, even j and nonnegative k. If there is reflection symmetry across the equatorial plane, then a_{ik} is real for even k and imaginary for odd k [30, 32–34]. However, as the present study remains general as far as the equatorial reflection symmetry is concerned, we restrain ourselves to make such assumptions. We assume that a_{ik} has both real and imaginary components for both even and odd values of k.

In terms of F and ψ , the metric components g_{tt} , $g_{t\phi}$ and $g_{\phi\phi}$, which will be of particular use, can be written as follows [30]

$$g_{tt} = -F, \quad g_{\phi\phi} = 1/g_{tt} \left\{ g_{t\phi}^2 - \rho^2 \right\},$$

$$g_{t\phi} = -F \int_{z=const} \frac{\rho}{F^2} \frac{\partial \psi}{\partial z} d\rho + F \int_{\rho=const} \frac{\rho}{F^2} \frac{\partial \psi}{\partial \rho} dz, \quad (4)$$

where, the detail calculations to arrive at the following expressions are shown in Appendix A.

3 Off-equatorial perturbation of the equatorial geodesics

In order to study the existence of equatorial circular orbits in a generic spacetime with metric given in Eq. (1), we start with the following geodesic equations:

$$\ddot{\rho} = \frac{1}{2} \frac{\partial \mathcal{V}_{\rho}}{\partial \rho}, \quad \ddot{z} = \frac{1}{2} \frac{\partial \mathcal{V}_{z}}{\partial z}, \tag{5}$$

where, 'dot' defines a derivative with respect to the affine parameter which we may call τ , \mathcal{V}_{ρ} and \mathcal{V}_z are radial and angular potential respectively. The above equations will determine locations of a orbit while the conserved energy and momentum are dictated by t and ϕ components. Given that we are interested in orbits confined on a plane and circular in nature, the above two equations would give $\ddot{\rho} = \ddot{z} = 0$ in principle. However, as the equatorial reflection symmetry is not respected, the potential \mathcal{V}_z is likely to contain terms with odd power of z such that $\mathcal{V}_z(z) \neq \mathcal{V}_z(-z)$. One should be mindful that condition of circularity is not expected to be affected by the equatorial reflection symmetry, and we may safely impose $\dot{\rho} = \ddot{\rho} = 0$. From the timelike constraint, $\mathcal{U}^{\alpha}\mathcal{U}_{\alpha} = -1$, we arrive at the following expression:

$$\mathcal{U}^t \mathcal{U}_t + \mathcal{U}^\rho \mathcal{U}_\rho + \mathcal{U}^z \mathcal{U}_z + \mathcal{U}^\phi \mathcal{U}_\phi = -1, \tag{6}$$

and finally [35]

$$\mathcal{V}_{z} = g_{zz} (\mathcal{U}^{z})^{2} = -1 - g^{tt} E^{2} - g^{\phi\phi} L_{z}^{2} - 2g^{t\phi} E L_{z}, \qquad (7)$$

where E and L_z are given as conserved energy and momentum respectively, appearing from the spacetime symmetries. We will expand \mathcal{V}_z about the equatorial plane, i.e., z = 0. Neglecting terms $\sim \mathcal{O}(\delta z)^3$ and beyond, i.e., $\mathcal{V}_z(\delta z) = \mathcal{V}_0 + (\delta z)\mathcal{V}_1 + (\delta z)^2\mathcal{V}_2$, we may rewrite z-equation in Eq. (5) as follows:

$$2\delta z = \mathcal{V}_1 + 2\mathcal{V}_2\delta z. \tag{8}$$

By solving the above equation, we arrive at

$$\delta z(\tau) = \frac{\mathcal{V}_1}{2\omega^2} + A \exp(-i\omega\tau) + B \exp(i\omega\tau), \qquad (9)$$

where we set $\mathcal{V}_2 = -\omega^2$, and ω can be both real and imaginary. If we assume that the initial conditions are, $\delta z(0) = \delta z(0) = 0$, the above equation may be rewritten as

$$\delta z(\tau) = -\frac{\mathcal{V}_1}{2\omega^2} + \frac{\mathcal{V}_1}{4\omega^2} \left\{ \exp(-i\omega\tau) + \exp(i\omega\tau) \right\}.$$
(10)

Depending on the nature of ω , the above equation either represent a hyperbola (Im[ω] \neq 0), and a oscillatory (Im[ω] = 0) motion. For the Kerr-NUT spacetime, as we have shown in Appendix B, Im[ω] = 0, and the solution is always oscillatory. For future purpose, we may note that in case of a oscillatory solution, we have,

$$\delta z(\tau) = -\frac{\mathcal{V}_1}{\omega^2} \sin^2(\omega \tau/2), \qquad (11)$$

which hints an interesting property of this motion written as follows. This perturbation is either positive or negative depending on the sign of \mathcal{V}_1 , but never switches sign. It indicates that it would never cross the equatorial plane, but approach it in each cycle. In short, the hobbling from the equatorial plane would be one-sided.

It should be mentioned that the above equation builtin with a condition where there is no external perturbation otherwise $\delta z(0) \neq 0$. Any additional external perturbation may result in some changes in the final expression which we have not studied in this article. Besides, one should also note how \mathcal{V}_1 and \mathcal{V}_2 are affecting the perturbation. While \mathcal{V}_1 is directly proportional to its value, \mathcal{V}_2 is responsible for engineering its nature. By setting $\mathcal{V}_2 = 0$, we obtain a diverging nature of the perturbation, as $\delta z(\tau) \sim \tau^2$, and may not appropriately model $\delta z(\tau)$. Keeping these points in mind we will keep \mathcal{V}_1 and \mathcal{V}_2 in the expression, and ignore higher-order corrections. Eventually, we will show that nonzero \mathcal{V}_1 is connected to the absence of equatorial reflection symmetry, which, as a result does not allow a circular orbit to exist in the equatorial plane, not even perturbatively.

4 Decomposition of the Ernst potential

Referring to Eq. (3), we may state that equatorial reflection symmetry in the potential comes through a_{jk} . For a clear exposition of our results, we separate out the real and imaginary part of a_{jk} , i.e.,

$$a_{jk} = \hat{a}_{jk} + i\breve{a}_{jk},\tag{12}$$

where \hat{a}_{jk} and \check{a}_{jk} are the real and imaginary part of the a_{jk} , respectively. If the equatorial reflection symmetry exists, then a_{jk} is real for even k and imaginary for odd k [30, 32–34],

$$\check{a}_{j(2m)} = 0, \quad \hat{a}_{j(2m+1)} = 0,$$
(13)

for all non-negative values of m. However, in the present context, we should note again that the above equations are not valid, and expected to be nonzero. With the above expressions in hand, we now attempt to connect the potential with a_{jk} , and start with decomposing $\tilde{\xi}$ in real and imaginary parts as follows,

$$\tilde{\xi} = \hat{\xi} + i\breve{\xi}.\tag{14}$$

Therefore, we gather

$$\hat{\xi} = \sum_{j,k=0}^{\infty} \hat{a}_{jk} \frac{\rho^j z^k}{(\rho^2 + z^2)^{j+k}}, \quad \breve{\xi} = \sum_{j,k=0}^{\infty} \breve{a}_{jk} \frac{\rho^j z^k}{(\rho^2 + z^2)^{j+k}}.$$
(15)

By using Eq. (3), Eq. (14) and Eq. (15), we may arrive at the following expression:

$$F = \frac{R - \hat{\xi}^2 - \breve{\xi}^2}{(R^{1/2} + \hat{\xi})^2 + \breve{\xi}^2}, \quad \psi = \frac{-2R^{1/2}\breve{\xi}}{(R^{1/2} + \hat{\xi})^2 + \breve{\xi}^2}, \quad (16)$$

where $R = \rho^2 + z^2$. For our purpose, we need to understand the properties of the $\partial \mathcal{V}_z / \partial z$ up to order of z, and considering \mathcal{V}_z up to order of z^2 . This would need the knowledge of $g^{tt}, g^{t\phi}, g^{\phi\phi}, F, g_{t\phi}$ up to the order of z^2 and ψ up to the order of z^3 . Therefore, we need contribution from $\hat{a}_{j0}, \hat{a}_{j1}$ and $\check{a}_{j0}, \check{a}_{j1}$ for our analysis. For the expansion, we take $z/\rho \ll 1$, and as a result, it is not necessary for z to be very small as long as $z \ll \rho$ is satisfied. Keeping up to order z^3/ρ^3 ,

$$\hat{\xi} = \sum_{j} \left[\frac{\hat{a}_{j0}}{\rho^{j}} \{ 1 - j \frac{z^{2}}{\rho^{2}} \} + \frac{\hat{a}_{j1}}{\rho^{j+1}} \{ \frac{z}{\rho} - (j+1) \frac{z^{3}}{\rho^{3}} \} \right],$$
$$\breve{\xi} = \sum_{j} \left[\frac{\breve{a}_{j0}}{\rho^{j}} \{ 1 - j \frac{z^{2}}{\rho^{2}} \} + \frac{\breve{a}_{j1}}{\rho^{j+1}} \{ \frac{z}{\rho} - (j+1) \frac{z^{3}}{\rho^{3}} \} \right], \quad (17)$$

Using the results found in this section we will find the $\partial \mathcal{V}_z/\partial z$ in the next section.

5 Results

In this section, we expressed the relevant quantities as a series expansion in z, and only keeping terms up to z^2 terms in the potential. Based on our earlier discussions, here, we will derive \mathcal{V}_1 and \mathcal{V}_2 for general stationery, axis-symmetric metric. By substituting Eq. (17) into Eq. (16), we arrive at

$$F = \frac{1 + \sum_{0}^{3} g_{i} z^{i} / \rho^{i}}{1 + \sum_{0}^{3} f_{i} z^{i} / \rho^{i}} \equiv \sum_{0}^{3} F_{i} z^{i}, \ \psi = \frac{\sum_{0}^{3} \psi_{i} z^{i} / \rho^{i}}{\rho^{2} (1 + \sum_{0}^{3} f_{i} z^{i} / \rho^{i})},$$
(18)

where, the coefficients can be expressed as follows:

$$\begin{split} f_{0} &= -g_{0} = \sum_{j} \hat{a}_{j0} \frac{2}{\rho^{j+1}} + \Big[\sum_{j,j'} \frac{1}{\rho^{j+j'+2}} \{ \hat{a}_{j0} \hat{a}_{j'0} \\ &\quad + \hat{a} \to \breve{a} \} \Big], \\ f_{1} &= -g_{1} = \sum_{j} \hat{a}_{j1} \frac{2}{\rho^{j+2}} + \Big[\sum_{j,j'} \frac{1}{\rho^{j+j'+3}} 2\{ \hat{a}_{j0} \hat{a}_{j'1} \\ &\quad + \hat{a} \to \breve{a} \} \Big] \\ f_{2} &= 2 - g_{2} = 1 + \sum_{j} - \hat{a}_{j0} \frac{2}{\rho^{j+1}} \left(j - \frac{1}{2} \right) + \Big[\sum_{j,j'} \frac{1}{\rho^{j+j'+2}} \\ &\quad \{ -\hat{a}_{j0} \hat{a}_{j'0} (j+j') + \frac{\hat{a}_{j1} \hat{a}_{j'1}}{\rho^{2}} + \hat{a} \to \breve{a} \} \Big] \\ \psi_{0} &= \sum_{j} \breve{a}_{j0} \frac{2}{\rho^{j-1}}, \ \psi_{1} = \sum_{j} \breve{a}_{j1} \frac{2}{\rho^{j}}, \\ \psi_{2} &= \sum_{i} - \breve{a}_{j0} \frac{2}{\rho^{j-1}} \left(j - \frac{1}{2} \right), \\ \psi_{3} = \sum_{i} - \breve{a}_{j1} \frac{2}{\rho^{j}} \left(j + \frac{1}{2} \right), \end{split}$$

$$F_{1} = -\frac{2f_{1}}{(f_{0}+1)^{2}\rho},$$

$$F_{2} = \frac{2\left[f_{0}^{2}+2f_{0}+f_{1}^{2}-(f_{0}+1)f_{2}+1\right]}{(f_{0}+1)^{3}\rho^{2}},$$

$$F_{3} = -\frac{2\left[f_{1}^{3}+(f_{0}+1)(f_{0}-2f_{2}+1)f_{1}+(f_{0}+1)^{2}f_{3}\right]}{(f_{0}+1)^{4}\rho^{3}},$$
(19)

which are function of ρ only, and independent of z. By employing these expressions, we can obtain the derivatives of the metric components, g_{tt} , $g_{t\phi}$ and $g_{\phi\phi}$ (given in Eq. (4)), which are essential for our study. We start with the following expression for $g_{t\phi}$ and obtain the derivative of $g_{t\phi}$ and $g_{\phi\phi}$:

$$g_{t\phi} = -F\sum_{i} z^{i} \int I_{i} d\rho' + N = -F\sum_{i} z^{i} \mathcal{I}_{i} + \sum_{i} N_{i} z^{i}, \quad (20)$$

$$\frac{\partial g^{t\phi}}{\partial z} = -\frac{F}{\rho^2} \int [I_1 + 2I_2 z] d\rho' + (N_1 + 2N_2 z),$$

$$\frac{\partial g^{\phi\phi}}{\partial z} = \frac{1}{\rho^2} [F_1 + 2F_2 z],$$
(21)

where we have used Eq. (18), and the expressions for N_1, N_2, I_0, I_1, I_2 are given below:

$$N_{1} = -\frac{2\psi_{0}}{f_{0}\rho^{2}(1+f_{0})},$$

$$N_{2} = \frac{1}{2\rho^{2}f_{0}(1+f_{0})} \left[\frac{3\psi_{0}f_{1}}{\rho(1+f_{0})} - \frac{3\psi_{1}}{\rho}\right]$$

$$I_{0} = \frac{(f_{0}+1)\psi_{1} - f_{1}\psi_{0}}{(f_{0}-1)^{2}\rho^{2}},$$

$$I_{1} = \frac{2}{(f_{0}-1)^{3}\rho^{3}} \left((f_{0}^{2}-1)\psi_{2} - f_{2}\psi_{0}(f_{0}-1) - f_{1}\psi_{1}(f_{0}+1) + f_{1}^{2}\psi_{0}\right),$$

$$I_{2} = \frac{1}{(f_{0}-1)^{4}\rho^{4}} \left(3f_{0}^{3}\psi_{3} - 3f_{3}f_{0}^{2}\psi_{0} - 3f_{2}f_{0}^{2}\psi_{1} + 4f_{0}^{2}\psi_{1} - 3f_{1}f_{0}^{2}\psi_{2} - 3f_{0}^{2}\psi_{3} - 4f_{1}f_{0}\psi_{0} + 6f_{1}f_{2}f_{0}\psi_{0} + 6f_{3}f_{0}\psi_{0} + 3f_{1}^{2}f_{0}\psi_{1} + 2f_{2}f_{0}\psi_{1} - 2f_{1}f_{0}\psi_{2} - 3f_{0}\psi_{3} - 3f_{1}^{3}\psi_{0} + 4f_{1}\psi_{0} - 6f_{1}f_{2}\psi_{0} - 3f_{3}\psi_{0} + 3f_{1}^{2}\psi_{1} + f_{2}\psi_{1} + 5f_{1}\psi_{2} - 4\psi_{1} + 3\psi_{3}\right).$$

It is now easy to evaluate the derivative of g_{tt} by using Eq. (21) and Eq. (22) as follows

$$\frac{\partial g^{tt}}{\partial z} = \frac{1}{\rho^2 F} \left(2g_{t\phi} \frac{\partial g_{t\phi}}{\partial z} \right) - \frac{1}{F^2} \frac{\partial F}{\partial z} \left(\frac{g_{t\phi}^2 - \rho^2}{\rho^2} \right) \\
= \left(-\frac{\mathcal{I}_0^2 F_1}{\rho^2} + \frac{2\mathcal{I}_1 \mathcal{I}_0 F_0}{\rho^2} + \frac{F_1}{F_0^2} \right) \\
+ \frac{1}{F_0^3 \rho^2} \left\{ 2z \left(F_2 F_0 \left(\rho^2 - \mathcal{I}_0^2 F_0^2 \right) \right) \\
+ \left(\mathcal{I}_1^2 + 2\mathcal{I}_0 \mathcal{I}_2 \right) F_0^4 - F_1^2 \rho^2 \right) \right\}$$
(22)

Finally, we can obtain the expression for $\partial \mathcal{V}_z(z)/\partial z$ by using Eq. (7) and the derivatives of $g_{t\phi}$, and, $g_{\phi\phi}$ given in Eq. (21), and derivative of g_{tt} evaluated in Eq. (22). The result is as follows:

$$\mathcal{V}_{1} = -E^{2} \left(-\frac{\mathcal{I}_{0}^{2}F_{1}}{\rho^{2}} + 2\frac{\mathcal{I}_{0}\mathcal{I}_{1}F_{0}}{\rho^{2}} + \frac{F_{1}}{F_{0}^{2}} \right) - \frac{L_{z}^{2}}{\rho^{2}}F_{1}$$
$$-2EL_{z}N_{1} + 2\frac{EL_{z}}{\rho^{2}}F_{0}\mathcal{I}_{1}$$
(23)

$$\mathcal{V}_{2} = -\frac{E^{2}}{F_{0}^{3}\rho^{2}} \left\{ F_{2}F_{0}(\rho^{2} - \mathcal{I}_{0}^{2}F_{0}^{2}) + (\mathcal{I}_{1}^{2} + 2\mathcal{I}_{0}\mathcal{I}_{2})F_{0}^{4} - F_{1}^{2}\rho^{2} \right\} - \frac{L_{z}^{2}}{\rho^{2}}F_{2} - 2EL_{z}N_{2} + \frac{EL_{z}}{\rho^{2}} \left(2\mathcal{I}_{2}F_{0} + \mathcal{I}_{1}F_{1} \right).$$

$$(24)$$

It is easy to notice that in general $\mathcal{V}_1 \neq 0$ from Eq. (23). The consequence of non-zero \mathcal{V}_1 has already been discussed in Section (3). It may be possible that even though the zindependent term is non-zero for each of the derivative, in some special cases they will add up to give a vanishing zindependent term. In particular, one may ask whether it is possible to choose conserved energy and momentum in such a way that it would result in $\mathcal{V}_z = 0$. Indeed, we find this can be a possibility, however, both the energy and momentum need to be in consonance with $\dot{\rho} = \ddot{\rho} = 0$ too, which can put further restriction in its motion. For example, in the Kerr-NUT spacetime, it is not possible to choose energy and momentum such a way that it would, not only cancel the off-equatorial push, but also describe a circular geodesic [21]. Therefore, while this is a valid possibility, it does not describe a general outcome of our findings.

For a quick follow up of the above analysis where the equatorial reflection symmetry is respected, we focus on the z independent term in Eq. (22) and Eq. (21). Here the z independent part depends only on F_0, F_1, I_0, I_1, N_1 . When equatorial reflection symmetry is present, $\check{a}_{j(2m)} = 0$ and $\hat{a}_{j(2m+1)} = 0$, hence, $f_1 = \psi_0 = \psi_2 = 0$. This implies $F_1 = I_1 = \mathcal{I}_1 = N_1 = 0$. Therefore, the \mathcal{V}_1 in Eq. (23) vanishes. It is remarkable that in the case of equatorial reflection symmetry, the z^0 terms in every term that constitutes $\frac{\partial \mathcal{V}_z}{\partial z}$ vanishes, and as a result of which \mathcal{V}_1 vanishes. This indicates that the absence of any oscillatory (across equatorial plane and under z perturbation) circular equatorial orbit implies absence of the reflection symmetry.

6 Observational prospects and possible constraints

We explore a possible connection between the equatorial reflection symmetry and equatorial circular orbits, and arrive at the conclusion that it is not possible to find "small oscillation" solution under a perturbation in the z-direction around a circular orbit in general. The parameters which break the reflection symmetry will engineer to elevate the orbit from equatorial plane and boosts it with a force. However, if you assume these parameters are small enough, we can still naively assume the orbits to be equatorial and still perform some astrophysical calculations [36]. Nonetheless, this would introduce a test-bed to execute several observational expedition to confirm the existence of stable orbits and equatorial reflection symmetry. In fact, the observation of stable circular equatorial orbit will be a tell-tale signature of the presence of reflection symmetry or a very mild violation of it. For the present purpose, we outline few of such examples where the violation of equatorial reflection symmetry may be detectable by observation.

6.1 Gravitational wave astronomy

Consider a binary system prepared in such a way that the spins of the components are either aligned or anti-aligned with the orbital angular momentum. But with time due to the push from \mathcal{V}_1 the spins will not stay aligned even if they were prepared in an aligned manner. This will result in a non-zero in-plane spin component (χ_p) (Check Ref. [37] for the definition). Therefore, for equatorial reflection symmetry violating bodies in a binary, χ_p measurement should be non-zero. Hence, non-zero χ_p can arise due to several different ways. One is due to the formation mechanism of binaries, which has components that respect equatorial reflection symmetry, yet introduce a nonzero χ_p possibly due to spin effects. Another reason for non-zero χ_p would be due to the absence of equatorial reflection symmetry. This means that there will be a degeneracy between the formation channel and equatorial reflection symmetry violation, and makes it uncertain to arrive at a unique conclusion.

The other investigation is consists with a possible test related to multipole moments of a compact object. It is well-known that for axis-symmetric bodies, there can be two sets multipole moments — one is mass moment (M_l) and another is current moment (S_l) . For metric solution with the equatorial reflection symmetry like Kerr, both the odd mass and even current moments would identically vanish. However, for a simple illustration of these moments in a Kerr-NUT spacetime which is known to break the equatorial reflection symmetry, one immediately notices that all the orders for mass and current multipole moments would survive [36]. From observation perspective, there now exists contemporary tools to measure the quadrupole moment in a binary [6, 10]. If reflection symmetry is violated then it is likely that the metric will have nonzero M_{2l+1} and/or S_{2l} , i.e. classes of fuzzball solutions [38-42]. Hence, measuring a nonzero S_2 will be a signature of breaking of equatorial reflection symmetry, and by the arguments laid before, it is expected that a nonzero χ_p should also be observed. But a simultaneous measurement of nonzero S_2 and M_3 along with zero χ_p , therefore will be a signature of deviation from general relativity.

6.2 Constraints from the galaxy structures

Most of the galaxies usually contain very compact supermassive objects at their centers. If the metric of these objects does not respect equatorial reflection symmetry then there should be some imprint of such violation on the matter distribution around it. It is observed that galaxies typically shows a thin disc structure located close to the equator plane [43], and any possible violation of equatorial reflection symmetry would invite a deviation from this. Depending on the value of \mathcal{V}_1 , we may expect the matter to be distributed in off-equatorial planes. However, as positive outlook, this can also give a possible opportunity to constraint \mathcal{V}_1 from observation. For example, if we assume that there is a deviation from the reflection symmetry, we gather from Eq. (11) that furthest a particle can go from the equatorial plane is $\mathcal{D} \sim |\frac{\mathcal{V}_1}{\omega^2}|$. Now if we assume that the observed thin-disk structure is retained, we need to ensure that the even the furthest flying particle is safely within the radius of influence, R_{inf} (here we have ignored the small scale structures of the disk, and only concentrating on the bulk properties of it). By assuming the presence of a supermassive body in the galactic centre, R_{inf} is expressed as $R_{\text{inf}} = GM/\sigma^2$, where M is the mass of the BH and σ is the velocity dispersion [44]. Therefore, to ensure a disk structure consistent with most of the observation, we have $\mathcal{D} < R_{\text{inf}}$ which translates into

$$\left|\frac{\mathcal{V}_1}{\omega^2}\right| < 3.3 \times 10^{15} \mathrm{m} \left(\frac{M}{10^6 M_{\odot}}\right) \left(\frac{200 \mathrm{\ km/sec}}{\sigma}\right)^2.$$
(25)

For a given mass of the central body and its velocity dispersion, we may be able to constrain the reflection breaking parameters from observation, which may provide a handful number of information about the central object.

7 Conclusion

We have studied the perturbation of the circular-equatorial orbits of a general stationary axis-symmetric metric in the zdirection. In the process, we have identified a set of parameters that are the potential source of the equatorial reflection symmetry violation, namely $\check{a}_{j(2m)}$ and $\hat{a}_{j(2m+1)}$. We have shown that in general, when these parameters are zero (nonzero) equatorial reflection symmetry is present (absent) and the small oscillation solution across equatorial plane is present (absent). This leads us to conclude that the very existence of equatorial circular orbit is an indication that the geometry respects equatorial reflection symmetry, while it may not be true other way round. To be precise, equatorial circular geodesics may not exist even if the spacetime respects the equatorial reflection symmetry. For example, this can be simply stemmed from the fact that the conserved momentum and energy are not favourable to host any bound circular geodesic.

In this paper, we argue that in case of equatorial reflection symmetry violation, it is unlikely to have a measurement with $\chi_p = 0$, as far as one is concerned with Einsteinian gravity. This can be addressed by properly identifying the orbital parameters that will be representative of symmetry violation, and possibly depend on $\check{a}_{j(2m)}$ and $\hat{a}_{j(2m+1)}$. Nonetheless, we claim that the breaking of equatorial reflection symmetry may also indicate different mechanisms associated with the binary formation — suggesting a degeneracy. Therefore, to break this stalemate, we need to confront the multipolar structure of the object which would consist of both odd mass multipole moments and even current multipole moments in case the symmetry is violated. By measuring the S_2 and M_3 components, it would be sufficed to confirm this claim. In passing, we should note that measuring non-vanishing S_2 and M_3 , along with the measurement of χ_p consistent with a vanishing value may be a signature of the violation of general relativity. We also constrained equatorial reflection symmetry violating parameters for the supermassive objects in the galaxy centers. As per our knowledge, this is the first time when such kind of constraint has been found.

Acknowledgments

Both the authors are indebted to Sukanta Bose, Sumanta Chakraborty, Naresh Dadhich, Sanjit Mitra and Kanak Saha, for useful comments, and also suggesting changes for the betterment of the article. S.D. would like to thank University Grants Commission (UGC), India, for providing a senior research fellowship, and, S.M is thankful to the Department of Science and Technology (DST), Government of India, for financial support.

A Metric components in terms of the Ernst potential

In this section we will derive $g_{t\phi}$ in terms of the Ernst potential. As per our knowledge this has not been computed explicitly in the literature. If the metric is stationary and axis-symmetric then there will exist a time like Killing vector field ξ^{α} and an axial Killing vactor field \mathfrak{A}^{α} . Then it is possible to define a vector field ω^{α} defined as,

$$\omega_{\alpha} = \epsilon_{\alpha\beta\gamma\delta} \xi^{\alpha} \nabla^{\beta} \xi^{\gamma} \tag{A.1}$$

which satisfies, $\nabla_{[\alpha}\omega_{\beta]} = 0$. Therefore, we can define a twist potential ψ as $\omega_{\alpha} = \partial_{\alpha}\psi$.

In the cylindrical co-ordinate system (t, ρ, z, ϕ) the nonzero components can be found as,

$$\begin{split} \omega_{\rho} &= -(g_{\rho\rho}g_{zz})^{1/2}g^{zz} \left[g^{t\phi}\partial_{z}g_{tt} + g^{\phi\phi}\partial_{z}g_{t\phi}\right] \\ &= \frac{1}{\rho}g_{tt}^{2}\partial_{z} \left(\frac{g_{t\phi}}{g_{tt}}\right), \\ \omega_{z} &= (g_{\rho\rho}g_{zz})^{1/2}g^{\rho\rho} \left[g^{\phi\phi}\partial_{\rho}g_{t\phi} + g^{t\phi}\partial_{\rho}g_{tt}\right] \\ &= -\frac{1}{\rho}g_{tt}^{2}\partial_{\rho} \left(\frac{g_{t\phi}}{g_{tt}}\right). \end{split}$$
(A.2)

From Eq. (A.2) it is simple to find $g_{t\phi}$ as,

$$\frac{g_{t\phi}}{g_{tt}} = \int \frac{\rho}{g_{tt}^2} \partial_\rho \psi \ dz - \int \frac{\rho}{g_{tt}^2} \partial_z \psi \ d\rho. \tag{A.3}$$

B Example of a equatorial reflection symmetry breaking spacetime: Kerr-NUT geometry

In order display the connection between equatorial symmetry and circular orbits explicitly, we consider an example, namely Kerr-NUT spacetime, where the reflection symmetry is known to be violated [21]. The non-existence of equatorial circular orbits in Kerr-NUT geometry is first claimed in Ref. [20], and recently explored further in Ref. [21]. In the present context though, we will be more involved in studying the perturbation equation in theta direction, and discuss the near equatorial plane behavior.

The Kerr-NUT spacetime is a solution to vacuum Einstein field equations and describes a stationary, axissymmetric and asymptotically non-flat spacetime. In order to discuss the nature of the angular perturbation, we need to consider the angular geodesic equation in Boyer-Lindquist coordinates (t, r, θ, ϕ) :

$$\mathcal{V}_{\theta} = (\dot{\theta})^2 = \left(\frac{d\theta}{d\tau_{\rm m}}\right)^2 = \left(\lambda \sin^2 \theta + (L_{\rm z} - aE)^2 \sin^2 \theta - (EP - L_{\rm z})^2 - \sin^2 \theta (l + a\cos\theta)^2\right),$$
(B.1)

where $\tau_{\rm m}$ is the Mino time, 'dot' defines a derivative with respect to it, and \mathcal{V}_{θ} can be defined as angular potential [21]. The quantities λ , E, L_z , l, a are defined as Carter constant, energy, momentum, NUT charge, angular momentum respectively; and ρ , P are given by, $\rho^2 = r^2 + (l + a \cos \theta)^2$, $P = a \sin^2 \theta - 2l \cos \theta$. Let us now assume that the particle is initially confined on the equatorial plane and we expect to study the perturbation originated from the term $\hat{\theta}$. By setting $\dot{\theta} = 0$ and $\theta = \pi/2$, we gather $\lambda = l^2$, which identically vanishes for zero NUT charge, i.e., in the case of Kerr spacetime. In addition to $\theta = 0$, to have a planner orbit, we need to ensure $\ddot{\theta} = 0$ too, which warrants that $\dot{\theta} = 0$ remains satisfied along the trajectory. This is where the NUT charge comes into play, and engineers to disobey $\ddot{\theta} = 0$. Even then, $\ddot{\theta}$ equation can be useful to extract information about the variation of θ near to the equatorial plane, which we will do next. Let us start by introducing the following equation,

$$\ddot{\theta} = \frac{1}{2} \frac{d\mathcal{V}(\theta)}{d\theta},\tag{B.2}$$

which we need to write in terms of the coordinate time, such that it can be useful for an asymptotic observer. To execute this task, we may unfold $\ddot{\theta}$ as follows:

$$\frac{d^2\theta}{d\tau^2} = \left(\frac{d^2\theta}{dt^2}\right) \left(\frac{dt}{d\tau}\right)^2 + \left(\frac{d\theta}{d\tau}\right) \left(\frac{dt}{d\tau}\right)^{-1} \left[\frac{d}{dt} \left(\frac{dt}{d\tau}\right) \frac{dt}{d\tau}\right],\tag{B.3}$$

and one easily notices that the second term goes to zero, as we assume $\dot{\theta} = 0$ in the first place. Bring together Eq. (B.2) and Eq. (B.3), and writing $\theta(t)$ as $\theta(t) = \pi/2 + \delta\theta(t)$, we arrive at the following expression:

$$\left(\frac{d^2\delta\theta(t)}{dt^2}\right)\left(\frac{dt}{d\tau}\right)^2 = \frac{1}{2}\frac{d\mathcal{V}(\theta)}{d\theta}.$$
 (B.4)

The expression of $\mathcal{U}^t = dt/d\tau$, can be derived from, $\mathcal{U}^t = -g^{tt}E + g^{t\phi}L_z$, where the metric components are explicitly written in the Appendix. Finally, assuming $\delta\theta(t) \ll 1$, and terms with $\delta\theta(t)^2$, $\delta\theta(t)\delta\theta'(t)$, $\delta\theta(t)\delta\theta''(t)$ and beyond are neglected, we gather:

$$\mathcal{C}_3 \delta \theta''(t) + \mathcal{C}_2 \delta \theta(t) + \mathcal{C}_1 = 0, \qquad (B.5)$$

where 'prime' denotes a differentiation with respect to t. The expressions for C_1 , C_2 and C_3 are given as follows:

$$C_{1} = l \left\{ a(2E^{2} - 1) - 2EL_{z} \right\},$$

$$C_{2} = L_{z}^{2} + 4E^{2}l^{2} - a^{2}(E^{2} - 1),$$

$$C_{3} = \frac{1}{\Delta^{2}} \left\{ E(r^{2} + l^{2})^{2} - 2aL_{z}(l^{2} + Mr) + a^{2}E\left(r(r + 2M) + 3l^{2}\right) \right\},$$

$$(B.6)$$

where, $\Delta = r^2 - 2Mr + a^2 - l^2$ becomes zero on the event horizon. The above equation has a generic solution of the form:

$$\delta\theta(t) = -\frac{\mathcal{C}_1}{\mathcal{C}_2} + A\cos\omega t + B\sin\omega t, \qquad (B.7)$$

where A and B are integration constants to be evaluated from the initial conditions, and $\omega = \sqrt{C_2/C_3}$. It is interesting to point out that C_3 diverges on the horizon, and ω becomes zero, which no longer describes a equatorial timelike circular orbits. This is in consonance with the fact that on the null surface of event horizon, no timelike circular orbit is possible to exist. With the initial condition $\delta\theta(t=0) = \delta\theta'(t=0) = 0$, the above equation turns out to be

$$\delta\theta(t) = -\frac{\mathcal{C}_1}{\mathcal{C}_2} \left(1 - \cos\omega t\right) = -\frac{2\mathcal{C}_1}{\mathcal{C}_2} \sin^2(\omega t/2).$$
(B.8)

which may oscillate on either side of $\theta = \pi/2$, depending on the signs of C_1 and C_2 . Finally, we should say some word regarding the nature of the oscillations and how it is different from the Kerr case. It is easy to realize that the oscillation solely depends on C_1 and C_2 , and among them C_2 is always positive as far as we are concerned with bound circular geodesics, i.e., $E \leq 1$. Coming to C_1 , it can only vanish if we have l = 0, or find a radius which satisfies $(2E^2 - 1) = 2EL_z$. Turns out that the later option is ruled out as far as one is interested with equatorial circular orbits in NUT spacetime [21], and one is left with no choice but to set l = 0 to stop the oscillation. Therefore, the NUT charge, which is entirely responsible to break the equatorial symmetry, also engineers to angular perturbation. In passing, we should also mention the similar scenario in connection to massless particles. In this case too, it is possible to arrive at an equation equivalent to Eq. (B.8), only with the expressions of C_1 , C_2 and C_3 changed as follows:

$$C_{1} = 2El \left\{ aE - L_{z} \right\},$$

$$C_{2} = L_{z}^{2} + E^{2}(4l^{2} - a^{2}) + l^{2},$$

$$C_{3} = \frac{1}{\Delta^{2}} \left\{ E(r^{2} + l^{2})^{2} - 2aL_{z}(l^{2} + Mr) + a^{2}E\left(r(r + 2M) + 3l^{2}\right) \right\},$$

$$(B.9)$$

The expression for C_1 can be set to zero by two possible ways, namely, l = 0, and $aE = L_z$. Like the earlier case, the later option can never gives rise to a circular geodesic, and we need to choose l = 0. Therefore, the massless case is also in agreement with our claim that the NUT charge is solely responsible for having no equatorial circular orbits. Note that the general study representing the same for an arbitrary spacetime with no equatorial symmetry is already presented in Section (5).

References

- R. P. Kerr, "Gravitational field of a spinning mass as an example of algebraically special metrics," *Phys. Rev. Lett.* **11** (1963) 237–238.
- [2] V. Cardoso, E. Franzin, A. Maselli, P. Pani, and G. Raposo, "Testing strong-field gravity with tidal Love numbers," *Phys. Rev.* D95 no. 8, (2017) 084014, arXiv:1701.01116 [gr-qc]. [Addendum: Phys. Rev.D95,no.8,089901(2017)].
- [3] N. Sennett, T. Hinderer, J. Steinhoff, A. Buonanno, and S. Ossokine, "Distinguishing Boson Stars from Black Holes and Neutron Stars from Tidal Interactions in Inspiraling Binary Systems," *Phys. Rev.* D96 no. 2, (2017) 024002, arXiv:1704.08651 [gr-qc].
- [4] A. Maselli, P. Pani, V. Cardoso, T. Abdelsalhin, L. Gualtieri, and V. Ferrari, "Probing Planckian corrections at the horizon scale with LISA binaries," *Phys. Rev. Lett.* **120** no. 8, (2018) 081101, arXiv:1703.10612 [gr-qc].
- [5] R. Brustein and Y. Sherf, "Quantum Love," arXiv:2008.02738 [gr-qc].
- [6] S. Datta and S. Bose, "Probing the nature of central objects in extreme-mass-ratio inspirals with

gravitational waves," *Phys. Rev. D* **99** no. 8, (2019) 084001, arXiv:1902.01723 [gr-qc].

- [7] S. Datta, R. Brito, S. Bose, P. Pani, and S. A. Hughes, "Tidal heating as a discriminator for horizons in extreme mass ratio inspirals," *Phys. Rev.* D101 no. 4, (2020) 044004, arXiv:1910.07841 [gr-qc].
- [8] S. Datta, K. S. Phukon, and S. Bose, "Recognizing black holes in gravitational-wave observations: Telling apart impostors in mass-gap binaries," arXiv:2004.05974 [gr-qc].
- [9] S. Datta, "Tidal heating of Quantum Black Holes and their imprints on gravitational waves," arXiv:2002.04480 [gr-qc].
- [10] N. Krishnendu, K. Arun, and C. K. Mishra, "Testing the binary black hole nature of a compact binary coalescence," *Phys. Rev. Lett.* **119** no. 9, (2017) 091101, arXiv:1701.06318 [gr-qc].
- [11] V. Cardoso, E. Franzin, and P. Pani, "Is the gravitational-wave ringdown a probe of the event horizon?," *Phys. Rev. Lett.* **116** no. 17, (2016) 171101, arXiv:1602.07309 [gr-qc]. [Erratum: Phys.Rev.Lett. 117, 089902 (2016)].
- [12] V. Cardoso, S. Hopper, C. F. B. Macedo, C. Palenzuela, and P. Pani, "Gravitational-wave signatures of exotic compact objects and of quantum corrections at the horizon scale," *Phys. Rev.* D94 no. 8, (2016) 084031, arXiv:1608.08637 [gr-qc].
- [13] K. W. Tsang, A. Ghosh, A. Samajdar,
 K. Chatziioannou, S. Mastrogiovanni, M. Agathos, and C. Van Den Broeck, "A morphology-independent search for gravitational wave echoes in data from the first and second observing runs of Advanced LIGO and Advanced Virgo," *Phys. Rev. D* 101 no. 6, (2020) 064012, arXiv:1906.11168 [gr-qc].
- [14] J. Abedi, H. Dykaar, and N. Afshordi, "Echoes from the Abyss: Tentative evidence for Planck-scale structure at black hole horizons," *Phys. Rev.* D96 no. 8, (2017) 082004, arXiv:1612.00266 [gr-qc].
- [15] J. Westerweck, A. Nielsen, O. Fischer-Birnholtz, M. Cabero, C. Capano, T. Dent, B. Krishnan,

G. Meadors, and A. H. Nitz, "Low significance of evidence for black hole echoes in gravitational wave data," *Phys. Rev.* **D97** no. 12, (2018) 124037, arXiv:1712.09966 [gr-qc].

- [16] V. Cardoso and P. Pani, "Testing the nature of dark compact objects: a status report," *Living Rev. Rel.* 22 no. 1, (2019) 4, arXiv:1904.05363 [gr-qc].
- [17] D. C. Wilkins, "Bound Geodesics in the Kerr Metric," *Phys. Rev. D* 5 (1972) 814–822.
- [18] B. O'Neill, The geometry of Kerr black holes. Courier Corporation, 2014.
- [19] S. Chandrasekhar, The mathematical theory of black holes, vol. 69. Oxford University Press, 1998.
- [20] P. Jefremov and V. Perlick, "Circular motion in NUT space-time," *Class. Quant. Grav.* 33 no. 24, (2016) 245014, arXiv:1608.06218 [gr-qc]. [Erratum: Class.Quant.Grav. 35, 179501 (2018)].
- [21] S. Mukherjee, S. Chakraborty, and N. Dadhich, "On some novel features of the Kerr-Newman-NUT spacetime," *Eur. Phys. J. C* 79 no. 2, (2019) 161, arXiv:1807.02216 [gr-qc].
- [22] C. Chakraborty and S. Bhattacharyya, "Circular orbits in Kerr-Taub-NUT spacetime and their implications for accreting black holes and naked singularities," *JCAP* 05 (2019) 034, arXiv:1901.04233 [astro-ph.HE].
- [23] E. Newman, L. Tamubrino, and T. Unti, "Empty space generalization of the Schwarzschild metric," *J. Math. Phys.* 4 (1963) 915.
- [24] D. Lynden-Bell and M. Nouri-Zonoz, "Classical monopoles: Newton, NUT space, gravimagnetic lensing and atomic spectra," *Rev. Mod. Phys.* **70** (1998) 427–446, arXiv:gr-qc/9612049 [gr-qc].
- [25] LISA Collaboration, P. Amaro-Seoane *et al.*, "Laser Interferometer Space Antenna," arXiv:1702.00786 [astro-ph.IM].
- [26] P. Amaro-Seoane, J. R. Gair, A. Pound, S. A. Hughes, and C. F. Sopuerta, "Research Update on Extreme-Mass-Ratio Inspirals," *J. Phys. Conf. Ser.* **610** no. 1, (2015) 012002, arXiv:1410.0958 [astro-ph.CO].

- [27] L. Barack and C. Cutler, "LISA capture sources: Approximate waveforms, signal-to-noise ratios, and parameter estimation accuracy," *Phys. Rev. D* 69 (2004) 082005, arXiv:gr-qc/0310125.
- [28] D. Merritt, T. Alexander, S. Mikkola, and C. M. Will, "Stellar Dynamics of Extreme-Mass-Ratio Inspirals," *Phys. Rev. D* 84 (2011) 044024, arXiv:1102.3180 [astro-ph.CO].
- [29] T. A. Apostolatos, C. Cutler, G. J. Sussman, and K. S. Thorne, "Spin induced orbital precession and its modulation of the gravitational wave forms from merging binaries," *Phys. Rev. D* 49 (1994) 6274–6297.
- [30] F. Ryan, "Gravitational waves from the inspiral of a compact object into a massive, axisymmetric body with arbitrary multipole moments," *Phys. Rev. D* 52 (1995) 5707–5718.
- [31] R. M. Wald, *General relativity*. University of Chicago press, 2010.
- [32] G. Fodor, C. Hoenselaers, and Z. Perjés, "Multipole moments of axisymmetric systems in relativity," *Journal of Mathematical Physics* **30** no. 10, (1989) 2252–2257.
- [33] F. J. Ernst, V. S. Manko, and E. Ruiz, "Equatorial symmetry / antisymmetry of stationary axisymmetric electrovac spacetimes," *Class. Quant. Grav.* 23 (2006) 4945–4952, arXiv:gr-qc/0701112.
- [34] F. J. Ernst, V. S. Manko, and E. Ruiz, "Equatorial symmetry/antisymmetry of stationary axisymmetric electrovac spacetimes. II.," *Class. Quant. Grav.* 24 (2007) 2193–2204, arXiv:gr-qc/0701113.
- [35] Y. Mino, "Perturbative approach to an orbital evolution around a supermassive black hole," *Phys. Rev. D* 67 (2003) 084027, arXiv:gr-qc/0302075.
- [36] S. Mukherjee and S. Chakraborty, "Multipole moments of compact objects with NUT charge: Theoretical and observational implications," arXiv:2008.06891 [gr-qc].
- [37] P. Schmidt, F. Ohme, and M. Hannam, "Towards models of gravitational waveforms from generic binaries II: Modelling precession effects with a single effective precession parameter," *Phys. Rev. D* 91 no. 2, (2015) 024043, arXiv:1408.1810 [gr-qc].

- [38] M. Bianchi, D. Consoli, A. Grillo, J. F. Morales, P. Pani, and G. Raposo, "The multipolar structure of fuzzballs," arXiv:2008.01445 [hep-th].
- [39] M. Bianchi, D. Consoli, A. Grillo, J. F. Morales, P. Pani, and G. Raposo, "Distinguishing fuzzballs from black holes through their multipolar structure," arXiv:2007.01743 [hep-th].
- [40] I. L. R. Bena, Black Holes, Black Rings and their Microstates. PhD thesis, Saclay, 2009.
- [41] G. Gibbons and N. Warner, "Global structure of five-dimensional fuzzballs,"

Class. Quant. Grav. **31** (2014) 025016, arXiv:1305.0957 [hep-th].

- [42] B. Bates and F. Denef, "Exact solutions for supersymmetric stationary black hole composites," *JHEP* 11 (2011) 127, arXiv:hep-th/0304094.
- [43] M. S. Longair, Galaxy formation. Springer Science & Business Media, 2007.
- [44] A. Feoli, "The m bh versus m $g\sigma 2$ relation and the accretion of supermassive black holes," *The* Astrophysical Journal **784** no. 1, (2014) 34.