

ANISOTROPIC DYNAMICAL HORIZONS ARISING IN GRAVITATIONAL COLLAPSE

XINLIANG AN AND QING HAN

ABSTRACT. In this paper, we study the evolution of the $3 + 1$ dimensional Einstein vacuum equations with anisotropic characteristic initial data. In a framework set up by Christodoulou and under a general anisotropic condition introduced by Klainerman, Luk and Rodnianski, we prove that *in the process of gravitational collapse, a smooth and spacelike apparent horizon (dynamical horizon) emerges from general (both isotropic and anisotropic) initial data and the corresponding black hole entropy is always strictly increasing toward the future.* This extends the theory of black hole thermodynamics along the apparent horizon to anisotropic scenarios. For the proof, we combine scale-critical hyperbolic method and non-perturbative elliptic techniques. Furthermore, we construct explicit single and multi-valley anisotropic apparent horizons. They are the first mathematical examples of the anisotropic marginally outer trapped surface (MOTS) and the anisotropic apparent horizon formed in dynamics.

CONTENTS

1. Introduction	1
2. Settings	13
3. Subsolutions and Supersolutions	17
4. Moser's Iteration and Schauder Estimates	22
5. The Linearized Equation	31
6. The Existence and Properties of Solutions	37
7. Anisotropic Apparent Horizons	42
Appendix A. Estimates of Geometric Quantities	48
References	50

1. INTRODUCTION

In this paper, we study the evolution of the $3 + 1$ dimensional Einstein vacuum equations (EVEs)

$$(1.1) \quad \text{Ric}_{\mu\nu} = 0$$

with anisotropic characteristic initial data. We will demonstrate that a *smooth spacelike anisotropic* apparent horizon will arise in the corresponding process of gravitational collapse.

Date: October 26, 2020.

1.1. Motivations. In 2009, Christodoulou published a breakthrough result. In a 589-page monumental work [11], Christodoulou showed that in the Einstein vacuum spacetime a trapped surface can form dynamically from arbitrarily dispersed initial data. A trapped surface is a two dimensional sphere with both future null expansions negative, and it indicates a strong field (black hole) region is formed. Together with the Penrose's incompleteness theorem, a trapped surface also predicts future geodesically incompleteness.

Later works by Klainerman and Rodnianski [21] and by the first author [1] simplified [11] in a systematical way by introducing signatures for short-pulse and signatures for decay rates. The work by the first author and Luk [4] further extended Christodoulou's result to the scale-critical regime. Recently, the first author [2] gave an alternative proof of [11] and [4] by using only signatures for decay rates and by employing a spacetime scaling argument.

In these aforementioned works, to form a trapped surface in the gravitational collapse, we require that the initial mass input of each angle of \mathbb{S}^2 satisfies a uniform non-zero lower bound and a uniform upper bound. In another word, the characteristic initial data of EVEs are not spherically symmetric, but they are also not very anisotropic. (The details of anisotropy will be explained in Theorem 7.1 and Theorem 7.4.) In [19], Klainerman, Luk, and Rodnianski studied the case of fully anisotropic initial data and showed that a trapped surface can also form dynamically. Their initial data allow that the initial mass input of most (almost all) angles of \mathbb{S}^2 to be *zero* and only along some angles they have a uniform non-zero lower bound.

So far, we have a relatively satisfactory theory of trapped surface formation. The next challenging question is to find the outer boundary of these trapped surfaces, which is called the apparent horizon. It is closely tied to the black hole thermodynamics in physics.

In 1970s, Bekenstein and Hawking found that black hole spacetimes are associated with an amazing property. At each fixed time t , the area of the black-hole event horizon could be viewed as an entropy of the spacetime (at time t) and it is increasing toward the future. In general relativity, we could define two types of boundaries for a black hole region: the event horizon and the apparent horizon. For dynamical spacetimes, these two horizons are not necessarily the same. To define the event horizon, global information of the whole spacetime up to $t = \infty$ is required.¹ On the other side, an apparent horizon could be defined and detected locally, and it is composed of marginally outer trapped surfaces (MOTSs), which is the outer boundary of the trapped region and where the outgoing future null expansion is zero. The apparent horizon is widely used in the mathematical and numerical explorations of gravitational collapse.

A natural task is to develop black hole thermodynamics along an apparent horizon. This remained to be an open question until 2003. In [9, 10], Ashtekar and Krishnan showed that if a smooth spacelike apparent horizon forms in gravitational collapse, then at each fixed time, one can define the area of the apparent horizon to be an entropy and

¹The event horizon is defined as the future boundary of the causal past set of the future null infinity.

it is indeed *increasing* toward the future. Hence, the second law of black hole mechanics holds.

However, a key mathematical step is missing in the argument by Ashtekar and Krishnan. *Does the smooth spacelike apparent horizon exist? Can it emerge in gravitational collapse from general initial data? These questions are important. Due to physical intuitions, it is conjectured that the smooth spacelike apparent horizon exists in gravitational collapse and it arises from general (both isotropic and anisotropic) initial data.*

The above physical thinking motives the first author to solve the apparent horizon based on the trapped surface formation results obtained in [11] and the other works aforementioned. By solving a new quasilinear elliptic equation with carefully designed gradient estimate, and by employing a limit argument to the scale-critical result in [4] established by An and Luk, the first author in [3] showed that smooth apparent horizon emerges from not-very-anisotropic characteristic initial data. If the initial data are moreover required to be close to isotropic, the corresponding apparent horizon is indeed spacelike and the second law of black hole mechanics (area law) along the apparent horizon is established.

In this paper, we generalize the above conclusion in [3] to the general and fully-anisotropic scenarios. We adopt a different form of the quasilinear elliptic equation for marginally outer trapped surfaces lying on an incoming null cone. By employing *non-perturbative* analytic tools, such as Moser's iteration, the Harnack inequality, Schauder estimates, and the strong maximal principle, via a method of continuity we show the existence, uniqueness, and smoothness of the apparent horizon arising from general (anisotropic) characteristic initial data and we furthermore prove that the apparent horizon obtained is always *spacelike*. Moreover, we construct apparent horizons with *isolated valleys* and demonstrate that the corresponding apparent horizons are also *anisotropic*.

Marginally outer trapped surfaces are also studied within a 3-dimensional spacelike hypersurface. Interested readers are referred to [5, 6, 7, 8, 9, 10, 14, 15, 16, 25, 26, 27].

1.2. The Trapped Surface Formation. Our mathematical explanation starts from [11] by Christodoulou. Whether a trapped surface could form in Einstein vacuum space-time was open for a long time. In a 589-page monumental work [11], with a double null foliation Christodoulou provided an affirmative answer.

In [11] characteristic initial data are prescribed along a truncated incoming cone \underline{H}_0 , where Minkowskian initial data are prescribed, and a truncated outgoing cone H_0 . The 2-sphere $S_{0,0}$ is the intersection of these two cones and it is a standard 2-sphere with radius 1. Along the outgoing null cone H_0 , we denote by χ its outgoing null second fundamental form. Define $\hat{\chi}$ to be the traceless part of χ . Then, $\hat{\chi}$ will serve as the free-prescribed data along H_0 for $0 \leq \underline{u} \leq \delta$, where δ is small. The EVEs are to be solved in the colored region of below Figure 1.

Along H_0 , setting $\hat{\chi}$ to be large, Christodoulou designed a short-pulse hierarchy, where different geometric quantities are required to be of different sizes in term of δ . This carefully chosen hierarchy is later shown to be preserved in the nonlinear evolution.




FIGURE 1

Thus, being a large data problem in nature, a semi-global existence theorem in the colored region can still be established. We hence could control the dynamics and geometry of EVEs from initial data quite far away to a region where a “black hole” is about to form. Requiring the incoming radiation per unit solid angle is bounded uniformly from below, a trapped surface is guaranteed to form in the colored region of Figure 1.

Here is the main theorem² in [11].

Theorem 1.1 (Christodoulou [11]). *Assume \underline{H}_0 coincides with a backwards light cone in the Minkowskian spacetime for $0 \leq u \leq 1$. Then for every $B > 0$ and $u_* \leq 1$, there exists a $\delta = \delta(B, u_*) > 0$ sufficiently small with the following property. If the initial $\hat{\chi}_0$, prescribed on H_0 for $0 \leq \underline{u} \leq \delta$, satisfies*

$$(1.2) \quad \sum_{i \leq 5, j \leq 3} \delta^{\frac{1}{2}+j} \|\nabla_{e_4}^j \nabla^i \hat{\chi}_0\|_{L_{\underline{u}}^\infty L^2(S_{0, \underline{u}})} \leq B,$$

where e_4 and ∇ are outgoing null vector and angular derivative on a 2-sphere $S_{u, \underline{u}}$ respectively, then the solution to EVEs remains regular in $0 \leq u \leq u_*$, $0 \leq \underline{u} \leq \delta$. In addition, if the initial data further verify the lower bound

$$(1.3) \quad \inf_{\omega \in \mathbb{S}^2} \int_0^\delta |\hat{\chi}_0|^2(\omega, \underline{u}') d\underline{u}' \geq M_* > 2(1 - u_*),$$

then for δ sufficiently small (depending on B , u_* and M_*), the sphere $S_{u_*, \delta}$ is a trapped surface.³

After [11], various simplifications and extensions of Theorem 1.1 have been obtained. Interested readers are also referred to [13, 20, 23]. In all these works, similar upper and lower bounds as in [11] are used and trapped surfaces formed are of radius close to 1.

²In Christodoulou’s original proof, outgoing initial data are prescribed at past null infinity. Here, we only mention its finite-region version.

³The initial data constructed in [11] satisfy both (1.2) and (1.3) at the same time. Moreover, the initial data can be chosen to obey $\inf_{\omega \in \mathbb{S}^2} \int_0^\delta |\hat{\chi}|^2(\omega, 0, \underline{u}') d\underline{u}' < 2$. Thus for δ sufficiently small, it can be proved that the initial hypersurface H_0 does not contain any trapped surfaces.

An important work [19] relaxed the lower bound assumption in [11] to allow for fully anisotropic initial data.

Theorem 1.2 (Klainerman-Luk-Rodnianski [19]). *Assume that the initial data for EVEs satisfy the condition (1.2) in Theorem 1.1 and the lower bound*

$$\sup_{\omega \in \mathbb{S}^2} \int_0^\delta |\hat{\chi}_0|^2(\omega, \underline{u}') d\underline{u}' \geq M_* > 0.$$

Then, for sufficiently small δ , a trapped surface forms in the future of the initial data.

To obtain this result, in [19] the authors employed a deformation of the double null foliation and solved a quasilinear elliptic inequality. See also a geometric approach later by Le in [22].

Another work [4] relaxed the upper bound assumption in [11]. The authors considered the colored region of Figure 1 with $u_* = 1 - b\delta a^{\frac{1}{2}}$ and improved [11] to the scale-critical regime.

Theorem 1.3 (An-Luk [4]). *Prescribe Minkowskian data along the initial incoming null hypersurface \underline{H}_0 with $0 \leq u \leq 1$. Given a fixed δ , for every B , there exist $a_0 = a_0(B)$ and $b_0 = b_0(B)$ sufficiently large with the following property. Choose any a and b verifying $a_0 \leq a \leq \delta^{-1}$ and $b_0 \leq b \leq a^{\frac{1}{2}} \leq \delta^{-\frac{1}{2}}$. Along H_0 , with $0 \leq \underline{u} \leq \delta$, if initially $\hat{\chi}_0$ satisfies*

$$(1.4) \quad \sum_{i \leq 5, j \leq 3} \delta^j a^{-\frac{1}{2}} \|\nabla_{e_4}^j \nabla^i \hat{\chi}_0\|_{L_{\underline{u}}^\infty L^2(S_{0, \underline{u}})} \leq B,$$

then the solution to EVEs (1.1) is regular in the spacetime region $0 \leq u \leq 1 - b\delta a^{\frac{1}{2}}$ and $0 \leq \underline{u} \leq \delta$. In addition, if the initial data also verify a lower bound

$$\inf_{\omega \in \mathbb{S}^2} \int_0^\delta |\hat{\chi}_0|^2(\omega, \underline{u}') d\underline{u}' \geq 4\delta a,$$

then the 2-sphere $S_{1-\delta a, \delta}$ is a trapped surface.

To prove the above result, smooth characteristic initial data with a different hierarchy to [11] are designed. Let a be a large universal constant. To form a trapped surface, it is assumed that the $H^{\frac{3}{2}}$ -norm of the metric is of size $a^{\frac{1}{2}}$ and all other H^s -norms are small for $s < \frac{3}{2}$. Note that the homogeneous Sobolev space $\dot{H}^{\frac{3}{2}}(\mathbb{R}^3)$ is the scale-critical space for the 3 + 1 dimensional EVEs. Hence, [4] is also the first scale-critical result for EVEs in the 3 + 1 dimensions.

Later, via a different method by only using signature for decay rates, requiring $b = a^{\frac{1}{2}}$, in [2] with less than 60 pages the first author reproved and extended Theorem 1.3 to past null infinity with sharper bounds. A spacetime rescaling argument is crucially used.

1.3. Apparent Horizons Arising from Non-Anisotropic Initial Data. In [3], the first author considered the following characteristic initial value problem for EVEs in the colored region of Figure 2.




FIGURE 2

- (A₁) The initial incoming null hypersurface \underline{H}_0 coincides with a backward light cone in the Minkowski space with $0 \leq u \leq 1$, and $S_{0,0}$ is the standard 2-sphere with radius 1.
- (A₂) Let δ , B , and a be positive constants. Along H_0 , for $0 \leq \underline{u} \leq \delta$, prescribe $\hat{\chi}_0$ such that

$$\sum_{i \leq 5, j \leq 3} a^{-\frac{1}{2}} \|\underline{u}^j \nabla_{e_4}^j \nabla^i \hat{\chi}_0\|_{L_{\underline{u}}^\infty L^2(S_{0, \underline{u}})} \leq B.$$

Theorem 1.4 (An [3]). *Given $\delta > 0$, for every $B > 0$, there exist $a_0 = a_0(B)$ and $b_0 = b_0(B)$ sufficiently large with the following property. Choose any a and b satisfying $a_0 \leq a \leq \delta^{-1}$ and $b_0 \leq b \leq a^{\frac{1}{2}} \leq \delta^{-\frac{1}{2}}$, and assume the characteristic initial data satisfy (A₁) and (A₂) above. Then, the EVEs (1.1) admit a unique solution in the region $0 < \underline{u} \leq \delta$ and $0 \leq u \leq 1 - b\underline{u}a^{\frac{1}{2}}$.*

1. Moreover, assume

$$(1.5) \quad \int_0^{\underline{u}} |\hat{\chi}_0|^2(\omega, \underline{u}') d\underline{u}' = f(\omega, \underline{u}) \underline{u} a \quad \text{for each } 0 < \underline{u} \leq \delta,$$

where $f(\omega, \underline{u})$ is a smooth function in $\mathbb{S}^2 \times (0, \delta]$ satisfying

$$(1.6) \quad 20/21 \leq f(\omega, \underline{u}) \leq 22/21.$$

Then along each $\underline{H}_{\underline{u}}$ ($0 < \underline{u} \leq \delta$), there exists a unique MOTS $M_{\underline{u}}$.

2. Furthermore, assume, for any $i, j \geq 0$,

$$a^{-\frac{1}{2}} \|\underline{u}^j \nabla_{e_4}^j \nabla^i \hat{\chi}_0\|_{L_{\underline{u}}^\infty L^2(S_{0, \underline{u}})} \leq B.$$

Then the collection of MOTSs $\{M_{\underline{u}}\}$ forms a smooth apparent horizon.

3. In addition, assume

$$(1.7) \quad 1 - \frac{1}{c} \leq f(\omega, \underline{u}) \leq 1 + \frac{1}{c},$$

with sufficiently large c . Then, the apparent horizon is spacelike.

Here we outline the approach of [3].

In the original double null foliation, on $S_{1-\underline{u}a, \underline{u}}$ the outgoing null expansion ($\text{tr}\chi|_{S_{1-\underline{u}a, \underline{u}}}$) is negative pointwisely. Hence, $S_{1-\underline{u}a, \underline{u}}$ is a trapped surface. For fixed \underline{u} , to find a 2-sphere $M_{\underline{u}} = \{(u, \underline{u}, \omega) : u = 1 - R(\omega, \underline{u})\}$ where the outgoing null expansion vanishes for all points (thus $M_{\underline{u}}$ is a marginally outer trapped surface), we need to deform the foliation on $\underline{H}_{\underline{u}}$. In this setting, the null expansion $\text{tr}\chi'$ has the following expression:

$$(1.8) \quad \begin{aligned} \text{tr}\chi' &= \text{tr}\chi - 2\Omega\Delta'_{M_{\underline{u}}}R(\omega, \underline{u}) - 4\Omega\eta_a\nabla^a R(\omega, \underline{u}) \\ &\quad - \Omega^2\text{tr}\underline{\chi}|\nabla R(\omega, \underline{u})|^2 - 8\Omega^2\underline{\omega}|\nabla R(\omega, \underline{u})|^2. \end{aligned}$$

Here, $\Delta'_{M_{\underline{u}}}$ and ∇ are the Laplacian operator and the angular derivative to the unknown 2-sphere $u = 1 - R(\omega, \underline{u})$ at the point $(\omega, 1 - R(\omega, \underline{u}), \underline{u})$; Ω is a geometric quantity satisfying $g(\frac{\partial}{\partial u}, \frac{\partial}{\partial \underline{u}}) = -2\Omega^2$; and η_a , $\underline{\chi}_{bc}$, and $\underline{\omega}$ are certain Ricci coefficients of g .

On fixed $\underline{H}_{\underline{u}}$, when there is no danger of confusion, for simplicity we use $R(\omega)$ to substitute $R(\omega, \underline{u})$. In [19], Klainerman, Luk, and Rodnianski derived (1.8) and first solved the quasilinear elliptic inequality $\text{tr}\chi' < 0$, i.e.,

$$\text{tr}\chi - 2\Omega\Delta'_{M_{\underline{u}}}R(\omega) - 4\Omega\eta_a\nabla^a R(\omega) - \Omega^2\text{tr}\underline{\chi}|\nabla R(\omega)|^2 - 8\Omega^2\underline{\omega}|\nabla R(\omega)|^2 < 0.$$

We proceed to solve the quasilinear elliptic equation $\text{tr}\chi' = 0$, i.e.,

$$(1.9) \quad \Delta'_{M_{\underline{u}}}R(\omega) + 2\eta_b\nabla^b R(\omega) + \frac{1}{2}\Omega\text{tr}\underline{\chi}|\nabla R(\omega)|^2 + 4\Omega\underline{\omega}|\nabla R(\omega)|^2 - \frac{\Omega^{-1}}{2}\text{tr}\chi = 0.$$

A key step in [3] is to derive C^1 (gradient) estimates by Bochner's formula. Define

$$(1.10) \quad h(R) = 1 + \frac{8}{\underline{u}^2 a^2} \left(R - \frac{\underline{u}a}{2} \right)^2.$$

By a careful analysis based on Bochner's formula and the explicit form of (1.9), in [3] we obtain

$$\Delta'_M(h(R)|\nabla R|^2) \geq \frac{1}{\underline{u}^2 a^2} |\nabla R|^4 + \frac{1}{8R^2} |\nabla R|^2 - \frac{1}{R^2} \cdot o(1).$$

Note that not only the sign in front of $|\nabla R|^4$ is *positive*, but the sign in front of $|\nabla R|^2$ is also *positive*! Then, via the maximum principle it implies

$$(1.11) \quad |\nabla R|(\omega) \ll 1 \quad \text{for all } \omega \in \mathbb{S}^2.$$

We point out that the exact form of $h(R)$ in (1.10) plays a *vital* role in the derivation of (1.11). The coefficient 8 is closely related to the assumption (1.6). The gradient estimate (1.11) is essential in subsequent studies. It demonstrates that the solution $R(\omega)$ is close to a constant under the condition (1.6). Physically, the resulting MOTS is close to a sphere.

To solve (1.9), we need to analyze the corresponding linearized equation. Denote by $G(R)$ the expression on the left-hand side of (1.9), for $R = R(\omega)$. By a lengthy computation, the linearized operator $G_R(R)$ is given by

$$(1.12) \quad G_R(R)[W] = \Delta'_M W - \frac{1}{R}(1 - |\nabla R|^2)(1 + o(1))W.$$

By (1.11), the coefficient of W in (1.12) is *negative*, and thus $G_R(R)[W]$ is invertible. Such invertibility implies the openness in the method of continuity to solve (1.9), or to prove the existence of MOTS. Similarly, (1.11) plays a crucial role in the proof of other assertions in Theorem 1.4.

We now make a simple but an important observation. The coefficient of zero order term in (1.12) does not have a desired sign if $|\nabla R|$ is not small as asserted in (1.11). Different methods are needed to solve (1.9) for the general anisotropic initial data!

1.4. Apparent Horizons Arising from General Anisotropic Initial Data. In this paper, we will study the equation (1.9) under a much relaxed assumption on the function f introduced in (1.5). We prove the following result.

Theorem 1.5. *With characteristic initial data satisfying (A_1) , (A_2) and with the choice of parameters a, b as in Theorem 1.4, the EVEs (1.1) admit a unique solution in the region $0 \leq \underline{u} \leq \delta$ and $0 \leq u \leq 1 - b\underline{u}a^{\frac{1}{2}}$.*

1. Assume

$$\int_0^{\underline{u}} |\hat{\chi}_0|^2(\omega, \underline{u}') d\underline{u}' = f(\omega, \underline{u})\underline{u}a \quad \text{for each } 0 < \underline{u} \leq \delta,$$

where $f(\omega, \underline{u})$ is a smooth function on $\mathbb{S}^2 \times (0, \delta]$ satisfying

$$(1.13) \quad 0 \leq f \leq 1 \quad \text{on } \mathbb{S}^2 \times (0, \delta],^4$$

and, for any $\underline{u} \in (0, \delta]$,

$$(1.14) \quad f(\cdot, \underline{u}) \geq m \quad \text{on } B_p(\epsilon),$$

for some point $p \in \mathbb{S}^2$ and some constants $m \in (0, 1)$ and $\epsilon \in (0, \pi/2)$. Then in a large region of each $\underline{H}_{\underline{u}}$ ($0 < \underline{u} \leq \delta$), there exists a unique MOTS $M_{\underline{u}}$. (See the details of considered regions in Theorem 6.1.)

2. In addition, if requiring, for any $i, j \geq 0$,

$$(1.15) \quad a^{-\frac{1}{2}} \|\underline{u}^j \nabla_{e_4}^j \nabla^i \hat{\chi}_0\|_{L_{\underline{u}}^\infty L^2(S_{0, \underline{u}})} \leq B \quad \text{and} \quad |\underline{u} \partial_{\underline{u}} f(\omega, \underline{u})| \leq a^{-\frac{1}{3}},$$

then the collection of MOTS $\{M_{\underline{u}}\}$ forms a smooth and spacelike hypersurface, i.e., the dynamical horizon.

Furthermore, we will provide explicit constructions of apparent horizons with isolated valleys, and by designing and employing a *new anisotropic lower bound* we prove that these apparent horizons are also anisotropic.

⁴The condition $0 \leq f(\underline{u}, \omega) \leq 1$ is very general. With the same method as in this paper, we could replace 1 by any fixed positive constant M . For simplicity of presentation, we always take $M = 1$.

Theorem 1.6. *Let p_1, \dots, p_n be fixed n points on \mathbb{S}^2 , ϵ be a positive constant much less than the distance between any two distinct p_i and p_j for $i, j = 1, \dots, n$, and f be a smooth function on \mathbb{S}^2 , such that $0 \leq f \leq \nu(n)$ and*

$$f \geq \nu(n)/2 \text{ on } \cup_{i=1}^n B_{p_i}(\epsilon/2), \quad f = 0 \text{ on } \mathbb{S}^2 \setminus \cup_{i=1}^n B_{p_i}(\epsilon),$$

for some constructed constant $\nu(n) \in (0, 1)$ depending only on n . Then for each \underline{u} , the equation $\mathcal{S}(\phi, \underline{u}) = 0$ admits a solution $\phi_\epsilon(\omega, \underline{u})$ satisfying, for any $r_0 \gg \epsilon$ and any $i = 1, \dots, n$,

$$(1.16) \quad \phi_\epsilon(p_i, \underline{u}) - \inf_{\omega \in \mathbb{S}^2 \setminus \cup_{i=1}^n B_{p_i}(r_0)} \phi_\epsilon(\omega, \underline{u}) \leq \frac{1}{2n} \log \epsilon + C,$$

where C is a positive constant, independent of ϵ and \underline{u} .

The estimate (1.16) demonstrates that the value of ϕ_ϵ at each p_i is significantly smaller than values away from those p_i 's. In this sense, we say ϕ_ϵ has a valley at p_i . The constant $\nu(n)$ can be computed explicitly. For example, for the case of a single valley, we can take $\nu(1) = 1$. The characteristic initial data leading to apparent horizons with valleys as stated in Theorem 1.6 satisfy the assumptions in Theorem 1.5. The corresponding apparent horizons are henceforth spacelike as well.

Theorem 1.5 infers that *smooth and spacelike apparent horizons arises from general and fully anisotropic initial data*. This certifies the assumption of the black hole thermodynamics theory. Following geometric computations in Section 11 of [3], the spacelike apparent horizons constructed in Theorem 1.5 also satisfy the following property.

Proposition 1.7. *Let $M_{\underline{u}}$ be as in Theorem 1.5. Then,*

$$\lim_{\underline{u} \rightarrow 0} \text{Area}(M_{\underline{u}}) = 0, \quad \text{and} \quad \text{Area}(M_{\underline{u}'}) > \text{Area}(M_{\underline{u}}) \text{ for } \underline{u}' > \underline{u}.$$

Inspired by the works of Ashtekar and Krishnan in [9, 10], we call the above proposition *the area law* along the dynamical horizon. Taking \underline{u} as a time parameter, $\text{Area}(M_{\underline{u}})$ could be viewed as the entropy along the apparent horizon at \underline{u} . It is indeed increasing toward the future!

1.5. An Overview of the Proof of Theorem 1.5. Under the assumptions (1.13) and (1.14), solutions are not expected to be close to constant. In fact, anisotropic solutions do exist according to Theorem 1.6. In the present far-away from close-to-isotropic case, the carefully designed gradient estimate as in [3] does not apply. We will employ more robust and powerful analytic methods to derive gradient estimates or, more generally, the $C^{2,\alpha}$ -estimates. These methods are significantly different from those employed in [3]. Moreover, the gradient estimate in this paper is not as restrictive as the one established in [3]. Due to the more general form of our gradient estimate, we have to employ entirely different methods to study the invertibility of the linearized equations and the smooth dependence of solutions on parameters.

Our proof of Theorem 1.5 is non-perturbative in nature. Moser's iteration, the Harnack inequality, Schauder estimates, and the strong maximum principle are crucially used in our paper. Moreover, we can trace all constants used in our proof explicitly.

The constants m and ϵ for f in (1.14) determine the upper bound constant of a priori estimates, which in turn determines all the rest of constants in later sections.

In the following, we outline our proof of Theorem 1.5. Key intermediate results are given by Theorem 4.1 for a priori $C^{2,1/3}$ -estimates, Lemma 5.3 for the invertibility of the linearized operator, Theorem 6.1 for the existence of solutions, Theorem 6.2 for the smooth dependence of solutions on the parameter \underline{u} , and Theorem 6.5 for the spacelikeness of the hypersurfaces.

1.5.1. *A New Quasilinear Elliptic Equation.* To derive the a priori estimates for the fully anisotropic case, we change the form of (1.9) and employ a more robust and powerful approach. We will use the function $\phi = \phi(\omega, \underline{u})$ defined by

$$R(\omega, \underline{u}) = \underline{u}ae^{-\phi(\omega, \underline{u})}.$$

A straightforward computation changes (1.9) to

$$\begin{aligned} \Delta_g \phi + \frac{e^{2\phi}}{\underline{u}^2 a^2} - \frac{e^{3\phi} f(\omega, \underline{u})}{2\underline{u}^2 a^2} - 2\Omega \hat{\chi}(\nabla_g \phi, \nabla_g \phi) \underline{u}ae^{-\phi} + 2g(\eta, \nabla_g \phi) \\ + \left[-\frac{1}{2}\Omega \text{tr}_g \chi - \frac{1}{\underline{u}ae^{-\phi}} - 4\Omega \underline{\omega} \right] \underline{u}ae^{-\phi} |\nabla_g \phi|^2 \\ + \frac{e^\phi}{\underline{u}a} \left[\frac{1}{2}\Omega^{-1} \text{tr}_g \chi - \frac{1}{\underline{u}ae^{-\phi}} + \frac{\underline{u}af(\omega, \underline{u})}{2(\underline{u}a)^2 e^{-2\phi}} \right] = 0. \end{aligned}$$

Set $g = (\underline{u}ae^{-\phi})^2 \gamma$. We further change the above equation to

$$\begin{aligned} \mathcal{S}(\phi) \equiv \Delta_\gamma \phi + 1 - \frac{1}{2}f(\omega, \underline{u})e^\phi - 2\Omega \hat{\chi}(\nabla_\gamma \phi, \nabla_\gamma \phi) \underline{u}ae^{-\phi} + 2\gamma(\eta, \nabla_\gamma \phi) \\ (1.17) \quad + \left[-\frac{1}{2}\Omega \text{tr}_g \chi - \frac{1}{\underline{u}ae^{-\phi}} - 4\Omega \underline{\omega} \right] \underline{u}ae^{-\phi} |\nabla_\gamma \phi|^2 \\ + \underline{u}ae^{-\phi} \left[\frac{1}{2}\Omega^{-1} \text{tr}_g \chi - \frac{1}{\underline{u}ae^{-\phi}} + \frac{\underline{u}af(\omega, \underline{u})}{2(\underline{u}a)^2 e^{-2\phi}} \right] = 0. \end{aligned}$$

Under the assumptions (1.13) and (1.14), in Section 3 we construct upper and lower barriers for solutions $\phi(\omega, \underline{u})$ to (1.17). We then seek the solution satisfying

$$(1.18) \quad 0 \leq \phi \leq \kappa.$$

Here, κ is a fixed large but specific constant related to the barriers. We should point out that the constants m and ϵ in (1.14) determine the constant κ in (1.18) explicitly, which then gives all the rest constants.

We would like to emphasize that the metric $g = (\underline{u}ae^{-\phi})^2 \gamma$ is degenerate as $\underline{u} \rightarrow 0$, but the conformal metric γ is uniformly bounded. As a consequence, equations in terms of the metric γ are uniformly elliptic. This fact plays a fundamental role in deriving various estimates for solutions to the elliptic equations of ϕ and $\partial_{\underline{u}}\phi$. We also point out that we will study three elliptic equations in this paper. The main equation is (1.17), which is the equation for ϕ . To solve it, we need to study its linearized equation. To prove the spacelikeness of the apparent horizon, we will also need to study the elliptic equation for $\partial_{\underline{u}}\phi$.

1.5.2. *Moser's Iteration and Schauder Estimates.* Note that (1.17) is a quasilinear elliptic equation and $\gamma = \gamma(\underline{u}, \phi, \omega)$. Assume ϕ is a solution to $\mathcal{S}(\phi) = 0$. We start our study with the L^∞ -bound for ϕ as in (1.18) and derive $C^{2,1/3}$ -estimates in three successive steps. First in Theorem 4.2, we prove a C^α -estimate of the form

$$|\phi|_{C^{0,\alpha}(\mathbb{S}^2)} \leq Ce^\kappa,$$

where α is a small positive number. This step from L^∞ to $C^{0,\alpha}$ is achieved through the weak Harnack inequality and Moser's iteration together with an important observation that the equation (1.17) satisfies the *natural growth* condition. Refer to Section 4.5 of [18] for the definition of natural growth condition. Next in Theorem 4.3, we prove a $C^{1,1/3}$ -estimate

$$|\phi|_{C^{1,1/3}(\mathbb{S}^2)} \leq (Ce^\kappa)^{Ce^{4\Lambda\kappa}}.$$

In this step, we use the characterization of Hölder continuous functions by Campanato and the technique of freezing coefficients. Finally, we use the Schauder estimate and obtain

$$(1.19) \quad |\phi|_{C^{2,1/3}(\mathbb{S}^2)} \leq (Ce^\kappa)^{Ce^{4\Lambda\kappa}},$$

where Λ is the ellipticity constant of the metric component (γ_{ij}) and C is a positive uniform constant. Refer to Theorem 4.1 for details.

1.5.3. *The Invertibility of the Linearized Operator via the Positivity of the First Eigenvalue and the Strong Maximum Principle.* We first compute the linearized operator $\partial_\phi \mathcal{S}(\phi)$. Let ϕ be a given function on \mathbb{S}^2 . In Proposition 5.1, we prove that, for any function w on \mathbb{S}^2 ,

$$(1.20) \quad \partial_\phi \mathcal{S}(\phi)[w] = \frac{d}{d\epsilon} \Big|_{\epsilon=0} \mathcal{S}(\phi + \epsilon w) = \Delta_\gamma w - \frac{1}{2} f e^\phi w + c^i \nabla_i w + cw,$$

where the connection is with respect to the metric γ , and $c = c(\omega, \phi, \underline{u})$ and $c^i = c^i(\omega, \phi, \underline{u})$ are functions satisfying

$$(1.21) \quad |c| + |c^i| \leq Ca^{-\frac{1}{3}} e^\phi (|\nabla_\gamma^2 \phi| + |\nabla_\gamma \phi|^2 + 1).$$

The estimate (1.21) plays an important role in the study of the invertibility of the linearized operator as in (1.20), and it is based on scale-invariant estimates of some geometric quantities associated with the Einstein vacuum equations established in [4]. In view of (1.20), we consider

$$L_0 w = \Delta_\gamma w - \frac{1}{2} f e^\phi w,$$

and view f and ϕ as two fixed nonnegative functions on \mathbb{S}^2 , with f satisfying (1.14).

In Lemma 5.2, we prove that the equation $L_0 w = 0$ admits only the trivial solution. Moreover, the first eigenvalue μ_1 of $-L_0$ is *positive* and simple, and has a *positive* eigenfunction ψ_1 . By the positivity of μ_1 and the positivity of ψ_1 , we have

$$\partial_\phi \mathcal{S}(\phi)[\psi_1] = -\mu_1 \psi_1 + c^i \nabla_i \psi_1 + c \psi_1 < 0,$$

by (1.19) and (1.21), and choosing a appropriately large depending on κ . By Theorem 2.11 in [18], the strong maximum principle then holds for the operator $\partial_\phi \mathcal{S}(\phi)$. In particular, for any $w \in C^2(\mathbb{S}^2)$, we have

$$\max_{\mathbb{S}^2} |w| \leq C \max_{\mathbb{S}^2} |\partial_\phi \mathcal{S}(\phi)[w]|,$$

where C is a positive constant independent of w . By the Schauder theory, we obtain the crucial invertibility of the linearized operator $\partial_\phi \mathcal{S}(\phi) : C^{2,\beta}(\mathbb{S}^2) \rightarrow C^{0,\beta}(\mathbb{S}^2)$ as in Lemma 5.3.

1.5.4. The Existence, Uniqueness and Regularity of Solutions. Now, we consider the equation $\mathcal{S}(\phi, \underline{u}) = 0$.

First, by the strong maximum principle, we establish the uniqueness of solutions in Lemma 5.5. For the existence of solutions, we employ the method of continuity. We obtain the closedness by the $C^{2,1/3}$ -estimates in Theorem 4.1 and the openness by the invertibility of the linearized operator in Lemma 5.3. As a consequence, for each fixed constant $\underline{u} \in (0, \delta]$, we find a smooth solution $\phi = \phi(\omega, \underline{u})$ to $\mathcal{S}(\phi, \underline{u}) = 0$ in Theorem 6.1.

Next, we view such a solution ϕ as a function of ω and \underline{u} . Using again the invertibility of the linearized operator and the implicit function theorem, we conclude that ϕ is smooth in both ω and \underline{u} in Theorem 6.2.

1.5.5. The Spacelikeness of Apparent Horizons. We further aim to prove that $\{u = 1 - \underline{u}ae^{-\phi(\omega, \underline{u})}\}$ is a spacelike hypersurface. A key step is to derive an upper bound for $\partial_{\underline{u}}\phi$. In Proposition 6.3, we deduce the equation satisfied by $\partial_{\underline{u}}\phi$. For each $\underline{u} \in (0, \delta]$, let $\phi = \phi(\omega, \underline{u})$ be the solution to $\mathcal{S}(\phi, \underline{u}) = 0$ as in Theorem 6.1. Then, $\partial_{\underline{u}}\phi$ satisfies

$$(1.22) \quad \Delta_\gamma(\partial_{\underline{u}}\phi) - \frac{1}{2}f(\omega, \underline{u})e^\phi \partial_{\underline{u}}\phi + c^i \nabla_i(\partial_{\underline{u}}\phi) + c \partial_{\underline{u}}\phi = \frac{1}{\underline{u}}h + \frac{1}{2}\partial_{\underline{u}}f e^\phi,$$

where the connection is with respect to the metric γ , and $c = c(\omega, \phi, \underline{u})$, $c^i = c^i(\omega, \phi, \underline{u})$ and $h = h(\omega, \phi, \underline{u})$ are functions satisfying

$$(1.23) \quad |c| + |c^i| + |h| \leq Ca^{-\frac{1}{3}}e^\phi(|\nabla_\gamma^2\phi| + |\nabla_\gamma\phi|^2 + 1).$$

We now compare (1.22) and (1.23) with (1.20) and (1.21), respectively. We point out that the linear operator acting on $\partial_{\underline{u}}\phi$ on the left-hand side of (1.22) is similar to the operator in (1.20), with similar estimates of coefficients. However, the computation leading to (1.22) and the derivation of (1.23) are much more complicated.

With the help of the positivity of the first eigenvalue μ_1 and the first eigenfunction ψ_1 again, in Corollary 6.4 we prove

$$(1.24) \quad \max_{\mathbb{S}^2} |\partial_{\underline{u}}\phi| \leq C[\underline{u}^{-1}a^{-\frac{1}{3}} \max_{\mathbb{S}^2} e^\phi(|\phi|_{C^2(\mathbb{S}^2)}^2 + 1) + \max_{\mathbb{S}^2} |\partial_{\underline{u}}f e^\phi|],$$

where C is a positive constant independent of a , \underline{u} and ϕ . By taking a sufficiently large, we have

$$\max_{\mathbb{S}^2} |\underline{u}\partial_{\underline{u}}\phi| \leq 1/2.$$

Let g' be the induced metric to the surface $u = 1 - R(\omega, \underline{u}) = 1 - \underline{u}ae^{-\phi(\omega, \underline{u})}$. We denote $\omega = (\theta_1, \theta_2)$. Let $X = \lambda_1\partial_{\theta_1} + \lambda_2\partial_{\theta_2} + \lambda_3\partial_{\underline{u}}$ be an arbitrary nonzero tangent vector with some constants $\lambda_1, \lambda_2, \lambda_3$. A straightforward computation yields

$$g'(X, X) = \underline{u}^2 a^2 e^{-2\phi} \lambda_i \lambda_j \gamma_{ij} - 4\underline{u}ae^{-\phi} \lambda_i \lambda_3 \partial_i \phi + 4\lambda_3^2 a e^{-\phi} (1 - \underline{u}\partial_{\underline{u}}\phi),$$

where the summation is over $i, j = 1, 2$. We then obtain $g'(X, X) > 0$ if a is sufficiently large. In other words, the metric g induced to $\{(\omega, u, \underline{u}); u = 1 - \underline{u}ae^{-\phi(\omega, \underline{u})}\}$ is *Riemannian* and the corresponding apparent horizon is spacelike, as asserted in Theorem 6.5.

1.5.6. *A Final Remark.* In analyzing both the quasilinear elliptic equation $\mathcal{S}(\phi, \underline{u}) = 0$ and its linearization, the variation of Laplacian plays an important role. We compute variations of Laplacian in three different circumstances for different purposes. In analyzing the nonlinear equation $\mathcal{S}(\phi, \underline{u}) = 0$, with the variation of Laplacian we calculate the difference between the Laplacian with respect to the general metric γ and the Laplacian with respect to the standard spherical metric. The purpose of the comparison is to construct an explicit supersolution. Second, we need to derive a variation of Laplacian in proving the invertibility of linearized equations. Third, we need to derive the variation of Laplacian with respect to \underline{u} in studying the equation satisfied by $\partial_{\underline{u}}\phi$. These variations are explicitly computed in terms of geometric quantities and solutions. With estimates of geometric quantities established in [4] and estimates of solutions established in Section 4, we are able to present precise estimates of these variations of the Laplacian operators as well as variations of other geometric quantities appearing in the equations. As a consequence, we can choose the parameter a explicitly in terms of the initial data, notably f , through m and ϵ defined in (1.14).

The paper is organized as follows. In Section 2, we demonstrate the settings. In Section 3, we construct subsolutions and supersolutions. In Section 4, we establish Schauder estimates. In Section 5, we analyze the linearized operator and prove its invertibility. In Section 6, we employ the method of continuity to prove the existence and properties of solutions. In Section 7, we construct explicit anisotropic apparent horizons with valleys. In Appendix A, we collect useful estimates for some geometric quantities from [4].

2. SETTINGS

In this section, we introduce the main equation to be studied and rearrange it into a preferred form.

2.1. The Double Null Foliation and the Coordinate System. All computations in this paper are based on a double null foliation, with which we decompose the spacetime with outgoing and incoming null cones, i.e., $u = \text{constant}$ and $\underline{u} = \text{constant}$, respectively. Here u and \underline{u} satisfy the eikonal equations

$$g^{\mu\nu} \partial_\mu u \partial_\nu u = 0, \quad g^{\mu\nu} \partial_\mu \underline{u} \partial_\nu \underline{u} = 0.$$

Given u' and \underline{u}' two fixed constants, the intersection of $u = u'$ and $\underline{u} = \underline{u}'$ is a 2-sphere, denoted by $S_{u', \underline{u}'}$.⁵

On $S_{u', \underline{u}'}$, we will use coordinates (θ^1, θ^2) . To decide (θ^1, θ^2) for each point on $S_{u', \underline{u}'}$, we follow Chapter 1 of [11]. First, $S_{0,0}$ is the standard 2-sphere in the Minkowskian spacetime with radius 1, and we define (θ^1, θ^2) to be coordinates on it via the stereographic projection. Then, we extend this (θ^1, θ^2) coordinate system to \underline{H}_0 ($\underline{u} = 0$) by requiring

$$\frac{\partial}{\partial u} \theta^A = 0 \text{ on } \underline{H}_0.⁶$$

In another word, fix a point p on $S_{0,0}$ and assume l_p is the incoming null geodesics on \underline{H}_0 emitting from p , and we require all the points along l_p have the same angular coordinate (θ^1, θ^2) . We further extend coordinate system (θ^1, θ^2) from \underline{H}_0 to the whole spacetime region by requiring

$$\not\!{L} \frac{\partial}{\partial \underline{u}} \theta^A = 0;⁷$$

namely, every point along the same outgoing null geodesics on H_u shares the same angular coordinate. We hence obtain a coordinate system in a neighborhood of $S_{0,0}$. With it, we take the ansatz of the Lorentzian metric g to be of the form

$$(2.1) \quad g = -2\Omega^2 (du \otimes d\underline{u} + d\underline{u} \otimes du) + \gamma_{AB} (d\theta^A - d^A du) \otimes (d\theta^B - d^B du).$$

We define e_3, e_4 by

$$e_3 = \Omega^{-1} \left(\frac{\partial}{\partial u} + d^A \frac{\partial}{\partial \theta^A} \right), \quad e_4 = \Omega^{-1} \frac{\partial}{\partial \underline{u}}.$$

It is straightforward to check that $\{e_3, e_4\}$ forms a null pair. Here e_3, e_4 are along incoming and outgoing null directions and

$$g(e_3, e_3) = 0, \quad g(e_4, e_4) = 0, \quad g(e_3, e_4) = -2.$$

Denote by $\{e_1, e_2\}$ a frame tangent to the 2-sphere $S_{u, \underline{u}}$. We then decompose curvature components and Ricci coefficients using frames $\{e_1, e_2, e_3, e_4\}$.

2.2. The Geometric Quantities. Set the indices A, B to be 1, 2 and let $D_A = D_{e_A}$. We define Ricci coefficients:

$$(2.2) \quad \begin{aligned} \chi_{AB} &= g(D_A e_4, e_B), & \underline{\chi}_{AB} &= g(D_A e_3, e_B), \\ \eta_A &= -\frac{1}{2} g(D_3 e_A, e_4), & \underline{\eta}_A &= -\frac{1}{2} g(D_4 e_A, e_3), \\ \tilde{\omega} &= -\frac{1}{4} g(D_4 e_3, e_4), & \underline{\omega} &= -\frac{1}{4} g(D_3 e_4, e_3), \\ \zeta_A &= \frac{1}{2} g(D_A e_4, e_3). \end{aligned}$$

⁵For more detailed description of the double null foliation, interested readers are referred to Chapter 1 and Chapter 2 of [11] and Section 2.1 of [2].

⁶Here we assume the geometry along \underline{H}_0 to be Minkowskian.

⁷Here $\not\!{L}$ is the restriction of the Lie derivative to $TS_{u', \underline{u}'}$.

Denote by $\not\phi_{AB}$ the induced metric on $S_{u,\underline{u}}$. We further decompose $\chi, \underline{\chi}$ as

$$\chi_{AB} = \frac{1}{2}\text{tr}\chi \cdot \not\phi_{AB} + \hat{\chi}_{AB}, \quad \underline{\chi}_{AB} = \frac{1}{2}\text{tr}\underline{\chi} \cdot \not\phi_{AB} + \hat{\underline{\chi}}_{AB}.$$

Note that $\hat{\chi}_{AB}$ and $\hat{\underline{\chi}}_{AB}$ are the corresponding traceless parts of χ_{AB} and $\underline{\chi}_{AB}$. In Appendix A, we will collect useful estimates for these quantities from [4]. Such estimates play important roles in this paper.

2.3. The Main Elliptic Equation. We now introduce the main equation to be solved and rearrange it in a preferred form. In Sections 3-5, we will view $g, \underline{\chi}, \chi, \eta, \Omega, \underline{\omega}$ defined above as functions of $\omega \in \mathbb{S}^2$ and u , parametrized by \underline{u} ; while in Section 6, we will view them as functions of ω, u , and \underline{u} .

We write $u = 1 - R$. For each fixed \underline{u} and any function $R = R(\omega, \underline{u})$ on \mathbb{S}^2 , we set

$$(2.3) \quad \begin{aligned} \mathcal{G}(R, \underline{u}) = & \Delta_g R + \frac{1}{2}\Omega \text{tr}_g \underline{\chi} |\nabla_g R|^2 - \frac{1}{2}\Omega^{-1} \text{tr}_g \chi \\ & + 4\Omega \underline{\omega} |\nabla_g R|^2 + 2\Omega \hat{\underline{\chi}}(\nabla_g R, \nabla_g R) + 2g(\eta, \nabla_g R). \end{aligned}$$

Here g is viewed as an induced metric on the 2-sphere $M = \{u = 1 - R\}$ along the incoming null cone $\underline{H}_{\underline{u}}$. Using the expression of $\text{tr}\chi'$ in (1.8), we have

$$u = 1 - R(\omega) \text{ is a } \begin{cases} \text{MOTS} & \text{if } \mathcal{G}(R, \underline{u}) = 0 \text{ on } \mathbb{S}^2, \\ \text{trapped surface} & \text{if } \mathcal{G}(R, \underline{u}) > 0 \text{ on } \mathbb{S}^2, \\ \text{untrapped surface} & \text{if } \mathcal{G}(R, \underline{u}) < 0 \text{ on } \mathbb{S}^2. \end{cases}$$

Our primary goal is to find a solution $R = R(\omega, \underline{u})$ of $\mathcal{G}(R, \underline{u}) = 0$ for each \underline{u} and then study the regularity of $R = R(\omega, \underline{u})$ in terms of \underline{u} . When there is no danger of confusion, we suppress \underline{u} and write $R = R(\omega)$ and $\mathcal{G}(R)$ for simplicity.

We proceed to rearrange $\mathcal{G}(R, \underline{u})$ in a preferred form. The rearrangement is based on estimates established in [4]. Set

$$(2.4) \quad \begin{aligned} \alpha_1 = & -\left(\frac{1}{2}\Omega \text{tr}_g \underline{\chi} + \frac{1}{R} + 4\Omega \underline{\omega}\right), \\ \alpha_2 = & \frac{1}{2}\Omega^{-1} \text{tr}_g \chi - \frac{1}{R} + \frac{\underline{u}a}{2R^2} f(\omega, \underline{u}), \end{aligned}$$

where $f = f(\omega, \underline{u})$ is the function introduced in (1.5). We now rewrite $\mathcal{G}(R, \underline{u})$ as

$$(2.5) \quad \begin{aligned} \mathcal{G}(R, \underline{u}) = & \Delta_g R - \frac{1}{R} |\nabla_g R|^2 - \frac{1}{R} + \frac{\underline{u}a}{2R^2} f \\ & + 2\Omega g^{ik} g^{jl} \hat{\underline{\chi}}_{kl} \nabla_i R \nabla_j R + 2g^{ij} \eta_i \nabla_j R - \alpha_1 |\nabla_g R|^2 - \alpha_2. \end{aligned}$$

The metric g depends on ω, u , and \underline{u} . Note that

$$\partial_u g_{kl} = 2\Omega \underline{\chi}_{kl}, \quad \partial_{\underline{u}} g_{kl} = 2\Omega \chi_{kl}.$$

By writing $\underline{\chi}$ and χ as combinations of trace parts and traceless parts, we have

$$(2.6) \quad \partial_u g_{kl} = \Omega \text{tr}_g \underline{\chi} g_{kl} + 2\Omega \hat{\underline{\chi}}_{kl},$$

and

$$(2.7) \quad \partial_{\underline{u}} g_{kl} = \Omega \text{tr}_g \chi g_{kl} + 2\Omega \hat{\chi}_{kl}.$$

Along fixed $\underline{H}_{\underline{u}}$, when no confusing would be caused, we suppress \underline{u} dependence and rewrite $g(\omega, u, \underline{u})$ as $g(\omega, u)$ with $u = 1 - R$. In local coordinates, the estimates of g in [4] and [2] imply

$$(2.8) \quad \frac{1}{2}R^2 \leq \sqrt{|g|}(\omega, u) \leq \frac{3}{2}R^2, \quad \Lambda^{-1}R^2I \leq g_{ij}(\omega, u) \leq \Lambda R^2I,$$

and

$$(2.9) \quad |D_{\omega} g_{ij}(\omega, u)| + R|\partial_u g_{ij}(\omega, u)| \leq \Lambda_1 R^2,$$

$$(2.10) \quad |D_{\omega}^2 g_{ij}(\omega, u)| + R|D_{\omega} \partial_u g_{ij}(\omega, u)| + R^2|\partial_u^2 g_{ij}(\omega, u)| \leq \Lambda_2 R^2,$$

for some positive constants Λ , Λ_1 , and Λ_2 . Estimates on Ω , χ , $\hat{\chi}$, η , α_1 , and α_2 were established in [4] and will be recalled in Appendix A.

Next, for each fixed \underline{u} we introduce a function $\phi = \phi(\omega, \underline{u})$ by

$$R(\omega, \underline{u}) = \underline{u} a e^{-\phi(\omega, \underline{u})}.$$

If there is no danger of confusion, we also write $\phi(\omega)$, $f(\omega)$ instead of $\phi(\omega, \underline{u})$, $f(\omega, \underline{u})$. We then define the operator $S(\phi, \underline{u})$ by

$$(2.11) \quad \mathcal{S}(\phi, \underline{u}) = -R\mathcal{G}(\phi, \underline{u}).$$

With a straightforward computation, we obtain

$$(2.12) \quad \mathcal{S}(\phi, \underline{u}) = R^2 \Delta_g \phi + 1 - \frac{1}{2} f e^{\phi} + F(\phi, \underline{u}),$$

where

$$(2.13) \quad F(\phi, \underline{u}) = -2R^3 \Omega g^{ik} g^{jl} \hat{\chi}_{kl} \nabla_i \phi \nabla_j \phi + 2R^2 g^{ij} \eta_i \nabla_j \phi + R^3 \alpha_1 |\nabla_g \phi|^2 + R\alpha_2.$$

Last, we introduce a metric $\gamma = \gamma(\omega, u)$ on \mathbb{S}^2 such that

$$(2.14) \quad g = R^2 \gamma.$$

By (2.8), (2.9) and (2.10), in local coordinates γ satisfies

$$(2.15) \quad \frac{1}{2} \leq \sqrt{|\gamma|}(\omega, u) \leq \frac{3}{2}, \quad \Lambda^{-1}I \leq \gamma_{ij}(\omega, u) \leq \Lambda I,$$

and

$$(2.16) \quad |D_{\omega} \gamma_{ij}(\omega, u)| + R|\partial_u \gamma_{ij}(\omega, u)| \leq \Lambda_1,$$

$$(2.17) \quad |D_{\omega}^2 \gamma_{ij}(\omega, u)| + R|D_{\omega} \partial_u \gamma_{ij}(\omega, u)| + R^2|\partial_u^2 \gamma_{ij}(\omega, u)| \leq \Lambda_2,$$

via renaming the positive constants Λ_1 and Λ_2 if necessary. Using γ , it is straightforward to check that

$$(2.18) \quad \mathcal{S}(\phi, \underline{u}) = \Delta_{\gamma} \phi + 1 - \frac{1}{2} f e^{\phi} + F(\phi, \underline{u}),$$

where

$$(2.19) \quad F(\phi, \underline{u}) = 2R^{-1} \Omega \hat{\chi}_{kl} \gamma^{ik} \gamma^{jl} \nabla_i \phi \nabla_j \phi + 2\gamma^{ij} \eta_i \nabla_j \phi + R\alpha_1 |\nabla_{\gamma} \phi|^2 + R\alpha_2,$$

and α_1, α_2 are defined in (2.4).

Note that $\mathcal{S}(\phi, \underline{u}) = 0$ is the main elliptic equation in the preferred form and can be expressed as

$$\begin{aligned} \Delta_\gamma \phi + 1 - \frac{1}{2} f(\omega, \underline{u}) e^\phi + 2\Omega \hat{\chi}(\nabla_\gamma \phi, \nabla_\gamma \phi) \underline{u} a e^{-\phi} + 2\gamma(\eta, \nabla_\gamma \phi) \\ + \left[-\frac{1}{2} \Omega \text{tr}_g \chi - \frac{1}{\underline{u} a e^{-\phi}} - 4\Omega \underline{\omega} \right] \underline{u} a e^{-\phi} |\nabla_\gamma \phi|^2 \\ + \underline{u} a e^{-\phi} \left[\frac{1}{2} \Omega^{-1} \text{tr}_g \chi - \frac{1}{\underline{u} a e^{-\phi}} + \frac{\underline{u} a f(\omega, \underline{u})}{2(\underline{u} a)^2 e^{-2\phi}} \right] = 0, \end{aligned}$$

which is (1.17) in the introduction.

By the relation (2.11), we further conclude that

$$u = 1 - \underline{u} a e^{-\phi(\omega)} \text{ is a } \begin{cases} \text{MOTS} & \text{if } \mathcal{S}(\phi, \underline{u}) = 0 \text{ on } \mathbb{S}^2, \\ \text{trapped surface} & \text{if } \mathcal{S}(\phi, \underline{u}) < 0 \text{ on } \mathbb{S}^2, \\ \text{untrapped surface} & \text{if } \mathcal{S}(\phi, \underline{u}) > 0 \text{ on } \mathbb{S}^2. \end{cases}$$

Note that we have

$$(2.20) \quad u = 1 - R(\omega, \underline{u}), \quad R(\omega, \underline{u}) = \underline{u} a e^{-\phi(\omega, \underline{u})}.$$

In the rest of the paper, we will study exclusively the equation $\mathcal{S}(\phi, \underline{u}) = 0$ and aim to find its solution $\phi(\omega, \underline{u})$.

3. SUBSOLUTIONS AND SUPERSOLUTIONS

In this section, we construct an untrapped surface and a trapped surface on $\underline{H}_{\underline{u}}$ for each small $\underline{u} > 0$. In later sections, we will prove that there exists a unique MOTS between these two surfaces.

Throughout this section, we fix a $\underline{u} \in (0, \delta]$. For any function $\phi = \phi(\omega)$ on \mathbb{S}^2 , let $\mathcal{S}(\phi)$ be the operator defined by (2.18), i.e.,

$$(3.1) \quad \mathcal{S}(\phi) = \Delta_\gamma \phi + 1 - \frac{1}{2} f e^\phi + F(\phi),$$

where F is given in (2.19). The aim of this section is to construct subsolutions and supersolutions for the operator \mathcal{S} . We always assume that f is a nonnegative smooth function on \mathbb{S}^2 , not identically zero. We point out that f is allowed to be zero in a large portion of \mathbb{S}^2 .

Remark 3.1. Throughout this paper, we will use parameters a and κ , which satisfy

$$(3.2) \quad (C e^\kappa)^{C e^{4\Lambda \kappa}} \leq a^{\frac{1}{4}} \ll a^{\frac{1}{3}}.$$

Here, C is a uniform positive constant defined in Theorem 4.1, and Λ is introduced in (2.15).

Remark 3.2. For any $m \in (0, 1)$, $\epsilon \in (0, \pi/2)$, $\tau > 2.2$, and $C(\tau)$ to be specified in Lemma 3.6, as to be discussed in Theorem 6.1 and Remark 3.4, we solve for MOTS lying in $0 \leq \underline{u} \leq \delta$ and $0 \leq u \leq 1 - m \epsilon^{\tau+2} C(\tau) \underline{u} a$. To guarantee that the MOTS is within

the existence region of Theorem 1.5, i.e., $0 \leq \underline{u} \leq \delta$ and $0 \leq u \leq 1 - b\underline{u}a^{\frac{1}{2}}$, we require $m\epsilon^{\tau+2}C(\tau) \geq ba^{-\frac{1}{2}}$. This requirement can easily be achieved by setting $1 \ll b \ll a^{\frac{1}{2}}$ and a being sufficiently large. Here allowing $b \neq a^{\frac{1}{2}}$ as in [4] is important.

We start from constructing subsolutions.

Lemma 3.3. *Let f be a nonnegative continuous function on \mathbb{S}^2 , with $f \leq 1$. Then, $\underline{\phi}(\omega) = \ln(3/2)$ satisfies $\mathcal{S}(\underline{\phi}) > 0$, for a sufficiently large.*

Proof. By (2.19), (2.4) and (A.1), we have

$$(3.3) \quad |F(\phi)| \leq Ce^{\phi}a^{-\frac{1}{3}}(|\nabla_{\gamma}\phi|^2 + 1).$$

Hence,

$$\mathcal{S}(\ln(3/2)) = 1 - \frac{3}{4}f + F(\ln(3/2)) \geq \frac{1}{4} + F(\ln(3/2)) > 0,$$

if a is sufficiently large. \square

Remark 3.4. There are other subsolutions. For example, the function $\underline{\phi} = \ln(\underline{u}a) < 0$ satisfies $\mathcal{S}(\underline{\phi}) > 0$. With $\underline{\phi} = \ln(\underline{u}a)$, we can prove a similar result to Theorem 6.1. For simplicity of presentations, we choose $\underline{\phi}(\omega) = \ln(3/2) > 0$ in the following sections.

Before we construct supersolutions, we estimate $\Delta_{\gamma}\phi - \Delta_{\gamma_0}\phi$. Here, γ_0 is the standard spherical metric on \mathbb{S}^2 . We first derive a well-known formula for the variation of the Laplacian operator.

Lemma 3.5. *Let $\{\gamma(s)\}$ be a family of metrics and $\{\phi(s)\}$ be a family of functions, both parametrized by s . Then,*

$$(3.4) \quad (\Delta_{\gamma}\phi)^{\cdot} = \Delta_{\gamma}\dot{\phi} - \gamma^{ij}\gamma^{kl}\dot{\gamma}_{jl}\nabla_{ik}\phi - \gamma^{ij}\gamma^{kl}\nabla_i\dot{\gamma}_{jl}\nabla_k\phi + \frac{1}{2}\gamma^{ij}\gamma^{kl}\nabla_l\dot{\gamma}_{ij}\nabla_k\phi,$$

where we denote $\dot{\cdot} = \frac{d}{ds}$.

Proof. First, we have

$$\dot{\gamma}^{ij} = -\gamma^{ik}\gamma^{jl}\dot{\gamma}_{kl}.$$

Next, with

$$\Gamma_{ij}^k = \frac{1}{2}\gamma^{kl}[\partial_i\gamma_{jl} + \partial_j\gamma_{il} - \partial_l\gamma_{ij}],$$

we get

$$\dot{\Gamma}_{ij}^k = \frac{1}{2}\gamma^{kl}[\partial_i\dot{\gamma}_{jl} + \partial_j\dot{\gamma}_{il} - \partial_l\dot{\gamma}_{ij}] + \gamma_{ml}\Gamma_{ij}^m\dot{\gamma}^{kl}.$$

With

$$\Delta_{\gamma}\phi = \gamma^{ij}\partial_{ij}\phi - \gamma^{ij}\Gamma_{ij}^k\partial_k\phi,$$

a straightforward calculation yields the desired result. \square

Lemma 3.6. *Let $\phi \in C^2(\mathbb{S}^2)$ be an arbitrary positive function. Then,*

$$(3.5) \quad |\Delta_{\gamma}\phi - \Delta_{\gamma_0}\phi| \leq C(1 + \phi)(|\nabla_{\gamma}^2\phi| + |\nabla_{\gamma}\phi| + |\nabla_{\gamma}\phi|^2)e^{\phi}a^{-\frac{1}{3}},$$

where C is a positive constant, independently of a and ϕ .

Proof. For any constants $\underline{u}, \underline{u}'$ and a function $\phi \in C^2(\mathbb{S}^2)$, set

$$S_{1-\underline{u}ae^{-\phi}, \underline{u}'} = \{(\omega, u, \underline{u}'); \omega \in \mathbb{S}^2, u = 1 - \underline{u}ae^{-\phi}\}.$$

In particular, when $\phi = 0$ we have

$$S_{1-\underline{u}a, \underline{u}'} = \{(\omega, u, \underline{u}'); \omega \in \mathbb{S}^2, u = 1 - \underline{u}a\}.$$

We now fix a positive function $\phi \in C^2(\mathbb{S}^2)$. Note that the Minkowski data are prescribed along $\underline{u} = 0$. Hence, $S_{1-\underline{u}a, 0}$ is the standard 2-sphere with radius $\underline{u}a$. Note that $S_{0,0}$ on \underline{H}_0 is the standard 2-sphere with radius 1 and that γ_0 is the standard spherical metric on it. Let $g|_{S_{1-\underline{u}a, 0}}$ be the metric g restricted to $S_{1-\underline{u}a, 0}$. Then, $g|_{S_{1-\underline{u}a, 0}} = \underline{u}^2 a^2 \gamma|_{S_{1-\underline{u}a, 0}}$, with $\gamma|_{S_{1-\underline{u}a, 0}} = \gamma_0$. Hence,

$$\Delta_\gamma \phi - \Delta_{\gamma_0} \phi = \Delta_\gamma|_{S_{1-\underline{u}ae^{-\phi}, \underline{u}}} \phi - \Delta_\gamma|_{S_{1-\underline{u}a, 0}} \phi.$$

For $s \in [0, 1]$, we set

$$(3.6) \quad R(s) = \underline{u}ae^{-s\phi}, \quad u(s) = 1 - R(s), \quad \underline{u}(s) = \underline{u}s.$$

In the following, we denote by dot the derivative with respect to s , and we also suppress the dependence on s . Hence, by (3.6) it follows

$$\dot{R} = -R\phi, \quad \dot{u} = -\dot{R}, \quad \dot{\underline{u}} = \underline{u}.$$

For simplicity, we also denote by $\gamma(s)$ the metric γ on $S_{1-\underline{u}ae^{-s\phi}, \underline{u}s}$. Then,

$$(3.7) \quad \Delta_\gamma \phi - \Delta_{\gamma_0} \phi = \Delta_{\gamma(1)} \phi - \Delta_{\gamma(0)} \phi = \int_0^1 (\Delta_\gamma \phi) \cdot ds.$$

By $g = R^2 \gamma$ and $\dot{u} = -\dot{R}$, via calculating $(g|_{S_{1-R(s), \underline{u}(s)}}) \cdot$, we get

$$(3.8) \quad R^2 \dot{\gamma}_{kl} = -2R\dot{R}\gamma_{kl} - \dot{R}\partial_u g_{kl} + \underline{u}\partial_{\underline{u}} g_{kl}.$$

Recall (2.6) and (2.7), i.e.,

$$\partial_u g_{kl} = \Omega \text{tr}_g \underline{\chi} g_{kl} + 2\Omega \hat{\underline{\chi}}_{kl}, \quad \partial_{\underline{u}} g_{kl} = \Omega \text{tr}_g \chi g_{kl} + 2\Omega \hat{\chi}_{kl}.$$

Hence, using (3.8) and $\dot{R} = -R\phi$, we get

$$\dot{\gamma}_{kl} = [\phi(R\Omega \text{tr}_g \underline{\chi} + 2) + \underline{u}\Omega \text{tr}_g \chi] \gamma_{kl} + [2\phi R^{-1}\Omega \hat{\underline{\chi}}_{kl} + 2\underline{u}R^{-2}\Omega \hat{\chi}_{kl}].$$

A substitution in (3.4) yields

$$\begin{aligned} (\Delta_\gamma \phi) \cdot &= -(R\Omega \text{tr}_g \underline{\chi} + 2)\phi \Delta_\gamma \phi - \underline{u}\Omega \text{tr}_g \chi \Delta_\gamma \phi - \gamma^{ij} \gamma^{kl} [2\phi R^{-1}\Omega \hat{\underline{\chi}}_{jl} + 2\underline{u}R^{-2}\Omega \hat{\chi}_{jl}] \nabla_{ik} \phi \\ &\quad - \gamma^{ik} [\nabla_i \phi (R\Omega \text{tr}_g \underline{\chi} + 2) + \underline{u}\nabla_i (\Omega \text{tr}_g \chi)] \nabla_k \phi - \phi \gamma^{ik} \nabla_i (R\Omega \text{tr}_g \underline{\chi}) \nabla_k \phi \\ &\quad - \gamma^{ij} \gamma^{kl} \nabla_i [2\phi R^{-1}\Omega \hat{\underline{\chi}}_{jl} + 2\underline{u}R^{-2}\Omega \hat{\chi}_{jl}] \nabla_k \phi \\ &\quad + \gamma^{kl} \nabla_l [\phi (R\Omega \text{tr}_g \underline{\chi} + 2) + \underline{u}\Omega \text{tr}_g \chi] \nabla_k \phi \\ &\quad + \frac{1}{2} \gamma^{ij} \gamma^{kl} \nabla_l [2\phi R^{-1}\Omega \hat{\underline{\chi}}_{ij} + 2\underline{u}R^{-2}\Omega \hat{\chi}_{ij}] \nabla_k \phi. \end{aligned}$$

We point out that the above identity is evaluated at $(\omega, 1 - \underline{u}ae^{-s\phi(\omega)}, \underline{u}s)$ and can be viewed as a linear combination of $\phi\nabla_{ij}\phi, \nabla_i\phi\nabla_j\phi, \phi\nabla_i\phi, \nabla_{ij}\phi, \nabla_i\phi$ with coefficients given by γ^{ij} and

$$R\Omega\text{tr}_g\underline{\chi} + 2, R^{-1}\Omega\hat{\underline{\chi}}_{kl}, \nabla_i(R\Omega\text{tr}_g\underline{\chi}), \nabla_i(R^{-1}\Omega\hat{\underline{\chi}}_{kl}), \\ \underline{u}\Omega\text{tr}_g\chi, \underline{u}R^{-2}\Omega\hat{\chi}_{kl}, \underline{u}\nabla_i(\Omega\text{tr}_g\chi), \underline{u}\nabla_i(R^{-2}\Omega\hat{\chi}_{kl}).$$

By (A.1), these quantities are bounded by $Ce^{s\phi}a^{-\frac{1}{3}}$. Therefore,

$$|(\Delta_\gamma\phi)'| \leq C(1 + \phi)(|\nabla^2\phi| + |\nabla\phi| + |\nabla\phi|^2)e^\phi a^{-\frac{1}{3}}.$$

With (3.7), this implies the desired result. \square

Next, we assume that f satisfies (1.14), i.e.,

$$f \geq m \quad \text{on } B_p(\epsilon),$$

for some point $p \in \mathbb{S}^2$ and some constants $m \in (0, 1)$ and $\epsilon \in (0, \pi/2)$.

Lemma 3.7. *Let f be a nonnegative continuous function on \mathbb{S}^2 , satisfying (1.14). Then, for any $\tau > 2.2$, there exists $\bar{\phi} \in C^\infty(\mathbb{S}^2)$ such that, for any sufficiently large a , $\mathcal{S}(\bar{\phi}) < 0$ and*

$$(3.9) \quad \begin{aligned} \bar{\phi}_{\min} &= \bar{\phi}(p) = -\log(m\epsilon^2) + C(\tau), \\ \bar{\phi}_{\max} &= \bar{\phi}(-p) = -\log(m\epsilon^{\tau+2}) + C(\tau), \end{aligned}$$

for some positive constant C depending only on τ .

Proof. The proof of this lemma is essentially a similar one in [19]. We provide some details, which will be used in a later section. We write $\mathcal{S}(\phi)$ as

$$(3.10) \quad \mathcal{S}(\phi) = \Delta_{\gamma_0}\phi + 1 - \frac{1}{2}fe^\phi + \Delta_\gamma\phi - \Delta_{\gamma_0}\phi + F(\phi),$$

where Δ_{γ_0} is the Laplacian with respect to the standard metric on \mathbb{S}^2 . We first construct a function $\phi_\epsilon = \phi_\epsilon(\omega)$ such that

$$(3.11) \quad \Delta_{\gamma_0}\phi_\epsilon + 1.1 < \frac{1}{2}f(\omega)e^{\phi_\epsilon}.$$

Here, we use 1.1 instead of 1 on the left-hand side to absorb the lower order terms in \mathcal{S} . Denote by λ the distance function from p . Set

$$w = \log \left[\sin \left(\frac{\lambda}{2} \right) \right].$$

Then,

$$\Delta_{\gamma_0}w + \frac{1}{2} = 2\pi\delta_p,$$

and

$$w = \log \lambda + O(1), \quad |\nabla_{\gamma_0}w| = O(\lambda^{-1}), \quad |\nabla_{\gamma_0}^2w| = O(\lambda^{-2}).$$

Let χ_ϵ be a smooth cut-off function, satisfying

$$\chi_\epsilon = \begin{cases} 0 & \text{on } B_p(\epsilon/2), \\ 1 & \text{on } \mathbb{S}^2 \setminus B_p(\epsilon). \end{cases}$$

Define $\tilde{w}_\epsilon = \chi_\epsilon w + (1 - \chi_\epsilon) \log \epsilon$. Then,

$$\tilde{w}_\epsilon = \begin{cases} \log \epsilon & \text{on } B_p(\epsilon/2), \\ \log \epsilon + O(1) & \text{on } B_p(\epsilon) \setminus B_p(\epsilon/2), \\ \log \lambda + O(1) & \text{on } \mathbb{S}^2 \setminus B_p(\epsilon), \end{cases}$$

and

$$|\nabla_{\gamma_0} \tilde{w}_\epsilon| = O(\epsilon^{-1}), \quad |\nabla_{\gamma_0}^2 \tilde{w}_\epsilon| = O(\epsilon^{-2}) \quad \text{on } \mathbb{S}^2 \setminus B_p(\epsilon/2).$$

Moreover,

$$\Delta_{\gamma_0} \tilde{w}_\epsilon + \frac{1}{2} = 0 \quad \text{on } \mathbb{S}^2 \setminus B_p(\epsilon).$$

Consider $\tilde{\phi}_\epsilon = \tau \tilde{w}_\epsilon$. Then we have

$$\tilde{\phi}_\epsilon = \begin{cases} \tau \log \epsilon & \text{on } B_p(\epsilon/2), \\ \tau \log \epsilon + O(1) & \text{on } B_p(\epsilon) \setminus B_p(\epsilon/2), \\ \tau \log \lambda + O(1) & \text{on } \mathbb{S}^2 \setminus B_p(\epsilon), \end{cases}$$

and

$$|\nabla_0 \tilde{\phi}_\epsilon| = O(\epsilon^{-1}), \quad |\nabla_0^2 \tilde{\phi}_\epsilon| = O(\epsilon^{-2}) \quad \text{on } \mathbb{S}^2 \setminus B_p(\epsilon/2).$$

Moreover, it holds

$$\Delta_{\gamma_0} \tilde{\phi}_\epsilon + 1.1 = -\frac{\tau}{2} + 1.1 < 0 \quad \text{on } \mathbb{S}^2 \setminus B_p(\epsilon).$$

For a constant $s_\epsilon > 0$ to be determined, set

$$\phi_\epsilon = \tilde{\phi}_\epsilon - \log s_\epsilon.$$

It is obvious that $\Delta_{\gamma_0} \phi_\epsilon = \Delta_{\gamma_0} \tilde{\phi}_\epsilon$. Hence,

$$\Delta_{\gamma_0} \phi_\epsilon + 1.1 = \Delta_{\gamma_0} \tilde{\phi}_\epsilon + 1.1 < 0 \quad \text{on } \mathbb{S}^2 \setminus B_p(\epsilon).$$

In addition, there exists a constant $C_0 > 0$ such that

$$\Delta_{\gamma_0} \phi_\epsilon = \Delta_{\gamma_0} \tilde{\phi}_\epsilon < \frac{C_0}{\epsilon^2} \quad \text{on } B_p(\epsilon).$$

By $f \geq m$ in $B_p(\epsilon)$ in (1.14), to have $C_0/\epsilon^2 \leq \frac{1}{2}f(\omega)e^{\phi_\epsilon}$ in $B_p(\epsilon)$, we need

$$\inf_{B_p(\epsilon)} \phi_\epsilon \geq \log \left(\frac{2C_0}{m\epsilon^2} \right).$$

By estimates on $\tilde{\phi}_\epsilon$, we have

$$\inf_{B_p(\epsilon)} \phi_\epsilon \geq -\log s_\epsilon + \tau \log \epsilon - \log C_1,$$

for some constant $C_1 > 0$. Therefore, it suffices to choose

$$s_\epsilon = \frac{m\epsilon^{\tau+2}}{2C_0C_1}.$$

Then, we obtain

$$\Delta_{\gamma_0}\phi_\epsilon + 1.1 < \frac{1}{2}f(\omega)e^{\phi_\epsilon} \quad \text{on } \mathbb{S}^2.$$

This is the desired inequality (3.11). Recall that ϕ_ϵ is given by

$$\phi_\epsilon = \tau[\chi_\epsilon w + (1 - \chi_\epsilon)\log \epsilon] - \log\left(\frac{m\epsilon^{\tau+2}}{2C_0C_1}\right).$$

We hence have (3.9) and

$$|\nabla_{\gamma_0}\phi_\epsilon| = O(\epsilon^{-1}), \quad |\nabla_{\gamma_0}^2\phi_\epsilon| = O(\epsilon^{-2}).$$

Moreover, by (3.3) and (3.5), $\bar{\phi} = \phi_\epsilon$ satisfies $\mathcal{S}(\bar{\phi}) < 0$ for any sufficiently large a . \square

The functions $\underline{\phi}$ and $\bar{\phi}$ in Lemma 3.3 and Lemma 3.7 are a subsolution and a supersolution of $\mathcal{S}(\phi) = 0$, respectively. In the following sections, we will construct a solution ϕ of $\mathcal{S}(\phi) = 0$ such that

$$(3.12) \quad \underline{\phi} \leq \phi \leq \bar{\phi} \quad \text{on } \mathbb{S}^2,$$

where the subsolution and supersolution serve as lower and upper barriers. Furthermore, we will prove that such a ϕ is unique.

4. MOSER'S ITERATION AND SCHAUDER ESTIMATES

In this section, we will derive Schauder estimates for solutions. We will present a detailed proof of $C^{1,1/3}(\mathbb{S}^2)$ -estimate, emphasizing on the explicit dependence of the L^∞ -bound of solutions. Again, we fix a \underline{u} and let $\mathcal{S}(\phi)$ be the operator defined by (3.1). We will prove the following result.

Theorem 4.1. *Let $\mathcal{S}(\phi)$ be defined by (3.1) and κ be a positive constant. Assume ϕ is a solution of $\mathcal{S}(\phi) = 0$, with $0 \leq \phi \leq \kappa$ on \mathbb{S}^2 . Then,*

$$|\phi|_{C^{2,1/3}(\mathbb{S}^2)} \leq (Ce^\kappa)^{Ce^{4\Lambda\kappa}},$$

where Λ is the ellipticity constant of the metric component (γ_{ij}) and C is a positive constant depending only on $\gamma, \hat{\chi}, \eta, \alpha_1, \alpha_2$ and their angular derivatives.

We will prove Theorem 4.1 in three steps. In the first step, we estimate the Hölder norms of the solutions. This step is based on the local boundedness and the weak Harnack inequality due to Moser. In the second step, we estimate the Hölder norms of the first derivatives. This step is based on an integral characterization of Hölder continuous functions. In the third step, we estimate the Hölder norms of the second derivatives. In the course of the proof, we keep track of the dependence on κ .

Let F be given by (2.19). We aim to establish a priori estimates for solutions $\phi = \phi(\omega)$ to the equation

$$(4.1) \quad -\Delta_\gamma \phi = 1 - \frac{1}{2} f e^\phi + F.$$

We point out that (4.1) is a quasilinear elliptic equation for ϕ , since F involves the gradient of ϕ and γ also depends on ϕ . We always assume (1.13), i.e.,

$$0 \leq f \leq 1 \text{ on } \mathbb{S}^2.$$

The equation (4.1) is defined globally on \mathbb{S}^2 . In the following, we will prove a more general result, valid in arbitrary dimensions.

In this section, to be aligned with the literature on elliptic equations, we let $B_r(p)$ be the standard ball in \mathbb{R}^n centered at p with radius r , for any integer $n \geq 2$. And sometimes we write B_r for short, when the center is at the origin. In other sections, to emphasize the radius r , we adopt the notation $B_p(r)$. We consider the nonlinear equation of the form

$$(4.2) \quad -\partial_j (a_{ij} \partial_i \phi) = \sqrt{|\gamma|} \left[1 - \frac{1}{2} f e^\phi + F \right].$$

This is the equation (4.1) in local coordinates. First, we assume

$$\Lambda^{-1} I \leq (a_{ij}(\omega, \phi)) \leq \Lambda I,$$

for some positive constant Λ , and

$$1/2 \leq \sqrt{|\gamma|}(\omega, \phi) \leq 2.$$

By abusing notations slightly, we set, for any $(\omega, \phi, p) \in B_1 \times \mathbb{R} \times \mathbb{R}^n$,

$$F(\omega, \phi, p) = a^{-\frac{1}{3}} e^\phi [c_{ij} p_i p_j + c_i p_i + c_0],$$

where c_{ij}, c_i, c_0 are functions of (ω, ϕ) on $B_1 \times \mathbb{R}_+$. By the Cauchy-Schwarz inequality, we have

$$|F| \leq C e^\phi a^{-\frac{1}{3}} (|D\phi|^2 + 1).$$

This is simply (3.3) in local coordinates. Next, we take a sufficiently large such that $a^{-\frac{1}{3}} e^\phi \ll 1$. Hence,

$$(4.3) \quad |F| \leq c(|D\phi|^2 + 1),$$

for some small positive constant c . In fact, we can take $c = 1$.

We now put (4.2) in divergence form as follows:

$$(4.4) \quad \int a_{ij}(\omega, \phi) D_i \phi D_j \psi \, d\omega = \int \sqrt{|\gamma|} \left[1 - \frac{1}{2} f e^\phi + F(\omega, \phi, D\phi) \right] \psi \, d\omega,$$

for any $\psi \in H_0^1(B_1) \cap L^\infty(B_1)$. We always assume $\phi = \phi(\omega) \in H^1(B_1)$ and

$$(4.5) \quad 0 \leq \phi \leq \kappa \text{ on } B_1,$$

for some positive constant κ .

In the following, we will derive estimates of the Hölder norms for the solution ϕ , its gradients, and its Hessians, successively, with a focus on explicit dependence in terms of κ .

4.1. Interior Estimates of Hölder Norms. Our first goal is to derive interior estimates of the Hölder norm of ϕ . In the proof, we will adapt some relevant results in [17] or [18].

Theorem 4.2. *Suppose $\phi \in H^1(B_1)$ is a positive solution to (4.4) with (4.5) satisfied. Then,*

$$(4.6) \quad [\phi]_{C^\alpha(B_{1/2})} \leq C e^\kappa,$$

where C is a positive constant depending only on n and Λ , and α is given by

$$(4.7) \quad \alpha = \epsilon_0 e^{-4\Lambda\kappa} \in (0, 1),$$

for some small constant $\epsilon_0 > 0$ depending on n and Λ .

Proof. Set, for $s \in (0, 1)$,

$$M(s) = \max_{B_s} \phi, \quad m(s) = \min_{B_s} \phi.$$

Fix an $s \in (0, 1)$ and consider ϕ defined in B_s . Write $m = m(s)$ for brevity and set

$$v = \phi - m.$$

For any $\psi \in H_0^1(B_s) \cap L^\infty(B_s)$ with $\psi \geq 0$, by (4.4) and (4.3), we have

$$\int a_{ij} D_i v D_j \psi \, d\omega \leq \int \sqrt{|\gamma|} \left[1 - \frac{1}{2} f e^\phi + |Dv|^2 + 1 \right] \psi \, d\omega \leq \int [4 + 2|Dv|^2] \psi \, d\omega,$$

and

$$\int a_{ij} D_i v D_j \psi \, d\omega \geq \int \sqrt{|\gamma|} \left[1 - \frac{1}{2} f e^\phi - |Dv|^2 - 1 \right] \psi \, d\omega \geq \int [-e^\phi - 2|Dv|^2] \psi \, d\omega.$$

First, we set $\underline{v} = \underline{\xi}(v)$ for some function $\underline{\xi}$ to be determined, with properties $\underline{\xi} \geq 0$, $\underline{\xi}' \geq 0$, and $\underline{\xi}'' \geq 0$. Then for $\psi \in H_0^1(B_s) \cap L^\infty(B_s)$ with $\psi \geq 0$, we have

$$\begin{aligned} \int a_{ij} D_i \underline{v} D_j \psi \, d\omega &= \int a_{ij} D_i v \underline{\xi}'(v) D_j \psi \, d\omega \\ &= \int a_{ij} D_i v D_j (\underline{\xi}'(v) \psi) \, d\omega - \int a_{ij} D_i v D_j v \underline{\xi}''(v) \psi \, d\omega \\ &\leq \int [4 + 2|Dv|^2] \underline{\xi}'(v) \psi \, d\omega - \int \Lambda^{-1} |Dv|^2 \underline{\xi}''(v) \psi \, d\omega \\ &= \int 4 \underline{\xi}'(v) \psi \, d\omega + \int |Dv|^2 [2 \underline{\xi}'(v) - \Lambda^{-1} \underline{\xi}''(v)] \psi \, d\omega. \end{aligned}$$

We take $\underline{\xi}$ such that $\underline{\xi}'' = 2\Lambda \underline{\xi}'$ with $\underline{\xi}(0) = 0$ and $\underline{\xi}'(0) = 1$. Then, it holds

$$(4.8) \quad \underline{\xi}(\tau) = \frac{1}{2\Lambda} (e^{2\Lambda\tau} - 1).$$

Hence, for $\psi \in H_0^1(B_s) \cap L^\infty(B_s)$ with $\psi \geq 0$, we have

$$\int a_{ij} D_i \underline{v} D_j \psi \, d\omega \leq \int 4 \underline{\xi}'(v) \psi \, d\omega \leq \int 4e^{2\Lambda\kappa} \psi \, d\omega,$$

where we used $\underline{\xi}'(v) = e^{2\Lambda(\phi-m)} \leq e^{2\Lambda\kappa}$.

By the local boundedness provided by Theorem 8.17 in [17] or Theorem 4.14 in [18], we obtain, for any $p > 0$,

$$\sup_{B_{s/2}} \underline{v} \leq C \left[s^{-\frac{n}{p}} \|\underline{v}\|_{L^p(B_s)} + e^{2\Lambda\kappa} s^2 \right].$$

With $\underline{v} = \underline{\xi}(v)$ and $1 \leq \underline{\xi}'(v) \leq e^{2\Lambda\kappa}$, by the mean value theorem, we have $v \leq \underline{v} \leq e^{2\Lambda\kappa} v$, and hence,

$$(4.9) \quad \sup_{B_{s/2}} v \leq C e^{2\Lambda\kappa} \left[s^{-\frac{n}{p}} \|v\|_{L^p(B_s)} + s^2 \right].$$

Next, we set $\bar{v} = \bar{\xi}(v)$ for some function $\bar{\xi}$ to be determined, with properties $\bar{\xi} \geq 0$, $\bar{\xi}' \geq 0$, and $\bar{\xi}'' \leq 0$. Then for $\psi \in H_0^1(B_s) \cap L^\infty(B_s)$ with $\psi \geq 0$, we have

$$\begin{aligned} \int a_{ij} D_i \bar{v} D_j \psi \, d\omega &= \int a_{ij} D_i v \bar{\xi}'(v) D_j \psi \, d\omega \\ &= \int a_{ij} D_i v D_j (\bar{\xi}'(v) \psi) \, d\omega - \int a_{ij} D_i v D_j v \bar{\xi}''(v) \psi \, d\omega \\ &\geq \int [-e^\phi - 2|Dv|^2] \bar{\xi}'(v) \psi \, d\omega - \int \Lambda^{-1} |Dv|^2 \bar{\xi}''(v) \psi \, d\omega. \end{aligned}$$

We take $\bar{\xi}$ such that $\bar{\xi}'' = -2\Lambda \bar{\xi}'$ with $\bar{\xi}(0) = 0$ and $\bar{\xi}'(0) = 1$. Then, it holds

$$\bar{\xi}(\tau) = \frac{1}{2\Lambda} (1 - e^{-2\Lambda\tau}).$$

Hence, for $\psi \in H_0^1(B_s) \cap L^\infty(B_s)$ with $\psi \geq 0$, we have

$$\int a_{ij} D_i \bar{v} D_j \psi \, d\omega \geq \int -e^\phi \bar{\xi}'(v) \psi \, d\omega \geq \int -e^\kappa \psi \, d\omega.$$

By the weak Harnack inequality provided by Theorem 8.18 in [17] or Theorem 4.15 in [18], we obtain, for any $p \in (0, \frac{n}{n-2})$,

$$s^{-\frac{n}{p}} \|\bar{v}\|_{L^p(B_s)} \leq C \left[\inf_{B_{s/2}} \bar{v} + e^\kappa s^2 \right].$$

With $\bar{v} = \bar{\xi}(v)$ and $e^{-2\Lambda\kappa} \leq \bar{\eta}'(\tau) \leq 1$, by the mean value theorem, we have $e^{-2\Lambda\kappa} v \leq \bar{v} \leq v$, and hence,

$$(4.10) \quad s^{-\frac{n}{p}} \|v\|_{L^p(B_s)} \leq C e^{2\Lambda\kappa} \left[\inf_{B_{s/2}} v + e^\kappa s^2 \right].$$

By combining (4.9) and (4.10), we get

$$\sup_{B_{s/2}} v \leq C e^{4\Lambda\kappa} \inf_{B_{s/2}} v + C e^{4\Lambda\kappa + \kappa} s^2.$$

With $v = \phi - m(s)$, we thus obtain

$$(4.11) \quad M\left(\frac{s}{2}\right) - m(s) \leq Ce^{4\Lambda\kappa} \left[m\left(\frac{s}{2}\right) - m(s) \right] + Ce^{4\Lambda\kappa + \kappa} s^2.$$

Write $M = M(s) = \max_{B_s} \phi$ for brevity and set

$$w = M - \phi.$$

For any $\psi \in H_0^1(B_s) \cap L^\infty(B_s)$ with $\psi \geq 0$, by (4.4) and (4.3), we have

$$\int a_{ij} D_i w D_j \psi \, d\omega \leq \int \sqrt{|\gamma|} \left[-1 + \frac{1}{2} f e^\phi + |Dw|^2 + 1 \right] \psi \, d\omega \leq \int [e^\phi + 2|Dw|^2] \psi \, d\omega,$$

and

$$\int a_{ij} D_i w D_j \psi \, d\omega \geq \int \sqrt{|\gamma|} \left[-1 + \frac{1}{2} f e^\phi - |Dv|^2 - 1 \right] \psi \, d\omega \geq \int [-4 - 2|Dw|^2] \psi \, d\omega.$$

Proceeding similarly as above for v , for $p \in (0, \frac{n}{n-2})$ we obtain

$$(4.12) \quad \sup_{B_{s/2}} w \leq Ce^{2\Lambda\kappa} \left[s^{-\frac{n}{p}} \|w\|_{L^p(B_s)} + e^\kappa s^2 \right]$$

and

$$(4.13) \quad s^{-\frac{n}{p}} \|w\|_{L^p(B_s)} \leq Ce^{2\Lambda\kappa} \left[\inf_{B_{s/2}} w + s^2 \right].$$

By combining (4.12) and (4.13), we have

$$\sup_{B_{s/2}} w \leq Ce^{4\Lambda\kappa} \inf_{B_{s/2}} w + Ce^{4\Lambda\kappa + \kappa} s^2,$$

and using $w = M(s) - \phi$, we then get

$$(4.14) \quad M(s) - m\left(\frac{s}{2}\right) \leq Ce^{4\Lambda\kappa} \left[M(s) - M\left(\frac{s}{2}\right) \right] + Ce^{4\Lambda\kappa + \kappa} s^2.$$

Set, for any $s \in (0, 1)$,

$$(4.15) \quad h(s) = M(s) - m(s).$$

Adding (4.11) and (4.14), we get

$$h(s) + h\left(\frac{s}{2}\right) \leq Ce^{4\Lambda\kappa} \left[h(s) - h\left(\frac{s}{2}\right) \right] + Ce^{4\Lambda\kappa + \kappa} s^2,$$

and hence

$$h\left(\frac{s}{2}\right) \leq \gamma h(s) + e^\kappa s^2,$$

where

$$\gamma = \frac{Ce^{4\Lambda\kappa} - 1}{Ce^{4\Lambda\kappa} + 1} < 1.$$

With this γ , we choose $\mu \in (0, 1)$ such that

$$\alpha := (1 - \mu) \log \gamma / \log(1/2) < 2\mu.$$

Since $\gamma = 1 - 2(1 + Ce^{4\Lambda\kappa})^{-1}$, where κ is large and C in (4.11) and (4.14) could be chosen to be large, we can write α as in (4.7), i.e., $\alpha = \epsilon_0 e^{-4\Lambda\kappa}$ for some small $\epsilon_0 > 0$. We then apply Lemma 8.23 in [17] or Lemma 4.19 in [18] and obtain, for any $s \in (0, 1/2]$,

$$(4.16) \quad h(s) \leq Cs^\alpha \{h(1) + e^\kappa s^{2\mu}\} \leq Ce^\kappa s^\alpha,$$

where we used $h(1) \leq M(1) \leq \kappa \leq e^\kappa$ for the second inequality. We then conclude (4.6). \square

4.2. Interior Gradient Estimates. Our next goal is to derive the gradient estimate of ϕ . We introduce an extra condition

$$(4.17) \quad |D_\omega a_{kl}(\omega, \phi)| + |\partial_\phi a_{kl}(\omega, \phi)| \leq \Lambda_1,$$

for some positive constant Λ_1 . In the proof below, we will adapt some relevant results in [18] and employ the characterization of Hölder continuous functions by Campanato.

Theorem 4.3. *Suppose $\phi \in H^1(B_1)$ is a positive solution to (4.4) with (4.5) satisfied. Then,*

$$(4.18) \quad |\phi|_{C^{1,1/3}(B_{1/2})} \leq (Ce^\kappa)^{Ce^{4\Lambda\kappa}},$$

where C is a positive constant depending only on n , Λ , and Λ_1 .

Proof. We adopt same notations as in the proof of Theorem 4.2. We first proceed as in the proof of Theorem 4.2 to obtain an interior estimate in the L^2 -norm for the gradient of ϕ . For any $\psi \in H_0^1(B_1) \cap L^\infty(B_1)$ with $\psi \geq 0$, by (4.4) and (4.3), we have

$$\int a_{ij} D_i \phi D_j \psi \, d\omega \leq \int \sqrt{|\gamma|} \left[1 - \frac{1}{2} f e^\phi + |D\phi|^2 + 1 \right] \psi \, d\omega \leq \int [4 + 2|D\phi|^2] \psi \, d\omega.$$

Taking $\underline{\xi}$ as in (4.8), we have

$$\int a_{ij} D_i (\underline{\xi}(\phi)) D_j \psi \, d\omega \leq \int 4\underline{\xi}'(\phi) \psi \, d\omega,$$

which is equivalent to

$$\int a_{ij} e^{2\Lambda\phi} D_i \phi D_j \psi \, d\omega \leq \int 4e^{2\Lambda\phi} \psi \, d\omega.$$

Take $\psi = \phi\mu^2$ for some $\mu \in C_0^1(B_1)$ with $\mu = 1$ on $B_{3/4}$. By the Cauchy-Schwarz inequality, we get

$$\int e^{2\Lambda\phi} |D\phi|^2 \mu^2 \, d\omega \leq C \int e^{2\Lambda\phi} (\phi^2 |D\mu|^2 + \phi\mu^2) \, d\omega,$$

and hence

$$(4.19) \quad \int_{B_{3/4}} |D\phi|^2 \, d\omega \leq C\kappa^2 e^{2\Lambda\kappa},$$

where C is a positive constant depending only on n and Λ .

We next derive an estimate of the Hölder norm of the gradient and prove

$$(4.20) \quad |D\phi|_{C^{1/3}(B_{1/2})} \leq (Ce^\kappa)^{Ce^{2\Lambda\kappa}}.$$

The proof below is similar to that of Theorem 4.24 in [18]. The difference is that in Theorem 4.24 of [18] $a_{ij} = a_{ij}(\omega)$, while here $a_{ij} = a_{ij}(\omega, \phi(\omega))$. We need to estimate $a_{ij}(\omega_0, \phi(\omega_0)) - a_{ij}(\omega, \phi(\omega))$ and to control $\phi(\omega) - \phi(\omega_0)$ with Theorem 4.2. In the following, C is a positive constant depending only on n and Λ . We denote by $\phi_{\omega, s}$ and $(D\phi)_{\omega, s}$ the average of ϕ and $D\phi$ over $B_s(\omega)$, respectively.

For any $B_s(\omega_0) \subset B_1$, take \hat{w} such that for any $\psi \in H_0^1(B_s(\omega_0))$ it holds

$$(4.21) \quad \int_{B_s(\omega_0)} a_{ij}(\omega_0, \phi(\omega_0)) D_i \hat{w} D_j \psi \, d\omega = 0,$$

with $\hat{w} - \phi \in H_0^1(B_s(\omega_0))$. The maximum principle implies

$$(4.22) \quad \inf_{B_s(\omega_0)} \phi \leq \hat{w} \leq \sup_{B_s(\omega_0)} \phi \quad \text{in } B_s(\omega_0).$$

For convenience, set $\hat{v} = \phi - \hat{w}$ and recall (4.15)

$$h(s) = h(\omega_0; s) = \sup_{B_s(\omega_0)} \phi - \inf_{B_s(\omega_0)} \phi.$$

Then, $\hat{v} \in H_0^1(B_s(\omega_0))$ and by (4.22) we get

$$(4.23) \quad \sup_{B_s(\omega_0)} |\hat{v}| \leq h(s).$$

By Corollary 3.11 in [18], we have for any $0 < \rho \leq s$

$$(4.24) \quad \int_{B_\rho(\omega_0)} |D\phi|^2 \, d\omega \leq c \left[\left(\frac{\rho}{s} \right)^n \int_{B_s(\omega_0)} |D\phi|^2 \, d\omega + \int_{B_s(\omega_0)} |D\hat{v}|^2 \, d\omega \right],$$

and

$$(4.25) \quad \int_{B_\rho(\omega_0)} |D\phi - (D\phi)_{\omega_0, \rho}|^2 \, d\omega \leq c \left[\left(\frac{\rho}{s} \right)^{n+2} \int_{B_s(\omega_0)} |D\phi - (D\phi)_{\omega_0, s}|^2 \, d\omega + \int_{B_s(\omega_0)} |D\hat{v}|^2 \, d\omega \right].$$

Moreover, by (4.21) and (4.4), we get for any $\psi \in H_0^1(B_s(\omega_0)) \cap L^\infty(B_s(\omega_0))$

$$\begin{aligned} & \int_{B_s(\omega_0)} a_{ij}(\omega_0, \phi(\omega_0)) D_i \hat{v} D_j \psi \, d\omega \\ &= \int_{B_s(\omega_0)} a_{ij}(\omega_0, \phi(\omega_0)) D_i \phi D_j \psi \, d\omega \\ &= \int_{B_s(\omega_0)} \sqrt{|\gamma|} \left[1 - \frac{1}{2} f(\omega) e^{\phi(\omega)} + F(\omega, \phi(\omega), D\phi(\omega)) \right] \psi \, d\omega \\ &+ \int_{B_s(\omega_0)} [a_{ij}(\omega_0, \phi(\omega_0)) - a_{ij}(\omega, \phi(\omega))] D_i \phi D_j \psi \, d\omega. \end{aligned}$$

Note that, for any $\omega \in B_s(\omega_0)$ it holds

$$|a_{ij}(\omega_0, \phi(\omega_0)) - a_{ij}(\omega, \phi(\omega))| \leq C[|\omega - \omega_0| + |\phi(\omega) - \phi(\omega_0)|] \leq C[s + h(s)].$$

Taking $\psi = \hat{v}$, together with (4.3), we obtain

$$\int_{B_s(\omega_0)} |D\hat{v}|^2 d\omega \leq C \int_{B_s(\omega_0)} \left[[s + h(s)] |D\phi| |D\hat{v}| + |D\phi|^2 |\hat{v}| + e^\kappa |\hat{v}| \right] d\omega.$$

We apply the Cauchy-Schwarz inequality to the first term in the right-hand side, substitute \hat{v} in the second term with (4.23), and apply the Sobolev inequality to \hat{v} in the third term as follows:

$$\int_{B_s(\omega_0)} |\hat{v}| d\omega \leq cs^{\frac{n+2}{2}} \left(\int_{B_s(\omega_0)} |D\hat{v}|^2 d\omega \right)^{\frac{1}{2}}.$$

Then, we obtain

$$(4.26) \quad \int_{B_s(\omega_0)} |D\hat{v}|^2 d\omega \leq C(h(s) + [h(s)]^2 + s^2) \int_{B_s(\omega_0)} |D\phi|^2 d\omega + Ce^{2\kappa} s^{n+2}.$$

Therefore, by (4.24) and (4.26), we have for any $0 < \rho \leq s$

$$\int_{B_\rho(\omega_0)} |D\phi|^2 d\omega \leq C \left[\left(\frac{\rho}{s} \right)^n + h(s) + [h(s)]^2 + s^2 \right] \int_{B_s(\omega_0)} |D\phi|^2 d\omega + Ce^{2\kappa} s^{n+2}.$$

Take any $\delta \in (0, 1)$. For a small $\epsilon_\delta > 0$, using (4.16) we can choose some s_δ small such that $B_{s_\delta}(\omega_0) \subset B_{3/4}$ and

$$(4.27) \quad h(s_\delta) + [h(s_\delta)]^2 + s_\delta^2 < \epsilon_\delta.$$

By Lemma 3.4 in [18], we have for any $s \leq s_\delta$

$$\int_{B_s(\omega_0)} |D\phi|^2 d\omega \leq Cs^{n-2+2\delta} \left[s_\delta^{-(n-2+2\delta)} \int_{B_{s_\delta}(\omega_0)} |D\phi|^2 d\omega + e^{2\kappa} \right].$$

By the choice of α in (4.7) and (4.6), using (4.16) together with $\kappa \gg 1$, we can take suitable positive constants C_1 and C_2 depending on n, Λ and

$$(4.28) \quad s_\delta = (C_1 e^\kappa)^{-C_2 e^{4\Lambda\kappa}},$$

so that (4.27) is satisfied. By (4.19), we get

$$(4.29) \quad \int_{B_s(\omega_0)} |D\phi|^2 d\omega \leq Cs^{n-2+2\delta} \left[\kappa^2 e^{2\Lambda\kappa} s_\delta^{-(n-2+2\delta)} + e^{2\kappa} \right].$$

By the Poincaré inequality, we have for any $s \leq s_\delta$ with $B_{s_\delta}(\omega_0) \subset B_{3/4}$

$$\int_{B_s(\omega_0)} |\phi - \phi_{\omega_0, s}|^2 d\omega \leq Cs^{n+2\delta} \left[\kappa^2 e^{2\Lambda\kappa} s_\delta^{-(n-2+2\delta)} + e^{2\kappa} \right].$$

Hence, by Theorem 3.1 in [18], we obtain for any $s \leq s_\delta$ with $B_{s_\delta}(\omega_0) \subset B_{3/4}$

$$(4.30) \quad h(\omega_0; s) = \sup_{B_s(\omega_0)} \phi - \inf_{B_s(\omega_0)} \phi \leq C \left[\kappa e^{\Lambda\kappa} s_\delta^{-(\frac{n}{2}-1+\delta)} + e^\kappa \right] s^\delta.$$

With (4.26), (4.29), and (4.30), we have for any $s \leq s_\delta$ with $B_{s_\delta}(\omega_0) \subset B_{3/4}$

$$\int_{B_s(\omega_0)} |D\hat{v}|^2 d\omega \leq C\kappa^3 e^{3\Lambda\kappa} s_\delta^{-\frac{3}{2}(n-2+2\delta)} s^{n-2+3\delta} + Ce^{2\kappa} s^{n+2}.$$

We take $\delta \in (2/3, 1)$ and write $\alpha' = 3\delta/2 - 1$. Note that $\alpha' \in (0, 1/2)$ and α' can be as close as to $1/2$ by taking δ as close as to 1. With (4.25) we obtain, for any $\rho \leq s \leq s_\delta$ with $B_{s_\delta}(\omega_0) \subset B_{3/4}$,

$$\begin{aligned} \int_{B_\rho(\omega_0)} |D\phi - (D\phi)_{\omega_0, \rho}|^2 d\omega &\leq C \left(\frac{\rho}{s}\right)^{n+2} \int_{B_s(\omega_0)} |D\phi - (D\phi)_{\omega_0, s}|^2 d\omega \\ &\quad + C\kappa^3 e^{3\Lambda\kappa} s_\delta^{-\frac{3}{2}(n-2+2\delta)} s^{n+2\alpha'} + C e^{2\kappa} s^{n+2}. \end{aligned}$$

By Lemma 3.4 in [18], together with (4.19), we have for any $s \leq s_\delta$ with $B_{s_\delta}(\omega_0) \subset B_{3/4}$

$$\int_{B_s(\omega_0)} |D\phi - (D\phi)_{\omega_0, s}|^2 d\omega \leq C\kappa^3 e^{3\Lambda\kappa} s_\delta^{-\frac{3}{2}(n-2+2\delta)} s^{n+2\alpha'}.$$

By Theorem 3.1 in [18], together with (4.19), we obtain for any $\omega \in B_{1/2}$

$$(4.31) \quad |D\phi(\omega)| \leq C \left[\kappa^{\frac{3}{2}} e^{\frac{3}{2}\Lambda\kappa} s_\delta^{-\frac{3}{4}(n-2+2\delta)} + \kappa e^{\Lambda\kappa} \right] \leq C\kappa^{\frac{3}{2}} e^{\frac{3}{2}\Lambda\kappa} s_\delta^{-\frac{3}{4}(n-2+2\delta)},$$

and for any $\omega_1, \omega_2 \in B_{1/2}$ with $|\omega_1 - \omega_2| < s_\delta$,

$$|D\phi(\omega_1) - D\phi(\omega_2)| \leq C\kappa^{\frac{3}{2}} e^{\frac{3}{2}\Lambda\kappa} s_\delta^{-\frac{3}{4}(n-2+2\delta)} |\omega_1 - \omega_2|^{\alpha'}.$$

Moreover, for any $\omega_1, \omega_2 \in B_{1/2}$ it holds

$$(4.32) \quad |D\phi(\omega_1) - D\phi(\omega_2)| \leq C\kappa^{\frac{3}{2}} e^{\frac{3}{2}\Lambda\kappa} s_\delta^{-\frac{3}{4}(n-2+2\delta)} s_\delta^{-\alpha'} |\omega_1 - \omega_2|^{\alpha'}.$$

Recall s_δ as in (4.28) and note $0 < \alpha' < 1/2$. We can choose $\alpha' = 1/3$ and, with (4.31) and (4.32), we hence conclude (4.20). \square

4.3. Interior Schauder Estimates. Our final goal is to prove the following Schauder estimates. We introduce an extra assumption

$$(4.33) \quad |D_\omega^2 a_{kl}(\omega, \phi)| + |D_\omega \partial_\phi a_{kl}(\omega, \phi)| + |\partial_\phi^2 a_{kl}(\omega, \phi)| \leq \Lambda_2,$$

for some positive constant Λ_2 .

Theorem 4.4. *Let $\phi \in H^1(B_1)$ be a positive solution to (4.4) with (4.5) satisfied, and $f \in C^{1/3}(B_1)$. Then,*

$$(4.34) \quad |\phi|_{C^{2,1/3}(B_{1/2})} \leq (C e^\kappa)^{C e^{4\Lambda\kappa}},$$

where C is a positive constant depending only on $n, \Lambda, \Lambda_1, \Lambda_2$, the C^1 -norms of $\sqrt{|\gamma|}$, c_{ij} , c_i , and c_0 on $B_1 \times \mathbb{R}_+$, and the $C^{1/3}$ -norm of f on B_1 .

Proof. We write (4.4) in nondivergence form

$$(4.35) \quad a_{ij}(\omega, \phi) \partial_{ij} \phi = h(\omega),$$

where

$$h(\omega) = -\partial_j a_{ij}(\omega, \phi) \partial_i \phi - \partial_\phi a_{ij}(\omega, \phi) \partial_i \phi \partial_j \phi - \sqrt{|\gamma|} \left[1 - \frac{1}{2} f e^\phi + F(\omega, \phi, D\phi) \right].$$

We treat the equation (4.35) as a linear equation of ϕ , with h as a given function defined in B_1 . By the explicit expression of F and Theorem 4.3, we have

$$|h|_{C^{1/3}(B_{3/4})} \leq (Ce^\kappa)^{Ce^{4\Lambda\kappa}}.$$

If we quote the standard interior Schauder estimate, we will get

$$|\phi|_{C^{2,1/3}(B_{1/2})} \leq C_* [|\phi|_{L^\infty(B_{3/4})} + |h|_{C^{1/3}(B_{3/4})}] \leq C_*(Ce^\kappa)^{Ce^{4\Lambda\kappa}},$$

where C_* is a positive constant depending only on n , Λ , and the $C^{1/3}$ -norm of a_{ij} on $B_{3/4}$. Since the leading coefficient a_{ij} is also a function of ϕ , i.e., $a_{ij} = a_{ij}(\omega, \phi)$, then C_* also depends on ϕ .

Instead of quoting the interior Schauder estimate, we need to examine its proof by the technique of *freezing* coefficients, and to keep track of the dependence on the Hölder norms of the leading coefficients. In this process, all contributions due to the dependence of a_{ij} on ϕ can be absorbed by $(Ce^\kappa)^{Ce^{4\Lambda\kappa}}$. Therefore, we obtain the desired estimate (4.34). \square

We are ready to prove Theorem 4.1.

Proof of Theorem 4.1. Fix an arbitrary ball B on \mathbb{S}^2 . In local coordinates within B , we write (4.1) as (4.2). Specifically, we set

$$a_{ij}(\omega, \phi) = \sqrt{|\gamma(\omega, u)|} \gamma^{ij}(\omega, u),$$

with $u = 1 - R = 1 - \underline{u}ae^{-\phi}$. By (2.15), (2.16), and (2.17), we obtain (4.17) and (4.33) by renaming Λ_1 and Λ_2 if necessary. Next, we examine the function $F(\phi, \underline{u})$ given by (2.19). Note that F can be viewed as a linear combination of $\nabla_i \phi \nabla_j \phi, \nabla_i \phi, 1$ with coefficients given by $\gamma^{ij}, R^{-1} \Omega \hat{\chi}_{kl}, \eta_i, R\alpha_1$, and $R\alpha_2$, which are functions of ω and $u = 1 - R = 1 - \underline{u}ae^{-\phi}$. In order to get an estimate of the $C^{1/3}$ -norm of F , we need to bound these coefficients and their derivatives with respect to ω and ϕ . Note $\partial_\phi = R\partial_u$. Hence, we need to bound

$$\begin{aligned} & R^{-1} \Omega \hat{\chi}_{kl}, \eta_k, R\alpha_1, R\alpha_2, \\ & R^{-1} \nabla_i (\Omega \hat{\chi}_{kl}), \nabla_i \eta_k, R\nabla_i \alpha_1, R\nabla_i \alpha_2, \\ & \partial_u (\Omega \hat{\chi}_{kl}), R\partial_u \eta_k, R^2 \partial_u \alpha_1, R^2 \partial_u \alpha_2. \end{aligned}$$

By (A.1) and (A.2), these quantities are bounded by $e^\phi a^{-\frac{1}{3}}$. Hence, we have an estimate of the $C^{1/3}$ -norm of F . Therefore, we can apply Theorem 4.2, Theorem 4.3, and Theorem 4.4 successively, and obtain the desired result. \square

5. THE LINEARIZED EQUATION

In this section, we derive and analyze the linearized equation. As in previous sections, we fix \underline{u} .

We first compute the linearized operator. The approach here is different from that in [3], where techniques via commutator formulas are used. Here we adopt a more direct calculation. Let $\mathcal{S}(\phi)$ be the operator defined by (2.18).

Proposition 5.1. *Let ϕ be a given function on \mathbb{S}^2 . The linearized operator $\partial_\phi \mathcal{S}(\phi)$ is of the form*

$$(5.1) \quad \partial_\phi \mathcal{S}(\phi)[w] = \Delta_\gamma w - \frac{1}{2}f(\omega)e^\phi w + c^i \nabla_i w + cw,$$

where the connection is with respect to the metric γ , and $c = c(\omega, \phi, \underline{u})$ and $c^i = c^i(\omega, \phi, \underline{u})$ are functions satisfying

$$|c| + |c^i| \leq Ca^{-\frac{1}{3}}e^\phi(|\nabla_\gamma^2 \phi| + |\nabla_\gamma \phi|^2 + 1).$$

Proof. The proof is divided into four steps.

Step 1. Setup. Recall the relations given by (2.14) and (2.20); namely, $g = R^2\gamma$, and

$$u = 1 - R, \quad R = \underline{u}ae^{-\phi}.$$

By (2.18) and (2.19), we write

$$(5.2) \quad \mathcal{S}(\phi) = I_1 + I_2 + I_3 + I_4 + I_5,$$

where

$$(5.3) \quad I_1 = \Delta_\gamma \phi + 1 - \frac{1}{2}fe^\phi$$

and

$$(5.4) \quad \begin{aligned} I_2 &= 2R^{-1}\Omega \hat{\underline{\chi}}_{kl} \gamma^{ik} \gamma^{jl} \nabla_i \phi \nabla_j \phi, & I_3 &= 2\gamma^{ij} \eta_i \nabla_j \phi, \\ I_4 &= R\alpha_1 |\nabla_\gamma \phi|^2, & I_5 &= R\alpha_2. \end{aligned}$$

Step 2. Parametrized functions. Now, we fix functions ϕ, R with $R = \underline{u}ae^{-\phi}, u = 1 - R$. Consider $\phi(\epsilon) = \phi + \epsilon w$. Then, $R(\epsilon) = Re^{-\epsilon w}$. We proceed to compute the linearized operator

$$\partial_\phi \mathcal{S}(\phi)[w] = \left. \frac{d}{d\epsilon} \right|_{\epsilon=0} \mathcal{S}(\phi + \epsilon w).$$

For brevity, we denote $\dot{\cdot} = \left. \frac{d}{d\epsilon} \right|_{\epsilon=0}$. Hence,

$$\dot{\phi} = w, \quad \dot{R} = -Rw, \quad \dot{u} = Rw.$$

Note that $\gamma, \underline{\chi}, \chi, \eta, \Omega, \underline{\omega}$ all depend on ϵ through $u = 1 - R$. We now derive several general formulas in terms of $\dot{\gamma}$.

Note that

$$\dot{\gamma}^{ij} = -\gamma^{ik} \gamma^{jl} \dot{\gamma}_{kl}.$$

By $|\nabla_\gamma \phi|^2 = \gamma^{ij} \nabla_i \phi \nabla_j \phi$, we obtain

$$(5.5) \quad (|\nabla_\gamma \phi|^2)^\cdot = 2\gamma^{ij} \nabla_i \phi \nabla_j w - \gamma^{ik} \gamma^{jl} \dot{\gamma}_{kl} \nabla_i \phi \nabla_j \phi.$$

By applying Lemma 3.5 to γ , together with $\dot{\phi} = w$, we have

$$(5.6) \quad (\Delta_\gamma \phi)^\cdot = \Delta_\gamma w - \gamma^{ik} \gamma^{jl} \dot{\gamma}_{kl} \nabla_{ij} \phi - \gamma^{ij} \gamma^{kl} \nabla_i \dot{\gamma}_{jl} \nabla_k \phi + \frac{1}{2} \gamma^{ij} \gamma^{kl} \nabla_l \dot{\gamma}_{ij} \nabla_k \phi.$$

Recall (2.6), i.e.,

$$\partial_u g_{kl} = \Omega \text{tr}_g \underline{\chi} g_{kl} + 2\Omega \hat{\underline{\chi}}_{kl}.$$

Then, with $g = R^2\gamma$,

$$\partial_u \gamma_{kl} = R^{-1}(R\Omega \text{tr}_g \underline{\chi} + 2)\gamma_{kl} + 2R^{-2}\Omega \hat{\underline{\chi}}_{kl},$$

and hence

$$\dot{\gamma}_{kl} = \dot{u} \partial_u \gamma_{kl} = w(R\Omega \text{tr}_g \underline{\chi} + 2)\gamma_{kl} + 2wR^{-1}\Omega \hat{\underline{\chi}}_{kl}.$$

A substitution in (5.5) and (5.6) yields

$$(5.7) \quad (|\nabla_\gamma \phi|^2)^\cdot = 2\gamma^{ij} \nabla_j \phi \nabla_i w - w(R\Omega \text{tr}_g \underline{\chi} + 2)|\nabla_\gamma \phi|^2 - 2wR^{-1}\Omega \hat{\underline{\chi}}_{kl} \gamma^{ik} \gamma^{jl} \nabla_i \phi \nabla_j \phi,$$

and

$$(5.8) \quad \begin{aligned} (\Delta_\gamma \phi)^\cdot &= \Delta_\gamma w - 2R^{-1}\Omega \hat{\underline{\chi}}_{kl} \gamma^{ik} \gamma^{jl} \nabla_j \phi \nabla_i w \\ &\quad - w[(R\Omega \text{tr}_g \underline{\chi} + 2)\Delta_\gamma \phi + 2R^{-1}\nabla_i(\Omega \hat{\underline{\chi}}_{kl})\gamma^{ik} \gamma^{jl} \nabla_j \phi \\ &\quad + 2R^{-1}\Omega \hat{\underline{\chi}}_{kl} \gamma^{ik} \gamma^{jl} (\nabla_{ij} \phi + \nabla_i \phi \nabla_j \phi)]. \end{aligned}$$

Step 3. Computations of \dot{I}_i , for $i = 1, \dots, 5$. We first compute \dot{I}_1 . By (5.3) and (5.8), we have

$$(5.9) \quad \begin{aligned} \dot{I}_1 &= \Delta_\gamma w - \frac{1}{2} f e^\phi w - 2R^{-1}\Omega \hat{\underline{\chi}}_{jl} \gamma^{ij} \gamma^{kl} \nabla_k \phi \nabla_i w - w(2 + R\Omega \text{tr}_g \underline{\chi}) \Delta_\gamma \phi \\ &\quad - 2w[R^{-1}\nabla_i(\Omega \hat{\underline{\chi}}_{jl})\gamma^{ij} \gamma^{kl} \nabla_k \phi + R^{-1}\Omega \hat{\underline{\chi}}_{kl} \gamma^{ik} \gamma^{jl} (\nabla_{ij} \phi + \nabla_i \phi \nabla_j \phi)]. \end{aligned}$$

We note that the first two terms in \dot{I}_1 are the first two terms on the right-hand side of (5.1). We will demonstrate that the rest terms in \dot{I}_1 and all other \dot{I}_i , for $i = 2, \dots, 5$, are in the forms of the last two terms on the right-hand side of (5.1).

Without presenting details of standard calculation process, here we list the expressions of \dot{I}_i , for $i = 2, \dots, 5$. We have

$$(5.10) \quad \begin{aligned} \dot{I}_2 &= 4R^{-1}\Omega \hat{\underline{\chi}}_{kl} \gamma^{ik} \gamma^{jl} \nabla_j \phi \nabla_i w \\ &\quad + 2w(\partial_u(\Omega \hat{\underline{\chi}}_{kl}) + R^{-1}\Omega \hat{\underline{\chi}}_{kl})\gamma^{ik} \gamma^{jl} \nabla_i \phi \nabla_j \phi \\ &\quad - 4wR^{-1}\Omega \hat{\underline{\chi}}_{kl} ((R\Omega \text{tr}_g \underline{\chi} + 2)\gamma^{ki} + 2R^{-1}\Omega \hat{\underline{\chi}}_{pq} \gamma^{ip} \gamma^{kq})\gamma^{lj} \nabla_i \phi \nabla_j \phi, \\ \dot{I}_3 &= 2\gamma^{ij} \eta_j \nabla_i w + 2wR \partial_u \eta_i \gamma^{ij} \nabla_j \phi \\ &\quad - 2w((R\Omega \text{tr}_g \underline{\chi} + 2)\gamma^{ij} + 2R^{-1}\Omega \hat{\underline{\chi}}_{kl} \gamma^{ik} \gamma^{jl})\eta_i \nabla_j \phi, \\ \dot{I}_4 &= 2R\alpha_1 \gamma^{ij} \nabla_i \phi \nabla_j w + w(R^2 \partial_u \alpha_1 - R\alpha_1)|\nabla_\gamma \phi|^2 \\ &\quad - wR\alpha_1 ((R\Omega \text{tr}_g \underline{\chi} + 2)\gamma^{ij} + 2R^{-1}\Omega \hat{\underline{\chi}}_{kl} \gamma^{ik} \gamma^{jl})\nabla_i \phi \nabla_j \phi, \\ \dot{I}_5 &= w(R^2 \partial_u \alpha_2 - R\alpha_2). \end{aligned}$$

Step 4. Expressions of coefficients. We now collect \dot{I}_i , for $1 \leq i \leq 5$, and write the sum in the form of (5.1). We obtain

$$(5.11) \quad c_i = 2R^{-1}\Omega \hat{\underline{\chi}}_{kl} \gamma^{ik} \gamma^{jl} \nabla_j \phi + 2\gamma^{ij} \eta_j + 2R\alpha_1 \gamma^{ij} \nabla_j \phi,$$

and

$$\begin{aligned}
(5.12) \quad c = & -(2 + R\Omega\text{tr}_g\chi)\Delta_\gamma\phi \\
& - 2[R^{-1}\Omega\hat{\chi}_{kl}\gamma^{ik}\gamma^{jl}(\nabla_{ij}\phi + \nabla_i\phi\nabla_j\phi) + R^{-1}\nabla_i(\Omega\hat{\chi}_{jl})\gamma^{ij}\gamma^{kl}\nabla_k\phi] \\
& - ((R\Omega\text{tr}_g\chi + 2)\gamma^{ij} + 2R^{-1}\Omega\hat{\chi}_{kl}\gamma^{ik}\gamma^{jl}) \\
& \quad \cdot (4R^{-1}\Omega\hat{\chi}_{ip}\gamma^{pq}\nabla_q\phi\nabla_j\phi + 2\eta_i\nabla_j\phi + R\alpha_1\nabla_i\phi\nabla_j\phi) \\
& + 2(\partial_u(\Omega\hat{\chi}_{kl}) + R^{-1}\Omega\hat{\chi}_{kl})\gamma^{ik}\gamma^{jl}\nabla_i\phi\nabla_j\phi + 2R\partial_u\eta_i\gamma^{ij}\nabla_j\phi \\
& + (R^2\partial_u\alpha_1 - R\alpha_1)|\nabla_\gamma\phi|^2 + R^2\partial_u\alpha_2 - R\alpha_2.
\end{aligned}$$

We point out that c_1, c_2 , and c can be viewed as linear combinations of $\nabla_{ij}\phi$, $\nabla_i\phi\nabla_j\phi$, $\nabla_i\phi$, and 1, with coefficients given by γ^{ij} and

$$\begin{aligned}
(5.13) \quad & 2 + R\Omega\text{tr}_g\chi, R^{-1}\Omega\hat{\chi}_{kl}, \eta_i, R\alpha_1, R\alpha_2, R^{-1}\nabla_i(\Omega\hat{\chi}_{kl}), \\
& \partial_u(\Omega\hat{\chi}_{kl}), R\partial_u\eta_i, R^2\partial_u\alpha_1, R^2\partial_u\alpha_2.
\end{aligned}$$

By (A.1) and (A.2), all expressions in (5.13) are controlled by $a^{-\frac{1}{3}}e^\phi$. \square

Next, we discuss the invertibility of the linearized operator $\partial_\phi S(\phi)$. We first examine the linear operator introduced by the first two terms on the left-hand side of (5.1). Set

$$(5.14) \quad L_0w = \Delta_\gamma w - \frac{1}{2}fe^\phi w.$$

Here, we view f and ϕ as two fixed nonnegative functions on \mathbb{S}^2 .

Lemma 5.2. *Let f and ϕ be given nonnegative functions on \mathbb{S}^2 , with (1.14) satisfied. Then the equation $L_0w = 0$ admits only the trivial solution. Moreover, the first eigenvalue μ_1 of $-L_0$ is positive and simple, and has a positive eigenfunction.*

Proof. Integrating $-wL_0w$ by parts on \mathbb{S}^2 , we have

$$\int_{\mathbb{S}^2} (|\nabla_\gamma w|^2 + \frac{1}{2}fe^\phi w^2) d\omega = - \int_{\mathbb{S}^2} wL_0w d\omega.$$

By restricting the second integral to $B_p(\epsilon)$, using $f \geq m$ in $B_p(\epsilon)$ in (1.14), we get

$$\int_{\mathbb{S}^2} |\nabla_\gamma w|^2 d\omega + \frac{1}{2}m \int_{B_p(\epsilon)} w^2 d\omega \leq - \int_{\mathbb{S}^2} wL_0w d\omega.$$

It is easy to verify

$$\int_{\mathbb{S}^2} w^2 d\omega \leq C \left[\int_{\mathbb{S}^2} |\nabla_\gamma w|^2 d\omega + \int_{B_p(\epsilon)} w^2 d\omega \right].$$

Hence,

$$\int_{\mathbb{S}^2} w^2 d\omega \leq C \int_{\mathbb{S}^2} |wL_0w| d\omega.$$

A simple application of the Cauchy-Schwarz inequality implies

$$\int_{\mathbb{S}^2} w^2 d\omega \leq C \int_{\mathbb{S}^2} |L_0w|^2 d\omega.$$

Hence, $L_0 w = 0$ implies $w = 0$.

Set

$$\langle w_1, w_2 \rangle = \int_{\mathbb{S}^2} \left(\nabla_\gamma w_1 \cdot \nabla_\gamma w_2 + \frac{1}{2} f e^\phi w_1 w_2 \right) d\omega.$$

The arguments above also demonstrate that $\langle w_1, w_2 \rangle$ is an inner-product on $H^1(\mathbb{S}^2)$, which induces a norm equivalent to the standard $H^1(\mathbb{S}^2)$ -norm. The standard L^2 -theory of elliptic equations applies to the operator L_0 . In particular, the first eigenvalue of $-L_0$ is characterized by

$$\mu_1 = \inf \left\{ \int_{\mathbb{S}^2} \left(|\nabla_\gamma w|^2 + \frac{1}{2} f e^\phi w^2 \right) d\omega; \int_{\mathbb{S}^2} w^2 d\omega = 1 \right\}.$$

Hence, $\mu_1 > 0$. Moreover, μ_1 is simple and has a positive eigenfunction. Refer to Theorem 8.38 in [17]. We need to point out that Theorem 8.38 in [17] is formulated for self-adjoint uniformly elliptic operators on bounded domains in the Euclidean space with zero Dirichlet data. However, the same conclusion holds for self-adjoint uniformly elliptic operators on closed manifolds. \square

We now prove the crucial invertibility of the linearized operators.

Lemma 5.3. *Let $\beta \in (0, 1)$ be a fixed constant, and f and ϕ be given nonnegative functions on \mathbb{S}^2 , with (1.14) satisfied and $|\phi|_{C^2(\mathbb{S}^2)} \leq K$ for some positive constant K . Then, $\partial_\phi \mathcal{S}(\phi) : C^{2,\beta}(\mathbb{S}^2) \rightarrow C^{0,\beta}(\mathbb{S}^2)$ is invertible, for a sufficiently large depending on K .*

Proof. Let L_0 be the operator as in (5.14). By Lemma 5.2, we can take a positive eigenfunction ψ_1 corresponding to the first eigenvalue $\mu_1 > 0$; namely,

$$\Delta_\gamma \psi_1 - \frac{1}{2} f e^\phi \psi_1 = -\mu_1 \psi_1.$$

By Proposition 5.1, we have

$$\begin{aligned} \partial_\phi \mathcal{S}(\phi)[\psi_1] &= \Delta_\gamma \psi_1 - \frac{1}{2} f e^\phi \psi_1 + c^i \nabla_i \psi_1 + c \psi_1 \\ &= -\mu_1 \psi_1 + c^i \nabla_i \psi_1 + c \psi_1. \end{aligned}$$

Hence, $\partial_\phi \mathcal{S}(\phi)[\psi_1] < 0$ on \mathbb{S}^2 by choosing a appropriately large depending on K . By Theorem 2.11 and its proof in [18], the strong maximum principle holds for the operator $\partial_\phi \mathcal{S}(\phi)$. Thus, for any $w \in C^2(\mathbb{S}^2)$,

$$(5.15) \quad \max_{\mathbb{S}^2} |w| \leq C \max_{\mathbb{S}^2} |\partial_\phi \mathcal{S}(\phi)[w]|,$$

where C is a positive constant independent of w . By the Schauder theory, we further obtain for any $w \in C^{2,\beta}(\mathbb{S}^2)$

$$|w|_{C^{2,\beta}(\mathbb{S}^2)} \leq C |\partial_\phi \mathcal{S}(\phi)[w]|_{C^{0,\beta}(\mathbb{S}^2)},$$

for some positive constant C independent of w . Therefore, $\partial_\phi \mathcal{S}(\phi) : C^{2,\beta}(\mathbb{S}^2) \rightarrow C^{0,\beta}(\mathbb{S}^2)$ is invertible. \square

In (5.15), the constant C may depend on the given function ϕ . The dependence of C on ϕ arises from the dependence of the eigenfunction ψ_1 on ϕ . This is because the operator L_0 depends on ϕ via the factor e^ϕ and the metric γ , which is parametrized by $u = 1 - \underline{u}ae^{-\phi}$. Moreover, this constance C may also depend on \underline{u} through the function $f = f(\omega, \underline{u})$. In the next result, we will eliminate such dependence on specific ϕ and \underline{u} .

Lemma 5.4. *Let f and ϕ be given nonnegative functions on \mathbb{S}^2 , with (1.14) satisfied and $|\phi|_{C^2(\mathbb{S}^2)} \leq K$ for some positive constant K . Then, for a sufficiently large depending on K and for any $w \in C^2(\mathbb{S}^2)$, it holds*

$$(5.16) \quad \max_{\mathbb{S}^2} |w| \leq C \max_{\mathbb{S}^2} |\partial_\phi \mathcal{S}(\phi)[w]|,$$

where C is a positive constant depending only on m and ϵ in (1.14) and K , independent of specific ϕ and $\underline{u} \in (0, \delta]$.

Proof. We first choose a smooth nonnegative function $f_0 = f_0(\omega)$ on \mathbb{S}^2 such that $f_0 \geq m/2$ on $B_p(\epsilon)$ and $f(\cdot, \underline{u}) \geq f_0$ on \mathbb{S}^2 for any $\underline{u} \in (0, \delta]$. We take the standard spherical metric γ_0 on \mathbb{S}^2 . Instead of L_0 in (5.14), we consider

$$L_* w = \Delta_{\gamma_0} w - \frac{1}{2} f_0 w.$$

By applying Lemma 5.2 to L_* , we can take a positive eigenfunction ψ_1 corresponding to the first eigenvalue $\mu_1 > 0$; namely,

$$\Delta_{\gamma_0} \psi_1 - \frac{1}{2} f \psi_1 = -\mu_1 \psi_1.$$

We point out that μ_1 and ψ_1 depend only on m and ϵ , independent of $\underline{u} \in (0, \delta]$. By $\phi \geq 0$, $f \geq f_0$, Lemma 3.6, and Proposition 5.1, we have

$$\begin{aligned} \partial_\phi \mathcal{S}(\phi)[\psi_1] &= \Delta_{\gamma_0} \psi_1 - \frac{1}{2} f e^\phi \psi_1 + \Delta_\gamma \psi_1 - \Delta_{\gamma_0} \psi_1 + c^i \nabla_i \psi_1 + c \psi_1 \\ &\leq \Delta_{\gamma_0} \psi_1 - \frac{1}{2} f_0 \psi_1 + \Delta_\gamma \psi_1 - \Delta_{\gamma_0} \psi_1 + c^i \nabla_i \psi_1 + c \psi_1 \\ &= -\mu_1 \psi_1 + \Delta_\gamma \psi_1 - \Delta_{\gamma_0} \psi_1 + c^i \nabla_i \psi_1 + c \psi_1 \leq -c(m, \epsilon, K) < 0, \end{aligned}$$

via choosing a appropriately large depending on m , ϵ , and K . By Theorem 2.11 and its proof in [18], the strong maximum principle holds, and in particular we obtain (5.16). \square

We further prove the following result.

Lemma 5.5. *Let f and ϕ_1, ϕ_2 be given nonnegative functions on \mathbb{S}^2 , with (1.14) satisfied and $|\phi_i|_{C^2(\mathbb{S}^2)} \leq K$ for some positive constant K and $i = 1, 2$. If $\mathcal{S}(\phi_1) \geq \mathcal{S}(\phi_2)$, then either $\phi_1 < \phi_2$ or $\phi_1 \equiv \phi_2$ on \mathbb{S}^2 .*

Proof. For the given ϕ_1 and ϕ_2 , we write

$$\mathcal{L}(\phi_1 - \phi_2) \equiv \mathcal{S}(\phi_1) - \mathcal{S}(\phi_2) = \int_0^1 \partial_\phi \mathcal{S}(t\phi_1 + (1-t)\phi_2)[\phi_1 - \phi_2] dt.$$

Here, \mathcal{L} is viewed as a linear operator acting on the difference $\phi_1 - \phi_2$. The operator $\int_0^1 \partial_\phi \mathcal{S}(t\phi_1 + (1-t)\phi_2) dt$ has a similar structure to the linearized operator $\partial_\phi \mathcal{S}(\phi)$ in

Proposition 5.1. The corresponding Lemmas 5.2-5.4 can be proved in the same manner. By the assumption, we get $\mathcal{L}(\phi_1 - \phi_2) \geq 0$. As in the proof of Lemma 5.3, by Theorem 2.11 in [18], the strong maximum principle holds for the operator \mathcal{L} . We hence obtain the desired result. \square

Remark 5.6. As a consequence, the equation $\mathcal{S}(\phi) = 0$ admits at most one solution ϕ with the property $|\phi|_{C^2(\mathbb{S}^2)} \leq K$.

6. THE EXISTENCE AND PROPERTIES OF SOLUTIONS

In this section, for each small $\underline{u} > 0$ we construct a MOTS along $\underline{H}_{\underline{u}}$, and we further prove that, with \underline{u} as a parameter, these MOTSs form a smooth apparent horizon. Mathematically, this is equivalent to first solving the elliptic equation $\mathcal{S}(\phi, \underline{u}) = 0$ with the operator \mathcal{S} defined in (2.18), and then to proving that the resulted solution is also smooth in terms of \underline{u} .

We first fix some positive constants. Let κ be the maximum of the supersolution $\bar{\phi}$ constructed in Lemma 3.7 and take an arbitrary constant K such that

$$K \geq \max\{C(\kappa), |\bar{\phi}|_{C^2(\mathbb{S}^2)}\},$$

where $C(\kappa)$ is the bound established in Theorem 4.1; namely,

$$C(\kappa) = (Ce^\kappa)^{Ce^{4\Lambda\kappa}}.$$

Then, we choose a sufficiently large depending on κ , according to Lemma 5.3.

We are now ready to prove the existence of MOTSs on each fixed $\underline{H}_{\underline{u}}$.

Theorem 6.1. *Let \underline{u} be a fixed constant in $(0, \delta]$, f be a given smooth function on \mathbb{S}^2 , satisfying (1.13) and (1.14), and $\underline{\phi}$ and $\bar{\phi}$ be as in Lemma 3.3 and Lemma 3.7, respectively. Then, there exists a unique smooth solution $\phi = \phi(\omega, \underline{u})$ of $\mathcal{S}(\phi, \underline{u}) = 0$, satisfying $\underline{\phi} < \phi(\cdot, \underline{u}) < \bar{\phi}$ on \mathbb{S}^2 .*

Proof. We suppress the dependence on \underline{u} in this proof. For any constant $\lambda \in [0, 1]$ and any C^2 -function ϕ on \mathbb{S}^2 , set

$$\mathcal{S}_\lambda(\phi) = \Delta_\gamma \phi + 1 - \frac{1}{2}[\lambda f + (1 - \lambda)]e^\phi + \lambda(I_2 + I_3 + I_4 + I_5),$$

where I_i , for $2 \leq i \leq 5$, are defined in (5.4). We have $\mathcal{S}_1(\phi) = \mathcal{S}(\phi)$ by (5.2). We will prove the existence of the solution to $\mathcal{S}_1(\phi) = 0$ via the method of continuity. For $\lambda = 0$, the operator $\mathcal{S}_0(\phi)$ is reduced to

$$\mathcal{S}_0(\phi) = \Delta_\gamma \phi + 1 - \frac{1}{2}e^\phi.$$

It is obvious that $\phi_0 = \ln 2$ is a solution to $\mathcal{S}_0(\phi) = 0$. For suitably small $\epsilon > 0$ as in (1.14), note that $\underline{\phi} < \phi_0 < \bar{\phi}$ on \mathbb{S}^2 .

Before we proceed to prove the openness and the closedness, we point out that results established in Sections 3-5 for $\mathcal{S}(\phi)$ also hold for $\mathcal{S}_\lambda(\phi)$ with all $\lambda \in [0, 1]$. The proofs are exactly the same with f replaced by $\lambda f + (1 - \lambda)$ and with I_i replaced by λI_i , where

$\lambda \in [0, 1]$ and $2 \leq i \leq 5$. In particular, the estimate in Theorem 4.1 holds for solutions to $\mathcal{S}_\lambda(\phi) = 0$, uniformly for $\lambda \in [0, 1]$.

For the openness, we fix a constant $\beta \in (0, 1)$, say $\beta = 1/3$. For some $\lambda \in [0, 1)$, assume there exists a solution $\phi_\lambda \in C^{2,\beta}(\mathbb{S}^2)$ to $\mathcal{S}_\lambda(\phi_\lambda) = 0$, satisfying $\underline{\phi} < \phi_\lambda < \bar{\phi}$ on \mathbb{S}^2 . By Theorem 4.1, we have $|\phi_\lambda|_{C^2(\mathbb{S}^2)} \leq K$. By Lemma 5.3, the operator $\partial_\phi \mathcal{S}_\lambda(\phi_\lambda) : C^{2,\beta}(\mathbb{S}^2) \rightarrow C^{0,\beta}(\mathbb{S}^2)$ is invertible. By the implicit function theorem, the equation $\mathcal{S}_{\lambda'}(\phi) = 0$ admits a $C^{2,\beta}$ -solution $\phi = \phi_{\lambda'}$ for every λ' close to λ . Moreover, $\underline{\phi} < \phi_{\lambda'} < \bar{\phi}$ on \mathbb{S}^2 , for any λ' even closer to λ .

The closedness follows from the estimate established in Theorem 4.1. Lemma 5.5 guarantees that the strict inequalities $\underline{\phi} < \phi < \bar{\phi}$ hold in the limiting process of λ , under the assumption $\underline{\phi} \leq \phi \leq \bar{\phi}$. The openness and closeness together prove the existence of the solution to $\mathcal{S}_\lambda(\phi) = 0$, satisfying $\underline{\phi} < \phi < \bar{\phi}$ on \mathbb{S}^2 , for any $\lambda \in [0, 1]$, and in particular for $\lambda = 1$.

Last, we prove the uniqueness of the solution to $\mathcal{S}(\phi) = 0$ with ϕ satisfying $\underline{\phi} < \phi < \bar{\phi}$ on \mathbb{S}^2 . Let ϕ be one such solution. By Theorem 4.1, we have $|\phi|_{C^2(\mathbb{S}^2)} \leq K$. As a consequence of Lemma 5.5, such a solution is unique. \square

Next, we view the solution ϕ in Theorem 6.1 as a function of ω and \underline{u} . We now study its regularity in terms of \underline{u} .

Theorem 6.2. *Under the assumptions in Theorem 6.1, for each $\underline{u} \in (0, \delta]$, let $\phi = \phi(\omega, \underline{u})$ be the solution of $\mathcal{S}(\phi, \underline{u}) = 0$ as in Theorem 6.1. Then, the function $\phi = \phi(\omega, \underline{u})$ is smooth in terms of the variables $(\omega, \underline{u}) \in \mathbb{S}^2 \times (0, \delta]$.*

Proof. We restore \underline{u} back to our notations. By Theorem 6.1, there exists a unique solution $\phi = \phi(\omega, \underline{u})$ to $\mathcal{S}(\phi, \underline{u}) = 0$, smooth in ω and satisfying $\underline{\phi} < \phi(\cdot, \underline{u}) < \bar{\phi}$ on \mathbb{S}^2 . By Theorem 4.1, we have $|\phi(\cdot, \underline{u})|_{C^2(\mathbb{S}^2)} \leq K$. We now let \underline{u} as a new parameter and will employ the method of continuity for one more time. Assume $\phi(\omega, \underline{u})$ is a solution to $\mathcal{S}(\phi, \underline{u}) = 0$, we will use the implicit function theorem to obtain a solution $\phi(\omega, \underline{u}')$ to $\mathcal{S}(\phi, \underline{u}') = 0$ with \underline{u}' close to \underline{u} . The *key* is the invertibility of $\partial_\phi \mathcal{S}(\phi, \underline{u}) : C^{2,\beta}(\mathbb{S}^2) \rightarrow C^{0,\beta}(\mathbb{S}^2)$ with any ϕ satisfying $|\phi|_{C^2(\mathbb{S}^2)} \leq K$. And this was *already* proved in Lemma 5.3. Hence, $\phi(\omega, \underline{u})$ is a smooth function of both ω and \underline{u} . \square

In the rest of this section, we show that $\{u = 1 - \underline{u}a e^{-\phi(\omega, \underline{u})}\}$ is a spacelike hypersurface. A crucial step in this part is to derive the equation satisfied by $\partial_{\underline{u}}\phi$ and to obtain an estimate of $\partial_{\underline{u}}\phi$.

Proposition 6.3. *For each $\underline{u} \in (0, \delta]$, let $\phi = \phi(\omega, \underline{u})$ be the solution to $\mathcal{S}(\phi, \underline{u}) = 0$ as in Theorem 6.1. Then, $\partial_{\underline{u}}\phi$ satisfies*

$$(6.1) \quad \Delta_\gamma(\partial_{\underline{u}}\phi) - \frac{1}{2}f(\omega, \underline{u})e^\phi \partial_{\underline{u}}\phi + c^i \nabla_i(\partial_{\underline{u}}\phi) + c \partial_{\underline{u}}\phi = \frac{1}{\underline{u}}h + \frac{1}{2}\partial_{\underline{u}}f e^\phi,$$

where the connection is with respect to the metric γ , and $c = c(\omega, \phi, \underline{u})$, $c^i = c^i(\omega, \phi, \underline{u})$ and $h = h(\omega, \phi, \underline{u})$ are functions satisfying

$$(6.2) \quad |c| + |c^i| + |h| \leq Ca^{-\frac{1}{3}}e^\phi(|\nabla_\gamma^2\phi| + |\nabla_\gamma\phi|^2 + 1).$$

Proof. The proof is similar to that of Proposition 5.1. By $\mathcal{S}(\phi, \underline{u}) = 0$, we have

$$(6.3) \quad \frac{d}{d\underline{u}} \mathcal{S}(\phi, \underline{u}) = 0.$$

We proceed to compute the left-hand side and express it in terms of $\partial_{\underline{u}}\phi$. Throughout the proof, we denote $\dot{\cdot} = \frac{d}{d\underline{u}}$.

Recall the relations given by (2.14) and (2.20), i.e.,

$$g = R^2\gamma, \quad u = 1 - R, \quad R = \underline{u}ae^{-\phi}.$$

Then,

$$(6.4) \quad \dot{u} = -\dot{R}, \quad \dot{R} = R(\underline{u}^{-1} - \dot{\phi}).$$

Note that $\gamma, \underline{\chi}, \chi, \eta, \Omega, \underline{\omega}$ are functions of ω, u, \underline{u} with $u = 1 - R$. We first derive a general formula. Consider a smooth function $\psi = \psi(\omega, u, \underline{u})$ defined along the apparent horizon. Then,

$$\dot{\psi} = \dot{u}\partial_u\psi + \partial_{\underline{u}}\psi = R(\dot{\phi} - \underline{u}^{-1})\partial_u\psi + \partial_{\underline{u}}\psi,$$

and hence

$$(6.5) \quad \dot{\psi} = R\partial_u\psi\dot{\phi} - \underline{u}^{-1}(R\partial_u\psi - \underline{u}\partial_{\underline{u}}\psi).$$

By $g = R^2\gamma$ and $\dot{u} = -\dot{R}$, we get

$$R^2\dot{\gamma}_{kl} = -2R\dot{R}\gamma_{kl} - \dot{R}\partial_u g_{kl} + \partial_{\underline{u}}g_{kl}.$$

Recall (2.6) and (2.7), i.e.,

$$\partial_u g_{kl} = \Omega \text{tr}_g \underline{\chi} g_{kl} + 2\Omega \hat{\underline{\chi}}_{kl}, \quad \partial_{\underline{u}} g_{kl} = \Omega \text{tr}_g \chi g_{kl} + 2\Omega \hat{\chi}_{kl}.$$

Hence,

$$(6.6) \quad \begin{aligned} \dot{\gamma}_{kl} = & \dot{\phi} [(R\Omega \text{tr}_g \underline{\chi} + 2)\gamma_{kl} + 2R^{-1}\Omega \hat{\underline{\chi}}_{kl}] \\ & - \underline{u}^{-1} [(R\Omega \text{tr}_g \chi + 2)\gamma_{kl} + 2R^{-1}\Omega \hat{\chi}_{kl} - \underline{u}\Omega \text{tr}_g \chi \gamma_{kl} - 2\underline{u}R^{-2}\Omega \hat{\chi}_{kl}]. \end{aligned}$$

We also note

$$(6.7) \quad \dot{\gamma}^{ij} = -\gamma^{ik}\gamma^{jl}\dot{\gamma}_{kl}.$$

In (5.2), (5.3), and (5.4), we decompose $\mathcal{S}(\phi, \underline{u})$ as

$$\mathcal{S}(\phi, \underline{u}) = \sum_{l=1}^5 I_l.$$

By (6.3), we have

$$(6.8) \quad \sum_{l=1}^5 \dot{I}_l = 0.$$

In the following, we compute each of \dot{I}_l , for $l = 1, \dots, 5$. We start with \dot{I}_1 . Recall that I_1 is given by (5.3), i.e.,

$$I_1 = \Delta_\gamma \phi + 1 - \frac{1}{2} f e^\phi.$$

Then,

$$(6.9) \quad \dot{I}_1 = (\Delta_\gamma \phi)^\cdot - \frac{1}{2} f e^\phi \dot{\phi} - \frac{1}{2} \dot{f} e^\phi.$$

By Lemma 3.5, we have

$$(6.10) \quad (\Delta_\gamma \phi)^\cdot = \Delta_\gamma \dot{\phi} - \gamma^{ik} \gamma^{jl} \dot{\gamma}_{kl} \nabla_{ij} \phi - \gamma^{ij} \gamma^{kl} \nabla_i \dot{\gamma}_{jl} \nabla_k \phi + \frac{1}{2} \gamma^{ij} \gamma^{kl} \nabla_l \dot{\gamma}_{ij} \nabla_k \phi.$$

By substituting (6.10) in (6.9), we have

$$(6.11) \quad \dot{I}_1 = \Delta_\gamma \dot{\phi} - \frac{1}{2} f e^\phi \dot{\phi} - \frac{1}{2} \partial_{\underline{u}} f e^\phi + c_1^i \nabla_i \dot{\phi} + c_1 \dot{\phi} - \frac{1}{\underline{u}} h_1,$$

where

$$\begin{aligned} c_1^i &= -2R^{-1} \Omega \hat{\chi}_{jl} \gamma^{ij} \gamma^{kl} \nabla_k \phi, \\ c_1 &= -(2 + R \Omega \text{tr}_g \underline{\chi}) \Delta_\gamma \phi - 2\gamma^{ij} \gamma^{kl} [R^{-1} \nabla_i (\Omega \hat{\chi}_{jl}) \nabla_k \phi + R^{-1} \Omega \hat{\chi}_{jl} (\nabla_{ik} \phi + \nabla_i \phi \nabla_k \phi)], \end{aligned}$$

and

$$\begin{aligned} h_1 &= -(2 + R \Omega \text{tr}_g \underline{\chi}) \Delta_\gamma \phi + \underline{u} \Omega \text{tr}_g \chi \Delta_\gamma \phi \\ &\quad - 2\gamma^{ij} \gamma^{kl} [R^{-1} \nabla_i (\Omega \hat{\chi}_{jl}) \nabla_k \phi + R^{-1} \Omega \hat{\chi}_{jl} (\nabla_{ik} \phi + \nabla_i \phi \nabla_k \phi)] \\ &\quad + 2\underline{u} \gamma^{ij} \gamma^{kl} [R^{-2} \nabla_i (\Omega \hat{\chi}_{jl}) \nabla_k \phi + R^{-2} \Omega \hat{\chi}_{jl} (\nabla_{ik} \phi + 2\nabla_i \phi \nabla_k \phi)]. \end{aligned}$$

We point out that c_1^i , c_1 , and h_1 can be viewed as linear combinations of $\nabla_{ij} \phi$, $\nabla_i \phi \nabla_j \phi$, and $\nabla_i \phi$, with coefficients given by γ^{ij} and

$$(6.12) \quad \begin{aligned} &2 + R \Omega \text{tr}_g \underline{\chi}, R^{-1} \Omega \hat{\chi}_{kl}, R^{-1} \nabla_i (\Omega \hat{\chi}_{kl}), \\ &\underline{u} \Omega \text{tr}_g \chi, \underline{u} R^{-2} \Omega \hat{\chi}_{kl}, \underline{u} R^{-2} \nabla_i (\Omega \hat{\chi}_{kl}). \end{aligned}$$

By (A.1), all expressions in (6.12) are controlled by $e^\phi a^{-\frac{1}{3}}$.

Next, we compute $\dot{I}_2, \dots, \dot{I}_5$. Recall that I_2, \dots, I_5 are given by (5.4), i.e.,

$$\begin{aligned} I_2 &= 2R^{-1} \Omega \hat{\chi}_{kl} \gamma^{ik} \gamma^{jl} \nabla_i \phi \nabla_j \phi, & I_3 &= 2\gamma^{ij} \eta_i \nabla_j \phi, \\ I_4 &= R \alpha_1 \gamma^{ij} \nabla_i \phi \nabla_j \phi, & I_5 &= R \alpha_2. \end{aligned}$$

Straightforward computations yield

$$\begin{aligned} \dot{I}_2 &= -2R^{-2} \dot{R} \Omega \hat{\chi}_{kl} \gamma^{ik} \gamma^{jl} \nabla_i \phi \nabla_j \phi + 2R^{-1} (\Omega \hat{\chi}_{kl})^\cdot \gamma^{ik} \gamma^{jl} \nabla_i \phi \nabla_j \phi \\ &\quad + 4R^{-1} \Omega \hat{\chi}_{kl} \dot{\gamma}^{ik} \gamma^{jl} \nabla_i \phi \nabla_j \phi + 4R^{-1} \Omega \hat{\chi}_{kl} \gamma^{ik} \gamma^{jl} \nabla_i \dot{\phi} \nabla_j \phi, \\ \dot{I}_3 &= 2\dot{\gamma}^{ij} \eta_i \nabla_j \phi + 2\gamma^{ij} \dot{\eta}_i \nabla_j \phi + 2\gamma^{ij} \eta_i \nabla_j \dot{\phi}, \end{aligned}$$

and

$$\begin{aligned}\dot{I}_4 &= \dot{R}\alpha_1\gamma^{ij}\nabla_i\phi\nabla_j\phi + R\dot{\alpha}_1\gamma^{ij}\nabla_i\phi\nabla_j\phi \\ &\quad + R\alpha_1\dot{\gamma}^{ij}\nabla_i\phi\nabla_j\phi + 2R\alpha_1\gamma^{ij}\nabla_i\dot{\phi}\nabla_j\phi, \\ \dot{I}_5 &= \dot{R}\alpha_2 + R\dot{\alpha}_2.\end{aligned}$$

In the above expressions, we substitute \dot{R} by (6.4), $\dot{\gamma}^{ij}$ by (6.6) and (6.7), and $\dot{\psi}$ by (6.5) for $\psi = \Omega\hat{\chi}_{kl}$, η_i , α_1 , and α_2 . For each $l = 2, 3, 4, 5$, we can write

$$(6.13) \quad \dot{I}_l = c_l^i\nabla_i\dot{\phi} + c_l\dot{\phi} - \frac{1}{\underline{u}}h_l,$$

where c_l^i, c_l , and h_l can be viewed as linear combinations of $\nabla_{ij}\phi$, $\nabla_i\phi\nabla_j\phi$, and $\nabla_i\phi$, with coefficients given by γ^{ij} and

$$(6.14) \quad \begin{aligned} &R\Omega\text{tr}_g\chi + 2, R^{-1}\Omega\hat{\chi}_{kl}, \eta_i, R\alpha_1, R\alpha_2, \underline{u}\Omega\text{tr}_g\chi, \underline{u}R^{-2}\Omega\hat{\chi}_{kl}, \\ &\partial_u(\Omega\hat{\chi}_{kl}), R\partial_u\eta_i, R^2\partial_u\alpha_1, R^2\partial_u\alpha_2, \\ &\underline{u}R^{-1}\partial_{\underline{u}}(\Omega\hat{\chi}_{kl}), \underline{u}\partial_{\underline{u}}\eta_i, \underline{u}R\partial_{\underline{u}}\alpha_1, \underline{u}R\partial_{\underline{u}}\alpha_2.\end{aligned}$$

By (A.1), (A.2), and (A.3), all expressions in (6.14) are controlled by $e^\phi a^{-\frac{1}{3}}$.

By combining (6.8), (6.11), and (6.13), we then have the desired result. \square

As a consequence, we have the following estimate.

Corollary 6.4. *Under the assumptions in Theorem 6.1, for each $\underline{u} \in (0, \delta]$, let $\phi = \phi(\omega, \underline{u})$ be the solution of $\mathcal{S}(\phi, \underline{u}) = 0$ as in Theorem 6.1. Then,*

$$(6.15) \quad \max_{\mathbb{S}^2} |\partial_{\underline{u}}\phi| \leq C[\underline{u}^{-1}a^{-\frac{1}{3}} \max_{\mathbb{S}^2} e^\phi (|\phi|_{C^2(\mathbb{S}^2)}^2 + 1) + \max_{\mathbb{S}^2} |\partial_{\underline{u}}f e^\phi|],$$

where C is a positive constant independent of a , \underline{u} , and ϕ .

Proof. The left-hand side of (6.1) can be viewed as a linear operator acting on $\partial_{\underline{u}}\phi$. This linear operator has the same structure as $\partial_\phi\mathcal{S}(\phi, \underline{u})$ in (5.1). Applying Lemma 5.4 with $\partial_\phi\mathcal{S}(\phi, \underline{u})$ replaced by the linear operator on the left hand side of (6.1), we then obtain the desired result. We emphasize that C can be chosen independent of a , \underline{u} , and ϕ . \square

Now, we are ready to prove another main result in this paper.

Theorem 6.5. *Under the assumptions in Theorem 6.1, for each $\underline{u} \in (0, \delta]$, let $\phi = \phi(\omega, \underline{u})$ be the solution of $\mathcal{S}(\phi, \underline{u}) = 0$ as in Theorem 6.1. Assume*

$$(6.16) \quad |\underline{u}\partial_{\underline{u}}f(\omega, \underline{u})| \leq a^{-\frac{1}{3}}.$$

Then, the metric g restricted to $\{(\omega, u, \underline{u}); u = 1 - \underline{u}ae^{-\phi(\omega, \underline{u})}\}$ is Riemannian.

Proof. With $R = \underline{u}ae^{-\phi(\omega, \underline{u})}$, we have

$$\partial_{\underline{u}}R = ae^{-\phi(\omega, \underline{u})}(1 - \underline{u}\partial_{\underline{u}}\phi).$$

By (6.15) and (6.16), taking a sufficiently large, we have

$$\max_{\mathbb{S}^2} |\underline{u} \partial_{\underline{u}} \phi| \leq 1/2.$$

Hence, $\partial_{\underline{u}} R \geq ae^{-\phi(\omega, \underline{u})}/2$.

Let g' be the induced metric to the hypersurface $u = 1 - R(\omega, \underline{u})$. Denote $\omega = (\theta_1, \theta_2)$. Then,

$$\begin{aligned} g'_{\theta_i \theta_j} &= g_{\theta_i \theta_j} + \partial_{\theta_i}(1 - R) \partial_{\theta_j}(1 - R) \cdot g(\partial u, \partial u) = g_{\theta_i \theta_j}, \\ g'_{\underline{u} \underline{u}} &= g_{\underline{u} \underline{u}} + 2 \partial_{\underline{u}}(1 - R) \partial_{\underline{u}} \underline{u} \cdot g(\partial u, \partial u) = 4 \partial_{\underline{u}} R, \\ g'_{\theta_i \underline{u}} &= g_{\theta_i \underline{u}} + \partial_{\theta_i}(1 - R) \partial_{\underline{u}} \underline{u} \cdot g(\partial u, \partial u) = 2 \partial_{\theta_i} R, \end{aligned}$$

where we used $g(\partial u, \partial u) = 0$, $g(\partial u, \partial_{\theta_i}) = -2$, and $g(\partial u, \partial_{\theta_i}) = 0$.

Let $X = \lambda_1 \partial_{\theta_1} + \lambda_2 \partial_{\theta_2} + \lambda_3 \partial_{\underline{u}}$ be an arbitrary nonzero tangent vector, for some constants $\lambda_1, \lambda_2, \lambda_3$. Then, we have

$$g'(X, X) = \lambda_i \lambda_j g_{ij} + 4 \lambda_i \lambda_3 \partial_i R + 4 \lambda_3^2 \partial_{\underline{u}} R,$$

where i, j are summed over 1, 2. If $\lambda_3 = 0$, then $g'(X, X) = \lambda_i \lambda_j g_{ij} > 0$. If $\lambda_3 \neq 0$, then by using $g_{ij} = R^2 \gamma_{ij}$ in (2.14), we obtain

$$\begin{aligned} g'(X, X) &\geq \underline{u}^2 a^2 e^{-2\phi} \lambda_i \lambda_j \gamma_{ij} - 4 \underline{u} a e^{-\phi} \lambda_i \lambda_3 \partial_i \phi + 2 \lambda_3^2 a e^{-\phi} \\ &\geq 2 \lambda_3^2 a e^{-\phi} - 4 \lambda_3^2 |\nabla_{\gamma} \phi|^2 > 0, \end{aligned}$$

if a is sufficiently large. □

Therefore, the corresponding apparent horizon is spacelike, and hence a dynamical horizon defined in [8, 9, 10].

7. ANISOTROPIC APPARENT HORIZONS

In previous sections, we constructed a unique MOTS in a certain range along each $\underline{H}_{\underline{u}}$, and gathering all the MOTSs we further proved that it is a smooth apparent horizon (except at the center). In this section, we will construct anisotropic apparent horizons. Specifically, we will construct MOTSs with isolated *valleys*. Via Theorem 6.2, these MOTSs also form a smooth apparent horizon with valleys. We now focus on the operator $\mathcal{S}(\phi, \underline{u})$ defined by (2.18) and construct single-valley and multi-valley solutions, which translates to solutions of $\mathcal{G}(R, \underline{u}) = 0$ with isolated *peaks*.

We allow f depends on \underline{u} , but has to satisfy assumptions imposed earlier on its \underline{u} -derivative in Theorem 1.5. In this section, when there is no danger of confusion, we suppress the dependence of f on \underline{u} . For fixed \underline{u} , we treat $f(\underline{u}, \omega)$ as a function of ω .

We start from constructing solution with one valley.

Theorem 7.1. *Let p be a point on \mathbb{S}^2 , ϵ be a sufficiently small positive constant, and f be a smooth function on \mathbb{S}^2 , such that $0 \leq f \leq 1$ and*

$$f \geq 1/2 \text{ on } B_p(\epsilon/2), \quad f = 0 \text{ on } \mathbb{S}^2 \setminus B_p(\epsilon).$$

Then, for each \underline{u} , the equation $\mathcal{S}(\phi, \underline{u}) = 0$ admits a solution $\phi_\epsilon(\omega, \underline{u})$ satisfying, for any $r_0 \gg \epsilon$,

$$(7.1) \quad \phi_\epsilon(p, \underline{u}) - \inf_{\omega \in \mathbb{S}^2 \setminus B_p(r_0)} \phi_\epsilon(\omega, \underline{u}) \leq \frac{1}{2} \log \epsilon + C,$$

where C is a positive constant depending on r_0 , independent of ϵ and \underline{u} .

Proof. We fix a \underline{u} and consider the operator \mathcal{S} as in (3.1). Note that f satisfies (1.14), with $m = 1/2$. Let $\underline{\phi}$ and $\overline{\phi}$ be the subsolution and the supersolution as in Lemma 3.3 and Lemma 3.7, with $m = 1/2$ and ϵ there replaced by $\epsilon/2$. By (3.9), we have

$$(7.2) \quad \begin{aligned} \overline{\phi}_{\min} &= \overline{\phi}(p) = -2 \log \epsilon + O(1), \\ \overline{\phi}_{\max} &= \overline{\phi}(-p) = -(\tau + 2) \log \epsilon + O(1). \end{aligned}$$

By Theorem 6.1, there exists a unique solution ϕ of $\mathcal{S}(\phi) = 0$, satisfying

$$(7.3) \quad \underline{\phi} < \phi < \overline{\phi}.$$

Recall that $\underline{\phi}$ constructed in Lemma 3.3 is constant. With constant $\underline{\phi}$, the estimate (7.3) is not sufficient to prove the desired result. In the following, we construct a new anisotropic subsolution $\underline{\phi}_*$ to improve the lower bound of ϕ such that $\underline{\phi}_*$, like $\overline{\phi}$, attains its minimum and maximum at two different points p and $-p$, respectively. Moreover, we will show $\overline{\phi}(p)$ is significantly less than $\underline{\phi}_*$ valued away from p .

For the given point $p \in \mathbb{S}^2$, denote by d the distance on \mathbb{S}^2 from p . Then, $d(p) = 0$ and $d(-p) = \pi$. For some positive constant $\mu \in (0.1, 0.9)$ and a small parameter δ to be determined, construct

$$(7.4) \quad \underline{\phi}_* = \mu \log \left[\frac{1 + \delta - \cos d}{\delta} \right].$$

Then,

$$\underline{\phi}_{*\min} = \underline{\phi}_*(p) = 0, \quad \underline{\phi}_{*\max} = \underline{\phi}_*(-p) = -\mu \log \left[\frac{\delta}{2 + \delta} \right] = -\mu \log \delta + O(1).$$

In fact, for any fixed $r_0 \in (0, \pi)$ and any $q \in \mathbb{S}^2 \setminus B_p(r_0)$, we have

$$(7.5) \quad \underline{\phi}_*(q) = -\mu \log \delta + O(1).$$

We claim

$$(7.6) \quad \Delta_{\gamma_0} \underline{\phi}_* + \mu = \frac{\mu \delta (2 + \delta)}{(1 + \delta - \cos d)^2},$$

where Δ_{γ_0} is the Laplacian with respect to the spherical metric γ_0 on \mathbb{S}^2 . We postpone the proof of (7.6) after (7.9) and it is a direct calculation. Then,

$$(\Delta_{\gamma_0} \underline{\phi}_* + \mu) e^{-\underline{\phi}_*} = \frac{1}{2} \underline{f},$$

where

$$(7.7) \quad \underline{f} = \frac{2\mu(2 + \delta)\delta^{1+\mu}}{(1 + \delta - \cos d)^{2+\mu}}.$$

We now claim $\underline{f} > f$ on \mathbb{S}^2 . Thus back to (3.10), with a sufficiently large, we have $\mathcal{S}(\underline{\phi}) > 0$, since $\mu < 0.9$. To prove the claim, we only need to prove $\underline{f} > 1$ on $B_p(\epsilon)$, since $f \leq 1$ on $B_p(\epsilon)$ and $f = 0$ outside $B_p(\epsilon)$. By $\cos d \geq 1 - d^2/2$ and (7.7), we have

$$\underline{f} \geq 2\mu(2 + \delta) \cdot \frac{\delta^{1+\mu}}{(\delta + \epsilon^2/2)^{2+\mu}} \quad \text{on } B_p(\epsilon).$$

For some $\beta > 0$ to be determined, set $\delta = \epsilon^\beta$. Then,

$$\underline{f} \geq 2\mu(2 + \epsilon^\beta) \cdot \frac{\epsilon^{\beta(1+\mu)}}{(\epsilon^\beta + \epsilon^2/2)^{2+\mu}} \quad \text{on } B_p(\epsilon).$$

If $\beta > 2$, then

$$\underline{f} \geq \mu 2^{4+\mu} \cdot \frac{1 + \epsilon^\beta/2}{(1 + 2\epsilon^{\beta-2})^{2+\mu}} \cdot \epsilon^{\beta(1+\mu)-2(2+\mu)} \quad \text{on } B_p(\epsilon).$$

Thus, $\underline{f} \geq \mu 2^3$ on $B_p(\epsilon)$ if $\beta(1 + \mu) = 2(2 + \mu)$ and ϵ is small. In the following, we take

$$(7.8) \quad \beta = \frac{2(2 + \mu)}{1 + \mu}.$$

Then, (7.5) reduces to, for any $q \in \mathbb{S}^2 \setminus B_p(r_0)$,

$$(7.9) \quad \underline{\phi}_*(q) = -\mu\beta \log \epsilon + O(1).$$

To prove (7.6), we position p at the north pole and denote by (θ, ψ) the spherical coordinates on \mathbb{S}^2 , with $\theta \in [0, \pi]$ and $\psi \in [0, 2\pi)$. Then, $d = \theta$. For a function $\phi = \phi(\theta)$, we have

$$\Delta_0 \phi(\theta) = \frac{1}{\sin \theta} \partial_\theta (\sin \theta \partial_\theta \phi(\theta)).$$

Then, a straightforward computation yields (7.6).

We now compare $\underline{\phi}_*$ and $\bar{\phi}$. By (7.2) and (7.9), we have for any $q \in \mathbb{S}^2 \setminus B_p(r_0)$

$$\bar{\phi}(p) - \underline{\phi}_*(q) \leq (\mu\beta - 2) \log \epsilon + C,$$

where we replaced $O(1)$ by C , a positive constant, independent of ϵ and \underline{u} . With β as in (7.8), we can take μ such that $\mu\beta > 2$, say $\mu\beta = 2.5$. This yields $2\mu(2 + \mu)/(1 + \mu) = 2.5$, and we have $\mu = (\sqrt{89} - 3)/8 \approx 0.80$.

By $\mathcal{S}(\phi) = 0$, $\mathcal{S}(\underline{\phi}_*) > 0$, and Lemma 5.5, we have $\underline{\phi}_* < \phi$. Here, in order to apply Lemma 5.5, we may need to enlarge K so that $|\underline{\phi}_*|_{C^2(\mathbb{S}^2)} \leq K$, if necessary. Then, for any $q \in \mathbb{S}^2 \setminus B_p(r_0)$, we obtain

$$\phi(p) - \phi(q) \leq \bar{\phi}(p) - \underline{\phi}_*(q) \leq \frac{1}{2} \log \epsilon + C.$$

We now restore ϵ and \underline{u} , and write $\phi_\epsilon(\cdot, \underline{u})$ for ϕ . Then $\phi_\epsilon(\omega, \underline{u})$ is our desired solution. \square

In Theorem 7.1, the solution ϕ has a value at p significantly less than the values away from p . In this sense, we say that ϕ has a valley at p .

Theorem 7.1 can be reformulated easily with the operator $\mathcal{G}(R, \underline{u})$ defined by (2.3).

Proposition 7.2. *Let p be a point on \mathbb{S}^2 , ϵ be a sufficiently small positive constant, and f be a smooth function on \mathbb{S}^2 , such that $0 \leq f \leq 1$ and*

$$f \geq 1/2 \text{ on } B_p(\epsilon/2), \quad f = 0 \text{ on } \mathbb{S}^2 \setminus B_p(\epsilon).$$

Then, for each \underline{u} , the equation $\mathcal{G}(R, \underline{u}) = 0$ admits a unique solution $R_\epsilon(\omega, \underline{u})$ satisfying

$$(7.10) \quad \sup_{\omega \in \mathbb{S}^2 \setminus B_p(r_0)} R_\epsilon(\omega, \underline{u}) \leq C\epsilon^{\frac{1}{2}} R_\epsilon(p, \underline{u}),$$

where C is a positive constant, independent of ϵ and \underline{u} .

As a consequence, the corresponding MOTS and apparent horizon are *anisotropic*.

In the rest of the section, we construct solutions with multiple valleys. We first modify the arguments in the proof of Lemma 3.7 to construct a supersolution with multiple valleys.

Lemma 7.3. *Let p_1, \dots, p_n be fixed n points on \mathbb{S}^2 , ϵ be a positive constant much less than the distance between any two distinct p_i and p_j for $i, j = 1, \dots, n$, and f be a nonnegative continuous function on \mathbb{S}^2 such that $f \geq m$ on $\cup_{i=1}^n B_{p_i}(\epsilon)$, for some constant $m > 0$. Then, for any $\tau > 2.2/n$, there exists $\bar{\phi} \in C^\infty(\mathbb{S}^2)$ such that $\mathcal{S}(\bar{\phi}) < 0$ and, for any $i = 1, \dots, n$,*

$$(7.11) \quad \begin{aligned} \bar{\phi}(p_i) &= -\log(m\epsilon^2) + C, \\ \bar{\phi}_{\max} &= -\log(m\epsilon^{\tau+2}) + C, \end{aligned}$$

for some positive constant C depending only on n and τ .

Proof. For each $i = 1, \dots, n$, denote by d_i the distance on \mathbb{S}^2 from p_i and take a cutoff function

$$\chi_{i,\epsilon} = \begin{cases} 0 & \text{on } B_{p_i}(\epsilon/2), \\ 1 & \text{on } \mathbb{S}^2 \setminus B_{p_i}(\epsilon). \end{cases}$$

Set

$$\tilde{\phi}_{i,\epsilon} = \tau \left[\chi_{i,\epsilon} \log \sin \frac{d_i}{2} + (1 - \chi_{i,\epsilon}) \log \epsilon \right].$$

Then, we have

$$\tilde{\phi}_{i,\epsilon} = \begin{cases} \tau \log \epsilon & \text{on } B_{p_i}(\epsilon/2), \\ \tau \log \epsilon + O(1) & \text{on } B_{p_i}(\epsilon) \setminus B_{p_i}(\epsilon/2), \\ \tau \log d_i + O(1) & \text{on } \mathbb{S}^2 \setminus B_{p_i}(\epsilon), \end{cases}$$

and

$$|\nabla_0 \tilde{\phi}_{i,\epsilon}| = O(\epsilon^{-1}), \quad |\nabla_0^2 \tilde{\phi}_{i,\epsilon}| = O(\epsilon^{-2}) \quad \text{on } \mathbb{S}^2 \setminus B_{p_i}(\epsilon/2).$$

Moreover, it holds

$$\Delta_0 \tilde{\phi}_{i,\epsilon} = -\frac{\tau}{2} \quad \text{on } \mathbb{S}^2 \setminus B_{p_i}(\epsilon).$$

Set

$$\tilde{\phi}_\epsilon = \sum_{i=1}^n \tilde{\phi}_{i,\epsilon}.$$

Then, it follows

$$\Delta_0 \tilde{\phi}_\epsilon + 1.1 = -\frac{n\tau}{2} + 1.1 < 0 \quad \text{on } \mathbb{S}^2 \setminus \cup_{i=1}^n B_{p_i}(\epsilon),$$

as long as $\tau > 2.2/n$. Next, for each $i = 1, \dots, n$, we have

$$e^{\tilde{\phi}_{i,\epsilon}} \geq \epsilon^\tau / C_1, \quad e^{\tilde{\phi}_{j,\epsilon}} \geq 1/C_1 \quad (j \neq i) \quad \text{on } B_{p_i}(\epsilon),$$

for some positive constant C_1 . Hence,

$$e^{\tilde{\phi}_\epsilon} \geq \epsilon^\tau / C_1^n \quad \text{on } \cup_{i=1}^n B_{p_i}(\epsilon).$$

We can now follow the same argument as in the proof of Lemma 3.7. Set

$$\phi_\epsilon = \tilde{\phi}_\epsilon - \log s_\epsilon,$$

where s_ϵ is a positive constant given by

$$s_\epsilon = \frac{m\epsilon^{\tau+2}}{2C_0C_1^n}.$$

We then have the desired result. \square

Theorem 7.4. *Let p_1, \dots, p_n be fixed n points on \mathbb{S}^2 , ϵ be a positive constant much less than the distance between any two distinct p_i and p_j for $i, j = 1, \dots, n$, and f be a smooth function on \mathbb{S}^2 , such that $0 \leq f \leq \nu(n)$ and*

$$f \geq \nu(n)/2 \quad \text{on } \cup_{i=1}^n B_{p_i}(\epsilon/2), \quad f = 0 \quad \text{on } \mathbb{S}^2 \setminus \cup_{i=1}^n B_{p_i}(\epsilon),$$

for some constructed constant $\nu(n) \in (0, 1)$ depending only on n . Then for each \underline{u} , the equation $\mathcal{S}(\phi, \underline{u}) = 0$ admits a solution $\phi_\epsilon(\omega, \underline{u})$ satisfying, for any $r_0 \gg \epsilon$ and any $i = 1, \dots, n$,

$$(7.12) \quad \phi_\epsilon(p_i, \underline{u}) - \inf_{\omega \in \mathbb{S}^2 \setminus \cup_{i=1}^n B_{p_i}(r_0)} \phi_\epsilon(\omega, \underline{u}) \leq \frac{1}{2n} \log \epsilon + C,$$

where C is a positive constant, independent of ϵ and \underline{u} .

This is simply Theorem 1.6.

Proof. We follow the proof of Theorem 7.1. We fix a \underline{u} and consider the operator \mathcal{S} as in (3.1). We take the subsolution $\underline{\phi}$ as in Lemma 3.3 and the supersolution $\overline{\phi}$ as in Lemma 7.3, with $m = \nu(n)/2$ and ϵ there replaced by $\epsilon/2$. By (7.11), we have, for each $i = 1, \dots, n$

$$(7.13) \quad \overline{\phi}(p_i) = -2 \log \epsilon + O(1).$$

By Theorem 6.1, there exists a unique solution ϕ of $\mathcal{S}(\phi) = 0$, satisfying

$$(7.14) \quad \underline{\phi} < \phi < \overline{\phi}.$$

We now construct an anisotropic subsolution $\underline{\phi}_*$ so that the value of $\overline{\phi}$ at each p_i is significantly less than the value of $\underline{\phi}_*$ at any point away from all p_i . For each $i = 1, \dots, n$, denote by d_i the distance on \mathbb{S}^2 from p_i . Then, $d_i(p_i) = 0$.

For some positive constant $\mu \in (0, 1)$ and a small positive constant δ to be determined, construct

$$(7.15) \quad \underline{\phi}_* = \mu \sum_{i=1}^n \log \left[\frac{1 + \delta - \cos d_i}{\delta} \right].$$

Then, for any fixed $r_0 \in (0, \pi)$ and any $q \in \mathbb{S}^2 \setminus \cup_{i=1}^n B_{p_i}(r_0)$, we have

$$(7.16) \quad \underline{\phi}_*(q) = -n\mu \log \delta + O(1).$$

Moreover,

$$\Delta_0 \underline{\phi}_* + n\mu = \sum_{i=1}^n \frac{\mu \delta (2 + \delta)}{(1 + \delta - \cos d_i)^2}.$$

Then, it holds

$$(\Delta_0 \underline{\phi}_* + n\mu) e^{-\underline{\phi}_*} = \frac{1}{2} \underline{f},$$

where

$$(7.17) \quad \underline{f} = 2 \left[\sum_{i=1}^n \frac{\mu \delta (2 + \delta)}{(1 + \delta - \cos d_i)^2} \right] \prod_{i=1}^n \left[\frac{\delta}{1 + \delta - \cos d_i} \right]^\mu.$$

In the following, we will take μ such that $n\mu < 1$.

We now claim $\underline{f} > f$ on \mathbb{S}^2 . Thus by (3.10), we have $\mathcal{S}(\underline{\phi}_*) > 0$, since $n\mu < 1$. To prove the claim, we fix an $i = 1, \dots, n$ and prove $\underline{f} > \nu(n)$ on $B_{p_i}(\epsilon)$, since $f \leq \nu(n)$ on $B_{p_i}(\epsilon)$ and $f = 0$ outside $\cup_{i=1}^n B_{p_i}(\epsilon)$. In the summation in the expression (7.17), we keep the term involving d_i and drop other terms; while in the product, we single out the factor involving d_i . Then,

$$\underline{f} \geq 2\mu(2 + \delta) \cdot \frac{\delta^{1+\mu}}{(1 + \delta - \cos d_i)^{2+\mu}} \prod_{j \neq i} \left[\frac{\delta}{1 + \delta - \cos d_j} \right]^\mu.$$

Note that there are $n - 1$ factors in the product. By $\cos d_i \geq 1 - d_i^2/2$ and $\cos d_j \geq -1$ for $j \neq i$, we have

$$\underline{f} \geq \frac{2\mu(2 + \delta)}{(2 + \delta)^{\mu(n-1)}} \cdot \frac{\delta^{1+n\mu}}{(\delta + d_i^2/2)^{2+\mu}}.$$

For some $\beta > 0$ to be determined, we set $\delta = \epsilon^\beta$. Then,

$$\underline{f} \geq 2\mu(2 + \epsilon^\beta)^{1-\mu(n-1)} \cdot \frac{\epsilon^{\beta(1+n\mu)}}{(\epsilon^\beta + \epsilon^2/2)^{2+\mu}} \quad \text{on } B_{p_i}(\epsilon).$$

If $\beta > 2$, we write

$$\epsilon^\beta + \epsilon^2/2 = \frac{1}{2} \epsilon^2 (1 + 2\epsilon^{\beta-2}),$$

and we hence have

$$\underline{f} \geq \mu 2^{4+2\mu-n\mu} \cdot \frac{(1 + \epsilon^\beta/2)^{1-\mu(n-1)}}{(1 + 2\epsilon^{\beta-2})^{2+\mu}} \cdot \epsilon^{\beta(1+n\mu)-2(2+\mu)} \quad \text{on } B_{p_i}(\epsilon).$$

We can take

$$(7.18) \quad \nu(n) = \mu 2^{1+2\mu-n\mu}.$$

Thus, $\underline{f} > \nu(n)$ on $B_{p_i}(\epsilon)$ if $\beta(1 + n\mu) = 2(2 + \mu)$ and ϵ is small. In the following, we take

$$(7.19) \quad \beta = \frac{2(2 + \mu)}{1 + n\mu}.$$

Then, (7.16) reduces to, for any $q \in \mathbb{S}^2 \setminus \cup_{i=1}^n B_{p_i}(r_0)$,

$$(7.20) \quad \underline{\phi}_*(q) = -n\mu\beta \log \epsilon + O(1).$$

At the moment, $\nu(n)$ also depends on μ . Later on, we will fix μ depending only on n .

We now compare $\underline{\phi}_*$ and $\bar{\phi}$. By (7.13) and (7.20), we have, for any $i = 1, \dots, n$ and any $q \in \mathbb{S}^2 \setminus \cup_{i=1}^n B_{p_i}(r_0)$,

$$\bar{\phi}(p_i) - \underline{\phi}_*(q) \leq (n\mu\beta - 2) \log \epsilon + C,$$

where we replaced $O(1)$ by C , a positive constant independent of ϵ . Note that the maximal μ is given by $1/n$. With β as in (7.19), if $\mu = 1/n$, then $n\mu\beta = 2 + 1/n$. Take any $\gamma \in (0, 1/n)$, say $\gamma = 1/(2n)$, and solve the algebraic equation $n\mu\beta = 2 + \gamma$, i.e.,

$$2n\mu^2 + (2 - \gamma)n\mu - (2 + \gamma) = 0.$$

Its positive solution is given by

$$\mu = \frac{1}{4n} \left[\sqrt{(2 - \gamma)^2 n^2 + 8n(2 + \gamma)} - (2 - \gamma)n \right].$$

We take such μ and fix $\nu(n)$ accordingly. For n large, we note that

$$\frac{1}{n} - \frac{1}{n^2} + o\left(\frac{1}{n^2}\right) < \mu < \frac{1}{n},$$

where the expression on the left-hand side corresponds to the solution for $\gamma = 0$.

By $\mathcal{S}(\phi) = 0$, $\mathcal{S}(\underline{\phi}_*) > 0$, and Lemma 5.5, we have $\underline{\phi}_* < \phi$. Here, in order to apply Lemma 5.5, we may need to enlarge K so that $|\underline{\phi}_*|_{C^2(\mathbb{S}^2)} \leq K$, if necessary. Hence, for any $i = 1, \dots, n$ and $q \in \mathbb{S}^2 \setminus \cup_{i=1}^n B_{p_i}(r_0)$, we prove have

$$\phi(p_i) - \phi(q) \leq \bar{\phi}(p_i) - \underline{\phi}_*(q) \leq \gamma \log \epsilon + C.$$

We then restore ϵ and \underline{u} , and write $\phi_\epsilon(\cdot, \underline{u})$ for ϕ . This $\phi_\epsilon(\omega, \underline{u})$ is our desired multi-valley solution. \square

APPENDIX A. ESTIMATES OF GEOMETRIC QUANTITIES

In this appendix, we collect useful estimates for geometric quantities $\Omega, \chi, \underline{\chi}, \eta, \underline{\omega}$ from [4], which play important roles in this paper. All these quantities are defined and pointwisely estimated as listed below at $(\omega, u, \underline{u})$ with $u = 1 - R$ and $R = \underline{u}ae^{-\phi}$. Recall $g = R^2\gamma$ in (2.14), Ω defined in (2.1), and $\chi, \underline{\chi}, \eta, \underline{\omega}$ defined in (2.2). Moreover, α_1, α_2 are defined in (2.4), i.e.,

$$\begin{aligned} \alpha_1 &= -\left(\frac{1}{2}\Omega \text{tr}_g \underline{\chi} + \frac{1}{R} + 4\Omega \underline{\omega}\right), \\ \alpha_2 &= \frac{1}{2}\Omega^{-1} \text{tr}_g \chi - \frac{1}{R} + \frac{\underline{u}a}{2R^2} f(\omega, \underline{u}). \end{aligned}$$

With C as a uniform constant, we have the following estimates.

- Estimates of geometric quantities and their angular derivatives:

$$(A.1) \quad \begin{aligned} R^2|\hat{\chi}_{ij}|_g + R^2|\eta_i|_g + R|\Omega - 1| &\leq Ce^\phi a^{-\frac{1}{3}}, \\ |R\Omega\text{tr}_g\chi + 2| + |\nabla_\gamma(R\Omega\text{tr}_g\chi)|_\gamma + R^{-1}|\Omega\hat{\chi}_{kl}|_\gamma + R^{-1}|\nabla_\gamma(\Omega\hat{\chi}_{kl})|_\gamma &\leq Ce^\phi a^{-\frac{1}{3}}, \\ \underline{u}|\Omega\text{tr}_g\chi| + \underline{u}|\nabla_\gamma(\Omega\text{tr}_g\chi)|_\gamma + \underline{u}R^{-2}|\Omega\hat{\chi}_{kl}|_\gamma + \underline{u}R^{-2}|\nabla_\gamma(\Omega\hat{\chi}_{kl})|_\gamma &\leq Ce^\phi a^{-\frac{1}{2}}, \\ |\eta_i|_\gamma + |\nabla_\gamma\eta_i|_\gamma + R|\alpha_1| + R|\nabla_\gamma\alpha_1|_\gamma + R|\alpha_2| + R|\nabla_\gamma\alpha_2|_\gamma &\leq Ce^\phi a^{-\frac{1}{3}}. \end{aligned}$$

These estimates can be read from norms defined in Section 2.6 and the statements of Theorem 3.1 and Proposition 5.1 in [4].

- Estimates of u -derivatives:

$$(A.2) \quad |\partial_u(\Omega\hat{\chi}_{kl})|_\gamma + R|\partial_u\eta_i|_\gamma + R^2|\partial_u\alpha_1| + R^2|\partial_u\alpha_2| \leq Ce^\phi a^{-\frac{1}{3}}.$$

We can derive the estimates for $\partial_u(\Omega\hat{\chi}_{kl})$, $\partial_u\eta_i$, $\partial_u\alpha_1$ via using equations in (2.4) and the statement of Theorem 3.1 in [4]. For the estimate for $\partial_u\alpha_2$, we apply (4.9) in [3] and get

$$R^2|\partial_u\alpha_2| = \left| \frac{1}{2}R^2\partial_u(\Omega^{-1}\text{tr}_g\chi) - 1 + R^{-1}\underline{u}af(\omega, \underline{u}) \right| \leq Ce^\phi a^{-\frac{1}{3}}.$$

- Estimates of \underline{u} -derivatives:

$$(A.3) \quad \underline{u}R^{-1}|\partial_{\underline{u}}(\Omega\hat{\chi}_{kl})|_\gamma + \underline{u}|\partial_{\underline{u}}\eta_i|_\gamma + \underline{u}R|\partial_{\underline{u}}\alpha_1| + \underline{u}R|\partial_{\underline{u}}\alpha_2| \leq Ce^\phi a^{-\frac{1}{3}}.$$

The estimates for $\partial_{\underline{u}}(\Omega\hat{\chi}_{kl})$, $\partial_{\underline{u}}\eta_i$, $\partial_{\underline{u}}\alpha_1$ can be derived via using equations in (2.4) and the statement of Theorem 3.1 in [4]. To obtain the estimate for $\partial_{\underline{u}}\alpha_2$, recall the definition of f in the statement of Theorem 1.5, i.e.,

$$\underline{u}af(\omega, \underline{u}) = \int_0^{\underline{u}} |\hat{\chi}_0|_g^2(\omega, \underline{u}') d\underline{u}'.$$

Hence, with $R = 1 - u$ and $\partial_{\underline{u}}u = 0$, it deduces

$$\underline{u}R\partial_{\underline{u}}\alpha_2 = \frac{\underline{u}R}{2}\partial_{\underline{u}}(\Omega^{-1}\text{tr}_g\chi) + \frac{\underline{u}}{2R}|\hat{\chi}_0|_g^2(\omega, \underline{u}).$$

Using the first and the last equations of (2.4) and the estimates ensured by Theorem 3.1 in [4], we get

$$\begin{aligned} |\partial_{\underline{u}}(\Omega^{-1}\text{tr}_g\chi)(\omega, 1 - R, \underline{u}) + |\hat{\chi}|_g^2(\omega, 1 - R, \underline{u})| &\leq \frac{Ca}{R^2}a^{-\frac{1}{3}}, \\ | - |\hat{\chi}|_g^2(\omega, 1 - R, \underline{u}) + R^{-2}|\hat{\chi}_0|_g^2(\omega, \underline{u})| &\leq \frac{Ca}{R^2}a^{-\frac{1}{3}}. \end{aligned}$$

Via the triangle inequality, it follows

$$|\partial_{\underline{u}}(\Omega^{-1}\text{tr}_g\chi)(\omega, 1 - R, \underline{u}) + R^{-2}|\hat{\chi}_0|_g^2(\omega, \underline{u})| \leq \frac{Ca}{R^2}a^{-\frac{1}{3}}.$$

Multiplying both sides of the above inequality by $\underline{u}R/2$, we hence obtain

$$\underline{u}R|\partial_{\underline{u}}\alpha_2| = \left| \frac{\underline{u}R}{2}\partial_{\underline{u}}(\Omega^{-1}\text{tr}_g\chi) + \frac{\underline{u}}{2R}|\hat{\chi}_0|_g^2(\omega, \underline{u}) \right| \leq \frac{C\underline{u}a}{R^2}a^{-\frac{1}{3}} = Ce^\phi a^{-\frac{1}{3}}.$$

REFERENCES

- [1] X. An, *Formation of trapped surfaces from past null infinity*, preprint (2012), arXiv:1207.5271.
- [2] X. An, *A scale-critical trapped surface formation criterion: a new proof via signature for decay rates*, preprint (2019), arXiv:1903.02967.
- [3] X. An, *Emergence of Apparent Horizon in Gravitational Collapse*, Ann. PDE 6 (2020), 89 pages. <https://doi.org/10.1007/s40818-020-00085-9>
- [4] X. An, J. Luk, *Trapped surfaces in vacuum arising dynamically from mild incoming radiation*, Adv. Theor. Math. Phys. 21 (2017), no.1, 1-120.
- [5] L. Andersson, M. Eichmair, J. Metzger, *Jang's equation and its applications to marginally trapped surfaces*, Complex analysis and dynamical systems IV. Part 2, 13-45, Contemp. Math., 554, Israel Math. Conf. Proc., Amer. Math. Soc., Providence, RI, 2011.
- [6] L. Andersson, M. Mars, W. Simon, *Local existence of dynamical and trapping horizons*, Phys. Rev. Lett. 95 (2005), 111102.
- [7] L. Andersson, J. Metzger, *The area of horizons and the trapped region*, Comm. Math. Phys. 290 (2009), no.3, 941-972.
- [8] A. Ashtekar, G. Galloway, *Some uniqueness results for dynamical horizons*, Adv. Theor. Math. Phys. 9 (2005), no.1, 1-30.
- [9] A. Ashtekar, B. Krishnan, *Dynamical horizons and their properties*, Phys. Rev. D. 68 (2003), 104030.
- [10] A. Ashtekar, B. Krishnan, *Isolated and dynamical horizons and their applications*, Living Rev. Relativ. 7 (2004), 10.
- [11] D. Christodoulou, *The formation of black holes in general relativity*, Monographs in Mathematics, European Mathematical Soc. (2009).
- [12] D. Christodoulou, S. Klainerman, *The global nonlinear stability of the Minkowski space*, Princeton mathematical series 41 (1993).
- [13] M. Dafermos, *The formation of black holes in general relativity*, Astérisque 352 (2013), 243-313.
- [14] M. Eichmair, *The plateau problem for marginally trapped surfaces*, J. Differential Geom. 83 (2009), no. 3, 551-583.
- [15] M. Eichmair, *Existence, regularity, and properties of generalized apparent horizons*, Comm. Math. Phys. 294 (2010), no.3, 745-760.
- [16] G. Galloway, R. Schoen, *A generalization of Hawking's black hole topology theorem to higher dimensions*, Comm. Math. Phys. 266 (2006), no. 2, 571-576.
- [17] D. Gilbarg, N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order* Grundlehren, Vol. 224, Springer-Verlag, Berlin, (1983).
- [18] Q. Han, F-H. Lin, *Elliptic partial differential equations (2nd ed.)*, Courant lecture notes, Vol. 1, (2011).
- [19] S. Klainerman, J. Luk, I. Rodnianski, *A fully anisotropic mechanism for formation of trapped surfaces in vacuum*, Invent. Math. 198 (2014), no. 1, 1-26.
- [20] S. Klainerman, I. Rodnianski, *On emerging scarred surfaces for the Einstein vacuum equations*, Discrete Contin. Dyn. Syst. 28 (2010), no. 3, 1007-1031.
- [21] S. Klainerman, I. Rodnianski, *On the the formation of trapped surfaces*, Acta Math. 208 (2012), no. 2, 211-333.
- [22] P. Le, *The intersection of a hyperplane with a lightcone in the Minkowski spacetime*, J. Differential Geom. 109 (2018), no. 3, 497-507.
- [23] J. Luk, I. Rodnianski, *Nonlinear interactions of impulsive gravitational waves for the vacuum Einstein equations*, Camb. J. Math. 5 (2017), no. 4, 435-570.
- [24] R. Penrose, *Gravitational collapse and space-time singularities*, Phys. Rev. Lett. 14 (1965), 57-59.
- [25] R. Schoen, Talk given at the Miami Waves conference, January 2004.
- [26] R. Schoen, S. T. Yau, *The existence of a black hole due to condensation of matter*, Comm. Math. Phys. 90 (1983), no.4, 575-579.

- [27] S. T. Yau, *Geometry of three manifolds and existence of black hole due to boundary effect*, Adv. Theor. Math. Phys. 5 (2001), no. 4, 755-767.

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, 119076, SINGAPORE
Email address: `matax@nus.edu.sg`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, NOTRE DAME, IN 46556, USA
Email address: `qhan@nd.edu`