arXiv:2010.12646v2 [math.AG] 10 Nov 2020

CURVATURE GRAFTED BY INSTANTONS


ELIZABETH GASPARIM AND BRUNO SUZUKI

In memory of Pushan Majumdar.

ABSTRACT. We show that an instanton with high charge can provoke the creation of extra curvature on the space that holds it. Geometrically, this corresponds to a new surgery operation, which we name grafting. Curvature around a sphere increases by grafting when the charge of an instanton decays.

1. PARTICLE DECAY AND LOCAL CHARACTERISTIC CLASSES


The concept of local characteristic classes has its motivation in the description of particle decay phenomena. Recent news on particle decay have brought to light the need to describe a general theory of local invariants. The new observation of decay was presented at the 40th International Conference on High Energy Physics. Scientists at CERN have reported on their first significant evidence for a process predicted by theory, paving the way for searches for evidence of new physics in particle processes that could explain dark matter and other mysteries of the universe. CERN NA62 presented the first significant experimental evidence for the ultra-rare decay of the charged kaon into a charged pion and two neutrinos.



Here we will discuss the concepts of local characteristic classes, in particular Chern class and local holomorphic Euler characteristic, and how these occur within the mathematical descriptions of particle decay. We will focus in 4 dimensions, recalling results obtained in joint work with Pushan Majumdar about instanton decay, and we will propose generalizations to higher dimensions. We are especially interested in the loss of charge that can be

B.S. acknowledges support of ANID 2019/13204-0, E.G. acknowledges support of VRIDT-UCN, Chile.

provoked by the contraction of a 2-sphere. The local Chern classes measure local manifestations of curvature concentrated around points or around subvarieties.



2. Instantons

We first review the basic mathematical setup of instantons. The original definition of instanton says it is a connection minimizing the Yang–Mills functional. Let us make this concept mathematically precise.

We start up by recalling the concept of connection on a principal bundle. Let P be a principal G-bundle over a smooth manifold M. A *connection* on P is a differential 1-form ω on P with values in the Lie algebra satisfying:

- $\mathrm{Ad}_g\, R_g^*\omega = \omega$ (G-equivariance),
- $\omega(X_\xi) = \xi$ (Fundamental vector fields),

where R_g denotes the right translation by g, upper $*$ the pull back of forms, and X_ξ denotes the fundamental vector field associated to ξ.

Let F_A denote the curvature of the connection A, and assume M has real dimension 4. An *instanton* on M is a connection minimizing the *Yang–Mills Functional*:

$$YM(A) = \int |F_A|^2 \mathrm{dvol}.$$

Note that the functional applies to 4 dimensions, since the integrand is a 4-form. The *Yang–Mills equations* are the Euler–Lagrange equations corresponding to the Yang–Mills Functional. Hence, an instanton is a solution of the YM equations. These are nonlinear partial differential equations, which are well known to be very difficult to solve.

One of the few cases where an explicit solution is written in terms of connections is the basic $SU(2)$ instanton on S^4:

$$A = \frac{1}{1+|x|^2}(\theta_1\mathbf{i} + \theta_2\mathbf{j} + \theta_3\mathbf{k})$$

where $\mathbf{i}, \mathbf{j}, \mathbf{k}$ is a basis for $\mathfrak{sl}(2)$ and

$$\begin{aligned}
\theta_1 &= x_1\mathrm{d}x_2 - x_2\mathrm{d}x_1 - x_3\mathrm{d}x_4 + x_4\mathrm{d}x_3,\\
\theta_2 &= x_1\mathrm{d}x_3 - x_3\mathrm{d}x_1 - x_4\mathrm{d}x_2 + x_2\mathrm{d}x_4,\\
\theta_3 &= x_1\mathrm{d}x_4 - x_4\mathrm{d}x_1 - x_2\mathrm{d}x_3 + x_3\mathrm{d}x_2.
\end{aligned}$$

However, for more general manifolds explicit connections solving the YM equations are not known. A simpler way to look for instantons is to consider the linearized version of the YM equations, namely:

$$\star F = -F,$$

where $\star$ is the Hodge operator. Connections whose curvatures satisfy this equation are called *anti-self-dual*. These ASD connections are solutions of the YM equations. Using ASD connections, alternative approaches to find instantons appeared. A celebrated construction of instantons in Euclidean space was obtained by Atiyah–Drinfeld–Hitchin–Manin in [ADHM] in terms of the following linear data:

- complex vector spaces V and W of dimension k and N,
- $k \times k$ complex matrices B_1, B_2,
- a $k \times N$ complex matrix I,
- a $N \times k$ complex matrix J,
- a real moment map $\mu_r = [B_1, B_1^\dagger] + [B_2, B_2^\dagger] + II^\dagger - J^\dagger J$,
- a complex moment map $\mu_c = [B_1, B_2] + IJ$.

The ADHM construction was explored and generalized in a variety of ways by several authors. Subsequently, this approach using complex vector spaces was significantly improved as to include a far wider variety of instantons by using holomorphic vector bundles, giving rise to a very general principle called the *Kobayashi–Hitchin correspondence*, see [LT], which provides an identification:

$$\text{ASD SU}(r) \text{ connections} \leftrightarrow \text{rank } r \text{ holomorphic bundles.}$$

The correspondence was proved by Donaldson [D] and Uhlenbeck–Yau [UY]. We will concern ourselves with the case of SU(2) instantons:

$$\left\{ \begin{array}{c} \text{irreducible SU(2)-instantons} \\ \text{of charge } n \end{array} \right\} \leftrightarrow \left\{ \begin{array}{c} \text{stable } \mathrm{SL}(2,\mathbb{C})\text{-bundles} \\ \text{with } c_2 = n \end{array} \right\}$$

If we represent the connection in terms of a covariant derivative on a vector bundle, then the association is:

$$\nabla = \partial + \bar\partial \quad \leftrightarrow \quad \bar\partial.$$

The method of using holomorphic bundles to study instantons is very fruitful, and has been used widely in the study of instanton moduli spaces, as it allows us to profit from all the technology of complex algebraic geometry to explore properties of instantons. In what follows we will identify instantons with holomorphic bundles via the Kobayashi–Hitchin correspondence, using the particular version of this conjecture proved in [GKM] for instantons on the surfaces Z_k. We can then talk about instanton charges in terms of characteristic classes.

3. Chern classes

Let us recall the basic properties of Chern classes of vector bundles. Let E be a vector bundle on a complex manifold M. The *Chern polynomial* of E is written as

$$c(E) = c_0(E) + c_1(E)t + c_2(E)t^2 + \cdots + c_n(E)t^n$$

where $c_i(E) \in H^{2i}(M, \mathbb{Z})$ is the i-th Chern class of E. These Chern classes satisfy the following 4 axioms:

(1) $c_0(E) = 1$

(2) $c_i(f^*(E)) = f^*(c_i(E))$
(3) $0 \to F_1 \to E \to F_2 \to 0 \Rightarrow c(E) = c(F_1)c(F_2)$
(4) $c_1(\gamma) = -1$,

where γ is the universal bundle on $\mathbb{P}^1$ whose fiber at a given point is the line in $\mathbb{C}^2$ it represents.


Observe that the single axiom setting the first Chern class of γ to -1 is the only one among the 4 axioms that guaranties nontriviality of the theory, else, without this axiom, the other properties would all have been satisfied by $c = 1$ for all bundles. The line bundle γ is usually denoted in algebraic geometry by $\mathcal{O}_{\mathbb{P}^1}(-1)$ and its total space is isomorphic to the blow-up of $\mathbb{C}^2$ at the origin: $\operatorname{Tot}\mathcal{O}_{\mathbb{P}^1}(-1) \simeq \widetilde{\mathbb{C}^2}$, where the blow-up of the complex plane is:

$$\widetilde{\mathbb{C}^2} := \{(z, l) \in \mathbb{C}^2 \times \mathbb{P}^1 : z \in l\}.$$

We will return to $\widetilde{\mathbb{C}^2}$ in the section about local surfaces, where it will be denoted by Z_1 being the first in the collection of surfaces $Z_k := \operatorname{Tot}\mathcal{O}_{\mathbb{P}^1}(-k)$ we will use, where $\mathcal{O}_{\mathbb{P}^1}(-k) = \mathcal{O}_{\mathbb{P}^1}(-1) \otimes \cdots \otimes \mathcal{O}_{\mathbb{P}^1}(-1)$ tensored k times has first Chern class $c_1(\mathcal{O}_{\mathbb{P}^1}(-k)) = -k$ by axiom 3. Given its importance in the context of characteristic classes and also for our calculations of local Chern classes, it is worth depicting the geometry of Z_1 in terms of the real analogue. The blow up of the real plane is

$$\widetilde{\mathbb{R}^2} := \{(z, l) \in \mathbb{R}^2 \times \mathbb{RP}^1 : z \in l\}$$

which can be depicted as



and is isomorphic to an infinite Möbius band (figure by C. Varea). Regarded as a vector bundle over $\mathbb{RP}^1$, the space $\widetilde{\mathbb{R}^2}$ is used within the axioms of characteristic classes of real vector bundles for the nontriviality axiom of Stiefel–Whitney classes.

Presented axiomatically the theory of Chern classes looks rather abstract. A more geometric approach is to write Chern classes in terms of polynomials on the curvature of a connection, as follows. Choosing any connection A on a complex vector bundle E, the first Chern class is actually the cohomology

class of the curvature of the connection, that is:

$$c_1(E) = \left[\frac{\sqrt{-1}}{2\pi}F_A\right] \in H^2(M)$$

and the higher Chern class c_i is given by the class of the elementary invariant polynomial P^i of degree i

$$c_i(E) = \left[P^i\left(\frac{\sqrt{-1}}{2\pi}F_A\right)\right] \in H^{2i}(M),$$

see [GH, p. 407].

The Gauss–Bonnet theorem then tells us that Chern classes measure obstruction to constructing nowhere vanishing sections. For example, the top Chern class gives the number of zeros of a generic section. In further generality, the i-th Chern class of a rank r vector bundle is Poincaré dual to the degeneracy cycle of $r - i + 1$ sections [GH, p. 413].

4. Local surfaces and local characteristic classes

We will consider the local surfaces $Z_k := \operatorname{Tot}\mathcal{O}_{\mathbb{P}^1}(-k)$. We begin by setting some notation:

We fix once and for all coordinate charts on Z_k, which we refer to as *canonical coordinates*, given by

$$U = \mathbb{C}^2_{z,u} = \big\{(z,u)\big\} \qquad \text{and} \qquad V = \mathbb{C}^2_{\xi,v} = \big\{(\xi,v)\big\},$$

such that on $U \cap V = \mathbb{C}^* \times \mathbb{C}$ we identify

$$\boxed{(\xi, v) = (z^{-1}, z^k u)}. \tag{1}$$

We denote by ℓ the subvariety of Z_k corresponding to the zero section of $\mathcal{O}(-k)$, thus $\ell \simeq \mathbb{P}^1$. Let X_k denote the surface obtained from Z_k by contracting ℓ to a point x, with

$$\pi\colon Z_k \to X_k$$

the contraction map. Then, $X_1 \simeq \mathbb{C}^2$ and X_k is singular at x for $k > 1$.

If E is a vector bundle on Z_k then the direct image π_*E happens to be a vector bundle on X_k only in the case when $E|_\ell$ is trivial, otherwise, it is a non locally free sheaf with the stalk at x strictly larger than at the smooth points of X_k.

We will consider rank 2 vector bundles and we wish to define the local second Chern class of a bundle E around ℓ using information from the sheaf π_*E. This local second Chern class will correspond to the local charge of an instanton on Z_k.

Considering π_*E then takes us to the realm of Chern classes of sheaves on singular varieties. Quoting T. Suwa we observe that for a singular complex algebraic or analytic variety X, there are at least three (in general different) kinds of Chern classes in the cohomology $H^*(X)$.

- the Chern–Schwartz–MacPherson class $c_*(X)$,
- the Chern–Mather class $c^M(X)$ and
- the canonical class or Fulton–Johnson's Chern class $c^{FJ}(X)$.

These three classes all reduce to the standard Chern classes $c(X)$ when the variety has no singularities. Since there is a priori no reason to prefer one of these definitions over the others, we chose instead to consider the concept of *local holomorphic Euler characteristic* $\chi(\ell, E)$ whose definition is uncontroversial.

If $\pi\colon (\widetilde{X}, \ell) \to (X, x)$ is a resolution of an isolated singularity, and $\mathcal{F}$ is a sheaf of rank n on $\widetilde{X}$, then the *local holomorphic Euler characteristic* of $\mathcal{F}$ is

$$\chi(x, \mathcal{F}) = \chi(\ell, \mathcal{F}) = h^0(x, Q) + \sum_{i=1}^{n} (-1)^i h^0(X, R^i\pi_*(\mathcal{F}))$$

where h^0 is the dimension of the 0-th Čech cohomology, $R^i\pi_*(\mathcal{F})$ is the i-th higher derived image of $\mathcal{F}$ (see [H, p. 250]) and Q is the skyscraper sheaf supported at x defined by the exact sequence

$$0 \to \pi_* F \to (\pi_*\mathcal{F})^{\vee\vee} \to Q \to 0.$$

If X is a compact orbifold and $\mathcal{F}$ a sheaf over X, there exists the global *holomorphic orbifold Euler characteristic* of $\mathcal{F}$ (see [Bl])

$$\chi_{orb}(X, \mathcal{F}) = \int_X \mathrm{ch}(\mathcal{F})\mathrm{td}(X).$$

It satisfies:

$$\chi(X, \mathcal{F}) = \chi_{orb}(X, \mathcal{F}) + \sum_{x \in \mathrm{Sing}(X)} \mu(x, \mathcal{F}).$$

Here $\mu(x, \mathcal{F})$ is a rational number that depends both on the class of the sheaf $\mathcal{F}$ in the Grothendieck group of the variety and on the order of singularity, with the denominator at an orbifold singularity given by the order of the corresponding group that gives the local quotient.

In the examples we wish to consider when E is a rank 2 vector bundle on Z_k then we refer to the local holomorphic Euler characteristic of E around ℓ as the *charge* of E for short, since it corresponds to the contribution to the charge in the language of instantons. It is given by

$$\chi(\ell, E) = h^0(Q) + h^0(R^1\pi_* E)$$

where

$$0 \to \pi_* E \to (\pi_* E)^{\vee\vee} \to Q \to 0.$$

The numbers $h^0(Q)$ and $h^0(R^1\pi_* E)$ are independent analytic invariants, called the *width* and the *height* of E respectively. These 2 invariants can be calculated using the theorem on formal functions (see [H, p. 277])

$$R^i f_*(E)^{\wedge}_x = \varprojlim H^i(\ell^n, E_n)$$

by calculating Čech cohomology in canonical coordinates as given in (1). It is the local holomorphic characteristic $\chi(\ell, E)$ that gets identified with the local instanton charge under the Kobayashi–Hitchin correspondence, and for this reason we call it the *charge* of the bundle, it gives the local contribution to the second Chern class $c_2^{loc}(X, E)$, which we will denote simply by $c_2^{loc}(E)$ when the base X is clear from the context.

To provide examples of these invariants, we first recall some properties of vector bundles on local surfaces. To start with, holomorphic vector bundles

on Z_k are filtrable and algebraic [G1, Lem. 3.1, Thm. 3.2]. Filtrability for a rank 2 bundle E means that it is an extension of line bundles. While filtrability always happens for bundles on complex curves (Riemann surfaces) this is a very unusual property for complex surfaces. For example, on the projective space $\mathbb{P}^2$ only split bundles are filtrable, so that the entire moduli spaces of stable vector bundles on $\mathbb{P}^2$ are made of nonfiltrable bundles.

If E is a rank 2 bundle on Z_k with $c_1(E) = 0$ then for some integer $j \geq 0$ there is a short exact sequence

$$0 \to \mathcal{O}(-j) \to E \to \mathcal{O}(j) \to 0, \tag{2}$$

where $E|_\ell = \mathcal{O}(-j) \oplus \mathcal{O}(j)$ splits by Grothendieck's splitting principle. The integer j is called the *splitting type* of E. Note that here we use the symbol $\mathcal{O}(j)$ for the line bundle with first Chern class j both on $\ell \simeq \mathbb{P}^1$ and on Z_k, but there should be no confusion given that one is obtained from the other by pullback.

Giving a short exact sequence $0 \to \mathcal{O}(-j) \to E \to \mathcal{O}(j) \to 0$ corresponds to giving an extension class $p \in \mathrm{Ext}^1(\mathcal{O}(j), \mathcal{O}(-j))$. This extension class can be written in canonical coordinates as a polynomial such that the transition matrix of E in canonical coordinates is $\begin{pmatrix} z^j & p \\ 0 & z^{-j} \end{pmatrix}$. Therefore, the bundle E is completely determined by the data (j, p).

The following table contains examples for the values of width, height and charge for vector bundles on Z_1 with splitting type $j = 3$ defined by the monomials in the first column:

monomial	width	height	charge
$z^{-1}u$	3	2	5
u	1	2	3
zu	1	2	3
z^2u	3	2	5
u^2	3	3	6
zu^2	2	3	5
z^2u^2	3	3	6
zu^3	4	3	7
z^2u^3	4	3	7
z^2u^4	5	3	8
zero	6	3	9

TABLE 1. $j = 3$ on Z_1

5. Moduli spaces

To define moduli of vector bundles, we first fix topological invariants, and then mod out by holomorphic isomorphisms. Next, as it is well known in algebraic geometry, we need to select a good subset of bundles to obtain moduli spaces. This means that we have to select a proper subset of the

set of all vector bundles with the chosen topology to obtain a well behaved quotient, else the quotient may be a stack, or something worse, but will not be an algebraic variety, at times not even a complex space. In geometric invariant theory, a well behaved moduli space is obtained using the concept of stable bundle. However, the concept of stability is defined for bundles on compact varieties and works well in such case, but it does not apply to bundles on noncompact varieties. For this reason, a somewhat ad-hoc choice of good moduli space was obtained for the surfaces Z_k by choosing those bundles that do not split on the first formal neighborhood of ℓ. In practice this means that the extension class has a nontrivial linear term on u, where $u = 0$ is the equation that cuts out ℓ inside the surface. Once this is done, then [BGK, Thm. 4.11] obtained: The *moduli space of holomorphic bundles* of rank 2 on Z_k with $c_1 = 0$ and splitting type $j \geq k \geq 2$ is a quasi-projective variety of dimension $2j - k - 2$.

Gasparim, Köppe, and Majumdar [GKM, Cor. 5.5] showed that a rank 2 bundle on Z_k corresponds to an SU(2) instanton if and only if its splitting type is a multiple of k. The correspondence also takes gauge transformation of instantons to isomorphism of holomorphic bundles. Furthermore, their choice of ad-hoc stability concept is such that *stable* instantons with splitting type j have topological charge j. It then follows that the *moduli space of* SU(2) *instantons* of charge $j = nk \geq 2$ on Z_k is a quasi-projective variety of dimension $2nk - k - 2$. There is a single instanton of charge 1 on Z_1 and, very surprisingly, there are no charge 1 instantons on Z_k for $k \geq 2$, as we shall see next.

6. Instanton decay

Sharp bounds for local charges of SU(2) instantons on Z_1 with splitting type j are as follows:

$$j \leq c_2^{loc}(Z_1, E) \leq j^2.$$

The minimum local charge of a splitting type j instanton on Z_1 equals j and is attained by those bundles that do not split on the first formal neighborhood, such as those with local data $(j, p = au + bzu)$ for any constants a, b not both 0. The maximum local charge is always attained by the split bundle, thus the case of the instanton with data $(j, p = 0)$ which has local charge j^2. Furthermore, every pair of admissible numerical values of $(width, height)$ occurs for some instanton on Z_1 [BG, Thm. 0.2]. Therefore, there exist instantons on Z_1 for any values of local charge, width and height.

In contrast, existence of instantons on Z_k with prescribed value of local charge is still an open problem. Indeed, even though sharp bounds are know for each fixed splitting type, it is not know whether all intermediate values of width and height actually occur. This issue is already featured in algebraic geometry, since the corresponding existence of vector bundles is unknown. The current state of affairs is as follows. Assume $k \geq 2$. Let E be a holomorphic rank 2 bundle on Z_k with $c_1 = 0$ and splitting type $j = nk + r$ with $0 \leq r < k$, then the following bounds are sharp [BGK, Cor. 2.18]:

$$j - 1 \leq c_2^{loc}(Z_k, E) \leq \begin{cases} n^2k + r(2n+1) - 1 & if \quad r \neq 0 \\ n^2k & if \quad r = 0. \end{cases} \tag{3}$$


Even though the question of existence of instantons on Z_k has not been completely understood when $k \geq 2$, the current partial knowledge generated strong conclusions regarding the possibilities for instanton decay. The main result obtained in [BGK, Thm. 6.8] is that if $k \geq 2$, then the minimal local charge of a nontrivial instanton on Z_k is $k-1$. As a corollary we conclude that instanton decay is obstructed by the self-intersection $\ell^2 = -k$ of the line in Z_k, given that there are no instantons of charge 1 on Z_k when $k > 2$.

Open question: For what admissible values of local charge c fitting into bounds (3) do there exist instantons on Z_k of charge c?

Open question: Assume A is an SU(2) instanton on Z_k of minimal charge, hence the local charge satisfies $c(A) = k - 1 > 1$ if $k > 2$. Then, such an instanton A ought to decay, because maintaining such local charge costs too much energy. However, A can not decay to other instantons on Z_k, because there are no nontrivial instantons of lower charge in this space. What is the result of the decay of such an instanton? We will give a partial solution to this question in the following section.

7. Holomorphic surgery and instanton decay

We define a new type of holomorphic surgery which we call *grafting*.



We will discuss 2 types of grafting: open and closed. Closed grafting is a specific new type of surgery on the base space such that, when the space holds an instanton (or a vector bundle), then the 4 dimensional manifold changes under grafting, but the instanton (or holomorphic bundle) on it remains essentially the same, although loosing some of its charge. We will carry out the closed grafting by replacing a Z_k with another $Z_{k'}$. Holomorphic patching of a Z_1 is possible at any smooth point, and corresponds to the operation of blowing up a point which is well known in algebraic geometry. However, the new operation considered here is not just that of changing the surface by adding a Z_1, but also carrying along with it an instanton (or holomorphic vector bundle).

We may regard the blow up $\pi\colon \widetilde{X} \to X$ of a point $x \in X$ as a holomorphic patching

$$\widetilde{X} = (X - \{x\}) \cup_{Z_1^o} Z_1.$$

Given holomorphic bundles E on X and F on Z_1 with the same rank, we may construct an open grafting of E and F, written $E \circledast F$, read "E graft F", by making $E \circledast F$ isomorphic to E on $\widetilde{X} - \ell$ and isomorphic to F on Z_1; this requires specifying a choice of gluing. So, the holomorphic type of the resulting bundle depends on the gluing, but the resulting second Chern class of $E \circledast F$ depends only on E and F, see [G2]. It is worth noticing that such grafting is always possible, because of the fact that every holomorphic bundle on Z_1 is trivial on $Z_1^o := Z_1 - \{\ell\}$, see [G1]. Similarly, we can graft a bundle locally by exchanging it on a Z_k type of neighborhood inside a complex surface. We have

$$c_2(E \circledast F) = c_2(E) + c_2^{loc}(F).$$

The second type of grafting, which is a somewhat more surprising operation is to essentially keep the vector bundle while changing the base manifold. Suppose that we have a complex surface X that contains a $-k$ line. Hence, X contains a line ℓ with a tubular neighborhood of the same holomorphic type as a neighborhood of ℓ in Z_k. We now replace k by k' by cutting out the $-k$ line and gluing in a $-k'$ line. If E is a bundle on X, we keep all of its transition matrices, and they now define a new bundle on $(X - \ell) \cup_{Z_k^o} Z_{k'}$ which we denote by $E \clubsuit_{k'}$ read "E graft k'". In general we will need to also modify other charts of X because of the cocycle condition in triple intersections. But, let us consider the simplest case when X is one of the Z_k themselves, so that we just exchange Z_k by $Z_{k'}$ and keep the transition matrix for the bundle. This might seem a trivial thing to do, nevertheless, the effect on the local charge is very strong, as the following examples illustrate.

First observe that over Z_k the bundle $E = \mathcal{O}(j) \oplus \mathcal{O}(-j)$, that is, the split bundle with type $j = nk$ and $p = 0$ over Z_k, has local charge $c_2^{loc}(Z_k, E) = n^2 k$, in fact, it realizes the upper bound in 3. Now, let us see the effect of grafting in a couple of cases, by considering the split bundles with type $j = 6$ over Z_1 and Z_2. So that, in each case, we are considering a bundle defined by transition matrix $T = \begin{pmatrix} z^6 & 0 \\ 0 & z^{-6} \end{pmatrix}$.

Case 1. When we consider the split bundle $E \to Z_1$ given by transition matrix T we are in the case $j = 6 \times 1$, so

$$c_2^{loc}(E) = 6^2 \times 1 = 36$$

by grafting to Z_3 we get the bundle $E \clubsuit_3$ also given by transition matrix T but now we are in the case $j = 2 \times 3$, so

$$c_2^{loc}(E \clubsuit_3) = 2^2 \times 3 = 12.$$

So that in this grafting the instanton bundles lost 24 units of charge.

Case 2. When we consider the split bundle $E \to Z_2$ given by transition matrix T we are in the case $j = 3 \times 2$, so

$$c_2^{loc}(E) = 3^2 \times 2 = 18$$

by grafting to Z_6 we get the bundle $E\clubsuit_6$ also given by transition matrix T but now we are in the case $j = 1 \times 6$, so

$$c_2^{loc}(E\clubsuit_6) = 1^2 \times 6 = 6.$$

So that in this grafting the instanton bundles lost 12 units of charge.

In both cases we can interpret the loss of charge as the energy used to provoke the grafting in the base.

Now let $j \geq 2$ and $E \to Z_1$ the bundle defined by the transition matrix $T' = \begin{pmatrix} z^j & zu \\ 0 & z^{-j} \end{pmatrix}$. As E does not split, we have that


$$c_2^{loc}(E) = j.$$

By grafting to Z_3 we have that

$$c_2^{loc}(E\clubsuit_3) = j - 1,$$

so in this grafting the instanton bundles lost only 1 unit of charge.

Here is a picture of closed grafting, as it exchanges the local surfaces Z_k and $Z_{k'}$ inside a complex surface, while essentially maintaining the instanton bundle globally. The grafted bundle is still given by the same transition matrices, however, loses global charge, that is, undergoes a loss of second Chern class.



In conclusion, we propose that an instanton with high charge provokes a grafting on the 4 dimensional base manifold which holds it. The grafting happens around a 2-sphere where the instanton charge is highly concentrated. The instanton looses charge in the process, while the curvature of the base manifold around the 2-sphere increases. In other words, we have explained that the decay of an instanton can happen when the instanton inflicts larger concentrations of curvature around 2-spheres, thereby losing charge. We then say that the addition to the curvature is grafted by the instanton.

8. Acknowledgment

I first met Pushan Majumdar in 1998 and we immediately engaged into scientific discussions. Shortly after that, I found him reading about moduli of instantons, and inquired why he was reading about such a theme, which was not part of his PhD project. He said that he wanted to be my friend, and if he did not learn about the themes that interested me, then soon we would not have anything to talk about. Decades later, this remains the most loyal offer of friendship I have ever received. E.G.

We thank Koushik Ray for inviting us to contribute to the memorial volume of Pushan and for suggesting several improvements to our article.

References

[ADHM] M. Atiyah, V.G. Drinfeld, N.J. Hitchin, Y.I. Manin, *Construction of instantons*, Physics Letters A, **65** n.3 (1978) 185–187.

[BG] E. Ballico, E. Gasparim, *Vector bundles on a neighborhood of a curve in a surface and elementary transformations*, Forum Math. **15** n.1 (2003) 115–122.

[BGK] E. Ballico, E. Gasparim, T. Köppe, *Vector bundles near negative curves: moduli and local Euler characteristic*, Comm. Algebra **37** (2009) 2688–2713.

[Bl] R. Blache, *Chern classes and Hirzebruch–Riemann–Roch theorem for coherent sheaves on complex projective orbifolds with isolated singularities*, Math. Z. **222** (1996) 7–57.

[D] S. K. Donaldson, *Anti-self-dual Yang-Mills connections over complex algebraic surfaces and stable vector bundles*, Proc. London Math. Soc. (3) **50** (1985), 1–26.

[G1] E. Gasparim, Holomorphic bundles on $\mathcal{O}(-k)$ are algebraic, Comm. Algebra **25** (1997) 3001–3009.

[G2] E. Gasparim, The Atiyah–Jones conjecture for rational surfaces, Adv. Math. **218** (2008) 1027–1050.

[GKM] E. Gasparim, T. Köppe, P. Majumdar, Local holomorphic Euler characteristic and instanton decay, Pure Appl. Math. Q. **4** (2008) 161–179.

[GH] P. Griffiths, J. Harris, *Principles of Algebraic Geometry*, Wiley & Sons Inc. (1978).

[H] R. Hartshorne, *Algebraic geometry*, Springer, Berlin 1977.

[LT] M. Lübke, A. Teleman, Andrei, *The Kobayashi–Hitchin correspondence*, River Edge, NJ: World Scientific Publishing Co. Inc. (1995).

[Se] C.S. Seshadri, *Theory of moduli*, in: Algebraic geometry (Arcata 1974), Proc. Sympos. Pure Math. **29**, Am. Math. Soc., Providence, RI (1975) 263–304.

[UY] K. Uhlenbeck, S.T. Yau, *On the existence of Hermitian–Yang–Mills connections in stable vector bundles*, Comm. Pure App. Math. **39** (1986) S257–S293.

Depto. Matemáticas, Universidad Católica del Norte, Chile, etgasparim@gmail.com, obrunosuzuki@gmail.com