

BIPARTITE EXTENSION GRAPHS AND THE DUFLO–SERGANOVA FUNCTOR

MARIA GORELIK

ABSTRACT. We consider several examples when the extension graph admits a bipartition compatible with the action of Duflo–Serganova functors.

0. INTRODUCTION

In this paper $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ is a complex Lie superalgebra and $x \in \mathfrak{g}_1$ is such that $[x, x] = 0$. This text is based on some results and ideas of [27], [18] and [33]. We give a more detailed presentation of some parts of [27], [18] which are slightly different for \mathfrak{osp} and \mathfrak{gl} -cases.

This text is not intended for publication; its different parts have been developed in other papers [14] and [12].

0.1. Let \mathcal{C} be a category of representations of a Lie superalgebra \mathfrak{g} and $\text{Irr}(\mathcal{C})$ be the set of isomorphism classes of simple modules in \mathcal{C} . Assume that the modules in \mathcal{C} are of finite length.¹ In many examples the extension graph of \mathcal{C} is bipartite, i.e. there exists a map $\text{dex} : \text{Irr}(\mathcal{C}) \rightarrow \mathbb{Z}_2$ such that

$$\text{(Dex1)} \quad \text{Ext}_{\mathcal{C}}^1(L_1, L_2) = 0 \quad \text{if} \quad \text{dex}(L_1) = \text{dex}(L_2).$$

In this paper we are interested in examples when the map dex is “compatible” with the Duflo–Serganova functors DS_x , i.e.

$$\text{(Dex2)} \quad \text{one has} \quad [\text{DS}_x(L) : L'] = 0 \quad \text{if} \quad \text{dex}(L) \neq \text{dex}(L').$$

Note that in (Dex2) we have to choose dex on $\text{Irr}(\mathcal{C})$ and on $\text{Irr}(\text{DS}_x(\mathcal{C}))$. If dex satisfies (Dex1) and (Dex2), then $\text{DS}_x(L)$ is completely reducible for each $L \in \mathcal{C}$.

We say that a module M is *pure* if for any subquotient L of M , $\Pi(L)$ is not a subquotient of M . Note that (Dex2) implies the purity of $\text{DS}_x(L)$ for each $L \in \text{Irr}(\mathcal{C})$.

In what follows $\mathcal{F}\text{in}(\mathfrak{g})$ stands for the full subcategory of finite-dimensional \mathfrak{g} -modules which are completely reducible over \mathfrak{g}_0 . In this paper we consider the case when \mathfrak{g} is a finite-dimensional Kac–Moody superalgebra and $\mathcal{C} = \mathcal{F}\text{in}(\mathfrak{g})$; for this case we require that (Dex2) holds for each $x \in X(\mathfrak{g})$, where

$$X(\mathfrak{g}) := \{x \in \mathfrak{g}_1 \mid [x, x] = 0\}.$$

¹this can be substituted by existence of local composition series constructed in [3].

For other cases it make sense to restrict (Dex2) to certain values of x : for instance, for affine superalgebras \mathfrak{g} it makes sense to consider x such that $\text{DS}_x(\mathfrak{g})$ is affine and for $\mathcal{C} = \mathcal{O}(\mathfrak{g})$ it makes sense to consider x "preserving" the category \mathcal{O} (see [11], Sections 7 and 8 respectively).

The examples with dex satisfying (Dex1) and (Dex2) include $\mathcal{F}\text{in}(\mathfrak{g})$ for $\mathfrak{g} = \mathfrak{gl}(m|n)$ (this follows from [21]) and the full subcategory of integrable modules in the category $\mathcal{O}(\mathfrak{gl}(1|n)^{(1)})$ (this follows from [16]). In Section 2 we will check the existence of dex for exceptional Lie superalgebras. We will introduce dex satisfying (Dex1) for $\mathfrak{osp}(m|n)$ -case; in [13] we will show that (Dex2) holds in this case too. It would be interesting to find other examples with dex satisfying (Dex1) and (Dex2). The strange superalgebras $\mathfrak{p}_n, \mathfrak{q}_n$ do not admit dex satisfying (Dex1), (Dex2). By [17], for $n \leq 3$, $\mathcal{F}\text{in}(\mathfrak{q}_n)$ admits dex satisfying (Dex1); this does not hold for larger n , see [26]. Moreover, the module $\text{DS}_x(L)$ is not pure for each atypical $L \in \text{Irr}(\mathcal{F}\text{in}(\mathfrak{q}_2))$ (and $x \neq 0$), see [11], 5.5.2. By contrast, the module $\text{DS}_x(L)$ is pure for $L \in \text{Irr}(\mathcal{F}\text{in}(\mathfrak{p}_n))$ if x is of rank 1, see [8].

In [18], C. Gruson and V. Serganova express the character of a simple finite-dimensional $\mathfrak{osp}(M|N)$ -module in terms of a basis consisting of "Euler characters". Using dex we will show that all coefficients in such formula have the same sign (equal to $\text{dex}(L)$) (we call this property "positiveness"). The similar formulae hold for the simple modules in $\mathcal{F}\text{in}(\mathfrak{g})$, where \mathfrak{g} is an exceptional Lie superalgebra or $\mathfrak{gl}(1|n)^{(1)}$ (see [24],[34]). For \mathfrak{q}_n the character formula of the above form was obtained in [30],[31] (see also [1]). In [14] we will prove a Gruson-Serganova type character formula for $\mathfrak{gl}(m|n)$ -case.

The reduced Grothendieck ring is the quotient of the Grothendieck ring modulo the relation $[N] + [\Pi N] = 0$. If \mathcal{C} is rigid, then $*$ induces an involution of the reduced Grothendieck ring. By Hinich's Lemma, DS-functor induces a homomorphism ds of the reduced Grothendieck rings; this homomorphism, introduced in [21], is compatible with the above involutions (in many cases the reduced Grothendieck ring is isomorphic to the ring of supercharacters so ds can be represented as the restriction of supercharacters to a subalgebra of \mathfrak{h} ; for the algebras from the list (1) the homomorphism ds was studied in [22]).

0.2. The map dex for finite-dimensional Kac-Moody superalgebras. Let \mathfrak{g} be one of the following superalgebras:

- (1) $\mathfrak{gl}(m|n), \mathfrak{osp}(m|2n)$ for $m, n \geq 0$, $D(2, 1|a)$, $F(4)$, $G(3)$, $\mathfrak{sl}(m|n)$ for $m \neq n$.

Note that the list (1) includes $\mathfrak{gl}(0|0) = \mathfrak{osp}(1|0) = \mathfrak{osp}(0|0) = 0$, $\mathfrak{osp}(2|0) = \mathbb{C}$ and the reductive Lie algebras $\mathfrak{gl}_m, \mathfrak{o}_m, \mathfrak{sp}_m$. For each value of x , the algebra \mathfrak{g}_x is again one of the algebras from the list (1).

Let \mathfrak{h} be a Cartan subalgebra of \mathfrak{g} . We denote by $\Lambda_{m|n}$ the integral weight lattice in \mathfrak{h}^* ; this lattice is equipped by the standard parity function $p : \Lambda_{m|n} \rightarrow \mathbb{Z}_2$ (for

$\mathfrak{gl}(m|n)$, $\mathfrak{osp}(M|N)$ and $G(3)$ the lattice $\Lambda_{m|n}$ is spanned by ε_i s and δ_j s with $p(\varepsilon_i) := \bar{0}$ and $p(\delta_j) := \bar{1}$. For our purposes the study of the category $\mathcal{F}\text{in}(\mathfrak{g})$ of finite dimensional representations of \mathfrak{g} reduces to study the category $\tilde{\mathcal{F}}(\mathfrak{g})$ with the modules whose weights lie in $\Lambda_{m|n}$. In its turn, the category $\tilde{\mathcal{F}}(\mathfrak{g})$ decomposes into a direct sum two equivalent categories

$$\tilde{\mathcal{F}}(\mathfrak{g}) = \mathcal{F}(\mathfrak{g}) \oplus \Pi(\mathcal{F}(\mathfrak{g})),$$

where the grading on the modules in $\mathcal{F}(\mathfrak{g})$ is induced by the parity function p . Our goal is to find a map

$$\text{dex} : \text{Irr}(\tilde{\mathcal{F}}(\mathfrak{g})) \rightarrow \mathbb{Z}_2$$

satisfying (Dex1) and (Dex2). By [33], for each $L \in \text{Irr}(\tilde{\mathcal{F}}(\mathfrak{g}))$ there exists x such that $\text{DS}_x(L)$ is a non-zero typical module. Therefore it is enough to define dex on the typical simple modules. If this is done in such a way that $\text{dex}(\Pi(L)) \neq \text{dex}(L)$, then dex satisfying (Dex2) is unique and satisfies

$$(2) \quad \text{dex}(\Pi(L)) \neq \text{dex}(L) \quad \text{for each } L \in \text{Irr}(\tilde{\mathcal{F}}(\mathfrak{g})).$$

0.3. Reduction to DS_1 . In many cases it is enough to check (Dex2) for one particular value of x . We continue to consider the case when \mathfrak{g} is as in (1) (the same reasoning work for symmetrizable affine Lie superalgebras for x as in Section 9 of [11]). We set

$$\text{rank } x := \text{defect } \mathfrak{g} - \text{defect } \mathfrak{g}_x \quad \text{for } x \in X(\mathfrak{g}).$$

By [5], $\mathfrak{g}_x \cong \mathfrak{g}_y$ if $x, y \in X(\mathfrak{g})$ are such that $\text{rank } x = \text{rank } y$; we set

$$X(\mathfrak{g})_r := \{x \in X(\mathfrak{g}) \mid \text{rank } x = r\}.$$

Using Lemma 2.4.1 in [11], one can reduce (Dex2) to the case $x \in X(\mathfrak{g})_1$ (see [13], Section 9 for details). Fix any $x \in X(\mathfrak{g})_1$ and denote DS_x by DS_1 . Using the results of [5], it is easy to see that for each $y \in X(\mathfrak{g})_1$ there exists an automorphism $\phi \in \text{Aut}(\mathfrak{g})$ satisfying $\phi(x) = y$; this automorphism induces an isomorphism $\bar{\phi} : \mathfrak{g}_x \xrightarrow{\sim} \mathfrak{g}_y$ and $\text{DS}_x(N^\phi) = (\text{DS}_y(N))^{\bar{\phi}}$. Thus DS_1 is ‘‘independent’’ from the choice of x ; in particular, if the formula (Dex2) holds for DS_x , then it holds for each $y \in X(\mathfrak{g})_1$. The argument of [13], Section 9 give

if dex satisfies (Dex1), (2) and (Dex2) holds for some x of rank 1, then

$$(3) \quad \text{DS}_{x'}(L) \cong \underbrace{\text{DS}_1(\text{DS}_1 \dots \text{DS}_1(L) \dots)}_{\text{rank } x' \text{ times}} \quad \text{for any } L \in \text{Irr}(\mathcal{F}\text{in}(\mathfrak{g})) \text{ and each } x' \in X(\mathfrak{g}).$$

In this way, the computation of $\text{DS}_x(L)$ reduces to the computation of the multiplicities $[\text{DS}_1(L') : L'']$ for each quadruple $(\mathfrak{g}', \mathfrak{g}'', L', L'')$, where

$$\mathfrak{g}' := \underbrace{\text{DS}_1(\text{DS}_1 \dots \text{DS}_1(\mathfrak{g}) \dots)}_{i \text{ times}}, \quad \mathfrak{g}'' := \text{DS}_1(\mathfrak{g}')$$

and $L' \in \text{Irr}(\mathcal{F}\text{in}(\mathfrak{g}'))$, $L'' \in \text{Irr}(\mathcal{F}\text{in}(\mathfrak{g}''))$.

0.3.1. The multiplicities $[\mathrm{DS}_1(L') : L'']$ were computed for $\mathfrak{gl}(m|n)$ in [21]. For the exceptional cases we compute the multiplicities in Section 2. For the remaining case $\mathfrak{osp}(m|n)$ the multiplicities are computed in [13]. In all these cases the following properties hold:

- the module $\mathrm{DS}_1(L')$ is pure;
- $[\mathrm{DS}_1(L') : L''] \leq 2$;
- $[\mathrm{DS}_1(L') : L''] \leq 1$ for $\mathfrak{g} = \mathfrak{gl}(m|n)$;
- there exists dex satisfying (Dex1), (2) and the formula (Dex2) for the case when $\mathrm{rank} x = 1$.

By above, this implies (Dex2) (for any $x \in X(\mathfrak{g})$) and shows that $\mathrm{DS}_x(L)$ is pure, semisimple and can be computed via the formula (3).

0.3.2. In 2.3.3 we compute $\mathrm{DS}_1(L)$ for $F(4)$. The results imply that the image of the Grothendieck ring of $\mathcal{F}(F(4))$ under the homomorphism ds coincides with σ -invariants in the Grothendieck ring of $\mathcal{F}(\mathfrak{sl}_3)$ for σ induced by a Dynkin diagram involution of \mathfrak{sl}_3 . For all other algebras from the list (1) a similar result is obtained in [22].

0.3.3. *Remark.* By [8], $\mathrm{DS}_1(L)$ is pure and multiplicity free for each $L \in \mathrm{Irr}(\mathcal{F}\mathrm{in}(\mathfrak{p}_n))$; however, $\mathrm{DS}_1(L)$ is not always semisimple, (3) does not hold and $\mathrm{DS}_1(\mathrm{DS}_1(L))$ is not always pure (see [11], Example 3.4.3).

0.3.4. *Question.* Let $\mathfrak{p} \subset \mathfrak{g}$ be a parabolic subalgebra containing \mathfrak{b} with the property that the defect of the Levi subalgebra of \mathfrak{p} is less by one than the defect of \mathfrak{g} . We denote by $L(\lambda)$ (resp., $L_{\mathfrak{p}}(\lambda)$) a simple \mathfrak{g} (resp., \mathfrak{p}) module of the highest weight λ . For $L(\lambda) \in \tilde{\mathcal{F}}(\mathfrak{g})$ we define $\Gamma_{\mathfrak{g},\mathfrak{p}}^i(L_{\mathfrak{p}}(\lambda))$ as in [18] (see 3.1 below) and consider the multiplicities

$$m_{\lambda,\mu}^{(i)} := [\Gamma_{\mathfrak{g},\mathfrak{p}}^i(L_{\mathfrak{p}}(\lambda)) : L_{\mathfrak{g}}(\mu)].$$

One has $m_{\lambda,\lambda}^{(0)} = 1$. In all our examples $m_{\lambda,\mu}^{(i)} \in \{0, 1\}$ and, except for the case $m_{\lambda,\lambda}^{(0)}$, one has

$$m_{\lambda,\mu}^{(i)} \neq 0 \implies \mathrm{dex}(L(\lambda)) - \mathrm{dex}(L(\mu)) \equiv i + 1 \pmod{2}.$$

It is interesting to know whether these properties hold in other cases.

0.4. **Content of the paper.** In Section 1 we recall the construction of DS-functor.

In Section 2 we consider the cases $\mathfrak{g} = D(2|1; a)$, $F(4)$, $G(3)$ and $\mathfrak{osp}(3|2)$. We compute $\mathrm{DS}_x(L)$ for $L \in \mathrm{Irr}(\tilde{\mathcal{F}}(\mathfrak{g}))$ and check that dex satisfies (Dex1), (Dex2) and (2). The modules $\mathrm{DS}_x(L)$ can be described as follows. To each atypical block \mathcal{B} in $\tilde{\mathcal{F}}(\mathfrak{g})$ we assign a \mathfrak{g}_x -module L' (this assignment is injective). By [10], for each atypical block of $\mathfrak{osp}(3|2)$, $G(3)$ the extension graph $\mathrm{Ext}(\mathcal{B})$ is D_{∞} ; for the cases $F(4)$, $D(2|1; a)$ the graphs of atypical blocks are A_{∞}^{∞} and D_{∞} (see [10], [25]). If $\mathrm{Ext}(\mathcal{B}) = D_{\infty}$, then L' is simple and $\mathrm{DS}_x(L) \cong L'$ if $L \in \mathrm{Irr}(\mathcal{B})$ is an ‘‘end vertex’’ of D_{∞} and $\mathrm{DS}_x(L) \cong \Pi^i(L')^{\oplus 2}$ if L is the i th vertex

counting from the ends. For $\text{Ext}(\mathcal{B}) = A_\infty^\infty$ one has $\text{DS}_x(L) \cong \Pi^i(L')$ for each $L \in \text{Irr}(\mathcal{B})$. For $\mathfrak{g} = F(4)$, $D(2|1; a)$ the module L' corresponding to D_∞ -graph is a simple \mathfrak{g}_x -module satisfying $(L')^* \cong L'$, whereas the module L' corresponding to A_∞^∞ -graph is of the form $L' = V \oplus V^*$, where V is a simple \mathfrak{g}_x -module with $V^* \not\cong V$.

In Section 3 we construct for each block \mathcal{B} in $\tilde{\mathcal{F}}(\mathfrak{g})$ a graph $\hat{\Gamma}^\times$ and its subgraph Γ^\times defined in terms of the functors Γ^i introduced in [29] and [18] (we follow the definition in [18]). The extension graph $\text{Ext}(\mathcal{B})$ is a subgraph of $\hat{\Gamma}^\times$; the graph Γ^\times is useful for Gruson-Serganova type character formulae. In 3.5.7 we introduce a notion of “parametric bipartition” on the graphs $\hat{\Gamma}^\times$, Γ^\times ; a parametric bipartition on $\hat{\Gamma}^\times$ induces a bipartition on $\text{Ext}(\mathcal{B})$; a parametric bipartition on Γ^\times gives a “positive” Gruson-Serganova type character formulae. In Corollary 3.6.3 we show that under a certain conditions (which hold in the \mathfrak{osp} -case) $\text{Ext}(\mathcal{B})$ is a subgraph of Γ^\times and a parametric bipartition on Γ^\times induces a bipartition on $\text{Ext}(\mathcal{B})$.

The graphs $\hat{\Gamma}^\times$, Γ^\times depend on the choice of triangular decomposition; for the case $\mathfrak{osp}(2|2) \cong \mathfrak{sl}(2|1)$ the graph $\text{Ext}(\mathcal{B})$ is a subgraph of Γ^\times for the “mixed” triangular decomposition and is not a subgraph of Γ^\times for the distinguished one, see Remark 3.9.4. By [27], for $\mathfrak{gl}(m|n)$ -case the map dex gives a parametric bipartition on $\hat{\Gamma}^\times$; we check that this also holds for $\mathfrak{osp}(2|2)$, $\mathfrak{osp}(3|2)$, $D(2|1; a)$, $F(4)$ and $G(3)$.

In Section 4 we consider the principal block \mathcal{B} for $\mathfrak{g} = \mathfrak{osp}(2n + t|2n)$ with $t = 0, 1, 2$. In this case the extension graph of \mathcal{B} is a subgraph of Γ^\times . We describe dex which gives a parametric bipartition on Γ^\times ; this implies (Dex1) and “positiveness” of the Gruson-Serganova character formula.

In Section 5 we give a description of Γ^\times in the \mathfrak{osp} -case using the language of “arch diagrams” introduced in [13]. The results of this section are not used in the rest of the paper.

In Appendix we explain why $\text{Ext}(\mathcal{B})$ is a subgraph of $\hat{\Gamma}^\times$ (this part essentially follows Sect. 6 of [27]).

0.5. Acknowledgments. The author is grateful to I. Entova-Aizenbud, V. Hinich, T. Heidersdorf, C. Hoyt, I. M. Musson, S. Reif, V. Serganova, A. Sherman and C. Stroppel for numerous helpful discussions.

0.6. Index of definitions and notation. Throughout the paper the ground field is \mathbb{C} ; \mathbb{N} stands for the set of non-negative integers. We will use the standard Kac’s notation [23] for the root systems.

dex, (Dex1), (Dex2), pure	0.1
$\Lambda_{m n}, \tilde{\mathcal{F}}(\mathfrak{g}), \mathcal{F}(\mathfrak{g})$	0.2
$\Gamma_{\mathfrak{p},\mathfrak{q}}^i, {}^i K_{\mathfrak{p},\mathfrak{q}}^{\lambda,\nu}$	3.1
increasing/descreeasing paths	3.4.1
decreasingly equivalent	3.4.2
(BB)	Lemma 3.4.4
$\Lambda^\chi, \text{tail}(\lambda), \mathfrak{p}_\lambda$	3.5
$\hat{\Gamma}^\chi, \kappa,$	3.5.1
$\Gamma^\chi, (\text{Tail})$	3.5.2
parametric bipartition	3.5.7
\mathcal{E}_λ	(13) in 3.7
weight diagram	4.2.1
$Diag_{k;t}, \lambda(f)$	4.2.4
τ	4.2.5
dex for \mathfrak{osp} -case, $\ \lambda\ , \ \lambda\ _{gr}$	4.3

1. DS-FUNCTOR

The DS-functor was introduced in [5]; see also [15] for an expanded exposition. We recall definitions and some results below. In this section \mathfrak{g} is any superalgebra; we set $X(\mathfrak{g}) := \{x \in \mathfrak{g}_1 \mid [x, x] = 0\}$.

1.1. Construction. For a \mathfrak{g} -module M and $g \in \mathfrak{g}$ we set $M^g := \text{Ker}_M g$. For $x \in X(\mathfrak{g})$ we set

$$\text{DS}_x(M) := M^x/xM.$$

Notice that \mathfrak{g}^x and $\mathfrak{g}_x := \text{DS}_x(\mathfrak{g}) = \mathfrak{g}^x/[x, \mathfrak{g}]$ are Lie superalgebras. Since M^x, xM are \mathfrak{g}^x -invariant and $[x, \mathfrak{g}]M^x \subset xM$, $\text{DS}_x(M)$ is a \mathfrak{g}^x -module and \mathfrak{g}_x -module. Thus

$$\text{DS}_x : M \mapsto \text{DS}_x(M)$$

is a functor from the category of \mathfrak{g} -modules to the category of $\text{DS}_x(\mathfrak{g})$ -modules.

There are canonical isomorphisms $\text{DS}_x(\Pi(N)) = \Pi(\text{DS}_x(N))$ and

$$\text{DS}_x(M) \otimes \text{DS}_x(N) = \text{DS}_x(M \otimes N).$$

For a finite-dimensional module L one has $\text{DS}_x(L^*) \cong \text{DS}_x(L)^*$.

1.2. Hinich's Lemma. The following result is called *Hinich's Lemma* (see [5]); a similar result is Lemma 2.1 in [21].

A short exact sequence of \mathfrak{g} -modules

$$0 \rightarrow M_1 \rightarrow N \rightarrow M_2 \rightarrow 0$$

induces a long exact sequence of \mathfrak{g}_x -modules

$$0 \rightarrow Y \rightarrow \text{DS}_x(M_1) \rightarrow \text{DS}_x(N) \rightarrow \text{DS}_x(M_2) \rightarrow \Pi(Y) \rightarrow 0$$

2.1. Proposition. *Take a non-zero $x \in \mathfrak{g}_1$ satisfying $x^2 = 0$.*

Let \mathcal{B} be a block and $\text{Irr}(\mathcal{B}) =: \{L^i\}_{i \in I}$. We assume that each $L^i \in \text{Irr}(\mathcal{B})$ has a projective cover with a three step radical filtration with the following subquotients

$$(5) \quad L^i; \bigoplus_{j \in \text{Adj}(i)} L^j; L^i, \quad \text{where } \text{Adj}(i) := \{j \in I \mid \text{Ext}^1(L^i, L^j) \neq 0\}.$$

We set $M_i := \text{DS}_x(L^i)$. Assume that for some s the module M_s is pure and

$$\text{Ext}_{\mathcal{F}(\mathfrak{g}_x)}^1(M_s, M_s) = 0.$$

- (i) *For each $i \in I$ the module M_i is pure. Moreover, if $M_s \neq 0$, then $M_i \neq 0$ for each $i \in I$.*
- (ii) *If $M_s \neq 0$, then the extension graph $\text{Ext}(\mathcal{B})$ is bipartite.*
- (iii) *Using the notation of (4) we have*

$$\begin{aligned} \text{if } \text{Ext}(\mathcal{B}) = A_\infty & \text{ then } M_j \cong \Pi^j(M_0)^{\oplus 2} \text{ for } j \geq 1; \\ \text{if } \text{Ext}(\mathcal{B}) = D_\infty & \text{ then } M_j \cong \Pi^{j-1}(M_0)^{\oplus 2} \text{ for } j \geq 2, M_1 \cong M_0; \\ \text{if } \text{Ext}(\mathcal{B}) = A_\infty^\infty & \text{ then } M_j \cong \Pi^j(M_0). \end{aligned}$$

Proof. By [5], the functor DS_x (for $x \neq 0$) kills the projective modules in $\mathcal{F}(\mathfrak{g})$. Using Corollary 1.2.1 and (5) we conclude that for any i the purity of M_i implies the existence of an exact sequence

$$(6) \quad 0 \rightarrow \Pi(M_i) \rightarrow \bigoplus_{j \in \text{Adj}(i)} M_j \rightarrow \Pi(M_i) \rightarrow 0$$

In particular, the purity of M_i implies the purity of M_j for each $j \in \text{Adj}(i)$ and $M_i = 0$ implies $M_j = 0$ for $j \in \text{Adj}(i)$. Since $\text{Ext}(\mathcal{B})$ is connected, this proves (i).

Let L' be a simple module L' such that $[M_s : L'] \neq 0$. For each $i \in I$ set $p_i := [M_i : L']$, $q_i := [M_i : \Pi(L')]$. By above, $p_s \neq 0$. Using (6) and (i) we obtain

$$(7) \quad p_i, q_i \geq 0, \quad 2q_i = \sum_{j \in \text{Adj}(i)} p_j, \quad 2p_i = \sum_{j \in \text{Adj}(i)} q_j, \quad p_i q_i = 0$$

for each $i \in I$. In particular, if $p_j = q_j = 0$ for some j , then $p_i = q_i = 0$ for each i , a contradiction. Hence for each i either $p_i \neq 0$ or $q_i \neq 0$. It is easy to see from (7) that $p_i = 0$ if L^i, L^s are connected by a path of odd length. Hence $\text{Ext}(\mathcal{B})$ does not have cycles of odd length; this gives (ii). For A_∞ and D_∞ , (iii) follows from (7) by induction. For the case A_∞^∞ observe that $m_i := p_i + q_i$ satisfies $2m_i = m_{i-1} + m_{i+1}$ for $i \in I = \mathbb{Z}$. Since $m_i \geq 0$, we get $m_i = m_s$ for each i . This completes the proof. \square

2.2. **DS-functor for small rank \mathfrak{g} .** Let \mathfrak{g} be one of the Lie superalgebras

$$\mathfrak{osp}(2|2), \mathfrak{osp}(3|2), D(2|1; a), G(3), F(4).$$

Let \mathfrak{h} be a Cartan subalgebra of \mathfrak{g}_0 . We denote by W the Weyl group of \mathfrak{g}_0 and by $(-|-)$ the symmetric non-degenerate form on \mathfrak{h}^* which is induced by a non-degenerate invariant form on \mathfrak{g} .

2.2.1. Let Σ be a base of \mathfrak{g} which contains an isotropic root β . Fix a non-zero $x \in \mathfrak{g}_\beta$.

Set $\Delta_x := (\beta^\perp \cap \Delta) \setminus \{\beta, -\beta\}$. By [5], \mathfrak{g}_x can be identified with a subalgebra of \mathfrak{g} generated by the root spaces \mathfrak{g}_α with $\alpha \in \Delta_x$ and a Cartan subalgebra $\mathfrak{h}_x \subset \mathfrak{h}$. If Δ_x is not empty, then Δ_x is the root system of the Lie superalgebra \mathfrak{g}_x and one can choose Σ_x in Δ_x such that $\Delta^+(\Sigma_x) = \Delta^+ \cap \Delta_x$. If $\mathfrak{g} = \mathfrak{osp}(m|2)$, then $\mathfrak{g}_x = \mathfrak{o}_{m-2}$; for $\mathfrak{g} = D(2|1; a), G_3, F_4$ one has $\mathfrak{g}_x = \mathbb{C}, \mathfrak{sl}_2, \mathfrak{sl}_3$ respectively.

2.2.2. **Lemma.** *Let $L := L(\lambda)$ be a finite-dimensional module and $(\lambda|\beta) = 0$. Set $L' := L_{\mathfrak{g}_x}(\lambda|_{\mathfrak{h}_x})$. One has*

$$DS_x(L) \cong \begin{cases} L' & \text{for } \mathfrak{osp}(2|2), \mathfrak{osp}(3|2), G(3) \\ L' & \text{for } D(2|1; a), F(4) \text{ if } L' \cong (L')^* \\ L' \oplus (L')^* & \text{for } D(2|1; a), F(4) \text{ if } L' \not\cong (L')^*. \end{cases}$$

Proof. It is easy to see that $[DS_x(L) : L'] = 1$. Set $\lambda' := \lambda|_{\mathfrak{h}_x}$. From [5], Sect. 7, $DS_x(L)$ is a typical module and each simple subquotient of $DS_x(L)$ is of the form $L_{\mathfrak{g}_x}(\nu)$ with $\nu \in \{\lambda', \sigma(\lambda')\}$, where $\sigma = \text{Id}$ for $\mathfrak{g} = \mathfrak{osp}(2|2), \mathfrak{osp}(3|2)$ and $G(3)$, $\sigma = -\text{Id}$ for $D(2|1; a)$ and σ is the Dynkin diagram automorphism of $\mathfrak{g}_x = \mathfrak{sl}_3$ in $F(4)$ -case. This gives the first formula. For $D(2|1; a), F(4)$ one has $L_{\mathfrak{g}_x}(\nu)^* \cong L_{\mathfrak{g}_x}(\sigma(\nu))$; this gives the second formula. For $\mathfrak{g} = D(2|1; a), F(4)$ the Weyl group contains $-\text{Id}$, so $L \cong L^*$ and thus $DS_x(L) \cong DS_x(L^*)$ by 1.1. This implies the third formula. \square

2.2.3. We fix a triangular decomposition of \mathfrak{g}_0 and denote by Δ_0^+ the corresponding set of positive roots. We consider all bases Σ for Δ which satisfy $\Delta_0^+ \subset \Delta^+(\Sigma)$. We say that an isotropic root β is of the *first type* if β lies in a base Σ with $\Delta_0^+ \subset \Delta^+(\Sigma)$.

Take any base Σ as above and denote by ρ the corresponding Weyl vector. It is easy to see that a simple atypical module $L = L(\nu)$ satisfies the assumptions of Lemma 2.2.2 for some Σ' and $\beta \in \Sigma'$ if and only if $\nu + \rho$ is orthogonal to an isotropic root of the first type.

2.3. **Blocks of atypicality 1.** The blocks of atypicality 1 for basic classical Lie superalgebras were studied by J. Germoni in [9],[10] and by L. Martirosyan in [25]. These blocks satisfy the assumption of Proposition 2.1 and have the following extension graphs:

$$A_\infty \text{ for } \mathfrak{g} = \mathfrak{gl}(m|n), \mathfrak{osp}(2m|2n), F(4), D(2|1; a) \text{ for } a \in \mathbb{Q};$$

$$D_\infty \text{ for } \mathfrak{g} = \mathfrak{osp}(2m+1|2n), \mathfrak{osp}(2m|2n), F(4), G(3), D(2|1; a).$$

Let \mathfrak{g} be as in 2.2 and \mathcal{B} be an atypical block.

We call a block containing the trivial module $L(0)$ a *principal block*. Clearly, $\mathrm{DS}_x(L(0))$ is the trivial \mathfrak{g}_x -module, so 2.1 gives $\mathrm{DS}_x(L)$ for each $L \in \mathrm{Irr}(\mathcal{B}_0)$. For $\mathfrak{osp}(2|2)$, $\mathfrak{osp}(3|2)$ the principal block is the only atypical block.

Combining 2.1 and 2.2.2, 2.2.3 we see that in order to compute $\mathrm{DS}_x(L)$ for each L in $\mathrm{Irr}(\mathcal{B})$, it is enough to find $L(\nu) \in \mathrm{Irr}(\mathcal{B})$ such that $\nu + \rho$ is orthogonal to an isotropic root of the first type. Below we will list such ν for each non-principal atypical block in the remaining cases $D(2|1; a)$, $F(4)$ and $G(3)$.

2.3.1. *Case $D(2|1; a)$.* For $\mathfrak{g} := D(2|1; a)$ one has $\mathfrak{g}_x = \mathbb{C}$. The atypical blocks were described in [10], Thm. 3.1.1.

The extension graph of the principal block \mathcal{B}_0 is D_∞ , so for $L^i \in \mathrm{Irr}(\mathcal{B}_0)$ we have $\mathrm{DS}_x(L^i) = \mathbb{C}$ for $i = 0, 1$ and $\mathrm{DS}_x(L^i) = \Pi^{i-1}(\mathbb{C})^{\oplus 2}$ for $i > 1$ (where \mathbb{C} stands for the trivial even \mathfrak{g}_x -module).

If a is irrational, the principal block is the only atypical block in $\mathcal{F}(\mathfrak{g})$. Consider the case when a is rational. Recall that \mathfrak{h}^* has an orthogonal basis $\{\varepsilon_1, \varepsilon_2, \varepsilon_3\}$ with

$$\|\varepsilon_1\|^2 = \frac{a}{2}, \quad \|\varepsilon_2\|^2 = \frac{1}{2}, \quad \|\varepsilon_3\|^2 = -\frac{1+a}{2};$$

let $\varepsilon_1^*, \varepsilon_2^*, \varepsilon_3^*$ be the dual basis in \mathfrak{h} . The lattice $\Lambda_{2|1}$ is spanned by $\varepsilon_i s$; the parity map is given by $p(\varepsilon_1) = p(\varepsilon_2) = \bar{0}$, $p(\varepsilon_3) = \bar{1}$. One has

$$D(2|1; 1) = \mathfrak{osp}(4|2), \quad D(2|1; a) \cong D(2|1; -1 - a) \cong D(2|1; a^{-1})$$

so we can assume that $0 < a < 1$ and write $a = \frac{p}{q}$, where p, q are relatively prime positive integers.

The atypical blocks are \mathcal{B}_k for $k \in \mathbb{N}$ (the principal block is \mathcal{B}_0). Consider the block \mathcal{B}_k with $k > 0$. The extension graph of \mathcal{B}_k is A_∞^∞ . By [10], Thm. 3.1.1, the block \mathcal{B}_k contains a simple module L with the highest weight $\lambda_{k;0}$ satisfying $(\lambda_{k;0} + \rho|\beta) = 0$ for

$$\beta := -\varepsilon_1 + \varepsilon_2 + \varepsilon_3.$$

Taking $x \in \mathfrak{g}_\beta$ we can identify \mathfrak{g}_x with $\mathbb{C}h$ for $h := q\varepsilon_1^* + p\varepsilon_2^*$. Combining 2.2.2 and 2.2.3 we get

$$\mathrm{DS}_x(L) = L_{\mathfrak{g}_x}(k) \oplus L_{\mathfrak{g}_x}(-k),$$

where $L_{\mathfrak{g}_x}(u)$ stands for the even one-dimensional \mathfrak{g}_x -module with h acting by $u(p^2 + q^2)$. By Proposition 2.1, $\mathrm{DS}_x(L^i) \cong \Pi^i(\mathrm{DS}_x(L))$ for each $L^i \in \mathrm{Irr}(\mathcal{B}_k)$ (for $k > 0$).

2.3.2. *Case $G(3)$.* For $\mathfrak{g} := G(3)$ the atypical blocks were described in [10], Thm. 4.1.1. The atypical blocks in $\mathcal{F}(\mathfrak{g})$ are \mathcal{B}_k for $k \in \mathbb{N}$; the extension graphs are D_∞ . The block \mathcal{B}_k contains a simple module with the highest weight $\lambda_{k;0}$ satisfying $(\lambda_{k;0} + \rho|\beta) = 0$ for

$$\beta := -\varepsilon_1 + \delta.$$

Taking $\Sigma := \{\delta - \varepsilon_1, \varepsilon_2 - \delta, \delta\}$ and $x \in \mathfrak{g}_\beta$ we can identify \mathfrak{g}_x with \mathfrak{sl}_2 -triple corresponding to the root $\alpha = \varepsilon_1 + 2\varepsilon_2$. One has $\lambda_{k;0} = k\alpha$. Combining 2.2.2 and 2.1 we get

$$\mathrm{DS}_x(L^0) \cong \mathrm{DS}_x(L^1) \cong L_{\mathfrak{sl}_2}(2k), \quad \mathrm{DS}_x(L^i) = \Pi^{i-1}(L_{\mathfrak{sl}_2}(2k))^{\oplus 2} \quad \text{for } i > 1.$$

2.3.3. *Case $F(4)$.* For $\mathfrak{g} := F(4)$ we have $\mathfrak{g}_x \cong \mathfrak{sl}_3$. The integral weight lattice is spanned by $\varepsilon_1, \varepsilon_2, \frac{1}{2}(\varepsilon_1 + \varepsilon_2 + \varepsilon_3)$ and $\frac{1}{2}\delta$; the parity is given by $p(\frac{\varepsilon_i}{2}) = \bar{0}$ and $p(\frac{\delta}{2}) = \bar{1}$.

The atypical blocks are described in [25], Thm. 2.1. These blocks are parametrized by the pairs (m_1, m_2) , where $m_1, m_2 \in \mathbb{N}$, $m_1 - m_2 \in 3\mathbb{N}$. We denote the corresponding block by $\mathcal{B}_{(m_1; m_2)}$.

The extension graph of $\mathcal{B}_{(i; i)}$ is D_∞ ; the block $\mathcal{B}_{(0; 0)}$ is principal. For $i > 0$ the block $\mathcal{B}_{(i; i)}$ contains a simple module $L(\lambda)$ with

$$\lambda + \rho = (i + 1)(\varepsilon_1 + \varepsilon_2) - \beta_1, \quad \text{where } \beta_1 := \frac{1}{2}(-\varepsilon_1 + \varepsilon_2 - \varepsilon_3 + \delta).$$

One has $(\lambda + \rho | \beta_1) = 0$. Take $x \in \mathfrak{g}_{\beta_1}$ and consider the base

$$\Sigma_1 := \{\beta_1; \frac{1}{2}(\varepsilon_1 + \varepsilon_2 - \varepsilon_3 - \delta); \varepsilon_3; \varepsilon_1 - \varepsilon_2\}.$$

Then \mathfrak{g}_x can be identified with \mathfrak{sl}_3 corresponding to the set of simple roots $\{\varepsilon_2 + \varepsilon_3; \varepsilon_1 - \varepsilon_3\}$ and Lemma 2.2.2 gives

$$\mathrm{DS}_x(L(\lambda)) = L_{\mathfrak{sl}_3}(i\omega_1 + i\omega_2),$$

where ω_1, ω_2 are the fundamental weights of \mathfrak{sl}_3 . By 2.1 we get for $L^j \in \mathrm{Irr}(\mathcal{B}_{(i; i)})$:

$$\mathrm{DS}_x(L^0) \cong \mathrm{DS}_x(L^1) \cong L_{\mathfrak{sl}_3}(i\omega_1 + i\omega_2), \quad \mathrm{DS}_x(L^j) \cong \Pi^{j-1}(L_{\mathfrak{sl}_3}(i\omega_1 + i\omega_2))^{\oplus 2} \quad \text{for } j > 1.$$

Consider a block $\mathcal{B}_{(i_1; i_2)}$ for $i_1 \neq i_2$. The extension graph of this block is A_∞^∞ and this block contains a simple module $L := L(\lambda')$ with

$$\lambda' + \rho = i_1\varepsilon_1 + i_2\varepsilon_2 + (i_1 - i_2)\varepsilon_3.$$

In particular, $(\lambda' + \rho | \beta_2) = 0$ for $\beta_2 := \frac{1}{2}(-\varepsilon_1 + \varepsilon_2 + \varepsilon_3 + \delta)$. Taking $x \in \mathfrak{g}_{\beta_1}$ and

$$\Sigma_2 := \{\beta_2; \varepsilon_2 - \varepsilon_3; -\beta_1; \frac{1}{2}(\varepsilon_1 - \varepsilon_2 - \varepsilon_3 + \delta)\}$$

we identify \mathfrak{g}_x with \mathfrak{sl}_3 corresponding to the set of simple roots $\{\varepsilon_2 - \varepsilon_3; \varepsilon_1 + \varepsilon_3\}$. Combining Lemma 2.2.2 and 2.1 we get

$$\mathrm{DS}_x(L) = L_{\mathfrak{sl}_3}(i_1\omega_1 + i_2\omega_2) \oplus L_{\mathfrak{sl}_3}(i_2\omega_1 + i_1\omega_2), \quad \mathrm{DS}_x(L^i) \cong \Pi^i(\mathrm{DS}_x(L))$$

for each L^i in the block $\mathcal{B}_{(i_1; i_2)}$.

Corollary. *The image of the Grothendieck ring of $\mathcal{F}(F(4))$ under the homomorphism ds coincides with σ -invariants in the Grothendieck ring of $\mathcal{F}(\mathfrak{sl}_3)$.*

Proof. The condition $m_1 - m_2$ divisible by 3 is equivalent to $m_1\omega_1 + m_2\omega_2$ lies in the root lattice of \mathfrak{sl}_3 . \square

2.4. Conclusion. Let \mathfrak{t} be one of the superalgebras in 2.2 or one of Lie algebras $0, \mathbb{C}, \mathfrak{sl}_2$ or \mathfrak{sl}_3 . We introduce the map dex for \mathfrak{t} by

- for a typical $L \in \text{Irr}(\tilde{\mathcal{F}}(\mathfrak{t}))$ we take $\text{dex}(L) := 0$ for $L \in \text{Irr}(\mathcal{F}(\mathfrak{t}))$;
- for an atypical $L \in \text{Irr}(\tilde{\mathcal{F}}(\mathfrak{t}))$ we set $\text{dex}(L) := 0$ if $\text{DS}_x(L)$ is an even vector space.
- $\text{dex}(\Pi(L)) := \text{dex}(L) + 1 \pmod{2}$.

One readily sees that dex satisfies (Dex1) and (Dex2).

3. FUNCTORS $\Gamma_{\mathfrak{g}, \mathfrak{q}}^i$

In this section \mathfrak{g} is one of the superalgebras $F(4), G(3), \mathfrak{gl}(m|n), \mathfrak{osp}(m|2n)$ for $m, n \geq 0$. We fix any triangular decomposition $\Delta = \Delta^+ \amalg (-\Delta^+)$ and denote by \mathfrak{b} the corresponding Borel subalgebra. We consider the standard partial order $\nu_1 \leq \nu_2$ for $\nu_2 - \nu_1 \in \mathbb{N}\Delta^+$.

3.1. Let $\mathfrak{q} \subset \mathfrak{p}$ be parabolic subalgebras containing \mathfrak{b} and \mathfrak{l} be the Levy factor of \mathfrak{p} . For a finite-dimensional \mathfrak{q} -module V denote by $\Gamma_{\mathfrak{p}, \mathfrak{q}}(V)$ the maximal finite-dimensional quotient of the induced module $\mathcal{U}(\mathfrak{p}) \otimes_{\mathcal{U}(\mathfrak{q})} V$. We denote by $\tilde{\mathcal{F}}(\mathfrak{p})$ the category of finite-dimensional \mathfrak{p} -modules with the restriction lying in $\tilde{\mathcal{F}}(\mathfrak{l})$ and by $\text{Ext}_{\mathfrak{p}}^1$ the functor Ext^1 in this category. For $\lambda \in \Lambda_{m|n}$ we denote by $L_{\mathfrak{p}}(\lambda)$ a simple \mathfrak{p} -module of the highest weight λ with the grading induced by the parity function on $\Lambda_{m|n}$.

In [18], Sect. 3 the authors introduce for $i = 0, 1, \dots$ an additive functor

$$\Gamma_{\mathfrak{g}, \mathfrak{p}}^i : \tilde{\mathcal{F}}(\mathfrak{p}) \rightarrow \tilde{\mathcal{F}}(\mathfrak{g})$$

(in [18] these functors are denoted by $\Gamma_i(G/P; -)$) in the following way. For each $V \in \tilde{\mathcal{F}}(\mathfrak{q})$ we take the vector bundle $G \times_P V$ over the generalized Grassmanian G/P and consider the cohomology groups $H^i(G/P, G \times_P V)$ as \mathfrak{g} -module. We set

$$\Gamma_{\mathfrak{g}, \mathfrak{p}}^i(V) := (H^i(G/P, G \times_P V^*))^*.$$

Below we recall several properties of the functors $\Gamma_{\mathfrak{g}, \mathfrak{p}}^i$; for the proofs and other properties see [18], Sections 3, 4.

3.1.1. One has $\Gamma_{\mathfrak{g}, \mathfrak{p}}^0(V) = \Gamma_{\mathfrak{g}, \mathfrak{p}}(V)$. For each i the module $\Gamma_{\mathfrak{g}, \mathfrak{p}}^i(L_{\mathfrak{p}}(\lambda))$ has the same central character as $L(\lambda)$.

3.1.2. Each short exact sequence of \mathfrak{q} -modules

$$0 \rightarrow U \rightarrow V \rightarrow U' \rightarrow 0$$

induces a long exact sequence

$$\dots \rightarrow \Gamma_{\mathfrak{g}, \mathfrak{q}}^1(V) \rightarrow \Gamma_{\mathfrak{g}, \mathfrak{q}}^1(U') \rightarrow \Gamma_{\mathfrak{g}, \mathfrak{q}}^0(U) \rightarrow \Gamma_{\mathfrak{g}, \mathfrak{q}}^0(V) \rightarrow \Gamma_{\mathfrak{g}, \mathfrak{q}}^0(U') \rightarrow 0.$$

3.1.3. If $[\Gamma_{\mathfrak{g},\mathfrak{p}}^i(L_{\mathfrak{p}}(\lambda)) : L(\nu)] \neq 0$, then there exist $I \subset \Delta_1^+$ and $w \in W$ of length i such that $\nu + \rho = w(\lambda + \rho) - \sum_{\alpha \in I} \alpha$.

3.2. **Poincaré polynomials.** Let $\mathfrak{h} \subset \mathfrak{q} \subset \mathfrak{p}$ be as in 3.1. We set

$$\mathfrak{q}' := \mathfrak{q} \cap \mathfrak{l}, \quad \mathfrak{h}' := \mathfrak{h} \cap \mathfrak{l}, \quad \mathfrak{h}'' = \{h \in \mathfrak{h} \mid [h, \mathfrak{l}] = 0\}$$

and notice that $\mathfrak{h} = \mathfrak{h}' \oplus \mathfrak{h}''$. Let $\lambda, \mu \in \Lambda_{m|n}$ be such that $L_{\mathfrak{q}}(\lambda) \in \mathcal{F}(\mathfrak{q})$ and $L_{\mathfrak{p}}(\mu) \in \mathcal{F}(\mathfrak{p})$. For $i = 0, 1, \dots$ we define

$${}^i K_{\mathfrak{p},\mathfrak{q}}^{\lambda,\mu} := \begin{cases} 0 & \text{if } \lambda|_{\mathfrak{h}''} \neq \nu|_{\mathfrak{h}''}; \\ [\Gamma_{\mathfrak{l},\mathfrak{q}'}^i(L_{\mathfrak{q}'}(\lambda|_{\mathfrak{h}'})) : L_{\mathfrak{l}}(\mu|_{\mathfrak{h}'})] & \text{if } \lambda|_{\mathfrak{h}''} = \nu|_{\mathfrak{h}''} \end{cases}$$

and introduce a Poincaré polynomial in the variable z by

$$K_{\mathfrak{p},\mathfrak{q}}^{\lambda,\mu}(z) := \sum_{i=0}^{\infty} {}^i K_{\mathfrak{p},\mathfrak{q}}^{\lambda,\mu} z^i.$$

(It is easy to see that this Poincaré polynomial is equal to the Poincaré polynomial defined in [18], Section 4.) When the term $K_{\mathfrak{p},\mathfrak{q}}^{\lambda,\mu}$ appears in a formula it is always assumed that $L_{\mathfrak{q}}(\lambda) \in \mathcal{F}(\mathfrak{q})$ and $L_{\mathfrak{p}}(\mu) \in \mathcal{F}(\mathfrak{p})$.

3.2.1. By 3.1.1 we have

$$[\Gamma_{\mathfrak{p},\mathfrak{q}}(L_{\mathfrak{q}}(\lambda)) : L_{\mathfrak{p}}(\mu)] = {}^0 K_{\mathfrak{p},\mathfrak{q}}^{\lambda,\mu} = K_{\mathfrak{p},\mathfrak{q}}^{\lambda,\mu}(0).$$

In particular, $K_{\mathfrak{p},\mathfrak{q}}^{\lambda,\lambda}(0) = 1$ and $K_{\mathfrak{p},\mathfrak{q}}^{\lambda,\mu}(0) \neq 0$ implies $\mu \leq \lambda$. By [18], Thm. 1 one has

$$(8) \quad K_{\mathfrak{g},\mathfrak{q}}^{\lambda,\mu}(-1) = \sum_{\nu} K_{\mathfrak{p},\mathfrak{q}}^{\lambda,\nu}(-1) K_{\mathfrak{g},\mathfrak{p}}^{\nu,\mu}(-1)$$

where the summation is taken on $\nu \in \mathfrak{h}^*$ with $\dim L_{\mathfrak{p}}(\nu) < \infty$.

3.3. **Euler characteristic formula.** Let ρ be the Weyl vector and R be the Weyl denominator, i.e.

$$2\rho = \sum_{\alpha \in \Delta^+} (-1)^{p(\alpha)} \alpha, \quad R = \prod_{\alpha \in \Delta^+} (1 - (-1)^{p(\alpha)} e^{-\alpha})^{(-1)^{p(\alpha)}}.$$

We denote by $\text{sgn} : W \rightarrow \mathbb{Z}_2$ the standard sign homomorphism and set

$$\mathcal{E}_{\lambda,\mathfrak{p}} := R^{-1} e^{-\rho} \sum_{w \in W} \text{sgn}(w) w \left(\frac{e^{\rho} \text{ch } L_{\mathfrak{p}}(\lambda)}{\prod_{\alpha \in \Delta_1^+(\mathfrak{l})} (1 + e^{-\alpha})} \right).$$

By [18], Prop.1, if $L_{\mathfrak{p}}(\lambda) \in \mathcal{F}(\mathfrak{p})$, then

$$\sum_{\mu} K_{\mathfrak{g},\mathfrak{p}}^{\lambda,\mu}(-1) \text{ch } L(\mu) = \mathcal{E}_{\lambda,\mathfrak{p}}.$$

3.3.1. Notice that $\mathcal{E}_{\lambda, \mathfrak{p}}$ can be zero. For instance, take $\mathfrak{g} = \mathfrak{osp}(m|2n)$ with $m \geq 4$ and \mathfrak{b} corresponding to the “mixed base”. Then $\mathcal{E}_{0, \mathfrak{b}} = R^{-1} e^{-\rho} \sum \text{sgn}(w) e^{w\rho} = 0$. Since $\text{ch } L(\mu)$ are linearly independent, we have $K_{\mathfrak{g}, \mathfrak{b}}^{0, \mu}(-1) = 0$ for all μ .

3.4. **Marked graphs.** Consider a directed graph (V, E) where V is at most countable and the number of edges between any two vertices is finite.

We say that $\iota : V \rightarrow \mathbb{N}$ (resp., $\iota : V \rightarrow \mathbb{Z}$) defines a \mathbb{N} -grading (resp., \mathbb{Z} -grading) on this graph if for each edge $\nu \xrightarrow{e} \lambda$ one has $\iota(\nu) < \iota(\lambda)$. Notice that for a \mathbb{Z} -graded graph the number of paths between any two vertices is finite.

Assume that the set of edges E is equipped by two functions b and κ , where $b : E \rightarrow \mathbb{Z}$ and κ is a function from E to a commutative ring.

3.4.1. For a path $P := \nu_1 \xrightarrow{e_1} \nu_2 \xrightarrow{e_2} \nu_3 \dots \xrightarrow{e_s} \nu_{s+1}$ we define

$$\text{length}(P) := s, \quad \kappa(P) := \prod_{i=1}^s \kappa(e_i).$$

We call the path P *b-decreasing* (resp., *b-increasing*) if $b(e_1) > b(e_2) > \dots > b(e_s)$ (resp., $b(e_1) < \dots < b(e_s)$). We consider $P = \nu$ as a *b-decreasing/increasing* path of zero length with $\kappa(P) = 1$. We denote the set of decreasing (resp., increasing) paths from ν to λ by $\mathcal{P}_b^>(\nu, \lambda)$ (resp., $\mathcal{P}_b^<(\nu, \lambda)$).

3.4.2. **Definition.** We call two functions $b, b' : E \rightarrow \mathbb{Z}$ *decreasingly-equivalent* if for each path $\nu_1 \xrightarrow{e_1} \nu_2 \xrightarrow{e_2} \nu_3$ one has

$$b(e_1) > b(e_2) \iff b'(e_1) > b'(e_2).$$

Notice that b, b' are decreasingly equivalent if and only if $\mathcal{P}_b^>(\nu, \lambda) = \mathcal{P}_{b'}^>(\nu, \lambda)$.

3.4.3. Let (V, E) be a \mathbb{Z} -graded graph. We introduce the square matrices $A^<(\kappa) = (a_{\lambda, \nu}^<)_{\lambda, \nu \in V}$ and $A^>(\kappa) = (a_{\lambda, \nu}^>)_{\lambda, \nu \in V}$ by

$$a_{\lambda, \nu}^> := \sum_{P \in \mathcal{P}_b^>(\nu, \lambda)} \kappa(P), \quad a_{\lambda, \nu}^< := \sum_{P \in \mathcal{P}_b^<(\nu, \lambda)} (-1)^{\text{length}(P)} \kappa(P).$$

Since the graph is a \mathbb{Z} -graded, these matrices are lower-triangular with $a_{\lambda, \lambda}^> = a_{\lambda, \lambda}^< = 1$.

3.4.4. **Lemma.** *Let (V, E) be a \mathbb{Z} -graded graph with a finite number of edges between any two vertices. Assume that $b : E \rightarrow \mathbb{Z}$ satisfies the property*

$$(BB) \quad \text{for each path } \nu_1 \xrightarrow{e_1} \nu_2 \xrightarrow{e_2} \nu_3 \text{ one has } b(e_1) \neq b(e_2).$$

Then $A^>(\kappa) \cdot A^<(\kappa) = A^<(\kappa) \cdot A^>(\kappa) = \text{Id}$.

Proof. The proof is similar to [18], Thm. 4. The entries of $A^{<}(\kappa) \cdot A^{>}(\kappa)$ are of the form

$$\sum_{\mu} \sum_{P \in \mathcal{P}_b^{<}(\nu, \mu)} \sum_{Q \in \mathcal{P}_b^{>}(\mu, \lambda)} (-1)^{\text{length}(Q)} \kappa(PQ),$$

where PQ stands for the concatenation of P and Q . The property (BB) implies that each path of non-zero length which can be presented as the concatenation PQ , where P is b -increasing and Q is b -decreasing, has exactly two presentations of this form: $PQ = P'Q'$ with $\text{length } Q' = \text{length } Q \pm 1$ (for instance, for a path of length 5 with

$$b(e_1) = 1, \quad b(e_2) = 2, \quad b(e_3) = 4, \quad b(e_4) = 2, \quad b(e_5) = 1$$

the increasing part can be either e_1, e_2 or e_1, e_2, e_3). This implies the statement. \square

3.5. Useful graphs. We fix a sequence of parabolic subalgebras in \mathfrak{g} :

$$(9) \quad \mathfrak{b} = \mathfrak{p}^{(0)} \subset \mathfrak{p}^{(1)} \subset \dots \subset \mathfrak{p}^{(k)} = \mathfrak{g}$$

and denote by $\mathfrak{l}^{(p)}$ the Levy subalgebra of $\mathfrak{p}^{(p)}$. We also fix a central character $\chi : \mathcal{Z}(\mathfrak{g}) \rightarrow \mathbb{C}$ and denote by Λ^χ the set of dominant weights corresponding to χ :

$$\Lambda^\chi := \{\lambda \in \Lambda_{m|n} \mid \dim L(\lambda) < \infty \text{ and } \text{Ann}_{\mathcal{Z}(\mathfrak{g})} L(\lambda) = \text{Ker } \chi\}.$$

For each $\lambda \in \Lambda^\chi$ we denote by $\text{tail}(\lambda)$ the maximal s such that $\lambda|_{\mathfrak{l}^{(s)} \cap \mathfrak{h}} = 0$. We set

$$\mathfrak{p}_\lambda := \mathfrak{p}^{\text{tail}(\lambda)}, \quad \mathfrak{l}_\lambda := \mathfrak{l}^{\text{tail}(\lambda)}.$$

3.5.1. Graph $\hat{\Gamma}^\chi$. Let $\hat{\Gamma}^\chi(z)$ be a graph with the set of vertices $V := \Lambda^\chi$ and the following edges: if $K_{\mathfrak{p}^{(s)}, \mathfrak{p}^{(s-1)}}^{\lambda, \nu} \neq \delta_{\nu, \lambda}$ (where $\delta_{\nu, \lambda}$ is the Kronecker symbol) we join ν, λ by the edge of the form

$$\nu \xrightarrow{e} \lambda \text{ with } b(e) = s, \quad \kappa(e) := K_{\mathfrak{p}^{(s)}, \mathfrak{p}^{(s-1)}}^{\lambda, \nu}(z) - \delta_{\nu, \lambda} \in \mathbb{Z}[z].$$

Note that each two vertices in $\hat{\Gamma}^\chi$ are connected by at most k edges.

For each $z_0 \in \mathbb{C}$ we denote by $\hat{\Gamma}^\chi(z_0)$ the subgraph where the edges with $\kappa(e)(z_0) = 0$ are deleted and the function $\kappa_{z_0} : E \rightarrow \mathbb{C}$ is given by

$$\kappa_{z_0}(e) := \kappa(e)(z_0).$$

Note that $\hat{\Gamma}^\chi(0)$ does not have loops, see 3.2.1.

3.5.2. Graph Γ^χ . Let Γ^χ (resp., $\Gamma^\chi(z_0)$) be the graph obtained from $\hat{\Gamma}^\chi$ (resp., from $\hat{\Gamma}^\chi(z_0)$) by removing the edges of the form $\nu \xrightarrow{e} \lambda$ with $b(e) \leq \text{tail}(\lambda)$. Thus we have

$$\text{tail}(\lambda) < b(e) \text{ for each edge } \nu \xrightarrow{e} \lambda \text{ in } \Gamma^\chi.$$

We will always assume that Γ^χ satisfies the following condition:

$$(Tail) \quad \text{tail}(\nu) \leq b(e) \text{ for each edge } \nu \xrightarrow{e} \lambda \text{ in } \Gamma^\chi$$

which is tautological for $k = 1$.

3.5.3. *Notation.* We denote by $\mathcal{P}_b^>(\nu, \lambda)$ (resp., by $\hat{\mathcal{P}}_b^>(\nu, \lambda)$) the set of b -decreasing paths from ν to λ in the graph Γ^x (resp., $\hat{\Gamma}^x$).

3.5.4. **Corollary.** *Take $\lambda, \nu \in \Lambda^x$.*

(i) *Assume that for each $\mu \in \Lambda^x$ one has*

$$(\exists i \text{ s.t. } K_{\mathfrak{p}^{(i+1)}, \mathfrak{p}^{(i)}}^{\mu, \eta}(z) \neq 0) \implies \eta \in \Lambda^x.$$

Then

$$K_{\mathfrak{g}, \mathfrak{b}}^{\lambda, \nu}(-1) = \sum_{P \in \hat{\mathcal{P}}_b^>(\nu, \lambda)} \kappa_{-1}(P).$$

(ii) *Assume that Γ^x satisfies (Tail) and that for each $\mu \in \Lambda^x$ one has*

$$(\exists i \geq \text{tail}(\mu) \text{ s.t. } K_{\mathfrak{p}^{(i+1)}, \mathfrak{p}^{(i)}}^{\mu, \eta}(z) \neq 0) \implies \eta \in \Lambda^x.$$

Then
$$K_{\mathfrak{g}, \mathfrak{p}_\lambda}^{\lambda, \nu}(-1) = \sum_{P \in \mathcal{P}_b^>(\nu, \lambda)} \kappa_{-1}(P).$$

Proof. The assertions follow from the formula (8). □

3.5.5. *Remark.* The graph Γ^x is useful for character formulae. Retain notation of 3.3 and notice that $\mathcal{E}_{\lambda, \mathfrak{p}}$ has a particularly nice formula if $\lambda|_{\mathfrak{l} \cap \mathfrak{h}} = 0$ (in this case $\text{ch } L_{\mathfrak{p}}(\lambda) = e^\lambda$). Thus it makes sense to express $\text{ch } L(\mu)$ in terms of $\mathcal{E}_{\lambda, \mathfrak{p}^{(j)}}$ for $j \leq \text{tail}(\lambda)$. By 3.3.1, $\mathcal{E}_{\lambda, \mathfrak{p}}$ can be zero if “ \mathfrak{p} is too small”; thus it makes sense to consider the maximal “nice” \mathfrak{p} for each λ , which is \mathfrak{p}_λ .

3.5.6. **Lemma.** *If the zero weight is a minimal dominant weight for $\mathfrak{l}^{(p)}$ for each p , then $\hat{\Gamma}^x(0) = \Gamma^x(0)$.*

Proof. By 3.1.1, as a $\mathfrak{l}^{(p)}$ -module $\Gamma_{\mathfrak{p}^{(p)}, \mathfrak{p}^{(p-1)}}^0(L_{\mathfrak{p}^{(p-1)}}(\lambda))$ is a finite-dimensional quotient of the Verma $\mathfrak{l}^{(p)}$ -module $M_{\mathfrak{l}^{(p)}}(\lambda)$. If the zero weight is a minimal dominant weight for $\mathfrak{l}^{(p)}$, then for each $p \leq \text{tail}(\lambda)$ the module $L_{\mathfrak{p}^{(p)}}(\lambda)$ is a unique finite-dimensional quotient of $M_{\mathfrak{l}^{(p)}}(\lambda)$, so $\Gamma_{\mathfrak{p}^{(p)}, \mathfrak{p}^{(p-1)}}^0(L_{\mathfrak{p}^{(p-1)}}(\lambda)) = L_{\mathfrak{p}^{(p)}}(\lambda)$. Therefore $\hat{\Gamma}^x(0) = \Gamma^x(0)$ as required. □

3.5.7. *Definitions.* Let $\text{dex} : V \rightarrow \mathbb{Z}_2 = \{0, 1\}$ be any map.

We say that $\Gamma^x(0)$ (resp., of $\hat{\Gamma}^x(0)$) is bipartite *with respect to* dex if for each edge $\nu \xrightarrow{e} \lambda$ in this graph $\text{dex}(\lambda) \neq \text{dex}(\nu)$.

Recall that $\kappa(e)$ is a polynomial with non-negative integral coefficients. We say that dex gives a *parametric bipartition* of (Γ^x, κ) if for each edge $\nu \xrightarrow{e} \lambda$ one has

$$(10) \quad z^{\text{dex}(\lambda) - \text{dex}(\nu) + 1} \kappa(e) \in \mathbb{Z}[z^2].$$

We say that dex gives a *signed bipartition* of $(\Gamma^\chi(-1), \kappa_{-1})$ if for each edge $\nu \xrightarrow{e} \lambda$

$$(11) \quad (-1)^{\text{dex}(\lambda) - \text{dex}(\nu) + 1} \kappa_{-1}(e) \in \mathbb{Z}_{\geq 0}$$

or, equivalently,

$$(12) \quad (-1)^{\text{dex}(\lambda) - \text{dex}(\nu)} (-1)^{\text{length}(P)} \kappa_{-1}(P) \in \mathbb{Z}_{\geq 0} \quad \text{for each path } P \text{ from } \nu \text{ to } \lambda.$$

We say that dex gives a *parametric bipartition* of $(\hat{\Gamma}^\chi, \kappa)$ if each edge $\nu \xrightarrow{e} \lambda$ with $\nu \neq \lambda$ satisfies (10). Similarly, we say that dex gives a *signed bipartition* of $(\hat{\Gamma}^\chi(-1), \kappa_{-1})$ if (11) holds for each $\nu \xrightarrow{e} \lambda$ with $\nu \neq \lambda$ or, equivalently if (12) holds for any paths without loops.

3.5.8. Recall that $\hat{\Gamma}^\chi(0)$ does not have loops. If dex is a parametric bipartition of Γ^χ (resp., $\hat{\Gamma}^\chi$), then dex is a signed bipartition of $(\Gamma^\chi(-1), \kappa_{-1})$ (resp., of $(\hat{\Gamma}^\chi(-1), \kappa_{-1})$) and $\Gamma^\chi(0)$ (resp., $\hat{\Gamma}^\chi$) is bipartite with respect to dex .

3.5.9. *Remark.* In the examples 3.9–3.9.5 below Γ^χ is a \mathbb{Z} -graded graph admitting a parametric bipartition; the same is true for the graph $\hat{\Gamma}^\chi$ for the dense flag for a distinguished Borel in $\mathfrak{gl}(n|n)$ -case, see [27]. The graph $\hat{\Gamma}^\chi$ has a loop $0 \xrightarrow{e} 0$ (and thus is not \mathbb{Z} -graded) for the dense flag for a mixed Borel for $\mathfrak{g} = \mathfrak{osp}(2|2)$ ($= \mathfrak{sl}(1|2)$) and for $\mathfrak{g} = \mathfrak{osp}(4|2)$ see 3.9.2 and 3.9.5. By [18], Lemma 26, $\kappa(e) = z$ for $\mathfrak{osp}(2|2)$ and $\kappa(e) = z^2$ for $\mathfrak{osp}(4|2)$ so the formula (10) holds for e if $\mathfrak{g} = \mathfrak{osp}(2|2)$ and does not hold if $\mathfrak{g} = \mathfrak{osp}(4|2)$. The graph $\hat{\Gamma}^\chi$ admits a parametric bipartition in both cases.

3.6. **Graph** $\text{Ext}(\chi)$. By 6.3 $\dim \text{Ext}^1(L(\lambda), L(\nu)) = \dim \text{Ext}^1(L(\nu), L(\lambda))$. We denote by $\text{Ext}(\chi)$ the graph without loops, with the set of vertices Λ^χ and $\dim \text{Ext}^1(L(\lambda), L(\nu))$ edges between ν and λ for $\nu \neq \lambda$ (we will usually consider the undirected edges).

3.6.1. We say that $\text{Ext}(\chi)$ is a subgraph of a directed graph if $\text{Ext}(\chi)$ is a subgraph of the “undirected version” of this graph (we forget the directions of edges).

3.6.2. Recall that $\Gamma_{\mathfrak{p}^{(p)}, \mathfrak{p}^{(p-1)}}^0(L_{\mathfrak{p}^{(p-1)}}(\lambda))$ is the maximal finite-dimensional quotient of $\text{Ind}_{\mathfrak{p}^{(p-1)}}^{\mathfrak{p}^{(p)}}(L_{\mathfrak{p}^{(p-1)}}(\lambda))$; this is indecomposable module with the cosocle is isomorphic to $L_{\mathfrak{p}^{(p)}}(\lambda)$. Using Corollary 6.5 we obtain the

Corollary. *The graph $\text{Ext}(\chi)$ is a subgraph of $\hat{\Gamma}^\chi(0)$.*

3.6.3. **Corollary.** *Assume that $\hat{\Gamma}^\chi$ admits a parametric bipartition dex .*

- (i) $\text{Ext}(\chi)$ is bipartite with respect to dex .
- (ii) $\Gamma_{\mathfrak{p}^{(p)}, \mathfrak{p}^{(p-1)}}^i(L_{\mathfrak{p}^{(p-1)}}(\lambda))$ is a semisimple $\mathfrak{p}^{(p)}$ -module for $i > 0$ and has a semisimple radical for $i = 0$;

- (iii) Assume that Γ^χ admits a parametric bipartition dex and that the zero weight is a minimal dominant weight for $\mathfrak{l}^{(p)}$ for each $p = 0, 1, \dots, k-1$. Then $\text{Ext}(\chi)$ is a subgraph of $\Gamma^\chi(0)$ and dex defines a bipartition of $\text{Ext}(\chi)$. Moreover, the claims of (ii) hold for $p > \text{tail}(\lambda)$.

Proof. Corollary 3.6.2 implies (i). For (ii) let $\text{Ext}_{(p)}$ be the “Ext”-graph for $\mathfrak{p}^{(p)}$: the set of vertices for this graph is Λ^χ and the multiplicity of the edge $\lambda \rightarrow \nu$ is $\dim \text{Ext}^1(L_{\mathfrak{p}^{(p)}}(\lambda), L_{\mathfrak{p}^{(p)}}(\nu))$. By Corollary 6.5, $\text{Ext}_{(p)}$ is a subgraph of $\hat{\Gamma}^\chi(0)$, so dex gives a bipartition on $\text{Ext}_{(p)}$. For $i > 0$ one has

$$[\Gamma_{\mathfrak{p}^{(p)}, \mathfrak{p}^{(p-1)}}^i(L_{\mathfrak{p}^{(p-1)}}(\lambda)) : L_{\mathfrak{p}^{(p)}}(\nu)] \neq 0 \implies \text{dex}(\nu) + \text{dex}(\lambda) \equiv i + 1 \pmod{2}.$$

Therefore there are no non-splitting extensions between the subquotients of the $\mathfrak{p}^{(p)}$ -module $\Gamma_{\mathfrak{p}^{(p)}, \mathfrak{p}^{(p-1)}}^i(L_{\mathfrak{p}^{(p-1)}}(\lambda))$; thus this module is completely reducible. For $i = 0$ the same holds for $\nu \neq \lambda$ and $L_{\mathfrak{p}^{(p)}}(\lambda)$ is a unique simple quotient of $\Gamma_{\mathfrak{p}^{(p)}, \mathfrak{p}^{(p-1)}}^0(L_{\mathfrak{p}^{(p-1)}}(\lambda))$. Hence the radical of $\Gamma_{\mathfrak{p}^{(p)}, \mathfrak{p}^{(p-1)}}^0(L_{\mathfrak{p}^{(p-1)}}(\lambda))$ is semisimple. This gives (ii). If the zero weight is a minimal dominant weight for $\mathfrak{l}^{(p)}$ for each p , then $\tilde{\Gamma}^\chi(0) = \Gamma^\chi(0)$ (see Lemma 3.5.6) and so (iii) has the same proof as (ii). \square

3.6.4. *Remark.* We see that in order to have a parametric partition on $\hat{\Gamma}^\chi$ one has to take a “dense enough” chain of the parabolic subalgebras, since if $\hat{\Gamma}^\chi$ admits such grading, then the maximal finite-dimensional quotient of $\text{Ind}_{\mathfrak{p}^{(p-1)}}^{\mathfrak{p}^{(p)}}(L_{\mathfrak{p}^{(p-1)}}(\lambda))$ has a Loewy filtration of length ≤ 2 . In the examples below we take $\mathfrak{l}^{(p)}$ of the defect p .

3.7. **The Gruson-Serganova algorithm.** We assume that $\Gamma^\chi = (\Lambda^\chi, E)$ is a \mathbb{Z} -graded graph which satisfies the assumptions of Corollary 3.5.4 (ii). The following construction is a slight reformulation of the construction described in [18], Sect. 12.

3.7.1. Recall that $\text{ch } L_{\mathfrak{p}^\lambda}(\lambda) = e^\lambda$. Set

$$(13) \quad \mathcal{E}_\nu := \mathcal{E}_{\nu, \mathfrak{p}^\nu} = R^{-1} e^{-\rho} \sum_{w \in W} \text{sgn}(w) w \left(\frac{e^{\nu+\rho}}{\prod_{\alpha \in \Delta(\mathfrak{l}_\nu)^+} (1 + e^{-\alpha})} \right).$$

Combining 3.3 and Corollary 3.5.4 (ii) we get

$$\sum_{\mu} a_{\lambda, \mu}^> \text{ch } L(\mu) = \mathcal{E}_\lambda,$$

for $A^>(\kappa_{-1}) = (a_{\lambda, \nu}^>)$ defined as in 3.4.3. The matrix $A := A^>(\kappa_{-1})$ is lower-triangular with $a_{\lambda, \lambda} = 1$. Thus A is invertible that is

$$\text{ch } L(\lambda) = \sum_{\mu} a'_{\lambda, \mu} \mathcal{E}_\mu,$$

for $(a'_{\lambda,\mu}) := A^{-1}$. In the light of Lemma 3.4.4 the entries of A^{-1} can be expressed in terms of b -increasing paths from ν to λ if $b : E \rightarrow \mathbb{Z}$ would satisfy the property (BB). Unfortunately, b almost never satisfy (BB); however, it is often possible to find a decreasingly-equivalent function b' satisfying (BB) (we do not require that b' satisfies (Tail)). For $\mathfrak{gl}(M|N)$ and $\mathfrak{osp}(M|N)$ the function b' is given in [27] and [18] respectively; in 4.4.3 below we describe b' for $\mathfrak{osp}(M|N)$ -case. Denoting by $\mathcal{P}_{b'}^<(\nu, \lambda)$ the set of b' -increasing paths in Γ^x we obtain

$$a_{\lambda,\mu}^> = \sum_{P \in \mathcal{P}_b^>(\mu, \lambda)} \kappa_{-1}(P) = \sum_{P \in \mathcal{P}_{b'}^>(\mu, \lambda)} \kappa_{-1}(P), \quad a'_{\lambda,\mu} = \sum_{P \in \mathcal{P}_{b'}^<(\mu, \lambda)} (-1)^{\text{length}(P)} \kappa_{-1}(P).$$

3.7.2. Assume that $\text{dex} : V \rightarrow \{0, 1\}$ is a signed bipartition of $(\Gamma^x(-1), \kappa_{-1})$ (see 3.5.7 for definition). By (12) the number $(-1)^{\text{dex}(\lambda) - \text{dex}(\mu)} a'_{\lambda,\mu}$ is a non-negative integer. (i.e. the Gruson-Serganova character formula is “positive”). These number can be interpreted as follows. Consider the following modification of the graph $\Gamma^x = (V, E)$: the graph D^x with the same set of vertices $V = \Lambda^x$ and the set of edges E' obtained from E by taking each edge $\nu \xrightarrow{e} \lambda$ with the multiplicity $(-1)^{\text{dex}(\lambda) - \text{dex}(\nu)} \kappa_{-1}(e)$ (this number is non-negative since dex is a signed bipartition). By above,

$$(-1)^{\text{dex}(\lambda) - \text{dex}(\mu)} a'_{\lambda,\mu} \text{ is the number of } b'\text{-increasing paths from } \mu \text{ to } \lambda \text{ in } D^x.$$

For $\mathfrak{osp}(M|N)$ -case the graph D^x is described in [18]; we give some details in 4.5 below; the case $\mathfrak{gl}(m|n)$ will be treated in [14].

3.7.3. The assumption that Γ^x is \mathbb{Z} -graded can be weakened using the following trick. Fix a set of “bad vertices” $\Lambda' \subset \Lambda(\chi)$ and consider a graph $\Gamma'(\chi)$ obtained from Γ^x by erasing all edges ending at $\lambda \in \Lambda'$. Assume that $\Gamma'(\chi)$ is \mathbb{Z} -graded. The above reasoning allows to express $\text{ch } L(\mu)$ in terms of \mathcal{E}_λ for $\lambda \in \Lambda(\chi) \setminus \Lambda'$ and $\text{ch } L(\nu)$ for $\nu \in \Lambda'$, see 3.9.5 for examples.

3.8. **Examples.** The Poincaré polynomials for certain chains of parabolic subalgebras were computed for the finite-dimensional Kac-Moody superalgebras in [32], [27], [10],[25] and for \mathfrak{q}_n in [30]. In all these cases the chain satisfies the following condition: $\mathfrak{l}^{(p)}$ has defect p . Below we list some properties of the corresponding graphs (for the \mathfrak{gl} -case we consider only the principal block in $\mathfrak{gl}(n|n)$).

In all these examples the Poincaré polynomials have the following property: the polynomial $K_{\mathfrak{p}^{(s)}, \mathfrak{p}^{(s-1)}}^{\lambda, \mu} - \delta_{\lambda, \mu}$ is non-zero for at most one value of $s > \text{tail}(\lambda)$. We denote by $\kappa^{\lambda, \mu}$ the corresponding non-zero polynomial (if it exists) and set $\kappa^{\lambda, \mu} = 0$ otherwise.

The above property implies that Γ^x does not have multi-edges (and that $\kappa(\mu \rightarrow \lambda) = \kappa^{\lambda, \mu}$). The graph Γ^x admits a \mathbb{N} -grading and satisfies (Tail).

For $\mathfrak{g} \neq \mathfrak{q}_n$ the graph Γ^x admits a parametric bipartition dex .

For $\mathfrak{g} \neq \mathfrak{q}_n$, $\mathfrak{osp}(2m|2n)$, the polynomials $\kappa^{\lambda,\mu}$ are monomials, so the condition on the parametric bipartition simply means that $\kappa^{\lambda,\mu}$ is zero or z^i for $i \equiv \text{dex}(\lambda) - \text{dex}(\mu) + 1$ modulo 2. In these cases $D^\chi = \Gamma^\chi$.

For $\mathfrak{g} = \mathfrak{osp}(2m|2n)$, \mathfrak{q}_n with χ of atypicality greater than one, $\kappa^{\lambda,\mu} \in \{0, z^i, z^i + z^j\}$ and the condition on the parametric bipartition takes the form $i \equiv \text{dex}(\lambda) - \text{dex}(\mu) + 1$ and $i \equiv j$ modulo 2. This holds for $\mathfrak{osp}(2m|2n)$ in this case D^χ is obtained from Γ^χ by doubling the edges with $\kappa^{\lambda,\mu} = z^i + z^j$.

An interesting example is the \mathfrak{q}_n -case. For χ of atypicality greater than one, $i - j$ can be odd, so Γ^χ does not admit a parametric bipartition and $\Gamma^\chi(0)$ is not bipartited. By [26] the Ext-graph is not bipartite. However, the $(\Gamma^\chi(-1), \kappa_{-1})$ admits a signed bipartition dex. In this case D^χ is obtained from Γ^χ by doubling the edges with even j and deleting the edges with odd j ($\kappa^{\lambda,\mu}(-1) = 0$ if j is odd). The Gruson-Serganova algorithm gives the Su-Zhang character formula [35].

3.9. Examples of defect one. We start from the examples when \mathfrak{g} has defect 1 and the chain is $\mathfrak{b} = \mathfrak{p}^{(0)} \subset \mathfrak{p}^{(1)} = \mathfrak{g}$.

In this case $b(e) = 1$ for each $e \in \hat{\Gamma}^\chi$, so the condition (BB) does not hold if Γ^χ contains paths of length two. Moreover, the decreasing paths in Γ^χ are the paths containing at most one edge. Thus any function $b' : E \rightarrow \mathbb{Z}$ satisfying $b'(e_1) < b'(e_2)$ for each path $\cdot \xrightarrow{e_1} \cdot \xrightarrow{e_2} \cdot$ in Γ^χ is decreasingly equivalent to b and satisfies (BB).

We will depict an edge e in Γ^χ (resp., in $\hat{\Gamma}^\chi$) as $\nu \xrightarrow{j;\kappa(e)} \lambda$ (resp., as $\nu \xrightarrow{\kappa(e)} \lambda$) with $j = b'(e)$.

Except for $\mathfrak{gl}(1|1)$ one has tail $\lambda = \delta_{0,\lambda}$; for $\mathfrak{gl}(1|1)$ one has tail $\lambda = 1$ for each atypical weight λ . Recall that the condition (Tail) in this case is tautological.

Except for $F(4), G(3)$ we take χ corresponding to the principal block (i.e., χ is the central character of the trivial module $L(0)$).

In all these examples the Poincaré polynomials $\kappa(e) \in \{1, z\}$; the graph Γ^χ admits a parametric partition dex (which means that $\kappa(e) = 1$ if e connects the vertices with different value of dex and $\kappa(e) = z$ otherwise). In particular, $D^\chi = \Gamma^\chi$.

3.9.1. Example: $\mathfrak{gl}(1|1)$. We take $\mathfrak{g} := \mathfrak{gl}(1|1)$ with $\Delta^+ = \{\alpha\}$.

The simple modules in the principal block are $\{L(s\alpha) | s \in \mathbb{Z}\}$. As a vector space $L(s\alpha) \cong \Pi^s(\mathbb{C})$, so $\text{sdim } L(s\alpha) = (-1)^s$; we define $\text{dex}(L(s\alpha)) := s$ modulo 2. For a non-zero $x \in \mathfrak{g}_1$ one has $\text{DS}_x(L(s\alpha)) = \Pi^s(\mathbb{C})$. We have

$$\begin{array}{cccccccc} \hat{\Gamma}^\chi & \dots & \xrightarrow{(1)} & -\alpha & \xrightarrow{(1)} & 0 & \xrightarrow{(1)} & \alpha & \xrightarrow{(1)} & \dots \\ \Gamma^\chi & \dots & & -\alpha & & 0 & & \alpha & & \dots \\ \text{Ext}(\chi) & \dots & \longleftrightarrow & -\alpha & \longleftrightarrow & 0 & \longleftrightarrow & \alpha & \longleftrightarrow & \dots \end{array}$$

Clearly, $\text{dex}(L(s\alpha)) := s$ modulo 2 defines a parametric partition on $\hat{\Gamma}^\chi$ and a bipartition on $\text{Ext}(\chi)$; the graph $\hat{\Gamma}^\chi$ does not admit a \mathbb{N} -grading.

3.9.2. *Example: $\mathfrak{osp}(2|2)$.* Take $\mathfrak{g} = \mathfrak{osp}(2|2) \cong \mathfrak{sl}(2|1)$ with the base $\Sigma = \{\delta_1 \pm \varepsilon_1\}$.

The simple modules in the principal block are $\{L(\lambda_s) | s \in \mathbb{Z}\}$, where $\lambda_s := |s|\delta_1 + s\varepsilon_1$. Note that 0 is a minimal dominant weight, so the assumption of Corollary 3.6.3 (iii) holds. The extension graph $\text{Ext}(\chi)$ is A_∞^∞ . The Poincaré polynomials $K_{\mathfrak{g},\mathfrak{b}}^{\lambda,\nu}$ were computed in [18], Sect. 12. One has

$$\begin{array}{cccccccc} \Gamma^\chi & & \dots & \xleftarrow{(2;1)} & \lambda_{-1} & \xleftarrow{(1;1)} & \lambda_0 & \xrightarrow{(1;1)} & \lambda_1 & \xrightarrow{(2;1)} & \dots \\ \text{Ext}(\chi) = A_\infty^\infty & & \dots & \longleftrightarrow & \lambda_{-1} & \longleftrightarrow & \lambda_0 & \longleftrightarrow & \lambda_1 & \longleftrightarrow & \dots \end{array}$$

By [18], Lemmata 25, 26 the graph $\hat{\Gamma}^\chi$ can be obtained from the graph Γ^χ by adding a loop around λ_0 which is marked by $(1; z)$. Observe that $\text{dex}(\lambda_j) := p(\lambda) \equiv j$ modulo 2 is a parametric partition on $\hat{\Gamma}^\chi$ and is a bipartition on $\text{Ext}(\chi)$. The function $\|\lambda_i\|_{gr} := |i|$ gives a \mathbb{N} -grading on Γ^χ ; the graph $\hat{\Gamma}^\chi$ does not admit a \mathbb{N} -grading (since it has a loop).

The Gruson-Serganova formula is

$$(14) \quad \text{ch } L(\lambda_{\pm j}) = \sum_{s=0}^j (-1)^{j-s} \mathcal{E}_{\lambda_{\pm s}} \quad \text{for } j \geq 0.$$

3.9.3. *Example: $\mathfrak{osp}(3|2)$.* Take $\mathfrak{g} = \mathfrak{osp}(3|2)$ with the base $\Sigma = \{\varepsilon_1 - \delta_1, \delta_1\}$. The simple modules in the principal block are $\{L(\lambda_s)\}_{s=0}^\infty$, where $\lambda_0 := 0$ and $\lambda_s := (s-1)\delta_1 + s\varepsilon_1$ for $s > 1$; 0 is a minimal dominant weight. The Poincaré polynomials $K_{\mathfrak{g},\mathfrak{b}}^{\lambda,\nu}$ were computed in [10]. One has

$$\hat{\Gamma}^\chi : \quad \begin{array}{ccccc} \lambda_1 & \xrightarrow{(1)} & \lambda_2 & \xrightarrow{(1)} & \lambda_3 & \xrightarrow{(1)} & \dots \\ & \searrow (z) & \uparrow (1) & & & & \\ & & \lambda_0 & & & & \end{array}$$

The map $\text{dex}(\lambda_0) := 0$, $\text{dex}(\lambda_i) := i - 1$ modulo 2 for $i \neq 0$ is a parametric partition on $\hat{\Gamma}^\chi$ (one has $\text{dex}(\lambda_i) = p(\lambda_i)$). The graph Γ^χ is

$$\begin{array}{ccccc} \lambda_1 & \xrightarrow{(2;1)} & \lambda_2 & \xrightarrow{(3;1)} & \lambda_3 & \xrightarrow{(4;1)} & \dots \\ & \searrow (1;z) & \uparrow (2;1) & & & & \\ & & \lambda_0 & & & & \end{array}$$

and the “undirected version” of $\text{Ext}(\chi)$ coincides with $\Gamma^\chi(0)$. The function $\|\lambda_i\|_{gr} := i$ gives an \mathbb{N} -grading Γ^χ (note that the graph $\hat{\Gamma}^\chi$ is not \mathbb{Z} -graded).

The Gruson-Serganova formula is $\text{ch } L(\lambda_0) = \mathcal{E}_0 = 1$ and

$$(15) \quad \text{ch } L(\lambda_1) = \mathcal{E}_{\lambda_0} + \mathcal{E}_{\lambda_1}, \quad \text{ch } L(\lambda_j) = 2(-1)^{j-1}\mathcal{E}_{\lambda_0} + \sum_{s=1}^j (-1)^{j-s}\mathcal{E}_{\lambda_s} \quad \text{for } j > 1.$$

3.9.4. *Remark.* For $\mathfrak{osp}(3|2)$ we have two bases: the “mixed” base $\{\varepsilon_1 - \delta_1; \delta_1\}$ and the base $\{\delta_1 - \varepsilon_1; \varepsilon_1\}$. The computations in [10] are performed for the second base; it is not hard to see that for the first base the results are the same.

For $\mathfrak{osp}(2|2) \cong \mathfrak{sl}(1|2)$ we have two bases: the “mixed” base $\{\delta_1 \pm \varepsilon_1\}$ and the distinguished base $\{\varepsilon_1 - \delta_1; 2\delta_1\}$. The graphs for the mixed base are given in 3.9.2; the graphs for the distinguished base are the same as in 3.9.1 (notice that 0 is a minimal dominant weight for the mixed base, whereas for the distinguished base the set of dominant weights does not have minimal elements).

3.9.5. *Cases $\mathfrak{osp}(4|2)$, $G(3)$ and $F(4)$.* Recall that $\text{Ext}(\chi)$ is either D_∞ or A_∞^∞ , see 2.3. The graphs $\hat{\Gamma}(\chi)$ for a certain distinguished Borel subalgebras were computed in [10], [25]: this graph is the same as for $\mathfrak{osp}(3|2)$ (resp., as for $\mathfrak{osp}(2|2)$) if $\text{Ext}(\chi) = D_\infty$ and (resp., if $\text{Ext}(\chi) = A_\infty^\infty$).

For the principal blocks the graph $\text{Ext}^1(\chi)$ is D_∞ and so the graph $\hat{\Gamma}(\chi)$ is the same as for $\mathfrak{osp}(3|2)$. Since $\text{tail } 0 = 1$ and $\text{tail } \lambda = 0$ for each $\lambda \neq 0$, the graph Γ^χ is the same as for $\mathfrak{osp}(3|2)$.

For the non-principal blocks one has $\hat{\Gamma}^\chi = \Gamma^\chi$ (since $\text{tail } \lambda = 0$ for each $\lambda \in \Lambda^\chi$).

The character formulae for these cases were obtained in [10],[25]. The above approach give other type of character formulae.

By above, for the principal block Γ^χ is the same as for $\mathfrak{osp}(3|2)$, so we obtain the same character formula (15). Consider a non-principal block. By above, $\Gamma^\chi = \hat{\Gamma}^\chi$ is the same as the graph $\hat{\Gamma}^\chi$ for $\mathfrak{osp}(2|2)$ or for $\mathfrak{osp}(3|2)$. In both cases Γ^χ have cycles and all these cycles contain λ_0 ; the graph Γ' which is obtained from Γ^χ by erasing all edges ending at λ^0 is \mathbb{N} -graded. Using 3.7.3, we get Gruson-Serganova type character formulae which can be obtained from (14) and (15) respectively by changing \mathcal{E}_0 by $\text{ch } L(\lambda_0)$ (notice that $\text{ch } L(\lambda_0)$ is given by the Kac-Wakimoto formula).

3.10. Remark on Gruson-Serganova type character formulae. Let \mathcal{B} one of the blocks of atypicality 1 considered in Section 2. Then Γ^χ is one of the graphs appeared in 3.9.2, 3.9.3 and we call λ_0 a *Kostant weight*; notice that λ_0 is uniquely defined in terms of Γ^χ which has fewer automorphisms than $\text{Ext}(\chi)$.

Take $L(\lambda_i) \in \text{Irr}(\mathcal{B})$ and write $\text{ch } L(\lambda_i) = \sum_i a_i \mathcal{E}_{\lambda_i}$ using (14), (15). By 2.3 one has $\text{DS}_x(L(\lambda_i)) = \Pi^s(\text{DS}_x(L(\lambda_0)))^{a_0}$ for $s = 0$ if $a_0 > 0$ and $s = 1$ if $a_0 < 0$. This can be translated to the language of supercharacters in the following manner.

Retain notation of 0.2. Denote by $\text{Sch}(\mathfrak{g})$ the image of the map $\text{sch} : \tilde{\mathcal{F}}(\mathfrak{g}) \rightarrow \mathbb{Z}[\Lambda_{m|n}]$. Since $\text{sch}(\Pi(V)) = -\text{sch}(V)$ one has $\text{sch}(\tilde{\mathcal{F}}(\mathfrak{g})) = \text{sch}(\mathcal{F}(\mathfrak{g}))$. For Lie superalgebras (1) the ring $\text{Sch}(\mathfrak{g})$ is isomorphic to the reduced Grothendieck ring of $\tilde{\mathcal{F}}(\mathfrak{g})$ and DS_x induces an algebra homomorphism $ds'_x : \text{Sch}(\mathfrak{g}) \rightarrow \text{Sch}(\mathfrak{g}_x)$ given by $f \mapsto f|_{\mathfrak{b}_x}$, see [22].

For $V \in \mathcal{F}(\mathfrak{g})$ one has $\text{sch} V = \pi(\text{ch } V)$, where $\pi : \mathbb{Z}[\Lambda_{m|n}] \rightarrow \mathbb{Z}[\Lambda_{m|n}]$ is the involution $\pi(e^\mu) := p(\mu)e^\mu$. In particular, $\{\pi(\mathcal{E}_\lambda) \mid \lambda \in \text{Irr}(\mathcal{F}(\mathfrak{g}))\}$ forms a basis of $\text{Sch}(\mathfrak{g})$. If \mathfrak{g} has defect 1, then the kernel of the map $ds'_x : \text{Sch}(\mathfrak{g}) \rightarrow \text{Sch}(\mathfrak{g}_x)$ is spanned by the basis elements $\pi(\mathcal{E}_\lambda)$ for λ s which are not Kostant weights. A similar property hold for the $\mathfrak{gl}(1|n)^{(1)}$ -case; in [14] we will show that this holds for $\mathfrak{osp}(m|n)$ -case as well (the situation is more complicated for $\mathfrak{gl}(m|n)$).

3.11. Example: $\mathfrak{g} = \mathfrak{gl}(k|k)$. Take $\mathfrak{g} = \mathfrak{gl}(k|k)$ with a distinguished Borel subalgebra \mathfrak{b} . For $p = 0, \dots, k$ we denote by $\mathfrak{p}^{(p)}$ a parabolic subalgebra containing \mathfrak{b} with the Levi factor $\mathfrak{l}^{(p)} \cong \mathfrak{gl}(p|p)$ ($\mathfrak{p}^{(p)}$ is unique since \mathfrak{b} is a distinguished Borel). We consider the corresponding chain of parabolic subalgebras (9).

The Poincaré polynomials $K_{\mathfrak{p}^{(p+1)}, \mathfrak{p}^{(p)}}^{\lambda, \nu}$ were computed in [32], [27] Cor. 3.8. The graph $\hat{\Gamma}^\chi$ is \mathbb{Z} -graded and does not have multi-edges. For the principal block the condition (Tail) holds. The map $p(\lambda)$ defines a parametric partition on $\hat{\Gamma}^\chi$. By [27], Sect. 6, $\text{Ext}(\chi) = \hat{\Gamma}^\chi(0)$ (this can be also deduced from [2]). One has $\text{Ext}(\chi) \neq \Gamma^\chi(0)$ (see the example of $\mathfrak{gl}(1|1)$ above).

4. CASE $\mathfrak{osp}(M|N)$

In this section $\mathfrak{g} = \mathfrak{osp}(M|N)$. The category $\mathcal{F}\text{in}(\mathfrak{g})$ was studied in [18] and [6]. In this section we deduce the existence of dex satisfying (Dex1) from the results of [18]. Another approach is developed in [6], [7]. By [18], each block of atypicality k in $\tilde{\mathcal{F}}(\mathfrak{g})$ is equivalent either to a principal $\mathfrak{osp}(2k+t|2k)$ -block \mathcal{B} or to $\Pi(\mathcal{B})$, where $t = 1$ for odd M and $t = 0, 2$ for even M , and this equivalence is “compatible with character formula”, see Remark 4.6.

We fix a “mixed” base consisting of odd roots, see 4.1.1 below. We denote by χ the central character of the principal block \mathcal{B} and retain notation of 3.5. For each $\lambda \in \Lambda^\chi$ we set $\text{dex}(L(\lambda)) = p(\lambda)$ if $t = 0, 1$; for $t = 2$ we define $\text{dex}(L(\lambda))$ via a one-to-one correspondence between the simple modules in the principal blocks for $\mathfrak{osp}(2n+1|2n)$ and $\mathfrak{osp}(2n+2|2n)$, see 4.3 below.

The multiplicities ${}^i K_{\mathfrak{p}^{(p)}, \mathfrak{p}^{(p-1)}}^{\lambda, \nu}$ were computed by C. Gruson and V. Serganova in [18] (see [20] for small rank examples). We will recall their results and describe the graphs Γ^χ, D^χ in 4.4. We will see that Γ^χ is \mathbb{N} -graded and satisfies (Tail). We will check that dex is a parametric partition and that b and b' are decreasingly-equivalent. As a result Corollary 3.6.3 holds for the block \mathcal{B} and the character formula (15) in [18] can be rewritten in the form (17).

Everywhere in this section, except for Remark 4.6, we take $\mathfrak{g} = \mathfrak{osp}(2k + t|2k)$ for $t = 0, 1, 2$.

4.1. Notation. We take $\mathfrak{g} := \mathfrak{osp}(2k + t|2k)$ for $t = 0, 1, 2$. The integral weight lattice $\Lambda_{k+\ell|k}$ is spanned by $\{\varepsilon_i\}_{i=1}^{k+\ell} \cup \{\delta_i\}_{i=1}^k$, where $\ell := 0$ for $t = 0, 1$ and $\ell := 1$ for $t = 2$; the parity function is given by $p(\varepsilon_i) = \bar{0}$, $p(\delta_j) = \bar{1}$ for all i, j .

4.1.1. We fix a triangular decomposition corresponding to the ‘‘mixed’’ base:

$$\Sigma := \begin{cases} \varepsilon_1 - \delta_1, \delta_1 - \varepsilon_2, \dots, \varepsilon_k - \delta_k, \delta_k & \text{for } \mathfrak{osp}(2k + 1|2k) \\ \delta_1 - \varepsilon_1, \varepsilon_1 - \delta_2, \dots, \varepsilon_{k-1} - \delta_k, \delta_k \pm \varepsilon_k & \text{for } \mathfrak{osp}(2k|2k) \\ \varepsilon_1 - \delta_1, \delta_1 - \varepsilon_2, \dots, \varepsilon_k - \delta_k, \delta_k \pm \varepsilon_{k+1} & \text{for } \mathfrak{osp}(2k + 2|2k). \end{cases}$$

We have $\rho = 0$ for $t = 0, 2$ and $\rho = \frac{1}{2} \sum_{i=1}^k (\delta_i - \varepsilon_i)$ for $t = 1$.

4.1.2. We consider the embeddings

$$\mathfrak{osp}(t|0) \subset \mathfrak{osp}(2 + t|2) \subset \mathfrak{osp}(4 + t|4) \subset \dots \subset \mathfrak{osp}(2k + t|2k) = \mathfrak{g}$$

where $\mathfrak{osp}(2p + t|2p)$ corresponds to the last $2p + \ell$ roots in Σ ; we denote the subalgebra $\mathfrak{osp}(2p + t|2p)$ by $\mathfrak{l}^{(p)}$. Note that $\mathfrak{l}^{(k)} = \mathfrak{g}$ and $\mathfrak{l}^{(0)} = 0$ for $t = 0, 1$, $\mathfrak{l}^{(0)} = \mathbb{C}$ to $t = 2$.

For $p = 0, \dots, k$ we consider the parabolic subalgebra $\mathfrak{p}^{(p)} := \mathfrak{l}^{(p)} + \mathfrak{b}$. Notice that $\mathfrak{l}^{(p)}$ is the Levi subalgebra of $\mathfrak{p}^{(p)}$; as in 3.5, we denote by $\text{tail}(\lambda)$ the maximal index q such that $\lambda|_{\mathfrak{b} \cap \mathfrak{t}_q} = 0$.

4.2. Highest weights in the principal block. For $\lambda \in \Lambda_{k+\ell|k}$ we set

$$a_i := -(\lambda|\delta_i)$$

and notice that $p(\lambda) = \sum_{i=1}^k a_i$. By [18], $\lambda \in \Lambda^\times$ if and only if a_1, \dots, a_k are non-negative integers with $a_{i+1} > a_i$ or $a_i = a_{i+1} = 0$, and

$$\lambda + \rho = \begin{cases} \sum_{i=1}^{k-1} a_i(\varepsilon_i + \delta_i) + a_k(\delta_k + \xi\varepsilon_k) & \text{for } t = 0 \\ \sum_{i=1}^k a_i(\varepsilon_i + \delta_i) & \text{for } t = 2 \\ \sum_{i=1}^{s-1} (a_i + \frac{1}{2})(\varepsilon_i + \delta_i) + \frac{1}{2}(\delta_s + \xi\varepsilon_s) + \sum_{i=s+1}^k \frac{1}{2}(\delta_i - \varepsilon_i) & \text{for } t = 1 \end{cases}$$

for $\xi \in \{\pm 1\}$. For $t = 1$ we have $1 \leq s \leq k + 1$ and $a_s = a_{s+1} = \dots = a_k = 0$ if $s \leq k$ (for $s = k + 1$ we have $\lambda + \rho = \sum_{i=1}^k (a_i + \frac{1}{2})(\varepsilon_i + \delta_i)$).

4.2.1. Take $\lambda \in \Lambda^\times$ and define a_i for $i = 1, \dots, k$ as above. We assign to λ a “weight diagram”, which is a number line with one or several symbols drawn at each position with non-negative integral coordinate:

we put the sign \times at each position with the coordinate a_i ;

for $t = 2$ we add $>$ at the zero position;

we add the “empty symbol” \circ to all empty positions.

For $t \neq 2$ a weight $\lambda \in \Lambda^\times$ is not uniquely determined by the weight diagram constructed by the above procedure. Therefore, for $t = 0$ with $a_k \neq 0$ and for $t = 1$ with $s \leq k$, we write the sign of ξ before the diagram ($+$ if $\xi = 1$ and $-$ if $\xi = -1$).

Notice that each position with a non-zero coordinate contains either \times or \circ . For $t = 0, 1$ the zero position is occupied either by \circ or by several symbols \times ; we write this as \times^i for $i \geq 0$. Similarly, for $t = 2$ the zero position is occupied by $>^{\times^i}$ with $i \geq 0$.

4.2.2. Notice that $\text{tail}(\lambda)$ is equal to the number of symbols \times at the zero position of the weight diagram for all cases except when $t = 1$ and the diagram has the sign $+$; in the latter case the number of symbols \times at the zero position is $\text{tail}(\lambda) + 1$.

4.2.3. *Examples.* The weight diagram of 0 is \times^k for $t = 0$, $-\times^k$ for $t = 1$ and $>^{\times^k}$ for $t = 2$; one has $\text{tail}(0) = k$.

The diagram $+\circ\times\times$ corresponds to the $\mathfrak{osp}(4|4)$ -weight $\lambda = \lambda + \rho = (\varepsilon_2 + \delta_2) + 2(\varepsilon_1 + \delta_1)$ with $\text{tail}(\lambda) = 0$.

The diagram $+\times^3$ corresponds to $\mathfrak{osp}(7|6)$ -weight $\lambda = \varepsilon_1$ with $\text{tail}(\varepsilon_1) = 2$.

The empty diagram correspond to $\mathfrak{osp}(0|0) = \mathfrak{osp}(1|0) = 0$; the diagram $>$ corresponds to the weight 0 for $\mathfrak{osp}(2|0) = \mathbb{C}$.

4.2.4. For $t = 0, 1, 2$ we denote by $\text{Diag}_{k;t}$ the set of (signed) weight diagrams. The above procedure gives a one-to-one correspondence between Λ^\times and $\text{Diag}_{k;t}$. For each diagram $f \in \text{Diag}_{k;t}$ we denote by $\lambda(f)$ the corresponding weight in Λ^\times .

In all cases the weight diagrams in $\text{Diag}_{k;t}$ contains k symbols \times .

For a diagram f and $a \in \mathbb{N}$ we denote by $f(a)$ the symbols at the position a . For $t = 0$ (resp., $t = 1$) a diagram in $\text{Diag}_{k;t}$ has a sign if and only if $f(0) = \circ$ (resp., $f(0) \neq \circ$).

4.2.5. *Map τ .* Following [18], we introduce a bijection $\tau : \text{Diag}_{k;2} \rightarrow \text{Diag}_{k;1}$. For $f \in \text{Diag}_{k;2}$ the diagram $\tau(f) \in \text{Diag}_{k;1}$ is constructed by the following procedure:

we remove $>$ and then shift all entires at the non-zero positions of f by one position to the left; then we add a sign in such a way that $\text{tail}(f) = \text{tail}(\tau(f))$: the sign $+$ if $f(1) = \times$

and the sign $-$ if $f(1) = \circ$ and $f(0) \neq >$. For instance,

$$\tau(\overset{\times}{>} \circ \times) = - \times \times, \quad \tau(\overset{\times}{>}) = - \times, \quad \tau(> \times) = + \times, \quad \tau(> \circ \times) = \circ \times.$$

One readily sees that τ is a one-to-one correspondence.

4.3. The maps $\|\lambda\|, \|\lambda\|_{gr}, \text{dex}$. Let $\{a_i\}_1^k$ be the coordinates of the symbols \times in a diagram of λ . We set

$$\|\lambda\| := \begin{cases} \sum_{i=1}^k a_i & \text{for } t = 0, 1 \\ \|\tau(f)\| & \text{for } t = 2, \end{cases} \quad \|\lambda\|_{gr} := \begin{cases} \sum_{i=1}^k a_i & \text{for } t = 0, 2 \\ \|\tau^{-1}(f)\| & \text{for } t = 1 \end{cases}$$

and

$$\text{dex}(\lambda) \equiv \|\lambda\| \pmod{2}.$$

Clearly, $\|\lambda\|, \|\lambda\|_{gr} \in \mathbb{N}$ and $\|\lambda\|_{gr} = 0$ if and only if $\lambda = 0$.

4.4. Graph Γ^\times . Retain notation of Section 3. Consider the chain of parabolic subalgebras (9) with $\mathfrak{p}^{(p)}$, $p = 0, 1, \dots, k$ defined in 4.1.2.

The Poincaré polynomials $K_{\mathfrak{p}^{(p)}, \mathfrak{p}^{(p-1)}}^{\lambda, \nu}(z)$ for $p > \text{tail } \lambda$ were computed in [18], Sect. 11. It is proven that the map $\tau : \text{Diag}_{k;2} \xrightarrow{\sim} \text{Diag}_{k;1}$ (see 4.2.5) preserves these polynomials (i.e., $K_{\mathfrak{p}^{(p)}, \mathfrak{p}^{(p-1)}}^{\tau(\lambda), \tau(\nu)} = K_{\mathfrak{p}^{(p)}, \mathfrak{p}^{(p-1)}}^{\lambda, \nu}$); the coefficients ${}^i K_{\mathfrak{p}^{(p)}, \mathfrak{p}^{(p-1)}}^{\lambda, \nu}$ are 0 or 1 and for $t = 0, 2$ one has ${}^i K_{\mathfrak{p}^{(p)}, \mathfrak{p}^{(p-1)}}^{\lambda, \nu} = 1$ if and only if the diagram of λ can be obtained from the diagram of ν by a “move” of degree i which ends at the p th symbol \times in the diagram of λ ; we will give some details in 4.4.1 below and give a description in terms of “arch diagrams” in Section 5.

4.4.1. Moves for $t = 0, 2$. Consider the cases $t = 0, 2$. Take $f \in \text{Diag}_{k;t}$. For each $p, q \in \mathbb{N}$ denote by $l_f(p, q)$ the number of symbols \times minus the number of symbols \circ strictly between the positions p and q in f .

A diagram $f \in \text{Diag}_{k;t}$ can be transformed to a diagram g by a “move” of degree d if f satisfies certain conditions, and g is obtained from f by moving either one symbol \times from a position p to an empty position q with $q > p$ or moving two symbols \times from the zero position to empty positions p, q with $p < q$. If f has a sign, then g has the same sign. In both cases we say that the move “ends at the position q ”. We will not specify all conditions on f , but notice that these conditions depend only on $l_f(s, q)$ for $s < q$.

By above, $\text{tail}(f) - \text{tail}(g)$ is 0, 1 or 2; the degree d satisfies the formula

$$(16) \quad d = \begin{cases} l_f(p, q) & \text{if } \text{tail}(f) - \text{tail}(g) \neq 1 \\ l_f(p, q) \text{ or } 2 \text{tail}(g) + l_f(p, q) & \text{if } t = 0, \text{tail}(f) - \text{tail}(g) = 1 \\ 2 \text{tail}(g) + l_f(p, q) + 1 & \text{if } t = 2, \text{tail}(f) - \text{tail}(g) = 1, \end{cases}$$

The conditions on f imply that $d \geq 0$. Except for the case $t = 0$ with $\text{tail}(f) - \text{tail}(g) = 1$, g can be obtained from f by at most one move; for $t = 0$ it is possible sometimes to obtain g from f by two moves of different degrees. We give below examples of several moves and their degrees

$$\begin{aligned} \dots \times \circ \dots &\longrightarrow \dots \circ \times \dots & d = 0 \\ \dots \times \times \times \circ \circ \dots &\longrightarrow \dots \circ \times \times \circ \times \dots & d = 1 \\ \times \circ \dots &\longrightarrow (\pm) \circ \times \dots & d = 0 \\ \times^2 \circ \dots &\longrightarrow \times \times \dots & d = 0, 2. \end{aligned}$$

4.4.2. Corollary. *Take $t = 0$ or $t = 2$. Let $\nu, \lambda \in \Lambda^x$ be two weights with the diagrams f and g respectively. Assume that g is obtained from f by a move of degree d . Then*

- (i) $\|\lambda\|_{gr} > \|\nu\|_{gr}$ and $\lambda > \nu$;
- (ii) if the move ends in the i th symbol \times in g , then $\text{tail } \nu \leq i$;
- (iii) $\text{dex}(\lambda) - \text{dex}(\nu) + d \equiv 1 \pmod{2}$.

Proof. The inequality $\|\lambda\|_{gr} > \|\nu\|_{gr}$ follows from the fact that τ preserves $\|\cdot\|_{gr}$ and that we move symbol(s) \times to the right. The inequality $\lambda > \nu$ follows from the fact that we move the symbol(s) \times to the right; (ii) is obvious. For (iii) retain notation of 4.4.1 and observe that

$$l_f(p, q) \equiv q - p + 1, \quad \text{dex}(\lambda) - \text{dex}(\nu) \equiv \|\lambda\| - \|\nu\| \pmod{2}.$$

For $t = 0$ one has $d \equiv l_f(p, q)$ by (16), and $\|\lambda\| - \|\nu\| = q - p$ if $\text{tail}(g) - \text{tail}(f) \neq 2$ and $q + p$ otherwise. For $t = 2$ the formula (16) gives $d \equiv l_f(p, q) + \text{tail}(\lambda) - \text{tail}(\nu)$; in this case $\|\lambda\| - \|\nu\| \equiv q - p + \text{tail}(\lambda) - \text{tail}(\nu)$. This gives (iii). \square

4.4.3. Retain notation of 3.5. For $t = 0, 2$ the graph $\Gamma^x = (\Lambda^x, E)$ has the edges $\nu \xrightarrow{e} \lambda$ with $b(e) = j$ if and only if the diagram of λ can be obtained from the diagram of ν by a move which ends at the j th symbol \times in the diagram of λ ; in this case we denote by $b'(e) = q$ the coordinate of the j th symbol \times in the diagram of λ . For the edge as above the Poincaré polynomial is the sum of z^d for all d such that λ can be obtained from the diagram of ν by a move of degree d . By (16) $\kappa(e) = z^d$ except for the case $t = 0$ with $\text{tail}(\nu) - \text{tail}(\lambda) = 1$; in the latter case $\kappa(e) = z^d$ or $z^d(1 + z^{2\text{tail}(\lambda)})$ (see Section 5 for details).

By above, τ gives a bijection between the graphs Γ^x for $t = 2$ and $t = 1$ and this bijection is compatible with the functions b and κ . For $t = 1$ we define b' on Γ^x using this bijection.

4.4.4. Corollary.

- (i) *The map $\text{dex}(\lambda)$ is a parametric bipartition on (Γ^x, κ) .*
- (ii) *If $\nu \rightarrow \lambda$ is an edge in Γ^x , then $\nu < \lambda$ and $\|\nu\|_{gr} < \|\lambda\|_{gr}$. In particular, $\|\lambda\|_{gr}$ defines a \mathbb{N} -grading on Γ^x .*

- (iii) *The graph Γ^x satisfies the assumption (Tail).*
- (iv) *The functions b, b' are decreasingly equivalent and b' satisfies the property (BB) of Lemma 3.4.4.*

Proof. Consider the cases $t = 0, 2$. Corollary 4.4.2 implies (i)–(iii). For (iv) take a path $\lambda(f_1) \xrightarrow{e_1} \lambda(f_2) \xrightarrow{e_2} \lambda(f_3)$ in Γ^x . Since f_3 is obtained from f_2 by a move which ends at the symbol \times with the coordinate $b'(e_2)$, the position with this coordinate in f_2 is empty, so $b'(e_1) \neq b'(e_2)$. Hence b' satisfies (BB). It remains to verify that b and b' are decreasingly equivalent. Set $j := b(e_1)$, $q := b'(e_1)$. Then q is the coordinate of the j th symbol \times in f_2 and $q > 0$. The condition $b(e_1) > b(e_2)$ means that for $i \geq j$ the i th symbols \times in f_2 and f_3 have the same coordinates, whereas the condition $q > b'(e_2)$ means that for $s \geq q$ one has $f_2(s) = f_3(s)$. Clearly, these conditions are equivalent, so b and b' are decreasingly equivalent.

Consider the remaining case $t = 1$. Since τ preserves dex, tail and $\|\cdot\|_{gr}$, almost all assertions for $t = 1$ follows from the corresponding assertions for $t = 2$. The only exception is the inequality $\nu < \lambda$ in (ii), which follows from the following observation: for an edge $\nu \rightarrow \lambda$ in Γ^x for $t = 1$, the diagram of λ is obtained from the diagram of ν either by moving symbol(s) \times to the right or by changing the sign from $-$ to $+$. \square

4.4.5. Corollary. *Take $\lambda, \nu \in \Lambda^x$ with $\text{Ext}^1(L(\lambda), L(\nu)) \neq 0$. Then*

- (i) *either λ can be obtained from ν by a move of zero degree or ν can be obtained from λ by a move of zero degree;*
- (ii) $\text{dex}(\lambda) \neq \text{dex}(\nu)$.

Proof. By Corollary 4.4.4, the graph Γ^x satisfies the assumptions of Corollary 3.6.3 (iii), which implies the assertions. \square

4.5. Gruson-Serganova character formula. We retain notation of 3.3; for $\nu \in \Lambda^x$ we introduce the “Euler character” \mathcal{E}_ν by (13).

4.5.1. A character formula is given by Theorem 4 in [18]. Using Corollary 4.4.4 we can write this formula for $\lambda \in \Lambda^x$ in the following way:

$$(17) \quad \text{ch } L(\lambda) = \sum_{\nu \in \Lambda^x} (-1)^{\text{dex}(\lambda) - \text{dex}(\nu)} d_{<}^{\lambda, \nu} \mathcal{E}_\nu,$$

where $d_{<}^{\lambda, \nu}$ is the number of increasing paths from ν to λ in the graph D^x , where D^x is obtained from Γ^x by doubling the edges e with $\kappa(e) \neq z^d$ (i.e., $D^x = \Gamma^x$ for $t \neq 0$, see 4.4.3).

Notice that $d_{<}^{\lambda, \nu}$ are non-negative integers, $d_{<}^{\lambda, \lambda} = 1$ and $d_{<}^{\lambda, \nu} \neq 0$ implies $\nu \leq \lambda$ and $\|\nu\| \leq \|\lambda\|$ (in particular, the right-hand side of (17) is finite).

4.6. *Remark.* Recall that each block of atypicality k for $\mathfrak{osp}(M|N)$ is equivalent to the principal block of $\mathfrak{osp}(2k+t|2k)$ where $t = 1$ for odd M and $t = 0, 2$ for even M . For a dominant weight λ of atypicality k let $\bar{\lambda}$ be the image of λ in Λ^\times (that is the $\mathfrak{osp}(2k+t|2k)$ -module $L(\bar{\lambda})$ is the image of $\mathfrak{osp}(M|N)$ -module $L(\lambda)$ under the above equivalence). It turns out that this equivalence “preserve tails”, i.e. $\text{tail}(\lambda) = \text{tail}(\bar{\lambda})$.

Introducing $\text{dex}(\lambda) := \text{dex}(\bar{\lambda})$ we obtain $\text{Ext}^1(L(\lambda), L(\nu)) = 0$ if $\text{dex}(\lambda) = \text{dex}(\nu)$. By [18], the formula (17) holds for an arbitrary dominant weight λ if we introduce \mathcal{E}_ν by the formula (13) and set $d_{<}^{\lambda, \nu} := d_{<}^{\bar{\lambda}, \bar{\nu}}$.

5. ARCH DIAGRAMS

In the cases when \mathfrak{g} is not exceptional and the flag of parabolic is standard, the description of Γ^\times in [27],[18] can be conveniently presented in terms of arc diagrams introduced in [19], [13], where the examples are presented. Below we will present this description of Γ^\times for the principal blocks in $\mathfrak{gl}(k|k)$, $\mathfrak{osp}(2k+t|2k)$.

Our diagrams differs from the arc- or cup diagrams of [6]; we will call these diagrams “arch diagram”.

We take $\mathfrak{g} = \mathfrak{gl}(k|k)$, $\mathfrak{osp}(2k+t|2k)$ and the central character χ corresponding to the principal block. For $\mathfrak{g} = \mathfrak{gl}(k|k)$ we take $\Sigma = \{\varepsilon_1 - \varepsilon_2, \dots, \varepsilon_k - \delta_1, \delta_1 - \delta_2, \dots, \delta_{k-1} - \delta_k\}$ and the flag (9) with $\mathfrak{l}^{(i)} \cong \mathfrak{gl}(i|i)$. For $\mathfrak{osp}(2k+t|2k)$ we retain notation of 4.1.

5.1. **Arch diagram.** Take $\lambda \in \Lambda^\times$.

For $\mathfrak{osp}(2k+t|2k)$ we assign to λ the weight diagram as in 4.2. For $\mathfrak{gl}(k|k)$ -case $\lambda + \rho = \sum_{i=1}^k a_i(\varepsilon_i - \delta_{k+1-i})$ and we assign to λ a weight diagram with the symbols \times at the positions a_1, \dots, a_k and the empty symbols \circ in other positions.

A *generalized arch diagram* is the following data:

- a weight diagram f , where the symbols \times at the zero position are drawn vertically and $>$ (if it is present) is drawn in the bottom,
- a collection of non-intersecting arches, where each arch is
 - either $\text{arc}(a; b)$ connecting the symbol \times with the empty symbol at the position b ;
 - or $\text{arc}(0; b, b')$ connecting the symbol \times at the zero position with two empty symbols at the positions $b < b'$;

An empty position is called *free* if this position is not an end of an arch.

We call $\text{arc}(a; b)$ a *two-legged arch supported at a* and $\text{arc}(0; b, b')$ a *three-legged arch supported at 0*.

A generalized arch diagram is called *arch diagram* if

- each symbol \times is the left end of exactly one arch;
- there are no free positions under the arches;
- for the \mathfrak{gl} -case all arches are two-legged;
- for the $\mathfrak{osp}(2k|2k)$, $\mathfrak{osp}(2k+1|2k)$ -cases the lowest \times at the zero position supports a two-legged arch and the other symbols \times at the zero position support three-legged arches;
- for the $\mathfrak{osp}(2k+2|2k)$ -case all symbols \times at the zero position support three-legged arches.

Each weight diagram f admits a unique arch diagram which we denote by $\text{Arc}(f)$; this diagram can be constructed in the following way: we pass from right to left through the weight diagram and connect each symbol \times with the next empty symbol(s) to the right by an arch.

5.1.1. *Partial order.* We consider a partial order on the set of arches by saying that one arch is smaller than another one if the first one is "below" the second one; one has

$$\begin{aligned} \text{arc}(a; b) > \text{arc}(a'; b') &\iff a < a' < b \\ \text{arc}(0; b_1, b_2) > \text{arc}(a'; b') &\iff a' < b_2, \end{aligned}$$

in addition, any two distinct three-legged arches are comparable.

5.1.2. For a weight diagram f we denote by $l_f(p, q)$ the number of \times – the number of \circ strictly between the positions p and q . We denote by $(f)_p^q$ the diagram which obtained from f by moving \times from the position p to a free position $q > p$; such diagram is defined only if

$$f(p) \in \{\times^i, \overset{\times^i}{>}\} \text{ for } s \geq 1, f(q) = \circ.$$

For instance, for $f = \times^2 \circ \times$ one has $(f)_0^3 = \times \circ \times \times$ and $(f)_0^2, (f)_1^5$ are not defined. If $f(0) = \times^i$ or $\overset{\times^i}{>}$ for $i > 1$, we denote by $(f)_{0,0}^{p,q}$ the diagram which obtained from f by moving two symbols \times from the zero position to free positions p and q with $p < q$; for example, $(\times^2 \times)_{0,0}^{3,4} = \circ \times \circ \times \times$.

5.2. Description of Γ^\times in diagrammatic terms. Below we present the description for the cases $\mathfrak{gl}(k|k)$, $\mathfrak{osp}(2k|2k)$ and $\mathfrak{osp}(2k+2|2k)$ (the map τ gives a bijection between the graphs for $\mathfrak{osp}(2k+1|2k)$ and $\mathfrak{osp}(2k+2|2k)$).

Take $\nu, \lambda \in \Lambda^\times$ and denote by f and g their weight diagrams. We denote by $l_f(a, b)$ the number of \times minus the number of \circ with the coordinates strictly between a and b . Note that $l_f(a, b) = l_{f'}(a, b)$ if $f' = (f)_a^b$.

The graph Γ^\times contains an edge $\nu \xrightarrow{e} \lambda$ if and only if $g = (f)_a^p$ or $g = (f)_{0,0}^{p,q}$ and the following conditions hold.

5.2.1. *Case $g = (f)_a^p$ and $f(a) = \times$.* In this case $\text{Arc}(f)$ contains a two-legged arch $\text{arc}(a; a_-)$ with $a < p \leq a_-$; one has $\kappa(e) = z^{l_f(a,p)}$ and $l_f(a, b) = l_g(a, b)$.

5.2.2. *Case $g = (f)_0^p$ and $f(0) \neq \times$.* In this case $\text{Arc}(f)$ contains three-legged arches (i.e., either $\mathfrak{g} = \mathfrak{osp}(2k|2k)$ and $\text{tail}(f) \geq 2$ or $\mathfrak{g} = \mathfrak{osp}(2k + 2|2k)$ and $\text{tail}(f) \geq 1$). For this case we denote by $\text{arc}(0; a', a_+)$ the highest three-legged arch in $\text{Arc}(f)$ (for example, for $\nu = 0$ this is $\text{arc}(0; 2k - 2, 2k - 1)$ if $\mathfrak{g} = \mathfrak{osp}(2k|2k)$, $k > 1$ and $\text{arc}(0; 2k - 1, 2k)$ if $\mathfrak{g} = \mathfrak{osp}(2k + 2|2k)$). In this case $p \leq a_+$.

If $f(0) = \times^{i+1}$ for $i \geq 0$ (then $t = 2$) we have $\kappa(e) = z^{2i+1+l_f(0,p)}$. In the remaining case $f(0) = \times^{i+1}$, $i > 0$ (then $t = 0$) there is a unique two-legged arch $\text{arc}(0; a_-)$ supported at the zero position and $\kappa(e) = z^{2i+l_f(0,p)}$ if $p > a_-$ and $\kappa(e) = z^{2i+l_f(0,p)} + z^{l_f(0,p)}$ if $p \leq a_-$.

For example, for $f = \times^3 \times \circ \circ \circ \times$ and $g = (f)_0^2 = \times^2 \times \times \circ \circ \times$ one has $\kappa(e) = z + z^5$ and for $g' = (f)_0^4 = \times^2 \times \circ \circ \times \times$ one has $\kappa(e') = z^3$.

5.2.3. *Case $g = (f)_{0,0}^{p,q}$.* In this case $\text{Arc}(f)$ contains a three-legged arch $\text{arc}(0; p, a_2)$ supported at the zero position which is not the highest arch and $p < q \leq a_2$; one has $\kappa(e) = z^{l_f(p,q)}$.

Examples.

For $t = 0$, $\text{tail}(f) \leq 2$ (resp., $t = 2$, $\text{tail}(f) \leq 1$) there are no suitable three-legged arches.

For $f = \times^3 \circ \circ \times \times$ there is only one suitable three-legged arch, which is $\text{arc}(0; 2, 7)$. Thus $p = 2$, $q \in \{5, 6, 7\}$ and $\kappa(e) = z^{7-q}$.

For $f = \times^2 \circ \times \times \circ \times$ there is only one suitable three-legged arch, which is $\text{arc}(0; 1, 8)$. Thus $p = 1$, $q \in \{4, 6, 7, 8\}$ with $\kappa(e) = 2$ for $q = 5, 6$ and $\kappa(e) = 8 - q$ for $q = 7, 8$.

5.3. **Applications to Ext-graph.** Take $\nu, \lambda \in \Lambda^\times$ with $\nu < \lambda$ and denote by f and g their weight diagrams.

By [27], in the \mathfrak{gl} -case $\text{Ext}^1(L(\lambda), L(\nu)) \neq 0$ if and only if $g = (f)_a^p$, where $\text{arc}(a, p) \in \text{Arc}(f)$ (i.e. g can be obtained from f by moving a symbol \times along the arch supported by this symbol).

By above, for $\mathfrak{g} = \mathfrak{osp}(2k + t|2k)$ with $t = 0, 2$ if $\text{Ext}^1(L(\lambda), L(\nu)) \neq 0$, then either $g = (f)_a^p$, where $\text{arc}(a, p) \in \text{Arc}(f)$ or $g = (f)_0^{a_+}$ where $\text{arc}(0; a', a_+)$ is the highest arch supported at 0, or $g = (f)_{0,0}^{a_1, a_2}$, where $\text{arc}(0; a_1, a_2)$ is not the highest arch supported at 0.

6. APPENDIX: USEFUL FACTS ABOUT Ext^1

6.1. We start from the following lemma which express $\dim \text{Ext}^1(L', L)$ in terms of indecomposable extensions.

Lemma. *Let A be an associative algebra and L, L' be simple non-isomorphic modules over A with $\text{End}_A(L) = \text{End}_A(L') = \mathbb{C}$. Let N be a module with*

$$(18) \quad \text{Soc } N = L^{\oplus m}, \quad N / \text{Soc } N = L'.$$

(i) *If N is indecomposable, then $m \leq \dim \text{Ext}^1(L', L)$.*

(ii) *If $m \leq \dim \text{Ext}^1(L', L)$, then there exists an indecomposable N satisfying (18).*

Proof. Consider any exact sequence of the form

$$(19) \quad 0 \rightarrow L^{\oplus m} \xrightarrow{\iota} N \xrightarrow{\phi} L' \rightarrow 0.$$

For each $i = 1, \dots, m$ let p_i be the projection from $L^{\oplus m}$ to the i th component and θ_i is the corresponding embedding $p_i \theta_i = \text{Id}_L$. Consider a commutative diagram

$$(20) \quad \begin{array}{ccccccccc} 0 & \longrightarrow & L^{\oplus m} & \xrightarrow{\iota} & N & \xrightarrow{\phi} & L' & \longrightarrow & 0 \\ & & \downarrow & & \downarrow \psi_i & & \downarrow \text{Id} & & \downarrow \\ & & \theta_i \downarrow & p_i \uparrow & & & & & \\ 0 & \longrightarrow & L & \xrightarrow{\iota_i} & M^i & \xrightarrow{\phi_i} & L' & \longrightarrow & 0 \end{array}$$

where $\psi_i : N \rightarrow M^i$ is a surjective map with $\text{Ker } \psi_i = \text{Ker } p_i$.

The bottom line is an element of $\text{Ext}^1(L', L)$, which we denote by Φ_i . If $m > \dim \text{Ext}^1(L', L)$, then $\{\Phi_i\}_{i=1}^m$ are linearly dependent and we can assume that $\Phi_1 = 0$, so Φ_1 splits. Let $\tilde{p} : M^1 \rightarrow L$ be the projection, i.e. $\iota_1 \tilde{p} = \text{Id}_L$. Consider the maps

$$L \xrightarrow{\iota \circ \theta_1} N \xrightarrow{\tilde{p} \circ \psi_1} L.$$

The composed map

$$\tilde{p} \circ \psi_1 \circ \iota \circ \theta_1 : L \rightarrow L$$

is surjective, so it is an isomorphism. Hence N is decomposable. This establishes (i).

For (ii) let $\{\Phi_i\}_{i=1}^m$ be linearly independent elements in $\text{Ext}^1(L', L)$, i.e.

$$\Phi_i : \quad 0 \rightarrow L \xrightarrow{\iota_i} M^i \xrightarrow{\phi_i} L' \rightarrow 0.$$

Consider the exact sequence

$$0 \rightarrow L^{\oplus m} \rightarrow \oplus M_i \rightarrow (L')^{\oplus m} \rightarrow 0.$$

Let $\text{diag}(L')$ be the diagonal copy of L' in $(L')^{\oplus m}$ and let N be the preimage of $\text{diag}(L')$ in $\oplus M^i$. This gives the exact sequence of the form (19) and the commutative diagram (20). Let us show that N is indecomposable. Assume that N decomposable, so $N = N_1 \oplus N_2$. Since $L' \not\cong L$ one has $\phi(N_1) = 0$ or $\phi(N_2) = 0$, so N_1 or N_2 lies in the socle of N . Therefore N can be written as $N = L^1 \oplus N''$ with $L^1 \cong L$. Since $\dim \text{Hom}(L, N) = m$ one has $\dim \text{Hom}(L, N'') = m - 1$. Changing the basis in the span of $\{\Phi_i\}_{i=1}^m$, we can assume that $\text{Ker } p_1 \subset N''$. Since $\text{Ker } \psi_1 = \text{Ker } p_1$, the exact sequence Φ_1 splits, a contradiction. Hence N is indecomposable. This completes the proof. \square

6.2. **Notation.** Let \mathfrak{g} be a Lie superalgebra of at most countable dimension.

Let $\mathfrak{h} \subset \mathfrak{g}_0$ be a finite-dimensional subalgebra satisfying

(H1) \mathfrak{h} acts diagonally on \mathfrak{g} and $\mathfrak{g}_0^{\mathfrak{h}} = \mathfrak{h}$.

We choose $h \in \mathfrak{h}$ satisfying

(H2) $\mathfrak{g}^h = \mathfrak{g}^{\mathfrak{h}}$ and each non-zero eigenvalue of $\text{ad } h$ has a non-zero real part.

(The assumption on $\dim \mathfrak{g}$ ensures the existence of h).

We write $\mathfrak{g} = \mathfrak{g}^{\mathfrak{h}} \oplus (\oplus_{\alpha \in \Delta(\mathfrak{g})} \mathfrak{g}_{\alpha})$, with $\Delta(\mathfrak{g}) \subset \mathfrak{h}^*$ and

$$\mathfrak{g}_{\alpha} := \{g \in \mathfrak{g} \mid [h, g] = \alpha(h)g \text{ for all } h \in \mathfrak{h}\}.$$

We introduce the triangular decomposition $\Delta(\mathfrak{g}) = \Delta^+(\mathfrak{g}) \amalg \Delta^-(\mathfrak{g})$, with

$$\Delta^{\pm}(\mathfrak{g}) := \{\alpha \in \Delta(\mathfrak{g}) \mid \pm \text{Re } \alpha(h) > 0\},$$

and define the partial order on \mathfrak{h}^* by $\lambda > \nu$ if $\nu - \lambda \in \mathbb{N}\Delta^-$. We set $\mathfrak{n}^{\pm} := \oplus_{\alpha \in \Delta^{\pm}} \mathfrak{g}_{\alpha}$ and consider the Borel subalgebra $\mathfrak{b} := \mathfrak{g}^{\mathfrak{h}} \oplus \mathfrak{n}^+$.

6.2.1. Take $z \in \mathfrak{h}$ satisfying

$$(21) \quad \alpha(z) \in \mathbb{R}_{\geq 0} \text{ for } \alpha \in \Delta^+ \text{ and } \alpha(z) \in \mathbb{R}_{\leq 0} \text{ for } \alpha \in \Delta^-.$$

Consider the superalgebras

$$\mathfrak{t} := \mathfrak{t}(z) := \mathfrak{g}^z, \quad \mathfrak{p} := \mathfrak{p}(z) := \mathfrak{g}^z + \mathfrak{b}.$$

Notice that $\mathfrak{p} = \mathfrak{t} \rtimes \mathfrak{m}$, where $\mathfrak{m} := \oplus_{\alpha \in \Delta: \alpha(z) > 0} \mathfrak{g}_{\alpha}$. Both triples $(\mathfrak{p}(z), \mathfrak{h}, h)$, $(\mathfrak{t}(z), \mathfrak{h}, h)$ satisfy (H1), (H2). One has $\mathfrak{t}^{\mathfrak{h}} = \mathfrak{p}^{\mathfrak{h}} = \mathfrak{g}^{\mathfrak{h}}$ and

$$\begin{aligned} \Delta^+(\mathfrak{p}) &= \Delta^+(\mathfrak{g}), & \Delta^+(\mathfrak{t}) &= \{\alpha \in \Delta^+(\mathfrak{g}) \mid \alpha(z) = 0\} \\ \Delta^-(\mathfrak{p}) &= \Delta^-(\mathfrak{t}) = \{\alpha \in \Delta^-(\mathfrak{g}) \mid \alpha(z) = 0\} \end{aligned}$$

6.2.2. *Modules* $M(\lambda), L(\lambda)$. For a semisimple \mathfrak{h} -module N we denote by N_{ν} the weight space of the weight ν and by $\Omega(N)$ the set of weights of N .

We denote by \mathcal{O} the full category of finitely generated modules with a diagonal action of \mathfrak{h} and locally nilpotent action of \mathfrak{n} .

By Dixmier generalization of Schur's Lemma (see [4]), up to a parity change, the simple $\mathfrak{g}^{\mathfrak{h}}$ -modules are parametrized by $\lambda \in \mathfrak{h}^*$; we denote by C_{λ} a simple $\mathfrak{g}^{\mathfrak{h}}$ -module, where \mathfrak{h} acts by λ . We view C_{λ} as a \mathfrak{b} -module with the zero action of \mathfrak{n} and set

$$M(\lambda) := \text{Ind}_{\mathfrak{b}}^{\mathfrak{g}} C_{\lambda};$$

this module has a unique simple quotient which we denote by $L(\lambda)$. (The module $M(\lambda)$ is a Verma module if $\mathfrak{g}^{\mathfrak{h}} = 0$). We introduce similarly the modules $M_{\mathfrak{p}}(\lambda), L_{\mathfrak{p}}(\lambda)$ for the algebra \mathfrak{p} and $M_{\mathfrak{t}}(\lambda), L_{\mathfrak{t}}(\lambda)$ for the algebra \mathfrak{t} .

6.2.3. Set $N(\mathfrak{g}; m)$. Let $\lambda \neq \nu \in \mathfrak{h}^*$ be such that $Re(\lambda - \nu)(h) \geq 0$. If

$$0 \rightarrow L(\nu) \rightarrow E \rightarrow L(\lambda) \rightarrow 0$$

is a non-split exact sequence, then E is generated by $E_\lambda \cong C_\lambda$, so E is a quotient of $M(\lambda)$ and $\nu < \lambda$.

For $\lambda, \nu \in \mathfrak{h}^*$ we denote by $\mathcal{N}(\mathfrak{g}; m)$ the set of indecomposable \mathfrak{g} -modules N such that

$$(22) \quad \text{Soc } N = L(\nu)^{\oplus m}, \quad N/\text{Soc } N = L(\lambda)$$

By Lemma 6.1 one has

$$(23) \quad \dim \text{Ext}^1(L(\lambda), L(\nu)) = \max\{m \mid \mathcal{N}(\mathfrak{g}; m) \neq \emptyset\}.$$

Note that each module $N \in \mathcal{N}(\mathfrak{g}; m)$ is a quotient of $M(\lambda) = \text{Ind}_{\mathfrak{b}}^{\mathfrak{g}} L_{\mathfrak{b}}(\lambda)$. We set

$$(24) \quad m(\mathfrak{g}; \mathfrak{p}; \lambda; \nu) := \max\{m \mid \exists N \in \mathcal{N}(\mathfrak{g}; m) \text{ which is a quotient of } \text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} L_{\mathfrak{p}}(\lambda)\}.$$

6.2.4. **Corollary.** Take $\lambda \neq \nu \in \mathfrak{h}^*$ with $Re(\lambda - \nu)(h) \geq 0$.

- (i) If $\text{Ext}^1(L(\lambda), L(\nu)) \neq 0$, then $\lambda > \nu$;
- (ii) $\text{Ext}^1(L(\lambda), L(\nu)) = \text{Ext}_{\mathcal{O}}^1(L(\lambda), L(\nu))$;
- (iii) $\dim \text{Ext}^1(L(\lambda), L(\nu)) = m(\mathfrak{g}; \mathfrak{b}; \lambda; \nu) \leq \dim M(\lambda)_{\nu}$.

6.3. **Remark.** If \mathfrak{g} is a Kac-Moody superalgebra, then \mathfrak{g} admits antiautomorphism which stabilizes the elements of \mathfrak{h} and the category $\mathcal{O}(\mathfrak{g})$ admits a duality functor $\#$ with the property $L^{\#} \cong L$ for each simple module $L \in \mathcal{O}(\mathfrak{g})$. In this case

$$\dim \text{Ext}^1(L(\lambda), L(\nu)) = \dim \text{Ext}^1(L(\nu), L(\lambda)).$$

6.4. The following lemma is a slight reformulation of Lemma 6.3 in [27].

Lemma. Take $\lambda, \nu \in \mathfrak{h}^*$ with $\lambda > \nu$.

- (i) $m(\mathfrak{g}; \mathfrak{b}; \lambda; \nu) \leq m(\mathfrak{p}; \mathfrak{b}; \lambda; \nu)$ if $\nu - \lambda \in \mathbb{N}\Delta^-(\mathfrak{p})$;
- (ii) $m(\mathfrak{g}; \mathfrak{b}; \lambda; \nu) = m(\mathfrak{g}; \mathfrak{p}; \lambda; \nu)$ if $\nu - \lambda \notin \mathbb{N}\Delta^-(\mathfrak{p})$.

Proof. Write $\mathfrak{p} = \mathfrak{g}^z \rtimes \mathfrak{m}$ as in 6.2.1. For each \mathfrak{g} -module M set

$$\text{Res}(M) := \{v \in M \mid zv = \lambda(z)v\}.$$

Note that $\text{Res}(M)$ is a \mathfrak{g}^z -module; we view $\text{Res}(M)$ as a \mathfrak{p} -module with the zero action of \mathfrak{m} . This defines an exact functor $\text{Res} : \mathfrak{g}\text{-Mod} \rightarrow \mathfrak{p}\text{-Mod}$. By the PBW Theorem

$$\text{Res}(M(\mu)) = \begin{cases} M_{\mathfrak{p}}(\mu) & \text{if } \mu(z) = \lambda(z) \\ 0 & \text{if } (\lambda - \mu)(z) < 0 \end{cases}$$

Let us show that

$$(25) \quad \text{Res}(L(\mu)) = \begin{cases} L_{\mathfrak{p}}(\mu) & \text{if } \mu(z) = \lambda(z) \\ 0 & \text{if } (\lambda - \mu)(z) < 0. \end{cases}$$

Indeed, since Res is exact, $\text{Res}(L(\mu))$ is a quotient of $\text{Res}(M(\mu))$; this gives the second formula. For the first formula assume that $\mu(z) = \lambda(z)$ and that E is a proper submodule of $\text{Res}(L(\mu))$. Since E is a \mathfrak{p} -module we have $\mathcal{U}(\mathfrak{g})E = \mathcal{U}(\mathfrak{n}^-)E$. Since $\text{Res}(L(\mu))$ is a quotient of $\text{Res}(M(\mu))$ and

$$(\text{Res}(M(\mu)))_\mu = (M_{\mathfrak{p}}(\mu))_\mu$$

is a simple $\mathfrak{g}^{\mathfrak{h}}$ -module, one has $\gamma < \mu$ for each $\gamma \in \Omega(E)$. Therefore $(\mathcal{U}(\mathfrak{n}^-)E)_\mu = 0$, so $\mathcal{U}(\mathfrak{g})E$ is a proper \mathfrak{g} -submodule of $L(\mu)$. Hence $E = 0$, so $\text{Res}(L(\mu))$ is simple. This establishes (25).

Now we fix a non-negative integer $m \leq m(\mathfrak{g}; \mathfrak{b}; \lambda; \nu)$ and $N \in \mathcal{N}(\mathfrak{g}; m)$.

Consider the case when $\nu - \lambda \in \mathbb{N}\Delta^-(\mathfrak{p})$. Then $\lambda(z) = \nu(z)$, so $\text{Res}(L(\nu)) = L_{\mathfrak{p}}(\nu)$. Since Res is exact one has

$$\text{Soc}(\text{Res } N) = L_{\mathfrak{p}}(\nu)^{\oplus m}, \quad \text{Res}(N)/\text{Soc}(\text{Res}(N)) = L_{\mathfrak{p}}(\lambda)$$

and $\text{Res}(N)$ is a quotient of $M_{\mathfrak{p}}(\lambda)$. Therefore $m \leq m(\mathfrak{p}; \mathfrak{b}; \lambda; \nu)$. This gives (i).

For (ii) $\nu - \lambda \notin \mathbb{N}\Delta^-(\mathfrak{p})$. Let us show that N is a quotient of $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} L_{\mathfrak{p}}(\lambda)$. Write

$$\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} L_{\mathfrak{p}}(\lambda) = M(\lambda)/J, \quad L_{\mathfrak{p}}(\lambda) = M_{\mathfrak{p}}(\lambda)/J'$$

where J (resp., J') is the corresponding submodule of $M(\lambda)$ (resp., of $M_{\mathfrak{p}}(\lambda)$). Since $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}}$ is exact and $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} M_{\mathfrak{p}}(\lambda) = M(\lambda)$ one has $J \cong \text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} J'$; in particular, each maximal element in $\Omega(J)$ lies in $\Omega(J')$. Note that

$$\Omega(J') \subset \lambda - \mathbb{N}\Delta^+(\mathfrak{l}).$$

Let $\phi : M(\lambda) \twoheadrightarrow N$ be the canonical surjection. Since $J_{\lambda} = 0$, $\phi(J)$ is a proper submodule of N , so $\phi(J)$ is a submodule of $\text{Soc}(N) = L(\nu)^{\oplus m}$.

If $\phi(J) \neq 0$, then ν is a maximal element in $\Omega(J)$ and so $\lambda - \nu \in \mathbb{N}\Delta^+(\mathfrak{l})$, which contradicts to $\nu - \lambda \in \mathbb{N}\Delta^-(\mathfrak{p})$. Therefore ϕ induces a map $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} L_{\mathfrak{p}}(\lambda) = M(\lambda)/J \twoheadrightarrow N$. Therefore

$$m(\mathfrak{g}; \mathfrak{b}; \lambda; \nu) \leq m(\mathfrak{g}; \mathfrak{p}; \lambda; \nu).$$

Since $\text{Ind}_{\mathfrak{p}}^{\mathfrak{g}} L_{\mathfrak{p}}(\lambda)$ is a quotient of $M(\lambda)$, we have $m(\mathfrak{g}; \mathfrak{b}; \lambda; \nu) \geq m(\mathfrak{g}; \mathfrak{p}; \lambda; \nu)$. Thus $m(\mathfrak{g}; \mathfrak{b}; \lambda; \nu) = m(\mathfrak{g}; \mathfrak{p}; \lambda; \nu)$ as required. \square

6.5. Corollary. *Let $z_1, \dots, z_{k-1} \in \mathfrak{h}$ satisfying (21) be such that $\mathfrak{t}^{z_i} \subset \mathfrak{t}^{z_{i+1}}$. Set $\mathfrak{t}^{(i)} := \mathfrak{t}^{z_i}$ and consider the chain*

$$\mathfrak{h} =: \mathfrak{t}^{(0)} \subset \mathfrak{p}^{(1)} \subset \mathfrak{p}^{(2)} \subset \dots \subset \mathfrak{t}^{(k)} := \mathfrak{g}.$$

For $\lambda \neq \nu \in \mathfrak{h}^*$ with $\lambda > \nu$ one has

$$\text{Ext}^1(L(\lambda), L(\nu)) \leq m(\mathfrak{t}^{(s)}; \mathfrak{p}; \lambda; \nu),$$

where s is minimal such that $\nu - \lambda \in \mathbb{N}\Delta^-(\mathfrak{t}^{(s)})$ and $\mathfrak{p} := (\mathfrak{t}^{(s-1)} + \mathfrak{b}) \cap \mathfrak{t}^{(s)}$.

Proof. Take $\mathfrak{p}^{(i)} := \mathfrak{t}^{(i)} + \mathfrak{b}$. Combining Corollary 6.2.4 and Lemma 6.4 we obtain

$$\mathrm{Ext}^1(L(\lambda), L(\nu)) = m(\mathfrak{g}; \mathfrak{b}; \lambda; \nu) \leq m(\mathfrak{p}^{(s)}; \mathfrak{p}^{(s-1)}; \lambda; \nu).$$

Recall that $\mathfrak{p}^{(i)} = \mathfrak{t}^{(i)} \rtimes \mathfrak{m}^{(i)}$, where $\mathfrak{m}^{(i)}$ is the maximal ideal in $\mathfrak{p}^{(i)}$ which lie in \mathfrak{n} . One has $\mathfrak{m}^{(s)} \subset \mathfrak{m}^{(s-1)}$. In particular, $\mathfrak{m}^{(s)}$ annihilates $\mathrm{Ind}_{\mathfrak{p}^{(s-1)}}^{\mathfrak{p}^{(s)}} L_{\mathfrak{p}^{(s-1)}}(\lambda)$, so

$$m(\mathfrak{p}^{(s)}; \mathfrak{p}^{(s-1)}; \lambda; \nu) = m(\mathfrak{t}^{(s)}; \mathfrak{p}^{(s-1)}/\mathfrak{m}^{(s)}; \lambda; \nu).$$

Since

$$\mathfrak{p}^{(s-1)} = \mathfrak{t}^{(s-1)} + \mathfrak{b} \subset \mathfrak{p}^{(s)} = \mathfrak{t}^{(s)} \rtimes \mathfrak{m}^{(s)}$$

the image of $\mathfrak{p}^{(s-1)}$ in $\mathfrak{t}^{(s)} = \mathfrak{p}^{(s)}/\mathfrak{m}^{(s)}$ coincides with $\mathfrak{p} \subset \mathfrak{t}^{(s)}$. \square

6.5.1. *Remark.* Consider the special case when $\mathfrak{g}^{\mathfrak{h}} = \mathfrak{h}$ and $\mathfrak{t}^{(s)} = \mathfrak{l}' \times \mathfrak{h}''$, where $\mathfrak{h}'' \subset \mathfrak{h}$. Take

$$\mathfrak{h}' := \mathfrak{l} \cap \mathfrak{h}, \quad \mathfrak{b}' := \mathfrak{l} \cap \mathfrak{b}, \quad \mathfrak{p}' := \mathfrak{l}' \cap \mathfrak{p}.$$

and let h' be the image of z_s in \mathfrak{h}' . Then the triple $(\mathfrak{l}'; \mathfrak{h}'; h')$ satisfies (H1), (H2) and \mathfrak{b}' is the Borel subalgebra of \mathfrak{l}' . For each $\lambda \in \mathfrak{h}^*$ we set

$$\lambda' := \lambda|_{\mathfrak{h}'}$$

Since $\nu - \lambda \in \mathbb{N}\Delta^-(\mathfrak{t}^{(s)})$ one has

$$m(\mathfrak{t}^{(s)}; \mathfrak{p}; \lambda; \nu) = m(\mathfrak{l}', \mathfrak{p}', \lambda', \nu').$$

Assume, in addition, that $L_{\mathfrak{l}'}(\lambda')$, $L_{\mathfrak{l}'}(\nu')$ are finite-dimensional. It is not hard to see that $\mathrm{Ind}_{\mathfrak{p}'}^{\mathfrak{l}'}(L_{\mathfrak{p}'}(\lambda'))$ admits a unique maximal finite-dimensional subquotient which we denote by $\Gamma_{\mathfrak{l}', \mathfrak{p}'}(L_{\mathfrak{p}'}(\lambda'))$ and that for any finite-dimensional quotient N' of $\mathrm{Ind}_{\mathfrak{p}'}^{\mathfrak{l}'}(L_{\mathfrak{p}'}(\lambda'))$ there exists an epimorphism $\Gamma_{\mathfrak{l}', \mathfrak{p}'}(L_{\mathfrak{p}'}(\lambda')) \rightarrow N'$. This implies

$$\mathrm{Ext}^1(L(\lambda), L(\nu)) \leq m(\mathfrak{l}', \mathfrak{p}', \lambda', \nu') \leq [\Gamma_{\mathfrak{l}', \mathfrak{p}'}(L_{\mathfrak{p}'}(\lambda')) : L_{\mathfrak{l}'}(\nu')].$$

REFERENCES

- [1] J. Brundan, *Kazhdan-Lusztig polynomials and character formulae for Lie superalgebra $\mathfrak{q}(n)$* , Adv. Math. **182** (2004), 28–77.
- [2] J. Brundan, C. Stroppel, *Highest weight categories arising from Khovanov’s diagram algebra. IV: the general linear supergroup.*, J. Eur. Math. Soc. (JEMS), **14**, 2, (2012).
- [3] V. Deodhar, O. Gabber, V. Kac, *Structure of some categories of representations of infinite-dimensional Lie algebras*, Adv. in Math. **45** (1982), 92–116.
- [4] J. Dixmier, *Représentations irréductibles des algèbres de Lie nilpotentes*, An. Acad. Brasil. Ci. **35**, 4(1963) 91–519 (French).
- [5] M. Duflo, V. Serganova, *On associated variety for Lie superalgebras*, arXiv:0507198.
- [6] M. Ehrig, C. Stroppel, *On the category of finite-dimensional representations of $OSP(r|2n)$, Part I*, Representation theory — current trends and perspectives, 109–170, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2017.
- [7] M. Ehrig, C. Stroppel, *On the category of finite-dimensional representations of $OSP(r|2n)$, Part II*, preprint.

- [8] I. Entova-Aizenbud, V. Serganova, *Duflo-Serganova functor and superdimension formula for the periplectic Lie superalgebra*, arXiv: 1910.02294, (2019)
- [9] J. Germoni, *Indecomposable representations of special linear Lie superalgebras*, J. of Algebra, **209**, (1998), 367–401.
- [10] J. Germoni, *Indecomposable representations of $\mathfrak{osp}(3, 2)$ $D(2, 1; \alpha)$ and $G(3)$* in Colloquium on Homology and Representation Theory, Vaqueras, 1998, Bol. Acad. Nac. Cienc. (Córdoba) **65** (2000), 147–163.
- [11] M. Gorelik, *Depths and cores in the light of DS-functors*, arXiv: 2010.05721.
- [12] M. Gorelik, *On modified extension graphs of a fixed atypicality*, arXiv:2204.02759.
- [13] M. Gorelik, T. Heidersdorf, *Semisimplicity of the DS functor for the orthosymplectic Lie superalgebra*, arXiv: 2010.14975.
- [14] M. Gorelik, T. Heidersdorf, *Gruson-Serganova character formulas and the Duflo-Serganova cohomology functor*, arXiv:2104.12634.
- [15] M. Gorelik, C. Hoyt, V. Serganova, A. Sherman, *The Duflo-Serganova functor, vingt ans après*, arXiv: 2203.00529.
- [16] M. Gorelik, V. Serganova, *Integrable modules over affine Lie superalgebras $\mathfrak{sl}(1|n)^{(1)}$* , Comm. Math. Phys. **364** (2018), no. 2, 635–654
- [17] N. Grantcharov, V. Serganova, *Extension quiver for Lie superalgebra \mathfrak{q}_3* , arXiv:2008.10649.
- [18] C. Gruson, V. Serganova, *Cohomology of generalized supergrassmannians and character formulae for basic classical Lie superalgebras*, Proc. London Math. Soc., (3), **101** (2010), 852–892.
- [19] C. Gruson, V. Serganova, *Bernstein-Gelfand-Gelfand reciprocity and indecomposable projective modules for classical algebraic supergroups*, Mosc. Math. J., **13** (2013), no. 2, 281–313.
- [20] C. Gruson, V. Serganova, *Category of finite-dimensional modules over an orthosymplectic Lie superalgebra: small rank examples*, in Symmetries, integrable systems and representations, 155–173, Springer Proc. Math. Stat., 40, Springer, Heidelberg, 2013.
- [21] T. Heidersdorf, R. Weissauer, *Cohomological tensor functors on representations of the general linear supergroup*, arXiv:1406.0321, to appear in Mem. Am. Math. Soc.
- [22] C. Hoyt, S. Reif, *Grothendieck rings for Lie superalgebras and the Duflo-Serganova functor*, Algebra Number Theory **12** (2018), no. 9, 2167–2184.
- [23] V. G. Kac, *Lie superalgebras*, Adv. in Math., **26**, no. 1 (1977), 8–96.
- [24] V. G. Kac, M. Wakimoto, *Integrable highest weight modules over affine superalgebras and Applel’s function*, Commun. Math. Phys. **215** (2001), 631–682.
- [25] L. Martirosyan, *The representation theory of the exceptional Lie superalgebras $F(4)$ and $G(3)$* , J. of Algebra, **419** (2014), 167–222.
- [26] V. Mazorchuk, V. Miemietz, *Serre functors for Lie algebras and superalgebras*, Ann. de l’Institut Fourier **62** (2012), no.1, 47–75.
- [27] I. Musson, V. Serganova, *Combinatorics of character formulas for the Lie superalgebra $\mathfrak{gl}(m|n)$* , Transform. Groups **16** (2011), no. 2, 555–578.
- [28] I. Penkov, *Characters of typical irreducible finite-dimensional $\mathfrak{q}(n)$ -modules*, Funct. Anal. Appl. **20** (1986), 30–37.
- [29] I. Penkov, *Borel-Weil-Bott theory for classical Lie supergroups*, J. Soviet Math. **51** (1990), 2108–2140 (Russian).
- [30] I. Penkov, V. Serganova, *Characters of finite-dimensional irreducible $\mathfrak{q}(n)$ -modules*, Lett. Math. Phys. **40** (1997), 147–158.
- [31] I. Penkov, V. Serganova, *Characters of irreducible G -modules and cohomology of G/P for the supergroup $G = Q(N)$* , J. Math. Sci., **84**, (1997), 1382–1412.
- [32] V. Serganova, *Kazhdan–Lusztig polynomials and character formula for the Lie superalgebra $\mathfrak{gl}(m|n)$* , (N.S.) **2**, Selecta Math. (1996), 607–651.

- [33] V. Serganova, *On a superdimension of an irreducible representation of a basic classical Lie superalgebras*, in *Supersymmetry in mathematics and physics*, 253–273, Lecture Notes in Math., 2027, Springer, Heidelberg, 2011.
- [34] V. Serganova, *Kac-Moody superalgebras and integrability*, in *Developments and trends in infinite-dimensional Lie theory*, 169–218, Progr. Math., 288, Birkhäuser Boston, Inc., Boston, MA, 2011.
- [35] Yucai Su, R. B. Zhang, *Character and dimension formulae for queer Lie superalgebra*, *Commun. Math. Phys.* 333, No. 3, 1465–1481 (2015).

DEPT. OF MATHEMATICS, THE WEIZMANN INSTITUTE OF SCIENCE, REHOVOT 76100, ISRAEL

Email address: maria.gorelik@weizmann.ac.il