

Yang-Mills sources in biconformal gravity

Davis W. Muhwezi* and James T. Wheeler†

October 10, 2021

Abstract

Biconformal gravity, based on gauging of the conformal group to $2n$ dimensions, reproduces n -dim scale-covariant general relativity on the co-tangent bundle in any dimension. We generalize this result to include Yang-Mills matter sources formulated as $SU(N)$ gauge theories with a twisted action on the full $2n$ -dimensional biconformal space. We show that the coupling of the sources to gravity does not stop the reduction of effective dimension $2n \rightarrow n$ of the gravity theory, and instead forces the Yang-Mills source to reduce to ordinary n -dimensional Yang-Mills theory on the gravitating cotangent bundle, with the usual Yang-Mills energy tensor as gravitational source. The results apply as well to gravity with sources on Kähler manifolds and in double field theories.

*Utah State University Department of Physics, dvsmuhwezi@gmail.com

†Utah State University Department of Physics, jim.wheeler@usu.edu

1 Introduction

In general relativity the coupling of matter sources to gravity is accomplished by making the matter action invariant under general coordinate transformations, then adding it to the Einstein-Hilbert action. In formulations of general relativity based on the conformal group, there may be additional conditions. Here we examine $SU(N)$ gauge theories as sources for gravity in a large class of spaces of doubled dimension.

Doubled dimension of spacetime arises in various contexts, frequently related to the idea of a relativistic phase space. Born reciprocity (M. Born, [1, 2]) was one early suggestion aimed at unifying relativity and quantum theory. The reciprocity involves the scaled symplectic exchange $x^\alpha \rightarrow ap^\alpha, p^\beta \rightarrow -bx^\beta$, thereby preserving Hamilton's equations. Further developments include the study of Kähler manifolds, with mutually compatible metric, symplectic, and complex structures such that any two of the structures yield the third.

In 1982, using a gauge theory approach to gravity, Ivanov and Niederle [3, 4] showed that general relativity can arise in a space of doubled dimension called biconformal space. Generalizing the 8-dimensional quotient of the conformal group of spacetime by its homogeneous Weyl subgroup led the authors to a class of curved geometries. From an action quadratic in the curvatures they found that suitable constraints reduced the field equations to the Einstein equation in 4-dimensions.

The group quotient approach used in [3, 4] generalizes to arbitrary dimension n and signature (p, q) . The quotient of conformal group of an n dimensional space by its Weyl subgroup yields a $2n$ dimensional biconformal manifold. In [5] it was shown that biconformal spaces of any dimension $2n$ admit an action *linear* in the curvature with field equations reducing to the vacuum Einstein equation in n dimensions. Unlike the other double field theories cited below, this reduction in both field count and the number of independent variables occurs *by virtue of the field equations*. Starting from the most general action linear in the curvatures—which takes essentially the same form in any dimension—it has now been shown that the field equations generically lead to scale invariant general relativity on the co-tangent bundle [5, 6]. As in Riemannian geometry, these biconformal spaces are taken to be torsion free.

Because biconformal spaces have natural symplectic structure [7] they give an arena appropriate to quantum problems [8], and when they are required to fully reproduce the properties of phase space, the $(3, 1)$ signature of spacetime emerges necessarily from an originally Euclidean space [9, 7, 10]. It is further shown in [6, 7, 10, 11] that these spaces are generically Kähler, and that they share the properties of double field theories.

Some years after the first biconformal spaces, another form of doubled dimension called double field theories arose as a means of making the $O(d, d)$ symmetry of T -duality manifest. By introducing scalars

to produce an additional d dimensions, Duff [12] doubled the $X(\sigma, \tau)$ string variables to make this $O(d, d)$ symmetry manifest. Siegel brought the idea to full fruition by deriving results from superstring theory [13, 14, 15]. Allowing fields to depend on all $2d$ coordinates, Siegel introduced generalized Lie brackets, gauge transformations, covariant derivatives, and a section condition on the full doubled space, thereby introducing torsions and curvatures in addition to the manifest T -duality. By restricting half the coordinates—called imposing a *section condition*—one recovers the d -dimensional theory.

In any of these doubled dimension gravity theories it is desirable to understand matter couplings, and preferable to introduce the matter fields *ab initio* in the doubled space. Carrying out such an investigation in biconformal spaces simultaneously gives results applicable to other doubled spaces. Because biconformal gravity is a gauge theory, it allows direct extension of the symmetry to include gravitational sources from $SU(N)$ gauge theories. However, such Yang-Mills type sources must be written in the space of doubled dimension. This means the introduction of far more potential source fields, $\frac{2n(2n-1)}{2} \times (N^2 - 1)$ instead of only $\frac{n(n-1)}{2} \times (N^2 - 1)$, each depending on $2n$ independent variables instead of only n . There are two principal questions we address here. First is the question of the correct form of the Yang-Mills action in the doubled dimension. We find that the usual $\int tr(\mathcal{F}^* \mathcal{F})$ form must be augmented with a “twist” to reproduce familiar results, similar to the twist found in studies of double field theories [16, 17, 18, 19]. The second question is whether the increase in fields and independent variables spoils the gravitational reduction, or at the other extreme, shares the reduction to n dimensions with gravity. We find that both gravity and sources reduce to n -dimensions. The reduction again occurs by virtue of the field equations, with no need for section conditions.

A further result arises from this investigation. With the exceptions of a study of biconformal supersymmetry [20], which necessarily includes matter fields, and of a brief study [21] with scalar field sources, previous solutions and further properties of biconformal gravity [4, 5, 6, 7, 10, 22, 23] have been based on pure gravity solutions. The field equations for pure biconformal gravity arise by variation of the gauge fields, and it has not been necessary to introduce a metric. However, the actions for Yang-Mills theories involve the Hodge dual and therefore a metric. But biconformal spaces possess both conformal and Kähler structures, it is not clear which of these should take precedence in defining the orthonormality of the basis gauge fields. The ambiguity in identifying the metric is confounded with the specification of the biconformal matter action. The outcome of the current investigation is that only the Killing form reproduces the expected coupling to general relativity.

Concretely, biconformal spaces are spanned by two sets of frame fields, $(\mathbf{e}^a, \mathbf{f}_b)$, called the solder form

and the co-solder form. In solutions, the solder form, $\mathbf{e}^a$, reduces via the field equations to the usual solder form on spacetime. In order to write any Yang-Mills action we are compelled to place an orthonormality condition on these forms,

$$\begin{aligned}\langle \mathbf{e}^a, \mathbf{e}^b \rangle &= M^{ab} \\ \langle \mathbf{e}^a, \mathbf{f}_b \rangle &= M^a{}_b \\ \langle \mathbf{f}_a, \mathbf{f}_b \rangle &= M_{ab}\end{aligned}$$

where some invariant matrix,

$$M^{AB} = \begin{pmatrix} M^{ab} & M^a{}_b \\ M_a{}^b & M_{ab} \end{pmatrix}$$

must be specified. A central hurdle in the course of our study was that there are two natural candidates for M^{AB} : the Killing form of the conformal group restricted to the base manifold, and the Kähler metric of the Kähler structure. We carry out the search for a suitable Yang-Mills action for each candidate symmetric form. Ultimately, we find that only the Killing form can give the expected coupling to gravitation. It is therefore the Killing form that provides the orthonormality of the solder and co-solder forms throughout the remainder of the paper.

Before proceeding with our investigation of $SU(N)$ sources in biconformal/double-field-theory/Kähler gravity, we look briefly at sources in other gravity theories. We first exhibit general relativity with Yang-Mills sources as a gauge theory. This displays our general approach to gauging, as well as the reduced result we hope to achieve. In the remainder of this Section, we look at the modifications of other conformally based gravity theories required in order to include sources. In general relativity, we must extend the symmetry of source actions to general coordinate from global Lorentz. By analogy, we expect that including matter in conformally based theories may require some additional conditions. We briefly discuss sources in scale-invariant gravity and Weyl gravity, before concluding the Section with a discussion of sources in doubled dimension theories.

1.1 Yang-Mills matter in general relativity

Defining a projection on the quotient of the $\frac{n(n+1)}{2}$ -dimensional Poincaré group $\mathcal{P}$ by its $\frac{n(n-1)}{2}$ -dimensional Lorentz subgroup $\mathcal{L}$ gives a principal fiber bundle with Lorentz fibers over an n -dimensional Minkowski spacetime. Generalizing the base space by changing the Maurer-Cartan connection forms of the Poincaré

group and perhaps changing the manifold, the fiber structure is maintained by demanding horizontality of the curvature and torsion. The result is a Riemann-Cartan geometry characterized by curvature and torsion with local Lorentz symmetry.

Concretely, the generalization of the connection $(\tilde{\mathbf{e}}^b, \tilde{\omega}^a{}_b) \Rightarrow (\mathbf{e}^b, \omega^a{}_b)$ takes the Maurer-Cartan equations of the Poincaré group,

$$\begin{aligned} d\tilde{\omega}^a{}_b &= \tilde{\omega}^c{}_b \wedge \tilde{\omega}^a{}_c \\ d\tilde{\mathbf{e}}^a &= \tilde{\mathbf{e}}^b \wedge \tilde{\omega}^a{}_b \end{aligned}$$

to the Cartan equations,

$$d\omega^a{}_b = \omega^c{}_b \wedge \omega^a{}_c + \mathbf{R}^a{}_b \quad (1)$$

$$d\mathbf{e}^a = \mathbf{e}^b \wedge \omega^a{}_b + \mathbf{T}^a \quad (2)$$

where horizontality of the curvature $\mathbf{R}^{ab}$ and the torsion $\mathbf{T}^a$ is captured by omitting any occurrence of the spin connection when writing them expressly as 2-forms

$$\begin{aligned} \mathbf{R}^a{}_b &= \frac{1}{2} R^a{}_{bcd} \mathbf{e}^c \wedge \mathbf{e}^d \\ \mathbf{T}^a &= \frac{1}{2} T^a{}_{bc} \mathbf{e}^b \wedge \mathbf{e}^c \end{aligned}$$

Horizontality insures the survival of the principal fiber bundle by guaranteeing that integrals of the curvatures over an area, or equivalently of the connection forms over closed curves, are independent of lifting.

Completing the description of the Riemann-Cartan geometry is the demand for integrability of the Cartan equations, which follows by exterior differentiation of Eqs.(1) and (2):

$$\begin{aligned} D\mathbf{R}^a{}_b &= 0 \\ D\mathbf{T}^a &= \mathbf{e}^b \wedge \mathbf{R}^a{}_b \end{aligned}$$

When torsion vanishes, this construction describes a general n -dimensional Riemannian spacetime with local Lorentz symmetry.

To include an additional $SU(N)$ Yang-Mills symmetry in the fiber bundle we extend the $\mathcal{P}/\mathcal{L}$ quotient

to the quotient of the product $\mathcal{P} \times SU(N)$ by the product $\mathcal{L} \times SU(N)$:

$$\mathcal{P} \times SU(N) / \mathcal{L} \times SU(N)$$

and carry out the same procedure. This still results in an n -dimensional spacetime but now the fibers of the principal bundle are isomorphic to $\mathcal{L} \times SU(N)$. The Cartan Eqs.(1) and (2) are augmented by a third equation,

$$d\mathbf{A}^i = -\frac{1}{2}c^i{}_{jk}\mathbf{A}^j \wedge \mathbf{A}^k + \mathbf{F}^i$$

where indices beginning with i have range $i, j, k, \dots = 1, 2, \dots, N^2 - 1$, and $\mathbf{F}^i$ is horizontal

$$\mathbf{F}^i = \frac{1}{2}F^i{}_{ab}\mathbf{e}^a \wedge \mathbf{e}^b$$

The integrability condition is $\mathcal{D}\mathbf{F}^i = 0$, where $\mathcal{D}$ is the $SU(N)$ covariant derivative. Here, indices from the first part of the alphabet have range $a, b, \dots = 1, \dots, n$.

To build a physical theory, we write an action functional using any of the tensor fields arising from the construction, $\mathbf{R}^a{}_b, \mathbf{T}^a, \mathbf{F}^i, \mathbf{e}^a, \eta_{ab}, e_{ab\dots c}$, together with any representations of the original group.

In any dimension of spacetime, the action coupling the $SU(N)$ Yang-Mills field to general relativity is written as

$$\begin{aligned} S &= S_{GR} + S_{YM} \\ &= \int \mathbf{R}^{ab} \wedge \mathbf{e}^c \wedge \dots \wedge \mathbf{e}^d e_{abc\dots d} - \frac{\kappa}{2} \int \mathbf{F}^i \wedge {}^* \mathbf{F}_i \end{aligned}$$

where ${}^*\mathbf{F}_i$ is the Hodge dual of the 2-form $\mathbf{F}_i$. We vary the action with respect to the solder form, $\mathbf{e}^a$, the spin connection, $\omega^a{}_b$, and the Yang-Mills connection $\mathbf{A}^i$. Making the usual assumptions for the gravity theory to reduce to general relativity, this results in

$$\begin{aligned} R_{ab} - \frac{1}{2}\eta_{ab}R &= \kappa \left(\eta^{cd}F^i{}_{ac}F_{i\,bd} - \frac{1}{4}\eta_{ab}F^{i\,cd}F_{i\,cd} \right) \\ \tilde{D}^c F^i{}_{ac} &= 0 \end{aligned} \tag{3}$$

where $\tilde{D}^c$ is covariant with respect to both local Lorentz and local $SU(N)$ transformations. These methods

generalize immediately to additional internal symmetries, such as the $SU(3) \times SU(2) \times U(1)$ of the standard model.

Our main result is to show that Eqs.(3) in n -dimensions follow from the field equations of biconformal gravity coupled to a twisted Yang-Mills matter action, formulated in $2n$ -dimensions.

1.2 Sources for scale-invariant gravity

In addition to general covariance, scale-invariant gravity typically requires tracelessness of the energy tensor. This condition arises when local scale invariance is a symmetry of the action. Let S_g be any locally scale invariant gravity action, with sources supplied by adding a matter action,

$$S_{matter} = \int \mathcal{L}_{matter} \sqrt{-g} d^4x$$

Then the source for gravity is the energy-momentum tensor given by the metric variation,

$$\begin{aligned} \delta_g S_{matter} &= \int \frac{\delta(\mathcal{L}_{matter} \sqrt{-g})}{\delta g^{\alpha\beta}} \delta g^{\alpha\beta} d^4x \\ &= \int T_{\alpha\beta} \delta g^{\alpha\beta} \sqrt{-g} d^4x \end{aligned}$$

If the energy-momentum tensor is scale *invariant*, we may apply Noether's theorem to the scaling symmetry transformation, $\delta g^{\alpha\beta} = -2\phi g^{\alpha\beta}$ immediately. This gives

$$0 \equiv \delta_{symmetry} S_{matter} = -2 \int T_{\alpha\beta} \phi g^{\alpha\beta} \sqrt{-g} d^4x = -2 \int T^\alpha{}_\alpha \phi \sqrt{-g} d^4x$$

and since this must hold for any ϕ , the energy tensor must be traceless, $T^\alpha{}_\alpha = 0$.

The tracelessness follows from the assumption that the only conformal dependence resides in the single metric factor. However, it is more often the case that the energy tensor $T_{\alpha\beta}$ is *covariant* under conformal transformations. For example, consider a simple massless scalar field ϕ with action

$$S_{scalar} = \frac{1}{2} \int g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi \sqrt{-g} d^4x$$

This action is scale invariant under the simultaneous transformations

$$\begin{aligned}\tilde{g}_{\alpha\beta} &= e^{2\alpha} g_{\alpha\beta} \\ \tilde{\phi} &= e^{-\alpha} \phi\end{aligned}$$

since then

$$\begin{aligned}\tilde{S}_{scalar} &= \frac{1}{2} \int \tilde{g}^{\mu\nu} \partial_\mu \tilde{\phi} \partial_\nu \tilde{\phi} \sqrt{-\tilde{g}} d^4x \\ &= \frac{1}{2} \int e^{-2\alpha} g^{\mu\nu} \partial_\mu (e^{-\alpha} \phi) \partial_\nu (e^{-\alpha} \phi) e^{4\alpha} \sqrt{-g} d^4x \\ &= S_{scalar}\end{aligned}$$

The energy-momentum tensor for ϕ is given by metric variation

$$\begin{aligned}\delta_g S_{scalar} &= \frac{1}{2} \int \left(\delta g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi \sqrt{-g} - \frac{1}{2} (g^{\alpha\beta} \partial_\alpha \phi \partial_\beta \phi) g_{\mu\nu} \delta g^{\mu\nu} \sqrt{-g} \right) d^4x \\ &= \frac{1}{2} \int \delta g^{\mu\nu} \left(\partial_\mu \phi \partial_\nu \phi - \frac{1}{2} g_{\mu\nu} (g^{\alpha\beta} \partial_\alpha \phi \partial_\beta \phi) \right) \sqrt{-g} d^4x\end{aligned}$$

as

$$T_{\mu\nu} = \partial_\mu \phi \partial_\nu \phi - \frac{1}{2} g_{\mu\nu} (g^{\alpha\beta} \partial_\alpha \phi \partial_\beta \phi)$$

This is clearly not traceless even though the action is scale invariant. Indeed, under conformal transformation $T_{\mu\nu}$ has conformal weight -2 ,

$$\tilde{T}_{\mu\nu} = e^{-2\alpha} T_{\mu\nu}$$

and this covariance is all that is required to make the action invariant.

In cases such as this, an energy-momentum tensor with definite, nonzero conformal weight may serve as the source for the curvature of a Weyl geometry. By including the Weyl vector in these geometries, even the Weyl-Ricci tensor takes on definite conformal weight:

$$\begin{aligned}\tilde{C}^a{}_{bcd} &= e^{-2\alpha} C^a{}_{bcd} \\ \tilde{\mathcal{R}}_{ab} &= e^{-2\alpha} \left(\mathcal{R}_{ab} + W_{(a;b)} - W_a W_b + \frac{1}{2} W^2 \eta_{ab} \right) \\ \tilde{\mathcal{R}} &= e^{-2\alpha} \left(\tilde{\mathcal{R}} + W^a{}_{;a} + W^2 \right)\end{aligned}$$

where $\mathcal{R}_{ab}$ is the Schouten tensor and W_a the Weyl vector. A scale-invariant action functional in a Weyl geometry differs from the Einstein-Hilbert action. A full discussion of such theories is given in [24].

1.3 Sources for Weyl gravity

We present two possibilities for the inclusion of sources in Weyl gravity.

Weyl, or conformal, gravity begins with the free action,

$$S_{Weyl} = \int C^{\alpha\beta\mu\nu} C_{\alpha\beta\mu\nu} \sqrt{-g} d^4x$$

which is quadratic in the conformal curvature, with metric variation leading to the Bach equation [25],

$$W^{\mu\nu} = D_\alpha D_\beta C^{\mu\alpha\nu\beta} - \frac{1}{2} C^{\mu\alpha\nu\beta} R_{\alpha\beta} = 0 \quad (4)$$

The Bach tensor $W^{\mu\nu}$ is automatically traceless by the tracelessness of the conformal curvature, $g_{\mu\nu} C^{\mu\alpha\nu\beta} = 0$. Because Eq.(4) arises by metric variation it is necessarily symmetric. Since the metric variation has the form

$$\delta_g S_{Weyl} = \int W^{\mu\nu} \delta g_{\mu\nu} \sqrt{-g} d^4x$$

the change of the metric, $\delta g_{\mu\nu} = h_{\mu;\nu} + h_{\nu;\mu}$ under a change of coordinates implies

$$\begin{aligned} 0 &= \int W^{\mu\nu} (h_{\mu;\nu} + h_{\nu;\mu}) \sqrt{-g} d^4x \\ &= -2 \int W^{\mu\nu}{}_{;\nu} h_\mu \sqrt{-g} d^4x \end{aligned}$$

and since h_μ is arbitrary, the Bach tensor is divergence free,

$$W^{\mu\nu}{}_{;\nu} = 0$$

By rewriting the conformal curvature $C^{\alpha\beta\mu\nu}$ in terms of the Riemann tensor minus its traces, and using the Gauss-Bonnet density for the Euler character to remove the Riemann squared term, S_{Weyl} becomes proportional to $S' = \int (R_{\alpha\beta} R^{\alpha\beta} - \frac{1}{3} R^2) \sqrt{-g} d^4x$. Variation of the metric now leads to a field equation

written entirely in terms of the Ricci tensor,

$$\begin{aligned}
0 = & R^{\mu\nu;\beta}{}_{;\beta} - R^{\mu\beta;\nu}{}_{;\beta} - R^{\nu\beta;\mu}{}_{;\beta} - 2R^{\mu\beta}R^\nu{}_\beta - \frac{1}{6}g^{\mu\nu}R^{;\beta}{}_{;\beta} \\
& + \frac{1}{2}g^{\mu\nu}R^{\alpha\beta}R_{\alpha\beta} + \frac{2}{3}R^{;\mu\nu} + 2RR^{\mu\nu} - \frac{1}{6}g^{\mu\nu}R^2
\end{aligned} \tag{5}$$

Clearly, vacuum solutions to the Einstein equation are also solutions to Eq.(5). Thus, it is fair to argue that evidence for vacuum general relativity is also evidence for vacuum Weyl gravity.

It is well known that the converse is false. There are explicit solutions to Eq.(5) that are not solutions to vacuum general relativity.

We cannot make a similar claim that solutions to general relativity with sources gives evidence for Weyl gravity with sources. As we show below, there are alternative candidates for sources for the Bach equation. Without a clear specification of the source, the question of the relationship between solutions in general relativity and solutions in Weyl gravity is ill-posed.

Nonetheless, many argue for or assume [26, 27] direct addition of a matter action with traceless energy tensor,

$$S = \int C^{\alpha\beta\mu\nu}C_{\alpha\beta\mu\nu}\sqrt{-g}d^4x + S_{matter} \tag{6}$$

The resulting metric variation leads to

$$W^{\mu\nu} = \kappa T^{\mu\nu} \tag{7}$$

which is consistent with the properties of the Bach tensor as long as the energy-momentum tensor is traceless.

However, the resulting solutions may differ substantially from the verified predictions of general relativity. Specifically, Flanagan has argued that the fourth order theory fails to agree with solar system values, and Yoon has criticized the Newtonian limit and the application to galactic rotation curves [for the critique and response see [28, 29, 30]]. On the other hand, Palatini-style variation has been shown for the vacuum case to reduce to scale covariant general relativity [31], and a novel coupling to matter might give better agreement with known results.

Because of the fourth order character of Eq.(7), the coupling constant must have units that differ from the Einstein coupling by some inverse length scale squared, L^{-2} . This is expressible as a (very small) mass, $m \sim \frac{\hbar}{Lc}$. Therefore, the nonvacuum solutions to Eq.(7) and solutions to general relativity with the same energy-momentum tensor will differ by terms proportional to m^2 . For sufficiently close agreement with general relativity, this mass must be extremely small. The length scale L might represent a radius of

inversion symmetry. Since inversion is a conformal symmetry, this is a possible interpretation. Consistency would require this length scale to be cosmologically large, with the mass m correspondingly small. The mass-dependent corrections might then be consistent with experiment.

As a second possible coupling of matter to Weyl gravity, we note that the introduction of a scale between general relativity and Weyl gravity is eliminated if the source field equations are also modified to fourth order equations. For example, we may modify the Klein-Gordon action to

$$S_{MKG} = \frac{1}{2} \int (D_\alpha D_\mu \phi D^\alpha D^\mu \phi + m^2 D_\mu \phi D^\mu \phi) \sqrt{-g} d^4x$$

so that the field equation becomes

$$D_\mu \square D^\mu \phi - m^2 \square \phi = 0$$

and commuting $\square D^\mu \phi = D^\mu \square \phi + R^{\mu\alpha} \phi_{;\alpha}$ this becomes

$$\square (\square \phi - m^2 \phi) + \frac{1}{2} R^{\alpha\alpha} \phi_{;\alpha} + R^{\mu\alpha} \phi_{;\alpha\mu} = 0$$

Because we have modified the mass term as well as the original kinetic term, solutions to the original Klein-Gordon equation satisfy this modified field equation up to terms proportional to the curvature. In regions of weak gravity, at least some solutions will be close to the corresponding solution fields in general relativity. Notice that if we identify $B_\alpha = D_\alpha \phi$, the action S_{MKG} describes a restriction of a Proca field.

Varying the metric, the energy tensor for this modified Klein-Gordon field is found to be

$$\mathcal{T}_{\alpha\beta} = g^{\mu\nu} D_\alpha D_\mu \phi D_\beta D_\nu \phi + \frac{1}{2} m^2 D_\alpha \phi D_\beta \phi - \frac{1}{4} (D_\nu D_\mu \phi D^\nu D^\mu \phi + m^2 D_\mu \phi D^\mu \phi) g_{\alpha\beta}$$

This is more transparent if we let $B_\alpha = D_\alpha \phi$,

$$\mathcal{T}_{\alpha\beta} = D_\alpha B^\mu D_\beta B_\mu + \frac{1}{2} m^2 B_\alpha B_\beta - \frac{1}{4} g_{\alpha\beta} (D_\nu B_\mu D^\nu B^\mu + m^2 B_\mu B^\mu)$$

Notice that $\mathcal{T}_{\alpha\beta}$ is traceless if we set $m^2 = 0$. To provide a conformal source, we therefore consider a massless modified Klein-Gordon field. The resulting conserved gravitational equation becomes

$$\begin{aligned} W_{\alpha\beta} &= \kappa \mathcal{T}_{\alpha\beta} \\ &= \kappa \left(g^{\mu\nu} D_\alpha D_\mu \phi D_\beta D_\nu \phi - \frac{1}{4} g_{\alpha\beta} (D_\nu D_\mu \phi D^\nu D^\mu \phi) \right) \end{aligned}$$

Additional curvature terms will occur for higher spin fields, but the basic conclusion remains the same. Solutions to the usual field equation continue to solve the field equation, up to terms proportional to the curvature. As a second example, the action for the general fourth order modification of the Maxwell field may be written as

$$S = \frac{1}{4} \int D_\alpha F_{\mu\nu} D^\alpha F^{\mu\nu} \sqrt{-g} d^4x$$

where $F_{[\alpha\beta;\mu]} = 0$ guarantees the existence of a potential and makes this form unique. This results in the higher order field equation

$$D^\nu \square F_{\mu\nu} = 0$$

which we may once again commute to express in terms of the usual lower order expression $D_\nu F^{\alpha\nu}$ plus curvature terms:

$$\square D_\nu F^{\alpha\nu} + \text{terms linear in } FR = 0$$

As we found for the scalar field, symmetry and vanishing divergence of the energy tensor,

$$\mathcal{T}_{\rho\sigma} = D_\rho F^{\alpha\beta} D_\sigma F_{\alpha\beta} + 2D^\beta F_\rho{}^\mu D_\beta F_{\sigma\mu} - \frac{1}{2} (D_\alpha F_{\mu\nu} D^\alpha F^{\mu\nu}) g_{\rho\sigma}$$

are guaranteed by the metric variation and diffeomorphism invariance.

Tracelessness of $\mathcal{T}_{\alpha\beta}$, however, requires an additional constraint on the derivatives of the field, $\mathcal{T}^\alpha{}_\alpha = D^\sigma F^{\alpha\beta} D_\sigma F_{\alpha\beta}$. Using the field equation, we may express this as a vanishing divergence,

$$\mathcal{T}^\alpha{}_\alpha = D_\sigma (F_{\alpha\beta} D^\sigma F^{\alpha\beta} - 2\square F^{\alpha\sigma} A_\alpha) \stackrel{!}{=} 0$$

indicating the presence of an additional conserved current.

Other field equations might be modified in a similar way. Of course, though these modifications do not necessarily introduce a new mass scale, they still require additional initial conditions.

1.4 Sources for biconformal gravity, Kähler gravity, and double field theory

The situation may not be as complicated for biconformal gravity [4, 23, 5, 6], double field theory [12, 13, 14, 15, 33, 35], or gravity on a Kähler manifold [7, 6]. Each of these cases starts as a fully $2n$ -dimensional theory but ultimately describes gravity on an n -dimensional submanifold. It is desirable to have a fully $2n$ -dimensional form of the matter action which nonetheless also reduces to the expected n -dimensional source

as a consequence of the field equations. It is this condition we address. We discuss the issue in biconformal space, since biconformal gravity is already a gauge theory and it generically includes the structures of both double field theory and Kähler manifolds [7, 6].

For matter fields we restrict our attention to Yang-Mills type sources. We find that although the usual form of $2n$ -dimensional Yang-Mills action gives nonstandard coupling to gravity, including a “twist” matrix in the action corrects the problem.

Biconformal gravity arises as follows. The quotient of the conformal group $\mathcal{C}_{p,q} = SO(p+1, q+1)$ of an $SO(p, q)$ -symmetric space ($p+q = n$, metric η_{ab}) by its homogeneous Weyl subgroup, $\mathcal{W}_{p,q} \equiv SO(p, q) \times SO(1, 1)$, leads to a $2n$ -dimensional homogeneous space with local $\mathcal{W}_{p,q}$ symmetry. This homogeneous space, discussed in [4] and [23] and studied extensively in [6, 7, 10], is found to have compatible symplectic, metric and complex structures, making it Kähler [7]. In addition, the restriction of the Killing form to the base manifold is nondegenerate and scale invariant, and the volume form of the base manifold is scale invariant. The homogeneous space and its curved generalizations are called biconformal spaces.

Ivanov and Niederle [4], wrote a gravity theory on an 8-dimensional biconformal space, using the curvature-quadratic action of Weyl gravity. By a suitable restriction of the coordinate transformations of the extra 4-dimensions, they showed that 4-dimensional general relativity describes the remaining subspace. Subsequently, Wehner and Wheeler [5] introduced a class of $\mathcal{W}$ -invariant actions *linear* in the curvatures, defining biconformal gravity. Curvature-linear actions are possible because the $2n$ -dimensional volume element is scale invariant. Unlike the 4-dimensional theories above with actions quadratic in the curvature, the linear action functionals take the same form in any dimension. The doubled dimension is understood in terms of the symplectic structure, leading to a phase space interpretation for generic solutions. Lagrangian submanifolds represent the physical spacetime and have the original dimension. The class of torsion-free biconformal spaces has been shown to reduce to general relativity on the cotangent bundle of spacetime [6]. These reductions of the model work for any signature (p, q) .

The most general action linear in the biconformal curvatures is given by

$$S = \lambda \int e_{ac \dots d}{}^{be \dots f} (\alpha \Omega^a{}_b + \beta \delta^a_b \Omega + \gamma \mathbf{e}^a \wedge \mathbf{f}_b) \wedge \mathbf{e}^c \wedge \dots \wedge \mathbf{e}^d \wedge \mathbf{f}_e \wedge \dots \wedge \mathbf{f}_f \quad (8)$$

where $\Omega^a{}_b$ is the curvature of the spin connection and Ω is the dilatational curvature. Here $\lambda = \frac{(-1)^n}{(n-1)!(n-1)!}$ is a convenient constant, chosen to eliminate a combinatoric factor and to make our sign conventions agree with [6]. The cotangent bundle is spanned by the pair, $(\mathbf{e}^a, \mathbf{f}_b)$, called the solder form and the co-solder form,

respectively. The variation is taken with respect to all $\frac{(n+1)(n+2)}{2}$ gauge fields.

The reduction of a fully $2n$ -dimensional gravity theory to dependence only on the fields of n -dimensional gravity is a remarkable feature of biconformal gravity. While it has been shown to be a double field theory [12, 13, 14, 15, 34, 35, 6], double field theories require the assumption that fields depend on only half the coordinates. This artificial constraint is called a section condition. In sharp contrast, biconformal solutions do *not* require a section condition, *reducing as a consequence of the field equations* of torsion-free biconformal spaces. Thus, using the torsion-free field equations, the components of the $\frac{(n+1)(n+2)}{2}$ curvatures—initially dependent on $2n$ independent coordinates—reduce to the usual Riemannian curvature tensor in n dimensions. Correspondingly, the n -dim solder form determines all fields, up to coordinate and gauge transformations. Generic, torsion-free, vacuum solutions describe n -dimensional scale-covariant general relativity on the co-tangent bundle.

As noted in the introduction, with the exceptions biconformal supergravity [20] and a scalar field example [32], studies of biconformal spaces [23, 5, 6, 9, 7, 10, 8, 22, 11] have considered pure gravity biconformal spaces, leading to vacuum general relativity.

Here we consider $SU(N)$ Yang-Mills fields as gravitational sources. The central issue is to show that even with a completely general $SU(N)$ gauge theory over a $2n$ -dimensional biconformal space, a full $2n \rightarrow n$ reduction occurs, both for gravity and the Yang-Mills field.

As with the Riemann-Cartan construction of general relativity above, the development of biconformal spaces from group symmetry makes it straightforward to include the additional symmetry of sources. By extending the quotient to

$$\mathcal{M}^{2n} = \mathcal{C}_{p,q} \times SU(N) / \mathcal{W}_{p,q} \times SU(N)$$

the local symmetry is enlarged by $SU(N)$ and we may add a Yang-Mills or similar action to Eq.(8). This construction gives the form of the Yang-Mills field in terms of the potentials, but not the form of the action.

There are then two basic parts to our investigation.

First, we must determine a suitable $2n$ -dimensional action functional for the sources. To accomplish this requires two interdependent specifications:

- Find a form of the Yang-Mills action which gives the usual n -dimensional Yang-Mills source to the Einstein tensor.
- Fix the orthonormality condition for the solder and co-solder form, $M^{AB} = \langle \tilde{\mathbf{e}}^A, \tilde{\mathbf{e}}^B \rangle$ where $\tilde{\mathbf{e}}^A =$

$(\mathbf{e}^a, \mathbf{f}_b)$. This determines the form of the Hodge dual and ultimately the metric variation of the matter action.

In accomplishing these steps we show that the standard Yang-Mills action

$$S_{YM}^0 = \int \text{tr} (\mathbf{F} \wedge^* \mathbf{F})$$

cannot give the right couplings, and find a satisfactory modification

$$S_{YM} = \int \text{tr} (\bar{\mathbf{F}} \wedge^* \mathbf{F})$$

where the twisted field, $\bar{\mathbf{F}}$ is defined below. The twisted action, together with the restricted Killing form as orthonormal metric, give the desired reduction.

The second challenge is then to use the field equations to show:

- The number of field components in $2n$ dimensions reduces to the expected number on n dimensional spacetime.
- The functional dependence of the fields reduces from $2n$ to n independent variables.
- The gravitational source is the usual Yang-Mills stress-energy tensor.
- The $SU(N)$ field equation is the usual n -dim Yang-Mills field equation.

The remainder of our presentation proceeds as follows. In the next Section, we introduce our notation and other conventions. In Section 3, we show that the usual form of Yang-Mills action, S_{YM}^0 , cannot produce the usual coupling to gravity. The twisted form $\bar{\mathbf{F}}$ is developed in Section 4, and the variation of S_{YM} carried out. The Yang-Mills potentials are identified and varied in Section 5. The next Section contains the reduction of the gravitational field equations, as far as possible with the presence of sources. This reduction closely follows reference [6]. We find that the field equations imply certain restrictions that must be applied to the matter sources.

The reduction of the number of fields and the number of independent variables is shown in Section 7 to follow from the full gravitational equations. We find that the reduction of fields that is necessary in the purely gravitational sector also forces reduction of the source fields. The Section concludes with the emergence of both the usual Yang-Mills gravitational source in n -dimensions, and the usual n -dimensional Yang-Mills equation. The final Section includes a brief review of the main results.

2 Notation and conventions

2.1 Conventions with biconformal tensors

2.1.1 Differential forms

The co-tangent space of biconformal manifolds are spanned by two sets of opposite conformal weight orthonormal frame fields, $\tilde{\mathbf{e}}^A = (\mathbf{e}^a, \mathbf{f}_b)$, with lowercase Latin indices $a, b, \dots = 1, 2, \dots, n$ indicating the use of these frames and upper case Latin $A, B, \dots = 1, 2, \dots, 2n$ to denote the pair. Coordinate indices are lower case Greek, $\mu, \nu, \dots = 1, 2, \dots, n$ so that we have, for example,

$$\mathbf{e}^a = e_{\mu}{}^a \mathbf{d}x^{\mu} + e^{\mu a} \mathbf{d}y_{\mu}$$

A general 2-form may be written in the orthonormal basis as

$$\mathcal{F} = \frac{1}{2} F_{ab} \mathbf{e}^a \wedge \mathbf{e}^b + F^a{}_b \mathbf{f}_a \wedge \mathbf{e}^b + \frac{1}{2} F^{ab} \mathbf{f}_a \wedge \mathbf{f}_b$$

It is important to realize that F_{ab} , $F^a{}_b$ and F^{ab} are distinct fields. Therefore, we cannot raise and lower indices in the usual way unless we choose different names for the separate independent components. As compensation for this, the raised or lowered position of an index reflects its conformal weight. Thus, F_{ab} has weight -2 while F^{ab} has weight $+2$. When practical these distinct fields will be given different names,

$$\mathcal{F} = \frac{1}{2} F_{ab} \mathbf{e}^a \wedge \mathbf{e}^b + G^a{}_b \mathbf{f}_a \wedge \mathbf{e}^b + H^{ab} \mathbf{f}_a \wedge \mathbf{f}_b$$

but this can lead to an unnecessary profusion of field names.

When we need to explicitly refer to internal $SU(N)$ indices, the field and its components will be given an additional index from the lower case Latin set $\{i, j, k\}$. Other lower case Latin indices refer to the null orthonormal frame field, $(\mathbf{e}^b, \mathbf{f}_a)$. Thus $F^i{}_{ab}$ represents the components of $\frac{1}{2} F^i{}_{ab} G_i \mathbf{e}^a \wedge \mathbf{e}^b$ where G_i is a generator of $SU(N)$ as i runs from 1 to $N^2 - 1$. In most cases the internal index is suppressed.

Differential forms are written in boldface and always multiplied with the wedge product. For brevity in some longer expressions we omit the explicit wedge between forms. Thus, for example,

$$\mathbf{f}_a \wedge \mathbf{f}_b \wedge \mathbf{f}_c \wedge \mathbf{e}^d \wedge \mathbf{e}^e \Longleftrightarrow \mathbf{f}_{abc} \mathbf{e}^{de}$$

The bold font shows that these are differential forms, and therefore are to be wedged together.

As a compromise between keeping track of conformal weights, while being able to assess the symmetry or antisymmetry of components, we introduce a weight +1 basis $\mathbf{e}^A \equiv (\mathbf{e}^a, \eta^{ab} \mathbf{f}_b)$, where η_{ab} is the n -dimensional metric of the original $SO(p, q)$ -symmetric space, *not* the metric of the biconformal space. Thus, we may write

$$\begin{aligned} \mathcal{F} &= \frac{1}{2} F_{AB} \mathbf{e}^A \wedge \mathbf{e}^B \\ &= \frac{1}{2} F_{ab} \mathbf{e}^a \wedge \mathbf{e}^b + \mathcal{G}_{ab} \eta^{ac} \mathbf{f}_c \wedge \mathbf{e}^b + \frac{1}{2} \mathcal{H}_{ab} \eta^{ac} \eta^{bd} \mathbf{f}_c \wedge \mathbf{f}_d \end{aligned}$$

where we have defined

$$\begin{aligned} \mathcal{G}_{ab} &\equiv \eta_{ac} G^c{}_b \\ \mathcal{H}_{ab} &\equiv \eta_{ac} \eta_{bd} H^{cd} \end{aligned}$$

The use of a different font is important because we are not using the biconformal metric, K_{AB} , to change index positions. It is the original index positions and font, $G^a{}_b$, H^{ab} , that are the proper field components.

The matrix components F_{AB} are then written as

$$\begin{aligned} F_{AB} &= \begin{pmatrix} F_{ab} & F_a{}^b \\ F^a{}_b & F^{ab} \end{pmatrix} \\ &= \begin{pmatrix} F_{ab} & -\mathcal{G}_{ba} \\ \mathcal{G}_{ab} & \mathcal{H}_{ab} \end{pmatrix} \end{aligned} \tag{9}$$

where the form of the upper right corner follows because

$$\begin{aligned} -\mathcal{G}_{ba} \mathbf{e}^a \wedge \eta^{bc} \mathbf{f}_c &= \mathcal{G}_{ba} \eta^{bc} \mathbf{f}_c \wedge \mathbf{e}^a \\ &= G^c{}_a \mathbf{f}_c \wedge \mathbf{e}^a \end{aligned}$$

With this convention, we can meaningfully define the transpose of matrices. Specifically, while the transpose of

$$\begin{pmatrix} F_{ab} & F_a{}^b \\ F^a{}_b & F^{ab} \end{pmatrix}$$

is ill-defined because of the mixed indices on the off-diagonal terms, the transpose of F_{AB} in the weight $+1$ basis is

$$\begin{pmatrix} F_{ab} & -\mathcal{G}_{ba} \\ \mathcal{G}_{ab} & \mathcal{H}_{ab} \end{pmatrix}^t = \begin{pmatrix} F_{ba} & \mathcal{G}_{ba} \\ -\mathcal{G}_{ab} & \mathcal{H}_{ba} \end{pmatrix} \quad (10)$$

For $\mathcal{H}_{ba} = -\mathcal{H}_{ab}$ and $F_{ba} = -F_{ab}$ this is manifestly antisymmetric, as befits a 2-form. Notice that the effect of two transposes is the identity, so this operation provides an involutive automorphism even though η_{ab} is not the biconformal metric.

From $[\mathcal{G}_{ab}]^t = \mathcal{G}_{ba}$ we have

$$[\eta_{ac} G^c{}_b]^t = [\eta_{bc} G^c{}_a]$$

and therefore, $[G^d{}_b]^t = \eta^{ad} \eta_{bc} G^c{}_a$.

2.1.2 Metric

Because all quadrants of the metric are used in the variation, it is desirable to retain both the name and index positions throughout. Since

$$K_{AB} = \begin{pmatrix} K_{ab} & K_a{}^b \\ K^a{}_b & K^{ab} \end{pmatrix}$$

contains all index positions, the inverse metric and its components are written with an overbar,

$$\bar{K}^{AB} = \begin{pmatrix} \bar{K}^{ab} & \bar{K}^a{}_b \\ \bar{K}_a{}^b & \bar{K}_{ab} \end{pmatrix}$$

Thus K_{ab} is the first quadrant of the metric, while $\bar{K}_{ab}$ is the final quadrant of the inverse metric. Here any changes of index position must be indicated with an explicit factor of η_{ab} or η^{ab}

2.1.3 Volume form

It is convenient to define a volume form as $\Phi \equiv *1$ but in defining the Hodge dual operation a number of ambiguities need to be clarified. Because up and down indices have distinct conformal weight, we may partially order the indices on the Levi-Civita tensor. We establish the following conventions:

1. All factors of $\mathbf{f}_a$ are written first, followed by all of the $\mathbf{e}^b$.
2. The Levi-Civita *tensor* is written as $e^{ab\dots c}{}_{de\dots f}$, with all n up indices first. The partially ordered

antisymmetric *symbol* is written as $\varepsilon^{ab\cdots c}_{de\cdots f}$.

3. When taking the dual of a p -form, we sum the components on the first p indices of the Levi-Civita tensor, then introduce factors of -1 to move indices to the default positions. For example, the dual of $\mathbf{H} = H^a{}_b \mathbf{f}_a \wedge \mathbf{e}^b$ is

$$\begin{aligned} {}^* \mathbf{H} &= \frac{1}{(n-1)!(n-1)!} H^a{}_b e_a{}^{bc\cdots d}{}_{e\cdots f} \mathbf{f}_c \wedge \cdots \wedge \mathbf{f}_d \wedge \mathbf{e}^e \wedge \cdots \wedge \mathbf{e}^f \\ &= \frac{(-1)^n}{(n-1)!(n-1)!} H^a{}_b e^{bc\cdots d}{}_{ae\cdots f} \mathbf{f}_c \wedge \cdots \wedge \mathbf{f}_d \wedge \mathbf{e}^e \wedge \cdots \wedge \mathbf{e}^f \\ &= \frac{(-1)^n}{(n-1)!(n-1)!} H^a{}_b e^{bc\cdots d}{}_{ae\cdots f} \mathbf{f}_{c\cdots d} \mathbf{e}^{e\cdots f} \end{aligned}$$

Notice that in the final step, the number of wedged forms in $\mathbf{f}_{c\cdots d}$ may be inferred from the Levi-Civita tensor. Since the Levi-Civita tensor always has n up and n down indices, the number of basis forms of each type is unambiguous. For example,

$$e^{abc\cdots d}{}_{e\cdots f} \mathbf{f}_{c\cdots d} \wedge \mathbf{e}^{e\cdots f} = e^{abc\cdots d}{}_{e\cdots f} \underbrace{\mathbf{f}_c \wedge \cdots \wedge \mathbf{f}_d}_{n-2} \wedge \underbrace{\mathbf{e}^e \wedge \cdots \wedge \mathbf{e}^f}_n$$

is a $2n - 2$ form that includes the wedge product of $n - 2$ factors of $\mathbf{f}_a$ and n factors of $\mathbf{e}^a$.

4. An m -form is a polynomial with each term having different numbers of $\mathbf{e}$'s and $\mathbf{f}$'s, we write the terms in order of increasing number of $\mathbf{f}$'s.
5. The partial ordering of indices on the Levi-Civita tensor reduces the normalization of the volume element from $\frac{1}{(2n)!}$ to $\frac{1}{n!n!}$.

It has been noted elsewhere [36] that there are alternative duals in biconformal space. For instance, we may use the symplectic form instead of the metric to connect indices. The difference resides in the relative signs between the $\mathbf{e}^a$, $\mathbf{f}_a$, and mixed terms. Here we use only the Hodge dual, taking care to keep the correct signs.

With these conventions in mind, we define

$$\begin{aligned} \Phi &\equiv {}^* 1 \\ &= \frac{1}{n!n!} e^{c\cdots d}{}_{e\cdots f} \mathbf{f}_{c\cdots d} \wedge \mathbf{e}^{e\cdots f} \\ &= \frac{1}{n!n!} \sqrt{K} \varepsilon^{c\cdots d}{}_{e\cdots f} \mathbf{f}_{c\cdots d} \wedge \mathbf{e}^{e\cdots f} \end{aligned} \tag{11}$$

and consequently,

$$\begin{aligned} \mathbf{f}_{c\dots d} \wedge \mathbf{e}^{e\dots f} &= \frac{1}{\sqrt{K}} \varepsilon^{e\dots f}{}_{c\dots d} \Phi \\ &= \bar{e}_{c\dots d}{}^{e\dots f} \Phi \end{aligned} \quad (12)$$

where the overbar denotes the contravariant form of the Levi-Civita tensor. The contravariant form satisfies

$$e^{a\dots b}{}_{c\dots d} \bar{e}_{a\dots b}{}^{c\dots d} = n!n!$$

We also need the reduction formulas,

$$\begin{aligned} e^{e\dots f}{}_{mnc\dots d} \bar{e}^{ghc\dots d}{}_{e\dots f} &= n!(n-2)! (\delta_m^g \delta_n^h - \delta_m^h \delta_n^g) \\ e^{mc\dots d}{}_{ne\dots f} \bar{e}^{he\dots f}{}_{gc\dots d} &= (n-1)!(n-1)! \delta_n^h \delta_g^m \end{aligned} \quad (13)$$

2.1.4 Further notation

The antisymmetric projection operator mapping for $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$ tensors,

$$\begin{aligned} \Delta_{db}^{ac} &\equiv \frac{1}{2} (\delta_d^a \delta_b^c - \eta^{ac} \eta_{db}) \\ &= \frac{1}{2} \eta^{ce} \eta_{bf} (\delta_d^a \delta_e^f - \delta_e^a \delta_d^f) \end{aligned}$$

arises frequently.

2.2 Conventions for invariant matrices

Possible actions can be constructed using curvatures naturally arising in a theory, together with any invariant tensors consistent with the gauging. The biconformal gauging of the conformal group has a surprising number of invariant objects. These invariant structures arise from internal symmetries of the conformal group are induced into generic biconformal spaces [7].

The conformally invariant Killing form, restricted to the base manifold:

$$K_{AB} = \begin{pmatrix} 0 & \delta_a{}^b \\ \delta^a{}_b & 0 \end{pmatrix} \quad (14)$$

The symplectic form, underlying dilatations,

$$\Omega_{AB} = \begin{pmatrix} 0 & \delta_a{}^b \\ -\delta^a{}_b & 0 \end{pmatrix} \quad (15)$$

Interestingly, this form manifests Born reciprocity [1, 2]. The complex structure, arising from the symmetry between translations of the origin and translations of the point at infinity (i.e., special conformal transformations),

$$J^A{}_B := \begin{pmatrix} 0 & -\eta^{ab} \\ \eta_{ab} & 0 \end{pmatrix} \quad (16)$$

The symmetric Kähler form, arising from the compatibility, $g(u, v) = \Omega(u, Jv)$, of the symplectic and complex structures,

$$\begin{aligned} g_{AB} &= \Omega_{AC} J^C{}_B \\ &= \begin{pmatrix} 0 & \delta_a{}^c \\ -\delta^a{}_c & 0 \end{pmatrix} \begin{pmatrix} 0 & -\eta^{cb} \\ \eta_{cb} & 0 \end{pmatrix} \\ &= \begin{pmatrix} \eta_{ab} & 0 \\ 0 & \eta^{ab} \end{pmatrix} \end{aligned} \quad (17)$$

These three Kähler structures satisfy

$$g(u, v) = \Omega(u, Jv) \quad (18)$$

Notice that the symmetric Kähler form is not invariant under the conformal structure.

2.3 Kähler or Killing?

As noted in the introduction, vacuum biconformal gravity depends only on the variation of the gauge fields and does not require introduction of a metric while Yang-Mills actions make use of the Hodge dual and a metric is required. Given the presence of two natural symmetric forms in biconformal spaces, i.e., the scale invariant Killing form, Eq.(14), and the Kähler metric, Eq.(17), we must make a choice of how to specify the orthonormality relation of the solder and co-solder forms. With $\tilde{\mathbf{e}}^A = (\mathbf{e}^a, \mathbf{f}_b)$, and using an overbar to

denote the inverse, we may specify orthonormality by either

$$\langle \tilde{\mathbf{e}}^A, \tilde{\mathbf{e}}^B \rangle \equiv \bar{K}^{AB} \quad (19)$$

or

$$\langle \tilde{\mathbf{e}}^A, \tilde{\mathbf{e}}^B \rangle \equiv \bar{g}^{AB} \quad (20)$$

but clearly not both. Once we choose either Eq.(19) or Eq.(20), the corresponding symmetric form becomes the metric, with its variation following from the variation of the solder and co-solder forms.

This choice between the symmetric Killing and Kähler forms is not arbitrary. We therefore studied both cases, ultimately showing that the correct matter couplings arise only if we use the *Killing form*, K_{AB} , to determine orthonormality of the basis forms. By inverting the components of the basis forms, we then have the metric in arbitrary coordinates (indicated by *coordinate* indices M, N , distinct from *orthonormal* indices A, B)

$$K_{MN} = \tilde{e}_M^A \tilde{e}_N^B K_{AB} \quad (21)$$

This is our final choice of metric, and is the only choice which yields the expected Yang-Mills source for general relativity.

The gravitational field equations follow by variation of the connection forms, including the solder and co-solder forms $\tilde{e}_M^A$. The variation of the metric then follows from Eq.(21). Until the variation is complete, we need the general form of the inverse metric,

$$\bar{K}^{AB} = \begin{pmatrix} \bar{K}^{ab} & \bar{K}^a{}_b \\ \bar{K}_a{}^b & \bar{K}_{ab} \end{pmatrix}$$

Once the variations are expressed in terms of $\delta \bar{K}^{AB}$ or $\delta \bar{g}^{AB}$ any remaining components may be returned to the orthonormal form, Eq.(14) or Eq.(17).

Let the solder and co-solder variations be given by

$$\delta \mathbf{e}^a = A^a{}_c \mathbf{e}^c + B^{ac} \mathbf{f}_c \quad (22)$$

$$\delta \mathbf{f}_c = C_{cd} \mathbf{e}^d + D_c{}^d \mathbf{f}_d \quad (23)$$

Then for variation of the components of the Killing metric we expand Eq.(19)

$$\begin{aligned}
\delta \bar{K}^{ab} &= \delta \langle \mathbf{e}^a, \mathbf{e}^b \rangle \\
&= \langle A^a{}_c \mathbf{e}^c + B^{ac} \mathbf{f}_c, \mathbf{e}^b \rangle + \langle \mathbf{e}^a, A^b{}_c \mathbf{e}^c + B^{bc} \mathbf{f}_c \rangle \\
&= A^a{}_c \bar{K}^{cb} + B^{ac} \bar{K}_c{}^b + A^b{}_c \bar{K}^{ac} + B^{bc} \bar{K}^a{}_c
\end{aligned}$$

Since the coefficients $A^a{}_c, B^{ac}, C_{cd}, D_c{}^d$ now represent the variation, we may return the remaining components of $\bar{K}^{AB}$ to the null orthonormal form of Eq.(14),

$$\delta \bar{K}^{ab} = B^{ab} + B^{ba}$$

Computing the remaining components in the same way, we arrive at the full set,

$$\begin{aligned}
\delta \bar{K}^{ab} &= B^{ab} + B^{ba} \\
\delta \bar{K}^a{}_b &= A^a{}_b + D_b{}^a \\
\delta \bar{K}_a{}^b &= D_a{}^b + A^b{}_a \\
\delta \bar{K}_{ab} &= C_{ab} + C_{ba}
\end{aligned} \tag{24}$$

We also considered the analogous calculation if we were to choose the Kähler inner product, Eq.(20). This variation gives

$$\begin{aligned}
\delta \bar{g}^{ab} &= A^a{}_c \eta^{cb} + A^b{}_c \eta^{ac} \\
\delta \bar{g}^a{}_b &= B^{ac} \eta_{bc} + C_{bc} \eta^{ac} \\
\delta \bar{g}_a{}^b &= C_{ac} \eta^{cb} + B^{bc} \eta_{ac} \\
\delta \bar{g}_{ab} &= D_a{}^c \eta_{cb} + D_b{}^c \eta_{ac}
\end{aligned} \tag{25}$$

As we indicate when we perform the gravitational variation, Eq.(63) in Section 6 below, it is the B^{ab} part of the variation which ultimately reduces to the Einstein equation. From Eqs.(24) and (25) above we see that the different possible choices for the metric lead to completely different sources for gravity. Using the Killing form to define orthonormality of the basis, it is the coefficient of the $\delta \bar{K}^{ab}$ variation that provides the gravitational source, while the opposite choice of the Kähler form couples the coefficients of the cross-terms

$\delta\bar{g}^a{}_b$ and $\delta\bar{g}_a{}^b$ to gravity. Only one of these can give the usual Yang-Mills energy tensor. To determine which, we checked each of the two inner products, for each proposed action functional below until it became clear that *we must use the Killing form*.

Rather than presenting these distinct variations here, we continue with the Killing metric and its variation, as given by Eqs.(14), (19), and (24). Details of the failure of the Kähler choice are given in Appendix A.

3 Constructing the Yang-Mills action

In spacetime, the action for a Yang-Mills field may be written as

$$S_{\text{YM}} = -\frac{\kappa}{2} \int \text{tr} \mathcal{F} \wedge^* \mathcal{F} \quad (26)$$

and it is natural to consider the same form in biconformal space. However, as we show in this Section, this usual form leads to nonstandard coupling to gravity. We show in the next Section that a twisted action is required to give the usual coupling.

Expanded into independent components in the $\tilde{\mathbf{e}}^A$ basis,

$$\mathcal{F}^i = \frac{1}{2} \mathcal{F}^i{}_{ab} \mathbf{e}^a \wedge \mathbf{e}^b + \mathcal{F}^{ia}{}_b \mathbf{f}_a \wedge \mathbf{e}^b + \frac{1}{2} \mathcal{F}^{iab} \mathbf{f}_a \wedge \mathbf{f}_b$$

where i is an index of the internal Lie algebra. This index can be suppressed without loss of generality in the action and gravitational field equations. Also, for the Yang-Mills field it proves more transparent to give the three coefficients distinct names. Finally, since we must insure antisymmetry of the twisted field, it is most transparent to use the uniform weight basis, $\mathbf{e}^A = (\mathbf{e}^a, \eta^{bc} \mathbf{f}_c) = (\mathbf{e}^a, \mathbf{f}^b)$. We therefore write

$$\mathcal{F} = \frac{1}{2} \mathcal{F}_{AB} \mathbf{e}^A \wedge \mathbf{e}^B = \frac{1}{2} F_{ab} \mathbf{e}^a \wedge \mathbf{e}^b + \mathcal{G}_{ab} \mathbf{f}^a \wedge \mathbf{e}^b + \frac{1}{2} \mathcal{H}_{ab} \mathbf{f}^a \wedge \mathbf{f}^b \quad (27)$$

In discussions of the Yang-Mills field equations (as opposed to the gravitational equations), the internal index becomes important and will be shown where necessary.

Note that F_{ab} and $\mathcal{H}_{ab}$ are antisymmetric. The cross-term may be written as

$$\begin{aligned} G^a{}_b \mathbf{f}_a \wedge \mathbf{e}^b &= \frac{1}{2} (G^a{}_b) \mathbf{f}_a \wedge \mathbf{e}^b + \frac{1}{2} (-G^b{}_a) \mathbf{e}^a \wedge \mathbf{f}_b \\ &= \mathcal{G}_{ab} \mathbf{f}^a \wedge \mathbf{e}^b \end{aligned}$$

where $\mathcal{G}_{ab}$ may be asymmetric. As a matrix, $\mathcal{F}_{AB}$ takes the form given in Eq.(9).

3.1 The Hodge dual and standard Yang-Mills action

With these factors in mind, we find the Hodge dual of the Yang-Mills field. In general terms the Hodge dual of a 2-form is given by

$$\begin{aligned} {}^* \mathcal{F} &= {}^* \left(\frac{1}{2} \mathcal{F}_{AB} \tilde{\mathbf{e}}^A \wedge \tilde{\mathbf{e}}^B \right) \\ &= \frac{1}{(2n-2)!} \frac{1}{2} \mathcal{F}_{AB} \bar{K}^{AC} \bar{K}^{BD} \varepsilon_{CDE\dots F} \tilde{\mathbf{e}}^E \wedge \dots \wedge \tilde{\mathbf{e}}^F \end{aligned}$$

However, we need to separate the distinct quadrants of each inverse metric. Expanding each upper case index $A, B, \dots$, as a raised, weight +1 index and a lowered weight -1 index in turn, then collecting like terms leads to

$$\begin{aligned} {}^* \mathcal{F} &= \frac{1}{n! (n-2)!} \left(\frac{1}{2} F_{ab} \bar{K}^{am} \bar{K}^{bn} + G^a{}_b \bar{K}_a{}^m \bar{K}^{bn} + \frac{1}{2} H^{ab} \bar{K}_a{}^m \bar{K}_b{}^n \right) \varepsilon^{c\dots d}{}_{mne\dots f} \mathbf{f}_{c\dots d} \mathbf{e}^{e\dots f} \\ &+ \frac{(-1)^{n-1}}{(n-1)! (n-1)!} \left(\frac{1}{2} F_{ab} \bar{K}^a{}_m \bar{K}^{bn} + G^a{}_b \bar{K}_{am} \bar{K}^{bn} + \frac{1}{2} H^{ab} \bar{K}_{am} \bar{K}_b{}^n \right) \varepsilon^{mc\dots d}{}_{ne\dots f} \mathbf{f}_{c\dots d} \mathbf{e}^{e\dots f} \\ &+ \frac{(-1)^n}{(n-1)! (n-1)!} \left(\frac{1}{2} F_{ab} \bar{K}^{am} \bar{K}^b{}_n + G^a{}_b \bar{K}_a{}^m \bar{K}^b{}_n + \frac{1}{2} H^{ab} \bar{K}_a{}^m \bar{K}_{bn} \right) \varepsilon^{nc\dots d}{}_{me\dots f} \mathbf{f}_{c\dots d} \mathbf{e}^{e\dots f} \\ &+ \frac{1}{n! (n-2)!} \left(\frac{1}{2} F_{ab} \bar{K}^a{}_m \bar{K}^b{}_n + G^a{}_b \bar{K}_{am} \bar{K}^b{}_n + \frac{1}{2} H^{ab} \bar{K}_{am} \bar{K}_{bn} \right) \varepsilon^{mnc\dots d}{}_{e\dots f} \mathbf{f}_{c\dots d} \mathbf{e}^{e\dots f} \quad (28) \end{aligned}$$

where one readily sees the advantage of omitting the wedge between forms, $\mathbf{e}^e \wedge \dots \wedge \mathbf{e}^f \longleftrightarrow \mathbf{e}^{e\dots f}$.

We form the usual Yang-Mills Lagrangian density as the wedge product, $\mathcal{F} \wedge {}^* \mathcal{F}$, eliminating the basis forms in favor of the volume form Φ . After a bit of algebra, we find

$$\begin{aligned}
\mathcal{F} \wedge {}^* \mathcal{F} = & \left(\frac{1}{2} F_{mn} \bar{K}^{am} \bar{K}^{bn} + G^m{}_n \bar{K}^a{}_m \bar{K}^{bn} + \frac{1}{2} H^{mn} \bar{K}^a{}_m \bar{K}^b{}_n \right) F_{ab} \Phi \\
& + (F_{mn} \bar{K}^a{}_m \bar{K}^{bn} + G^m{}_n (\bar{K}_{am} \bar{K}^{bn} - \bar{K}^a{}_n \bar{K}^b{}_m) + H^{mn} \bar{K}_{am} \bar{K}^b{}_n) G^a{}_b \Phi \\
& + \left(\frac{1}{2} F_{mn} \bar{K}^a{}_m \bar{K}^b{}_n + G^m{}_n \bar{K}_{am} \bar{K}^b{}_n + \frac{1}{2} H^{mn} \bar{K}_{am} \bar{K}_{bn} \right) H^{ab} \Phi
\end{aligned} \tag{29}$$

and the matter action is given by Eq.(26). The full action is the combination of Eq.(8) and Eq.(26),

$$S = S_G + S_{YM}$$

Substituting the null orthonormal form of the Killing metric, Eq.(14) into Eq.(29) the lagrange density reduces to

$$\mathcal{F}^* \mathcal{F} = (H^{ab} F_{ab} - G^a{}_b G^a{}_b) \Phi \tag{30}$$

We find that this Hodge dual form of the action is identical to the form of the action given in [20], despite the claim in [20] that the action is independent of the metric. The presence of the metric is concealed in [20] because the Killing metric in this basis is comprised of Kronecker deltas.

Although varying the Yang-Mills potentials in Eq.(29), or equivalently in Eq.(30), yields the usual Yang-Mills field equation, we show below that varying the metric gives a nonstandard coupling to gravity. In the next Section we define a twisted action that gives the usual coupling to gravity, while still leading to the correct Yang-Mills field equations.

3.2 The failure of the $\mathcal{F} \wedge {}^* \mathcal{F}$ action

Variation of the usual Lagrangian Eq.(29) gives

$$\begin{aligned}
\delta_g \int \mathcal{F} \wedge {}^* \mathcal{F} = & \int (F_{mn} F_{ab} \delta \bar{K}^{am} \eta^{bn}) \Phi \\
& + (G^m{}_n F_{ab} \eta^{bn} \delta \bar{K}^a{}_m + F_{mn} G^a{}_b \eta^{bn} \delta \bar{K}^a{}_m) \Phi \\
& + (G^m{}_n G^a{}_b \eta^{bn} \delta \bar{K}_{am} + G^m{}_n G^a{}_b \eta_{am} \delta \bar{K}^{bn}) \Phi \\
& + (H^{mn} G^a{}_b \eta_{am} \delta \bar{K}^b{}_n + G^m{}_n H^{ab} \eta_{am} \delta \bar{K}_b{}^n) \Phi \\
& + (H^{mn} H^{ab} \eta_{bn} \delta \bar{K}_{am}) \Phi
\end{aligned}$$

where after variation we returned the remaining inverse metric components to the orthonormal form. The metric variations are now given by Eq.(24). Substituting, the variation yields the gravitational field equations,

$$\begin{aligned} \alpha \Omega^n{}_b{}^b{}_m - \alpha \Omega^a{}_b{}^b{}_a \delta_m^n + \beta \Omega^n{}_m - \beta \Omega^a{}_a \delta_m^n + \Lambda \delta_m^n &= -\kappa (H^{bn} F_{bm} - G^b{}_m G^n{}_b) \\ &\quad - \kappa \left(\frac{1}{2} \delta_m^n (F_{bc} H^{bc} - G^b{}_c G^c{}_b) \right) \end{aligned} \quad (31)$$

$$\begin{aligned} \alpha \Omega^a{}_n{}^m{}_a - \alpha \Omega^a{}_b{}^b{}_a \delta_n^m + \beta \Omega^m{}_n - \beta \Omega^a{}_a \delta_n^m + \Lambda \delta_n^m &= -\kappa (H^{bm} F_{bn} - G^b{}_n G^m{}_b) \\ &\quad - \kappa \left(\frac{1}{2} \delta_n^m (F_{bc} H^{bc} - G^b{}_c G^c{}_b) \right) \end{aligned} \quad (32)$$

$$\alpha \Omega^a{}_{nam} + \beta \Omega_{nm} = \kappa (F_{am} G^a{}_n + F_{an} G^a{}_m) \quad (33)$$

$$\alpha \Omega^n{}_b{}^{bm} + \beta \Omega^{nm} = -\kappa (H^{nb} G^m{}_b + H^{mb} G^n{}_b) \quad (34)$$

The problem with this coupling to gravity becomes evident when the gravitational equations require vanishing momentum curvatures on the left side of Eq.(34). With this, Eq.(34) implies

$$H^{nb} G^m{}_b + H^{mb} G^n{}_b = 0$$

The failure of this result is not immediate, but the reduction of Eqs.(31) and (32) gives a second constraint on H^{ab} and $G^a{}_b$. The two constraints lead, at the most general, to vanishing H^{ab} and symmetric $G^a{}_b$, leaving the source for the Einstein equation, Eq.(33) linear in F_{ab} and therefore incompatible with the usual energy source for general relativity.

The problem cannot be altered by a different choice of the inner product (see Appendix A), but must lie in the use of the usual action. In the context of other double field theories, a twist allows dimensional reduction to preserve gauging of supersymmetries [16, 18, 19]. Here we find that including a twist insures consistency under dimensional reduction not only for supersymmetry, but also gives the correct coupling of Yang-Mills sources to gravity.

We now turn to the definition of the twisted Yang-Mills action, and its metric variation. Then, in Section 5, we vary the Yang-Mills potentials. The remainder of our presentation details the reduction of the full set of field equations to reproduce scale covariant general relativity with Yang-Mills sources in the usual n -dimensional form.

4 Metric variation of the twisted Yang-Mills Action

Instead of the usual spacetime action, Eq.(26), we consider a biconformal Yang-Mills theory with an action functional of the form

$$S_{\text{TYM}} = -\frac{\kappa}{2} \int \text{tr} \bar{\mathcal{F}} \wedge^* \mathcal{F} \quad (35)$$

where $*$ is the usual Hodge dual, $\mathcal{F}$ is a curvature 2-form, $\bar{\mathcal{F}}$ is a twisted conjugate curvature, and the trace is over the $SU(N)$ generators. The twist matrix is formed using both the Killing metric and the Kähler form, $K^A{}_B \equiv \bar{K}^{AC} g_{CB}$. While the twist matrix is similar to that used to preserve supersymmetry in other double field theories, [16, 18, 19], we define the twisted Yang-Mills field by

$$\bar{\mathcal{F}}_{AB} = \frac{1}{2} (K_A{}^C \mathcal{F}_{CB} + \mathcal{F}_{AC} K^C{}_B)$$

We find that this form is necessary to preserve the antisymmetry of the field while giving the required interchange of source fields. Variation of the inverse Killing form $\bar{K}^{AB}$ then gives the source for the gravitational field equations.

4.1 Details of the twist

The twist is accomplished using $K^A{}_B \equiv \bar{K}^{AC} g_{CB}$ where $K^A{}_B = K_B{}^A$, since both g_{AB} and K_{AB} are symmetric. In the null-orthonormal form and the $\mathbf{e}^A$ basis, this matrix is simply

$$K^A{}_B = K_B{}^A = \begin{pmatrix} 0 & \delta_b^a \\ \delta_b^a & 0 \end{pmatrix}$$

and the required form of the field is

$$\begin{aligned} \bar{\mathcal{F}}_{AB} &= \frac{1}{2} (K_A{}^C \mathcal{F}_{CB} + \mathcal{F}_{AC} K^C{}_B) \\ &= \begin{pmatrix} \mathcal{G}_{[ab]} & \frac{1}{2} (F_{ab} + \mathcal{H}_{ab}) \\ \frac{1}{2} (F_{ab} + \mathcal{H}_{ab}) & \mathcal{G}_{[ab]} \end{pmatrix} \end{aligned} \quad (36)$$

where $\mathcal{F}_{AB}$ is given by Eq.(9). Note that this transformation maintains the antisymmetry while interchanging the diagonal and anti-diagonal elements.

However, the reduced form of the twisted field in the orthonormal frame given in Eq.(36) is insufficient.

Until we complete the metric variation we must use the generic form of the metric in computing the twist matrix,

$$\begin{aligned}
K^A{}_B &= \begin{pmatrix} \bar{K}^{ac} & \bar{K}^a{}_e \eta^{ec} \\ \eta^{ae} \bar{K}_e{}^c & \eta^{ae} \eta^{ef} \bar{K}_{ef} \end{pmatrix} \begin{pmatrix} \eta_{cb} & 0 \\ 0 & \eta_{bc} \end{pmatrix} \\
&= \begin{pmatrix} \bar{K}^{ac} \eta_{cb} & \bar{K}^a{}_b \\ \bar{K}^a{}_b & \eta^{ae} \bar{K}_{eb} \end{pmatrix} \\
K_A{}^B &= \begin{pmatrix} \eta_{ad} \bar{K}^{db} & \bar{K}_a{}^b \\ \bar{K}_a{}^b & \bar{K}_{ae} \eta^{eb} \end{pmatrix}
\end{aligned}$$

with symmetry given by $\bar{K}^a{}_b = \eta^{ae} \bar{K}_e{}^c \eta_{cb}$ (see Appendix B for details of the symmetry). Then $\bar{\mathcal{F}}_{AB}$ becomes

$$\begin{aligned}
\bar{\mathcal{F}}_{AB} &= \frac{1}{2} (K_A{}^C \mathcal{F}_{CB} + \mathcal{F}_{AC} K^C{}_B) \\
&= \frac{1}{2} \begin{pmatrix} \eta_{ad} \bar{K}^{dc} F_{cb} + \bar{K}_a{}^c \mathcal{G}_{cb} & -\eta_{ad} \bar{K}^{dc} \mathcal{G}_{bc} + \bar{K}_a{}^c \mathcal{H}_{cb} \\ \eta_{ad} \bar{K}^d{}_e \eta^{ec} F_{cb} + \bar{K}_{ae} \eta^{ec} \mathcal{G}_{cb} & -\eta_{ad} \bar{K}^d{}_e \eta^{ec} \mathcal{G}_{bc} + \bar{K}_{ae} \eta^{ec} \mathcal{H}_{cb} \end{pmatrix} \\
&\quad + \frac{1}{2} \begin{pmatrix} F_{ac} \bar{K}^{ce} \eta_{eb} - \mathcal{G}_{ca} \eta^{cd} \bar{K}_d{}^e \eta_{eb} & F_{ac} \bar{K}^c{}_b - \mathcal{G}_{ca} \eta^{ce} \bar{K}_{eb} \\ \mathcal{G}_{ac} \bar{K}^{ce} \eta_{eb} + \mathcal{H}_{ac} \eta^{cd} \bar{K}_d{}^e \eta_{eb} & \mathcal{G}_{ac} \bar{K}^c{}_b + \mathcal{H}_{ac} \eta^{ce} \bar{K}_{eb} \end{pmatrix} \quad (37)
\end{aligned}$$

and we check that 37 agrees with Eq.(36) when we restore the null orthonormal frame for the metric. The twisted field is simpler when written as a 2-form,

$$\begin{aligned}
\bar{\mathcal{F}} &= \frac{1}{2} \bar{\mathcal{F}}_{AB} \mathbf{e}^A \wedge \mathbf{e}^B \\
&= \frac{1}{2} (F_{ac} \bar{K}^{ce} \eta_{eb} + \bar{K}_a{}^c \mathcal{G}_{cb}) \mathbf{e}^a \wedge \mathbf{e}^b \\
&\quad + \frac{1}{2} (F_{ac} \bar{K}^c{}_b \eta^{bf} + \bar{K}_a{}^c \mathcal{H}_{cb} \eta^{bf} - \mathcal{G}_{ca} \eta^{ce} \bar{K}_{eb} \eta^{bf} - \mathcal{G}_{bc} \bar{K}^{ce} \eta_{ea} \eta^{bf}) \mathbf{e}^a \wedge \mathbf{f}_f \\
&\quad + \frac{1}{2} (\mathcal{H}_{ac} \eta^{ce} \bar{K}_{eb} + \mathcal{G}_{ac} \bar{K}^c{}_b) \eta^{af} \mathbf{f}_f \wedge \eta^{bg} \mathbf{f}_g \quad (38)
\end{aligned}$$

We may interchange $\bar{K}_a{}^b$ and $\bar{K}^b{}_a$ when convenient since these have identical variations and both restrict to δ_b^a in the null orthonormal basis.

4.2 The action

The dual field is given by Eq.(28). To avoid duplication of indices when we wedge the dual field together with the twisted field, we rename the indices in the twisted field

$$\begin{aligned}\bar{\mathcal{F}} = & \frac{1}{2} (F_{rq} \bar{K}^{qt} \eta_{ts} + \bar{K}_r{}^q \mathcal{G}_{qs}) \mathbf{e}^r \wedge \mathbf{e}^s \\ & + \frac{1}{2} (F_{rc} \bar{K}^c{}_s + \bar{K}_r{}^c \mathcal{H}_{cs}) \eta^{sw} \mathbf{e}^r \wedge \mathbf{f}_w \\ & + \frac{1}{2} (-\mathcal{G}_{sq} \bar{K}^{qt} \eta_{tr} - \mathcal{G}_{qr} \eta^{qt} \bar{K}_{ts}) \eta^{sw} \mathbf{e}^r \wedge \mathbf{f}_w \\ & + \frac{1}{2} (\mathcal{G}_{rq} \bar{K}^q{}_s + \mathcal{H}_{rq} \eta^{qt} \bar{K}_{ts}) \eta^{rw} \eta^{sx} \mathbf{f}_w \wedge \mathbf{f}_x\end{aligned}$$

Each term of the wedge product is proportional to the volume form, using $\mathbf{f}_{c\dots d} \wedge \mathbf{e}^{e\dots f} = \bar{e}_{c\dots d}{}^{e\dots f} \Phi$. Then, replacing the double Levi-Civita tensors with Kronecker deltas according to Eqs.(13), we fully distribute the lengthy expression. We summarize the essential features here, but details including the full wedge product $\bar{\mathcal{F}} \wedge {}^* \mathcal{F}$ and its reduction to $\bar{\mathcal{F}} \wedge {}^* \mathcal{F}|_{contributing}$ below are given in Appendix C.

Since we are interested only in the variation, and will return the metric to the null orthonormal form of Eq.(14) after variation, certain terms clearly do not contribute to the field equations. For example, in the product

$$(-\mathcal{G}_{sq} \bar{K}^{qt} \eta_{tr} - \mathcal{G}_{qr} \eta^{qt} \bar{K}_{ts}) \eta^{sw} \left(\frac{1}{2} F_{ab} \bar{K}^a{}_m \bar{K}^{bn} + \frac{1}{2} \mathcal{H}_{gh} \eta^{ga} \eta^{hb} \bar{K}_{am} \bar{K}_b{}^n \right)$$

none of the four terms after distribution will contribute to the field equation because the variation of any one factor of metric components always leaves an unvaried $\bar{K}_{ts}$ or $\bar{K}^{bn}$,

$$\delta (\bar{K}^a{}_m \bar{K}_{ts} \bar{K}^{bn}) = \delta \bar{K}^a{}_m \bar{K}_{ts} \bar{K}^{bn} + \bar{K}^a{}_m \delta \bar{K}_{ts} \bar{K}^{bn} + \bar{K}^a{}_m \bar{K}_{ts} \delta \bar{K}^{bn}$$

Dropping such terms, we are only required to vary terms linear in $\bar{K}_{ab}$ or $\bar{K}^{ab}$, or cubic in the off diagonal components $\bar{K}^a{}_b$. Collecting these and using the symmetries of the fields, finally yields,

$$\begin{aligned}\bar{\mathcal{F}} \wedge {}^* \mathcal{F}|_{contributing} = & \left(\frac{1}{2} F_{bc} F_{da} \eta^{ac} + F_{bc} \mathcal{H}^{ac} \eta_{ad} + \frac{1}{2} (\mathcal{G}_{ad} - 2\mathcal{G}_{da}) \eta_{bc} \mathcal{G}^{ac} \right) \bar{K}^{bd} \Phi \\ & + (\mathcal{G}_{ba} H^{dc} + F_{ab} \mathcal{G}^{cd}) \bar{K}^b{}_e \bar{K}^e{}_d \bar{K}^a{}_c \Phi \\ & + \left(\frac{1}{2} \mathcal{H}_{ac} \mathcal{H}^{ab} \eta^{cd} + F_{ac} \mathcal{H}^{ab} \eta^{cd} + \frac{1}{2} (\mathcal{G}^{ba} - 2\mathcal{G}^{ab}) \eta^{cd} \mathcal{G}_{ca} \right) \bar{K}_{bd} \Phi\end{aligned}\quad (39)$$

We may now vary the metric.

To carry out the variation of the Yang-Mills potentials we may write the action in the null orthonormal frame. This form is still contained in the expression above, given by the purely off-diagonal terms, cubic in $\bar{K}^a{}_b$. This simpler form of the action follows immediately as

$$S_{TYM} = \kappa \int tr (\bar{\mathcal{F}} \wedge {}^* \mathcal{F}) = \kappa \int tr (\mathcal{G}^{ab} (\mathcal{H}_{ab} + F_{ab})) \Phi \quad (40)$$

The variation of the potentials is carried out in Section 5.

4.3 Metric variation

Using the variation of the Killing metric given in Eq.(24) and the variation of the volume form given by

$$\begin{aligned} \delta \Phi &= -\frac{1}{2} K_{AB} \delta \bar{K}^{AB} \Phi \\ &= -\frac{1}{2} [K_{ab} (B^{ab} + B^{ba}) + K^a{}_b (D_a{}^b + A^b{}_a) + K_b{}^a (A^b{}_a + D_a{}^b) + K^{ab} (C_{ab} + C_{ba})] \Phi \\ &= -\delta^a{}_b (A^b{}_a + D_a{}^b) \Phi \end{aligned}$$

the variation of Eq.(39) yields

$$\begin{aligned} \delta (\bar{\mathcal{F}} \wedge {}^* \mathcal{F}) &= \left(\frac{1}{2} F_{bc} F_{da} \eta^{ac} + F_{bc} \mathcal{H}^{ac} \eta_{ad} + \frac{1}{2} (\mathcal{G}_{ad} - 2\mathcal{G}_{da}) \eta_{bc} \mathcal{G}^{ac} \right) 2B^{(bd)} \Phi \\ &\quad + ((2\mathcal{G}^{ad} - \mathcal{G}^{da}) F_{ab} - (2\mathcal{G}^{da} - \mathcal{G}^{ad}) \mathcal{H}_{ab} - (F_{ac} + \mathcal{H}_{ac}) \mathcal{G}^{ac} \delta^d{}_b) (A^b{}_d + D_d{}^b) \Phi \\ &\quad + \left(\frac{1}{2} \mathcal{H}_{ac} \mathcal{H}^{ab} \eta^{cd} + F_{ac} \mathcal{H}^{ab} \eta^{cd} + \frac{1}{2} (\mathcal{G}^{ba} - 2\mathcal{G}^{ab}) \eta^{cd} \mathcal{G}_{ca} \right) 2C_{(bd)} \Phi \end{aligned} \quad (41)$$

This variation couples to the $(\mathbf{e}^a, \mathbf{f}_a)$ variation of the gravity action, Eq.(8). The resulting field equations and their reduction to the gravity theory are given in Section (6) below. Before considering reduction of the field equations, we turn to variation of the Yang-Mills potentials to find the Yang-Mills equations.

5 Yang-Mills potentials and field equations

We also need to vary the potential to find the Yang-Mills field equations. We start with the action of Eq.(40) in the null orthonormal basis,

$$S_{TYM} = \kappa \int \text{tr} \bar{\mathcal{F}}^i \wedge * \mathcal{F}^i = \kappa \int \mathcal{G}^{iab} (\mathcal{H}_{i\ ab} + F_{i\ ab}) \Phi \quad (42)$$

In this Section, we make the internal symmetry explicit, varying S_{YM} with respect to the $SU(N)$ potentials. It is most convenient to work in the $\bar{\mathbf{e}}^A = (\mathbf{e}^a, \mathbf{f}_b)$ basis. Internal indices are labeled with letters i, j, k , while frame indices are chosen from the beginning of the alphabet, $a, b, c, \dots$

5.1 The field components and potentials

The $SU(N)$ field is given in the $\bar{\mathbf{e}}^A$ basis by Eq.(43),

$$\mathcal{F}^i = \frac{1}{2} F^i_{\ ab} \mathbf{e}^a \wedge \mathbf{e}^b + \mathcal{G}^i_{\ cb} \eta^{ac} \mathbf{f}_a \wedge \mathbf{e}^b + \frac{1}{2} \mathcal{H}^i_{\ ab} \eta^{ac} \eta^{bd} \mathbf{f}_c \wedge \mathbf{f}_d \quad (43)$$

where as a matrix,

$$\mathcal{F}^i_{\ AB} = \begin{pmatrix} F^i_{\ ab} & G^i_{\ a\ b} \\ G^i_{\ a\ b} & \mathcal{H}^i_{\ ab} \end{pmatrix}$$

with $G^i_{\ a\ b} = -G^i_{\ b\ a}$.

The field is given in terms of its $U(1)$ or $SU(N)$ potential by the Cartan equation,

$$\mathcal{F}^i = d\mathcal{A}^i - \frac{1}{2} c^i_{\ jk} \mathcal{A}^j \wedge \mathcal{A}^k$$

where the potentials are biconformal 1-forms,

$$\mathcal{A}^i = A^i_{\ a} \mathbf{e}^a + B^i_{\ a} \mathbf{f}_a$$

In terms of $A^i_{\ a}, B^i_{\ a}$, the field becomes

$$\begin{aligned}
\mathcal{F}^i &= \mathbf{D}A^i{}_a \wedge \mathbf{e}^a - \frac{1}{2} \mathcal{A}^i{}_k \wedge A^k{}_b \mathbf{e}^b \\
&\quad + \mathbf{D}B^{i\,a} \wedge \mathbf{f}_a + B^{i\,a} \mathbf{S}_a - \frac{1}{2} \mathcal{A}^i{}_k \wedge B^k{}_b \mathbf{f}_b
\end{aligned} \tag{44}$$

where $\mathcal{A}^i{}_k$ is the $SU(N)$ connection in the adjoint representation,

$$\mathcal{A}^i{}_k \equiv c^i{}_{jk} \mathcal{A}^j$$

and the Weyl-covariant derivatives of the potentials are given by

$$\begin{aligned}
\mathbf{D}A^i{}_a &= \mathbf{d}A^i{}_a - \omega^a{}_c A^i{}_a + A^i{}_a \omega \\
\mathbf{D}B^{i\,a} &= \mathbf{d}B^{i\,a} - B^{i\,a} \omega^c{}_a - B^{i\,a} \omega
\end{aligned}$$

Note that the covariant derivative of η^{ab} does not necessarily vanish, $\mathbf{D}\eta^{ab} = \mathbf{d}\eta^{ab} - 2\omega\eta^{ab}$ where $\mathbf{d}\eta^{ab}$ takes into account the conformal equivalence class, $\eta_{ab} \in \{e^{2\varphi}\eta_{ab}^0 \mid \varphi = \varphi(x, y)\}$.

Now we separate Eq.(44) into its independent $\mathbf{e}^a \wedge \mathbf{e}^b$, $\mathbf{f}_a \wedge \mathbf{e}^b$ and $\mathbf{f}_a \wedge \mathbf{f}_b$ parts. We begin by expanding the exterior derivatives in the orthonormal basis

$$\begin{aligned}
\mathbf{d}A^i{}_a &= A^i{}_{a,b} \mathbf{e}^b + A^i{}_a{}^{;b} \mathbf{f}_b \\
\mathbf{d}B^{i\,a} &= B^{i\,a}{}_{;b} \mathbf{e}^b + B^{i\,a,b} \mathbf{f}_b
\end{aligned}$$

and similarly the covariant exterior derivatives

$$\begin{aligned}
\mathbf{D}A^i{}_a &= A^i{}_{a;b} \mathbf{e}^b + A^i{}_a{}^{;b} \mathbf{f}_b \\
\mathbf{D}B^{i\,a} &= B^{i\,a}{}_{;b} \mathbf{e}^b + B^{i\,a,b} \mathbf{f}_b
\end{aligned}$$

We also expand the $SU(N)$ connection, $\mathcal{A}^j G_j$, in the adjoint representation:

$$\begin{aligned}
\mathcal{A}^i{}_k &= c^i{}_{jk} A^j{}_a \mathbf{e}^a + c^i{}_{jk} B^{j\,a} \mathbf{f}_a \\
&= \alpha^i{}_k + \beta^i{}_k
\end{aligned}$$

Finally, writing the general form of the co-torsion

$$\mathbf{S}_a = \frac{1}{2}S_{abc}\mathbf{e}^b \wedge \mathbf{e}^c + S_a{}^b{}_c \mathbf{f}_b \wedge \mathbf{e}^c + \frac{1}{2}S_a{}^{bc}\mathbf{f}_b \wedge \mathbf{f}_c$$

we arrive the component fields in terms of the potentials:

$$F^i{}_{ab} = A^i{}_{b;a} - A^i{}_{a;b} - c^i{}_{jk}A^j{}_a A^k{}_b + B^i{}^c S_{cab} \quad (45)$$

$$G^i{}^a{}_b = A^i{}^a{}_{;b} - B^i{}^a{}_{;b} - c^i{}_{jk}B^j{}^a A^k{}_b + B^i{}^c S_c{}^a{}_b \quad (46)$$

$$H^i{}^{ab} = B^i{}^{b;a} - B^i{}^{a;b} - c^i{}_{jk}B^j{}^a B^k{}^b + B^i{}^c S_c{}^{ab} \quad (47)$$

These expressions are covariant with respect to both $SU(N)$ and Weyl group transformations.

5.2 The field equations for the potentials

Using Eqs.(45)-(47) in the action, Eq.(42), the variation of $A^i{}_a$ gives

$$\begin{aligned} \delta_A S_{TYM} &= \kappa \int (\delta_A \mathcal{G}^{iab} (\mathcal{H}_{iab} + F_{iab})) \Phi \\ &\quad + \kappa \int \mathcal{G}^{iab} (\delta_A \mathcal{H}_{iab} + \delta_A F_{iab}) \Phi \\ &= \kappa \int (\delta A^k{}_b \eta^{bc} (-(\mathcal{H}_{kac} + F_{kac})^{;a} - c^i{}_{jk} B^j{}^a (\mathcal{H}_{iac} + F_{iac}))) \Phi \\ &\quad + \kappa \int \delta A^k{}_b \left(-(\mathcal{G}_k{}^{ab} - \mathcal{G}_k{}^{ba})_{;a} - c^i{}_{kj} A^j{}_a (\mathcal{G}_i{}^{ba} - \mathcal{G}_i{}^{ab}) \right) \Phi \end{aligned}$$

Therefore the field equation becomes

$$\begin{aligned} 0 &= \eta^{bc} (\mathcal{H}_{kac} + F_{kac})^{;a} + \eta^{bc} (\mathcal{H}_{iac} + F_{iac}) \beta^i{}_k{}^a \\ &\quad + (\mathcal{G}_k{}^{ab} - \mathcal{G}_k{}^{ba})_{;a} + (\mathcal{G}_i{}^{ab} - \mathcal{G}_i{}^{ba}) \alpha^i{}_{ka} \end{aligned}$$

For the $B^i{}^a$ variation, recalling that i, j , are internal indices and a, b, c, e orthonormal indices, we find

$$\begin{aligned} 0 &= \eta^{bc} \left((\mathcal{H}_{jac} + F_{jac})_{;b} + \alpha^i{}_{jb} (\mathcal{H}_{iac} + F_{iac}) + S_a{}^e{}_b (\mathcal{H}_{jec} + F_{jec}) \right) \\ &\quad + (\mathcal{G}_{jab} - \mathcal{G}_{jba})^{;b} + (\mathcal{G}_{iab} - \mathcal{G}_{iba}) \beta^i{}_j{}^b + \frac{1}{2} (\mathcal{G}_{jbc} - \mathcal{G}_{jcb}) S_a{}^{bc} \\ &\quad + \frac{1}{2} (\mathcal{G}_j{}^{bc} - \mathcal{G}_j{}^{cb}) S_{abc} \end{aligned}$$

Notice that there is no dynamical equation for the symmetric part of $\mathcal{G}^k_{cd}$. Moreover, the action depends only on the antisymmetric part of $\mathcal{G}^k_{cd}$. Therefore, from here on we assume that like F^i_{ab} and H^{iab} , $\mathcal{G}^k_{cd}$ is antisymmetric, $\mathcal{G}^k_{ab} \equiv \mathcal{G}^k_{[ab]}$. Also notice that there is no separate Yang-Mills field equation for F_{ab} and H_{ab} . Both field equations contain only their sum, $\mathcal{H}^k_{ab} + F^k_{ab}$, although F_{ab} and $\mathcal{H}_{ab}$ enter the gravitational equations separately. We therefore define a new field,

$$K^k_{ab} \equiv \frac{1}{2} (\mathcal{H}^k_{ab} + F^k_{ab})$$

In terms of these, the field equations become

$$0 = \eta^{bc} K_{kac}{}^{;a} + \eta^{bc} K_{iac} \beta^i_{k}{}^a + \mathcal{G}_k{}^{ab}{}_{;a} + \mathcal{G}_i{}^{ab} \alpha^i_{ka} \quad (48)$$

$$0 = \eta^{bc} (K_{jac;b} + K_{iac} \alpha^i_{jb} + K_{jec} S_a{}^e{}_b) \\ + \mathcal{G}_{jab}{}^{;b} + \mathcal{G}_{iab} \beta^i_{j}{}^b + \frac{1}{2} \mathcal{G}_{jbc} S_a{}^{bc} + \frac{1}{2} \mathcal{G}_j{}^{bc} S_{abc} \quad (49)$$

These have the expected form of a divergence of the field strength.

The field equations of an $SU(N)$ gauge theory on a $2n$ -dimensional space given by Eqs.(48) and (49) together with the gravitational sources of Eqs.(41) complete the first stage of our investigation. These expressions give a satisfactory formulation of biconformal Yang-Mills theory.

The second stage of this study is to understand how the $SU(N)$ gravitational sources together with their field equations affect the gravitational and Yang-Mills solutions. Specifically, we want to know if the reduction to the co-tangent bundle of n -dimensional spacetime still occurs, and if so, whether the usual Yang-Mills field equations and gravitational sources result.

This is indeed what happens. Once we have reduced the gravitational field equations to more simply describe the underlying geometry, we will return to Eqs.(48) and (49). Our final result is to show that the reduction of the underlying geometry to the co-tangent bundle of general relativity simultaneously forces the reduction of the source equations to the usual Yang-Mills sources for general relativity.

In the next Section we carry out the gravitational reduction. Since the pure gravitational field case is presented in detail elsewhere [6], we are able to simply state some of the conclusions, focussing on those features which are different in the presence of matter. Subsequently, in Sec.(7), we give particular attention to the resulting twofold reduction of the $SU(N)$ fields: (1) from $(N^2 - 1) \times \frac{2n(2n-1)}{2}$ field components to $(N^2 - 1) \times \frac{n(n-1)}{2}$ components, and (2) the restriction of the number of effective independent variables,

$$F(x, y) \rightarrow F(x).$$

6 Reduction of the gravitational field equations

Although the reduction of the gravitational field equations has been presented in detail elsewhere [6] and we only highlight the features which differ in the presence of matter, the presentation is still somewhat lengthy. To aid in following the discussion, we begin with a short description of the steps before providing details. The basic steps are the following:

- Present the structure equations and Bianchi identities and their immediate consequences for the form of the torsion-free biconformal curvatures (Subsections 6.1 - 6.2.1). By manipulating the field equations, we arrive at a reduced form for the components of the curvature tensors (Subsections 6.2.1 - 6.2.4). (These calculations involve the sources in essential ways that require more detailed presentation here.)
- The Cartan structure equations show that the solder form $\mathbf{e}^a$ is in involution. The Frobenius theorem therefore allows us to set $\mathbf{e}^a = e_\mu^a \mathbf{d}x^\mu$ and study the restricted solution on the $x^\mu = \text{constant}$ submanifolds spanned by n additional coordinates y_μ . Because of the reduced form of the curvature components, this restricted solution completely determines the $\mathbf{e}^a = 0$ pieces of the connection forms. We then extend the connection forms back to the full biconformal space. At this point, the connection forms have far fewer than their original number of components (Summarized in Subsection 6.2.5).
- Substitute the reduced connection forms into the structure equations and impose the field equations. This is done one structure equation at a time, each time reducing the degrees of freedom of the full system. Ultimately, all connection and curvature components are determined by the solder form, $\mathbf{e}^a$, and the components of the solder form itself only depend on half the original coordinates, $e_\mu^a(x, y) \Rightarrow e_\mu^a(x)$ (Subsections 6.2.6, 6.2.7).

6.1 Field equations for the twisted action

The full action is

$$\begin{aligned}
S &= S_G + S_{YM} \\
&= \lambda \int e_{ac \dots d}{}^{be \dots f} (\alpha \Omega^a{}_b + \beta \delta^a{}_b \Omega + \gamma \mathbf{e}^a \wedge \mathbf{f}_b) \wedge \mathbf{e}^c \wedge \dots \wedge \mathbf{e}^d \wedge \mathbf{f}_e \wedge \dots \wedge \mathbf{f}_f \\
&\quad - \frac{\kappa}{2} \int \text{tr} \overline{\mathcal{F}} \wedge^* \mathcal{F}
\end{aligned} \tag{50}$$

We choose the combinatoric factor λ so the final coupling is κ .

6.1.1 Curvatures, Bianchi identities, and gravity variation

The gravitational field equations are given by varying S_G in Eq.(50) with respect to all connection 1-forms, then combining with the Yang-Mills sources found in Eq.(41). The curvature components are given in terms of the connection by the Cartan structure equations,

$$\mathbf{d}\omega^a{}_b = \omega^c{}_b \omega^a{}_c + 2\Delta_{db}^{ac} \mathbf{f}_c \mathbf{e}^d + \Omega^a{}_b \quad (51)$$

$$\mathbf{d}\mathbf{e}^a = \mathbf{e}^c \omega^a{}_c + \omega \mathbf{e}^a + \mathbf{T}^a \quad (52)$$

$$\mathbf{d}\mathbf{f}_a = \omega^c{}_a \mathbf{f}_c + \mathbf{f}_a \omega + \mathbf{S}_a \quad (53)$$

$$\mathbf{d}\omega = \mathbf{e}^c \mathbf{f}_c + \Omega \quad (54)$$

We also have the integrability conditions—generalized Bianchi identities—which follow from the Poincarè lemma, $\mathbf{d}^2 \equiv 0$. Exterior differentiation of the Cartan equations, Eqs.(51)-(54) yields conditions which must be satisfied in order for solutions to exist. These take the form

$$\mathbf{D}\Omega^a{}_b = 2\Delta_{db}^{ac} \mathbf{f}_c \mathbf{T}^d - 2\Delta_{db}^{ac} \mathbf{S}_c \mathbf{e}^d \quad (55)$$

$$\mathbf{D}\mathbf{T}^a = \mathbf{e}^c \Omega^a{}_c - \Omega \mathbf{e}^a \quad (56)$$

$$\mathbf{D}\mathbf{S}_a = -\Omega^c{}_a \mathbf{f}_c + \mathbf{f}_a \Omega \quad (57)$$

$$\mathbf{D}\Omega = -\mathbf{T}^c \mathbf{f}_c + \mathbf{e}^c \mathbf{S}_c \quad (58)$$

Conversely, the converse to the Poincarè lemma tells us that in a star-shaped region the integrability conditions Eqs.(55)-(58) imply the original Cartan equations, up to boundary terms.

For example, consider a Newtonian force written as minus the gradient of a potential, $\mathbf{F} = -\mathbf{d}V$. The Poincarè lemma shows that this equation has no solution unless

$$0 \equiv \mathbf{d}^2 V = -\mathbf{d}\mathbf{F}$$

so the force must be curl-free. Conversely, we may solve the this “Bianchi identity” first. In a star-shaped region any curl-free force must be a gradient, $\mathbf{F} = \mathbf{d}U$. This reproduces the potential up to a “boundary”

term satisfying

$$\mathbf{d}(U + V) = 0$$

$$U + V = \text{constant}$$

This in turn provides the usual additive constant to the potential. These principles hold for any set of equations where we may apply the Poincarè lemma and its converse, allowing us use both the original equations and their integrability conditions throughout the development of a solution. In the end, one set of equations cannot be satisfied without simultaneously satisfying the other up to boundary terms.

The variation of the gravitational part of the action, Eq.(50), is discussed in detail in [6], and involves only variation of the connection forms, so we simply state the result. The variation of the spin connection $\omega^a{}_b$ and Weyl vector ω give

$$T^{ae}{}_e - T^{ea}{}_e - S_e{}^{ae} = 0 \quad (59)$$

$$T^a{}_{ca} + S_c{}^a{}_a - S_a{}^a{}_c = 0 \quad (60)$$

$$\alpha \Delta_{sb}^{ar} (T^{mb}{}_a - \delta_a^m T^{eb}{}_e - \delta_a^m S_c{}^{bc}) = 0 \quad (61)$$

$$\alpha \Delta_{sb}^{ar} (\delta_c^b T^d{}_{ad} + S_c{}^b{}_a - \delta_c^b S_d{}^d{}_a) = 0 \quad (62)$$

and these acquire no sources since the Yang-Mills action is independent of the these connection forms, as noted in [5]. The variation of the solder and co-solder forms lead to

$$\begin{aligned} & [\alpha (\Omega^n{}_b{}^b{}_m - \Omega^a{}_b{}^b{}_a \delta_m^n) + \beta (\Omega^n{}_m - \Omega^a{}_a \delta_m^n) + \Lambda \delta_m^n] A^m{}_n \\ & [\alpha (\Omega^a{}_n{}^m{}_a - \Omega^a{}_b{}^b{}_a \delta_n^m) + \beta (\Omega^m{}_n - \Omega^a{}_a \delta_n^m) + \Lambda \delta_n^m] D_n{}^m \\ & - [\alpha \Omega^a{}_{nam} + \beta \Omega_{nm}] B^{mn} \\ & [\alpha \Omega^n{}_b{}^bm + \beta \Omega^{nm}] C_{mn} \end{aligned} \quad (63)$$

with the arbitrary variations $A^m{}_n, B^{mn}, C_{mn}$ and $D_n{}^m$ defined in Eqs.(22) and (23). In [6] the expressions above are equated to zero, but they now acquire sources.

6.1.2 Combined equations

Equating corresponding parts of Eqs.(41) and Eqs.(63) and symmetrizing appropriately,

$$[\alpha (\Omega^a{}_c{}^c{}_b - \Omega^c{}_d{}^d{}_c \delta_b^a) + \beta (\Omega^a{}_b - \Omega^c{}_c \delta_b^a) + \Lambda \delta_b^a] = \kappa W^a{}_b \quad (64)$$

$$[\alpha (\Omega^c{}_b{}^a{}_c - \Omega^c{}_d{}^d{}_c \delta_b^a) + \beta (\Omega^a{}_b - \Omega^c{}_c \delta_b^a) + \Lambda \delta_b^a] = \kappa W^a{}_b \quad (65)$$

$$-[\alpha \Omega^c{}_{bca} + \beta \Omega_{ba}] = \kappa T_{ab} \quad (66)$$

$$[\alpha \Omega^b{}_c{}^{ca} + \beta \Omega^{ba}] = \kappa S^{ab} \quad (67)$$

where, recalling the antisymmetry of $\mathcal{G}^{ab}$,

$$T_{ab} \equiv \text{tr} (F_{ac} F_{bd} \eta^{cd} + H_{ac} F_{bd} \eta^{cd} + H_{bc} F_{ad} \eta^{cd} + 3\eta^{cd} \mathcal{G}_{ac} \mathcal{G}_{bd}) \quad (68)$$

$$S^{ab} \equiv \eta^{bd} \eta^{ac} \text{tr} (\eta^{ef} H_{ec} H_{fd} + \eta^{ef} H_{ec} F_{fd} + \eta^{ef} H_{ed} F_{fc} + 3\eta^{fe} \mathcal{G}_{ec} \mathcal{G}_{fd}) \quad (69)$$

$$W^a{}_b \equiv \text{tr} (3\mathcal{G}^{ca} F_{cb} + 3\mathcal{G}^{ca} \mathcal{H}_{cb} - (F_{cd} + \mathcal{H}_{cd}) \mathcal{G}^{cd} \delta^a{}_b) \quad (70)$$

In addition, we have the field equations for the $U(1)$ field,

$$0 = \eta^{bc} K_{kac}{}^{;a} + \eta^{bc} K_{iac} \beta^i{}_k{}^a + \mathcal{G}_k{}^{ab}{}_{;a} + \mathcal{G}_i{}^{ab} \alpha^i{}_{ka} \quad (71)$$

$$0 = \eta^{bc} (K_{jac;b} + K_{iac} \alpha^i{}_{jb} + K_{jec} S_a{}^e{}_b) \\ + \mathcal{G}_{jab}{}^{;b} + \mathcal{G}_{iab} \beta^i{}_j{}^b + \frac{1}{2} \mathcal{G}_{jbc} S_a{}^{bc} + \frac{1}{2} \mathcal{G}_j{}^{bc} S_{abc} \quad (72)$$

Our gravitational solution now follows many of the steps presented in detail in [6].

6.2 Solving the field equations for the twisted action

For the remainder of the gravitational solution, the particular forms of T_{ab} , S^{ab} and $W^a{}_b$ make little difference; indeed, the solution of this Section holds for the metric variation of any sources at all. While the form of these source tensors varies with the fields and the matter action, the positions in which they occur in the field equations and their symmetries follow knowing only the variation of $\bar{K}^{AB}$. For the present, this is all we need.

We first turn to the consequences of vanishing torsion, $\mathbf{T}^a = 0$.

6.2.1 Vanishing torsion

Similarly to general relativity, with vanishing torsion the torsion Bianchi identity, Eq.(56), simplifies to an algebraic relation,

$$0 = \mathbf{e}^c \Omega^a{}_c - \Omega \mathbf{e}^a$$

which must hold independently of any sources. The algebraic condition $\mathbf{e}^c \Omega^a{}_c = \Omega \mathbf{e}^a$ expands to three independent component equations,

$$\Omega^a{}_{[bcd]} = \delta^a_{[b} \Omega_{cd]} \quad (73)$$

$$\Omega^a{}_b{}^c{}_d - \Omega^a{}_d{}^c{}_b = \delta^a_b \Omega^c{}_d - \delta^a_d \Omega^c{}_b \quad (74)$$

$$\Omega^a{}_b{}^{cd} = \delta^a_b \Omega^{cd} \quad (75)$$

Since $\eta_{ea} \Omega^a{}_b{}^{cd} = -\eta_{ba} \Omega^a{}_e{}^{cd}$, the ab trace of Eq.(75) leaves $\Omega^{cd} = 0$. Therefore each term vanishes separately,

$$\Omega^a{}_b{}^{cd} = 0 \quad (76)$$

$$\Omega^{cd} = 0 \quad (77)$$

The ad contraction of Eq.(74) gives

$$\Omega^a{}_b{}^c{}_a = -(n-1) \Omega^c{}_b \quad (78)$$

Combining Eqs.(76) and (77) with Eq.(67) we immediately find that the gravitational fields force a constraint on the source fields. This is our *first source constraint*:

$$S^{ab} = 0 \quad (79)$$

We next look at the field equations for the curvature.

6.2.2 Curvature equations

We now combine the vanishing torsion simplifications with the curvature and dilatation field equations, Eqs.(64) and (65). The reduction of these equations begins by noting that the difference between Eq.(64)

and Eq.(65) immediately gives equality of the traces,

$$\Omega^c{}_{ba}{}^c = \Omega^a{}_{cb}{}^b \quad (80)$$

Next, formally lowering an index in Eq.(74)

$$\eta_{ea}\Omega^a{}_{bc}{}^c{}_d - \eta_{ea}\Omega^a{}_{dc}{}^c{}_b = \eta_{eb}\Omega^c{}_{da}{}^a{}_c - \eta_{ed}\Omega^c{}_{ba}{}^a{}_c$$

we cycle ebd , then add the first two and subtract the third. Using the antisymmetry of the curvature on the first two indices, $\eta_{ea}\Omega^a{}_{dc}{}^c{}_b = -\eta_{da}\Omega^a{}_{ec}{}^c{}_b$ we find

$$\Omega^a{}_{ec}{}^c{}_d = -2\Delta_{de}^{ab}\Omega^c{}_{ba}{}^a{}_c \quad (81)$$

Substituting Eq.(81) into the trace symmetry, Eq.(80) to the two contractions of Eq.(81) constrains the cross-dilatation,

$$-\delta_d^a\Omega^c{}_{ca}{}^a{}_c + \eta^{ae}\eta_{cd}\Omega^c{}_{ea}{}^a{}_c = -(n-1)\Omega^a{}_{dc}{}^c{}_c$$

Contracting with η_{ba} , we see that the antisymmetric part vanishes,

$$(n-2)(\eta_{bc}\Omega^c{}_{da}{}^a{}_c - \eta_{cd}\Omega^c{}_{ba}{}^a{}_c) = 0$$

in dimensions greater than 2, while an explicit check confirms the vanishing antisymmetry in 2-dimensions as well. Therefore, the symmetric part, $\eta_{bc}\Omega^c{}_{da}{}^a{}_c + \eta_{dc}\Omega^c{}_{ba}{}^a{}_c = \frac{2}{n}\eta_{bd}\Omega^c{}_{ca}{}^a{}_c$, becomes a solution for the full cross-dilatation in terms of its trace,

$$\Omega^c{}_{da}{}^a{}_c = \frac{1}{n}\delta_b^a\Omega^c{}_{ca}{}^a{}_c \quad (82)$$

This, in turn, combines with Eq.(81) to give the cross-curvature in terms of the trace of the dilatation,

$$\Omega^a{}_{bc}{}^c{}_d = -\frac{2}{n}\Delta_{db}^{ac}\Omega^e{}_{ea}{}^a{}_c \quad (83)$$

We have one remaining cross-curvature field equation, Eq.(64), which couples the cross-dilatation trace, $\Omega^a{}_{ac}{}^c{}_c$, to the Yang-Mills source fields. Using Eqs.(82) and (83) to replace the cross-curvature and the cross-

dilatation in Eq.(64), and simplifying,

$$\frac{1}{n} (n-1) [((n-1)\alpha - \beta)\Omega^c{}_c + n\Lambda] \delta_b^a = W^a{}_b$$

so that $W^a{}_b = f\delta_b^a$ for some function f . The constant Λ is given by $\Lambda = (n-1)\alpha - \beta + n^2\gamma$. Contracting then substituting back, the gravitational field equations force a *second source constraint*:

$$W^a{}_b = \frac{1}{n} W^c{}_c \delta_b^a \quad (84)$$

where

$$W^c{}_c = (n-1) [((n-1)\alpha - \beta)\Omega^c{}_c + n\Lambda]$$

The traced source tensor on the right, $W^c{}_c$, therefore drives the entire cross-curvature and cross-dilatation. It is striking that the only source dependence for these components is the Yang-Mills Lagrangian density,

$$W^a{}_a = 3\mathcal{G}^{i\,ac}(\mathcal{H}_{i\,ac} + F_{i\,ac}) = 3\mathcal{L}$$

6.2.3 Spacetime terms

Finally, we combine the remaining field equation, Eq.(66),

$$\alpha\Omega^c{}_{bca} + \beta\Omega_{ba} = -\kappa T_{ab}$$

and the corresponding part of the vanishing torsion Bianchi, Eq.(73), which expanded becomes

$$\Omega^a{}_{bcd} + \Omega^a{}_{cdb} + \Omega^a{}_{dbc} = \delta_b^a \Omega_{cd} + \delta_c^a \Omega_{db} + \delta_d^a \Omega_{bc}$$

The ac trace reduces this to

$$\Omega^c{}_{bcd} - \Omega^c{}_{dcb} = -(n-2)\Omega_{bd}$$

Combining this with the antisymmetric part of the field equation, $\alpha(\Omega^a{}_{nam} - \Omega^a{}_{man}) = -2\beta\Omega_{nm}$ shows that

$$((n-2)\alpha - 2\beta)\Omega_{ab} = 0$$

so that generically (i.e., unless $(n-2)\alpha = 2\beta$), the spacetime dilatation vanishes. Note that this is true for any symmetric source tensor, so spacetime dilatation is never driven by ordinary matter. As a result,

$$\Omega^c{}_{acb} = \Omega^c{}_{(a|c|b)} = -\frac{\kappa}{\alpha}T_{ab} \quad (85)$$

$$\Omega_{nm} = 0 \quad (86)$$

6.2.4 Dilatation

Having reduced the dilatational curvature to a single function,

$$\Omega = \chi \mathbf{e}^a \wedge \mathbf{f}_a \quad (87)$$

where $\chi \equiv -\frac{1}{n}\Omega^a{}_{a}$, we can now use the dilatational integrability condition, Eq.(58), to press further. Substituting Eq.(87) into the Bianchi identity, Eq.(58),

$$\begin{aligned} 0 &= \mathbf{d}\Omega - \mathbf{e}^b \wedge \mathbf{S}_b \\ &= \mathbf{d}\chi \wedge \mathbf{e}^a \wedge \mathbf{f}_a - (1 + \chi) \mathbf{e}^a \wedge \mathbf{S}_a \end{aligned}$$

where we have used $\mathbf{d}(\mathbf{e}^a \wedge \mathbf{f}_a) = \mathbf{D}(\mathbf{e}^a \wedge \mathbf{f}_a) = \mathcal{T}^a \wedge \mathbf{f}_a - \mathbf{e}^a \wedge \mathbf{S}_a$. Setting $\mathbf{d}\chi = \chi_c \mathbf{e}^c + \chi^c \mathbf{f}_c$, expanding the co-torsion into components, and combining like forms yields three independent equations,

$$(1 + \chi) S_{[acd]} = 0 \quad (88)$$

$$(1 + \chi) (S_c{}^a{}_d - S_d{}^a{}_c) = \chi_d \delta_c^a - \chi_c \delta_d^a \quad (89)$$

$$(1 + \chi) S_a{}^{cd} = \chi^d \delta_a^c - \chi^c \delta_a^d \quad (90)$$

We may now use the co-torsion field equations to gain insight into χ .

With vanishing torsion, the co-torsion field equations Eqs.(59)-(62) reduce to

$$S_e^{ae} = 0 \quad (91)$$

$$S_c^a{}_a - S_a^a{}_c = 0 \quad (92)$$

$$\alpha \Delta_{sb}^{ar} (S_c^b{}_a - \delta_c^b S_d^d{}_a) = 0 \quad (93)$$

Using the field equation Eq.(92) to replace the co-torsion terms in the trace of the Bianchi identity, Eq.(89), gives

$$\begin{aligned} (1 + \chi) (S_c^a{}_a - S_a^a{}_c) &= -(n-1) \chi_c \\ \chi_c &= 0 \end{aligned}$$

Then, combining Eq.(91) with the *ad* trace of Eq.(90),

$$\begin{aligned} (1 + \chi) S_a^{ca} &= -(n-1) \chi^c \\ \chi^c &= 0 \end{aligned}$$

and therefore,

$$\mathbf{d}\chi = 0 \quad (94)$$

The dilatation therefore takes the form

$$\Omega = \chi \mathbf{e}^a \wedge \mathbf{f}_a$$

with χ *constant*. The remainder of the development of the solution continues as in the homogeneous case but with a different constant value,

$$\chi = \frac{1}{(n-1)\alpha - \beta} \left(\frac{1}{n-1} \Lambda - \frac{\kappa}{n(n-1)} W^c{}_c \right)$$

for the magnitude of the dilatation cross-term.

Importantly, the constancy of χ implies the constancy of $W^c{}_c$, and via the second source constraint, Eq.(84), the (mostly zero) constancy of all of $W^a{}_b$.

6.2.5 The Frobenius theorem and the final reduction

With vanishing torsion, 52 shows that the solder form becomes involute, and we may write

$$\mathbf{e}^a = e_\alpha{}^a \mathbf{d}x^\alpha$$

where x^α comprise n of the $2n$ coordinates. Holding x^α constant, $\mathbf{e}^a = 0$, and the residual field equations describe a submanifold. Here the discussion exactly parallels that of [6]:

1. Solve for the connection on the $\mathbf{e}^a = 0$ submanifold. Here, because the curvature and dilatation vanish (see Eqs.(76) and (77)), we may gauge the restricted components of the spin connection and Weyl vector to zero. A careful coordinate choice puts the submanifold basis in the form $\mathbf{h}_a = e_a{}^\mu \mathbf{d}y_\mu$.
2. Now let x^α vary, extending the solution back to the full biconformal space. This allows all connection forms to acquire an additional $\mathbf{d}x^\alpha$ or $\mathbf{e}^a$ term,

$$\begin{aligned} \omega^a{}_b &= \omega^a{}_{bc} \mathbf{e}^c \\ \mathbf{e}^a &= e_\alpha{}^a \mathbf{d}x^\alpha \\ \mathbf{f}_a &= e_a{}^\mu \mathbf{d}y_\mu + c_{ab} \mathbf{e}^b \\ \omega &= W_a \mathbf{e}^a \end{aligned} \tag{95}$$

3. Note that Eq.(52) is now purely quadratic in $\mathbf{e}^a$, and therefore requires the coefficients to depend only on the x -coordinates, $e_\alpha{}^a = e_\alpha{}^a(x)$. Solving for the connection separates it into a compatible piece and a Weyl vector piece,

$$\omega^a{}_b = \alpha^a{}_b - 2\Delta_{db}^{ac} W_c \mathbf{e}^d$$

where $\mathbf{d}\mathbf{e}^a = \mathbf{e}^b \alpha^a{}_b$.

4. Substitute these reduced forms of $\mathbf{e}^a$, $\mathbf{f}_b$ into the dilatation, Eq.(54) and solve for the Weyl vector. This yields

$$\omega = -(1 + \chi) y_a \mathbf{e}^a$$

where $y_a = e_a{}^\mu y_\mu$.

These steps give the final expressions for the connection forms, except for the form of $c_{ab} = c_{ba}$ in the expansion of the co-solder form $\mathbf{f}_a$.

We note that the submanifolds found by setting either $\mathbf{e}^a = 0$ by holding x^μ constant, or $\mathbf{h}_a = 0$ by holding y_μ constant are Lagrangian submanifolds.

6.2.6 The curvature

The final steps in the gravitational reduction are to substitute the partial solution for the connection forms Eq.(95) into Eqs.(51) and (53) to impose the final field equations.

To express the remaining undetermined component of the curvature, $\Omega^a{}_{bcd}$, we define the Schouten tensor

$$\mathcal{R}_a = \frac{1}{n-2} \left(R_{ab} - \frac{1}{2(n-1)} \eta_{ab} R \right) \mathbf{e}^b$$

where $R_{ab} = (n-2) \mathcal{R}_{ab} + \eta_{ab} \mathcal{R}$ is the Ricci tensor. The generalization of the Schouten tensor to an integrable Weyl geometry is then (see [24])

$$\mathcal{R}_a = \mathcal{R}_a + \mathbf{D}_{(\alpha, x)} W_a + W_a \omega - \frac{1}{2} \eta_{ab} W^2 \mathbf{e}^b$$

In terms of the Schouten tensor, the decomposition of the Riemann curvature 2-form into the Weyl curvature 2-form and trace parts is

$$\mathbf{R}^a{}_b = \mathbf{C}^a{}_b - 2\Delta_{db}^{ac} \mathbf{R}_c \wedge \mathbf{e}^d$$

Because of the manifest involution of $\mathbf{h}_a = e_a{}^\mu(x) \mathbf{d}y_\mu$, the subspace spanned by the solder form, $\mathbf{e}^a$, is a submanifold. Because $\Omega_{ab} = 0$ the submanifold geometry is always an *integrable* Weyl geometry, so the Weyl vector may be removed from the spacetime submanifold by a gauge transformation. The spacetime submanifold is simply a Riemannian geometry with local scale invariance.

Now, introducing the reduced form of the connection into Eq.(51) and imposing the corresponding field equation, Eq.(66) shows that

$$\frac{1}{1+\chi} \mathcal{R}_{ab} + c_{ab} = -\frac{\kappa}{\alpha} T_{ab} \quad (96)$$

with the full spacetime component of the biconformal curvature given by the Weyl curvature, $\Omega^a{}_{bcd} = C^a{}_{bcd}$.

6.2.7 The co-torsion

A similar introduction of the reduced connection into Eq.(53) for the co-torsion shows that the momentum and cross-terms vanish, while (following the somewhat intricate calculation of [6]) the remaining component

is given by

$$\mathbf{S}_a = \mathbf{d}_{(x)}\mathbf{c}_a + \mathbf{c}_b \wedge \boldsymbol{\omega}^b_a + \boldsymbol{\omega} \wedge \mathbf{c}_a \quad (97)$$

where $\mathbf{c}_a$, in turn, is determined by Eq.(96).

Expanding $\mathbf{c}_a$ fully to separate the Weyl vector parts,

$$\begin{aligned} \mathbf{c}_a &= -\frac{1}{1+\chi} \left(\mathcal{R}_a + \mathbf{D}_{(\alpha,x)} W_a + W_a \boldsymbol{\omega} - \frac{1}{2} \eta_{ab} W^2 \mathbf{e}^b \right) - \frac{\kappa}{\alpha} \mathbf{T}_a \\ &= \mathbf{b}_a - \frac{1}{1+\chi} \left(\mathbf{D}_{(\alpha,x)} W_a + W_a \boldsymbol{\omega} - \frac{1}{2} \eta_{ab} W^2 \mathbf{e}^b \right) \end{aligned}$$

where $\mathbf{T}_a = T_{ab} \mathbf{e}^b$ is the remaining source field and $\mathbf{b}_a \equiv -\frac{1}{1+\chi} \mathcal{R}_a - \frac{\kappa}{\alpha} \mathbf{T}_a$. Then substituting into Eq.(97), after multiple cancellations the co-torsion becomes

$$\begin{aligned} \mathbf{S}_a &= \frac{1}{1+\chi} W_b \mathbf{R}^b_a - \mathbf{D}_{(\alpha,x)} \left(\frac{1}{1+\chi} \mathcal{R}_a + \frac{\kappa}{\alpha} \mathbf{T}_a \right) \\ &\quad + 2\Delta_{ca}^{db} W_d \left(\frac{1}{1+\chi} \mathcal{R}_b + \frac{\kappa}{\alpha} \mathbf{T}_b \right) \wedge \mathbf{e}^c \end{aligned} \quad (98)$$

with the cross term and momentum term of the co-torsion vanishing.

This result is quite similar to an integrability condition. It is shown in [24] that the condition for the existence of a conformal gauge in which the Einstein equation, $G_{ab} = \kappa T_{ab}$, holds is

$$0 = \phi_{,b} \mathbf{R}^b_a - \mathbf{D}_{(\alpha,x)} (\mathcal{R}_a - \kappa \mathcal{T}_a) + 2\Delta_{ca}^{db} \phi_{,d} (\mathcal{R}_b - \kappa \mathcal{T}_b) \wedge \mathbf{e}^c \quad (99)$$

where

$$\mathcal{T}_a = \frac{1}{n-2} \left(T_{ab} - \frac{1}{n-1} T \eta_{ab} \right)$$

When $\mathcal{T}_a = 0$, Eq.(99) reduces to the well-known condition, $\mathbf{D}_{(\alpha,x)} \mathcal{R}_a - \phi_{,b} \mathbf{C}^b_a = 0$, for the existence of a Ricci flat conformal gauge.

There are two differences between Eq.(98) and Eq.(99). First, the co-torsion on the left hand side of Eq.(98) obstructs the integrability condition, Eq.(99), and we cannot set $\mathbf{S}_a = 0$ because the Triviality Theorem shows that when both torsion and co-torsion vanish, biconformal space must be trivial. The second difference is that the Weyl vector on the right is not integrable on the *full* biconformal space.

These issues have a common solution. The part of structure equation for the co-solder form involving $\mathbf{h}_a$

is

$$\mathbf{d}\mathbf{h}_a = \omega^c{}_a \wedge \mathbf{h}_c + \mathbf{h}_a \wedge \omega \quad (100)$$

Therefore, as briefly noted above, $\mathbf{h}_a = e_a{}^\mu \mathbf{d}y_\mu$ is in involution. Holding $y_\mu = y_\mu^0$ constant shows that $\mathbf{e}^a$ spans a submanifold. On that submanifold, the Weyl vector becomes exact,

$$\omega = W_a \mathbf{e}^a = \mathbf{d}(y_\mu^0 x^\mu)$$

This means that on the $y_\mu = y_\mu^0$ spacetime submanifold, the right side takes the form of the integrability condition.

At the same time, we may use the form of $\mathbf{S}_a$ as the covariant derivative of $\mathbf{c}_a$, Eq.(97) with a suitable choice of c_{ab} . In [6] it is shown that c_{ab} is symmetric and divergence free. While the interpretation given in [6] of a phenomenological energy tensor is consistent, it is at odds with the more fundamental interpretation of sources given here. Instead, we identify c_{ab} as proportional to the Minkowski metric—the only invariant, symmetric tensor available. It is also divergence free with respect to the compatible connection, since $D_{(\alpha,x)}\eta_{ab} = 0$. However, as noted in the previous Section the fully biconformal-covariant derivative of η_{ab} does not necessarily vanish. Since by Eq.(97) the co-torsion is given by the full biconformal derivative of c_{ab} , the identification $c_{ab} = \Lambda_0 \eta_{ab}$, implies

$$\mathbf{S}_a = 2(1 + \chi) \Delta_{ca}^{bc} y_b \Lambda_0 \eta_{cd} \mathbf{e}^d \wedge \mathbf{e}^e$$

thereby avoiding the Triviality Theorem. This residual form of the co-torsion may now be combined into the right hand side of Eq.(98).

Combining these observations, on the $\mathbf{h}_a = 0$ spacetime submanifold with $c_{ab} = \Lambda_0 \eta_{ab}$, setting $\phi_{,a} = y_a^0$, and using $\mathbf{D}_{(\alpha,x)}(\Lambda_0 \eta_{ab}) = 0$, it follows that

$$\begin{aligned} 0 = & \frac{1}{1 + \chi} \phi_{,b} \mathbf{R}^b{}_a - \mathbf{D}_{(\alpha,x)} \left(\frac{1}{1 + \chi} \mathbf{R}_a + \frac{\kappa}{\alpha} \mathbf{T}_a + \Lambda_0 \eta_{ab} \mathbf{e}^b \right) \\ & + 2 \Delta_{ca}^{db} \phi_{,d} \left(\frac{1}{1 + \chi} \mathbf{R}_b + \frac{\kappa}{\alpha} \mathbf{T}_b + \Lambda_0 \eta_{be} \mathbf{e}^e \right) \wedge \mathbf{e}^c \end{aligned} \quad (101)$$

This is now the condition for the existence of a conformal transformation such that

$$\frac{1}{1+\chi}\mathbf{R}_b + \Lambda_0\eta_{be}\mathbf{e}^e = -\frac{\kappa}{\alpha}\mathbf{T}_b$$

Expressed in terms of the Einstein tensor,

$$G_{ab} + \Lambda_C\eta_{ab} = -(n-2)\frac{\kappa(1+\chi)}{\alpha}(T_{ab} - \eta_{ab}T) \quad (102)$$

where the net effect of c_{ab} is a cosmological constant, $\Lambda_C = -(n-1)(n-2)(1+\chi)\Lambda_0$.

If we make the conformal transformation that produces Eq.(102), the co-torsion equation (101) reduces to $\phi_{,b}\mathbf{R}^b_a = 0$. In generic spacetimes this requires $W_\mu = y_\mu^0 = 0$.

7 The source for gravity

7.1 The reduction of sources forced by coupling to gravity

The necessary source constraints from the gravitational couplings, Eqs.(79) and (84), may be written as

$$0 = \mathcal{H}^i_{ac}\mathcal{H}_{i\,bd}\eta^{cd} + (\mathcal{H}^i_{ac}F_{i\,bd} + \mathcal{H}^i_{bc}F_{i\,ad})\eta^{cd} + 3\mathcal{G}^i_{ac}\mathcal{G}_{i\,bd}\eta^{cd} \quad (103)$$

$$0 = \mathcal{G}^{i\,ca}(F^i_{cb} + \mathcal{H}^i_{cb}) - \frac{1}{n}\delta^a_b\mathcal{G}^{i\,cd}(F^i_{cd} + \mathcal{H}^i_{cd}) \quad (104)$$

where the full contraction of the second, $\mathcal{G}^{i\,cd}(F^i_{cd} + \mathcal{H}^i_{cd})$, is constant.

These conditions must continue to hold for small physical variations of the independent potentials, A^i_a and B^{ia} . We may imagine two nearby solutions differing only in one or both of the potentials and look at their difference. The change in F^i_{ab} as we change A^i_a is given by

$$\begin{aligned} \delta F^i_{ab} &= \delta A^i_{b;a} - \delta A^i_{a;b} - c^i_{jk}\delta A^j_a A^k_b - c^i_{jk}A^j_a \delta A^k_b \\ &= (\delta A^i_{b;a} - \alpha^i_{ka}\delta A^k_b) - (\delta A^i_{a;b} - \alpha^i_{jb}\delta A^j_a) \\ &= \mathcal{D}_a(\delta A^i_b) - \mathcal{D}_b(\delta A^i_a) \end{aligned}$$

where $\mathcal{D}_a$ is covariant with respect to local Lorentz, dilatational and $SU(N)$ transformations. Similarly we

find for $G^{i\ a}_{\ b}$ and $H^{i\ ab}$,

$$\begin{aligned}\delta_A G^{i\ a}_{\ b} &= \mathcal{D}^a (\delta A^i_{\ b}) \\ \delta_A H^{i\ ab} &= 0\end{aligned}$$

Of course, under changes of gauge, these fields are invariant.

The conditions (103) and (104) must continue to hold throughout such small changes. Substituting these variations into the first constraint,

$$0 = \delta F^i_{\ ed} (\mathcal{H}_{i\ ac} \delta_b^e + \mathcal{H}_{i\ bc} \delta_a^e) \eta^{cd} + 3 (\eta_{ae} \delta G^{i\ e}_{\ c} \mathcal{G}_{i\ bd} + \mathcal{G}_{i\ ac} \eta_{be} \delta G^{i\ e}_{\ d}) \eta^{cd} \quad (105)$$

The first term of Eq.(103) has dropped out because $H^{i\ ab}$ is independent of $A^i_{\ a}$.

Now we expand Eq.(105) in terms of the variation $\delta A^i_{\ a}$ and its derivatives,

$$\begin{aligned}0 &= \delta A^i_{\ d,e} (\mathcal{H}_{i\ ac} \delta_b^e \eta^{cd} + \mathcal{H}_{i\ bc} \delta_a^e \eta^{cd} - \mathcal{H}_{i\ ac} \delta_b^d \eta^{ce} - \mathcal{H}_{i\ bc} \delta_a^d \eta^{ce}) \\ &\quad + \delta A^i_{\ c\ \prime e} 3 (\eta_{ae} \mathcal{G}_{i\ bd} + \mathcal{G}_{i\ ad} \eta_{be}) \eta^{cd} \\ &\quad + \delta A^k_{\ f} \left(-\omega^f_{\ de} + W_e \delta_d^f + \alpha^i_{\ kd} \delta_e^f \right) (\mathcal{H}_{k\ ac} \delta_b^e \eta^{cd} + \mathcal{H}_{k\ bc} \delta_a^e \eta^{cd} - \mathcal{H}_{k\ ac} \delta_b^d \eta^{ce} - \mathcal{H}_{k\ bc} \delta_a^d \eta^{ce}) \\ &\quad + \delta A^k_{\ f} (3c^i_{\ kj} B^{j\ e}) (\eta_{ae} \mathcal{G}_{i\ bd} \eta^{fd} + \mathcal{G}_{i\ ad} \eta_{be} \eta^{fd})\end{aligned}$$

where we collect terms proportional to $\delta A^k_{\ f}$, $\delta A^k_{\ f,e}$ and $\delta A^k_{\ f\ \prime e}$ separately, noting that the gravitational solution reduces the y -covariant derivative to a partial, $\delta A^i_{\ b\ \prime a} = \delta A^i_{\ b\ \prime a}$.

While the field equations determine the second derivatives of the potentials, the potential itself and its first derivative are arbitrary initial conditions on any Cauchy surface. Therefore, the three variations $\delta A^k_{\ f}$, $\delta A^k_{\ f,e}$ and $\delta A^k_{\ f\ \prime e}$ are independent, and the coefficient of each must vanish separately:

$$0 = \mathcal{H}_{i\ ac} \delta_b^e \eta^{cd} + \mathcal{H}_{i\ bc} \delta_a^e \eta^{cd} - \mathcal{H}_{i\ ac} \delta_b^d \eta^{ce} - \mathcal{H}_{i\ bc} \delta_a^d \eta^{ce} \quad (106)$$

$$0 = 3 (\eta_{ae} \mathcal{G}_{i\ bd} + \mathcal{G}_{i\ ad} \eta_{be}) \eta^{cd} \quad (107)$$

$$\begin{aligned}0 &= \left(-\omega^f_{\ de} + W_e \delta_d^f + \alpha^i_{\ kd} \delta_e^f \right) (\mathcal{H}_{k\ ac} \delta_b^e \eta^{cd} + \mathcal{H}_{k\ bc} \delta_a^e \eta^{cd} - \mathcal{H}_{k\ ac} \delta_b^d \eta^{ce} - \mathcal{H}_{k\ bc} \delta_a^d \eta^{ce}) \\ &\quad + (3c^i_{\ kj} B^{j\ e}) (\eta_{ae} \mathcal{G}_{i\ bd} \eta^{fd} + \mathcal{G}_{i\ ad} \eta_{be} \eta^{fd})\end{aligned} \quad (108)$$

For the x -derivative part of the constraint, Eq.(106), we contract eb and lower the d index to show that $\mathcal{H}_{i\ ac}$

must vanish,

$$0 = n\mathcal{H}_{i\,ac}$$

Similarly, contracting ac in Eq.(107) expressing the independence of the y -derivative, shows that $\mathcal{G}_{i\,be}$ must also vanish.

$$0 = 3\mathcal{G}_{i\,be}$$

With these two conditions, the final equation Eq.(108) is identically satisfied.

These conditions satisfy both gravitational conditions on the sources, Eqs.(79) and (84).

7.2 The source for gravity

We have shown that

$$\frac{1}{1+\chi}\mathcal{R}_b + \Lambda_0\eta_{be}\mathbf{e}^e = -\frac{\kappa}{\alpha}\mathbf{T}_b$$

Expressed in terms of the Einstein tensor this is

$$G_{ab} + \Lambda_C\eta_{ab} = -(n-2)(1+\chi)\frac{\kappa}{\alpha}(T_{ab} - \eta_{ab}T)$$

where the vanishing of $\mathcal{H}_{i\,ab}$ and $\mathcal{G}_{i\,ab}$ leave us with

$$T_{ab} = F^i{}_{ca}F_{i\,db}\eta^{cd}$$

The trace of this is well-known to be gauge dependent, and the conformal symmetry requires the energy tensor to be trace free. Therefore, we are justified in adjusting the $SU(N)$ gauge to give

$$G_{ab} + \Lambda_C\eta_{ab} = -\lambda\left(F^i{}_{ca}F_{i\,db}\eta^{cd} - \frac{1}{4}\eta_{ab}(\eta^{ce}\eta^{df}F_{cd}F_{i\,ef})\right)$$

where

$$\lambda = (n-2)(1+\chi)\frac{\kappa}{\alpha}$$

7.3 The Yang-Mills equation

With $\mathcal{H}^k_{ab} = G^i{}^a{}_b = 0$, Eqs.(48) and (49) reduce to

$$\begin{aligned} 0 &= \eta^{bc} F_{kac}{}^{;a} + \eta^{bc} F_{iac} \beta^i{}_{k}{}^a \\ 0 &= \eta^{bc} (F_{jac;b} + F_{iac} \alpha^i{}_{jb}) \end{aligned}$$

where

$$\begin{aligned} \mathcal{A}^i{}_{k}{}^a &= c^i{}_{jk} A^j{}_a \mathbf{e}^a + c^i{}_{jk} B^{j}{}^a \mathbf{f}_a \\ &= \alpha^i{}_{k}{}^a + \beta^i{}_{k}{}^a \end{aligned}$$

We may use these results and the form of the co-torsion,

$$\mathbf{S}_a = 2(1 + \chi) \Delta_{ea}^{bc} y_b \Lambda \eta_{cd} \mathbf{e}^d \wedge \mathbf{e}^e$$

to solve for the potentials,

$$\begin{aligned} F^i{}_{ab} &= A^i{}_{b;a} - A^i{}_{a;b} - c^i{}_{jk} A^j{}_a A^k{}_b + (1 + \chi) \Lambda B^{i}{}^c \left(\Delta_{bc}^{gf} y_g \eta_{fa} - \Delta_{ac}^{gf} y_g \eta_{fb} \right) \\ 0 &= A^i{}_{b}{}^{;a} - B^{i}{}^a{}_{;b} - c^i{}_{jk} B^{j}{}^a A^k{}_b \\ 0 &= B^{i}{}^{b,a} - B^{i}{}^{a,b} - c^i{}_{jk} B^{j}{}^a B^{k}{}^b \end{aligned} \tag{109}$$

The third equation is the vanishing of the Yang-Mills field strength on the y -submanifold,

$$\mathbf{d}_{(y)} \mathbf{B}^i = -\frac{1}{2} c^i{}_{jk} \mathbf{B}^j \wedge \mathbf{B}^k$$

so that $\mathbf{B}^k$ is a pure-gauge connection for any fixed x^α . Therefore, for each x_0^α we may choose an $SU(N)$ gauge $\Lambda(x_0^\alpha, y_\beta)$ such that $\mathbf{B}^k = 0$. But this makes the value of $\mathbf{B}^k$ independent of x^α as well, so $\mathbf{B}^k = 0$ everywhere. As a result, the fields in terms of the potentials reduce to

$$\begin{aligned} F^i{}_{ab} &= A^i{}_{b;a} - A^i{}_{a;b} - c^i{}_{jk} A^j{}_a A^k{}_b \\ A^i{}_{b}{}^{;a} &= 0 \end{aligned}$$

Now, when we write the field equations in terms of the potentials and set $B^{ka} = 0$, we have

$$\begin{aligned} 0 &= F_{kab}{}^{,a} \\ 0 &= \eta^{bc} (F_{jac;b} + F_{iac}{}^i{}_{jk} A^j{}_a) \end{aligned}$$

The first shows that F_{kab} is independent of y_α and the second shows it to be covariantly divergence free.

8 Conclusions

Gravitational field theories in doubled dimensions include biconformal gravity [4, 23, 5, 6], double field theory [13, 14, 15, 12, 33, 35], and gravity on a Kähler manifold [7, 6]. Each of these cases starts as a fully $2n$ -dimensional theory but ultimately is intended to describe gravity on an n -dimensional submanifold. We have found a satisfactory $2n$ -dimensional form of Yang-Mills matter sources and shown that they also reduce to the expected n -dimensional sources *as a consequence of the field equations*. Our gravitational reduction and the consequent reduction of the Yang-Mills fields and field equations *does not require a section condition*.

While we discussed the issue in biconformal space, our results hold in the related forms of double field theory and Kähler manifolds [7, 6].

For matter fields we restrict our attention to gauged $SU(N)$ sources (Yang-Mills type). While we find that the usual form of $2n$ -dimensional Yang-Mills action gives incorrect coupling to gravity, including a “twist” matrix in the action corrects the problem.

For the gravitational fields we use the most general action linear in the biconformal curvatures. The variation is taken with respect to all $\frac{(n+1)(n+2)}{2}$ conformal gauge fields. In the absence of sources, the use of the gravitational field equations to reduce fully $2n$ -dimensional gravity theory to dependence only on the fields of n -dimensional gravity is well established. The field equations of torsion-free biconformal space restrict the $\frac{1}{2}(n+1)(n+2)$ curvature components, each initially dependent on $2n$ independent coordinates, to the usual locally scale covariant Riemannian curvature tensor in n dimensions. Ultimately, the n -dimensional solder form determines all fields, up to coordinate and gauge transformations. Generic, torsion-free, vacuum solutions describe n -dimensional scale-covariant general relativity on the co-tangent bundle.

Here we have shown that the same reduction occurs when gauged $SU(N)$ field strengths are included as matter sources. The result goes well beyond any previous work. With two exceptions [32, 20] studies of biconformal spaces [23, 5, 20, 8, 9, 7, 10, 6, 22, 11] have considered the pure gravity biconformal spaces,

leading to vacuum general relativity. With $SU(N)$ Yang-Mills fields as gravitational sources, the central issue was to show that a completely general $SU(N)$ gauge theory over a $2n$ -dimensional biconformal space does not disrupt the gravitational reduction to general relativity, but rather itself reduces to a suitable n -dim gravitational source and Yang-Mills field equation.

As with the Riemann-Cartan construction of general relativity above, the development of biconformal spaces from group symmetry made it straightforward to include the additional symmetry of sources. By extending the quotient to

$$\mathcal{M}^{2n} = [SO(p+1, q+1) \times SU(N)] / [SO(p, q) \times SO(1, 1) \times SU(N)]$$

the local symmetry is enlarged by $SU(N)$. We considered the effects of adding an $SU(N)$ action to the gravitational action Eq.(8). As central results we successfully showed:

1. The number of field components in $2n$ dimensions reduces by a factor of $\frac{n-2}{2(2n-1)}$ to the expected number $\frac{n(n-1)}{2} (N^2 - 1)$ on n -dimensional spacetime.
2. The functional dependence of the fields reduces from $2n$ to n independent variables.
3. The usual form of Yang-Mills stress-energy tensor provides the source for the scale-covariant Einstein equation on n -dimensional spacetime.
4. The usual Yang-Mills field equation holds on the spacetime submanifold.

To accomplish these goals we required two interdependent intermediate steps:

1. We considered alternate forms of $2n$ -dimensional Yang-Mills action, showing that the usual action, $S_{YM}^0 = \int tr(\mathbf{F} \wedge^* \mathbf{F})$ gives nonstandard coupling to gravity. Instead, including a “twist” matrix in the action $S_{YM} = \int tr(\bar{\mathbf{F}} \wedge^* \mathbf{F})$ with twisted form

$$\bar{\mathcal{F}}_{AB} = \frac{1}{2} (K_A^C \mathcal{F}_{CB} + \mathcal{F}_{AC} K_B^C)$$

leads to both the usual n -dimensional Yang-Mills source to the Einstein tensor and the usual Yang-Mills equation for the $SU(N)$ fields. A similar twist has been found in other double field theory studies [16, 18, 19] in order to enforce supersymmetry. Here, the twist is required for the bosonic fields alone. Interestingly, the twist matrix $K^A_B = \bar{K}^{AC} g_{CB}$ makes use of both the Kähler and Killing forms, g_{AB} and K_{AB} , respectively.

2. We considered two naturally occurring inner products for the orthonormal frame fields: the restriction to the base manifold of the Killing form, and the Kähler metric. We showed the Kähler form cannot lead to the usual field equations while the variation of the Killing form in the twisted action gives usual Yang-Mills equations and usual coupling to gravity. Previous results in biconformal gravity did not require the inner product.

We also gave a brief discussion of sources in Weyl geometry and in Weyl gravity. For Weyl gravity we indicated the possibility of alternative, fourth order source fields. We give two examples—scalar and electromagnetic fields—of fourth order generalizations for sources. The fourth order fields satisfy field equations which differ from the field equations for the corresponding second order fields only by terms linear in the product of the fields and the curvature. The difference between solutions to Weyl gravity with fourth order sources and general relativity with the corresponding second order sources need not introduce a mass scale.

9 Appendices

Appendix A: Failure of the Kähler inner product

In this Appendix, we show that using the Kähler metric to define orthonormality of the solder and co-solder forms cannot lead to the usual Yang-Mills gravitational coupling to gravity, for either the usual Yang-Mills action or the twisted Yang-Mills action.

We first find the Hodge dual of the Yang-Mills field here using the inverse Kähler metric $\bar{g}^{AB}$. In general terms the Hodge dual of a 2-form is given by Eq.(28), where we now substitute the Kähler metric,

$$\begin{aligned}
{}^*\mathcal{F} = & \frac{1}{n!(n-2)!} \left(\frac{1}{2} F_{ab} \bar{g}^{am} \bar{g}^{bn} + G^a{}_b \bar{g}_a{}^m \bar{g}^{bn} + \frac{1}{2} H^{ab} \bar{g}_a{}^m \bar{g}_b{}^n \right) \varepsilon^{c\dots d}{}_{mne\dots f} \mathbf{f}_{c\dots d} \mathbf{e}^{e\dots f} \\
& + \frac{(-1)^{n-1}}{(n-1)!(n-1)!} \left(\frac{1}{2} F_{ab} \bar{g}^a{}_m \bar{g}^{bn} + G^a{}_b \bar{g}_{am} \bar{g}^{bn} + \frac{1}{2} H^{ab} \bar{g}_{am} \bar{g}_b{}^n \right) \varepsilon^{mc\dots d}{}_{ne\dots f} \mathbf{f}_{c\dots d} \mathbf{e}^{e\dots f} \\
& + \frac{(-1)^n}{(n-1)!(n-1)!} \left(\frac{1}{2} F_{ab} \bar{g}^{am} \bar{g}^b{}_n + G^a{}_b \bar{g}_a{}^m \bar{g}^b{}_n + \frac{1}{2} H^{ab} \bar{g}_a{}^m \bar{g}_{bn} \right) \varepsilon^{nc\dots d}{}_{me\dots f} \mathbf{f}_{c\dots d} \mathbf{e}^{e\dots f} \\
& + \frac{1}{n!(n-2)!} \left(\frac{1}{2} F_{ab} \bar{g}^a{}_m \bar{g}^b{}_n + G^a{}_b \bar{g}_{am} \bar{g}^b{}_n + \frac{1}{2} H^{ab} \bar{g}_{am} \bar{g}_{bn} \right) \varepsilon^{mnc\dots d}{}_{e\dots f} \mathbf{f}_{c\dots d} \mathbf{e}^{e\dots f} \quad (110)
\end{aligned}$$

Forming the usual Yang-Mills Lagrangian density as the wedge product, $\mathcal{F} \wedge {}^*\mathcal{F}$, and eliminating the basis forms in favor of the volume form Φ yields

$$\begin{aligned}
\mathcal{F} \wedge {}^* \mathcal{F} = & \left(\frac{1}{2} F_{mn} \bar{g}^{am} \bar{g}^{bn} + G^m{}_n \bar{g}^a{}_m \bar{g}^{bn} + \frac{1}{2} H^{mn} \bar{g}^a{}_m \bar{g}^b{}_n \right) F_{ab} \Phi \\
& + (F_{mn} \bar{g}^a{}_m \bar{g}^{bn} + G^m{}_n (\bar{g}_{am} \bar{g}^{bn} - \bar{g}^a{}_n \bar{g}^b{}_m) + H^{mn} \bar{g}_{am} \bar{g}^b{}_n) G^a{}_b \Phi \\
& + \left(\frac{1}{2} F_{mn} \bar{g}^a{}_m \bar{g}^b{}_n + G^m{}_n \bar{g}_{am} \bar{g}_b{}^n + \frac{1}{2} H^{mn} \bar{g}_{am} \bar{g}_{bn} \right) H^{ab} \Phi
\end{aligned} \tag{111}$$

For the diagonal form of the Kähler metric, Eq.(17), this Lagrange density reduces to

$$\mathcal{F} \wedge {}^* \mathcal{F} = \left(\frac{1}{2} \eta^{am} \eta^{bn} F_{ab} F_{mn} + \eta_{am} \eta^{bn} G^m{}_n G^a{}_b + \frac{1}{2} \eta_{am} \eta_{bn} H^{mn} H^{ab} \right) \Phi \tag{112}$$

Varying the potentials in Eq.(112) yields the usual Yang-Mills field equation. However, metric variation of Eq.(111) gives a nonstandard coupling to gravity. Varying and reducing the integral of Eq.(111), the resulting gravitational field equations are:

$$\alpha (\Omega^n{}_b{}^b{}_m - \Omega^a{}_b{}^b{}_a \delta_m^n) + \beta (\Omega^n{}_m - \Omega^a{}_a \delta_m^n) + \Lambda \delta_m^n = - (2\eta^{be} F_{me} F_{ab} + 2G^d{}_a G^e{}_m \eta_{de}) \eta^{an} \tag{113}$$

$$[\alpha (\Omega^a{}_n{}^m{}_a - \Omega^a{}_b{}^b{}_a \delta_n^m) + \beta (\Omega^m{}_n - \Omega^a{}_a \delta_n^m) + \Lambda \delta_n^m] = - (2G^d{}_c G^n{}_b \eta_{dm} \eta^{bc} + 2\eta_{bc} \eta_{dm} H^{dc} H^{nb}) \tag{114}$$

$$\alpha \Omega^a{}_{nam} + \beta \Omega_{nm} = (2\eta^{ca} G^b{}_a F_{mc} + 2\eta_{ac} G^c{}_m H^{ab}) \eta_{bn} \tag{115}$$

$$\alpha \Omega^n{}_b{}^{bm} + \beta \Omega^{nm} = - (2\eta^{ca} G^m{}_a F_{bc} + 2\eta_{ac} G^c{}_b H^{am}) \eta^{nb} \tag{116}$$

The remaining four field equations involving the torsion and co-torsion are unchanged.

Notice that the sources on the right sides of Eqs.(115) and (116) differ only in the overall sign. This means that both the spacetime curvature, $\Omega^a{}_{nam}$, and the momentum space curvature, $\Omega^n{}_b{}^{bm}$, are driven with equal strength. Thus, if spacetime curvature is nonzero, the momentum space must also be correspondingly curved. In torsion-free solutions the left hand side of Eq.(116) vanishes independently of the sources, implying a constraint on the source fields. Thus, for the $\int \mathcal{F} \wedge {}^* \mathcal{F}$ source and the Kähler case,

$$\eta^{ca} G^m{}_a F_{bc} + \eta_{ac} G^c{}_b H^{am} = 0$$

and this immediately shows that the source for the Einstein equation, Eq.(115), vanishes.

The necessity for equal spacetime and momentum curvatures suggests the possibility implementing Born reciprocity. This idea will be explored elsewhere. However, momentum curvature also requires some part of the torsion to be nonvanishing, and this in turn requires a different gravitational solution than that known

to reproduce general relativity. Thus, we cannot maintain vanishing torsion without forcing the spacetime source to vanish.

In addition to the inescapability of torsion and momentum space curvature, the independence of the sources to Eqs.(113) and (114) also raises issues with the Kähler variation, because the method of solution employed in [6] makes use of the near identity of these two equations. At the very least, an entirely different form of reduction of the equations would be required, with no guarantee that the Einstein equation would emerge.

A similar calculation shows that the same difficulties arise from the twisted form of the action using the Kähler metric.

These issues do not arise with the Killing variation, Eqs.(24)—the source for the spacetime curvature and momentum space curvature are independent while remaining two variations are identical. The use of the Killing form as metric also makes good geometric sense, since it arises directly as a symmetric form in the Lie algebra and thus as metric of the co-tangent space. As such, it respects the conformal invariance of the full model. The Kähler structure, by contrast, reflects symmetries and dynamical properties within the conformal group and depend for their existence on the solution on the biconformal space. It is not conformally invariant.

Appendix B: Symmetry of the twist matrix

Writing the twist matrix while keeping factors of η^{ab} explicit, we have

$$\begin{aligned}
K^A{}_B &= \begin{pmatrix} \bar{K}^{ac} & \bar{K}^a{}_e \eta^{ec} \\ \eta^{ae} \bar{K}_e{}^c & \eta^{ae} \eta^{ef} \bar{K}_{ef} \end{pmatrix} \begin{pmatrix} \eta_{cb} & 0 \\ 0 & \eta_{cm} \eta_{bn} \eta^{mn} \end{pmatrix} \\
&= \begin{pmatrix} \bar{K}^{ac} \eta_{cb} & \bar{K}^a{}_b \\ \eta^{ae} \bar{K}_e{}^c \eta_{cb} & \eta^{ae} \bar{K}_{eb} \end{pmatrix} \\
K_A{}^B &= \begin{pmatrix} \eta_{ad} \bar{K}^{db} & \bar{K}_a{}^b \\ \eta_{ad} \bar{K}^d{}_e \eta^{eb} & \bar{K}_{ae} \eta^{eb} \end{pmatrix}
\end{aligned}$$

Symmetry of $K^A{}_B$ follows from the symmetry of the Killing and Kähler forms, $K_{AB} = K_{BA}, g_{AB} = g_{BA}$:

$$K^A{}_B \equiv \bar{K}^{AC} g_{CB} = g_{BC} \bar{K}^{CA} = K_B{}^A$$

Comparing the expressions,

$$\begin{pmatrix} K^A{}_B & = & K_B{}^A \\ \left(\begin{array}{cc} \bar{K}^{ac} \eta_{cb} & \bar{K}^a{}_b \\ \eta^{ae} \bar{K}_e{}^c \eta_{cb} & \eta^{ae} \bar{K}_{eb} \end{array} \right) & = & \left(\begin{array}{cc} \eta_{bc} \bar{K}^{ca} & \eta_{bc} \bar{K}^c{}_e \eta^{ea} \\ \bar{K}_b{}^a & \bar{K}_{bc} \eta^{ca} \end{array} \right) \end{pmatrix}$$

Therefore, from the upper right quadrant we must have $\bar{K}^a{}_b = \eta_{bc} \bar{K}^c{}_e \eta^{ea}$ and similarly $\eta^{ae} \bar{K}_e{}^c \eta_{cb} = \bar{K}_b{}^a$ from the lower left. However we also have symmetry $\bar{K}^{AB} = \bar{K}^{BA}$ of $\bar{K}^{AB}$ itself:

$$\begin{aligned} \bar{K}^{AB} &= \begin{pmatrix} \bar{K}^{ab} & \bar{K}^a{}_e \eta^{eb} \\ \eta^{ae} \bar{K}_e{}^b & \eta^{ae} \eta^{bf} \bar{K}_{ef} \end{pmatrix} \\ [\bar{K}^t]^{BA} &= \begin{pmatrix} \bar{K}^{ba} & \eta^{be} \bar{K}_e{}^a \\ \bar{K}^b{}_e \eta^{ea} & \eta^{ae} \eta^{bf} \bar{K}_{ef} \end{pmatrix} \end{aligned}$$

This shows that $\bar{K}^a{}_e \eta^{eb} = \eta^{be} \bar{K}_e{}^a$ from which it follows that

$$\bar{K}^a{}_b = \bar{K}_b{}^a$$

Combine this with $\eta^{ae} \bar{K}_e{}^c \eta_{cb} = \bar{K}_b{}^a$ and we see that all forms are equivalent,

$$\eta^{ae} \bar{K}_e{}^c \eta_{cb} = \bar{K}_b{}^a = \bar{K}^a{}_b = \eta_{bc} \bar{K}^c{}_e \eta^{ea}$$

With this, we may write

$$\begin{aligned} K^A{}_B &= \begin{pmatrix} \bar{K}^{ac} \eta_{cb} & \bar{K}^a{}_b \\ \bar{K}^a{}_b & \eta^{ae} \bar{K}_{eb} \end{pmatrix} \\ K_B{}^A &= \begin{pmatrix} \eta_{bd} \bar{K}^{da} & \bar{K}_b{}^a \\ \bar{K}_b{}^a & \bar{K}_{be} \eta^{ea} \end{pmatrix} = K^A{}_B \end{aligned}$$

Appendix C: Details of the metric variation of the action

The variation of the twisted action is lengthy and includes some subtleties, so we include details here.

We have the dual field,

$$\begin{aligned}
{}^*\mathcal{F} &= \frac{1}{n!(n-2)!} \left(\frac{1}{2} F_{ab} \bar{K}^{am} \bar{K}^{bn} + \mathcal{G}_{gb} \eta^{ga} \bar{K}_a{}^m \bar{K}^{bn} + \frac{1}{2} \mathcal{H}_{gh} \eta^{ga} \bar{K}_a{}^m \eta^{hb} \bar{K}_b{}^n \right) \varepsilon^{c\dots d}{}_{mne\dots f} \mathbf{f}_{c\dots d} \mathbf{e}^{e\dots f} \\
&+ \frac{(-1)^{n-1}}{(n-1)!(n-1)!} \left(\frac{1}{2} F_{ab} \bar{K}^a{}_m \bar{K}^{bn} + \mathcal{G}_{gb} \eta^{ga} \bar{K}_{am} \bar{K}^{bn} + \frac{1}{2} \mathcal{H}_{gh} \eta^{ga} \eta^{hb} \bar{K}_{am} \bar{K}_b{}^n \right) \varepsilon^{mc\dots d}{}_{ne\dots f} \mathbf{f}_{c\dots d} \mathbf{e}^{e\dots f} \\
&+ \frac{(-1)^n}{(n-1)!(n-1)!} \left(\frac{1}{2} F_{ab} \bar{K}^{am} \bar{K}^b{}_n + \mathcal{G}_{gb} \eta^{ga} \bar{K}_a{}^m \bar{K}^b{}_n + \frac{1}{2} \mathcal{H}_{gh} \eta^{ga} \eta^{hb} \bar{K}_a{}^m \bar{K}_{bn} \right) \varepsilon^{nc\dots d}{}_{me\dots f} \mathbf{f}_{c\dots d} \mathbf{e}^{e\dots f} \\
&+ \frac{1}{n!(n-2)!} \left(\frac{1}{2} F_{ab} \bar{K}^a{}_m \bar{K}^b{}_n + \mathcal{G}_{gb} \eta^{ga} \bar{K}_{am} \bar{K}^b{}_n + \frac{1}{2} \mathcal{H}_{gh} \eta^{ga} \eta^{hb} \bar{K}_{am} \bar{K}_{bn} \right) \varepsilon^{mnc\dots d}{}_{e\dots f} \mathbf{f}_{c\dots d} \mathbf{e}^{e\dots f}
\end{aligned}$$

and, changing indices to avoid duplication, the barred field,

$$\begin{aligned}
\bar{\mathcal{F}} &= \frac{1}{2} (F_{rq} \bar{K}^{qt} \eta_{ts} + \bar{K}_r{}^q \mathcal{G}_{qs}) \mathbf{e}^r \wedge \mathbf{e}^s \\
&+ \frac{1}{2} (F_{rc} \bar{K}^c{}_s + \bar{K}_r{}^c \mathcal{H}_{cs}) \eta^{sw} \mathbf{e}^r \wedge \mathbf{f}_w \\
&+ \frac{1}{2} (-\mathcal{G}_{sq} \bar{K}^{qt} \eta_{tr} - \mathcal{G}_{qr} \eta^{qt} \bar{K}_{ts}) \eta^{sw} \mathbf{e}^r \wedge \mathbf{f}_w \\
&+ \frac{1}{2} (\mathcal{G}_{rq} \bar{K}^q{}_s + \mathcal{H}_{rq} \eta^{qt} \bar{K}_{ts}) \eta^{rw} \eta^{sx} \mathbf{f}_w \wedge \mathbf{f}_x
\end{aligned}$$

We must wedge these together, then vary the metric.

Wedging,

$$\begin{aligned}
\bar{\mathcal{F}} \wedge^* \mathcal{F} &= \frac{1}{2} \frac{1}{n!(n-2)!} (F_{rq} \bar{K}^{qt} \eta_{ts} + \bar{K}_r{}^q \mathcal{G}_{qs}) \\
&\times \left(\mathcal{G}_{gb} \eta^{ga} \bar{K}_a{}^m \bar{K}^{bn} + \frac{1}{2} \mathcal{H}_{gh} \eta^{ga} \bar{K}_a{}^m \eta^{hb} \bar{K}_b{}^n \right) \varepsilon^{c\dots d}{}_{mne\dots f} \mathbf{f}_{c\dots d} \mathbf{e}^{rse\dots f} \\
&- \frac{1}{2} \frac{1}{(n-1)!(n-1)!} (F_{rc} \bar{K}^c{}_s + \bar{K}_r{}^c \mathcal{H}_{cs}) \eta^{sw} \\
&\times \left(\frac{1}{2} F_{ab} \bar{K}^a{}_m \bar{K}^{bn} + \frac{1}{2} \mathcal{H}_{gh} \eta^{ga} \eta^{hb} \bar{K}_{am} \bar{K}_b{}^n \right) \varepsilon^{mc\dots d}{}_{ne\dots f} \mathbf{f}_{wc\dots d} \mathbf{e}^{re\dots f} \\
&+ \frac{1}{2} \frac{1}{(n-1)!(n-1)!} (F_{rc} \bar{K}^c{}_s + \bar{K}_r{}^c \mathcal{H}_{cs}) \eta^{sw} \\
&\times \left(\frac{1}{2} F_{ab} \bar{K}^{am} \bar{K}^b{}_n + \mathcal{G}_{gb} \eta^{ga} \bar{K}_a{}^m \bar{K}^b{}_n + \frac{1}{2} \mathcal{H}_{gh} \eta^{ga} \eta^{hb} \bar{K}_a{}^m \bar{K}_{bn} \right) \varepsilon^{nc\dots d}{}_{me\dots f} \mathbf{f}_{wc\dots d} \mathbf{e}^{re\dots f} \\
&- \frac{1}{2} \frac{1}{(n-1)!(n-1)!} (-\mathcal{G}_{sq} \bar{K}^{qt} \eta_{tr} - \mathcal{G}_{qr} \eta^{qt} \bar{K}_{ts}) \eta^{sw} \\
&\times \left(\frac{1}{2} F_{ab} \bar{K}^a{}_m \bar{K}^{bn} + \frac{1}{2} \mathcal{H}_{gh} \eta^{ga} \eta^{hb} \bar{K}_{am} \bar{K}_b{}^n \right) \varepsilon^{mc\dots d}{}_{ne\dots f} \mathbf{f}_{wc\dots d} \mathbf{e}^{re\dots f} \\
&+ \frac{1}{2} \frac{1}{(n-1)!(n-1)!} (-\mathcal{G}_{sq} \bar{K}^{qt} \eta_{tr} - \mathcal{G}_{qr} \eta^{qt} \bar{K}_{ts}) \eta^{sw} \\
&\times \left(\frac{1}{2} F_{ab} \bar{K}^{am} \bar{K}^b{}_n + \mathcal{G}_{gb} \eta^{ga} \bar{K}_a{}^m \bar{K}^b{}_n + \frac{1}{2} \mathcal{H}_{gh} \eta^{ga} \eta^{hb} \bar{K}_a{}^m \bar{K}_{bn} \right) \varepsilon^{nc\dots d}{}_{me\dots f} \mathbf{f}_{wc\dots d} \mathbf{e}^{re\dots f} \\
&+ \frac{1}{2} \frac{1}{n!(n-2)!} (\mathcal{G}_{rq} \bar{K}^q{}_s + \mathcal{H}_{rq} \eta^{qt} \bar{K}_{ts}) \eta^{rw} \eta^{sx} \\
&\times \left(\frac{1}{2} F_{ab} \bar{K}^a{}_m \bar{K}^b{}_n + \mathcal{G}_{gb} \eta^{ga} \bar{K}_{am} \bar{K}^b{}_n \right) \varepsilon^{mnc\dots d}{}_{e\dots f} \mathbf{f}_{wxc\dots d} \mathbf{e}^{e\dots f}
\end{aligned}$$

Now use the relation between the basis forms and the volume element, and the Kronecker reduction of pairs of Levi-Civita tensors,

$$\begin{aligned}
\mathbf{f}_{c\dots d} \wedge \mathbf{e}^{e\dots f} &= \bar{e}_{c\dots d}{}^{e\dots f} \Phi \\
\varepsilon^{mnc\dots d}{}_{e\dots f} \bar{e}_{pqc\dots d}{}^{e\dots f} &= n! (n-2)! (\delta_p^m \delta_q^n - \delta_p^n \delta_q^m) \\
\varepsilon^{mc\dots d}{}_{ne\dots f} \bar{e}_{pc\dots d}{}^{qe\dots f} &= (n-1)! (n-1)! \delta_p^m \delta_n^q
\end{aligned}$$

to reduce the Lagrange density to

$$\begin{aligned}
\bar{\mathcal{F}} \wedge {}^* \mathcal{F} = & \frac{1}{2} (F_{rq} \bar{K}^{qt} \eta_{ts} + \bar{K}_r{}^q \mathcal{G}_{qs}) \left(\mathcal{G}_{gb} \eta^{ga} \bar{K}_a{}^m \bar{K}^{bn} + \frac{1}{2} \mathcal{H}_{gh} \eta^{ga} \bar{K}_a{}^m \eta^{hb} \bar{K}_b{}^n \right) (\delta_m^r \delta_n^s - \delta_m^s \delta_n^r) \Phi \\
& - \frac{1}{2} (F_{rc} \bar{K}^c{}_s + \bar{K}_r{}^c \mathcal{H}_{cs}) \eta^{sw} \left(\frac{1}{2} F_{ab} \bar{K}^a{}_m \bar{K}^{bn} + \frac{1}{2} \mathcal{H}_{gh} \eta^{ga} \eta^{hb} \bar{K}_{am} \bar{K}_b{}^n \right) \delta_w^m \delta_n^r \Phi \\
& + \frac{1}{2} (F_{rc} \bar{K}^c{}_s + \bar{K}_r{}^c \mathcal{H}_{cs}) \eta^{sw} \left(\frac{1}{2} F_{ab} \bar{K}^{am} \bar{K}^b{}_n + \mathcal{G}_{gb} \eta^{ga} \bar{K}_a{}^m \bar{K}^b{}_n + \frac{1}{2} \mathcal{H}_{gh} \eta^{ga} \eta^{hb} \bar{K}_a{}^m \bar{K}_{bn} \right) \delta_w^m \delta_m^r \Phi \\
& - \frac{1}{2} (-\mathcal{G}_{sq} \bar{K}^{qt} \eta_{tr} - \mathcal{G}_{qr} \eta^{qt} \bar{K}_{ts}) \eta^{sw} \left(\frac{1}{2} F_{ab} \bar{K}^a{}_m \bar{K}^{bn} + \frac{1}{2} \mathcal{H}_{gh} \eta^{ga} \eta^{hb} \bar{K}_{am} \bar{K}_b{}^n \right) \delta_w^m \delta_n^r \Phi \\
& + \frac{1}{2} (-\mathcal{G}_{sq} \bar{K}^{qt} \eta_{tr} - \mathcal{G}_{qr} \eta^{qt} \bar{K}_{ts}) \eta^{sw} \left(\frac{1}{2} F_{ab} \bar{K}^{am} \bar{K}^b{}_n + \mathcal{G}_{gb} \eta^{ga} \bar{K}_a{}^m \bar{K}^b{}_n + \frac{1}{2} \mathcal{H}_{gh} \eta^{ga} \eta^{hb} \bar{K}_a{}^m \bar{K}_{bn} \right) \delta_w^n \delta_m^r \Phi \\
& + \frac{1}{2} (\mathcal{G}_{rq} \bar{K}^q{}_s + \mathcal{H}_{rq} \eta^{qt} \bar{K}_{ts}) \eta^{rw} \eta^{sx} \left(\frac{1}{2} F_{ab} \bar{K}^a{}_m \bar{K}^b{}_n + \mathcal{G}_{gb} \eta^{ga} \bar{K}_{am} \bar{K}^b{}_n \right) (\delta_w^m \delta_x^n - \delta_w^n \delta_x^m) \Phi
\end{aligned}$$

Then, absorbing the η_{ab} and δ_b^a factors,

$$\begin{aligned}
\bar{\mathcal{F}} \wedge {}^* \mathcal{F} = & \frac{1}{2} (F_{mq} \bar{K}^{qt} \eta_{tn} - F_{nq} \bar{K}^{qt} \eta_{tm} + \bar{K}_m{}^q \mathcal{G}_{qn} - \bar{K}_n{}^q \mathcal{G}_{qm}) \left(\mathcal{G}_{gb} \eta^{ga} \bar{K}_a{}^m \bar{K}^{bn} + \frac{1}{2} \mathcal{H}_{gh} \eta^{ga} \bar{K}_a{}^m \eta^{hb} \bar{K}_b{}^n \right) \Phi \\
& - \frac{1}{2} (F_{nc} \bar{K}^c{}_s + \bar{K}_n{}^c \mathcal{H}_{cs}) \eta^{sm} \left(\frac{1}{2} F_{ab} \bar{K}^a{}_m \bar{K}^{bn} + \frac{1}{2} \mathcal{H}_{gh} \eta^{ga} \eta^{hb} \bar{K}_{am} \bar{K}_b{}^n \right) \Phi \\
& + \frac{1}{2} (F_{mc} \bar{K}^c{}_s + \bar{K}_m{}^c \mathcal{H}_{cs}) \eta^{sn} \left(\frac{1}{2} F_{ab} \bar{K}^{am} \bar{K}^b{}_n + \mathcal{G}_{gb} \eta^{ga} \bar{K}_a{}^m \bar{K}^b{}_n + \frac{1}{2} \mathcal{H}_{gh} \eta^{ga} \eta^{hb} \bar{K}_a{}^m \bar{K}_{bn} \right) \Phi \\
& - \frac{1}{2} (-\mathcal{G}_{sq} \bar{K}^{qt} \eta_{tn} - \mathcal{G}_{qn} \eta^{qt} \bar{K}_{ts}) \eta^{sm} \left(\frac{1}{2} F_{ab} \bar{K}^a{}_m \bar{K}^{bn} + \frac{1}{2} \mathcal{H}_{gh} \eta^{ga} \eta^{hb} \bar{K}_{am} \bar{K}_b{}^n \right) \Phi \\
& + \frac{1}{2} (-\mathcal{G}_{sq} \bar{K}^{qt} \eta_{tm} - \mathcal{G}_{qm} \eta^{qt} \bar{K}_{ts}) \eta^{sn} \left(\frac{1}{2} F_{ab} \bar{K}^{am} \bar{K}^b{}_n + \mathcal{G}_{gb} \eta^{ga} \bar{K}_a{}^m \bar{K}^b{}_n + \frac{1}{2} \mathcal{H}_{gh} \eta^{ga} \eta^{hb} \bar{K}_a{}^m \bar{K}_{bn} \right) \Phi \\
& + \frac{1}{2} (\mathcal{G}_{rq} \bar{K}^q{}_s \eta^{rm} \eta^{sn} - \mathcal{G}_{rq} \bar{K}^q{}_s \eta^{rn} \eta^{sm} + \mathcal{H}_{rq} \eta^{qt} \bar{K}_{ts} \eta^{rm} \eta^{sn} - \mathcal{H}_{rq} \eta^{qt} \bar{K}_{ts} \eta^{rn} \eta^{sm}) \\
& \times \left(\frac{1}{2} F_{ab} \bar{K}^a{}_m \bar{K}^b{}_n + \mathcal{G}_{gb} \eta^{ga} \bar{K}_{am} \bar{K}^b{}_n \right) \Phi
\end{aligned}$$

After varying the metric, the null-orthonormal form of the metric is restored, so we may anticipate this and drop terms in the product such as $\mathcal{G}_{gb} \eta^{ga} \bar{K}_a{}^m \bar{K}^{bn} \mathcal{G}_{sq} \bar{K}^{qt} \eta_{tr}$ which will ultimately vanish. Terms with two or more factors of $\bar{K}^{ab}$ and/or $\bar{K}_{ab}$ will vanish, so we have only terms with one of these and two off diagonal factors such as $\bar{K}_a{}^b$, or terms with three off diagonal factors. In all cases where there is one factor of $\bar{K}^{ab}$ or $\bar{K}_{ab}$, it is this factor that must be varied so the off diagonal factors may be replaced. For example, once the null orthonormal basis is restored, the only surviving term of the variation of

$$F_{mq} \eta_{tn} \frac{1}{2} \mathcal{H}_{gh} \eta^{ga} \eta^{hb} \bar{K}^{qt} \bar{K}_a{}^m \bar{K}_b{}^n$$

will be

$$F_{mq}\eta_{tn}\frac{1}{2}\mathcal{H}_{gh}\eta^{ga}\eta^{hb}\delta\bar{K}^{qt}\bar{K}_a{}^m\bar{K}_b{}^n$$

which we may immediately write as

$$F_{mq}\eta_{tn}\frac{1}{2}\mathcal{H}_{gh}\eta^{ga}\eta^{hb}\delta\bar{K}^{qt}\delta_a{}^m\delta_b{}^n = \frac{1}{2}(F_{aq}\eta_{tb}\mathcal{H}_{gh}\eta^{ga}\eta^{hb})\delta\bar{K}^{qt}$$

Terms with three off-diagonal components of the metric must be retained until after variation.

Distributing fully, then making these reductions where possible, we collect terms

$$\begin{aligned}\bar{\mathcal{F}} \wedge {}^*\mathcal{F} &= \frac{1}{2}\left(\frac{1}{2}F_{ec}\eta^{cb}F_{db} - \frac{1}{2}F_{ec}\eta^{ca}F_{ad} + \frac{1}{2}F_{ad}\eta_{eb}H^{ab} + \frac{1}{2}\mathcal{H}_{ec}\eta^{cb}F_{db} - \frac{1}{2}F_{bd}\eta_{ea}H^{ab} - \frac{1}{2}\mathcal{H}_{ec}\eta^{ca}F_{ad}\right)\bar{K}^{de}\Phi \\ &+ \frac{1}{2}\left(-\mathcal{G}_{ea}\mathcal{G}_{cd}\eta^{ca} + \mathcal{G}_{ae}\mathcal{G}_{cd}\eta^{ca} - \mathcal{G}_{cd}\eta_{ea}\eta^{cb}\mathcal{G}_{gb}\eta^{ga}\right)\bar{K}^{de}\Phi \\ &+ \frac{1}{2}\left(-\frac{1}{2}F_{bc}\eta^{ce}H^{db} - \mathcal{H}_{bc}\eta^{ce}\frac{1}{2}H^{db} + \frac{1}{2}F_{ac}\eta^{ce}H^{ad} + \frac{1}{2}\mathcal{H}_{ac}\eta^{ce}H^{ad} + \frac{1}{2}H^{ad}\eta^{eb}F_{ab} - \frac{1}{2}H^{bd}\eta^{ea}F_{ab}\right)\bar{K}_{de}\Phi \\ &+ \frac{1}{2}\left(\mathcal{G}_{rs}\eta^{re}\eta^{sb}\mathcal{G}_{gb}\eta^{gd} - \mathcal{G}_{rs}\eta^{rb}\eta^{se}\mathcal{G}_{gb}\eta^{gd} - \mathcal{G}_{qa}\eta^{qd}\eta^{eb}\mathcal{G}_{gb}\eta^{ga}\right)\bar{K}_{de}\Phi \\ &+ \frac{1}{2}\left(\frac{1}{2}\mathcal{G}_{qn}H^{ab}\delta_m^c - \frac{1}{2}\mathcal{G}_{qm}H^{ab}\delta_n^c + F_{mq}\eta^{cb}\mathcal{G}_{gn}\eta^{ga}\right)\bar{K}^q{}_c\bar{K}_a{}^m\bar{K}_b{}^n\Phi \\ &+ \frac{1}{2}\left(\mathcal{H}_{qs}\eta^{sb}\mathcal{G}_{gn}\eta^{ga}\delta_m^c + \frac{1}{2}\mathcal{G}_{rq}\eta^{ra}\eta^{cb}F_{mn} - \frac{1}{2}\mathcal{G}_{rq}\eta^{rb}\eta^{ca}F_{mn}\right)\bar{K}^m{}_a\bar{K}^n{}_b\bar{K}^q{}_c\Phi\end{aligned}$$

and consolidate using symmetries,

$$\begin{aligned}\bar{\mathcal{F}} \wedge {}^*\mathcal{F} &= \frac{1}{2}(F_{dc}\eta^{cb}F_{eb} + 2F_{ad}H^{ab}\eta_{be})\bar{K}^{de}\Phi \\ &+ \frac{1}{2}(\mathcal{G}_{ae}\mathcal{G}_{cd}\eta^{ca} - 2\mathcal{G}_{ea}\mathcal{G}_{cd}\eta^{ca})\bar{K}^{de}\Phi \\ &+ \frac{1}{2}(\mathcal{H}_{ab}H^{ad}\eta^{be} + 2F_{ac}H^{ad}\eta^{ce})\bar{K}_{de}\Phi \\ &+ \frac{1}{2}(\mathcal{G}_{ca}\mathcal{G}_{db}\eta^{ab} - 2\mathcal{G}_{da}\mathcal{G}_{bc}\eta^{ab})\eta^{de}\eta^{cf}\bar{K}_{ef}\Phi \\ &+ \frac{1}{2}\left(\frac{1}{2}\mathcal{G}_{dn}H^{ab}\delta_m^c - \frac{1}{2}\mathcal{G}_{dm}H^{ab}\delta_n^c + F_{md}\eta^{cb}\mathcal{G}_{gn}\eta^{ga}\right)\bar{K}_a{}^m\bar{K}_b{}^n\bar{K}^d{}_c\Phi \\ &+ \frac{1}{2}\left(\mathcal{H}_{ds}\eta^{sb}\mathcal{G}_{gn}\eta^{ga}\delta_m^c + \frac{1}{2}\mathcal{G}_{rd}\eta^{ra}\eta^{cb}F_{mn} - \frac{1}{2}\mathcal{G}_{rd}\eta^{rb}\eta^{ca}F_{mn}\right)\bar{K}^m{}_a\bar{K}^n{}_b\bar{K}^d{}_c\Phi\end{aligned}$$

Checking the limit in the orthonormal basis, we find the correct form, $\bar{\mathcal{F}} \wedge {}^*\mathcal{F} = (\mathcal{G}_{ab}H^{ab} + F_{ab}\eta^{ac}\eta^{bd}\mathcal{G}_{cd})\Phi$.

Proceeding, we vary the metric

$$\begin{aligned}
\delta(\bar{\mathcal{F}} \wedge {}^* \mathcal{F}) = & \frac{1}{2} (F_{dc} \eta^{cb} F_{eb} + 2F_{ad} H^{ab} \eta_{be} + (\mathcal{G}_{ae} - 2\mathcal{G}_{ea}) \mathcal{G}_{cd} \eta^{ca}) \delta \bar{K}^{de} \Phi \\
& + \frac{1}{2} \bar{K}_b{}^n \bar{K}^d{}_c \left(\frac{1}{2} \mathcal{G}_{db} H^{ab} \delta_m^c - \frac{1}{2} \mathcal{G}_{dm} H^{ab} \delta_b^c + F_{md} \eta^{cb} \mathcal{G}_{gb} \eta^{ga} \right) \delta \bar{K}_a{}^m \Phi \\
& + \frac{1}{2} \left(\frac{1}{2} \mathcal{G}_{dn} H^{ab} \delta_m^c - \frac{1}{2} \mathcal{G}_{dm} H^{ab} \delta_n^c + F_{md} \eta^{cb} \mathcal{G}_{gn} \eta^{ga} \right) \bar{K}_a{}^m \delta \bar{K}_b{}^n \bar{K}^d{}_c \Phi \\
& + \frac{1}{2} \left(\frac{1}{2} \mathcal{G}_{dn} H^{ab} \delta_m^c - \frac{1}{2} \mathcal{G}_{dm} H^{ab} \delta_n^c + F_{md} \eta^{cb} \mathcal{G}_{gn} \eta^{ga} \right) \bar{K}_a{}^m \bar{K}_b{}^n \delta \bar{K}^d{}_c \Phi \\
& + \frac{1}{2} \left(\mathcal{H}_{ds} \eta^{sb} \mathcal{G}_{gn} \eta^{ga} \delta_m^c + \frac{1}{2} \mathcal{G}_{rd} \eta^{ra} \eta^{cb} F_{mn} - \frac{1}{2} \mathcal{G}_{rd} \eta^{rb} \eta^{ca} F_{mn} \right) \delta \bar{K}^m{}_a \bar{K}^n{}_b \bar{K}^d{}_c \Phi \\
& + \frac{1}{2} \left(\mathcal{H}_{ds} \eta^{sb} \mathcal{G}_{gn} \eta^{ga} \delta_m^c + \frac{1}{2} \mathcal{G}_{rd} \eta^{ra} \eta^{cb} F_{mn} - \frac{1}{2} \mathcal{G}_{rd} \eta^{rb} \eta^{ca} F_{mn} \right) \bar{K}^m{}_a \delta \bar{K}^n{}_b \bar{K}^d{}_c \Phi \\
& + \frac{1}{2} \left(\mathcal{H}_{ds} \eta^{sb} \mathcal{G}_{gn} \eta^{ga} \delta_m^c + \frac{1}{2} \mathcal{G}_{rd} \eta^{ra} \eta^{cb} F_{mn} - \frac{1}{2} \mathcal{G}_{rd} \eta^{rb} \eta^{ca} F_{mn} \right) \bar{K}^m{}_a \bar{K}^n{}_b \delta \bar{K}^d{}_c \Phi \\
& + \frac{1}{2} (\mathcal{H}_{ab} H^{ad} \eta^{be} + 2F_{ac} H^{ad} \eta^{ce}) \bar{K}_{de} \Phi + \frac{1}{2} \mathcal{G}_{ca} \eta^{ab} (\mathcal{G}_{db} - 2\mathcal{G}_{bd}) \eta^{de} \eta^{cf} \bar{K}_{ef} \Phi \\
& + (\mathcal{G}_{ab} H^{ab} + F_{ab} \eta^{ac} \eta^{bd} \mathcal{G}_{cd}) \delta \Phi
\end{aligned}$$

At this point we may replace remaining unvaried metric components, leaving

$$\begin{aligned}
\delta(\bar{\mathcal{F}} \wedge {}^* \mathcal{F}) = & \frac{1}{2} (F_{dc} \eta^{cb} F_{eb} + 2F_{ad} H^{ab} \eta_{be} + (\mathcal{G}_{ae} - 2\mathcal{G}_{ea}) \mathcal{G}_{cd} \eta^{ca}) \delta \bar{K}^{de} \Phi \\
& + \frac{1}{2} \left(\frac{1}{2} \mathcal{G}_{cb} H^{ab} \delta_m^c - \frac{1}{2} \mathcal{G}_{cm} H^{ab} \delta_b^c + F_{mc} \eta^{cb} \mathcal{G}_{gb} \eta^{ga} \right) \delta \bar{K}_a{}^m \Phi \\
& + \frac{1}{2} \left(\frac{1}{2} \mathcal{G}_{an} H^{ab} - \frac{1}{2} \mathcal{G}_{na} H^{ab} + F_{ac} \eta^{cn} \mathcal{G}_{gb} \eta^{ga} \right) \delta \bar{K}_b{}^n \Phi \\
& + \frac{1}{2} \left(\frac{1}{2} \mathcal{G}_{db} H^{cb} - \frac{1}{2} \mathcal{G}_{da} H^{ac} + F_{ad} \eta^{cb} \mathcal{G}_{gb} \eta^{ga} \right) \bar{K}_b{}^n \delta \bar{K}^d{}_c \Phi \\
& + \frac{1}{2} \left(\mathcal{H}_{cs} \eta^{sb} \mathcal{G}_{gb} \eta^{ga} \delta_m^c + \frac{1}{2} \mathcal{G}_{rc} \eta^{ra} \eta^{cb} F_{mb} - \frac{1}{2} \mathcal{G}_{rc} \eta^{rb} \eta^{ca} F_{mb} \right) \delta \bar{K}^m{}_a \Phi \\
& + \frac{1}{2} \left(\mathcal{H}_{cs} \eta^{sb} \mathcal{G}_{gn} \eta^{gc} + \frac{1}{2} \mathcal{G}_{rc} \eta^{ra} \eta^{cb} F_{an} - \frac{1}{2} \mathcal{G}_{rc} \eta^{rb} \eta^{ca} F_{an} \right) \delta \bar{K}^n{}_b \Phi \\
& + \frac{1}{2} \left(\mathcal{H}_{ds} \eta^{sb} \mathcal{G}_{gb} \eta^{gc} + \frac{1}{2} \mathcal{G}_{rd} \eta^{ra} \eta^{cb} F_{ab} - \frac{1}{2} \mathcal{G}_{rd} \eta^{rb} \eta^{ca} F_{ab} \right) \delta \bar{K}^d{}_c \Phi \\
& + \frac{1}{2} ((\mathcal{H}_{ab} H^{ae} \eta^{bf} + 2F_{ac} H^{ae} \eta^{cf}) + \mathcal{G}_{ca} \eta^{ab} (\mathcal{G}_{db} - 2\mathcal{G}_{bd}) \eta^{de} \eta^{cf}) \bar{K}_{ef} \Phi \\
& + (\mathcal{G}_{ab} H^{ab} + F_{ab} \eta^{ac} \eta^{bd} \mathcal{G}_{cd}) \delta \Phi
\end{aligned}$$

Finally, we collect all terms by type of variation,

$$\begin{aligned}
\delta (\bar{\mathcal{F}} \wedge {}^* \mathcal{F}) &= \frac{1}{2} (F_{dc} \eta^{cb} F_{eb} + 2F_{ad} H^{ab} \eta_{be} + (\mathcal{G}_{ae} - 2\mathcal{G}_{ea}) \mathcal{G}_{cd} \eta^{ca}) \delta \bar{K}^{de} \Phi \\
&+ \frac{1}{2} \left(\frac{1}{2} \mathcal{G}_{cb} H^{nb} \delta_m^c - \frac{1}{2} \mathcal{G}_{cm} H^{nb} \delta_b^c + F_{mc} \eta^{cb} \mathcal{G}_{gb} \eta^{gn} \right) \delta \bar{K}_n{}^m \Phi \\
&+ \frac{1}{2} \left(\frac{1}{2} \mathcal{G}_{am} H^{an} - \frac{1}{2} \mathcal{G}_{ma} H^{an} + F_{ac} \eta^{cn} \mathcal{G}_{gm} \eta^{ga} \right) \delta \bar{K}_n{}^m \Phi \\
&+ \frac{1}{2} \left(\frac{1}{2} \mathcal{G}_{mb} H^{nb} - \frac{1}{2} \mathcal{G}_{ma} H^{an} + F_{am} \eta^{nb} \mathcal{G}_{gb} \eta^{ga} \right) \delta \bar{K}_n{}^m \Phi \\
&+ \frac{1}{2} \left(\mathcal{H}_{cs} \eta^{sb} \mathcal{G}_{gb} \eta^{gn} \delta_m^c + \frac{1}{2} \mathcal{G}_{rc} \eta^{rn} \eta^{cb} F_{mb} - \frac{1}{2} \mathcal{G}_{rc} \eta^{rb} \eta^{cn} F_{mb} \right) \delta \bar{K}_n{}^m \Phi \\
&+ \frac{1}{2} \left(\mathcal{H}_{cs} \eta^{sn} \mathcal{G}_{gm} \eta^{gc} + \frac{1}{2} \mathcal{G}_{rc} \eta^{ra} \eta^{cn} F_{am} - \frac{1}{2} \mathcal{G}_{rc} \eta^{rn} \eta^{ca} F_{am} \right) \delta \bar{K}_n{}^m \Phi \\
&+ \frac{1}{2} \left(\mathcal{H}_{ms} \eta^{sb} \mathcal{G}_{gb} \eta^{gn} + \frac{1}{2} \mathcal{G}_{rm} \eta^{ra} \eta^{nb} F_{ab} - \frac{1}{2} \mathcal{G}_{rm} \eta^{rb} \eta^{na} F_{ab} \right) \delta \bar{K}_n{}^m \Phi \\
&+ \frac{1}{2} ((\mathcal{H}_{ab} H^{ae} \eta^{bf} + 2F_{ac} H^{ae} \eta^{cf}) + \mathcal{G}_{ca} \eta^{ab} (\mathcal{G}_{db} - 2\mathcal{G}_{bd}) \eta^{de} \eta^{cf}) \bar{K}_{ef} \Phi \\
&+ (\mathcal{G}_{ab} H^{ab} + F_{ab} \eta^{ac} \eta^{bd} \mathcal{G}_{cd}) \delta \Phi
\end{aligned}$$

This allows further simplifications, then including the variation of the volume form we have the final result,

$$\begin{aligned}
\delta (\bar{\mathcal{F}} \wedge {}^* \mathcal{F}) &= \frac{1}{2} (F_{dc} \eta^{cb} F_{eb} + 2F_{ad} H^{ab} \eta_{be} + (\mathcal{G}_{ae} - 2\mathcal{G}_{ea}) \mathcal{G}_{cd} \eta^{ca}) \delta \bar{K}^{de} \Phi \\
&+ H^{na} (2\mathcal{G}_{ma} - \mathcal{G}_{am}) \delta \bar{K}_n{}^m \Phi \\
&+ F_{ma} (\mathcal{G}^{na} - 2\mathcal{G}^{an}) \delta \bar{K}_n{}^m \Phi \\
&+ \frac{1}{2} ((\mathcal{H}_{ab} H^{ae} \eta^{bf} + 2F_{ac} H^{ae} \eta^{cf}) + \mathcal{G}_{ca} \eta^{ab} (\mathcal{G}_{db} - 2\mathcal{G}_{bd}) \eta^{de} \eta^{cf}) \bar{K}_{ef} \Phi \\
&+ (\mathcal{G}_{ab} H^{ab} + F_{ab} \eta^{ac} \eta^{bd} \mathcal{G}_{cd}) \delta \Phi
\end{aligned}$$

References

- [1] M. Born, A suggestion for unifying quantum theory and relativity, Proceedings of the Royal Society London A (1938), vol. 165, pp. 291–303, doi:10.1098/rspa.1938.0060
- [2] M. Born, Reciprocity Theory of Elementary Particles, Review of Modern Physics (1949) vol. 21, no. 3, pp. 463–473 doi:10.1103/RevModPhys.21.463
- [3] Ivanov, E.A. and Niederle, J., Gauge formulation of gravitation theories. I. The Poincaré, de Sitter, and conformal cases, Phys. Rev. D **25** 4 (1982) 976.
- [4] Ivanov, E.A. and Niederle, J., Gauge formulation of gravitation theories. II. The special conformal case, Phys. Rev. D **25** 4 (1982) 988.
- [5] Wehner, André. and James T. Wheeler, *Conformal actions in any dimension*, Nucl. Phys. B **557** (1999) 380-406, arXiv:hep-th/9812099.
- [6] James T. Wheeler, *General relativity as a biconformal gauge theory*, Nucl.Phys. B (2019) 56pp 114624.
- [7] Hazboun, Jeffrey S. and James T. Wheeler, *Time and dark matter from the conformal symmetries of Euclidean space*, Class. Quantum Gravity **31** (2014) 215001.
- [8] Lara B. Anderson, James T. Wheeler, *Quantum mechanics as a measurement theory on biconformal space*, Int. J. Geom. Meth. Mod. Phys. 3:315-340, 2006 DOI: 10.1142/S0219887806001168.
- [9] Spencer, Joseph Andrew and James T. Wheeler, *The existence of time*, International Journal of Geometric Methods in Modern Physics, Vol. 8 No. 2 (2011) 273-301. <http://arxiv.org/abs/0811.0112>
- [10] Hazboun, Jeffrey Shafiq, "Conformal Gravity and Time" (2014). All Graduate Theses and Dissertations. 3856. <https://digitalcommons.usu.edu/etd/3856>.
- [11] Jeffrey S Hazboun, *Constructing an Explicit AdS/CFT Correspondence with Cartan Geometry*, Nuclear Physics B, Volume 929, April 2018, Pages 254-265 DOI: 10.1016/j.nuclphysb.2018.02.006
- [12] M. J. Duff, *Duality rotations in string theory*, Nuclear Physics B Volume 335, Issue 3, 14 May 1990, Pages 610-620.
- [13] W. Siegel, *Manifest Duality in Low-Energy Superstrings*, arXiv:hep-th/9308133.
- [14] W. Siegel, *Superspace Duality in Low-Energy Superstrings*, Phys.Rev. D48 (1993) 2826-2837.

- [15] W. Siegel, *Two-Vierbein Formalism for String-Inspired Axionic Gravity*, Phys.Rev. D47 (1993) 5453-5459.
- [16] Hazboun, Jeffrey S. and Wheeler, James T., Proceedings of the Loops11: *Non-Perturbative/Background Independent Quantum Gravity* (Madrid, 2011), Volume 360, IOP Journal of Physics: Conference Series, http://iopscience.iop.org/1742-6596/360/1/012013/pdf/1742-6596_360_1_012013.pdf
- [17] Cho, W., Fernández-Melgarejo, J.J., Jeon, I. et al. Supersymmetric gauged double field theory: systematic derivation by virtue of twist. J. High Energ. Phys. 2015, 84 (2015). [https://doi.org/10.1007/JHEP08\(2015\)084](https://doi.org/10.1007/JHEP08(2015)084)
- [18] Özer, A.Ç. *Duality twisted reductions of Double Field Theory of Type II strings*. J. High Energ. Phys. 2017, 44 (2017). [https://doi.org/10.1007/JHEP09\(2017\)044](https://doi.org/10.1007/JHEP09(2017)044)
- [19] Graña, M., Marques, D. *Gauged double field theory*. J. High Energ. Phys. 2012, 20 (2012). [https://doi.org/10.1007/JHEP04\(2012\)020](https://doi.org/10.1007/JHEP04(2012)020)
- [20] Anderson, Lara B. and James T. Wheeler, *Biconformal supergravity and the AdS/CFT conjecture*, J. T., Nucl. Phys. B **686** (2004) 285-309, arXiv:hep-th/0309111.
- [21] Wehner, Andre and James T. Wheeler, *Biconformal Matter Actions*, (2000) arXiv:hep-th/0001061v2
- [22] Lovelady, B. C., and Wheeler, J. T. (2016). *Dynamical spacetime symmetry*. Phys. Rev. D, 93(8), 085002. <http://doi.org/10.1103/PhysRevD.93.085002>
- [23] Wheeler, J.T., *New conformal gauging and the electromagnetic theory of Weyl*, J. Math. Phys. **39** (1) (Jan., 1998) 299-328. [arXiv:hep-th/9706214]
- [24] Wheeler, J. T., *Weyl geometry*, Gen Relativ Gravit (2018) 50:80 DOI: 10.1007/s10714-018-2401-5
- [25] Bach, R. Zur Weylschen Relativitätstheorie und der Weylschen Erweiterung des Krümmungstensorbegriffs. Math Z 9, 110–135 (1921). <https://doi.org/10.1007/BF01378338>
- [26] Mannheim, P.D. Making the Case for Conformal Gravity. Found Phys 42, 388–420 (2012). <https://doi.org/10.1007/s10701-011-9608-6>
- [27] P. D. Mannheim, Prog. Part. Nucl. Phys. 56, 340 (2006).
- [28] É. É. Flanagan, Phys. Rev. D 74, 023002 (2006).

- [29] Youngsub Yoon Phys. Rev. D 88, 25 July 2013 , 027504
- [30] Comment on “Problems with Mannheim’s conformal gravity program” Philip D. Mannheim Phys. Rev. D 93, 068501 (2016)
- [31] James T. Wheeler, Phys. Rev. D 90, 025027 (2014) DOI: 10.1103/PhysRevD.90.025027
- [32] James Thomas Wheeler and Andre Wehner. "Biconformal matter actions" International Journal of Pure and Applied Mathematics Vol. 30 Iss. No. 2 (2006) p. 207 - 224 ISSN: ISSN 1311-8080; arXiv:hep-th/0001061v2.
- [33] Robert Brandenberger, Renato Costa Santos, Guilherme Franzmann, Amanda Weltman, *T-dual cosmological solutions in double field theory*, Phys Rev D.99.023531, January 29, 2019, DOI: 10.1103
- [34] David S. Berman, Chris D. A. Blair, Emanuel Malek, Malcolm J. Perry, *The $O_{-}\{D,D\}$ Geometry of String Theory*, arXiv:1303.6727v2 [hep-th], DOI: 10.1142/S0217751X14500808.
- [35] Gerardo Aldazabal Diego Marqués, and Carmen Núñez, *Double Field Theory: A Pedagogical Review*, Class. Quant. Grav. 30 (2013) 163001 DOI: 10.1088/0264-9381/30/16/163001
- [36] Lara B. Anderson and James T. Wheeler, *Yang-Mills gravity in biconformal space*, Class.Quant.Grav. 24 (2007) 475-496 DOI: 10.1088/0264-9381/24/2/012; arXiv:hep-th/0412293.