

A NOTE ON GLOBAL SMOOTHABILITY OF SMALL L^p -CONNECTION WITH PRESCRIBED CURVATURE

SIRAN LI

ABSTRACT. We show that if Ω is a connection 1-form on a vector bundle η over a closed n -dimensional Riemannian manifold \mathcal{M} with L^p -regularity ($p > n$) and smooth curvature 2-form \mathcal{F} , then it can be approximated in the L^p -norm by smooth connections of the same curvature, provided that $\|\Omega\|_{L^p(\mathcal{M})}$ is smaller than a uniform constant depending only on p and \mathcal{M} .

1. INTRODUCTION

In this note, we prove a result concerning the structure of the space of L^p -affine connections on vector bundles over an n -dimensional Riemannian manifold, where $p > n$. The set up of our problem closely follows the seminal work [16] by Uhlenbeck; nevertheless, in our note less regularity assumption will be imposed.

Let η be a vector bundle with structure group G over a Riemannian n -dimensional manifold \mathcal{M} . Let $\mathfrak{A}(\mathcal{M})$ be the affine space of smooth connections on \mathcal{M} , i.e.,

$$\mathfrak{A}(\mathcal{M}) = \left\{ D_0 + \Omega : \Omega \in C^\infty \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^1 T^* \mathcal{M} \right) \right\} \quad (1.1)$$

where $D_0 \in \mathfrak{A}(\mathcal{M})$ is a base connection and \mathfrak{g} is the Lie algebra of G . In other words, $\mathfrak{A}(\mathcal{M})$ is the affine space modelled over \mathfrak{g} -valued smooth 1-forms. Each fibre of the adjoint bundle $\text{Ad } \eta$ is isomorphic of \mathfrak{g} . We also denote by $\mathfrak{A}_0^p(\mathcal{M})$ the connection 1-forms of Sobolev L^p -regularity:

$$\mathfrak{A}_0^p(\mathcal{M}) = \left\{ D_0 + \Omega : \Omega \in L^p \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^1 T^* \mathcal{M} \right) \right\}. \quad (1.2)$$

This definition is independent of the choice of D_0 . The curvature 2-form of a connection $\Omega \in \mathfrak{A}(\mathcal{M})$ is given by

$$F_\Omega := d\Omega + \Omega \wedge \Omega \in C^\infty \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right). \quad (1.3)$$

Note that the definition of curvature in Eq. (1.3) extends naturally to $\Omega \in \mathfrak{A}_0^p(\mathcal{M})$ with $p \geq 2$: $F_\Omega \in [W^{-1,p} + L^{p/2}] \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right)$, which is well defined as a distribution. We shall refer to Eq. (1.3) as the *gauge equation* (for F_Ω prescribed).

Our main result of this note is as follows:

Theorem 1.1. *Let η be a vector bundle with structure group G over an n -dimensional closed Riemannian manifold \mathcal{M} ; $n \geq 2$. Consider the indices $p > n$, $q \in [\frac{p}{2}, \infty[$, and $s \in [0, \infty[$. Let $\mathcal{F} \in W^{s,q} \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right)$ be a prescribed 2-form. Then, there exists a constant $\kappa_0 > 0$*

Date: January 11, 2022.

2010 Mathematics Subject Classification. 35F50, 53C07.

Key words and phrases. Connection form; prescribed curvature; smooth approximation.

depending only on p and \mathcal{M} such that the following space is smoothable:

$$\mathfrak{A}_0^p(\mathcal{M}, \kappa_0, \mathcal{F}) := \left\{ \Omega \in \mathfrak{A}_0^p(\mathcal{M}) : \|\Omega\|_{L^p(\mathcal{M})} < \kappa_0 \text{ and } F_\Omega = \mathcal{F} \right\}.$$

That is, there exists a family of smooth connections $\{\Omega^\epsilon\} \subset \mathfrak{A}(\mathcal{M})$ such that $\Omega^\epsilon \rightarrow \Omega$ in $L^p(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^1 T^* \mathcal{M})$ as $\epsilon \rightarrow 0$ and $F_{\Omega^\epsilon} \equiv \mathcal{F}$ for each ϵ .

The argument in our note essentially follows S. Mardare [8, 9] on solubility of the Pfaff equations and extension of the fundamental theorem of surface theory to $W_{\text{loc}}^{1,p}$ -metrics for $p > 2$.

Our usage of Sobolev norms and other analytic/geometric tools on Riemannian manifolds and vector bundles is standard; cf. [6, 7, 12].

2. THE PROOF

This section is devoted to the proof of Theorem 1.1. We shall prove the following variant:

Theorem 2.1. *Let η be a vector bundle with structure group G over a Riemannian n -dimensional manifold \mathcal{M} ; $n \geq 2$. Assume that $\Omega \in \mathfrak{A}_0^p$ for some $p > n$ and that curvature \mathcal{F} of Ω is in the regularity class $W^{s,q}$, where $s \geq 0$ and $\frac{p}{2} \leq q < \infty$. Then, for every smooth, bounded subdomain $U \Subset \mathcal{M}$, there is a uniform constant $\kappa_0 > 0$ depending only on U and p , such that if*

$$\|\Omega\|_{L^p(U)} < \kappa_0,$$

then one can find a one-parameter family $\{\Omega^\epsilon\}_{\epsilon \in]0,1]} \subset W^{1+s,q}(U; \text{Ad } \eta \otimes \bigwedge^1 T^ \mathcal{M})$ such that $\Omega^\epsilon \rightarrow \Omega$ on U in the L^p -topology, and that the curvature of Ω^ϵ coincides with \mathcal{F} for each $\epsilon \in]0,1]$.*

Note that given the smallness condition $\|\Omega\|_{L^p(U)} < \kappa_0$, our smoothability Theorem 2.1 does not require any assumption on the topology of U . Also, the proof easily carries over to closed manifolds (compact and without boundary).

Proof. We divide our arguments into nine steps. In Steps 1–8 we prove for the case $\mathcal{M} = \mathbb{R}^n$, and in Step 9 we generalise to Riemannian manifolds.

Step 1. We first establish

Claim A. Let $U \subset \mathcal{M}$ be a smooth subdomain. We can find a scalar function $\varphi \in W^{1,p}(U; \text{Ad } \eta)$ and a 2-form $\psi \in W^{1,p}(U; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M})$ such that $\mathfrak{n}\psi = 0$ and $\mathfrak{n}d\psi = 0$ on ∂U , and that

$$\Omega = d\varphi + d^*\psi \quad \text{in } U.$$

Here and throughout, \mathfrak{t} and \mathfrak{n} denote the tangential and normal trace operators, respectively.

Proof of Claim A. This follows from the classical Hodge decomposition theorem on manifolds-with-boundary. We first solve for ψ from the boundary value problem

$$\begin{cases} \Delta\psi = \mathcal{F} - \Omega \wedge \Omega & \text{in } U, \\ \mathfrak{n}\psi = 0 \quad \text{and} \quad \mathfrak{n}d\psi = 0 & \text{on } \partial U, \end{cases} \quad (2.1)$$

where $\Delta = dd^* + d^*d$. For $p > n$, the Sobolev embedding theorem gives us

$$W_{\text{loc}}^{1,p'}(\mathcal{M}) \hookrightarrow L_{\text{loc}}^{\frac{p}{\frac{n-1}{n}, p-1}}(\mathcal{M}) \hookrightarrow L_{\text{loc}}^{\frac{p}{p-2}}(\mathcal{M});$$

hence, by duality,

$$L_{\text{loc}}^{\frac{p}{2}}(\mathcal{M}) = L_{\text{loc}}^{\left(\frac{p}{p-2}\right)'}(\mathcal{M}) \hookrightarrow W_{\text{loc}}^{-1,p}(\mathcal{M}).$$

Then we can solve the boundary value problem (2.1) from the standard elliptic theory to get a solution ψ in $W^{1,p}$ on U , as Eq. (2.1) is elliptic in the sense of Agmon–Douglis–Nirenberg [1, 2] (or Šapiro–Lopatinskiĭ). See Schwarz [12] for details.

Moreover, since

$$d(\mathcal{F} - \Omega \wedge \Omega) = dd\Omega = 0$$

and that d commutes with Δ , we have $\Delta d\psi = 0$ in U . This together with the boundary condition shows that ψ is closed on U . It then follows from Eq. (2.1) that

$$d(\Omega - d^*\psi) = d\Omega - \Delta\psi = d\Omega + \Omega \wedge \Omega - \mathcal{F} = 0 \quad \text{in } U.$$

Finally, we can find $\phi \in W^{1,p}(U; \text{Ad } \eta)$ such that $d\phi = \Omega - d^*\psi$ on U . This is always possible as we do not impose any boundary conditions on ϕ . \square

Step 2. Next, take $\{\phi^\epsilon\} \subset C^\infty(U; \text{Ad } \eta)$ such that

$$\phi^\epsilon \longrightarrow \phi \quad \text{in } W^{1,p}(U; \text{Ad } \eta). \quad (2.2)$$

This can be obtained, e.g., via a standard mollification process (there is no obstructions on density of Sobolev spaces here, as ϕ is a scalar function). The above convergence is only required to take place in the interior of U .

Step 3. To proceed, let us find ψ^ϵ , the smooth approximates of ψ , from the following nonlinear elliptic system:

$$\begin{cases} \Delta\psi^\epsilon = \mathcal{F} - (d\phi^\epsilon + d^*\psi^\epsilon) \wedge (d\phi^\epsilon + d^*\psi^\epsilon) & \text{in } U, \\ \mathbf{n}\psi^\epsilon = 0 \quad \text{and} \quad \mathbf{n}d\psi^\epsilon = 0 & \text{on } \partial U \end{cases} \quad (2.3)$$

This system can be solved from a standard Schauder’s fixed-point argument. Denote by Δ^{-1} the solution operator for Laplace–Beltrami subject to the same boundary conditions. Consider the operator for each fixed $\epsilon \in]0, 1[$:

$$\begin{aligned} \mathcal{T}_\epsilon : \quad W^{1,p} \left(U; \mathfrak{g} \otimes \bigwedge^2 T^* \mathcal{M} \right) &\longrightarrow W^{1,p} \left(U; \mathfrak{g} \otimes \bigwedge^2 T^* \mathcal{M} \right) \\ \zeta &\longmapsto \Delta^{-1} \left\{ \mathcal{F} - (d\phi^\epsilon + d^*\zeta) \wedge (d\phi^\epsilon + d^*\zeta) \right\}. \end{aligned} \quad (2.4)$$

Claim B. For each $\epsilon \in [0, 1[$, the operator \mathcal{T}_ϵ has a fixed point ψ^ϵ .

Proof of Claim B. First, we observe that \mathcal{T}_ϵ indeed maps into the $W^{1,p}$ -space. This is because for $\zeta \in W^{1,p} \left(U; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right)$, the same Sobolev embedding argument as above yields that

$$\mathcal{F} - (d\phi^\epsilon + d^*\zeta) \wedge (d\phi^\epsilon + d^*\zeta) \in W^{-1,p} \left(U; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right).$$

Hence $\mathcal{T}_\epsilon(\zeta) \in W^{1,p} \left(U; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right)$ by standard elliptic estimates.

In addition, we can easily bound

$$\|\mathcal{T}_\epsilon \zeta\|_{W^{1,p}(U)} \leq C \|\mathcal{F} - (d\phi^\epsilon + d^*\zeta) \wedge (d\phi^\epsilon + d^*\zeta)\|_{W^{-1,p}(U)}$$

$$\begin{aligned}
&\leq C \left(\|\mathcal{F}\|_{W^{-1,p}(U)} + \|(d\phi^\epsilon + d^*\zeta) \wedge (d\phi^\epsilon + d^*\zeta)\|_{L^{\frac{np}{n+p}}(U)} \right) \\
&\leq C \left(\|\mathcal{F}\|_{W^{-1,p}(U)} + \|d\phi^\epsilon + d^*\zeta\|_{L^{\frac{2np}{n+p}}(U)}^2 \right) \\
&\leq C \left(\|\mathcal{F}\|_{W^{-1,p}(U)} + \|d\phi^\epsilon + d^*\zeta\|_{L^p(U)}^2 \right) \\
&\leq C \left(\|\mathcal{F}\|_{W^{-1,p}(U)} + \|\phi^\epsilon\|_{W^{1,p}(U)}^2 + \|\zeta\|_{W^{1,p}(U)}^2 \right),
\end{aligned}$$

where C depends only on U and p . The first line follows from standard elliptic estimates, the second line holds by the Sobolev embedding

$$W_{\text{loc}}^{1, \frac{np}{n+p}} \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right) \hookrightarrow L_{\text{loc}}^p \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right),$$

the third line holds by Hölder, the fourth line holds by $p \geq n$, and the last line is trivial.

As $\|\phi^\epsilon\|_{W^{1,p}(U)}$ is uniformly bounded due to Eq. (2.2), we readily conclude that \mathcal{T}_ϵ is a bounded operator for each ϵ . This follows from the estimate

$$\|\mathcal{T}_\epsilon \zeta\|_{W^{1,p}(U)} \leq C(U, p) \left(\|\mathcal{F}\|_{W^{-1,p}(U)} + \|\phi\|_{W^{1,p}(U)}^2 + \|\zeta\|_{W^{1,p}(U)}^2 + \mathfrak{o}(\epsilon) \right) \quad (2.5)$$

with $\mathfrak{o}(\epsilon) \rightarrow 0$ as $\epsilon \rightarrow 0$.

Note that by assumption we get $\mathcal{F} \in W_{\text{loc}}^{s,q} \left(\mathcal{M}; \mathfrak{g} \otimes \bigwedge^2 T^* \mathcal{M} \right)$ for $s \geq 0$ and $q \geq \frac{p}{2}$. But one has the compact embedding $W_{\text{loc}}^{s,q} \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right) \hookrightarrow W_{\text{loc}}^{-1,p} \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right)$ for $p > n$. So Eq. (2.5) indeed gives us the boundedness of \mathcal{T}_ϵ .

Next we check that \mathcal{T}_ϵ is a compact operator. Take an arbitrary sequence $\{\zeta_j \equiv \zeta_j^\epsilon\}_{j=1,2,\dots}$ which converges weakly in the $W_{\text{loc}}^{1,p}$ -topology to $\bar{\zeta} \equiv \bar{\zeta}^\epsilon$. Then

$$\begin{aligned}
&\left\{ \mathcal{F} - (d\phi^\epsilon + d^*\zeta_j) \wedge (d\phi^\epsilon + d^*\zeta_j) \right\} - \left\{ \mathcal{F} - (d\phi^\epsilon + d^*\bar{\zeta}) \wedge (d\phi^\epsilon + d^*\bar{\zeta}) \right\} \\
&= d^* (\bar{\zeta} - \zeta_j) \wedge (d\phi^\epsilon + d^*\bar{\zeta}) + (d\phi^\epsilon + d^*\zeta_j) \wedge d^* (\bar{\zeta} - \zeta_j) \\
&\rightarrow 0 \quad \text{in } L_{\text{loc}}^{\frac{p}{2}-\delta} \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right),
\end{aligned}$$

thanks to the Hölder's inequality, the weak convergence $\zeta_j \rightharpoonup \bar{\zeta}$ in $W_{\text{loc}}^{1,p} \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right)$, and the uniform boundedness of $\{d\phi^\epsilon + d^*\bar{\zeta}\}$ and $\{d\phi^\epsilon + d^*\zeta_j\}$ (in fact, independent of both j and ϵ) in $L_{\text{loc}}^p \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^1 T^* \mathcal{M} \right)$. As we have the strict inequality $p > n$, we can choose a $\delta > 0$ such that the compact embedding holds:

$$L_{\text{loc}}^{\frac{p}{2}-\delta} \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right) \hookrightarrow W_{\text{loc}}^{-1,p} \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right) \quad (2.6)$$

Therefore, one may select a subsequence $\{\zeta_{j_\nu} \equiv \zeta_{j_\nu}^\epsilon\}_{\nu=1,2,\dots}$ such that $\zeta_{j_\nu} \rightarrow \bar{\zeta}$ strongly in $W_{\text{loc}}^{-1,p} \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right)$ as $\nu \rightarrow \infty$. This together with the standard elliptic estimates applied to $(\mathcal{T}_\epsilon \zeta_{j_\nu} - \mathcal{T}_\epsilon \bar{\zeta})$ via Eq. (2.3) gives us

$$\lim_{\nu \rightarrow \infty} \|\mathcal{T}_\epsilon \zeta_{j_\nu} - \mathcal{T}_\epsilon \bar{\zeta}\|_{W^{1,p}(U)} = 0.$$

We can now conclude that \mathcal{T}_ϵ is compact.

The existence (for each $\epsilon \in]0, 1[$) of a solution $\psi^\epsilon \in W^{1,p} \left(U; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right)$ for Eq. (2.1) follows now from Schauder's fixed point theorem. \square

Step 4. Two remarks are in order concerning Claim B:

- (1) The solution operator \mathcal{T}_ϵ in Eq. (2.4) is uniformly bounded in ϵ . That is, there is a uniform constant K_0 independent of ϵ such that

$$\sup_{\epsilon \in]0, 1[} \|\mathcal{T}_\epsilon\| \leq K_0, \quad (2.7)$$

where $\|\bullet\|$ is the $W^{1,p} \rightarrow W^{1,p}$ operator norm. This indeed follows from Eq. (2.5); K_0 can be chosen to depend only on U , p , $\|\mathcal{F}\|_{W^{-1,p}(U)}$, and $\|\Omega\|_{L^p(U)}$ (in view of Step 1 of the same proof).

- (2) Given a general L^p -connection Ω , we are unable to show the uniqueness of the solution ψ^ϵ in Eq. (2.1). In fact, this remains the case even if the connection is smooth. By now we can only prove uniqueness for the case of small data.

Claim C. For each $\epsilon \in [0, 1[$, the operator \mathcal{T}_ϵ has a fixed point ψ^ϵ , provided that

$$\|\mathcal{F}\|_{W^{-1,p}(U)} + \|\Omega\|_{L^p(U)} \leq \kappa_0 \quad (2.8)$$

for a small uniform number κ_0 depending only on p and U , and that the convergence of ϕ^ϵ to ϕ in Eq. (2.2) is rapid enough.

We remark the smallness of $\|\mathcal{F}\|_{W^{-1,p}(U)}$ follows from the smallness of $\|\Omega\|_{L^p(U)}$, thanks to the gauge equation $d\Omega + \Omega \wedge \Omega = \mathcal{F}$. Hence, the above smallness hypothesis (2.8) is equivalent to the condition in the statement of Theorem 2.1. Indeed, $\|\Omega\|_{L^p(U)} \leq \kappa$ implies that

$$\|d\Omega\|_{W^{-1,p}(U)} \leq \kappa$$

and that

$$\|\Omega \wedge \Omega\|_{W^{-1,p}(U)} \leq C(p, U) \|\Omega \wedge \Omega\|_{L^{\frac{p}{2}}(U)} \leq C(p, U) \|\Omega\|_{L^p(\Omega)}^2 \leq C(p, U) \kappa^2,$$

where $C(p, U)$ is the Sobolev constant corresponding to

$$L^{\frac{p}{2}} \left(U; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right) \hookrightarrow W^{-1,p} \left(U; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right).$$

This compact embedding holds when $p > n$; see the proof of Claim A in Step 1.

Proof of Claim C. Denote by $K_1 = C(U, p)$ the uniform constant in Eq. (2.5) in the proof of Claim B above. First we fix $\kappa_1 > 0$ such that

$$K_1 \cdot \kappa_1 \leq \frac{1}{3}.$$

Then, choose $\kappa_0 \in]0, 1[$ such that

$$2K_1 \cdot \kappa_0 \leq \frac{1}{3} \cdot \kappa_1.$$

Finally, by passing to a subsequence in Eq. (2.2) if necessary, we assume that

$$\mathfrak{o}(\epsilon) = \|\phi^\epsilon - \phi\|_{W^{1,p}(U)} \leq \kappa_0 \quad \text{for all } \epsilon \in]0, 1[.$$

In view of the Hodge decomposition in Claim A, we can infer from Eq. (2.5) that with

$$\mathfrak{C}_{\kappa_1} := \left\{ \zeta \in W^{1,p} \left(U; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right) : \|\zeta\|_{W^{1,p}(U)} \leq \kappa_1 \right\},$$

the restricted operator

$$\mathcal{T}_\epsilon|_{\mathfrak{C}_{\kappa_1}} : \mathfrak{C}_{\kappa_1} \longrightarrow \mathfrak{C}_{\kappa_1}$$

is coercive under the smallness condition (2.8). That is, in comparison with Eq. (2.7), we have

$$\sup_{\epsilon \in]0,1]} \left\| \mathcal{T}_\epsilon|_{\mathfrak{C}_{\kappa_1}} \right\| < 1 \quad (2.9)$$

whenever Eq. (2.8) is valid. Our choice of κ_1 (and hence of κ_0) again depends only on p and U .

The uniqueness of ψ^ϵ follows from Eq. (2.9) and the Banach fixed point theorem. \square

Step 5. In Steps 3 and 4 above we have proved the solubility of the equation (2.3) for ψ^ϵ , and the uniqueness of solution under the smallness assumption in Eq. (2.8). The regularity of ψ^ϵ follows directly from the standard elliptic (Calderón–Zygmund or $W^{2,p}$ -) theory.

Claim D. Assume that the prescribed curvature 2-form \mathcal{F} has $W^{s,q}$ -regularity for $s \geq 0$ and $q \geq \frac{p}{2} > \frac{n}{2}$. Then we have

$$\psi^\epsilon \in W_{\text{loc}}^{s+2,q} \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right), \quad (2.10)$$

with $\|\psi^\epsilon\|_{W^{s+2,q}(U)} \leq K_2$ depending only on U and p .

Proof of Claim D. Starting with a solution ψ^ϵ for Eq. (2.3), we find that the right-hand side

$$\mathcal{F} - (d\phi^\epsilon + d^* \psi^\epsilon) \wedge (d\phi^\epsilon + d^* \psi^\epsilon)$$

lies in

$$\left[W_{\text{loc}}^{s,q} + L_{\text{loc}}^{\frac{p}{2}} \right] \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right) \hookrightarrow L_{\text{loc}}^{\frac{p}{2}} \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right)$$

for $s \geq 0$ and $q \geq p/2$. Thus $\psi^\epsilon \in W_{\text{loc}}^{2,\frac{p}{2}} \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right)$ by the standard elliptic theory. Then, by Sobolev embedding one may further infer that

$$\mathcal{F} - (d\phi^\epsilon + d^* \psi^\epsilon) \wedge (d\phi^\epsilon + d^* \psi^\epsilon) \in \begin{cases} \left[W_{\text{loc}}^{s,q} + L_{\text{loc}}^{\frac{np}{4n-2p}} \right] \left(\mathcal{M}; \mathfrak{g} \otimes \bigwedge^2 T^* \mathcal{M} \right) & \text{in } n < p \leq 2n, \\ \left[W_{\text{loc}}^{s,q} + L_{\text{loc}}^\infty \right] \left(\mathcal{M}; \mathfrak{g} \otimes \bigwedge^2 T^* \mathcal{M} \right) & \text{in } p > 2n; \end{cases}$$

so on and so forth.

Repeating the above procedure for finitely many steps, one arrives at

$$\mathcal{F} - (d\phi^\epsilon + d^* \psi^\epsilon) \wedge (d\phi^\epsilon + d^* \psi^\epsilon) \in W_{\text{loc}}^{s,q} \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right),$$

namely that the right-hand side is as regular as the prescribed curvature \mathcal{F} . We now conclude from Eq. (2.3) that $\psi^\epsilon \in W_{\text{loc}}^{s+2,q} \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right)$. The uniform $W_{\text{loc}}^{s+2,q}$ -boundedness of ψ^ϵ follows from explicit estimates in the standard Calderón–Zygmund theory. \square

Step 6. As a consequence of Step 5 above, we deduce the following:

Claim E. Assume that the prescribed curvature 2-form \mathcal{F} has $W^{s,q}$ -regularity for $s \geq 0$ and $q \geq \frac{p}{2} > \frac{n}{2}$. Define

$$\Omega^\epsilon := d\phi^\epsilon + d^*\psi^\epsilon. \quad (2.11)$$

Then, under the smallness condition (2.8) and modulo subsequences, we have the strong convergence

$$\Omega^\epsilon \longrightarrow \Omega \quad \text{in } L^p \left(U; \text{Ad } \eta \otimes \bigwedge^1 T^* \mathcal{M} \right),$$

where Ω is the connection in the statement of Theorem 2.1.

Proof of Claim E. By Claim D, $\{d\psi^\epsilon\}$ is uniformly bounded in $W^{1+s,q} \left(U; \text{Ad } \eta \otimes \bigwedge^1 T^* \mathcal{M} \right)$. We have the compact Sobolev embedding $W_{\text{loc}}^{1+s,q}(\mathcal{M}) \hookrightarrow L_{\text{loc}}^p(\mathcal{M})$ for $q \geq \frac{p}{2} > \frac{n}{2}$. In addition, under the smallness condition (2.8) we have, thanks to Claim C, the unique solution ψ^ϵ for each $\epsilon \in]0, 1]$. Sending further $\epsilon \rightarrow 0$ and noting Eq. (2.7), we deduce that a weak $W_{\text{loc}}^{1+s,q}$ -limit of $d\psi^\epsilon$ modulo subsequences — hence the strong L_{loc}^p -limit, after passing to a further subsequence — coincides with $d\psi$. This together with the convergence of ϕ^ϵ in Eq. (2.2) implies the strong L^p -convergence of Ω^ϵ over $U \subset \mathcal{M}$. \square

Step 7. Define μ_U to be the smallest positive number such that the following holds:

$$\|\Xi\|_{L^2(U)} \leq \mu_U \|\nabla \Xi\|_{L^2(U)} \quad \forall \Xi \in W^{1,2} \left(U; \text{Ad } \eta \otimes \bigwedge^3 T^* \mathcal{M} \right) \text{ with vanishing normal trace.} \quad (2.12)$$

Let us prove the following:

Claim F. In the setting of Claim E and under the smallness condition (2.8) as in Claim C, ψ^ϵ is closed (*i.e.*, $d\psi^\epsilon = 0$) in U for each $\epsilon > 0$.

Proof of Claim F. Let us take d to both sides of Eq. (2.1) and write $\Omega^\epsilon := d\phi^\epsilon + d^*\psi^\epsilon$ as in Step 6 above. Since d commutes with Δ , we get

$$\begin{cases} \Delta d\psi^\epsilon = \Omega^\epsilon \wedge d^*d\psi^\epsilon - d^*d\psi^\epsilon \wedge \Omega^\epsilon & \text{in } U, \\ \mathbf{n}(d\psi^\epsilon) = 0 & \text{on } \partial U. \end{cases} \quad (2.13)$$

Since $\psi^\epsilon \in W_{\text{loc}}^{s+2,q} \left(\mathcal{M}; \text{Ad } \eta \otimes \bigwedge^2 T^* \mathcal{M} \right)$ for $s \geq 0$, the above boundary value problem (2.13) for the variable $d\psi^\epsilon$ is well defined, and one can apply the Stokes' or the Gauss–Green theorem to deduce that

$$\langle \langle d\psi^\epsilon, \Delta d\psi^\epsilon \rangle \rangle_U = \|\nabla d\psi^\epsilon\|_{L^2(U)}^2 + \int_{\partial U} \mathbf{t}(d^*d\psi^\epsilon) \wedge \star \mathbf{n}(d\psi^\epsilon) = \|\nabla d\psi^\epsilon\|_{L^2(U)}^2, \quad (2.14)$$

where the final equality holds by the boundary condition in Eq. (2.13). Here and hereafter,

$$\langle \langle \alpha, \beta \rangle \rangle_U := \int_U \alpha \wedge \star \beta$$

with \star being the Hodge star operator associated to the Riemannian volume measure $U \subset \mathcal{M}$. The wedge product \wedge is intertwined with the product on the Lie algebra \mathfrak{g} , if α and β are \mathfrak{g} -valued differential forms.

On the other hand, Eq. (2.13) and Hölder's inequality give us

$$\langle \langle d\psi^\epsilon, \Delta d\psi^\epsilon \rangle \rangle_U \leq C_1 \|\Omega^\epsilon\|_{L^p(U)} \|d\psi^\epsilon\|_{L^{\frac{2p}{p-2}}(U)} \|\nabla d\psi^\epsilon\|_{L^2(U)}$$

where C_1 depends only on p and U . We have the compact embedding

$$W^{1,2} \left(U; \text{Ad } \eta \otimes \bigwedge^3 T^* \mathcal{M} \right) \hookrightarrow L^{\frac{2p}{p-2}} \left(U; \text{Ad } \eta \otimes \bigwedge^3 T^* \mathcal{M} \right);$$

thus for another constant $C_2 = C(p, U)$ it holds that

$$\langle \langle d\psi^\epsilon, \Delta d\psi^\epsilon \rangle \rangle_U \leq C_2 \|\Omega^\epsilon\|_{L^p(U)} \left(\|\nabla d\psi^\epsilon\|_{L^2(U)} + \|d\psi^\epsilon\|_{L^2(U)} \right)^2. \quad (2.15)$$

Moreover, by Claim E in Step 6 of the same proof, we have

$$\|\Omega^\epsilon\|_{L^p(U)} \leq \|\Omega\|_{L^p(U)} + \mathfrak{o}(\epsilon).$$

Putting the above estimate together and using Cauchy–Schwarz, one obtains

$$\left[1 - \frac{3}{2} C_2 \left(\|\Omega\|_{L^p(U)} + \mathfrak{o}(\epsilon) \right) \right] \|\nabla d\psi^\epsilon\|_{L^2(U)}^2 \leq \frac{3}{2} C_2 \left(\|\Omega\|_{L^p(U)} + \mathfrak{o}(\epsilon) \right) \|d\psi^\epsilon\|_{L^2(U)}^2,$$

By passing to a further subsequence of $\{\phi^\epsilon\}$ if necessary, we can take $\mathfrak{o}(\epsilon)$ to be as small as we would like. So, the smallness condition (2.8) in Claim C is tantamount to assuming that

$$\frac{3}{2} C_2 \left(\|\Omega\|_{L^p(U)} + \mathfrak{o}(\epsilon) \right) \leq \kappa'_0$$

is arbitrarily small, which reduces the previous estimate to

$$\|\nabla d\psi^\epsilon\|_{L^2(U)}^2 \leq \frac{\kappa'_0}{1 - \kappa'_0} \|d\psi^\epsilon\|_{L^2(U)}^2. \quad (2.16)$$

Therefore, by further shrinking κ_0 (hence κ'_0) if necessary such that

$$\mu_U \cdot \frac{\kappa'_0}{1 - \kappa'_0} < 1, \quad (2.17)$$

where μ_U is the Poincaré constant, the Poincaré Inequality (2.12) implies that $\|\nabla d\psi^\epsilon\|_{L^2(U)} = 0$. Thus $d\psi^\epsilon$ equals to a constant almost everywhere. Under the boundary condition $\mathfrak{n}(d\psi^\epsilon) = 0$ on ∂U , one immediately concludes that ψ^ϵ is closed for each ϵ on U . \square

Step 8. Now we are ready to conclude the proof for the case $\mathcal{M} = \text{Euclidean space}$.

Claim F in Step 7 establishes that $d\psi^\epsilon \equiv 0$ in U . So Eq. (2.1) can be recast into

$$\Delta \phi^\epsilon = dd^* \psi^\epsilon \equiv d\Omega^\epsilon = \mathcal{F} - \Omega^\epsilon \wedge \Omega^\epsilon,$$

with $\Omega^\epsilon := d\phi^\epsilon + d^* \psi^\epsilon$ as before. That is, $\{\Omega^\epsilon\}$ satisfies the gauge equation on U . By the convergence proved in Claim E, we conclude that $\{\Omega^\epsilon\}$ is indeed a one-parameter family of smooth approximation for Ω .

Step 9. Finally, let \mathcal{M} be a Riemannian manifold. The arguments in Steps 1–6 carry over verbatim, and once Claim F in Step 7 gets established, we can conclude the proof as in Step 8.

In Step 7, Eqs. (2.14) and (2.15) can be replaced respectively by

$$\langle \langle d\psi^\epsilon, \Delta d\psi^\epsilon \rangle \rangle_U = \|d^* d\psi^\epsilon\|_{L^2(U)}^2$$

and

$$\langle d\psi^\epsilon, \Delta d\psi^\epsilon \rangle_U \leq C_3 \|\Omega^\epsilon\|_{L^p(U)} \|d\psi^\epsilon\|_{W^{1,2}(U)}^2.$$

Here C_3 again depends on p and U . On the other hand, the following version of the Gaffney's inequality holds:

$$\|d\psi^\epsilon\|_{W^{1,2}(U)} \leq C_4 \|d^* d\psi^\epsilon\|_{L^2(U)},$$

where C_4 depends on the geometry of $U \subset \mathcal{M}$. This holds by the boundary condition $\mathbf{n}(d\psi^\epsilon) = 0$, as well as the fact that $d\psi^\epsilon$ is L^2 -orthogonal to any harmonic field \mathcal{H} on U with the Neumann boundary data $\mathbf{n}\mathcal{H} = 0$ on ∂U (which follows easily from the Stokes'/Gauss–Green's theorem). See Schwarz [12], p.88, Lemma 2.4.10 (iii). Thus, for κ_0 in Eq. (2.8) chosen sufficiently small, we conclude as in Step 7 that $d^* d\psi^\epsilon \equiv 0$, and then deduce from the Gaffney's inequality that $d\psi^\epsilon \equiv \text{constant}$. Hence, by the boundary condition $\mathbf{n}(d\psi^\epsilon) = 0$, we have $d\psi^\epsilon \equiv 0$. \square

A straightforward adaption of the proof of Theorem 2.1 gives us the following:

Corollary 2.2. *Let η be a vector bundle with structure group G over a Riemannian n -dimensional manifold \mathcal{M} , $n \geq 2$. Assume that $\Omega \in \mathfrak{A}_0^p$ for some $p > n$ and that curvature \mathcal{F} of Ω is in the regularity class $W^{s,q}$, where $s \geq 0$ and $\frac{p}{2} \leq q < \infty$. Then, for every point $x \in \mathcal{M}$, we can find an open neighbourhood $U \subset \mathcal{M}$ containing x and a one-parameter family $\{\Omega^\epsilon\}_{\epsilon \in]0,1]} \subset W^{1+s,q}(U; \text{Ad } \eta \otimes \bigwedge^1 T^* \mathcal{M})$, such that $\Omega^\epsilon \rightarrow \Omega$ on $U \subset \mathcal{M}$ in the L^p -topology, and that the curvature of Ω^ϵ coincides with \mathcal{F} for each $\epsilon \in]0,1]$.*

Proof. The proof follows almost verbatim as for Theorem 2.1, modulo the following modifications. First, since Ω has bounded L^p -norm and hence \mathcal{F} has bounded $W^{-1,p}$ -norm on \mathcal{M} , by restricting to a small enough domain $U \subset \mathcal{M}$ we can achieve the smallness condition (2.8). Second, the Poincaré constant μ_U decreases as we shrink U (as it is controlled by the diameter of U), so we can easily achieve Eq. (2.17). \square

3. REMARKS

- (1) It remains unknown if Theorem 2.1 and/or Corollary 2.2 holds for the critical case $p = n$, even under the assumption that $\|\Omega\|_{L^n}$ is small or U is small.
- (2) Corollary 2.2 has been utilised by S. Mardare [8] (in dimension 2 and codimension 1) and Chen–Li [4] (in arbitrary dimension and codimension) to prove the existence of $W_{\text{loc}}^{2,p}$ -isometric immersions of $W^{1,p}$ -metrics on n -dimensional Riemannian manifolds ($p > n$) into Euclidean spaces, provided that the Gauss–Codazzi–Ricci equations (*i.e.*, compatibility equations for curvatures) hold in the distributional sense. Let us summarise this result in the general setting following Tenenblat [15] (see also [4]).

The convention for indices is that $1 \leq i, j, k \leq n$, $n+1 \leq \alpha, \beta \leq n+k$, and $1 \leq a, b, c \leq n+k$. That is, i, j, k index for the tangent bundle $T\mathcal{M}$, and α, β index for the (putative) normal bundle E . A Sobolev map $\iota : (\mathcal{M}, g) \rightarrow (\mathbb{R}^{n+k}, \text{Euclidean})$ is said to be an *isometric immersion* iff $d\iota$ is one-to-one outside a null set of \mathcal{M} , and that g coincides almost everywhere with the pullback of the Euclidean metric on \mathbb{R}^{n+k} under ι .

Theorem 3.1. *Let (\mathcal{M}, g) be an n -dimensional simply-connected closed Riemannian manifold with metric $g \in W^{1,p} \cap L^\infty$. Let E be a vector bundle of rank k over \mathcal{M} . Assume that E is equipped with a $W^{1,p} \cap L^\infty$ -metric g^E and an L^p -connection ∇^E compatible with*

g^E , where $p > n$ throughout. Suppose that there is an L^p -tensor field

$$\mathcal{S} : \Gamma(E) \times \Gamma(T\mathcal{M}) \longrightarrow \Gamma(T\mathcal{M}), \quad \mathcal{S}(\eta, X) \equiv \mathcal{S}_\eta X$$

such that

$$g(X, \mathcal{S}_\eta Y) = g(\mathcal{S}_\eta X, Y) \quad (3.1)$$

for all $X, Y \in \Gamma(T\mathcal{M})$ and $\eta \in \Gamma(E)$. Then define

$$\Pi : \Gamma(T\mathcal{M}) \times \Gamma(T\mathcal{M}) \rightarrow \Gamma(E)$$

by

$$g^E(\Pi(X, Y), \eta) := -g(\mathcal{S}_\eta X, Y). \quad (3.2)$$

The following are equivalent:

- (a) There exists a global isometric immersion $\iota : (\mathcal{M}, g) \rightarrow (\mathbb{R}^{n+k}, \text{Euclidean})$ in $W^{2,p}$ whose normal bundle $T\mathbb{R}^{n+k}/T(\iota\mathcal{M})$, Levi-Civita connection on the normal bundle, and second fundamental form can be identified with E , ∇^E , and Π , respectively.
- (b) The Cartan formalism holds in the sense of distributions:

$$d\omega^i = \sum_j \omega^j \wedge \Omega_j^i; \quad (3.3)$$

$$0 = d\Omega_b^a + \sum_c \Omega_b^c \wedge \Omega_c^a, \quad (3.4)$$

where $\{\omega^i\}_{1 \leq i \leq n}$ is an orthonormal coframe for $(T^*\mathcal{M}, g)$, and $\{\Omega_b^a\}_{1 \leq a, b \leq n+k}$ is the connection 1-form given by

$$\Omega_j^i(\partial_k) := g(\nabla_{\partial_k} \partial_i, \partial_j); \quad (3.5)$$

$$\Omega_\alpha^i(\partial_j) \equiv -\Omega_i^\alpha(\partial_j) := g^E(\Pi(\partial_i, \partial_j), \eta_\alpha); \quad (3.6)$$

$$\Omega_\beta^\alpha(\partial_j) := g^E(\nabla_{\partial_j}^E \eta_\alpha, \eta_\beta). \quad (3.7)$$

In the above, $\{\partial_i\}$ is the orthonormal frame for $(T\mathcal{M}, g)$ dual to $\{\omega^i\}$, and $\{\eta_\alpha\}_{n+1 \leq \alpha \leq n+k}$ is an orthonormal frame for (E, g^E) .

- (c) The Gauss–Codazzi–Ricci equations hold in the sense of distributions:

$$g(\Pi(X, Z), \Pi(Y, W)) - g(\Pi(X, W), \Pi(Y, Z)) = R(X, Y, Z, W); \quad (3.8)$$

$$\bar{\nabla}_Y \Pi(X, Z) - \bar{\nabla}_X \Pi(Y, Z) = 0; \quad (3.9)$$

$$g([\mathcal{S}_\eta, \mathcal{S}_\zeta]X, Y) = R^E(X, Y, \eta, \zeta), \quad (3.10)$$

for all $X, Y, Z, W \in \Gamma(T\mathcal{M})$ and $\eta, \zeta \in \Gamma(E)$. Here, $[\bullet, \bullet]$ is the commutator of operators, R and R^E are respectively the Riemann curvature tensors for $(T\mathcal{M}, g)$ and (E, g^E) , and $\bar{\nabla}$ is the Levi-Civita connection on Euclidean space \mathbb{R}^{n+k} .

Moreover, in (1) the isometric immersion ι is unique up to the Euclidean rigid motions in \mathbb{R}^{n+k} modulo null sets.

From the PDE point of view, in Eqs. (3.8), (3.9), and (3.10), g is given and (Π, ∇^E) are unknown.

Sketch of Proof. It is well-known that (2) \Leftrightarrow (3). Also, (3) is classically known to be a necessary condition for (1); see do Carmo [6], Chapter 6. All the above hold as purely algebraic (namely, pointwise) identities, which can be easily validated in the sense of distributions too.

It remains to show that (2) \Rightarrow (1). Since \mathcal{M} is simply-connected, it suffices to prove on a local chart, as the general case follows from a standard monodromy argument. Adapting almost verbatim the arguments in the proof of Theorem 5.2, [4] (also see [15, 8, 9]), we can reduce the proof of (1) to solving, in the distributional sense, a Pfaff system:

$$dP = -\Omega P. \quad (3.11)$$

Then, the isometric immersion ι is solved from

$$d\iota = \omega P, \quad (3.12)$$

where

$$\omega := (\omega^1, \dots, \omega^n, 0, \dots, 0)^\top.$$

The compatibility condition $d\Omega + \Omega \wedge \Omega = 0$ is given precisely by the *second structural equation* (3.4). Hence, in view of Theorem 2.1, Eq. (3.11) has a weak solution $P \in W^{1,p}$: We can first approximate Ω on sufficiently small balls by smooth connections Ω^ϵ , then deduce the existence of P^ϵ such that $dP^\epsilon = -\Omega^\epsilon P^\epsilon$, and finally conclude by passing to $W^{1,p}$ -limits. Thus, the right-hand side of Eq. (3.12) is in $W^{1,p}$, hence by a Hodge decomposition argument (or a Poincaré lemma of weak regularity; see [8, 9]) we have $\iota \in W^{2,p}$. One may now proceed as in [4] to check that ι is indeed an isometric immersion. \square

- (3) The weak stability of the gauge equation $d\Omega + \Omega \wedge \Omega = \mathcal{F}$ can be established for $\Omega \in L^r_{\text{loc}}(\mathcal{M}; \text{Ad } \eta \otimes \wedge^2 T^* \mathcal{M})$ for any $r > 2$, in contrast to $r > n = \dim \mathcal{M}$. This has been essentially proven in Chen–Slemrod–Wang [5] using the div-curl lemma of Murat [10] and Tartar [13, 14].

Theorem 3.2. *Let η be a vector bundle with structure group G over a Riemannian n -dimensional manifold \mathcal{M} , $n \geq 2$. Assume that $\Omega \in \mathfrak{A}_0^r$ for some $r > 2$ and that curvature \mathcal{F} of Ω is in the regularity class $L^{\frac{r}{2}}$. Suppose there is a sequence $\{\Omega_j\} \subset L^r(\mathcal{M}; \text{Ad } \eta \otimes \wedge^2 T^* \mathcal{M})$ such that*

- $\Omega_j \rightharpoonup \overline{\Omega}$ in the L^r -topology; and
- $d\Omega_j + \Omega_j \wedge \Omega_j = \mathcal{F} + \mathfrak{o}_j$ weakly, with the error term $\mathfrak{o}_j \rightarrow 0$ in the $W^{-1,\sigma}$ -topology for some $1 < \sigma \leq 2$ as $j \rightarrow \infty$.

Then $d\overline{\Omega} + \overline{\Omega} \wedge \overline{\Omega} = \mathcal{F}$ in the sense of distributions.

Sketch of proof. It suffices to pass to the weak limits in the quadratic nonlinear term; i.e., we prove that

$$\Omega_j \wedge \Omega_j \longrightarrow \overline{\Omega} \wedge \overline{\Omega} \quad \text{in the sense of distributions.}$$

In view of the wedge product compensated compactness theorem of Robbin–Rogers–Temple [11] (which is a generalisation of the classical div-curl lemma; also see [4]), it is enough to show that $\{d\Omega_j\}$ is precompact in the $W_{\text{loc}}^{-1,2}$ -topology. This follows from the (approximate) gauge equation $d\Omega_j = -\Omega_j \wedge \Omega_j + \mathcal{F} + \mathfrak{o}_j$.

Indeed, $\Omega_j \wedge \Omega_j + \mathcal{F}$ is precompact in $L^{r/2}(\mathcal{M}; \text{Ad } \eta \otimes \wedge^2 T^* \mathcal{M})$ by Hölder’s inequality and the assumption on \mathcal{F} . Thus, by the Sobolev embedding and/or the Rellich–Kondrachov lemma together with the assumption on \mathfrak{o}_j , one can find a $\sigma' \in [1, 2[$ such that $\{d\Omega_j\}$ is precompact in $W^{-1,\sigma'}(\mathcal{M}; \text{Ad } \eta \otimes \wedge^3 T^* \mathcal{M})$. On the other hand, since $\Omega_j \rightharpoonup \overline{\Omega}$ in the L^r -topology, we deduce that $\{d\Omega_j\}$ is bounded in $W^{-1,r}(\mathcal{M}; \text{Ad } \eta \otimes \wedge^3 T^* \mathcal{M})$

where $r > 2$. A simple interpolation theorem (see, *e.g.*, Theorem 3 in [3]) shows that $\{d\Omega_j\}$ is precompact in $W_{\text{loc}}^{-1,2}(\mathcal{M}; \text{Ad } \eta \otimes \wedge^3 T^* \mathcal{M})$. \square

REFERENCES

- [1] S. Agmon, A. Douglis, and L. Nirenberg, Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. I, *Comm. Pure Appl. Math.* **12** (1959), 623–727
- [2] S. Agmon, A. Douglis, and L. Nirenberg, Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. II, *Comm. Pure Appl. Math.* **17** (1964), 35–92
- [3] X. Ding, G.-Q. G. Chen, and P. Luo, Convergence of the fractional step Lax–Friedrichs scheme and Godunov scheme for the isentropic system of gas dynamics, *Comm. Math. Phys.* **121** (1989), 63–84
- [4] G.-Q. G. Chen and S. Li, Global weak rigidity of the Gauss–Codazzi–Ricci equations and isometric immersions of Riemannian manifolds with lower regularity, *J. Geom. Anal.* **28** (2018), 1957–2007
- [5] G.-Q. G. Chen, M. Slemrod, and D. Wang, Weak continuity of the Gauss–Codazzi–Ricci system for isometric embedding, *Proc. Amer. Math. Soc.* **138** (2010), 1843–1852
- [6] M. P. do Carmo, *Riemannian Geometry*. Translated from the second Portuguese edition by Francis Flaherty. Mathematics: Theory & Applications. Birkhäuser Boston, Inc., Boston, MA, 1992
- [7] E. Hebey, *Nonlinear analysis on manifolds: Sobolev spaces and inequalities*. Courant Lecture Notes in Mathematics, 5. New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 1999. x+309 pp. ISBN: 0-9658703-4-0
- [8] S. Mardare, On Pfaff systems with L^p coefficients and their applications in differential geometry, *J. Math. Pures Appl.* **84** (2005), 1659–1692
- [9] S. Mardare, On systems of first order linear partial differential equations with L^p coefficients, *Adv. Differential Equations* **12** (2007), 301–360
- [10] F. Murat, Compacité par compensation, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* **5** (1978), 489–507
- [11] J. W. Robbin, R. C. Rogers, and B. Temple, On weak continuity and the Hodge decomposition, *Trans. Amer. Math. Soc.* **303** (1987), 609–618
- [12] G. Schwarz, *Hodge decomposition—a method for solving boundary value problems*. Lecture Notes in Mathematics, 1607. Springer-Verlag, Berlin, 1995
- [13] L. Tartar, Compensated compactness and applications to partial differential equations, *Nonlinear Analysis and Mechanics: Heriot-Watt Symposium 4* (R. J. Knops, ed.), Research Notes in Math., Pitman, 1979
- [14] L. Tartar, The compensated compactness method applied to systems of conservation laws, *Systems of Nonlinear Partial Differential Equations* (J. M. Ball, ed.), NATO ASI Series, Reidel, 1983
- [15] K. Tenenblat, On isometric immersions of Riemannian manifolds, *Bol. Soc. Brasil. Mat.* **2** (1971), 23–36
- [16] K. K. Uhlenbeck, Connections with L^p bounds on curvature, *Comm. Math. Phys.* **83** (1982), 31–42

SIRAN LI: NEW YORK UNIVERSITY–SHANGHAI, OFFICE 1146, 1555 CENTURY AVENUE, PUDONG DISTRICT, SHANGHAI, CHINA

Email address: `s14025@nyu.edu`