

A remark on optimal data spaces for classical solutions of $\bar{\partial}$

Martino Fassina, Yifei Pan and Yuan Zhang*

Abstract

We study the minimal regularity required on the datum to guarantee the existence of classical C^1 solutions to the inhomogeneous Cauchy-Riemann equations on planar domains.

1 Introduction

Let Ω be a bounded domain in \mathbb{C} with $C^{1,\alpha}$ boundary, where $\alpha > 0$, and let \mathbf{f} be a $(0, 1)$ form on Ω . Consider the Cauchy-Riemann equation

$$\bar{\partial}u = \mathbf{f} \quad \text{on } \Omega. \quad (1)$$

The standard singular integral theory (see [3, Chapter 1]) implies the solvability of (1) in several function spaces. For instance, if $\mathbf{f} \in L^p(\Omega)$, $1 < p \leq 2$, then there exists a weak solution $u \in L^q(\Omega)$, $q < \frac{2p}{2-p}$ to (1). Moreover, if $\mathbf{f} \in L^p(\Omega)$, $p > 2$, then there exists a weak solution $u \in C^\gamma(\Omega)$, $\gamma = \frac{p-2}{p}$.

On the other hand, if $\mathbf{f} \in C^\alpha(\Omega)$ for some $0 < \alpha < 1$, then (1) admits a classical (i.e., $C^1(\bar{\Omega})$) solution $u \in C^{1,\alpha}(\Omega)$. The purpose of the note is to study the minimal regularity required on the datum \mathbf{f} to guarantee the existence of a classical solution to (1). The following example shows that continuity is not sufficient.

Example 1.1. Consider the equation $\bar{\partial}u = \mathbf{f}_\nu = f_\nu d\bar{z}$ on the disk $D(0, \frac{1}{2}) := \{z \in \mathbb{C} : |z| < \frac{1}{2}\}$, where $\nu > 0$ is fixed and

$$f_\nu(z) := \begin{cases} \frac{z}{\bar{z} \ln^\nu |z|^2} & z \neq 0 \\ 0 & z = 0. \end{cases} \quad (2)$$

Clearly, $\mathbf{f}_\nu \in C(\overline{D(0, \frac{1}{2})})$. However (see the proof at the end of the paper) if $\nu \leq 1$, then there exists no solution $u \in C^1(\overline{D(0, \frac{1}{2})})$.

We consider subspaces of $C(\bar{\Omega})$ consisting of functions satisfying a logarithmic continuity condition.

*partially supported by NSF DMS-1501024

2010 Mathematics Subject Classification. Primary 45E05; Secondary 45P05.

Keywords and phrases. $\bar{\partial}$ -equation, weak solutions, Log-continuous spaces.

Definition 1.2. Let Ω be a bounded domain in \mathbb{R}^n , $k \in \mathbb{Z}^+ \cup \{0\}$, $\nu \in \mathbb{R}^+$. A function $f \in C^k(\Omega)$ is said to be in $C^{k, \text{Log}^\nu L}(\Omega)$ if

$$\|f\|_{C^{k, \text{Log}^\nu L}(\Omega)} := \sum_{|\gamma|=0}^k \sup_{w \in \Omega} |D^\gamma f(w)| + \sum_{|\gamma|=k} \sup_{w, w+h \in \Omega} |D^\gamma f(w+h) - D^\gamma f(w)| |\ln |h||^\nu < \infty.$$

A $(0, 1)$ form \mathbf{f} is said to be in $C^{k, \text{Log}^\nu L}(\Omega)$ if all its components are in $C^{k, \text{Log}^\nu L}(\Omega)$.

When $k = 0$, the space $C^{0, \text{Log}^\nu L}(\Omega)$ is abbreviated as $C^{\text{Log}^\nu L}(\Omega)$. For any $k \in \mathbb{Z}^+ \cup \{0\}$, $0 < \nu < \mu$, and $0 < \alpha < 1$, we have

$$C^{k, \alpha}(\Omega) \hookrightarrow C^{k, \text{Log}^\mu L}(\Omega) \hookrightarrow C^{k, \text{Log}^\nu L}(\Omega) \hookrightarrow C^k(\overline{\Omega}),$$

where every inclusion map is a continuous embedding. In our main result we prove, in particular, that a classical solution to (1) exists whenever \mathbf{f} is in $C^{\text{Log}^\nu L}(\Omega)$ for some $\nu > 1$. Here is the precise statement.

Theorem 1.3. *Let $\Omega \subset \mathbb{C}$ be a bounded domain with $C^{1, \alpha}$ boundary, where $\alpha > 0$. Assume that $\mathbf{f} = fd\bar{z} \in C^{\text{Log}^\nu L}(\Omega)$, $\nu > 1$. Then there exists a solution $u \in C^{1, \text{Log}^{\nu-1} L}(\Omega)$ to $\bar{\partial}u = \mathbf{f}$ such that $\|u\|_{C^{1, \text{Log}^{\nu-1} L}(\Omega)} \leq C\|\mathbf{f}\|_{C^{\text{Log}^\nu L}(\Omega)}$, where C depends only on Ω and ν . In particular, $u \in C^1(\overline{\Omega})$, with $\|u\|_{C^1(\overline{\Omega})} \leq C\|\mathbf{f}\|_{C^{\text{Log}^\nu L}(\Omega)}$.*

Example 1.1 shows that the assumption $\nu > 1$ in Theorem 1.3 cannot be relaxed. In this sense, Theorem 1.3 identifies the largest possible data set guaranteeing the existence of classical solutions to (1).

The following example shows another way in which Theorem 1.3 is sharp: the loss of 1 in the order of Log-continuity of the solution is optimal.

Example 1.4. Let $\nu > 0$ be fixed, and consider the equation $\bar{\partial}u = \mathbf{f}_\nu = f_\nu d\bar{z}$ on the disk $D(0, \frac{1}{2})$, where f_ν is defined by (2). Then $\mathbf{f}_\nu \in C^{\text{Log}^\nu L}(D(0, \frac{1}{2}))$. However (see the proof at the end of the paper) if $\nu > 1$, there does not exist a weak solution in $C^{1, \text{Log}^\mu L}(D(0, \frac{1}{2}))$ for any $\mu > \nu - 1$.

2 Preliminaries on the integral operators T and 2T

Let Ω be a bounded domain in \mathbb{C} with $C^{1, \alpha}$ boundary, where $\alpha > 0$. Given a function $f \in C(\overline{\Omega})$, define

$$Tf(z) := \frac{-1}{2\pi i} \int_{\Omega} \frac{f(\zeta)}{\zeta - z} d\bar{\zeta} \wedge d\zeta, \quad z \in \Omega.$$

It is well known that T is a solution operator to $\bar{\partial}$ on planar domains in several function spaces (see for instance [1] and [3]). For $f \in C(\overline{\Omega})$, we have

$$\frac{\partial}{\partial \bar{z}} Tf = f, \quad \frac{\partial}{\partial z} Tf = p.v. \frac{-1}{2\pi i} \int_{\Omega} \frac{f(\zeta)}{(\zeta - \cdot)^2} d\bar{\zeta} \wedge d\zeta =: Hf \quad (3)$$

in Ω in the sense of distributions [3, Theorem 1.32]. Here $p.v.$ represents the principal value.

In their inspiring paper [2], Nijenhuis and Woelf introduced the related integral operator 2T . For functions $f \in C^\alpha(\Omega)$, where $0 < \alpha < 1$, define

$${}^2Tf(z) := \frac{-1}{2\pi i} \int_{\Omega} \frac{f(\zeta) - f(z)}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta, \quad z \in \Omega.$$

2T is a bounded operator from the space $C^\alpha(\Omega)$ to itself whenever $0 < \alpha < 1$ (see [2, Appendix 6.1.e] for a proof in the case of Ω being a disk, and [3, Theorem 1.32] for the general case). The next proposition shows that 2Tf is well defined also for functions f in Log-continuous spaces, and describes the connection between the integral operators H and 2T .

Proposition 2.1. *Let $\Omega \subset \mathbb{C}$ be a bounded domain with $C^{1,\alpha}$ boundary, where $\alpha > 0$. For every $f \in C^{Log^\nu L}(\Omega)$, with $\nu > 1$, the function 2Tf is well defined in Ω . Moreover, letting*

$$\Phi(z) := \frac{1}{2\pi i} \int_{b\Omega} \frac{d\bar{\zeta}}{\zeta - z}, \quad z \in \Omega. \quad (4)$$

we have

$$Hf(z) = {}^2Tf(z) - f(z)\Phi(z), \quad z \in \Omega. \quad (5)$$

In the special case of Ω being a disk centered at 0, then

$$Hf(z) = {}^2Tf(z), \quad z \in \Omega. \quad (6)$$

To prove Proposition 2.1 we need the two elementary lemmas below. Throughout the paper, unless otherwise specified, we use C to represent a positive constant which depends only on Ω or ν , and which may be different at each occurrence.

Lemma 2.2. *Let $\Omega \subset \mathbb{C}$ be a bounded domain with $C^{1,\alpha}$ boundary, where $\alpha > 0$, and let Φ be defined as in (4). Then $\Phi \in C^\alpha(\Omega)$. Moreover, if Ω is a disk centered at 0, then $\Phi \equiv 0$ on Ω .*

Proof. By Stokes' Theorem,

$$\int_{b\Omega} \frac{d\bar{\zeta}}{\zeta - z} = \int_{b\Omega} \frac{\bar{\zeta} d\zeta}{(\zeta - z)^2} = \left(- \int_{b\Omega} \frac{\bar{\zeta} d\zeta}{\zeta - z} \right)' = (S\bar{z})',$$

where the operator S is defined for functions $f \in C(\bar{\Omega})$ by

$$Sf(z) := - \int_{b\Omega} \frac{f(\zeta)}{\zeta - z} d\zeta, \quad z \in \Omega.$$

Recall that S sends the space $C^{1,\alpha}(\Omega)$ into itself, with $\|Sf\|_{C^{1,\alpha}(\Omega)} \leq C\|f\|_{C^{1,\alpha}(\Omega)}$ for some constant C depending only on Ω (see [3, Theorem 1.10]). Hence $\Phi \in C^\alpha(\Omega)$.

If $\Omega \subset \mathbb{C}$ is a disk centered at 0 with radius R , then using the relation $\bar{\zeta} = R^2/\zeta$ on $b\Omega$ and the Residue Theorem, we get

$$\Phi(z) = -\frac{1}{2\pi i} \int_{b\Omega} \frac{R^2 d\zeta}{\zeta^2(\zeta - z)} = \begin{cases} R^2(\frac{1}{z^2} - \frac{1}{z^2}) & \text{if } z \in \Omega, z \neq 0 \\ 0 & \text{if } z = 0. \end{cases}$$

Hence $\Phi \equiv 0$ in Ω . ■

Lemma 2.3. *Let $\nu \in \mathbb{R}^+$. There exists a constant C depending only on ν such that, for every choice of h, h_0 with $0 < h \leq h_0 < 1$, the following hold:*

1. $\int_h^{h_0} s^{-2} |\ln s|^{-\nu} ds \leq h^{-1} |\ln h|^{-\nu}$.
2. If $\nu > 1$, then $\int_0^h s^{-1} |\ln s|^{-\nu} ds \leq C |\ln h|^{1-\nu}$.

Proof. 1. Integration by parts yields, when $\nu > 0$ and $0 < h \leq h_0 < 1$,

$$\int_h^{h_0} s^{-2} |\ln s|^{-\nu} ds = h^{-1} |\ln h|^{-\nu} - h_0^{-1} |\ln h_0|^{-\nu} - \nu \int_h^{h_0} s^{-2} |\ln s|^{-\nu-1} ds.$$

In particular,

$$\int_h^{h_0} s^{-2} |\ln s|^{-\nu} ds \leq h^{-1} |\ln h|^{-\nu}.$$

2. Direct integration gives, for $\nu > 1$ and $0 < h < 1$,

$$\int_0^h s^{-1} |\ln s|^{-\nu} ds = \frac{1}{\nu-1} |\ln h|^{-\nu+1},$$

which proves the second part of the lemma. ■

Proof of Proposition 2.1: Fix $z \in \Omega$, and let h_0 be such that $0 < h_0 < 1$. By Lemma 2.3 part 2,

$$\begin{aligned} |2\pi^2 T f(z)| &= \left| \int_{\Omega} \frac{f(\zeta) - f(z)}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta \right| \\ &\leq \left| \int_{D(z, h_0) \cap \Omega} \frac{f(\zeta) - f(z)}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta \right| + \left| \int_{\Omega \setminus D(z, h_0)} \frac{f(\zeta) - f(z)}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta \right| \\ &\leq C \|f\|_{C^{Log^\nu L}(\Omega)} \int_0^{h_0} |\ln s|^{-\nu} s^{-1} ds + C h_0^{-2} \|f\|_{C(\Omega)} \\ &\leq C \|f\|_{C^{Log^\nu L}(\Omega)} |\ln h_0|^{-\nu+1} \\ &\leq C \|f\|_{C^{Log^\nu L}(\Omega)}. \end{aligned} \tag{7}$$

In particular, 2Tf is well defined pointwise in Ω .

A direct computation gives

$$\begin{aligned} p.v. \int_{\Omega} \frac{f(\zeta)}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta &= \lim_{\epsilon \rightarrow 0} \int_{\Omega \setminus D(z, \epsilon)} \frac{f(\zeta)}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta \\ &= \lim_{\epsilon \rightarrow 0} \left(\int_{\Omega \setminus D(z, \epsilon)} \frac{f(\zeta) - f(z)}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta + f(z) \int_{\Omega \setminus D(z, \epsilon)} \frac{1}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta \right). \end{aligned} \tag{8}$$

Note that

$$\left| \frac{f(\zeta) - f(z)}{(\zeta - z)^2} \right| \leq C \|f\|_{C^{Log^\nu L}(\Omega)} |\zeta - z|^{-2} |\ln |\zeta - z||^{-\nu}. \tag{9}$$

By Lemma 2.3 part 2, the function on the right side of (9) belongs to $L^1(\Omega)$. Hence the dominated convergence theorem implies

$$\lim_{\epsilon \rightarrow 0} \int_{\Omega \setminus D(z, \epsilon)} \frac{f(\zeta) - f(z)}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta = -2\pi i^2 T f(z). \quad (10)$$

On the other hand,

$$\int_{\Omega \setminus D(z, \epsilon)} \frac{1}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta = \int_{b\Omega} \frac{1}{\zeta - z} d\bar{\zeta} - \int_{bD(z, \epsilon)} \frac{1}{\zeta - z} d\bar{\zeta} = \int_{b\Omega} \frac{1}{\zeta - z} d\bar{\zeta}. \quad (11)$$

Here the first equality makes use of Stokes' Theorem, and the second equality follows from Lemma 2.2. Combining (8), (10), and (11), we conclude

$$Hf(z) = {}^2Tf(z) - f(z)\Phi(z), \quad z \in \Omega,$$

which proves (5).

In the case of Ω being a disk, (6) follows from (5) together with Lemma 2.2. ■

3 Optimal bounds for 2T and T in Log-continuous spaces

In this section we study the boundedness of the operator 2T in the space $C^{Log^\nu L}(\Omega)$. We will show in Theorem 3.4 that 2T is a bounded linear operator from $C^{Log^\nu L}(\Omega)$ into $C^{Log^{\nu-1} L}(\Omega)$ when $\nu > 1$. As a consequence, we derive our Main Theorem 1.3.

We begin by pointing out that 2T does not send $C(\overline{\Omega})$ into itself, as shown by the following example.

Example 3.1. Let f be the function defined on the disk $D(0, \frac{1}{2})$ by

$$f(z) = \begin{cases} \frac{z}{\bar{z} \ln |z|} & z \neq 0 \\ 0 & z = 0. \end{cases}$$

Then $f \in C^{Log^1 L}(D(0, \frac{1}{2})) \subset C(\overline{D(0, \frac{1}{2})})$. However,

$${}^2Tf(0) = \int_{D(0, \frac{1}{2})} \frac{1}{|\zeta|^2 \ln |\zeta|} d\zeta \wedge d\bar{\zeta} = C \int_0^{\frac{1}{2}} \frac{1}{r \ln r} dr = \infty,$$

and therefore ${}^2Tf(0)$ is not defined. In particular, ${}^2Tf \notin C(\overline{D(0, \frac{1}{2})})$.

The following important inequality is proved in [2].

Lemma 3.2. [2, Appendix 6.1d] Let $z \in D(0, R)$ and $r > 0$. Then

$$\left| \int_{D(0, R) \setminus D(z, r)} \frac{1}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta \right| \leq 8\pi. \quad (12)$$

Note that the disk $D(z, r)$ may or may not be completely contained in the ambient disk $D(0, R)$. Moreover, the bound in (12) is independent of both R and r . The proof of Lemma 3.2 in [2] relies on the symmetries of the disk, and unfortunately does not carry through when the disk $D(0, R)$ is replaced by more general domains. In the next result, using Lemma 2.2, we achieve a generalization of Lemma 3.2 to arbitrary smoothly bounded planar domains.

Lemma 3.3. *Let $\Omega \subset \mathbb{C}$ be a bounded domain with $C^{1,\alpha}$ boundary, where $\alpha > 0$. Let $z \in \Omega$ and $r > 0$. There exists a positive constant C depending only on Ω such that*

$$\left| \int_{\Omega \setminus D(z,r)} \frac{1}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta \right| \leq C.$$

Proof. Let $R > 0$ be such that $\Omega \subset\subset D(0, R)$. Since $D(0, R) \setminus D(z, r) = (D(0, R) \setminus \Omega) \sqcup (\Omega \setminus D(z, r))$, Lemma 3.2 yields

$$\begin{aligned} \left| \int_{\Omega \setminus D(z,r)} \frac{1}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta \right| &= \left| \int_{D(0,R) \setminus D(z,r)} \frac{1}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta - \int_{D(0,R) \setminus \Omega} \frac{1}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta \right| \\ &\leq 8\pi + \left| \int_{D(0,R) \setminus \Omega} \frac{1}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta \right|. \end{aligned} \quad (13)$$

By Stokes' Theorem and Lemma 2.2,

$$\int_{D(0,R) \setminus \Omega} \frac{1}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta = \int_{bD(0,R)} \frac{1}{\zeta - z} d\bar{\zeta} - \int_{b\Omega} \frac{1}{\zeta - z} d\bar{\zeta} = - \int_{b\Omega} \frac{1}{\zeta - z} d\bar{\zeta} \in C^\alpha(\Omega). \quad (14)$$

In particular,

$$\left| \int_{D(0,R) \setminus \Omega} \frac{1}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta \right| \leq C$$

for some constant C independent of r . ■

We are now ready to prove the main theorem of this section, following the ideas in [2, Appendix, 6.1e].

Theorem 3.4. *Let $\Omega \subset \mathbb{C}$ be a bounded domain with $C^{1,\alpha}$ boundary, where $\alpha > 0$. Then 2T is a bounded linear operator from $C^{\text{Log}^\nu L}(\Omega)$ into $C^{\text{Log}^{\nu-1} L}(\Omega)$, $\nu > 1$.*

Proof. Let $f \in C^{\text{Log}^\nu L}(\Omega)$ and $z \in \Omega$. Let h_0 be fixed with $0 < h_0 < \frac{1}{2}$. By (7),

$$|{}^2Tf(z)| \leq C \|f\|_{C^{\text{Log}^\nu L}(\Omega)}.$$

Next, given $z, z+h \in \Omega$, where $|h| \leq h_0$, set $D_0 := \Omega \cap D(z, 2|h|)$. Then

$$\begin{aligned} &|{}^2Tf(z) - {}^2Tf(z+h)| \\ &= \left| \int_{\Omega} \frac{f(\zeta) - f(z)}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta - \int_{\Omega} \frac{f(\zeta) - f(z+h)}{(\zeta - z-h)^2} d\bar{\zeta} \wedge d\zeta \right| \\ &= \left| \int_{\Omega \setminus D_0} (f(\zeta) - f(z+h)) \left[\frac{1}{(\zeta - z)^2} - \frac{1}{(\zeta - z-h)^2} \right] d\bar{\zeta} \wedge d\zeta - \int_{\Omega \setminus D_0} \frac{f(z) - f(z+h)}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta \right. \\ &\quad \left. + \int_{D_0} \frac{f(\zeta) - f(z)}{(\zeta - z)^2} d\bar{\zeta} \wedge d\zeta - \int_{D_0} \frac{f(\zeta) - f(z+h)}{(\zeta - z-h)^2} d\bar{\zeta} \wedge d\zeta \right| =: |I_1 + I_2 + I_3 + I_4|. \end{aligned}$$

Let γ be the segment connecting z and $z+h$. We can rewrite $|I_1|$ as follows:

$$\begin{aligned} |I_1| &= 2 \left| \int_{\Omega \setminus D_0} (f(\zeta) - f(z+h)) \left[\int_{\gamma} \frac{dw}{(\zeta - w)^3} \right] d\bar{\zeta} \wedge d\zeta \right| \\ &= 2 \left| \int_{\gamma} dw \int_{\Omega \setminus D_0} \frac{f(\zeta) - f(z+h)}{(\zeta - w)^3} d\bar{\zeta} \wedge d\zeta \right|. \end{aligned}$$

Note, if $\zeta \in \Omega \setminus D_0$ and $w \in \gamma$, then $|\zeta - w| \geq |\zeta - z| - |z - w| \geq |h|$. Thus

$$|\zeta - z - h| \leq |\zeta - w| + |w - z - h| \leq |\zeta - w| + |h| \leq 2|\zeta - w|.$$

Writing $\zeta = w + se^{i\theta}$ we see, in particular, that $\Omega \setminus D_0 \subset \{\zeta \in \mathbb{C} : |h| < s < 2R\}$, where R is the diameter of Ω . By Lemma 2.3 part 1 and the fact that $f \in C^{Log^\nu L}(\Omega)$,

$$\begin{aligned} |I_1| &\leq C\|f\|_{C^{Log^\nu L}(\Omega)}|h| \left| \int_{|h|}^{2R} |\ln s|^{-\nu} s^{-2} ds \right| \\ &\leq C\|f\|_{C^{Log^\nu L}(\Omega)}|h| \left(\left| \int_{|h|}^{h_0} |\ln s|^{-\nu} s^{-2} ds \right| + \left| \int_{h_0}^{2R} |\ln s|^{-\nu} s^{-2} ds \right| \right) \\ &\leq C\|f\|_{C^{Log^\nu L}(\Omega)}(|\ln |h||^{-\nu} + |h|) \\ &\leq C\|f\|_{C^{Log^\nu L}(\Omega)}|\ln |h||^{-\nu+1}. \end{aligned}$$

The estimate for $|I_2|$ follows directly from Lemma 3.3, and the estimate of $|I_3|$ is a straightforward consequence of Lemma 2.3 part 2, as shown below:

$$|I_3| \leq C\|f\|_{C^{Log^\nu L}(\Omega)} \left| \int_0^{|h|} |\ln s|^{-\nu} s^{-1} ds \right| \leq C\|f\|_{C^{Log^\nu L}(\Omega)} |\ln |h||^{-\nu+1}.$$

Finally, I_4 is estimated in the same way as I_3 . ■

Combining Proposition 2.1 with Theorem 3.4, we obtain the next corollary on the solution operator T , from which our Main Theorem 1.3 follows immediately.

Corollary 3.5. *Let Ω be a bounded domain in \mathbb{C} with $C^{1,\alpha}$ boundary for some $\alpha > 0$. Then T is a bounded linear operator from $C^{Log^\nu L}(\Omega)$ into $C^{1,Log^{\nu-1}L}(\Omega)$, $\nu > 1$.*

Proof. Let $f \in C^{Log^\nu L}(\Omega)$. Then $Hf \in C^{Log^{\nu-1}L}(\Omega)$ by Proposition 2.1 and Theorem 3.4. Hence, by (3), Tf is a continuous function whose weak derivatives are in $C^{Log^{\nu-1}L}(\Omega)$. Using a standard mollifier argument, we further know that (3) holds pointwise in Ω , and in particular $Tf \in C^1(\Omega)$. The $C^{1,Log^{\nu-1}L}(\Omega)$ estimate for Tf is again a consequence of Theorem 3.4. ■

Example 3.1 implies that the assumption $\nu > 1$ in Theorem 3.4 cannot be dropped. Furthermore, the following example shows that the loss by 1 in the order of Log-continuity in Theorem 3.4 and Corollary 3.5 is optimal in a very precise sense.

Example 3.6. Let f_ν be defined by (2) for some $\nu > 1$. Then $f_\nu \in C^{Log^\nu L}(D(0, \frac{1}{2}))$. However,

1. $Tf_\nu \notin C^{1,Log^\mu L}(D(0, \frac{1}{2}))$ for any $\mu > \nu - 1$.
2. ${}^2Tf_\nu \notin C^{Log^\mu L}(D(0, \frac{1}{2}))$ for any $\mu > \nu - 1$.

Proof. For the first statement, first check that $u_\nu = \frac{z}{(1-\nu)\ln^{\nu-1}|z|^2}$ satisfies $\frac{\partial}{\partial \bar{z}}u = f_\nu$ on $D(0, \frac{1}{2})$. Hence there exists a holomorphic function h on $D(0, \frac{1}{2})$ such that $Tf_\nu = u_\nu + h$. In particular, Tf_ν has the same regularity at $z = 0$ as u_ν , which is not in $C^{1,Log^\mu L}$ near 0 for any $\mu > \nu - 1$. The second statement is a direct consequence of the first, in view of the identity ${}^2Tf_\nu = \frac{\partial}{\partial z}Tf_\nu$ on $D(0, \frac{1}{2})$. ■

Proof of Example 1.1 and Example 1.4: One can check that

$$u_\nu(z) := \begin{cases} \frac{z}{(1-\nu)\ln^{\nu-1}|z|^2} & \nu \neq 1 \\ z \ln(|\ln|z||^2) & \nu = 1 \end{cases}$$

is a solution to $\bar{\partial}u = \mathbf{f}_\nu$ on $D(0, \frac{1}{2})$. Let u be any weak solution to $\bar{\partial}u = \mathbf{f}_\nu$ on $D(0, \frac{1}{2})$. Then u differs from u_ν by a holomorphic function, which is always smooth near 0. However, when $\nu \leq 1$, u_ν fails to be C^1 near 0. Moreover, for any $\mu > \nu - 1 > 0$, u_ν (and therefore u) is not in $C^{1, \text{Log}^\mu L}$ near 0. ■

References

- [1] ASTALA, K.; IWANIEC, T.; MARTIN, G.: *Elliptic partial differential equations and quasiconformal mappings in the plane*. Princeton Mathematical Series, 48. Princeton University Press, Princeton, NJ, 2009. xviii+677 pp.
- [2] NIJENHUIS, A.; WOOLF, W. B.: *Some integration problems in almost-complex and complex manifolds*. Ann. of Math. (2) **77** (1963), 424–489.
- [3] VEKUA, I. N.: *Generalized analytic functions*, vol. 29, Pergamon Press Oxford, 1962.

Martino Fassina, martino.fassina@univie.ac.at, Fakultät für Mathematik, Universität Wien, Oskar-Morgenstern-Platz 1, 1090 Wien, Austria

Yifei Pan, pan@pfw.edu, Department of Mathematical Sciences, Purdue University Fort Wayne, Fort Wayne, IN 46805-1499, USA

Yuan Zhang, zhangyu@pfw.edu, Department of Mathematical Sciences, Purdue University Fort Wayne, Fort Wayne, IN 46805-1499, USA