

Existence of solutions for a quasilinear elliptic system with local nonlinearity on \mathbb{R}^N

Xingyong Zhang^{1,2,*}; Cuiling Liu¹

¹Faculty of Science, Kunming University of Science and Technology, Kunming, Yunnan, 650500, P.R. China.

²School of Mathematics and Statistics, Central South University, Changsha, Hunan, 410083, P.R. China.

Abstract: In this paper, we investigate the existence of solutions for a class of quasilinear elliptic system

$$\begin{cases} -\operatorname{div}(\phi_1(|\nabla u|)\nabla u) + V_1(x)\phi_1(|u|)u = \lambda F_u(x, u, v), & x \in \mathbb{R}^N, \\ -\operatorname{div}(\phi_2(|\nabla v|)\nabla v) + V_2(x)\phi_2(|v|)v = \lambda F_v(x, u, v), & x \in \mathbb{R}^N, \\ u \in W^{1,\Phi_1}(\mathbb{R}^N), v \in W^{1,\Phi_2}(\mathbb{R}^N), \end{cases}$$

where $N \geq 2$, $\inf_{\mathbb{R}^N} V_i(x) > 0$, $i = 1, 2$, and $\lambda > 0$. We obtain that when the nonlinear term F satisfies some growth conditions only in a circle with center 0 and radius 4, system has a nontrivial solution (u_λ, v_λ) with $\|(u_\lambda, v_\lambda)\|_\infty \leq 2$ for every λ large enough, and the families of solutions $\{(u_\lambda, v_\lambda)\}$ satisfy that $\|(u_\lambda, v_\lambda)\| \rightarrow 0$ as $\lambda \rightarrow \infty$. Moreover, a corresponding result for a quasilinear elliptic equation is also obtained, which is better than the result for the elliptic system.

Keywords: Quasilinear elliptic system; Local nonlinearity near origin; Mountain pass theorem; Cut-off technique; Moser iteration technique

2010 Mathematics Subject Classification. 35A15, 35J62

1. Introduction

In the past years, the existence and multiplicity of solutions for the quasilinear elliptic problem with the following form

$$\begin{cases} -\operatorname{div}(\phi(|\nabla u|)\nabla u) + V(x)\phi(|u|)u = f(x, u), & x \in \Omega, \\ u = 0, & x \in \partial\Omega \end{cases} \quad (1.1)$$

has been investigated extensively (for example, see [2], [3], [5], [6], [13], [14], [15], [18], [21] and references therein), where $\Omega \subset \mathbb{R}^N$ is an open set, $N \geq 2$, V, f are continuous functions and $\phi : [0, \infty) \rightarrow [0, \infty)$ satisfies some

*Corresponding author, E-mail address: zhangxingyong1@163.com

suitable monotonicity and growth conditions. Equation (1.1) arises from some fields of physics, for example,

- (1) nonlinear elasticity: $\Phi(t) = (1 + t^2)^\gamma - 1, \gamma > \frac{1}{2}$;
- (2) plasticity: $\Phi(t) = t^\alpha (\log(1 + t))^\beta, \alpha \geq 1, \beta > 0$;
- (3) generalized Newtonian fluids: $\Phi(t) = \int_0^t s^{1-\alpha} (\sinh^{-1} s)^\beta ds, 0 \leq \alpha \leq 1, \beta > 0$,

where $\Phi(t) = \int_0^t \phi(s) s ds$ (see [2], [11], [12], [13], [14]).

Specifically, in [2], Alves-Figueiredo-Santos considered equation (1.1) with $\Omega = \mathbb{R}^N$. They assumed that $V \in C(\mathbb{R}^N, \mathbb{R})$, $\inf_{x \in \mathbb{R}^N} V(x) = V_0 > 0$, V is a radial function or is a \mathbb{Z}^N periodic function and $f \in C(\mathbb{R}^N, \mathbb{R})$ and satisfies

$$\lim_{|t| \rightarrow 0} \frac{f(t)}{\phi(|t|)|t|} = 0, \quad \lim_{|t| \rightarrow +\infty} \frac{f(t)}{\phi_*(|t|)|t|} = 0 \quad (1.2)$$

and there exists $\nu > m$ such that

$$0 < \nu F(t) = \int_0^t f(s) ds \leq t f(t) \quad \text{for all } t \in \mathbb{R} \setminus \{0\} \quad (1.3)$$

which is usually called as Ambrosseti-Rabinowitz condition ((AR) for short). After developing a Strauss-type result and a Lions-type result, they obtained equation (1.1) has a nontrivial solution. Recently, in [18], Liu considered the case that V has an infinite potential well, that is,

$$(\mathcal{V}1) \text{ for all } M > 0, \mu(V^{-1}(-\infty, M]) < \infty, \text{ where } \mu \text{ is the Lebesgue measure,}$$

or V has a finite potential well, that is,

$$(\mathcal{V}1)' \text{ for all } x \in \mathbb{R}^N, V(x) < \lim_{|x| \rightarrow \infty} V(x) < \infty.$$

He also considered the case that V is a steep potential well, that is, $V(x) = \lambda a(x) + 1$, where λ is a parameter and $a \in C(\mathbb{R}^N, \mathbb{R})$. For all these cases, he assumed that (1.2) and (AR) hold. Then he obtained some existence and multiplicity results of solutions for system (1.1).

It is easy to see that (1.2) and (AR) imply that f satisfies some conditions near both 0 and ∞ . So it is natural to ask if it is possible to restrict those conditions for f to either of them. To this end, in [8], Costa and Wang used a cut-off technique together with energy estimates to study the multiplicity of both signed and sign-changing solutions for one-parameter family of elliptic problems (1.1) with $\phi = 1$, $V = 0$, $f(x, u) = \lambda f(u)$, where $\lambda > 0$ is a parameter, Ω is a bounded smooth domain in $\mathbb{R}^N (N \geq 3)$ and $f \in C^1(\mathbb{R}, \mathbb{R})$. By such ingenious method, the nonlinearity $f(u)$ was assumed to satisfy the superlinear growth only in a neighborhood of $u = 0$. Afterwards, in [20], Medeiros and Severo applied the idea in [8] to the problem (1.1) with $\phi(t) = |t|^{p-2}$ and $f(x, u) = \lambda f(u)$ on the whole space \mathbb{R}^N , i.e. the following p -Laplacian equation

$$-\Delta_p u + V(x)|u|^{p-2}u = \lambda f(u) \quad \text{in } \mathbb{R}^N, \quad (1.4)$$

where $1 < p < N$ and $\lambda > 0$. They assumed that $V \in C(\mathbb{R}^N, \mathbb{R})$, $\inf_{x \in \mathbb{R}^N} V(x) = V_0 > 0$ and (V1) holds. Moreover, f satisfies the following conditions:

(f1) there exists $r \in (p, p^*)$ such that

$$\limsup_{|s| \rightarrow 0} \frac{f(s)s}{|s|^r} < +\infty;$$

(f2) there exists $q \in (p, p^*)$ such that

$$\liminf_{|s| \rightarrow 0} \frac{F(s)}{|s|^q} > 0;$$

(f3) there exists $\nu \in (p, p^*)$ such that

$$0 < \nu F(s) \leq sf(s) \text{ for } |s| \neq 0 \text{ small,}$$

where $p^* = \frac{Np}{N-p}$. With developing Moser iteration technique, they proved that equation (1.4) has one positive solution and one negative solution for all λ large enough. (f1)-(f3) show that $f(s)$ satisfies the superlinear growth only in a neighborhood of $s = 0$. Here, it needs to be emphasized that (f1)-(f3) with $p = 2$ were given first in [8]. The idea in [8] has been applied to various differential equations and we cite [9], [16] and [24] as some examples.

Inspired by [8] and [20], in this paper, we investigate the existence of solutions for the following quasilinear elliptic problem with a parameter

$$\begin{cases} -\operatorname{div}(\phi_1(|\nabla u|)\nabla u) + V_1(x)\phi_1(|u|)u = \lambda F_u(x, u, v), & x \in \mathbb{R}^N, \\ -\operatorname{div}(\phi_2(|\nabla v|)\nabla v) + V_2(x)\phi_2(|v|)v = \lambda F_v(x, u, v), & x \in \mathbb{R}^N, \\ u \in W^{1, \Phi_1}(\mathbb{R}^N), v \in W^{1, \Phi_2}(\mathbb{R}^N), \end{cases} \quad (1.5)$$

where $N \geq 2$, λ is a parameter with $\lambda > 0$, $F \in C^1(\mathbb{R}^N \times \mathbb{R} \times \mathbb{R}, \mathbb{R})$, ϕ_i and $V_i \in C(\mathbb{R}^N, \mathbb{R}^+)$, $i = 1, 2$ satisfy the following assumptions:

(ϕ_1) $\phi_i \in C^1(0, \infty)$, $t \rightarrow \phi_i(t)t$ are strictly increasing, $i = 1, 2$;

(ϕ_2) $1 < l_i := \inf_{t>0} \frac{t^2 \phi_i(t)}{\Phi_i(t)} \leq \sup_{t>0} \frac{t^2 \phi_i(t)}{\Phi_i(t)} =: m_i < \min\{N, l_i^*\}$, where $\Phi_i(t) := \int_0^{|t|} s \phi_i(s) ds$, $t \in \mathbb{R}$ and $l_i^* := \frac{l_i N}{N-l_i}$, $i = 1, 2$.

Recently, in [25] and [26], Wang-Zhang-Fang investigated the quasilinear elliptic system (1.5) with $\lambda = 1$. In [25], when F has a sub-linear growth, by using the least action principle, they obtained that system has at least one nontrivial solution and when F satisfies an additional symmetric condition, by using the genus theory, they obtained that system has infinitely many solutions. In [26], by using the mountain pass theorem, when F satisfies some superlinear conditions, they obtained that system has a ground state solution. Especially, they obtained the following theorem.

Theorem A. ([26]) *Assume that (ϕ_1) , (ϕ_2) and the following conditions hold:*

(V1) V_i are 1-periodic functions in x_1, \dots, x_N for all $x \in \mathbb{R}^N$, $i = 1, 2$ (called 1-periodic for short);

(V2) there exist two positive constants α_1 and α_2 such that

$$\alpha_1 \leq \min\{V_1(x), V_2(x)\} \leq \max\{V_1(x), V_2(x)\} \leq \alpha_2$$

for all $x \in \mathbb{R}^N$;

(H0) $F \in C^1(\mathbb{R}^N \times \mathbb{R} \times \mathbb{R}, \mathbb{R})$, F is 1-periodic in $x \in \mathbb{R}^N$ and $F(x, 0, 0) = 0$ for all $x \in \mathbb{R}^N$;

(H1)

$$\begin{aligned} \lim_{|(t,s)| \rightarrow 0} \frac{F_t(x, t, s)}{\phi_1(|t|) + \tilde{\Phi}_1^{-1}(\Phi_2(|s|))} &= 0, & \lim_{|(t,s)| \rightarrow 0} \frac{F_s(x, t, s)}{\tilde{\Phi}_2^{-1}(\Phi_1(|t|)) + \phi_2(|s|)} &= 0, \\ \lim_{|(t,s)| \rightarrow \infty} \frac{F_t(x, t, s)}{\Phi_{1*}'(|t|) + \tilde{\Phi}_{1*}^{-1}(\Phi_{2*}(|s|))} &= 0, & \lim_{|(t,s)| \rightarrow \infty} \frac{F_s(x, t, s)}{\tilde{\Phi}_{2*}^{-1}(\Phi_{1*}(|t|)) + \Phi_{2*}'(|s|)} &= 0 \end{aligned}$$

uniformly in $x \in \mathbb{R}^N$, where $\tilde{\Phi}_i$ and Φ_{i*} , $i = 1, 2$ are defined in section 2 below;

(H2) there exist $\mu_i > m_i$ ($i = 1, 2$) such that

$$0 < F(x, t, s) \leq \frac{1}{\mu_1} t F_t(x, t, s) + \frac{1}{\mu_2} s F_s(x, t, s), \quad \text{for all } (t, s) \neq 0.$$

Then system (1.5) with $\lambda = 1$ has a ground state solution.

It is easy to see (H0)-(H2) imply F satisfies some growth near both 0 and ∞ . In this paper, by applying the method in [8] and developing the Moser iteration technique, we only need to make some assumptions on the nonlinearity $F(x, t, s)$ in a circle with center 0 and radius 4. If we assume that $V_i, i = 1, 2$ satisfies (V1) instead of (V1) in Theorem 1.1 below, then Theorem 1.1 can be seen as an extension of the result in [20] to system (1.5) in some sense. Moreover, we shall find that the elliptic system (1.5) is more complex and more general than the scalar equation (1.1) with $x \in \mathbb{R}^N$, which directly leads to some stronger restrictions for the nonlinearity F in Theorem 1.1 because of a different Moser iteration result, see section 5 for more details. To be precise, we obtain the following theorem for system (1.5):

Theorem 1.1. Assume that (ϕ_1) -(ϕ_2), (V1) and the following conditions hold:

(ϕ_3) there exist positive constants q_i such that $t^2 \phi_i(|t|) \geq q_i |t|^{l_i}$ for all $t \in \mathbb{R}$, $i = 1, 2$;

(V0) $V_i \in C(\mathbb{R}^N, \mathbb{R}^+)$ and $V_{i,\infty} := \inf_{\mathbb{R}^N} V_i(x) > 0$, $i = 1, 2$;

(F0) $F \in C^1(\mathbb{R}^N \times \mathbb{R} \times \mathbb{R}, \mathbb{R})$ and F is 1-periodic in $x \in \mathbb{R}^N$;

(F1) there exist $k_i \in (m_i, l_i^*)$ and $M_i > 0$, $i = 1, 2$ such that

$$F(x, t, s) \geq M_1 |t|^{k_1} + M_2 |s|^{k_2}$$

for all $|(t, s)| < 4$ and $x \in \mathbb{R}^N$;

(F2) there exist $M_3 > 0, M_4 > 0, r_1 \in \left(\max \left\{ m_1, m_2, 1 + \frac{(1+m_1)(l_2-1)(\Theta_2-1)}{l_2 \Theta_2} \right\}, \min \left\{ l_1^*, 1 + \frac{l_1^*(l_1-1)}{l_1} - \frac{(l_1-1)(1+m_1)}{\Theta_1 l_1} \right\} \right)$

and $r_2 \in \left(\max \left\{ m_1, m_2, 1 + \frac{(1+m_2)(l_1-1)(\Theta_1-1)}{l_1\Theta_1} \right\}, \min \left\{ l_2^*, 1 + \frac{l_2^*(l_2-1)}{l_2} - \frac{(l_2-1)(1+m_2)}{\Theta_2 l_2} \right\} \right)$ for some $\Theta_1 > 1$ and $\Theta_2 > 1$, such that

$$|F_t(x, t, s)| \leq M_3|t|^{r_1-1} + M_4|s|^{r_2-1}, \quad |F_s(x, t, s)| \leq M_3|t|^{r_1-1} + M_4|s|^{r_2-1}$$

for all $|(t, s)| < 4$ and $x \in \mathbb{R}^N$;

(F3) there exist $\mu_i > m_i$, $i = 1, 2$ such that

$$0 < F(x, t, s) \leq \frac{1}{\mu_1} t F_t(x, t, s) + \frac{1}{\mu_2} s F_s(x, t, s)$$

for all $x \in \mathbb{R}^N$ and $|(t, s)| < 4$ with $(t, s) \neq (0, 0)$.

Then there exists $\Lambda_* > 0$ such that system (1.5) has a nontrivial solution (u_λ, v_λ) with $\|(u_\lambda, v_\lambda)\|_\infty \leq 2$ for each $\lambda > \Lambda_*$ and $\|(u_\lambda, v_\lambda)\| \rightarrow 0$ as $\lambda \rightarrow \infty$.

Remark 1.1. There exist examples satisfying Theorem 1.1. For example, let

$$F(x, t, s) = \sigma(t, s)b(x)G_1(t, s) + (1 - \sigma(t, s))b(x)G_2(t, s)$$

for all $(x, t, s) \in \mathbb{R}^N \times \mathbb{R} \times \mathbb{R}$, where $b(x)$ satisfies (V0), $G_1 \in C^1(\overline{B_8}, \mathbb{R})$ satisfies (F1)-(F3), $G_2 \in C^1(\mathbb{R}^2/B_4, \mathbb{R})$, B_R denotes a circle with center 0 and radius R , and

$$\sigma(t, s) = \begin{cases} 1, & \text{if } |(t, s)| < 4, \\ \sin \frac{\pi(t^2 + s^2 - 64)^2}{4608}, & \text{if } 4 \leq |(t, s)| \leq 8, \\ 0, & \text{if } |(t, s)| > 8. \end{cases} \quad (1.6)$$

In particular, F satisfies Theorem 1.1 but not satisfying Theorem A if we let $N = 6$, $\phi_1(t) = 4|t|^2 + 5|t|^3$ and $\phi_2(t) = 4|t|^2 \log(2 + |t|) + \frac{|t|^3}{1 + |t|}$, $b(x) = \left(1 + \sum_{i=1}^6 \cos^2 \pi x_i\right)$ (or $b(x) \equiv 1$), $G_1(t, x) = |t|^{\frac{47}{2}} + |s|^{\frac{47}{2}} + |t|^7|s|^7$ and $G_2(t, s) = |t|^3 + |s|^3$. The readers can see the detail computation in section 4.

Remark 1.2. In Theorem 1.1, we only assume that $V_i (i = 1, 2)$ are periodic functions. In fact, it is also possible that similar results can be established if $V_i (i = 1, 2)$ are radial functions or satisfy (V1) or (V1)' by combing those arguments in [2] and [18].

We organize the paper as follows. In section 2, we recall some knowledge for Orlicz and Orlicz-Sobolev spaces. In section 3, we complete the proof of Theorem 1.1. In section 4, we present some detail arguments for the example mentioned in Remark 1.1. In section 5, corresponding to Theorem 1.1, we present a result for a quasilinear elliptic equation with a parameter λ , which shows some differences between equation (1.1) with $\Omega = \mathbb{R}^N$ and system (1.5).

2. Preliminaries

In this section, we recall some notions and properties about Orlicz and Orlicz-Sobolev spaces and some useful lemmas. The readers can see these details in [1, 2, 4, 14, 17, 22, 23, 27].

We will start with some properties about N -function. Assume that $a : [0, \infty) \rightarrow [0, \infty)$ is a right continuous, monotone increasing function with

- (i) $a(0) = 0$;
- (ii) $\lim_{t \rightarrow \infty} a(t) = \infty$;
- (iii) $a(t) > 0$ whenever $t > 0$.

Then the integral $A(t) = \int_0^t a(s)ds$ is called an N -function which is defined on $[0, \infty)$.

Define the complement of A by

$$\tilde{A}(t) = \max_{s \geq 0} \{ts - A(s)\}, \quad \text{for } t \geq 0.$$

Then \tilde{A} is also an N -function and $\tilde{\tilde{A}} = A$, and the Young's inequality holds, that is

$$st \leq A(s) + \tilde{A}(t), \quad \text{for all } s, t \geq 0, \quad (2.1)$$

If

$$\sup_{t > 0} \frac{A(2t)}{A(t)} < \infty,$$

then we call A satisfies a Δ_2 -condition globally. When A satisfies Δ_2 -condition globally,

$$A(t) > A^\beta(t) \quad (2.2)$$

for any $\beta > 1$ (see [17]) and the Orlicz space $L^A(\Omega)$ is defined by the vectorial space consisting of the measurable functions $u : \Omega \rightarrow \mathbb{R}$ satisfying

$$\int_{\Omega} A(|u|)dx < \infty,$$

where $\Omega \subset \mathbb{R}^N$ is an open set. Define

$$\|u\|_A := \inf \left\{ \alpha > 0 : \int_{\Omega} A\left(\frac{|u|}{\alpha}\right) dx < 1 \right\}, \quad \text{for } u \in L^A(\Omega),$$

which is called Luxemburg norm. Then $(L^A(\Omega), \|\cdot\|_A)$ is a Banach space. If $A(t) = |t|^p$ ($1 < p < +\infty$), $(L^A(\Omega), \|\cdot\|_A)$ corresponds to the classical Lebesgue space $L^p(\Omega)$ with the norm

$$\|u\|_p := \left(\int_{\Omega} |u(x)|^p dx \right)^{\frac{1}{p}}.$$

Define

$$W^{1,A}(\Omega) = \left\{ u \in L^A(\Omega) : \frac{\partial u}{\partial x_i} \in L^A(\Omega), i = 1, \dots, N \right\}$$

with the norm

$$\|u\|_{1,A} = \|u\|_A + \|\nabla u\|_A.$$

Then $W^{1,A}(\Omega)$ is a Banach space which is called Orlicz-Sobolev space. Denote the closure of $C_0^\infty(\Omega)$ in $W^{1,A}(\Omega)$ by $W_0^{1,A}(\Omega)$. $W_0^{1,A}(\mathbb{R}^N) = W^{1,A}(\mathbb{R}^N)$ if $\Omega = \mathbb{R}^N$.

Lemma 2.1. ([14]) *If A is an N -function, then the following conditions are equivalent:*

(i)

$$l = \inf_{t>0} \frac{ta(t)}{A(t)} \leq \sup_{t>0} \frac{ta(t)}{A(t)} = m; \quad (2.3)$$

(ii) *let $\zeta_0(t) = \min\{t^l, t^m\}$, $\zeta_1(t) = \max\{t^l, t^m\}$, for $t \geq 0$. A satisfies*

$$\zeta_0(t)A(\rho) \leq A(\rho t) \leq \zeta_1(t)A(\rho), \quad \text{for all } \rho, t \geq 0;$$

(iii) *A satisfies a Δ_2 -condition globally.*

Lemma 2.2. ([14]) *If A is an N -function and (2.3) holds, then A satisfies*

$$\zeta_0(\|u\|_A) \leq \int_{\mathbb{R}^N} A(|u|)dx \leq \zeta_1(\|u\|_A), \quad \text{for all } u \in L^A(\mathbb{R}^N).$$

Lemma 2.3. ([14]) *If A is an N -function, $l, m \in (1, \infty)$ and (2.3) holds. Let \tilde{A} be the complement of A and $\zeta_2(t) = \min\{t^{\tilde{l}}, t^{\tilde{m}}\}$, $\zeta_3(t) = \max\{t^{\tilde{l}}, t^{\tilde{m}}\}$, for $t \geq 0$, where $\tilde{l} := \frac{l}{l-1}$ and $\tilde{m} := \frac{m}{m-1}$. Then \tilde{A} satisfies*

(i)

$$\tilde{m} = \inf_{t>0} \frac{t\tilde{A}'(t)}{\tilde{A}(t)} \leq \sup_{t>0} \frac{t\tilde{A}'(t)}{\tilde{A}(t)} = \tilde{l};$$

(ii)

$$\zeta_2(t)\tilde{A}(\rho) \leq \tilde{A}(\rho t) \leq \zeta_3(t)\tilde{A}(\rho), \quad \text{for all } \rho, t \geq 0;$$

(iii)

$$\zeta_2(\|u\|_{\tilde{A}}) \leq \int_{\mathbb{R}^N} \tilde{A}(|u|)dx \leq \zeta_3(\|u\|_{\tilde{A}}), \quad \text{for all } u \in L^{\tilde{A}}(\mathbb{R}^N).$$

Lemma 2.4. ([14]) *If A is an N -function, $l, m \in (1, N)$ and (2.3) holds. Let $\zeta_4(t) = \min\{t^{l^*}, t^{m^*}\}$, $\zeta_5(t) = \max\{t^{l^*}, t^{m^*}\}$, for $t \geq 0$, where $l^* := \frac{lN}{N-l}$, $m^* := \frac{mN}{N-m}$. Then A_* satisfies*

(i)

$$l^* = \inf_{t>0} \frac{tA'_*(t)}{A_*(t)} \leq \sup_{t>0} \frac{tA'_*(t)}{A_*(t)} = m^*;$$

(ii)

$$\zeta_4(t)A_*(\rho) \leq A_*(\rho t) \leq \zeta_5(t)A_*(\rho), \quad \text{for all } \rho, t \geq 0;$$

(iii)

$$\zeta_4(\|u\|_{A_*}) \leq \int_{\mathbb{R}^N} A_*(|u|) dx \leq \zeta_5(\|u\|_{A_*}), \quad \text{for all } u \in L^{A_*}(\mathbb{R}^N),$$

where A_* is the Sobolev conjugate function of A , which is defined by

$$A_*^{-1}(t) = \int_0^t \frac{A^{-1}(s)}{s^{\frac{N+1}{N}}} ds, \quad \text{for } t \geq 0.$$

Proposition 2.5. ([1]) Under the assumptions of Lemma 2.4, the embedding

$$W^{1,A}(\mathbb{R}^N) \hookrightarrow L^B(\mathbb{R}^N)$$

is continuous for any N -function B satisfying

$$\limsup_{r \rightarrow 0} \frac{B(r)}{A(r)} < \infty \quad \text{and} \quad \limsup_{r \rightarrow \infty} \frac{B(r)}{A_*(r)} < \infty.$$

Therefore, there exists a constant C such that

$$\|u\|_B \leq C\|u\|_{1,A}, \quad \text{for all } u \in W^{1,A}(\mathbb{R}^N).$$

Remark 2.1. By Lemma 2.1 and Lemma 2.3, (ϕ_1) – (ϕ_2) imply that Φ_i and $\tilde{\Phi}_i$, $i = 1, 2$, are N -functions that satisfy Δ_2 -condition globally. Thus, $L^{\Phi_i}(\mathbb{R}^N)$ and $W^{1,\Phi_i}(\mathbb{R}^N)$ are separable and reflexive Banach spaces (see [1, 23]). By (ii) in Lemma 2.1, (ii) in Lemma 2.4 and Proposition 2.5, it is easy to obtain the embedding

$$W^{1,\Phi_i}(\mathbb{R}^N) \hookrightarrow L^{p_i}(\mathbb{R}^N)$$

is continuous with $p_i \in [m_i, l_i^*]$, $i = 1, 2$, if we let $A(r) := \Phi_i(r)$ and $B(r) := r^{p_i}$. Hence, there exist positive constants $C_{0,i}$, $i = 1, 2$, such that

$$\|u\|_{p_i} \leq C_{0,i}\|u\|_{1,\Phi_i}, \tag{2.4}$$

whenever $p_i \in [m_i, l_i^*]$, $i = 1, 2$.

Next, we recall a variant of mountain pass theorem. Let X be a Banach space. $\varphi \in C^1(X, \mathbb{R})$ and $c \in \mathbb{R}$. A sequence $\{u_n\} \subset X$ is called $(PS)_c$ sequence if the sequence $\{u_n\}$ satisfies

$$\varphi(u_n) \rightarrow c, \quad \varphi'(u_n) \rightarrow 0.$$

Lemma 2.5. (Mountain Pass Theorem [4, 22, 27]) Let X be a Banach space, $\varphi \in C^1(X, \mathbb{R})$, $w \in X$ and $r > 0$ be such that $\|w\| > r$ and

$$b := \inf_{\|u\|=r} \varphi(u) > \varphi(0) \geq \varphi(w).$$

Then there exists a $(PS)_c$ sequence with

$$c := \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} \varphi(\gamma(t)), \quad (2.5)$$

and

$$\Gamma := \{\gamma \in ([0,1], X) : \gamma(0) = 0, \gamma(1) = w\}.$$

3. Proofs

Define $W := W^{1,\Phi_1}(\mathbb{R}^N) \times W^{1,\Phi_2}(\mathbb{R}^N)$ with the norm

$$\|(u, v)\| = \|u\|_{1,\Phi_1} + \|v\|_{1,\Phi_2} = \|\nabla u\|_{\Phi_1} + \|u\|_{\Phi_1} + \|\nabla v\|_{\Phi_2} + \|v\|_{\Phi_2}.$$

Then $(W, \|\cdot\|)$ is a separable and reflexive Banach space.

Next we use the idea in [8] to prove our theorem, on the whole, which is the cut-off technique together with energy estimates. In order to adapt system (1.5), we make an extension to \mathbb{R}^2 for the cut-off function in [8]. For some $\delta > 0$, let $\rho_\delta \in C^1(\mathbb{R} \times \mathbb{R}, [0, 1])$ be a cut-off function defined by

$$\rho_\delta(t, s) = \begin{cases} 1, & \text{if } |(t, s)| \leq \delta/2, \\ 0, & \text{if } |(t, s)| \geq \delta \end{cases} \quad (3.1)$$


and $t\rho'_t(t, s) + s\rho'_s(t, s) \leq 0$ for all $(t, s) \in \mathbb{R}^2$. We give some examples and their figures for the cut-off functions as follows

$$(1) \quad \rho_\delta(t, s) = \begin{cases} 1, & \text{if } |(t, s)| \leq \frac{\delta}{2}, \\ \sin \frac{8\pi(t^2 + s^2 - \delta^2)^2}{9\delta^4}, & \text{if } \frac{\delta}{2} < |(t, s)| < \delta, \\ 0, & \text{if } |(t, s)| \geq \delta; \end{cases} \quad (3.2)$$


$$(2) \quad \rho_\delta(t, s) = \begin{cases} 1, & \text{if } |(t, s)| \leq \frac{\delta}{2}, \\ \sin^2 \frac{2\pi(t^2 + s^2 - \delta^2)}{3\delta^2}, & \text{if } \frac{\delta}{2} < |(t, s)| < \delta, \\ 0, & \text{if } |(t, s)| \geq \delta; \end{cases} \quad (3.3)$$

$$(3) \quad \rho_\delta(t, s) = \begin{cases} 1, & \text{if } |(t, s)| \leq \frac{\delta}{2}, \\ \cos \frac{8\pi(t^2 + s^2 - \frac{\delta^2}{4})^2}{9\delta^4}, & \text{if } \frac{\delta}{2} < |(t, s)| < \delta, \\ 0, & \text{if } |(t, s)| \geq \delta; \end{cases} \quad (3.4)$$


$$(4) \quad \rho_\delta(t, s) = \begin{cases} 1, & \text{if } |(t, s)| \leq \frac{\delta}{2}, \\ \cos^2 \frac{2\pi(t^2 + s^2 - \frac{\delta^2}{4})}{3\delta^2}, & \text{if } \frac{\delta}{2} < |(t, s)| < \delta, \\ 0, & \text{if } |(t, s)| \geq \delta. \end{cases} \quad (3.5)$$




(a) Example (1) with $\delta = 4$.



(b) Example (1) with $\delta = 4$.



(c) Example (3) with $\delta = 4$.



(d) Example (4) with $\delta = 4$.

Remark 3.1. For the examples of ρ_δ , it seems to be natural to use exponential functions as a link between 1 and 0 because of their infinite differentiability. However, we can not find such examples with exponential functions because it seems to be difficult to ensure that ρ is differentiable at both $|(t, s)| = \frac{\delta}{2}$ and $|(t, s)| = \delta$. From the characteristic or shape of ρ_δ , sine functions and cosine functions seem to be better choices.

By (F0)-(F3) and similar to the argument of Remark 2.8 in [25], there exist positive constants M_5 and M_6 such that

$$|F(x, t, s)| \leq M_5 |t|^{r_1} + M_6 |s|^{r_2} \quad (3.6)$$

for all $|(t, s)| < 2$ and $x \in \mathbb{R}^N$. In fact, it follows from (F0) and (F3) that $F(x, 0, 0) = 0$ for all $x \in \mathbb{R}^N$. Then if $r_2 \geq r_1$, $t \in (-2, 0)$ and $s \in (-\sqrt{4 - t^2}, 0)$, by (F2), we have

$$|F(x, t, s)| \leq \int_t^0 |F_\tau(x, \tau, s)| d\tau + \int_s^0 |F_\varsigma(x, 0, \varsigma)| d\varsigma$$

$$\begin{aligned}
&\leq \int_t^0 (M_3|\tau|^{r_1-1} + M_4|s|^{r_2-1}) d\tau + \int_s^0 M_4|\varsigma|^{r_2-1} d\varsigma \\
&\leq \frac{M_3}{r_1}|t|^{r_1} + M_4|s|^{r_2-1}|t| + \frac{M_4}{r_2}|s|^{r_2} \\
&\leq \frac{M_3}{r_1}|t|^{r_1} + \frac{M_4(r_2-1)}{r_2}|s|^{r_2} + \frac{M_4}{r_2}|t|^{r_2} + \frac{M_4}{r_2}|s|^{r_2} \\
&= \left(\frac{M_3}{r_1} + \frac{M_4}{r_2}|t|^{r_2-r_1} \right) |t|^{r_1} + M_4|s|^{r_2} \\
&\leq \left(\frac{M_3}{r_1} + \frac{2^{r_2-r_1}M_4}{r_2} \right) |t|^{r_1} + M_4|s|^{r_2}
\end{aligned} \tag{3.7}$$

for all $x \in \mathbb{R}^N$. If $r_1 > r_2$, $t \in (-2, 0)$ and $s \in (-\sqrt{4-t^2}, 0)$, we have

$$\begin{aligned}
|F(x, t, s)| &\leq \int_s^0 |F_\varsigma(x, t, \varsigma)| d\varsigma + \int_t^0 |F_\tau(x, \tau, 0)| d\tau \\
&\leq \int_s^0 (M_3|t|^{r_1-1} + M_4|\varsigma|^{r_2-1}) d\varsigma + \int_t^0 M_3|\tau|^{r_1-1} d\tau \\
&\leq M_3|t|^{r_1-1}|s| + \frac{M_4}{r_2}|s|^{r_2} + \frac{M_4}{r_1}|t|^{r_1} \\
&\leq \frac{M_3(r_1-1)}{r_1}|t|^{r_1} + \frac{M_3}{r_1}|s|^{r_1} + \frac{M_4}{r_2}|s|^{r_2} + \frac{M_3}{r_1}|t|^{r_1} \\
&= \frac{M_3}{r_1}|t|^{r_1} + \left(\frac{M_4}{r_2} + \frac{M_3}{r_1}|s|^{r_1-r_2} \right) |s|^{r_2} \\
&\leq M_3|t|^{r_1} + \left(\frac{M_4}{r_2} + \frac{2^{r_1-r_2}M_3}{r_1} \right) |s|^{r_2}
\end{aligned} \tag{3.8}$$

for all $x \in \mathbb{R}^N$. Combing (3.7) and (3.8), let $M_5 = M_3 + 2^{|r_1-r_2|}M_4$ and $M_6 = M_4 + 2^{|r_1-r_2|}M_3$. Then (3.6) holds. Similar arguments can be done for other cases that $|(t, s)| < 2$.

Let $\delta = 4$ in (3.1). Define $\tilde{F} : \mathbb{R}^N \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ by

$$\tilde{F}(x, t, s) = \rho_4(t, s)F(x, t, s) + (1 - \rho_4(t, s))M_5|t|^{r_1} + (1 - \rho_4(t, s))M_6|s|^{r_2}.$$

Then by (F0)-(F3), the definition of ρ_4 and a direct computation, it is easy to obtain the following lemma:

Lemma 3.1. *Assume that (F0)-(F3) hold. Then*

(F0)' $\tilde{F} \in C^1(\mathbb{R}^N \times \mathbb{R} \times \mathbb{R}, \mathbb{R})$, \tilde{F} is 1-periodic in $x \in \mathbb{R}^N$ and $\tilde{F}(x, 0, 0) = 0$ for all $x \in \mathbb{R}^N$;

(F1)'

$$0 \leq \tilde{F}(x, t, s) \leq M_5|t|^{r_1} + M_6|s|^{r_2}, \quad \text{for all } (t, s) \in \mathbb{R}^2 \text{ and } x \in \mathbb{R}^N;$$

(F2)' there exist $M_7 > 0$ and $M_8 > 0$ such that

$$|\tilde{F}_t(x, t, s)| \leq M_7|t|^{r_1-1} + M_8|s|^{r_2-1}, \quad |\tilde{F}_s(x, t, s)| \leq M_7|t|^{r_1-1} + M_8|s|^{r_2-1}$$

for all $(t, s) \in \mathbb{R}^2$ and $x \in \mathbb{R}^N$;

(F3)'

$$\theta \tilde{F}(x, t, s) \leq \tilde{F}_t(x, t, s)t + \tilde{F}_s(x, t, s)s, \quad \text{for all } (t, s) \in \mathbb{R}^2 / \{(0, 0)\} \text{ and } x \in \mathbb{R}^N,$$

where $\theta = \min\{r_1, r_2, \mu_1, \mu_2\}$.

Consider the modified problem

$$\begin{cases} -\operatorname{div}(\phi_1(|\nabla u|)\nabla u) + V_1(x)\phi_1(|u|)u = \lambda\tilde{F}_u(x, u, v), & x \in \mathbb{R}^N, \\ -\operatorname{div}(\phi_2(|\nabla v|)\nabla v) + V_2(x)\phi_2(|v|)v = \lambda\tilde{F}_v(x, u, v), & x \in \mathbb{R}^N, \\ u \in W^{1, \Phi_1}(\mathbb{R}^N), v \in W^{1, \Phi_2}(\mathbb{R}^N). \end{cases} \quad (3.9)$$

Define the functional $\tilde{J}_\lambda : W \rightarrow \mathbb{R}$ by

$$\tilde{J}_\lambda(u, v) = \int_{\mathbb{R}^N} \Phi_1(|\nabla u|)dx + \int_{\mathbb{R}^N} \Phi_2(|\nabla v|)dx + \int_{\mathbb{R}^N} V_1(x)\Phi_1(|u|)dx + \int_{\mathbb{R}^N} V_2(x)\Phi_2(|v|)dx - \lambda \int_{\mathbb{R}^N} \tilde{F}(x, u, v)dx.$$

By (F1)' and a standard procedure, we can obtain that \tilde{J}_λ is well defined and $\tilde{J}_\lambda \in C^1(W, \mathbb{R})$ and

$$\begin{aligned} \langle \tilde{J}'_\lambda(u, v), (\tilde{u}, \tilde{v}) \rangle &= \int_{\mathbb{R}^N} (\phi_1(|\nabla u|)\nabla u, \nabla \tilde{u})dx + \int_{\mathbb{R}^N} V_1(x)\phi_1(|u|)u\tilde{u}dx \\ &\quad + \int_{\mathbb{R}^N} (\phi_2(|\nabla v|)\nabla v, \nabla \tilde{v})dx + \int_{\mathbb{R}^N} V_2(x)\phi_2(|v|)v\tilde{v}dx \\ &\quad - \lambda \int_{\mathbb{R}^N} \tilde{F}_u(x, u, v)\tilde{u}dx - \lambda \int_{\mathbb{R}^N} \tilde{F}_v(x, u, v)\tilde{v}dx \end{aligned}$$

for all $(\tilde{u}, \tilde{v}) \in W$.

Lemma 3.2. \tilde{J}_λ satisfies the mountain pass geometry, that is,

- (i) there exist two positive constants γ, η such that $\tilde{J}_\lambda(u, v) \geq \eta$ for all $\|(u, v)\| = \gamma$;
- (ii) there exists $(u_0, v_0) \in C_0^\infty(\mathbb{R}^N)/\{0\} \times C_0^\infty(\mathbb{R}^N)/\{0\}$ with $u_0 > 0, v_0 > 0$ and $\|(u_0, v_0)\|_\infty := \max_{x \in \mathbb{R}^N} (u_0^2 + v_0^2)^{\frac{1}{2}} < 1$ such that $\tilde{J}_\lambda(u_0, v_0) < 0$.

Proof. If $|t| \leq 1$, by (F2)', (ϕ_2) and Lemma 2.1, we have

$$\begin{aligned} \frac{\tilde{F}_t(x, t, s)}{\phi_1(|t|)|t| + \tilde{\Phi}_1^{-1}(\Phi_2(|s|))} &\leq \left| \frac{\tilde{F}_t(x, t, s)|t|}{\phi_1(|t|)t^2 + |t|\tilde{\Phi}_1^{-1}(\Phi_2(|s|))} \right| \\ &\leq \frac{M_7|t|^{r_1} + M_8|t||s|^{r_2-1}}{\phi_1(|t|)t^2} \\ &\leq \frac{M_7|t|^{r_1} + M_8|t||s|^{r_2-1}}{l_1\Phi_1(|t|)} \\ &\leq \frac{M_7|t|^{r_1} + M_8|t||s|^{r_2-1}}{l_1 \min\{|t|^{l_1}, |t|^{m_1}\}\Phi(1)} \\ &= \frac{M_7|t|^{r_1} + M_8|t||s|^{r_2-1}}{l_1\Phi(1)|t|^{m_1}} \\ &= \frac{M_7|t|^{r_1-1} + M_8|s|^{r_2-1}}{l_1\Phi(1)|t|^{m_1-1}}. \end{aligned} \quad (3.10)$$

Since $r_i > \max\{m_1, m_2\}, i = 1, 2$, (3.10) implies that

$$\lim_{|(t,s)| \rightarrow 0} \frac{\tilde{F}_t(x, t, s)}{\phi_1(|t|)|t| + \tilde{\Phi}_1^{-1}(\Phi_2(|s|))} = 0.$$

Similarly, we also get that

$$\lim_{|(t,s)| \rightarrow 0} \frac{\tilde{F}_s(x, t, s)}{\tilde{\Phi}_2^{-1}(\Phi_1(|t|)) + \phi_2(|s|)|s|} = 0.$$

Next, we prove that

$$\lim_{|(t,s)| \rightarrow \infty} \frac{\tilde{F}_t(x, t, s)}{\Phi'_{1*}(|t|) + \tilde{\Phi}_{1*}^{-1}(\Phi_{2*}(|s|))} = 0. \quad (3.11)$$

We divide three cases. For the case that $|t| \rightarrow \infty$ and $|s|$ is bounded, we assume that $|t| \geq 1$. Then by (F2)', Lemma 2.4 and $l_1^* \leq m_1^*$, we have

$$\begin{aligned} \frac{\tilde{F}_t(x, t, s)}{\Phi'_{1*}(|t|) + \tilde{\Phi}_{1*}^{-1}(\Phi_{2*}(|s|))} &\leq \left| \frac{\tilde{F}_t(x, t, s)|t|}{\Phi'_{1*}(|t|)|t| + |t|\tilde{\Phi}_{1*}^{-1}(\Phi_{2*}(|s|))} \right| \\ &\leq \frac{M_7|t|^{r_1} + M_8|t||s|^{r_2-1}}{\Phi'_{1*}(|t|)|t|} \\ &\leq \frac{M_7|t|^{r_1} + M_8|t||s|^{r_2-1}}{l_1^* \min\{|t|^{l_1^*}, |t|^{m_1^*}\} \Phi_{1*}(1)} = \frac{M_7|t|^{r_1-1} + M_8|s|^{r_2-1}}{l_1^* \Phi_{1*}(1)|t|^{l_1^*-1}}. \end{aligned} \quad (3.12)$$

Note that $l_1^* > r_1$. Then (3.11) holds. For the case that $|t|$ is bounded and $|s| \rightarrow \infty$, we assume that $|s| \geq 1$. Since $\frac{\tilde{\Phi}_1^{-1}(s)}{s^{\frac{N+1}{N}}}$ is nondecreasing on $(0, \infty)$, we have $\frac{\tilde{\Phi}_1^{-1}(s)}{s^{\frac{N+1}{N}}} \geq \tilde{\Phi}_1^{-1}(1)$ for all $s \geq 1$. Hence, by Lemma 2.4, we have

$$\begin{aligned} \tilde{\Phi}_{1*}^{-1}(\Phi_{2*}(|s|)) &\geq \tilde{\Phi}_{1*}^{-1}(\Phi_{2*}(1) \min\{|s|^{l_2^*}, |s|^{m_2^*}\}) \\ &= \tilde{\Phi}_{1*}^{-1}(\Phi_{2*}(1)|s|^{l_2^*}) \\ &= \int_0^{\Phi_{2*}(1)|s|^{l_2^*}} \frac{\tilde{\Phi}_1^{-1}(s)}{s^{\frac{N+1}{N}}} ds \\ &\geq \tilde{\Phi}_1^{-1}(1) \Phi_{2*}(1) |s|^{l_2^*}. \end{aligned} \quad (3.13)$$

Then by (F2)', Lemma 2.4, (3.13) and $l_1^* \leq m_1^*$, similar to the argument of (3.12), we have

$$\frac{\tilde{F}_t(x, t, s)}{\Phi'_{1*}(|t|) + \tilde{\Phi}_{1*}^{-1}(\Phi_{2*}(|s|))} \leq \frac{M_7|t|^{r_1-1} + M_8|s|^{r_2-1}}{\Phi_1^{-1}(1) \Phi_{2*}(1) |s|^{l_2^*}}.$$

Note that $l_2^* > r_2$. Then (3.11) holds. For the case that $|t| \rightarrow \infty$ and $|s| \rightarrow \infty$, we assume that $|t| \geq 1$ and $|s| \geq 1$. Then

$$\begin{aligned} \frac{\tilde{F}_t(x, t, s)}{\Phi'_{1*}(|t|) + \tilde{\Phi}_{1*}^{-1}(\Phi_{2*}(|s|))} &\leq \left| \frac{\tilde{F}_t(x, t, s)|t|}{\Phi'_{1*}(|t|)|t| + |t|\tilde{\Phi}_{1*}^{-1}(\Phi_{2*}(|s|))} \right| \\ &\leq \frac{M_7|t|^{r_1} + M_8|t||s|^{r_2-1}}{\Phi'_{1*}(|t|)|t| + |t|\tilde{\Phi}_{1*}^{-1}(\Phi_{2*}(|s|))} \\ &\leq \frac{M_7|t|^{r_1} + M_8|t||s|^{r_2-1}}{l_1^* \min\{|t|^{l_1^*}, |t|^{m_1^*}\} \Phi_{1*}(1) + |t|\Phi_1^{-1}(1) \Phi_{2*}(1) |s|^{l_2^*}} \\ &= \frac{M_7|t|^{r_1-1} + M_8|s|^{r_2-1}}{l_1^* \Phi_{1*}(1) |t|^{l_1^*-1} + \Phi_1^{-1}(1) \Phi_{2*}(1) |s|^{l_2^*}}. \end{aligned}$$

Note that $l_1^* > r_1$ and $l_2^* > r_2$. Then (3.11) holds. Hence, (F2)' implies (F2) of Lemma 3.14 in [26]. So the conclusion holds by Lemma 3.14 and Lemma 3.15 in [26]. \square

Remark 3.1. There is a similar result of Lemma 3.2 in [26] (see [26], Corollary 3.3) where the authors did not show the detail proof. Here, we present the proof for readers' convenience.

Since (F0)'-(F3)' imply those conditions of Theorem 3.1 in [26], it follows from Lemma 3.14-Lemma 3.16 in [26] that system (3.9) has a nontrivial solution (u_λ, v_λ) such that $\tilde{J}_\lambda(u_\lambda, v_\lambda) = c_\lambda$ with

$$c_\lambda := \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} \tilde{J}_\lambda(\gamma(t)),$$

and

$$\Gamma := \{\gamma \in ([0, 1], W) : \gamma(0) = (0, 0), \gamma(1) = (u_0, v_0)\}.$$

Next, we make an estimate for $\|u_\lambda\|_{1, \Phi_1}$ and $\|v_\lambda\|_{1, \Phi_2}$. We introduce the functional $\bar{J}_\lambda : W \rightarrow \mathbb{R}$ as follows

$$\begin{aligned} \bar{J}_\lambda(u) &= \int_{\mathbb{R}^N} \Phi_1(|\nabla u|)dx + \int_{\mathbb{R}^N} V_1(x)\Phi_1(|u|)dx + \int_{\mathbb{R}^N} \Phi_2(|\nabla v|)dx + \int_{\mathbb{R}^N} V_2(x)\Phi_2(|v|)dx \\ &\quad - \lambda M_1 \int_{\mathbb{R}^N} |u|^{k_1}dx - \lambda M_2 \int_{\mathbb{R}^N} |v|^{k_2}dx. \end{aligned}$$

Lemma 3.3. There exist $\Lambda_0 > 0$ such that for each $\lambda > \Lambda_0$, there exists $C_* > 0$ such that

$$\begin{aligned} \|u_\lambda\|_{1, \Phi_1} &\leq C_* \max \left\{ \lambda^{-\frac{1}{k_1-l_1}} + \lambda^{-\frac{l_2}{l_1(k_2-l_2)}}, \lambda^{-\frac{l_1}{m_1(k_1-l_1)}} + \lambda^{-\frac{l_2}{m_1(k_2-l_2)}} \right\}, \\ \|v_\lambda\|_{2, \Phi_2} &\leq C_* \max \left\{ \lambda^{-\frac{l_1}{l_2(k_1-l_1)}} + \lambda^{-\frac{1}{k_2-l_2}}, \lambda^{-\frac{l_1}{m_2(k_1-l_1)}} + \lambda^{-\frac{l_2}{m_2(k_2-l_2)}} \right\}. \end{aligned}$$

Proof. By (V0) and (V1), we know that $V_i^\infty := \max_{x \in \bar{\Omega}} V_i(x), i = 1, 2$ exist. Then by Lemma 2.2, we have

$$\begin{aligned} \bar{J}_\lambda(su_0, sv_0) &= \int_{\mathbb{R}^N} \Phi_1(s|\nabla u_0|)dx + \int_{\mathbb{R}^N} V_1(x)\Phi_1(s|u_0|)dx + \int_{\mathbb{R}^N} \Phi_2(s|\nabla v_0|)dx \\ &\quad + \int_{\mathbb{R}^N} V_2(x)\Phi_2(s|v_0|)dx - \lambda M_1 s^{k_1} \int_{\mathbb{R}^N} |u_0|^{k_1}dx - \lambda M_2 s^{k_2} \int_{\mathbb{R}^N} |v_0|^{k_2}dx \\ &\leq \max\{s^{l_1}\|\nabla u_0\|_{\Phi_1}^{l_1}, s^{m_1}\|\nabla u_0\|_{\Phi_1}^{m_1}\} + V_1^\infty \max\{s^{l_1}\|u_0\|_{\Phi_1}^{l_1}, s^{m_1}\|u_0\|_{\Phi_1}^{m_1}\} \\ &\quad + \max\{s^{l_2}\|\nabla v_0\|_{\Phi_2}^{l_2}, s^{m_2}\|\nabla v_0\|_{\Phi_2}^{m_2}\} + V_2^\infty \max\{s^{l_2}\|v_0\|_{\Phi_2}^{l_2}, s^{m_2}\|v_0\|_{\Phi_2}^{m_2}\} \\ &\quad - \lambda M_1 s^{k_1} \int_{\mathbb{R}^N} |u_0|^{k_1}dx - \lambda M_2 s^{k_2} \int_{\mathbb{R}^N} |v_0|^{k_2}dx \\ &\leq s^{l_1} \left(\|\nabla u_0\|_{\Phi_1}^{l_1} + V_1^\infty \|u_0\|_{\Phi_1}^{l_1} + \|\nabla u_0\|_{\Phi_1}^{m_1} + V_1^\infty \|u_0\|_{\Phi_1}^{m_1} \right) \\ &\quad + s^{l_2} \left(\|\nabla v_0\|_{\Phi_2}^{l_2} + V_2^\infty \|v_0\|_{\Phi_2}^{l_2} + \|\nabla v_0\|_{\Phi_2}^{m_2} + V_2^\infty \|v_0\|_{\Phi_2}^{m_2} \right) \\ &\quad - \lambda M_1 s^{k_1} \int_{\mathbb{R}^N} |u_0|^{k_1}dx - \lambda M_2 s^{k_2} \int_{\mathbb{R}^N} |v_0|^{k_2}dx \end{aligned}$$

for $s \in [0, 1]$ and (u_0, v_0) obtained in Lemma 3.2. Let

$$g_1(s) = s^{l_1} \left(\|\nabla u_0\|_{\Phi_1}^{l_1} + V_1^\infty \|u_0\|_{\Phi_1}^{l_1} + \|\nabla u_0\|_{\Phi_1}^{m_1} + V_1^\infty \|u_0\|_{\Phi_1}^{m_1} \right) - \lambda M_1 s^{k_1} \int_{\mathbb{R}^N} |u_0|^{k_1}dx,$$

$$g_2(s) = s^{l_2} \left(\|\nabla v_0\|_{\Phi_2}^{l_2} + V_2^\infty \|v_0\|_{\Phi_2}^{l_2} + \|\nabla v_0\|_{\Phi_2}^{m_2} + V_2^\infty \|v_0\|_{\Phi_2}^{m_2} \right) - \lambda M_2 s^{k_2} \int_{\mathbb{R}^N} |v_0|^{k_2} dx.$$

Obviously, there exist $\Lambda_1 > 0$ and $\Lambda_2 > 0$ such that $s_{\Lambda_1} := \left(\frac{l_1 (\|\nabla u_0\|_{\Phi_1}^{l_1} + V_1^\infty \|u_0\|_{\Phi_1}^{l_1} + \|\nabla u_0\|_{\Phi_1}^{m_1} + V_1^\infty \|u_0\|_{\Phi_1}^{m_1})}{\lambda M_1 k_1 \int_{\mathbb{R}^N} |u_0|^{k_1} dx} \right)^{\frac{1}{k_1 - l_1}} \in (0, 1)$ and $s_{\Lambda_2} := \left(\frac{l_2 (\|\nabla v_0\|_{\Phi_2}^{l_2} + V_2^\infty \|v_0\|_{\Phi_2}^{l_2} + \|\nabla v_0\|_{\Phi_2}^{m_2} + V_2^\infty \|v_0\|_{\Phi_2}^{m_2})}{\lambda M_2 k_2 \int_{\mathbb{R}^N} |v_0|^{k_2} dx} \right)^{\frac{1}{k_2 - l_2}} \in (0, 1)$ if $\lambda > \max\{\Lambda_1, \Lambda_2\}$. Then when $s = s_{\Lambda_1}$, $g_1(s)$ attains the maximum on $[0, 1]$. Then there exists $C_{1,*} > 0$ such that

$$\max_{s \in [0,1]} g_1(s) = \lambda^{-\frac{l_1}{k_1 - l_1}} k_1^{-\frac{k_1}{k_1 - l_1}} l_1^{\frac{l_1}{k_1 - l_1}} (k_1 - l_1) A^{\frac{k_1}{k_1 - l_1}} \left(M_1 \int_{\mathbb{R}^N} |u_0|^{k_1} dx \right)^{-\frac{l_1}{k_1 - l_1}} \leq C_{1,*} \lambda^{-\frac{l_1}{k_1 - l_1}},$$

where $A = \|\nabla u_0\|_{\Phi_1}^{l_1} + V_1^\infty \|u_0\|_{\Phi_1}^{l_1} + \|\nabla u_0\|_{\Phi_1}^{m_1} + V_1^\infty \|u_0\|_{\Phi_1}^{m_1}$. When $s = s_{\Lambda_2}$, $g_2(s)$ attains the maximum on $[0, 1]$ and there exists $C_{2,*} > 0$ such that

$$\max_{s \in [0,1]} g_2(s) = \lambda^{-\frac{l_2}{k_2 - l_2}} k_2^{-\frac{k_2}{k_2 - l_2}} l_2^{\frac{l_2}{k_2 - l_2}} (k_2 - l_2) B^{\frac{k_2}{k_2 - l_2}} \left(M_2 \int_{\mathbb{R}^N} |v_0|^{k_2} dx \right)^{-\frac{l_2}{k_2 - l_2}} \leq C_{2,*} \lambda^{-\frac{l_2}{k_2 - l_2}},$$

where $B = \|\nabla v_0\|_{\Phi_2}^{l_2} + V_2^\infty \|v_0\|_{\Phi_2}^{l_2} + \|\nabla v_0\|_{\Phi_2}^{m_2} + V_2^\infty \|v_0\|_{\Phi_2}^{m_2}$. Notice that $|s(u_0(x), v_0(x))| \leq \|(u_0, v_0)\|_\infty < 1$ for all $x \in \mathbb{R}^N$ and $s \in [0, 1]$. Then by the definition of \tilde{F} and (F1), we have $\tilde{F}(su_0, sv_0) = F(su_0, sv_0) \geq M_1 |su_0(x)|^{k_1} + M_2 |sv_0(x)|^{k_2}$. Thus,

$$c_\lambda := \inf_{\gamma \in \Gamma} \max_{s \in [0,1]} \tilde{J}_\lambda(\gamma(s)) \leq \max_{s \in [0,1]} \tilde{J}_\lambda(su_0, sv_0) \leq \max_{s \in [0,1]} \bar{J}_\lambda(su_0, sv_0) \leq C_{1,*} \lambda^{-\frac{l_1}{k_1 - l_1}} + C_{2,*} \lambda^{-\frac{l_2}{k_2 - l_2}}. \quad (3.14)$$

Note that (u_λ, v_λ) is a critical point of \tilde{J}_λ with critical value c_λ . Since $\langle \tilde{J}'_\lambda(u_\lambda, v_\lambda), (u_\lambda, v_\lambda) \rangle = 0$ and $\theta \geq \max\{m_1, m_2\}$, by (ϕ_2) , (F3)' and Lemma 2.2, we have

$$\begin{aligned} \theta c_\lambda &= \theta \tilde{J}_\lambda(u_\lambda, v_\lambda) \\ &= \theta \langle \tilde{J}_\lambda(u_\lambda, v_\lambda), (u_\lambda, v_\lambda) \rangle - \langle \tilde{J}'_\lambda(u_\lambda, v_\lambda), (u_\lambda, v_\lambda) \rangle \\ &= \theta \int_{\mathbb{R}^N} \Phi_1(|\nabla u_\lambda|) dx + \theta \int_{\mathbb{R}^N} V_1(x) \Phi_1(|u_\lambda|) dx + \theta \int_{\mathbb{R}^N} \Phi_2(|\nabla v_\lambda|) dx + \theta \int_{\mathbb{R}^N} V_2(x) \Phi_2(|v_\lambda|) dx \\ &\quad - \theta \lambda \int_{\mathbb{R}^N} \tilde{F}(x, u_\lambda, v_\lambda) dx - \int_{\mathbb{R}^N} \phi_1(|\nabla u_\lambda|) |\nabla u_\lambda|^2 dx - \int_{\mathbb{R}^N} V_1(x) \phi_1(|u_\lambda|) u_\lambda^2 dx \\ &\quad - \int_{\mathbb{R}^N} \phi_2(|\nabla v_\lambda|) |\nabla v_\lambda|^2 dx - \int_{\mathbb{R}^N} V_2(x) \phi_2(|v_\lambda|) v_\lambda^2 dx + \lambda \int_{\mathbb{R}^N} \tilde{F}_u(x, u_\lambda, v_\lambda) u_\lambda dx \\ &\quad + \lambda \int_{\mathbb{R}^N} \tilde{F}_v(x, u_\lambda, v_\lambda) v_\lambda dx \\ &\geq (\theta - m_1) \int_{\mathbb{R}^N} \Phi_1(|\nabla u_\lambda|) dx + (\theta - m_1) \int_{\mathbb{R}^N} V_1(x) \Phi_1(|u_\lambda|) dx + (\theta - m_2) \int_{\mathbb{R}^N} \Phi_2(|\nabla v_\lambda|) dx \\ &\quad + (\theta - m_2) \int_{\mathbb{R}^N} V_2(x) \Phi_2(|v_\lambda|) dx \\ &\geq (\theta - m_1) \min\{\|\nabla u_\lambda\|_{\Phi_1}^{l_1}, \|\nabla u_\lambda\|_{\Phi_1}^{m_1}\} + (\theta - m_1) V_{1,\infty} \min\{\|u_\lambda\|_{\Phi_1}^{l_1}, \|u_\lambda\|_{\Phi_1}^{m_1}\} \\ &\quad + (\theta - m_2) \min\{\|\nabla v_\lambda\|_{\Phi_2}^{l_2}, \|\nabla v_\lambda\|_{\Phi_2}^{m_2}\} + (\theta - m_2) V_{2,\infty} \min\{\|v_\lambda\|_{\Phi_2}^{l_2}, \|v_\lambda\|_{\Phi_2}^{m_2}\}, \end{aligned}$$

which, together with (3.14), implies that

$$\begin{aligned} & \min\{\|\nabla u_\lambda\|_{\Phi_1}^{l_1}, \|\nabla u_\lambda\|_{\Phi_1}^{m_1}\} + V_{1,\infty} \min\{\|u_\lambda\|_{\Phi_1}^{l_1}, \|u_\lambda\|_{\Phi_1}^{m_1}\} \\ & \leq \frac{\theta}{\theta - m_1} c_\lambda \leq \frac{\theta}{\theta - m_1} C_{1,*} \lambda^{-\frac{l_1}{k_1 - l_1}} + \frac{\theta}{\theta - m_1} C_{2,*} \lambda^{-\frac{l_2}{k_2 - l_2}} \end{aligned}$$

and

$$\begin{aligned} & \min\{\|\nabla v_\lambda\|_{\Phi_2}^{l_2}, \|\nabla v_\lambda\|_{\Phi_2}^{m_2}\} + V_{2,\infty} \min\{\|v_\lambda\|_{\Phi_2}^{l_2}, \|v_\lambda\|_{\Phi_2}^{m_2}\} \\ & \leq \frac{\theta}{\theta - m_2} c_\lambda \leq \frac{\theta}{\theta - m_2} C_{1,*} \lambda^{-\frac{l_1}{k_1 - l_1}} + \frac{\theta}{\theta - m_2} C_{2,*} \lambda^{-\frac{l_2}{k_2 - l_2}}. \end{aligned}$$

Hence, there exists $C_* > 0$ such that

$$\begin{aligned} \|u_\lambda\|_{1,\Phi_1} &= \|\nabla u_\lambda\|_{\Phi_1} + \|u_\lambda\|_{\Phi_1} \\ &\leq \left(\frac{1}{V_{1,\infty}} + 1\right) \max \left\{ \left(\frac{\theta}{\theta - m_1} C_{1,*}\right)^{\frac{1}{l_1}} \lambda^{-\frac{1}{k_1 - l_1}} + \left(\frac{\theta}{\theta - m_1} C_{2,*}\right)^{\frac{1}{l_1}} \lambda^{-\frac{l_2}{l_1(k_2 - l_2)}}, \right. \\ &\quad \left. \left(\frac{\theta}{\theta - m_1} C_{1,*}\right)^{\frac{1}{m_1}} \lambda^{-\frac{l_1}{m_1(k_1 - l_1)}} + \left(\frac{\theta}{\theta - m_1} C_{2,*}\right)^{\frac{1}{m_1}} \lambda^{-\frac{l_2}{m_1(k_2 - l_2)}} \right\} \\ &\leq C_* \max \left\{ \lambda^{-\frac{1}{k_1 - l_1}} + \lambda^{-\frac{l_2}{l_1(k_2 - l_2)}}, \lambda^{-\frac{l_1}{m_1(k_1 - l_1)}} + \lambda^{-\frac{l_2}{m_1(k_2 - l_2)}} \right\} \end{aligned}$$

and

$$\begin{aligned} \|v_\lambda\|_{2,\Phi_2} &= \|\nabla v_\lambda\|_{\Phi_2} + \|v_\lambda\|_{\Phi_2} \\ &\leq \left(\frac{1}{V_{2,\infty}} + 1\right) \max \left\{ \left(\frac{\theta}{\theta - m_2} C_{1,*}\right)^{\frac{1}{l_2}} \lambda^{-\frac{l_1}{l_2(k_1 - l_1)}} + \left(\frac{\theta}{\theta - m_2} C_{2,*}\right)^{\frac{1}{l_2}} \lambda^{-\frac{1}{k_2 - l_2}}, \right. \\ &\quad \left. \left(\frac{\theta}{\theta - m_2} C_{1,*}\right)^{\frac{1}{m_2}} \lambda^{-\frac{l_1}{m_2(k_1 - l_1)}} + \left(\frac{\theta}{\theta - m_2} C_{2,*}\right)^{\frac{1}{m_2}} \lambda^{-\frac{l_2}{m_2(k_2 - l_2)}} \right\} \\ &\leq C_* \max \left\{ \lambda^{-\frac{l_1}{l_2(k_1 - l_1)}} + \lambda^{-\frac{1}{k_2 - l_2}}, \lambda^{-\frac{l_1}{m_2(k_1 - l_1)}} + \lambda^{-\frac{l_2}{m_2(k_2 - l_2)}} \right\}. \end{aligned}$$

Let $\Lambda_0 = \max\{\Lambda_1, \Lambda_2\}$. Then we complete the proof. \square

Lemma 3.4. *There exists $\Lambda_* > \Lambda_0$ such that for all $\lambda > \Lambda_*$,*

$$\|u_\lambda\|_\infty \leq 1, \|v_\lambda\|_\infty \leq 1.$$

Proof. The proof is motivated by Lemma 2.6 in [20], which originates from [19] and is often called Moser iteration technique (see also [7] and [10] for some related arguments). However, because of the interaction of u and v , our proof is more difficult than those in [20], [7] and [10], where some scalar equations were investigated.

Next, we start the proof. Since (u_λ, v_λ) is a critical point of \tilde{J}_λ , we have

$$\int_{\mathbb{R}^N} (\phi_1(|\nabla u_\lambda|) \nabla u_\lambda, \nabla \tilde{u}) dx + \int_{\mathbb{R}^N} V_1(x) \phi_1(|u_\lambda|) u_\lambda \tilde{u} dx = \lambda \int_{\mathbb{R}^N} \tilde{F}_{u_\lambda}(x, u_\lambda, v_\lambda) \tilde{u} dx, \quad (3.15)$$

$$\int_{\mathbb{R}^N} (\phi_2(|\nabla v_\lambda|) \nabla v_\lambda, \nabla \tilde{v}) dx + \int_{\mathbb{R}^N} V_2(x) \phi_2(|v_\lambda|) v_\lambda \tilde{v} dx = \lambda \int_{\mathbb{R}^N} \tilde{F}_{v_\lambda}(x, u_\lambda, v_\lambda) \tilde{v} dx.$$

for all $(\tilde{u}, \tilde{v}) \in W$. Next, we prove $\|u_\lambda\|_\infty \leq 1$. Without loss of generality, for each $k > 0$, we define

$$u_k = \begin{cases} u_\lambda, & \text{if } u_\lambda \leq k, \\ k, & \text{if } u_\lambda > k, \end{cases}$$

$\varphi_k = |u_k|^{l_1(\beta_1-1)} u_\lambda$ and $w_k = u_\lambda |u_k|^{\beta_1-1}$ with $\beta_1 > 1$. By (ϕ_3) , we have $\phi_1(|\nabla u_\lambda|) |\nabla u_\lambda|^2 \geq q_1 |\nabla u_\lambda|^{l_1}$. Then it follows from (2.1), (2.2), Hölder inequality, Lemma 2.2 and (2.4) that

$$\begin{aligned} & q_1 \int_{\mathbb{R}^N} |\nabla u_\lambda|^{l_1} |u_k|^{l_1(\beta_1-1)} dx \\ & \leq \int_{\mathbb{R}^N} \phi_1(|\nabla u_\lambda|) (\nabla u_\lambda, \nabla \varphi_k) dx - l_1(\beta_1-1) \int_{\mathbb{R}^N} |u_k|^{l_1(\beta_1-1)-2} u_k u_\lambda \phi_1(|\nabla u_\lambda|) (\nabla u_\lambda, \nabla u_k) dx \\ & \leq - \int_{\mathbb{R}^N} V_1(x) \phi(|u_\lambda|) u_\lambda \varphi_k dx + \lambda \int_{\mathbb{R}^N} \tilde{F}_{u_\lambda}(x, u_\lambda, v_\lambda) \varphi_k dx \\ & \leq \lambda M_7 \int_{\mathbb{R}^N} |u_\lambda|^{r_1} |u_k|^{l_1(\beta_1-1)} dx + \lambda M_8 \int_{\mathbb{R}^N} |v_\lambda|^{r_2-1} |u_\lambda| |u_k|^{l_1(\beta_1-1)} dx \\ & = \lambda M_7 \int_{\mathbb{R}^N} |u_\lambda|^{r_1-l_1} |w_k|^{l_1} dx + \lambda M_8 \int_{\mathbb{R}^N} |v_\lambda|^{r_2-1} |w_k| |u_\lambda|^{(l_1-1)(\beta_1-1)} dx \\ & \leq \lambda M_7 \int_{\mathbb{R}^N} |u_\lambda|^{r_1-l_1} |w_k|^{l_1} dx + \lambda M_8 \int_{\mathbb{R}^N} |v_\lambda|^{r_2-1} |w_k| |u_\lambda|^{(l_1-1)(\beta_1-1)} dx \\ & = \lambda M_7 \int_{\mathbb{R}^N} |u_\lambda|^{r_1-l_1} |w_k|^{l_1} dx + \lambda M_8 \int_{\mathbb{R}^N} |v_\lambda|^{r_2-1} |u_\lambda|^{(l_1-1)(\beta_1-1)-\frac{r_1-l_1}{l_1}} |u_\lambda|^{\frac{r_1-l_1}{l_1}} |w_k| dx \\ & \leq \lambda M_7 \int_{\mathbb{R}^N} |u_\lambda|^{r_1-l_1} |w_k|^{l_1} dx + \lambda M_8 \left(\int_{\mathbb{R}^N} |v_\lambda|^{\frac{l_1(r_2-1)}{l_1-1}} |u_\lambda|^{l_1(\beta_1-1)-\frac{r_1-l_1}{l_1-1}} dx \right)^{1-\frac{1}{l_1}} \left(\int_{\mathbb{R}^N} |w_k|^{l_1} |u_\lambda|^{r_1-l_1} dx \right)^{\frac{1}{l_1}} \\ & \leq \lambda M_8 \left(\int_{\mathbb{R}^N} |v_\lambda|^{\frac{\Theta_1 l_1(r_2-1)}{(\Theta_1-1)(l_1-1)}} dx \right)^{\frac{(l_1-1)(\Theta_1-1)}{\Theta_1 l_1}} \left(\int_{\mathbb{R}^N} |u_\lambda|^{\Theta_1 l_1(\beta_1-1)-\frac{\Theta_1(r_1-l_1)}{l_1-1}} dx \right)^{\frac{l_1-1}{\Theta_1 l_1}} \left(\int_{\mathbb{R}^N} |w_k|^{l_1} |u_\lambda|^{r_1-l_1} dx \right)^{\frac{1}{l_1}} \\ & \quad + \lambda M_7 \int_{\mathbb{R}^N} |u_\lambda|^{r_1-l_1} |w_k|^{l_1} dx \\ & = \lambda M_8 \left(\int_{\mathbb{R}^N} |v_\lambda|^{\frac{\Theta_1 l_1(r_2-1)}{(\Theta_1-1)(l_1-1)}-m_2} |v_\lambda|^{m_2} dx \right)^{\frac{(l_1-1)(\Theta_1-1)}{\Theta_1 l_1}} \left(\int_{\mathbb{R}^N} |u_\lambda|^{\Theta_1 l_1(\beta_1-1)-\frac{\Theta_1(r_1-l_1)}{l_1-1}-m_1} |u_\lambda|^{m_1} dx \right)^{\frac{l_1-1}{\Theta_1 l_1}} \\ & \quad \times \left(\int_{\Omega} |w_k|^{l_1} |u_\lambda|^{r_1-l_1} dx \right)^{\frac{1}{l_1}} + \lambda M_7 \int_{\mathbb{R}^N} |u_\lambda|^{r_1-l_1} |w_k|^{l_1} dx \\ & \leq \lambda M_8 \left(\int_{\mathbb{R}^N} |w_k|^{l_1} |u_\lambda|^{r_1-l_1} dx \right)^{\frac{1}{l_1}} \left(\int_{\mathbb{R}^N} (\Phi_2(|v_\lambda|) + \Phi_2^{-1}(1))^{\frac{\Theta_1 l_1(r_2-1)}{(\Theta_1-1)(l_1-1)}-m_2} |v_\lambda|^{m_2} dx \right)^{\frac{(l_1-1)(\Theta_1-1)}{\Theta_1 l_1}} \\ & \quad \times \left(\int_{\mathbb{R}^N} (\Phi_1(|u_\lambda|) + \Phi_1^{-1}(1))^{\Theta_1 l_1(\beta_1-1)-\frac{\Theta_1(r_1-l_1)}{l_1-1}-m_1} |u_\lambda|^{m_1} dx \right)^{\frac{l_1-1}{\Theta_1 l_1}} + \lambda M_7 \int_{\mathbb{R}^N} |u_\lambda|^{r_1-l_1} |w_k|^{l_1} dx \\ & \leq \lambda M_8 2^{r_2-1-\frac{(m_2+1)(l_1-1)(\Theta_1-1)}{\Theta_1 l_1}} 2^{(l_1-1)(\beta_1-1)-\frac{r_1-l_1}{l_1}-\frac{(m_1+1)(l_1-1)}{\Theta_1 l_1}} \left(\int_{\mathbb{R}^N} |w_k|^{l_1} |u_\lambda|^{r_1-l_1} dx \right)^{\frac{1}{l_1}} \\ & \quad \times \left(\int_{\mathbb{R}^N} \left(\Phi_2^{\frac{\Theta_1 l_1(r_2-1)}{(\Theta_1-1)(l_1-1)}-m_2} (|v_\lambda|) + (\Phi_2^{-1}(1))^{\frac{\Theta_1 l_1(r_2-1)}{(\Theta_1-1)(l_1-1)}-m_2} \right) |v_\lambda|^{m_2} dx \right)^{\frac{(l_1-1)(\Theta_1-1)}{\Theta_1 l_1}} \end{aligned}$$

$$\begin{aligned}
& \times \left(\int_{\mathbb{R}^N} \left(\Phi_1^{\Theta_1 l_1 (\beta_1 - 1) - \frac{\Theta_1 (r_1 - l_1)}{l_1 - 1} - m_1} (|u_\lambda|) + (\Phi_1^{-1}(1))^{\Theta_1 l_1 (\beta_1 - 1) - \frac{\Theta_1 (r_1 - l_1)}{l_1 - 1} - m_1} \right) |u_\lambda|^{m_1} dx \right)^{\frac{l_1 - 1}{\Theta_1 l_1}} \\
& + \lambda M_7 \int_{\Omega} |u_\lambda|^{r_1 - l_1} |w_k|^{l_1} dx \\
< & \lambda M_8 2^{r_2 - 1 - \frac{(m_2 + 1)(l_1 - 1)(\Theta_1 - 1)}{\Theta_1 l_1}} 2^{(l_1 - 1)(\beta_1 - 1) - \frac{r_1 - l_1}{l_1} - \frac{(m_1 + 1)(l_1 - 1)}{\Theta_1 l_1}} \left(\int_{\mathbb{R}^N} |w_k|^{l_1} |u_\lambda|^{r_1 - l_1} dx \right)^{\frac{1}{l_1}} \\
& \times \left(\int_{\mathbb{R}^N} \left(\Phi_2(|v_\lambda|) |v_\lambda|^{m_2} + (\Phi_2^{-1}(1))^{\frac{\Theta_1 l_1 (r_2 - 1)}{(\Theta_1 - 1)(l_1 - 1)} - m_2} |v_\lambda|^{m_2} \right) dx \right)^{\frac{(l_1 - 1)(\Theta_1 - 1)}{\Theta_1 l_1}} \\
& \times \left(\int_{\mathbb{R}^N} \left(\Phi_1(|u_\lambda|) |u_\lambda|^{m_1} + (\Phi_1^{-1}(1))^{\Theta_1 l_1 (\beta_1 - 1) - \frac{\Theta_1 (r_1 - l_1)}{l_1 - 1} - m_1} |u_\lambda|^{m_1} \right) dx \right)^{\frac{l_1 - 1}{\Theta_1 l_1}} \\
& + \lambda M_7 \int_{\Omega} |u_\lambda|^{r_1 - l_1} |w_k|^{l_1} dx \\
= & \lambda M_8 2^{r_2 - 1 - \frac{(m_2 + 1)(l_1 - 1)(\Theta_1 - 1)}{\Theta_1 l_1}} 2^{(l_1 - 1)(\beta_1 - 1) - \frac{r_1 - l_1}{l_1} - \frac{(m_1 + 1)(l_1 - 1)}{\Theta_1 l_1}} \left(\int_{\mathbb{R}^N} |w_k|^{l_1} |u_\lambda|^{r_1 - l_1} dx \right)^{\frac{1}{l_1}} \\
& \times \left(\int_{\mathbb{R}^N} \Phi_2(|v_\lambda|) |v_\lambda|^{m_2} dx + \int_{\mathbb{R}^N} (\Phi_2^{-1}(1))^{\frac{\Theta_1 l_1 (r_2 - 1)}{(\Theta_1 - 1)(l_1 - 1)} - m_2} |v_\lambda|^{m_2} dx \right)^{\frac{(l_1 - 1)(\Theta_1 - 1)}{\Theta_1 l_1}} \\
& \times \left(\int_{\mathbb{R}^N} \Phi_1(|u_\lambda|) |u_\lambda|^{m_1} dx + \int_{\mathbb{R}^N} (\Phi_1^{-1}(1))^{\Theta_1 l_1 (\beta_1 - 1) - \frac{\Theta_1 (r_1 - l_1)}{l_1 - 1} - m_1} |u_\lambda|^{m_1} dx \right)^{\frac{l_1 - 1}{\Theta_1 l_1}} \\
& + \lambda M_7 \int_{\mathbb{R}^N} |u_\lambda|^{r_1 - l_1} |w_k|^{l_1} dx \\
\leq & \lambda M_8 2^{r_2 - 1 - \frac{(m_2 + 1)(l_1 - 1)(\Theta_1 - 1)}{\Theta_1 l_1}} 2^{(l_1 - 1)(\beta_1 - 1) - \frac{r_1 - l_1}{l_1} - \frac{(m_1 + 1)(l_1 - 1)}{\Theta_1 l_1}} \left(\int_{\mathbb{R}^N} |w_k|^{l_1} |u_\lambda|^{r_1 - l_1} dx \right)^{\frac{1}{l_1}} \\
& \times \left(\left(\int_{\mathbb{R}^N} \Phi_2^{\frac{\sigma_1}{\sigma_1 - 1}} (|v_\lambda|) dx \right)^{\frac{\sigma_1 - 1}{\sigma_1}} \left(\int_{\mathbb{R}^N} |v_\lambda|^{m_2 \sigma_1} dx \right)^{\frac{1}{\sigma_1}} \right. \\
& + \left. \int_{\mathbb{R}^N} (\Phi_2^{-1}(1))^{\frac{\Theta_1 l_1 (r_2 - 1)}{(\Theta_1 - 1)(l_1 - 1)} - m_2} |v_\lambda|^{m_2} dx \right)^{\frac{(l_1 - 1)(\Theta_1 - 1)}{\Theta_1 l_1}} \\
& \times \left(\left(\int_{\mathbb{R}^N} \Phi_1^{\frac{\sigma_2}{\sigma_2 - 1}} (|u_\lambda|) dx \right)^{\frac{\sigma_2 - 1}{\sigma_2}} \left(\int_{\mathbb{R}^N} |u_\lambda|^{m_1 \sigma_2} dx \right)^{\frac{1}{\sigma_2}} \right. \\
& + \left. \int_{\mathbb{R}^N} (\Phi_1^{-1}(1))^{\Theta_1 l_1 (\beta_1 - 1) - \frac{\Theta_1 (r_1 - l_1)}{l_1 - 1} - m_1} |u_\lambda|^{m_1} dx \right)^{\frac{l_1 - 1}{\Theta_1 l_1}} \\
& + \lambda M_7 \int_{\mathbb{R}^N} |u_\lambda|^{r_1 - l_1} |w_k|^{l_1} dx \\
\leq & \lambda M_8 2^{r_2 - 1 - \frac{(m_2 + 1)(l_1 - 1)(\Theta_1 - 1)}{\Theta_1 l_1}} 2^{(l_1 - 1)(\beta_1 - 1) - \frac{r_1 - l_1}{l_1} - \frac{(m_1 + 1)(l_1 - 1)}{\Theta_1 l_1}} \left(\int_{\mathbb{R}^N} |w_k|^{l_1} |u_\lambda|^{r_1 - l_1} dx \right)^{\frac{1}{l_1}} \\
& \times \left(\left(\int_{\mathbb{R}^N} \Phi_2^{\frac{\sigma_1}{\sigma_1 - 1}} (|v_\lambda|) dx \right)^{\frac{(\sigma_1 - 1)(l_1 - 1)(\Theta_1 - 1)}{\sigma_1 \Theta_1 l_1}} \left(\int_{\mathbb{R}^N} |v_\lambda|^{m_2 \sigma_1} dx \right)^{\frac{(l_1 - 1)(\Theta_1 - 1)}{\sigma_1 \Theta_1 l_1}} \right. \\
& + \left. (\Phi_2^{-1}(1))^{r_2 - 1 - \frac{m_2 (l_1 - 1)(\Theta_1 - 1)}{\Theta_1 l_1}} \left(\int_{\mathbb{R}^N} |v_\lambda|^{m_2} dx \right)^{\frac{(l_1 - 1)(\Theta_1 - 1)}{\Theta_1 l_1}} \right) \\
& \times \left(\left(\int_{\mathbb{R}^N} \Phi_1^{\frac{\sigma_2}{\sigma_2 - 1}} (|u_\lambda|) dx \right)^{\frac{(\sigma_2 - 1)(l_1 - 1)}{\sigma_2 \Theta_1 l_1}} \left(\int_{\mathbb{R}^N} |u_\lambda|^{m_1 \sigma_2} dx \right)^{\frac{l_1 - 1}{\sigma_2 \Theta_1 l_1}} \right.
\end{aligned}$$

$$\begin{aligned}
& + (\Phi_1^{-1}(1))^{(\beta_1-1)(l_1-1) - \frac{r_1-l_1}{l_1} - \frac{m_1(l_1-1)}{\Theta_1 l_1}} \left(\int_{\mathbb{R}^N} |u_\lambda|^{m_1} dx \right)^{\frac{l_1-1}{\Theta_1 l_1}} \\
& + \lambda M_7 \int_{\mathbb{R}^N} |u_\lambda|^{r_1-l_1} |w_k|^{l_1} dx \\
\leq & \lambda M_8 2^{r_2-1 - \frac{(m_2+1)(l_1-1)(\Theta_1-1)}{\Theta_1 l_1}} 2^{(l_1-1)(\beta_1-1) - \frac{r_1-l_1}{l_1} - \frac{(m_1+1)(l_1-1)}{\Theta_1 l_1}} \left(\int_{\mathbb{R}^N} |w_k|^{l_1} |u_\lambda|^{r_1-l_1} dx \right)^{\frac{1}{l_1}} \\
& \times C_{0,2}^{\frac{m_2(l_1-1)(\Theta_1-1)}{\Theta_1 l_1}} \|v_\lambda\|_{\Phi_2}^{\frac{m_2(l_1-1)(\Theta_1-1)}{\Theta_1 l_1}} \left((\Phi_2^{-1}(1))^{r_2-1 - \frac{m_2(l_1-1)(\Theta_1-1)}{\Theta_1 l_1}} \right. \\
& \left. + \max \left\{ \|v_\lambda\|_{\Phi_2}^{\frac{l_2(\sigma_1-1)(l_1-1)(\Theta_1-1)}{\sigma_1 \Theta_1 l_1}}, \|v_\lambda\|_{\Phi_2}^{\frac{m_2(\sigma_1-1)(l_1-1)(\Theta_1-1)}{\sigma_1 \Theta_1 l_1}} \right\} \right) \\
& \times C_{0,1}^{\frac{m_1(l_1-1)}{\Theta_1 l_1}} \|u_\lambda\|_{\Phi_1}^{\frac{m_1(l_1-1)}{\Theta_1 l_1}} \left((\Phi_1^{-1}(1))^{(\beta_1-1)(l_1-1) - \frac{r_1-l_1}{l_1} - \frac{m_1(l_1-1)}{\Theta_1 l_1}} \right. \\
& \left. + \max \left\{ \|u_\lambda\|_{\Phi_1}^{\frac{l_1(\sigma_2-1)(l_1-1)}{\sigma_2 \Theta_1 l_1}}, \|u_\lambda\|_{\Phi_1}^{\frac{m_1(\sigma_2-1)(l_1-1)}{\sigma_2 \Theta_1 l_1}} \right\} \right) \\
& + \lambda M_7 \int_{\mathbb{R}^N} |u_\lambda|^{r_1-l_1} |w_k|^{l_1} dx \\
= & \lambda M_8 C_0 C_{0,2}^{\frac{m_2(l_1-1)(\Theta_1-1)}{\Theta_1 l_1}} \|v_\lambda\|_{\Phi_2}^{\frac{m_2(l_1-1)(\Theta_1-1)}{\Theta_1 l_1}} C_{0,1}^{\frac{m_1(l_1-1)}{\Theta_1 l_1}} \|u_\lambda\|_{\Phi_1}^{\frac{m_1(l_1-1)}{\Theta_1 l_1}} \\
& \times \left(\max \left\{ \|v_\lambda\|_{\Phi_2}^{\frac{l_2(\sigma_1-1)(l_1-1)(\Theta_1-1)}{\sigma_1 \Theta_1 l_1}}, \|v_\lambda\|_{\Phi_2}^{\frac{m_2(\sigma_1-1)(l_1-1)(\Theta_1-1)}{\sigma_1 \Theta_1 l_1}} \right\} + (\Phi_2^{-1}(1))^{r_2-1 - \frac{m_2(l_1-1)(\Theta_1-1)}{\Theta_1 l_1}} \right) \\
& \times \left(\max \left\{ \|u_\lambda\|_{\Phi_1}^{\frac{l_1(\sigma_2-1)(l_1-1)}{\sigma_2 \Theta_1 l_1}}, \|u_\lambda\|_{\Phi_1}^{\frac{m_1(\sigma_2-1)(l_1-1)}{\sigma_2 \Theta_1 l_1}} \right\} + (\Phi_1^{-1}(1))^{(\beta_1-1)(l_1-1) - \frac{r_1-l_1}{l_1} - \frac{m_1(l_1-1)}{\Theta_1 l_1}} \right) \\
& \times \left(\int_{\mathbb{R}^N} |w_k|^{l_1} |u_\lambda|^{r_1-l_1} dx \right)^{\frac{1}{l_1}} + \lambda M_7 \int_{\mathbb{R}^N} |u_\lambda|^{r_1-l_1} |w_k|^{l_1} dx \\
\leq & \lambda M_7 \int_{\mathbb{R}^N} |u_\lambda|^{r_1-l_1} |w_k|^{l_1} dx + \lambda M_8 C_1 \left(\int_{\mathbb{R}^N} |w_k|^{l_1} |u_\lambda|^{r_1-l_1} dx \right)^{\frac{1}{l_1}} \tag{3.16}
\end{aligned}$$

for some $\sigma_1 \in \left(1, \frac{l_2^*}{m_2}\right]$, $\sigma_2 \in \left(1, \frac{l_1^*}{m_1}\right]$ and any given $\beta_1 \in I_\beta := \left(1 + \frac{r_1-l_1}{l_1(l_1-1)} + \frac{m_1+1}{\Theta_1 l_1}, +\infty\right)$, where

$$\Phi_1^{\Theta_1 l_1 (\beta_1-1) - \frac{\Theta_1(r_1-l_1)}{l_1-1} - m_1}(|u_\lambda|) < \Phi_1(|u_\lambda|)$$

which holds by (2.2). Moreover, for the validity of the interval of Θ_1 , we need the following restriction

$$r_2 > 1 + \frac{(1+m_2)(l_1-1)(\Theta_1-1)}{l_1 \Theta_1}$$

to make

$$\Phi_2^{\frac{\Theta_1 l_1 (r_2-1)}{(\Theta_1-1)(l_1-1)} - m_2}(|v_\lambda|) < \Phi_2(|v_\lambda|)$$

hold by (2.2). In addition, the range of σ_1 and σ_2 is determined by (2.4), and we assume that

$$\begin{aligned}
C_1 &= C_0 C_{0,1}^{\frac{m_1(l_1-1)}{\Theta_1 l_1}} C_{0,2}^{\frac{m_2(l_1-1)(\Theta_1-1)}{\Theta_1 l_1}} \|v_\lambda\|_{\Phi_2}^{\frac{m_2(l_1-1)(\Theta_1-1)}{\Theta_1 l_1}} \|u_\lambda\|_{\Phi_1}^{\frac{m_1(l_1-1)}{\Theta_1 l_1}} \\
& \times \left(\max \left\{ \|v_\lambda\|_{\Phi_2}^{\frac{l_2(\sigma_1-1)(l_1-1)(\Theta_1-1)}{\sigma_1 \Theta_1 l_1}}, \|v_\lambda\|_{\Phi_2}^{\frac{m_2(\sigma_1-1)(l_1-1)(\Theta_1-1)}{\sigma_1 \Theta_1 l_1}} \right\} + (\Phi_2^{-1}(1))^{r_2-1 - \frac{m_2(l_1-1)(\Theta_1-1)}{\Theta_1 l_1}} \right) \\
& \times \left(\max \left\{ \|u_\lambda\|_{\Phi_1}^{\frac{l_1(\sigma_2-1)(l_1-1)}{\sigma_2 \Theta_1 l_1}}, \|u_\lambda\|_{\Phi_1}^{\frac{m_1(\sigma_2-1)(l_1-1)}{\sigma_2 \Theta_1 l_1}} \right\} + (\Phi_1^{-1}(1))^{(\beta_1-1)(l_1-1) - \frac{r_1-l_1}{l_1} - \frac{m_1(l_1-1)}{\Theta_1 l_1}} \right),
\end{aligned}$$

where $C_0 = 2^{r_2-1+(l_1-1)(\beta_1-1)-\frac{(m_2+1)(l_1-1)(\Theta_1-1)+(m_1+1)(l_1-1)}{\Theta_1 l_1^*}-\frac{r_1-l_1}{l_1^*}}$.

By Gagliard-Nirenberg-Sobolev inequality, (3.16) and Hölder inequality, we have

$$\begin{aligned}
\left(\int_{\mathbb{R}^N} |w_k|^{l_1^*} dx \right)^{\frac{l_1}{l_1^*}} &\leq D_1 \int_{\mathbb{R}^N} |\nabla w_k|^{l_1} dx \\
&\leq \lambda D_2 \beta_1^{l_1} \int_{\mathbb{R}^N} |u_\lambda|^{r_1-l_1} |w_k|^{l_1} dx + \lambda D_3 \beta_1^{l_1} \left(\int_{\mathbb{R}^N} |u_\lambda|^{r_1-l_1} |w_k|^{l_1} dx \right)^{\frac{1}{l_1}} \\
&\leq \lambda D_2 \beta_1^{l_1} \left(\int_{\mathbb{R}^N} |u_\lambda|^{l_1^*} dx \right)^{\frac{r_1-l_1}{l_1^*}} \left(\int_{\mathbb{R}^N} |w_k|^{\frac{l_1 l_1^*}{l_1^*-r_1+l_1}} dx \right)^{\frac{l_1^*-r_1+l_1}{l_1^*}} \\
&\quad + \lambda D_3 \beta_1^{l_1} \left(\int_{\mathbb{R}^N} |u_\lambda|^{l_1^*} dx \right)^{\frac{r_1-l_1}{l_1 l_1^*}} \left(\int_{\mathbb{R}^N} |w_k|^{\frac{l_1 l_1^*}{l_1^*-r_1+l_1}} dx \right)^{\frac{l_1^*-r_1+l_1}{l_1 l_1^*}} \\
&\leq \lambda D_2 \beta_1^{l_1} \left(\int_{\mathbb{R}^N} |u_\lambda|^{l_1^*} dx \right)^{\frac{r_1-l_1}{l_1^*}} \left(\int_{\mathbb{R}^N} |u_\lambda|^{\frac{\beta_1 l_1 l_1^*}{l_1^*-r_1+l_1}} dx \right)^{\frac{l_1^*-r_1+l_1}{l_1^*}} \\
&\quad + \lambda D_3 \beta_1^{l_1} \left(\int_{\mathbb{R}^N} |u_\lambda|^{l_1^*} dx \right)^{\frac{r_1-l_1}{l_1 l_1^*}} \left(\int_{\mathbb{R}^N} |u_\lambda|^{\frac{\beta_1 l_1 l_1^*}{l_1^*-r_1+l_1}} dx \right)^{\frac{l_1^*-r_1+l_1}{l_1 l_1^*}} \\
&\leq \lambda D_2 \beta_1^{l_1} \|u_\lambda\|_{1, \Phi_1}^{r_1-l_1} \left(\int_{\mathbb{R}^N} |u_\lambda|^{\frac{\beta_1 l_1 l_1^*}{l_1^*-r_1+l_1}} dx \right)^{\frac{l_1^*-r_1+l_1}{l_1^*}} \\
&\quad + \lambda D_3 \beta_1^{l_1} \|u_\lambda\|_{1, \Phi_1}^{\frac{r_1-l_1}{l_1}} \left(\int_{\mathbb{R}^N} |u_\lambda|^{\frac{\beta_1 l_1 l_1^*}{l_1^*-r_1+l_1}} dx \right)^{\frac{l_1^*-r_1+l_1}{l_1 l_1^*}}.
\end{aligned}$$

Note that $r_1 \in \left(\max \left\{ m_1, m_2, 1 + \frac{(1+m_1)(l_2-1)(\Theta_2-1)}{l_2 \Theta_2} \right\}, \min \left\{ l_1^*, 1 + \frac{l_1^*(l_1-1)}{l_1} - \frac{(l_1-1)(1+m_1)}{\Theta_1 l_1} \right\} \right)$. Let $\beta_1 = 1 + \frac{l_1^*-r_1}{l_1}$. Then $\beta_1 \in I_\beta$. Thus $\frac{\beta_1 l_1 l_1^*}{l_1^*-r_1+l_1} = l_1^*$. Thus, we have

$$\left(\int_{\mathbb{R}^N} |w_k|^{l_1^*} dx \right)^{\frac{l_1}{l_1^*}} \leq \lambda D_2 \beta_1^{l_1} \|u_\lambda\|_{1, \Phi_1}^{r_1-l_1} \|u_\lambda\|_{\beta_1 \alpha_1^*}^{l_1 \beta_1} + \lambda D_3 \beta_1^{l_1} \|u_\lambda\|_{1, \Phi_1}^{\frac{r_1-l_1}{l_1}} \|u_\lambda\|_{\beta_1 \alpha_1^*}^{\beta_1}, \quad (3.17)$$

where $\alpha_1^* = \frac{l_1 l_1^*}{l_1^*-r_1+l_1}$. Lemma 3.3 implies that there exists sufficient large $\Lambda_3 > \Lambda_0$ such that $\|u_\lambda\|_{1, \Phi_1} < \min\{1, \frac{1}{C_{0,1}}\}$ for all $\lambda > \Lambda_3$ and then (2.4) implies that $\|u_\lambda\|_{\beta_1 \alpha_1^*} = \|u_\lambda\|_{l_1^*} < \min\{C_{0,1}, 1\}$. Since $r_1 - l_1 > \frac{r_1-l_1}{l_1}$ and $l_1 \beta_1 > \beta_1$. Hence, by (3.17), we have

$$\left(\int_{\mathbb{R}^N} |w_k|^{l_1^*} dx \right)^{\frac{l_1}{l_1^*}} \leq \lambda (D_2 + D_3) \beta_1^{l_1} \|u_\lambda\|_{1, \Phi_1}^{\frac{r_1-l_1}{l_1}} \|u_\lambda\|_{\beta_1 \alpha_1^*}^{\beta_1}. \quad (3.18)$$

Then it follows from the definition of w_k , Fatou's Lemma and (3.18) that

$$\|u_\lambda\|_{\beta_1 l_1^*} \leq \left(\lambda (D_2 + D_3) \beta_1^{l_1} \|u_\lambda\|_{1, \Phi_1}^{\frac{r_1-l_1}{l_1}} \right)^{\frac{1}{l_1 \beta_1}} \|u_\lambda\|_{\beta_1 \alpha_1^*}^{\frac{1}{l_1}}.$$

Next, we start the iteration process. For each $n = 0, 1, 2, \dots$, we define $\beta_1^{(n+1)} = \frac{l_1^*}{\alpha_1^*} \beta_1^{(n)}$, where $\beta_1^{(0)} = \beta_1$.

$$\|u_\lambda\|_{\beta_1^{(1)} l_1^*} \leq \left(\lambda (D_2 + D_3) \beta_1^{(1) l_1} \|u_\lambda\|_{1, \Phi_1}^{\frac{r_1-l_1}{l_1}} \right)^{\frac{1}{l_1 \beta_1^{(1)}}} \|u_\lambda\|_{\beta_1^{(1)} \alpha_1^*}^{\frac{1}{l_1}}$$

$$\begin{aligned}
&\leq \left(\lambda(D_2 + D_3) \beta_1^{(1)l_1} \|u_\lambda\|_{1, \Phi_1}^{\frac{r_1 - l_1}{l_1}} \right)^{\frac{1}{l_1 \beta_1^{(1)}}} \left(\left(\lambda(D_2 + D_3) \beta_1^{l_1} \|u_\lambda\|_{1, \Phi_1}^{\frac{r_1 - l_1}{l_1}} \right)^{\frac{1}{l_1 \beta_1}} \|u_\lambda\|_{\beta_1 \alpha_1^*}^{\frac{1}{l_1}} \right)^{\frac{1}{l_1}} \\
&= (\lambda(D_2 + D_3) \|u_\lambda\|_{1, \Phi_1}^{\frac{r_1 - l_1}{l_1}})^{\frac{1}{l_1 \beta_1^{(1)}} + \frac{1}{l_1^2 \beta_1}} \beta_1^{\frac{1}{l_1 \beta_1}} \beta_1^{(1) \frac{1}{\beta_1^{(1)}}} \|u_\lambda\|_{\beta_1 \alpha_1^*}^{\frac{1}{l_1^2}}
\end{aligned}$$

and then

$$\begin{aligned}
\|u_\lambda\|_{\beta_1^{(2)} l_1^*} &\leq \left(\lambda(D_2 + D_3) \beta_1^{(2)l_1} \|u_\lambda\|_{1, \Phi_1}^{\frac{r_1 - l_1}{l_1}} \right)^{\frac{1}{l_1 \beta_1^{(2)}}} \|u_\lambda\|_{\beta_1^{(2)} \alpha_1^*}^{\frac{1}{l_1}} \\
&\leq \left((\lambda(D_2 + D_3) \|u_\lambda\|_{1, \Phi_1}^{\frac{r_1 - l_1}{l_1}})^{\frac{1}{l_1 \beta_1^{(1)}} + \frac{1}{l_1^2 \beta_1}} \beta_1^{\frac{1}{l_1 \beta_1}} \beta_1^{(1) \frac{1}{\beta_1^{(1)}}} \|u_\lambda\|_{\beta_1 \alpha_1^*}^{\frac{1}{l_1^2}} \right)^{\frac{1}{l_1}} \\
&\quad \left(\lambda(D_2 + D_3) \beta_1^{(2)l_1} \|u_\lambda\|_{1, \Phi_1}^{\frac{r_1 - l_1}{l_1}} \right)^{\frac{1}{l_1 \beta_1^{(2)}}} \\
&= \left(\lambda(D_2 + D_3) \|u_\lambda\|_{1, \Phi_1}^{\frac{r_1 - l_1}{l_1}} \right)^{\frac{1}{l_1 \beta_1^{(2)}} + \frac{1}{l_1^2 \beta_1^{(1)}} + \frac{1}{l_1^3 \beta_1}} \beta_1^{\frac{1}{l_1^2 \beta_1}} \beta_1^{(1) \frac{1}{l_1 \beta_1^{(1)}}} \beta_1^{(2) \frac{1}{\beta_1^{(2)}}} \|u_\lambda\|_{\beta_1 \alpha_1^*}^{\frac{1}{l_1^3}}.
\end{aligned}$$

Repeating such process, we obtain that

$$\begin{aligned}
&\|u_\lambda\|_{\beta_1^{(n)} l_1^*} \\
&\leq \left(\lambda(D_2 + D_3) \|u_\lambda\|_{1, \Phi_1}^{\frac{r_1 - l_1}{l_1}} \right)^{\sum_{i=0}^n \frac{1}{l_1^{n+1-i} \beta_1^{(i)}}} \prod_{i=0}^n \beta_1^{(i) \frac{1}{l_1^{n-i} \beta_1^{(i)}}} \|u_\lambda\|_{\beta_1 \alpha_1^*}^{\frac{1}{l_1^{n+1}}} \\
&= D_4 \left(\lambda(D_2 + D_3) \|u_\lambda\|_{1, \Phi_1}^{\frac{r_1 - l_1}{l_1}} \right)^{\sum_{i=0}^n \frac{1}{\beta_1 l_1 l_1^{n-i} \left(\frac{l_1^*}{\alpha_1^*} \right)^i}} \|u_\lambda\|_{\beta_1 \alpha_1^*}^{\frac{1}{l_1^{n+1}}}, \tag{3.19}
\end{aligned}$$

$$\text{where } D_4 = \beta_1^{\frac{1}{\beta_1} \sum_{i=0}^n \left(\frac{1}{l_1} \right)^{n-i} \left(\frac{\alpha_1^*}{l_1^*} \right)^i} \left(\frac{l_1^*}{\alpha_1^*} \right)^{\frac{1}{\beta_1} \sum_{i=0}^n i \left(\frac{1}{l_1} \right)^{n-i} \left(\frac{\alpha_1^*}{l_1^*} \right)^i}.$$

Notice that

$$\alpha_1^* < l_1^*, \quad \lim_{n \rightarrow \infty} \sum_{i=0}^n \frac{1}{\beta_1 l_1 l_1^{n-i} \left(\frac{l_1^*}{\alpha_1^*} \right)^i} = 0, \quad \lim_{n \rightarrow \infty} \sum_{i=0}^n i \left(\frac{1}{l_1} \right)^{n-i} \left(\frac{\alpha_1^*}{l_1^*} \right)^i = 0, \quad \lim_{n \rightarrow \infty} \frac{1}{l_1^{n+1}} = 0.$$

So (3.19) implies that $\|u_\lambda\|_\infty \leq 1$ for all $\lambda > \Lambda_3$.

Similarly, there exists $\Lambda_4 > \Lambda_0$ such that $\|v_\lambda\|_\infty \leq 1$ for all $\lambda > \Lambda_4$. Let $\Lambda_* = \max\{\Lambda_3, \Lambda_4\}$. Then we complete the proof. \square

Proof of Theorem 1.1. By Lemma 3.4, for each $\lambda > \Lambda_*$, we have

$$\|(u_\lambda, v_\lambda)\|_\infty \leq \|u_\lambda\|_\infty + \|v_\lambda\|_\infty \leq 2$$

which implies that $\tilde{F}(x, u_\lambda, v_\lambda) = F(x, u_\lambda, v_\lambda)$ for all $x \in \mathbb{R}^N$. Hence, (u_λ, v_λ) is a nontrivial weak solution of system (1.5) and Lemma 3.3 implies that $\|u_\lambda\|_{1, \Phi_1} \rightarrow 0$ and $\|v_\lambda\|_{1, \Phi_2} \rightarrow 0$ as $\lambda \rightarrow \infty$. \square

4. Example

$$\begin{cases} -\operatorname{div}[(4|\nabla u|^2 + 5|\nabla u|^3)\nabla u] + \left(1 + \sum_{i=1}^6 \cos^2 \pi x_i\right) (4|u|^2 + 5|u|^3)u \\ = \lambda F_u(x, u, v), \quad x \in \mathbb{R}^6, \\ -\operatorname{div}\left[\left(4|\nabla v|^2 \log(2 + |\nabla v|) + \frac{|\nabla v|^3}{1 + |\nabla v|}\right) \nabla v\right] + \left(1 + \sum_{i=1}^6 \sin^2 \pi x_i\right) \left(4|v|^2 \log(1 + |v|) + \frac{|v|^3}{1 + |v|}\right) v \\ = \lambda F_v(x, u, v), \quad x \in \mathbb{R}^6, \end{cases} \quad (4.1)$$

where

$$F(x, t, s) = \sigma(t, s)b(x) \left(|t|^{\frac{17}{2}} + |s|^{\frac{17}{2}} + |t|^7|s|^7\right) + (1 - \sigma(t, s))b(x) (|t|^3 + |s|^3) \quad (4.2)$$

for all $(x, t, s) \in \mathbb{R}^N \times \mathbb{R} \times \mathbb{R}$ with $\sigma(t, s)$ defined by (1.6) and $b(x) = \left(1 + \sum_{i=1}^6 \cos^2 \pi x_i\right)$ (or $b(x) \equiv 1$). So,

$$F(x, t, s) = \begin{cases} b(x) \left(|t|^{\frac{17}{2}} + |s|^{\frac{17}{2}} + |t|^7|s|^7\right), & \text{if } |(t, s)| \leq 4, \\ \sin \frac{\pi(t^2 + s^2 - 64)^2}{4608} b(x) \left(|t|^{\frac{17}{2}} + |s|^{\frac{17}{2}} + |t|^7|s|^7\right) \\ + \left(1 - \sin \frac{\pi(t^2 + s^2 - 64)^2}{4608}\right) b(x) (|t|^3 + |s|^3), & \text{if } 4 < |(t, s)| \leq 8, \\ b(x) (|t|^3 + |s|^3), & \text{if } |(t, s)| > 8, \end{cases}$$

and we also draw the figure of F (see figure (e)-(h) below).

Let $N = 6$, $\phi_1(t) = 4|t|^2 + 5|t|^3$ and $\phi_2(t) = 4|t|^2 \log(2 + |t|) + \frac{|t|^3}{1 + |t|}$. Then $\phi_i (i = 1, 2)$ satisfy (ϕ_1) -(ϕ_3) and $l_1 = l_2 = 4$, $m_1 = m_2 = 5$ (see [25]). So, $l_1^* = l_2^* = 12$.


Let $V_1(x) = 1 + \sum_{i=1}^6 \cos^2 \pi x_i$ and $V_2(x) = 1 + \sum_{i=1}^6 \sin^2 \pi x_i$ for all $(x, t, s) \in \mathbb{R}^N \times \mathbb{R} \times \mathbb{R}$. Then it is obvious that V_i satisfies (V0) and (V1).

Let


$$\begin{aligned} A(t, s) &:= \frac{t\pi(t^2 + s^2 - 64)}{1152} \cos \frac{\pi(t^2 + s^2 - 64)^2}{4608} b(x) \left(|t|^{\frac{17}{2}} + |s|^{\frac{17}{2}} + |t|^7|s|^7\right) \\ &+ \sin \frac{\pi(t^2 + s^2 - 64)^2}{4608} b(x) \left(\frac{17}{2}|t|^{\frac{13}{2}}t + 7|t|^5|s|^7t\right) \\ &- \frac{t\pi(t^2 + s^2 - 64)}{1152} \cos \frac{\pi(t^2 + s^2 - 64)^2}{4608} b(x) (|t|^3 + |s|^3) + 3 \left(1 - \sin \frac{\pi(t^2 + s^2 - 64)^2}{4608}\right) a(x)|t|t. \end{aligned}$$

and


$$\begin{aligned} B(t, s) &:= \frac{s\pi(t^2 + s^2 - 64)}{1152} \cos \frac{\pi(t^2 + s^2 - 64)^2}{4608} b(x) \left(|t|^{\frac{17}{2}} + |s|^{\frac{17}{2}} + |t|^7|s|^7\right) \\ &+ \sin \frac{\pi(t^2 + s^2 - 64)^2}{4608} b(x) \left(\frac{17}{2}|s|^{\frac{13}{2}}s + 7|s|^5|t|^7s\right) \end{aligned}$$




(e) $(t, s) \in ((-0.5, 0.5), (-0.5, 0.5))$



(f) $(t, s) \in ((-4, 4), (-4, 4))$



(g) $(t, s) \in ((-8, 8), (-8, 8))$



(h) $(t, s) \in ((-2000, 2000), (-2000, 2000))$

$$-\frac{s\pi(t^2 + s^2 - 64)}{1152} \cos \frac{\pi(t^2 + s^2 - 64)^2}{4608} b(x) (|t|^3 + |s|^3) + 3 \left(1 - \sin \frac{\pi(t^2 + s^2 - 64)^2}{4608} \right) b(x) |s|s.$$

It is easy to see that F satisfies (F0) and

$$F_t(x, t, s) = \begin{cases} b(x) \left(\frac{17}{2} |t|^{\frac{13}{2}} t + 7 |t|^5 |s|^7 t \right), & \text{if } |(t, s)| \leq 4, \\ A(t, s), & \text{if } 4 < |(t, s)| \leq 8 \\ 3b(x) |t|t, & \text{if } |(t, s)| > 8, \end{cases}$$

$$F_s(x, t, s) = \begin{cases} b(x) \left(\frac{17}{2} |s|^{\frac{13}{2}} s + 7 |s|^5 |t|^7 s \right), & \text{if } |(t, s)| \leq 4, \\ B(t, s), & \text{if } 4 < |(t, s)| \leq 8 \\ 3b(x) |s|s, & \text{if } |(t, s)| > 8. \end{cases}$$

Thus, we get

$$\begin{aligned} F(x, t, s) &= b(x) \left(|t|^{\frac{17}{2}} + |s|^{\frac{17}{2}} + |t|^7 |s|^7 \right) \\ &\geq \begin{cases} |t|^9 + |s|^9, & \text{if } |(t, s)| < 1, \\ \frac{1}{16} (|t|^2 + |s|^2) (|t|^7 + |s|^7), & \text{if } 1 \leq |(t, s)| \leq 4, \end{cases} \\ &\geq \frac{1}{16} |t|^9 + \frac{1}{16} |s|^9, \quad \text{for all } x \in \mathbb{R}^N \text{ and } |(t, s)| \leq 4. \end{aligned}$$

Then (F1) holds with $k_1 = k_2 = 9 \in (m_1, l_1^*) = (m_2, l_2^*) = (5, 12)$, $M_1 = M_2 = \frac{1}{16}$. We also have

$$\begin{aligned} |F_t(x, t, s)| &= b(x) \left(\frac{17}{2} |t|^{\frac{15}{2}} + 7 |t|^6 |s|^7 \right) \\ &\leq \begin{cases} \left(\frac{17}{2} + 7 \right) \times 7 (|t|^6 + |s|^6), & \text{if } |(t, s)| < 1 \\ \left(\frac{17}{2} + 7 \right) \times 7 (t^2 + s^2)^4 (|t|^6 + |s|^6), & \text{if } 1 \leq |(t, s)| \leq 4 \end{cases} \\ &\leq 2^{41} (|t|^6 + |s|^6), \quad \text{for all } x \in \mathbb{R}^N \text{ and } |(t, s)| \leq 4. \end{aligned}$$

Similarly,

$$|F_s(x, t, s)| \leq 2^{41} (|t|^6 + |s|^6), \quad \text{for all } x \in \mathbb{R}^N \text{ and } |(t, s)| \leq 4.$$

Then (F2) holds with $M_3 = M_4 = 2^{41}$ and $r_1 = 7 \in \left(\max \left\{ m_1, m_2, 1 + \frac{(1+m_1)(l_2-1)(\Theta_2-1)}{l_2\Theta_2} \right\}, 1 + \frac{l_1^*(l_1-1)}{l_1} - \frac{(l_1-1)(1+m_1)}{\Theta_1 l_1} \right) = \left(\max \left\{ 5, 1 + \frac{9(\Theta_2-1)}{2\Theta_2} \right\}, 10 - \frac{9}{2\Theta_1} \right)$ and $r_2 = 7 \in \left(\max \left\{ m_1, m_2, 1 + \frac{(1+m_2)(l_1-1)(\Theta_1-1)}{l_1\Theta_1} \right\}, 1 + \frac{l_2^*(l_2-1)}{l_2} - \frac{(l_2-1)(1+m_2)}{\Theta_2 l_2} \right) = \left(\max \left\{ 5, 1 + \frac{9(\Theta_1-1)}{2\Theta_1} \right\}, 10 - \frac{9}{2\Theta_2} \right)$ for some $\Theta_1, \Theta_2 > 1$. In particular, taking $\Theta_1 = \Theta_2 = 6$, it is easy to see that $r_1 = r_2 = 7 \in \left(5, \frac{37}{4} \right)$. Note that

$$b(x) \left(|t|^{\frac{17}{2}} + |s|^{\frac{17}{2}} + |t|^7 |s|^7 \right) \leq b(x) \left(|t|^{\frac{17}{2}} + |s|^{\frac{17}{2}} + \frac{28}{17} |t|^7 |s|^7 \right).$$

So

$$F(x, t, s) \leq \frac{2}{17} t F_t(x, t, s) + \frac{2}{17} s F_s(x, t, s) = \frac{1}{\mu_1} t F_t(x, t, s) + \frac{1}{\mu_2} s F_s(x, t, s)$$

for all $x \in \mathbb{R}^N$ and $|(t, s)| \leq 4$, where $\mu_1 = \mu_2 = \frac{17}{2} > 5 = m_1 = m_2$. Thus, we have verified that system (4.1) satisfies all the conditions of Theorem 1.1. Hence, there exists $\Lambda_* > 0$ such that system (1.5) has a nontrivial solution (u_λ, v_λ) with $\|(u_\lambda, v_\lambda)\|_\infty \leq 2$ for each $\lambda > \Lambda_*$ and $\|(u_\lambda, v_\lambda)\| \rightarrow 0$ as $\lambda \rightarrow \infty$.

Next, we show that (4.2) does not satisfy Theorem A. In fact, for (4.2), we have

$$\begin{aligned} &\frac{1}{\mu_1} t F_t(x, t, s) + \frac{1}{\mu_2} s F_s(x, t, s) \\ &\leq \frac{1}{5} [t F_t(x, t, s) + s F_s(x, t, s)] \\ &\leq \frac{3}{5} b(x) |t|^3 + \frac{3}{5} b(x) |s|^3 \\ &< b(x) |t|^3 + b(x) |s|^3 \end{aligned}$$

for all $x \in \mathbb{R}^N$, $|(t, s)| > 4$ and any $\mu_i \in (5, +\infty)$, $i = 1, 2$, which shows that F does not satisfy (H2) in Theorem A.

5. A result for elliptic equation

Notice that (F1)-(F3) hold for all $|(t, s)| \leq 2$, which are stronger than (f1)-(f3) where $|t| \leq \delta$ for some positive constant δ . This is because we can not obtain an estimate for $\|u\|_\infty$ and $\|v\|_\infty$ like Lemma 2.6 in [20] where $\|u\|_\infty$ grows with $\|u\|$. Instead, we obtain $\|u\|_\infty \leq 1$ and $\|v\|_\infty \leq 1$ (see Lemma 3.4), which are caused by the relation of u and v from the proof of Lemma 3.4. Hence, in this section, for the following quasilinear scalar equation

$$\begin{cases} -\operatorname{div}(\phi(|\nabla u|)\nabla u) + V(x)\phi(|u|)u = \lambda f(x, u), & x \in \mathbb{R}^N, \\ u \in W^{1,\Phi}(\mathbb{R}^N), \end{cases} \quad (5.1)$$

where $N \geq 2$ and $\lambda > 0$, we shall obtain a result similar to Theorem 1 in [8] and Theorem 1.2 in [20]. To be precise, we have the following theorem:

Theorem 5.1. *Assume that ϕ satisfies (ϕ_1) , (ϕ_2) and the following conditions hold:*

(ϕ_3) there exists a positive constant q such that $t^2\phi(|t|) \geq q|t|^l$ for all $t \in \mathbb{R}$;

$(V0)$ $V \in C(\mathbb{R}^N, \mathbb{R}^+)$, $V_\infty := \inf_{\mathbb{R}^N} V(x) > 0$ and V is a 1-periodic function;

$(C1)$ $f \in C(\mathbb{R}^N \times \mathbb{R}, \mathbb{R})$ and f is 1-periodic in $x \in \mathbb{R}^N$;

$(C2)$ there exist $\delta > 0$, $k \in (m, K)$ and $D_1 > 0$ such that

$$F(x, t) \geq D_1|t|^k$$

for all $t \in (-\delta, \delta)$ and $x \in \mathbb{R}^N$, where $K = \min \left\{ l^, \frac{ml-l+l^*}{m} \right\}$ and $F(x, t) = \int_0^t f(x, \xi)d\xi$;*

$(C3)$ there exist $r \in (m, l^)$ and $D_2 > 0$ such that*

$$|f(x, t)| \leq D_2|t|^{r-1}$$

for all $t \in (-\delta, \delta)$ and $x \in \mathbb{R}^N$;

$(C4)$ there exists $\mu > m$ such that

$$0 < F(x, t) \leq \frac{1}{\mu}tf(x, t)$$

for all $x \in \mathbb{R}^N$ and $t \in (-\delta, \delta)$ with $t \neq 0$.

Then there exists $\Lambda_0 > 0$ such that equation (5.1) has at least nontrivial solution u_λ for each $\lambda > \Lambda_0$, $\|u_\lambda\| \rightarrow 0$ and $\|u_\lambda\|_\infty \rightarrow 0$ as $\lambda \rightarrow \infty$.

The proof is similar to Theorem 1.1. The main difference is the result of Moser iteration. Next, we outline the proof. We work on the space $W^{1,\Phi}(\mathbb{R}^N)$ with the norm $\|\cdot\|_{1,\Phi}$. Define the cut-off function $\rho \in C^1(\mathbb{R}, [0, 1])$

$$\rho(t) = \begin{cases} 1, & \text{if } |t| \leq \delta/2, \\ 0, & \text{if } |t| \geq \delta \end{cases}$$

for all $t \in \mathbb{R}$ and $t\rho'(t) \leq 0$. Similar to (3.2)-(3.5), some examples of $\rho(t)$ can also be given, for example,

$$\rho(t) = \begin{cases} 1, & \text{if } |t| < \frac{\delta}{2}, \\ \sin \frac{8\pi(t^2 - \delta^2)^2}{9\delta^4}, & \text{if } \frac{\delta}{2} \leq |t| \leq \delta, \\ 0, & \text{if } |t| > \delta. \end{cases}$$

Let

$$\tilde{F}(x, t) = \rho(t)F(x, t) + (1 - \rho(t))D_3|t|^r.$$

Lemma 5.1. *Assume that (C1)-(C4) hold. Then*

(C1)' $\tilde{F} \in C^1(\mathbb{R}^N \times \mathbb{R}, \mathbb{R})$, \tilde{F} is 1-periodic in $x \in \mathbb{R}^N$ and $\tilde{F}(x, 0) = 0$ for all $x \in \mathbb{R}^N$;

(C2)'

$$0 \leq \tilde{F}(x, t) \leq D_3|t|^r, \quad \text{for all } t \in \mathbb{R} \text{ and } x \in \mathbb{R}^N;$$

(C3)' there exists $D_4 > 0$ such that

$$|\tilde{f}(x, t)| \leq D_4|t|^{r-1}$$

for all $t \in \mathbb{R}$ and $x \in \mathbb{R}^N$;

(C3)'

$$\theta \tilde{F}(x, t) \leq \tilde{f}(x, t)t, \quad \text{for all } t \in \mathbb{R}/\{0\} \text{ and } x \in \mathbb{R}^N,$$

where $\theta = \min\{r, \mu\}$.

Consider the modified problem

$$\begin{cases} -\operatorname{div}(\phi(|\nabla u|)\nabla u) + V(x)\phi(|u|)u = \lambda \tilde{f}(x, u), & x \in \mathbb{R}^N, \\ u \in W^{1, \Phi}(\mathbb{R}^N). \end{cases} \quad (5.2)$$

Define the functional $\tilde{J}_\lambda : W^{1, \Phi}(\mathbb{R}^N) \rightarrow \mathbb{R}$ by

$$\tilde{J}_\lambda(u) = \int_{\mathbb{R}^N} \Phi(|\nabla u|)dx + \int_{\mathbb{R}^N} V(x)\Phi(|u|)dx - \lambda \int_{\mathbb{R}^N} \tilde{F}(x, u)dx.$$

It is easy to see that \tilde{J}_λ is well defined and $\tilde{J}_\lambda \in C^1(W^{1, \Phi}(\mathbb{R}^N), \mathbb{R})$ and

$$\langle \tilde{J}'_\lambda(u), \tilde{u} \rangle = \int_{\mathbb{R}^N} (\phi(|\nabla u|)\nabla u, \nabla \tilde{u})dx + \int_{\mathbb{R}^N} V(x)\phi(|u|)u\tilde{u}dx - \lambda \int_{\mathbb{R}^N} \tilde{F}_u(x, u)\tilde{u}dx.$$

Lemma 5.2. \tilde{J}_λ satisfies the mountain pass geometry, that is,

(i) there exist two positive constants γ, η such that $\tilde{J}_\lambda(u) \geq \eta$ for all $\|u\| = \gamma$;

(ii) there exists $u_0 \in C_0^\infty(\mathbb{R}^N)/\{0\}$ with $u_0 > 0$ and $0 < \|u_0\|_\infty < \frac{\delta}{2}$ such that $\tilde{J}_\lambda(u_0) < 0$.

Proof. Notice that (C1)'-(C2)' imply those conditions of Theorem 1.5 in [2]. Then the proof is completed easily. \square

By Lemma 4.1-Lemma 4.3 in [2] that system (5.1) has a nontrivial solution u_λ such that $\tilde{J}_\lambda(u_\lambda) = c_\lambda$ with

$$c_\lambda := \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} \tilde{J}_\lambda(\gamma(t)),$$

and

$$\Gamma := \{\gamma \in ([0,1], W^{1,\Phi}(\mathbb{R}^N)) : \gamma(0) = (0), \gamma(1) = u_0\}.$$

Lemma 5.3. For each $\lambda > 0$, there exists $C_{**} > 0$ such that

$$\|u_\lambda\|_{1,\Phi} \leq C_{**} \max \left\{ \lambda^{-\frac{1}{k-l}}, \lambda^{-\frac{l}{m(k-l)}} \right\}.$$

Proof. It is easy to obtain the conclusion from the proof of Lemma 3.3.

Lemma 5.4. For each $\lambda > 0$, there exist positive constants C which only depends on q, r, l, N , respectively, such that

$$\|u_\lambda\|_\infty \leq C(\lambda \|u_\lambda\|_{1,\Phi}^{r-l})^{\frac{1}{l^*-r}} \|u_\lambda\|_{1,\Phi}.$$

Proof. The proof is a direct generalization of Lemma 2.6 in [20]. Since u_λ is a critical point of \tilde{J}_λ , we have

$$\int_{\mathbb{R}^N} (\phi(|\nabla u_\lambda|) \nabla u_\lambda, \nabla \tilde{u}) dx + \int_{\mathbb{R}^N} V(x) \phi(|u_\lambda|) u_\lambda \tilde{u} dx = \lambda \int_{\mathbb{R}^N} \tilde{f}(x, u_\lambda) \tilde{u} dx \quad (5.3)$$

for all $\tilde{u} \in W^{1,\Phi}(\mathbb{R}^N)$. Without loss of generality, for each $k > 0$, define

$$u_k = \begin{cases} u_\lambda, & \text{if } u_\lambda \leq k, \\ k, & \text{if } u_\lambda > k, \end{cases}$$

$\varphi_k = |u_k|^{l(\beta-1)} u_\lambda$ and $w_k = u_\lambda |u_k|^{\beta-1}$ with $\beta > 1$. By (ϕ_4) , we have $\phi(|\nabla u_\lambda|) |\nabla u_\lambda|^2 \geq q |\nabla u_\lambda|^l$. Then taking $\tilde{u} = \varphi_k$ in (5.3), by the definition of u_k and (C3)', we have

$$\begin{aligned} & q \int_{\mathbb{R}^N} |\nabla u_\lambda|^l |u_k|^{l(\beta-1)} dx \\ & \leq \int_{\mathbb{R}^N} \phi(|\nabla u_\lambda|) \nabla u_\lambda, \nabla \varphi_k dx - l(\beta-1) \int_{\mathbb{R}^N} |u_k|^{l(\beta-1)-2} u_k u_\lambda \phi(|\nabla u_\lambda|) (\nabla u_\lambda, \nabla u_k) dx \\ & \leq - \int_{\mathbb{R}^N} V(x) \phi(|u_\lambda|) u_\lambda \varphi_k dx + \lambda \int_{\mathbb{R}^N} \tilde{f}(x, u_\lambda) \varphi_k dx \\ & \leq \lambda D_4 \int_{\mathbb{R}^N} |u_\lambda|^r |u_k|^{l(\beta-1)} dx \\ & = \lambda D_4 \int_{\mathbb{R}^N} |u_\lambda|^{r-l} |w_k|^l dx. \end{aligned}$$

The rest proof is the same as Lemma 2.6 in [20] with replacing p with l and p^* with l^* . \square

Proof of Theorem 5.1. By Lemma 5.3 and Lemma 5.4, we have

$$\|u_\lambda\|_\infty \leq C\lambda^{\frac{1}{l^*-r}} C_{**}^{\frac{l^*-l}{l^*-r}} \max \left\{ \lambda^{-\frac{1}{k-l} \cdot \frac{l^*-l}{l^*-r}}, \lambda^{-\frac{l}{m(k-l)} \cdot \frac{l^*-l}{l^*-r}} \right\}.$$

Notice that $k > l, l^* > l, l^* > r$ and

$$k < K = \min \left\{ l^*, \frac{ml - l + l^*}{m} \right\}.$$

Then there exists a large $\Lambda_0 > 0$ such that $\|u_\lambda\|_\infty < \frac{\delta}{2}$ for all $\lambda > \Lambda_0$, which implies that $\tilde{F}(x, u_\lambda) = F(x, u_\lambda)$ for all $x \in \mathbb{R}^N$. Hence, u_λ is a nontrivial weak solution of system (5.1) and Lemma 5.3 and Lemma 5.4 imply that $\|u_\lambda\|_{1,\Phi} \rightarrow 0$ and $\|u_\lambda\|_\infty \rightarrow 0$ as $\lambda \rightarrow \infty$, respectively. \square

Remark 5.1. Comparing Theorem 1.1 with Theorem 5.1, it is easy to see Theorem 5.1 for the scalar equation (5.1) is better because f satisfies growth conditions just for $|t| \leq \delta$ with some positive constant δ rather than for $|t| \leq 2$. The proof of Theorem 1.1 for the elliptic system (1.5) present more complex derivation. Especially, Moser iteration for system (1.5) is more difficulties than that for the scalar equation (5.1). Moreover, in Theorem 1.1, we assume that F satisfies (F1)-(F3) for all $|(t, s)| \leq 4$ which is a circular domain. However, if we assume that there exist two positive constants δ_1 and δ_2 such that F satisfies (F1)-(F3) for all $|t| \leq \delta_1$ and $|s| \leq \delta_2$ which is a rectangular domain, the arguments will become more complex and it is unknown if Theorem 1.1 holds in such rectangular domain. One can consider the proof of Lemma 3.1 and the examples of cut-off functions to see such complexity. Finally, we would like to mention that if we let

$$\tilde{F}^+(u) = \begin{cases} \tilde{F}(u) & \text{if } u \geq 0 \\ 0 & \text{if } u < 0 \end{cases}, \quad \tilde{F}^-(u) = \begin{cases} \tilde{F}(u) & \text{if } u \leq 0 \\ 0 & \text{if } u > 0 \end{cases} \quad (5.4)$$

and consider the functional

$$\tilde{J}_\lambda(u) = \int_{\mathbb{R}^N} \Phi(|\nabla u|)dx + \int_{\mathbb{R}^N} V(x)\Phi(|u|)dx - \lambda \int_{\mathbb{R}^N} \tilde{F}^+(x, u)dx$$

and

$$\tilde{J}_\lambda(u) = \int_{\mathbb{R}^N} \Phi(|\nabla u|)dx + \int_{\mathbb{R}^N} V(x)\Phi(|u|)dx - \lambda \int_{\mathbb{R}^N} \tilde{F}^-(x, u)dx,$$

respectively, then we can obtain equation (5.2) has a positive solution and a negative solution. Thus Theorem 5.1 can be seen as the generalization of Theorem 1.2 in [20] if we assume that (V1) holds instead of the periodicity of V .

Acknowledgments

This work is supported by Yunnan Ten Thousand Talents Plan Young & Elite Talents Project and Candidate Talents Training Fund of Yunnan Province (No: 2017HB016). This work was completed partially during the

first author's visit to the Western University. The first author would like to thank the support of the China Scholarship Council for study abroad and the faculties in the Department of Applied Mathematics of the Western University for their warm help.

References

- [1] R.A. Adams and J.F. Fournier, Sobolev Spaces, Academic Press, New York, 2003.
- [2] C.O. Alves, G.M. Figueiredo and J.A. Santos, Strauss and Lions type results for a class of Orlicz-Sobolev spaces and applications, *Topol. Method. Nonl. An.* 44 (2014), 435-456.
- [3] K. Ait-Mahiout, Multiplicity and concentration behavior of positive solutions for a quasilinear problem in Orlicz-Sobolev spaces without Ambrosetti-Rabinowitz condition via penalization method, *J. Elliptic Parabol. Equ.* 2020, <https://doi.org/10.1007/s41808-020-00054-0>.
- [4] H. Brezis and L. Nirenberg, Remarks on finding critical points, *Commun. Pur. Appl. Math.* 44 (1991), 939-963.
- [5] G. Bonanno, G.M. Bisci and V.D. Rădulescu, Quasilinear elliptic non-homogeneous Dirichlet problems through Orlicz-Sobolev spaces, *Nonlinear Anal.* 75 (2012), 4441-4456.
- [6] G. Bonanno, G.M. Bisci and V.D. Rădulescu, Arbitrarily small weak solutions for a nonlinear eigenvalue problem in Orlicz-Sobolev spaces, *Monatsh. Math.* 165 (2012), 305-318.
- [7] J. Chabrowski and J. Yang, Existence theorems for elliptic equations involving supercritical Sobolev exponent, *Adv. Differential Equations*, 2 (1997), 231-256.
- [8] D.G. Costa and Z. Wang, Multiplicity results for a class of superlinear elliptic problems, *P. Am. Math. Soc.* 133 (2005), 787-794.
- [9] J.M. do ó, E. Medeiros and U. Severo, On the existence of signed and sign-changing solutions for a class of superlinear Schrödinger equations, *J. Math. Anal. Appl.* 342 (2008) 432-445.
- [10] G.M. Figueiredo and M. Furtado, Positive solutions for some quasilinear equations with critical and supercritical growth, *Nonlinear Anal.* 66 (2007), 1600-1616.
- [11] M. Fuchs and G. Li, Variational inequalities for energy functionals with nonstandard growth conditions, *Abstr. Appl. Anal.* 3 (2015), 41-64.

- [12] M. Fuchs and V. Osmolovski, Variational integrals on Orlicz-Sobolev spaces, *Z. Anal. Anwend.* 17(1998), 393-415.
- [13] N. Fukagai and K. Narukawa, Nonlinear eigenvalue problem for a model equation of an elastic surface, *Hiroshima Math. J.* 25 (1995), 19-41.
- [14] N. Fukagai, M. Ito and K. Narukawa, Positive solutions of quasilinear elliptic equations with critical Orlicz-Sobolev nonlinearity on \mathbb{R}^N , *Funkc. Ekvacioj*, 49 (2006), 235-267.
- [15] N. Fukagai, M. Ito and K. Narukawa, Quasilinear elliptic equations with slowly growing principal part and critical Orlicz-Sobolev nonlinear term, *Proc. Math. Roy. Soc. Edinb.* 139 (2009), 73-106.
- [16] C. Huang and J. Gao, Existence of positive solutions for supercritical quasilinear Schrödinger elliptic equations, *J. Math. Anal. Appl.* 472 (2019), 705-727.
- [17] M.A. Krasnosel'skiĭ, Ja. B. Rutickiĭ, Convex functions and Orlicz spaces, P. Noordhoff, 1961.
- [18] S. Liu, On quasilinear elliptic problems with finite or infinite potential wells, arXiv:1909.02822.
- [19] J. Moser, A new proof of De Giorgi's theorem concerning the regularity problem for elliptic differential equations, *Comm. Pure Appl. Math.* 13 (1960), 457-468.
- [20] E.S. Medeiros and U.B. Severo, On the existence of signed solutions for a quasilinear elliptic problem in \mathbb{R}^N , *Mat. Contemp.* 32 (2007), 193-205.
- [21] M. Mihăilescu and D. Repovš, Multiple solutions for a nonlinear and non-homogeneous problems in Orlicz-Sobolev spaces, *Appl. Math. Comput.* 217 (2011), 6624-6632.
- [22] P.H. Rabinowitz, Minimax methods in critical point theory with applications to differential equations, American Mathematical Society, Providence, RI, 1986.
- [23] M.M. Rao and Z.D. Ren, Applications of Orlicz Spaces, Marcel Dekker, New York, 2002.
- [24] Y. Xu, Z. Tan and D. Sun, Multiplicity results for a nonlinear elliptic problem involving the fractional Laplacian, *Acta Math. Sci.* 36 (2016), 1793-1803.
- [25] L. Wang, X. Zhang and H. Fang, Existence and multiplicity of solutions for a class of (φ_1, φ_2) -Laplacian elliptic system in \mathbb{R}^N via genus theory, *Comput. Math. with Appl.* 72 (2016), 110-130.
- [26] L. Wang, X. Zhang and H. Fang, Existence of ground state solutions for a class of quasilinear elliptic systems in Orlicz-Sobolev spaces, *Bound. Value Probl.* (2017) 2017: 106.
- [27] M. Willem, Minimax Theorems, Birkhäuser: Boston, MA, USA, 1996.