

Tangent fields, intrinsic stationarity, and self-similarity (with a supplement on Matheron Theory)

Jinqi Shen, Stilian Stoev, and Tailen Hsing

¹*Department of Statistics*

The University of Michigan, Ann Arbor

e-mail: jqshen@umich.edu; sstoev@umich.edu; thsing@umich.edu

Dedicated to the Memory of Mark Marvin Meerschaert (1955–2020)

Abstract: This paper studies the local structure of continuous random fields on \mathbb{R}^d taking values in a complete separable linear metric space \mathbb{V} . Extending seminal work of Falconer, we show that the generalized $(1+k)$ -th order increment tangent fields are self-similar and almost everywhere intrinsically stationary in the sense of Matheron. These results motivate the further study of the structure of \mathbb{V} -valued intrinsic random functions of order k (IRF $_k$, $k = 0, 1, \dots$). To this end, we focus on the special case where \mathbb{V} is a Hilbert space. Building on the work of Sasvari and Berschneider, we establish the spectral characterization of all second order \mathbb{V} -valued IRF $_k$'s, extending the classical Matheron theory. Using these results, we further characterize the class of Gaussian, *operator self-similar* \mathbb{V} -valued IRF $_k$'s, generalizing results of Dobrushin and Didier, Meerschaert and Pipiras, among others. These processes are the Hilbert-space-valued versions of the general k -th order operator fractional Brownian fields and are characterized by their self-similarity operator exponent as well as a finite trace class operator valued spectral measure. We conclude with several examples motivating future applications to probability and statistics.

In a technical [Supplement](#) of independent interest, we provide a unified treatment of the Matheron spectral theory for second-order stationary and intrinsically stationary processes taking values in a separable Hilbert space. We give the proofs of the Bochner-Neeb and Bochner-Schwartz theorems.

Keywords and phrases: tangent field, IRF $_k$, operator self-similarity, spectral theory, functional data analysis.

1. Introduction.

The tangent process of a random field is the stochastic process obtained in the limit of the suitably normalized increments of the random field at a fixed location. A pair of papers, [Falconer \[2002, 2003\]](#), discovered a remarkable property about the structure of the tangent process. Briefly speaking, Falconer proved that the tangent processes must be self-similar and have stationary increments (a.e.). The self-similarity of the tangent field is not surprising, which is a consequence of scaling. It is akin to what is shown in many limit theorems in the literature, such as those in the seminal work of Lamperti [cf. [Lamperti, 1962](#)]; see also [Davydov and Paulauskas \[2017\]](#) and a host of results on (univariate and multivariate) regular variation in [Gnedenko \[1943\]](#), [Meerschaert \[1984\]](#), [Hult and Lindskog \[2006\]](#). The property of stationary increments for the tangent process, however, is less expected. The proof of this property in Falconer's works involves a remarkable Lebesgue-density argument and ideas from geometric measure theory [cf. [Falconer, 1986](#), [Preiss, 1987](#)].

The starting point of our paper is extending Falconer’s results in two directions. Firstly, we consider generalized tangent processes obtained by taking local $(1 + k)$ -th order increments, $k \geq 0$. This is necessary if one wants to study the local behavior of many models arising in spatial statistics. Secondly, we consider random fields taking values in a linear complete separable metric space \mathbb{V} such as but not limited to a separable Banach space. The resulting limit processes, will be referred to as k -th order tangent processes. In Falconer [2002, 2003], $k = 0$ and $\mathbb{V} = \mathbb{R}$. The self-similarity property continues to hold for k -th order tangent processes, where self-similarity is in the sense of a general class of scaling actions, including operator scaling [cf. Meerschaert and Scheffler, 2001], that commensurate with the generality of the state space \mathbb{V} . To establish the generalized stationary-increment property, we introduce a new proof strategy based on the Lusin and Egorov theorems as well as some core ideas in Falconer [2002, 2003].

Interestingly, the stationarity of the higher-order increments of the k -th order tangent processes is related to the notion of *intrinsic random functions* of order k (IRF_k) introduced by Matheron [1973]. In the special case of real-valued processes ($\mathbb{V} = \mathbb{R}$), the classic results of Matheron as well as Gel’fand and Vilenkin [1964] lead to a concrete formula for *all possible* covariance structures of the k -th order tangent fields. It involves the local self-similarity exponent $H \in (0, k + 1]$ and the local *spectral measure* σ . Such results have been established by Dobrushin [1979] in the setting where the paths of the stochastic processes are generalized functions, i.e., random elements in $\mathcal{S}'(\mathbb{R}^d)$ – the topological dual of the Schwartz space $\mathcal{S}(\mathbb{R}^d)$. Our study, motivated by applications to spatial statistics and functional data analysis, considers random fields taking values in a separable Hilbert space \mathbb{V} . We follow the approach of Matheron rather than Dobrushin and realize the notion of a higher-order increment by integrating the process against signed measures with finite supports.

To this end, in Section 4, we develop an extension of Matheron’s theory to the case of processes taking values in a separable Hilbert space \mathbb{V} . Our theoretical development for Hilbert-space-valued IRF_k ’s is of independent interest and builds on a large body of existing although somewhat scattered work. With no intention to provide a complete list, we refer to Bochner [1948] and Khintchine [1934] for Bochner’s theorem and Neeb [1998] for extensions to general spaces; Cramér [1942] for the spectral representation of stationary random fields; Matheron [1973], Sasvári [2009] for the existence of general covariance of IRF_k and its integral representation; Berschneider [2012] for the integral representation of IRF_k in an abstract space. A more comprehensive summary on this line of literature can be found in Berschneider and Sasvári [2018]. Our proofs in this regard are contained in the [Supplement](#), which aims to be self-contained and only uses arguments that are common in probability and statistics.

The developed theory is then utilized in Section 5 to characterize the covariance structure of self-similar intrinsic random functions taking values in a separable Hilbert space. General linear operator scaling actions are considered as well as the classic multiplication by a scalar. In particular, our results provide characterizations of Gaussian operator self-similar IRF_k ’s, which can be viewed as infinite-dimensional versions of the k -th order fractional Brownian fields. The 0-th order operator fractional Brownian fields taking values in $\mathbb{V} = \mathbb{R}^m$ have been the subject of active investigation and numerous applications [see e.g. Mason and Xiao, 2001, Amblard and Coeurjolly, 2011, Kechagias and Pipiras, 2015, Abry and Didier, 2018, Abry et al., 2019, Düker, 2020, Biermé and Lacaux, 2020, and the references therein]. Most if not all of the existing work, however, focuses on random fields taking values in \mathbb{R}^m . In this paper, we provide a first comprehensive treatment of Hilbert space valued operator fractional Brownian fields and their higher order stationary increment counterparts – the Gaussian IRF_k ’s. This leads to infinite-dimensional extensions of seminal results due to Didier and Pipiras [2011], Didier et al. [2017], Perrin et al. [2001] among others.

This paper also contributes to statistical research in several ways. Matheron’s work on \mathbb{R} -valued IRF $_k$ ’s has had a substantial impact on the field of spatial statistics [see e.g. the monographs of [Stein, 1999](#), [Chilès and Delfiner, 2012](#)]. Our extension of the Matheron theory to the case of Hilbert-space-valued random fields provides novel tools and framework for spatially dependent functional data analysis – an active area in statistics [see, e.g., the monographs [Ramsay and Silverman, 2005](#), [Horváth and Kokoszka, 2012](#), [Hsing and Eubank, 2015](#), and references therein]. Unfortunately, the details of Matheron’s theory have been elusive to the broader community. The [Supplement](#) will be a useful resource for those who are interested in learning those details and their novel generalizations. The self-similar IRF $_k$ is itself a flexible model for spatial statistics. The self-similarity exponent operator \mathcal{H} and spectral measure σ , which now takes values in the space of positive trace-class operators on \mathbb{V} characterize the covariance structure. The pair (\mathcal{H}, σ) may be object of further modeling and inference in the context of in-fill asymptotics [cf. [Stein, 1999](#)]. The tangent process connection also provides guidance for building flexible random field models with desired local properties.

The paper is organized as follows. Section 2 introduces a suitable topology on the path space and scaling actions needed to define and study higher-order tangent fields. In Section 3, we establish the main results on the structure of higher-order tangent fields, namely their self-similarity and almost everywhere intrinsic stationarity. Section 4 develops the spectral theory for second-order stationary and intrinsically stationary random fields taking values in a separable Hilbert space \mathbb{V} . This treatment unifies and extends results of Bochner, Cramér, Gelfand-Vilenkin, Matheron, Neeb, Sasvári, and Berschneider. The covariance structure of the self-similar \mathbb{V} -valued IRF $_k$ is characterized in Section 5. General linear operator-scaling actions (Section 5.1) and the classic scalar scaling actions (Section 5.3) are studied. Open problems, examples and connections to the existing literature are presented in Section 5.2. Some technical proofs are relegated to the Appendix. Further background and details are given in the [Supplement](#).

2. Preliminaries.

This section develops some tools that will be useful for the study of tangent fields. We commence by defining some key spaces and operations.

For $d = 1, 2, \dots$, let Λ denote the collection of complex-valued measures on \mathbb{R}^d supported on finitely many points, i.e.,

$$\lambda(du) = \sum_{i=1}^n c_i \delta_{t_i}(du), \quad (2.1)$$

where $n = 1, 2, \dots$, $c_i \in \mathbb{C}$, $t_i \in \mathbb{R}^d$ for all i , and δ_a is the Dirac measure at a . Without loss of generality, we always assume the t_i ’s in the representation (2.1) to be distinct. In this paper we will be concerned with functions defined on Λ or a subspace of Λ and take values in some vector space \mathbb{V} . A special case of that is

$$g(\lambda) := \int g d\lambda \equiv \sum_{i=1}^n c_i g(t_i),$$

for any $g : \mathbb{R}^d \mapsto \mathbb{V}$.

For any $k \in \mathbb{N} := \{0, 1, 2, \dots\}$, a monomial of degree k on \mathbb{R}^d is any function of the form $u = (u_1, \dots, u_d) \mapsto u_1^{j_1} \cdots u_d^{j_d}$, where j_1, \dots, j_d are non-negative integer powers such that $j_1 + \cdots + j_d = k$.

More generally, a polynomial of degree k is any complex linear combination of monomials of degree less than or equal to k with at least one non-zero degree k term.

Definition 2.1. For any $k = 0, 1, 2, \dots$, let Λ_k be the class of $\lambda \in \Lambda$ such that $\int_{\mathbb{R}^d} f d\lambda = 0$ for polynomials f with degree $\deg(f) \leq k$. Thus, measures in Λ_k “annihilate” all polynomials of degree up to k . We also let $\Lambda_{-1} := \Lambda$.

Next, we define two operations pertinent to the definition of tangent fields. As usual, for any set $B \subset \mathbb{R}^d$, $c \in \mathbb{R}$ and $s \in \mathbb{R}^d$, let $c \cdot B = \{ct : t \in B\}$ and $s + B = B + s = \{s + t : t \in B\}$. Also, define the scaling and translation operations on Λ :

$$r \cdot \lambda := \lambda(r^{-1} \cdot), \quad r \neq 0, \quad \text{and} \quad s + \lambda := \lambda(\cdot - s), \quad s \in \mathbb{R}^d. \quad (2.2)$$

Clearly, Λ_k is closed with respect to both of these operations.

Assume that the random elements considered in the paper take values in a complete and separable metric linear space $(\mathbb{V}, d_{\mathbb{V}})$ over \mathbb{C} . Recall that $(\mathbb{V}, d_{\mathbb{V}})$ is said to be a metric linear space [cf. [Rolewicz, 1985](#)] if scalar multiplication and addition are continuous with respect to $d_{\mathbb{V}}$. Namely, for all $x_n, y_n, x, y \in \mathbb{V}$ and $c_n, c \in \mathbb{C}$, such that $|c_n - c| + d_{\mathbb{V}}(x_n, x) + d_{\mathbb{V}}(y_n, y) \rightarrow 0$, we have

$$d_{\mathbb{V}}(c_n x_n, cx) \rightarrow 0 \quad \text{and} \quad d_{\mathbb{V}}(x_n + y_n, x + y) \rightarrow 0.$$

By the Birkhoff-Kakutani Theorem [cf. Theorem 1.1.1 of [Rolewicz, 1985](#)], without loss of generality, we can and do assume that the metric $d_{\mathbb{V}}$ is translation invariant, that is,

$$d_{\mathbb{V}}(x, y) = d_{\mathbb{V}}(x - y, 0) \quad \text{for any } x, y \in \mathbb{V}. \quad (2.3)$$

A typical example of \mathbb{V} in our applications is a separable Banach or even Hilbert space. However, we do not restrict to only Banach spaces for now. We also assume throughout that the following continuity condition holds: For any $K > 0$,

$$\lim_{\delta \rightarrow 0} \sup_{|c| \leq K, d_{\mathbb{V}}(x, 0) < \delta} d_{\mathbb{V}}(cx, 0) \rightarrow 0. \quad (2.4)$$

Note that (2.4) readily holds if \mathbb{V} is a normed space and $d_{\mathbb{V}}$ is induced by the norm.

2.1. The spaces $S(\Lambda_k, \mathbb{V})$ and $\check{S}(\Lambda_k, \mathbb{V})$.

A function f from Λ to \mathbb{V} is linear if $f(c_1 \lambda_1 + c_2 \lambda_2) = c_1 f(\lambda_1) + c_2 f(\lambda_2)$, $c_1, c_2 \in \mathbb{C}$, $\lambda_1, \lambda_2 \in \Lambda$. Denote by $S(\Lambda_k, \mathbb{V})$ the set of all linear functions from Λ_k to \mathbb{V} . In this section, we focus on obtaining a representation of functions in $S(\Lambda_k, \mathbb{V})$ in terms of functions from \mathbb{R}^d to \mathbb{V} as well as a topological structure for a subspace of $S(\Lambda_k, \mathbb{V})$.

Next we discuss the important notion of *representation* introduced by [Matheron \[1973\]](#). A function $\check{f} : \mathbb{R}^d \mapsto \mathbb{V}$ is said to be a representation of $f \in S(\Lambda_k, \mathbb{V})$ if

$$f(\lambda) = \int \check{f} d\lambda, \quad \lambda \in \Lambda_k. \quad (2.5)$$

Consider the following construction of a representation. Denote by m_i , $i = 1, \dots, M_k := \binom{k+d}{k}$ all the monomials of degree less than or equal to k on \mathbb{R}^d , where the ordering is arbitrary. Define the \mathbb{R}^{M_k} -valued function

$$\mathbf{b}(t) := (m_1(t), \dots, m_{M_k}(t))^{\top}, \quad t \in \mathbb{R}^d. \quad (2.6)$$

Pick the points $t_i \in \mathbb{R}^d, i = 1, \dots, M_k$, such that the $M_k \times M_k$ matrix $B := (\mathbf{b}(t_1), \dots, \mathbf{b}(t_{M_k}))$ has full rank. Such points always exist. For instance, if $d = 1$ then B has full rank for arbitrary distinct t_i 's. However, for $d > 1$ some care is needed in selecting the t_i to ensure that B has full rank. For $t \in \mathbb{R}^d$, define the measure

$$\lambda_t = \delta_t - (\delta_{t_1}, \dots, \delta_{t_{M_k}})B^{-1}\mathbf{b}(t), \quad (2.7)$$

where, for any $\mathbf{c} = (c_1, \dots, c_{M_k})^\top \in \mathbb{C}^{M_k}$, $(\delta_{t_1}, \dots, \delta_{t_{M_k}})\mathbf{c}$ denotes the measure $\sum_{j=1}^{M_k} c_j \delta_{t_j}$. Below, for convenience, we adopt such matrix notation when there is no ambiguity. It follows from (2.6) and (2.7) that

$$\left(\int m_1 d\lambda_t, \dots, \int m_{M_k} d\lambda_t \right)^\top = (I - BB^{-1})\mathbf{b}(t) = \mathbf{0},$$

and so $\lambda_t \in \Lambda_k$ for all t . For any $f \in S(\Lambda_k, \mathbb{V})$, consider

$$\check{f}(t) := f(\lambda_t). \quad (2.8)$$

Note that $\lambda_{t_i} = 0$, the null measure, for all i . Thus,

$$\left(\check{f}(t_1), \dots, \check{f}(t_{M_k}) \right) = \mathbf{0}. \quad (2.9)$$

Moreover, for $\lambda \in \Lambda_k$, by linearity,

$$\int \check{f} d\lambda = f \left(\lambda - (\delta_{t_1}, \dots, \delta_{t_{M_k}})B^{-1}\mathbf{b}(\lambda) \right) = f(\lambda), \quad (2.10)$$

since $\mathbf{b}(\lambda) = \mathbf{0}$, and therefore (2.5) holds showing that \check{f} is a representation of f .

Clearly, the function \check{f} defined by (2.8) is not the only possible representation of f . However, any two representations g_1, g_2 of f differ by a polynomial of degree k , since, for all t ,

$$0 = \int (g_1 - g_2) d\lambda_t = g_1(t) - g_2(t) - (g_1(t_1) - g_2(t_1), \dots, g_1(t_{M_k}) - g_2(t_{M_k}))B^{-1}\mathbf{b}(t), \quad (2.11)$$

The difference will not affect any of the results in this paper. Thus, from now on, we will adhere to the representation \check{f} defined by (2.8).

It follows from (2.9) and (2.10) that there is a one-to-one correspondence between $S(\Lambda_k, \mathbb{V})$ and $\check{S}(\Lambda_k, \mathbb{V})$, where $\check{S}(\Lambda_k, \mathbb{V})$ denotes the set of functions from \mathbb{R}^d to \mathbb{V} that are equal to zero at each $t_i, i = 1, \dots, M_k$, where the isomorphism is determined by the bijection

$$\begin{aligned} (\mathcal{J}f)(t) &= f(\lambda_t), \quad f \in S(\Lambda_k, \mathbb{V}), \\ (\mathcal{J}^{-1}\check{f})(\lambda) &= \int \check{f} d\lambda, \quad \check{f} \in \check{S}(\Lambda_k, \mathbb{V}). \end{aligned} \quad (2.12)$$

A linear function $f \in S(\Lambda_k, \mathbb{V})$ is said to be continuous if its representation \check{f} is a continuous function from \mathbb{R}^d to \mathbb{V} . By (2.11), this property is “intrinsic” to f and does not depend on the representation. Let $S_c(\Lambda_k, \mathbb{V})$ denote the subset of continuous linear functions from Λ_k to \mathbb{V} , and $\check{S}_c(\Lambda_k, \mathbb{V})$ the corresponding set of representations. The mapping \mathcal{J} in (2.12) continues to be an isomorphism between $S_c(\Lambda_k, \mathbb{V})$ and $\check{S}_c(\Lambda_k, \mathbb{V})$.

We now proceed to define a topology on $S_c(\Lambda_k, \mathbb{V})$ and $\check{S}_c(\Lambda_k, \mathbb{V})$. A convenient metric on $\check{S}_c(\Lambda_k, \mathbb{V})$ is

$$\rho(\check{f}, \check{g}) = \sum_{n \geq 1} 2^{-n} \left(1 - \exp \left\{ - \sup_{\|t\| \leq n} d_{\mathbb{V}}(\check{f}(t), \check{g}(t)) \right\} \right), \quad (2.13)$$

where, as before, $\|\cdot\|$ denotes the Euclidean norm on \mathbb{R}^d and $d_{\mathbb{V}}$ denotes the metric on \mathbb{V} . Clearly, ρ metrizes the local uniform convergence topology, i.e., uniform convergence on compact sets. We also have the following simple but important fact, proved in Section A.1 in Appendix.

Lemma 2.2. *The metric space $(\check{S}_c(\Lambda_k, \mathbb{V}), \rho)$ is complete and separable.*

Using the canonical bijection \mathcal{J} , we define the corresponding metric on $S_c(\Lambda_k, \mathbb{V})$ as $\rho(f, g) := \rho(\mathcal{J}f, \mathcal{J}g)$, for $f, g \in S_c(\Lambda_k, \mathbb{V})$, where the same symbol ρ is adopted for convenience. Again, by (2.3), (2.4) and (2.11), the topology so defined does not depend on the particular representation used to define $(\check{S}_c(\Lambda_k, \mathbb{V}), \rho)$. It follows that \mathcal{J} is an isometry and both $(S_c(\Lambda_k, \mathbb{V}), \rho)$ and $(\check{S}_c(\Lambda_k, \mathbb{V}), \rho)$ are separable and complete. Thus, weak convergence of probability measures on these spaces can be defined in the usual manner [see, e.g., Billingsley, 1999]. Specifically, by Prokhorov's theorem, convergence in distribution in $(S_c(\Lambda_k, \mathbb{V}), \rho)$ or equivalently $(\check{S}_c(\Lambda_k, \mathbb{V}), \rho)$ is equivalent to the convergence of the finite-dimensional distributions and tightness. The following result provides a general criterion [see also Proposition 2.1 in Falconer, 2002].

Proposition 2.3. *Let X_n, X be random elements in $(S_c(\Lambda_k, \mathbb{V}), \rho)$ and let $\check{X}_n = \mathcal{J}(X_n)$ and $\check{X} = \mathcal{J}(X)$. Then $X_n \xrightarrow{d} X$ in $(S_c(\Lambda_k, \mathbb{V}), \rho)$, or equivalently $\check{X}_n \xrightarrow{d} \check{X}$ in $(\check{S}_c(\Lambda_k, \mathbb{V}), \rho)$, if and only if the following two conditions hold:*

(i) *For all $m > 0$ and $s_1, \dots, s_m \in \mathbb{R}^d$,*

$$(\check{X}_n(s_1), \dots, \check{X}_n(s_m)) \xrightarrow{d} (\check{X}(s_1), \dots, \check{X}(s_m)) \quad (2.14)$$

(ii) *For every compact set $K \subset \mathbb{R}^d$, \check{X}_n is strongly stochastically equicontinuous on K , namely, for all $\eta, \epsilon > 0$, there exists $\delta > 0$ such that*

$$\limsup_{n \rightarrow \infty} \mathbb{P} \left(\sup_{\|s-t\| < \delta, s, t \in K} d_{\mathbb{V}}(\check{X}_n(s), \check{X}_n(t)) > \eta \right) < \epsilon.$$

Proof. Since $(\check{S}_c(\Lambda_k, \mathbb{V}), \rho)$ is separable and complete, the result is a direct consequence of Theorem 14.5 and Proposition 14.6 in Kallenberg [1997]. \square

The convergence of the finite-dimensional distributions (2.14) is often easier to establish, while the challenge is to prove tightness. The following result provides a simple sufficient condition, which also implies the Hölder continuity of the limit. It is a restatement of Corollary 14.9 in Kallenberg [1997].

Proposition 2.4. *Suppose that $\check{X}_n, n \in \mathbb{N}$ take values in $(\check{S}_c(\Lambda_k, \mathbb{V}), \rho)$ and let the sequence of random variables $\{\check{X}_n(s_0), n \in \mathbb{N}\}$ be tight, for some $s_0 \in \mathbb{R}^d$.*

(i) *If for some $p > 0$ and $\alpha > 0$, and all $M > 0$, there exist $C_M < \infty$, such that*

$$\sup_{n \in \mathbb{N}} \mathbb{E}[d_{\mathbb{V}}(\check{X}_n(s), \check{X}_n(t))^p] \leq C_M \|s - t\|^{d+\alpha}, \quad \text{for all } \|s\|, \|t\| \leq M, s, t \in \mathbb{R}^d,$$

then the laws of the processes $\{\check{X}_n, n \in \mathbb{N}\}$ are tight in $(\check{S}_c(\Lambda_k, \mathbb{V}), \rho)$.

(ii) If, moreover, $\check{X}_n \xrightarrow{d} \check{X}$, in $(\check{S}_c(\Lambda_k, \mathbb{V}), \rho)$, then with probability one, \check{X} has γ -Hölder continuous paths, for all $\gamma \in (0, \alpha/p)$. That is, there is an event Ω^* , with $\mathbb{P}(\Omega^*) = 1$, such that for all $M > 0$, we have

$$d_{\mathbb{V}}(\check{X}(s, \omega), \check{X}(t, \omega)) \leq C_M(\omega) \|s - t\|^\gamma, \quad \text{for all } \|s\|, \|t\| \leq M, \omega \in \Omega^*,$$

with some $C_M(\omega) < \infty$.

Remark 2.1. By taking $\check{X}_n \equiv \check{X}$ in Proposition 2.4, we recover an extension of the well-known Kolmogorov-Chentsov path-regularity criterion. See also Theorem 2.23 in [Kallenberg \[1997\]](#).

Remark 2.2. If $k \geq 0$, recall that by (2.9), we have $\check{X}_n(t_i) = 0$, $i = 1, \dots, M_k$. Thus, in Proposition 2.4, one can trivially take $s_0 = t_1$ and the required tightness of the random variables $\{\check{X}_n(s_0), n \in \mathbb{N}\}$ is immediate. This condition is non-trivial only when $k = -1$ (the case of stationary processes) where by convention \mathcal{J} is the identity and $\check{S}_c(\Lambda_{-1}, \mathbb{V}) \equiv S_c(\Lambda_{-1}, \mathbb{V})$.

The above moment-based criterion is used in Section 5.2 to furnish examples of tangent processes.

2.2. Scaling actions on \mathbb{V} .

When considering limit theorems for \mathbb{V} -valued processes, one may need to rescale the process using an operator different from the usual scalar multiplication. This is particularly relevant for the case where \mathbb{V} is an infinite dimensional space of functions. The next definition introduces the natural conditions that such rescaling operators should possess. It is similar to the one considered in [Hult and Lindskog \[2006\]](#) in their abstract treatment of regular variation.

Definition 2.5. A family of (possibly non-linear) operators $T_a : \mathbb{V} \rightarrow \mathbb{V}$, indexed by the multiplicative group $\mathbb{R}_+ := (0, \infty)$ is said to be a scaling action on \mathbb{V} if the following conditions hold:

- (i) for all $a_1 > 0$ and $a_2 > 0$, we have $T_{a_1} \circ T_{a_2} = T_{a_1 a_2}$,
- (ii) T_1 is the identity, and $T_a(0) = 0$ for all $a > 0$,
- (iii) $\{T_a\}$ is continuous, i.e., $d_{\mathbb{V}}(T_{a_n}(x_n), T_a(x)) \rightarrow 0$, whenever $a_n \rightarrow a > 0$ and $d_{\mathbb{V}}(x_n, x) \rightarrow 0$,
- (iv) $\{T_a\}$ is radially monotone, i.e., $d_{\mathbb{V}}(T_{a_1}(x), 0) < d_{\mathbb{V}}(T_{a_2}(x), 0)$, for all $0 \neq x \in \mathbb{V}$ and $0 < a_1 < a_2$, and
- (v) $d_{\mathbb{V}}(T_a(x), 0) \rightarrow 0$ as $a \downarrow 0$, for all $x \in \mathbb{V}$.

The above definition readily implies that $T_a, a > 0$ are bijections and in particular $T_a(x) \neq 0$ for all $x \neq 0$.

Remark 2.3. Property (v) in Definition 2.5 can be replaced by the equivalent condition of

$$\bigcup_{n=1}^{\infty} T_n(B_r) = \mathbb{V} \text{ for all } r \in \mathbb{R}_+, \quad (2.15)$$

where $B_r := \{x \in \mathbb{V} : d_{\mathbb{V}}(x, 0) < r\}$ is the open ball centered at the origin with radius r . To see the equivalence, first assume that (2.15) holds and, by (iv), verifying (v) then amounts to showing that $d_{\mathbb{V}}(T_{1/n}(x), 0) \rightarrow 0$ for all $x \in \mathbb{V}$. For every $r := \epsilon > 0$, however, (2.15) entails that $x \in T_n(B_\epsilon)$ for all sufficiently large n . By (i) and (ii), this implies that $T_{1/n}(x) \in B_\epsilon$, or $d_{\mathbb{V}}(T_{1/n}(x), 0) < \epsilon$, for all sufficiently large n . The converse argument showing (v) implies (2.15) is similar.

Remark 2.4. Definition 2.5 does not require the space \mathbb{V} to be linear. The linearity of \mathbb{V} , however, is needed in the definition of the path-spaces $S(\Lambda_k, \mathbb{V})$ and $S_c(\Lambda_k, \mathbb{V})$.

Many common examples of scaling operations readily satisfy the above conditions. For instance, if \mathbb{V} is a linear normed space, a natural scaling action is scalar multiplication itself:

$$T_a(x) := a \cdot x. \quad (2.16)$$

More generally, the scalar multiplication is a scaling action if the underlying metric is homogeneous, e.g., $d_{\mathbb{V}}(a \cdot x, a \cdot y) = a^{\gamma} d_{\mathbb{V}}(x, y)$, $\gamma > 0$.

Observe that since $T_1 = \text{I}$ by property (ii), where I stands for the identity operator, we have $T_a^{-1} = T_{1/a}$, $a > 0$ by property (i). Hence the mappings T_a are homeomorphisms and map open (closed) sets to open (closed) sets. The radial monotonicity property (iv) implies that $T_{a_2}^{-1}(B_r) \subset T_{a_1}^{-1}(B_r)$, for all $0 < a_1 < a_2$. This, since $T_a^{-1} = T_{1/a}$, entails

$$T_{a_1}(B_r) \subset T_{a_2}(B_r), \quad \text{for all } 0 < a_1 < a_2. \quad (2.17)$$

Note that the metric $d_{\mathbb{V}}$ and the action need not be “compatible”, that is, $d_{\mathbb{V}}(T_a(x), 0)$ is in general not equal to $a \cdot d_{\mathbb{V}}(x, 0)$ and therefore, $T_a(B_r)$ is in general not B_{ar} .

Remark 2.5. In the case where \mathbb{V} is a Hilbert space and $\mathcal{H} : \mathbb{V} \rightarrow \mathbb{V}$ is a fixed bounded linear operator, one can consider the action $T_c(x) := c^{\mathcal{H}} x$, $x \in \mathbb{V}$ for $c > 0$, where $c^{\mathcal{H}} := e^{\log(c)\mathcal{H}}$ (see (5.1)). Lemma A.1 shows that $c^{\mathcal{H}}$ is a scaling action in the sense of Definition 2.5, under certain natural conditions on the operator \mathcal{H} .

Limit theorems under linear operator scaling on $\mathbb{V} := \mathbb{R}^m$ have been studied extensively in the literature [see e.g. Meerschaert and Scheffler, 2001, and the references therein and thereof]. Such actions for a general separable Hilbert space \mathbb{V} will be considered in Section 5.1.

Given a scaling action $\{T_a, a \in \mathbb{R}_+\}$ on \mathbb{V} , it is natural to consider its coordinate-wise extension on the space of \mathbb{V} -valued functions $S(\Lambda_k, \mathbb{V})$. Namely, the action $\tilde{T}_a : S_c(\Lambda_k, \mathbb{V}) \rightarrow S_c(\Lambda_k, \mathbb{V})$ is defined such that for all $f \in S_c(\Lambda_k, \mathbb{V})$, and any $\lambda \in \Lambda_k$

$$\tilde{T}_a(f)(\lambda) = T_a(f(\lambda)). \quad (2.18)$$

The following result shows that the coordinate-wise action is in fact a scaling action on $S_c(\Lambda_k, \mathbb{V})$. Its proof is given in Section A.1, below.

Lemma 2.6. *For any scaling action $\{T_a, a \in \mathbb{R}_+\}$ on $(\mathbb{V}, d_{\mathbb{V}})$, the coordinate-wise action $\{\tilde{T}_a, a \in \mathbb{R}_+\}$ in (2.18) is a scaling action on the linear space $S_c(\Lambda_k, \mathbb{V})$ equipped with the metric ρ in (2.13).*

In view of Lemma 2.6, from now on we will use the same notation $\{T_a\}$ for the scaling action on \mathbb{V} and its coordinate-wise extensions on $S_c(\Lambda_k, \mathbb{V})$.

3. Tangent fields and their properties.

Throughout this section, suppose that $X = \{X(\lambda), \lambda \in \Lambda_k\}$ is a random element in $(S_c(\Lambda_k, \mathbb{V}), \rho)$. That is, for some probability space $(\Omega, \mathcal{F}, \mathbb{P})$, we have that $X : \Omega \rightarrow S_c(\Lambda_k, \mathbb{V})$ is an $\mathcal{F}|\mathcal{B}_{(S_c(\Lambda_k, \mathbb{V}), \rho)}$ -measurable map, where $\mathcal{B}_{(S_c(\Lambda_k, \mathbb{V}), \rho)}$ stands for the Borel σ -field on $S_c(\Lambda_k)$. For $s \in \mathbb{R}^d$, $\lambda \in \Lambda_k$ and $r > 0$, define

$$X(s, r \cdot \lambda) = X(s + r \cdot \lambda), \quad (3.1)$$

where $s + r \cdot \lambda$ is as defined in (2.2). With some abuse of notation, an example of $X(\lambda)$ is $X(\lambda) := \int_{\mathbb{R}^d} X(u) \lambda(du)$ for some random field $X = \{X(s), s \in \mathbb{R}^d\}$ with continuous sample paths. (Note that if \check{X} is the representation of X as defined in Section 2.1, then $X(s, r \cdot \lambda) = \int_{\mathbb{R}^d} \check{X}(s + ru) \lambda(du) = \int_{\mathbb{R}^d} X(s + ru) \lambda(du)$, since $\lambda \in \Lambda_k$ annihilates all polynomials of degree up to k .)

One can interpret $X(s, r \cdot \lambda)$ as a generalized $(1 + k)$ -th order increment of X at location s and scale $r > 0$, relative to λ . Indeed, consider for example the case $d = 1$ and let $\lambda(du) = \sum_{j=0}^{k+1} \binom{k+1}{j} (-1)^j \delta_{\{j\}}(du)$. Then, for $s \in \mathbb{R}$ and $r > 0$,

$$X(s, r \cdot \lambda) = \sum_{j=0}^{k+1} \binom{k+1}{j} (-1)^j X(s + rj) \equiv \Delta_r^{k+1} X(s),$$

where $\Delta_r X(s) := X(s + r) - X(s)$ and $\Delta_r^{k+1} X(s) := \Delta_r(\Delta_r^k X(\cdot))(s)$, $k = 0, 1, \dots$ is the usual $(1 + k)$ -th order regular difference operator.

Thus, considering the process X as a function of $(s + r \cdot \lambda)$, for all (any) signed measures $\lambda \in \Lambda_k$ effectively amounts to zooming in on its $(1 + k)$ th order increments at location s and scale $r > 0$. By letting $r \downarrow 0$, one can examine the local behavior of X and arrive at (generalized) tangent fields as detailed next. This indeed extends the setting of Falconer [2002] who focused on $\mathbb{V} = \mathbb{R}, k = 0$, and studied the increment process $X(s + rt) - X(s) \equiv X(s + r \cdot t\lambda)$, $t \in \mathbb{R}$.

One of the goals of the paper is to study the asymptotic behavior of the generalized increments $X(s, r \cdot \lambda)$ as $r \downarrow 0$ for fixed $s \in \mathbb{R}^d$, where $r \cdot \lambda$ is as in (2.2). The normalization of the asymptotics will be facilitated by scaling actions $T_s = \{T_{s,a}, a > 0\}$ as described by the next definition. In this context, we use that the process $\{T_{s,c(s,r)}(X(s, r \cdot \lambda)), \lambda \in \Lambda_k\}$ is a random element in $S_c(\Lambda_k, \mathbb{V})$ for all s, r , which is easy to verify by (3.1) and the continuity of the scaling action. This remark applies to similar situations below and will not be repeated.

Definition 3.1. *Let $s \in \mathbb{R}^d$. A random process $Y_s = \{Y_s(\lambda), \lambda \in \Lambda_k\} \in S_c(\Lambda_k, \mathbb{V})$ is said to be a k -th order tangent field (or tangent process) to X at s based on the scaling action $T_s = \{T_{s,a}, a > 0\}$, if it is non-zero and for some normalizing function $c(s, r) > 0$, we have*

$$\{T_{s,c(s,r)}(X(s, r \cdot \lambda)), \lambda \in \Lambda_k\} \xrightarrow{d} \{Y_s(\lambda), \lambda \in \Lambda_k\}, \quad \text{as } r \downarrow 0, \quad (3.2)$$

where the convergence in distribution takes place in $(S_c(\Lambda_k, \mathbb{V}), \rho)$.

The role of the function $c(s, r)$ is to provide flexibility in the choice of normalization without having to change the scaling action. For example, in the special setting $\mathbb{V} = \mathbb{R}, k = 0$ with the simple scalar scaling action $T_{s,a}(x) \equiv T_a(x) = a \cdot x$, and λ replaced by $\lambda_t(dx) := \delta_{\{t\}}(dx) - \delta_{\{0\}}(dx)$, Relation (3.2) implies

$$\{c(s, r)(X(s + rt) - X(s)), t \in \mathbb{R}\} \xrightarrow{d} \{Y_s(\lambda_t), t \in \mathbb{R}\}, \quad \text{as } r \downarrow 0.$$

This recovers the classic setting, of tangent processes, where $c(s, r)$ plays the role of a normalizing constant. In this case, Falconer [2002] showed that tangent fields must be self-similar and have stationary increments. In the following two subsections, we extend Falconer's results to the general setting of this paper.

3.1. Tangent fields are self-similar.

Self-similarity is a distributional invariance phenomenon, which is ubiquitous in the study of stochastic process limit theory. Recall that a real-valued stochastic process $\xi = \{\xi(t), t \in \mathbb{R}^d\}$

is said to be self-similar with self-similarity exponent $H > 0$, if for all $r > 0$, we have $\{\xi(rt), t \in \mathbb{R}^d\} \stackrel{fdd}{=} \{r^H \xi(t), t \in \mathbb{R}^d\}$, where $\stackrel{fdd}{=}$ means equality of all finite-dimensional distributions. For \mathbb{V} -valued processes, we have the following natural extension of the notion of self-similarity.

Definition 3.2. An \mathbb{V} -valued stochastic process $\xi = \{\xi(\lambda), \lambda \in \Lambda_k\}$ is said to be self-similar relative to the scaling action $\{T_a\}$ if for some $\alpha \in \mathbb{R}$, we have

$$\{\xi(r \cdot \lambda), \lambda \in \Lambda_k\} \stackrel{fdd}{=} \{T_{r^\alpha}(\xi(\lambda)), \lambda \in \Lambda_k\}, \quad \text{for all } r > 0. \quad (3.3)$$

Remark 3.1. The above definition views ξ in the wide sense as a measurable map $\xi : \Omega \rightarrow \mathbb{V}^{\Lambda_k}$, where \mathbb{V}^{Λ_k} is equipped with the product σ -field $\mathcal{B}_{\mathbb{V}^{\Lambda_k}}$ generated by the class \mathcal{C} all finite-dimensional cylinder sets $C := \{x \in \mathbb{V}^{\Lambda_k} : x(\lambda_i) \in B_i, i = 1, \dots, m\}$, $B_i \in \mathcal{B}_{(\mathbb{V}, d_{\mathbb{V}})}$, $\lambda_i \in \Lambda_k, i = 1, \dots, m, m \in \mathbb{N}$. When the paths of ξ are continuous, i.e., ξ is a random element in $(S_c(\Lambda_k, \mathbb{V}), \rho)$, it can be shown that the equality of the finite-dimensional distributions “ $\stackrel{fdd}{=}$ ” in (3.3) is equivalent to equality in distribution “ $\stackrel{d}{=}$ ” between $S_c(\Lambda_k, \mathbb{V})$ -valued random elements. Indeed, firstly, all finite-dimensional projections $\pi_{\lambda_1, \dots, \lambda_m} : S_c(\Lambda_k, \mathbb{V}) \rightarrow \mathbb{V}^m$, $m \in \mathbb{N}$ for $\lambda_i \in \Lambda_k, i = 1, \dots, m$ are continuous and hence measurable. This shows that $\mathcal{B}_{\mathbb{V}^{\Lambda_k}} \subset \mathcal{B}_{(S_c(\Lambda_k, \mathbb{V}), \rho)}$ and hence “ $\stackrel{d}{=}$ ” implies “ $\stackrel{fdd}{=}$ ”. On the other hand, the fact that $(S_c(\Lambda_k, \mathbb{V}), \rho)$ is second-countable, entails that $\mathcal{B}_{(S_c(\Lambda_k, \mathbb{V}), \rho)}$ is generated by the class of all closed balls, for example. Since each such ball is a countable intersection of cylinder sets [e.g., as in the proof of Proposition 12.2.2 in [Dudley, 2002](#)] it follows that the Borel σ -field $\mathcal{B}_{(S_c(\Lambda_k, \mathbb{V}), \rho)}$ is determined by the π -system \mathcal{C} of all finite-dimensional cylinder sets. Thus, appealing to the π - λ Theorem, we see that “ $\stackrel{fdd}{=}$ ” implies also “ $\stackrel{d}{=}$ ”.

The seminal work of [Lamperti \[1962\]](#) shows that all non-trivial large-scale limits of stochastically continuous processes are self-similar. From this perspective, it is expected that tangent fields (as small-scale limits) be self-similar. [Falconer \[2002, 2003\]](#) has shown that this is indeed the case for $k = 0$. The next result addresses the general case of k -th order tangent fields of \mathbb{V} -valued processes.

Theorem 3.3. Assume that, for some location $s \in \mathbb{R}^d$, $\{Y_s(\lambda), \lambda \in \Lambda_k\}$ is a k -th order tangent field to X at s with respect to the scaling action T_s . That is, Relation (3.2) holds for some $c(s, r)$.

(i) Then, for all $r > 0$, we have

$$\{Y_s(r \cdot \lambda), \lambda \in \Lambda_k\} \stackrel{d}{=} \{T_{r^{\alpha(s)}} Y_s(\lambda), \lambda \in \Lambda_k\}, \quad (3.4)$$

where $\alpha(s) > 0$ is some positive constant. We have, moreover, that

$$c(s, r) = r^{-\alpha(s)} \ell_s(r), \quad (3.5)$$

where $\ell_s(r)$ is a slowly varying function at 0, i.e., for every fixed $h > 0$, $\ell_s(hr)/\ell_s(r) \rightarrow 1$, $r \downarrow 0$.

(ii) The tangent process is unique up to rescaling. That is, if (3.2) also holds with $c(s, r)$ and $Y_s = \{Y_s(\lambda)\}$ replaced by $\tilde{c}(s, r)$ and $\tilde{Y}_s = \{\tilde{Y}_s(\lambda)\}$, respectively, then we have

$$\lim_{r \downarrow 0} \frac{\tilde{c}(s, r)}{c(s, r)} = a \in (0, \infty) \quad \text{and} \quad \tilde{Y}_s \stackrel{d}{=} T_{s, a}(Y_s). \quad (3.6)$$

Remark 3.2. Relations (3.4) and (3.5) show that the normalization used to define a tangent field may differ from the scaling action that characterizes the self-similarity of the tangent field by a slowly varying factor, which cannot be dropped in general. This is akin to the fundamental role of slowly varying functions in the normalization of the partial sums in the non-Gaussian Central Limit Theorem.

Proof of Theorem 3.3: *Proof of part (i).* For all fixed $h > 0$, by (3.2), as $r \downarrow 0$, we have

$$\tilde{\xi}_r := \{T_{s,c(s,r)}(X(s, (hr) \cdot \lambda)), \lambda \in \Lambda_k\} \xrightarrow{d} \tilde{\xi} := \{Y_s(h \cdot \lambda), \lambda \in \Lambda_k\}.$$

On the other hand, as $r \downarrow 0$,

$$\xi_r := \{T_{s,c(s,hr)}(X(s, (hr) \cdot \lambda)), \lambda \in \Lambda_k\} \xrightarrow{d} \xi := \{Y_s(\lambda), \lambda \in \Lambda_k\}.$$

By assumption both ξ and $\tilde{\xi}$ are non-zero. Observe that

$$T_{s,c(s,r)}(X(s, (hr) \cdot \lambda)) = T_{s, \frac{c(s,r)}{c(s,hr)}} \circ T_{s,c(s,hr)}(X(s, (hr) \cdot \lambda))$$

and hence

$$T_{s, \frac{c(s,r)}{c(s,hr)}}(\xi_r) = \tilde{\xi}_r \xrightarrow{d} \tilde{\xi}.$$

Applying Lemma A.5 (with $\mathbb{X} := S_c(\Lambda_k, \mathbb{V})$ – recall Lemma 2.6) gives

$$\frac{c(s,r)}{c(s,hr)} \rightarrow a(s,h), \quad \text{as } r \downarrow 0, \quad (3.7)$$

for some positive $a(s,h) > 0$. We have, moreover, $\tilde{\xi} \stackrel{d}{=} T_{s,a(s,h)}(\xi)$, which reads

$$\{Y_s(h \cdot \lambda), \lambda \in \Lambda_k\} \stackrel{d}{=} \{T_{s,a(s,h)}(Y_s(\lambda)), \lambda \in \Lambda_k\}. \quad (3.8)$$

We will next show that $a(s,h) = h^{\alpha(s)}$, for some $\alpha(s) > 0$. First, Relation (3.8) readily implies that for all $h_1 > 0$ and $h_2 > 0$

$$a(s, h_1 h_2) = a(s, h_1) a(s, h_2). \quad (3.9)$$

Indeed, by (3.8),

$$\begin{aligned} \{T_{s,a(s,h_1 h_2)}(Y_s(\lambda))\} &\stackrel{d}{=} \{Y_s((h_1 h_2) \cdot \lambda)\} \stackrel{d}{=} \{T_{s,a(s,h_1)}(Y_s(h_2 \cdot \lambda))\} \\ &\stackrel{d}{=} \{T_{s,a(s,h_1)} \circ T_{s,a(s,h_2)}(Y_s(\lambda))\} = \{T_{s,a(s,h_1)a(s,h_2)}(Y_s(\lambda))\}. \end{aligned}$$

Since Y_s is nonzero, the last relation implies (3.9) by (i) of Lemma A.5.

The function $a(s,h)$ is also continuous in $h \in (0, \infty)$. Indeed, for any sequence $h_n \rightarrow h$, $h_n, h \in (0, \infty)$, by Lemma A.6 (applied with $X_n := Y_s$, $v_n := 0$, and $r_n := h_n$), we have $\{Y_s(h_n \cdot \lambda)\} \xrightarrow{d} \{Y_s(h \cdot \lambda)\}$. Therefore, by (3.8),

$$\{T_{s,a(s,h_n)}(Y_s(\lambda))\} \stackrel{d}{=} \{Y_s(h_n \cdot \lambda)\} \xrightarrow{d} \{Y_s(h \cdot \lambda)\} \stackrel{d}{=} \{T_{s,a(s,h)}(Y_s(\lambda))\}. \quad (3.10)$$

Since Y_s is nonzero, applying (ii) of Lemma A.5, we obtain

$$a(s, h_n) \rightarrow a(s, h), \quad (3.11)$$

which shows the desired continuity.

Combining (3.9), (3.11), the continuity of $a(s, \cdot)$ and the fact that, trivially, $a(s, 1) = 1$, it is straightforward to conclude that $a(s, h) = h^{\alpha(s)}$, $h > 0$, for some $\alpha(s) \in (-\infty, \infty)$, which is a special example of Cauchy's functional equation [cf. Theorem 5.2.1 of Kuczma, 2009].

We will show next that $a(s, h_n) \rightarrow 0$ as $h_n \downarrow 0$, which necessarily implies $\alpha(s) > 0$. Indeed, with $h = 0$, (3.10) implies that $\{T_{s,a(s,h_n)}(Y_s(\lambda))\} \xrightarrow{d} 0 = \{Y_s(0 \cdot \lambda)\}$, as $n \rightarrow \infty$. This, by (iii) of Lemma A.5, yields $a(s, h_n) \rightarrow 0$.

To conclude the proof of part (i), letting $c(s, r) =: r^{-\alpha(s)}\ell_s(r)$, we see from Equation (3.7) that, for all $h > 0$, $\ell_s(r)/\ell_s(hr) \rightarrow 1$, as $r \downarrow 0$, which shows ℓ_s is a slowly varying function at 0.

Proof of part (ii). Assume now that in addition to (3.2), we have

$$\tilde{\eta}_r := \{T_{s,\tilde{c}(s,r)}(X(s, r \cdot \lambda)), \lambda \in \Lambda_k\} \xrightarrow{d} \tilde{Y}_s = \{\tilde{Y}_s(\lambda), \lambda \in \Lambda_k\},$$

as $r \downarrow 0$. By the properties of the scaling action, with $\eta_r := \{T_{s,c(s,r)}(X(s, r \cdot \lambda)), \lambda \in \Lambda_k\}$, we have

$$\tilde{\eta}_r = T_{s, \frac{\tilde{c}(s,r)}{c(s,r)}}(\eta_r) \xrightarrow{d} \tilde{Y}_s, \quad \text{as } r \downarrow 0.$$

On the other hand, Relation (3.2) reads $\eta_r \xrightarrow{d} Y_s$, as $r \downarrow 0$. Since both limits \tilde{Y}_s and Y_s are non-zero, Lemma A.5 entails $\tilde{c}(s, r)/c(s, r) \rightarrow a > 0$ and $T_{s,a}(Y_s) \stackrel{d}{=} \tilde{Y}_s$, which proves (3.6), i.e., the essential uniqueness of the tangent process. \square

3.2. Tangent fields are intrinsically stationary.

One of the key results in Falconer [2002] is that (almost all) tangent fields have stationary increments (cf. Theorem 3.6 therein). The proof of that is based on a delicate measure-theoretic argument. Below, we show that this phenomenon extends to higher order tangent fields to processes taking values in a linear separable metric space \mathbb{V} .

Let (3.2) hold and let

$$F_n(s) := \text{Law of } \left\{ T_{s,c(s,1/n)} X(s, (1/n) \cdot \lambda), \lambda \in \Lambda_k \right\} \quad (3.12)$$

be the probability distribution of the rescaled version of $\{X(s, \lambda), \lambda \in \Lambda_k\}$ in $(S_c(\Lambda_k, \mathbb{V}), \rho)$. Similarly, let

$$G(s) := \text{Law of } \{Y_s(\lambda), \lambda \in \Lambda_k\}. \quad (3.13)$$

In this notation, the convergence in (3.2) (with $r := 1/n$) is simply

$$F_n(s) \xrightarrow{w} G(s), \quad n \rightarrow \infty, \quad (3.14)$$

where ' \xrightarrow{w} ' denotes the weak convergence of probability measures. An important result that will be utilized below is Proposition 3.6 in the Appendix. In that regard, we first equip the space $\mathcal{P}(S_c(\Lambda_k, \mathbb{V}), \rho)$ of probability measures on $(S_c(\Lambda_k, \mathbb{V}), \rho)$ with a separable metric that metrizes the weak convergence (3.14). Since $(S_c(\Lambda_k, \mathbb{V}), \rho)$ is complete and separable, a suitable metric is d_{LP} , the Lévy-Prokhorov distance [cf. Theorem 6.8 of Billingsley, 1999]. Thus, (3.14) can be re-expressed as

$$F_n(s) \rightarrow G(s) \text{ in } (\mathcal{P}(S_c(\Lambda_k, \mathbb{V}), \rho), d_{\text{LP}}), \quad (3.15)$$

namely, $d_{\text{LP}}(F_n(s), G(s)) \rightarrow 0$.

Definition 3.4. A process $Y = \{Y(\lambda), \lambda \in \Lambda_k\}$ is said to be strictly intrinsically stationary if

$$\{Y(w + \lambda), \lambda \in \Lambda_k\} \stackrel{fdd}{=} \{Y(\lambda), \lambda \in \Lambda_k\}, \quad \text{for all } w \in \mathbb{R}^d.$$

Note that this is different from the usual notion of weak or second-order intrinsic stationarity in the literature [cf. [Sasvári, 2009](#)]. The latter is the topic of Section 4.

Remark 3.3. Observe that the notion of strict intrinsic stationarity like that of self-similarity in (3.3) is stated in greater generality using equality in the sense of finite-dimensional distributions. As discussed in Remark 3.1, when the processes therein take values in the path space $S_c(\mathbb{R}^d, \mathbb{V})$, the equality of the finite-dimensional distributions is equivalent to that of the probability distributions of the processes. In the next result, the processes are understood as random elements in $S_c(\mathbb{R}^d, \mathbb{V})$.

Theorem 3.5. Let B be a Borel set of \mathbb{R}^d . Assume that $X = \{X(\lambda), \lambda \in \Lambda_k\}$ is a random element in $S_c(\mathbb{R}^d, \mathbb{V})$ and it has a k -th order tangent field $Y_s = \{Y_s(\lambda), \lambda \in \Lambda_k\}$ at every $s \in B$ in the sense of Definition 3.1. Also assume that for any r , the normalization $c(s, r)$ is Borel measurable in s , and for any s and any sequence $w_n \rightarrow w \in \mathbb{R}^d$,

$$T_{s_n, c(s_n, 1/n)} \circ T_{s, c(s, 1/n)}^{-1} \rightarrow \mathbf{I}, \quad (3.16)$$

where \mathbf{I} is the identity operator and $s_n := s + n^{-1}w_n$. Then, there exists a set U with zero Lebesgue measure such that for all $s \in B \setminus U$, the tangent field Y_s is strictly intrinsically stationary. That is, at almost all locations s , tangent fields are strictly intrinsically stationary.

The proof of this result uses the following proposition established in Section A.2 below.

Proposition 3.6. Let $B \subset \mathbb{R}^d$ be a Borel set with finite Lebesgue measure $\text{Leb}(B) < \infty$. Suppose that $F_n : B \rightarrow E$ is a sequence of Borel measurable functions into the separable metric space (E, ρ_E) such that

$$F_n(s) \xrightarrow[n \rightarrow \infty]{} G(s), \quad \text{for almost all } s \in B.$$

Then, for every $\epsilon > 0$, there exists a compact set $K_\epsilon \subset B$, such that $\text{Leb}(B \setminus K_\epsilon) < \epsilon$, the function G being continuous on K_ϵ , and

$$F_n(s_n) \rightarrow G(s), \quad \text{whenever } s_n \rightarrow s, \text{ for } s_n, s \in K_\epsilon. \quad (3.17)$$

Proof of Theorem 3.5: By the σ -additivity of the Lebesgue measure on \mathbb{R}^d , it suffices to establish the result for the case $\text{Leb}(B) < \infty$.

The assumption implies that (3.15) holds for all $s \in B$. The continuity of X and the Borel-measurability of $s \mapsto c(s, 1/n)$ entail that $s \mapsto F_n(s)$ is a sequence of Borel measurable functions in $(E, \rho_E) := (\mathcal{P}(S_c(\Lambda_k, \mathbb{V}), \rho), d_{\text{LP}})$. Therefore, the assumptions of Proposition 3.6 are fulfilled and for any $\epsilon > 0$, there is a compact set $K_\epsilon \subset B$ with $\text{Leb}(B \setminus K_\epsilon) < \epsilon$ such that $F_n(s_n) \rightarrow G(s)$ so long as $s_n, s \in K_\epsilon$ and $s_n \rightarrow s$.

It follows from Lebesgue's density theorem that there is a subset K'_ϵ of K_ϵ on which the Lebesgue density is equal to 1 and $\text{Leb}(K_\epsilon \setminus K'_\epsilon) = 0$. By Lemma 3.5 of [Falconer \[2002\]](#), for any $s \in K'_\epsilon$ and $w \in \mathbb{R}^d$, there exists a sequence w_n such that $w_n \rightarrow w$ and $s_n := s + w_n/n \in K_\epsilon$. Thus, for any $s \in K'_\epsilon$, $F_n(s_n) \rightarrow G(s)$ as $n \rightarrow \infty$ or, equivalently,

$$\xi_n := \{T_{s_n, c(s_n, 1/n)}(X(s_n, (1/n) \cdot \lambda)), \lambda \in \Lambda_k\} \xrightarrow{d} \xi := \{Y_s(\lambda), \lambda \in \Lambda_k\}, \quad (3.18)$$

On the other hand, we have

$$\begin{aligned}
\xi_n &= \{T_{s_n, c(s_n, 1/n)}(X(s_n, (1/n) \cdot \lambda)), \lambda \in \Lambda_k\} \\
&= \left\{T_{s_n, c(s_n, 1/n)} \circ T_{s, c(s, 1/n)}^{-1} \circ T_{s, c(s, 1/n)}(X(s_n, (1/n) \cdot \lambda)), \lambda \in \Lambda_k\right\} \\
&= \left\{T_{s_n, c(s_n, 1/n)} \circ T_{s, c(s, 1/n)}^{-1} \circ T_{s, c(s, 1/n)}(X(s, (1/n) \cdot (w_n + \lambda))), \lambda \in \Lambda_k\right\}.
\end{aligned} \tag{3.19}$$

Since by (3.16), we have $T_{s_n, c(s_n, 1/n)} \circ T_{s, c(s, 1/n)}^{-1} \rightarrow \text{I}$, Lemma A.6 implies

$$\xi_n \xrightarrow{d} \tilde{\xi} := \{Y_s(w + \lambda), \lambda \in \Lambda_k\}, \tag{3.20}$$

which implies that $\xi \stackrel{d}{=} \tilde{\xi}$.

Finally, we take $U := \cap_{k=1}^{\infty} (B \setminus K'_{1/n})$, which is a set with measure 0. This concludes the proof. \square

We make next an important observation that Condition (3.16) holds automatically in the case when the scaling actions $T_{s, c}$ can be expressed through a single scaling action independent of the location s . This is the case in particular for the usual scalar multiplication actions (2.16) considered for example in [cf. Falconer, 2002].

Corollary 3.7. *Assume the conditions of Theorem 3.5. If the scaling action does not depend on location, i.e., $T_{s, c(s, r)} = T_{c(s, r)}$, $c > 0$, then Condition (3.16) always holds.*

The proof is given in Section A.3. We conclude this section with several remarks.

Remark 3.4. The null set U in Theorem 3.5 cannot be dropped in general. While all tangent fields are self-similar, not all of them are intrinsically stationary. Indeed, one can consider the simple example $X(t) = \|t\|^H Z$, $t \in \mathbb{R}^d$, where Z is a fixed random variable and $H > 0$. Consider the usual scalar multiplication action and observe that with $s := 0$, and $\lambda = \sum_i c_i \delta_{t_i} \in \Lambda_k$, for all $r > 0$, we have

$$X(s + r \cdot \lambda) = \sum_i c_i X(rt_i) = r^H \sum_i c_i X(t_i) = r^H X(s + \lambda).$$

That is, X is its own tangent field at $s = 0$ for all $k \geq 0$. Note that X is not intrinsically stationary if $H \notin \mathbb{N}$.

Remark 3.5. Notice that $\Lambda_{k_2} \subset \Lambda_{k_1}$ for all $0 \leq k_1 < k_2$. Therefore, all k_1 -order tangent fields are also k_2 -order tangent fields. Specifically, if (3.2) holds with $k = k_1$, then it also holds with k_2 .

Remark 3.6. As in Falconer [2002], we focus here on random fields with continuous paths. One can study the structure of generalized tangent fields for processes with discontinuous paths and potentially extend the results in Falconer [2003] which focus on the space of càdlàg functions equipped with the Skorokhod J_1 -topology. The key challenge is coming up with a suitable topology on the path-space in question which is separable and complete. Provided that this is the case, we believe that versions of Theorems 3.3 and 3.5 will continue to hold.

Remark 3.7. In principle, in the definition of the tangent field (3.2) one could apply a general scaling action on both the domain \mathbb{R}^d of the stochastic process as well as on its range \mathbb{V} . In this case, we anticipate that an analog of Theorem 3.3 will hold, where the limits are scale-invariant processes similar to the ones studied in Bierné et al. [2007], Didier et al. [2017]. Here, for simplicity, we chose to apply a general scaling action only on the range of the process and retain the usual rescaling by scalars in the domain \mathbb{R}^d .

4. Spectral theory for Hilbert space valued IRF_k's.

In this section, we develop the general correlation theory for stationary and intrinsically stationary processes taking values in a separable Hilbert space \mathbb{V} over \mathbb{C} equipped with the inner product $\langle \cdot, \cdot \rangle$. In the following section, we present a generalization of the celebrated Bochner Theorem and then in Section 4.2, we extend the Matheron spectral characterization to the class of \mathbb{V} -valued intrinsic random functions. The applications of these results to the characterization of Gaussian \mathbb{V} -valued stationary and intrinsically stationary processes requires us to carefully consider *both* real and complex Hilbert spaces (discussed in Section 4.3).

Throughout this paper, a random element X in \mathbb{V} is said to have mean zero and finite variance, together referred to as *second order*, if $\mathbb{E}[X] = 0$ and $\mathbb{E}[\|X\|^2] < \infty$, where, for definiteness, all expectations here are defined in the Bochner sense (see Section S.2.1). A process is said to be second order if each element is second order.

Denote by \mathbb{T} the collection of trace-class operators on \mathbb{V} . That is, linear operators $\mathcal{T} : \mathbb{V} \rightarrow \mathbb{V}$, with finite *trace norm*:

$$\|\mathcal{T}\|_{\text{tr}} = \sum_{j=1}^{\infty} \langle (\mathcal{T}^* \mathcal{T})^{1/2} e_j, e_j \rangle,$$

where $\{e_j\}$ is an arbitrary complete orthonormal system (CONS) on \mathbb{V} , and where \mathcal{T}^* denotes the adjoint operator of \mathcal{T} . One can show that the trace norm does not depend on the choice of the CONS and the space \mathbb{T} equipped with the trace norm is a Banach space [cf. Simon, 2015].

Recall that \mathcal{T} is self-adjoint if $\mathcal{T} = \mathcal{T}^*$. Also \mathcal{T} is positive definite (or just positive), denoted $\mathcal{T} \geq 0$, if \mathcal{T} is self-adjoint and $\langle f, \mathcal{T} f \rangle \geq 0$, for all $f \in \mathbb{V}$. The class of positive and trace-class operators will be denoted by \mathbb{T}_+ .

4.1. The Bochner Theorem.

The aim of this subsection is to review the basic properties of second order covariance-stationary processes on \mathbb{R}^d taking values in the separable Hilbert space \mathbb{V} over \mathbb{C} . We start with the important notion of positive definiteness.

Definition 4.1. A collection of operators $\{\mathcal{K}(t), t \in \mathbb{R}^d\}$ on the complex Hilbert space \mathbb{V} is said to be positive definite in the weak sense if for all $c_j \in \mathbb{C}$, $t_j \in \mathbb{R}^d$, $j = 1, \dots, n$, we have

$$\sum_{j=1}^n \sum_{j'=1}^n c_j \bar{c}_{j'} \mathcal{K}(t_j - t_{j'}) \geq 0 \text{ (operator positivity)}. \quad (4.1)$$

The classical Bochner's Theorem [cf. Bochner, 1948, Khintchine, 1934] which connects the space of positive-definite functions with range in \mathbb{C} and finite positive measures has provided a fundamental tool for constructing useful models for stationary random fields. Below we state an extension of that for the infinite-dimensional setting. To do so we need the notion of integration with respect to a \mathbb{T}_+ -valued measure.

We say that $\mu : \mathcal{B}(\mathbb{R}^d) \mapsto \mathbb{T}_+$ is a \mathbb{T}_+ -valued measure if μ is σ -additive, where $\mathcal{B}(\mathbb{R}^d)$ denotes the σ -field of Borel sets in \mathbb{R}^d . Note that *a fortiori* $\mu(\emptyset) = 0$ and μ is finite in the sense that $0 \leq \mu(B) \leq \mu(\mathbb{R}^d) \in \mathbb{T}_+$, $B \in \mathcal{B}(\mathbb{R}^d)$ as positive operators. Integration of a \mathbb{C} -valued measurable function with respect to such μ can be defined along the line of Lebesgue integration; see Section A.4.

Theorem 4.2. Let $\{\mathcal{K}(t), t \in \mathbb{R}^d\} \subset \mathbb{T}$ be a positive-definite set of trace-class operators in the sense of Definition 4.1. If \mathcal{K} is continuous at 0 in the trace norm, i.e., $\|\mathcal{K}(t) - \mathcal{K}(0)\|_{\text{tr}} \rightarrow 0$, as $t \rightarrow 0$, then there exists a unique finite \mathbb{T}_+ -valued measure μ such that

$$\mathcal{K}(t) = \int_{\mathbb{R}^d} e^{it^\top x} \mu(dx), \quad \text{for all } t \in \mathbb{R}^d. \quad (4.2)$$

Conversely, for every finite \mathbb{T}_+ -valued measure μ , Relation (4.2) yields a positive-definite set of trace class operators.

We note that Theorem 4.2 or variations of it have been mentioned in the literature. See, for instance, Kallianpur and Mandrekar [1971], Holmes [1979], Neeb [1998], Durand and Roueff [2020] and van Delft and Eichler [2020]. In Section S.4.1 of Supplement, we provide a detailed proof that uses standard arguments familiar to the readers in the statistics and probability community.

Both Neeb [1998] and van Delft and Eichler [2020] present Bochner's Theorem in terms of the following natural but stronger version of positivity.

Definition 4.3. A collection of operators $\{\mathcal{K}(t), t \in \mathbb{R}^d\}$ on \mathbb{V} is said to be completely positive definite, or just positive definite, if

$$\sum_{j=1}^n \sum_{j'=1}^n \langle f_j, \mathcal{K}(t_j - t_{j'}) f_{j'} \rangle \geq 0, \quad (4.3)$$

for all $f_j \in \mathbb{V}, t_j \in \mathbb{R}^d, j = 1, \dots, n$ and $n \in \mathbb{N}$.

Definition 4.3 simply means that the matrices $(\mathcal{K}(t_j - t_{j'}))_{n \times n}$ with operator \mathbb{T} -valued entries are self-adjoint positive definite operators on the product Hilbert space \mathbb{V}^n . For more mathematical insight into this condition, see abstract literature on Hilbert C^* -modules, e.g., Murphy [1997] and Pellonpää and Ylinen [2011]. Clearly (4.3) implies (4.1). However, observe that for every finite \mathbb{T}_+ -valued measure μ , Relation (4.2) defines a completely positive definite kernel $\mathcal{K}(\cdot)$. This entails the following curious result, already noted in Durand and Roueff [2020].

Corollary 4.4. Let $\mathcal{K} : \mathbb{R}^d \rightarrow \mathbb{T}$ be continuous at 0 in $\|\cdot\|_{\text{tr}}$. The collection of operators $\{\mathcal{K}(t), t \in \mathbb{R}^d\}$ is positive definite in the sense of Definition 4.1 if and only if it is completely positive definite in the sense of Definition 4.3.

Let now $\{X(t), t \in \mathbb{R}^d\}$ be a \mathbb{V} -valued, second order random field. The cross covariance operator for X is then well-defined as

$$\mathcal{C}_X(s, t) := \mathbb{E}[X(s) \otimes X(t)], \quad s, t \in \mathbb{R}^d,$$

and takes values in the space of trace-class operators \mathbb{T} equipped with the trace norm Lemma S.2.2 in Supplement. The latter expectation is understood to be defined in the sense of Bochner integral in the separable Banach space $(\mathbb{T}, \|\cdot\|_{\text{tr}})$, and for $f, g \in \mathbb{V}$, the outer product operator is by definition $(f \otimes g)h := \langle h, g \rangle f$. Observe that $\mathcal{C}_X(t, t)$ is positive definite, and

$$\mathcal{C}_X(s, t) = \mathcal{C}_X^*(t, s), \quad \text{for all } s, t \in \mathbb{R}^d.$$

The process X is said to be mean-square or L^2 -continuous if

$$\mathbb{E}\|X(s) - X(t)\|^2 \rightarrow 0 \text{ as } s \rightarrow t, \text{ for all } t \in \mathbb{R}^d. \quad (4.4)$$

It is easy to see (Section S.2.2 or Proposition S.2.3 in Supplement) that X is mean-square continuous if and only if $\|\mathcal{C}_X(s', t') - \mathcal{C}_X(t, t)\|_{\text{tr}} \rightarrow 0$ as $(s', t') \rightarrow (t, t)$.

Definition 4.5. A process X is said to be weakly or covariance-stationary if it is second order and its cross covariance is shift invariant, i.e.,

$$\mathcal{K}_X(h) := \mathcal{C}_X(s, s+h), \quad \text{for all } h \in \mathbb{R}^d,$$

does not depend on $s \in \mathbb{R}^d$. The function $\mathcal{K}_X(h)$, $h \in \mathbb{R}^d$, will be referred to as the stationary covariance function of X .

Observe that every stationary covariance function \mathcal{K}_X is positive definite in the sense of (4.3) (and hence (4.1)). Also, the L^2 -continuity of a stationary process is equivalent to the continuity of its stationary covariance function at 0. Thus, the characterization in Theorem 4.2 readily holds for the stationary covariance of a stationary process that is L^2 -continuous.

We conclude this section with a version of the classical Cramér stochastic representation of stationary Hilbert-space-valued random fields. Recall that, for $\mathbb{V} = \mathbb{C}$, the well-known integral representation [cf. Cramér, 1942] of a covariance stationary random process X on \mathbb{R}^d states that

$$X(t) = \int_{\mathbb{R}^d} e^{it^\top x} \xi(dx), \quad \text{almost surely, } t \in \mathbb{R}^d, \quad (4.5)$$

where ξ is a second order random measure with orthogonal increments. To extend this result to a general \mathbb{V} , we first have to define integration with respect to a random measure with orthogonal increments in that setting. This is done in Section A.4 of Appendix. Here, we only give the main ideas.

Let $\mathbb{L}^2(\Omega)$ be the L^2 space of all \mathbb{V} -valued random elements η on the probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with $\mathbb{E}[\|\eta\|^2] < \infty$, equipped with the inner product

$$\langle \eta_1, \eta_2 \rangle_\Omega := \mathbb{E} \langle \eta_1, \eta_2 \rangle.$$

Definition 4.6. Let μ be a \mathbb{T}_+ -valued measure on $\mathcal{B}(\mathbb{R}^d)$. A second order stochastic process $\xi = \{\xi(A), A \in \mathcal{B}(\mathbb{R}^d)\} \subset \mathbb{L}^2(\Omega)$ indexed by the Borel sets is said to be a \mathbb{V} -valued orthogonal random measure on \mathbb{R}^d with control measure μ if the following conditions hold:

- (i) $\mathbb{E}[\|\xi(A_n)\|^2] \rightarrow 0$ if $A_n \rightarrow \emptyset$,
- (ii) $\mu(A \cap B) = \mathbb{E}[\xi(A) \otimes \xi(B)]$, for all $A, B \in \mathcal{B}(\mathbb{R}^d)$, where the expectation is in the sense of Bochner on $(\mathbb{T}, \|\cdot\|_{\text{tr}})$.

It is straightforward to see that (ii) implies for disjoint A and B , that $\xi(A \cup B) = \xi(A) + \xi(B)$, almost surely, so that ξ is in fact an additive set-function. This, combined with the continuity property (i), yields the σ -additivity of ξ (for more details, see Section A.4).

For a simple function $f(t) = \sum_{i=1}^n c_i I_{A_i}(t)$, with $c_i \in \mathbb{C}$ and pairwise disjoint A_i 's, we naturally define $\mathcal{J}_\xi(f) := \int_{\mathbb{R}^d} f(t) \xi(dt) := \sum_{i=1}^n c_i \xi(A_i)$. Letting $\|\mu\|_{\text{tr}}(A) := \|\mu(A)\|_{\text{tr}}$ be the trace measure of μ , we see that

$$\mathbb{E}[\|\mathcal{J}_\xi(f)\|^2] = \sum_{i=1}^n |c_i|^2 \mathbb{E}[\|\xi(A_i)\|^2] = \int_{\mathbb{R}^d} |f(t)|^2 \|\mu\|_{\text{tr}}(dt).$$

That is, the linear operator \mathcal{J}_ξ is an isometry from the space of simple functions in $L^2(\mathbb{R}^d, \|\mu\|_{\text{tr}})$ into the Hilbert space $\mathbb{L}^2(\Omega)$. Thus, one can extend the definition of $\mathcal{J}_\xi(f)$, by continuity, to all $f \in L^2(\mathbb{R}^d, \|\mu\|_{\text{tr}})$. We have moreover that, for all $f, g \in L^2(\mathbb{R}^d, \|\mu\|_{\text{tr}})$,

$$\mathbb{E}[\mathcal{J}_\xi(f) \otimes \mathcal{J}_\xi(g)] = \int_{\mathbb{R}^d} f(t) \overline{g(t)} \mu(dt),$$

where the latter integral is in the sense of Bochner (cf Section A.4 of Appendix).

Theorem 4.7. *Let $X = \{X(t), t \in \mathbb{R}^d\}$ be an L^2 -continuous, weakly stationary process taking values in the separable Hilbert space \mathbb{V} and having stationary covariance function \mathcal{K} .*

Then, (4.2) holds and there exists an a.s unique orthogonal \mathbb{V} -valued random measure ξ with control measure μ , such that (4.5) holds.

The proof of this result can be found in Section S.4.3.

4.2. Spectral theory for general IRF_k .

Gel'fand and Vilenkin [1964] provide an illuminating treatment of the spectral theory of generalized stochastic processes, i.e., processes with paths in the space of generalized functions. In this setting, the paths of the stochastic process have derivatives of all orders and one can naturally study processes with stationary $(1+k)$ -th order derivatives. One drawback of this treatment is that it is difficult to use generalized process models in practice. Motivated by fundamental problems in spatial statistics, Matheron [1973] developed the framework of intrinsic stationary functions, which allows one to study classical random field models with stationary increments.

In a series of works, Matheron developed the theory of intrinsic random functions, which has become the *lingua franca* of spatial statistics [see e.g. Chilès and Delfiner, 2012]. Our goal here is to extend the Matheron theory to the functional setting, where the underlying stochastic processes take values in a separable Hilbert space \mathbb{V} . This is not straightforward and new *covariance asymmetry* phenomena arise that reflect the potential *irreversibility* of multivariate IRF's (see Remark 5.10).

Following Matheron [1973], in this section we will focus on second order linear processes $Y = \{Y(\lambda), \lambda \in \Lambda_k\}$, viewed in the weak sense as stochastic processes indexed by Λ_k . That is, Y is a random element in $S(\Lambda_k, \mathbb{V})$ equipped with the product σ -field $\mathcal{B}_{\mathbb{V}^{\Lambda_k}}$ generated by all finite-dimensional cylinder sets. We emphasize that, in contrast to Section 3, here we no longer require that Y has continuous paths. The (cross) covariance operator of Y is defined as

$$\mathcal{C}_Y(\lambda, \mu) := \mathbb{E}[Y(\lambda) \otimes Y(\mu)], \quad \lambda, \mu \in \Lambda_k.$$

Denote by $\check{Y} = \{\check{Y}(t), t \in \mathbb{R}^d\}$ the representation of Y (cf. Section 2.1) in $\check{S}(\Lambda_k, \mathbb{V})$, i.e., $\check{Y}(t) = Y(\lambda_t)$, so that

$$Y(\lambda) := \int \check{Y}(t) \lambda(dt), \quad \lambda \in \Lambda_k. \quad (4.6)$$

We say that Y is mean-square continuous if \check{Y} is mean-square continuous in the sense of (4.4).

Definition 4.8. *A second order process $\{Y(\lambda), \lambda \in \Lambda_k\} \in S(\Lambda_k, \mathbb{V})$ is said to be an intrinsic random function of order k (IRF_k), $k = -1, 0, 1, \dots$, if*

$$\mathcal{C}_Y(\lambda, \mu) \equiv \mathcal{C}_Y(w + \lambda, w + \mu), \quad w \in \mathbb{R}^d, \lambda, \mu \in \Lambda_k. \quad (4.7)$$

Note that (4.7) is equivalent to

$$\mathcal{C}_Y(\lambda, \lambda) \equiv \mathcal{C}_Y(w + \lambda, w + \lambda), \quad w \in \mathbb{R}^d, \lambda \in \Lambda_k, \quad (4.8)$$

by Lemma A.8, and, in turn, to the weak stationarity of $\{Y(t + \lambda), t \in \mathbb{R}^d\}$ in t for all $\lambda \in \Lambda_k$. Indeed, if $Y(t + \lambda)$ is stationary in t for all λ then (4.8) holds, and if (4.7) holds then $Y(t + \lambda)$ is stationary in t for all λ .

Definition 4.9. A collection of trace-class operators $\{\mathcal{K}(h), h \in \mathbb{R}^d\} \subset \mathbb{T}$ is said to be conditionally positive definite of degree k , $k = -1, 0, 1, \dots$, if for all $n \geq 1$,

$$\sum_{j=1}^n \sum_{j'=1}^n c_j \bar{c}_{j'} \mathcal{K}(t_j - t_{j'}) \geq 0 \quad (\text{operator positivity}) \quad (4.9)$$

for all $c_i \in \mathbb{C}, t_j \in \mathbb{R}^d, 1 \leq i \leq n$, such that $\lambda(du) = \sum_j c_j \delta_{t_j}(du) \in \Lambda_k$.

Relation (4.9) can be succinctly written as $\mathcal{K}(\lambda * \tilde{\lambda}) \geq 0$, where

$$\tilde{\lambda}(du) := \bar{\lambda}(-du) = \sum_j \bar{c}_j \delta_{-t_j}(du),$$

and $\lambda * \mu$ denotes the usual convolution. More generally, with $\lambda = \sum_j c_j \delta_{t_j}$ and $\mu = \sum_{j'} d_{j'} \delta_{s_{j'}}$ $\in \Lambda_k$,

$$\mathcal{K}(\lambda * \tilde{\mu}) = \sum_j \sum_{j'} c_j \bar{d}_{j'} \mathcal{K}(t_j - s_{j'}). \quad (4.10)$$

Interestingly, since $(w + \lambda) * \widetilde{(w + \mu)} \equiv \lambda * \tilde{\mu}$, for all $w \in \mathbb{R}^d$, the map $(\lambda, \mu) \mapsto \mathcal{K}(\lambda * \tilde{\mu})$ is automatically shift invariant. This motivates the following definition.

Definition 4.10. A collection of operators $\mathcal{K} : \mathbb{R}^d \rightarrow \mathbb{T}$ is said to be a generalized covariance of Y with degree k if

$$\mathcal{C}_Y(\lambda, \mu) = \mathcal{K}(\lambda * \tilde{\mu}), \quad \text{for all } \lambda, \mu \in \Lambda_k. \quad (4.11)$$

Again, by Lemma A.8, (4.11) is equivalent to

$$\mathcal{C}_Y(\lambda, \lambda) = \mathcal{K}(\lambda * \tilde{\lambda}), \quad \text{for all } \lambda \in \Lambda_k.$$

The following result describes the connections between the notions in Definitions 4.8-4.10, and gives a *spectral representation* of a conditionally positive definite \mathcal{K} . As a terminology, a polynomial in \mathbb{T} refers to a linear combination of d -dimensional monomials with coefficients in \mathbb{T} , where the degree is equal to the highest degree of the monomials in the linear combination.

Theorem 4.11. Let $k \geq -1$ and the process $Y = \{Y(\lambda), \lambda \in \Lambda_k\} \in S(\Lambda_k, \mathbb{V})$ be second order.

- (i) If Y has a generalized covariance \mathcal{K} of degree k , then Y is IRF_k and \mathcal{K} must be conditionally positive definite of degree k . Conversely, if Y is a mean-square continuous IRF_k , then it has a continuous generalized covariance of degree k .
- (ii) A continuous function $\mathcal{K} : \mathbb{R}^d \rightarrow \mathbb{T}$ is conditionally positive definite of degree k if and only if it can be represented as

$$\mathcal{K}(h) = \int_{\mathbb{R}^d} \frac{e^{\mathbf{i}u^\top h} - I_B(u)P(u^\top h)}{1 \wedge \|u\|^{2k+2}} \chi(du) + \mathcal{Q}(h), \quad (4.12)$$

where $P(x) = \sum_{j=0}^{2k+1} (\mathbf{i}x)^j / j!$, B is some arbitrary bounded neighborhood of 0, $\mathcal{Q}(h)$ is a conditionally positive definite polynomial with degree no more than $2k + 2$ and χ is a finite \mathbb{T}_+ -valued measure with no point mass at 0. The measure χ in (4.12) is unique and does not depend on the choice of the set B . The polynomial \mathcal{Q} therein is unique modulo an additive polynomial of degree $2k + 1$.

The detailed proof of this result can be found in Section S.4.2 of Supplement. Note that the proof follows closely the general and elegant treatment of Sasvári [2009].

Remark 4.1. In the notation of Sasvári [2009], our situation corresponds to having a single multiplicative function (character) $\gamma_1 \equiv 1$ and $y_1 := 0$ and $k_1 := k + 1$ and their measure σ is our $(1 \wedge \|u\|^{2k+2})^{-1} \chi(du)$. Observe also that Relation (4.3) in Theorem 4.2 of Sasvári [2009] appears to be missing the non-ignorable degree $2k+2$ polynomial component in \mathcal{Q} of (4.12). This omission can be attributed to the fact that the spectral measure of a stationary process in the Bochner theorem could have an atom at $\{0\}$, while σ and χ do not. See Section S.4.2 for more details.

The measure χ and polynomial \mathcal{Q} in (4.12) will be referred to as the *spectral characteristics* of an IRF_k with generalized covariance \mathcal{K} . Note that the *spectral characteristics* pair (χ, \mathcal{Q}) is unique modulo an additive polynomial of degree $2k+1$ in the component \mathcal{Q} . That is, the generalized covariance in (4.12) is unique up to an additive polynomial of order $2k+1$. This implies that $\mathcal{K}(\lambda * \tilde{\mu})$ is uniquely determined for $\lambda, \mu \in \Lambda_k$, where $\mathcal{K}(\nu) := \int \mathcal{K}(h) \nu(dh)$, $\nu \in \Lambda$. Notice that $\lambda * \tilde{\mu} \in \Lambda_{2k+1}$, for $\lambda, \mu \in \Lambda_k$. Thus, in view of (4.11), the covariance structure of an IRF_k process is completely determined by the linear measure-indexed \mathbb{T} -valued function $\mathcal{K}(\nu)$, $\nu \in \Lambda_{2k+1}$. By integrating (4.12) with respect to $\nu \in \Lambda_{2k+1}$, we obtain

$$\mathcal{K}(\nu) = \int_{\mathbb{R}^d} \frac{\widehat{\nu}(u)}{1 \wedge \|u\|^{2k+2}} \chi(du) + \mathcal{Q}(\nu), \quad \nu \in \Lambda_{2k+1}, \quad (4.13)$$

where $\widehat{\nu}(u) = \int e^{iu^\top x} \nu(dx)$ is the Fourier transform of ν . Since $\widehat{\lambda * \tilde{\mu}} = \widehat{\lambda} \widehat{\tilde{\mu}}$, the cross covariance operator $\mathcal{C}_Y(\lambda, \mu)$ of Y can be uniquely expressed as

$$\mathcal{C}_Y(\lambda, \mu) = \mathcal{K}(\lambda * \tilde{\mu}) = \int_{\mathbb{R}^d} \frac{\widehat{\lambda}(u) \widehat{\tilde{\mu}}(u)}{1 \wedge \|u\|^{2k+2}} \chi(du) + \mathcal{Q}(\lambda * \tilde{\mu}). \quad (4.14)$$

Now, consider the following counterpart to Definition 4.3.

Definition 4.12. A collection of trace-class operators $\{\mathcal{K}(h), h \in \mathbb{R}^d\} \subset \mathbb{T}$ is said to be conditionally complete positive definite of degree k , $k = -1, 0, 1, \dots$, if

$$\sum_{j=1}^n \sum_{j'=1}^n \langle f_j, \mathcal{K}(\mu_j * \tilde{\mu}_{j'}) f_{j'} \rangle \geq 0, \quad (4.15)$$

for all $f_j \in \mathbb{V}$, $\mu_j \in \Lambda_k$, $j = 1, \dots, n$ and $n \in \mathbb{N}$.

Since (4.9) is the special case of (4.15) with $n = 1$, conditional complete positive definiteness implies conditional positive definiteness. However, as seen from (4.14), $\mathcal{K}(\lambda * \tilde{\mu})$, $\lambda, \mu \in \Lambda_k$ is a valid cross-covariance, and hence the operator function \mathcal{K} in (4.12) is conditionally complete positive definite. Thus, Theorem 4.11 implies the following parallel of Corollary 4.4.

Corollary 4.13. If $\mathcal{K} : \mathbb{R}^d \rightarrow \mathbb{T}$ is continuous, then Definitions 4.9 and 4.12 are equivalent.

We end this section with a stochastic representation result for continuous IRF_k , which parallels the Cramér representation in Theorem 4.7. The proof is given in Section S.4.4.

Theorem 4.14. Let $k \geq -1$ and the process $Y = \{Y(\lambda), \lambda \in \Lambda_k\}$ in $S(\Lambda_k, \mathbb{V})$ be mean-square continuous. Then Y is IRF_k if and only if it can be uniquely represented as

$$Y(\lambda) = \sum_{(j_1, \dots, j_d) \in J} \frac{\partial^{k+1} \widehat{\lambda}}{\partial^{j_1} \dots \partial^{j_d}}(0) \cdot Z_{j_1 \dots j_d} + \int_{\mathbb{R}^d} \frac{\widehat{\lambda}(u)}{1 \wedge \|u\|^{k+1}} \xi(du), \quad \lambda \in \Lambda_k, \quad a.s. \quad (4.16)$$

where

- (i) $J = \{(j_1, \dots, j_d) : j_1, \dots, j_d \geq 0 \text{ and } j_1 + \dots + j_d = k + 1\}$,
- (ii) ξ is an a.s. unique random orthogonal measure ξ on $(\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d))$ with control measure χ , where χ is a finite \mathbb{T}_+ -valued measure with no point mass at 0, and
- (iii) the $Z_{j_1 \dots j_d}$ are uncorrelated random variables with values in \mathbb{V} and are uncorrelated with ξ .

Remark 4.2. [Berschneider \[2012\]](#) also obtains the stochastic representation of IRF_k with the more abstract setting of locally compact Abelian domains. Our result here can be considered as an extension to the case of \mathbb{V} -valued processes connecting $\xi(dx)$ with the covariance operator functions in Theorem [4.11](#) explicitly.

4.3. Real and complex IRF_k 's: Covariance (ir)reversibility.

The general treatment in the previous two subsections involves an abstract separable Hilbert space \mathbb{V} over the field of complex numbers \mathbb{C} . In practice, however, one often deals with Hilbert spaces of real-valued functions and it is useful to know how our results specialize to this setting. Furthermore, the distribution of a zero-mean Gaussian process taking values in a *complex* Hilbert space \mathbb{V} cannot be *directly* characterized using their covariance structure, alone. To this end one needs to consider both real and complex Hilbert spaces.

Real and imaginary parts in a complex Hilbert space. In an abstract complex Hilbert space \mathbb{V} the notions of a real and imaginary part of an element $z \in \mathbb{V}$ are not well defined unless one fixes a basis. Let $\mathcal{E} := \{e_j, j \in \mathbb{N}\}$ be a *fixed* CONS of \mathbb{V} . Then one can postulate that the CONS \mathcal{E} is *real* and for each $z = \sum_j z_j e_j \in \mathbb{V}$, with coordinates $z_j := \langle z, e_j \rangle$, we can define

$$\Re(z) \equiv \Re_{\mathcal{E}}(z) := \sum_j \Re(z_j) e_j \quad \text{and} \quad \Im(z) \equiv \Im_{\mathcal{E}}(z) := \sum_j \Im(z_j) e_j, \quad (4.17)$$

as the *real* and *imaginary* parts of z , relative to the CONS \mathcal{E} . (Should one change the basis \mathcal{E} the notions of real and imaginary part may change.) Notice that $\mathbb{V}_{\mathbb{R}} := \{z \in \mathbb{V} : \Im(z) = 0\}$ is invariant to addition and multiplication by real scalars and it becomes a real Hilbert space, with the inner product inherited from \mathbb{V} . All elements of \mathbb{V} that belong to $\mathbb{V}_{\mathbb{R}}$ will be referred to as *real*.

For $z \in \mathbb{V}$, we shall write $z = \Re(z) + i\Im(z)$ and naturally define the *complex conjugate* $\bar{z} := \Re(z) - i\Im(z)$. The complex conjugate operation as well as the real and imaginary part operators extend to \mathbb{V} -valued random elements in a straightforward manner and we shall say that $x \in \mathbb{V}$ is *real* if $x \in \mathbb{V}_{\mathbb{R}}$, i.e., if its imaginary part is zero.

The complex conjugate of a linear operator $\mathcal{A} : \mathbb{V} \rightarrow \mathbb{V}$ is defined as: $\overline{\mathcal{A}}(x) := \overline{\mathcal{A}(\bar{x})}$, $x \in \mathbb{V}$. This implies that $\overline{\mathcal{A}(x)} = \overline{\mathcal{A}}(\bar{x})$, the operator $\overline{\mathcal{A}}$ is also *linear* and one can define the real and imaginary parts of \mathcal{A} in as: $\Re(\mathcal{A}) := (\mathcal{A} + \overline{\mathcal{A}})/2$ and $\Im(\mathcal{A}) := (\mathcal{A} - \overline{\mathcal{A}})/2i$. Thus, $\mathcal{A} = \Re(\mathcal{A}) + i\Im(\mathcal{A})$ and the usual operations with complex numbers and vectors extend to the operator Banach algebra over the complex Hilbert space \mathbb{V} . Note that the real and imaginary parts of \mathcal{A} can be equivalently defined in terms of the real and imaginary parts of the coordinates of \mathcal{A} in the fixed CONS \mathcal{E} . We shall say that an operator \mathcal{A} is *real* if $\mathcal{A} = \Re(\mathcal{A})$.

Real IRF_k 's. The above discussion shows how one can specialize and interpret the results in Sections [4.1](#) and [4.2](#) for the case of real Hilbert spaces $\mathbb{V}_{\mathbb{R}}$. Indeed, let $\Lambda_k(\mathbb{R})$ be the set of all real $\lambda \in \Lambda_k$. It is easy to see that $\Lambda_k = \Lambda_k(\mathbb{R}) + i\Lambda_k(\mathbb{R})$.

Suppose now that Y is a $\mathbb{V}_{\mathbb{R}}$ -valued IRF_k . That is, Definition 4.8 holds with Λ_k replaced by $\Lambda_k(\mathbb{R})$. Then, by linearity, Y extends uniquely to a \mathbb{V} -valued IRF_k as follows

$$Y(\lambda) := Y(\Re(\lambda)) + \mathbf{i}Y(\Im(\lambda)), \quad \lambda \in \Lambda_k, \quad (4.18)$$

where in fact $Y(\Re(\lambda))$ and $Y(\Im(\lambda))$ are real (belong to $\mathbb{V}_{\mathbb{R}}$). This leads us to the following

Definition 4.15. A \mathbb{V} -valued IRF_k Y is said to be real if $Y(\lambda)$ is real for all $\lambda \in \Lambda_k(\mathbb{R})$.

Thus, there is a one-to-one correspondence between the real IRF_k 's in \mathbb{V} and the $\mathbb{V}_{\mathbb{R}}$ -valued IRF_k 's as processes indexed by $\Lambda_k(\mathbb{R})$.

Proposition 4.16. Let Y be a mean-square continuous IRF_k taking values in \mathbb{V} and having spectral characteristics (χ, \mathcal{Q}) .

- (i) Y is real if and only if in Relation (4.16) the vectors $(\mathbf{i})^{k+1}Z_{j_1 \dots j_d}$ are real and the orthogonal measure ξ is Hermitian, i.e., $\xi(-A) = \overline{\xi(A)}$, almost surely, for all $A \in \mathcal{B}(\mathbb{R}^d \setminus \{0\})$.
- (ii) If Y is real, then \mathcal{Q} has real (operator) coefficients (modulo polynomials of degree up to $2k+1$) and the spectral measure χ is Hermitian, i.e., $\overline{\chi(-A)} = \chi(A)$, for all $A \in \mathcal{B}(\mathbb{R}^d \setminus \{0\})$.
- (iii) Conversely, if χ is Hermitian and \mathcal{Q} real, then there is a real IRF_k Y with spectral characteristics (χ, \mathcal{Q}) . Let now Y and Y' be two real IRF_k with the same spectral characteristics (χ, \mathcal{Q}) and such that $\mathbb{E}[Y(\lambda) \otimes Y'(\mu)] = \mathbb{E}[Y'(\lambda) \otimes Y(\mu)]$, $\lambda, \mu \in \Lambda_k$. Then, for any $a, b \in \mathbb{R}$, with $a^2 + b^2 = 1$, the IRF_k defined as $\tilde{Y}(\lambda) = aY(\lambda) + \mathbf{i}bY'(\lambda)$ has the same spectral characteristics as Y and Y' .

The proof is given in Section A.4 of the Appendix.

Covariance (ir)reversibility. Next, we comment on an important covariance irreversibility phenomenon, which arises in the case of vector valued processes. It extends the notion of *time reversibility* for vector time series.

Definition 4.17. We shall say that an IRF_k Y with generalized operator covariance $\mathcal{K}(\cdot)$ is covariance-symmetric or -reversible if

$$\mathcal{K}(\nu) = \mathcal{K}((-1) \cdot \nu), \quad \text{for all } \nu \in \Lambda_{2k+1}. \quad (4.19)$$

Observe that the symmetry of the generalized covariance is equivalent to the fact that the IRF_k processes $\{Y((-1) \cdot \lambda)\}$ and $\{Y(\lambda)\}$ have the same covariance structure. Indeed, recall that

$$\mathbb{E}[Y(\lambda) \otimes Y(\mu)] = \mathcal{C}_Y(\lambda, \mu) = \mathcal{K}(\lambda * \tilde{\mu}), \quad \lambda, \mu \in \Lambda_k$$

and observe that $(-1) \cdot \lambda * \tilde{\mu} = ((-1) \cdot \lambda) * (\widetilde{(-1) \cdot \mu})$. Thus, the IRF_k process $\tilde{Y}(\lambda) := Y((-1) \cdot \lambda)$ has covariance $\mathcal{K}((-1) \cdot \nu)$.

In the simple case, where Y takes real scalar values, all IRF_k 's are automatically covariance-symmetric. This is perhaps why symmetry is often taken for granted. In the multivariate and especially function-valued case, however, covariance-symmetry is an exception rather than a rule. Naturally, in view of (4.13) and the uniqueness of the spectral measure, (4.19) holds if and only if $\chi(-A) = \chi(A)$, for all $A \in \mathcal{B}(\mathbb{R}^d \setminus \{0\})$. This simple observation and Proposition 4.16 yield the following fact [see also Theorem 5.1 in Didier and Pipiras, 2011].

Proposition 4.18. A real mean-square continuous IRF_k Y is covariance symmetric, if and only if its spectral measure χ is real.

We end this section with a comment on the use of the results from Sections 4.1 and 4.2 in the context of Gaussian processes. Recall that a \mathbb{V} -valued random element Y is said to be Gaussian, if $\langle Y, f \rangle$ is a complex Gaussian variable, for each $f \in \mathbb{V}$. This means that the joint distribution of $(\Re(\langle Y, f \rangle), \Im(\langle Y, f \rangle))$ is bivariate normal, for all $f \in \mathbb{V}$. Equivalently, Y is Gaussian in \mathbb{V} if and only if $\tilde{Y} := (\Re(Y), \Im(Y))$ is a Gaussian element in the *real* Hilbert space $\mathbb{V}_{\mathbb{R}}^2 := \mathbb{V}_{\mathbb{R}} \times \mathbb{V}_{\mathbb{R}}$.

Remark 4.3 (Characterization of Gaussian IRF_k 's). Part (iii) of Proposition 4.16 is a manifestation of the fact that the covariance structure alone does not determine the distribution of zero-mean Gaussian processes taking values in *complex* Hilbert spaces (cf Example S.5.1 in Supplement). To determine the distribution of a zero-mean Gaussian IRF_k $Y = \{Y(\lambda), \lambda \in \Lambda_k\}$, one needs to know both the *cross-covariance* and *pseudo cross-covariance* operators: $\mathcal{C}_Y(\lambda, \mu) = \mathbb{E}[Y(\lambda) \otimes Y(\mu)]$ and $\mathcal{C}_{Y, \bar{Y}}(\lambda, \mu) = \mathbb{E}[Y(\lambda) \otimes \bar{Y}(\mu)]$ (see, e.g., Section S.5.1 and Corollary S.5.3 in Supplement).

Equivalently, the distribution of a \mathbb{V} -valued Gaussian IRF_k is completely determined by the real IRF_k $\tilde{Y}(\cdot) := (\Re(Y)(\cdot), \Im(Y)(\cdot))$ in the product space \mathbb{V}^2 . Since the law of the real Gaussian IRF_k \tilde{Y} is determined by its cross-covariance, the results of Sections 4.1 and 4.2 provide a complete characterization of the \mathbb{V} -valued Gaussian IRF_k 's.

5. Second order covariance self-similar IRF_k 's.

In view of Theorems 3.3 and 3.5, essentially all tangent fields are self-similar IRF_k . This motivates a more in-depth study of self-similar IRF_k 's. In this section, we focus on second order covariance self-similar IRF_k 's with respect to *linear* operator-scaling actions. We establish their covariance structure and spectral representation. Section 5.1 addresses the general case, Section 5.2 discusses examples and related work, while Section 5.3 deals with the scalar scaling action, where the generalized covariance can be written in closed form.

5.1. Linear operator scaling.

Let $\mathcal{H} : \mathbb{V} \rightarrow \mathbb{V}$ be a bounded *linear operator* on the Hilbert space \mathbb{V} . Consider the operator scaling actions $T_c := c^{\mathcal{H}}$, $c > 0$, where $c^{\mathcal{H}}$ is interpreted as $\exp\{\log(c)\mathcal{H}\}$ and as usual,

$$c^{\mathcal{H}} := \sum_{n=0}^{\infty} \frac{\mathcal{H}^n}{n!}. \quad (5.1)$$

The latter series converges in operator norm and $\|c^{\mathcal{H}}\|_{\text{op}} \leq e^{\|\mathcal{H}\|_{\text{op}}}$. We have moreover that if the bounded operators \mathcal{H}_1 and \mathcal{H}_2 *commute*, i.e., $\mathcal{H}_1\mathcal{H}_2 = \mathcal{H}_2\mathcal{H}_1$, then $e^{\mathcal{H}_1}e^{\mathcal{H}_2} = e^{\mathcal{H}_1+\mathcal{H}_2} = e^{\mathcal{H}_2}e^{\mathcal{H}_1}$. This readily implies that $e^{\mathcal{H}}$ has a bounded inverse $(e^{\mathcal{H}})^{-1} = e^{-\mathcal{H}}$. Consequently, $c^{\mathcal{H}}$, $c > 0$ is a strongly (operator) continuous and invertible group action on \mathbb{V} , i.e., $c_1^{\mathcal{H}}c_2^{\mathcal{H}} = (c_1c_2)^{\mathcal{H}}$, $c_1, c_2 > 0$. In fact, using the power-series representation (5.1), one can readily show that $c \mapsto c^{\mathcal{H}}$ is continuously Fréchet differentiable with derivative $c^{\mathcal{H}-1}\mathcal{H}$, i.e.,

$$\left\| \frac{1}{h}((c+h)^{\mathcal{H}} - c^{\mathcal{H}}) - c^{\mathcal{H}-1}\mathcal{H} \right\|_{\text{op}} \rightarrow 0, \quad \text{as } h \rightarrow 0, \quad (5.2)$$

where here and below $\mathcal{H} - a$, $a \in \mathbb{R}$ is interpreted as $\mathcal{H} - a\text{I}$, so that $c^{\mathcal{H}-1} = c^{-1}c^{\mathcal{H}}$.

As in Definition 3.2, we consider the following notion of *covariance* operator self-similarity.

Definition 5.1. Fix an arbitrary bounded linear operator \mathcal{H} on \mathbb{V} . A second order IRF_k Y is said to be *covariance \mathcal{H} -self-similar*, if $\{Y(c \cdot \lambda), \lambda \in \Lambda_k\}$ and $\{c^{\mathcal{H}}Y(\lambda), \lambda \in \Lambda_k\}$ have the same operator cross-covariance function for all $c > 0$.

Remark 5.1. If the IRF_k process Y is *real* and Gaussian, then Y is covariance \mathcal{H} -self-similar if and only if it is \mathcal{H} -self-similar in the following stronger sense:

$$\{Y(c \cdot \lambda), \lambda \in \Lambda_k\} \stackrel{fdd}{=} \{c^{\mathcal{H}} Y(\lambda), \lambda \in \Lambda_k\}, \quad \text{for all } c > 0. \quad (5.3)$$

We emphasize that \mathcal{H} in Definition 5.1 and (5.3) is an arbitrary bounded linear operator and we do not require that $c \mapsto c^{\mathcal{H}}$ be a scaling action on \mathbb{V} in the sense of Definition 2.5 (see also Remark 2.5). If $T_c := c^{\mathcal{H}}$ is a scaling action, however, then (5.3) recovers the notion of self-similarity in Definition 3.2.

Let now Y be a second order, mean-square continuous IRF_k with operator auto-covariance \mathcal{H} and spectral characteristics (χ, \mathcal{Q}) . By Theorem 4.14, we have the decomposition

$$Y(\lambda) = Y_{(0, \mathcal{Q})}(\lambda) + Y_{(\chi, 0)}(\lambda), \quad \text{almost surely,} \quad (5.4)$$

for all $\lambda \in \Lambda_k$, where $\{Y_{(0, \mathcal{Q})}(\lambda)\}$ and $\{Y_{(\chi, 0)}(\lambda)\}$ are orthogonal mean-square continuous IRF_k 's with spectral characteristics $(0, \mathcal{Q})$ and $(\chi, 0)$, respectively. This decomposition is second order unique. Therefore, it follows that Y is covariance \mathcal{H} -self-similar *if and only if* both the components $Y_{(0, \mathcal{Q})}$ and $Y_{(\chi, 0)}$ are covariance \mathcal{H} -self-similar. More precisely, we have the following general characterization of covariance \mathcal{H} -self-similar IRF_k 's. For convenience, write

$$\chi_k(dx) = \frac{\chi(dx)}{1 \wedge \|x\|^{2k+2}}. \quad (5.5)$$

Theorem 5.2. *Let Y be an L^2 -continuous IRF_k with spectral characteristics (χ, \mathcal{Q}) . Let also \mathcal{H} be a bounded linear operator.*

(i) *We have that Y is covariance \mathcal{H} -self-similar if and only if for all $c > 0$ and $\lambda, \mu \in \Lambda_k$*

$$\chi_k(dx) = c^{-\mathcal{H}} \chi_k(dx/c) c^{-\mathcal{H}^*} \quad \text{and} \quad \mathcal{Q}(\lambda * \tilde{\mu}) = c^{k+1-\mathcal{H}} \mathcal{Q}(\lambda * \tilde{\mu}) c^{k+1-\mathcal{H}^*}, \quad (5.6)$$

where χ_k is as in (5.5).

Suppose henceforth that (5.6) holds and consider the polar coordinates $(r, \theta) := (\|x\|, x/\|x\|)$ in $\mathbb{R}^d \setminus \{0\}$.

(ii) *There exists a finite \mathbb{T}_+ -valued measure σ on the unit sphere $\mathbb{S} = \{\|x\| = 1\}$ such that*

$$\chi_k(D) = \int_0^\infty r^{-\mathcal{H}} \left(\int_{\mathbb{S}} 1_D(r\theta) \sigma(d\theta) \right) r^{-\mathcal{H}^*} \frac{dr}{r}, \quad (5.7)$$

for all Borel sets $D \in \mathcal{B}(\mathbb{R}^d \setminus \{0\})$ that are bounded away from 0. If (5.7) holds, we simply write

$$\chi_k(dr d\theta) = r^{-\mathcal{H}} \sigma(d\theta) r^{-\mathcal{H}^*} \frac{dr}{r} \quad (5.8)$$

and refer to (5.8) as a disintegration formula for χ_k .

(iii) *The measure σ in (5.8) is uniquely determined by the measure χ_k and it does not depend on the possibly non-unique operator \mathcal{H} in (5.6).*

(iv) The component $Y_{(\chi,0)}$ of Y admits the Cramér-type stochastic integral representation

$$Y_{(\chi,0)}(\lambda) = \int_0^\infty \int_{\mathbb{S}} \widehat{\lambda}(r\theta) W(dr, d\theta), \quad \text{almost surely,} \quad (5.9)$$

$\lambda \in \Lambda_k$, where $W(dr, d\theta)$ is an orthogonal \mathbb{V} -valued random measure on $(0, \infty) \times \mathbb{S}$, such that

$$\mathbb{E}[W(dr, d\theta) \otimes W(dr, d\theta)] = r^{-(\mathcal{H}+1/2)} \sigma(d\theta) r^{-(\mathcal{H}^*+1/2)} dr. \quad (5.10)$$

The proof of this result is deferred to Section A.5, below.

Remark 5.2 (The support of an \mathcal{H} -self-similar IRF_k is \mathcal{H} -invariant). The self-similarity exponent operator \mathcal{H} can in principle be arbitrary outside the support of the IRF_k process Y . The support of Y , denoted $\text{supp}(Y)$, is the smallest closed linear subspace of \mathbb{V} , which contains all $Y(\lambda)$'s almost surely. One can show that $\mathcal{H}(\text{supp}(Y))$ is a dense subset of $\text{supp}(Y)$. This allows one to essentially restrict the operator \mathcal{H} to $\text{supp}(Y)$ (see Section S.6 in Supplement for more details).

Remark 5.3. If Y is \mathcal{H} -self-similar, so are its components $Y_{(\chi,0)}$ and $Y_{(0,\varrho)}$ in (5.4). While this decomposition is unique in law, the operator \mathcal{H} need not be unique. See for example [Didier et al. \[2017\]](#) and Remark 5.7 below. For example, the polynomial component $Y_{(0,\varrho)}$ is always $(k+1) \cdot \text{I}$ -self-similar. In general, however, we cannot conclude that $\mathcal{H} = (k+1) \cdot \text{I}$, where I is the identity. The non-uniqueness of the operator self-similarity exponent in the general setting of \mathbb{V} -valued IRF_k 's is an interesting problem of future research.

For simplicity, in the rest of this section we suppose that $Y \stackrel{d}{=} Y_{(\chi,0)}$ has a *trivial* polynomial component $Y_{(0,\varrho)} = 0$. We will examine two classes of operators \mathcal{H} which can serve as self-similarity exponents of Y . We begin with a simple criterion.

Corollary 5.3. *A finite \mathbb{T}_+ -valued measure χ is the spectral measure of an L^2 -continuous, covariance \mathcal{H} -self-similar IRF_k if and only if for some finite \mathbb{T}_+ -valued measure σ on \mathbb{S} , we have*

$$\chi_k(dx) \equiv \frac{1}{1 \wedge \|x\|^{2k+2}} \chi(dx) = r^{-\mathcal{H}} \sigma(d\theta) r^{-\mathcal{H}^*} r^{-1} dr \quad (5.11)$$

and

$$\int_0^\infty r^{-1} (1 \wedge r^{2k+2}) \text{trace} \left(r^{-\mathcal{H}} \sigma(\mathbb{S}) r^{-\mathcal{H}^*} \right) dr < \infty, \quad (5.12)$$

where $(r, \theta) := (\|x\|, x/\|x\|)$ are the polar coordinates in $\mathbb{R}^d \setminus \{0\}$.

Proof. ‘only if’ Let Y be a continuous covariance \mathcal{H} -self-similar IRF_k . Then, by Theorem 5.2, χ_k satisfies the disintegration formula in (5.11). We have moreover that

$$\chi(\mathbb{R}^d) = \int_0^\infty r^{-1} (1 \wedge r^{2k+2}) r^{-\mathcal{H}} \sigma(\mathbb{S}) r^{-\mathcal{H}^*} dr \in \mathbb{T}_+.$$

Since $\chi(\mathbb{R}^d)$ is self-adjoint and positive definite, $\|\chi(\mathbb{R}^d)\|_{tr} = \text{trace}(\chi(\mathbb{R}^d)) < \infty$, which proves (5.12).

‘if’: Conversely, suppose that σ is a finite \mathbb{T}_+ -valued measure on \mathbb{S} such that (5.12) holds. Then, the fact that for all $B \in \mathcal{B}(\mathbb{S})$ and $r > 0$, $r^{-\mathcal{H}} \sigma(B) r^{-\mathcal{H}^*} \leq r^{-\mathcal{H}} \sigma(\mathbb{S}) r^{-\mathcal{H}^*}$ as positive operators in \mathbb{T}_+ , implies that

$$\chi(D) := \int_0^\infty r^{-1} (1 \wedge r^{2k+2}) r^{-\mathcal{H}} \int_{\mathbb{S}} 1_D(r\theta) \sigma(d\theta) r^{-\mathcal{H}^*} dr, \quad D \in \mathcal{B}(\mathbb{R}^d \setminus \{0\})$$

is well-defined in the sense of Bochner. The so-defined χ is a finite \mathbb{T}_+ -valued Borel measure on $\mathbb{R}^d \setminus \{0\}$, which can be taken as the spectral measure of an IRF_k process $Y = Y_{(\chi,0)}$ with trivial polynomial component. Clearly, χ_k defined as in (5.11) satisfies the scaling property (5.6), which entails the covariance \mathcal{H} -self-similarity of Y . \square

• *Normal diagonalizable exponents.* Corollary 5.3 allows us to provide a complete characterization of the valid pairs (\mathcal{H}, σ) of operator exponents and spectral measures in the important case where \mathcal{H} is *normal* and diagonalizable operator. Namely, suppose \mathcal{H} is a normal operator with

$$\mathcal{H} = \sum_{j=1}^{\infty} \lambda_j e_j \otimes e_j, \quad (5.13)$$

where $\lambda_j \in \mathbb{C}$ and where $\{e_j\}$ is a CONS of $\text{Range}(\mathcal{H}) = \text{Range}(\mathcal{H}^*)$. The convergence of the last series is understood in the weak operator topology.

Theorem 5.4. *Let \mathcal{H} be a normal diagonalizable operator as in (5.13) and let σ be a finite \mathbb{T}_+ -valued measure on \mathbb{S} . The measure $\chi(drd\theta) = (1 \wedge r^{2k+2})r^{-\mathcal{H}}\sigma(d\theta)r^{-\mathcal{H}^*}r^{-1}dr$ is the spectral measure of an \mathcal{H} -self-similar IRF_k , if and only if*

$$0 < \Re(\lambda_j) < k + 1, \quad \text{whenever } \langle \sigma(\mathbb{S})e_j, e_j \rangle > 0 \quad (5.14)$$

and

$$\sum_j \left(\frac{1}{k+1-\Re(\lambda_j)} + \frac{1}{\Re(\lambda_j)} \right) \langle \sigma(\mathbb{S})e_j, e_j \rangle < \infty. \quad (5.15)$$

Proof. First, we prove the ‘if’ part. Let $\sigma \in \mathbb{T}_+$ satisfy (5.14) and define (in polar coordinates) $\chi(drd\theta) = (1 \wedge r^{2k+2})r^{-\mathcal{H}}\sigma(d\theta)r^{-\mathcal{H}^*}r^{-1}dr$. By the simple fact

$$\text{trace} \left(r^{-\mathcal{H}} \sigma(\mathbb{S}) r^{-\mathcal{H}^*} \right) = \sum_j r^{-2\Re(\lambda_j)} \langle \sigma(\mathbb{S})e_j, e_j \rangle,$$

we obtain

$$\begin{aligned} & \int_0^\infty r^{-1} (1 \wedge r^{2k+2}) \text{trace} \left(r^{-\mathcal{H}} \sigma(\mathbb{S}) r^{-\mathcal{H}^*} \right) dr \\ &= \sum_j \langle \sigma(\mathbb{S})e_j, e_j \rangle \int_0^\infty (1 \wedge r^{2k+2}) r^{-2\Re(\lambda_j)-1} dr \\ &= \frac{1}{2} \sum_j \left(\frac{1}{k+1-\Re(\lambda_j)} + \frac{1}{\Re(\lambda_j)} \right) \langle \sigma(\mathbb{S})e_j, e_j \rangle < \infty \end{aligned} \quad (5.16)$$

where the integration is justified by (5.14) and (5.15). By Corollary 5.3, χ is the spectral measure of a covariance \mathcal{H} -self-similar IRF_k .

Conversely, suppose χ is the spectral measure of a covariance \mathcal{H} -self-similar IRF_k . In order for the integral in (5.12) to be finite, the calculations in (5.16) show that both (5.14) and (5.15) must hold. \square

• *General bounded operator exponents.* Suppose now that \mathcal{H} is a *general* bounded operator, which need not be normal nor diagonalizable. In this case, we cannot provide a complete characterization of the covariance \mathcal{H} -self-similar IRF_k 's, but still furnish a general sufficient condition using Riesz functional calculus [see e.g., Ch. VII.4 in [Conway, 2007](#)]. Recall that the spectrum $\text{sp}(\mathcal{H})$ of a bounded operator consists of all $z \in \mathbb{C}$ such that $(\mathcal{H} - z \cdot \text{I})$ has no bounded inverse. The spectrum $\text{sp}(\mathcal{H})$ is always a non-empty compact subset of \mathbb{C} and $\text{sp}(\mathcal{H}^*) = \{\bar{z} : z \in \text{sp}(\mathcal{H})\}$ consists of the complex conjugates of the elements in the spectrum of \mathcal{H} . If Γ is a rectifiable curve containing $\text{sp}(\mathcal{H})$ in its interior then for every holomorphic function f on an open set containing the curve Γ along with its interior, we define

$$f(\mathcal{H}) := \frac{1}{2\pi i} \oint_{\Gamma} \frac{f(z)}{z - \mathcal{H}} dz, \quad (5.17)$$

where the latter integral over Γ is considered in the positive direction and $1/(z - \mathcal{H}) := (z \cdot \text{I} - \mathcal{H})^{-1}$ is a bounded operator since $z \in \Gamma \subset \mathbb{C} \setminus \text{sp}(\mathcal{H})$. Since $f(z) = \exp\{-\log(r)z\}$ is analytic for all $r > 0$, we can use the above Riesz functional calculus tool to study the operator $r^{-\mathcal{H}}$.

Proposition 5.5. *Let \mathcal{H} be a bounded operator and let $\Re(\text{sp}(\mathcal{H}))$ denote the set of real parts of its spectrum. If*

$$\Re(\text{sp}(\mathcal{H})) \subset (0, k+1), \quad (5.18)$$

for some $k \geq 0$, $k \in \mathbb{Z}$, then for all finite \mathbb{T}_+ -valued measures σ on \mathbb{S} , we have that

$$\chi(dr d\theta) := (1 \wedge r^{2k+2}) r^{-\mathcal{H}} \sigma(d\theta) r^{-\mathcal{H}^*} r^{-1} dr,$$

is the spectral measure of a covariance \mathcal{H} -self-similar IRF_k .

Proof. We will show first that, for some $\epsilon > 0$ and $C_{\mathcal{H}} > 0$,

$$\|r^{-\mathcal{H}}\|_{\text{op}} \leq C_{\mathcal{H}} \cdot \left(r^{-(k+1)+\epsilon} 1_{(0,1)}(r) + r^{-\epsilon} 1_{[1,\infty)}(r) \right). \quad (5.19)$$

Recall that $\text{sp}(\mathcal{H})$ is a compact subset of \mathbb{C} . This fact and the assumption (5.18) on the spectrum of \mathcal{H} implies that $\Re(\text{sp}(\mathcal{H})) \subset (\epsilon, k+1-\epsilon)$, for some $\epsilon > 0$. Since $\text{sp}(\mathcal{H})$ is compact, one can take a closed curve Γ containing $\text{sp}(\mathcal{H})$ in its interior, such that $\epsilon \leq \Re(z) \leq k+1-\epsilon$ for all $z \in \Gamma$. Note that $z \mapsto \|(z \cdot \text{I} - \mathcal{H})^{-1}\|_{\text{op}}$ is a continuous function of z for all $z \in \Gamma \subset \mathbb{C} \setminus \text{sp}(\mathcal{H})$. Thus, since Γ is a compact set, we have that

$$\max_{z \in \Gamma} \|(z \cdot \text{I} - \mathcal{H})^{-1}\|_{\text{op}} =: C_{\mathcal{H}}(\Gamma) < \infty.$$

Now, by applying (5.17) to $f(z) := r^{-\mathcal{H}} = e^{-\log(r)\mathcal{H}}$, for all $r > 0$, we obtain

$$\begin{aligned} \|r^{-\mathcal{H}}\|_{\text{op}} &\leq \frac{1}{2\pi} \int_{\Gamma} |e^{-\log(r)z}| \|(z \cdot \text{I} - \mathcal{H})^{-1}\|_{\text{op}} |dz| \\ &\leq \frac{C_{\mathcal{H}}(\Gamma)}{2\pi} \text{Len}(\Gamma) \sup_{z \in \Gamma} |e^{-\log(r)z}|, \end{aligned} \quad (5.20)$$

where $\text{Len}(\Gamma) = \int_{\Gamma} |dz|$ is the length of Γ .

Observe now that $|e^{-\log(r)z}| = r^{-\Re(z)}$ and recall that $\epsilon \leq \Re(z) \leq k+1-\epsilon$, for all $z \in \Gamma$. This, implies that

$$|e^{-\log(r)z}| \leq r^{-(k+1)+\epsilon} 1_{(0,1)}(r) + r^{-\epsilon} 1_{[1,\infty)}(r),$$

which in view of (5.20) yields (5.19).

Now, by Corollary 5.3, the measure χ in (5.11) is the spectral measure of a covariance \mathcal{H} -self-similar IRF $_k$, provided (5.12) holds. This, however, readily follows from (5.19). Indeed, by (A.19), the integral in (5.12) is bounded above by

$$\|\sigma(\mathbb{S})\|_{\text{tr}} \int_0^1 r^{2k+2} \|r^{-\mathcal{H}}\|_{\text{op}}^2 r^{-1} dr + \|\sigma(\mathbb{S})\|_{\text{tr}} \int_1^\infty \|r^{-\mathcal{H}}\|_{\text{op}}^2 r^{-1} dr, \quad (5.21)$$

where we used the fact that $\|r^{-\mathcal{H}}\|_{\text{op}} = \|r^{-\mathcal{H}^*}\|_{\text{op}}$. By (5.19), the integrals in (5.21) are finite and the proof is complete. \square

Remark 5.4. The sufficient condition in (5.18) may appear restrictive. In particular, it implies that \mathcal{H} has a bounded inverse (since $0 \notin \text{sp}(\mathcal{H})$). This condition is not all that restrictive when the Hilbert space \mathbb{V} is finite-dimensional and our sufficient conditions are precisely the same as the existing literature in the special case $k = 0$ [see e.g., [Didier and Pipiras, 2011](#), [Didier et al., 2018](#)].

5.2. Related work and examples.

Here, we first specialize the results from the previous section and discuss existing related work when \mathbb{V} is finite-dimensional. Then, we consider a class of stationary infinite-dimensional processes, which admit higher-order tangent fields under operator scaling.

Example 5.1 (IRF $_0$ or operator fractional Brownian motions). When $k = 0, d = 1$, and $\mathbb{V} = \mathbb{R}^m$, the IRF $_k$ processes can be identified with the well-studied class of vector-valued stationary increment processes. The seminal paper of the [Didier and Pipiras \[2011\]](#) established the spectral representation and stochastic integral representations for essentially all Gaussian operator self-similar processes with stationary increments taking values in \mathbb{R}^m . We demonstrate next how these processes, known as operator fractional Brownian motions (OFBM), can be recovered from our Theorem 5.2. In this setting the operator \mathcal{H} is a real $m \times m$ matrix with eigenvalues $\lambda_i \in \mathbb{C}$, $i = 1, \dots, m$ such that

$$0 < \Re(\lambda_i) < 1, \quad (5.22)$$

[see e.g., (1.4) in [Didier and Pipiras, 2011](#)]. Observe that the last condition coincides with (5.18) of Proposition 5.5 for $k = 0$.

Let $\{Y(\lambda)\}$ be a zero-mean Gaussian \mathcal{H} -self-similar IRF $_0$. Then, if one considers

$$\lambda_t(du) := \delta_t(du) - \delta_0(du), \quad t \in \mathbb{R},$$

the process $B(t) := Y(\lambda_t)$, $t \in \mathbb{R}$ has stationary increments. The \mathcal{H} -self-similarity of $\{B(t)\}$ follows readily from the self-similarity of $\{Y(\lambda)\}$ and the fact that $c \cdot \lambda_t = \lambda_{ct}$, $c > 0$, $t \in \mathbb{R}$. Conversely, every \mathcal{H} -self-similar stationary increment process $\{B(t), t \in \mathbb{R}\}$ can be taken as a representer of an \mathcal{H} -self-similar IRF $_0$ process.

Since $\hat{\lambda}_t(x) = e^{itx} - 1$ and $\mathbb{S} = \{-1, 1\}$, Relation (5.9) yields

$$\begin{aligned} B(t) \equiv Y(\lambda_t) &= \int_0^\infty \int_{\{-1, 1\}} (e^{itr\theta} - 1) W(dr, d\theta) \\ &= \int_0^\infty (e^{ixt} - 1) W(dx, \{1\}) + \int_0^\infty (e^{-ixt} - 1) W(dx, \{-1\}). \end{aligned} \quad (5.23)$$

Now, following the notation in Theorem 3.1 of [Didier and Pipiras \[2011\]](#), let $\tilde{B}(dx) = \tilde{B}_1(dx) + i\tilde{B}_2(dx)$, where \tilde{B}_i , $i = 1, 2$ are independent zero-mean Gaussian \mathbb{R}^m -valued measures such that $\tilde{B}_1(dx) = \tilde{B}_1(-dx)$, $\tilde{B}_2(dx) = -\tilde{B}_2(-dx)$, and

$$\mathbb{E}[\tilde{B}(dx)\tilde{B}(dx)^*] \equiv \mathbb{E}[\tilde{B}(dx)\overline{\tilde{B}(dx)}^\top] = I_m dx. \quad (5.24)$$

Observe that, by (5.10),

$$\left\{ W(dx, \{\pm 1\}), dx \in (0, \infty) \right\} \stackrel{d}{=} \left\{ x^{-(\mathcal{H}+1/2)} A_{\pm 1} \tilde{B}(\pm dx), dx \in (0, \infty) \right\},$$

where $A_{\pm 1} A_{\pm 1}^* = \sigma(\{\pm 1\})$. Therefore, Relation (5.23) yields

$$\{B(t)\} \stackrel{d}{=} \left\{ \int_{-\infty}^{\infty} \frac{(e^{ixt} - 1)}{ix} \left(x_+^{-(\mathcal{H}-1/2)} A_1 + x_-^{-(\mathcal{H}-1/2)} A_{-1} \right) \tilde{B}(dx) \right\}.$$

This is precisely the representation established in Theorem 3.1 of [Didier and Pipiras \[2011\]](#), wherein $A_{-1} = \overline{A_1}$ is the complex conjugate of A_1 since they consider real-valued processes. Indeed, the last stochastic integral is real-valued if and only if the integrand $f_t(x)$ is a Hermitian function of x , i.e., $f_t(-x) = \overline{f_t(x)}$. This is the case, if and only if $A_1 = \overline{A_{-1}}$.

Remark 5.5. Note that the condition (5.22) on the eigenvalues of the matrix \mathcal{H} does not imply in general that $T_c := c^{\mathcal{H}}$, $c > 0$ are scaling actions in the Euclidean norm of $\mathbb{V} \equiv \mathbb{R}^m$. This is because the monotonicity of the function $c \mapsto \|c^{\mathcal{H}}\|$ may be violated except when the matrix \mathcal{H} is normal (i.e., diagonalizable in an orthonormal basis). In particular, Lemma A.1 is not applicable. Nevertheless, as shown in [Lemma 6.1.5 in [Meerschaert and Scheffler, 2001](#)], there is a suitable norm in \mathbb{V} , with respect to which the latter are monotone increasing and in this new (equivalent norm) $\{c^{\mathcal{H}}, c > 0\}$ is a scaling action in the sense of Definition 2.5. See also [Jurek \[1984\]](#) for the case where \mathbb{V} is a Banach space.

Remark 5.6. By Proposition 5.5 (with $k = 0$), Condition (5.22) implies that the stochastic integrals in (5.23) are well-defined.

Example 5.2 (Operator fractional Brownian fields). Stationary increment vector-valued random fields (IRF $_k$ with $k = 0$) where $d \geq 2$ have been actively studied by many authors [see e.g., [Biermé et al., 2007](#), [Li and Xiao, 2011](#), [Baek et al., 2014](#), [Didier et al., 2018](#), among others.] In the latter references, self-similarity is considered under operator rescaling of *both* the range and the domain of the process. Here, we consider only scalar rescaling of the domain. In this setting, in the special case of processes taking values in \mathbb{R}^m and $k = 0$, Theorem 5.2 recovers Proposition 3.1 of [Didier et al. \[2018\]](#).

Interestingly, using Fréchet differentiability in Theorem 5.2, we extend the *disintegration formula* in Relation (3.10) of [Didier et al. \[2018\]](#) to the case of processes taking values in a separable Hilbert space as well as to the general case of intrinsic random functions of order k . We anticipate that a version of our Theorem 5.2 holds under operator scaling of both the range and the domain of Y .

Remark 5.7 (The non-uniqueness of the operator exponent \mathcal{H}). Suppose that Y is an operator \mathcal{H} -self-similar zero-mean Gaussian continuous IRF $_k$ taking values in the *real* Hilbert space $\mathbb{V}_{\mathbb{R}}$ (recall Section 4.3). Then, the distribution of Y is determined by its covariance structure, i.e., by the unique pair of its spectral characteristics (χ, \mathcal{Q}) or equivalently (σ, \mathcal{Q}) . The operator exponent \mathcal{H} , however, is not necessarily unique even when \mathcal{H} is restricted to the support of the process Y . For the notion of a support of Y and its relation to the operator exponent \mathcal{H} , see Section S.6 in

Supplement. To gain some intuition, suppose that for an operator \mathcal{A} on $\mathbb{V} := \text{supp}(Y)$, we have that $\{c^{\mathcal{A}}Y(\lambda)\} \stackrel{d}{=} \{Y(\lambda)\}$, for all $c > 0$. If \mathcal{H} and \mathcal{A} commute, then $c^{\mathcal{H}+\mathcal{A}} = c^{\mathcal{H}}c^{\mathcal{A}}$, $c > 0$ and hence Y is also $(\mathcal{H} + \mathcal{A})$ -self-similar.

As shown in [Didier et al. \[2017\]](#) such non-uniqueness can arise even in the finite-dimensional case with $k = 0$, where a wealth of interesting phenomena emerge. Specifically, Theorem 2.4 therein characterizes all possible operator exponents and shows that one can always choose a *commuting exponent* \mathcal{H}_0 such that $\mathcal{H}_0\mathcal{A} = \mathcal{A}\mathcal{H}_0$. In their terminology, the operator \mathcal{A} belongs to the tangent space of the group of symmetries of the process. Notice that \mathcal{A} can indeed be viewed as a tangent since it is the Fréchet derivative of $f(c) = c^{\mathcal{A}}$ at $c = 1$.

Understanding the non-uniqueness of the operator self-similarity exponent in the general infinite-dimensional case is a challenging problem. We anticipate that the extension of the important characterization results of [Didier et al. \[2017\]](#) to the infinite-dimensional case is possible but considerably beyond the scope of this paper.

We end this section with an example of stationary Gaussian \mathbb{V} -valued processes, which admit a large class of tangent fields.

Example 5.3 (Higher-order tangent fields in infinite dimensions). In this example, we shall assume that $\mathbb{V}_{\mathbb{R}}$ is a real Hilbert space and through the method of complexification define $\mathbb{V} = \mathbb{V}_{\mathbb{R}} + i\mathbb{V}_{\mathbb{R}}$, with the natural inclusion $\mathbb{V}_{\mathbb{R}} \subset \mathbb{V}$.

Consider polar coordinates in $\mathbb{R}^d \setminus \{0\}$, where $u = \|x\|$, $\theta := x/\|x\|$, are the radial and angular components of $x \in \mathbb{R}^d \setminus \{0\}$ and let $\mu(d\theta)$ be a finite, real, \mathbb{T}_+ -valued measure on the unit sphere $\mathbb{S} = \{\theta \in \mathbb{R}^d : \|\theta\| = 1\}$. Define the real, σ -finite \mathbb{T}_+ -valued measure

$$\nu(dx) = \nu(du, d\theta) = du\mu(d\theta), \quad (u, \theta) \in (0, \infty) \times \mathbb{S}.$$

Let $W_{\mathbb{R}} = \{W_{\mathbb{R}}(A), A \in \mathcal{B}(\mathbb{R}^d \setminus \{0\})\}$ and $W_{\mathbb{I}} = \{W_{\mathbb{I}}(A), A \in \mathcal{B}(\mathbb{R}^d \setminus \{0\})\}$ be two independent, real (i.e., $\mathbb{V}_{\mathbb{R}}$ -valued) orthogonal Gaussian measures with the same control measure $2^{-1}\nu$ (in the sense of Definition 4.6). Construct

$$W(A) = W_{\mathbb{R}}(A) + iW_{\mathbb{I}}(A). \quad (5.25)$$

It is straightforward to see that $W = \{W(A)\}$ is an orthogonal Gaussian \mathbb{V} -valued random measure with control measure ν , i.e., for all bounded Borel $A, B \in \mathcal{B}(\mathbb{R}^d \setminus \{0\})$, the random vectors $W(A)$ and $W(B)$ are such that

$$\mathbb{E}[W(A) \otimes W(B)] = \nu(A \cap B) = \int_0^\infty \int_{\mathbb{S}} 1_{A \cap B}(u\theta) du \mu(d\theta). \quad (5.26)$$

Note, moreover, that W is also independently scattered, i.e., $W(A_i)$, $i = 1, \dots, n$ are independent for all disjoint bounded Borel sets $A_i \subset \mathbb{R}^d \setminus \{0\}$, $i = 1, \dots, n$, which is not necessarily true for all orthogonal Gaussian random measures taking values in a complex Hilbert space.

By analogy with the scalar \mathbb{C} -valued case the Gaussian random measure W in (5.25) will be referred to as standard. Since its real and imaginary components are iid, the distribution of the process $W = \{W(A)\}$ is completely determined by its cross-covariance operators in (5.26). Moreover, W has circular symmetry and self-similarity properties:

$$\{e^{i\varphi}W(du, d\theta)\} \stackrel{fdd}{=} \{W(du, d\theta)\} \quad \text{and} \quad \{r^{1/2}W(du, d\theta)\} \stackrel{fdd}{=} \{W(d(r \cdot u), d\theta)\}, \quad (5.27)$$

for all $\varphi \in \mathbb{R}$ and $r > 0$.

Let \mathcal{H} be a bounded linear operator on \mathbb{V} such that

$$\Re(\text{sp}(\mathcal{H})) \subset (\epsilon, k+1), \quad (5.28)$$

for some $\epsilon > 0$ and $k \geq 0$, $k \in \mathbb{Z}$. Suppose also that $\mathcal{A}(\theta)$, $\theta \in \mathbb{S}$ is a collection of bounded linear operators such that $\theta \mapsto \mathcal{A}(\theta)$ is Borel measurable in θ and such that

$$\int_{\mathbb{S}} \|\mathcal{A}(\theta)\|_{\text{op}}^2 \|\mu\|_{\text{tr}}(d\theta) < \infty, \quad (5.29)$$

where $\|\mu\|_{\text{tr}}$ denoted the (finite) trace measure $\|\mu\|_{\text{tr}}(A) := \|\mu(A)\|_{\text{tr}}$.

Proposition 5.6. *Suppose that (5.28) and (5.29) hold, for some $k \geq 0$, $k \in \mathbb{Z}$.*

(i) *For all $s \in \mathbb{R}^d$, the stochastic integral*

$$X(s) := \int_0^\infty \int_{\mathbb{S}} f_s(u, \theta) W(du, d\theta), \quad \text{where } f_s(u, \theta) := e^{i u s^\top \theta} (1 \wedge u)^{k+1} u^{-(\mathcal{H}+1/2)} \mathcal{A}(\theta) \quad (5.30)$$

exists and defines a stationary \mathbb{V} -valued Gaussian random field.

(ii) *The process $X = \{X(s), s \in \mathbb{R}^d\}$ has a version with γ -Hölder continuous paths for all $\gamma \in (0, 1 \wedge \epsilon)$, where ϵ is as in (5.28).*

(iii) *The continuous-path version of the process $\{X(s)\}$ has a k -th order tangent field at each (any) s_0 . More precisely,*

$$\left\{ r^{-\mathcal{H}} X(s_0 + r \cdot \lambda), \lambda \in \Lambda_k \right\} \xrightarrow{d} Y = \{Y(\lambda), \lambda \in \Lambda_k\}, \quad \text{as } r \downarrow 0, \quad (5.31)$$

where the tangent process is an \mathcal{H} -self-similar IRF $_k$ given by

$$Y(\lambda) = \int_{\mathbb{R}^d} \widehat{\lambda}(u\theta) u^{-\mathcal{H}-1/2} \mathcal{A}(\theta) W(du, d\theta).$$

The proof of this result is given in Section A.6, below.

Remark 5.8. Notice that when \mathbb{V} is infinite-dimensional in (5.30) one *cannot* consider Gaussian measures W with the control measure equal to the Lebesgue measure times the identity operator I_m as in (5.24). Indeed, for $W(A)$ to be a bona fide random element in \mathbb{V} the control measure of W must take values in \mathbb{T}_+ . This is the key reason why we consider control measures of the type $du\mu(d\theta)$. In the *finite-dimensional case*, one can obtain more familiar, but ultimately equivalent stochastic integral representations, in terms of Gaussian \mathbb{C}^m -valued Gaussian random measures with the Lebesgue control measure times the identity by considering $\mathbb{E}[W_{\text{Leb}}(dr, d\theta) \otimes W_{\text{Leb}}(dr, d\theta)] = v_d r^{d-1} dr \times I_m$, where $v_d := \pi^{d/2}/\Gamma(1+d/2)$ is the volume of the unit sphere in \mathbb{R}^d . In this case, the stochastic integral in (5.30) can be equivalently written in Cartesian coordinates as follows:

$$X(s) := v_d^{-1/2} \int_{\mathbb{R}^d} e^{i s^\top x} (1 \wedge \|x\|)^{k+1} \|x\|^{-(\mathcal{H}+d/2)} \mathcal{A}(x/\|x\|) W_{\text{Leb}}(dx), \quad s \in \mathbb{R}^d.$$

5.3. Scalar actions.

In this section, we characterize the spectral measure of covariance self-similar IRF_k 's with respect to the usual scalar scaling action. In this special but important case we obtain a more complete picture of the H -self-similar IRF_k 's, where now H is a scalar exponent.

Proposition 5.7. *Let Y be a non-constant continuous IRF_k , $k \geq 0$ with operator auto-covariance function \mathcal{K} and spectral characteristics (χ, \mathcal{Q}) . If Y is covariance self-similar with exponent H , then $H \in (0, k+1]$ and we have the following dichotomy:*

- (i) *If $H = k+1$, then $\chi \equiv 0$ and if $0 < H < k+1$, then \mathcal{Q} is trivial, i.e., $\mathcal{Q}(\nu) = 0$, for all $\nu \in \Lambda_{2k+1}$.*
- (ii) *In the case $0 < H < k+1$, the measure χ_k in (5.5) satisfies the scaling property in (5.6) and consequently, the disintegration formula in (5.8) reads:*

$$\chi_k(dr d\theta) = r^{-2H-1} dr \sigma(d\theta), \quad (r, \theta) \in (0, \infty) \times \mathbb{S}, \quad (5.32)$$

for some finite \mathbb{T}_+ -valued measure σ on \mathbb{S} .

- (iii) *Conversely, for every $0 < H < k+1$ and any finite \mathbb{T}_+ -valued measure σ on \mathbb{S} , there exists a covariance H -self-similar IRF_k with spectral measure χ such that (5.32) holds, which can be written as in (5.9).*

The proof is given in Section A.7, below.

In view of (4.13) and (5.32), one can obtain explicit formulae for the generalized covariance \mathcal{K} of all covariance H -self-similar IRF_k 's. This is done next.

Theorem 5.8. *Let $\mathcal{K}(\cdot)$ be the generalized covariance of a covariance self-similar IRF_k , $k \geq 0$ with exponent $H \in (0, k+1)$. Then, with σ as in (5.32), we have:*

- (i) *If $2H \notin \{1, \dots, k\}$ is non-integer, then for all $\nu \in \Lambda_{2k+1}$,*

$$\mathcal{K}(\nu) = I(H) \int_{\mathbb{S}^{d-1}} |(\theta, \cdot)|^{2H}(\nu) \sigma(d\theta) + \mathfrak{i} J(H) \int_{\mathbb{S}^{d-1}} (\theta, \cdot)^{<2H>}(\nu) \sigma(d\theta), \quad (5.33)$$

where $(\theta, t) = \theta^\top t$ denotes the Euclidean inner product, $x^{<H>} := \text{sign}(x)|x|^H$, and $f(\cdot)(\nu) := \int f(t) \nu(dt)$. Here the real functions $I(H)$ and $J(H)$ are such that

$$I(H) + \mathfrak{i} J(H) := \int_0^\infty \left(e^{\mathfrak{i}r} - \sum_{j=0}^{\lfloor 2H \rfloor} \frac{(\mathfrak{i}r)^j}{j!} \right) \frac{dr}{r^{2H+1}}. \quad (5.34)$$

- (ii) *If $2H \in \{1, \dots, k\}$ is integer, then*

$$\mathcal{K}(\nu) = \int_{\mathbb{S}^{d-1}} \left[|(\theta, \cdot)|^{2H} \left(\frac{(-1)^{H+1}}{(2H)!} \log |(\theta, \cdot)| + \mathfrak{i} J(H) \text{sign}(\theta, \cdot) \right) \right] (\nu) \sigma(d\theta), \quad \text{if } 2H \text{ is even,}$$

and

$$\mathcal{K}(\nu) = \int_{\mathbb{S}^{d-1}} \left[|(\theta, \cdot)|^{2H} \left(I(H) + \mathfrak{i} \frac{(-1)^{H+1/2}}{(2H)!} \text{sign}(\theta, \cdot) \log |(\theta, \cdot)| \right) \right] (\nu) \sigma(d\theta), \quad \text{if } 2H \text{ is odd.}$$

The proof of this result is given in Section A.7, below.

Remark 5.9. Gel'fand and Vilenkin [1964] provide spectral theory for generalized random fields taking values in the dual of the Schwartz space on \mathbb{R}^d with homogeneous $(1+k)$ th-order increments (denoted as G-IRF $_k$ here). Dobrushin [1979] then studied the self-similar G-IRF $_k$ and obtained results similar to Proposition 5.7 where the self-similarity parameter H can take any value in $(-\infty, k+1]$. The G-IRF $_k$ class of processes is more general than the IRF $_k$'s studied by Matheron [1973] and they do not always have a representation on \mathbb{R}^d . Specifically, it can be shown that a Gaussian self-similar G-IRF $_k$ has a representation as in (4.6) on \mathbb{R}^d only if $H > 0$ [see Shen, 2019].

Remark 5.10 (On symmetry and covariance (ir)reversibility). Observe that for σ in (5.33), one can write $\sigma = \sigma_s + \sigma_a$, where $\sigma_s(A) := (\sigma(A) + \sigma(-A))/2$ and $\sigma_a(A) := (\sigma(A) - \sigma(-A))/2$, are the symmetric and anti-symmetric components of σ . Thus,

$$\int_{\mathbb{S}^{d-1}} |(\theta, \cdot)|^{2H}(\nu) \sigma_a(dt) = \int_{\mathbb{S}^{d-1}} (\theta, \cdot)^{<2H>}(\nu) \sigma_s(dt) = 0,$$

and (5.33) can be equivalently written as:

$$\mathcal{K}(\nu) = I(H) \int_{\mathbb{S}^{d-1}} |(\theta, \cdot)|^{2H}(\nu) \sigma_s(d\theta) + \mathfrak{i}J(H) \int_{\mathbb{S}^{d-1}} (\theta, \cdot)^{<2H>}(\nu) \sigma_a(d\theta). \quad (5.35)$$

This shows that unless $\sigma_a \equiv 0$, we have that $\mathcal{K}(\nu) \neq \mathcal{K}(-\nu)$, for some $\nu \in \Lambda_{2k+1}$. Recall that by $-\nu$ we understand $((-1) \cdot \nu)(dx) := \nu(-dx)$.

Recall Definition 4.17; Y is covariance reversible, i.e., $\{Y(-\lambda)\}$ and $\{Y(\lambda)\}$ have the same covariance structure, if and only if $\mathcal{K}(\nu) = \mathcal{K}(-\nu)$, $\forall \nu \in \Lambda_{2k+1}$ or equivalently if and only if $\sigma_a \equiv 0$ [see also Proposition 4.18 above as well as Theorem 5.1 in Didier and Pipiras, 2011, for a related result].

Remark 5.11 (Real H -self-similar IRF $_k$'s). Recall that $\mathbb{V} = \mathbb{V}_{\mathbb{R}} + \mathfrak{i}\mathbb{V}_{\mathbb{R}}$ (cf Section 4.3). Thus for an H -self-similar IRF $_k$ Y , we have

$$Y(\lambda) = \Re Y(\lambda) + \mathfrak{i}\Im Y(\lambda),$$

where the real and imaginary parts $\Re Y$ and $\Im Y$ are real, i.e., $\mathbb{V}_{\mathbb{R}}$ -valued. Thus, in view of (5.35), one can see that Y is real-valued (i.e., $\Im Y \equiv 0$) if and only if σ_s is real and σ_a imaginary, i.e., if σ is Hermitian, $\sigma(A) = \overline{\sigma(-A)}$ for all $A \in \mathcal{B}(\mathbb{S})$. Observe that Y need not be covariance-reversible for it to be real (see Section 4.3.)

Since the covariance structure characterizes completely the zero-mean Gaussian processes taking values in *real* Hilbert spaces, Theorem 5.8 with Hermitian σ provides a complete characterization of all H -self-similar $\mathbb{V}_{\mathbb{R}}$ -valued IRF $_k$'s.

Remark 5.12 (n -th order fractional Brownian motion). Perrin et al. [2001] have studied the so-called n -th order fractional Brownian motion defined (in Remark 2 therein) as

$$B_H^{(n)}(t) := \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{1}{(\mathfrak{i}\omega)^{H+1/2}} \left(e^{\mathfrak{i}t\omega} - \sum_{\ell=0}^{n-1} \frac{(\mathfrak{i}t\omega)^\ell}{\ell!} \right) W(d\omega),$$

with $n-1 < H < n$, $n \in \mathbb{N}$, where $W(d\omega)$ is a zero mean complex Gaussian measure on \mathbb{R} with the Lebesgue control measure and such that $W(-d\omega) = \overline{W(d\omega)}$. Notice, however, that the above integral representation is well defined only if $H \in (n-1, n)$. While one can always put $n := \lceil H \rceil$,

the integer values of H have to be dealt with separately. Using our abstract approach, we can handle *all* values of $H \in (0, k+1)$, $k := n-1$, in a unified manner.

Indeed, observe that with $k = n-1$, for any $\lambda \in \Lambda_k$, we have

$$\begin{aligned} B_H^{(n)}(\lambda) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \widehat{\lambda}(\omega)(i\omega)^{-(H+1/2)} W(d\omega) \\ &= \frac{1}{2\pi} \int_0^{\infty} \int_{\mathbb{S}} \widehat{\lambda}(u\theta) u^{-(H+1/2)} e^{-i\theta \frac{\pi}{2}(H+1/2)} W(du, d\theta) \\ &=: \frac{1}{2\pi} \int_0^{\infty} \int_{\mathbb{S}} \widehat{\lambda}(u\theta) u^{-(H+1/2)} \widetilde{W}(du, d\theta), \end{aligned} \quad (5.36)$$

where we used the change to polar coordinates $(u, \theta) := (|\omega|, \text{sign}(\omega))$, with $\mathbb{S} = \{\pm 1\}$ and the fact that λ annihilates all polynomials of degree up to k . This latter integral is defined for all $0 < H < k+1$. Notice that $\widetilde{W}(du, d\theta) := e^{-i\theta \frac{\pi}{2}(H+1/2)} W(du, d\theta)$ is equal in law to $W(du, d\theta)$ and (5.36) is a particular case of the stochastic representation of the self-similar IRF $_k$'s characterized in Proposition 5.7.

We end with an example outlining the general form of the \mathbb{R} -valued H -self-similar Gaussian IRF $_k$'s in \mathbb{R}^d , which may be viewed as generalized fractional Brownian fields with $(1+k)$ -th order stationary increments.

Example 5.4 (Real n -th order fractional Brownian fields). Fix an integer $k := n-1 \geq 0$, and let λ_t be as in (2.7). Let also $\sigma(d\theta)$ be a finite symmetric measure on the unit sphere $\mathbb{S} \subset \mathbb{R}^d$. For all $H \in (0, k+1)$, it can be shown that $f_t(x, \theta) := (\langle \cdot, \theta \rangle - x)_+^{H-1/2}(\lambda_t)$ belongs to $\mathcal{L}^2(dx, \sigma(d\theta))$, where dx is the Lebesgue measure on \mathbb{R} . Thus, one can define the \mathbb{R} -valued Gaussian random field

$$Y(t) := \int_{-\infty}^{\infty} \int_{\mathbb{S}} \left(\langle \cdot, \theta \rangle - x \right)_+^{H-1/2}(\lambda_t) W(dx, d\theta), \quad t \in \mathbb{R}^d,$$

where $W(dx, d\theta)$ is a zero-mean Gaussian real-valued random measure on $\mathbb{R} \times \mathbb{S}$ with control measure $dx\sigma(d\theta)$. Then, it is easy to see with a simple change of variables that $Y(\lambda) := \int Y d\lambda$, $\lambda \in \Lambda_k$ is an H -self-similar IRF $_k$ with *real* and symmetric generalized covariance K . In this case Theorem 5.8 yields:

$$K(\nu) = C_H \int_{\mathbb{S}} \left[|\langle \cdot, \theta \rangle|^{2H} \left(I(H) + \frac{(-1)^{H+1}}{(2H)!} 1_{\mathbb{N}}(H) \log |\langle \cdot, \theta \rangle| \right) \right] (\nu) \sigma(d\theta).$$

For more examples and further insights, see the PhD thesis of Shen [2019].

Appendix A: Proofs and auxiliary results.

A.1. Proofs and tools for Section 2.

Proof of Lemma 2.2: Notice that $S_c(\check{\Lambda}_k, \mathbb{V})$ is a closed set in the metric space $(C(\mathbb{R}^d, \mathbb{V}), \rho)$ of continuous \mathbb{V} -valued functions on \mathbb{R}^d , equipped with the metric ρ in (2.13). Thus, it is enough to show that $(C(\mathbb{R}^d, \mathbb{V}), \rho)$ is a complete separable metric space [cf. Theorems 1 and 2 in Chapter XIV.2 in Kuratowski, 1977].

It is known that the metric ρ generates the compact-open topology [see, e.g., Theorem 46.8 in Munkres, 2000]. Therefore, to prove separability it is enough to demonstrate that this topology has a countable base. Recall that the compact-open topology on $C(\mathbb{R}^d, \mathbb{V})$ has a sub-base comprising

all sets $V(K, U) = \{f \in C(\mathbb{R}^d, \mathbb{V}) : f(K) \subset U\}$, where $K \subset \mathbb{R}^d$ is compact and $U \subset \mathbb{V}$ is open. Since $(\mathbb{V}, d_{\mathbb{V}})$ is separable and \mathbb{R}^d is locally compact, the compact-open topology on $C(\mathbb{R}^d, \mathbb{V})$ is second countable [cf. Theorem 5.2, page 265 in [Dugundji, 1966](#)]. This entails the separability of $(C(\mathbb{R}^d, \mathbb{V}), \rho)$.

Completeness is established in a standard manner. Let $\{f_n\}$ be a Cauchy sequence in $(C(\mathbb{R}^d, \mathbb{V}), \rho)$. In view of (2.13), for each $t \in \mathbb{R}^d$, $\{f_n(t)\}$ is a Cauchy sequence in the complete metric space $(\mathbb{V}, d_{\mathbb{V}})$. Thus, $f_n(t) \rightarrow f(t) \in \mathbb{V}$. It remains to show that f is continuous and $\rho(f_n, f) \rightarrow 0$. Fix an arbitrary compact $K \subset \mathbb{R}^d$ and $\epsilon > 0$. Since $\{f_n\}$ is Cauchy in ρ , there exists an N_{ϵ} such that

$$\sup_{\tau \in K} d_{\mathbb{V}}(f_n(\tau), f_m(\tau)) \leq \epsilon/3, \quad \text{for all } m, n \geq N_{\epsilon}.$$

On the other hand, for every fixed $t \in K$, we have

$$d_{\mathbb{V}}(f_n(t), f(t)) = \lim_{m \rightarrow \infty} d_{\mathbb{V}}(f_n(t), f_m(t)) \leq \epsilon/3, \quad n \geq N_{\epsilon}.$$

Since the latter bound is uniform in t , we also obtain $\sup_{t \in K} d_{\mathbb{V}}(f_n(t), f(t)) \leq \epsilon/3$, $n \geq N_{\epsilon}$. That is, f_n converge to f uniformly on all compact K . It remains to establish that f is continuous. For all $s, t \in K$, we have by the triangle inequality that

$$d_{\mathbb{V}}(f(t), f(s)) \leq 2 \sup_{\tau \in K} d_{\mathbb{V}}(f_n(\tau), f(\tau)) + d_{\mathbb{V}}(f_n(t), f_n(s)) \leq \epsilon,$$

provided that $\|t - s\| < \delta$ for some sufficiently small $\delta > 0$. Here, we used the uniform continuity of f_n on K . Since $\epsilon > 0$ was arbitrary, this completes the proof of the (uniform) continuity of f on K . \square

Proof of Lemma 2.6: Properties (i) and (ii) in Definition 2.5 are immediate. We now verify (iii). Consider the coordinate-wise action on $\check{S}_c(\Lambda_k, \mathbb{V})$, also denoted as $\{\tilde{T}_a, a \in \mathbb{R}_+\}$ for convenience. One can easily verify that $\mathcal{J}(\tilde{T}_a(f)) = \tilde{T}_a(\mathcal{J}(f))$. Let $f_n \rightarrow f$ in $S_c(\Lambda_k, \mathbb{V})$ and $a_n \rightarrow a > 0$. To show that $\rho(\tilde{T}_{a_n}(f_n), \tilde{T}_a(f)) \rightarrow 0$, it is enough to verify that for every $K > 0$, we have

$$\sup_{\|t\| \leq K} d_{\mathbb{V}}(\tilde{T}_{a_n}(\check{f}_n)(t), \tilde{T}_a(\check{f})(t)) \rightarrow 0, \quad \text{as } n \rightarrow \infty,$$

where $\check{f}_n = \mathcal{J}(f_n)$ and $\check{f} = \mathcal{J}(f)$.

In view of Lemma A.2, below, it is enough to show that $\tilde{T}_{a_n}(\check{f}_n)(t_n) \rightarrow \tilde{T}_a(\check{f})(t)$, whenever $t_n \rightarrow t$ in $\overline{B}_K := \{t : \|t\| \leq K, t \in \mathbb{R}^d\}$. Notice, however, that $\tilde{T}_{a_n}(\check{f}_n)(t_n) = T_{a_n}(y_n)$ and $\tilde{T}_a(\check{f})(t) = T_a(y)$, where $y_n := \check{f}_n(t_n)$ and $y := \check{f}(t)$. By applying Lemma A.2 again, but now to the locally converging functions \check{f}_n and \check{f} , we have that $y_n = \check{f}_n(t_n) \rightarrow y = \check{f}(t)$ in \mathbb{V} , whenever $t_n \rightarrow t$ in \overline{B}_K . Hence, the continuity of the scaling action $\{T_a\}$, yields $T_{a_n}(y_n) \rightarrow T_a(y)$ in \mathbb{V} , which completes the proof of property (iii).

Let now $0 \neq f \in S_c(\Lambda_k, \mathbb{V})$. Proving property (iv) of Definition 2.5 amounts to showing that $\rho(\tilde{T}_{a_1}(f), 0) < \rho(\tilde{T}_{a_2}(f), 0)$, for all $0 < a_1 < a_2$. Observe that by property (iv) for $\{T_a\}$, we have

$$d_{\mathbb{V}}(\tilde{T}_{a_1}(\check{f})(t), 0) \equiv d_{\mathbb{V}}(T_{a_1}(\check{f}(t)), 0) \leq d_{\mathbb{V}}(T_{a_2}(\check{f}(t)), 0) \equiv d_{\mathbb{V}}(\tilde{T}_{a_2}(\check{f})(t), 0).$$

This implies that $\rho(\tilde{T}_{a_1}(f), 0) \leq \rho(\tilde{T}_{a_2}(f), 0)$. We next argue that the inequality is strict. Since $f \neq 0$, we have $0 \neq \check{f}(t) \in \mathbb{V}$ for some $t \in \mathbb{R}^d$. Let $t \in \overline{B}_j$ for some large enough j , where B_j is as

defined in (2.15). Since the suprema therein are attained, it is enough to show that

$$\begin{aligned} d_{\mathbb{V}}(T_{a_1}(\check{f}(t_1)), 0) &:= \max_{t \in \bar{B}_j} d_{\mathbb{V}}(T_{a_1}(\check{f}(t)), 0) \\ &< \max_{t \in \bar{B}_j} d_{\mathbb{V}}(T_{a_2}(\check{f}(t)), 0) =: d_{\mathbb{V}}(T_{a_2}(\check{f}(t_2)), 0). \end{aligned} \quad (\text{A.1})$$

Observe that, $0 < d_{\mathbb{V}}(T_{a_1}(\check{f}(t)), 0) \leq d_{\mathbb{V}}(T_{a_1}(\check{f}(t_1)), 0)$ and hence $\check{f}(t_1) \neq 0$. Thus, by the radial monotonicity of the action $\{T_a\}$, we have

$$d_{\mathbb{V}}(T_{a_1}(\check{f}(t_1)), 0) < d_{\mathbb{V}}(T_{a_2}(\check{f}(t_1)), 0) \leq d_{\mathbb{V}}(T_{a_2}(\check{f}(t_2)), 0),$$

which yields (A.1) and completes the proof of (iv).

We now verify property (v). In view of Remark 2.3, it is equivalent to show that for all $f \in S_c(\Lambda_k, \mathbb{V})$, we have $\tilde{T}_{1/n}(f) \rightarrow 0$ in $S_c(\Lambda_k, \mathbb{V})$. Suppose that this is not the case. Then, for some compact $K \subset \mathbb{R}^d$, some $\epsilon_0 > 0$ and a sequence $t_n \in K$, we have

$$d_{\mathbb{V}}(T_{1/n}(\check{f}(t_n)), 0) \geq \epsilon_0 > 0.$$

Since K is compact, for some $n' \rightarrow \infty$, we have $t_{n'} \rightarrow$ some $t_* \in K$, and by the continuity of \check{f} , we have $\check{f}(t_n) \rightarrow \check{f}(t_*)$ in \mathbb{V} . For all $\delta > 0$, fixed, the radial monotonicity implies that

$$0 < \epsilon_0 \leq \limsup_{n' \rightarrow \infty} d_{\mathbb{V}}(T_{1/n'}(\check{f}(t_{n'})), 0) \leq \lim_{n' \rightarrow \infty} d_{\mathbb{V}}(T_{\delta}(\check{f}(t_{n'})), 0) = d_{\mathbb{V}}(T_{\delta}(\check{f}(t_*)), 0).$$

Property (v), for the scaling action $\{T_a\}$, however, entails that $T_{\delta}(\check{f}(t_*)) \rightarrow 0$ in \mathbb{V} , as $\delta \downarrow 0$, which yields a contradiction with the above inequality and completes the proof. \square

The following result shows that the linear operator actions considered in Remark 2.5 are in fact actions under mild natural conditions on the operator exponent \mathcal{H} .

Lemma A.1. *Let \mathbb{V} be a Hilbert space and $\mathcal{H} : \mathbb{V} \rightarrow \mathbb{V}$ a bounded linear operator such that*

$$\Re(\text{sp}(\mathcal{H})) \subset (0, \infty) \quad \text{and} \quad 2\Re\langle \mathcal{H}x, x \rangle_{\mathbb{V}} = \langle (\mathcal{H} + \mathcal{H}^*)x, x \rangle_{\mathbb{V}} > 0, \quad (\text{A.2})$$

for all $x \neq 0$. Then, $T_c := c^{\mathcal{H}}$, $c > 0$ is a scaling action in the sense of Definition 2.5. Here $\Re(\text{sp}(\mathcal{H}))$ denotes the set of real parts of the elements in the spectrum of \mathcal{H} .

Proof. Properties (i)–(iii) of Definition 2.5 are immediate. Indeed, we have $(c_1 c_2)^{\mathcal{H}} = c_1^{\mathcal{H}} c_2^{\mathcal{H}}$, $c_1, c_2 > 0$ and $1^{\mathcal{H}} = \text{I}$, while Property (iii) follows from the strong continuity (in operator norm) of $c \mapsto c^{\mathcal{H}}$. To establish the radial monotonicity (Property (iv) in Definition 2.5), it is enough to show that for all $x \neq 0$, the function $\varphi(c) := \langle c^{\mathcal{H}} x, c^{\mathcal{H}} x \rangle_{\mathbb{V}}$ is strictly increasing in $c > 0$. To this end, we will show that $\varphi'(c) > 0$ for all $c > 0$ and $x \neq 0$. By (5.2), $h^{-1}((c+h)^{\mathcal{H}} - c^{\mathcal{H}}) \rightarrow \mathcal{H} c^{\mathcal{H}-1}$, in operator norm, as $h \rightarrow 0$. Thus, for $c > 0$,

$$\begin{aligned} \varphi'(c) &= \lim_{h \rightarrow 0} \frac{1}{h} \left(\langle (c+h)^{\mathcal{H}} x, (c+h)^{\mathcal{H}} x \rangle_{\mathbb{V}} - \langle c^{\mathcal{H}} x, (c+h)^{\mathcal{H}} x \rangle_{\mathbb{V}} \right. \\ &\quad \left. + \langle c^{\mathcal{H}} x, (c+h)^{\mathcal{H}} x \rangle_{\mathbb{V}} - \langle c^{\mathcal{H}} x, c^{\mathcal{H}} x \rangle_{\mathbb{V}} \right) \\ &= \langle \mathcal{H} c^{\mathcal{H}-1} x, c^{\mathcal{H}} x \rangle_{\mathbb{V}} + \langle c^{\mathcal{H}}, \mathcal{H} c^{\mathcal{H}-1} x \rangle_{\mathbb{V}} = 2c^{-1} \Re\langle \mathcal{H} c^{\mathcal{H}} x, c^{\mathcal{H}} x \rangle_{\mathbb{V}}, \end{aligned}$$

which is strictly positive, by assumption. Finally, property (v) follows from the first condition in (A.2) as in the proof of Relation (5.19). \square

A.2. Proof of Proposition 3.6.

Proposition 3.6 is the key to establishing the a.e. intrinsic stationarity of the tangent fields in Theorem 3.5. This section outlines its proof, which is based on the following lemma and the Egorov and Lusin Theorems.

Lemma A.2. *Let (K, ρ_K) be a compact metric space and (E, ρ_E) be a metric space. Suppose that f_n and $f : K \rightarrow E$ are Borel measurable functions.*

If the function f is continuous, then

$$\sup_{x \in K} \rho_E(f_n(x), f(x)) \rightarrow 0, \quad \text{as } n \rightarrow \infty, \quad (\text{A.3})$$

if and only if

$$\rho_E(f_n(x_n), f(x)) \rightarrow 0, \quad \text{whenever } \rho_K(x_n, x) \rightarrow 0. \quad (\text{A.4})$$

Proof. ('if') Suppose that (A.4) holds and assume that (A.3) fails. Then, for some $\epsilon_0 > 0$, there exist an infinite sequence $n_k \in \mathbb{N}$ and $x_{n_k} \in K$, such that $\rho_E(f_{n_k}(x_{n_k}), f(x_{n_k})) \geq \epsilon_0 > 0$. It is easy to see that $\{x_{n_k}\}$ is also an infinite sequence, since for every $k_0 \in \mathbb{N}$, by (A.4), we have $f_{n_k}(x_{n_{k_0}}) \rightarrow f(x_{n_{k_0}})$, as $n_k \rightarrow \infty$. Indeed, had $\{x_{n_k}\}$ been a finite set, for some infinite subsequence $\{n_k\} \subset \{n_{k_0}\}$, we would have $x_{n_k} = x_{n_{k_0}}$ and hence $\rho_E(f_{n_k}(x_{n_k}), f(x_{n_k})) = \rho_E(f_{n_k}(x_{n_{k_0}}), f(x_{n_{k_0}})) \rightarrow 0$, contradicting the construction of the x_{n_k} 's.

The infinite sequence $\{x_{n_k}\}$ is included in the compact K , and hence it has a converging subsequence $x_{n_k(m)} \rightarrow x$. This, in view of (A.4), implies that $f_{n_k(m)}(x_{n_k(m)}) \rightarrow f(x)$ in \mathbb{V} . Since f is continuous at x , however, $f(x_{n_k(m)}) \rightarrow f(x)$. This, by the triangle inequality, implies

$$\begin{aligned} & \rho_E(f_{n_k(m)}(x_{n_k(m)}), f(x_{n_k(m)})) \\ & \leq \rho_E(f_{n_k(m)}(x_{n_k(m)}), f(x)) + \rho_E(f(x), f(x_{n_k(m)})) \rightarrow 0, \text{ as } n_k(m) \rightarrow \infty. \end{aligned}$$

This contradicts the assumption that $\rho_E(f_{n_k(m)}(x_{n_k(m)}), f(x_{n_k(m)})) \geq \epsilon_0$.

('only if') Let $x_n \rightarrow x$. By the triangle inequality, we have that

$$\begin{aligned} \rho_E(f_n(x_n), f(x)) & \leq \rho_E(f_n(x_n), f(x_n)) + \rho_E(f(x_n), f(x)) \\ & \leq \sup_{x' \in K} \rho_E(f_n(x'), f(x')) + \rho_E(f(x_n), f(x)), \end{aligned}$$

which converges to zero by (A.3) and the continuity of f . □

The next result is a restatement of Theorem 7.5.1 in [Dudley \[2002\]](#).

Theorem A.3 (Egorov). *Let (B, \mathcal{B}, μ) be a finite measure space and (Y, ρ_Y) be a separable metric space. Suppose that $f_n : B \rightarrow Y$, $n = 1, \dots$ are measurable functions such that, for μ -almost all $x \in B$,*

$$f_n(x) \rightarrow f(x), \quad \text{as } n \rightarrow \infty.$$

Then, for all $\epsilon > 0$, there exists a measurable set $B_\epsilon \subset B$, such that

$$\mu(B \setminus B_\epsilon) < \epsilon \quad \text{and} \quad \sup_{x \in B_\epsilon} \rho_Y(f_n(x), f(x)) \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

We present next a relatively general form of the classic Lusin's theorem stating that every Borel function is nearly continuous. The proof follows the elegant 3-line argument given in Theorem 1 on page 56 in [Loeb and Talvila \[2004\]](#). We provide a bit more detail and tailor the result to the case of metric spaces.

Theorem A.4 (Lusin). *Let (X, ρ_X) be a metric space and (Y, ρ_Y) be a separable metric space. Let also $f : X \rightarrow Y$ be a Borel measurable function and μ be a finite Borel measure μ on X .*

For every $\epsilon > 0$, there exists a closed set $F \subset X$, such that $\mu(X \setminus F) < \epsilon$ and $f : F \rightarrow Y$ is continuous. If (X, ρ_X) is separable and complete, then the set F can be taken to be compact.

Proof. We will essentially unpack the argument on page 56 of [Loeb and Talvila \[2004\]](#) with small modifications.

By Theorem 7.1.3 on page 175 in [Dudley \[2002\]](#) every finite Borel measure μ on (X, ρ_X) is *closed regular*, that is, for every Borel set A in X , we have

$$\mu(A) = \sup\{\mu(F) : F \subset A, F \text{ is closed}\}. \quad (\text{A.5})$$

Recall that μ is called regular if the sets F above can be taken to be compact. Ulam's Theorem implies that if (X, ρ_X) is separable and complete, then μ is regular [cf. Theorem 7.1.4 in [Dudley, 2002](#)].

We now fix an $\epsilon > 0$ and construct the closed set F . Since (Y, ρ_Y) is separable, it is second countable, i.e., its topology has a countable base. Namely, there exists a countable collection of open sets $\{V_n, n \in \mathbb{N}\}$ in Y such that every open set $V \subset Y$ can be represented as a union of V_n 's, i.e., $V = \cup\{V_n : V_n \subset V, n \in \mathbb{N}\}$.

Following [Loeb and Talvila \[2004\]](#), by (A.5) since μ is finite, we can find closed sets $F_n \subset f^{-1}(V_n)$ and $F'_n \subset X \setminus f^{-1}(V_n)$ in X (compact if μ is regular), such that

$$\mu(f^{-1}(V_n) \setminus F_n) < \frac{\epsilon}{2^{n+1}} \quad \text{and} \quad \mu([X \setminus f^{-1}(V_n)] \setminus F'_n) < \frac{\epsilon}{2^{n+1}}.$$

Observe that

$$\mu(X \setminus (F_n \cup F'_n)) = \mu(f^{-1}(V_n) \setminus F_n) + \mu([X \setminus f^{-1}(V_n)] \setminus F'_n) < \frac{\epsilon}{2^n}.$$

Define $F := \cap_{n \in \mathbb{N}} (F_n \cup F'_n)$ and notice that F is closed and in fact compact if (X, ρ_X) is separable and complete. The above relation implies moreover that

$$\mu(X \setminus F) \leq \sum_{n \in \mathbb{N}} \mu(X \setminus (F_n \cup F'_n)) < \sum_{n \in \mathbb{N}} \frac{\epsilon}{2^n} = \epsilon.$$

To complete the proof, it remains to show that $f : F \rightarrow Y$ is continuous. To this end, it is enough to show that for every $x \in F$ and every V_n such that $f(x) \in V_n$, there is an open set $U \ni x$ such that $f(U \cap F) \subset V_n$. Suppose $f(x) \in V_n$ and consider the open set $U := X \setminus F'_n$. Since $F \subset F_n \cup F'_n$ and $U \cap F'_n = \emptyset$, we have

$$U \cap F \subset U \cap F_n \subset f^{-1}(V_n),$$

which implies $f(U \cap F) \subset V_n$. We have thus established the desired continuity of f on F . \square

Remark A.1. Loeb and Talvila's proof of Lusin's Theorem A.4 is not constructive and it does not use approximation arguments based on the Tietze–Uryson Lemma and Egorov's theorem as many other proofs in the literature [see, e.g., Theorem 7.5.2 in [Dudley, 2002](#)]. This makes it possible to extend Lusin's theorem to functions taking values in an arbitrary separable metric space.

We conclude this section with the proof of Proposition 3.6.

Proof of Proposition 3.6: By Egorov's Theorem (see Theorem A.3, above), there is a Borel set $B_\epsilon \subset B$ such that $\mu(B \setminus B_\epsilon) < \epsilon/2$ and

$$\sup_{s \in B_\epsilon} \rho_E(F_n(s), G(s)) \rightarrow 0, \quad (\text{A.6})$$

as $n \rightarrow \infty$. Observe that since the Lebesgue measure is closed regular (recall (A.5)), one can choose the set B_ϵ to be closed. Therefore, B_ϵ with the usual metric in \mathbb{R}^d is a complete and separable metric space. Hence, we can apply Lusin's Theorem A.4 to $X := B_\epsilon \subset \mathbb{R}^d$ and $Y := E$ to conclude that there is a further compact set $K_\epsilon \subset B_\epsilon$, such that $\mu(B_\epsilon \setminus K_\epsilon) < \epsilon/2$ and the function $G : K_\epsilon \rightarrow E$ is *continuous*.

Observe that

$$\mu(B \setminus K_\epsilon) = \mu(B \setminus B_\epsilon) + \mu(B_\epsilon \setminus K_\epsilon) < \epsilon.$$

By Lemma A.2, the continuity of G on K_ϵ and the uniform convergence (A.6) imply (3.17). \square

A.3. Supplementary results and proofs for Section 3.

The following convergence to types lemma is rather useful.

Lemma A.5. *Let $\{T_a, a > 0\}$ be a scaling action on some complete separable (not necessarily linear) metric space $(\mathbb{X}, d_{\mathbb{X}})$. Let also $\xi, \tilde{\xi}$ and ξ_n be random elements taking values in \mathbb{X} . Then the following hold.*

- (i) *If ξ is non-zero, then $T_{a'}(\xi) \stackrel{d}{=} T_{a''}(\xi)$ implies $a' = a''$.*
- (ii) *Suppose that $\xi_n \xrightarrow{d} \xi$ and $\tilde{\xi}_n := T_{a_n}(\xi_n) \xrightarrow{d} \tilde{\xi}$, for some sequence $a_n > 0$, where both ξ and $\tilde{\xi}$ are non-zero. Then $a_n \rightarrow a$ for some $a > 0$ and $\tilde{\xi} \stackrel{d}{=} T_a(\xi)$.*
- (iii) *If ξ is non-zero and $T_{a_n}(\xi) \xrightarrow{d} 0$, then $a_n \rightarrow 0$.*

Proof. Part (i). Suppose that $a' < a''$. Then,

$$\xi \stackrel{d}{=} T_{a''}^{-1} \circ T_{a'}(\xi) \equiv T_{a'/a''}(\xi),$$

which implies $\xi \stackrel{d}{=} T_{c_n}(\xi)$, for all $n \in \mathbb{N}$, where $c_n = (a'/a'')^n \downarrow 0$. Thus, for every B_r , we obtain

$$\mathbb{P}(\xi \in B_r) = \mathbb{P}(T_{c_n}(\xi) \in B_r) = \mathbb{P}(\xi \in T_{1/c_n}(B_r)).$$

Since $1/c_n \rightarrow \infty$, applying Property (2.15) applied with \mathbb{V} replaced by \mathbb{X} (recall (2.17)), we see that $\mathbb{P}(\xi \in B_r) = 1$, for all $r > 0$, which contradicts the assumption that ξ is non-zero.

Part (ii). We will first show that $\{a_n\}$ is bounded away from 0 and ∞ . Indeed, suppose that $a_{n'} \rightarrow \infty$ for some $n' \rightarrow \infty$. Consider the balls

$$B_r := \{x \in \mathbb{X} : d_{\mathbb{X}}(x, 0) < r\}$$

and observe that all but countably many of them are continuity sets for the distribution of ξ . Indeed, the sets $\partial B_r := \overline{B_r} \setminus B_r$, $r > 0$ are pairwise disjoint in r and for each $\epsilon > 0$, there are at most $1/\epsilon$ distinct values for r , such that $\mathbb{P}(\xi \in \partial B_r) > \epsilon$.

For every $r > 0$ such that $\mathbb{P}(\xi \in \partial B_r) = 0$, since $\xi_{n'} \xrightarrow{d} \xi$, we have

$$\begin{aligned} \mathbb{P}(\xi \in B_r) &= \lim_{n' \rightarrow \infty} \mathbb{P}(\xi_{n'} \in B_r) = \lim_{n' \rightarrow \infty} \mathbb{P}(T_{a_{n'}}(\xi_{n'}) \in T_{a_{n'}}(B_r)) \\ &\geq \limsup_{n' \rightarrow \infty} \mathbb{P}(\tilde{\xi}_{n'} \in T_m(B_r)), \end{aligned} \quad (\text{A.7})$$

where $\tilde{\xi}_{n'} := T_{a_{n'}}(\xi_{n'})$ and m is an arbitrary fixed integer. Here, we used the fact that $T_m(B_r) \subset T_{a_{n'}}(B_r)$, for all large enough n' , by (2.15) and (2.17).

Now, since T_m is a homeomorphism, we have $\partial T_m(B_r) = T_m(\partial B_r)$ are disjoint in $r > 0$, and by the above argument, for all but countably many r 's, we have $\mathbb{P}(\xi \in \partial T_m(B_r)) = 0$ and hence $\mathbb{P}(\tilde{\xi}_{n'} \in T_m(B_r)) \rightarrow \mathbb{P}(\tilde{\xi} \in T_m(B_r))$, as $n' \rightarrow \infty$. Therefore, in view of (A.7), we obtain

$$\mathbb{P}(\xi \in B_r) \geq \mathbb{P}(\tilde{\xi} \in T_m(B_r)), \quad \text{for all } m \text{ and all but countably many } r > 0.$$

Relation (2.15), however, implies that $T_m(B_r) \uparrow E$ as $m \rightarrow \infty$, which implies

$$\mathbb{P}(\xi \in B_r) = 1 = \lim_{m \rightarrow \infty} \mathbb{P}(\tilde{\xi} \in T_m(B_r)),$$

for all but countably many r . This implies that $\mathbb{P}(\xi = 0) = 1$, which is a contradiction.

We have thus shown that the sequence $\{a_n\}$ is bounded above. One can similarly show that $\{a_n\}$ is bounded away from 0. Indeed, by defining $\tilde{a}_n := 1/a_n$, we see that $\xi_n = T_{\tilde{a}_n}(\tilde{\xi}_n) \xrightarrow{d} \xi$. Therefore, repeating the above argument with a_n, ξ_n and ξ replaced by $\tilde{a}_n, \tilde{\xi}_n$ and $\tilde{\xi}$, respectively, we see that $\{\tilde{a}_n \equiv 1/a_n\}$ is bounded.

We have thus shown that $\{a_n\}$ can only have positive cluster points. Suppose that $a_{n'} \rightarrow a' > 0$ and $a_{n''} \rightarrow a'' > 0$, for some sub-sequences $n', n'' \rightarrow \infty$. Since the space $(\mathbb{X}, d_{\mathbb{X}})$ is separable, by the Skorokhod-Dudley representation [cf. Theorem 3.30 of Kallenberg, 1997], on a suitable probability space we can define ξ^* and ξ_n^* such that

$$\xi_n^* \xrightarrow{d} \xi_n, \quad \xi^* \xrightarrow{d} \xi, \quad \text{and} \quad \xi_n^* \rightarrow \xi^*, \quad \text{almost surely.}$$

Thus, the continuity property (iii) in Definition 2.5, implies that

$$T_{a_{n'}}(\xi_{n'}^*) \rightarrow T_{a'}(\xi^*) \quad \text{and} \quad T_{a_{n''}}(\xi_{n''}^*) \rightarrow T_{a''}(\xi^*),$$

almost surely. Since also $T_{a_n}(\xi_n^*) \xrightarrow{d} T_{a_n}(\xi_n) \xrightarrow{d} \tilde{\xi}$, and $\xi \xrightarrow{d} \xi^*$, we obtain

$$T_{a'}(\xi) \xrightarrow{d} \tilde{\xi} \xrightarrow{d} T_{a''}(\xi). \quad (\text{A.8})$$

By part (i), this is only possible if $a' = a''$. We have thus shown that the sequence $\{a_n\}$ has a unique cluster point $a = a' = a'' > 0$ and in view of (A.8), that $T_a(\xi) \xrightarrow{d} \tilde{\xi}$.

Part (iii). Suppose that $\limsup_{n \rightarrow \infty} a_n > 0$, i.e., for some subsequence $n' \rightarrow \infty$, we have $a_{n'} \geq \epsilon_0 > 0$, for all n' . Then, in view of (2.17), for all $r > 0$, we have

$$\mathbb{P}(T_{\epsilon_0}(\xi) \in B_r^c) \leq \mathbb{P}(T_{a_{n'}}(\xi) \in B_r^c).$$

Since $T_{a_{n'}}(\xi) \xrightarrow{d} 0$, the right-hand side vanishes, as $n' \rightarrow \infty$. On the other hand, since ξ is nonzero, so is $T_{\epsilon_0}(\xi)$ and the left-hand side is positive for sufficiently small $r > 0$. This contradiction yields $\limsup_{n \rightarrow \infty} a_n = 0$. \square

The next result is used in the proof of Corollary 3.7, given below.

Lemma A.6. *Let $X := \{X(\lambda), \lambda \in \Lambda_k\}$ and $X_n := \{X_n(\lambda), \lambda \in \Lambda_k\}$ be random fields in $S_c(\Lambda_k, \mathbb{V})$ such that $X_n \xrightarrow{d} X$. Then for any sequences $v_n \rightarrow 0$ and $r_n \rightarrow 1$, we have*

$$\{X_n(v_n + r_n \cdot \lambda), \lambda \in \Lambda_k\} \xrightarrow{d} \{X(\lambda), \lambda \in \Lambda_k\}.$$

Proof. Let $Y_n(\lambda) = X_n(v_n + r_n \cdot \lambda)$ and $\check{Y}_n = Y_n(\lambda_t)$. By Proposition 2.3, conditions (i) and (ii) hold for \check{X}_n . We need to show that they also hold for \check{Y}_n . The proof for (i) is an easy application of the Skorokhod-Dudley representation Theorem 3.30 on page 56 in Kallenberg [1997]. We focus on proving (ii).

First consider the relationship between \check{X}_n and \check{Y}_n . It follows that

$$\check{Y}_n(t) = X_n(v_n + r_n \cdot \lambda_t) = \int \check{X}_n(v_n + r_n u) \lambda_t(du),$$

where one can write

$$\lambda_t(du) = \sum_{i=1}^{M_k+1} c_i(t) \delta_{s_i(t)}(du),$$

for some continuous functions $c_i(t) \in \mathbb{C}$ and $s_i(t) \in \mathbb{R}^d$, $i = 1, \dots, M_k + 1$ (recall (2.7)).

Since $\check{X}_n(t) = X_n(\lambda_t)$, for any K and n large enough,

$$\begin{aligned} \sup_{\|t\|_2 \leq K} d_{\mathbb{V}}(\check{Y}_n(t), \check{X}_n(t)) &\leq \sup_{\|t\|_2 \leq K} \sum_{i=1}^{M_k+1} d_{\mathbb{V}}\left(c_i(t) \check{X}_n(v_n + r_n s_i(t)), c_i(t) \check{X}_n(s_i(t))\right) \\ &\leq (M_k + 1) \sup_{\substack{\|s\|, \|t\| \leq 2K \\ |c| \leq C_K, \|s-t\| \leq \delta_n}} d_{\mathbb{V}}(c \check{X}_n(s), c \check{X}_n(t)), \end{aligned} \tag{A.9}$$

where $\delta_n := |r_n - 1|S_K + |v_n| \rightarrow 0$ with

$$S_K := \sup_{\substack{\|t\| \leq K \\ i=1, \dots, M_k+1}} \|s_i(t)\| \quad \text{and} \quad C_K := \sup_{\substack{\|t\| \leq K \\ i=1, \dots, M_k+1}} |c_i(t)|.$$

Therefore, for any $\eta, \epsilon > 0$, by (ii) in Proposition 2.3 there exists n large enough such that, $\mathbb{P}(A_n < \eta) > 1 - \epsilon$, where

$$A_n := \sup_{\substack{\|s\|, \|t\| \leq 2K \\ \|s-t\| \leq \delta_n}} d_{\mathbb{V}}(\check{X}_n(s), \check{X}_n(t)).$$

Then according to (2.3), (2.4) and (A.9), we have on the event $\{A_n < \eta\}$, that

$$\sup_{\|t\| \leq K} d_{\mathbb{V}}(\check{Y}_n(t), \check{X}_n(t)) \leq (M_k + 1) f_{C_K}(\eta),$$

where for any $C_K > 0$, $f_{C_K}(\eta) := \sup_{d_{\mathbb{V}}(x,y) < \eta, |c| < C_K} d_{\mathbb{V}}(cx, cy) \rightarrow 0$ as $\eta \rightarrow 0$. Thus, on $\{A_n < \eta\}$, we have

$$\begin{aligned} \sup_{\substack{\|s\|, \|t\| \leq K \\ \|s-t\| \leq \delta_n}} d_{\mathbb{V}}(\check{Y}_n(s), \check{Y}_n(t)) &\leq 2 \sup_{\|t\| \leq K} d_{\mathbb{V}}(\check{Y}_n(t), \check{X}_n(t)) + \sup_{\substack{\|s\|, \|t\| \leq K \\ \|s-t\| \leq \delta_n}} d_{\mathbb{V}}(\check{X}_n(s), \check{X}_n(t)) \\ &\leq 2(M_k + 1) f_{C_K}(\eta) + \eta. \end{aligned}$$

Thus, the second condition of Proposition 2.3 for \check{Y}_n follows. \square

Proof of Corollary 3.7: Consider the context of the proof of Theorem 3.5. When $T_{s,c} = T_c$, $c > 0$, for a *fixed* scaling action T_c . Relation (3.19) becomes

$$\begin{aligned}\xi_n &= \left\{ T_{c_n(s_n)} \circ T_{c_n(s)}^{-1} \circ T_{c_n(s)} X(s, (1/n) \cdot (w_n + \lambda)), \lambda \in \Lambda_k \right\} \\ &=: T_{c_n(s_n)/c_n(s)} \tilde{\xi}_n.\end{aligned}\tag{A.10}$$

where $c_n(s) := c(s, 1/n)$, and we used the fact that $T_{c_n(s)}^{-1} = T_{1/c_n(s)}$.

Relation (3.18) implies that

$$\xi_n \xrightarrow{d} \xi := \{Y_s(\lambda), \lambda \in \Lambda_k\}.$$

On the other hand, by Lemma A.6, we have

$$\tilde{\xi}_n = \{T_{c_n(s)} X(s, (1/n) \cdot (w_n + \lambda)), \lambda \in \Lambda_k\} \xrightarrow{d} \tilde{\xi} := \{Y_s(w + \lambda), \lambda \in \Lambda_k\}.$$

Thus, in view (A.10), we have $\tilde{\xi}_n = T_{c_n(s)/c_n(s_n)} \xi_n \xrightarrow{d} \tilde{\xi}$ and since both ξ and $\tilde{\xi}$ are non-zero, Lemma A.5 implies that

$$\frac{c_n(s)}{c_n(s_n)} \equiv \frac{c_n(s)}{c_n(s + w_n/n)} \rightarrow a_s(\{w_n\}) > 0.$$

One can verify that the limit $a_s(\{w_n\})$ is independent of the choice of the sequence $\{w_n\}$. Indeed, if there exists another $w'_n \rightarrow w$ and $s + w'_n/n \in K_\epsilon$ then we will have

$$\{Y_s(w + \lambda), \lambda \in \Lambda_k\} \stackrel{d}{=} \{T_{a_s(\{w_n\})}(Y_s(\lambda)), \lambda \in \Lambda_k\} \stackrel{d}{=} \{T_{a_s(\{w'_n\})}(Y_s(\lambda)), \lambda \in \Lambda_k\},$$

which shows $a_s(\{w_n\}) = a_s(\{w'_n\})$ by (i) of Lemma A.5. Thus, we can just use the notation $a_s(w)$ and we have

$$\{Y_s(w + \lambda), \lambda \in \Lambda_k\} \stackrel{d}{=} \{T_{a_s(w)} Y_s(\lambda), \lambda \in \Lambda_k\}.\tag{A.11}$$

To prove that (3.16) holds, or equivalently $T_{c_n(s)/c_n(s_n)} \rightarrow T_1 \equiv I$, we only need to verify $a_s(w) = 1$, which we do next. By (A.11), it is easy to see that $a_s(0) = 1$ and $a_s(w + u) = a_s(w)a_s(u)$. By Theorem 3.3, there is a positive scalar $\alpha = \alpha(s) > 0$, such that

$$\{Y_s(r \cdot \lambda), \lambda \in \Lambda_k\} \stackrel{d}{=} \{T_{r^\alpha}(Y_s(\lambda)), \lambda \in \Lambda_k\}, \text{ for all } r > 0.\tag{A.12}$$

Consider $Y_s(r \cdot (w + \lambda))$ with $r \in \mathbb{N}$. On one hand, (A.11) and (A.12) imply that

$$\{Y_s(r \cdot (w + \lambda)), \lambda \in \Lambda_k\} \stackrel{d}{=} \{T_{r^\alpha}(Y_s(w + \lambda)), \lambda \in \Lambda_k\} \stackrel{d}{=} \{T_{r^\alpha a_s(w)}(Y_s(\lambda)), \lambda \in \Lambda_k\}.\tag{A.13}$$

On the other hand, since $a_s(rw) = a_s^r(w)$, for all $r \in \mathbb{N}$, viewing $Y_s(r \cdot (w + \lambda))$ as $Y_s(rw + r \cdot \lambda)$, by (A.11), we have

$$\{Y_s(r \cdot (w + \lambda)), \lambda \in \Lambda_k\} \stackrel{d}{=} \{T_{a_s(rw)}(Y_s(r \cdot \lambda)), \lambda \in \Lambda_k\} \stackrel{d}{=} \{T_{a_s^r(w) r^\alpha}(Y_s(\lambda)), \lambda \in \Lambda_k\}.\tag{A.14}$$

Thus, by Lemma A.5 applied to (A.13) and (A.14), we obtain $a_s(w) = a_s^r(w)$ for all $r \in \mathbb{N}$, which, since $a_s(w) > 0$, implies $a_s(w) = 1$. \square

A.4. Supplemental background and some proofs for Section 4.

In Bochner's Theorem, we need the notion of integration of a \mathbb{C} -valued function on \mathbb{R}^d with respect to a *finite* \mathbb{T}_+ -valued measure. A finite \mathbb{T}_+ -valued measure is a mapping $\mu : \mathcal{B}(\mathbb{R}^d) \rightarrow \mathbb{T}_+$ from the class of Borel sets in \mathbb{R}^d to \mathbb{T}_+ that is *countably additive*. Notice that this readily implies that $\mu(\emptyset) = 0$; that μ is monotone, i.e., $\mu(A) \leq \mu(B)$ as positive operators for $A \subset B$; and that μ is finite since $\mu(\mathbb{R}^d) \in \mathbb{T}_+$. This notion can be developed along the line of ordinary Lebesgue integration, making use of the completeness of \mathbb{T} . We will provide a brief outline below and leave the details to Section S.3 in Supplement. (The construction naturally extends to the case of σ -finite \mathbb{T}_+ -valued measures.)

Let μ be a finite \mathbb{T}_+ -valued measure. We follow the development of ordinary Lebesgue integration:

- (i) For any real nonnegative simple function $f = \sum_{i=1}^k c_i I_{A_i}$, define $\int f d\mu = \sum_{i=1}^k c_i \mu(A_i)$.
- (ii) For nonnegative measurable functions f , let

$$\int f d\mu = \lim_{n \rightarrow \infty} \int f_n d\mu \quad (\text{A.15})$$

in $(\mathbb{T}, \|\cdot\|, \|\cdot\|_{\text{tr}})$ where $\{f_n\}$ is any sequence of simple functions such that

- (a) $f_n \leq f_{n+1}$,
- (b) $f_n(x) \uparrow f(x)$ for all x ,
- (c) $\int f_n d\mu \leq \mathcal{B}$ for all n and some fixed $\mathcal{B} \in \mathbb{T}_+$.

The existence of $\{f_n\}$ satisfying (a) and (b) for any given nonnegative measurable f follows from standard measure theory. However, we need the extra condition (c) (along with the completeness of \mathbb{T}_+) to ensure that the limit on rhs of (A.15) exists and does not depend on the choice of $\{f_n\}$. Clearly, (c) is automatically fulfilled if f is bounded.

- (iii) For a general real measurable f , let

$$\int f d\mu = \int f_+ d\mu - \int f_- d\mu$$

provided both terms on the right is finite. For a general complex f , let

$$\int f d\mu = \int f_{\text{re}} d\mu + \mathbf{i} \int f_{\text{im}} d\mu$$

where $f_{\text{re}}, f_{\text{im}}$ be the real and imaginary parts, respectively.

It is immediate that $\|\mu\|_{\text{tr}}(A) := \|\mu(A)\|_{\text{tr}}$, $A \in \mathcal{B}(\mathbb{R}^d)$ defines a finite Borel measure referred to as the *trace measure* of μ . The following useful integrability criterion is straightforward.

Proposition A.7. *The integral $\int_{\mathbb{R}^d} f d\mu \in \mathbb{T}$ is well-defined, for all $f \in L^1(\mathbb{R}^d, \|\mu\|_{\text{tr}})$, and*

$$\int_{\mathbb{R}^d} f(x) \|\mu\|_{\text{tr}}(dx) = \text{trace} \left(\int_{\mathbb{R}^d} f(x) \mu(dx) \right).$$

The integral can be readily extended to σ -finite \mathbb{T} -valued signed measures. One can also naturally consider finite signed \mathbb{T} -valued measures, namely $\mu = \mu_+ - \mu_-$ where μ_+, μ_- are both finite \mathbb{T}_+ -valued measures, by $\int f d\mu := \int f d\mu_+ - \int f d\mu_-$. In this case, the trace measure of μ is the (scalar) signed measure defined as $\|\mu\|_{\text{tr}}(\cdot) := \|\mu_+\|_{\text{tr}}(\cdot) - \|\mu_-\|_{\text{tr}}(\cdot)$.

Next we turn to the stochastic integral in Cramér's representation. Recall that $\mathbb{L}^2(\Omega)$ is the L^2 -space of all \mathbb{V} -valued random elements η on the probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with $\mathbb{E}\|\eta\|^2 < \infty$, equipped with the inner product

$$\langle \eta_1, \eta_2 \rangle_\Omega := \mathbb{E}\langle \eta_1, \eta_2 \rangle.$$

We briefly discuss the properties of the orthogonal random measure $\xi = \{\xi(A), A \in \mathcal{B}(\mathbb{R}^d)\}$ in Definition 4.6. Condition (ii) therein readily implies the orthogonality as well as finite additivity of the measure ξ . Indeed,

$$\mathbb{E}[\xi(A) \otimes \xi(B)] = \mu(\emptyset) = 0, \quad \text{for all disjoint } A, B \in \mathcal{B}(\mathbb{R}^d), \quad (\text{A.16})$$

where 0 stands for the zero operator in \mathbb{T} . This implies in particular that $\langle \xi(A), \xi(B) \rangle_\Omega = 0$ (cf Remark A.2). We have, moreover, that for all (orthogonal) projection operators $\Pi : \mathbb{V} \rightarrow \mathbb{V}$, the random measure $\eta(\cdot) := \Pi \xi(\cdot)$ is also an orthogonal measure with control measure $\Pi \mu(\cdot) \Pi$.

Now, to show finite additivity, observe that for disjoint Borel sets A and B ,

$$\begin{aligned} \mathbb{E}\|\xi(A \cup B) - \xi(A) - \xi(B)\|^2 \\ &= \mathbb{E} \text{trace} \left((\xi(A \cup B) - \xi(A) - \xi(B)) \otimes (\xi(A \cup B) - \xi(A) - \xi(B)) \right) \\ &= \text{trace} \left(\mu(A \cup B) - \mu(A) - \mu(B) \right) = 0, \end{aligned}$$

by the fact that the \mathbb{E} and trace operators can be exchanged and the finite additivity of μ . This shows $\xi(A \cup B) = \xi(A) + \xi(B)$, almost surely.

Finally, Condition (i) in Definition 4.6 along with the established finite additivity implies the σ -additivity of ξ , in the sense that for any sequence of pairwise disjoint $A_n \in \mathcal{B}(\mathbb{R}^d)$, $n = 1, 2, \dots$, we have

$$\xi \left(\bigcup_{n=1}^{\infty} A_n \right) = \sum_{n=1}^{\infty} \xi(A_n), \quad \text{almost surely,}$$

where the latter series converges in $\mathbb{L}^2(\Omega)$.

Remark A.2. Observe that for random vectors $X, Y \in \mathbb{L}^2(\Omega)$, we have that

$$\mathbb{E}[X \otimes Y] = 0 \quad \text{implies} \quad \langle X, Y \rangle_\Omega = 0,$$

but the converse implication is not always true. Thus, the orthogonality condition in (A.16) is stronger than requiring simply $\langle \xi(A), \xi(B) \rangle_\Omega = 0$.

Introduce the finite Borel scalar measure

$$\|\mu\|_{\text{tr}}(A) := \text{trace}(\mu(A)) \equiv \|\mu(A)\|_{\text{tr}}, \quad A \in \mathcal{B}(\mathbb{R}^d).$$

It is easy to verify (cf Lemma S.2.2 in Supplement) that, for all $A \in \mathcal{B}(\mathbb{R}^d)$,

$$\|\xi(A)\|_\Omega^2 = \|\mu(A)\|_{\text{tr}} = \|\mu\|_{\text{tr}}(A).$$

Next, for an orthogonal random measure ξ with control measure μ , we sketch the construction of the stochastic integral $\mathcal{I}_\xi f := \int_{\mathbb{R}^d} f(t) \xi(dt)$ defined for all functions $f \in \mathbb{L}^2(\mathbb{R}^d, \|\mu\|_{\text{tr}})$, i.e., all measurable $f : \mathbb{R}^d \rightarrow \mathbb{C}$ with $\int_{\mathbb{R}^d} |f(t)|^2 \|\mu\|_{\text{tr}}(dt) < \infty$. For any simple function $\psi(x) = \sum_{j=1}^n c_j I_{A_j}(x)$, where $A_j \in \mathcal{B}(\mathbb{R}^d)$, $A_j \cap A_{j'} = \emptyset$ when $j \neq j'$, define the integral

$$\mathcal{I}_\xi(\psi) = \sum_{j=1}^n c_j \xi(A_j).$$

The integral $\mathcal{I}_\xi(\psi)$ takes values in $\mathbb{L}_\xi^2(\Omega)$, defined as the closure of

$$\text{span}(\xi) := \left\{ \sum_{j=1}^n c_j \xi(A_j), c_j \in \mathbb{C}, A_j \in \mathcal{B}(\mathbb{R}^d), n = 1, 2, \dots \right\}$$

in $\mathbb{L}^2(\Omega)$. Property (ii) in Definition 4.6 entails

$$\|\mathcal{I}_\xi(\psi)\|_\Omega^2 = \int_{\mathbb{R}^d} |\psi(x)|^2 \|\mu\|_{\text{tr}}(dx).$$

Therefore, \mathcal{I}_ξ is an isometric linear mapping between the class of simple functions in the L^2 -space $\mathbb{L}^2(\mathbb{R}^d, \|\mu\|_{\text{tr}})$ and the Hilbert space $\mathbb{L}_\xi^2(\Omega)$.

As the class of simple functions is dense in $\mathbb{L}^2(\mathbb{R}^d, \|\mu\|_{\text{tr}})$ and the integrals $\mathcal{I}_\xi(\psi)$ are dense in $\mathbb{L}_\xi^2(\Omega)$, the linear operator \mathcal{I}_ξ can be uniquely extended to an isometric linear mapping between $\mathbb{L}^2(\mathbb{R}^d, \|\mu\|_{\text{tr}})$ and $\mathbb{L}_\xi^2(\Omega)$. This completes the construction of the stochastic integral

$$\int_{\mathbb{R}^d} f(x) \xi(dx) := \mathcal{I}_\xi(f), \quad f \in \mathbb{L}^2(\mathbb{R}^d, \|\mu\|_{\text{tr}}).$$

Observe, moreover that for all $f, g \in \mathbb{L}^2(\mathbb{R}^d, \|\mu\|_{\text{tr}})$, we have

$$\mathbb{E}[\mathcal{I}_\xi(f) \otimes \mathcal{I}_\xi(g)] = \int_{\mathbb{R}^d} f(x) \overline{g}(x) \mu(dx),$$

where the last integral is well-defined in view of Proposition A.7.

Proof of Proposition 4.16: Part (i): If Y is a real IRF_k then Y is also a real $\text{IRF}_{k+1}^{\widehat{\lambda}}$ with trivial polynomial spectral characteristic. Indeed, since for all $\lambda \in \Lambda_{k+1}$, we have $\partial_{j_1, \dots, j_d}^{k+1} \widehat{\lambda}(0) = 0$, Relation (4.16) becomes

$$Y(\lambda) = \int_{\mathbb{R}^d} \frac{\widehat{\lambda}(u)}{1 \wedge \|u\|^{k+1}} \xi(du) =: \int_{\mathbb{R}^d} \frac{\widehat{\lambda}(u)}{1 \wedge \|u\|^{k+2}} \eta(du),$$

where $\eta(du) := (1 \wedge \|u\|^{k+2})(1 \wedge \|u\|^{k+1})^{-1} \xi(du)$. By taking $\lambda \in \Lambda_{k+1}(\mathbb{R})$, since $Y(\lambda) = \overline{Y(\lambda)}$ and $\widehat{\lambda}(u) = \widehat{\lambda}(-u)$, we obtain that

$$Y(\lambda) = \int_{\mathbb{R}^d} \frac{\widehat{\lambda}(u)}{1 \wedge \|u\|^{k+2}} \overline{\eta(-du)}, \quad \lambda \in \Lambda_{k+1}(\mathbb{R}).$$

By (4.18), the last relation continues to hold for all complex $\lambda \in \Lambda_{k+1}$. Hence, appealing to the uniqueness of the representation (4.16) of Y viewed as an IRF_{k+1} (with k replaced by $k+1$ and ξ by η), we obtain

$$\eta(du) = \overline{\eta(-du)}, \quad \text{almost surely,}$$

or equivalently $\xi(du) = \overline{\xi(-du)}$, a.s., which shows that the orthogonal measure ξ is Hermitian.

The fact that ξ is Hermitian and Y a real IRF_k , imply that for all $\lambda \in \Lambda_k(\mathbb{R})$,

$$Y(\lambda) - \int_{\mathbb{R}^d} \frac{\widehat{\lambda}(u)}{1 \wedge \|u\|^{k+1}} \xi(du) = \sum_{j_1, \dots, j_d} \partial_{j_1, \dots, j_d}^{k+1} \widehat{\lambda}(0) Z_{j_1, \dots, j_d} \quad \text{is real.}$$

By taking suitable *real* λ 's for which $\partial_{j_1, \dots, j_d}^{k+1} \widehat{\lambda}(0)$ vanish for all but each one term in the sum, we obtain that all $(\mathfrak{i})^{k+1} Z_{j_1, \dots, j_d}$ must be real.

Conversely, if ξ is Hermitian and the $(\mathfrak{i})^{k+1} Z_{j_1, \dots, j_d}$'s are real, it is immediate that $Y(\lambda)$ is real for all $\lambda \in \Lambda_k(\mathbb{R})$. This completes the proof of part (i). Part (ii) is an immediate consequence of (i).

Part (iii): One can define Y as in (4.18), where all $(\mathfrak{i})^{k+1} Z_{j_1, \dots, j_d}$'s are real and ξ Hermitian. By part (i), this entails that Y is a real IRF_k . Suppose now that $\tilde{Y}(\lambda) = aY(\lambda) + \mathfrak{i}bY'(\lambda)$ and observe that

$$\begin{aligned} \mathcal{C}_{\tilde{Y}}(\lambda, \mu) &= a^2 \mathbb{E}[Y(\lambda) \otimes Y(\mu)] + b^2 \mathbb{E}[Y'(\lambda) \otimes Y'(\mu)] + ab\mathfrak{i} \left[\mathbb{E}Y'(\lambda) \otimes Y(\mu) - \mathbb{E}Y(\lambda) \otimes Y'(\mu) \right] \\ &= \mathbb{E}[Y(\lambda) \otimes Y(\mu)] = \mathcal{C}_Y(\lambda, \mu), \end{aligned}$$

since $a^2 + b^2 = 1$ and the cross-covariance terms cancel. This shows that \tilde{Y} and Y have the same covariance structure. \square

Lemma A.8. *Let $T(x, y)$ be a bivariate mapping from $\mathbb{D} \times \mathbb{D}$ to \mathbb{F} , where both \mathbb{D} and \mathbb{F} are linear spaces over \mathbb{C} . Assume that T is sesquilinear form with $T(cx, y) = cT(x, y)$ and $T(x, cy) = \bar{c}T(x, y)$, $x, y \in \mathbb{D}$, $c \in \mathbb{C}$. Then, for $x, y \in \mathbb{D}$,*

$$T(x, y) = \frac{\mathfrak{i} - 1}{2}(T(x, x) + T(y, y)) + \frac{1}{2}T(x + y, x + y) - \frac{\mathfrak{i}}{2}T(\mathfrak{i}x + y, \mathfrak{i}x + y).$$

Proof. The proof is trivial by noticing the identities:

$$\begin{aligned} T(x + y, x + y) &= T(x, x) + T(y, y) + T(x, y) + T(y, x), \\ T(\mathfrak{i}x + y, \mathfrak{i}x + y) &= T(x, x) + T(y, y) + \mathfrak{i}T(x, y) - \mathfrak{i}T(y, x). \end{aligned}$$

\square

A.5. Proofs for Section 5.1.

We start with an auxiliary result needed for the proof of Theorem 5.2 below.

Lemma A.9. *For any $\mathcal{A} \in \mathbb{T}$ and any bounded linear operator \mathcal{H} , define $f(c) := c^{-\mathcal{H}} \mathcal{A} c^{-\mathcal{H}*}$, $c > 0$. The function $f : (0, \infty) \rightarrow \mathbb{T}$ is continuously Fréchet differentiable in $(\mathbb{T}, \|\cdot\|_{\text{tr}})$ with derivative*

$$f'(c) = -c^{-1}(\mathcal{H}f(c) + f(c)\mathcal{H}^*) = -c^{-1}c^{-\mathcal{H}}(\mathcal{H}\mathcal{A} + \mathcal{A}\mathcal{H}^*)c^{-\mathcal{H}*}, c > 0.$$

That is,

$$\|f(c + h) - f(c) - f'(c)h\|_{\text{tr}} = o(h), \quad \text{as } h \rightarrow 0. \quad (\text{A.17})$$

Proof. Observe that $(c + h)^{-\mathcal{H}} - c^{-\mathcal{H}} = \sum_{n=1}^{\infty} (-\mathcal{H})^n [(\log(c + h))^n - (\log c)^n] / n!$. Applying the mean value theorem to the terms $(\log(c + h))^n - (\log c)^n$, and using the triangle inequality for $\|\cdot\|_{\text{op}}$, one can show that for all $c > 0$,

$$\left\| (c + h)^{-\mathcal{H}} - c^{-\mathcal{H}} - c^{-\mathcal{H}} (-\mathcal{H}) c^{-1} h \right\|_{\text{op}} = o(h), \quad \text{as } h \rightarrow 0. \quad (\text{A.18})$$

That is, $c \mapsto c^{-\mathcal{H}}$ and similarly $c \mapsto c^{-\mathcal{H}*}$ are Fréchet differentiable in the Hilbert space \mathbb{V} .

On the other hand, by Proposition IV.5.4 on page 62 of [Gohberg et al. \[2000\]](#), for any two bounded operators \mathcal{B} and \mathcal{C} and a trace class operator $\mathcal{A} \in \mathbb{T}$, we have that $\mathcal{B}\mathcal{A}\mathcal{C} \in \mathbb{T}$ and moreover

$$\|\mathcal{B}\mathcal{A}\mathcal{C}\|_{\text{tr}} \leq \|\mathcal{B}\|_{\text{op}} \|\mathcal{A}\|_{\text{tr}} \|\mathcal{C}\|_{\text{op}}. \quad (\text{A.19})$$

This inequality can be used to show that the Fréchet differentiability of $c \mapsto c^{-\mathcal{H}}$ and $c \mapsto c^{-\mathcal{H}^*}$ in the operator norm induced by the Hilbert space norm in \mathbb{V} entails the Fréchet differentiability in trace-norm.

Indeed, for all $c > 0$ and $h > -c$, we have

$$\begin{aligned} f(c+h) - f(c) &= (c+h)^{-\mathcal{H}} \mathcal{A} (c+h)^{-\mathcal{H}^*} - c^{-\mathcal{H}} \mathcal{A} c^{-\mathcal{H}^*} \\ &= ((c+h)^{-\mathcal{H}} - c^{-\mathcal{H}}) \mathcal{A} (c+h)^{-\mathcal{H}^*} + c^{-\mathcal{H}} \mathcal{A} ((c+h)^{-\mathcal{H}^*} - c^{-\mathcal{H}^*}). \end{aligned}$$

Now, in view of (A.18) and using the inequality (A.19), we obtain

$$\|f(c+h) - f(c) - hc^{-\mathcal{H}-1}(-\mathcal{H})\mathcal{A}(c+h)^{-\mathcal{H}^*} - h(c+h)^{-\mathcal{H}}\mathcal{A}(-\mathcal{H}^*)c^{-\mathcal{H}^*-1}\|_{\text{tr}} = o(h),$$

as $h \rightarrow 0$. The continuity of $c \mapsto c^{-\mathcal{H}}$ and $c \mapsto c^{-\mathcal{H}^*}$ in the operator norm and another application of (A.19) entails (A.17). \square

Proof of Theorem 5.2: Fix $c > 0$ and define the rescaled IRF $_k$ process $\tilde{Y}(\lambda) := c^{-\mathcal{H}} Y(c \cdot \lambda)$, $\lambda \in \Lambda_k$. Observe that Y is covariance \mathcal{H} -self-similar if and only if \tilde{Y} and Y have the same covariance structure or, equivalently, the same spectral characteristics (χ, \mathcal{Q}) .

In view of (4.14), for all $\lambda, \mu \in \Lambda_k$, we have

$$\begin{aligned} \mathcal{C}_{\tilde{Y}}(\lambda, \mu) &= \mathbb{E}[\tilde{Y}(\lambda) \otimes \tilde{Y}(\mu)] = c^{-\mathcal{H}} \mathbb{E}[Y(c \cdot \lambda) \otimes Y(c \cdot \mu)] c^{-\mathcal{H}^*} \\ &= \int_{\mathbb{R}^d} \widehat{c \cdot \lambda}(u) \overline{\widehat{c \cdot \mu}(u)} c^{-\mathcal{H}} \chi_k(du) c^{-\mathcal{H}^*} + c^{-\mathcal{H}} \mathcal{Q}(c \cdot \lambda * \widetilde{c \cdot \mu}) c^{-\mathcal{H}^*} \\ &= \int_{\mathbb{R}^d} \widehat{\lambda}(x) \overline{\widehat{\mu}(x)} c^{-\mathcal{H}} \chi_k(dx/c) c^{-\mathcal{H}^*} + c^{-\mathcal{H}} c^{2k+2} \mathcal{Q}(\lambda * \widetilde{\mu}) c^{-\mathcal{H}^*}, \end{aligned} \quad (\text{A.20})$$

where in the last relation we used that $\widehat{c \cdot \lambda}(u) = \widehat{\lambda}(cu)$, the change of variables $u := x/c$, and the fact that $\mathcal{Q}(c \cdot \nu) = c^{2k+2} \mathcal{Q}(\nu)$ for all $\nu \in \Lambda_{2k+1}$.

Relation (A.20) shows that the spectral characteristics of \tilde{Y} are

$$(c^{-\mathcal{H}} \chi_k(dx/c) c^{-\mathcal{H}^*}, c^{k+1-\mathcal{H}} \mathcal{Q}(\cdot) c^{k+1-\mathcal{H}^*}).$$

Hence, by the uniqueness of the spectral representation in (4.14), the IRF $_k$ process Y and \tilde{Y} have the same covariance structure, if and only if Relation (5.6) holds. This completes the proof of part (i).

Part (iv) is an immediate consequence of part (ii) and Theorem 4.14, where the random measure $\xi(dx)/(1 \wedge |x|^{k+1})$ therein, is now written in polar coordinates as $W(dr d\theta)$. Thus, in the remainder of the proof we focus on establishing the *disintegration formula* (5.8) (part (ii)) and the uniqueness of the measure σ (part (iii)).

Define the \mathbb{T} -valued set-mapping

$$\sigma(B) := \mathcal{H}\chi_k((1, \infty) \times B) + \chi_k((1, \infty) \times B)\mathcal{H}^*, \quad B \in \mathcal{B}(\mathbb{S}), \quad (\text{A.21})$$

where in short, we write $\chi_k(A \times B)$ for $\chi_k(\{(r, \theta) \in A \times B\})$, with $A \subset (0, \infty)$ and $B \subset \mathbb{S}$. The fact that χ_k is a \mathbb{T}_+ -valued measure readily implies $\sigma(\emptyset) = 0$ and the countable additivity of σ . Note also that σ is finite, since for all $B \in \mathcal{B}(\mathbb{S})$, by (A.19),

$$\|\sigma(B)\|_{\text{tr}} \leq (\|\mathcal{H}\|_{\text{op}} + \|\mathcal{H}^*\|_{\text{op}})\|\chi_k((1, \infty) \times \mathbb{S})\|_{\text{tr}} < \infty.$$

We will argue next that $\sigma(B)$ is positive and hence it defines a \mathbb{T}_+ -valued measure on \mathbb{S} . We will also show that for all $c > 0$ and $B \in \mathcal{B}(\mathbb{S})$, we have

$$\chi_k((c, \infty) \times B) = \int_c^\infty u^{-\mathcal{H}} \sigma(B) u^{-\mathcal{H}^*} \frac{du}{u}. \quad (\text{A.22})$$

This fact and a standard $\pi - \lambda$ argument then entail that (5.7) holds. Indeed, the right-hand side of (5.7) defines a σ -finite \mathbb{T}_+ -valued measure, say $\tilde{\chi}_k$, on $\mathcal{B}(\mathbb{R}^d \setminus \{0\})$. By Relation (A.22), the measures χ_k and $\tilde{\chi}_k$ agree on the semi-ring of rectangle sets $(c, \infty) \times B$. Since the latter generates the σ -field $\mathcal{B}(\mathbb{R}^d \setminus \{0\})$, by considering projections on a fixed CONS, it can be seen that the two \mathbb{T}_+ -valued measures coincide.

We now prove that σ is a finite \mathbb{T}_+ -valued measure and show (A.22). In view of Lemma A.9 (above), it follows that the function

$$f(u) := u^{-\mathcal{H}} \chi_k((1, \infty) \times B) u^{-\mathcal{H}^*}.$$

is Fréchet continuously differentiable in $(\mathbb{T}, \|\cdot\|_{\text{tr}})$ with derivative:

$$\begin{aligned} f'(u) &= -u^{-1} u^{-\mathcal{H}} \left(\mathcal{H}\chi_k((1, \infty) \times B) + \chi_k((1, \infty) \times B)\mathcal{H}^* \right) u^{-\mathcal{H}^*} \\ &= -u^{-1} u^{-\mathcal{H}} \sigma(B) u^{-\mathcal{H}^*}, \quad u > 0. \end{aligned} \quad (\text{A.23})$$

Observe that by the operator-scaling property for χ_k in (5.6), we have

$$f(u) = \chi_k((u, \infty) \times B), \quad u > 0.$$

The monotonicity of the \mathbb{T}_+ -valued measure χ_k then implies that

$$\chi_k((u, u+h] \times B) = f(u) - f(u+h) \in \mathbb{T}_+,$$

for all $h > 0$. This shows that $-f'(u) \in \mathbb{T}_+$ and by setting $u = 1$, we obtain $-f'(1) = \sigma(B) \in \mathbb{T}_+$, which shows that $\sigma(B) \in \mathbb{T}_+$, completing the proof that the so-defined set-mapping in (A.21) is a finite \mathbb{T}_+ -valued measure.

Now, using (A.23) and a straightforward extension of the fundamental theorem of calculus to Bochner integrals, we obtain

$$\begin{aligned} \int_c^\infty u^{-\mathcal{H}} \sigma(B) u^{-\mathcal{H}^*} \frac{du}{u} &= - \int_c^\infty f'(u) du = f(c) \\ &= c^{-\mathcal{H}} \chi_k((1, \infty) \times B) c^{-\mathcal{H}^*}. \end{aligned}$$

(Note that $f(u) = \chi_k((u, \infty) \times B) \downarrow 0$ as $u \uparrow \infty$.) The latter, in view of the scaling property of χ_k , equals $\chi_k((c, \infty) \times B)$, completing the proof of (A.22) and part (ii).

Part (iii). We now show that σ is uniquely determined by χ_k , alone. By definition, we have

$$\sigma(B) = -\frac{d}{dc}\Big|_{c=1} \chi_k((c, \infty) \times B), \quad B \in \mathcal{B}(\mathbb{S}),$$

where the latter is interpreted as the Fréchet derivative in $(\mathbb{T}, \|\cdot\|)$ of the function $c \mapsto \chi_k((c, \infty) \times B)$, evaluated at $c = 1$. This shows that σ is uniquely determined in terms of the measure χ_k and it does not depend on the choice of the exponent operator \mathcal{H} , which need not be unique (see e.g., Remark 5.7). \square

A.6. Proofs for Section 5.2.

Proof of Proposition 5.6: Stationarity is immediate, provided that the stochastic integral in (5.30) is well-defined. To this end, it suffices to show that

$$\begin{aligned} \int_0^\infty \int_{\mathbb{S}} \|f_s(u, \theta)\|_{\text{op}}^2 du \|\mu\|_{\text{tr}}(d\theta) &:= \int_0^\infty \int_{\mathbb{S}} (1 \wedge u)^{2(k+1)} \|u^{-\mathcal{H}} \mathcal{A}(\theta)\|_{\text{op}}^2 u^{-1} du \|\mu\|_{\text{tr}}(d\theta) \\ &\leq \int_{\mathbb{S}} \|\mathcal{A}(\theta)\|_{\text{op}}^2 \|\mu\|_{\text{tr}}(d\theta) \int_0^\infty (1 \wedge u)^{2(k+1)} \|u^{-\mathcal{H}}\|_{\text{op}}^2 u^{-1} du. \end{aligned} \quad (\text{A.24})$$

By the assumption (5.29), the first integral in (A.24) is finite. It remains to show that the second one therein is also finite. This, however, readily follows from the inequality

$$(1 \wedge u)^{2(k+1)} u^{-1} \|u^{-\mathcal{H}}\|_{\text{op}}^2 \leq C \times \left(1_{(0,1)}(u) u^{2\delta-1} + 1_{[1,\infty)}(u) u^{-2\epsilon-1}\right), \quad (\text{A.25})$$

where $\epsilon > 0$ is as in (5.28) and $\delta > 0$ is such that $\Re(\text{sp}(\mathcal{H})) \subset (\epsilon, k+1-\delta)$. Relation (A.25) can be established exactly as in the proof of (5.19) using (5.28) and Riesz functional calculus. This completes the proof of (A.25) and part (i).

To prove part (ii), it suffices to establish that, for all $s, t \in B(0, M)$, $M > 0$,

$$\mathbb{E}[\|X(s) - X(t)\|^2] \leq \int_0^\infty \int_{\mathbb{S}} \|f_s(u, \theta) - f_t(u, \theta)\|_{\text{op}}^2 du \|\mu\|_{\text{tr}}(d\theta) \leq C \|s - t\|^{2\zeta}. \quad (\text{A.26})$$

Indeed, consider the Gaussian \mathbb{V} -valued variables $\xi_{s,t} := (X(s) - X(t))/\sigma_{s,t}$, where $\sigma_{s,t}^2 := \mathbb{E}[\|X(s) - X(t)\|^2]$, and where by convention $\xi_{s,t} := 0$ if $\sigma_{s,t} = 0$. By Corollary S.5.4 (with $\theta := 1/4$ therein) we have that

$$\mathbb{E}[\|\xi_{s,t}\|^p] \leq c_p \mathbb{E} \exp\{\|\xi_{s,t}\|^2/4\} \leq c_p \sqrt{2}$$

for all $p > 0$ and some finite universal constant c_p . The last bound and Relation (A.26) can be equivalently written as

$$\mathbb{E}[\|X(s) - X(t)\|^p] \leq C_p \sqrt{2} \|s - t\|^{p\zeta}.$$

This, in view of Proposition 2.4, implies the existence of a γ -Hölder continuous version of $\{X(s), s \in B(0, M)\}$ for all $\gamma \in (0, \zeta - d/p)$. Taking p large, we see that every $\gamma \in (0, \zeta)$ is a possible Hölder

exponent. We shall continue to denote this continuous-path version of the process by $\{X(s)\}$.

Now, we turn to proving (A.26). We have

$$\|f_s(u, \theta) - f_t(u, \theta)\|_{\text{op}}^2 \leq 2|e^{is^\top u\theta} - e^{it^\top u\theta}|^2 (1 \wedge u)^{2(k+1)} u^{-1} \|\mathcal{A}(\theta)\|_{\text{op}}^2 \|u^{-\mathcal{H}}\|_{\text{op}}^2 \quad (\text{A.27})$$

By Relations (5.29) and (A.25), we have

$$\begin{aligned} I_{s,t} &:= \int_0^\infty \int_{\mathbb{S}} \|f_s(u, \theta) - f_t(u, \theta)\|_{\text{op}}^2 du \|\mu\|_{\text{tr}}(d\theta) \\ &\leq C \int_0^\infty \sup_{\theta \in \mathbb{S}} |e^{i(s-t)^\top u\theta} - 1|^2 \left(u^{2(\delta)-1} 1_{[0,1]}(u) + u^{-2\epsilon-1} 1_{[1,\infty)}(u) \right) du. \end{aligned}$$

Thus, Lemma A.10 applied with $\Delta := (s-t)^\top \theta$, $\gamma := 2\delta > 0$ and $\epsilon := 2\epsilon$, yields

$$I_{s,t} \leq C \|s-t\|^{2(\epsilon \wedge 1)} (1 + |\log(\|s-t\|)|^2 1_{\{\epsilon=1\}}).$$

This implies that (A.26) holds with any $\zeta < 1 \wedge \epsilon$.

Part (iii). Consider the measures in (2.7) and observe that

$$\lambda_t = \delta_t - \sum_j c_j(t) \delta_{t_j}, \quad (\text{A.28})$$

where the $c_j(t)$'s are polynomials in t of degrees up to k and the t_j 's are some fixed points in \mathbb{R}^d . In view of Proposition 2.4, to prove (5.31) it is enough to work with

$$\begin{aligned} \check{X}_r(t) &:= r^{-\mathcal{H}} X(s_0 + r \cdot \lambda_t) \\ &= r^{-\mathcal{H}} \int_0^\infty \int_{\mathbb{S}} e^{is_0^\top u\theta} \left(e^{irs^\top u\theta} - \sum_j c_j(s) e^{irt_j^\top u\theta} \right) (1 \wedge u)^{k+1} u^{-\mathcal{H}-1/2} \mathcal{A}(\theta) W(du, d\theta) \\ &= \int_0^\infty \int_{\mathbb{S}} \widehat{\lambda}_t(ru\theta) (1 \wedge u)^{k+1} (ru)^{-\mathcal{H}-1/2} \mathcal{A}(\theta) \widetilde{W}(du, d\theta), \end{aligned}$$

where $\widetilde{W}(du, d\theta) := e^{ius_0^\top \theta} r^{1/2} W(du, d\theta)$. We will show that since the \mathbb{V} -valued Gaussian random measure W is circularly symmetric and self-similar (recall (5.27)), we have

$$\{\widetilde{W}(du, d\theta)\} \stackrel{fdd}{=} \{W(d(r \cdot u), d\theta)\}. \quad (\text{A.29})$$

Indeed, since W is standard Gaussian, it is not only orthogonal but also an independently scattered measure. This readily implies that \widetilde{W} is also independently scattered. It is straightforward that \widetilde{W} has control measure $rdu\mu(d\theta)$. It remains to show that \widetilde{W} is standard, i.e., its real and imaginary components are independent and identically distributed. Indeed, for all bounded Borel $A \subset \mathbb{R}^d \setminus \{0\}$, in view of (5.25), we have

$$\begin{aligned} \widetilde{W}(A) &= \int_A \cos(\varphi_{u,\theta}) r^{1/2} W_{\mathbb{R}}(du, d\theta) - \int_A \sin(\varphi_{u,\theta}) r^{1/2} W_{\mathbb{I}}(du, d\theta) \\ &\quad + \mathbf{i} \left(\int_A \sin(\varphi_{u,\theta}) r^{1/2} W_{\mathbb{R}}(du, d\theta) + \int_A \cos(\varphi_{u,\theta}) r^{1/2} W_{\mathbb{I}}(du, d\theta) \right) \\ &=: \Re(\widetilde{W}(A)) + \mathbf{i} \Im(\widetilde{W}(A)), \end{aligned}$$

where $\varphi_{u,\theta} := us_0^\top \theta$. Using the fact that $W_{\mathbb{R}}$ and $W_{\mathbb{I}}$ are independent and identically distributed real orthogonal measures, we obtain that $\Re(\widetilde{W}(A))$ and $\Im(\widetilde{W}(A))$ are independent and identically distributed. This, since \widetilde{W} is independently scattered and Gaussian, completes the proof of (A.29).

Therefore, by (A.29),

$$\begin{aligned} \{\check{X}_r(t), t \in \mathbb{R}^d\} &\stackrel{d}{=} \left\{ \int_0^\infty \int_{\mathbb{S}} \widehat{\lambda}_t(v\theta) (1 \wedge (v/r))^{k+1} v^{-\mathcal{H}-1/2} \mathcal{A}(\theta) W(dv, d\theta), t \in \mathbb{R}^d \right\} \\ &=: \{X_r(t), t \in \mathbb{R}^d\}, \end{aligned} \quad (\text{A.30})$$

where we made the change of variables $v := ru$.

This relation readily implies that the convergence in (5.31) holds in the sense of finite-dimensional distributions. Indeed, writing $\check{Y}(t) := \int_0^\infty \int_{\mathbb{S}} \widehat{\lambda}_t(v\theta) v^{-\mathcal{H}-1/2} \mathcal{A}(\theta) W(dv, d\theta)$, with the *same* measure defining the $X_r(t)$'s in (A.30), it is enough to show that

$$\begin{aligned} &\mathbb{E} \|X_r(t) - \check{Y}(t)\|^2 \\ &\leq \int_0^\infty \int_{\mathbb{S}} |\widehat{\lambda}_t(v\theta)|^2 \cdot |(1 \wedge (v/r))^{k+1} - 1|^2 \cdot \left\| v^{-\mathcal{H}-1/2} \mathcal{A}(\theta) \right\|_{\text{op}}^2 \|\mu\|_{\text{tr}}(d\theta) dv \rightarrow 0, \end{aligned} \quad (\text{A.31})$$

as $r \rightarrow 0$. It is easy to see that since $\lambda_t \in \Lambda_k$, we have $|\widehat{\lambda}_t(x)| \leq C(1 \wedge \|x\|)^{k+1}$, $x \in \mathbb{R}^d$. Therefore, in view of Relations (5.29) and (A.25), the fact that $|(1 \wedge (v/r))^{k+1} - 1| \rightarrow 0$, $r \rightarrow 0$ and the Dominated Convergence Theorem imply (A.31), proving the convergence of the finite-dimensional distributions. To complete the proof of (5.31), we will establish the tightness of $\{\check{X}_r(\cdot), r \in (0, 1)\}$. Since the \check{X}_r 's are Gaussian, as argued above, in view of Proposition 2.4 it is enough to show that for all $M > 0$, there exist $C > 0$ and $\zeta > 0$ such that

$$\sup_{r \in (0, 1)} \mathbb{E} \|\check{X}_r(s) - \check{X}_r(t)\|^2 \leq C \|s - t\|^{2\zeta}, \quad \text{for all } s, t \in B(0, M).$$

This is because for $k \geq 0$, the tightness condition on $\{\check{X}_r(s_0), r \in (0, 1)\}$ is automatically fulfilled if one takes $s_0 = t_1$ (recall (2.9) and Remark 2.2).

To this end, we begin with some key observations about the measures λ_t . Since the $c_j(t)$'s in (A.28) are fixed polynomials, we have

$$\begin{aligned} |\widehat{\lambda}_s(v\theta) - \widehat{\lambda}_t(v\theta)|^2 &\leq |e^{is^\top v\theta} - e^{it^\top v\theta}|^2 + \sum_j |c_j(s) - c_j(t)|^2 \\ &\leq Cv^2 \|s - t\|^2 + C_M \|s - t\|^2 \leq C_M (1 \vee v^2) \|s - t\|^2, \end{aligned} \quad (\text{A.32})$$

for all $v > 0$ and $s, t \in B(0, M)$, where the constant C_M does not depend on v, s, t and $\theta \in \mathbb{S}$.

On the other hand, in view of (A.28), the $\widehat{\lambda}_t$'s are uniformly bounded for $t \in B(0, M)$ and since $\lambda_t \in \Lambda_k$ they annihilate polynomials of degree up to k and hence $\widehat{\lambda}_t(v\theta) = \mathcal{O}(v^{k+1})$. This implies that

$$\sup_{t, s \in B(0, M), \theta \in \mathbb{S}} |\widehat{\lambda}_s(v\theta) - \widehat{\lambda}_t(v\theta)|^2 \leq C_M (1 \wedge v^{2(k+1)}). \quad (\text{A.33})$$

We are now ready to estimate the difference moments. For all $s, t \in B(0, M)$, we have

$$\begin{aligned} \mathbb{E} \|\check{X}_r(s) - \check{X}_r(t)\|^2 &\leq \int_0^\infty \int_{\mathbb{S}} |\widehat{\lambda}_s(v\theta) - \widehat{\lambda}_t(v\theta)|^2 \left\| v^{-\mathcal{H}-1/2} \mathcal{A}(\theta) \right\|_{\text{op}}^2 \|\mu\|_{\text{tr}}(d\theta) dv \\ &=: \int_0^\infty \int_{\mathbb{S}} g_{s,t}(v, \theta) \|\mu\|_{\text{tr}}(d\theta) dv. \end{aligned}$$

Observe first that by (A.33) the bound in (A.25) applies and by (5.29), we have

$$\int_{\mathbb{S}} g_{s,t}(v, \theta) \|\mu\|_{\text{tr}}(d\theta) \leq C \left(v^{2\delta-1} 1_{(0,1)}(v) + v^{-2\epsilon-1} 1_{[1,\infty)}(v) \right), \quad (\text{A.34})$$

where the latter function is integrable in v over $(0, \infty)$.

The rest of the strategy is as follows. We will consider the integral of $g_{s,t}$ over three regions $(v, \theta) \in (0, \alpha] \times \mathbb{S}$, $(\alpha, 1/\alpha) \times \mathbb{S}$ and $[1/\alpha, \infty) \times \mathbb{S}$. We will choose $\alpha = 1 \wedge \|s - t\|^\kappa$ for some $\kappa > 0$, such that each of the three integrals can be dominated by $C\alpha^\eta$, for some positive η . Namely, let

$$I_1(s, t) + I_2(s, t) + I_3(s, t) := \left(\int_{(0, \alpha] \times \mathbb{S}} + \int_{(\alpha, 1/\alpha) \times \mathbb{S}} + \int_{[1/\alpha, \infty) \times \mathbb{S}} \right) g_{s,t}(v, \theta) \|\mu\|_{\text{tr}}(d\theta) dv.$$

In view of (A.34), we have

$$I_1(s, t) \leq C \int_0^\alpha v^{2\delta-1} dv \leq C\alpha^{2\delta}$$

and

$$I_3(s, t) \leq C \int_{1/\alpha}^\infty v^{-2\epsilon-1} dv \leq C\alpha^{2\epsilon}.$$

Now, for the middle piece, using the bound in (A.32), we obtain

$$I_2(s, t) \leq C \|s - t\|^2 \int_\alpha^{1/\alpha} (1 \vee v^2) dv \leq C \|s - t\|^2 \alpha^{-3}.$$

Setting $\|s - t\|^2 \alpha^{-3} = \alpha^{2(\epsilon \wedge \delta)}$, we see that $\alpha = 1 \wedge \|s - t\|^{2/(3+2(\epsilon \wedge \delta))}$ yields the desired bound $\mathbb{E} \|\check{X}_r(s) - \check{X}_r(t)\|^2 \leq \sum_{i=1}^3 I_i(s, t) \leq C \|s - t\|^{2\zeta}$, uniformly in $r \in (0, 1)$, where $\zeta = 2(\epsilon \wedge \delta)/(3 + 2(\epsilon \wedge \delta)) > 0$. \square

Lemma A.10. (i) *There exists a constant C such that*

$$\int_1^\infty |e^{i\Delta u} - 1|^2 u^{-1-\epsilon} du \leq C \begin{cases} |\Delta|^2 & , \text{ if } \epsilon > 2 \\ |\Delta|^2 |\log |\Delta|| & , \text{ if } \epsilon = 2 \\ |\Delta|^\epsilon & , \text{ if } 0 < \epsilon < 2. \end{cases}$$

(ii) *Also, for any $\gamma > 0$, there is a constant $C_\gamma > 0$, such that $\int_0^1 |e^{i\Delta u} - 1|^2 u^{\gamma-1} du \leq C_\gamma |\Delta|^2$.*

Proof. Noting that $|e^{i\Delta u} - 1|^2 = 2(1 - \cos(\Delta u)) = \mathcal{O}(\Delta^2)$, as $\Delta \rightarrow 0$, the claim in part (ii) is immediate. Part (i) follows by straightforward calculus by considering the change of variables $x := |\Delta|u$. \square

A.7. Proofs for Section 5.3.

Proof of Proposition 5.7: Theorem 5.2 applied to $\mathcal{H} := H \cdot \mathbf{I}$ implies the scaling property of χ_k and the disintegration formula (5.32) in part (ii). Part (iii) follows from Proposition 5.5. We need only prove that $0 < H \leq k + 1$ and the dichotomy claim (i).

Recall the decomposition in (5.4) and observe that both components $Y_{(\chi,0)}$ and $Y_{(0,\mathcal{Q})}$ are co-variance H -self-similar. Notice, however, that $Y_{(0,\mathcal{Q})}$ is always either $(k+1)$ -self-similar or zero. In particular, for all $c > 0$,

$$c^H Y_{(0,\mathcal{Q})}(\lambda) \stackrel{d}{=} Y_{(0,\mathcal{Q})}(c \cdot \lambda) \stackrel{d}{=} c^{k+1} Y_{(0,\mathcal{Q})}(\lambda), \quad \lambda \in \Lambda_k$$

This, since H is scalar, implies $H = k+1$ unless $Y_{(0,\mathcal{Q})} \equiv 0$ (recall Lemma A.5).

Thus, the polynomial spectral component \mathcal{Q} is non-trivial only if $H = k+1$. To complete the proof of (i), it remains to show that if $Y_{(\chi,0)}$ is non-zero, then its self-similarity exponent is in the range $0 < H < k+1$. Firstly, note that $Y_{(\chi,0)}(c \cdot \lambda) \stackrel{d}{=} c^H Y_{(\chi,0)}(\lambda) \rightarrow 0$, in probability, as $c \downarrow 0$, by the continuity of Y . This implies that $H > 0$.

Now, observe that by the scaling property of χ_k , we have, for all $r \in (0, 1)$,

$$\begin{aligned} r^{-2H} \chi_k((1, \infty) \times \mathbb{S}) &= \chi_k((r, \infty) \times \mathbb{S}) = \int_{\{r < \|u\|\}} \frac{1}{1 \wedge \|u\|^{2k+2}} \chi(du) \\ &= \int_{\{r < \|u\| \leq 1\}} \frac{1}{\|u\|^{2k+2}} \chi(du) + \chi(\{\|u\| > 1\}) \\ &= r^{-(2k+2)} \int_{\{r < \|u\| \leq 1\}} \left(\frac{r}{\|u\|}\right)^{2k+2} \chi(du) + \chi(\{\|u\| > 1\}). \end{aligned}$$

Since χ is a finite \mathbb{T}_+ -valued measure, however, the last integral vanishes as $r \downarrow 0$, by the Dominated Convergence Theorem. Thus, by multiplying the last expression by r^{2k+2} , we obtain

$$r^{2(k+1-H)} \chi_k((1, \infty) \times \mathbb{S}) \rightarrow 0,$$

as $r \rightarrow 0$, which means that $H < k+1$, since by the scaling property $\chi_k((r, \infty) \times \mathbb{S}) \neq 0$, for any (all) $r > 0$. This completes the proof. \square

Proof of Theorem 5.8: The proof of *Part (i)* is as follows. Since $H < k+1$, the polynomial \mathcal{Q} in the spectral characteristic of \mathcal{K} is zero and χ satisfies (5.32) in polar coordinates. Let $\nu \in \Lambda_{2k+1}$. Using that $\widehat{\nu}(r\theta) = \int_{\mathbb{R}^d} e^{ir\theta^\top t} \nu(dt)$, Relation (4.13) in polar coordinates becomes

$$\begin{aligned} \mathcal{K}(\nu) &= \int_{\mathbb{S}^{d-1}} \int_0^\infty \int_{\mathbb{R}^d} e^{ir\theta^\top t} \nu(dt) r^{-(2H+1)} dr \sigma(d\theta) \\ &= \int_{\mathbb{S}^{d-1}} \int_0^\infty \int_{\mathbb{R}^d} \left(e^{ir\theta^\top t} - \sum_{j=0}^{[2H]} \frac{(\mathbf{i}r\theta^\top t)^j}{j!} \right) \nu(dt) r^{-(2H+1)} dr \sigma(d\theta) \\ &= \int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}^d} \int_0^\infty \left(e^{ir\theta^\top t} - \sum_{j=0}^{[2H]} \frac{(\mathbf{i}r\theta^\top t)^j}{j!} \right) r^{-(2H+1)} dr \nu(dt) \sigma(d\theta) \end{aligned} \tag{A.35}$$

where the second relation follows from the fact that $\nu \in \Lambda_{2k+1}$ and therefore we could add a polynomial in t of degree $[2H] \leq 2k+1$ without changing the integral. We will justify next the interchange of the inner two integrals leading to (A.35) and compute the inner integral, therein.

For all integer $m \geq 0$, one can show that

$$\left| e^{\mathbf{i}z} - \sum_{j=0}^m \frac{(\mathbf{i}z)^j}{j!} \right| \leq \min \left\{ \frac{|z|^{m+1}}{(m+1)!}, \frac{2|z|^m}{m!} \right\}, \quad z \in \mathbb{R} \tag{A.36}$$

[see e.g., page 298 in [Resnick, 1999](#)]. Thus, for $0 \leq \lfloor 2H \rfloor < 2H + 1 < \lfloor 2H \rfloor + 1$ (recall $2H$ is not integer), we obtain

$$\int_0^\infty \left| e^{\mathrm{i}r\theta^\top t} - \sum_{j=0}^{\lfloor 2H \rfloor} \frac{(\mathrm{i}r\theta^\top t)^j}{j!} \right| r^{-(2H+1)} dr < \infty.$$

This allows us to interchange the order of the two inner integrals in (A.35). Doing so and making the change of variables $z := r|\theta^\top t|$, we obtain

$$\begin{aligned} & \int_0^\infty \left(e^{\mathrm{i}r\theta^\top t} - \sum_{j=0}^{\lfloor 2H \rfloor} \frac{(\mathrm{i}r\theta^\top t)^j}{j!} \right) r^{-(2H+1)} dr \\ &= |\theta^\top t|^{2H} \int_0^\infty \left(e^{\mathrm{i}z \operatorname{sign}(\theta^\top t)} - \sum_{j=0}^{\lfloor 2H \rfloor} \frac{(\mathrm{i}z \operatorname{sign}(\theta^\top t))^j}{j!} \right) z^{-(2H+1)} dz \\ &= |\theta^\top t|^{2H} \int_0^\infty \left(\cos(z) - \sum_{0 \leq 2j \leq \lfloor 2H \rfloor, j \in \mathbb{N}} \frac{(-1)^j z^{2j}}{2j!} \right) z^{-(2H+1)} dz \\ &\quad + \mathrm{i} \operatorname{sign}(\theta^\top t) |\theta^\top t|^{2H} \int_0^\infty \left(\sin(z) - \sum_{1 \leq 2j+1 \leq \lfloor 2H \rfloor, j \in \mathbb{N}} \frac{(-1)^j z^{2j+1}}{(2j+1)!} \right) z^{-(2H+1)} dz \\ &= |\theta^\top t|^{2H} I(H) + \mathrm{i}(\theta^\top t)^{<2H>} J(H), \end{aligned}$$

where $I(H)$ and $J(H)$ are in (5.34). Note that this argument also demonstrates that $I(H)$ and $J(H)$ are well-defined.

By substituting the last expression in the right-hand side of (A.35), we obtain

$$\begin{aligned} \mathcal{K}(\nu) &= I(H) \int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}^d} |\theta^\top t|^{2H} \nu(dt) \sigma(d\theta) \\ &\quad + \mathrm{i} J(H) \int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}^d} (\theta^\top t)^{<2H>} \nu(dt) \sigma(d\theta) \\ &= I(H) \int_{\mathbb{S}^{d-1}} |(\theta, \cdot)|^{2H}(\nu) \sigma(dt) + \mathrm{i} J(H) \int_{\mathbb{S}^{d-1}} (\theta, \cdot)^{<2H>}(\nu) \sigma(dt). \end{aligned}$$

This completes the proof of (5.33).

Part (ii): Suppose now that $2H \in \mathbb{Z}$, where $1 \leq 2H \leq 2k+1$ and observe that $2H$ may be either even or odd. Using the fact that $\nu \in \Lambda_{2k+1}$, we get

$$\begin{aligned} \widehat{\nu}(r\theta) &= \int_{\mathbb{R}^d} \left(e^{\mathrm{i}r\theta^\top t} - \sum_{j=0}^{2H-1} \frac{(\mathrm{i}r\theta^\top t)^j}{j!} - 1_{[0,1]}(r) \frac{(\mathrm{i}r\theta^\top t)^{2H}}{(2H)!} \right) \nu(dt) \\ &=: \int_{\mathbb{R}^d} f(r\theta^\top t, r) \nu(dt). \end{aligned}$$

Relation (A.36) implies that $|f(r\theta^\top t, r)| = \mathcal{O}(|r|^{2H+1})$ as $r \rightarrow 0$ and because of the presence of the indicator function $1_{[0,1]}(r)$, we have $|f(r\theta^\top t, r)| = \mathcal{O}(|r|^{2H-1})$ as $r \rightarrow \infty$. Therefore,

$$\int_0^\infty |f(r\theta^\top t, r)| r^{-(2H+1)} dr < \infty,$$

and by Fubini, we obtain

$$\begin{aligned}
& \int_0^\infty \widehat{\nu}(r\theta) r^{-(2H+1)} dr = \int_{\mathbb{R}^d} \int_0^\infty f(r\theta^\top t, r) r^{-(2H+1)} dr \nu(dt) \\
&= \int_{\mathbb{R}^d} \left[|\theta^\top t|^{2H} \int_0^\infty \left(e^{\mathbf{i}z\kappa} - \sum_{j=0}^{2H-1} \frac{(\mathbf{i}z\kappa)^j}{j!} - 1_{[0,1]}(z/|\theta^\top t|) \frac{(\mathbf{i}z\kappa)^{2H}}{(2H)!} \right) z^{-(2H+1)} dz \right] \nu(dt) \\
&=: \int_{\mathbb{R}^d} I_H(t) \nu(dt)
\end{aligned} \tag{A.37}$$

where we made the change of variables $z := r|\theta^\top t|$ and where $\kappa := \text{sign}(\theta^\top t)$.

We now consider two cases. Suppose first that $2H$ is even. Upon separating the real and imaginary parts the integral in the r.h.s. of (A.37) becomes:

$$\begin{aligned}
I_H(t) &= |\theta^\top t|^{2H} \int_0^\infty \left(\cos(z) - \sum_{0 \leq 2j \leq 2H-1, j \in \mathbb{Z}} \frac{(-1)^j z^{2j}}{(2j)!} - 1_{[0,1]}(z/|\theta^\top t|) \frac{(-1)^H z^{2H}}{(2H)!} \right) z^{-(2H+1)} dz \\
&\quad + \mathbf{i} \kappa |\theta^\top t|^{2H} \int_0^\infty \left(\sin(z) - \sum_{1 \leq 2j+1 \leq 2H-1, j \in \mathbb{Z}} \frac{(-1)^j z^{2j+1}}{(2j+1)!} \right) z^{-(2H+1)} dz \\
&=: (\theta^\top t)^{2H} I_{H,\cos}(t) + \mathbf{i} (\theta^\top t)^{<2H>} J(H).
\end{aligned} \tag{A.38}$$

Observe that the integral in the imaginary part above equals $J(H)$ in (5.34) and where in the real part we dropped the absolute value around $\theta^\top t$ since $2H$ is even.

Since $(\theta^\top t)^{2H}$ is a polynomial in t of degree $2H < 2k+1$, we have $\int_{\mathbb{R}^d} (\theta^\top t)^{2H} \nu(dt) = 0$. Therefore,

$$\int_{\mathbb{R}^d} (\theta^\top t)^{2H} I_{H,\cos}(t) \nu(dt) = \int_{\mathbb{R}^d} (\theta^\top t)^{2H} (I_{H,\cos}(t) - \mathcal{C}_{H,\cos}) \nu(dt), \tag{A.39}$$

where

$$\mathcal{C}_{H,\cos} := \int_0^\infty \left(\cos(z) - \sum_{0 \leq 2j \leq 2H-1, j \in \mathbb{Z}} \frac{(-1)^j z^{2j}}{(2j)!} - 1_{[0,1]}(z) \frac{(-1)^H z^{2H}}{(2H)!} \right) z^{-(2H+1)} dz,$$

which is almost the same as the first integral in (A.38) except that the indicator function $1_{[0,1]}(z)$ no longer depends on t . Notice again that $\mathcal{C}_{H,\cos}$ is well-defined. We thus obtain

$$\begin{aligned}
I_{H,\cos}(t) - \mathcal{C}_{H,\cos} &= \int_0^\infty \left(1_{[0,1]}(z) - 1_{[0,|\langle \theta, t \rangle|]}(z) \right) \frac{(-1)^H z^{2H}}{(2H)!} z^{-(2H+1)} dz \\
&= \frac{(-1)^H}{(2H)!} \int_{|\theta^\top t|}^1 z^{-1} dz = \frac{(-1)^{H+1}}{(2H)!} \log(|\theta^\top t|).
\end{aligned} \tag{A.40}$$

In view of (A.38), (A.39), and (A.40), we get

$$\begin{aligned}
\mathcal{K}(\nu) &= \int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}^d} I_H(t) \nu(dt) \sigma(d\theta) \\
&= \frac{(-1)^{H+1}}{(2H)!} \int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}^d} (\theta^\top t)^{2H} \log |\theta^\top t| \nu(dt) \sigma(d\theta) \\
&\quad + \mathbf{i} J(H) \int_{\mathbb{S}^{d-1}} \int_{\mathbb{R}^d} (\theta^\top t)^{<2H>} \nu(dt) \sigma(d\theta) \\
&= \frac{(-1)^{H+1}}{(2H)!} \int_{\mathbb{S}^{d-1}} \left[(\theta, \cdot)^{2H} \log |(\theta, \cdot)| \right] (\nu) \sigma(d\theta) + \mathbf{i} J(H) \int_{\mathbb{S}^{d-1}} (\theta, \cdot)^{<2H>} (\nu) \sigma(d\theta),
\end{aligned}$$

which completes the expression in the case when $2H$ is an even integer.

Suppose now that $2H$ is odd. With a very similar argument, using the fact that $\nu \in \Lambda_{2k+1}$ annihilates polynomials of degree $2k+1 \geq 2H$, we obtain that the right-hand side of (A.37) equals

$$\begin{aligned} & \int_{\mathbb{R}^d} |\theta^\top t|^{2H} I(H) \nu(dt) + \mathfrak{i} \int_{\mathbb{R}^d} \kappa |\theta^\top t|^{2H} \int_0^\infty \left(\sin(z) - \sum_{1 \leq 2j+1 \leq 2H-1} \frac{(-1)^j z^{2j+1}}{(2j+1)!} \right. \\ & \quad \left. - 1_{[0,1]}(z/|\theta^\top t|) \frac{(-1)^{H-1/2} z^{2H}}{(2H)!} \right) z^{-(2H+1)} dz \nu(dt) \\ &= I(H) |(\theta, \cdot)|^{2H}(\nu) + \mathfrak{i} \int_0^\infty \left[\int_{\mathbb{R}^d} (\theta^\top t)^{2H} (1_{[0,1]}(z) - 1_{[0,1/|\theta^\top t|]}(z)) \frac{(-1)^{H-1/2} z^{-1}}{(2H)!} \nu(dt) \right] dz, \end{aligned} \quad (\text{A.41})$$

where we used the fact that $\kappa |\theta^\top t|^{2H} = (\theta^\top t)^{2H}$ and the last relation is obtained with the same strategy as in (A.39). More precisely, applying Fubini and using that $\int_{\mathbb{R}^d} (\theta^\top t)^{2H} \nu(dt) = 0$, allows us to eliminate the terms involving $\sin(z)$ and z^{2j+1} . At the same time, we add the term $(\theta^\top t)^{2H} 1_{[0,1]}(z) z^{-1}$, which is a polynomial in t also annihilated by ν .

Now, since the inner integrand in the r.h.s. of (A.41) is integrable with respect to z , another application of Fubini shows that right-hand side of (A.37) equals

$$\begin{aligned} & I(H) |(\theta, \cdot)|^{2H}(\nu) + \frac{\mathfrak{i}(-1)^{H-1/2}}{(2H)!} \int_{\mathbb{R}^d} (\theta^\top t)^{2H} \left[\int_{1/|\theta^\top t|}^1 z^{-1} dz \right] \nu(dt) \\ &= I(H) |(\theta, \cdot)|^{2H}(\nu) + \frac{\mathfrak{i}(-1)^{H+1/2}}{(2H)!} \int_{\mathbb{R}^d} (\theta^\top t)^{2H} \log(|\theta^\top t|) \nu(dt). \end{aligned}$$

This leads to the desired expression

$$\mathcal{K}(\nu) = I(H) \int_{\mathbb{S}^{d-1}} |(\theta, \cdot)|^{2H}(\nu) \sigma(d\theta) + \mathfrak{i} \frac{(-1)^{H+1/2}}{(2H)!} \int_{\mathbb{S}^{d-1}} \left[|(\theta, \cdot)|^{2H} \log |(\theta, \cdot)| \right] (\nu) \sigma(d\theta),$$

completing the proof. \square

Acknowledgements.

SS and TH were partially supported by the NSF Grant DMS-1916226 *The Argo Data and Functional Spatial Processes*. We dedicate this paper to the memory of Mark Marvin Meerschaert (1955-2020). Mark was a great visionary, mentor, and friend to us. His work has inspired and guided us in this paper and elsewhere. We are very grateful to Rafail Kartsioukas for his exceptionally careful reading of the manuscript and help with fixing a number of important mathematical errors. We thank Yimin Xiao for an inspiring discussion and pointing out important references to the available literature. We are also very indebted to two anonymous referees for their exceptionally detailed and insightful comments, which helped us correct a number of errors and improve the manuscript.

References

P. Abry and G. Didier. Wavelet estimation for operator fractional Brownian motion. *Bernoulli*, 24(2):895–928, 2018.

- P. Abry, H. Wendt, S. Jaffard, and G. Didier. Multivariate scale-free temporal dynamics: From spectral (Fourier) to fractal (wavelet) analysis. *Comptes Rendus Physique*, 20(5):489 – 501, 2019.
- P. Amblard and J. Coeurjolly. Identification of the multivariate fractional Brownian motion. *IEEE Transactions on Signal Processing*, 59(11):5152–5168, 2011. .
- C. Baek, G. Didier, and V. Pipiras. On integral representations of operator fractional Brownian fields. *Statist. Probab. Lett.*, 92:190–198, 2014. ISSN 0167-7152. . URL <https://doi.org/10.1016/j.spl.2014.05.015>.
- G. Berschneider. Spectral representation of intrinsically stationary fields. *Stochastic Process. Appl.*, 122(12):3837–3851, 2012. ISSN 0304-4149. . URL <https://doi.org/10.1016/j.spa.2012.07.005>.
- G. Berschneider and Z. Sasvári. Spectral theory of stationary random fields and their generalizations. A short historical survey. In *Indefinite Inner Product Spaces, Schur analysis, and Differential Equations*, volume 263 of *Oper. Theory Adv. Appl.*, pages 217–235. Birkhäuser/Springer, Cham, 2018.
- H. Biermé and C. Lacaux. Fast and exact synthesis of some operator scaling Gaussian random fields. *Appl. Comput. Harmon. Anal.*, 48(1):293–320, 2020. ISSN 1063-5203. . URL <https://doi.org/10.1016/j.acha.2018.05.004>.
- H. Biermé, M. M. Meerschaert, and H.-P. Scheffler. Operator scaling stable random fields. *Stochastic Process. Appl.*, 117(3):312–332, 2007. ISSN 0304-4149. . URL <https://doi.org/10.1016/j.spa.2006.07.004>.
- P. Billingsley. *Convergence of Probability Measures*. Wiley Series in Probability and Statistics: Probability and Statistics. John Wiley & Sons, Inc., New York, second edition, 1999. ISBN 0-471-19745-9. . URL <https://doi.org/10.1002/9780470316962>. A Wiley-Interscience Publication.
- S. Bochner. *Vorlesungen über Fouriersche integrale*. Chelsea Publishing Company, 1948.
- J.-P. Chilès and P. Delfiner. *Geostatistics*. Wiley Series in Probability and Statistics. John Wiley & Sons, Inc., Hoboken, NJ, second edition, 2012. ISBN 978-0-470-18315-1. URL <https://doi.org/10.1002/9781118136188>. Modeling spatial uncertainty.
- J. B. Conway. *A Course in Functional Analysis*. Springer, 2007.
- H. Cramér. On harmonic analysis in certain functional spaces. *Ark. Mat. Astr. Fys.*, 28B(12):7, 1942. ISSN 0004-2080.
- Y. Davydov and V. Paulauskas. Lamperti type theorems for random fields, 2017. arXiv:1705.00182.
- G. Didier and V. Pipiras. Integral representations and properties of operator fractional Brownian motions. *Bernoulli*, 17(1):1–33, 2011. ISSN 1350-7265. . URL <https://doi.org/10.3150/10-BEJ259>.
- G. Didier, M. M. Meerschaert, and V. Pipiras. Exponents of operator self-similar random fields. *J. Math. Anal. Appl.*, 448(2):1450–1466, 2017. ISSN 0022-247X. . URL <https://doi.org/10.1016/j.jmaa.2016.11.055>.
- G. Didier, M. M. Meerschaert, and V. Pipiras. Domain and range symmetries of operator fractional Brownian fields. *Stochastic Process. Appl.*, 128(1):39–78, 2018. ISSN 0304-4149. . URL <https://doi.org/10.1016/j.spa.2017.04.003>.
- J. Diestel and J. Uhl. *Vector Measures*. Mathematical Surveys and Monographs. American Mathematical Society, 1977. ISBN 9780821873748. URL <https://books.google.com/books?id=EQFjD90fXWAC>.
- R. L. Dobrushin. Gaussian and their subordinated self-similar random generalized fields. *The Annals of Probability*, 7:1–28, 1979.
- R. M. Dudley. *Real Analysis and Probability*, volume 74 of *Cambridge Studies in Advanced*

- Mathematics*. Cambridge University Press, Cambridge, 2002. ISBN 0-521-00754-2. . URL <https://doi.org/10.1017/CB09780511755347>. Revised reprint of the 1989 original.
- J. Dugundji. *Topology*. Allyn and Bacon, Inc., Boston, Mass., 1966.
- M.-C. Düker. Limit theorems in the context of multivariate long-range dependence. *Stochastic Process. Appl.*, 130(9):5394–5425, 2020. ISSN 0304-4149. . URL <https://doi.org/10.1016/j.spa.2020.03.011>.
- A. Durand and F. Roueff. Spectral analysis of weakly stationary processes valued in a separable Hilbert space. 2020. arXiv: 1910.08491.
- K. J. Falconer. *The geometry of fractal sets*, volume 85 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1986. ISBN 0-521-25694-1; 0-521-33705-4.
- K. J. Falconer. Tangent fields and the local structure of random fields. *J. Theoret. Probab.*, 15(3): 731–750, 2002. ISSN 0894-9840. . URL <https://doi.org/10.1023/A:1016276016983>.
- K. J. Falconer. The local structure of random processes. *J. London Math. Soc. (2)*, 67(3):657–672, 2003. ISSN 0024-6107. . URL <https://doi.org/10.1112/S0024610703004186>.
- I. M. Gel’fand and N. Y. Vilenkin. *Generalized Functions. Vol. 4: Applications of harmonic analysis*. Translated by Amiel Feinstein. Academic Press, New York - London, 1964, 1964.
- B. Gnedenko. Sur la distribution limite du terme maximum d’une série aléatoire. *Ann. of Math. (2)*, 44:423–453, 1943. ISSN 0003-486X.
- I. Gohberg, S. Goldberg, and N. Krupnik. *Traces and Determinants of Linear Operators*, volume 116 of *Operator Theory: Advances and Applications*. Birkhäuser Verlag, Basel, 2000. ISBN 3-7643-6177-8. . URL <https://doi.org/10.1007/978-3-0348-8401-3>.
- R. B. Holmes. Mathematical foundations of signal processing. *SIAM Review*, 21(3):361–388, 1979.
- L. Horváth and P. Kokoszka. *Inference for functional data with applications*. Springer Series in Statistics. Springer, New York, 2012. ISBN 978-1-4614-3654-6. . URL <https://doi.org/10.1007/978-1-4614-3655-3>.
- T. Hsing and R. Eubank. *Theoretical Foundations of Functional Data Analysis, with an Introduction to Linear Operators*. Wiley Series in Probability and Statistics. John Wiley & Sons, Ltd., Chichester, 2015. ISBN 978-0-470-01691-6. . URL <https://doi.org/10.1002/9781118762547>.
- H. Hult and F. Lindskog. Regular variation for measures on metric spaces. *Publ. Inst. Math. (Beograd) (N.S.)*, 80(94):121–140, 2006. ISSN 0350-1302. . URL <http://dx.doi.org/10.2298/PIM0694121H>.
- Z. J. Jurek. Polar coordinates in Banach spaces. *Bull. Polish Acad. Sci. Math.*, 32(1-2):61–66, 1984. ISSN 0239-7269.
- O. Kallenberg. *Foundations of Modern Probability*. Probability and its Applications (New York). Springer-Verlag, 1997.
- G. Kallianpur. Zero-one laws for Gaussian processes. *Transactions of the American Mathematical Society*, 149:199–211, 1970.
- G. Kallianpur and V. Mandrekar. Spectral theory of stationary H-valued processes. *Journal of Multivariate Analysis*, 1(1):1–16, 1971.
- S. Kechagias and V. Pipiras. Definitions and representations of multivariate long-range dependent time series. *J. Time Series Anal.*, 36(1):1–25, 2015. ISSN 0143-9782. . URL <https://doi.org/10.1111/jtsa.12086>.
- A. Khintchine. Korrelationstheorie der stationären stochastischen prozesse. *Mathematische Annalen*, 109(1):604–615, 1934.
- M. Kuczma. *An introduction to the Theory of Functional Equations and Inequalities*. Birkhäuser Verlag, Basel, second edition, 2009. ISBN 978-3-7643-8748-8. . URL

- <https://doi.org/10.1007/978-3-7643-8749-5>. Cauchy’s equation and Jensen’s inequality, Edited and with a preface by Attila Gilányi.
- K. Kuratowski. *Introduction to Set Theory and Topology*. PWN—Polish Scientific Publishers, Warsaw; Pergamon Press, Oxford-New York-Toronto, Ont., 1977. Containing a supplement, “Elements of algebraic topology” by Ryszard Engelking, Translated from the Polish by Leo F. Boron, International Series of Monographs in Pure and Applied Mathematics, 101.
- S. Kwapień and N. A. Woyczyński. *Random Series and Stochastic Integrals: Single and Multiple*. Birkhäuser, Boston, 1992.
- J. Lamperti. Semi-stable stochastic processes. *Trans. Amer. Math. Soc.*, 104:62–78, 1962. ISSN 0002-9947. . URL <https://doi.org/10.2307/1993933>.
- M. Ledoux and M. Talagrand. *Probability in Banach Spaces: Isoperimetry and Processes*. Springer-Verlag, New York, 1991.
- Y. Li and Y. Xiao. Multivariate operator-self-similar random fields. *Stochastic Process. Appl.*, 121(6):1178–1200, 2011. ISSN 0304-4149. . URL <https://doi.org/10.1016/j.spa.2011.02.005>.
- P. A. Loeb and E. Talvila. Lusin’s theorem and Bochner integration. *Sci. Math. Jpn.*, 60(1): 113–120, 2004. ISSN 1346-0862.
- J. D. Mason and Y. Xiao. Sample path properties of operator-self-similar Gaussian random fields. *Teor. Veroyatnost. i Primenen.*, 46(1):94–116, 2001. ISSN 0040-361X. . URL <https://doi.org/10.1137/S0040585X97978749>.
- G. Matheron. The intrinsic random functions and their applications. *Advances in Appl. Probability*, 5:439–468, 1973. ISSN 0001-8678. . URL <https://doi.org/10.2307/1425829>.
- M. M. Meerschaert. *Multivariate Domains of Attraction and Regular Variation*. PhD thesis, University of Michigan, Ann Arbor, 1984.
- M. M. Meerschaert and H.-P. Scheffler. *Limit distributions for sums of independent random vectors*. Wiley Series in Probability and Statistics: Probability and Statistics. John Wiley & Sons Inc., New York, 2001. ISBN 0-471-35629-8. Heavy tails in theory and practice.
- R. Melrose. *Functional Analysis*. MIT, <http://math.mit.edu/~rbm/18.102-S13/FunctAnal.pdf>, 2013. Lecture Notes for 18.102.
- J. R. Munkres. *Topology*. Prentice Hall, Inc., Upper Saddle River, NJ, 2000. ISBN 0-13-181629-2. Second edition of [MR0464128].
- G. J. Murphy. Positive definite kernels and Hilbert C^* -modules. *Proc. Edinburgh Math. Soc.* (2), 40(2):367–374, 1997. ISSN 0013-0915. . URL <https://doi.org/10.1017/S0013091500023804>.
- K.-H. Neeb. Operator-valued positive definite kernels on tubes. *Monatsh. Math.*, 126(2):125–160, 1998. ISSN 0026-9255. . URL <https://doi.org/10.1007/BF01473583>.
- J.-P. Pellonpää and K. Ylinen. Modules, completely positive maps, and a generalized KSGNS construction. *Positivity*, 15(3):509–525, 2011. ISSN 1385-1292. . URL <https://doi.org/10.1007/s11117-010-0104-6>.
- E. Perrin, R. Harba, C. Berzin-Joseph, I. Iribarren, and A. Bonami. nth-order fractional Brownian motion and fractional Gaussian noises. *IEEE Transactions on Signal Processing*, 49(5):1049–1059, 2001.
- D. Preiss. Geometry of measures in \mathbf{R}^n : distribution, rectifiability, and densities. *Ann. of Math.* (2), 125(3):537–643, 1987. ISSN 0003-486X. . URL <https://doi.org/10.2307/1971410>.
- J. O. Ramsay and B. W. Silverman. *Functional data analysis*. Springer Series in Statistics. Springer, New York, second edition, 2005. ISBN 978-0387-40080-8; 0-387-40080-X.
- S. I. Resnick. *A Probability Path*. Birkhäuser Boston Inc., Boston, MA, 1999. ISBN 0-8176-4055-X.
- S. Rolewicz. *Metric Linear Spaces*, volume 20 of *Mathematics and its Applications (East European*

- Series*). D. Reidel Publishing Co., Dordrecht; PWN—Polish Scientific Publishers, Warsaw, second edition, 1985. ISBN 90-277-1480-0.
- Z. Sasvári. Correlation functions of intrinsically stationary random fields. In *Modern Analysis and Applications. The Mark Krein Centenary Conference. Vol. 1: Operator theory and related topics*, volume 190 of *Oper. Theory Adv. Appl.*, pages 451–470. Birkhäuser Verlag, Basel, 2009. . URL https://doi.org/10.1007/978-3-7643-9919-1_28.
- Z. Sasvári. *Multivariate Characteristic and Correlation Functions*, volume 50 of *De Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, 2013. ISBN 978-3-11-022398-9; 978-3-11-022399-6. . URL <https://doi.org/10.1515/9783110223996>.
- J. Shen. *Local Structure of Random Fields - Properties and Inference*. ProQuest LLC, Ann Arbor, MI, 2019. ISBN 978-1687-99522-3. URL http://gateway.proquest.com/openurl?url_ver=Z39.88-2004&rft_val_fmt=info:ofi/fmt:kev:mtx:dis Thesis (Ph.D.)—University of Michigan.
- B. Simon. *A comprehensive Course in Analysis*. American Mathematical Society Providence, Rhode Island, 2015.
- M. L. Stein. *Interpolation of Spatial Data*. Springer Series in Statistics. Springer-Verlag, New York, 1999. ISBN 0-387-98629-4. . URL <https://doi.org/10.1007/978-1-4612-1494-6>. Some theory for Kriging.
- A. van Delft and M. Eichler. A note on Herglotz’s theorem for time series on function spaces. *Stochastic Process. Appl.*, 130(6):3687–3710, 2020.
- K. Yosida. *Functional Analysis*. Classics in Mathematics. Springer Berlin Heidelberg, 2012. ISBN 9783642618598. URL <https://books.google.com/books?id=yj4mBQAAQBAJ>.

SUPPLEMENT

The rest of the appendix contains the supplement to the main paper. In order to differentiate sections and results in the supplement from those in the main paper, we add a character “S” in front of sections, lemmas, etc., in the supplement.

Appendix S.1: Notation and preliminaries on Hilbert spaces.

The purpose of this section is to fix some notation and collect basic facts on Hilbert spaces used in the main paper and the proofs. The details can be found in most standard functional analysis texts such as [Conway \[2007\]](#).

Fix a separable Hilbert space \mathbb{V} over the field of complex numbers \mathbb{C} with inner product $\langle f, g \rangle$ and norm $\|f\| := \sqrt{\langle f, f \rangle}$, for $f, g \in \mathbb{V}$. We will focus on bounded linear operators $\mathcal{A} : \mathbb{V} \rightarrow \mathbb{V}$, namely linear operator with a bounded operator norm:

$$\|\mathcal{A}\|_{\text{op}} = \sup_{f: \|f\|=1} \|\mathcal{A}f\|.$$

Observe that for any two bounded operators \mathcal{A} and \mathcal{B} , we have $\|\mathcal{A}\mathcal{B}\|_{\text{op}} \leq \|\mathcal{A}\|_{\text{op}}\|\mathcal{B}\|_{\text{op}}$.

The adjoint of \mathcal{A} is denoted as \mathcal{A}^* , and we have

$$\langle \mathcal{A}f, g \rangle = \langle f, \mathcal{A}^*g \rangle = \overline{\langle \mathcal{A}^*g, f \rangle}, \quad \text{for all } f, g \in \mathbb{V}.$$

\mathcal{A} is self adjoint if $\mathcal{A} = \mathcal{A}^*$, which holds if and only if $\langle \mathcal{A}f, f \rangle \in \mathbb{R}$ for all $f \in \mathbb{V}$.

For self-adjoint operators \mathcal{A} and \mathcal{B} , write $\mathcal{A} \leq \mathcal{B}$ or $\mathcal{B} \geq \mathcal{A}$ whenever

$$\langle f, \mathcal{A}f \rangle \leq \langle f, \mathcal{B}f \rangle, \quad \text{for all } f \in \mathbb{V}.$$

In particular, \mathcal{A} is positive definite (or just positive) if $\mathcal{A} \geq 0$.

The outer (or tensor) product of two elements $f, g \in \mathbb{V}$, denoted by $f \otimes g$, is the operator that maps h to $\langle h, g \rangle f$ on \mathbb{V} . Clearly, $(f \otimes g)^* = g \otimes f$. For a compact, self-adjoint operator \mathcal{A} with spectral decomposition

$$\mathcal{A} = \sum_{j=1}^{\infty} \lambda_j e_j \otimes e_j,$$

where the $\lambda_j \in \mathbb{R}$, $e_j \in \mathbb{V}$ are the eigenvalues and eigenfunctions of \mathcal{A} , let

$$\mathcal{A}^{\pm} = \sum_{j=1}^{\infty} \lambda_j^{\pm} e_j \otimes e_j, \tag{S.1.1}$$

where $a^{\pm} = \max\{0, \pm a\}$, $a \in \mathbb{R}$, and

$$|\mathcal{A}| := \mathcal{A}^+ + \mathcal{A}^-.$$

The trace of an operator $\mathcal{A} : \mathbb{V} \rightarrow \mathbb{V}$ is

$$\text{trace}(\mathcal{A}) = \sum_{j=1}^{\infty} \langle \mathcal{A}e_j, e_j \rangle$$

if it is well defined, where $\{e_j\}$ is any CONS of \mathbb{V} .

An operator $\mathcal{A} : \mathbb{V} \rightarrow \mathbb{V}$ is trace class (or nuclear) if the self-adjoint positive operator $(\mathcal{A}^* \mathcal{A})^{1/2} =$ the square-root operator of $\mathcal{A}^* \mathcal{A}$ has finite trace, in which case the trace norm of \mathcal{A} is defined as

$$\|\mathcal{A}\|_{\text{tr}} := \text{trace}(\mathcal{B}) = \sum_{j=1}^{\infty} \lambda_j,$$

where $\{\lambda_j\}$ is the set of eigenvalues of $(\mathcal{A}^* \mathcal{A})^{1/2}$ or singular values of \mathcal{A} (counting multiplicities). The space of trace-class operators equipped with the trace norm will be denoted by \mathbb{T} , which is a Banach space. The collection of positive definite trace-class operators is denoted by \mathbb{T}_+ .

Appendix S.2: Cross covariance operators.

For a fixed probability space $(\Omega, \mathcal{F}, \mathbb{P})$, the class of \mathbb{V} -valued random elements will be denoted by $\mathcal{L}^0(\mathbb{V})$. We shall also work with the class $\mathcal{L}^2(\mathbb{V})$ of all $X \in \mathcal{L}^0(\mathbb{V})$ such that $\mathbb{E}\|X\|^2 < \infty$. The space $\mathcal{L}^2(\mathbb{V})$ becomes a Hilbert space with respect to the inner product $\mathbb{E}[\langle X, Y \rangle]$.

The expectation of \mathbb{V} -valued random elements can be defined in the sense of Bochner. See, for example, [Diestel and Uhl \[1977\]](#) or [Yosida \[2012\]](#) for details, or Section 2.5 in [Hsing and Eubank \[2015\]](#) for a brief treatment on Bochner's integral.

For two zero-mean random elements $X, Y \in \mathcal{L}^2(\mathbb{V})$, it is also natural to consider the cross covariance operator

$$\mathcal{C}(X, Y) := \mathbb{E}[X \otimes Y],$$

where $X \otimes Y$ is a random element taking values in the space of trace-class operators \mathbb{T} . The latter expectation can also be defined in the sense of Bochner in the Banach space $(\mathbb{T}, \|\cdot\|_{\text{tr}})$. For completeness, we briefly review next the Bochner integral in this setting.

S.2.1. The Bochner integral in $(\mathbb{T}, \|\cdot\|_{\text{tr}})$.

The set of trace class operators $(\mathbb{T}, \|\cdot\|_{\text{tr}})$ is a separable Banach space. We will need to consider integrals of measurable functions $f : (\Omega, \mathcal{F}, \mu) \rightarrow (\mathbb{T}, \mathcal{B}(\mathbb{T}))$, for a finite measure space $(\Omega, \mathcal{F}, \mu)$, where $\mathcal{B}(\mathbb{T})$ is the Borel σ -field on \mathbb{T} .

Such integrals can be defined in a standard way in the sense of Bochner as discussed in the references mentioned above. Here, we only give a simple criterion needed for our purposes, akin to Theorem 2.6.5 in [Hsing and Eubank \[2015\]](#).

Theorem S.2.1. *Let $f : \Omega \rightarrow \mathbb{T}$ be a measurable function. If $\int_{\Omega} \|f\|_{\text{tr}} d\mu < \infty$, then f is Bochner integrable and*

$$\left\| \int_{\Omega} f d\mu \right\|_{\text{tr}} \leq \int_{\Omega} \|f\|_{\text{tr}} d\mu.$$

Proof. Fix a CONS $\{e_j, j \in \mathbb{N}\}$ of \mathbb{V} , let $\Pi_n := \sum_{j=1}^n e_j \otimes e_j$ be the projection operator onto $\text{span}\{e_j, j = 1, \dots, n\}$ and define $\mathcal{A}_n := \Pi_n \mathcal{A} \Pi_n$. It is easy to see that

$$\|\mathcal{A} - \mathcal{A}_n\|_{\text{tr}} \leq 2\|\mathcal{A}\|_{\text{tr}} \quad \text{and} \quad \|\mathcal{A} - \mathcal{A}_n\|_{\text{tr}} \rightarrow 0, \quad \text{as } n \rightarrow \infty. \quad (\text{S.2.1})$$

Indeed, Proposition IV.2.3 on page 51 in [Gohberg et al. \[2000\]](#) implies that

$$\|\mathcal{A}_n\|_{\text{tr}} = \|\Pi_n \mathcal{A} \Pi_n\|_{\text{tr}} \leq \|\Pi_n\|_{\text{op}}^2 \|\mathcal{A}\|_{\text{tr}} = \|\mathcal{A}\|_{\text{tr}}$$

and hence the inequality in (S.2.1) follows from the triangle inequality. The convergence in (S.2.1) is a consequence of Theorem IV.5.5 on page 63 in Gohberg et al. [2000].

In view of Theorem 2.6.4 in Hsing and Eubank [2015], since $\int_{\Omega} \|f\|_{\text{tr}} d\mu < \infty$, to prove the Bochner integrability of f , it suffices to show that

$$\int_{\Omega} \|f - f_n\|_{\text{tr}} d\mu \rightarrow 0, \quad \text{as } n \rightarrow \infty, \quad (\text{S.2.2})$$

where $f_n(t) := \Pi_n f(t) \Pi_n$. This, however, follows from (S.2.1) and the Lebesgue dominated convergence theorem. \square

Remark S.2.1. It is well-known that if $\mathcal{O} : \mathbb{T} \rightarrow \mathbb{F}$ is a bounded linear operator into another Banach space $(\mathbb{F}, \|\cdot\|_{\mathbb{F}})$, and $f : \Omega \rightarrow \mathbb{T}$ is Bochner integrable, then $\mathcal{O}f$ is Bochner integrable and

$$\mathcal{O} \int_{\mathbb{R}^d} f d\mu = \int_{\mathbb{R}^d} \mathcal{O}f d\mu.$$

In particular, since $\text{trace} : \mathbb{T} \rightarrow \mathbb{R}$ is a continuous linear functional, we obtain

$$\text{trace} \left(\int_{\mathbb{R}^d} f d\mu \right) = \int_{\mathbb{R}^d} \text{trace}(f) d\mu, \quad (\text{S.2.3})$$

for all Bochner integrable f .

S.2.2. Existence and continuity of cross covariance operators.

For $X, Y \in \mathcal{L}^2(\mathbb{V})$, using Theorem S.2.1, we will show next that the *cross covariance* operator is well-defined the sense of Bochner.

Lemma S.2.2. *Let $X, Y \in \mathcal{L}^2(\mathbb{V})$, then $X \otimes Y \in \mathcal{L}^1(\mathbb{T})$. This implies that*

$$\mathcal{C}(X, Y) := \mathbb{E}[X \otimes Y], \quad (\text{S.2.4})$$

is a well-defined element of \mathbb{T} . We have moreover that

$$\|\mathcal{C}(X, Y)\|_{\text{tr}} \leq \mathbb{E}\|X \otimes Y\|_{\text{tr}} = \mathbb{E}[\|X\| \|Y\|] \leq (\mathbb{E}\|X\|^2)^{1/2} (\mathbb{E}\|Y\|^2)^{1/2}, \quad (\text{S.2.5})$$

and

$$\|\mathcal{C}(X, X)\|_{\text{tr}} = \mathbb{E}[\|X\|^2]. \quad (\text{S.2.6})$$

Proof. Note that $X \otimes Y \in \mathbb{T}$ and since $(X \otimes Y)^*(X \otimes Y) = \|X\|^2(Y \otimes Y)$, we have that $\|X \otimes Y\|_{\text{tr}}^2 = \text{trace}(\|X\|^2(Y \otimes Y)) = \|X\|^2\|Y\|^2$. This shows that

$$\|X \otimes Y\|_{\text{tr}} = \|X\| \|Y\|,$$

and hence by the Cauchy-Schwartz inequality,

$$\mathbb{E}\|X \otimes Y\|_{\text{tr}} \leq \mathbb{E}\|X\| \|Y\| \leq (\mathbb{E}\|X\|^2)^{1/2} (\mathbb{E}\|Y\|^2)^{1/2} < \infty$$

Theorem S.2.1 then implies that $X \otimes Y$ is Bochner integrable, $\mathcal{C}(X, Y)$ is well-defined, and (S.2.5) holds.

Finally, note $X \otimes X$ and $\mathcal{C}(X, X)$ are both positive operators and hence

$$\|X \otimes X\|_{\text{tr}} = \text{trace}(X \otimes X) \quad \text{and} \quad \|\mathcal{C}(X, X)\|_{\text{tr}} = \text{trace}(\mathcal{C}(X, X)). \quad (\text{S.2.7})$$

Since the trace is a bounded linear functional, we have

$$\text{trace}(\mathbb{E}(X \otimes X)) = \mathbb{E}(\text{trace}(X \otimes X)) = \mathbb{E}\|X\|^2,$$

which in view of (S.2.7) yields (S.2.6). \square

For zero-mean random variables in $\mathcal{L}^2(\mathbb{V})$, i.e., $\mathbb{E}[X] = \mathbb{E}[Y] = 0$, the operator $\mathcal{C}(X, Y)$ defined by (S.2.4) is referred to as the cross-covariance operator of X, Y .

Continuity of the covariance is a key element in Bochner's Theorem. The following result is the counterpart to the classical fact that for cross covariance functions, continuity at the diagonal implies continuity everywhere.

Proposition S.2.3. *Let $X = \{X(t), t \in \mathbb{R}^d\}$ be a \mathbb{V} -valued process. Then X is \mathcal{L}^2 -continuous, namely, $\mathbb{E}\|X(t') - X(s')\|^2 \rightarrow 0$ as $s', t' \rightarrow t$, if and only if, for every $t \in \mathbb{R}^d$,*

$$\|\mathcal{C}_X(s', t') - \mathcal{C}_X(t, t)\|_{\text{tr}} \rightarrow 0, \quad \text{as } (s', t') \rightarrow (t, t). \quad (\text{S.2.8})$$

In this case, we also have $\|\mathcal{C}_X(s', t') - \mathcal{C}_X(s, t)\|_{\text{tr}} \rightarrow 0$, as $(s', t') \rightarrow (s, t)$.

Proof. Suppose first that $\{X(t)\}$ is \mathcal{L}^2 -continuous. By (S.2.5)

$$\|\mathcal{C}_X(s', t) - \mathcal{C}_X(s, t)\|_{\text{tr}} \leq (\mathbb{E}\|X(t)\|^2)^{1/2} (\mathbb{E}\|X(s') - X(s)\|^2)^{1/2} \rightarrow 0,$$

as $s' \rightarrow s$. Thus, the triangle inequality implies

$$\|\mathcal{C}_X(s', t') - \mathcal{C}_X(s, t)\|_{\text{tr}} \leq \|\mathcal{C}_X(s', t') - \mathcal{C}_X(s, t')\|_{\text{tr}} + \|\mathcal{C}_X(s, t') - \mathcal{C}_X(s, t)\|_{\text{tr}} \rightarrow 0,$$

as $(s', t') \rightarrow (s, t)$, where we also used the elementary fact that \mathcal{L}^2 -continuity implies the continuity of $t \mapsto \mathbb{E}\|X(t)\|$. We have thus shown the continuity of the cross covariance operator in the trace norm.

Conversely, assume (S.2.8), i.e., the continuity of \mathcal{C}_X on the diagonal. In the sense of Bochner on the space $(\mathbb{T}, \|\cdot\|_{\text{tr}})$, we have

$$\begin{aligned} \mathcal{D}(s', t') &:= \mathbb{E}(X(t') - X(s')) \otimes (X(t') - X(s')) \\ &= \mathcal{C}_X(t', t') + \mathcal{C}_X(s', s') - (\mathcal{C}_X(t', s') + \mathcal{C}_X(s', t')), \end{aligned}$$

which converges to zero in trace norm, by (S.2.8). Relation (S.2.6) in Lemma S.2.2, however, yields $\|\mathcal{D}(s', t')\|_{\text{tr}} = \mathbb{E}\|X(t') - X(s')\|^2 \rightarrow 0$ as $s', t' \rightarrow t$, proving the desired continuity of $\{X(t)\}$. \square

Appendix S.3: Integration Theory for \mathbb{T}_+ -valued measures.

Let μ be a \mathbb{T}_+ -valued measure on $\mathcal{B}(\mathbb{R}^d)$, i.e., μ is a countably additive function on $\mathcal{B}(\mathbb{R}^d)$ taking values in \mathbb{T}_+ . Below we develop integration of a real or complex valued Borel measurable function with an operator-valued measure μ .

We follow the development of ordinary Lebesgue integration:

- (1) For any real nonnegative simple function $f = \sum_{i=1}^k c_i I_{A_i}$, define $\int f d\mu = \sum_{i=1}^k c_i \mu(A_i)$.

(2) For nonnegative measurable functions f , let

$$\int f d\mu = \lim_{n \rightarrow \infty} \int f_n d\mu \quad (\text{S.3.1})$$

in $(\mathbb{T}, \|\cdot\|, \|\cdot\|_{\text{tr}})$ where $\{f_n\}$ is any sequence of simple functions such that

- (a) $f_n \leq f_{n+1}$,
- (b) $f_n(x) \uparrow f(x)$ for all x ,
- (c) $\int f_n d\mu \leq \mathcal{B}$ for all n and some fixed $\mathcal{B} \in \mathbb{T}_+$.

The existence of $\{f_n\}$ satisfying (a) and (b) for any given nonnegative measurable f follows from standard measure theory. However, we need the extra condition (c) (along with the completeness of \mathbb{T}_+) to ensure that the limit on rhs of (S.3.1) exists and does not depend on the choice of $\{f_n\}$. Since $\mu(\mathbb{R}^d) \in \mathbb{T}_+$, (c) is automatically fulfilled if f is bounded.

(3) For a general real measurable f , let

$$\int f d\mu = \int f_+ d\mu - \int f_- d\mu$$

provided both terms on the right is finite. For a general complex f , let

$$\int f d\mu = \int f_{\text{Re}} d\mu + \mathbf{i} \int f_{\text{Im}} d\mu$$

where f_{Re} and f_{Im} are the real and imaginary parts, respectively.

The integral can be further extended to \mathbb{T} -valued signed measures, namely $\mu = \mu_+ - \mu_-$ where μ_+, μ_- are both \mathbb{T}_+ -valued measures, by $\int f d\mu := \int f d\mu_+ - \int f d\mu_-$.

The following propositions justify the definition of the integration described above.

Proposition S.3.1. *Let \mathcal{B} and $\mathcal{T}_n, n \geq 1$, be operators in \mathbb{T}_+ such that $\mathcal{T}_n \leq \mathcal{T}_{n+1} \leq \mathcal{B}$ for all n . Then \mathcal{T}_n converges to a limit in \mathbb{T}_+ .*

Proof. For $i \geq j$,

$$\|\mathcal{T}_i - \mathcal{T}_j\|_{\text{tr}} = \text{tr}(\mathcal{T}_i - \mathcal{T}_j) = \|\mathcal{T}_i\|_{\text{tr}} - \|\mathcal{T}_j\|_{\text{tr}}.$$

Now, $\|\mathcal{T}_n\|_{\text{tr}}$ is nondecreasing and bounded by $\|\mathcal{B}\|_{\text{tr}}$ for all n . Hence, $\|\mathcal{T}_n\|_{\text{tr}}$ converges to some finite nonnegative limit. Consequently,

$$\lim_{n \rightarrow \infty} \sup_{i \geq j \geq n} \|\mathcal{T}_i - \mathcal{T}_j\|_{\text{tr}} = 0.$$

This shows that $\{\mathcal{T}_n\}$ is Cauchy and has a limit by the completeness of \mathbb{T}_+ . □

Proposition S.3.2. *Let f_n be a sequence of nonnegative simple functions with $f_n \leq f_{n+1}$ for each n and such that $\int f_n d\mu \leq \mathcal{B}$ for some $\mathcal{B} \in \mathbb{T}$.*

- (i) $\int f_n d\mu$ converges in the space of trace class operators.
- (ii) If $\lim_{n \rightarrow \infty} f_n(x) \geq g(x)$ for all x for some simple function g , then $\lim_{n \rightarrow \infty} \int f_n d\mu \geq \int g d\mu$.

Proof. Note that nonnegative simple functions $g \leq f$ can be formulated as $f = \sum_{i=1}^k c_i I_{A_i}$, $g = \sum_{i=1}^k d_i I_{A_i}$ where $0 \leq c_i \leq d_i$ and the A_i are disjoint. Then it is easy to conclude that

$$\int g d\mu = \sum_{i=1}^k c_i \mu(A_i) \leq \sum_{i=1}^k d_i \mu(A_i) = \int f d\mu.$$

Thus, part (i) follows readily from Proposition S.3.1.

To prove (ii), let $g = \sum_{i=1}^k d_i I_{A_i}$, $A = \cup_{i=1}^k A_i$, and

$$E_n = \{x : f_n(x) + \epsilon > g(x)\}$$

for some fixed $\epsilon > 0$. Since

$$f_n \geq f_n I_{E_n \cap A} \geq (g - \epsilon) I_{E_n \cap A},$$

we have

$$\begin{aligned} \int f_n d\mu &\geq \int (g - \epsilon) I_{E_n \cap A} d\mu \\ &\geq \int g I_{E_n \cap A} d\mu - \epsilon \mu(A) \\ &= \sum_{i=1}^k c_i \mu(E_n \cap A_i) - \epsilon \mu(A). \end{aligned}$$

Letting $n \rightarrow \infty$, by the fact $E_n \cap A_i \uparrow A_i$,

$$\lim_{n \rightarrow \infty} \int f_n d\mu \geq \sum_{i=1}^k c_i \mu(A_i) - \epsilon \mu(A) = \int g d\mu - \epsilon \mu(A)$$

Since $\epsilon > 0$ is arbitrary, (ii) follows. □

Proposition S.3.3. *Let f be a nonnegative measurable function and $\{f_n\}$ be an increasing sequence of nonnegative simple functions satisfying (a)-(c) above. Then the limit $\lim_{n \rightarrow \infty} \int f_n d\mu$ does not depend on the particular sequence $\{f_n\}$.*

Proof. Suppose there are two sequences $\{f_n\}$ and $\{g_n\}$ of simple functions both satisfying (a)-(c). Then we have

$$\lim_{n \rightarrow \infty} f_n \geq g_m, \quad \lim_{n \rightarrow \infty} g_n \geq f_m$$

for all m , and, by Proposition S.3.2,

$$\lim_{n \rightarrow \infty} \int f_n d\mu \geq \int g_m d\mu, \quad \lim_{n \rightarrow \infty} \int g_n d\mu \geq \int f_m d\mu.$$

The result follows by letting $m \rightarrow \infty$. □

Appendix S.4: Proofs and auxiliary results for Section 4.

S.4.1. Proof of Theorem 4.2 (Bochner).

We begin by defining the notion of tightness for \mathbb{T}_+ -valued measures as follows.

Definition S.4.1. A sequence of finite \mathbb{T}_+ -valued measures $\{\mu_n\}$ is said to be tight if the following two conditions hold:

- (i) There exists some $\mathcal{B} \in \mathbb{T}_+$ such that $\mu_n(\mathbb{R}^d) \leq \mathcal{B}$ for all n .
- (ii) For any $\epsilon > 0$, there is a compact set $K_\epsilon \subset \mathbb{R}^d$ such that

$$\sup_n \|\mu_n(\mathbb{R}^d \setminus K_\epsilon)\|_{\text{tr}} < \epsilon.$$

We now proceed to prove Theorem 4.2. By Lemma S.4.2 below, $\mathcal{K}(t)$ is continuous in trace norm. We first consider the case

$$\int_{\mathbb{R}^d} \|\mathcal{K}(t)\|_{\text{tr}} dt < \infty, \quad (\text{S.4.1})$$

which will be relaxed in the second part of the proof. Let $\Pi_n = \sum_{i=1}^n e_i \otimes e_i$ be the projection operator onto $\text{span}\{e_1, \dots, e_n\}$ where $\{e_i\}$ is CONS of \mathbb{V} . Define $\mathcal{K}_n(t) = \Pi_n \mathcal{K}(t) \Pi_n$. By Proposition IV.2.3 on page 51 in Gohberg et al. [2000], $\|\mathcal{K}_n(t) - \mathcal{K}_n(s)\|_{\text{tr}} \leq \|\mathcal{K}(t) - \mathcal{K}(s)\|_{\text{tr}}$ and therefore $\mathcal{K}_n(t)$ is also continuous in trace norm. The same argument entails that

$$\int_{\mathbb{R}^d} \|\mathcal{K}_n(t)\|_{\text{tr}} dt \leq \int_{\mathbb{R}^d} \|\mathcal{K}(t)\|_{\text{tr}} dt < \infty. \quad (\text{S.4.2})$$

By Theorem S.2.1, the following Bochner's integrals in \mathbb{T} are well defined:

$$\widehat{\mathcal{K}}_n(x) := \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{-it^\top x} \mathcal{K}_n(t) dt, \quad \widehat{\mathcal{K}}(x) := \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{-it^\top x} \mathcal{K}(t) dt. \quad (\text{S.4.3})$$

It follows from Theorem IV.5.5 on page 63 in Gohberg et al. [2000], Theorem S.2.1, (S.4.2), and the Dominated Convergence Theorem, that

$$\|\mathcal{K}_n(t) - \mathcal{K}(t)\|_{\text{tr}} \rightarrow 0 \quad \text{and} \quad \|\widehat{\mathcal{K}}_n(x) - \widehat{\mathcal{K}}(x)\|_{\text{tr}} \rightarrow 0 \quad (\text{S.4.4})$$

for all t, x . For each $f, g \in \mathbb{V}$, define the complex-valued functions

$$\mathcal{K}_{f,g}(t) = \langle \mathcal{K}(t)g, f \rangle \quad \text{and} \quad \mathcal{K}_f(t) = \mathcal{K}_{f,f}(t).$$

Since $\mathcal{K}(\cdot)$ is positive definite and continuous at 0 in operator norm (entailed by continuity in trace norm), $\mathcal{K}_f(\cdot)$ is a positive-definite function and continuous at 0. It is also integrable by (S.4.1). We obtain, by the classical Bochner Theorem, that

$$\mathcal{K}_f(t) = \int_{\mathbb{R}^d} e^{it^\top x} \widehat{\mathcal{K}}_f(x) dx \quad \text{and} \quad \widehat{\mathcal{K}}_f(x) := \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{-it^\top x} \mathcal{K}_f(t) dt, \quad (\text{S.4.5})$$

where $\widehat{\mathcal{K}}_f(x) \geq 0$ for all x . Since $\langle \widehat{\mathcal{K}}(x)f, f \rangle = \widehat{\mathcal{K}}_f(x) \geq 0$, for all $f \in \mathbb{V}$, it follows that $\widehat{\mathcal{K}}(x)$ is a positive operator for all x . Notice that, by (S.4.5) and Lemma A.8,

$$\begin{aligned} \mathcal{K}_n(t) &= \sum_{i=1}^n \sum_{j=1}^n \mathcal{K}_{e_i, e_j}(t) e_i \otimes e_j \\ &= \sum_{i=1}^n \sum_{j=1}^n \left(\frac{\mathbf{i}-1}{2} (\mathcal{K}_{e_i}(t) + \mathcal{K}_{e_j}(t)) - \frac{\mathbf{i}}{2} \mathcal{K}_{\mathbf{i}e_j + e_i}(t) + \frac{1}{2} \mathcal{K}_{e_i + e_j}(t) \right) e_i \otimes e_j \\ &= \sum_{i=1}^n \sum_{j=1}^n \int_{\mathbb{R}^d} e^{\mathbf{i}t^\top x} \left(\frac{\mathbf{i}-1}{2} (\widehat{\mathcal{K}}_{e_i}(x) + \widehat{\mathcal{K}}_{e_j}(x)) - \frac{\mathbf{i}}{2} \widehat{\mathcal{K}}_{\mathbf{i}e_j + e_i}(x) + \frac{1}{2} \widehat{\mathcal{K}}_{e_i + e_j}(x) \right) dx e_i \otimes e_j \\ &= \int_{\mathbb{R}^d} e^{\mathbf{i}t^\top x} \widehat{\mathcal{K}}_n(x) dx, \end{aligned}$$

which, by (S.4.4), implies

$$\mathcal{K}(t) = \int_{\mathbb{R}^d} e^{\mathbf{i}t^\top x} \widehat{\mathcal{K}}(x) dx. \quad (\text{S.4.6})$$

To summarize, we have shown under (S.4.1) that

$$\mathcal{K}(t) = \int_{\mathbb{R}^d} e^{\mathbf{i}t^\top x} \widehat{\mathcal{K}}(x) dx \quad \text{and} \quad \widehat{\mathcal{K}}(x) = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{-\mathbf{i}t^\top x} \mathcal{K}(t) dt \in \mathbb{T}_+. \quad (\text{S.4.7})$$

Next, relax (S.4.1) and define

$$\mathcal{K}_\sigma(t) = e^{-\sigma^2 \|t\|^2/2} \mathcal{K}(t) = \frac{1}{(\sqrt{2\pi}\sigma)^d} \int_{\mathbb{R}^d} \mathcal{K}(t) e^{\mathbf{i}t^\top y} e^{-\|y\|^2/(2\sigma^2)} dy, \quad \sigma \geq 0. \quad (\text{S.4.8})$$

Note that $\mathcal{K}(t)e^{\mathbf{i}t^\top y}$ is positive definite in t . As a convex combination of positive definite functions, $\mathcal{K}_\sigma(\cdot)$ is also positive definite. Since $\int_{\mathbb{R}^d} \|\mathcal{K}_\sigma(t)\|_{\text{tr}} dt < \infty$, it follows from (S.4.7) that

$$\mathcal{K}_\sigma(t) = \int_{\mathbb{R}^d} e^{\mathbf{i}t^\top x} \widehat{\mathcal{K}}_\sigma(x) dx \quad \text{and} \quad \widehat{\mathcal{K}}_\sigma(x) = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{-\mathbf{i}t^\top x} \mathcal{K}_\sigma(t) dt \in \mathbb{T}_+ \quad (\text{S.4.9})$$

for any $\sigma \geq 0$. Define $\mu_\sigma(\cdot) := \int \widehat{\mathcal{K}}_\sigma(x) dx$ and suppose now that $\{\mu_\sigma\}$ is tight, which will be established in the last step of the proof. For any \mathbb{T}_+ -valued measure μ and e_i, e_j in CONS, write $\mu^{(i,j)}(A) = \langle \mu(A)e_j, e_i \rangle$ for any Borel set A . By Lemma S.4.3, for any sequence $\sigma_n \rightarrow 0$, the tightness of $\{\mu_{\sigma_n}\}$ implies there exist a \mathbb{T}_+ -valued measure μ and a subsequence sequence n' of n such that

$$\int_{\mathbb{R}^d} h(x) \mu_{\sigma_{n'}}^{(i,j)}(dx) \rightarrow \int_{\mathbb{R}^d} h(x) \mu^{(i,j)}(dx)$$

for all i, j and all bounded and continuous functions h . Since $\|\mathcal{K}_\sigma(t) - \mathcal{K}(t)\|_{\text{tr}} \rightarrow 0$ by the definition of \mathcal{K}_σ , we have, for all i, j ,

$$\begin{aligned} \langle \mathcal{K}(t)e_j, e_i \rangle &= \lim_{n' \rightarrow \infty} \langle \mathcal{K}_{\sigma_{n'}}(t)e_j, e_i \rangle = \lim_{n' \rightarrow \infty} \left\langle \int_{\mathbb{R}^d} e^{\mathbf{i}t^\top x} \mu_{\sigma_{n'}}(dx) e_j, e_i \right\rangle \\ &= \lim_{n' \rightarrow \infty} \int_{\mathbb{R}^d} e^{\mathbf{i}t^\top x} \langle \mu_{\sigma_{n'}}(dx) e_j, e_i \rangle = \lim_{n' \rightarrow \infty} \int_{\mathbb{R}^d} e^{-\mathbf{i}t^\top x} \mu_{\sigma_{n'}}^{(i,j)}(dx) \\ &= \int_{\mathbb{R}^d} e^{\mathbf{i}t^\top x} \mu^{(i,j)}(dx) = \left\langle \int_{\mathbb{R}^d} e^{\mathbf{i}t^\top x} \mu(dx) e_j, e_i \right\rangle, \end{aligned}$$

where the interchange of inner product and integration can be easily justified by the properties of Bochner's integral. Thus, for any $f, g \in \mathbb{V}$,

$$\langle \mathcal{K}(t)g, f \rangle = \left\langle \int_{\mathbb{R}^d} e^{it^\top x} \mu(dx) g, f \right\rangle,$$

which entails that $\mathcal{K}(t) = \int_{\mathbb{R}^d} e^{it^\top x} \mu(dx)$. Finally, we have

$$\mathcal{K}_f(t) = \int_{\mathbb{R}^d} e^{it^\top x} \langle \mu(dx) f, f \rangle.$$

Since this is a Fourier transform which does not depend on the sequence σ_n or $\sigma_{n'}$, we conclude that μ is unique.

It remains to show that $\{\mu_\sigma\}$ is tight. By (S.4.8) and (S.4.9), we have

$$\mu_\sigma(\mathbb{R}^d) = \mathcal{K}_\sigma(0) = \mathcal{K}(0) \in \mathbb{T}_+,$$

which implies (i) of the tightness definition. Next,

$$\begin{aligned} & \frac{1}{(2T)^d} \int_{t \in [-T, T]^d} \mathcal{K}_\sigma(t) dt \\ &= \frac{1}{(2T)^d} \int_{t \in [-T, T]^d} \int_{x \in \mathbb{R}^d} e^{it^\top x} \widehat{\mathcal{K}_\sigma}(x) dx dt \\ &= \int_{x \in \mathbb{R}^d} \prod_{j=1}^d \frac{\sin T x_j}{T x_j} \widehat{\mathcal{K}_\sigma}(x) dx \\ &\leq \int_{x \in [-b, b]^d} \prod_{j=1}^d \left| \frac{\sin T x_j}{T x_j} \right| \widehat{\mathcal{K}_\sigma}(x) dx + \int_{x \notin [-b, b]^d} \prod_{j=1}^d \left| \frac{\sin T x_j}{T x_j} \right| \widehat{\mathcal{K}_\sigma}(x) dx \\ &\leq \mu_\sigma([-b, b]^d) + \frac{1}{(Tb)^d} \mu_\sigma([-b, b]^d)^c \\ &= \mu_\sigma(\mathbb{R}^d) - (1 - 1/(Tb)^d) \mu_\sigma([-b, b]^d)^c, \end{aligned} \tag{S.4.10}$$

where “ \leq ” here is the operator inequality. By (S.4.10), with $b = T^{-1}(1 - 2^{-d})^{-1/d}$ so that $1 - 1/(Tb)^d = 2^{-d}$, we have

$$\begin{aligned} \mu_\sigma([-b, b]^d)^c &\leq 2^d \left(\mu_\sigma(\mathbb{R}) - \frac{1}{(2T)^d} \int_{[-T, T]^d} \mathcal{K}_\sigma(t) dt \right) \\ &= \frac{1}{T^d} \int_{[-T, T]^d} (\mathcal{K}(0) - \mathcal{K}_\sigma(t)) dt. \end{aligned}$$

Thus,

$$\left\| \mu_\sigma([-b, b]^d)^c \right\|_{\text{tr}} \leq \frac{1}{T^d} \int_{[-T, T]^d} \|\mathcal{K}(0) - \mathcal{K}_\sigma(t)\|_{\text{tr}} dt.$$

By the triangle inequality,

$$\|\mathcal{K}(0) - \mathcal{K}_\sigma(t)\|_{\text{tr}} \leq (1 - e^{-\sigma^2 t^2/2}) \|\mathcal{K}(0)\|_{\text{tr}} + e^{-\sigma^2 t^2/2} \|\mathcal{K}(0) - \mathcal{K}(t)\|_{\text{tr}} \rightarrow 0,$$

by the continuity of $\mathcal{K}(t)$ at 0. This show that $\left\| \mu_\sigma([-b, b]^d)^c \right\|_{\text{tr}} \rightarrow 0$ as $T \rightarrow 0$ and establishes the tightness of $\{\mu_\sigma\}$.

□

Lemma S.4.2. Assume that $\{\mathcal{K}(t), t \in \mathbb{R}\}$ is a collection of operators satisfying the assumptions of Theorem 4.2. Then we have

$$\lim_{\delta \rightarrow 0} \sup_{|t-s| < \delta} \|\mathcal{K}(t) - \mathcal{K}(s)\|_{\text{tr}} \rightarrow 0.$$

Proof. For any element $f \in \mathbb{V}$, denote $\mathcal{K}_f = \langle \mathcal{K}f, f \rangle$. It follows that $\mathcal{K}_f(t)$ is positive definite. By taking $c_1 = c_2 = 1$ and $t_1 = 0, t_2 = t$, one will have $\text{Im}(\mathcal{K}_f(t)) = -\text{Im}(\mathcal{K}_f(-t))$. Similarly, by taking $c_1 = 1, c_2 = i, t_1 = 0, t_2 = t$, we will have $\text{Re}(\mathcal{K}_f(t)) = \text{Re}(\mathcal{K}_f(-t))$. Therefore $\mathcal{K}_f(t) = \overline{\mathcal{K}_f(-t)}$. By taking $t_1 = t, t_2 = s, t_3 = 0$, we will have the following matrix to be nonnegative definite

$$\begin{pmatrix} \mathcal{K}_f(0) & \mathcal{K}_f(t-s) & \mathcal{K}_f(t) \\ \overline{\mathcal{K}_f(t-s)} & \mathcal{K}_f(0) & \mathcal{K}_f(s) \\ \overline{\mathcal{K}_f(t)} & \overline{\mathcal{K}_f(s)} & \mathcal{K}_f(0) \end{pmatrix}$$

As a result, its determinant will be nonnegative, that is

$$\begin{aligned} 0 &\leq \mathcal{K}_f^3(0) - \mathcal{K}_f(0)[|\mathcal{K}_f(t) - \mathcal{K}_f(s)|^2 + |\mathcal{K}_f(t-s)|^2] \\ &\quad - 2 \text{Re}([\mathcal{K}_f(s)\mathcal{K}_f(t)(\mathcal{K}_f(0) - \mathcal{K}_f(t-s))]) \\ &\leq \mathcal{K}_f^3(0) - \mathcal{K}_f(0)[|\mathcal{K}_f(t) - \mathcal{K}_f(s)|^2 + |\mathcal{K}_f(t-s)|^2] \\ &\quad - 2\mathcal{K}_f^2(0)|\mathcal{K}_f(0) - \mathcal{K}_f(t-s)|, \end{aligned}$$

where in the last line we applied the fact $|\mathcal{K}_f(t)| \leq \mathcal{K}_f(0)$ for all t . Rearranging terms then gives

$$\begin{aligned} |\mathcal{K}_f(t) - \mathcal{K}_f(s)|^2 &\leq \mathcal{K}_f^2(0) - |\mathcal{K}_f(t-s)|^2 + 2\mathcal{K}_f(0)|\mathcal{K}_f(0) - \mathcal{K}_f(t-s)| \\ &\leq 4\mathcal{K}_f(0)|\mathcal{K}_f(0) - \mathcal{K}_f(t-s)|, \end{aligned}$$

where we again use the fact that $|\mathcal{K}_f(t-s)| \leq \mathcal{K}_f(0)$.

Then, for CONS $\{e_i\}$ that is the system of eigenfunction of $\mathcal{K}(t) - \mathcal{K}(s)$, we have

$$\begin{aligned} \|\mathcal{K}(t) - \mathcal{K}(s)\|_{\text{tr}} &= \sum_i |\mathcal{K}_{e_i}(t) - \mathcal{K}_{e_i}(s)| \\ &\leq \sum_i 2\sqrt{\mathcal{K}_{e_i}(0)|\mathcal{K}_{e_i}(0) - \mathcal{K}_{e_i}(t-s)|} \\ &\leq 2\sqrt{\sum_i \mathcal{K}_{e_i}(0) \sum_i |\mathcal{K}_{e_i}(0) - \mathcal{K}_{e_i}(t-s)|} \\ &\leq 2\sqrt{\|\mathcal{K}(0)\|_{\text{tr}} \|\mathcal{K}(0) - \mathcal{K}(t-s)\|_{\text{tr}}}, \end{aligned}$$

where the last expression converges to 0 uniformly as $|t-s| \rightarrow 0$. □

Lemma S.4.3. Let $\{\mu_n\}$ be a class of finite \mathbb{T}_+ -valued measures. If $\{\mu_n\}$ is tight in the sense of Definition S.4.1, then there exists a finite \mathbb{T}_+ -valued measure μ and an infinite subsequence n' such that

$$\int_{\mathbb{R}^d} h(x) \mu_{n'}^{(i,j)}(dx) \rightarrow \int_{\mathbb{R}^d} h(x) \mu^{(i,j)}(dx), \quad (\text{S.4.11})$$

for all i, j and all bounded and continuous functions h , where $\mu^{(i,j)}(A) = \langle \mu(A)e_j, e_i \rangle$.

Proof. In this proof, the term measure without any qualification refers to the usual \mathbb{R}_+ -valued measure. Let $\{e_j\}$ be a CONS which we fix as the eigenfunctions of the operator \mathcal{B} in Definition S.4.1. First consider the signed measures $\mu_n^{(i,j)}(A) = \langle \mu_n(A)e_j, e_i \rangle$. The tightness of $\{\mu_n\}$ implies that, for any f , the sequence of finite measures $\{\mu_{n,f}(A) := \langle \mu_n(A)f, f \rangle\}$ is tight. By Prokhorov's Theorem, there exist a finite measure μ_f and an infinite subsequence n' such that

$$\int_{\mathbb{R}^d} h(x) \mu_{n',f}(dx) \rightarrow \int_{\mathbb{R}^d} h(x) \mu_f(dx)$$

for all bounded and continuous functions h . In particular, for each i there exist a finite measure $\mu^{[i,i]}$ and an infinite subsequence n' such that

$$\int_{\mathbb{R}^d} h(x) \mu_{n'}^{(i,i)}(dx) \rightarrow \int_{\mathbb{R}^d} h(x) \mu^{[i,i]}(dx)$$

for all bounded and continuous functions h . Applying the identity (Lemma S.4.2 of supplement)

$$\begin{aligned} \mu_n^{(i,j)}(A) = & \frac{\mathfrak{i} - 1}{2} (\mu_n^{(i,i)}(A) + \mu_n^{(j,j)}(A)) + \frac{1}{2} \langle e_i + e_j, \mu_n(A)(e_i + e_j) \rangle \\ & - \frac{\mathfrak{i}}{2} \langle e_i + \mathfrak{i}e_j, \mu_n(A)(e_i + \mathfrak{i}e_j) \rangle, \end{aligned} \quad (\text{S.4.12})$$

we also conclude that, for each i, j , there is an infinite subsequence n' such that

$$\int_{\mathbb{R}^d} h(x) \mu_{n'}^{(i,j)}(dx) \rightarrow \int_{\mathbb{R}^d} h(x) \mu^{[i,j]}(dx)$$

for all bounded and continuous functions h , where $\mu^{[i,j]}$ is defined as in (S.4.12) with $\mu_n^{(i,i)}$ replaced by $\mu^{[i,i]}$. Note that $\mu^{[i,j]}$ is in general a complex-valued set-function for $i \neq j$. Since the set of pairs (i, j) is countable, a straightforward argument shows that the convergence holds along a common subsequence n' for all $i, j \in \mathbb{N}$.

Construct, for a fixed m ,

$$\mu(A; m) := \sum_{i=1}^m \sum_{j=1}^m \mu^{[i,j]}(A) e_i \otimes e_j, \quad A \in \mathcal{B}(\mathbb{R}^d).$$

We consider the properties of $\mu(A; m)$ for any m and Borel set A .

- (i) $\mu(\cdot; m)$ is σ -additive as a finite sum of countably additive set-functions $\mu^{[i,j]}$.
- (ii) $\mu(A; m) \geq 0$.

Proof. We know that $\mu(A; m) \geq 0$ for $A \in \mathcal{F}$, where

$$\mathcal{F} = \{A \in \mathcal{B}(\mathbb{R}^d) : \mu^{[i,i]}(\partial A) = 0 \text{ for all } i\}. \quad (\text{S.4.13})$$

Indeed, the weak convergence $\mu_{n'}^{(i,i)} \xrightarrow{w} \mu^{[i,i]}$ implies that $\mu_{n'}^{(i,i)}(A) \rightarrow \mu^{[i,i]}(A)$ for all $A \in \mathcal{F}$, $i \in \mathbb{N}$, and hence

$$\Pi_m \mu_{n'}(A) \Pi_m = \sum_{i,j=1}^m \mu_{n'}^{(i,j)}(A) e_i \otimes e_j \rightarrow \mu(A; m),$$

as finite rank operators, where $\Pi_m := \sum_{i=1}^m e_i \otimes e_i$. This shows that $\mu(A; m) \geq 0$. Note that \mathcal{F} is a field (i.e., a nonempty collection of sets containing the whole space and closed under

finite unions and complements). Consider the measure $\mu_f(\cdot; m) := \langle f, \mu(\cdot; m)f \rangle$ on \mathcal{F} for any fixed $f \in \mathbb{V}$. By the measure extension theorem, there exists a unique measure $\widetilde{\mu}_f(\cdot; m)$ on $\mathcal{B}(\mathbb{R}^d)$ that agrees with $\mu_f(\cdot; m)$ on \mathcal{F} . Since both $\widetilde{\mu}_f(\cdot; m)$ and $\mu_f(\cdot; m)$ are countably additive on $\mathcal{B}(\mathbb{R}^d) = \sigma(\mathcal{F})$, Proposition S.4.4 implies that $\widetilde{\mu}_f(\cdot; m) \equiv \mu_f(\cdot; m)$ on $\mathcal{B}(\mathbb{R}^d)$. Thus, $\mu_f(A; m) \geq 0$ for all Borel sets A and all $f \in \mathbb{V}$. This together with (i) prove that $\mu(\cdot; m)$ is a \mathbb{T}_+ -valued measure.

(iii) $\mu(A; m) \leq \mathcal{B}$.

Proof. By (ii), weak convergence and the tightness assumption, for all m , Borel set A , and $f \in \mathbb{V}$,

$$\langle f, \mu(A; m)f \rangle \leq \langle f, \mu(\mathbb{R}^d; m)f \rangle = \lim_{n' \rightarrow \infty} \langle f, \mu_{n'}(\mathbb{R}^d; m)f \rangle \leq \langle f, \mathcal{B}f \rangle.$$

(iv) $\mu(A; m) = \Pi_m \mu(A; m') \Pi_m$ for all $m' > m$.

Fix any Borel set A . By Lemma S.4.6, for any infinite sequence m there exists a subsequence m' such that $\|\mu(A; m') - \mu(A)\|_{\text{HS}} \rightarrow 0$ for some $\mu(A) \in \mathbb{T}_+$ along m' . It follows that

$$\langle \mu(A)e_j, e_i \rangle = \langle \mu(A; \ell)e_j, e_i \rangle = \mu^{[i,j]}(A) \text{ for } \ell \geq i \vee j.$$

Thus, $\mu(A)$ does not depend on m' and we have

$$\lim_{m \rightarrow \infty} \|\mu(A; m) - \mu(A)\|_{\text{HS}} = 0. \quad (\text{S.4.14})$$

This implies, in particular, that $\mu(A) \geq 0$.

It remains to show $\mu(A)$ is countably additive. Suppose $B_k \downarrow \emptyset$. It follows that

$$\begin{aligned} \sup_m \|\mu(B_k; m)\|_{\text{tr}} &= \sup_m \sum_{j=1}^{\infty} \langle \mu(B_k; m)e_j, e_j \rangle \\ &\leq \sum_{j=1}^{\infty} \sup_m \langle \mu(B_k; m)e_j, e_j \rangle = \sum_{j=1}^{\infty} \langle \mu(B_k; j)e_j, e_j \rangle, \end{aligned}$$

since, by Property (iv), $\langle \mu(B_k; m)e_j, e_j \rangle = 0$ or $\langle \mu(B_k; j)e_j, e_j \rangle$ depending on $m < j$ or $\geq j$. Note that $\mu(B_k; j) \leq \mathcal{B} \in \mathbb{T}_+$ for all j by Property (iii) above. Therefore,

$$\langle \mu(B_k; j)e_j, e_j \rangle \leq B_j := \langle \mathcal{B}e_j, e_j \rangle,$$

where $\|\mathcal{B}\|_{\text{tr}} = \sum_j B_j < \infty$. We also have $\langle \mu(B_k; j)e_j, e_j \rangle \rightarrow 0$ as $k \rightarrow \infty$ for fixed j by the continuity property of measure. Thus, by the DCT,

$$\sup_m \|\mu(B_k; m)\|_{\text{tr}} \rightarrow 0 \text{ as } k \rightarrow \infty. \quad (\text{S.4.15})$$

By (S.4.14), (S.4.15) and the triangle inequality,

$$\|\mu(B_k)\|_{\text{HS}} = \lim_{m \rightarrow \infty} \|\mu(B_k; m)\|_{\text{HS}} \leq \sup_m \|\mu(B_k; m)\|_{\text{tr}} \rightarrow 0 \text{ as } k \rightarrow \infty. \quad (\text{S.4.16})$$

Let $A_i, i \geq 1$, be disjoint Borel sets. Since $\mu(\cdot)$ is finitely additive,

$$\left\| \mu(\cup_{i=1}^{\infty} A_i) - \sum_{i=1}^k \mu(A_i) \right\|_{\text{HS}} = \left\| \mu(\cup_{i=k+1}^{\infty} A_i) \right\|_{\text{HS}} \rightarrow 0, \quad (\text{S.4.17})$$

by (S.4.16). Letting $\mathcal{T}_k := \sum_{i=1}^k \mu(A_i) \leq \mathcal{B} \in \mathbb{T}_+$, since $\mathcal{T}_k \leq \mathcal{T}_{k+1}$, $k \in \mathbb{N}$, Proposition S.3.1 implies that there exists an operator $\mathcal{T} \in \mathbb{T}_+$, such that $\|\mathcal{T}_k - \mathcal{T}\|_{\text{tr}} \rightarrow 0$. This also implies that $\|\mathcal{T}_k - \mathcal{T}\|_{\text{HS}} \rightarrow 0$ and hence by (S.4.17), $\mathcal{T} = \mu(\cup_{i=1}^\infty A_i)$ and we have

$$\left\| \mu(\cup_{i=1}^\infty A_i) - \sum_{i=1}^k \mu(A_i) \right\|_{\text{tr}} \rightarrow 0, \quad \text{as } k \rightarrow \infty,$$

which proves the σ -additivity of μ in $(\mathbb{T}_+, \|\cdot\|_{\text{tr}})$. □

Proposition S.4.4. *Let \mathcal{F} be a field of subsets of \mathbb{R}^d and μ_1, μ_2 be finite countably additive set functions on $\sigma(\mathcal{F})$ that agree on \mathcal{F} . Then $\mu_1 \equiv \mu_2$.*

Proof. Let

$$\mathcal{L} = \{B \in \sigma(\mathcal{F}) : \mu_1(B) = \mu_2(B)\}.$$

First, $\mathcal{F} \subset \mathcal{L}$ by assumption. For $B \in \mathcal{L}$, since $\mathbb{R}^d \in \mathcal{F} \subset \mathcal{L}$,

$$\mu_1(B^c) = \mu_1(\mathbb{R}^d) - \mu_1(B) = \mu_2(\mathbb{R}^d) - \mu_2(B) = \mu_2(B^c).$$

So \mathcal{L} is closed under the operation of complement. We can similarly show that \mathcal{L} is closed under countable disjoint unions. Thus, \mathcal{L} is a λ -system containing \mathcal{F} , and we conclude $\mathcal{L} = \sigma(\mathcal{F})$ by the $\pi - \lambda$ Theorem. □

As usual, we say that a sequence $\{x_n\} \subset \mathbb{V}$ is relatively compact if there is an $x \in \mathbb{V}$ such that $\|x_{n'} - x\| \rightarrow 0$ along some subsequence. The following result is straightforward; see e.g., Proposition 30 in Melrose [2013].

Lemma S.4.5. *Let \mathbb{V} be a separable Hilbert space. A sequence $\{x_n\} \subset \mathbb{V}$ is relatively compact if and only if (i) $\sup_n \|x_n\| < \infty$ and (ii) for some (any) CONS $\{e_j\}$, we have*

$$\lim_{N \rightarrow \infty} \sup_n \sum_{i \geq N} |\langle x_n, e_i \rangle|^2 = 0.$$

Lemma S.4.6. *Let $0 \leq \mathcal{A}_n \leq \mathcal{B}$, where $\text{trace}(\mathcal{B}) < \infty$. Then, there exists a subsequence $n' \rightarrow \infty$ and an operator $\mathcal{A} \in \mathbb{T}_+$, such that*

$$\|\mathcal{A}_{n'} - \mathcal{A}\|_{\text{HS}} \rightarrow 0$$

where $\|\cdot\|_{\text{HS}}$ stands for the Hilbert-Schmidt norm.

Proof. Fix a CONS $\{e_i\}$ and consider the positive, self-adjoint Hilbert-Schmidt operators \mathcal{C}_n such that $\mathcal{A}_n = \mathcal{C}_n^2$. We have

$$\langle \mathcal{A}_n e_i, e_i \rangle = \|\mathcal{C}_n e_i\|^2 \leq \langle \mathcal{B} e_i, e_i \rangle =: B_i, \quad \text{where } \|\mathcal{B}\|_{\text{tr}} = \sum_i B_i < \infty. \quad (\text{S.4.18})$$

We will first show that for all e_j the sequence $\{\mathcal{C}_n e_j\}$ is relatively compact in \mathbb{V} . Indeed, $\sup_n \|\mathcal{C}_n e_j\| < \infty$ and by the Cauchy-Schwartz inequality and (S.4.18), we have

$$\sup_n \sum_{i \geq N} |\langle \mathcal{C}_n e_j, e_i \rangle|^2 = \sup_n \sum_{i \geq N} |\langle e_j, \mathcal{C}_n e_i \rangle|^2 \leq \sup_n \sum_{i \geq N} \|\mathcal{C}_n e_i\|^2 \leq \sum_{i \geq N} B_i.$$

Lemma S.4.5 implies the desired relative compactness. Thus, for some $n' \rightarrow \infty$, we have $\|\mathcal{C}_{n'}e_j - \mathcal{C}(e_j)\| \rightarrow 0$ for some $\mathcal{C}(e_j) \in \mathbb{V}$. By passing to a further subsequence, we can assume that the latter convergence holds for all j . In view of (S.4.18), we have

$$\|\mathcal{C}(e_j)\|^2 \leq B_j \quad \text{and hence} \quad \sum_j \|\mathcal{C}(e_j)\|^2 \leq \|\mathcal{B}\|_{\text{tr}} < \infty. \quad (\text{S.4.19})$$

We shall argue next that \mathcal{C} can be extended by linearity to a bounded linear Hilbert-Schmidt operator and show that $\|\mathcal{C}_{n'} - \mathcal{C}\|_{\text{HS}} \rightarrow 0$. For $x = \sum_i x_i e_i \in \mathbb{V}$, consider

$$\mathcal{C}(x) := \sum_j x_j \mathcal{C}(e_j). \quad (\text{S.4.20})$$

The latter series converges in \mathbb{V} . Indeed, by the Cauchy-Schwartz inequality and (S.4.19), for any $J \subset \mathbb{N}$, we have

$$\left(\sum_{j \in J} \|x_j \mathcal{C}(e_j)\| \right)^2 \leq \left(\sum_{j \in J} |x_j|^2 \right) \left(\sum_{j \in J} \|\mathcal{C}(e_j)\|^2 \right) \leq \|\mathcal{B}\|_{\text{tr}} \sum_{j \in J} |x_j|^2.$$

This proves that (S.4.20) converges in norm and in fact implies that $\mathcal{C}(x)$ is a bounded linear operator with $\|\mathcal{C}\|_{\text{op}} \leq \|\mathcal{B}\|_{\text{tr}}^{1/2}$. Furthermore, clearly \mathcal{C} is self-adjoint and by Relation (S.4.19) we have that \mathcal{C} is a Hilbert-Schmidt operator and \mathcal{C}^2 is trace class.

Observe now that

$$\|\mathcal{C}_{n'} - \mathcal{C}\|_{\text{HS}}^2 = \sum_i \|(\mathcal{C}_{n'} - \mathcal{C})e_i\|^2 \leq \sum_{i=1}^N \|\mathcal{C}_{n'}e_i - \mathcal{C}e_i\|^2 + 4 \sum_{i>N} B_i, \quad (\text{S.4.21})$$

where we used (S.4.18), (S.4.19), the triangle inequality for the norm in \mathbb{V} and the simple bound $(a+b)^2 \leq 2a^2 + 2b^2$, $a, b \geq 0$. For every $\epsilon > 0$, one can pick N large enough such that $\sum_{i>N} B_i < \epsilon/8$. Holding N fixed and appealing to the convergence $\|\mathcal{C}_{n'}e_i - \mathcal{C}e_i\| \rightarrow 0$, as $n' \rightarrow \infty$, one can make the first term in the right-hand side of (S.4.21) smaller than $\epsilon/2$, for all n' large enough. This argument shows that

$$\|\mathcal{C}_{n'} - \mathcal{C}\|_{\text{HS}} \rightarrow 0.$$

Now,

$$\mathcal{C}_{n'}^2 - \mathcal{C}^2 = (\mathcal{C}_{n'} - \mathcal{C})(\mathcal{C}_{n'} + \mathcal{C}) + \mathcal{C}(\mathcal{C}_{n'} - \mathcal{C}) + (\mathcal{C} - \mathcal{C}_{n'})\mathcal{C}.$$

Thus, $\|\mathcal{C}_{n'} - \mathcal{C}\|_{\text{HS}} = \|\mathcal{C}_{n'}^2 - \mathcal{C}^2\|_{\text{HS}} \rightarrow 0$. □

S.4.2. Proof of Theorem 4.11.

We first prove (ii). For simplicity, we only illustrate proofs for $d = 1$. The extension to the general case of d is straightforward. Also, since the “if” part is obvious, it suffices to focus on the “only if” part by assuming that $\mathcal{K} : \mathbb{R}^d \mapsto \mathbb{T}$ is continuous and conditionally positive definite of degree k .

Define

$$C_\mu(h) = \mathcal{K}(\mu * \tilde{\mu} + h), \quad h \in \mathbb{R}, \mu \in \Lambda_k. \quad (\text{S.4.22})$$

We first observe that C_μ is positive definite, i.e.,

$$\sum_{i=1}^n \sum_{j=1}^n c_i \bar{c}_j C_\mu(t_i - t_j) \geq 0 \quad \text{for all } c_i \in \mathbb{C}, t_i \in \mathbb{R}, i = 1, \dots, n.$$

This follows simply from

$$\sum_{i=1}^n \sum_{j=1}^n c_i \bar{c}_j C_\mu(t_i - t_j) = \mathcal{K} \left((\mu * \lambda) * \widetilde{\mu * \lambda} \right) \geq 0,$$

where $\lambda := \sum_{i=1}^n c_i \delta_{t_i}$, since $\mu * \lambda \in \Lambda_k$ and \mathcal{K} is conditionally positive definite. Thus, C_μ admits a spectral representation by Theorem 4.2 (Bochner's Theorem) where we denote by τ_μ the spectral measure.

Note also that

$$C_{\mu * \nu}(h) = C_\mu(\nu * \tilde{\nu} + h) = C_\nu(\mu * \tilde{\mu} + h), \quad h \in \mathbb{R}, \mu, \nu \in \Lambda_k. \quad (\text{S.4.23})$$

If $\mu = \sum_{k=1}^m c_k \delta_{x_k}$ then

$$C_\nu(\mu * \tilde{\mu} + h) = \int \sum_{k=1}^m \sum_{l=1}^m c_k \bar{c}_l e^{i(h+x_k-x_l)u} \tau_\nu(du) = \int e^{ihu} |\hat{\mu}(u)|^2 \tau_\nu(du)$$

where $\hat{\mu}(u) = \int e^{iux} \mu(dx)$. By (S.4.23) and the uniqueness of spectral measure,

$$|\hat{\mu}(u)|^2 \tau_\nu(du) \equiv |\hat{\nu}(u)|^2 \tau_\mu(du), \quad \mu, \nu \in \Lambda_k. \quad (\text{S.4.24})$$

Let $\{\mu_n\} \in \Lambda_k$ and μ_0 be as in Lemma S.4.9 where $\mu_n \rightarrow \mu_0$ weakly and $\hat{\mu}_0(u) \neq 0$ for $u \neq 0$. Since \mathcal{K} is continuous, we define a positive definite C_μ by taking a limit in (S.4.22), and conclude that (S.4.24) holds for any $\mu \in \Lambda_k$ and $\nu = \mu_0$. Define the measure

$$\chi(du) = \begin{cases} \frac{\tau_{\mu_0}(du)}{|\hat{\mu}_0(u)|^2} (1 \wedge |u|^{2k+2}) & u \neq 0, \\ 0 & u = 0. \end{cases} \quad (\text{S.4.25})$$

We will show χ is a \mathbb{T}_+ -valued measure in Proposition S.4.7. For $\mu \in \Lambda_k$, let $S_\mu := \{u : |\hat{\mu}(u)|^2 = 0\}$, which contains 0. By Lemma S.4.9, $\hat{\mu}_0(u) \neq 0$ for $u \neq 0$. Thus, for $u \notin S_\mu$, both $\hat{\mu}(u)$ and $\hat{\mu}_0(u)$ are nonzero and hence (S.4.24) entails that

$$\frac{\tau_{\mu_0}(du)}{|\hat{\mu}_0(u)|^2} = \frac{\tau_\mu(du)}{|\hat{\mu}(u)|^2}.$$

This implies that

$$\int e^{ihu} \tau_\mu(du) = \int \frac{e^{ihu} |\hat{\mu}(u)|^2}{1 \wedge |u|^{2k+2}} \chi(du) + \int_{u \in S_\mu} e^{ihu} \tau_\mu(du). \quad (\text{S.4.26})$$

Since $\hat{\mu}(z), z \in \mathbb{C}$, is an entire function, the set of zeros is countable by the Identity Theorem. Also, for $u \in S_\mu \setminus \{0\}$, we have $\hat{\mu}(u) = 0$ and $\hat{\mu}_0(u) \neq 0$, which, by (S.4.24) with $\nu = \mu_0$, entails $\tau_\mu(\{u\}) = 0$. Therefore, (S.4.26) becomes

$$\int e^{ihu} \tau_\mu(du) = \int \frac{e^{ihu} |\hat{\mu}(u)|^2}{1 \wedge |u|^{2k+2}} \chi(du) + \tau_\mu(\{0\}). \quad (\text{S.4.27})$$

Define

$$\mathcal{K}_0(h) = \int \frac{e^{\mathfrak{i}hu} - I_B(u)P(uh)}{1 \wedge |u|^{2k+2}} \chi(du), \quad (\text{S.4.28})$$

where B is a bounded interval containing 0 and

$$P(x) = \sum_{j=0}^{2k+1} (\mathfrak{i}x)^j / j!.$$

We will establish in Proposition S.4.7 that $\mathcal{K}_0(h)$ is well defined for every h . Notice that in (S.4.27), LHS = $C_\mu(h)$ and hence RHS is equal to

$$\begin{aligned} C_\mu(h) &= \int \frac{\sum_{i=1}^m \sum_{j=1}^m c_i \bar{c}_j e^{\mathfrak{i}(h+x_i-x_j)u}}{1 \wedge |u|^{2k+2}} \chi(du) + \tau_\mu(\{0\}) \\ &= \sum_{i=1}^m \sum_{j=1}^m c_i \bar{c}_j \int \frac{e^{\mathfrak{i}(h+x_i-x_j)u} - I_B(u)P(u(h+x_i-x_j))}{1 \wedge |u|^{2k+2}} \chi(du) + \tau_\mu(\{0\}) \\ &= \mathcal{K}_0(\mu * \tilde{\mu} + h) + \tau_\mu(\{0\}), \end{aligned} \quad (\text{S.4.29})$$

since $P(\mu * \tilde{\mu}) = 0$. The rest of the proof focuses on the property of $\tau_\mu(\{0\})$.

For any $\mu, \nu \in \Lambda_k$, define

$$C_{\mu,\nu}(h) = \mathcal{K}(\mu * \tilde{\nu} + h)$$

By Lemma A.8,

$$C_{\mu,\nu}(h) = \frac{\mathfrak{i}-1}{2}(C_\mu(h) + C_\nu(h)) + \frac{1}{2}C_{\mu+\nu}(h) - \frac{\mathfrak{i}}{2}C_{\mathfrak{i}\mu+\nu}(h), \quad (\text{S.4.30})$$

and

$$\begin{aligned} \mathcal{K}_0(\mu * \tilde{\nu} + h) &= \frac{\mathfrak{i}-1}{2}(\mathcal{K}_0(\mu * \tilde{\mu} + h) + \mathcal{K}_0(\nu * \tilde{\nu} + h)) \\ &\quad + \frac{1}{2}\mathcal{K}_0((\mu + \nu) * (\widetilde{\mu + \nu}) + h) - \frac{\mathfrak{i}}{2}\mathcal{K}_0((\mathfrak{i}\mu + \nu) * (\widetilde{\mathfrak{i}\mu + \nu}) + h). \end{aligned} \quad (\text{S.4.31})$$

Thus, it follows from (S.4.29) that

$$C_{\mu,\nu}(h) - \mathcal{K}_0(\mu * \tilde{\nu} + h) = \frac{\mathfrak{i}-1}{2}(\tau_\mu(\{0\}) + \tau_\nu(\{0\})) + \frac{1}{2}\tau_{\mu+\nu}(\{0\}) - \frac{\mathfrak{i}}{2}\tau_{\mathfrak{i}\mu+\nu}(\{0\}). \quad (\text{S.4.32})$$

Define

$$F(h_1, h_2) = \mathcal{K}(h_1 - h_2) - \mathcal{K}_0(h_1 - h_2),$$

and

$$F(\mu, \nu) = \iint F(h_1, h_2) d\mu(h_1) d\nu(h_2), \quad \mu, \nu \in \Lambda_k.$$

Applying (S.4.32), we obtain

$$\begin{aligned} F(\mu + h_1, \nu + h_2) &= C_{\mu,\nu}(h_1 - h_2) - \mathcal{K}_0(\mu * \tilde{\nu} + h_1 - h_2) \\ &= \frac{\mathfrak{i}-1}{2}(\tau_\mu(\{0\}) + \tau_\nu(\{0\})) + \frac{1}{2}\tau_{\mu+\nu}(\{0\}) - \frac{\mathfrak{i}}{2}\tau_{\mathfrak{i}\mu+\nu}(\{0\}). \end{aligned} \quad (\text{S.4.33})$$

Since this expression does not depend on h_1, h_2 for all $\mu, \nu \in \Lambda_k$, Lemma S.4.11 can be invoked to give

$$F(h_1, h_2) = \sum_{l=1}^k G_l^{(1)}(h_1)(h_2)^l + \sum_{l=1}^k G_l^{(2)}(h_2)(h_1)^l + C_0(h_1 - h_2)^{2k+2}, \quad (\text{S.4.34})$$

where the $G_l^{(i)}$ are arbitrary functions. Then, (S.4.33) and (S.4.34) entail that

$$F(\mu, \mu) = \tau_\mu(\{0\}) = C_0 w(\mu * \tilde{\mu})$$

where $w(h) = h^{2k+2}$ and μ is any arbitrary measure in Λ_k such that $w(\mu * \tilde{\mu}) \neq 0$. Thus, we conclude that

$$C_0 = \frac{\tau_\mu(\{0\})}{w(\mu * \tilde{\mu})},$$

which does not depend on $\mu \in \Lambda_k$. It follows that

$$\mathcal{K}_1(h) := \int \frac{e^{ihu} - I_B(u)P(uh)}{1 \wedge \|u\|^{2k+2}} \chi(du) + C_0 h^{2k+2}$$

is an equivalent version of \mathcal{K} in the sense that $\mathcal{K}_1(\mu * \tilde{\nu}) = \mathcal{K}(\mu * \tilde{\nu})$, $\mu, \nu \in \Lambda_k$. To prove uniqueness of χ , assume there is another χ' that can be used in the representation of the generalized covariance operator. Then, for any $\mu \in \Lambda_k$, we have

$$\int \frac{e^{ihu} |\hat{\mu}(u)|^2}{1 \wedge |u|^{2k+2}} \chi(du) \equiv \int \frac{e^{ihu} |\hat{\mu}(u)|^2}{1 \wedge |u|^{2k+2}} \chi'(du). \quad (\text{S.4.35})$$

By Lemma S.4.9 and weak convergence, (S.4.35) holds for $\mu = \mu_0$ as well. Since $\hat{\mu}_0(u) \neq 0$ for $u \neq 0$, it follows immediately that $\chi = \chi'$. This completes the proof of (ii).

We next prove (i). The first statement is obviously true. The proof of the second statement follows the exact same line of arguments as the proof of (ii), except we let

$$C_\mu(h) = C_Y(\mu + h, \mu) \quad \text{and} \quad C_{\mu, \nu}(h) = C_Y(\mu + h, \nu).$$

□

Proposition S.4.7. $\chi(du)$ defined in (S.4.25) is a finite \mathbb{T}_+ -valued measure and \mathcal{K}_0 defined in (S.4.28) is well defined and \mathbb{T} -valued function whenever X is mean-square continuous.

Proof. It suffices to show that there exists a neighborhood B containing 0 such that

$$\|\chi(B)\|_{\text{tr}} = \left\| \int_B \frac{1 \wedge |u|^{2k+2}}{|\hat{\mu}_0(u)|^2} \tau_{\mu_0}(du) \right\|_{\text{tr}} < \infty, \quad (\text{S.4.36})$$

$$\|\chi(B^c)\|_{\text{tr}} = \left\| \int_{B^c} \frac{1 \wedge |u|^{2k+2}}{|\hat{\mu}_0(u)|^2} \tau_{\mu_0}(du) \right\|_{\text{tr}} < \infty \quad (\text{S.4.37})$$

$$\left\| \int_B \frac{e^{ihu} - P(uh)}{1 \wedge |u|^{2k+2}} \chi(du) \right\|_{\text{tr}} < \infty, \quad (\text{S.4.38})$$

and

$$\left\| \int_{B^c} \frac{1}{1 \wedge |u|^{2k+2}} \chi(du) \right\|_{\text{tr}} < \infty. \quad (\text{S.4.39})$$

By Taylor's expansion,

$$\left\| \int_B |e^{ihu} - P(uh)| \chi(du) \right\|_{\text{tr}} \leq \left\| \frac{|h|^{2k+2}}{(2k+2)!} \int_B |u|^{2k+2} \chi(du) \right\|_{\text{tr}}$$

and

$$2(1 - \cos u) \geq u^2 - u^4/12 > u^2/2, \quad |u| < \sqrt{6}.$$

Thus, taking $B = (-\sqrt{6}, \sqrt{6})$, $\mu = (\delta_1 - \delta_0)^{*(k+1)}$, the $(k+1)$ -convolution power of the measure $\delta_1 - \delta_0$, we have $|\widehat{\mu}(u)| = |1 - e^{iu}|^{k+1}$, and since $\tau_{\mu_0}(\{0\}) = 0$, we obtain

$$\begin{aligned} \left\| \int_B \frac{1 \wedge |u|^{2k+2}}{|\widehat{\mu}_0(u)|^2} \tau_{\mu_0}(du) \right\|_{\text{tr}} &\leq \left\| \int_B \frac{|u|^{2k+2}}{|\widehat{\mu}_0(u)|^2} \tau_{\mu_0}(du) \right\|_{\text{tr}} \\ &\leq \left\| \int_B \frac{|u|^{2k+2}}{|\widehat{\mu}(u)|^2} \tau_{\mu}(du) \right\|_{\text{tr}} \leq \left\| \int_B \tau_{\mu}(du) \right\|_{\text{tr}} < \infty, \end{aligned}$$

and

$$\begin{aligned} \left\| \int_B \frac{|e^{ihu} - P(uh)|}{1 \wedge |u|^{2k+2}} \chi(du) \right\|_{\text{tr}} &\leq \left\| \frac{|h|^{2k+2}}{(2k+2)!} \int_B \frac{|u|^{2k+2}}{1 \wedge |u|^{2k+2}} \chi(du) \right\|_{\text{tr}} \\ &\leq \frac{|h|^{2k+2}}{2(2k+2)!} \left\| \int_B \frac{|\widehat{\mu}(u)|^2}{1 \wedge |u|^{2k+2}} \chi(du) \right\|_{\text{tr}} < \infty, \end{aligned}$$

where the last inequality follows from the facts that both sides of (S.4.26) are finite for any h (and in particular for $h = 0$) and $\mu \in \Lambda_k$. This establishes (S.4.36) and (S.4.38).

To prove (S.4.39), take μ_0 as the measure in Lemma S.4.9 and in the proof of Theorem 4.11. By Lemma S.4.9, we can pick δ such that $\inf_{u \in B^c} |\widehat{\mu}_0(u)| > \delta > 0$. Therefore,

$$\begin{aligned} \left\| \int_{B^c} \frac{(1 \wedge |u|^{2k+2})}{|\widehat{\mu}_0(u)|^2} \tau_{\mu_0}(du) \right\|_{\text{tr}} &\leq \left\| \delta^{-2} \int_{B^c} \tau_{\mu_0}(du) \right\|_{\text{tr}} < \infty, \\ \left\| \int_{B^c} \frac{1}{1 \wedge |u|^{2k+2}} \chi(du) \right\|_{\text{tr}} &\leq \left\| \delta^{-2} \int \frac{|\widehat{\mu}_0(u)|^2}{1 \wedge |u|^{2k+2}} \chi(du) \right\|_{\text{tr}} < \infty \end{aligned}$$

which proves (S.4.37) and (S.4.39). □

Let

$$M_f(k) = \{\mu = \mu_1 * \cdots * \mu_{k+1} : \mu_i \in \Lambda_0\}.$$

Also, define the larger class

$$M_c(k) = \left\{ \mu_1 * \cdots * \mu_{k+1} : 1(\mu_i) = 0, \right. \\ \left. \mu_i \text{'s are finite signed measures with compact supports} \right\}.$$

Note that $M_c(k)$ contains discrete as well as diffuse measures. It is easy to see that any $\mu \in M_c(k)$ annihilates polynomials of degree k .

Lemma S.4.8. (i) For any $k \in \mathbb{N}$, $M_f(k)$ is a dense set in $M_c(k)$ with respect to the weak topology.

(ii) For any open set O containing 0, there exists $\mu \in M_c(k)$ and $\delta > 0$ such that μ is symmetric, $\|\mu\|_{\text{TV}} < \infty$, where $\|\mu\|_{\text{TV}}$ stands for the total variation of μ , and for any $u \in O^c$, $\widehat{\mu}(u) > \delta$.

(iii) Let $f(h) : \mathbb{R} \mapsto \mathbb{T}$ be a continuous function in trace norm. If for some $k \in \mathbb{N}$, $f(\mu) = 0$ for any $\mu \in M_f(k)$, then $f(\lambda) = 0$ for any $\lambda \in \Lambda_k$.

Proof. Part (i) follows readily from Lemma 2.5 of Sasvári [2009]. The proof of (ii) is almost exactly the same as the proof for Lemma 3.2 of Sasvári [2009]. One only needs to notice that μ in the proof of their original lemma can be made to be symmetric under our scenario. Part (iii) is a trivial extension of Lemma C.9.4 of Sasvári [2013]. \square

Lemma S.4.9. There exists a measure μ on \mathbb{R} that satisfies

(i) $\|\mu\|_{\text{TV}} < \infty$.

(ii) For any open set O containing 0, there exists $\delta > 0$ such that for any $u \in O^c$, $\widehat{\mu}(u) > \delta$.

(iii) There exists a sequence of $\{\mu_n\} \subset M_f(k)$ converging to μ weakly.

Proof. For $V_n = [-1/n, 1/n]$, by (ii) of Lemma S.4.8, there exists $\{\mu_n\} \subset M_c(k)$ be a set of symmetric measures satisfying $\|\mu_n\|_{\text{TV}} \leq 1/2^n$ and $\mu_n(u) > 0$ for $u \notin V_n$. Then define $\mu = \sum_n \mu_n$, which obviously satisfies properties (i) and (ii). Property (iii) is guaranteed by (i) of Lemma S.4.8 and the fact $\sum_{n=1}^M \mu_n \rightarrow \mu$ weakly. \square

The following result can be easily obtained from Proposition 3 in Berschneider [2012] and its proof.

Lemma S.4.10. Let $f(h) : \mathbb{R} \mapsto \mathbb{T}$. If for some $k \in \mathbb{N}$, $f(\lambda) = 0$ for any $\lambda \in \Lambda_k$, then f has the following decomposition

$$f(h) = \sum_{i=0}^k a_i h^i,$$

where $a_i = \sum_{j=0}^{M_k} b_{ij} f(x_j)$ for some constants $b_{ij}, x_j, i = 0, \dots, k+1, j = 0, \dots, M_k$.

Lemma S.4.11. Assume that the bivariate function $F(h_1, h_2)$ from \mathbb{R}^2 to \mathbb{T} is continuous in trace norm. If $F(h_1 + \mu, h_2 + \nu)$ does not depend on h_1, h_2 for any two measures $\mu, \nu \in \Lambda_k$, then $F(h_1, h_2)$ has the following decomposition

$$F(h_1, h_2) = \sum_{l=1}^k G_l^{(1)}(h_1)(h_2)^l + \sum_{l=1}^k G_l^{(2)}(h_2)(h_1)^l + C_0(h_1 - h_2)^{2k+2},$$

where C_0 is an arbitrary operator in \mathbb{T} and the $G_l^{(i)}$'s are arbitrary \mathbb{T} -valued functions.

Proof. Fix $h_1, \mu \in \Lambda_k$. By (iii) of Lemma S.4.8, we obtain for any $\nu \in \Lambda_{k+1}$, $F(h_1 + \mu, h_2 + \nu) = 0$. Thus, by Lemma S.4.10,

$$F(h_1 + \mu, h_2) = \sum_{i=0}^{k+1} G_{1,i}(h_1 + \mu) h_2^i \quad \text{where} \quad G_{1,i}(h) = \sum_{j=0}^{M_{k+1}} b_{ij} F(h, x_j).$$

Therefore, again by Lemma S.4.10,

$$F(h_1, h_2) = \sum_{i=0}^{k+1} G_{1,i}(h_1) h_2^i + \sum_{i=0}^k G_{2,i}(h_2) h_1^i.$$

Notice that for any $\mu, \nu \in \Lambda_k$, $F(h_1 + \mu, h_2 + \nu) = G_{1,k+1}(h_1 + \mu)C(\nu)$, where $C(\nu) = \int h^{k+1}\nu(dx)$ is some constant. Since, by assumption $G_{1,k+1}(h_1 + \mu)$ does not depend on h_1 , we conclude that $G_{1,k+1}(\mu) = 0$ for all $\mu \in M_f(k+1)$. Applying (iii) of Lemma S.4.8 and Lemma S.4.10 again, we get $G_{1,k+1}(h_1) = \sum_{i=0}^{k+1} a'_i h_1^i$. The result follows after reorganizing the terms. \square

S.4.3. Proof of Theorem 4.7.

Recall the notation in Section A.4. Since

$$\begin{aligned} \text{trace}(X(s) \otimes X(t)) &= \sum_j \langle X(s) \otimes X(t) e_j, e_j \rangle \\ &= \sum_j \langle X(s), e_j \rangle \overline{\langle X(t), e_j \rangle} = \langle X(s), X(t) \rangle, \end{aligned}$$

we have, by Bochner's Theorem,

$$\langle X(s), X(t) \rangle_\Omega = \mathbb{E} \langle X(s), X(t) \rangle = \text{trace}(C(t-s)) = \int e^{i(t-s)^\top x} \|\mu\|_{\text{tr}}(dx), \quad (\text{S.4.40})$$

where μ the spectral measure of the stationary covariance operator, and $\|\mu\|_{\text{tr}}(A) := \text{trace}(\mu(A))$.

Let \mathbb{L} be the linear span of the functions $\{e^{it^\top \cdot}, t \in \mathbb{R}^d\}$. First, note that \mathbb{L} is dense in $\mathbb{L}^2(\mathbb{R}^d, \|\mu\|_{\text{tr}})$. To see this, assume $g \in \mathbb{L}^2(\mathbb{R}^d, \|\mu\|_{\text{tr}})$ and

$$\int e^{it^\top x} \overline{g(x)} \|\mu\|_{\text{tr}}(dx) = 0 \text{ for all } t.$$

Then $\nu(dx) := \overline{g(x)} \|\mu\|_{\text{tr}}(dx)$ is a finite measure and thus the above implies that g corresponds to the zero element in $\mathbb{L}^2(\mathbb{R}^d, \|\mu\|_{\text{tr}})$.

Define $\mathbb{H}(X) = \overline{\text{span}(X(t), t \in \mathbb{R}^d)}$, the closed linear span in $\mathbb{L}^2(\Omega)$, and the linear mapping

$$\mathcal{J}_X : \sum_{k=1}^n c_k e^{it_k^\top x} \mapsto \sum_{k=1}^n c_k X(t_k), \quad \mathbb{L} \mapsto \mathbb{H}(X).$$

By (S.4.40),

$$\|\mathcal{J}_X(f)\|_\Omega = \|f\|_{\mathbb{L}^2(\mathbb{R}^d, \|\mu\|_{\text{tr}})}, \quad f \in \mathbb{L}.$$

By the denseness of \mathbb{L} , \mathcal{J}_X can be readily extended to an isometric isomorphism, still denoted as \mathcal{J}_X , from $\mathbb{L}^2(\mathbb{R}^d, \|\mu\|_{\text{tr}})$ to $\mathbb{H}(X)$.

For $f, g \in \mathbb{L}^2(\mathbb{R}^d, \|\mu\|_{\text{tr}})$ and $e \in \mathbb{V}$, we have

$$\langle \mathbb{E}[\mathcal{J}_X(f) \otimes \mathcal{J}_X(g)] e, e \rangle = \int f(x) \overline{g(x)} \langle \mu(dx) e, e \rangle.$$

Indeed, this obviously holds for $f, g \in \mathbb{L}$, and, by the denseness of \mathbb{L} , the extension to $\mathbb{L}^2(\mathbb{R}^d, \|\mu\|_{\text{tr}})$ is straightforward. Then, applying Lemma A.8, it follows that

$$\mathbb{E}[\mathcal{J}_X(f) \otimes \mathcal{J}_X(g)] = \int f(x) \overline{g(x)} \mu(dx) \text{ for all } f, g \in \mathbb{L}^2(\mathbb{R}^d, \|\mu\|_{\text{tr}}). \quad (\text{S.4.41})$$

Define ξ by

$$\xi(A) = \mathcal{J}_X(\mathbf{1}_A), \quad A \in \mathcal{B}(\mathbb{R}^d).$$

By isometry, for any sequence of Borel sets A_n satisfying $A_n \rightarrow \emptyset$,

$$\|\xi(A_n)\|_\Omega = \|\mathbf{1}_{A_n}\|_{\mathbb{L}^2(\mathbb{R}^d, \|\mu\|_{\text{tr}})} = \|\mu(A_n)\|_{\text{tr}} \rightarrow 0.$$

For any disjoint Borel sets A and B , we have by (S.4.41) that

$$\mathbb{E}[\xi(A) \otimes \xi(B)] = 0.$$

Finally, by (S.4.41), for any Borel set A ,

$$\mathbb{E}[\xi(A) \otimes \xi(A)] = \mu(A).$$

Thus, we have proved that ξ is a random orthogonal measure with control measure μ (cf Definition 4.6) and therefore $\int e^{itx} d\xi(x)$ is well-defined by the construction in Section A.4.

It remains to show

$$X(t) = \int_{\mathbb{R}^d} e^{it^\top x} \xi(dx).$$

By (S.4.41), for any Borel set A ,

$$\mathbb{E}[X(t) \otimes \xi(A)] = \int e^{it^\top x} \mathbf{1}_A(x) \mu(dx).$$

Next, observe that, for any fixed s , $\int_{\mathbb{R}^d} e^{is^\top x} \xi(dx) \in \mathbb{H}(X)$ since it is the limit of $\mathcal{J}_X(g_n)$ where $g_n(x) = \sum_k c_k^n \mathbf{1}_{A_k^n}(x)$, $n \in \mathbb{N}$, $A_k \cap A_l = \emptyset$ when $k \neq l$, are step functions converging to $e^{is^\top x}$ in $\mathbb{L}^2(\mathbb{R}^d, \|\mu\|_{\text{tr}})$ as $n \rightarrow \infty$. Consequently,

$$\mathbb{E} \left[X(t) \otimes \int_{\mathbb{R}^d} e^{is^\top x} \xi(dx) \right] = \int e^{i(t-s)^\top x} \mu(dx) = \mathbb{E}[X(t) \otimes X(s)].$$

Thus, we have

$$\mathbb{E} \left[\sum_{i=1}^m c_i X(t_i) \otimes Y(s) \right] = 0$$

for $Y(s) := \int_{\mathbb{R}^d} e^{is^\top x} \xi(dx) - X(s)$, which is in $\mathbb{H}(X)$. Since linear combinations $\sum_{i=1}^m c_i X(t_i)$ are dense in $\mathbb{H}(X)$, we conclude that $\mathbb{E}(Y \otimes Y) = 0$ which implies that $Y = 0$ in $\mathbb{H}(X)$.

To prove the uniqueness of ξ , let η be a random orthogonal measure on \mathbb{R}^d both with structure measure χ . Then we will have

$$\int e^{it^\top x} \eta(dx) = \int e^{it^\top x} \xi(dx),$$

and, applying again the denseness of \mathbb{L} , we obtain

$$\int h(x) \eta(dx) = \int h(x) \xi(dx), \quad \text{almost surely, } \forall h \in \mathbb{L}^2(\mathbb{R}^d, \|\mu\|_{\text{tr}}).$$

Taking $h = \mathbf{1}_B$ for any Borel set B immediately leads to the conclusion $\eta \equiv \xi$.

□

S.4.4. Proof of Theorem 4.14.

We will focus on the “only if” part since the “if” part is obvious. By Theorem 4.7, for any $\lambda \in \Lambda_k$,

$$Y(\lambda + t) = \int e^{\mathbf{i}t^\top x} \xi_\lambda(dx)$$

for an orthogonal random measure ξ_λ with control measure τ_λ . Let μ_0 be the measure defined in the proof of Theorem 4.11. By the proof of Theorem 4.11, we also have

$$Y(\mu_0 + t) = \int e^{\mathbf{i}t^\top x} \xi_{\mu_0}(dx)$$

for an orthogonal random measure ξ_{μ_0} with control measure τ_{μ_0} . Thus,

$$Y(\lambda * \mu_0 + t) = \int e^{\mathbf{i}t^\top x} \hat{\mu}_0(x) \xi_\lambda(dx) = \int e^{\mathbf{i}t^\top x} \hat{\lambda}(x) \xi_{\mu_0}(dx).$$

By the uniqueness of the orthogonal random measure in the integral representation of a stationary process, we have

$$\hat{\mu}_0(x) \xi_\lambda(dx) \equiv \hat{\lambda}(x) \xi_{\mu_0}(dx). \quad (\text{S.4.42})$$

By (S.4.27), for any $\mu = \mu_0$ and λ , we have

$$\int \frac{e^{\mathbf{i}t^\top x} |\hat{\mu}(x)|^2}{1 \wedge |x|^{2k+2}} \chi(dx) = \int e^{\mathbf{i}t^\top x} \tau_\mu(dx) - \tau_\mu(\{0\}).$$

Now, define $\xi(\{0\}) = 0$, and for any Borel set B ,

$$\xi(B) = \int_{\mathbb{R}^d \setminus \{0\}} \mathbf{1}_B(x) \frac{1 \wedge |x|^{k+1}}{\hat{\mu}_0(x)} \xi_{\mu_0}(dx).$$

Then ξ is a well-defined orthogonal random measure with control measure χ . Set

$$U(t) := Y(t) - \int \frac{e^{\mathbf{i}t^\top x} - P(t^\top x)}{1 \wedge |x|^{k+1}} \xi(dx),$$

where $P(x) = \sum_{j=0}^k (\mathbf{i}x)^j / j!$, $x \in \mathbb{R}$. The integral exists by a similar argument to that of Proposition S.4.7. Therefore, for any $\lambda \in \Lambda_k$,

$$\begin{aligned} U(\lambda + t) &= \int e^{\mathbf{i}t^\top x} \xi_\lambda(dx) - \int_{\mathbb{R}^d \setminus \{0\}} e^{\mathbf{i}t^\top x} \frac{\hat{\lambda}(x)}{1 \wedge |x|^{k+1}} \xi(dx) \\ &= \int e^{\mathbf{i}t^\top x} \xi_\lambda(dx) - \int_{\mathbb{R}^d \setminus \{0\}} e^{\mathbf{i}t^\top x} \frac{\hat{\lambda}(x)}{\hat{\mu}_0(x)} \xi_{\mu_0}(dx) \\ &= \int e^{\mathbf{i}t^\top x} \xi_\lambda(dx) - \int_{\mathbb{R}^d \setminus \{0\}} e^{\mathbf{i}t^\top x} \xi_\lambda(dx) \\ &= \xi_\lambda(\{0\}) \end{aligned} \quad (\text{S.4.43})$$

where we applied (S.4.42). By a trivial extension of the proof of Lemma C.9.4 in Sasvári [2013], it follows that U is a random polynomial of degree no greater than $k + 1$. The uniqueness of ξ and Properties (iii) of the theorem are also easily established.

□

Appendix S.5: Gaussian elements in a Hilbert space.

Gaussian distributions in general Banach spaces have been studied extensively (see e.g. the monographs of [Ledoux and Talagrand \[1991\]](#) and [Kwapień and Woyczyński \[1992\]](#), and the references therein.) In this section, we provide a simple self-contained review of some basic results in the special case when the Banach space is a *separable Hilbert space* \mathbb{V} . The treatment may be of independent interest since it does not use advanced tools.

S.5.1. Real and complex Hilbert spaces.

It is well-known that the mean and covariance structure determine the distribution of *real* Gaussian vectors. Perhaps surprisingly, the mean and covariance alone are not enough to determine the distribution of complex Gaussian vectors. To clarify the issue, we start with a simple example in the case $\mathbb{V} = \mathbb{C}^m$.

Example S.5.1. Let $Z = X + iY$, where X and Y are independent and Normally distributed random vectors in \mathbb{R}^m having zero means and variance-covariance matrices $\Sigma_X = \mathbb{E}[XX^\top]$ and $\Sigma_Y = \mathbb{E}[YY^\top]$. Observe that the covariance operator of Z in \mathbb{C}^m , in the standard basis, can be expressed as the matrix:

$$\mathcal{C}_Z = \mathbb{E}[Z \otimes Z] \equiv \mathbb{E}[Z \bar{Z}^\top] = \Sigma_X + \Sigma_Y.$$

On the other hand, the real vector $\tilde{Z} := X + Y$ has the same covariance operator in \mathbb{C}^m as Z . This clearly shows that the distribution of zero-mean Gaussian random elements in complex Hilbert spaces are *not determined* by their covariance operators alone. To determine their distribution, one needs the additional information about the cross-covariance between the real and imaginary parts, that is, the pseudo-covariance $\mathcal{C}_{Z, \bar{Z}} = \mathbb{E}[Z \otimes \bar{Z}]$.

The purpose of this section is to clarify how one can deal with this issue in an abstract complex Hilbert space. We start by recalling and expanding on the notion of a real and imaginary part in a complex Hilbert space outlined in Section 4.3 of the main paper.

Real and imaginary parts in a complex Hilbert space. Note that in an abstract complex Hilbert space (over the field \mathbb{C}) the notion of a real and imaginary part is not well-defined unless one fixes a basis. Let $\mathcal{E} := \{e_j, j \in \mathbb{N}\}$ be a *fixed* CONS of \mathbb{V} . Then one can postulate that the CONS \mathcal{E} is *real* and for each $z = \sum_j z_j e_j \in \mathbb{V}$, with coordinates $z_j := \langle z, e_j \rangle$, we can define

$$\Re(z) \equiv \Re_{\mathcal{E}}(z) := \sum_j \Re(z_j) e_j \quad \text{and} \quad \Im(z) \equiv \Im_{\mathcal{E}}(z) := \sum_j \Im(z_j) e_j, \quad (\text{S.5.1})$$

as the *real* and *imaginary* parts of z , relative to the CONS \mathcal{E} . (Should one change the basis \mathcal{E} the notions of real and imaginary part will change.) Notice that $\mathbb{V}_{\mathbb{R}} := \{z \in \mathbb{V} : \Im(z) = 0\}$ is invariant to addition and multiplication by real scalars and it becomes a real Hilbert space, with the inner product inherited from \mathbb{V} . All elements of \mathbb{V} that belong to $\mathbb{V}_{\mathbb{R}}$ will be referred to as *real*.

For $z \in \mathbb{V}$, we shall write $z = \Re(z) + i\Im(z)$ and naturally define the *complex conjugate* $\bar{z} := \Re(z) - i\Im(z)$. The complex conjugate operation as well as the real and imaginary part operators extend to \mathbb{V} -valued random elements in a straightforward manner and we shall say that $x \in \mathbb{V}$ is *real* if $x \in \mathbb{V}_{\mathbb{R}}$, i.e., if its imaginary part is zero.

The complex conjugate of a linear operator $\mathcal{A} : \mathbb{V} \rightarrow \mathbb{V}$ is defined as: $\overline{\mathcal{A}}(x) := \overline{\mathcal{A}(\bar{x})}$, $x \in \mathbb{V}$. This implies that $\overline{\mathcal{A}(x)} = \overline{\mathcal{A}}(\bar{x})$, the operator $\overline{\mathcal{A}}$ is also *linear* and one can define the real and

imaginary parts of \mathcal{A} in as:

$$\Re(\mathcal{A}) := \frac{\mathcal{A} + \overline{\mathcal{A}}}{2} \quad \text{and} \quad \Im(\mathcal{A}) := \frac{\mathcal{A} - \overline{\mathcal{A}}}{2i}. \quad (\text{S.5.2})$$

Thus, $\mathcal{A} = \Re(\mathcal{A}) + i\Im(\mathcal{A})$ and the usual operations with complex numbers and vectors extend to the operator Banach algebra over the complex Hilbert space \mathbb{V} . We shall say that an operator \mathcal{A} is *real* if $\mathcal{A} = \Re(\mathcal{A})$ (i.e., $\Im(\mathcal{A}) = 0$). Observe that this is the case if and only if $\mathcal{A}(\mathbb{V}_{\mathbb{R}}) \subset \mathbb{V}_{\mathbb{R}}$.

Complexification of a real Hilbert space. On the other hand, suppose that one starts with a *real* Hilbert space $\mathbb{V}_{\mathbb{R}}$ with inner product $\langle \cdot, \cdot \rangle_{\mathbb{V}_{\mathbb{R}}}$. Then to be able to apply the results in Sections 4.1 and 4.2, one needs to extend $\mathbb{V}_{\mathbb{R}}$ to a Hilbert space over \mathbb{C} . This can be done with the standard method of *complexification*. Namely, consider the set $\mathbb{V} := \mathbb{V}_{\mathbb{R}} + i\mathbb{V}_{\mathbb{R}}$ of all pairs of $(x, y) \in \mathbb{V}_{\mathbb{R}} \times \mathbb{V}_{\mathbb{R}}$ written as $z := x + iy$, $x, y \in \mathbb{V}_{\mathbb{R}}$, where by definition

$$\Re(z) := x \quad \text{and} \quad \Im(z) := y \quad (\text{S.5.3})$$

are the real and imaginary parts of z . The complex conjugate operation is $\bar{z} := \Re(z) - i\Im(z)$ and the scalar multiplication is $(\alpha + i\beta) \cdot z := (\alpha \cdot x - \beta \cdot y) + i(\alpha \cdot y + \beta \cdot x)$, $\alpha, \beta \in \mathbb{R}$, $x, y \in \mathbb{V}_{\mathbb{R}}$. The inner product in \mathbb{V} is defined as $\langle z, z' \rangle := (\langle x, x' \rangle_{\mathbb{V}_{\mathbb{R}}} + \langle y, y' \rangle_{\mathbb{V}_{\mathbb{R}}}) + i(\langle y, x' \rangle_{\mathbb{V}_{\mathbb{R}}} - \langle x, y' \rangle_{\mathbb{V}_{\mathbb{R}}})$, where $z' = x' + iy'$, $x', y' \in \mathbb{V}_{\mathbb{R}}$. Thus, it is easy to see that \mathbb{V} becomes a Hilbert space over \mathbb{C} and $\mathbb{V}_{\mathbb{R}} = \{z : \Im(z) = 0\}$ is trivially embedded in \mathbb{V} . The complex conjugate operation as well as the real and imaginary part operators extend to \mathbb{V} -valued random elements in a straightforward manner and we shall say that $x \in \mathbb{V}$ is *real* if $x \in \mathbb{V}_{\mathbb{R}}$, i.e., if its imaginary part is zero.

If one fixes a CONS $\mathcal{E} := \{e_j\}$ in $\mathbb{V}_{\mathbb{R}}$, then it readily follows that \mathcal{E} this is also a CONS of \mathbb{V} and the definition (S.5.1) of the real and imaginary parts operators relative to \mathcal{E} coincides with (S.5.3). Thus, the notions of real, imaginary parts, as well as complex conjugate of \mathbb{V} -valued random elements and linear operators $\mathcal{A} : \mathbb{V} \rightarrow \mathbb{V}$ are exactly the same as outlined in (S.5.2) above, for example.

S.5.2. Fundamentals.

We shall assume that as in Section S.5.1, \mathbb{V} is a complex Hilbert space with a fixed real CONS so that we can consider the complex conjugates, real, and imaginary parts of the elements and operators on \mathbb{V} . Sometimes, it is helpful to view \mathbb{V} as $\mathbb{V}_{\mathbb{R}} + i\mathbb{V}_{\mathbb{R}}$ obtained by the method of complexification from the real Hilbert space $\mathbb{V}_{\mathbb{R}}$.

Definition S.5.1. A \mathbb{V} -valued random element X is said to be *Gaussian* if $\langle X, f \rangle$ is a (complex) Gaussian random variable, for all $f \in \mathbb{V}$. A random variable in \mathbb{C} is said to be *complex Gaussian* if its real and imaginary parts are jointly real Gaussian variables.

We shall need the following simple but fundamental result, which entails well-known classic bounds on the tail behavior of the norm of a Gaussian vector (see Remark S.5.2 below).

Proposition S.5.2. Let Z_j , $j \in \mathbb{N}$ be jointly Gaussian zero-mean real random variables with $\text{Var}(Z_j) = \sigma_j^2$ and let

$$\xi := \sum_{j \in \mathbb{N}} Z_j^2 \quad \text{and} \quad \sigma^2 := \sum_{j \in \mathbb{N}} \sigma_j^2,$$

which may take infinite values. Then, the following are equivalent:

$$(i) \sigma^2 \equiv \sum_{j \in \mathbb{N}} \sigma_j^2 < \infty$$

$$(ii) \mathbb{P}(\xi < \infty) = 1$$

$$(iii) \mathbb{P}(\xi < \infty) > 0.$$

If one (and hence all) of the above conditions holds, then

$$e^{\theta \sigma^2} \leq \mathbb{E}[e^{\theta \xi}] \leq \frac{1}{\sqrt{1 - 2\theta \sigma^2}}, \quad \text{for all } -\infty < \theta < \frac{1}{2\sigma^2}. \quad (\text{S.5.4})$$

We emphasize that the above inequality is valid, in particular, for all negative θ .

Before we give the proof, we make a few comments on the important consequences of this result.

Remark S.5.1. Proposition S.5.2 is a type of zero-one law for zero-mean Gaussian sequences stating that their squares are summable with probability zero or one. Many general zero-one laws for Gaussian sequences exist [see, e.g., Kallianpur, 1970].

Remark S.5.2. Let $\{e_j, j \in \mathbb{N}\}$ be a CONS of the Hilbert space \mathbb{V} . Then, provided one (and hence all) of the conditions in Proposition S.5.2 hold, we have that

$$X := \sum_{j \in \mathbb{N}} Z_j e_j$$

is a well-defined random element in \mathbb{V} . Since \mathbb{V} is separable, all probability distributions on \mathbb{V} are tight (cf Ulam's tightness theorem, or more generally, Theorem 11.5.1 in Dudley [2002]). Thus, X is *automatically* a Radon random element in the terminology in Section 2 of Ledoux and Talagrand [1991]. That is, for every $\epsilon > 0$, there is a compact set $\mathcal{K}_\epsilon \subset E$, such that $\mathbb{P}(X \in \mathcal{K}_\epsilon) \geq 1 - \epsilon$.

In this context, Proposition S.5.2 implies that every Gaussian random element X in \mathbb{V} is square integrable, in the sense that $\mathbb{E}[\|X\|^2] < \infty$. Moreover, the upper bound in (S.5.4) recovers the well-known result that $\mathbb{E}[e^{\theta \|X\|^2}] < \infty$, for all $\theta < 1/2\sigma^2$. This result is valid in much greater generality for (Radon) Gaussian random variables taking values in a Banach space [see, e.g., Corollary 3.2 on page 59-60 in Ledoux and Talagrand, 1991].

Proof of Proposition S.5.2. Since $\mathbb{E}[Z_j^2] = \sigma_j^2$, part (i) implies (ii) by the Tonelli-Fubini theorem. Trivially, (ii) \Rightarrow (iii).

We now prove the implication (iii) \Rightarrow (i). Suppose that $\xi = \sum_{j=1}^{\infty} Z_j^2$, which is finite with some positive probability. For all fixed n , using the joint Gaussianity of the Z_j 's, we have that

$$\xi_n := \sum_{j=1}^n Z_j^2 = \sum_{j=1}^n \lambda_{n,j} Z_{n,j}^2,$$

where $Z_{n,j}$, $j = 1, \dots, n$ are independent standard normal and where $\lambda_{n,j} \geq 0$ are such that

$$v_n := \sum_{j=1}^n \sigma_j^2 = \sum_{j=1}^n \lambda_{n,j}. \quad (\text{S.5.5})$$

Since $0 \leq \xi_n \leq \xi$ and $\mathbb{P}[\xi < \infty] > 0$, for all $t > 0$, we have

$$0 < \mathbb{E}[e^{-t\xi}] \leq \mathbb{E}[e^{-t\xi_n}] = \prod_{j=1}^n \mathbb{E}[e^{-t\lambda_{n,j} Z_{n,j}^2}] = \prod_{j=1}^n \frac{1}{\sqrt{1 + 2t\lambda_{n,j}}}, \quad (\text{S.5.6})$$

where in the last two equalities we used the independence of the $Z_{n,j}$'s and their normality. Upon taking logs, and changing the sign, we obtain

$$\frac{1}{2} \sum_{j=1}^n \log(1 + 2t\lambda_{n,j}) \leq -\log \mathbb{E}[e^{-t\xi}] < \infty. \quad (\text{S.5.7})$$

The function $f(\lambda_1, \dots, \lambda_n) := 2^{-1} \sum_{j=1}^n \log(1 + 2\lambda_j)$, $\lambda_j \geq 0$ is concave. Therefore, its minimum in the simplex

$$\left\{ \lambda = (\lambda_i)_{i=1}^n : \lambda_i \geq 0, i = 1, \dots, n, \sum_{i=1}^n \lambda_i = tv_n \right\}$$

is attained at one of the extremal points $(tv_n, 0, \dots, 0), \dots, (0, \dots, 0, tv_n)$. Since

$$f(tv_n, 0, \dots, 0) = \dots = f(0, \dots, 0, tv_n) = \frac{1}{2} \log(1 + 2tv_n),$$

from (S.5.7), we obtain

$$\sup_{n \in \mathbb{N}} \frac{1}{2} \log(1 + 2tv_n) \leq -\log \mathbb{E}[e^{-t\xi}] < \infty, \quad (\text{S.5.8})$$

which in turn implies that $\sup_{n \in \mathbb{N}} v_n = \sum_{n=1}^{\infty} \sigma_n^2 < \infty$, completing the proof of (i).

Assume now that one and hence all of the conditions (i)–(iii) hold. Since $\mathbb{E}[\xi] = \sigma^2 < \infty$, the lower bound in (S.5.4) follows by appealing to the Jensen's inequality for the convex function $x \mapsto e^{\theta x}$.

We now prove the upper bound in (S.5.4). Notice first that the case $\theta := -t < 0$ follows by taking $n \rightarrow \infty$ in (S.5.8). For the case $0 \leq \theta < 1/(2\sigma^2)$, as above let $\xi_n = \sum_{j=1}^n Z_j^2$ and as in (S.5.6), we obtain

$$\mathbb{E}[e^{\theta \xi_n}] = \prod_{j=1}^n \mathbb{E}[e^{\theta \lambda_{n,j} Z_{n,j}^2}] = \prod_{j=1}^n \frac{1}{\sqrt{1 - 2\theta \lambda_{n,j}}},$$

where now $-t$ is replaced by θ such that $\theta \in [0, 1/(2v_n))$. As before, consider the concave function $g(\lambda_1, \dots, \lambda_n) := 2^{-1} \sum_{i=1}^n \log(1 - 2\lambda_i)$, over the simplex

$$\left\{ \lambda = (\lambda_i)_{i=1}^n : \lambda_i \geq 0, \sum_{i=1}^n \lambda_i = \theta v_n < 1 \right\}.$$

Since the minimum of g is attained at an extremal point of the simplex, we obtain

$$-\log(\mathbb{E}[e^{\theta \xi_n}]) = g(\theta \lambda_{n,1}, \dots, \theta \lambda_{n,n}) \geq g(\theta v_n, 0, \dots, 0) = \frac{1}{2} \log(1 - 2\theta v_n)$$

which entails

$$\mathbb{E}[e^{\theta \xi_n}] \leq \frac{1}{\sqrt{1 - 2\theta v_n}}.$$

Since $\xi_n \uparrow \xi$ and $v_n \uparrow \sigma^2 < \infty$, by letting $n \rightarrow \infty$ and appealing to the Monotone Convergence Theorem, we obtain the upper bound in (S.5.4). \square

Proposition S.5.2 implies the following natural result.

Corollary S.5.3. *If X is a Gaussian random element in the separable Hilbert space \mathbb{V} , then:*

(i) $\mathbb{E}[\|X\|^2] < \infty$ and consequently the mean vectors $\mu_X := \mathbb{E}[X]$, the covariance $\mathcal{C}_X := \mathbb{E}[(X - \mu_X) \otimes (X - \mu_X)]$ and pseudo-covariance operators $\mathcal{C}_{X, \overline{X}} := \mathbb{E}[(X - \mu_X) \otimes (\overline{X} - \overline{\mu_X})]$ are well-defined elements in \mathbb{V} and \mathbb{T}_+ , respectively, in the sense of Bochner.

(ii) The distribution of X is determined by the mean vector μ_X and the pair of covariance and pseudo-covariance operators

$$\mathcal{C}_X = \mathbb{E}[(X - \mu_X) \otimes (X - \mu_X)] \quad \text{and} \quad \mathcal{C}_{X, \overline{X}} = \mathbb{E}[(X - \mu_X) \otimes (\overline{X} - \overline{\mu_X})]. \quad (\text{S.5.9})$$

Equivalently, the distribution of X is determined by μ_X and the covariance operator of the vector $Y := (X, \overline{X})^\top$ in the product Hilbert space $\mathbb{V}^2 := \mathbb{V} \times \mathbb{V}$.

Proof. Let $\{e_j, j \in \mathbb{N}\}$ be a real CONS of \mathbb{V} , i.e., $\Im(e_j) = 0, j \in \mathbb{N}$. We have that

$$X = \sum_{j \in \mathbb{N}} \xi_j e_j,$$

where $\xi_j := \langle X, e_j \rangle, j \in \mathbb{N}$ are (complex) jointly Gaussian (Definition S.5.1).

Suppose that $\mu_j := \mathbb{E}[\xi_j]$ and $\sigma_j^2 := \mathbb{E}[|\xi_j - \mu_j|^2], j \in \mathbb{N}$. To prove that $\mathbb{E}[\|X\|^2] < \infty$, it suffices to show that $\sum_{j \in \mathbb{N}} (|\mu_j|^2 + \sigma_j^2) < \infty$.

Let $\tilde{X} := \sum_{j \in \mathbb{N}} \tilde{\xi}_j e_j$ be an independent copy of X and observe that $Y := X - \tilde{X}$ is also a Gaussian \mathbb{V} -valued random element. The random variables $Z_j := \xi_j - \tilde{\xi}_j, j \in \mathbb{N}$ are (complex-valued) jointly Gaussian. Since $\mathbb{E}[Z_j] = 0$ and $\mathbb{P}(\|Y\|^2 < \infty) = 1$, by Proposition S.5.2 applied to the real and imaginary parts of the Z_j 's, we obtain

$$\mathbb{E}[\|Y\|^2] = \sum_{j \in \mathbb{N}} \mathbb{E}[|Z_j|^2] = 2 \sum_{j \in \mathbb{N}} \sigma_j^2 < \infty.$$

This, since $\sigma_j^2 = \mathbb{E}|\xi_j - \mu_j|^2$, implies that $X_0 := \sum_{j \in \mathbb{N}} (\xi_j - \mu_j) e_j$ takes values in \mathbb{V} , with probability one. But then $X - X_0 = \sum_{j \in \mathbb{N}} \mu_j e_j$ takes values in \mathbb{V} and hence $\sum_{j \in \mathbb{N}} |\mu_j|^2 < \infty$. This completes the proof of part (i).

Since $\mathbb{E}[\|X\|^2] = \mathbb{E}[\|\overline{X}\|^2] < \infty$ it follows that $\mathbb{E}[\|X\|] < \infty$ and hence the expectation $\mu := \mathbb{E}[X]$, the covariance, and pseudo-covariance operators in (S.5.9) are well-defined in the sense of Bochner [cf. Theorem 2.6.5 in Hsing and Eubank, 2015, and Theorem S.2.1] in the spaces $(\mathbb{V}, \|\cdot\|)$ and $(\mathbb{T}, \|\cdot\|_{\text{tr}})$, respectively. From the properties of the Bochner integral, we readily obtain that

$$\mu = \sum_{j \in \mathbb{N}} \mu_j e_j, \quad \mathcal{C}_X = \sum_{i, j \in \mathbb{N}} \mathbb{E}[(\xi_i - \mu_i) \overline{(\xi_j - \mu_j)}] e_i \otimes e_j,$$

and

$$\mathcal{C}_{X, \overline{X}} = \sum_{i, j \in \mathbb{N}} \mathbb{E}[(\xi_i - \mu_i)(\xi_j - \mu_j)] e_i \otimes e_j.$$

The distribution of X on \mathbb{V} is determined by the finite-dimensional distributions of the real and imaginary parts of its coordinates $\xi_j, j \in \mathbb{N}$ in any fixed CONS [see, e.g., Theorem 7.1.2 in Hsing and Eubank, 2015]. The latter are, in turn, determined by the coordinates of μ, \mathcal{C}_X , and $\mathcal{C}_{X, \overline{X}}$, completing the proof of (ii). \square

The next result provides uniform tail bounds for Gaussian vectors in a Hilbert space under the minimal condition that their norms are tight. Note that this does not mean in general that the Gaussian vectors are (uniformly) tight.

Corollary S.5.4. *Suppose that $\{X_n, n \in \mathbb{N}\}$ is a collection of zero-mean Gaussian random elements taking values in the separable Hilbert space \mathbb{V} . If the set of real random variables $\{\|X_n\|, n \in \mathbb{N}\}$ has uniformly tight distributions, then*

$$\sigma^2 := \sup_{n \in \mathbb{N}} \mathbb{E}[\|X_n\|^2] < \infty$$

and

$$\sup_{n \in \mathbb{N}} \mathbb{E} \left[\exp\{\theta \|X_n\|^2\} \right] \leq \frac{1}{\sqrt{1 - 2\theta\sigma^2}} < \infty, \quad \text{for all } \theta < \frac{1}{2\sigma^2}. \quad (\text{S.5.10})$$

Proof. Fix a CONS $\{e_j, j \in \mathbb{N}\}$ of \mathbb{V} . Let

$$U_{n,i} := \Re(\langle X_n, e_i \rangle) \quad \text{and} \quad V_{n,i} := \Im(\langle X_n, e_i \rangle),$$

be the real and imaginary parts of $\langle X_n, e_i \rangle$. Since X_n is a valid \mathbb{V} -valued random variable, we have that

$$\|X_n\|^2 = \sum_{i \in \mathbb{N}} (U_{n,i}^2 + V_{n,i}^2) < \infty, \quad (\text{S.5.11})$$

and hence by Proposition S.5.2 applied to the jointly Gaussian zero-mean random variables $\{U_{n,i}, V_{n,i}, i \in \mathbb{N}\}$, we obtain $\sigma_n^2 := \mathbb{E}[\|X_n\|^2] < \infty$.

Since $\{\|X_n\|, n \in \mathbb{N}\}$ are uniformly tight, there is an $M > 0$ such that

$$\inf_{n \in \mathbb{N}} \mathbb{P}(\|X_n\|^2 \leq M) \geq 1/2 > 0.$$

This implies that

$$0 < \frac{1}{2}e^{-M} \leq \inf_{n \in \mathbb{N}} \mathbb{E}[\exp\{-\|X_n\|^2\}] \leq \inf_{n \in \mathbb{N}} \frac{1}{\sqrt{1 + 2\sigma_n^2}}, \quad (\text{S.5.12})$$

where the last inequality follows by applying the upper bound in (S.5.4) with $\theta := -1$ to each $\|X_n\|^2$ in (S.5.11).

Relation (S.5.12) shows that $\sigma^2 := \sup_{n \in \mathbb{N}} \sigma_n^2 < \infty$. The inequality in (S.5.4) is a simple application of the upper bound in (S.5.4). \square

S.5.3. Convergence in distribution.

We start with the following natural criterion for convergence in distribution in the space $\mathcal{L}^2(\mathbb{V})$ of all \mathbb{V} -valued random elements with $\mathbb{E}[\|X\|^2] < \infty$.

For a sequence of $X_n \in \mathcal{L}^2(\mathbb{V})$, let

$$\mu_n := \mathbb{E}[X_n] \quad \text{and} \quad \mathcal{C}_n := \mathbb{E}[(X_n - \mu_n) \otimes (X_n - \mu_n)]$$

be the mean vectors and covariance operators, respectively. As we know (cf Lemma S.2.2), $\mathcal{C}_n \in \mathbb{T}_+$ are necessarily positive trace-class operators.

Theorem S.5.5. *Fix some (any) CONS $\{e_j, j \in \mathbb{N}\}$ of \mathbb{V} and let $\Pi_k := \sum_{j=1}^k e_j \otimes e_j$. Let $\{X_n\} \subset \mathcal{L}^2(\mathbb{V})$.*

(i) *We have $X_n \xrightarrow{d} X_\infty$, as $n \rightarrow \infty$, if*

- (1) $\langle X_n, f \rangle \xrightarrow{d} \langle X_\infty, f \rangle$, for all $f \in \mathbb{V}$
- (2) $\|\mu_n - \mu_\infty\| \rightarrow 0$, and
- (3) $\|\mathcal{C}_n - \mathcal{C}_\infty\|_{\text{op}} \rightarrow 0$ and

$$\limsup_{n \rightarrow \infty} (\text{tr}(\mathcal{C}_n) - \text{tr}(\mathcal{C}_{n,k})) \rightarrow 0 \text{ as } k \rightarrow \infty,$$

where $\mathcal{C}_{n,k} := \Pi_k \mathcal{C}_n \Pi_k$.

(ii) Conversely, if $\{\|X_n\|^2, n \in \mathbb{N}\}$ are uniformly integrable, then the convergence $X_n \xrightarrow{d} X_\infty$, as $n \rightarrow \infty$ implies (1), (2) and (3) in part (i), above.

Remark S.5.3. Notice that the above result applies to general not necessarily Gaussian \mathbb{V} -valued random elements.

Proof. Part (ii) is immediate since uniform integrability and convergence in distribution imply convergence of the moments.

Part (i): If $\|\mu_n - \mu_\infty\| \rightarrow 0$, then by Slutsky's theorem, the convergence $X_n - \mu_n \xrightarrow{d} X_\infty - \mu_\infty$ implies $X_n \xrightarrow{d} X_\infty$. Thus, without loss of generality, we will assume that $\mu_n = \mu_\infty = 0$.

Condition (1), in view of the Wold device, is equivalent to the finite-dimensional convergence

$$(\langle X_n, f_1 \rangle, \dots, \langle X_n, f_k \rangle) \xrightarrow{d} (\langle X_\infty, f_1 \rangle, \dots, \langle X_\infty, f_k \rangle). \quad (\text{S.5.13})$$

To prove $X_n \xrightarrow{d} X_\infty$, it remains to establish tightness. For convenience of notation, assume that the eigenspaces of $\mathcal{C}_n, \mathcal{C}_\infty$ are all one dimensional. If this is not true, we need to work with eigenspaces and the notation becomes more complicated. Let $e_{n,j}$ and $e_{\infty,j}$ be the eigenfunctions of \mathcal{C}_n and \mathcal{C}_∞ , respectively, that correspond to the j -th descending eigenvalues. Let $S_{n,k} = \text{span}(e_{n,j}, j \leq k)$ and $S_{\infty,k} = \text{span}(e_{\infty,j}, j \leq k)$. For any $S \subset \mathbb{V}$, define

$$S^\epsilon = \{f \in \mathbb{V} : \|f - g\| < \epsilon \text{ for some } g \in S\}.$$

Write

$$X_n = X_{n,k} + \tilde{X}_{n,k}.$$

where $X_{n,k} = \Pi_{S_{n,k}} X_n$. Clearly, if $X_{n,k} \in S_{\infty,k}^\epsilon$ and $\|\tilde{X}_{n,k}\| \leq \epsilon$, then $X_n \in S_{\infty,k}^{2\epsilon}$. Thus,

$$\begin{aligned} \mathbb{P}(X_n \in S_{\infty,k}^{2\epsilon}) &\geq \mathbb{P}(X_{n,k} \in S_{\infty,k}^\epsilon, \|\tilde{X}_{n,k}\| \leq \epsilon) \\ &= \mathbb{P}(X_{n,k} \in S_{\infty,k}^\epsilon) - \mathbb{P}(X_{n,k} \in S_{\infty,k}^\epsilon, \|\tilde{X}_{n,k}\| > \epsilon). \end{aligned} \quad (\text{S.5.14})$$

By (S.5.13) and the fact that the eigenvalues and eigenfunctions of \mathcal{C}_n converge to those of \mathcal{C}_∞ under $\|\mathcal{C}_n - \mathcal{C}_\infty\|_{\text{op}} \rightarrow 0$ [cf. Theorems 5.1.6 and 5.1.8 in [Hsing and Eubank, 2015](#)], we have $X_{n,k} = \Pi_{S_{n,k}} X_n \xrightarrow{d} \Pi_{S_{\infty,k}} X_\infty$. Since $S_{\infty,k}^\epsilon$ is open, by the Portmanteau Theorem of weak convergence,

$$\liminf_{n \rightarrow \infty} \mathbb{P}(X_{n,k} \in S_{\infty,k}^\epsilon) \geq \mathbb{P}(\Pi_{S_{\infty,k}} X_\infty \in S_{\infty,k}^\epsilon) = 1 \text{ for each } k.$$

It follows that, for any k ,

$$\begin{aligned} &\limsup_{n \rightarrow \infty} \mathbb{P}(X_{n,k} \in S_{\infty,k}^\epsilon, \|\tilde{X}_{n,k}\| > \epsilon) \\ &\leq \limsup_{n \rightarrow \infty} \mathbb{P}(\|\tilde{X}_{n,k}\| > \epsilon) \\ &\leq \epsilon^{-2} \limsup_{n \rightarrow \infty} (\text{tr}(\mathcal{C}_n) - \text{tr}(\mathcal{C}_{n,k})). \end{aligned} \quad (\text{S.5.15})$$

By (S.5.14)-(S.5.15) and assumption (ii), we conclude that for any $\epsilon, \delta > 0$, there exists $k = k(\epsilon, \delta)$ and $n(\epsilon, \delta)$, such that for all $n > n(\epsilon, \delta)$, we have

$$\mathbb{P}(X_n \in S_{\infty, k}^{2\epsilon}) \geq 1 - \delta.$$

For $n \leq n(\epsilon, k)$, we can create another finite dimensional set S by enlarging $S_{\infty, k}$ (e.g., adding eigenfunctions of \mathcal{C}_n 's to the spanning set) so that

$$\inf_n \mathbb{P}(X_n \in S^{2\epsilon}) \geq 1 - \delta.$$

Thus, the *flat concentration* condition [cf. Hsing and Eubank, 2015, Theorem 7.7.5] is fulfilled. \square

The following is a necessary and sufficient condition for the weak convergence of Gaussian distributions on a *real* separable Hilbert space. Applying this result to the concatenation of the real and imaginary parts, one can extend this criterion to Gaussian distributions in complex Hilbert spaces.

Theorem S.5.6. *Let X_n , $n \in \mathbb{N}$ be Gaussian elements taking values in the real separable Hilbert space $\mathbb{V}_{\mathbb{R}}$. Suppose that the X_n 's have means μ_n and covariance operators \mathcal{C}_n .*

We have $X_n \xrightarrow{d} X_{\infty}$ as $n \rightarrow \infty$ in $\mathbb{V}_{\mathbb{R}}$ if and only if X_{∞} is Gaussian and

$$\|\mu_n - \mu_{\infty}\| \rightarrow 0 \quad \text{and} \quad \|\mathcal{C}_n - \mathcal{C}_{\infty}\|_{\text{tr}} \rightarrow 0, \quad (\text{S.5.16})$$

where μ_{∞} and \mathcal{C}_{∞} are the mean vector and covariance operator of X_{∞} .

Proof. The ‘if’ part follows from Theorem S.5.5. Indeed, by (S.5.16) it follows that $\langle X_n, f \rangle \xrightarrow{d} \langle X_{\infty}, f \rangle$ for all $f \in \mathbb{V}$. Relation (S.5.16) implies also that $\|\mathcal{C}_n\|_{\text{tr}} \rightarrow \|\mathcal{C}_{\infty}\|_{\text{tr}}$ and for each k , $\|\mathcal{C}_{n, k}\|_{\text{tr}} \rightarrow \|\mathcal{C}_{\infty, k}\|_{\text{tr}}$. Thus,

$$\limsup_{n \rightarrow \infty} (\|\mathcal{C}_n\|_{\text{tr}} - \|\mathcal{C}_{n, k}\|_{\text{tr}}) = \|\mathcal{C}_{\infty}\|_{\text{tr}} - \|\mathcal{C}_{\infty, k}\|_{\text{tr}},$$

vanishes as $k \rightarrow \infty$, which shows that the conditions of Theorem S.5.5 are fulfilled.

We now prove the ‘only if’ part. The weak convergence $X_n \xrightarrow{d} X_{\infty}$ implies that for all $f \in \mathbb{V}$, we have $\langle X_n, f \rangle \xrightarrow{d} \langle X_{\infty}, f \rangle$, where the limit is a (complex) Gaussian random variable. This proves that X_{∞} is a Gaussian element in \mathbb{V} with some mean vector $\mu_{\infty} \in \mathbb{V}$ and covariance operator \mathcal{C}_{∞} .

We will first prove that $\|\mu_n - \mu_{\infty}\| \rightarrow 0$. To this end, it is enough to show that $\{\mu_n\}$ is relatively compact, since the convergence of the finite-dimensional distributions implies $\langle \mu_n, f \rangle \rightarrow \langle \mu_{\infty}, f \rangle$.

Let the \tilde{X}_n 's be independent copies of the X_n 's and observe that

$$(Y_n, Z_n) := (X_n - \tilde{X}_n, X_n + \tilde{X}_n) \xrightarrow{d} (Y_{\infty}, Z_{\infty}). \quad (\text{S.5.17})$$

Notice that by using the symmetry of the Gaussian distribution

$$Z_n \stackrel{d}{=} 2\mu_n + Y_n. \quad (\text{S.5.18})$$

By Prokhorov's theorem, the convergence in distribution in (S.5.17) entails the tightness in distribution of both the sequences $\{Z_n\}$ and $\{Y_n\}$. On the other hand (S.5.18) shows that $\{2\mu_n + Y_n\}$ is also tight. This implies that $\{\mu_n\}$ is relatively compact. Indeed, by tightness, there exist compact sets \mathcal{K}_1 and \mathcal{K}_2 , such that $\mathbb{P}(Y_n \in \mathcal{K}_1) \geq 2/3$ and $\mathbb{P}(2\mu_n + Y_n \in \mathcal{K}_2) \geq 2/3$. This implies that

$$\begin{aligned} \frac{1}{3} &\leq \mathbb{P}(Y_n \in \mathcal{K}_1) + \mathbb{P}(2\mu_n + Y_n \in \mathcal{K}_2) - 1 \\ &\leq \mathbb{P}(Y_n \in \mathcal{K}_1, 2\mu_n + Y_n \in \mathcal{K}_2) \leq \mathbb{P}(2\mu_n \in \mathcal{K}_2 - \mathcal{K}_1), \end{aligned}$$

which means the deterministic sequence $\{\mu_n\}$ belongs to the compact set $2^{-1}(\mathcal{K}_2 - \mathcal{K}_1)$ with positive probability. Hence, $\{\mu_n\} \subset 2^{-1}(\mathcal{K}_2 - \mathcal{K}_1)$ is a relatively compact sequence completing the proof of the convergence $\|\mu_n - \mu_\infty\| \rightarrow 0$.

We next show that $\|\mathcal{C}_n - \mathcal{C}_\infty\|_{\text{tr}} \rightarrow 0$ and without loss of generality suppose that $\mu_n = \mu_\infty = 0$ so that $\mathcal{C}_n = \mathbb{E}[X_n \otimes X_n]$. Fix a CONS $\{e_i\}$ of \mathbb{V} and define the projection operators

$$\Pi_m := \sum_{i=1}^m e_i \otimes e_i \quad \text{and} \quad \tilde{\Pi}_m := \sum_{i=m+1}^{\infty} e_i \otimes e_i.$$

Since $X_n \xrightarrow{d} X_\infty$, $\|X_n\| \xrightarrow{d} \|X_\infty\|$ and since \mathbb{V} is separable it follows (Prokhorov) that $\{X_n\}$ is tight. Hence, by Corollary S.5.4, $\{\|X_n\|^2\}$ is uniformly integrable and the convergence $\|X_n\| \xrightarrow{d} \|X_\infty\|$ implies

$$\|\mathcal{C}_n\|_{\text{tr}} = \mathbb{E}[\|X_n\|^2] \rightarrow \|\mathcal{C}_\infty\|_{\text{tr}} = \mathbb{E}[\|X_\infty\|^2], \quad \text{as } n \rightarrow \infty. \quad (\text{S.5.19})$$

Fix an arbitrary $\epsilon > 0$ and pick $m = m(\epsilon)$ large enough so that

$$\|\tilde{\Pi}_m \mathcal{C} \tilde{\Pi}_m\|_{\text{tr}} < \epsilon.$$

Recall that every self-adjoint trace-class operator \mathcal{A} can be decomposed as follows

$$\mathcal{A} = \mathcal{A}^+ - \mathcal{A}^-,$$

where \mathcal{A}^\pm are self-adjoint, positive and such that $\mathcal{A}^+ \mathcal{A}^- = \mathcal{A}^- \mathcal{A}^+ = 0$. Thus, $(\mathcal{A} \mathcal{A}^*)^{1/2} = \mathcal{A}^+ + \mathcal{A}^-$ and consequently,

$$\|\mathcal{A}\|_{\text{tr}} = \text{trace}(\mathcal{A}^+) + \text{trace}(\mathcal{A}^-) \equiv \|\mathcal{A}^+\|_{\text{tr}} + \|\mathcal{A}^-\|_{\text{tr}}.$$

Therefore, for the self-adjoint trace-class operator $\mathcal{C}_n - \mathcal{C}_\infty$, we have

$$\|\mathcal{C}_n - \mathcal{C}_\infty\|_{\text{tr}} = \text{trace}((\mathcal{C}_n - \mathcal{C}_\infty)^+) + \text{trace}((\mathcal{C}_n - \mathcal{C}_\infty)^-). \quad (\text{S.5.20})$$

Notice that for every positive self-adjoint operator \mathcal{A} , we have

$$\begin{aligned} \|\mathcal{A}\|_{\text{tr}} &= \text{trace}(\mathcal{A}) \\ &= \text{trace}(\Pi_m \mathcal{A} \Pi_m) + \text{trace}(\tilde{\Pi}_m \mathcal{A} \tilde{\Pi}_m) \\ &= \|\Pi_m \mathcal{A} \Pi_m\|_{\text{tr}} + \|\tilde{\Pi}_m \mathcal{A} \tilde{\Pi}_m\|_{\text{tr}}. \end{aligned} \quad (\text{S.5.21})$$

Thus, by writing $\mathcal{A}_n^\pm := (\mathcal{C}_n - \mathcal{C}_\infty)^\pm$, in view of (S.5.20), we obtain

$$\begin{aligned} \|\mathcal{C}_n - \mathcal{C}_\infty\|_{\text{tr}} &= \text{trace}(\Pi_m \mathcal{A}_n^+ \Pi_m) + \text{trace}(\tilde{\Pi}_m \mathcal{A}_n^+ \tilde{\Pi}_m) \\ &\quad + \text{trace}(\Pi_m \mathcal{A}_n^- \Pi_m) + \text{trace}(\tilde{\Pi}_m \mathcal{A}_n^- \tilde{\Pi}_m). \end{aligned} \quad (\text{S.5.22})$$

Since $X_n \xrightarrow{d} X_\infty$, by Lemma S.5.8, we have that $\|\mathcal{C}_n - \mathcal{C}_\infty\|_{\text{op}} \rightarrow 0$. On the other hand, since $\text{span}\{e_1, \dots, e_m\}$ is finite-dimensional, Lemma S.5.7 applied to $\mathcal{A} := \mathcal{C}_n - \mathcal{C}_\infty$ yields

$$\|\Pi_m (\mathcal{C}_n - \mathcal{C}_\infty)^\pm \Pi_m\|_{\text{tr}} \leq m \|\mathcal{C}_n - \mathcal{C}_\infty\|_{\text{op}} \rightarrow 0, \quad (\text{S.5.23})$$

as $n \rightarrow \infty$.

On the other hand, since $0 \leq (\mathcal{C}_n - \mathcal{C}_\infty)^+ \leq \mathcal{C}_n$, for the positive operators $(\mathcal{C}_n - \mathcal{C}_\infty)^+$ and \mathcal{C}_n , we obtain

$$\|\tilde{\Pi}_m(\mathcal{C}_n - \mathcal{C}_\infty)^+ \tilde{\Pi}_m\|_{\text{tr}} \leq \|\tilde{\Pi}_m \mathcal{C}_n \tilde{\Pi}_m\|_{\text{tr}} = \text{trace}(\tilde{\Pi}_m \mathcal{C}_n \tilde{\Pi}_m). \quad (\text{S.5.24})$$

Similarly, $0 \leq (\mathcal{C}_n - \mathcal{C}_\infty)^- \leq \mathcal{C}$, and hence

$$\|\tilde{\Pi}_m(\mathcal{C}_n - \mathcal{C}_\infty)^- \tilde{\Pi}_m\|_{\text{tr}} \leq \|\tilde{\Pi}_m \mathcal{C}_\infty \tilde{\Pi}_m\|_{\text{tr}} = \text{trace}(\tilde{\Pi}_m \mathcal{C}_\infty \tilde{\Pi}_m). \quad (\text{S.5.25})$$

The last two bounds imply that

$$\|\tilde{\Pi}_m(\mathcal{C}_n - \mathcal{C}_\infty)^\pm \tilde{\Pi}_m\|_{\text{tr}} \leq \|\tilde{\Pi}_m \mathcal{C}_n \tilde{\Pi}_m\|_{\text{tr}} + \|\tilde{\Pi}_m \mathcal{C}_\infty \tilde{\Pi}_m\|_{\text{tr}} \leq 3\epsilon, \quad (\text{S.5.26})$$

for all sufficiently large n . Indeed, $\|\tilde{\Pi}_m \mathcal{C}_\infty \tilde{\Pi}_m\|_{\text{tr}} < \epsilon$, by the choice of m . Whereas, by (S.5.19) and (S.5.21), we have

$$\begin{aligned} \|\tilde{\Pi}_m \mathcal{C}_n \tilde{\Pi}_m\|_{\text{tr}} &= \|\mathcal{C}_n\|_{\text{tr}} - \|\Pi_m \mathcal{C}_n \Pi_m\|_{\text{tr}} \\ &\rightarrow \|\mathcal{C}_\infty\|_{\text{tr}} - \|\Pi_m \mathcal{C}_\infty \Pi_m\|_{\text{tr}} = \|\tilde{\Pi}_m \mathcal{C}_\infty \tilde{\Pi}_m\|_{\text{tr}} \leq \epsilon. \end{aligned}$$

This ensures that for all n large $\|\tilde{\Pi}_m \mathcal{C}_n \tilde{\Pi}_m\|_{\text{tr}} \leq 2\epsilon$ and hence (S.5.26) holds.

Since ϵ was arbitrary, in view of (S.5.22), Relations (S.5.23) and (S.5.26) yield $\|\mathcal{C}_n - \mathcal{C}_\infty\|_{\text{tr}} \rightarrow 0$, as $n \rightarrow \infty$. \square

Lemma S.5.7. *Let $e_i \in \mathbb{V}$, $i = 1, \dots, m$ be orthonormal and $\Pi_m = \sum_{i=1}^m e_i \otimes e_i$.*

For every bounded self-adjoint linear operator $\mathcal{A} : \mathbb{V} \rightarrow \mathbb{V}$, we have

$$\|\Pi_m \mathcal{A}^\pm \Pi_m\|_{\text{tr}} \leq m \sup_{\|f\|=1} |\langle f, \mathcal{A} f \rangle|.$$

Proof. This crude bound follows by observing that the self-adjoint operator $\mathcal{B} := \Pi_m \mathcal{A}^\pm \Pi_m$ has at most m non-zero eigenvalues, which are all bounded above in absolute value by the spectral norm of \mathcal{A} . \square

Lemma S.5.8. *Let X_n and X be zero-mean Gaussian \mathbb{V} -valued vectors with covariance operators \mathcal{C}_n and \mathcal{C} , respectively. If $X_n \xrightarrow{d} X$, then*

$$\|\mathcal{C}_n - \mathcal{C}\|_{\text{op}} \equiv \sup_{\|f\|=1} |\langle f, (\mathcal{C}_n - \mathcal{C}) f \rangle| \rightarrow 0,$$

as $n \rightarrow \infty$.

Proof. By Skorokhod's representation, we can define X_n, X on the same probability space where $X_n \xrightarrow{a.s.} X$. Notice that for $a, b \in \mathbb{C}$, we have

$$\left| |a|^2 - |b|^2 \right| = \left| \Re(a - b) \overline{(a + b)} \right| \leq |a - b| |a + b|.$$

By applying this inequality to $a := \langle X_n, f \rangle$ and $b := \langle X, f \rangle$, for all $f \in E$, $\|f\| = 1$, we obtain

$$\begin{aligned} \left| \langle f, \mathcal{C}_n f \rangle - \langle f, \mathcal{C} f \rangle \right| &= \left| \mathbb{E} |\langle X_n, f \rangle|^2 - \mathbb{E} |\langle X, f \rangle|^2 \right| \\ &\leq \mathbb{E} \left| |\langle X_n, f \rangle|^2 - |\langle X, f \rangle|^2 \right| \\ &\leq \mathbb{E} |\langle X_n - X, f \rangle \langle X_n + X, f \rangle| \\ &\leq (\mathbb{E} \|X_n - X\|^2 \mathbb{E} \|X_n + X\|^2)^{1/2}, \end{aligned} \quad (\text{S.5.27})$$

where the last bound follows from the Cauchy-Schwartz inequality.

The fact that $\{\|X_n - X\|\}$ and $\{\|X_n + X\|\}$ are tight, in view of Corollary S.5.4, implies the uniform integrability of both $\{\|X_n - X\|^2\}$ and $\{\|X_n + X\|^2\}$. This, since $\|X_n - X\| \xrightarrow{a.s.} 0$, shows that the right-hand side of (S.5.27) vanishes, which completes the proof. \square

S.5.4. Hilbert space valued Gaussian processes.

Let $\{X(t), t \in T\} \subset \mathcal{L}^2(\mathbb{V})$ be a second-order stochastic process taking values in the Hilbert space \mathbb{V} , i.e., $\mathbb{E}\|X(t)\|^2 < \infty$. Then, as seen above (cf Lemma S.2.2), its cross covariance operator

$$\mathcal{C}_X(t, s) := \mathbb{E}[X(t) \otimes X(s)], \quad t, s \in T, \quad (\text{S.5.28})$$

is well-defined and takes values in \mathbb{T} . Recall that the latter expectation is understood in the sense of Bochner in the Banach space $\mathbb{T} := \mathbb{T}(\mathbb{V})$ of trace-class operators on \mathbb{V} equipped with the trace norm $\|\cdot\|_{\text{tr}}$. The next definition is a natural extension of a classic concept (see also Definition 4.12).

Definition S.5.9. Fix an arbitrary index set T . A function $\mathcal{C} : T \times T \mapsto \mathbb{T}(\mathbb{V})$, is said to be (completely) positive definite, if for all $f_i \in E$, $t_i \in T$, $i = 1, \dots, n$ and $n \in \mathbb{N}$,

$$\sum_{i,j} \langle f_i, \mathcal{C}(t_i, t_j) f_j \rangle \geq 0. \quad (\text{S.5.29})$$

Remark S.5.4. By taking $(t_1, t_2) := (t, s)$, $(f_1, f_2) = (f, \mathfrak{ig})$, for some (any) $f, g \in \mathbb{V}$, and $(f_1, f_2) = (f, g)$ and applying Relation (S.5.29) twice, one obtains that

$$\langle f, \mathcal{C}(t, s) g \rangle = \langle f, \mathcal{C}(s, t)^* g \rangle, \quad t, s \in T,$$

and hence $\mathcal{C}(t, s) = \mathcal{C}(s, t)^*$. That is, positive definite functions are necessarily Hermitian.

It is straightforward to see that every cross covariance function as in (S.5.28) is positive definite in the sense of the above definition. Naturally, the converse is also true and formally shown next using Gaussian processes. Recall Definition S.5.1.

Definition S.5.10. A \mathbb{V} -valued stochastic process $\{X(t), t \in T\}$ is said to be Gaussian, if $\sum_{i=1}^n \langle f_i, X(t_i) \rangle$ is a (complex) Gaussian random variable, for all $f_i \in \mathbb{V}$, $t_i \in T$, $i = 1, \dots, n$.

Proposition S.5.11. A function $\mathcal{C} : T \times T \rightarrow \mathbb{T}(\mathbb{V})$ is positive definite in the sense of Definition S.5.9, if and only if

$$\mathcal{C}(t, s) = \mathbb{E}[X(t) \otimes X(s)], \quad t, s \in T$$

for some second-order \mathbb{V} -valued stochastic process $X = \{X(t), t \in T\}$. In this case, the process X can be taken to be zero-mean Gaussian.

Proof. The ‘if’ part is immediate, by observing that

$$\mathbb{E} \left| \sum_{i=1}^n \langle f_i, X(t_i) \rangle \right|^2 = \sum_{i,j=1}^n \langle f_i, \mathcal{C}(t_i, t_j) f_j \rangle,$$

where $\mathcal{C}(t, s) = \mathbb{E}[X(t) \otimes X(s)]$, $t, s \in T$.

To prove the ‘only if’ part, consider the direct-sum Hilbert space $\mathbb{V}^{\oplus n} = \{(f_i)_{i=1}^n, f_i \in \mathbb{V}\}$, which is equipped with the inner product

$$\langle f, g \rangle_n := \sum_{i=1}^n \langle f_i, g_i \rangle.$$

For any $\{t_i\}_{i=1}^n \subset T$, define the operator $C_n := (\mathcal{C}(t_i, t_j))_{n \times n}$ acting on $\mathbb{V}^{\oplus n}$ as follows:

$$\langle f, C_n g \rangle_n := \sum_{i,j=1}^n \langle f_i, \mathcal{C}(t_i, t_j) g_j \rangle,$$

for all $f = (f_i)_{i=1}^n$ and $g = (g_i)_{i=1}^n$ with $f_i, g_i \in \mathbb{V}$. Relation (S.5.29) implies that C_n is self-adjoint and positive. Therefore, to establish that $C_n \in \mathbb{T}_+(\mathbb{V}^{\oplus n})$, it is enough to show that its trace is finite. Let $\{e_j\}$ be a CONS of \mathbb{V} and define the elements in $\mathbb{V}^{\oplus n}$:

$$\psi_{1,j} = (e_j, 0, \dots, 0), \psi_{2,j} = (0, e_j, 0, \dots, 0), \dots, \psi_{n,j} := (0, \dots, 0, e_j),$$

for $j \in \mathbb{N}$. Then, it is easy to see that $\{\psi_{i,j}, i = 1, \dots, n, j \in \mathbb{N}\}$ is a CONS of the direct-sum Hilbert space $\mathbb{V}^{\oplus n}$. Since C_n is self-adjoint and positive, we have

$$\begin{aligned} \|C_n\|_{\text{tr}} &= \text{trace}(C_n) = \sum_{i=1}^n \sum_{j \in \mathbb{N}} \langle \psi_{i,j}, C_n \psi_{i,j} \rangle_n \\ &= \sum_{i=1}^n \sum_{j \in \mathbb{N}} \langle e_j, \mathcal{C}(t_i, t_i) e_j \rangle = \sum_{i=1}^n \text{trace}(\mathcal{C}(t_i, t_i)) \\ &= \sum_{i=1}^n \|\mathcal{C}(t_i, t_i)\|_{\text{tr}} < \infty, \end{aligned}$$

where the terms in the last sum are finite since $\mathcal{C}(t_i, t_i) \in \mathbb{T}_+(\mathbb{V})$ are positive trace-class operators.

The spectral theorem for the positive trace-class operator C_n yields the decomposition

$$C_n = \sum_{j \in \mathbb{N}} \sigma_j \varphi_j \otimes \varphi_j,$$

where $\sigma_j \geq 0$ and $\{\varphi_j\}$ is an orthonormal set of eigenvectors of the operator C_n . Now, taking independent standard normal random variables $Z_j, j \in \mathbb{N}$, we define

$$X_n := \sum_j \sqrt{\sigma_j} Z_j \varphi_j. \quad (\text{S.5.30})$$

Since $\sum_j \sigma_j = \|C_n\|_{\text{tr}} < \infty$, by the Itô-Nisio Theorem, the above series converges in norm with probability one and defines a zero-mean Gaussian vector in $\mathbb{V}^{\oplus n}$. Noting that $X_n = (X_n(i))_{i=1}^n$, where $X_n(i) \in \mathbb{V}$, and using the fact that

$$e \otimes f = (e_i \otimes f_j)_{i,j=1}^n,$$

for all $e = (e_i)_{i=1}^n$ and $f = (f_i)_{i=1}^n$ in $E^{\oplus n}$, it readily follows that the collection of Gaussian vectors $\{X_n(i), i = 1, \dots, n\}$ has cross covariance operators

$$\mathcal{C}(t_i, t_j) = \mathbb{E}[X_n(i) \otimes X_n(j)], \quad i, j = 1, \dots, n.$$

The joint distribution of the components of the vector X_n defines a probability distribution F_{t_1, \dots, t_n} on the *product* Hilbert space \mathbb{V}^n . Since the law of the Gaussian vectors X_n in (S.5.30) is (by definition) completely determined by $\{(\sigma_j, \varphi_j), j = 1, \dots, n\}$, it is easy to see that the family $\{F_{t_1, \dots, t_n}, t_i \in T\}$ is consistent and hence the generalized version of the Kolmogorov existence theorem applies (see, e.g., Theorem 5.16 in Kallenberg [1997]). Thus, on a suitable probability space one can have an \mathbb{V} -valued stochastic process $\{X(t), t \in T\}$ with finite-dimensional distributions F_{t_1, \dots, t_n} , and in particular such that $\mathbb{E}[X(t) \otimes X(s)] = \mathcal{C}(t, s), t, s \in T$. \square

Caution. Suppose that $C : T \times T \rightarrow \mathbb{T}(\mathbb{V})$ is such that $\sum_{i,j=1}^n a_i \bar{a}_j \mathcal{C}(t_i, t_j)$ is a positive self-adjoint operator, for all $a_i \in \mathbb{C}$ and $t_i \in T$, $i = 1, \dots, n$. One may be tempted to conclude that $\{\mathcal{C}(t, s)\}$ is then positive definite in the sense of Definition S.5.9. This is not the case in general according to the following classic example borrowed from the literature of C^* -algebras. (See, however, Definitions 4.1, 4.3, and Corollary 4.4 in the main paper.)

Example S.5.2. Consider the simple case $T = \{1, 2\}$ and $\mathbb{V} = \mathbb{R}^2$, where $\mathcal{C}(i, j) = C_{i,j}$, $i, j \in T$ are as follows

$$C := \begin{pmatrix} C_{1,1} & C_{1,2} \\ C_{2,1} & C_{2,2} \end{pmatrix} := \left(\begin{array}{cc|cc} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ \hline 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right).$$

It is easy to see that

$$\sum_{i,j=1}^2 a_i \bar{a}_j C_{i,j} = \begin{pmatrix} |a_1|^2 & \bar{a}_1 a_2 \\ a_1 \bar{a}_2 & |a_2|^2 \end{pmatrix},$$

which is positive semidefinite. However, since $\det(C) = -1$ the matrix C is not positive semidefinite.

Appendix S.6: On the support of operator self-similar IRF_k's.

In this section, we focus on zero-mean second-order operator self-similar IRF_k's taking values in a separable Hilbert space \mathbb{V} . We provide miscellaneous results about their supports and operator exponents complementing Section 5 of the main paper.

Definition S.6.1. For a second-order IRF_k process Y , we write

$$\text{supp}(Y) := \overline{\text{span}} \left(\bigcup_{\lambda \in \Lambda_k} \text{Im}(\mathcal{C}_Y(\lambda, \lambda)) \right),$$

where $\text{Im}(\mathcal{C}_Y(\lambda, \lambda))$ stands for the range of the cross covariance operator $\mathcal{C}_Y(\lambda, \mu) = \mathbb{E}[Y(\lambda) \otimes Y(\mu)]$ and $\overline{\text{span}}$ denotes the closure of the linear span. The closed subspace $\text{supp}(Y)$ of \mathbb{V} will be referred to as the support of Y . If $\text{supp}(Y) = \mathbb{V}$, then Y will be referred to as proper.

Since for the self-adjoint positive operator $\mathcal{C}_Y(\lambda, \lambda)$, we have $\text{Im}(\mathcal{C}_Y(\lambda, \lambda)) = \text{Ker}(\mathcal{C}_Y(\lambda, \lambda))^\perp$, one can equivalently write

$$\begin{aligned} \text{supp}(Y) &= \left(\bigcap_{\lambda \in \Lambda_k} \text{Ker}(\mathcal{C}_Y(\lambda, \lambda)) \right)^\perp \\ &= \left\{ e \in \mathbb{V} : \mathbb{E}|\langle e, Y(\lambda) \rangle|^2 = 0, \text{ for all } \lambda \in \Lambda_k \right\}^\perp. \end{aligned}$$

The last relation follows from the fact that $\mathbb{E}|\langle e, Y(\lambda) \rangle|^2 = \langle \mathcal{C}_Y e, e \rangle = \langle \mathcal{A} e, \mathcal{A} e \rangle$, where \mathcal{A} is a self-adjoint positive operator such that $\mathcal{C}_Y = \mathcal{A}^2$.

Since for all $\lambda_i \in \Lambda_k$ and $c_i \in \mathbb{C}$, $i = 1, \dots, n$, we have $Y(\lambda) = \sum_{i=1}^n c_i Y(\lambda_i)$, for $\lambda := \sum_{i=1}^n c_i \cdot \lambda_i$, it follows $\text{supp}(Y)$ is the smallest closed linear subspace W of \mathbb{V} such that $Y(\lambda) \in W$, almost surely, for all λ .

By analogy with the notion of a support of an IRF, we shall introduce the notion of a range of a \mathbb{T}_+ -valued measure.

Definition S.6.2. Let χ be a σ -finite \mathbb{T}_+ -valued measure on $\mathcal{B}(\mathbb{R}^d \setminus \{0\})$. For $a \in \mathbb{V}$, consider the σ -finite measure $\mu_a(du) := \langle a, \chi(du)a \rangle$. The range of χ , denoted $\text{Range}(\chi)$ is defined as:

$$\text{Range}(\chi) := \left\{ a \in \mathbb{V} : \mu_a(du) = 0 \right\}^\perp.$$

It is easy to see that $\text{Range}(\chi) = \overline{\text{span}} \cup_{A: \|\chi(A)\|_{\text{tr}} < \infty} \text{Im}(\chi(A))$. The intuition behind the above definition is that $\text{Range}(\chi)$ is the minimal subspace of \mathbb{V} such that the projected measures μ_a are non-trivial. The support of an IRF with trivial polynomial component is precisely the range of its spectral measure.

Proposition S.6.3. Let Y be a continuous second-order IRF $_k$ with spectral representation (χ_k, \mathcal{Q}) as in (4.14) (in the main paper), with trivial polynomial component \mathcal{Q} . Then, with χ_k as in (5.5),

$$\text{supp}(Y) = \text{Range}(\chi_k) \equiv \text{Range}(\chi).$$

Proof. By (4.14), we can write $\mathcal{C}(\lambda, \lambda) = \int_{\mathbb{R}^d} |\hat{\lambda}(u)|^2 \chi_k(du)$, for all $\lambda \in \Lambda_k$, since $\mathcal{Q}(\lambda * \tilde{\lambda}) = 0$ by assumption. Thus, for all $a \in \mathbb{V}$,

$$\langle a, \mathcal{C}(\lambda, \lambda)a \rangle = \int_{\mathbb{R}^d} |\hat{\lambda}(u)|^2 \mu_a(du), \quad (\text{S.6.1})$$

where $\mu_a(du) = \langle a, \chi_k(du)a \rangle$.

Recall that $\text{supp}(Y) = \mathbb{W}^\perp$, where $\mathbb{W} = \bigcap_{\lambda \in \Lambda_k} \text{Ker}(\mathcal{C}(\lambda, \lambda))$. Thus, (S.6.1) implies that $a \in \mathbb{W}$ if and only if $\mu_a(du) = 0$. Indeed, if μ_a is the zero measure, then $a \in \text{Ker}(\mathcal{C}(\lambda, \lambda))$ for all $\lambda \in \Lambda_k$. Conversely, if $\int_{\mathbb{R}^d} |\hat{\lambda}(u)|^2 \mu_a(du) = 0$ for all $\lambda \in \Lambda_k$, then it follows that μ_a is the zero measure. Indeed, this follows from the fact that one can choose a judicious sequence of measures $\lambda_n \in \Lambda_k$, such that $|\hat{\lambda}_n(u)|^2 \rightarrow g(u)$, where $g(u) > 0$ for all $u \neq 0$ (see e.g. Lemma S.4.9). The Fatou Lemma then entails $\int g(u) \mu_a(du) = 0$, and hence $\mu_a = 0$, since $\mu_a(\{0\}) = 0$. This completes the proof since $\mu_a(du) = (1 \wedge \|u\|)^{-2k-2} \langle a, \chi(du)a \rangle$ and the facts that χ_k and χ put no mass at $\{0\}$ entail $\text{Range}(\chi_k) = \text{Range}(\chi)$. \square

Let now $\mathcal{H} : \mathbb{V} \rightarrow \mathbb{V}$ be a bounded linear operator on the Hilbert space \mathbb{V} . Recall that a zero-mean IRF $_k$ Y is said to be second-order \mathcal{H} -self-similar if $\{c^{\mathcal{H}} Y(\lambda)\}$ and $\{Y(c \cdot \lambda)\}$ have the same cross covariance operators, for all $c > 0$. In particular, if Y is Gaussian, this entails the \mathcal{H} -self-similarity of Y .

The operator \mathcal{H} can in principle be rather arbitrary outside the support Y . This is perhaps why a *standard* assumption adopted in the literature on operator self-similarity in the finite-dimensional setting is that Y be *proper*, i.e., $\text{supp}(Y) = \mathbb{V}$ so that its support cannot be confined to a proper linear subspace of \mathbb{V} (see e.g., Didier et al. [2017] and the references therein). The following result allows us to restrict the operator \mathcal{H} to $\text{supp}(Y)$.

Proposition S.6.4. If Y is a continuous second-order \mathcal{H} -self-similar IRF $_k$, then:

- (i) $\mathcal{H}(\text{supp}(Y)) \subset \text{supp}(Y)$,
- (ii) $\overline{\mathcal{H}(\text{supp}(Y))} = \text{supp}(Y)$.

Proof. Let $\mathbb{W} := \bigcap_{\lambda \in \Lambda_k} \text{Ker}(\mathcal{C}_Y(\lambda, \lambda))$, where $\mathcal{C}_Y(\lambda, \mu) = \mathbb{E}[Y(\lambda) \otimes Y(\mu)]$, $\lambda, \mu \in \Lambda_k$. For every $r > 0$, and $\lambda \in \Lambda_k$, the operator self-similarity Relation (5.3) implies that

$$r^{\mathcal{H}} \mathcal{C}_Y(r^{-1} \lambda, r^{-1} \lambda) r^{\mathcal{H}*} = \mathcal{C}_Y(\lambda, \lambda). \quad (\text{S.6.2})$$

This implies that

$$r^{\mathcal{H}^*} \text{Ker}(\mathcal{C}_Y(\lambda, \lambda)) \subset \text{Ker}(\mathcal{C}_Y(r^{-1}\lambda, r^{-1}\lambda)) \subset \mathbb{W}.$$

Since λ was arbitrary, it follows that $r^{\mathcal{H}^*}(\mathbb{W}) \subset \mathbb{W}$. Note however that $r^{\mathcal{H}^*}$ is invertible with bounded inverse $r^{-\mathcal{H}^*}$ such that $r^{-\mathcal{H}^*} r^{\mathcal{H}^*} = r^{\mathcal{H}^*} r^{-\mathcal{H}^*} = \text{I}$, $r > 0$. Therefore,

$$\mathbb{W} = r^{-\mathcal{H}^*}(r^{\mathcal{H}^*} \mathbb{W}) \subset r^{-\mathcal{H}^*}(\mathbb{W}), \quad \text{for all } r > 0.$$

By replacing r with r^{-1} above we obtain $\mathbb{W} \subset r^{\mathcal{H}^*} \mathbb{W}$ and hence

$$r^{\mathcal{H}^*}(\mathbb{W}) = \mathbb{W}, \quad \text{for all } r > 0. \quad (\text{S.6.3})$$

Observe that $\text{supp}(Y) = \mathbb{W}^\perp$. Therefore, Relation (S.6.3) implies $r^{\mathcal{H}} \text{supp}(Y) = \text{supp}(Y)$, for all $r > 0$. Indeed, this can be seen by writing

$$\langle r^{\mathcal{H}} x, y \rangle = \langle x, r^{\mathcal{H}^*} y \rangle \quad (\text{S.6.4})$$

Taking $y \in \mathbb{W}$ and $x \in \text{supp}(Y) \equiv \mathbb{W}^\perp$, we see that $r^{\mathcal{H}^*} y \in \mathbb{W}$ by (S.6.3) and hence $\langle r^{\mathcal{H}} x, y \rangle = 0$, for all $x \in \text{supp}(Y)$ and $y \in \mathbb{W}$. This shows that $r^{\mathcal{H}}(\text{supp}(Y)) \subset \text{supp}(Y)$, for all $r > 0$. As argued above, by applying the inverse $r^{-\mathcal{H}} = (r^{\mathcal{H}})^{-1}$, we get

$$r^{\mathcal{H}}(\text{supp}(Y)) = \text{supp}(Y), \quad \text{for all } r > 0.$$

Now, we argue that this relation entails $\mathcal{H}(\text{supp}(Y)) \subset \text{supp}(Y)$. Observe that $r \mapsto r^{\mathcal{H}}$ is Fréchet differentiable in the Banach space of bounded operators on \mathbb{V} equipped with the operator norm. That is, for all $r > 0$, we have

$$\left\| \frac{(r+h)^{\mathcal{H}} - r^{\mathcal{H}}}{h} - r^{\mathcal{H}-1} \mathcal{H} \right\| \rightarrow 0, \quad \text{as } h \rightarrow 0.$$

By (S.6.4), for every $x \in \text{supp}(Y)$, we have $y_h := h^{-1}((r+h)^{\mathcal{H}} - r^{\mathcal{H}})(x) \in \text{supp}(Y)$, thus the Fréchet differentiability relation above implies that $\|y_h - r^{\mathcal{H}-1} \mathcal{H} x\| \rightarrow 0$, as $h \rightarrow 0$, which, since $\text{supp}(Y)$ is closed entails $r^{\mathcal{H}-1} \mathcal{H} x \in \text{supp}(Y)$. Taking $r = 1$ and considering that $x \in \text{supp}(Y)$ was arbitrary, we obtain $\mathcal{H}(\text{supp}(Y)) \subset \text{supp}(Y)$, which completes the proof of (i). Note that unless \mathcal{H} is invertible with bounded inverse, we cannot readily conclude that $\mathcal{H}(\text{supp}(Y)) = \text{supp}(Y)$.

Since $\text{supp}(Y)$ is a closed linear subspace of \mathbb{V} , it is itself a Hilbert space. Part (i) allows us to consider the restriction $\mathcal{H}_Y := \mathcal{H}|_{\text{supp}(Y)}$ of the operator \mathcal{H} to $\text{supp}(Y)$. To emphasize that the domain is now restricted to $\text{supp}(Y)$, we denote by \mathcal{H}_Y^* the adjoint of \mathcal{H}_Y in $\text{supp}(Y)$.

Note that $\text{Im}(\mathcal{H}_Y)^\perp = \text{Ker}(\mathcal{H}_Y^*)$ and since $\text{Im}(\mathcal{H}_Y)$ is a linear space, it follows that $\overline{\text{Im}(\mathcal{H}_Y)} = \text{Ker}(\mathcal{H}_Y^*)^\perp$. Thus, to prove (ii), it is enough to show that $\text{Ker}(\mathcal{H}_Y^*) = \{0\}$. To this end, suppose that $\mathcal{H}_Y^* x = 0$ for some $x \in \text{supp}(Y)$. We have that $r^{\mathcal{H}_Y^*} x = x$, for all $r > 0$ since

$$r^{\mathcal{H}_Y^*}(x) = e^{\log(r)\mathcal{H}_Y^*} x = x + \sum_{n=1}^{\infty} \frac{\log^n(r)(\mathcal{H}_Y^*)^n x}{n!} = x.$$

This, in view of (S.6.2), implies that

$$\mathbb{E}|\langle x, Y(r\lambda) \rangle|^2 = \langle r^{\mathcal{H}_Y^*} x, \mathcal{C}_Y(\lambda, \lambda) r^{\mathcal{H}_Y^*} x \rangle = \langle x, \mathcal{C}_Y(\lambda, \lambda) x \rangle,$$

for all $r > 0$ and $\lambda \in \Lambda_k$. Now, the continuity of the IRF Y implies that the left-hand side of the last expression vanishes as $r \downarrow 0$. This means that $x \in W$, but $\text{supp}(Y) = W^\perp$ (in \mathbb{V}) and hence $x \in W \cap W^\perp = \{0\}$. This shows that $\text{Ker}(\mathcal{H}_Y^*) = \{0\}$ in $\text{supp}(Y)$ completing the proof of (ii). \square

Contents

1	Introduction.	1
2	Preliminaries.	3
2.1	The spaces $S(\Lambda_k, \mathbb{V})$ and $\check{S}(\Lambda_k, \mathbb{V})$	4
2.2	Scaling actions on \mathbb{V}	7
3	Tangent fields and their properties.	8
3.1	Tangent fields are self-similar.	9
3.2	Tangent fields are intrinsically stationary.	12
4	Spectral theory for Hilbert space valued IRF_k 's.	15
4.1	The Bochner Theorem.	15
4.2	Spectral theory for general IRF_k	18
4.3	Real and complex IRF_k 's: Covariance (ir)reversibility.	21
5	Second order covariance self-similar IRF_k 's.	23
5.1	Linear operator scaling.	23
5.2	Related work and examples.	28
5.3	Scalar actions.	32
A	Proofs and auxiliary results.	34
A.1	Proofs and tools for Section 2.	34
A.2	Proof of Proposition 3.6.	37
A.3	Supplementary results and proofs for Section 3.	39
A.4	Supplemental background and some proofs for Section 4.	43
A.5	Proofs for Section 5.1.	46
A.6	Proofs for Section 5.2.	49
A.7	Proofs for Section 5.3.	52
	Acknowledgements	56
	References	56
S	Supplement	61
S.1	Notation and preliminaries on Hilbert spaces.	61
S.2	Cross covariance operators.	62
S.2.1	The Bochner integral in $(\mathbb{T}, \ \cdot\ _{\text{tr}})$	62
S.2.2	Existence and continuity of cross covariance operators.	63
S.3	Integration Theory for \mathbb{T}_+ -valued measures.	64
S.4	Proofs and auxiliary results for Section 4.	67
S.4.1	Proof of Theorem 4.2 (Bochner).	67
S.4.2	Proof of Theorem 4.11.	74
S.4.3	Proof of Theorem 4.7.	80
S.4.4	Proof of Theorem 4.14.	82
S.5	Gaussian elements in a Hilbert space.	83
S.5.1	Real and complex Hilbert spaces.	83
S.5.2	Fundamentals.	84
S.5.3	Convergence in distribution.	88
S.5.4	Hilbert space valued Gaussian processes.	93
S.6	On the support of operator self-similar IRF_k 's.	95