

# Bernoulli-Fibonacci Polynomials

Oktay K Pashaev and Merve Özvatan

Department of Mathematics  
Izmir Institute of Technology  
Urla-Izmir, 35430, Turkey

December 23, 2021

## Abstract

By using definition of Golden derivative, corresponding Golden exponential function and Fibonomial coefficients, we introduce generating functions for Bernoulli-Fibonacci polynomials and related numbers. Properties of these polynomials and numbers are studied in parallel with usual Bernoulli counterparts. Fibonacci numbers and Golden ratio are intrinsically involved in formulas obtained.

## 1 Introduction

The Bernoulli numbers and related polynomials can be introduced on the basis of the so called  $h$ -calculus [1]. Several generalizations of these numbers in  $q$ -calculus are known and listed in papers [2], [3]. Here we are going to introduce analogs of Bernoulli polynomials in the so called Golden calculus, which is based on the Golden ratio dilatations and Binet formula for Fibonacci numbers [4].

The paper is organized in following way. Firstly, in Section 2 we briefly review main properties of Bernoulli polynomials. In Section 3 we introduce main objects of Golden calculus, as Golden derivative, exponential function and Fibonomials. In the main part, Section 4, we introduce Bernoulli-Fibonacci polynomials and corresponding numbers, formulate and proof related theorems.

## 2 Bernoulli polynomials

Here we review briefly main definitions and properties of Bernoulli polynomials and Bernoulli numbers [1].

**Definition 2.0.1** *The generating function for Bernoulli polynomials is defined by Taylor series expansion,*

$$\frac{ze^{zx}}{e^z - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{z^n}{n!}, \quad (1)$$

where  $B_n(x)$  are the Bernoulli polynomials in  $x$ , for all  $n > 0$ .

Bernoulli numbers are a special values of the Bernoulli polynomials  $B_n(x)$ ,  $b_n = B_n(0)$ . The generating function for Bernoulli numbers is

$$\frac{z}{e^z - 1} = \sum_{n=0}^{\infty} b_n \frac{z^n}{n!}. \quad (2)$$

First few Bernoulli polynomials

$$\begin{aligned} B_0(x) &= 1 \\ B_1(x) &= x - \frac{1}{2} \\ B_2(x) &= x^2 - x + \frac{1}{6} \\ B_3(x) &= x^3 - \frac{3}{2}x^2 + \frac{1}{2}x \\ B_4(x) &= x^4 - 2x^3 + x^2 - \frac{1}{30} \\ B_5(x) &= x^5 - \frac{5}{2}x^4 + \frac{5}{3}x^3 - \frac{1}{6}x \\ B_6(x) &= x^6 - 3x^5 + \frac{5}{2}x^4 - \frac{1}{2}x^2 + \frac{1}{42}, \end{aligned}$$

determine corresponding Bernoulli numbers

$$b_0 = 1, \quad b_1 = -\frac{1}{2}, \quad b_2 = \frac{1}{6}, \quad b_3 = 0, \quad b_4 = -\frac{1}{30}, \quad b_5 = 0, \quad b_6 = \frac{1}{42}.$$

Bernoulli polynomials and numbers poses following properties[1].

**Proposition 2.0.2** *For odd Bernoulli numbers*

$$b_{2n+1} = 0, \quad (3)$$

where  $n = 1, 2, \dots$

**Proposition 2.0.3** *Derivative of  $n^{\text{th}}$  Bernoulli polynomial gives,*

$$\frac{d}{dx} B_n(x) = B'_n(x) = n B_{n-1}(x). \quad (4)$$

**Proposition 2.0.4** *For any  $n \geq 0$ , we have expansion*

$$B_n(x) = \sum_{j=0}^n \binom{n}{j} b_j x^{n-j}. \quad (5)$$

**Proposition 2.0.5** *For any  $n \geq 1$ , we have summation formula*

$$\sum_{j=0}^{n-1} \binom{n}{j} B_j(x) = n x^{n-1}.$$

**Corollary 2.0.6** *From previous proposition at  $x = 0$ , for any  $n \geq 2$  we have*

$$\sum_{j=0}^{n-1} \binom{n}{j} b_j = 0.$$

**Proposition 2.0.7** *For any  $n \geq 2$ ,*

$$B_n(1) = b_n. \quad (6)$$

### 3 Golden Derivative and Golden Exponential

The main ingredients of Golden calculus are described in [4]. It is based on Fibonacci numbers and Binet representation of these numbers:

$$F_n = \frac{\varphi^n - \varphi'^n}{\varphi - \varphi'}, \quad (7)$$

where  $\varphi, \varphi'$  are positive and negative roots of equation  $x^2 - x - 1 = 0$ . These roots explicitly are

$$\varphi = \frac{1 + \sqrt{5}}{2}, \quad \varphi' = \frac{1 - \sqrt{5}}{2} = -\frac{1}{\varphi}. \quad (8)$$

Number  $\varphi$  is known as the Golden ratio or the Golden section. The Golden derivative operator is defined as

$$D_F f(x) = \frac{f(\varphi x) - f(-\frac{x}{\varphi})}{(\varphi + \frac{1}{\varphi})x}. \quad (9)$$

For analytic function  $f(x)$  it can be represented by Binet formula for dilatation operator

$$F_{x \frac{d}{dx}} = \frac{\varphi^{x \frac{d}{dx}} - \varphi'^{x \frac{d}{dx}}}{\varphi - \varphi'}, \quad (10)$$

so that

$$x F_{x \frac{d}{dx}} f(x) = D_F f(x).$$

The Golden exponential function is entire analytic function, defined as

$$e_F^x \equiv \sum_{n=0}^{\infty} \frac{x^n}{F_n!}, \quad (11)$$

and satisfying

$$D_F e_F^{kx} = k e_F^{kx}. \quad (12)$$

The Golden binomial, introduced in [4], can be expanded

$$(x+y)_F^n \equiv (x+y)_{\varphi, -\frac{1}{\varphi}}^n = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_F (-1)^{\frac{k(k-1)}{2}} x^{n-k} y^k \quad (13)$$

in terms of the Golden binomial coefficients, called as Fibonomials,

$$\begin{bmatrix} n \\ k \end{bmatrix}_F = \frac{F_n!}{F_{n-k}! F_k!}, \quad (14)$$

with  $n$  and  $k$  being nonnegative integers,  $n \geq k$ . They satisfy next recursion formulas

$$\begin{bmatrix} n \\ k \end{bmatrix}_F = \left(-\frac{1}{\varphi}\right)^k \begin{bmatrix} n-1 \\ k \end{bmatrix}_F + \varphi^{n-k} \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_F \quad (15)$$

$$= \varphi^k \begin{bmatrix} n-1 \\ k \end{bmatrix}_F + \left(-\frac{1}{\varphi}\right)^{n-k} \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_F. \quad (16)$$

## 4 Bernoulli-Fibonacci polynomials

By using definition of Golden exponential function  $e_F^x$  in (11) we introduce generating function for Bernoulli-Fibonacci polynomials.

**Definition 4.0.1** *Generating function for Bernoulli-Fibonacci polynomials  $B_n^F(x)$  is defined by series expansion,*

$$\frac{z e_F^{zx}}{e_F^z - 1} = \sum_{n=0}^{\infty} B_n^F(x) \frac{z^n}{F_n!}. \quad (17)$$

The Bernoulli-Fibonacci numbers are special values of these polynomials,

$$b_n^F = B_n^F(0), \quad (18)$$

with generating function,

$$\frac{z}{e_F^z - 1} = \sum_{n=0}^{\infty} b_n^F \frac{z^n}{F_n!}. \quad (19)$$

For the first few polynomials we have,

$$\begin{aligned} B_0^F(x) &= 1, \\ B_1^F(x) &= \frac{x}{F_1!} - \frac{1}{F_2!}, \\ B_2^F(x) &= x^2 - x \frac{1}{F_1!} + \frac{1}{F_2!} - \frac{1}{F_3!}, \\ B_3^F(x) &= x^3 - x^2 \frac{F_3!}{(F_2!)^2} + x \left( \frac{F_3!}{F_1!(F_2!)^2} - \frac{1}{F_1!} \right) + \frac{2}{F_2!} - \frac{1}{F_4!} - F_3, \\ B_4^F(x) &= x^4 - x^3 \left( \frac{F_4!}{F_3!F_2!} \right) + x^2 \left( \frac{F_4!}{(F_2!)^3} - \frac{F_4!}{F_2!F_3!} \right) \\ &\quad + x \left( -\frac{F_4!}{F_1!(F_2!)^3} + 2 \frac{F_4!}{F_1!F_2!F_3!} - \frac{1}{F_1!} \right) \\ &\quad + \left( \frac{F_4!}{(F_2!)^4} - 3 \frac{F_4!}{(F_2!)^2F_3!} + 2 \frac{1}{F_2!} + \frac{F_4!}{(F_3!)^2} - \frac{1}{F_5!} \right), \\ B_5^F(x) &= x^5 + x^4 \left( -\frac{F_5!}{F_4!F_2!} \right) + x^3 \left( -\frac{F_5!}{(F_3!)^2} + \frac{F_5!}{F_3!(F_2!)^2} \right) \end{aligned}$$

$$\begin{aligned}
& + x^2 \left( -\frac{F_5!}{F_2!F_4!} + 2\frac{F_5!}{(F_2!)^2F_3!} - \frac{F_5!}{(F_2!)^4} \right) \\
& + x \left( -\frac{1}{F_1!} + \frac{F_5!}{F_1!(F_3!)^2} + 2\frac{F_5!}{F_1!F_2!F_4!} - 3\frac{F_5!}{F_1!(F_2!)^2F_3!} + \frac{F_5!}{F_1!(F_2!)^4} \right) \\
& + \left( -\frac{F_5!}{(F_2!)^5} + 4\frac{F_5!}{(F_2!)^3F_3!} - 3\frac{F_5!}{(F_3!)^2F_2!} - 3\frac{F_5!}{(F_2!)^2F_4!} + 2\frac{F_5!}{F_3!F_4!} + \frac{2}{F_2!} - \frac{F_5!}{F_6!} \right)
\end{aligned}$$

It is noticed that all coefficients of Bernoulli-Fibonacci polynomials are rational Fibonacci numbers.

**Proposition 4.0.2** (*Compare with Proposition (2.0.3)*)

The Golden derivative, applied to Bernoulli-Fibonacci polynomials  $B_n^F(x)$  gives Fibonacci numbers,

$$D_F^x(B_n^F(x)) = F_n B_{n-1}^F(x). \quad (20)$$

**Proof 4.0.3** Taking the derivative from both sides of equation (17) gives,

$$\begin{aligned}
D_F^x \left( \frac{z e_F^{zx}}{e_F^z - 1} \right) &= D_F^x \left( \sum_{n=0}^{\infty} B_n^F(x) \frac{z^n}{F_n!} \right) \Rightarrow \\
\frac{z D_F^x(e_F^{zx})}{e_F^z - 1} &= D_F^x \left( B_0^F(x) + B_1^F(x) \frac{z}{F_1!} + B_2^F(x) \frac{z^2}{F_2!} + \dots \right)
\end{aligned}$$

In the l.h.s., Golden derivative can be calculated from equation (12). In the r.h.s. we use

$$D_F^x(B_0^F(x)) = D_F^x(1) = 0.$$

Then,

$$\begin{aligned}
z \cdot \frac{z e_F^{zx}}{e_F^z - 1} &= \sum_{k=1}^{\infty} D_F^x(B_k^F(x)) \frac{z^k}{F_k!} \Rightarrow \\
z \sum_{n=0}^{\infty} B_n^F(x) \frac{z^n}{F_n!} &= \sum_{k=1}^{\infty} D_F^x(B_k^F(x)) \frac{z^k}{F_k!} \Rightarrow \\
\sum_{n=0}^{\infty} B_n^F(x) \frac{z^{n+1}}{F_n!} &= \sum_{k=1}^{\infty} D_F^x(B_k^F(x)) \frac{z^k}{F_k!}.
\end{aligned}$$

In the r.h.s., by shifting  $k - 1 = n$  we have

$$\begin{aligned}\sum_{n=0}^{\infty} B_n^F(x) \frac{z^{n+1}}{F_n!} &= \sum_{n=0}^{\infty} D_F^x \left( B_{n+1}^F(x) \right) \frac{z^{n+1}}{F_{n+1}!} \Rightarrow \\ \sum_{n=0}^{\infty} B_n^F(x) \frac{z^{n+1}}{F_n!} &= \sum_{n=0}^{\infty} D_F^x \left( B_{n+1}^F(x) \right) \frac{1}{F_{n+1}} \frac{z^{n+1}}{F_n!}.\end{aligned}$$

From equality of these two series, finally we get

$$B_n^F(x) = \frac{D_F^x \left( B_{n+1}^F(x) \right)}{F_{n+1}}.$$

From Bernoulli-Fibonacci polynomials, due to  $b_n^F = B_n^F(0)$ , we find Bernoulli-Fibonacci numbers as,

$$b_0^F = 1, \quad b_1^F = -1, \quad b_2^F = \frac{1}{2}, \quad b_3^F = -\frac{1}{3}, \quad b_4^F = \frac{3}{10}, \quad b_5^F = -\frac{5}{8}, \quad b_6^F = \frac{101}{39}, \dots$$

**Proposition 4.0.4** (Compare with Proposition (2.0.4))

Bernoulli-Fibonacci Polynomials can be represented by sum,

$$B_n^F(x) = \sum_{j=0}^n \left[ \begin{matrix} n \\ j \end{matrix} \right]_F b_j^F x^{n-j}. \quad (21)$$

**Proof 4.0.5** It is evident from this formula that  $B_n^F(0) = b_n^F$ . So it suffices to show that  $B_n^F(x)$  are satisfying equation (20):

$$\begin{aligned}D_F^x \left[ B_n^F(x) \right] \quad (21) &= D_F^x \left[ \sum_{j=0}^n \left[ \begin{matrix} n \\ j \end{matrix} \right]_F b_j^F x^{n-j} \right] \\ &= \sum_{j=0}^{n-1} \left[ \begin{matrix} n \\ j \end{matrix} \right]_F b_j^F D_F^x \left( x^{n-j} \right) \\ &= \sum_{j=0}^{n-1} \left[ \begin{matrix} n \\ j \end{matrix} \right]_F b_j^F F_{n-j} x^{n-j-1} \\ &= \sum_{j=0}^{n-1} \frac{F_n!}{F_{n-j}! F_j!} b_j^F F_{n-j} x^{n-j-1}\end{aligned}$$

$$\begin{aligned}
&= \sum_{j=0}^{n-1} \frac{F_n!}{F_{n-j-1}! F_j!} b_j^F x^{n-j-1} \\
&= F_n \sum_{j=0}^{n-1} \frac{F_{n-1}!}{F_{n-j-1}! F_j!} b_j^F x^{n-j-1} \\
(21) &= F_n B_{n-1}^F(x).
\end{aligned}$$

**Proposition 4.0.6** (Compare with Proposition (2.0.5))

For  $n \geq 1$ , Bernoulli-Fibonacci polynomials can be calculated recursively by formula:

$$\sum_{l=0}^{n-1} \begin{bmatrix} n \\ l \end{bmatrix}_F B_l^F(x) = F_n x^{n-1}.$$

**Proof 4.0.7** Starting from the generating function,

$$\frac{z e_F^{zx}}{e_F^z - 1} = \sum_{n=0}^{\infty} B_n^F(x) \frac{z^n}{F_n!},$$

multiplying it by  $e_F^z$ ;

$$\frac{z e_F^{zx} e_F^z}{e_F^z - 1} = \sum_{n=0}^{\infty} B_n^F(x) e_F^z \frac{z^n}{F_n!},$$

and taking difference, we get

$$\begin{aligned}
\frac{z e_F^{zx}}{e_F^z - 1} (e_F^z - 1) &= \sum_{n=0}^{\infty} (B_n^F(x) e_F^z - B_n^F(x)) \frac{z^n}{F_n!} \Rightarrow \\
z e_F^{zx} &= \sum_{n=0}^{\infty} (B_n^F(x) e_F^z - B_n^F(x)) \frac{z^n}{F_n!} \Rightarrow \\
D_F^x (e_F^{zx}) &= \sum_{n=0}^{\infty} (B_n^F(x) e_F^z - B_n^F(x)) \frac{z^n}{F_n!} \Rightarrow \\
D_F^x \left( \sum_{n=0}^{\infty} \frac{(zx)^n}{F_n!} \right) &= \sum_{n=0}^{\infty} (B_n^F(x) e_F^z - B_n^F(x)) \frac{z^n}{F_n!} \Rightarrow \\
D_F^x \left( \sum_{n=1}^{\infty} \frac{z^n x^n}{F_n!} \right) &= \sum_{n=0}^{\infty} (B_n^F(x) e_F^z - B_n^F(x)) \frac{z^n}{F_n!} \Rightarrow \\
\sum_{n=1}^{\infty} \frac{z^n D_F^x (x^n)}{F_n!} &= \sum_{n=0}^{\infty} (B_n^F(x) e_F^z - B_n^F(x)) \frac{z^n}{F_n!} \Rightarrow \\
\sum_{n=1}^{\infty} \frac{z^n F_n x^{n-1}}{F_n!} &= \sum_{l=0}^{\infty} B_l^F(x) e_F^z \frac{z^l}{F_l!} - \sum_{n=0}^{\infty} B_n^F(x) \frac{z^n}{F_n!}.
\end{aligned}$$



For the first sum in r.h.s. of this equation,

$$\begin{aligned}
\sum_{l=0}^{\infty} B_l^F(x) e_F^z \frac{z^l}{F_l!} &= \sum_{l=0}^{\infty} \sum_{k=0}^{\infty} B_l^F(x) \frac{z^k}{F_k!} \frac{z^l}{F_l!} = \sum_{l=0}^{\infty} \sum_{k=0}^{\infty} B_l^F(x) \frac{z^{k+l}}{F_k! F_l!} \\
&= \sum_{l=0}^{\infty} \sum_{k=0}^{\infty} B_l^F(x) \frac{z^{k+l}}{F_{k+l}!} \\
k+l=n &= \sum_{n=0}^{\infty} \frac{1}{F_n!} \sum_{l=0}^n B_l^F(x) \frac{z^n F_n!}{F_{n-l}! F_l!} \\
&= \sum_{n=0}^{\infty} \frac{z^n}{F_n!} \left( \sum_{l=0}^n \begin{bmatrix} n \\ l \end{bmatrix}_F B_l^F(x) \right).
\end{aligned}$$

After substituting this,

$$\begin{aligned}
\sum_{n=1}^{\infty} \frac{z^n F_n x^{n-1}}{F_n!} &= \sum_{n=0}^{\infty} \frac{z^n}{F_n!} \left( \sum_{l=0}^n \begin{bmatrix} n \\ l \end{bmatrix}_F B_l^F(x) \right) - \sum_{n=0}^{\infty} B_n^F(x) \frac{z^n}{F_n!} \Rightarrow \\
\sum_{n=1}^{\infty} \frac{z^n F_n x^{n-1}}{F_n!} &= B_0^F(x) \begin{bmatrix} 0 \\ 0 \end{bmatrix}_F + \sum_{n=1}^{\infty} \frac{z^n}{F_n!} \left( \sum_{l=0}^n \begin{bmatrix} n \\ l \end{bmatrix}_F B_l^F(x) \right) - B_0^F(x) - \sum_{n=1}^{\infty} B_n^F(x) \frac{z^n}{F_n!} \Rightarrow \\
\sum_{n=1}^{\infty} \frac{z^n}{F_n!} F_n x^{n-1} &= \sum_{n=1}^{\infty} \frac{z^n}{F_n!} \left( \sum_{l=0}^n \begin{bmatrix} n \\ l \end{bmatrix}_F B_l^F(x) - B_n^F(x) \right).
\end{aligned}$$

Then, we have

$$\sum_{l=0}^n \begin{bmatrix} n \\ l \end{bmatrix}_F B_l^F(x) - B_n^F(x) = F_n x^{n-1},$$

where  $n \geq 1$ . Cancelling  $n^{\text{th}}$  terms,

$$\sum_{l=0}^{n-1} \begin{bmatrix} n \\ l \end{bmatrix}_F B_l^F(x) + B_n^F(x) \begin{bmatrix} n \\ n \end{bmatrix}_F - B_n^F(x) = F_n x^{n-1},$$

finally, we get desired equality,

$$\sum_{l=0}^{n-1} \begin{bmatrix} n \\ l \end{bmatrix}_F B_l^F(x) = F_n x^{n-1}.$$

**Corollary 4.0.8** (*Compare with Corollary (2.0.6)*)

*From previous proposition, by taking  $x = 0$ , we have the formula, allowing us to compute Bernoulli-Fibonacci numbers inductively,*

$$\sum_{j=0}^{n-1} \begin{bmatrix} n \\ j \end{bmatrix}_F b_j^F = 0, \quad (22)$$

where  $n \geq 2$ .

**Proof 4.0.9** *Proof will be done by equating expansions of  $B_n^F(x) + F_n x^{n-1}$ , obtained in two ways. By the first way we obtain,*

$$B_n^F(x) + F_n x^{n-1} = x^n + \sum_{j=2}^n \begin{bmatrix} n \\ j \end{bmatrix}_F b_j^F x^{n-j}, \quad (23)$$

and by the second one,

$$B_n^F(x) + F_n x^{n-1} \equiv H_n(x) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_F B_{n-k}^F(x). \quad (24)$$

*These equations are derived in Appendix. Comparison of these two expansions gives,*

$$\sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_F B_{n-k}^F(x) = x^n + \sum_{j=2}^n \begin{bmatrix} n \\ j \end{bmatrix}_F b_j^F x^{n-j}.$$

*Then we have,*

$$\sum_{k=0}^{n-1} \begin{bmatrix} n \\ k \end{bmatrix}_F B_{n-k}^F(x) + B_0^F(x) = x^n + \sum_{j=2}^n \begin{bmatrix} n \\ j \end{bmatrix}_F b_j^F x^{n-j} \Rightarrow$$

$$x^n - 1 = \sum_{k=0}^{n-1} \begin{bmatrix} n \\ k \end{bmatrix}_F B_{n-k}^F(x) - \sum_{j=2}^n \begin{bmatrix} n \\ j \end{bmatrix}_F b_j^F x^{n-j} \Rightarrow$$

$$x^n - 1 = \sum_{k=0}^{n-1} \begin{bmatrix} n \\ k \end{bmatrix}_F B_{n-k}^F(x) - \left( \begin{bmatrix} n \\ 2 \end{bmatrix}_F b_2^F x^{n-2} + \dots + \begin{bmatrix} n \\ n-1 \end{bmatrix}_F b_{n-1}^F x + \begin{bmatrix} n \\ n \end{bmatrix}_F b_n^F \right).$$

For  $x = 0$  it gives,

$$-1 = \sum_{k=0}^{n-1} \begin{bmatrix} n \\ k \end{bmatrix}_F b_{n-k}^F - b_n^F.$$

After simplifying,

$$-1 = \sum_{k=1}^{n-1} \begin{bmatrix} n \\ k \end{bmatrix}_F b_{n-k}^F + \begin{bmatrix} n \\ 0 \end{bmatrix}_F b_n^F - b_n^F,$$

we get

$$\sum_{k=1}^{n-1} \begin{bmatrix} n \\ k \end{bmatrix}_F b_{n-k}^F + 1 = 0.$$

By using symmetry of Fibonomial coefficients, we rewrite it as,

$$\sum_{k=1}^{n-1} \begin{bmatrix} n \\ n-k \end{bmatrix}_F b_{n-k}^F + 1 = 0,$$

then by denoting  $n - k = j$ ,

$$\sum_{j=n-1}^1 \begin{bmatrix} n \\ j \end{bmatrix}_F b_j^F + 1 = 0.$$

Substituting  $1 = b_0^F$ ,

$$\sum_{j=1}^{n-1} \begin{bmatrix} n \\ j \end{bmatrix}_F b_j^F + \begin{bmatrix} n \\ 0 \end{bmatrix}_F b_0^F = 0$$

we obtain desired result

$$\sum_{j=0}^{n-1} \begin{bmatrix} n \\ j \end{bmatrix}_F b_j^F = 0.$$

**Proposition 4.0.10** (Compare with Proposition (2.0.7))

For any  $n \geq 2$ , Bernoulli-Fibonacci polynomials and corresponding numbers satisfy the following equation,

$$B_n^F(1) = b_n^F. \tag{25}$$

**Proof 4.0.11** *Starting with,*

$$\begin{aligned}
\frac{e_F^{zx} - 1}{e_F^z - 1} &= \left( \frac{e_F^{zx}}{e_F^z - 1} - \frac{1}{e_F^z - 1} \right) = \left( \frac{z}{e_F^z - 1} \frac{e_F^{zx}}{z} - \frac{z}{e_F^z - 1} \right) \frac{1}{z} \\
(17) &= \left( \sum_{n=0}^{\infty} B_n^F(x) \frac{z^n}{F_n!} - \frac{z}{e_F^z - 1} \right) \frac{1}{z} \\
(19) &= \left( \sum_{n=0}^{\infty} B_n^F(x) \frac{z^n}{F_n!} - \sum_{n=0}^{\infty} b_n^F \frac{z^n}{F_n!} \right) \frac{1}{z} \\
&= \sum_{n=0}^{\infty} (B_n^F(x) - b_n^F) \frac{z^{n-1}}{F_n!}.
\end{aligned}$$

*Thus, we have,*

$$\begin{aligned}
\frac{e_F^{zx} - 1}{e_F^z - 1} &= (B_0^F(x) - b_0^F) \frac{z^{-1}}{F_0!} + (B_1^F(x) - b_1^F) \frac{1}{F_1!} + \sum_{n=2}^{\infty} (B_n^F(x) - b_n^F) \frac{z^{n-1}}{F_n!} \Rightarrow \\
\frac{e_F^{zx} - 1}{e_F^z - 1} &= (1 - 1) \frac{z^{-1}}{F_0!} + (x - 1 - (-1)) \frac{1}{F_1!} + \sum_{n=2}^{\infty} (B_n^F(x) - b_n^F) \frac{z^{n-1}}{F_n!} \Rightarrow \\
\frac{e_F^{zx} - 1}{e_F^z - 1} &= x + \sum_{n=2}^{\infty} (B_n^F(x) - b_n^F) \frac{z^{n-1}}{F_n!}. \tag{26}
\end{aligned}$$

*From this expansion, for  $x = 1$ ,*

$$\begin{aligned}
\frac{e_F^z - 1}{e_F^z - 1} &= 1 + \sum_{n=2}^{\infty} (B_n^F(1) - b_n^F) \frac{z^{n-1}}{F_n!} \\
&\Rightarrow \sum_{n=2}^{\infty} (B_n^F(1) - b_n^F) \frac{z^{n-1}}{F_n!} = 0.
\end{aligned}$$

*Then, the coefficients at every power of  $z$  are zero and*

$$B_n^F(1) - b_n^F = 0, \tag{27}$$

*for  $n = 2, 3, 4, \dots$*

## 5 Acknowledgements

This work is supported by TUBITAK grant 116F206.

## References

- [1] Kac, V. and Cheung, P., 2002. Quantum Calculus, Springer, New York.
- [2] Nalci S and Pashaev O K 2012. q-Bernoulli numbers and zeros of q-Sine function. *arXiv:1202.2265v1* **math.QA**
- [3] Ismail Mourad E H and Mansour Zeinab S I 2019. q-Analogs of Lidstone expansion theorem, two-point Taylor expansion theorem, and Bernoulli polynomials. *Analysis and Applications* **17** pp.853-895
- [4] Pashaev O K and Nalci S 2012. Golden quantum oscillator and Binet-Fibonacci calculus. *J Phys A:Math Theor* **45** 015303

## 6 Appendix

### 6.1 Getting $B_n^F(x) + F_n x^{n-1}$ in two ways

Our aim is to get equation;

$$B_n^F(x) + F_n x^{n-1} = H_n(x) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_F B_{n-k}^F(x).$$

Starting with,

$$\begin{aligned} B_n^F(x) + F_n x^{n-1} &= H_n(x) \Rightarrow \\ \sum_{n=0}^{\infty} B_n^F(x) \frac{z^n}{F_n!} + \sum_{n=0}^{\infty} F_n x^{n-1} \frac{z^n}{F_n!} &= \sum_{n=0}^{\infty} H_n(x) \frac{z^n}{F_n!} \Rightarrow \\ \sum_{n=0}^{\infty} H_n(x) \frac{z^n}{F_n!} - \sum_{n=0}^{\infty} B_n^F(x) \frac{z^n}{F_n!} &= \sum_{n=0}^{\infty} F_n x^{n-1} \frac{z^n}{F_n!}. \end{aligned}$$

For the r.h.s., we get

$$z e_F^{zx} = D_F^x(e_F^{zx}) = D_F^x \left( \sum_{n=1}^{\infty} \frac{x^n z^n}{F_n!} \right) = \sum_{n=1}^{\infty} \frac{F_n x^{n-1} z^n}{F_n!} = \sum_{n=0}^{\infty} \frac{F_n x^{n-1} z^n}{F_n!}.$$

Then, we have;

$$\begin{aligned} \sum_{n=0}^{\infty} H_n(x) \frac{z^n}{F_n!} - \sum_{n=0}^{\infty} B_n^F(x) \frac{z^n}{F_n!} &= z e_F^{zx} \Rightarrow \\ \sum_{n=0}^{\infty} H_n(x) \frac{z^n}{F_n!} &= z e_F^{zx} + \sum_{n=0}^{\infty} B_n^F(x) \frac{z^n}{F_n!} \Rightarrow \end{aligned}$$

$$\begin{aligned}
\sum_{n=0}^{\infty} H_n(x) \frac{z^n}{F_n!} &= z e_F^{zx} + \frac{z e_F^{zx}}{e_F^z - 1} \Rightarrow \\
\sum_{n=0}^{\infty} H_n(x) \frac{z^n}{F_n!} &= z e_F^{zx} \left( 1 + \frac{1}{e_F^z - 1} \right) = z e_F^{zx} \frac{e_F^z}{e_F^z - 1} \Rightarrow \\
\sum_{n=0}^{\infty} H_n(x) \frac{z^n}{F_n!} &= \sum_{n=0}^{\infty} B_n^F(x) \frac{z^n}{F_n!} \cdot e_F^z \Rightarrow \\
\sum_{n=0}^{\infty} H_n(x) \frac{z^n}{F_n!} &= \sum_{n=0}^{\infty} B_n^F(x) \frac{z^n}{F_n!} \cdot \sum_{k=0}^{\infty} \frac{z^k}{F_k!} \Rightarrow \\
\sum_{n=0}^{\infty} H_n(x) \frac{z^n}{F_n!} &= \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{B_n^F(x)}{F_n!} \frac{z^{n+k}}{F_k!} \Rightarrow \\
\sum_{n=0}^{\infty} H_n(x) \frac{z^n}{F_n!} \quad (n+k=N) &= \sum_{N=0}^{\infty} \sum_{k=0}^N \frac{B_{N-k}^F(x)}{F_{N-k}!} \frac{z^N}{F_k!} \left( \frac{F_N!}{F_N!} \right) \Rightarrow \\
\sum_{n=0}^{\infty} H_n(x) \frac{z^n}{F_n!} &= \sum_{N=0}^{\infty} \frac{z_N}{F_N!} \sum_{k=0}^N \frac{F_N!}{F_{N-k}! F_k!} B_{N-k}^F(x) \Rightarrow \\
\sum_{n=0}^{\infty} H_n(x) \frac{z^n}{F_n!} &= \sum_{N=0}^{\infty} \frac{z_N}{F_N!} \sum_{k=0}^N \begin{bmatrix} N \\ k \end{bmatrix}_F B_{N-k}^F(x) \Rightarrow \\
\sum_{n=0}^{\infty} H_n(x) \frac{z^n}{F_n!} \quad (N=n) &= \sum_{n=0}^{\infty} \frac{z_n}{F_n!} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_F B_{n-k}^F(x).
\end{aligned}$$

By equating two series we get,

$$H_n(x) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_F B_{n-k}^F(x). \quad (28)$$

Following in another way we show that,

$$B_n^F(x) + F_n x^{n-1} = H_n(x) = x^n + \sum_{j=2}^n \begin{bmatrix} n \\ j \end{bmatrix}_F b_j^F x^{n-j}.$$

Starting with,

$$\begin{aligned}
B_n^F(x) + F_n x^{n-1} \quad (21) &= \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix}_F b_j^F x^{n-j} + F_n x^{n-1} \\
&= \begin{bmatrix} n \\ 0 \end{bmatrix}_F b_0^F x^n + \begin{bmatrix} n \\ 1 \end{bmatrix}_F b_1^F x^{n-1} + \sum_{j=2}^n \begin{bmatrix} n \\ j \end{bmatrix}_F b_j^F x^{n-j} + F_n x^{n-1}
\end{aligned}$$

$$\begin{aligned}
&= x^n - \frac{F_n!}{F_{n-1}!F_1!}x^{n-1} + \sum_{j=2}^n \begin{bmatrix} n \\ j \end{bmatrix}_F b_j^F x^{n-j} + F_n x^{n-1} \\
&= x^n + \sum_{j=2}^n \begin{bmatrix} n \\ j \end{bmatrix}_F b_j^F x^{n-j},
\end{aligned}$$

thus, we obtain

$$B_n^F(x) + F_n x^{n-1} = H_n(x) = x^n + \sum_{j=2}^n \begin{bmatrix} n \\ j \end{bmatrix}_F b_j^F x^{n-j}. \quad (29)$$