

LUIGI ACCARDI

*Centro Vito Volterra, Università di Roma "Tor Vergata",
Roma I-00133, Italy*

first_accardi@volterra.uniroma2.it

ABDESSATAR SOUISSI

¹ *Department of Accounting, College of Business Management
Qassim University, Ar Rass, Saudi Arabia*

² *Preparatory Institute for Scientific and Technical Studies La Marsa,
Carthage University, Tunisia*

second_a.souaissi@qu.edu.sa, abdessattar.souissi@ipest.rnu.tn

EI GHETEB SOUEIDY

*Department of Mathematics, Nouakchott University,
Nouakchott, Mauritania.*

third_elkotobmedsalem@gmail.com

On the limit of superposition states

February 8, 2022

Contents

1	Introduction	3
2	Preliminaries	4
3	Schur kernels	5
4	The structure of superposition states	9
5	The infinite volume limiting state	12
6	Homogeneous superposition states	18
7	Mixing property for superposition states	21

Abstract

In this paper, we study the structure of a family of superposition states on tensor algebras. The correlation functions of the considered states are described through a new kind of positive definite kernels valued in the dual of C^* -algebras, so-called Schur kernels. Mainly, we show the existence of the limiting state of a net of superposition states over an arbitrary locally finite graph. Furthermore, we show that this limiting state enjoys a mixing property and an α -mixing property in the case of the multi-dimensional integer lattice \mathbf{Z}^{ν} .

Keywords: Schur kernels, superposition states, mixing property, graphs.

1 Introduction

It is well known that, in statistical mechanics and quantum field theory superposition states allow the description of the subsystem relative to bounded regions in terms of its degrees of freedom [20].

The superposition states in quantum theory are convex combinations of many states. In many important cases, a complex quantum system is identified with a graph $V = (V, E)$ with vertex set V and edge set E . To each bounded region $\Lambda \subset V$, we associated a finite dimensional Hilbert space $\mathcal{H}_\Lambda := \bigotimes_{x \in \Lambda} \mathcal{H}_x$. The local superposition states under consideration are states on the algebras $\mathcal{B}(\mathcal{H}_\Lambda)$ indexed by bounded regions Λ has the form

$$\psi_\Lambda(\cdot) := \langle \Psi_\Lambda, (\cdot) \Psi_\Lambda \rangle \quad (1)$$

for some state vector $\Psi_\Lambda := \sum_{i \in I} \bigotimes_{x \in \Lambda} h_{x,i} \in \mathcal{H}_\Lambda$ where I is a finite set and $\{h_{x,i}\}_{i \in I} \subset \mathcal{H}_x$.

The present paper mainly deals with the structure of limiting state of a net of superposition states. Namely, we describe the correlation functions in terms of Schur multiplication. This leads to a class of positive definite Kernels with values in the dual of C^* -algebras, so-called *Schur kernels*.

Notice that positive definite kernels valued in C^* -algebras were studied in connection with operator-valued cocycles, see for instance [6]. It turns out that, the Schur multiplication appears naturally in the explicit form of the states under consideration. These results might open new perspective to the study of ergodic properties of this kind of states, such as entanglement and mixing.

In [9], it was proven that the states of the form (1) have rich entanglement properties. Furthermore, it was shown that they define a class of quantum Markov chains in the sense of [2], [3].

The finite volume states (1) are states on type I factors, with finite degree of freedom. In local quantum physics [17], it was proven that these states play a key role in the description of quantum systems with an infinite degree of freedom, naturally appearing in quantum fields theory and quantum statistical mechanics. Moreover, some relevance of such kind of states in quantum information theory, quantum random walks and quantum hidden Markov models were clarified (see for instance [8], [10], [11], [18], [16]).

It is crucial to know certain ergodic properties of the limiting state of the superposition states (1), which may reveal some physical characteristics of the system. With this aim, we prove a mixing property for the inhomogeneous

superposition states under consideration in the case of the multidimensional integer lattice. Moreover, we exhibit a definition for α -mixing property and show that the limit of superposition states is indeed α -mixing.

A main problem in entanglement theory, is to find an easily applicable criteria to distinguish quantum states according to their degree of entanglement, or at least to discriminate entangled from non entangled states. The first step to attack this problem is to have a sufficiently detailed description of the class of superposition states under consideration.

The paper is organized as follows. We devote Section 3 to Schur kernels and their properties. In section 4, we describe the correlations functions of finite volume superposition states through a class of Schur kernels. Section 5 is devoted to the existence (Theorem 4) of the infinite volume-limiting state. Section 6 is devoted to the homogeneous case. Section 7 is devoted to the mixed property for the inhomogeneous limiting state. This will be the aim of some futures works.

2 Preliminaries

In this section we introduce some basic notions and notations. Let $d \in \mathbf{N}$, the space of $d \times d$ matrices over the complex field \mathbf{C} will be denoted $M_d(\mathbf{C})$. Let us consider two C^* -algebras \mathcal{A} and \mathcal{B} . Recall that a map $\Phi : \mathcal{A} \rightarrow \mathcal{B}$ is called *positive* if it maps \mathcal{A}^+ into \mathcal{B}^+ . A linear map Φ is called k -positive if the mapping $\Phi \otimes Id_k$ is positive from $M_k(\mathcal{A}) \equiv \mathcal{A} \otimes M_k(\mathbf{C})$ into $M_k(\mathcal{B}) \equiv \mathcal{B} \otimes M_k(\mathbf{C})$. If Φ is k -positive for each integer k then it is called *completely positive* (CP). It is well known that a linear map Φ is completely positive if and only if

$$\sum_{i,j=1}^k b_i^* \Phi(a_i^* a_j) b_j \geq 0; \quad a_1, \dots, a_k \in \mathcal{A}, b_1, \dots, b_k \in \mathcal{B} \quad (2)$$

for every $k \in \mathbf{N}$. The reader is referred to [19] for a detailed discussion about CP maps. The dual space of \mathcal{B} will be denoted by \mathcal{B}^* (see for instance [21]). Let $G = (V, E)$ be a simple graph. To each vertex $x \in V$ we associate a finite dimensional Hilbert space \mathcal{H}_x . Let $\Lambda \subset V$ be a bounded region. To which we associate a Hilbert space of observable \mathcal{H}_Λ and the associated algebra of observable

$$\mathcal{B}_\Lambda := \mathcal{B}(\mathcal{H}_\Lambda) \equiv \bigotimes_{x \in \Lambda} \mathcal{B}(\mathcal{H}_x) =: \bigotimes_{x \in \Lambda} \mathcal{B}_x. \quad (3)$$

Let \mathcal{H} be a Hilbert space. The adjoint of a vector $\xi \in \mathcal{H}$ is defined by $\xi^*(\eta) = \langle \eta, \xi \rangle$. The notation $\eta\xi^*$ refers to the rank-one operator over \mathcal{H} defined by $\eta\xi^*(\psi) = \langle \psi, \xi \rangle \eta$.

Throughout this paper, by $\text{Tr}_{\mathcal{H}}$, we denote the trace on the algebra $\mathcal{B}(\mathcal{H})$ of all bounded operators over \mathcal{H} . Notice that for any orthonormal basis $\{e_j\}_{1 \leq j \leq d}$ of the Hilbert space \mathcal{H}

$$\text{Tr}_{\mathcal{H}}(a) = \sum_{j=1}^d \langle ae_j, e_j \rangle, \quad \forall a \in \mathcal{B}(\mathcal{H})$$

where $d = \dim(\mathcal{H})$.

The Schur multiplication of two matrices $A = (a_{i,j}), B = (b_{i,j}) \in M_d(\mathbf{C})$ is defined by

$$A \diamond B_{i,j} = a_{i,j} b_{i,j}. \quad (4)$$

The Schur algebra $(M_d(\mathbf{C}), \diamond)$ is abelian, with identity the matrix J with all entries are equal 1.

3 Schur kernels

This section we introduce a new kind of positive definite kernels valued in dual of C^* -algebras. These kernels will be referred as Schur kernels. We will see later that these kernels play a crucial role in the description of correlation functions of the superposition states.

Definition 1 Let I be a finite set and \mathcal{B} a C^* -algebra. A map

$$E : I \times I \rightarrow \mathcal{B}^*$$

is called **Schur kernel** associated to the pair (I, \mathcal{B}) if the family of linear functionals $\{E_{j,i}\}_{j,i \in I}$ enjoys the following property.

Property S: for any $n \in \mathbf{N}$ and any function

$$b : h \in \{1, \dots, n\} \rightarrow b_h \in \mathcal{B}$$

the complex valued kernel on $I \times \{1, \dots, n\}$ given by

$$K_{(j,h),(i,k)} := E_{j,i}(b_h^* b_k) \quad ; \quad (j,h), (i,k) \in I \times \{1, \dots, n\} \quad (5)$$

is positive definite.

Remark. The Schur kernels in the sense of Definition 1 are valued in the dual of the C^* -algebra. Property (S) implies that for each fixed element $b \in \mathcal{B}$ the map

$$K_b : (i, j) \in I^2 \rightarrow E_{i,j}(b^*b)$$

is a complex valued positive definite kernel. And for any fixed $i \in I$ the map

$$K_i : (h, k) \in \{1, 2, \dots, n\}^2 \rightarrow E_{i,i}(b_h^*b_k)$$

is also a complex positive definite kernel.

Theorem 1 Let I be a **finite set** ($d_I = |I|$) and \mathcal{B} a C^* -algebra. A map

$$E : I \times I \rightarrow \mathcal{B}^*$$

is a (I, \mathcal{B}) -**Schur kernel** iff the map \hat{E} defined by

$$\hat{E} : b \in \mathcal{B} \rightarrow \hat{E}(b) := (E_{i,j}(b))_{i,j \in I} \in M_{d_I}(\mathbf{C}) \quad (6)$$

is **completely positive**.

Proof. Necessity. If \hat{E} is completely positive then for any $a_1, a_2, \dots, a_n \in M_{d_I}(\mathbf{C})$ and $b_1, b_2, \dots, b_n \in \mathcal{B}$

$$\sum_{k,j=0}^n a_j^* \hat{E}(b_j^*b_k) a_k \geq 0 \Leftrightarrow \sum_{k,j=0}^n a_j^* [E \ . \ . \ . (b_j^*b_k)] a_k \geq 0$$

where $\hat{E}(b_j^*b_k) = [E \ . \ . \ . (b_j^*b_k)]$.

By taking $a_j = (a_{k,l}^{(j)})_{k,l \in I}$, we have for each $X = (x_1, \dots, x_d) \in M_{1 \times d_I}(\mathbf{C})$ at

$$\begin{aligned} & X \left(\sum_{k,j=0}^n a_j^* [E \ . \ . \ . (b_j^*b_k)] a_k \right) X^* \geq 0 \\ \Leftrightarrow & \sum_{k,j=0}^n X \left(\sum_{s=1}^d \sum_{l=1}^d \bar{a}_{l,p}^{(j)} E_{l,s}(b_j^*b_k) a_{s,q}^{(k)} \right)_{p,q \in I} X^* \geq 0 \\ \Leftrightarrow & \sum_{k,j=0}^n \left(\sum_{t,s,l,r=1}^d x_t \bar{a}_{l,t}^{(j)} E_{l,s}(b_j^*b_k) a_{s,r}^{(k)} x_r^* \right) \geq 0 \end{aligned} \quad (7)$$

For $X = E_1 := (1, 0, \dots, 0) \in M_{1 \times d_I}(\mathbf{C})$ one gets

$$\sum_{t,s,l,r=1}^d x_t \bar{a}_{l,t}^{(j)} E_{l,s}(b_j^* b_k) a_{s,r}^{(k)} x_r^* = \sum_{s,l=1}^d \bar{a}_{l,1}^{(j)} E_{l,s}(b_j^* b_k) a_{s,1}^{(k)}.$$

By taking $n = d$ and denote $z_{j,l} = a_{l,1}^{(j)}$, then (7) becomes

$$\sum_{(l,j),(s,k) \in I \times I} \bar{z}_{j,l} E_{l,s}(b_j^* b_k) z_{k,s} \geq 0$$

Since $a_{l,1}^{(j)}$ can be arbitrary chosen, one concludes that E is a Schur kernel. **Sufficiency.** Conversely, if E is a Schur kernel, let $a_1, a_2, \dots, a_n \in M_{d_I}(\mathbf{C})$ and $b_1, b_2, \dots, b_n \in \mathcal{B}$ be such that

$$\sum_{k,j=0}^n a_j^* \hat{E}(b_j^* b_k) a_k = \sum_{k,j=0}^n a_j^* [E \ . \ . \ (b_j^* b_k)] a_k$$

Let $X = (x_1, \dots, x_d) \in M_{1 \times d_I}(\mathbf{C})$, then

$$\begin{aligned} X \left(\sum_{k,j=0}^n a_j^* [E_{i,m}(b_j^* b_k)]_{i,m} a_k \right) X^* &= \sum_{k,j=0}^n \left(\sum_{s,l=1}^d \left(\sum_{t=1}^d x_t \bar{a}_{l,t}^{(j)} \right) E_{l,s}(b_j^* b_k) \left(\sum_{r=1}^d a_{s,r}^{(k)} \bar{x}_r \right) \right) \\ &= \sum_{k,j=0}^n \left(\sum_{s,l=1}^d \bar{z}_{j,l} E_{l,s}(b_j^* b_k) z_{k,s} \right) = \sum_{(j,l),(s,k) \in I \times I} \bar{z}_{j,l} E_{l,s}(b_j^* b_k) z_{k,s} \\ &= \sum_{(j,l),(s,k) \in I \times I} \bar{z}_{j,l} K_{(j,l),(k,s)} z_{k,s} \geq 0 \end{aligned}$$

where $z_{k,s} = \sum_{r=1}^d a_{s,r}^{(k)} \bar{x}_r$. \square

Definition 2 Let I be a set, \mathcal{B}, \mathcal{C} two C^* -algebras and $E_{\mathcal{B}; \cdot}, E_{\mathcal{C}; \cdot}$ two Schur kernels respectively associated to the pairs $(I, \mathcal{B}), (I, \mathcal{C})$. The **tensor product of the kernels** $E_{\mathcal{B}; \cdot}$ and $E_{\mathcal{C}; \cdot}$, denoted $E_{\mathcal{B}; \cdot} \otimes E_{\mathcal{C}; \cdot}$, is defined by

$$(j, i) \in I \times I \mapsto E_{\mathcal{B}; j, i} \otimes E_{\mathcal{C}; j, i} \in \mathcal{B}^* \otimes \mathcal{C}^* \equiv (\mathcal{B} \otimes \mathcal{C})^*$$

where $E_{\mathcal{B}; j, i} \otimes E_{\mathcal{C}; j, i}$ is the usual tensor product of duals of C^* -algebras.

Remark. Definition 2, as well as Theorem 2 below, do not depend on the choice of the C^* -cross norm used to define $\mathcal{B} \otimes \mathcal{C}$.

Theorem 2 In the notations and assumptions of Definition 2, $E_{\mathcal{B}; \cdot} \otimes E_{\mathcal{C}; \cdot}$ is an $(I, \mathcal{B} \otimes \mathcal{C})$ -Schur kernel.

Proof. We have to prove that for any $n \in \mathbf{N}$ and any function

$$X : h \in \{1, \dots, n\} \rightarrow X_h \in \mathcal{B} \otimes \mathcal{C}$$

the complex valued kernel on $I \times \{1, \dots, n\}$ given by

$$K_{(j,h),(i,k)} := E_{x;j,i}(X_h^* X_k) \quad ; \quad (j,h), (i,k) \in I \times \{1, \dots, n\} \quad (8)$$

is positive definite. Since elements of the form

$$X_k = \sum_{m \in F} b_{k,m} \otimes c_{k,m}$$

where F is a finite set, are dense in $\mathcal{B} \otimes \mathcal{C}$, it is sufficient to prove the statement for these elements. For any function $z : (j,h) \in I \times \{1, \dots, n\} \rightarrow z_{j,h} \in \mathbf{C}$, one has

$$\begin{aligned} & \sum_{(j,h),(i,k) \in I \times \{1, \dots, n\}} \bar{z}_{j,h} K_{(j,h),(i,k)} z_{i,k} = \sum_{(j,h),(i,k) \in I \times \{1, \dots, n\}} \bar{z}_{j,h} E_{j,i}(X_{x,h}^* X_{x,k}) z_{i,k} \\ &= \sum_{(j,h),(i,k) \in I \times \{1, \dots, n\}} \bar{z}_{j,h} E_{j,i} \left(\left(\sum_{m' \in F} b_{h,m'}^* \otimes c_{h,m'}^* \right) \left(\sum_{m \in F} b_{k,m} \otimes c_{k,m} \right) \right) z_{i,k} \\ &= \sum_{(j,h),(i,k) \in I \times \{1, \dots, n\}} \sum_{m', m \in F} \bar{z}_{j,h} E_{j,i} \left((b_{h,m'}^* b_{k,m} \otimes c_{h,m'}^* c_{k,m}) \right) z_{i,k} \\ &= \sum_{(j,h),(i,k) \in I \times \{1, \dots, n\}} \sum_{m', m \in F} \bar{z}_{j,h} E_{\mathcal{B};j,i}(b_{h,m'}^* b_{k,m}) E_{\mathcal{C};j,i}(c_{h,m'}^* c_{k,m}) z_{i,k}. \quad (9) \end{aligned}$$

Now define the new function

$$(i, k, m) \in I \times \{1, \dots, n\} \times F \mapsto z_{i,k,m} := z_{i,k}.$$

Then (9) becomes

$$\sum_{j,i \in I} \sum_{(h,m'), (k,m) \in \{1, \dots, n\} \times F} \bar{z}_{j,h,m'} E_{\mathcal{B};j,i}(b_{h,m'}^* b_{k,m}) E_{\mathcal{C};j,i}(c_{h,m'}^* c_{k,m}) z_{i,k,m}. \quad (10)$$

Since

$$|\{1, \dots, n\} \times F| = n \times |F|$$

one can re-number the pairs $(k, m) \in \{1, \dots, n\} \times F$ so that they become in one-to-one correspondence with the set $\{1, \dots, n|F|\}$. With this numeration, (10) becomes

$$\sum_{(j,h'),(i,k') \in I \times \{1, \dots, n|F|\}} \bar{z}_{j,h'} E_{\mathcal{B};j,i}(b_{h'}^*, b_{k'}) E_{\mathcal{C};j,i}(c_{h'}^*, c_{k'}) z_{i,k'} \geq 0 \quad (11)$$

because, since n in (5) is arbitrary, both kernels $(E_{\mathcal{B};j,i}(b_{h'}^*, b_{k'}))$ and $(E_{\mathcal{C};j,i}(c_{h'}^*, c_{k'}))$ are positive definite, hence their Schur product is by Schur lemma. \square

4 The structure of superposition states

To each $x \in V$, we associate:

- a finite set D_x of **non-zero** vectors in \mathcal{H}_x ,
- a finite set I of cardinality d_I ,
- a map

$$h_{x,(\cdot)} : i \in I \rightarrow h_{x,i} \in D_x$$

and, for $\Lambda \subset_{fin} V$, define the vector

$$\Psi_\Lambda := \sum_{i \in I} \bigotimes_{x \in \Lambda} h_{x,i} \in \bigotimes_{x \in \Lambda} \mathcal{H}_x =: \mathcal{H}_\Lambda \quad (12)$$

and the associated positive linear functional on \mathcal{B}_Λ given by

$$\psi_\Lambda = \langle \Psi_\Lambda, (\cdot) \Psi_\Lambda \rangle = \text{Tr}_{\mathcal{H}_\Lambda}(\Psi_\Lambda \Psi_\Lambda^* (\cdot)). \quad (13)$$

Lemma 1 In the notations above

$$\Psi_\Lambda \Psi_\Lambda^* := \sum_{i,j \in I} \bigotimes_{x \in \Lambda} h_{x,i} h_{x,j}^* \quad (14)$$

and, for

$$\Lambda_1 \supset \bar{\Lambda}$$

$$b_\Lambda \equiv \left(\bigotimes_{x \in \Lambda} b_x \right) \otimes 1_{\Lambda_1 \setminus \Lambda} \in \mathcal{B}_\Lambda \equiv \mathcal{B}_\Lambda \otimes 1_{\Lambda_1 \setminus \Lambda}$$

one has

$$\psi_{\Lambda_1}(b_\Lambda) = \sum_{i,j \in I} \prod_{x \in \Lambda} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^* b_x) \prod_{y \in \Lambda_1 \setminus \Lambda} \text{Tr}_{\mathcal{H}_y} (h_{y,i} h_{y,j}^*). \quad (15)$$

In particular

$$\psi_\Lambda(b_\Lambda) = \sum_{i,j \in I} \prod_{x \in \Lambda} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^* b_x). \quad (16)$$

Proof. (14) follows from (12) by distributivity.

$$\begin{aligned} \psi_{\Lambda_1}(b_\Lambda) &= \text{Tr}_{\mathcal{H}_{\Lambda_1}} \left(\Psi_{\Lambda_1} \Psi_{\Lambda_1}^* \bigotimes_{x \in \Lambda} b_x \right) = \sum_{i,j \in I} \text{Tr}_{\mathcal{H}_{\Lambda_1}} \left(\prod_{x \in \Lambda_1} h_{x,i} h_{x,j}^* \prod_{x \in \Lambda} b_x \right) \\ &= \sum_{i,j \in I} \text{Tr}_{\mathcal{H}_\Lambda} \left(\bigotimes_{x \in \Lambda} h_{x,i} h_{x,j}^* b_x \right) \text{Tr}_{\mathcal{H}_{\Lambda_1 \setminus \Lambda}} \left(\bigotimes_{y \in \Lambda_1 \setminus \Lambda} h_{y,i} h_{y,j}^* \right) \\ &= \sum_{i,j \in I} \prod_{x \in \Lambda} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^* b_x) \prod_{y \in \Lambda_1 \setminus \Lambda} \text{Tr}_{\mathcal{H}_y} (h_{y,i} h_{y,j}^*). \end{aligned}$$

which is (15). And (16) follows from (15) replacing Λ_1 by Λ in ψ_{Λ_1} . \square

Theorem 3 The family of linear functionals

$$E_{x;j,i} : b_x \in \mathcal{B}_x \rightarrow \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^* b_x) \in \mathbf{C} \quad ; \quad x \in V, j, i \in I \quad (17)$$

enjoys **property (S)** (5). Equivalently, for each $x \in V$ the map $E_x : (j, i) \mapsto E_{x;i,j}$ is an (I, \mathcal{B}_x) -Schur kernel.

Proof. In the notation of Definition 1, for any $n \in \mathbf{N}$ and any function $z : (j, h) \in I \times \{1, \dots, n\} \rightarrow z_{j,h} \in \mathbf{C}$, one has

$$\begin{aligned} \sum_{(j,h),(i,k) \in I \times \{1, \dots, n\}} \bar{z}_{j,h} K_{x;(j,h),(i,k)} z_{i,k} &= \sum_{(j,h),(i,k) \in I \times \{1, \dots, n\}} \bar{z}_{j,h} E_{x;j,i} (b_{x,h}^* b_{x,k}) z_{i,k} \\ &= \sum_{(j,h),(i,k) \in I \times \{1, \dots, n\}} E_{x;j,i} (\bar{z}_{j,h} b_{x,h}^* z_{i,k} b_{x,k}) \end{aligned}$$

$$\begin{aligned}
&= \sum_{(j,h),(i,k) \in I \times \{1, \dots, n\}} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^* \bar{z}_{j,h} b_{x,h}^* z_{i,k} b_{x,k}) \\
&= \sum_{(j,h),(i,k) \in I \times \{1, \dots, n\}} \langle h_{x,j}, \bar{z}_{j,h} b_{x,h}^* z_{i,k} b_{x,k} h_{x,i} \rangle \\
&= \sum_{(j,h),(i,k) \in I \times \{1, \dots, n\}} \langle z_{j,h} b_{x,h} h_{x,j}, z_{i,k} b_{x,k} h_{x,i} \rangle \\
&= \left\langle \sum_{(j,h) \in I \times \{1, \dots, n\}} z_{j,h} b_{x,h} h_{x,j}, \sum_{(i,k) \in I \times \{1, \dots, n\}} z_{i,k} b_{x,k} h_{x,i} \right\rangle \\
&= \left\| \sum_{(i,k) \in I \times \{1, \dots, n\}} z_{i,k} b_{x,k} h_{x,i} \right\|^2 \geq 0.
\end{aligned}$$

□

Corollary 1 Let I be a finite set and, in the notation (3), let be given, for each $x \in V$ a Schur kernel $E_x; \cdot$ associated to the pair (I, \mathcal{B}_x) . Then, for each $\Lambda \subset_{fin} V$, the map

$$\hat{E}_\Lambda : \mathcal{B}_\Lambda \rightarrow (M_{d_I}(\mathbf{C}), \diamond)$$

associated to the tensor product kernel $\bigotimes_{x \in \Lambda} E_x; \cdot$ according to Theorem 1 is characterized by

$$\hat{E}_\Lambda \left(\bigotimes_{x \in \Lambda} b_x \right) = \diamond_{x \in \Lambda} \hat{E}_x(b_x) \quad ; \quad b_x \in \mathcal{B}_x, x \in \Lambda \quad (18)$$

where each $\hat{E}_x : b \in \mathcal{B}_x \mapsto (E_{x,i,j}(b)) \in M_{d_I}(\mathbf{C})$ is the map associated to according to Theorem 1.

Proof. From Theorem 2 it follows by induction that $E_\Lambda; \cdot := \bigotimes_{x \in \Lambda} E_x; \cdot$ is a Schur kernel and that the associated map $\hat{E}_\Lambda : \mathcal{B}_\Lambda \rightarrow M_{d_I}(\mathbf{C})$ has the form (18). □

Corollary 2 In the identification of $M_{d_I}(\mathbf{C})$ with $\mathcal{B}(\mathbf{C}^{d_I})$ obtained by fixing a linear basis $e_I \equiv (e_{I,j})_{j \in I}$, for each $\Lambda \subset_{fin} V$, the state ψ_Λ defined by (13) has the form

$$\psi_\Lambda(b_\Lambda) = \langle \hat{e}_I, \left(\diamond_{x \in \Lambda} \hat{E}_x(b_x) \right) \hat{e}_I \rangle \quad ; \quad b_\Lambda := \bigotimes_{x \in \Lambda} b_x, b_x \in \mathcal{B}_x \quad (19)$$

where \hat{E}_x is as in Corollary 1 with E_x given by (17) and

$$\hat{e}_I := \sum_{j \in I} e_{I,j}$$

is the non-normalized maximally entangled vector in the e_I -basis of \mathbf{C}^{d_I} .

Proof. In the above notations

$$\begin{aligned} \langle \hat{e}_I, \left(\diamond_{x \in \Lambda} \hat{E}_x(b_x) \right) \hat{e}_I \rangle &= \sum_{j,i \in I} \langle e_{I,j}, \left(\diamond_{x \in \Lambda} \hat{E}_x(b_x) \right) e_{I,i} \rangle \\ &= \sum_{j,i \in I} \prod_{x \in \Lambda} E_{x;j,i}(b_x) = \sum_{j,i \in I} \prod_{x \in \Lambda} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^* b_x) = \psi_\Lambda(b_\Lambda). \end{aligned}$$

□

Remark. Formula (19) suggests a natural **extension** of the superposition states obtained by replacing the Schur product's kernel in (17) by arbitrary Schur kernels.

5 The infinite volume limiting state

If Λ_α is an increasing net of finite sub-sets in V , we write

$$\Lambda_\alpha \uparrow\uparrow V$$

to mean that the net (Λ_α) definitively absorbs every finite set in V . This implies

$$\bigcup_{\alpha} \Lambda_\alpha = V.$$

The simplified notation

$$\Lambda_1 \uparrow\uparrow V$$

means that Λ_1 denotes the generic element of an increasing net $\Lambda_\alpha \uparrow\uparrow V$.

Lemma 2 The limit

$$\lim_{\Lambda_1 \uparrow\uparrow V} \psi_{\Lambda_1}(b_\Lambda) = \sum_{i,j \in I} \prod_{x \in \Lambda} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^* b_x) \prod_{x \in \Lambda_1 \setminus \Lambda} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^*) \quad (20)$$

exists for all $b_\Lambda \in \mathcal{B}_\Lambda$ if, for each $i, j \in I$, the limit

$$\lim_{\Lambda_1 \uparrow \uparrow V} \prod_{x \in \Lambda_1 \setminus \Lambda} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^*) \quad (21)$$

exists. The converse is true if, for each $x \in V$, the $(h_{x,i})_{i \in I}$ are linearly independent **but not a linear basis of \mathcal{H}_x** .

Proof. The implication (21) \Rightarrow (20) is clear because I is a finite set. Conversely, if the limit (20) exists for all $b_\Lambda \in \mathcal{B}_\Lambda$, the assumption on the vectors $h_{x,i}$ implies that one can fix an $\bar{x} \in \Lambda$ and a pair $\bar{i}, \bar{j} \in I$ and choose the b_x so that

$$\text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^* b_x) = \begin{cases} \delta_{i,\bar{i}} \delta_{j,\bar{j}} c_{ij}, & \text{if } x = \bar{x}, \\ 0, & \text{if } x \neq \bar{x} \end{cases} \quad (22)$$

with $c_{ij} \neq 0$. In fact, since the $(h_{x,i})_{i \in I}$ are not a linear basis of \mathcal{H}_x , for each $x \in V$ there is a non-zero vector h_x orthogonal to all the $(h_{x,i})_{i \in I}$. Choosing $b_x := h_x h_x^*$ for $x \neq \bar{x}$, the second identity in (22) is satisfied. To satisfy the first identity in (22) we use the linear independence of the $h_{\bar{x},k}$. This guarantees that each such vector has the form

$$h_{\bar{x},k} = \hat{h}_{\bar{x},k} + h_{\bar{x},k}^\perp$$

where $\hat{h}_{\bar{x},k}$ is a linear combination of the set $(\hat{h}_{\bar{x},j})_{j \in I \setminus \{k\}}$ and $h_{\bar{x},k}^\perp$ is orthogonal to the linear span of this set. Therefore, with the choice $b_{\bar{x}} := h_{\bar{x},\bar{j}}^\perp (h_{\bar{x},\bar{i}}^\perp)^*$, the first identity in (22) is satisfied.

Lemma 3 If, for each $x \in V$, the $(h_{x,i})_{i \in I}$ are an ortho-normal set, condition (21) is satisfied and the limit positive functional is a state and has the form

$$\lim_{\Lambda_1 \uparrow \uparrow V} \psi_{\Lambda_1}(b_\Lambda) = \sum_{i \in I} \prod_{x \in \Lambda} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,i}^* b_x) =: \hat{\psi}_\Lambda(b_\Lambda). \quad (23)$$

The family $(\hat{\psi}_\Lambda)_{\Lambda \subset_{fin} V}$ is projective.

Proof. The assumption implies that, for each $i, j \in I$, the right hand side of (21) is equal to δ_{ij} . Moreover, for each $i \in I$, $\text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,i}^* (\cdot))$ is a state on

\mathcal{B}_x . Hence $\hat{\psi}_\Lambda$ is a sum of homogeneous product states on \mathcal{B}_Λ . This implies projectivity. \square

Define, for $x \in V$, the **interaction matrix at x** by

$$H_{x,i,j} := \log \left(\text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^*) \right) \iff \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^*) = e^{H_{x,i,j}} \quad ; \quad i, j \in I$$

where \log denotes the principal value logarithm. Then for any $i, j \in I$

$$\lim_{\Lambda_1 \uparrow \uparrow V} \prod_{x \in \Lambda_1 \setminus \Lambda} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^*) = \lim_{\Lambda_1 \uparrow \uparrow V} e^{\sum_{x \in \Lambda_1 \setminus \Lambda} H_{x,i,j}} \quad (24)$$

in the sense that each limit exists iff the other one does and the equality holds.

Lemma 4 Suppose that, for each $i, j \in I$, the limit (24) exists for each $\Lambda \subseteq_{fin} V$ and $\Lambda_0 \subseteq \Lambda$ define:

$$\beta_{\Lambda^c; i, j} := \lim_{\Lambda_1 \uparrow \uparrow V} \prod_{x \in \Lambda_1 \setminus \Lambda} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^*) = \lim_{\Lambda_1 \uparrow \uparrow V} e^{\sum_{x \in \Lambda_1 \setminus \Lambda} H_{x,i,j}}, \quad (25)$$

$$\beta_{\Lambda, \Lambda_0; i, j} := \prod_{x \in \Lambda \setminus \Lambda_0} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^*) = e^{\sum_{x \in \Lambda \setminus \Lambda_0} H_{x,i,j}}. \quad (26)$$

Then the matrices $(\beta_{\Lambda^c; i, j})$ and $(\beta_{\Lambda, \Lambda_0; i, j})$ are positive definite and satisfy

$$\beta_{\Lambda^c; i, j} \beta_{\Lambda, \Lambda_0; i, j} = \beta_{\Lambda_0^c; i, j}. \quad (27)$$

Moreover, the limit (21) exists and is equal to

$$\hat{\psi}_\Lambda(b_\Lambda) := \sum_{i, j \in I} \prod_{x \in \Lambda} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^* b_x) \beta_{\Lambda; i, j} \quad (28)$$

and the family $(\hat{\psi}_\Lambda)$ is projective.

Conversely let, for each $\Lambda \subseteq_{fin} V$ be given a positive definite matrix $(\beta_{\Lambda^c; i, j})$ which satisfies (27) with $(\beta_{\Lambda, \Lambda_0; i, j})$ given by (26). Then, for each $\Lambda \subseteq_{fin} V$, the linear functional (28) is positive and the family $(\hat{\psi}_\Lambda)$ is projective.

Proof. If the limit (24) exists, the fact that the matrices $(\beta_{\Lambda; i, j})$ and $(\beta_{\Lambda, \Lambda_0; i, j})$ are positive definite follows from Schur lemma. Equation (27) follows from (25) and (26). If $\Lambda_0 \subseteq \Lambda$ and $b_\Lambda = b_{\Lambda_0} \otimes 1_{\Lambda \setminus \Lambda_0}$, then (25) and (26) imply

$$\hat{\psi}_\Lambda(b_{\Lambda_0} \otimes 1_{\Lambda \setminus \Lambda_0}) = \sum_{i, j \in I} \prod_{x \in \Lambda} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^* b_x) \beta_{\Lambda^c; i, j}$$

$$\begin{aligned}
&= \sum_{i,j \in I} \prod_{x \in \Lambda_0} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^* b_x) \prod_{x \in \Lambda \setminus \Lambda_0} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^*) \beta_{\Lambda^c; i,j} \\
&= \sum_{i,j \in I} \prod_{x \in \Lambda_0} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^*) \beta_{\Lambda, \Lambda_0; i,j} \beta_{\Lambda^c; i,j} \\
&= \sum_{i,j \in I} \prod_{x \in \Lambda_0} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^*) \beta_{\Lambda_0^c; i,j} = \hat{\psi}_{\Lambda_0}(b_{\Lambda_0}).
\end{aligned}$$

Conversely, if $(\beta_{\Lambda^c; i,j})$ is a positive definite matrix, then by Schur lemma $\hat{\psi}_{\Lambda}(b_{\Lambda})$, defined by the right hand side of (28), is a positive linear functional and the projectivity of the family $(\hat{\psi}_{\Lambda})$ is proved as in the first part of the theorem.

Remark. Lemma 4 suggests a way to construct families of vectors $(\Psi_{\Lambda})_{\Lambda \subset_{fin} V}$ which, in the limit $\Lambda \uparrow V$, produce states on the algebra $\mathcal{A} = \bigotimes_{x \in V} \mathcal{B}_x$. In the following we discuss a way to realize this construction.

Lemma 5 Given any positive definite matrix $T := (T_{i,j})_{i,j \in I}$, if $h = (h_{i,j})_{i,j \in I}$ is any **right square root** of T , in the sense that

$$hh^* = T \tag{29}$$

the column vectors of h

$$h_i := (h_{i,m})_m \quad ; \quad i, m \in I \tag{30}$$

satisfy the relation

$$\langle h_j, h_i \rangle = T_{i,j} \quad ; \quad \forall i, j \in I. \tag{31}$$

Proof. By assumption T is positive definite. Let $h = (h_{i,j})$ be any right square root of T . Define the column vectors of h as in (30). Then

$$\langle h_j, h_i \rangle = \sum_h \overline{h_{j,h}} h_{i,h} = \sum_h h_{i,h} (h^*)_{h,j} = (hh^*)_{i,j} = T_{i,j}. \square$$

Lemma 6 For any Hermitean positive definite matrix $T = (T_{i,j})_{i,j \in I}$ on \mathbf{C}^d , there exists an Hermitean matrix $H = (H_{i,j})$ on \mathbf{C}^d such that T has the form

$$T = e^H = e^{U^* D_H U} \tag{32}$$

where U is a unitary matrix and

$$D_H = \text{diag}(h_1^0, \dots, h_d^0) \quad ; \quad h_j^0 \in \mathbf{R}, \quad j \in \{1, \dots, d\}. \quad (33)$$

Moreover, for any isometry W on \mathbf{C}^d , the matrix

$$h := e^{H/2} W^* \quad (34)$$

is a right square root of T and the entries of H satisfy the inequality

$$|H_{i,j}| \leq \text{Tr}(|D_H|) \quad ; \quad \forall i, j \in I \quad (35)$$

where

$$|D_H| := \text{diag}(|h_1^0|, \dots, |h_d^0|).$$

Proof. Since T is Hermitean positive definite, by the spectral theorem it has the form

$$T = U^* \text{diag}(t_1, \dots, t_d) U \quad ; \quad t_j \in \mathbf{R}_+, \quad j \in \{1, \dots, d\}$$

where U is a unitary matrix. Defining

$$D_H := \text{diag}(h_1^0, \dots, h_d^0) := \text{diag}(\log(t_1), \dots, \log(t_d)).$$

$$H := U^* D_H U. \quad (36)$$

One sees that (32) and (33) hold. And that h , defined by (34), is a right square root of T follows from

$$hh^* := e^{H/2} W^* W e^{H/2} = e^H = T.$$

Finally (32) implies that, for any $i, j \in I$.

$$H_{i,j} = \sum_{m,n} (U^*)_{i,m} h_m^0 \delta_{m,n} U_{n,j} = \sum_m \overline{U_{m,i}} h_m^0 U_{m,j}.$$

Therefore

$$|H_{i,j}| \leq \sum_m |\overline{U_{m,i}}| |h_m^0| |U_{m,j}| \leq \sum_m |h_m^0| = \text{Tr}(|D_H|).$$

which is (35).

Theorem 4 For each $x \in V$ fix:

– the dimension of the fiber independent of x

$$d := \dim(\mathcal{H}_x),$$

– a set I such that

$$d = |I| \quad ; \quad I := \{1, \dots, d\},$$

– a diagonal $d \times d$ matrix with real entries D_{H_x} ,

– a unitary $d \times d$ matrix U_x .

So that the family of matrices $(D_{H_x})_{x \in V}$ satisfies the additional condition

$$\sum_{x \in V} \text{Tr}(|D_{H_x}|) < +\infty. \quad (37)$$

Define

$$H_x := U_x^* D_{H_x} U_x$$

and fix a right square root h_x of $e^{H_x} = U_x^* e^{D_{H_x}} U_x$ as described in Lemma (6).

If $(h_{x;i,m})_{m \in I}$ is the i -th column vectors of h_x ($i \in I$), define the vector

$$h_{x;i} := \sum_{m \in I} h_{x;i,m} e_{x;m} \quad (38)$$

where $(e_{x;m})_{m \in I}$ is an ONB of \mathcal{H}_x . Then the family of vectors associated to the vectors (38) as in (12), i.e.

$$\Psi_\Lambda := \sum_{i \in I} \bigotimes_{x \in \Lambda} h_{x,i} \in \bigotimes_{x \in \Lambda} \mathcal{H}_x =: \mathcal{H}_\Lambda \quad ; \quad \Lambda \subset_{fin} V \quad (39)$$

is such that the limit

$$\hat{\psi}(b_\Lambda) := \lim_{\Lambda_1 \uparrow \uparrow V} \langle \Psi_{\Lambda_1}, b_\Lambda \Psi_{\Lambda_1} \rangle \quad (40)$$

exists for all $b_\Lambda \in \mathcal{B}_\Lambda$ and the family $(\hat{\psi}(b_\Lambda))_{\Lambda \subset_{fin} V}$ is projective.

Proof. We know from Lemma 2 that a sufficient condition for the existence of the limit (40) is that, for each $i, j \in I$, the limit

$$\lim_{\Lambda_1 \uparrow \uparrow V} \prod_{x \in \Lambda_1 \setminus \Lambda} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^*) = \lim_{\Lambda_1 \uparrow \uparrow V} e^{\sum_{x \in \Lambda_1 \setminus \Lambda} H_{x,i,j}} \quad (41)$$

exists. From (35) and (37), we know that for all $i, j \in I$

$$\sum_{x \in \Lambda_1 \setminus \Lambda} |H_{x,i,j}| \leq \sum_{x \in V} |H_{x,i,j}| \leq \sum_{x \in V} \text{Tr}(|D_{H_x}|) < +\infty.$$

Therefore the series $\sum_{x \in \Lambda_1 \setminus \Lambda} H_{x,i,j}$ is absolutely convergent. Therefore for each $i, j \in I$, the limit (41) exists. The projectivity of the family $(\hat{\psi}_\Lambda(b_\Lambda))_{\Lambda \subset_{fin} V}$ follows from the convergence of the series $\sum_{x \in \Lambda_1 \setminus \Lambda} H_{x,i,j}$ with the same argument used in Lemma 4.

6 Homogeneous superposition states

Homogeneous bundles on V are characterized by the fact that there is a Hilbert space \mathcal{H} and for each $x \in V$, a unitary isomorphism

$$J_x : \mathcal{H}_x \rightarrow \mathcal{H}$$

a finite set I_0 and a set of vectors $(h_j)_{j \in I_0}$ in \mathcal{H} such that

$$h_{x,j} = J_x^* h_j \quad ; \quad \forall j \in I_0, \forall x \in V.$$

In this case (15) becomes

$$\begin{aligned} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^*) &= \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^*) = \text{Tr}_{\mathcal{H}_x} (J_x h_i h_j^* J_x^*) \\ &= \text{Tr}_{\mathcal{H}} (h_i h_j^*) = \langle h_j, h_i \rangle =: \beta_{ij}. \end{aligned} \quad (42)$$

Therefore

$$\psi_{\Lambda_1}(b_\Lambda) = \sum_{i,j \in I_0} \prod_{x \in \Lambda} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^* b_x) \beta_{ij}^{|\Lambda_1 \setminus \Lambda|}. \quad (43)$$

Remark. The following Lemma shows that the condition that the β_{ij} in (42) are constant in (i, j) , leads to non-interesting states.

Lemma 7 Let \mathcal{K} be a Hilbert space, $n \in \mathbf{N}$ and $c \in \mathbf{R}$ with $c > 0$. Then two vectors $h_1, h_2 \in \mathcal{K}$ satisfy

$$\langle h_j, h_i \rangle = c \quad ; \quad \forall i, j \in \{1, 2\} \quad (44)$$

then they are proportional.

In particular, if the β_{ij} in (42) are independent of (i, j) , the state ψ_{Λ_1} in (43) is a product linear functional.

Proof. If the vectors h_1, h_2 satisfy (44), then

$$\langle h_1, h_2 \rangle^2 = c^2 = \langle h_1, h_1 \rangle \langle h_2, h_2 \rangle \quad (45)$$

and by the Schwartz inequality they are proportional.

If the $\beta_{ij} = \beta$ in (42) are independent of (i, j) , then by the first part of the lemma, the vectors h_i are proportional to a single vector $h \in \mathcal{H}$, hence for some $t_i \in \mathbf{C}$, $h_{x,i} = t_i h_x = t_i J_x^{-1} h$. In this case (43) becomes

$$\psi_{\Lambda_1}(b_\Lambda) = \sum_{i,j \in I_0} \bar{t}_j^{|\Lambda|} t_i^{|\Lambda|} \beta^{|\Lambda_1 \setminus \Lambda|} \prod_{x \in \Lambda} \text{Tr}_{\mathcal{H}_x}(h_x h_x^* b_x)$$

which shows that ψ_{Λ_1} is a product linear functional.

Remark. Allowing a dependence of the t_i on $x \in V$ only changes the homogeneous product linear functional in the proof of Lemma 7 into a non-homogeneous one, but still of product type.

Remark. If we consider a weaker condition for which (44) is satisfied only for $i \neq j$ then the vectors $(h_i)_{i \in I}$ are not necessary proportional. For the convenience of the reader, let $(e_j)_{1 \leq j \leq p}$ be an orthonormal family of vectors in \mathcal{H} . If $(\alpha_i)_{1 \leq i \leq p}$ is the sequence of real number defined by

$$\alpha_1 = c; \quad \alpha_j = c - (\alpha_1^2 + \alpha_2^2 + \cdots + \alpha_{j-1}^2); \quad j = 2, \cdots, p-1. \quad (46)$$

Then the following vectors

$$h_1 = e_1, \quad h_j = \alpha_1 e_1 + \alpha_2 e_2 + \cdots + \alpha_{j-1} e_{j-1} + e_j; \quad j = 2, \cdots, p$$

satisfy $\langle h_i, h_j \rangle = c$ for $i \neq j$. However, they are not proportional since they form a linearly independent family in \mathcal{H} . In this case the state in (43) becomes

$$\psi_{\Lambda_1}(a_\Lambda) = \sum_{i \in I} \prod_{x \in \Lambda} \text{Tr}(h_{x,i} h_{x,i}^* a_x) \beta_{ii}^{|\Lambda_1 \setminus \Lambda|} + \sum_{i \neq j} \prod_{x \in \Lambda} \text{Tr}(h_{x,i} h_{x,j}^* a_x) c^{|\Lambda_1 \setminus \Lambda|}, \quad (47)$$

which is a combination of product positive linear functionals.

Theorem 5 Let $x \in V \mapsto \mathcal{H}_x = J_x^* \mathcal{H}$ be an homogeneous bundle on V and let be given a finite set I_0 and a set of vectors $(h_j)_{j \in I_0}$ in \mathcal{H} . Define

$$h_{x,j} = J_x^* h_j \quad ; \quad \forall j \in I_0, \forall x \in V. \quad (48)$$

and suppose that the vectors h_j satisfy the **generic condition**

$$i \neq j \Rightarrow |\langle h_j, h_i \rangle| = \beta_{ij} < \|h_j\| \|h_i\| = \sqrt{\beta_{ii}} \sqrt{\beta_{jj}}. \quad (49)$$

For $\Lambda \subset_{fin} V$, define the vector

$$\Psi_\Lambda := \sum_{i \in I} \bigotimes_{x \in \Lambda} h_{x,i} \quad (50)$$

and the associated positive linear functional on \mathcal{B}_Λ given by

$$\psi_\Lambda = \langle \Psi_\Lambda, (\cdot) \Psi_\Lambda \rangle = \text{Tr}_{\mathcal{H}_\Lambda} (\Psi_\Lambda \Psi_\Lambda^* (\cdot)). \quad (51)$$

Then, for each $b_\Lambda := \bigotimes_{x \in \Lambda} b_x \in \mathcal{B}_\Lambda$, the limit

$$\lim_{\Lambda_n \uparrow V} \frac{\psi_{\Lambda_n}(b_\Lambda)}{\psi_{\Lambda_n}(1_{\Lambda_n})} \quad (52)$$

exists, is independent of the sequence (Λ_n) with the above property and is equal to

$$\frac{1}{\beta^\Lambda \sum_{i \in I_0, \beta_{ii} = \beta} 1} \sum_{i \in I_0, \beta_{ii} = \beta} \prod_{x \in \Lambda} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,i}^* b_x) \quad (53)$$

for some $\beta > 0$. In particular, it is a convex combination of product states.

Proof. From (43) it is known that

$$\psi_{\Lambda_1}(b_\Lambda) = \sum_{i,j \in I_0} \prod_{x \in \Lambda} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^* b_x) \beta_{ij}^{|\Lambda_1 \setminus \Lambda|}. \quad (54)$$

Therefore

$$\psi_{\Lambda_1}(1_{\Lambda_1}) = \sum_{i,j \in I_0} \beta_{ij}^{|\Lambda_1|}.$$

Denote

$$\beta := \max\{|\beta_{ij}| : i, j \in I_0\} = \max\{|\beta_{ii}| : i \in I_0\} \quad (55)$$

where the second identity follows from the Schwartz inequality. Then

$$\frac{\psi_{\Lambda_1}(b_\Lambda)}{\psi_{\Lambda_1}(1_{\Lambda_1})} = \frac{1}{\sum_{i,j \in I_0} \beta_{ij}^{|\Lambda_1|}} \sum_{i,j \in I_0} \prod_{x \in \Lambda} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^* b_x) \beta_{ij}^{|\Lambda_1 \setminus \Lambda|}$$

$$\begin{aligned}
&= \frac{1}{\beta^{|\Lambda_1|} \sum_{i,j \in I_0} (\beta_{ij}/\beta)^{|\Lambda_1|}} \sum_{i,j \in I_0} \prod_{x \in \Lambda} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^* b_x) \beta_{ij}^{|\Lambda_1 \setminus \Lambda|} \\
&= \frac{1}{\beta^{|\Lambda|} \sum_{i,j \in I_0} (\beta_{ij}/\beta)^{|\Lambda_1|}} \sum_{i,j \in I_0} \prod_{x \in \Lambda} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^* b_x) (\beta_{ij}/\beta)^{|\Lambda_1 \setminus \Lambda|}.
\end{aligned}$$

From (42) and (49) one knows that, for $i \neq j$,

$$|\beta_{ij}| < \beta.$$

Therefore

$$\lim_{\Lambda_1 \uparrow \uparrow V} \frac{\psi_{\Lambda_1}(b_\Lambda)}{\psi_{\Lambda_1}(1_{\Lambda_1})} = \frac{1}{\beta^{|\Lambda|} \sum_{i \in I_0, \beta_{ii} = \beta} 1} \sum_{i \in I_0, \beta_{ii} = \beta} \prod_{x \in \Lambda} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,i}^* b_x).$$

This proves (56).

Remark Theorem 6 below propose a similar result to 5 replacing the generic condition (49) by the weaker assumption that the β_{ij} are real numbers but under the assumptions of Lemma 4.

Theorem 6 Under the assumptions and the notations of Lemma 4 and if ψ_Λ is given by (51). For each $b_\Lambda := \bigotimes_{x \in \Lambda} b_x \in \mathcal{B}_\Lambda$, the limit (52) exists, is independent of the sequence (Λ_n) and is equal to

$$\frac{1}{\beta^{|\Lambda|} \sum_{i,j \in I_0, \beta_{ij} = \beta} 1} \sum_{i,j \in I_0, \beta_{ij} = \beta} \prod_{x \in \Lambda} \text{Tr}_{\mathcal{H}_x} (h_{x,i} h_{x,j}^* b_x) \quad (56)$$

where β is given by (55).

Proof. Lemma 4 guarantees the existence of the limit (52), the rest of the proof can be done similarly to the proof of Theorem 5. \square

7 Mixing property for superposition states

In this section, we consider the multidimensional integer lattice for which $V = \mathbf{Z}^\nu$. A finite Hilbert space \mathcal{H} is fixed and for each finite bounded region Λ , unitary isomorphisms

$$J_\Lambda : \mathcal{H}^{\otimes |\Lambda|} \rightarrow \mathcal{H}_\Lambda.$$

If Λ and Λ' are bounded regions w $|\Lambda| = |\Lambda'|$, we have the following identification

$$J_{\Lambda}^{-1}(J_{\Lambda}(a)) \equiv J_{\Lambda'}^{-1}(J_{\Lambda'}(a)); \quad a \in \mathcal{H}^{\otimes |\Lambda|}.$$

For $z = (z_1, z_2, \dots, z_{\nu}) \in \mathbf{Z}^{\nu}$ we denote

$$|z| = |z_1| + |z_2| + \dots + |z_{\nu}|.$$

Denote for each $r > 0$

$$D_r = \{z \in \mathbf{Z}^{\nu} : |z| \leq r\}.$$

For each $\Lambda' \subset_{\text{fin}} V$ and $t > 0$, then notation $J_t(\Lambda')$ means any embedding of Λ' in D_t^c .i.e.

$$J_t(\Lambda') \subset D_t^c; \quad |J_t(\Lambda')| = |\Lambda'|. \quad (57)$$

The set of all embedding of Λ' in D_t^c is identified to the set of all injective maps from Λ' into D_t^c . It will be denoted by $\mathcal{E}(\Lambda', D_t^c)$. For each $\Lambda \subset_{\text{fin}} \mathbf{Z}^{\nu}$, since

$$\beta_{\Lambda^c, i, j} = \prod_{x \in \Lambda^c} \text{Tr}(h_{x, i} h_{x, j}^*) \in \mathbf{C}.$$

For $a_{\Lambda} = \bigotimes_{x \in \Lambda} a_x \in \mathcal{B}(\mathcal{H}_{\Lambda})$

$$\psi(a_{\Lambda}) = \sum_{i, j \in I} \prod_{x \in \Lambda} (\text{Tr}(h_{x, i} h_{x, i}^*) \text{Tr}(h_{x, j} h_{x, j}^*)) \beta_{\Lambda^c, i, j} \quad (58)$$

the linear functional ψ is positive and we assume furthermore that it is normalized .i.e

$$\sum_{i, j \in I} \beta_{V, i, j} = 1. \quad (59)$$

Definition 3 A state φ on \mathcal{A}_V is said to enjoy the mixing property if for each bounded regions $\Lambda, \Lambda' \subset V$ and $a_{\Lambda} \in \mathcal{B}_{\Lambda}, b_{\Lambda'} \in \mathcal{B}_{\Lambda'}$ the following property holds:

$$\forall \varepsilon > 0, \exists t > 0, J_t \in \mathcal{E}(\Lambda', D_t^c) \quad |\varphi(a_{\Lambda} J_t(b_{\Lambda'})) - \varphi(a_{\Lambda}) \varphi(J_t(b_{\Lambda'}))| \leq \varepsilon. \quad (60)$$

Definition 4 Let $(\alpha_p)_p$ be a net of states on $\mathcal{B}(\mathcal{H}^{\otimes p})$. A state φ on \mathcal{A}_V is said to enjoy the α -mixing property if for each bounded regions $\Lambda, \Lambda' \subset V$ and $a_{\Lambda} \in \mathcal{B}_{\Lambda}, b_{\Lambda'} \in \mathcal{B}_{\Lambda'}$ the following property holds:

$$\forall \varepsilon > 0, \exists t > 0, \forall J_t \in \mathcal{E}(\Lambda', D_t^c) \quad \left| \varphi(a_{\Lambda}) J_t(b_{J_t^{-1}(\Lambda')}) - \varphi(a_{\Lambda}) \alpha_{|\Lambda'|}(b_{\Lambda'}) \right| \leq \varepsilon \quad (61)$$

Theorem 7 *Under the notations and assumptions of Lemma 4. Assume that for each finite region $\Lambda' \subset_{\text{fin}} V$ and $b_{\Lambda'} = \bigotimes_{y \in \Lambda'} b_y \in \mathcal{B}_{\Lambda'}$ the limit*

$$\lim_{t \rightarrow +\infty} \prod_{y \in \Lambda'} \text{Tr}(h_{y,i} h_{y,j}^* b_y) =: \alpha^{(|\Lambda'|)}(J_{\Lambda'}^{-1}(b_{\Lambda'})) \quad (62)$$

exists and the limit $\alpha^{(|\Lambda'|)}(J_{\Lambda'}^{-1}(b_{\Lambda'}))$ is independent of $i, j \in I$. Then $\alpha_{|\Lambda'|}$ is a state of product type on the algebra $\mathcal{B}(\mathcal{H}^{|\Lambda'|})$. The state $\hat{\psi}$ is given by (40), enjoys the α -mixing property (61) and the mixing property (60).

Proof. The functional $\alpha^{(|\Lambda'|)}$ is a point-wise limit of positive functional of product type. This make it positive functionals. Moreover, since the left hand side of (62) is independent of the one-to-one embedding J_t of $\mathcal{B}(\mathcal{H}^{|\Lambda'|}) \equiv \mathcal{B}(\mathcal{H})^{|\Lambda'|}$ into $\mathcal{B}(\mathcal{H}_{\mathcal{J}_t(\cdot)})$. And since $\alpha^{(|\Lambda'|)}$ is independent of the indices i, j then for $b_{\Lambda'} \geq 0$ one has

$$\alpha^{(|\Lambda'|)}(b'_{\Lambda'}) = \lim_{t \rightarrow \infty} \prod_{y \in \Lambda'} \text{Tr}(h_{J_t(y),i} b_y h_{J_t(y),i}^*) \geq 0$$

and

$$\alpha^{(|\Lambda'|)}(I_{\Lambda'}) = \lim_{t \rightarrow \infty} \prod_{y \in \Lambda'} \text{Tr}(h_{J_t(y),i} h_{J_t(y),i}^*) = 1$$

whenever $\text{Tr}(h_{y,i} h_{y,i}^*) = 1$ for each $y \in V$ and $i \in I$. Thus $\alpha^{(|\Lambda'|)}$ is a state on $\mathcal{B}(\mathcal{H})^{|\Lambda'|}$.

One has

$$\hat{\psi}(a_{\Lambda} J_t(b_{\Lambda'})) = \sum_{i,j} \left(\prod_{x \in \Lambda} \text{Tr}(h_{x,i} h_{x,j}^* a_x) \right) \left(\prod_{y \in \Lambda'} \text{Tr}(h_{J_t(y),i} b_y h_{J_t(y),j}^*) \right) \beta_{(\Lambda \cup J_t(\Lambda'))^c, i, j}$$

and

$$\psi(a_{\Lambda}) \psi(J_t(b_{\Lambda'})) = \left(\sum_{i,j} \prod_{x \in \Lambda} \text{Tr}(h_{x,i} h_{x,j}^* a_x) \beta_{\Lambda^c, i, j} \right) \left(\sum_{k,l} \prod_{y \in \Lambda'} \text{Tr}(h_{J_t(y),k} b_y h_{J_t(y),l}^*) \beta_{(J_t(\Lambda'))^c, k, l} \right)$$

Since $\sum_{z \in \mathbf{Z}^d} |H_{z,i,j}| < +\infty$ then as $t \rightarrow +\infty$ one gets

$$\begin{cases} \beta_{(\Lambda \cup J_t(\Lambda'))^c, i, j} \longrightarrow \beta_{\Lambda^c, i, j}; \\ \beta_{(J_t(\Lambda'))^c, k, l} \longrightarrow \beta_{V, k, l}. \end{cases}$$

From (62) one gets

$$\begin{aligned} \lim_{t \rightarrow \infty} \hat{\psi}(a_\Lambda J_t(b_{\Lambda'})) &= \sum_{i,j} \left(\prod_{x \in \Lambda} \text{Tr}(h_{x,i} h_{x,j}^* a_x) \right) \beta_{\Lambda^c, i, j} \alpha^{(|\Lambda'|)}(J_{\Lambda'}^{-1}(b_{\Lambda'})) \quad (63) \\ &= \psi(a_\Lambda) \alpha^{(|\Lambda'|)}(J_{\Lambda'}^{-1}(b_{\Lambda'})). \end{aligned}$$

Therefore, the state ψ satisfies the α -mixing property (61). On the other hand, on has

$$\begin{aligned} \lim_{t \rightarrow \infty} \psi(a_\Lambda) \psi(J_t(b_{\Lambda'})) &= \left(\sum_{i,j} \prod_{x \in \Lambda} \text{Tr}(h_{x,i} h_{x,j}^* a_x) \beta_{\Lambda^c, i, j} \right) \left(\sum_{k,l} \beta_{V, k, l} \alpha^{(|\Lambda'|)}(J_{\Lambda'}^{-1}(b_{\Lambda'})) \right) \\ &= \left(\sum_{i,j} \prod_{x \in \Lambda} \text{Tr}(h_{x,i} h_{x,j}^* a_x) \beta_{\Lambda^c, i, j} \right) \left(\sum_{k,l} \beta_{V, k, l} \right) \alpha^{(|\Lambda'|)}(J_{\Lambda'}^{-1}(b_{\Lambda'})). \end{aligned}$$

And since

$$\sum_{k,l} \beta_{V, k, l} = \psi(1) = 1.$$

Then

$$\lim_{t \rightarrow \infty} \psi(a_\Lambda) \psi(J_t(b_{\Lambda'})) = \psi(a_\Lambda) \alpha^{(|\Lambda'|)}(J_{\Lambda'}^{-1}(b_{\Lambda'})). \quad (64)$$

From (63) and (64) for each $\varepsilon > 0$ there exists $t \geq 0$ such that for each $J_t \in \mathcal{E}(\Lambda', D_t^c)$

$$\left| \psi(a_\Lambda J_t(b_{\Lambda'})) - \psi(a_\Lambda) \alpha^{(|\Lambda'|)}(J_{\Lambda'}^{-1}(b_{\Lambda'})) \right| < \varepsilon/2$$

and

$$\left| \psi(a_\Lambda) \psi(J_t(b_{\Lambda'})) - \psi(a_\Lambda) \alpha^{(|\Lambda'|)}(J_{\Lambda'}^{-1}(b_{\Lambda'})) \right| < \varepsilon/2.$$

Then, for each $J_t \in (\Lambda', D_t^c)$

$$|\psi(a_\Lambda J_t(b_{\Lambda'})) - \psi(a_\Lambda) \psi(J_t(b_{\Lambda'}))| < \varepsilon.$$

Therefore the state ψ satisfies the mixing property (60). \square

Remark If the right hand side in (62) depends in the indices i, j , then it defines a state of product type on $\mathcal{B}(\mathcal{H}_x)$ denoted by $\alpha_{i,j}^{(|\Lambda'|)}(J_{\Lambda'}^{-1}(b_{\Lambda'}))$. In this case,

$$\hat{\psi}(a_\Lambda J_t(b_{\Lambda'})) \longrightarrow \sum_{i,j} \prod_{x \in \Lambda} \text{Tr}(h_{x,i} h_{x,j}^* a_x) \beta_{\Lambda^c, i, j} \alpha_{i,j}^{(|\Lambda'|)}(J_{\Lambda'}^{-1}(b_{\Lambda'}))$$

while

$$\psi(a_\Lambda)\psi(J_t(b_{\Lambda'})) \longrightarrow \left(\sum_{i,j} \prod_{x \in \Lambda} \text{Tr}(h_{x,i} h_{x,j}^* a_x) \beta_{\Lambda^c, i, j} \right) \left(\sum_{k,l} \beta_{V, k, l} \alpha_{k, l}^{(|\Lambda'|)}(J_{\Lambda'}^{-1}(b_{\Lambda'})) \right).$$

Therefore, the mixing properties (61) and (60) are not necessarily satisfied.

References

- [1] Choi, M. D. *Completely positive linear maps on Complex matrices*, Linear algebra and its applications 10, 285-290 (1975).
- [2] Accardi L.: Non-commutative Markov chains, In: Proceedings International School of Mathematical Physics Universita' di Camerino 30 Sept. - 12 Oct. (1974) 268-295
- [3] Accardi L.: The noncommutative markovian property, Func. Anal. Appl. 9 (1) (1975) 1-8 (in russian), submitted 31-1-1974, translated in: Funct. Anal. and its Appl. 9 (1) (1975) 1-8
- [4] Accardi L.: Some trends and problems in quantum probability, In: Quantum probability and applications to the quantum theory of irreversible processes
- [5] Gustafson S., Sigal I. M. *Mathematical Concepts of Quantum Mechanics*, Springer-Verlag Berlin Heidelberg 2011
- [6] J. Heo, Reproducing kernel Hilbert -modules and kernels associated with cocycles, Journal of Mathematical Physics, 49 (2008), 103507-103519
- [7] Accardi, L., Quantum theory and nonkolmogorovian probability, In: Stochastic Processes in quantum theory and Statistical Physics, eds. S. Albeverio, Ph. Combe, M. Sirugue-Collin; Springer Lecture Notes in Physics N. 173
- [8] Accardi L., Ohya M.: Compound channels, transition expectations and liftings, Applied Mathematics Optimization 39 (1999) 33-59
Volterra preprint N. 75 (1991)

- [9] L. Accardi, F. Fidaleo: Entangled Markov Chains *Annali di Matematica Pura e Applicata*, 184 (3) (2005) 327-346 Preprint Volterra N.556 (2003)
- [10] Accardi L., Ohya M., Matsuoka T.: Entangled Markov chains are indeed entangled, *Infin. Dimens. Anal. Quantum Probab. Relat. Top. (IDA-QP)* 9 (3) (2006) 379-390
- [11] Accardi L., Matsuoka T., Ohya M.: On Entangled Markov Chains, *Publ. RIMS, Kyoto Univ.*, 1609 (2008) 75-87
- [12] Accardi L, Souissi A, Soueidy El: Quantum Markov chains, A unification approach, to appear in: *Infin. Dimens. Anal. Quantum Probab. Relat. Top. (IDA-QP)* (2020)
- [13] Bennett G. Schur multipliers, *Duke Math. J.* 44 (1977), 603–639
- [14] Kruszynski P.: A problem in quantum probability, National Aeronautics and Space Administration, REPT-87-06, N88-10573 preprint 1987
- [15] Bezhaeva, Z.I., Oseledets, V.I. Remarks on quantum Markov states, *Funct. Anal. Appl.*, 3, 205-209 (2015)
- [16] Bezhaeva, Z.I., Oseledets, V.I. Quantum Markov States and Quantum Hidden Markov States. *J Math Sci* 240, 507-514 (2019)
- [17] Haag, R.: *Local Quantum Physics: Fields, Particles, Algebras.* Springer, Berlin (1996)
- [18] Kang, Y.B. Quantum Markov semigroups for continuous-time open quantum random walk. *Quantum Inf Process* 18, 196 (2019)
- [19] Paulsen V.: *Completely Bounded Maps and Operator Algebras*, Cambridge University Press, Cambridge (2002)
- [20] G.G. Emch, *Algebraic Methods in Statistical Mechanics and Quantum Field Theory*, Dover Publications, 2014.
- [21] Sakai, S.: *C^* -Algebras and W^* -Algebras*, Springer (1971)
- [22] Stout Q. F.: Schur multiplication on $\mathcal{B}(\ell_p, \ell_q)$, *J. Operator Theory* 5 (1981) 231–243