Adiabatic index for relativistic stars as a coefficient of non-thermal dissipation

Seema Satin

Indian Institute for Science Education and Research , Pune , India.*

The adiabatic index of a relativistic star (modeled by a perfect fluid) is shown to act as a dissipation constant due to considerations of mesoscopic scale stochastic effects. This dissipative effect arises without taking heat flux in the fluid model and is termed as adiabatic or non-thermal for the system under consideration. A basic formalism for introducing stochastic effects in interiors of massive stars has recently been proposed via a classical Einstein-Langevin equation. The origin of stochasticity is associated with the pressure variable (due to degeneracy of constituent particles) and its fluctuations, and is not viable for pressureless fluids. The fluctuations dissipate their fermi energy into perturbing the system, adiabatically. This is shown by a fluctuation dissipation relation, and a Langevin fermi temperature associated with these fluctuations is defined. The high magnitude of background field fluctuations which we get here, is feasible to account for perturbing the strong gravity regions with dense matter. Nevertheless, the perturbations themselves are consistently and reasonably small as first order deviations from equilibrium.

Exploring the interiors of relativistic/compact stars and the properties of matter in highly dense state of matter in strong gravity regions, poses a challenge for astronomers as well as for theoretical physicists for modeling them [1-3]. With the advent of gravitational wave detection methods [4, 5], one can probe into these, by developing more detailed formalism that would help to delve into deeper layers in one way or the other. Hence this area is open for enhancing the basic models and extending towards new methods of analysis in order to decipher the interiors of such astrophysical objects, theoretically as well as for observational purposes. In this letter we show that the adiabatic index can be identified as a coefficient of non-thermal dissipation for cold matter relativistic star models, enhanced with stochastic effects. These stochastic fluctuations which partially capture the microscopic/quantum properties resulting in a mesoscopic description of the matter in the interiors, act as source of induced perturbations. It is not just the magnitude of these mesoscopic fluctuations that is of interest to emphasise the importance of such additions, but new underlying physics which can be explored due to such considerations is the aim of such an endeavour. The basic structure of such a theory begins with a classical Einstein-Langevin formalism in recent work [6, 7]. While in this letter we obtain a fluctuation-dissipation relation and give estimates over magnitude of background fluctuations for a simple model, which are obtained due to known approximate values of the adiabatic index of the neutron star. This also opens up new avenues in statistical physics, both equilibrium and non-equilibrium, to extend such developments and studies for compact stars and objects governed by general relativity.

Considering a zero temperature equation of state to describe matter $\epsilon = \epsilon(p)$, say for a neutron star (one-component perfect fluid) where pressures are quantum mechanical in origin due to degeneracy of neutrons in

order to encounter strong gravity, we model noise in the system. The stress tensor for such a system is that of a the perfect fluid,

$$T^{ab} = u^a u^b \epsilon + q^{ab} p \tag{1}$$

where $q^{ab} = u^a u^b + g^{ab}$.

The identification of the adiabatic index of the massive star, given by

$$\gamma = \frac{(\epsilon + p)}{p} \frac{dp}{d\epsilon} \tag{2}$$

and defined as the fractional change in pressure per fractional change in comoving volume at constant entropy and composition for a massive star, with a dissipative term is seen to follow from the F-D relation that we obtain here. It is known that coefficient of friction/dissipation in thermal systems, depends on details of molecular size and their kinetic energy in the medium and is linked to temperature and pressure of the system [8]. However it is not a thermodynamic property, and can be linked to other parameters than the thermal temperature of the system. Here we connect the adiabatic index as a coefficient of non-thermal dissipation with the background noise (fluctuations) due to degeneracy pressure of neutrons. The fermi energy which is associated with the degeneracy pressure in The neutron stars after they cool down to have thermal temperatures $\sim 1 \text{ MeV}$, while the fermi temperature is more than $E_F \approx 60 \text{ MeV}$ [10]. The stochastic effects which we model, thus encode these effects of quantum origin and are not thermodynamical.

However since the equation of state is not known correctly for such cases, we do not claim any new estimates over magnitudes of fermi temperature or polytropic index, but work with known estimates only, introducing in the model these additional effects in terms of fluctuations.

On the other hand, study of oscillations and perturbations in relativistic stars, forms the base of asteroseismology and theoretical developments start with perturbed

^{*} seemasatin@iiserpune.ac.in

Einstein's equations [9, 10].

$$\delta G^{ab}(x) = 8\pi \delta T^{ab}(x) \tag{3}$$

However all through the well established literature, the perturbed Einstein's equations lack a source term, which may induce such perturbations in the system. The perturbations in an isolated relativistic star or a more involved configuration of binary/accreting systems are usually assumed to arise due to internal or external effects [11] which are often overlooked and not specified. The focus is rather over calculating and classifying the perturbations and modes of oscillations in an astrophysical system [12]. Taking into consideration a source which induces perturbations in the system, is desirable, specially when these sources are due to some mechanism in the interior of the star itself. These could also arise as response to some external agency which triggers internal mechanism in the matter content to start off the oscillations. The characteristics of the perturbations depend on the properties of the stellar structure, rather than the details of the external source.

Our analysis is useful for cold neutron stars and other compact matter objects in a different way than just calculating estimates over TOV equations and mass radius relation, though exploring these could also be of significance in some specific cases. It is thus useful to lay a new strategy to explore the interiors of these objects based on enhancement of fluid properties as we do here.

In this letter we state the basic framework for radial perturbations which has been worked out in [6, 7], and show the first example of a parameter namely the adiabatic index that can be viewed in a new way.

The perturbations are regarded as deformation of the original configuration of the star, with each fluid element mapped to a corresponding fluid element by a Lagrangian displacement vector ξ^a and by metric perturbation h_{ab} . Thus fundamental variables describing the perturbations are ξ^a and h_{ab} [10]. With a source term acting as driving mechanism (in the background spacetime) to induce stochastic perturbations, the classical Einstein-Langevin equation reads [6],

$$\delta G^{ab}[h,\xi;x) = 8\pi (\delta T^{ab}[h,\xi;x) + \tau^{ab}[g;x)) \qquad (4)$$

where the Langevin noise $\tau^{ab}[g;x) = T^{ab}[g;x) - \langle T^{ab}[g;x) \rangle$ is defined on the background metric g_{ab} , and satisfies $\langle \tau^{ab}(x) \rangle = 0$. The two point noise kernel is given by $\langle \tau^{ab}(x)\tau^{cd}(x') \rangle = N^{abcd}(x,x')$, a bitensor. Such a treament of the gravitating system, calls for a system-bath separation. One can consider the spacetime metric to act as the system while the matter fields as the environment/bath.

Modes of oscillations which can be detectable from such isolated objects for realistic cases, and originating in the interiors can travel through the compact object to the surface and further , in principle, in the form of gravitational waves, for non-radial perturbations. It is possible that these stochastic effects set stage for new stochastic modes of oscillations and finding them. However one has to correctly work out the corresponding analysis and show details of such modes. This can be carried out in detailed upcoming work.

The pressures inside neutron stars are of the order of $10^{31} - 10^{34}$ Pa and they increase from the outer radius to the core modeled by perfect fluids. A correspondence between quantum field fluctuations in classicalized limit to hydrodymanic approximation in terms of macroscopic variables of the perfect fluid, has recently been obtained in [13]. Thus one can see, how any quantum fluctuations can be partially captured via a fluid model approximation of matter.

For a relativistic star, the equation of state in simplest form is

p

$$=w\epsilon^{\Gamma}$$
 (5)

where ϵ is the energy density of the fluid, and Γ the polytropic index. We give the above relation just for the sake of completion of describing the system, and the results are not affected by unsettled issues over polytropic index for neutron stars in this letter. One may as well consider a simple case of $p = w\epsilon$ here for a toy model that can be used for such a relativistic star, as is used in literature for complete gravitational collapse of massive stars to singularities [14]. It is apparent that the pressure fluctuations extend the stochastic nature to the energy density via the equation of state. However we assign the origin of fluctuations in the pressure variable, as it is pressure in the neutron star which balances gravity. One can consistently add the fluctuations of the stress tensor given by τ^{ab} as Langevin noise in the system. It is to be noted that the background spacetime with g_{ab} is treated deterministically while the perturbations h_{ab} , ξ^a show a stochastic nature, hence $G^{ab}[g] = 8\pi T^{ab}[g]$ still remains the unaltered Einstein's equation while its perturbations $\langle \delta G^{ab} \rangle = 8\pi \langle \delta T^{ab} \rangle$ are stochastic and have an added noise on the rhs to give it a Langevin form. The perturbed metric is then of the form $g_{ab} + \langle h_{ab} \rangle_s, \langle \rangle_s$ being the stochastic average.

For the present analysis we restrict ourselves to linearized equations, thus limiting the perturbations of the Einstein's equations to first order in h_{ab} and ξ^a . This can be obtained by second order variation of the action

$$S = -\int (\frac{1}{16\pi}R - \epsilon)\sqrt{g}d^4x \tag{6}$$

and is well established in literature, giving the perturbed Einstien's equation (4), to which we add the Langevin term and write the E-L equation with explicit form of δT^{ab} as

$$\delta G^{ab} = 8\pi (W^{abcd} \Delta g_{cd} - \mathcal{L}_{\xi} T^{ab}) + 8\pi \tau^{ab} \tag{7}$$

where $\Delta g_{ab} = h_{ab} + \nabla_a \xi_b + \nabla_b \xi_a$. The Lagrangian change in a quantity is given by Δ , while Eulerian changes are respresented by δ , the usual convention followed by [9, 10] We split and write the term

$$W^{abcd} = E^{abcd} + D^{abcd} \tag{8}$$

with

$$E^{abcd} = \frac{1}{2} (\epsilon + p) u^a u^b u^c u^d + \frac{1}{2} p (g^{ab} g^{cd} - g^{ac} g^{bd} - g^{ad} g^{bd})$$
$$D^{abcd} = -\frac{1}{2} \gamma p q^{ab} q^{cd} \tag{9}$$

here γ stands for the adiabatic index of the star as defined in equation (2). We claim that the term D^{abcd} shows a dissipative effect and in absence of any heat flux. The claim is based on the reasoning that, since noise here is sourced in pressure fluctuations of the fluid fundamentally, the dissipative effect would be seen in the pressure perturbations $\Delta p = -1/2\gamma pq^{cd}\Delta g_{cd}$ which in the per-turbed stress tensor accounts for the $q^{ab}\Delta p$ term only, contributing to $D^{abcd}\Delta q_{cd}$ in the E-L equation. More specifically, the degeneracy pressure fluctuations in the compact object dissipate their fermi (non-thermal) energy into perturbing the background fluid variables and the geometry given in terms of Δg_{ab} . Also Δg_{cd} carries a harmonic time dependence here and hence we get a local dissipation effect, for the temporal coordinate.

The nature of noise in the static background model is such that stochasticity is seen prominently in the radial coordinate, which means pressure fluctuations vary randomly with some distribution along the radial vector, for the spherically symmetric geometry containing the perfect fluid.

$$ds^{2} = -e^{2\nu(r)}dt^{2} + e^{2\lambda(r)} + r^{2}(d\Omega^{2})$$
(10)

Thus the radial dependence of noise is an important point to note here in the Langevin term, for different layers in the star. It is this nature of noise (having radial dependence) at different depths, that such models can help to probe the interiors in a compact star. The temporal dependence of the terms is taken harmonic (as in [6]) for this static background configuration for simplicity.

We associate the above said dissipative term with the noise in the system having an F-D relation as discussed further.

The general form of a fluctuation dissipation relation, is known to be given as

$$N^{abcd}(x,x') = \int K(x,x'')\tilde{D}^{abcd}(x'',x')d^4x'' \qquad (11)$$

where $N^{abcd}(x, x')$ and $\tilde{D}^{abcd}(x'', x')$ are the noise and dissipation kernels respectively, K(x, x'') being the fluctuation-dissipation kernel. Such a general form of FD theorem also follows for semiclassical stochastic Einstein-Langevin equations [15, 16] though the details are obtained through formulations specific to quantum open systems. However the above relation is valid for classical systems as well, relating the fluctuations in a system to the dissipation.

For the astrophysical model that we consider here with metric given by equation (10) and a perfect fluid stress tensor, the noise kernel has been defined and obtained in all its details in [6], which is of the form

$$N^{abcd}(x, x') = \text{Cov}[T^{ab}(\vec{x}), T^{cd}(\vec{x}')]e^{-i\omega(t-t')}$$
(12)

with fluctuations $\tau^{ab}(x) \equiv (T^{ab}(\vec{x}) - \langle T^{ab}(\vec{x}) \rangle)e^{i\omega t}$. The fundamental elements of perturbations are given by $h_{ab}(x) \equiv h_{ab}(\vec{x})e^{i\omega t}$ and $\xi^a(x) \equiv \xi^a(\vec{x})e^{i\omega t}$, oscillating with frequency ω over the static background. The components of the point separated (two point) noise kernel, as one can see in the above reference, are composed of two point covariances of pressure and energy density. In the coincidence limit of spatial points, where we take $\vec{x}' \rightarrow \vec{x}$ these reduce to the variances of pressure and energy density, keeping the temporal dependence separate.

As one can clearly see from the above relations, dissipation kernel $D^{abcd}(x) = D^{abcd}(\vec{x})$, is local in spatial coordinates, for the static background fluid, while the factor Δg_{cd} carries the perturbations and also the time dependence. This leads to ascertain taking coincidence limit in spatial coordinates for the two point noise kernel, with $\vec{x} \to \vec{x}'$. Denoting $(t - t') \equiv \mathbf{T}$, the F-D theorem here takes the form

$$N^{abcd}(\vec{x})e^{-i\omega\mathbf{T}} = \int K(\vec{x}, \vec{x}'; \mathbf{T})D^{abcd}(\vec{x}')d^3\vec{x}' \qquad (13)$$

The simplest choice for the fluctuation-dissipation kernel

is $K(\vec{x}, \vec{x}', \mathbf{T}) = -\frac{\langle p(x) \rangle}{2} \delta(\vec{x}, \vec{x}') e^{-i\omega \mathbf{T}}$. For the given case $q^{ab} = 0$ when $\{ab\} = \{tt, tr, t\theta, t\phi\}$ (since $u^a \equiv \{e^{-\nu}, 0, 0, 0\}$) and $q^{\alpha\beta} = g^{\alpha\beta}$ for indices spanning over $\{r, \theta, \phi\}$. It then follows that $D^{tttt} = D^{tt\zeta\delta} =$ $D^{\alpha\beta tt} = 0$ (where $\alpha, \beta, \zeta, \delta$ span over spatial coordinates only), which means there is no dissipation in the system with the corresponding noise components. The non-zero components of dissipation are then given by $D^{\alpha\beta\zeta\delta}(\vec{x})$, and one can write the F-D relation as

$$N^{\alpha\beta\zeta\delta}(\vec{x})e^{-i\omega\mathbf{T}} = -\frac{\langle p(x) \rangle}{2}D^{\alpha\beta\zeta\delta}(\vec{x})e^{-i\omega\mathbf{T}}$$
(14)

which reduces to

$$\operatorname{Var}[\mathbf{p}(\mathbf{r})] = \frac{1}{4}\gamma \langle p(r) \rangle^2 \tag{15}$$

The factor $\gamma/4$ here, acts as coefficient of adiabatic dissipation and has values 0.33 and 0.41 for $\gamma = 4/3$ and 5/3 respectively. The above gives standard deviation $\sigma = \sqrt{\gamma/2} < p(r) >$ with coefficient of variance CV $\equiv\sigma/=\frac{\sqrt{\gamma}}{2}=0.57$ for an adibatic index $\gamma=4/3$ and a value of CV = 0.64 for $\gamma=5/3$. Thus the fluctuations are about 57% to 64% of the degeneracy pressure. This value is w.r.t radial (spatial) coordinate hence it shows the amplitude (or rms) of normalised fluctuations which are oscillating harmonically w.r.t time, in the otherwise static equilibrium configuration of the star. This results in dispersion in the pressure values in the background configuration and also feeds the perturbations.

Most of the equations of state for neutron stars, estimate the energy density at the core to be nonrelativisitic (for neutrons), so that degeneracy pressure $p \sim np_F^2/m = n2E_F = 2nk_BT_F$, with p_F being the fermi momentum of neutrons, m, n being the mass and number density and E_F, T_F being the fermi energy and fermi temperature respectively. As one can clearly see the fermi temperature fluctuations will follow on similiar lines that of the pressure from the relation for CV and standard deviation is around 34.2 MeV for a background fermi temperature 60 MeV. We term these fluctuations of fermi temperature as the Langevin temperature associated with the noise, and one can see that it has a radial dependence as well for the model here. It is to be noted that the statistical average of these random fluctuations will be vanishing, as is required by a Langevin noise.

From equation (14) we see that it can be written in full form of a fluctuation dissipation relation as

$$N^{\alpha\beta\zeta\delta}(\vec{x},t,t') = \frac{\gamma}{4}q^{\alpha\beta}q^{\zeta\delta} < p(r) >^2 e^{-i\omega(t-t')}$$
(16)

It would be further interesting to work out nonequilibrium and dynamical configurations via the Einstein-Langevin equations and obtain corresponding results.

It is with such strong magnitudes of fluctuations of pressure in the interiors of the relativisitic star, that these can perturb the configuration in terms of h_{ab} and ξ^a , by dissipating their fermi energy.

Other way to look at this result is due to Kubo's formalism for linear response theory, where any external agency causing perturbations in the system, is equilivalent to stochastic fluctuation inside the system causing the same effect, in absence of the external agency [8]. Hence stochastic fluctuations inside the neutron stars of such high magnitudes that we obtain in this letter, can also be compared as an effect of some equivalent external agency which can perturb the system. The F-D theorem is a two way statement, which also states that perturbations caused due to an external agency in the system can dissipate their energy in the isolated system (once the external source is removed) in the form of such stochastic fluctuations. From our result we see that perturbations induced in the star, can adiabatically dissipate into large magnitudes of stochastic fluctuations of pressure inside the star. This may also be an interesting way to study stability issues (for radial fluctuations) or if there is a possiblity of kicking the static equilibrium configuration into a dynamically non-equilibrium state. Such fluctuations in the interiors have the potential to kick a dynamically equilibrium body out of its equilibrium state, at critical points.

Whether the system is perturbed due to an external agency or noise originating in the system, the E-L formalism gives access to probe better for statistical properties of the unknown matter content in the interiors using a first principles approach.

One can thus see in this simple formulation, we have not used any information of the unknown matter content of the compact object, except for pressures, rough estimates of fermi temperature and the spacetime metric. Hence we do not in advance claim over any specifications of the interiors which are yet unknown. Rather with this simple approach, we try to analyse such systems in more detail in a new way.

Usually the adiabatic index is taken to be constant for a given star and is approximated for the outer layers, however γ can have dependence on the spatial coordinates, or more specifically a radial dependence due to different layers inside the star given by $\gamma(r)$. This is in fact an explorable direction on adiabatic index and composition of relativistic stars.

To summarize this letter, we have given a fluctuation dissipation relation, and assigned a new dissipative effect with the adiabatic index in a very crude model, but quite relevant as the first step. This paves way for stationary and dynamical models of relativistic stars which are more interesting for observational consequences, on such lines of thought. It is expected that in collapsing stars, the parameters of collapse can find association with such fluctuations of matter fields comprising them. The role of fluctuations in still unknown in this regard. However indications towards usefulness of such an approach comes from the theory of dynamical systems, critical phenomena and role that fluctuations play in other areas of Physics [17] . This area still awaits even the basic developments at present.

It is just at the beginning stages of developing statistical mechanics and showing importance of fluctuations for astrophysical systems governed by Einstein's equations that we present the F-D relation here with emphasis on the adiabatic index, the first parameter of the star to be veiwed in this fashion.

ACKNOWLEDGEMENTS

The author is thankful to K.D. Kokkotas, Rajesh Nayak, Deepak Dhar and Dipankar Bhattacharya for useful comments and discussions over the ideas presented here. Also acknowledgements are due to Jasjeet Bagla for encouraging comments over occurrence of dissipation in models of relativistic stars for an earlier draft on this idea. This work is part of the project funded by DST India over grant no. WoSA/PM-100/2016.

- Norman K. Glendenning. Compact Stars. Springer-Verlag. (1996).
- [2] James M.Lattimer, M.Prakash. Phys. Reports, 442(1-6) (2007)

- [4] Geraint Pratten, P.Schmidt, Tanja Hinderer, Nat Commun, 11, 2553 (2020).
- [5] Michele Maggiore, Gravitational Waves: Volume 1: Theory and Experiments. Oxford University Press .(2007)
- $\left[6\right]$ Seema Satin . GRG 50, 97 (2018)
- [7] Seema Satin . GRG 51, 52 (2019)
- [8] R.Kubo et.at Statistical Mechanics. North Holland (1990).
- [9] John L.Friedman & Bernard F.Shutz. The Astr.J 200:204-220 (1975)
- [10] Friedman & Stergioulas. Rotating Relativistic Stars. Cambridge University Press. (2013)

- [11] Jolien D.E. Creighton, Warren G.Anderson. Gravitational Wave Physics and Astronomy: An introduction to Theory, Experiment and Data Analysis. John Wiley & Sons (2011).
- [12] K.D.Kokkotas & Bernard G.Schmidt. Living Rev.Rel 2,2 (1999)
- [13] Seema Satin . PRD 100, 044031 (2019)
- [14] Charles W.Misner & David H.Sharp, Phys.Rev 136, 2B (1964).
- [15] B.L.Hu & Sukanya Sinha PRD 51, 1587 (1995)
- [16] B.L.Hu, E.Verdaguer Living Rev.Rel 11: 3 (2008)
- [17] R.Klages & W. Just & C.Jarzynski. Nonequilibrium Statistical Physics of Small Systems: Fluctuation Relations and Beyond. Wiley-VCH Verlag. (2013)