

# Stability of a class of entropies based on fractional calculus

Rui A. C. Ferreira

**Abstract.** In this work we argue about the Lesche stability of some systems, that are motivated by the use of fractional derivatives.

**Mathematics Subject Classification (2010).** 82A05, 94A17.

**Keywords.** Fractional derivatives, stability, entropy.

## 1. Preamble

This text is primarily motivated by the work of Ubriaco [5], namely, with the proof of the Lesche stability of a certain entropic functional presented therein. Specifically, based on the concept of the Liouville–fractional derivative, the author defined the following entropy (see [5, Section 3])

$$S(p) = \sum_{i=1}^n p_i (-\ln(p_i))^\alpha, \quad 0 < \alpha \leq 1. \quad (1.1)$$

In 1982, B. Lesche [4] introduced a way of measuring the stability of a system (see also [1]). Concretely, a system  $S$  over a probability distribution  $P = \{p_1, \dots, p_n\}$  with  $n \geq 2$  is said to be *Lesche stable* iff

$$\forall \varepsilon > 0 \exists \delta > 0 \text{ s.t. } \forall n, \forall p, p' \in P : \|p - p'\|_1 \leq \delta \Rightarrow \frac{|S(p) - S(p')|}{S_{\max}} < \varepsilon.$$

In [5, Section 4.1] the author *showed* that (1.1) is Lesche stable, using the main result of [2]. However his proof has an inconsistency, as we shall now explain: having in mind the notation of [2], Ubriaco defined in [5, formula (30)] the following function,

$$f^{-1}(t) = (-\ln(t))^\alpha - \alpha(-\ln(t))^{\alpha-1}, \quad \alpha \in (0, 1], t \in (0, 1),$$

---

The author was supported by the “Fundação para a Ciência e a Tecnologia (FCT)” through the program “Stimulus of Scientific Employment, Individual Support-2017 Call” with reference CEECIND/00640/2017.

which is nothing else but the derivative of  $t(-\ln(t))^\alpha$ . We see that  $f^{-1}(1) = \lim_{t \rightarrow 1^-} f^{-1}(t) = -\infty$ . Now, the departure formula for the general entropic expression in [2] is

$$S(p) = \sum_{i=1}^n \left[ p_i f^{-1}(p_i) - \int_{f^{-1}(0)}^{f^{-1}(p_i)} f(t) dt \right] + \int_{f^{-1}(0)}^{f^{-1}(1)} f(t) dt - f^{-1}(1),$$

where  $f$  is a monotonically decreasing function with range  $(0, 1)$ . Therefore, for  $p_i < 1$ , we see that  $S(p) \geq +\infty$  which is just nonsense. We also note that the authors assumed differentiability of  $f^{-1}$  in order to obtain [2, formula (6)], which will be used somewhere after in their analysis. Summing up, we cannot conclude from the reasoning in [5, Section 4.1] that (1.1) is Lesche stable. However, we can conclude it from the results in [3]. Indeed, from [3, Example 3] we know that, defining  $g(x) = x^\alpha$  for  $x \in [0, \infty)$ , then  $g(0) = 0$  and the function  $xg(-\ln(x))$  is concave. Therefore, the entropy (1.1) is Lesche stable.

In the next section we will introduce a function that generalizes the entropy given by (1.1). Some properties of it are presented as well.

## 2. A novel entropic function

We use the same motivation as in [5] to define a new entropy, but now by considering the *tempered fractional derivative*.

Let us start by introducing the concept (see e.g. [6]). For  $0 < \sigma < 1$  and  $\lambda \geq 0$ , the Liouville tempered fractional derivative of order  $\sigma$  is defined by

$$D^{\sigma, \lambda} u(t) = \frac{e^{-\lambda t}}{\Gamma(1 - \sigma)} \frac{d}{dt} \int_{-\infty}^t (t - s)^{-\sigma} e^{\lambda s} u(s) ds. \quad (2.1)$$

For  $\lambda = 0$  the operator (2.1) is just the Liouville fractional derivative used in [5, formula (16)]. Now we wish to calculate (2.1) at  $t = -1$  with  $u(s) = e^{-s \ln(p_i)}$ ,  $p_i < 1$  (see [5, Section 3]). We have,

$$\begin{aligned} \frac{d}{dt} \int_{-\infty}^t (t - s)^{-\sigma} e^{\lambda s} e^{-s \ln(p_i)} ds &= \frac{d}{dt} e^{t(\lambda - \ln(p_i))} \int_0^\infty u^{-\sigma} e^{-u(\lambda - \ln(p_i))} du \\ &= \frac{d}{dt} e^{t(\lambda - \ln(p_i))} (\lambda - \ln(p_i))^{\sigma-1} \Gamma(1 - \sigma) \\ &= e^{t(\lambda - \ln(p_i))} (\lambda - \ln(p_i))^\sigma \Gamma(1 - \sigma), \end{aligned}$$

where we used the definition of the gamma function  $\Gamma(t) = \int_0^\infty x^{t-1} e^{-x} dx$ . Therefore,

$$\begin{aligned} D^{\sigma, \lambda}(s \rightarrow e^{-s \ln(p_i)})(-1) &= \frac{e^{-\lambda(-1)}}{\Gamma(1 - \sigma)} e^{-1(\lambda - \ln(p_i))} (\lambda - \ln(p_i))^\sigma \Gamma(1 - \sigma) \\ &= p_i (\lambda - \ln(p_i))^\sigma, \quad p_i < 1. \end{aligned}$$

For a probability distribution  $P = \{p_1, \dots, p_n\}$  with  $n \geq 2$ , we postulate the following two parameter entropy:

$$S_{\sigma,\lambda}(p) = \sum_{i=1}^n p_i [(\lambda - \ln(p_i))^\sigma - \lambda^\sigma], \quad \sigma \in (0, 1], \lambda \geq 0. \quad (2.2)$$

Note that, for  $\sigma = 1$ , we are adopting the usual convention that  $x \ln(x)|_{x=0}$ . It is clear that  $S_{\sigma,0}$  reduces to the entropy defined by Ubriaco in (1.1). The entropy defined above satisfies the following three Shannon–Khinchin axioms: let  $\Delta_n = \{(p_1, \dots, p_n) \in \mathbb{R}^n \mid p_i \geq 0, p_1 + \dots + p_n = 1\}$ . Then,

1.  $S_n$  is nonnegative and continuous on  $\Delta_n$ .
2. For all  $(p_1, \dots, p_n) \in \Delta_n$ :  $S_n(p_1, \dots, p_n) \leq S_n(1/n, \dots, 1/n)$ .
3. For all  $(p_1, \dots, p_n) \in \Delta_n$ :  $S_{n+1}(p_1, \dots, p_n, 0) = S_n(p_1, \dots, p_n)$ .

Essentially that is due to the fact that the function defined by

$$f_{\sigma,\lambda}(x) = x [(\lambda - \ln(x))^\sigma - \lambda^\sigma], \quad x \in [0, 1],$$

where we understand for the value  $f(0) = \lim_{x \rightarrow 0^+} x [(\lambda - \ln(x))^\sigma - \lambda^\sigma] = 0$ , is concave. Moreover, defining  $g(x) = (\lambda + x)^\sigma - \lambda^\sigma$ ,  $x \geq 0$ , we get that  $g(0) = 0$  and  $g$  is concave. Therefore, using again [3, Example 3], we conclude that  $S_{\sigma,\lambda}$  given by (2.2) is Lesche stable<sup>1</sup>.

We would like to end this work remarking that  $S_{\sigma,\lambda}(p) \leq S_{\sigma,0}(p)$ . Indeed, this inequality follows from the elementary inequality  $(x+y)^\alpha \leq x^\alpha + y^\alpha$  for  $x, y \geq 0$  and  $\alpha \in (0, 1)$ .

## References

- [1] S. Abe, Stability of Tsallis entropy and instabilities of Rényi and normalized Tsallis entropies: a basis for  $q$ -exponential distributions, *Phys. Rev. E* (3) **66** (2002), no. 4, 046134, 6 pp.
- [2] S. Abe, G. Kaniadakis and A. M. Scarfone, Stabilities of generalized entropies, *J. Phys. A* **37** (2004), no. 44, 10513–10519.
- [3] X. Cao and S. Luo, On the stability of generalized entropies, *J. Phys. A* **42** (2009), no. 7, 075205, 9 pp.
- [4] B. Lesche, Instabilities of Rényi entropies, *J. Statist. Phys.* **27** (1982), no. 2, 419–422.
- [5] M. R. Ubriaco, Entropies based on fractional calculus, *Phys. Lett. A* **373** (2009), no. 30, 2516–2519.
- [6] F. Sabzikar, M. M. Meerschaert and J. Chen, Tempered fractional calculus, *J. Comput. Phys.* **293** (2015), 14–28.

Rui A. C. Ferreira

Grupo Física-Matemática, Faculdade de Ciências, Universidade de Lisboa,  
Av. Prof. Gama Pinto, 2, 1649-003, Lisboa, Portugal.

e-mail: [raferreira@fc.ul.pt](mailto:raferreira@fc.ul.pt)

---

<sup>1</sup>We note that, for  $\lambda > 0$ , the function  $f_{\sigma,\lambda}(x)$  is differentiable at  $x = 1$ . Therefore, if we define by  $f^{-1}(x) = \frac{d}{dx} x [(\lambda - \ln(x))^\sigma - \lambda^\sigma]$ , then we could actually use the results of [2] to conclude that (2.2) is Lesche stable.