# How Likely Are Large Elections Tied?

Lirong Xia, Rensselaer Polytechnic Institute, Troy, NY 12180, USA, xialirong@gmail.com

#### Abstract

Understanding the likelihood for an election to be tied is a classical topic in many disciplines including social choice, game theory, political science, and public choice. The problem is important not only as a fundamental problem in probability theory and statistics, but also because it plays a critical role in many other important issues such as indecisiveness of voting, strategic voting, privacy of voting, voting power, voter turnout, etc. Despite a large body of literature and the common belief that ties are rare, little is known about how rare ties are in large elections except for a few simple positional scoring rules under the i.i.d. uniform distribution over the votes, known as the *Impartial Culture (IC)* in social choice. In particular, little progress was made after Marchant explicitly posed the likelihood of k-way ties under IC as an open question in 2001 [38].

We give an asymptotic answer to this open question for a wide range of commonly studied voting rules under a more general and realistic model, called the *smoothed social choice framework* [66], which was inspired by the celebrated smoothed complexity analysis Spielman and Teng [59]. We prove dichotomy theorems on the smoothed likelihood of ties under positional scoring rules, edge-order-based rules, and some multi-round score-based elimination rules, which include commonly studied voting rules such as plurality, Borda, veto, maximin, Copeland, ranked pairs, Schulze, STV, and Coombs as special cases. We also complement the theoretical results by experiments on synthetic data and real-world rank data on Preflib [41]. Our main technical tool is an improved characterization of the smoothed likelihood for a Poisson multinomial variable to be in a polyhedron, by exploring the interplay between the V-representation and the matrix representation of polyhedra and might be of independent interest.

# 1 Introduction

Suppose a presidential election between two alternatives (candidates) a and b will be held soon, and there are n agents (voters). Each agent independently votes for an alternative with probability 0.5, and the alternative with more votes wins. How likely will the election end up with a tie? What if there are more than two alternatives, agents rank the alternatives, and a rank-based voting rule is used to choose the winner? What if the distribution is not independent and identically distributed (i.i.d.) and is controlled by an adversary?

Understanding the likelihood of tied elections is an important and classical topic in many disciplines including social choice, game theory, political science, and public choice, not only because it is a fundamental problem in probability theory and statistics, but also because it plays a critical role in many important issues. For example, ties are undesirable in the context of indecisiveness of voting [27], strategic voting [26, 56], privacy of voting [35], etc. On the other hand, ties are desirable in the context of voting power [3], voter turnout [19, 55], etc.

While the likelihood of ties for two alternatives is well-understood [3, 5], we are not aware of a rigorous mathematical analysis for elections with three or more alternatives except for a few simple voting rules. Previous studies were mostly done along three dimensions: (1) the voting rule used in the election, (2) the indecisiveness of the outcome, measured by the number of tied alternatives k, and (3) the statistical model for generating votes. See Section 1.1 for more discussions.

Despite these efforts, the following question largely remains open.

#### How likely are large elections tied under realistic models?

Specifically, Marchant [38] posed the likelihood of ties beyond certain positional scoring rules under the i.i.d. uniform distribution, known as the *Impartial Culture (IC)* in social choice, as an open question in 2001, but we are not aware of any progress afterwards. While IC has been a popular choice in social choice theory, it has also been widely criticized of being unrealistic [34].

In fact, the question is already highly challenging under IC as illustrated in Example 1 below. Consider the probability of 3-way ties under the Borda rule for 3 alternatives and n agents. Borda is a positional scoring rule, which scores every alternative according to its rank. Under Borda, each agent uses a linear order over the alternatives to represent his/her preferences, and the *i*-th ranked alternative gets m - i points. The winners are the alternatives with maximum total points.

**Example 1.** Let  $\mathcal{A} = \{1, 2, 3\}$  denote the set of alternatives. For each linear order R over  $\mathcal{A}$ , let  $X_R$  denote the random variable that represents the number of agents whose votes are R, when their votes are generated uniformly at random (i.e., IC). For example,  $X_{123}$  represents the multiplicity of  $1 \succ 2 \succ 3$ . Then, the probability of 3-way ties under Borda w.r.t. IC can be represented by the probability for the following system of linear equations to hold, where (1) states that alternatives 1 and 2 are tied, (2) states that 2 and 3 are tied, and (3) states that 1 and 3 are tied.

$$2X_{123} + 2X_{132} + X_{213} + X_{312} = 2X_{213} + 2X_{231} + X_{123} + X_{321}$$
(1)

$$2X_{213} + 2X_{231} + X_{123} + X_{321} = 2X_{312} + 2X_{321} + X_{132} + X_{231}$$
<sup>(2)</sup>

$$2X_{123} + 2X_{132} + X_{213} + X_{312} = 2X_{312} + 2X_{321} + X_{132} + X_{231}$$
(3)

The difficulty in accurately bounding the likelihood of ties comes from two types of statistical correlations. The first type consists of correlations among components of X. That is, for any pairs of linear orders R and W,  $X_R$  and  $X_W$  are statistically dependent. The second type consists of correlations among equations, and more generally, inequalities as we will see in the general problem studied in this paper. For example, while it is straightforward to see that (1) and (2) implies (3) in Example 1, it is unclear how much correlation exists between (1) and (2). Existing asymptotic tools

such as multivariate Central Limit Theorems and Berry-Esseen-type Theorems [6, 61, 16, 17, 54] (a.k.a. Lyapunov-type bounds) contain an  $O(n^{-0.5})$  or higher error bound, which are too coarse and do not match the lower bound that will be proved in this paper. The problem becomes more challenging for inequalities, other voting rules, other number of alternatives, other k's, and non-i.i.d. distributions over votes.

**The Model.** We address the likelihood of ties under the smoothed social choice framework [66], which is inspired by the celebrated smoothed complexity analysis [59]. We believe that the framework is more general and realistic than the extensively studied i.i.d. models, especially IC. In the framework, agents' "ground truth" preferences can be arbitrarily correlated and are chosen by an adversary, and then independent noises are added to form their votes. Mathematically, the adversary chooses a distribution  $\pi_j$  for each agent j from a set  $\Pi$  of distributions over all linear orders over the alternatives, under which the probability of various events of interest are studied, for example Condorcet's paradox and satisfaction of axioms [66].

**Our Contributions.** In this paper, we adopt the statistical model in [66] to formulate and study the smoothed likelihood of ties. Given an (irresolute) voting rule  $r, 2 \le k \le m$ , and  $n \in \mathbb{N}$  agents, the *max-adversary* aims to maximize the likelihood of k-way ties, denoted by  $\widetilde{\text{Tie}}_{\Pi}^{\max}(r,k,n)$ , by choosing  $\vec{\pi} = (\pi_1, \ldots, \pi_n) \in \Pi^n$ . Formally,

$$\widetilde{\mathrm{Tie}}_{\Pi}^{\max}(r,k,n) \triangleq \sup_{\vec{\pi} \in \Pi^n} \Pr_{P \sim \vec{\pi}} \left( |r(P)| = k \right) \tag{4}$$

Similarly, the *min-adversary* aims to minimize the likelihood of k-way ties defined as follows:

$$\widetilde{\mathrm{Tie}}_{\Pi}^{\min}(r,k,n) \triangleq \inf_{\vec{\pi} \in \Pi^n} \mathrm{Pr}_{P \sim \vec{\pi}} \left( |r(P)| = k \right)$$
(5)

We call  $\widetilde{\text{Tie}}_{\Pi}^{\max}(r, k, n)$  (respectively,  $\widetilde{\text{Tie}}_{\Pi}^{\min}(r, k, n)$ ) the max (respectively, min) smoothed likelihood of ties. When  $\Pi$  consists of a single distribution  $\pi$ ,  $\widetilde{\text{Tie}}_{\Pi}^{\max}(r, k, n)$  and  $\widetilde{\text{Tie}}_{\Pi}^{\min}(r, k, n)$  coincide with each other and become the likelihood of ties under i.i.d. distribution  $\pi$ . In particular, when  $\Pi = \{\pi_{\text{uni}}\}$ , where  $\pi_{\text{uni}}$  is the uniform distribution,  $\widetilde{\text{Tie}}_{\Pi}^{\max}(r, k, n)$  and  $\widetilde{\text{Tie}}_{\Pi}^{\min}(r, k, n)$  become the classical analysis of ties under IC. As discussed in [66], the smoothed social choice framework allows agents' ground truth preferences to be arbitrarily correlated, while the noises are independent, which is a standard assumption in many literatures such as behavior science, economics, statistics, and smoothed complexity analysis.

Our main technical results are asymptotic characterizations of the smoothed likelihood of ties for a fixed number of at least three alternatives  $(m \ge 3)$  in large elections  $(n \to \infty)$ . Informally, our main results can be summarized as follows.

Main results: smoothed likelihood of ties, informally put. Under mild assumptions on  $\Pi$ , for many commonly studied voting rules r, for any fixed  $m \ge 3$ , any  $2 \le k \le m$ , and any  $n \in \mathbb{N}$ ,  $\widetilde{Tie}_{\Pi}^{\max}(r, k, n)$  (respectively,  $\widetilde{Tie}_{\Pi}^{\min}(r_{\vec{s}}, k, n)$ ) is either 0,  $\exp(-\Theta(n))$ , or  $\Theta(poly^{-1}(n))$ .

More precisely, we prove Theorem 3 for integer positional scoring rules (including plurality, Borda, veto), Theorem 4 for edge-order-based rules (including maximin, Schulze, ranked pairs, and Copeland, see Definition 11), and Theorem 5 for STV and Coombs. The formal statements of the theorems also characterize the condition for each (0, exponential, or polynomial) case as well as asymptotically tight bounds in the polynomial cases.

When ties are undesirable, e.g., in the context of indecisiveness of voting, strategic voting, or privacy, a low max smoothed likelihood is good news. When ties are desirable, e.g., in the context

of voting power and voter turnout, a high min smoothed likelihood is good news. Our theorems therefore completely characterize conditions for good news in different contexts.

Straightforward applications of our theorems answer the open question by Marchant [38] for many commonly studied voting rules as summarized in Table 1 below.

Table 1: Probability of k-way ties  $(2 \le k \le m)$  under some commonly studied voting rules w.r.t. IC. For Copeland<sub> $\alpha$ </sub>,  $l_{\alpha}$  is the minimum positive integer s.t.  $\alpha l_{\alpha} \in \mathbb{Z}$ .

Int. Pos. scoring (Coro. 1) STV and Coombs (Prop. 6)		maximin (Prop. 3) Schulze (Prop. 4)	ranked pairs (Prop. 5)			
ſ	if no profile of $n$ votes		Lower	Upper		
K .	contains a $k$ -way tie	$\Theta(n^{-\frac{k-1}{2}})$	$\int \Omega(n^{-\frac{k-1}{2}})$	$-\Omega(\frac{\log k}{\log\log k})$		
$\bigcup \Theta(n^{-\frac{k-1}{2}})$	otherwise		$\left[ \Omega(n^{-\frac{\lceil \log k \rceil}{2}}) \text{ if } m \ge k + 5 \lceil \log k \rceil \right]$	$n$ $\log \log \kappa$		
Copeland <sub><math>\alpha</math></sub> $(0 \le \alpha \le 1)$ (Prop. 2)						
	( 0 if	$2 \nmid n, 2 \mid k, \text{and } k \geq m$ -	- 1			
	$\left\{ \begin{array}{ll} \Theta(n^{-\frac{k}{4}}) & \text{ if } \end{array} \right.$	$\begin{array}{c c} 1 & 2 &   & n, 2 &   & k, and \\ \hline & (2) & k \\ & (3) & k \end{array}$	x = m, or $x = m - 1$ and $\alpha \ge \frac{1}{2}$ , or $x = m - 1$ and $k \le l_{\alpha}(l_{\alpha} + 1)$			
	$\Theta\left(n^{-\frac{l_{\alpha}(l_{\alpha}+1)}{4}}\right)$ if	$2 \mid n, 2 \mid k, k = m - 1, c$	$\alpha < \frac{1}{2}$ , and $k > l_{\alpha}(l_{\alpha} + 1)$			
	Ο(1)Ο	therwise (i.e., if $2 \nmid k$ or	$k \le m - 2)$			

Roughly speaking, Table 1 reveals the following ranking over the voting rules w.r.t. their likelihood of k-way ties under IC, for every  $2 \le k \le m$ .

{Int. Pos. Scoring, STV, Coombs}  $\leq$  {maximin, Schulze}  $\leq$  ranked pairs  $\leq$  Copeland

A closely related question is the likelihood of any-way ties, sometimes referred to as indecisiveness [38], under IC, i.e., the election admits a k-way tie for any  $2 \le k \le m$ . It is not hard to see from Table 1 that such likelihood is dominated by the probability of 2-way ties and is either 0 or  $\Theta(\frac{1}{\sqrt{n}})$ for all rules in the table except ranked pairs and Copeland, which are covered by Proposition 5 and Proposition 2, respectively. To the best of our knowledge, these results are new, except for plurality and Borda. Experiments on synthetic data generated from IC confirm these observations, while experiments on Preflib data [41] reveal a difference order, where ties are rare under Borda (1.6%) and Copeland (2.6%), and are quite common under veto (31.3%) due to situations where m > n.

**Technical Innovations.** The proofs of the smoothed likelihood of ties in this paper follow the same high-level idea. We first model the existence of a k-way tie by systems of linear inequalities that are similar to the ones in Example 1. In this way, the likelihood of ties becomes the likelihood for the histogram of the randomly generated profile, which is a *Poisson multivariate variable (PMV)*, to be in the polyhedron  $\mathcal{H}$  represented by the linear inequalities. Then, we prove a dichotomous characterization (Theorem 1) for a PMV to be in  $\mathcal{H}$ , and finally apply Theorem 1 (more precisely, its extension Theorem 2 to unions of multiple polyhedra) to characterize the smoothed likelihood of ties.

More precisely, given  $n, q \in \mathbb{N}$  and a vector  $\vec{\pi} = (\pi_1, \ldots, \pi_n)$  of n distributions over  $\{1, \ldots, q\}$ , an (n, q)-PMV is denoted by  $\vec{X}_{\vec{\pi}}$ , which represents the histogram of n independent random variables whose distributions are  $\{\pi_1, \ldots, \pi_n\}$ , respectively.

**Theorem 1.** (The PMV-in-polyhedron theorem, informally put). Let  $\mathcal{H}$  denote a polyhedron and  $\Pi$  denote a set of distributions that satisfy some mild conditions, for any  $n \in \mathbb{N}$ ,

$$\sup_{\vec{\pi}\in\Pi^n} \Pr(\vec{X}_{\vec{\pi}}\in\mathcal{H}) \text{ is } 0, \exp(-\Theta(n)), \text{ or } \Theta(poly(n)), \text{ and} \\ \inf_{\vec{\pi}\in\Pi^n} \Pr(\vec{X}_{\vec{\pi}}\in\mathcal{H}) \text{ is } 0, \exp(-\Theta(n)), \text{ or } \Theta(poly(n)) \end{cases}$$

The bounds are asymptotically tight and the formal statement of the theorem also characterizes conditions for the 0, exponential, and polynomial cases, respectively. As commented after Example 1, we do not see a way to prove Theorem 1 by straightforward applications of existing asymptotic tools. We also believe that Theorem 1 is a useful tool to study the smoothed likelihood of many events of interest in social choice as commented in [66].

### 1.1 Related Work and Discussions

**Ties in elections.** The importance of estimating likelihood of ties has been widely acknowledged, for example, as Mulligan and Hunter commented: "Perhaps it is common knowledge that civic elections are not often decided by one vote...a precise calculation of the frequency of a pivotal vote can contribute to our understanding of how many, if any, votes might be rationally and instrumentally cast" [47]. In practice, however, ties are not as rare as commonly believed, even in high-stakes elections, and have led to pitfalls and consequent modifications of electoral systems and constitutional laws. For example, in the 1800 US presidential election, Jefferson and Burr tied in the electoral college votes. By the Constitution, the House of Representatives should vote until a candidate wins the majority. However, in the subsequent 35 rounds of deadlocked voting, none of the two candidates got the majority. Eventually, Jefferson won the 36th revote to become the president. This "had demonstrated a fundamental flaw with the Constitution. As a result, the Twelfth Amendment to the Constitution was introduced and ratified" [9].

Three-or-more-way ties. Nowadays, legislators are well-aware of the possibility of two-way ties in elections and have specified tie-breaking mechanisms to handle them. However, three-or-moreway ties have not received their deserved attention and are sometimes overlooked. For example, the 2019 Code of Alabama Section 17-12-23 states: "In all elections where there is a tie between the two highest candidates for the same office, for all county or precinct offices, it shall be decided by lot by the sheriff of the county in the presence of the candidates". The Code does not specify what action should be taken when three or more candidates are tied for the first place. Results in this paper characterize how rare this happens, so that the legislators can make an informed decision about whether the loophole is a significant concern in practice, and in case it is, how to fix it.

**Smoothed analysis.** There is a large body of literature on the applications of smoothed analysis to mathematical programming, machine learning, numerical analysis, discrete math, combinatorial optimization, etc., see [59] for a survey. Smoothed analysis has also been applied to various problems in economics, for example price of anarchy [11] and market equilibrium [32]. In a recent position paper, Baumeister, Hogrebe, and Rothe [4] proposed a Mallows-based model to conduct smoothed analysis on computational aspects of social choice and commented that the model can be used to analyze voting paradoxes and ties, but the paper does not contain technical results. [66] independently proposed to conduct smoothed analysis for paradoxes and impossibility theorems in social choice, characterized the smoothed likelihood of Condorcet's voting paradox and the ANR impossibility theorem, and proposed a new tie-breaking mechanism. We only use the probabilistic model in [66] and topic-wise, our paper is different from [66], because we formulate and study the smoothed likelihood of ties under commonly studied voting rules, which was not studied in [66].

**Technical novelty.** We believe that the main technical tool of this paper (Theorem 1) is a significant and non-trivial extension of Lemma 1 in [66] to arbitrary polyhedron represented by an

integer matrix, every n, and the min-adversary. More discussions can be found in the remark after Theorem 1. We believe that Theorem 1 is a useful tool to analyze smoothed likelihood of many other problems of interest in social choice. For example, all results in [66] can be immediately strengthened by Theorem 1.

**Previous work on likelihood of ties.** The following table summarizes previous works that are closest to this paper, whose main contributions are characterizations of likelihood of ties.

Paper	m	k	Voting rule	Distribution
[5]	2	2	majority	two groups, i.i.d. within each group
[39] [10]	2	2	majority	i.i.d. w.r.t. an uncertain distribution
[27]	any $m \in \mathbb{N}$	$2 \le k \le m$	plurality	uniformly i.i.d. (IC)
[28]	3	$2 \le k \le m$	Borda	uniformly i.i.d. (IC)
[38]	any $m \in \mathbb{N}$	k = m	certain scoring rules	uniformly i.i.d. over all score vectors

More precisely, Beck [5] studied the probability of ties under the majority rule (over two alternatives) with two groups of agents, whose votes are i.i.d. within each group. Margolis [39] and Chamberlain and Rothschild [10] focused on the majority rule for two alternatives, where agents' preferences are i.i.d. according to a randomly generated distribution. Gillett [27] studied probability of all-way ties (i.e., k = m) under plurality w.r.t. IC for arbitrary numbers of alternatives and agents. Gillett [28] obtained a closed-form formula for Borda indecisiveness (two or more alternatives being tied) for m = 3 w.r.t. IC. Marchant [38] considered a class of scoring rules where each agent can choose a score vector from a given scoring vectors set (SVS), and characterized the asymptotic probability of m-way ties under a class of scoring rule to be  $\Theta(n^{\frac{1-m}{2}})$  w.r.t. the i.i.d. uniform distribution over all SVS. This result can be applied to Borda and approval voting but cannot be applied to plurality. Marchant [38] also noted that Domb [18] obtained equivalent formulas for m = 3 under Borda. As discussed earlier, the smoothed social choice framework used in our paper is more general. In particular, corollaries of our theorems answer the open questions proposed by Marchant [38] and reveal a ranking over these rules according to the likelihood of ties under IC as summarized in Table 1.

**Previous work related to likelihood of ties.** [33] studied the setting where the agents are partitioned into multiple groups, and within each group, agents' votes are generated from the *impartial anonymous culture (IAC)* model. The smoothed social choice framework and IAC are not directly comparable. The former is more general in the sense that agents' "ground truth" preferences are arbitrarily correlated. The latter is more general in the sense that agents' "noises" are not independent. There is also a line of empirical and mixed empirical-theoretical work on the likelihood of ties under the US electoral college system [24, 25]. Studying the smoothed likelihood of ties under these settings are left for future work.

The probability of tied elections is closely related to the probability for a single voter to be pivotal, sometimes called *voting power*, which plays an important role in the paradox of voting [19] and in definitions of power indices in cooperative game theory. There is a large body of work on voting games, where the probability for a voter to be pivotal, which is equivalent to the likelihood of ties among other voters, plays a central role in the analysis of voters' strategic behavior. Examples include seminal works [1, 21, 48], and more recent work [49]. It is not hard to see that the voting power for two alternatives or for multiple alternatives under the plurality rule almost equals to

the probability of tied elections with one less vote under certain tie-breaking rules, as pointed out in [29]. For three or more alternatives the two problems are closely related but technically different, which we leave for future work.

The likelihood of ties is also related to the manipulability of voting rules [14]—if an election is not tied, then no single agent can change the outcome, therefore no agent alone has incentive to cast a manipulative vote. Our results are related to but different from the typical-case analysis of manipulability in the literature [53, 67, 45, 65] and the quantitative Gibbard-Satterthwaite theorem [23, 46], where votes are assumed to be i.i.d. Likelihood of ties are also related to but different from the margin of victory [37, 64] and more broadly, bribery and control in elections [20].

Adding noise to study ties. We are not aware of a previous work that characterized the smoothed likelihood of ties as we do in this paper. The idea of adding noise to study ties is not new. In the definition of *resolvability* in [60], an additional vote is added to break ties. In [22], irresolute voting rules were defined by taking the union of winners under profiles around a given profile. Another *resolvability* studied in the literature (see e.g., Formulation#1 in Section 4.2 of  $[58]^1$ ) requires that the probability of ties goes to 0 under the voting rule w.r.t. IC, which is closely related to the literature in the indecisiveness of voting. Our setting and results are more general because IC is a special case of the smoothed social choice framework, and our results also characterize the rate of convergence.

**Computational aspects of tie-breaking.** There is a large body of recent work on computational aspects of tie-breaking. [13] proposed the *parallel universe tie-breaking (PUT)* for multi-stage voting rules and characterized the complexity of the STV rule. [7] characterize the complexity of PUT under ranked pairs, whose smoothed likelihood of ties is studied in this paper. [40] characterized complexity of PUT under other multi-stage voting rules such as Baldwin and Coombs. [51, 50, 2, 52] investigated the effect of different tie-breaking mechanisms to the complexity of manipulation. [22] propose a general way of defining ties under generalized scoring rules. [62] proposed practical AI algorithms for computing PUT under STV and ranked pairs.

# 2 Preliminaries

**Basic Setting.** For any  $q \in \mathbb{N}$ , we let  $[q] = \{1, \ldots, q\}$ . Let  $\mathcal{A} = [m]$  denote the set of  $m \geq 3$ alternatives. Let  $\mathcal{L}(\mathcal{A})$  denote the set of all linear orders over  $\mathcal{A}$ . Let  $n \in \mathbb{N}$  denote the number of agents. Each agent uses a linear order to represent his or her preferences, called a *vote*. The vector of n agents' votes, denoted by P, is called a *(preference) profile*, sometimes called an n-profile. A fractional profile is a preference profile P together with a possibly non-integer and possibly negative weight vector  $\vec{\omega}_P = (\omega_R : R \in P) \in \mathbb{R}^n$  for the votes in P. It follows that a non-fractional profile is a fractional profile with uniform weight, namely  $\vec{\omega}_P = \vec{1}$ . Sometimes we slightly abuse the notation by omitting the weight vector when it is clear from the context or when  $\vec{\omega}_P = \vec{1}$ .

For any (fractional) profile P, let  $\operatorname{Hist}(P) \in \mathbb{Z}_{\geq 0}^{m!}$  denote the anonymized profile of P, also called the *histogram* of P, which contains the total weight of every linear order in  $\mathcal{L}(\mathcal{A})$  according to P. An *(irresolute) voting rule* r is a mapping from a profile to a non-empty set of winners in  $\mathcal{A}$ . Below we recall the definitions of several commonly studied voting rules.

Integer positional scoring rules. An *(integer)* positional scoring rule is characterized by an integer scoring vector  $\vec{s} = (s_1, \ldots, s_m) \in \mathbb{Z}^m$  with  $s_1 \ge s_2 \ge \cdots \ge s_m$  and  $s_1 > s_m$ . For any

<sup>&</sup>lt;sup>1</sup>Wikipedia [63] contributes this definition to Douglas R. Woodall but we were not able to find a formal reference.

alternative a and any linear order  $R \in \mathcal{L}(\mathcal{A})$ , we let  $\vec{s}(R, a) = s_i$ , where *i* is the rank of *a* in *R*. Given a profile *P* with weights  $\vec{\omega}_P$ , the positional scoring rule  $r_{\vec{s}}$  chooses all alternatives *a* with maximum  $\sum_{R \in P} \omega_R \cdot s(R, a)$ . For example, *plurality* uses the scoring vector  $(1, 0, \ldots, 0)$ , *Borda* uses the scoring vector  $(m - 1, m - 2, \ldots, 0)$ , and veto uses the scoring vector  $(1, \ldots, 1, 0)$ .

Weighted Majority Graphs. For any (fractional) profile P and any pair of alternatives a, b, let  $P[a \succ b]$  denote the total weight of votes in P where a is preferred to b. Let WMG(P) denote the weighted majority graph of P, whose vertices are  $\mathcal{A}$  and whose weight on edge  $a \rightarrow b$ is  $w_P(a, b) = P[a \succ b] - P[b \succ a]$ . Sometimes a distribution  $\pi$  over  $\mathcal{L}(\mathcal{A})$  is viewed as a fractional profile, where for each  $R \in \mathcal{L}(\mathcal{A})$  the weight on R is  $\pi(R)$ . In this case we let WMG( $\pi$ ) denote the weighted majority graph of the fractional profile represented by  $\pi$ .

A voting rule is said to be *weighted-majority-graph-based (WMG-based)* if its winners only depend on the WMG of the input profile. In this paper we consider the following commonly studied WMG-based rules.

- Copeland. The Copeland rule is parameterized by a number  $0 \le \alpha \le 1$ , and is therefore denoted by Copeland<sub> $\alpha$ </sub>, or Cd<sub> $\alpha$ </sub> for short. For any fractional profile *P*, an alternative *a* gets 1 point for each other alternative it beats in their head-to-head competition, and gets  $\alpha$  points for each tie. Copeland<sub> $\alpha$ </sub> chooses all alternatives with the highest total score as the winners.
- Maximin. For each alternative a, its min-score is defined to be  $\min_{b \in \mathcal{A}} w_P(a, b)$ . Maximin, denoted by MM, chooses all alternatives with the max min-score as the winners.
- Ranked pairs. Given a profile P, an alternative a is a winner under ranked pairs (denoted by RP) if there exists a way to fix edges in WMG(P) one by one in a non-increasing order w.r.t. their weights (and sometimes break ties), unless it creates a cycle with previously fixed edges, so that after all edges are considered, a has no incoming edge. This is known as the parallel-universes tie-breaking (PUT) [13].
- Schulze. For any directed path in the WMG, its strength is defined to be the minimum weight on any single edge along the path. For any pair of alternatives a, b, let s[a, b] be the highest weight among all paths from a to b. Then, we write  $a \succeq b$  if and only if  $s[a, b] \ge s[b, a]$ , and Schulze [58] proved that the strict version of this binary relation, denoted by  $\succ$ , is transitive. The Schulze rule, denoted by Sch, chooses all alternatives a such that for all other alternatives b, we have  $a \succeq b$ .

Multi-round score-based elimination (MRSE) rules. Another large class of voting rules studied in this paper select the winner(s) in m - 1 rounds. In each round, an integer positional scoring rule is used to rank the remaining alternatives, and a *loser* (an alternative with the minimum total score) is removed from the election. Like in ranked pairs, PUT is used to select winners—an alternative *a* is a winner if there is a way to break ties among losers so that *a* is the remaining alternative after m - 1 rounds. For example, the STV rule uses the plurality rule in each round and the Coombs rule uses the veto rule in each round.

We now recall the statistical model used in the smoothed social choice framework [66].

**Definition 1** (Single-Agent Preference Model [66]). A single-agent preference model is denoted by  $\mathcal{M} = (\Theta, \mathcal{L}(\mathcal{A}), \Pi)$ , where  $\Theta$  is the parameter space,  $\mathcal{L}(\mathcal{A})$  is the sample space, and  $\Pi$ consists of distributions indexed by  $\Theta$ .  $\mathcal{M}$  is strictly positive if there exists  $\epsilon > 0$  such that the probability of any linear order under any distribution in  $\Pi$  is at least  $\epsilon$ .  $\mathcal{M}$  is closed if  $\Pi$  (which is a subset of the probability simplex in  $\mathbb{R}^{m!}$ ) is a closed set in  $\mathbb{R}^{m!}$ . For example, given  $0 < \varphi \leq \overline{\varphi} \leq 1$ , in the single-agent Mallows model [66, Example 2 in the appendix], denoted by  $\mathcal{M}_{[\varphi,\overline{\varphi}]}$ , we have  $\Theta = \mathcal{L}(\mathcal{A}) \times [\varphi,\overline{\varphi}]$ . For any  $\varphi \in [\varphi,\overline{\varphi}]$  and any  $R \in \mathcal{L}(\mathcal{A})$ ,  $\pi_{(R,\varphi)} \in \Pi$  is the Mallows distribution with central ranking R and dispersion parameter  $\varphi$ . That is, for any  $W \in \mathcal{L}(\mathcal{A})$ ,  $\pi_{(R,\varphi)}(W) = \varphi^{\mathrm{KT}(R,W)}/Z_{\varphi}$ , where  $\mathrm{KT}(R,W)$  is the Kendall Tau distance between R and W, namely the number of pairwise disagreements between R and W, and  $Z_{\varphi}$  is the normalization constant. It follows that  $\mathcal{M}_{[\varphi,\overline{\varphi}]}$  is strictly positive, closed, and CH(\Pi) contains the uniform distribution over all rankings, denoted by  $\pi_{\mathrm{uni}}$ .

**Definition 2** (Smoothed likelihood of ties). Given a voting rule r, a single-agent preference model  $\mathcal{M} = (\Theta, \mathcal{L}(\mathcal{A}), \Pi), 2 \leq k \leq m$ , and  $n \in \mathbb{N}$ , the max (respectively, min) smoothed likelihood of (k-way) ties is defined as  $\widetilde{Tie}_{\Pi}^{\max}(r, k, n)$  in (4) (respectively,  $\widetilde{Tie}_{\Pi}^{\min}(r, k, n)$  in (5)).

# **3** PMV-in-Polyhedron Problem and Main Technical Theorems

We first formally define PMV and the PMV-in-polyhedron problem studied in this paper.

**Definition 3** (Poisson multivariate variables (PMVs)). Given any  $q, n \in \mathbb{N}$  and any vector  $\vec{\pi}$  of n distributions over [q], we let  $\vec{X}_{\vec{\pi}}$  denote the (n, q)-PMV that corresponds to  $\vec{\pi}$ . That is, let  $Y_1, \ldots, Y_n$  denote n independent random variables over [q] such that for any  $j \leq n, Y_j$  is distributed as  $\pi_j$ . For any  $1 \leq i \leq q$ , the *i*-th component of  $\vec{X}_{\vec{\pi}}$  is the number of  $Y_j$ 's that take value *i*.

**Definition 4** (The PMV-in-polyhedron problem). Given  $q \in \mathbb{N}$ , a polyhedron  $\mathcal{H} \subseteq \mathbb{R}^q$ , and a set  $\Pi$  of distributions over [q], we are interested in

the upper bound  $\sup_{\vec{\pi}\in\Pi^n} \Pr(\vec{X}_{\vec{\pi}}\in\mathcal{H})$ , and the lower bound  $\inf_{\vec{\pi}\in\Pi^n} \Pr(\vec{X}_{\vec{\pi}}\in\mathcal{H})$ 

In words, the former (respectively, latter) is the maximum (respectively, minimum) probability for the (n, q)-PMV to be in  $\mathcal{H}$ , where the distribution for each of the *n* variables is chosen from  $\Pi$ .

To present the theorem, we introduce some notation followed by an example. Given  $q \in \mathbb{N}, L \in \mathbb{N}$ , an  $L \times q$  integer matrix **A**, a q-dimensional row vector  $\vec{b}$ , and an  $n \in \mathbb{N}$ , we define  $\mathcal{H}, \mathcal{H}_{\leq 0}, \mathcal{H}_n$ , and  $\mathcal{H}_n^{\mathbb{Z}}$  as follows.

$$\mathcal{H} \triangleq \left\{ \vec{x} \in \mathbb{R}^q : \mathbf{A} \cdot (\vec{x})^\top \leq \left( \vec{b} \right)^\top \right\}, \quad \mathcal{H}_{\leq 0} \triangleq \left\{ \vec{x} \in \mathbb{R}^q : \mathbf{A} \cdot (\vec{x})^\top \leq \left( \vec{0} \right)^\top \right\}, \\ \mathcal{H}_n \triangleq \left\{ \vec{x} \in \mathcal{H} \cap \mathbb{R}^q_{\geq 0} : \vec{x} \cdot \vec{1} = n \right\}, \quad \mathcal{H}_n^\mathbb{Z} \triangleq \mathcal{H}_n \cap \mathbb{Z}^q_{\geq 0}.$$

That is,  $\mathcal{H}$  is the polyhedron represented by  $\mathbf{A}$  and  $\vec{b}$ ;  $\mathcal{H}_{\leq 0}$  is the *characteristic cone* of  $\mathcal{H}$ ,  $\mathcal{H}_n$  consists of non-negative vectors in  $\mathcal{H}$  whose  $L_1$  norm is n, and  $\mathcal{H}_n^{\mathbb{Z}}$  consists of non-negative integer vectors in  $\mathcal{H}_n$ . By definition,  $\mathcal{H}_n^{\mathbb{Z}} \subseteq \mathcal{H}_n \subseteq \mathcal{H}$ . Let dim $(\mathcal{H}_{\leq 0})$  denote the *dimension* of  $\mathcal{H}_{\leq 0}$ , i.e., the dimension of the minimal linear subspace of  $\mathbb{R}^q$  that contains  $\mathcal{H}_{\leq 0}$ .

Throughout the paper, we assume that the set of distribution  $\Pi$  is *strictly positive* and *closed*, which are mild assumptions as discussed in [66], formally defined as follows.

**Definition 5** (Strictly positive and closed  $\Pi$ ). Given any  $q \in \mathbb{N}$ . A probability distribution  $\pi$ over [q] is strictly positive (by  $\epsilon$ ) for some  $\epsilon > 0$ , if for all  $i \in [q]$ ,  $\pi(i) \ge \epsilon$ . We say that a set  $\Pi$ of distributions over [q] is strictly positive (by  $\epsilon$ ) for some  $\epsilon > 0$ , if every  $\pi \in \Pi$  is strictly positive by  $\epsilon$ .  $\Pi$  is closed if it is a closed set in  $\mathbb{R}^q$ .

Let  $CH(\Pi)$  denote the convex hull of  $\Pi$  and let  $Cone(\Pi)$  denote the convex cone generated by  $\Pi$ .

**Example 2.** Figure 1 illustrates two examples with q = 2 and  $\Pi = \{\pi_1, \pi_2\}$ , where  $\pi_1 = (\frac{1}{3}, \frac{2}{3})$ and  $\pi_2 = (\frac{1}{2}, \frac{1}{2})$ . In both examples,  $CH(\Pi)$  is the line segment between  $\pi_1$  and  $\pi_2$ ,  $Cone(\Pi)$  is the shaded area,  $\mathcal{H}$  is the red area,  $\mathcal{H}_{\leq 0}$  is the blue area, the intersection of  $\mathcal{H}$  and  $\mathcal{H}_{\leq 0}$  is the purple area, and  $\mathcal{H}_n$  is the green line segment. A key difference between Figure 1 (a) and (b) is whether  $CH(\Pi) \cap \mathcal{H}_{\leq 0} = \emptyset$  (which is true in Figure 1 (a) but not in (b)). Also it is possible that  $\mathcal{H}_{\leq 0} \nsubseteq \mathcal{H}$ , as can be seen in Figure 1 (b).



Figure 1: Two examples of  $\mathcal{H}$ ,  $\mathcal{H}_{\leq 0}$ ,  $\mathcal{H}_n$ ,  $CH(\Pi)$ , and  $Cone(\Pi)$ .

A high-level attempt at the PMV-in-Polyhedron problem. Before formally presenting the theorem, let us take a high-level attempt to develop intuition. Take the upper bound  $\sup_{\vec{\pi}\in\Pi^n} \Pr(\vec{X}_{\vec{\pi}}\in\mathcal{H})$  for example, there are three cases.

• The 0 case. Clearly, if  $\mathcal{H}$  does not contain any non-negative integer whose  $L_1$  norm is n, which is equivalent to  $\mathcal{H}_n^{\mathbb{Z}} = \emptyset$ , then  $\sup_{\vec{\pi} \in \Pi^n} \Pr(\vec{X}_{\vec{\pi}} \in \mathcal{H}) = 0$ .

• The exponential case (Figure 1 (a)). Suppose  $\mathcal{H}_n^{\mathbb{Z}} \neq \emptyset$ . For any  $\vec{\pi} = (\pi_1, \ldots, \pi_n) \in \Pi^n$  chosen by the max-adversary, we have  $\vec{\pi} \cdot \vec{1} = \sum_{j=1}^n \pi_j \in \text{Cone}(\Pi)$ . According to various multivariate central limit theorems,  $\vec{X}_{\vec{\pi}}$  is "centered" around an  $\Theta(\sqrt{n})$  neighborhood of  $\vec{\pi} \cdot \vec{1}$  with high probability. Therefore, if  $\vec{\pi} \cdot \vec{1}$  is  $\Theta(n)$  away from  $\mathcal{H}$ , then  $\sup_{\vec{\pi} \in \Pi^n} \Pr(\vec{X}_{\vec{\pi}} \in \mathcal{H})$  is exponentially small. This happens when  $\text{CH}(\Pi) \cap \mathcal{H}_{\leq 0} = \emptyset$  as shown in Figure 1 (a).

• The polynomial case (Figure 1 (b)). Otherwise we have  $\operatorname{CH}(\Pi) \cap \mathcal{H}_{\leq 0} \neq \emptyset$  as shown in Figure 1 (b). In this case, the max-adversary can choose  $\vec{\pi} \in \Pi^n$  such that  $\vec{\pi} \cdot \vec{1}$  is either in  $\operatorname{Cone}(\Pi)$  or close to it, which means that  $\sup_{\vec{\pi} \in \Pi^n} \Pr(\vec{X}_{\vec{\pi}} \in \mathcal{H})$  should be larger than that in the exponential case. However, it is not immediately clear that the probability is polynomial, because  $\vec{\pi} \cdot \vec{1}$  being close to  $\operatorname{Cone}(\Pi)$  and  $\mathcal{H}_n^{\mathbb{Z}} \neq \emptyset$  does not immediately imply that  $\vec{X}_{\vec{\pi}}$  is close to  $\mathcal{H}_n^{\mathbb{Z}}$ . Even if we assume that  $\vec{X}_{\vec{\pi}}$  is close to some (integer) vectors in  $\mathcal{H}_n^{\mathbb{Z}}$ , it is unclear how "dense" such vectors are in the  $\Theta(\sqrt{n})$  neighborhood of  $\vec{\pi} \cdot \vec{1}$ , which  $\vec{X}_{\vec{\pi}}$  falls into with high probability. In fact, accurately bounding the probability in the polynomial case is the most challenging part of the problem, because existing asymptotic tools fail to work due to their  $O(n^{-0.5})$  error terms.

The main technical theorem below confirms the intuition developed above when  $\Pi$  is closed and strictly positive (see Definition 5), and the answer to the polynomial case is  $\Theta\left(n^{\frac{\dim(\mathcal{H}_{\leq 0})-q}{2}}\right)$ , which is often much smaller than  $n^{-0.5}$ . **Theorem 1** (Smoothed Likelihood of PMV-in-polyhedron). Given any  $q \in \mathbb{N}$ , any closed and strictly positive  $\Pi$  over [q], and any polyhedron  $\mathcal{H}$  characterized by an integer matrix  $\mathbf{A}$ , for any  $n \in \mathbb{N}$ ,

$$\sup_{\vec{\pi}\in\Pi^{n}}\Pr\left(\vec{X}_{\vec{\pi}}\in\mathcal{H}\right) = \begin{cases} 0 & \text{if } \mathcal{H}_{n}^{\mathbb{Z}} = \emptyset \\ \exp(-\Theta(n)) & \text{if } \mathcal{H}_{n}^{\mathbb{Z}} \neq \emptyset \text{ and } \mathcal{H}_{\leqslant 0} \cap CH(\Pi) = \emptyset \\ \Theta\left(n^{\frac{\dim(\mathcal{H}_{\leqslant 0})-q}{2}}\right) & \text{otherwise } (i.e. \ \mathcal{H}_{n}^{\mathbb{Z}} \neq \emptyset \text{ and } \mathcal{H}_{\leqslant 0} \cap CH(\Pi) \neq \emptyset) \end{cases}$$
$$\inf_{\vec{\pi}\in\Pi^{n}}\Pr\left(\vec{X}_{\vec{\pi}}\in\mathcal{H}\right) = \begin{cases} 0 & \text{if } \mathcal{H}_{n}^{\mathbb{Z}} = \emptyset \\ \exp(-\Theta(n)) & \text{if } \mathcal{H}_{n}^{\mathbb{Z}} \neq \emptyset \text{ and} \\ \Theta\left(n^{\frac{\dim(\mathcal{H}_{\leqslant 0})-q}{2}}\right) & \text{otherwise } (i.e. \ \mathcal{H}_{n}^{\mathbb{Z}} \neq \emptyset \text{ and } CH(\Pi) \subseteq \mathcal{H}_{\leqslant 0}) \end{cases}$$

Remarks on the power of Theorem 1. We believe that the main power of Theorem 2 is that it provides a systematic way of reducing probabilistic analysis (asymptotically tight upper and lower bounds for the PMV-in-Polyhedron problem) to worst-case non-probabilistic analysis, which are often easy to verify. In particular, when  $CH(\Pi)$  can be represented by the convex hull of a finite number of vectors, whether  $\mathcal{H}_{\leq 0} \cap CH(\Pi) = \emptyset$  and/or  $CH(\Pi) \not\subseteq \mathcal{H}_{\leq 0}$  can be verified by linear programming. Take the sup part of Theorem 1 in the setting of Example 2 for instance. Suppose  $\mathcal{H}_n^{\mathbb{Z}} \neq \emptyset$ .

• In Figure 1 (a), it is easy to see that  $\mathcal{H}_{\leq 0} \cap CH(\Pi) = \emptyset$ . Therefore,

$$\forall \vec{\pi} \in \Pi^n, \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}\right) \le \exp(-\Theta(n))$$

• In Figure 1 (b), we have  $\mathcal{H}_{\leq 0} \cap \operatorname{CH}(\Pi) = \{\pi_2\} \neq \emptyset$ ,  $\operatorname{CH}(\Pi) \not\subseteq \mathcal{H}_{\leq 0}$ , and  $\dim(\mathcal{H}_{\leq 0}) = 2$ . Therefore, for any sufficiently large n (for which it is not hard to prove that  $\mathcal{H}_n^{\mathbb{Z}} \neq \emptyset$ ), we have

$$\forall \vec{\pi} \in \Pi^n, \exp(-\Theta(n)) \le \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}\right) \le \Theta(n^{-\frac{2-2}{2}}) = \Theta(1)$$

Notice that both bounds are asymptotically tight. For example, the lower bound can be achieved by  $\vec{\pi} = {\pi_1}^n$  and the upper bound can be achieved by  $\vec{\pi} = {\pi_2}^n$ .

As an example of the inf part of Theorem 1, suppose  $\pi_1$  is replaced by  $\pi'_1 = (\frac{2}{3}, \frac{1}{3})$  in Figure 1 (b). Then,  $CH(\Pi) \subseteq \mathcal{H}_{\leq 0}$ , which means that  $\forall \vec{\pi} \in \Pi^n, \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}\right) \geq \Theta(1)$  and the lower bound is asymptotically tight.

Remarks on the generality and limitations of Theorem 1. We believe that Theorem 1 is quite general, because first, it provides a dichotomy (more precisely, trichotomy) for the PMV-in-Polyhedron problem. Second, the upper and lower bounds are asymptotically tight. And third, the theorem works for arbitrary  $\mathcal{H}$  characterized by an integer matrix  $\mathcal{A}$  and arbitrary  $\vec{b}$ , and any closed and strictly positive II. As a notable special case, when II contains a single distribution  $\pi$ , the sup and inf parts of the theorem coincide, and the theorem characterizes the PMV-in-Polyhedron problem for i.i.d. PMVs.

The main limitations are, first, the constants in the asymptotic bounds depend on q,  $\Pi$ , and  $\mathcal{H}$ , which are assumed to be fixed; and second,  $\Pi$  must be strictly positive. Nevertheless, we believe that the two limitations are mild at least in the social choice context, because as can be seen in the next section as well as in [66], applications of Theorem 1 (or more precisely, its extension to unions of multiple polyhedra in Theorem 2 in Section 3.1) answer open questions in social choice under a more general and realistic model than IC. Moreover, as commented in [66], many classical models, such as Mallows model and random utility models, are strictly positive.

**Remarks on the comparison with [66, Lemma 1].** We first recall an equivalent and simplified version of [66, Lemma 1] as Lemma\* below for easy reference.

**Lemma\* ([66, Lemma 1]).** Let  $\mathcal{H} = \{\vec{x} \in \mathbb{R}^q : \mathbf{E} \cdot (\vec{x})^\top = (\vec{0})^\top \text{ and } \mathbf{S} \cdot (\vec{x})^\top < (\vec{0})^\top \}$ , where  $\mathbf{E}$  and  $\mathbf{S}$  are integer matrices and  $\mathbf{E} \cdot (\vec{1})^\top = (\vec{0})^\top$  and  $\mathbf{S} \cdot (\vec{1})^\top = (\vec{0})^\top$ . Then,  $\sup_{\vec{\pi} \in \Pi^n} \Pr(\vec{X}_{\vec{\pi}} \in \mathcal{H})$  is 0,  $\exp(-\Omega(n))$ , or  $O(n^{-Rank(\mathbf{E})/2})$ , and the poly bound is asymptotically tight for infinitely many  $n \in \mathbb{N}$ .

We believe that our Theorem 1 is a non-trivial and significant improvement of Lemma \* in the following three aspects.

First, Theorem 1 works for any polyhedron  $\mathcal{H} = \left\{ \vec{x} : \mathbf{A} \cdot (\vec{x})^{\top} \leq (\vec{b})^{\top} \right\}$  with arbitrary integer matrix  $\mathbf{A}$ , while Lemma\* requires  $\mathbf{A} \cdot (\vec{1})^{\top} = (\vec{0})^{\top}$  and also essentially requires that elements in  $\vec{b}$  to be either 0 or -1, which correspond to the  $\mathbf{E}$  part and the  $\mathbf{S}$  part in Lemma\*, respectively.

Second, Theorem 1 provides asymptotically tight bounds, while Lemma\* only claims that the bounds are asymptotically tight for infinitely many n's.

Third, Theorem 1 characterizes smoothed likelihood for the min-adversary, while Lemma\* only works for the max-adversary. While the proof of the min-adversary part of Theorem 1 is similar to its max-adversary part, it is due to the improved techniques and lemmas (Lemma 1 and 2 in the appendix). Without them we do not see an easy way to generalize Lemma\* to the min-adversary.

The proof can be found in Appendix A.1, where a proof sketch is presented in Appendix A.1.1 and the full proof is presented in Appendix A.1.2.

### 3.1 An Extension of Theorem 1 to Unions of Polyhedra

In this subsection, we present an extension of Theorem 1 to the union of  $I \in \mathbb{N}$  polyhedra, denoted by  $\mathcal{C} = \bigcup_{i \leq I} \mathcal{H}_i$ , where  $\mathcal{H}_i = \{\vec{x} \in \mathbb{R}^q : \mathbf{A}_i \cdot (\vec{x})^\top \leq (\vec{b}_i)^\top\}$  and  $\mathbf{A}_i$  is an integer matrix of q columns. We define the *PMV-in-C* problem similarly as the PMV-in-Polyhedron problem (Definition 4), except that  $\mathcal{H}$  is replaced by  $\mathcal{C}$ .

**Definition 6** (The PMV-in-C problem). Given  $q, I \in \mathbb{N}$ ,  $C = \bigcup_{i \leq I} \mathcal{H}_i$ , where  $\forall i \leq I$ ,  $\mathcal{H}_i \subseteq \mathbb{R}^q$  is a polyhedron, and a set  $\Pi$  of distributions over [q], we are interested in

the upper bound  $\sup_{\vec{\pi}\in\Pi^n} \Pr(\vec{X}_{\vec{\pi}}\in\mathcal{C})$ , and the lower bound  $\inf_{\vec{\pi}\in\Pi^n} \Pr(\vec{X}_{\vec{\pi}}\in\mathcal{C})$ 

The key observation is the following straightforward inequality for every PMV  $\vec{X}_{\vec{\pi}}$ :

$$\max_{i \leq I} \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i\right) \leq \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{C}\right) \leq \sum_{i \leq I} \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i\right)$$
(6)

See Figure 2 for an illustration of I = 3. Notice that the right hand side of (6) is no more than  $I \cdot \max_{i \leq I} \Pr(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i)$ , which is  $\Theta(\max_{i \leq I} \Pr(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i))$  because I is a constant.

The high-level idea behind the extension is based on a weighted complete bipartite *activation* graph defined as follows, which represents the relationship between  $CH(\Pi)$  and polyhedra in C in light of Theorem 1. Let  $\mathcal{H}_{i,\leq 0}$  denote the characteristic cone of  $\mathcal{H}_i$ .



Figure 2: Illustration of inequality (6), where  $C = H_1 \cup H_2 \cup H_3$ .

**Definition 7** (Activation graph  $\mathcal{G}_{\Pi,\mathcal{C},n}$ ). For any set of distributions  $\Pi$  over [q], any  $\mathcal{C} = \bigcup_{i=1}^{I} \mathcal{H}_i$ , and any  $n \in \mathbb{N}$ ,  $\mathcal{H}_i$  is said to be active (at n) if  $\mathcal{H}_{i,n}^{\mathbb{Z}} \neq \emptyset$ ; otherwise  $\mathcal{H}_i$  is said to be inactive (at n). Moreover, we define the activation graph  $\mathcal{G}_{\Pi,\mathcal{C},n}$  as follows.

• Vertices. The vertices are  $CH(\Pi)$  and  $\{\mathcal{H}_i : 1 \leq i \leq I\}$ .

• Edges and weights. There is an edge between each  $\pi \in CH(\Pi)$  and each  $\mathcal{H}_i$ , whose weight is

$$w_n(\pi, \mathcal{H}_i) \triangleq \begin{cases} -\infty & \text{if } \mathcal{H}_i \text{ is inactive at } n \\ -\frac{n}{\log n} & \text{otherwise, if } \pi \notin \mathcal{H}_{i,\leqslant 0} \\ \dim(\mathcal{H}_{i,\leqslant 0}) & \text{otherwise} \end{cases}$$

For example, in Figure 3,  $\mathcal{H}_1$  is inactive at n and both  $\mathcal{H}_2$  and  $\mathcal{H}_3$  are active,  $\pi \in \mathcal{H}_{1,\leq 0} \cap \mathcal{H}_{2,\leq 0}$ and  $\pi \notin \mathcal{H}_{3,\leq 0}$ . Notice that the weight on  $(\pi, \mathcal{H}_2)$  is dim $(\mathcal{H}_{2,\leq 0})$  instead of dim $(\mathcal{H}_2)$ .



Figure 3: Illustration of an activation graph  $\mathcal{G}_{\Pi,\mathcal{C},n}$ .

Intuitively, the 0, exponential, and polynomial cases of Theorem 1 (applied to  $\Pi = \{\pi\}$ ) corresponds to the  $-\infty$  edge, the  $-\frac{n}{\log n}$  edge, and the dim $(\mathcal{H}_{i,\leqslant 0})$  edge, respectively. That is, for any  $\vec{\pi} \in \Pi^n$  with  $\sum_{j=1}^n \pi_j = n \cdot \pi$ ,  $\Pr(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i)$  is roughly *n* raise to the power of the weight between  $\pi$  and  $\mathcal{H}_i$  in the activation graph, i.e.,  $n^{w_n(\pi,\mathcal{H}_i)}$ . In particular,  $n^{-\infty} = 0$  and  $n^{-\frac{n}{\log n}-q} = \exp(-\Theta(n))$ .

Therefore, according to (6),  $\Pr(\vec{X}_{\vec{\pi}} \in C)$  is primarily determined by the largest  $w_n(\pi, \mathcal{H}_i)$ , i.e., the maximum weight of all edges connected to  $\pi$  in the activation graph. This is formally defined as follows.

**Definition 8 (Active dimension).** Given  $\mathcal{C}$ , n, and  $\pi \in \mathbb{R}^q$ , we define maximum active dimension of  $\mathcal{C}$  at  $\pi$  and n (active dimension at  $\pi$  for short, when  $\mathcal{C}$  and n are clear from the context), denoted by  $\dim_{\mathcal{C},n}^{\max}(\pi)$ , as follows.

$$\dim_{\mathcal{C},n}^{\max}(\pi) \triangleq \max_{i \leq I} w_n(\pi, \mathcal{H}_i)$$

<sup>&</sup>lt;sup>2</sup>This is the reason behind using  $-\frac{n}{\log n}$ . Theorem 2 still holds if  $-\frac{n}{\log n}$  is replace by any finite negative number.

Consequently, a max- (respectively, min-) adversary aims to choose  $\vec{\pi} = (\pi_1, \ldots, \pi_n) \in \Pi^n$  to maximize (respectively, minimize)  $\dim_{\mathcal{C},n}^{\max}(\frac{1}{n}\sum_{j=1}^n \pi_j)$ , which are characterized by  $\alpha_n$  (respectively,  $\beta_n$ ) defined as follows.

$$\alpha_n \triangleq \max_{\pi \in CH(\Pi)} \dim_{\mathcal{C},n}^{\max}(\pi)$$
$$\beta_n \triangleq \min_{\pi \in CH(\Pi)} \dim_{\mathcal{C},n}^{\max}(\pi)$$

We note that  $\alpha_n$  and  $\beta_n$  depend on  $\Pi$  and C, which are often clear from the context. Also, by definition,  $\alpha_n = -\infty$  is equivalent to  $\beta = -\infty$ , which is equivalent to  $C_n^{\mathbb{Z}} = \emptyset$ . We are now ready to use  $\alpha_n$  and  $\beta_n$  to present the extension of Theorem 1 to the PMV-in-C problem.

**Theorem 2** (Smoothed Likelihood of PMV-in-C). Given any  $q, I \in \mathbb{N}$ , any closed and strictly positive  $\Pi$  over [q], and any  $C = \bigcup_{i \in I} \mathcal{H}_i$  characterized by integer matrices, for any  $n \in \mathbb{N}$ ,

$$\sup_{\vec{\pi}\in\Pi^n} \Pr\left(\vec{X}_{\vec{\pi}}\in\mathcal{C}\right) = \begin{cases} 0 & \text{if } \alpha_n = -\infty\\ \exp(-\Theta(n)) & \text{if } -\infty < \alpha_n < 0\\ \Theta\left(n^{\frac{\alpha_n - q}{2}}\right) & \text{otherwise } (i.e. \ \alpha_n > 0) \end{cases}$$
$$\inf_{\vec{\pi}\in\Pi^n} \Pr\left(\vec{X}_{\vec{\pi}}\in\mathcal{C}\right) = \begin{cases} 0 & \text{if } \beta_n = -\infty\\ \exp(-\Theta(n)) & \text{if } -\infty < \beta_n < 0\\ \Theta\left(n^{\frac{\beta_n - q}{2}}\right) & \text{otherwise } (i.e. \ \beta_n > 0) \end{cases}.$$

Roughly speaking, the max- (respectively, min-) smoothed likelihood for an (n, q)-PMV to be in C is approximately  $n^{\frac{\alpha_n-q}{2}}$  (respectively,  $n^{\frac{\beta_n-q}{2}}$ ). The proof is done by combining the applications of Theorem 1 to  $\Pi$  and every  $\mathcal{H}_i$ , and can be found in Appendix A.2.

Remarks on the applications of Theorem 2. We believe that Theorem 2 is a useful and general tool to study the smoothed likelihood of many events and properties in social choice, as shown in [66] as well as in the rest of this paper. Like Theorem 1, the power of Theorem 2 is that it provides a systematic way of reducing probabilistic analysis to worst-case and non-probabilistic analysis, i.e., the characterizations of  $\alpha_n$ , and  $\beta_n$ . Nevertheless, characterizing  $\alpha_n$  and  $\beta_n$  can still be challenging, which is equivalent to characterizing active  $\mathcal{H}_i$ ,  $\mathcal{H}_{i,\leqslant 0}$ , and dim $(\mathcal{H}_{i,\leqslant 0})$ , as we will see in the next section.

# 4 Smoothed Likelihood of Ties

In this section, we apply Theorem 2 to provide dichotomous characterizations of the smoothed likelihood of ties (Definition 2) under some commonly studied voting rules.

#### 4.1 Integer Positional Scoring Rules

We first apply Theorem 2 to polyhedra that are similar to those in Example 1 and obtain the following theorem for integer positional scoring rules.

**Theorem 3** (Smoothed likelihood of ties: integer positional scoring rules). For any fixed  $m \geq 3$ , let  $\mathcal{M} = (\Theta, \mathcal{L}(\mathcal{A}), \Pi)$  be a strictly positive and closed single-agent preference model and let

 $\vec{s}$  be an integer scoring vector. For any  $2 \leq k \leq m$  and any  $n \in \mathbb{N}$ ,

$$\begin{split} \widetilde{Tie}_{\Pi}^{\max}(r_{\vec{s}},k,n) &= \begin{cases} 0 & if \,\forall P \in \mathcal{L}(\mathcal{A})^n, |r_{\vec{s}}(P)| \neq k \\ \exp(-\Theta(n)) & otherwise, \, if \,\forall \pi \in CH(\Pi), |r_{\vec{s}}(\pi)| < k \\ \Theta(n^{-\frac{k-1}{2}}) & otherwise \end{cases} \\ \widetilde{Tie}_{\Pi}^{\min}(r_{\vec{s}},k,n) &= \begin{cases} 0 & if \,\forall P \in \mathcal{L}(\mathcal{A})^n, |r_{\vec{s}}(P)| \neq k \\ \exp(-\Theta(n)) & otherwise, \, if \,\exists \pi \in CH(\Pi) \, s.t. \, |r_{\vec{s}}(\pi)| < k \\ \Theta(n^{-\frac{k-1}{2}}) & otherwise \end{cases}$$

Take the max smoothed likelihood of ties in Theorem 2 for example. Like Theorem 1, the condition for the 0 case is trivial. Assuming that the 0 case does not happen, the exponential case happens if no distribution (viewed as a fraction profile) in the convex hull of  $\Pi$  has at least k winners under  $r_{\vec{s}}$ . Otherwise, the polynomial case happens. That is, there exists an n-profile P with exactly k winners under  $r_{\vec{s}}$ , and there exists  $\pi \in CH(\Pi)$  that has at least k winners. Notice that the existence of such P (which depends on n but not  $\Pi$ ) does not imply the existence of such  $\pi$  (which depends on  $\Pi$  but not n), nor vice versa. All proofs in this section are delegated to Appendix C.

We immediately have the follow corollary of Theorem 3 when the uniform distribution  $\pi_{\text{uni}}$  is in CH(II), because  $r_{\vec{s}}(\pi_{\text{uni}}) = \mathcal{A}$  and  $|\mathcal{A}| = m \geq k$ , which means that the exponential case never happends. Notice that the corollary does not require  $\pi_{\text{uni}} \in \Pi$ .

Corrollary 1 (Max smoothed likelihood of ties: positional scoring rules). For any fixed  $m \geq 3$ , let  $\mathcal{M} = (\Theta, \mathcal{L}(\mathcal{A}), \Pi)$  be a strictly positive and closed single-agent preference model with  $\pi_{uni} \in CH(\Pi)$ . For any integer scoring vector  $\vec{s}$  and any  $k \leq m$ , for any  $n \in \mathbb{N}$ ,

$$\widetilde{Tie}_{\Pi}^{\max}(r_{\vec{s}}, k, n) = \begin{cases} 0 & \text{if } \forall P \in \mathcal{L}(\mathcal{A})^n, |r_{\vec{s}}(P)| \neq k \\ \Theta(n^{-\frac{k-1}{2}}) & \text{otherwise} \end{cases}$$

The 0 case can indeed happen, for example, when r is the plurality rule, k = m, and  $m \nmid n$ . As commented in [66],  $\pi_{\text{uni}} \in \text{CH}(\Pi)$  is a natural assumption that holds for many single-agent preference models. In particular, Corollary 1 works for IC, which corresponds to  $\Pi = \{\pi_{\text{uni}}\}$ .

#### 4.2 Edge-Order-Based Rules

The characterization for edge-order-based rules is more complicated due to the hardness in characterizing active  $\mathcal{H}_i$ ,  $\mathcal{H}_{i,\leq 0}$ , and dim $(\mathcal{H}_{i,\leq 0})$  in the polyhedra representation of k-way ties. We first introduce necessary notation to formally define edge-order-based rules, whose winners only depend on the order over all edges in WMG w.r.t. their weights, called *palindromic orders*.

**Definition 9 (Palindromic orders).** A total preorder O over  $\mathcal{E} = \{(a, b) \in \mathcal{A} \times \mathcal{A} : a \neq b\}$  is palindromic, if for any pair of edges (a, b), (c, d) in  $\mathcal{E}, (a, b) \triangleright_O (c, d)$  if and only if  $(d, c) \triangleright_O (b, a)$ , where  $e_1 \triangleright_O e_2$  means that  $e_1$  is ranked strictly above  $e_2$  in O, and  $e_1 \equiv_O e_2$  means that  $e_1$  and  $e_2$  are tied in O. Let  $\mathcal{O}_{\mathcal{A}}$  denote the set of all palindromic orders over  $\mathcal{E}$ .

For any weighted directed graph G over  $\mathcal{A}$  with weights  $\{w(a,b) : a \neq b\}$  such that w(a,b) = -w(b,a), let  $EO(G) \in \mathcal{O}_{\mathcal{A}}$  denote the palindromic order w.r.t. the decreasing order of weights in G. For any profile P, let EO(P) = EO(WMG(P)).

In this paper we often use the *tier representation* of palindromic orders, which partition edges into equivalent classes (tiers).

**Definition 10** (Tier representation and refinement of palindromic orders). Any palindromic order  $O \in \mathcal{O}_{\mathcal{A}}$  can be partitioned into tiers:

$$O = T_1 \triangleright \cdots \triangleright T_t \triangleright T_0 \triangleright T_{t+1} \triangleright \cdots \triangleright T_{2t},$$

where for each  $1 \leq i \leq t$ , edges in  $T_i$  are tied, edges in  $T_{2t+1-i}$  are tied, and edges in  $T_{2t+1-i}$  are obtained by flipping edges in  $T_i$ .  $T_0$  is called the middle tier, which consists of all edges e with  $e \equiv_O \bar{e}$ , where  $\bar{e}$  represents flipped e. Only  $T_0$  is allowed to be empty. Let  $Ties(O) = \sum_{i=1}^t (|T_i| - 1) + |T_0|/2$ .  $O_1 \in \mathcal{O}_A$  refines  $O_2 \in \mathcal{O}_A$ , if for all pair of elements  $(e_1, e_2)$ ,  $e_1 \triangleright_{O_2} e_2$  implies  $e_1 \triangleright_{O_1} e_2$ .

**Example 3.** Figure 4 illustrates an example of a profile P, its WMG, and its corresponding palindromic order. In Figure 4 (b) only edges with positive weights are shown.

$$\{1 \succ 2 \succ 3, 1 \succ 3 \succ 2\}$$
(a) Profile P.
$$\begin{array}{c} 1 \\ 2 \\ (b) WMG(P). \end{array}$$

$$\begin{array}{c} 1 \\ 2 \\ (b) WMG(P). \end{array}$$

$$\begin{array}{c} 1 \\ (c) EO(P). \end{array}$$

$$\begin{array}{c} 1 \\ (c) EO(P). \end{array}$$

$$\begin{array}{c} 1 \\ (c) EO(P). \end{array}$$

Figure 4: An example of a profile, its WMG, and its palindromic order.

$$Let \ O = \underbrace{\{(1,2)\}}_{T_1} \triangleright \underbrace{\{(1,3)\}}_{T_2} \triangleright \underbrace{\{(2,3), (3,2)\}}_{T_0} \triangleright \underbrace{\{(3,1)\}}_{T_3} \triangleright \underbrace{\{(2,1)\}}_{T_4}. We \ have \ Ties(EO(P)) = 3,$$
  
$$Ties(O) = 1, \ and \ O \ refines \ EO(P).$$

We are now ready to formally define edge-order-based rules using palindromic orders.

**Definition 11 (Edge-order-based rules).** A voting rule r is said to be edge-order-based, if for every pair of profiles  $P_1, P_2$  with  $EO(P_1) = EO(P_2)$ , we have  $r(P_1) = r(P_2)$ .

Many WMG-based rules, such as Copeland, Maximin, Schulze, and ranked pairs, are edgeorder-based. The domain of any edge-order-based rule r can be naturally extended to palindromic orders. When applying Theorem 2 to edge-order-based rules, each polyhedron  $\mathcal{H}_i$  in  $\mathcal{C}$  is indexed by a palindromic order O with k co-winners, such that  $\mathcal{H}_{i,n}^{\mathbb{Z}}$  corresponds to the histograms of n-profiles whose edge orders are O.

We now define and characterize palindromic orders obtained from n-profiles.

**Definition 12.** For any  $n \in \mathbb{N}$ , let  $\mathcal{O}_{\mathcal{A}}^n \triangleq \{EO(P) : P \in \mathcal{L}(\mathcal{A})^n\}$ . Let  $\mathcal{O}_{\mathcal{A}}' \subset \mathcal{O}_{\mathcal{A}}$  denote the set of palindromic orders O whose middle tier is empty.

**Proposition 1.** For any  $\mathcal{A}$  and any  $n \ge m^4$ ,  $\mathcal{O}_{\mathcal{A}}^n = \begin{cases} \mathcal{O}_{\mathcal{A}} & \text{if } 2 \mid n \\ \mathcal{O}_{\mathcal{A}}' & \text{if } 2 \nmid n \end{cases}$ .

The proof of Proposition 1 is delegated to Appendix C.1. Next, we define  $\mathcal{O}_{r,k,n}^{\pi}$  as the set of palindromic orders O that satisfies three conditions: (1)  $O \in \mathcal{O}_{\mathcal{A}}^{n}$ ; (2) there are exactly kwinners in O under r, and (3) O refines EO( $\pi$ ), where  $\pi$  is viewed as a fractional profile. Let  $\mathcal{O}_{r,k,n}^{\Pi} \triangleq \bigcup_{\pi \in CH(\Pi)} \mathcal{O}_{r,k,n}^{\pi}$ . When  $\mathcal{O}_{r,k,n}^{\Pi} \neq \emptyset$ , we let  $\ell_{\min}$  denote the minimum number of ties in palindromic orders in  $\mathcal{O}_{r,k,n}^{\Pi}$ . When  $\mathcal{O}_{r,k,n}^{\Pi} \neq \emptyset$ , we let  $\ell_{\min}$  denote the maximin number of ties, where the maximum is taken for all  $\pi \in CH(\Pi)$ , and for any given  $\pi$ , the minimum is taken for all palindromic orders in  $\mathcal{O}_{r,k,n}^{\pi}$ . We note that  $\ell_{\min}$  and  $\ell_{\min}$  depend on  $\Pi$ , r, k, and n, which are clear from the context. The formal definitions can be found in Appendix C.2. In fact,  $\ell_{\min}$  and  $\ell_{\min}$  correspond to  $m! - \alpha_n$  and  $m! - \beta_n$  in Theorem 2. We are now ready to present the theorem for EO-based rules.

**Theorem 4** (Smoothed likelihood of ties: edge-order-based rules). For any fixed  $m \ge 3$ , let  $\mathcal{M} = (\Theta, \mathcal{L}(\mathcal{A}), \Pi)$  be a strictly positive and closed single-agent preference model and let r be an edge-order-based rule. For any  $2 \le k \le m$  and any  $n \in \mathbb{N}$ ,

$$\begin{split} \widetilde{Tie}_{\Pi}^{\max}(r,k,n) &= \begin{cases} 0 & if \,\forall O \in \mathcal{O}_{\mathcal{A}}^{n}, |r(O)| \neq k \\ \exp(-\Theta(n)) & otherwise \, if \, \mathcal{O}_{r,k,n}^{\Pi} = \emptyset \\ \Theta\left(n^{-\frac{\ell_{\min}}{2}}\right) & otherwise \end{cases} \\ \widetilde{Tie}_{\Pi}^{\min}(r,k,n) &= \begin{cases} 0 & if \,\forall O \in \mathcal{O}_{\mathcal{A}}^{n}, |r(O)| \neq k \\ \exp(-\Theta(n)) & otherwise \, if \,\exists \pi \in CH(\Pi) \, s.t. \, \mathcal{O}_{r,k,n}^{\pi} = \emptyset \\ \Theta\left(n^{-\frac{\ell_{\min}}{2}}\right) & otherwise \end{cases}$$

The proof can be found in Appendix C.3. In the remainder of this section, we apply Theorem 4 to provide dichotomous characterizations of max-smooth likelihood of ties under Copeland<sub> $\alpha$ </sub>, maximin, Schulze, and ranked pairs for the model in Corollary 1, which includes IC as a special case.

**Proposition 2** (Max smoothed likelihood of ties: Copeland<sub> $\alpha$ </sub>). For any fixed  $m \geq 3$ , let  $\mathcal{M} = (\Theta, \mathcal{L}(\mathcal{A}), \Pi)$  be a strictly positive and closed single-agent preference model with  $\pi_{uni} \in CH(\Pi)$ . Let  $l_{\alpha} = \min\{t \in \mathbb{N} : t\alpha \in \mathbb{Z}\}$ . For any  $2 \leq k \leq m$  and any  $n \in \mathbb{N}$ ,

$$\widetilde{Tie}_{\Pi}^{\max}(Cd_{\alpha},k,n) = \begin{cases} 0 & \text{if } 2 \nmid n, 2 \mid k, \text{ and } k \geq m-1 \\ \Theta(n^{-\frac{k}{4}}) & \text{if } 2 \mid n, 2 \mid k, \text{ and } \begin{cases} (1)k = m, \text{ or} \\ (2)k = m-1 \text{ and } \alpha \geq \frac{1}{2}, \text{ or} \\ (3)k = m-1 \text{ and } k \leq l_{\alpha}(l_{\alpha}+1) \end{cases} \\ \Theta\left(n^{-\frac{l_{\alpha}(l_{\alpha}+1)}{4}}\right) & \text{if } 2 \mid n, 2 \mid k, k = m-1, \alpha < \frac{1}{2}, \text{ and } k > l_{\alpha}(l_{\alpha}+1) \\ \Theta(1) & \text{otherwise } (i.e., \text{ if } 2 \nmid k \text{ or } k \leq m-2) \end{cases}$$

The  $\Theta(1)$  case appears most typical, which happens when k is odd or  $k \leq m-2$ . The  $\Theta(n^{-\frac{l_{\alpha}(l_{\alpha}+1)}{4}})$  case appears most interesting, because its degree depends on the smallest natural number  $l_{\alpha}$  such that  $\alpha l_{\alpha}$  is an integer. For example,  $l_0 = 1$ ,  $l_{1/3} = 3$ ,  $l_{2/5} = 5$ , and  $l_{\alpha} = \infty$  for any irrational number  $\alpha$  (which means that the  $\Theta(n^{-\frac{l_{\alpha}(l_{\alpha}+1)}{4}})$  case does not happen because  $k < \infty = l_{\alpha}(l_{\alpha} + 1)$ ). While the  $\Theta(1)$  case can probably be proved by standard central limit theorem and the union bound, we are not aware of a previous work on it. Standard techniques are too coarse for other cases.

**Proposition 3** (Max smoothed likelihood of ties: maximin). For any fixed  $m \geq 3$ , let  $\mathcal{M} = (\Theta, \mathcal{L}(\mathcal{A}), \Pi)$  be a strictly positive and closed single-agent preference model with  $\pi_{uni} \in CH(\Pi)$ . For any  $2 \leq k \leq m$  and any  $n \in \mathbb{N}$ ,  $\widetilde{Tie}_{\Pi}^{\max}(MM, k, n) = \Theta(n^{-\frac{k-1}{2}})$ .

**Proposition 4** (Max smoothed likelihood of ties: Schulze). For any fixed  $m \geq 3$ , let  $\mathcal{M} = (\Theta, \mathcal{L}(\mathcal{A}), \Pi)$  be a strictly positive and closed single-agent preference model with  $\pi_{uni} \in CH(\Pi)$ . For any  $2 \leq k \leq m$  and any  $n \in \mathbb{N}$ ,  $\widetilde{Tie}_{\Pi}^{\max}(Sch, k, n) = \Theta(n^{-\frac{k-1}{2}})$ .

**Proposition 5** (Max smoothed likelihood of ties: ranked pairs). For any fixed  $m \ge 3$ , let  $\mathcal{M} = (\Theta, \mathcal{L}(\mathcal{A}), \Pi)$  be a strictly positive and closed single-agent preference model with  $\pi_{uni} \in CH(\Pi)$ .

For any  $2 \leq k \leq m$  and any  $n \in \mathbb{N}$ ,  $\Omega(n^{-\frac{k-1}{2}}) \leq \widetilde{Tie}_{\Pi}^{\max}(RP, k, n) \leq n^{-\Omega(\frac{\log k}{\log \log k})}$ . Moreover, when  $m \geq k + 5\lceil \log k \rceil$ ,  $\widetilde{Tie}_{\Pi}^{\max}(RP, k, n) = \Omega(n^{-\frac{\lceil \log k \rceil}{2}})$ . When k = 2, we have  $\widetilde{Tie}_{\Pi}^{\max}(RP, 2, n) = \Theta(n^{-0.5})$ .

**Proof sketches of Propositions 2, 3, 4, and 5.** The proofs are done by applying Theorem 4. For any EO-based rules r studied in this paper, the condition for the 0 case can be verified efficiently using Proposition 1 for any sufficiently large n. If the 0 case does not happen, then the exponential case does not happen either, because for any  $O \in \mathcal{O}^n_{\mathcal{A}}$  such that |r(O)| = k, O extends  $EO(\pi_{uni})$ , which is the palindromic order that only has the middle tier  $T_0$ . This also means that for every  $\pi \in CH(\Pi)$ , we have  $\mathcal{O}^{\pi}_{r,k,n} \subseteq \mathcal{O}^{\pi_{uni}}_{r,k,n}$ . Consequently,  $\ell_{\min}$  is achieved at  $\pi_{uni}$ .

The bulk of proof then focuses on characterizing  $O \in \mathcal{O}_{\mathcal{A}}^n$  with the minimum number of ties such that |r(O)| = k. This can be more complicated than it appears, for example for ranked pairs and for Copeland<sub> $\alpha$ </sub> when  $2 \nmid$  and  $\alpha$  is not 0 or 1. In particular, for ranked pairs we were only able to obtain (non-tight) upper and lower bounds. The full proofs of Propositions 2, 3, 4, and 5 can be found in Appendix C.4, C.5, C.6, and C.7, respectively.

#### 4.3 STV and Coombs

**Theorem 5** (Smoothed likelihood of ties: STV and Coombs). For any fixed  $m \geq 3$ , let  $r \in \{STV, Coombs\}$  and let  $\mathcal{M} = (\Theta, \mathcal{L}(\mathcal{A}), \Pi)$  be a strictly positive and closed single-agent preference model. For any  $2 \leq k \leq m$  and  $n \in \mathbb{N}$ ,  $\widetilde{Tie}_{\Pi}^{\max}(r, k, n)$  (respectively,  $\widetilde{Tie}_{\Pi}^{\min}(r_{\vec{s}}, k, n)$ ) is either 0,  $\exp(-\Theta(n))$ , or  $\Theta(poly(n))$ .

The formal statement of the theorem and its proof are delegated to Appendix D. To accurately characterize the degree in the polynomial case, we introduce *PUT structures* (Definition 21) as the counterpart of palindromic orders to define and analyze active polyhedra and the dimensions of their characteristic cones. See Appendix D for its formal definitions and an example. Like in Section 4.2, the theorem can be applied to characterize max smoothed likelihood of ties for STV and Coombs for distributions  $\Pi$  where  $\pi_{uni} \in CH(\Pi)$ , as shown in the following proposition.

**Proposition 6** (Max smoothed likelihood of ties: STV and Coombs). For any fixed  $m \ge 3$ , let  $r \in \{STV, Coombs\}$  and let  $\mathcal{M} = (\Theta, \mathcal{L}(\mathcal{A}), \Pi)$  be a strictly positive and closed single-agent preference model with  $\pi_{uni} \in CH(\Pi)$ . For any  $2 \le k \le m$  and any  $n \in \mathbb{N}$ ,

$$\widetilde{Tie}_{\Pi}^{\max}(r,k,n) = \begin{cases} \Theta(n^{-\frac{k-1}{2}}) & \text{if } \begin{cases} (1)m \ge 4, \text{ or } \\ (2)m = 3 \text{ and } k = 2, \text{ or } \\ (3)m = k = 3 \text{ and } (2 \mid n \text{ or } 3 \mid n) \\ 0 & \text{otherwise } (i.e., m = k = 3, 2 \nmid n, \text{ and } 3 \nmid n) \end{cases}$$

To prove the proposition, we prove a McGarvey [42]-type result for STV and Coombs (Lemma 3) to characterize active polyhedra. The lemma might be of independent interest.

## 5 Experimental Studies

We examine the fraction of profiles with two-or-more-way ties using simulated data and Preflib data [41] under Borda, plurality, veto, maximin, ranked pairs, Schulze, Copeland<sub>0.5</sub>, STV, and Coombs. All experiments were implemented in Python 3 and were conducted on a MacOS laptop

with 3.1 GHz Intel Core i7 CPU and 16 GB memory.

Simulated data. We generate profiles of m = 4 alternatives under IC. *n* ranges from 20 to 200. In each setting we generate 100000 profiles.

The goal of experiments on synthetic data is to provide a sanity check for the theoretical results in this paper. For clarity, we present the results for Borda, STV, maximin, ranked pairs, and Copeland<sub>0.5</sub> in Figure 5. Results for all voting rules described above can be found in Figure 15 in Appendix E. Figure 5 confirms Corollary 1 and Propositions 2, 3, 5, and 6 under IC as discussed in the Introduction: the probability of any-way ties is  $\Theta(1)$ for Copeland<sub>0.5</sub> and is  $\Theta(\frac{1}{\sqrt{n}})$  for other rules.



Figure 5: Fraction of tied profiles under IC.

**Preflib data.** Because the PUT versions of STV and ranked pairs are NP-hard to compute [13, 7], we used AI-search-based implementations of STV and ranked pairs [62] and computed the fraction of tied profiles among the 307 profiles from Strict Order-Complete Lists (SOC) under election data category at Preflib [41].<sup>3</sup> The results are summarized in Table 2 below. We emphasize that the observations are drawn only from Preflib data and should not be interpreted as general conclusions on the likelihood of ties in presidential elections.

Table 2: Percentage of tied profiles in Preflib data in weakly increasing order.

	Borda	$Copeland_{0.5}$	Plurality	Maximin	Schulze	Ranked pairs	STV	Coombs	Veto
Ties	1.6%	2.6%	4.6%	6.8%	6.8%	6.8%	7.5%	10.4%	31.3%

Table 2 shows that ties occur least frequently under Borda (1.6% of the profiles), which is consistent with the experiments on synthetic data in Figure 5. Two interesting observations are: first, ties are rare under Copeland<sub>0.5</sub> (2.6%); and second, ties occur frequently under veto (31.3%), which mostly happen when the number of alternatives is larger than the number of voters—in such cases the election is guaranteed to be tied under veto. The two observations are quite different from Figure 5, which is probably because real-life preference data can be significantly different from IC, as widely acknowledged in the literature [34].

## 6 Future work

We see three immediate directions for future work. First, technically, how can we improve the results for more general models, especially by dropping the strictness assumption on  $\Pi$ ? Second, how can we extend the study to other events of interest in voting, for example, stability and margin of victory of voting rules, and more generally other topics such as multi-winner elections, judgement aggregation, matching, and resource allocation? Third, what are the smoothed complexity

<sup>&</sup>lt;sup>3</sup>Preflib mentioned that this category can be interpreted as election data, though not all of them come from real-life elections. See some statistics in Appendix E. We used the same dataset in [62], where PUT ranked pairs can finish in one hour.

in various computational aspects of voting [4], such as winner determination [68], manipulation, bribery and control?

# 7 Acknowledgements

We thank Rupert Freeman, Qishen Han, Ao Liu, Marcus Pivato, Sikai Ruan, Rohit Vaish, Weiqiang Zheng, Bill Zwicker, participants of the COMSOC video seminar, and anonymous reviewers for helpful comments. This work is supported by NSF #1453542, ONR #N00014-17-1-2621, and a gift fund from Google.

# References

- David Austen-Smith and Jeffrey S. Banks. 1996. Information Aggregation, Rationality, and the Condorcet Jury Theorem. *The American Political Science Review* 90, 1 (1996), 34–45.
- [2] Haris Aziz, Serge Gaspers, Nicholas Mattei, Nina Narodytska, and Toby Walsh. 2013. Ties Matter: Complexity of Manipulation when Tie-Breaking with a Random Vote. In *Proceedings* of IJCAI.
- [3] John F. Banzhaf III. 1968. One Man, 3.312 Votes: A Mathematical Analysis of the Electoral College. Villanova Law Review 13, 2 (1968), Article 3.
- [4] Dorothea Baumeister, Tobias Hogrebe, and Jörg Rothe. 2020. Towards Reality: Smoothed Analysis in Computational Social Choice. In *Proceedings of AAMAS*. 1691–1695.
- [5] Nathaniel Beck. 1975. A note on the probability of a tied election. Public Choice 23, 1 (1975), 75–79.
- [6] Vidmantas Bentkus. 2005. A Lyapunov-type bound in R<sup>d</sup>. Theory of Probability & Its Applications 49, 2 (2005), 311-323.
- [7] Markus Brill and Felix Fischer. 2012. The Price of Neutrality for the Ranked Pairs Method. In Proceedings of the National Conference on Artificial Intelligence (AAAI). Toronto, Canada, 1299–1305.
- [8] James M. Buchanan. 1974. Hegel on the Calculus of Voting. Public Choice 11 (1974), 99–101.
- [9] Noel Campbell and Marcus Witcher. 2015. Political entrepreneurship: Jefferson, Bayard, and the election of 1800. Journal of Entrepreneurship and Public Policy 4, 3 (2015), 298–312.
- [10] Gary Chamberlain and Michael Rothschild. 1981. A note on the probability of casting a decisive vote. Journal of Economic Theory 25, 1 (1981), 152–162.
- [11] Christine Chung, Katrina Ligett, Kirk Pruhs, and Aaron Roth. 2008. The Price of Stochastic Anarchy. In *International Symposium on Algorithmic Game Theory*. 303–314.
- [12] Marquis de Condorcet. 1785. Essai sur l'application de l'analyse à la probabilité des décisions rendues à la pluralité des voix. Paris: L'Imprimerie Royale.
- [13] Vincent Conitzer, Matthew Rognlie, and Lirong Xia. 2009. Preference Functions That Score Rankings and Maximum Likelihood Estimation. In Proceedings of the Twenty-First International Joint Conference on Artificial Intelligence (IJCAI). Pasadena, CA, USA, 109–115.

- [14] Vincent Conitzer and Toby Walsh. 2016. Barriers to Manipulation in Voting. In Handbook of Computational Social Choice, Felix Brandt, Vincent Conitzer, Ulle Endriss, Jérôme Lang, and Ariel Procaccia (Eds.). Cambridge University Press, Chapter 6.
- [15] William J. Cook, Albertus M. H. Gerards, Alexander Schrijver, and Eva Tardos. 1986. Sensitivity theorems in integer linear programming. *Mathematical Programming* 34, 3 (1986), 251–264.
- [16] Constantinos Daskalakis, Anindya De, Gautam Kamat, and Christos Tzamos. 2016. A Size-Free CLT for Poisson Multinomials and its Applications. In *Proceedings of STOC*. 1074–1086.
- [17] Ilias Diakonikolas, Daniel Mertz Kane, and Alistair Stewart. 2016. The fourier transform of poisson multinomial distributions and its algorithmic applications. In *Proceedings of STOC*. 1060–1073.
- [18] Cyril Domb. 1960. On the theory of cooperative phenomena in crystals. Advances in Physics 9, 34 (1960), 149–244.
- [19] Anthony Downs. 1957. An Economic Theory of Democracy. New York: Harper & Row.
- [20] Piotr Faliszewski and Jörg Rothe. 2016. Control and bribery in voting. In *Handbook of Computational Social Choice*. Cambridge University Press, Chapter 7.
- [21] Timothy Feddersen and Wolfang Pesendorfer. 1996. The swing voter's curse. American Economic Review 86 (1996), 408–424.
- [22] Rupert Freeman, Markus Brill, and Vincent Conitzer. 2015. General Tiebreaking Schemes for Computational Social Choice. In Proceedings of the 2015 International Conference on Autonomous Agents and Multiagent Systems. 1401–1409.
- [23] Ehud Friedgut, Gil Kalai, Nathan Keller, and Noam Nisan. 2011. A Quantitative Version of the Gibbard-Satterthwaite theorem for Three Alternatives. SIAM J. Comput. 40, 3 (2011), 934–952.
- [24] Andrew Gelman, Gary King, and John Boscardin. 1998. Estimating the Probability of Events that Have Never Occurred: When Is Your Vote Decisive? J. Amer. Statist. Assoc. 93 (1998), 1–9.
- [25] Andrew Gelman, Nate Silver, and Aaron Edlin. 2012. What is the probability your vote will make a difference? *Economic Inquiry* 50, 2 (2012), 321–326.
- [26] Allan Gibbard. 1973. Manipulation of voting schemes: A general result. Econometrica 41 (1973), 587–601.
- [27] Raphael Gillett. 1977. Collective Indecision. Behavioral Science 22, 6 (1977), 383–390.
- [28] Raphael Gillett. 1980. The Comparative Likelihood of an Equivocal Outcome under the Plurality, Condorcet, and Borda Voting Procedures. *Public Choice* 35, 4 (1980), 483–491.
- [29] I. J. Good and Lawrence S. Mayer. 1975. Estimating the efficacy of a vote. Behavioral Science 20, 1 (1975), 25–33.
- [30] Wassily Hoeffding. 1956. On the Distribution of the Number of Successes in Independent Trials. The Annals of Mathematical Statistics 27, 3 (1956), 713–721.

- [31] Wassily Hoeffding. 1963. Probability inequalities for sums of bounded random variables. J. Amer. Statist. Assoc. 58, 301 (1963), 13-30.
- [32] Li-Sha Huang and Shang-Hua Teng. 2007. On the Approximation and Smoothed Complexity of Leontief Market Equilibria. In *Proceedings of FAW*. 96–107.
- [33] Michel Le Breton, Dominique Lepelley, and Hatem Smaoui. 2016. Correlation, partitioning and the probability of casting a decisive vote under the majority rule. *Journal of Mathematical Economics* 64 (2016), 11–22.
- [34] Aki Lehtinen and Jaakko Kuorikoski. 2007. Unrealistic Assumptions in Rational Choice Theory. Philosophy of the Social Sciences 37, 2 (2007), 115–138.
- [35] Ao Liu, Yun Lu, Lirong Xia, and Vassilis Zikas. 2020. How Private Is Your Voting?. Presented at WADE-18 workshop. In *Proceedings of UAI*.
- [36] L. Lovász and M.D. Plummer. 2009. Matching Theory. North-Holland; Elsevier Science Publishers B.V.; Sole distributors for the U.S.A. and Canada, Elsevier Science Publishing Company.
- [37] Thomas R. Magrino, Ronald L. Rivest, Emily Shen, and David Wagner. 2011. Computing the Margin of Victory in IRV Elections. In *Proceedings of 2011 EVT/WOTE Conference*.
- [38] Thierry Marchant. 2001. The probability of ties with scoring methods: Some results. Social Choice and Welfare 18, 4 (2001), 709–735.
- [39] Howard Margolis. 1977. Probability of a Tie Election. Public Choice 31 (1977), 135–138.
- [40] Nicholas Mattei, Nina Narodytska, and Toby Walsh. 2014. How hard is it to control an election by breaking ties?. In Proceedings of the Twenty-first European Conference on Artificial Intelligence. 1067–1068.
- [41] Nicholas Mattei and Toby Walsh. 2013. PrefLib: A Library of Preference Data. In Proceedings of Third International Conference on Algorithmic Decision Theory (Lecture Notes in Artificial Intelligence).
- [42] David C. McGarvey. 1953. A Theorem on the Construction of Voting Paradoxes. *Econometrica* 21, 4 (1953), 608–610.
- [43] Iain McLean and Fiona Hewitt (Eds.). 1994. Condorcet: Foundations of Social Choice and Political Theory. Edward Elgar Publishing.
- [44] Carl D. Meyer. 2000. Matrix analysis and applied linear algebra. SIAM.
- [45] Elchanan Mossel, Ariel D. Procaccia, and Miklos Z. Racz. 2013. A Smooth Transition From Powerlessness to Absolute Power. *Journal of Artificial Intelligence Research* 48, 1 (2013), 923–951.
- [46] Elchanan Mossel and Miklos Z. Racz. 2015. A quantitative Gibbard-Satterthwaite theorem without neutrality. *Combinatorica* 35, 3 (2015), 317–387.
- [47] Casey B. Mulligan and Charles G. Hunter. 2003. The Empirical Frequency of a Pivotal Vote. Public Choice 116, 1/2 (2003), 31–54.

- [48] Roger B. Myerson. 2000. Large Poisson Games. Journal of Economic Theory 84 (2000), 7–45.
- [49] Matías Núñez and Marcus Pivato. 2019. Truth-revealing voting rules for large populations. Games and Economic Behavior 113 (2019), 285–305.
- [50] Svetlana Obraztsova and Edith Elkind. 2011. On the Complexity of Voting Manipulation under Randomized Tie-Breaking. In Proceedings of the Twenty-Second International Joint Conference on Artificial Intelligence (IJCAI). Barcelona, Catalonia, Spain, 319–324.
- [51] Svetlana Obraztsova, Edith Elkind, and Noam Hazon. 2011. Ties Matter: Complexity of Voting Manipulation Revisited. In Proceedings of the Tenth International Joint Conference on Autonomous Agents and Multi-Agent Systems (AAMAS). Taipei, Taiwan, 71–78.
- [52] Svetlana Obraztsova, Yair Zick, and Edith Elkind. 2013. On manipulation in multi-winner elections based on scoring rules. In *Proceedings of AAMAS*. 359—366.
- [53] Ariel D. Procaccia and Jeffrey S. Rosenschein. 2007. Junta Distributions and the Average-Case Complexity of Manipulating Elections. Journal of Artificial Intelligence Research (JAIR) 28 (2007), 157–181.
- [54] Martin Raič. 2019. A multivariate Berry-Esseen theorem with explicit constants. Bernoulli 25, 4A (2019), 2824–2853.
- [55] William H. Riker and Peter C. Ordeshook. 1968. A Theory of the Calculus of Voting. The American Political Science Review 62, 1 (1968), 25–42.
- [56] Mark Satterthwaite. 1975. Strategy-proofness and Arrow's conditions: Existence and correspondence theorems for voting procedures and social welfare functions. *Journal of Economic Theory* 10 (1975), 187–217.
- [57] Alexander Schrijver. 1998. Theory of Linear and Integer Programming. Wiley.
- [58] Markus Schulze. 2011. A new monotonic, clone-independent, reversal symmetric, and condorcet-consistent single-winner election method. *Social Choice and Welfare* 36, 2 (2011), 267–303.
- [59] Daniel A. Spielman and Shang-Hua Teng. 2009. Smoothed Analysis: An Attempt to Explain the Behavior of Algorithms in Practice. *Commun. ACM* 52, 10 (2009), 76–84.
- [60] Thorwald Nicolaus Tideman. 1987. Independence of clones as a criterion for voting rules. Social Choice and Welfare 4, 3 (1987), 185–206.
- [61] Gregory Valiant and Paul Valiant. 2011. Estimating the unseen: an n/log(n)-sample estimator for entropy and support size, shown optimal via new CLTs. In *Proceedings of STOC*. 685–694.
- [62] Jun Wang, Sujoy Sikdar, Tyler Shepherd, Zhibing Zhao, Chunheng Jiang, and Lirong Xia. 2019. Practical Algorithms for STV and Ranked Pairs with Parallel Universes Tiebreaking. In Proceedings of AAAI.
- [63] Wikipedia. [n. d.]. https://en.wikipedia.org/wiki/Resolvability\_criterion.
- [64] Lirong Xia. 2012. Computing The Margin of Victory for Various Voting Rules. In Proceedings of the ACM Conference on Electronic Commerce (EC). Valencia, Spain, 982–999.

- [65] Lirong Xia. 2015. Generalized Decision Scoring Rules: Statistical, Computational, and Axiomatic Properties. In Proceedings of the Sixteenth ACM Conference on Economics and Computation. Portland, Oregon, USA, 661–678.
- [66] Lirong Xia. 2020. The Smoothed Possibility of Social Choice. In Proceedings of NeurIPS.
- [67] Lirong Xia and Vincent Conitzer. 2008. Generalized Scoring Rules and the Frequency of Coalitional Manipulability. In Proceedings of the ACM Conference on Electronic Commerce. 109–118.
- [68] Lirong Xia and Weiqiang Zheng. 2021. The Smoothed Complexity of Computing Kemeny and Slater Rankings. In *Proceedings of AAAI*.

# Contents

1	Introduction         1.1       Related Work and Discussions	<b>1</b> 4				
<b>2</b>	Preliminaries					
3	PMV-in-Polyhedron Problem and Main Technical Theorems         3.1       An Extension of Theorem 1 to Unions of Polyhedra	<b>8</b> 11				
4	Smoothed Likelihood of Ties         4.1 Integer Positional Scoring Rules         4.2 Edge-Order-Based Rules         4.3 STV and Coombs	<b>13</b> 13 14 17				
<b>5</b>	Experimental Studies	17				
6	3 Future work					
7	' Acknowledgements					
A	Materials for Section 3         A.1 Proof of Theorem 1         A.1.1 Proof Sketch of Theorem 1         A.1.2 Full Proof of Theorem 1         A.2 Proof of Theorem 2	<ul> <li>25</li> <li>25</li> <li>27</li> <li>44</li> </ul>				
в	Appendix for Section 4.1: Integer Positional Scoring Rules         B.1 Proof of Theorem 3	<b>48</b> 48				
С	Appendix for Section 4.2: EO-Based RulesC.1 Proof of Proposition 1C.2 Formal Definition of Edge-Order-Based Rules (Section 4.2)C.3 Proof of Theorem 4C.4 Proof of Proposition 2C.5 Proof of Proposition 3C.6 Proof of Proposition 4C.7 Proof of Proposition 5	<b>50</b> 51 51 53 58 59 61				
D	Appendix for Section 4.3: STV and Coombs         D.1 Formal Definitions and Statements of Results         D.2 Proof of Lemma 3         D.3 Proof of Theorem 5         D.4 Proof of Proposition 6         Experimental Besults for All Bules	<ul> <li>62</li> <li>62</li> <li>65</li> <li>68</li> <li>70</li> <li>72</li> </ul>				
Ľ	Experimental Results for All Rules	( 4				

# A Materials for Section 3

#### A.1 Proof of Theorem 1

**Theorem 1.** (The main technical theorem). Given any  $q \in \mathbb{N}$ , any closed and strictly positive  $\Pi$  over [q], and any polyhedron  $\mathcal{H}$  with integer matrix  $\mathbf{A}$ , for any  $n \in \mathbb{N}$ ,

$$\sup_{\vec{\pi}\in\Pi^{n}}\Pr\left(\vec{X}_{\vec{\pi}}\in\mathcal{H}\right) = \begin{cases} 0 & \text{if } \mathcal{H}_{n}^{\mathbb{Z}} = \emptyset \\ \exp(-\Theta(n)) & \text{if } \mathcal{H}_{n}^{\mathbb{Z}} \neq \emptyset \text{ and } \mathcal{H}_{\leq 0} \cap CH(\Pi) = \emptyset \\ \Theta\left(n^{\frac{\dim(\mathcal{H}_{\leq 0})-q}{2}}\right) & \text{otherwise } (i.e. \ \mathcal{H}_{n}^{\mathbb{Z}} \neq \emptyset \text{ and } \mathcal{H}_{\leq 0} \cap CH(\Pi) \neq \emptyset) \end{cases}$$
$$\inf_{\vec{\pi}\in\Pi^{n}}\Pr\left(\vec{X}_{\vec{\pi}}\in\mathcal{H}\right) = \begin{cases} 0 & \text{if } \mathcal{H}_{n}^{\mathbb{Z}} = \emptyset \\ \exp(-\Theta(n)) & \text{if } \mathcal{H}_{n}^{\mathbb{Z}} \neq \emptyset \text{ and } CH(\Pi) \not\subseteq \mathcal{H}_{\leq 0} \\ \Theta\left(n^{\frac{\dim(\mathcal{H}_{\leq 0})-q}{2}}\right) & \text{otherwise } (i.e. \ \mathcal{H}_{n}^{\mathbb{Z}} \neq \emptyset \text{ and } CH(\Pi) \subseteq \mathcal{H}_{\leq 0}) \end{cases}.$$

#### A.1.1 Proof Sketch of Theorem 1

In this subsection we present a proof sketch for the exponential and polynomial cases of the sup part, because the 0 case is trivial.

Intuition and proof sketch for the exponential bounds on Sup. We first note that  $\vec{X}_{\vec{\pi}}$  is an integer-vector-valued random variable. Therefore,  $\Pr(\vec{X}_{\vec{\pi}} \in \mathcal{H}) = \Pr(\vec{X}_{\vec{\pi}} \in \mathcal{H}_n^{\mathbb{Z}})$  and is mainly determined by two factors: (1) the distance between  $\mathbb{E}(\vec{X}_{\vec{\pi}})$  and  $\mathcal{H}_n^{\mathbb{Z}}$ , and (2) the density of integer vectors in  $\mathcal{H}_n^{\mathbb{Z}}$ . Standard concentration bounds, e.g., Hoeffding's inequality, tell us that when  $\mathbb{E}(\vec{X}_{\vec{\pi}})$  and  $\mathcal{H}_n^{\mathbb{Z}}$  are  $\Theta(n)$  away, the probability for  $\vec{X}_{\vec{\pi}}$  to be in  $\mathcal{H}_n^{\mathbb{Z}}$  is exponentially small. This is the intuition behind the exponential case, as illustrated in Figure 1 (a).

Intuition behind the polynomial bounds on Sup. As illustrated in Figure 1 (b), in the polynomial case, Cone(II) is O(1) away from  $\mathcal{H}$ , and one may expect that  $\sup_{\vec{\pi}\in\Pi^n} \Pr(\vec{X}_{\vec{\pi}} \in \mathcal{H})$  is achieved when  $\mathbb{E}(\vec{X}_{\vec{\pi}})$  is close to  $\mathcal{H}$ . Then,  $\Pr(\vec{X}_{\vec{\pi}} \in \mathcal{H})$  is mostly determined by the density of integer vectors in  $\mathcal{H}_n^{\mathbb{Z}}$ . A natural conjecture is that the density can be measured by the dimension of  $\mathcal{H}$ , but this is not true as illustrated in Figure 6 (a) below, where dim $(\mathcal{H}) = 2$ , which is the same as dim $(\mathcal{H})$  in Figure 1 (b). However,  $\Pr(\vec{X}_{\vec{\pi}} \in \mathcal{H})$  in Figure 6 (a) is smaller than that in Figure 1 (b) as  $n \to \infty$ , because the "volume" of  $\mathcal{H}_n^{\mathbb{Z}}$  in Figure 6 (a) does not increase as n increases.

It turns out that the dimension of  $\mathcal{H}_{\leq 0}$  is the right measure. For example,  $\dim(\mathcal{H}_{\leq 0}) = 1$  in Figure 6 (a) and  $\dim(\mathcal{H}_{\leq 0}) = 2$  in Figure 1 (b). The rest of the proof leverages the interplay between the matrix representation and the *V*-representation of  $\mathcal{H}$ , which is defined by the Minkowski-Weyl theorem (see, e.g., [57, p. 100]). More precisely, the V-representation of  $\mathcal{H}$  is  $\mathcal{H} = \mathcal{V} + \mathcal{H}_{\leq 0}$ , where  $\mathcal{V}$  is a finitely generated polyhedron and  $\mathcal{H}_{\leq 0}$  is the characterization cone of  $\mathcal{H}$ . See Figure 6 (b) and (c) for the V-representations of  $\mathcal{H}$  in Figure 6 (a) and in Figure 1 (b), respectively.

**Polynomial upper bound on Sup.** To accurately upper-bound  $Pr(\vec{X}_{\vec{\pi}} \in \mathcal{H})$ , we partition the q dimensions of  $\vec{X}_{\vec{\pi}}$  into two sets:  $I_0$  and  $I_1$ , such that vectors in  $\mathcal{H}_n^{\mathbb{Z}}$  can be enumerated by first enumerating their  $I_1$  components with high flexibility, conditioned on which the  $I_0$  components are more or less determined. More precisely, the following two conditions are satisfied.



Figure 6: An example of  $\dim(\mathcal{H}) > \dim \mathcal{H}_{\leq 0}$  and V-representations of  $\mathcal{H}$  in (a) and  $\mathcal{H}$  in Figure 1 (b).

**Condition (1).** For any  $\vec{h}_1 \in \mathbb{Z}_{\geq 0}^{I_1}$ , the restriction of  $\mathcal{H}_n^{\mathbb{Z}}$  on  $\vec{h}_1$ , denoted by  $\mathcal{H}_n^{\mathbb{Z}}|_{\vec{h}_1} = \{\vec{h}_0 \in \mathbb{Z}_{\geq 0}^{I_0} : (\vec{h}_0, \vec{h}_1) \in \mathcal{H}_n^{\mathbb{Z}}\}$ , contains a constant number (in n) of integer vectors.

**Condition (2).** With high (marginal) probability on the  $I_1$  components of  $\vec{X}_{\vec{\pi}}$ , the conditional probability for the  $I_0$  components of  $\vec{X}_{\vec{\pi}}$  to be in  $\mathcal{H}_n^{\mathbb{Z}}|_{\vec{h}_1}$  is  $O\left(n^{\frac{\dim(\mathcal{H}_{\leq 0})-q}{2}}\right)$ .

Once such  $I_0$  and  $I_1$  are defined, the upper bound follows after applying the law of total probability.

We use the matrix representation of  $\mathcal{H}$  to define  $I_0$  and  $I_1$  as follows, which is similar to the definitions in the proof of [66, Lemma 1] except that our definition works for general **A**. Let  $\mathbf{A}^=$  denote the *implicit equalities* of **A**, which is the maximum set of rows of **A** such that for all  $\vec{x} \in \mathcal{H}_{\leq 0}$ , we have  $\mathbf{A}^= \cdot (\vec{x})^\top = (\vec{0})^\top$ . We note that  $\mathbf{A}^=$  does not depend on  $\vec{b}$ , and  $\operatorname{Rank}(\mathbf{A}^=) = q - \dim(\mathcal{H}_{\leq 0})$  [57, Equation (9), p. 100]. For example, in Figure 1 (b),  $\mathbf{A}^= = \emptyset$  and  $\dim(\mathcal{H}_{\leq 0}) = 2$ .

We then use the reduced row echelon form (a.k.a. row canonical form) [44] of  $\begin{bmatrix} \mathbf{A}^{=} \\ \vec{1} \end{bmatrix}$  to define  $I_0$  and  $I_1$ . More precisely, we apply the Gauss-Jordan elimination method to convert the system of linear equations  $\begin{bmatrix} \mathbf{A}^{=} \\ \vec{1} \end{bmatrix} \cdot (\vec{x})^{\top} = (\vec{0}, n)^{\top}$  to another system of linear equations  $(\vec{x}_{I_0})^{\top} = \mathbf{D} \cdot (\vec{x}_{I_1}, n)^{\top}$ , where  $\vec{x}_{I_0}$  are the  $I_0$  components of  $\vec{x}, I_0 \cup I_1 = [q], |I_0| = \operatorname{Rank}(\mathbf{A}^{=}) + 1 = q - \dim(\mathcal{H}_{\leq 0}) + 1$ , and  $\mathbf{D}$  is an  $(q - \dim(\mathcal{H}_{\leq 0}) + 1) \times \dim(\mathcal{H}_{\leq 0})$  rational matrix that does not depend on n. For example, in Figure 1 (b),  $I_0 = \{1\}, I_1 = \{2\}$ , and  $\mathbf{D} = \begin{bmatrix} -1 & 1 \end{bmatrix}$ . See [66, Example 4 in the Appendix] for a more informative example.

Then, we prove in Claim 1 in Appendix A.1.2 that Condition (1) above holds. Condition (2) is proved by applying the point-wise anti-concentration bound [66, Lemma 3 in the Appendix] and an alternative representation of the PMV as a simple Bayesian network as done in [66].

**Polynomial lower bound on Sup.** The proof of this part is the hardest and drastically different from the proofs in [66]. We will specify  $\vec{\pi}$  and a  $(\dim(\mathcal{H}_{\leq 0}) - 1)$ -dimensional region  $B^n$  in  $\mathcal{H}_n$  that is  $O(\sqrt{n})$  away from  $\mathbb{E}(\vec{X}_{\vec{\pi}})$  based on the V-representation of  $\mathcal{H}$ , as illustrated in Figure 7, which

continues the setting of Figure 1 (b). More precisely, we first choose the following three vectors arbitrarily and then fix them throughout the proof: let  $\vec{x}^* \in \mathcal{H}_{\leq 0} \cap \operatorname{CH}(\Pi), \, \vec{x}^{\#} \in \mathcal{H} \cap \mathbb{Z}_{\geq 0}^q$ , and let  $\vec{x}^{@}$  be an inner point of  $\mathcal{H}_{\leq 0}$ . For example, in Figure 7 we let  $\vec{x}^{*} = \pi_{2}$  because  $\pi_{2}$  is the only vector in  $\mathcal{H}_{\leq 0} \cap \mathrm{CH}(\Pi)$ .

Given any  $n \in \mathbb{N}$ , we define  $\vec{x}^{\ell} \triangleq \ell \vec{x}^* +$  $\vec{x}^{\#} + \sqrt{\ell} \vec{x}^{@}$ , where  $\ell \in \mathbb{R}_{\geq 0}$  is chosen to guarantee that  $\vec{x}^{\ell} \cdot \vec{1} = n$ . Then, we define the following vectors.

• An integer vector  $\vec{y}^n \in \mathcal{H}_n^{\mathbb{Z}}$  that is O(1) away from  $\vec{x}^{\ell}$  in  $L_{\infty}$ . The existence of such  $\vec{y}^n$  is guaranteed by the sensitivity analysis of integer programming ([15, Theorem 1(i)] with w = 0).

• A vector  $\vec{\pi} = (\pi_1, \dots, \pi_n) \in \Pi^n$  such that  $\mathbb{E}(\vec{X}_{\vec{\pi}}) = \sum_{j=1}^{n} \pi_j$  is  $O(\sqrt{n})$  away from  $\vec{x}^{\ell}$ . This is done by rounding *n* multiplied by the representation of  $\vec{x}^*$  as the convex combination of no more than q distribu-



Figure 7: Proof of the poly lower bound in Theorem 1.

tions in  $\Pi$ , which is guaranteed by the Carathéodory's theorem for convex hulls.

• An  $\Omega(\sqrt{n})$  neighborhood of  $\vec{y}^n$  in  $\mathcal{H}_n$ , denoted by  $B^n$ . This is done by first defining an  $\gamma$ neighborhood of  $\vec{x}^{@}$  in  $\mathcal{H}_{\leq 0}$ , denoted by  $B_{\gamma}$ , and then letting  $B^n \triangleq \vec{y}^n + \sqrt{\ell}(B_{\gamma} - \vec{x}^{@})$ .

The construction guarantees that  $B^n$  contains  $\Omega\left(n^{\frac{\dim(\mathcal{H}_{\leq 0})-1}{2}}\right)$  many integer vectors, each of

which is  $O(\sqrt{n})$  away from  $\mathbb{E}(\vec{X}_{\vec{\pi}})$ . Then, we prove a point-wise concentration bound in Lemma 1 (which works for arbitrary strictly positive PMVs and is thus stronger than [66, Lemma 4 in the Appendix], which only holds for i.i.d. PMVs) to show that for each integer vector  $\vec{x} \in B^n$ , the probability for  $\vec{X}_{\vec{\pi}}$  to take  $\vec{x}$  is  $\Omega(n^{\frac{1-q}{2}})$ . The lower bound then becomes

$$\Omega\left(n^{\frac{\dim(\mathcal{H}_{\leq 0})-1}{2}}\right) \times \Omega\left(n^{\frac{1-q}{2}}\right) = \Omega\left(n^{\frac{\dim(\mathcal{H}_{\leq 0})-q}{2}}\right),$$

which matches the upper bound asymptotically.

Lemma 1. (Point-wise concentration bound for PMVs). For any  $q \in \mathbb{N}$ , any  $\epsilon > 0$ , and any  $\alpha > 0$ , there exists  $C_{q,\epsilon,\alpha} > 0$  such that for any  $n \in \mathbb{N}$ , any (n,q)-PMV  $\vec{X}_{\vec{\pi}}$  where  $\vec{\pi}$  is above  $\epsilon$ , 

$$\Pr(\vec{X}_{\vec{\pi}} = \vec{x}) > C_{q,\epsilon,\alpha} \cdot n^{\frac{1}{2}}$$

Lemma 1 is proved by extending the idea of solving a constrained optimization problem for Poisson binomial variables [30] to PMVs. Lemma 1 might be of independent interest.

#### Full Proof of Theorem 1 A.1.2

*Proof.* It suffices to prove the theorem holds for all sufficiently large n. In other words, we will prove that given  $\mathcal{H}$  and  $\Pi$ , there exists  $N \in \mathbb{N}$  such that the theorem holds for all n > N. This is because given any constant  $N \in \mathbb{N}$ , the theorem trivially holds for and any  $n \leq N$ —notice that any  $\sup_{\vec{\pi}\in\Pi^n} \Pr\left(\vec{X}_{\vec{\pi}}\in\mathcal{H}\right) \neq 0$  (respectively,  $\inf_{\vec{\pi}\in\Pi^n} \Pr\left(\vec{X}_{\vec{\pi}}\in\mathcal{H}\right) \neq 0$ ) can be viewed as  $\exp(-\Theta(n))$ or  $\Theta\left(n^{\frac{\dim(\mathcal{H}_{\leq 0})-q}{2}}\right)$ , and the zero case is true for any n.

**Proof of the exponential upper bound on Sup.** For any  $n \in \mathbb{N}$  and any  $\vec{\pi} \in \Pi^n$ , let  $\vec{\mu}_{\vec{\pi}} = (\mu_{\vec{\pi},1}, \ldots, \mu_{\vec{\pi},q}) = \mathbb{E}(\vec{X}_{\vec{\pi}}/n)$  denote the mean of  $\vec{X}_{\vec{\pi}}/n$  and let  $\vec{\sigma}_{\vec{\pi}} = (\sigma_{\vec{\pi},1}, \ldots, \sigma_{\vec{\pi},q})$  denote the standard deviations of each component in  $\vec{X}_{\vec{\pi}}/n$ . Because  $\Pi$  is strictly positive, there exists  $\epsilon_1 > 0, \epsilon_2 > 0$  such that for all n, all  $\vec{\pi} \in \Pi^n$ , and all  $i \leq q$ , we have  $\epsilon_1 < \mu_{\vec{\pi},i} < \epsilon_2$  and  $\frac{\epsilon_1}{\sqrt{n}} < \sigma_{\vec{\pi},i} < \frac{\epsilon_2}{\sqrt{n}}$ .

We first prove that  $\mathcal{H}_{\leq 0}$  is sufficiently separated from  $\operatorname{CH}(\Pi)$ . Notice that  $\mathcal{H}_{\leq 0}$  is convex and closed by definition. Because  $\Pi$  is closed and bounded,  $\operatorname{CH}(\Pi)$  is convex, closed and compact. Because  $\mathcal{H}_{\leq 0} \cap \operatorname{CH}(\Pi) = \emptyset$ , by the strict hyperplane separation theorem, there exists a hyperplane that strictly separates  $\mathcal{H}_{\leq 0}$  and  $\operatorname{CH}(\Pi)$ . In other words, there exists  $\epsilon' > 0$  such that for any  $\vec{x}_1 \in \mathcal{H}_{\leq 0}$  and any  $\vec{x}_2 \in \operatorname{CH}(\Pi)$ , we have  $|\vec{x}_1 - \vec{x}_2|_{\infty} > \epsilon'$ , where  $|\cdot|_{\infty}$  is the  $L_{\infty}$  norm.

We then prove that any  $\vec{x} \in \mathcal{H}_n^{\mathbb{Z}}$  is  $\Theta(n)$  away from  $\mathbb{E}(\vec{X}_{\vec{\pi}}) = n\vec{\mu}_{\vec{\pi}}$  when  $n = \vec{x} \cdot \vec{1}$  is sufficiently large. By the Minkowski-Weyl theorem, we can write  $\mathcal{H} = \mathcal{V} + \mathcal{H}_{\leq 0} = \{\vec{v} + \vec{h} : \vec{v} \in \mathcal{V}, \vec{h} \in \mathcal{H}_{\leq 0}\}$ . Let  $C_{\max} = \max_{\vec{x}' \in \mathcal{V}} |\vec{x}'|_{\infty}$ . For any  $\vec{x} \in \mathcal{H}$ , let  $\vec{x} = \vec{x}' + \vec{x}_1$  where  $\vec{x}' \in \mathcal{V}$  and  $\vec{x}_1 \in \mathcal{H}_{\leq 0}$ . We have:

$$|\vec{x} - \mathbb{E}(\vec{X}_{\vec{\pi}})|_{\infty} = |\vec{x} - n\vec{\mu}_{\vec{\pi}}|_{\infty} = |\vec{x}' + \vec{x}_1 - n\vec{\mu}_{\vec{\pi}}|_{\infty} \ge n|\frac{\vec{x}_1}{n} - \vec{\mu}_{\vec{\pi}}| - C_{\max} \ge n\epsilon' - C_{\max},$$

Therefore, when  $\vec{X}_{\vec{\pi}} \in \mathcal{H}_n^{\mathbb{Z}}$ ,  $\vec{X}_{\vec{\pi}}$  must be away from  $\mathbb{E}(\vec{X}_{\vec{\pi}})$  by at least  $n\epsilon' - C_{\max}$  in  $\mathcal{L}_{\infty}$ . For any  $n > \frac{2C_{\max}}{\epsilon'}$ , we have  $n\epsilon' - C_{\max} > \frac{\epsilon'}{2}n$ . Therefore,

$$\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_{n}^{\mathbb{Z}}\right) \leq \Pr\left(|\vec{X}_{\vec{\pi}} - \mathbb{E}(\vec{X}_{\vec{\pi}})|_{\infty} > n\epsilon' - C_{\max}\right) \leq \Pr\left(|\vec{X}_{\vec{\pi}} - n\vec{\mu}_{\vec{\pi}}|_{\infty} > \frac{\epsilon'}{2}n\right)$$
$$\leq \sum_{i=1}^{q} \Pr\left(|X_{\vec{\pi},i} - n\mu_{\vec{\pi},i}| > \frac{\epsilon'}{2}n\right) \leq 2q \exp\left(-\frac{(\epsilon')^{2}n}{4(1-2\epsilon)^{2}}\right)$$

1

The last inequality follows after Hoeffding's inequality (Theorem 2 in [31]), where recall that  $\epsilon$  is a constant such that  $\Pi$  is above  $\epsilon$ .

**Proof of the exponential lower bound on Sup.** In fact the lower bound can be achieved by any  $\vec{\pi} \in \Pi^n$ . Let  $\vec{y} \in [q]^n$  denote an arbitrary vector such that  $\text{Hist}(\vec{y}) \in \mathcal{H}_n^{\mathbb{Z}}$ . Because  $\Pi$  is above  $\epsilon$ , we have  $\Pr(\vec{X}_{\vec{\pi}} \in \mathcal{H}) \ge \Pr(\vec{Y} = \vec{y}) \ge \epsilon^n = \exp(n \log \epsilon)$ , which is  $\exp(-O(n))$ .

**Proof of the polynomial upper bound on Sup.** We use the V-representation of  $\mathcal{H} = \mathcal{V} + \mathcal{H}_{\leq 0}$  in this part of the proof. Moreover, we will use the equivalent representation of **A** as the *implicit* equalities, denoted by  $\mathbf{A}^{=}$ , and other inequalities, denoted by  $\mathbf{A}^{+}$ , formally defined as follows.

**Definition 13** ((2) on page 99 of [57]). For any integer matrix **A**, let  $\mathbf{A}^{=}$  denote the implicit equalities, which is the maximal set of rows of **A** such that for all  $\vec{x} \in \mathcal{H}_{\leq 0}$ , we have  $\mathbf{A}^{=} \cdot (\vec{x})^{\top} = (\vec{0})^{\top}$ . Let  $\mathbf{A}^{+}$  denote the remaining rows of **A**.

We note that  $\mathbf{A}^{=}$  and  $\mathbf{A}^{+}$  do not depend on  $\vec{b}$ . As we will see soon,  $\mathbf{A}^{=}$  is the main constraint for  $\vec{X}_{\vec{\pi}}$  to be in  $\mathcal{H}_{\leq 0}$ . To simplify notation, throughout the proof we let  $o = \operatorname{Rank}(\mathbf{A}^{=}) = q - \dim(\mathcal{H}_{\leq 0})$ , where the equation holds due to [57, p. 100, Equation (9)]. The following running example illustrates these notions and its setting will be used throughout this proof.

**Example 4** (Running example:  $\mathcal{H}$  and  $A^{=}$  for Borda winners being  $\{1, 2\}$  under IC). We use a sub-case of k = 2 way ties over m = 3 alternatives under Borda w.r.t. IC for example. Notice that q = m! = 6. Each of the six outcomes is a linear order. Let  $1 \succ 2 \succ 3, 1 \succ 3 \succ 2, 2 \succ 1 \succ 3, 2 \succ 3 \succ 1, 3 \succ 1 \succ 2, 3 \succ 2 \succ 1$  denote outcomes 1, 2, 3, 4, 5, 6, respectively. For any  $n \in \mathbb{N}$ , let  $\vec{X}_{IC}$  denote the histogram of random profile under IC. Then, the Borda co-winners are  $\{1,2\}$  if and only if  $\vec{X}_{IC}$  is in polyhedron  $\mathcal{H}$  represented by the following linear inequalities, where the variables are  $\vec{x} = (x_{123}, x_{132}, x_{213}, x_{231}, x_{312}, x_{321})$ :

$$x_{123} + 2x_{132} - x_{213} - 2x_{231} + x_{312} - x_{321} \le 0 \tag{7}$$

$$-x_{123} - 2x_{132} + x_{213} + 2x_{231} - x_{312} + x_{321} \le 0$$
(8)

$$-2x_{123} - x_{132} - x_{213} + x_{231} + x_{312} + 2x_{321} \le -1 \tag{9}$$

Equation (7) states that the Borda score of alternative 1 is no more than the Borda score of alternative 2. Equation (8) states that the Borda score of 2 is no more than the Borda score of 1. Equation (9) states that the Borda score of 3 is at least one less than the Borda score of alternative 1. Because Borda scores are always integers, Equation (9) is equivalent to requiring that the Borda score of 3 is strictly smaller than the Borda score of 1. Then, we have:

$$\mathbf{A} = \begin{bmatrix} 1 & 2 & -1 & -2 & 1 & -1 \\ -1 & -2 & 1 & 2 & -1 & 1 \\ -2 & -1 & -1 & 1 & 1 & 2 \end{bmatrix} \text{ and } \vec{b} = \begin{bmatrix} 0 \\ 0 \\ -1 \end{bmatrix}$$

It is not hard to verify that  $\mathbf{A}^{=}$  consists of the first two rows, i.e.,

$$\mathbf{A}^{=} = \begin{bmatrix} 1 & 2 & -1 & -2 & 1 & -1 \\ -1 & -2 & 1 & 2 & -1 & 1 \end{bmatrix}, \text{ and } o = \operatorname{Rank}(\mathbf{A}^{=}) = 1$$

To calculate  $\operatorname{Pr}(\vec{X}_{\vec{\pi}} \in \mathcal{H})$ , we will focus on the reduced row echelon form (a.k.a. row canonical form) [44] of  $\begin{bmatrix} \mathbf{A}^{=} \\ \vec{1} \end{bmatrix}$ , which can be computed by Gauss-Jordan elimination: there exists  $I_0 \subseteq [q]$  with  $|I_0| = \operatorname{Rank}\left(\begin{bmatrix} \mathbf{A}^{=} \\ \vec{1} \end{bmatrix}\right)$  and a rational matrix  $\mathbf{D}$  such that  $\begin{bmatrix} \mathbf{A}^{=} \\ \vec{1} \end{bmatrix} \cdot (\vec{x})^{\top} = \begin{bmatrix} \begin{pmatrix} \vec{0} \\ n \end{pmatrix}^{\top} \end{bmatrix}$  if and only if  $(\vec{x}_{I_0})^{\top} = \mathbf{D} \cdot (\vec{x}_{I_1}, n)^{\top}$ , where  $I_1 = [q] \setminus I_0$ . In other words,  $\vec{x}_{I_1}$  can be viewed as "free" variables whose value can be quite flexible, and for any  $\vec{x} \in \mathcal{H}_{\leq 0}$ ,  $\vec{x}_{I_0}$  is completely determined by  $\vec{x}_{I_1}$ .

We have  $|I_0| = \operatorname{Rank}\left(\begin{bmatrix} \mathbf{A}^{=} \\ \vec{1} \end{bmatrix}\right) = \operatorname{Rank}(\mathbf{A}^{=}) + 1$  because  $\vec{1}$  is linearly independent with the

rows in  $\mathbf{A}^{=}$ . To see why this is true, suppose for the sake of contradiction that  $\vec{1}$  is linear dependent with rows in  $\mathbf{A}^{=}$ . Then, by the definition of  $\mathbf{A}^{=}$ , for all  $\vec{x} \in \mathcal{H}_{\leq 0}$  we have  $\vec{x} \cdot \vec{1} = 0$ . Therefore, for any  $\vec{x}' \in \mathcal{H}$ , according to Minkowski-Weyl theorem, we can write  $\vec{x}' = \vec{v} + \vec{h}$ , where  $\vec{h} \in \mathcal{H}_{\leq 0}$  and  $\vec{v}$ is in a finitely generated polyhedron. This means that

$$\vec{x}' \cdot \vec{1} = \vec{v} \cdot \vec{1} + \vec{h} \cdot \vec{1} = \vec{v} \cdot \vec{1},$$

which means that  $\vec{x}' \cdot \vec{l}$  is upper bounded by a constant. However, this contradicts the premise of the polynomial case, because when n is sufficiently large,  $\mathcal{H}_n = \mathcal{H}_n^{\mathbb{Z}} = \emptyset$ . W.l.o.g. let  $I_0 = \{1, \ldots, o+1\}$  and  $I_1 = \{o+2, \ldots, q\}$ . We also note that  $\mathbf{D}$  does not depend on n, which means that for any  $\vec{x} \in \mathbb{R}^q$ ,  $\mathbf{A}^{=} \cdot (\vec{x})^{\top} = (\vec{0})^{\top}$  if and only if  $(\vec{x}_{I_0})^{\top} = \mathbf{D} \cdot (\vec{x}_{I_1}, \vec{x} \cdot \vec{1})^{\top}$ .

The following example illustrates Gauss-Jordan elimination in the setting in Example 4.

Example 5 (Running example: Gauss-Jordan elimination, D,  $I_0$ , and  $I_1$ ). Continuing Example 4, we run Gauss-Jordan elimination as follows.

$$\begin{bmatrix} \mathbf{A} & 0 \\ 1 & n \end{bmatrix} = \begin{bmatrix} 1 & 2 & -1 & -2 & 1 & -1 & 0 \\ -1 & -2 & 1 & 2 & -1 & 1 & 0 \\ 1 & 1 & 1 & 1 & 1 & 1 & n \end{bmatrix}$$

$$\xrightarrow{R1;R2+R1;R3-R1} \begin{bmatrix} 1 & 2 & -1 & -2 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 2 & 3 & 0 & 2 & n \end{bmatrix}$$

$$\xrightarrow{R1+2R3;R2;-R3} \begin{bmatrix} 1 & 0 & 3 & 4 & 1 & 3 & 2n \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -2 & -3 & 0 & -2 & -n \end{bmatrix}$$

Let  $\vec{x} = [x_{123}, x_{132}, x_{213}, x_{231}, x_{312}, x_{321}]$ , we have

$$\begin{bmatrix} 1 & 0 & 3 & 4 & 1 & 3 \\ 0 & 1 & -2 & -3 & 0 & -2 \end{bmatrix} \cdot (\vec{x})^{\top} = \begin{bmatrix} 2n \\ -n \end{bmatrix},$$

which is equivalent to

$$\begin{bmatrix} x_{123} \\ x_{132} \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \times \begin{bmatrix} x_{123} \\ x_{132} \end{bmatrix} = \begin{bmatrix} -3 & -4 & -1 & -3 & 2 \\ 2 & 3 & 0 & 2 & -1 \end{bmatrix} \times \begin{bmatrix} x_{213} \\ x_{231} \\ x_{312} \\ x_{321} \\ n \end{bmatrix}$$

Therefore, we have  $\mathbf{D} = \begin{bmatrix} -3 & -4 & -1 & -3 & 2\\ 2 & 3 & 0 & 2 & -1 \end{bmatrix}$ ,  $I_0 = \{1, 2\}$ ,  $I_1 = \{3, 4, 5, 6\}$ .

As in [66], we adopt the following alternative representation of  $Y_1, \ldots, Y_n$ . For each  $j \leq n$ , we use a random variable  $Z_j \in \{0, 1\}$  to represent whether the outcome of  $Y_j$  is in  $I_0$  (corresponding to  $Z_j = 0$ ) or is in  $I_1$  (corresponding to  $Z_j = 1$ ). Then, we use another random variable  $W_j \in [q]$  to represent the outcome of  $Y_j$  conditioned on  $Z_j$ .

**Definition 14** (Alternative representation of  $Y_1, \ldots, Y_n$  [66]). For each  $j \leq n$ , we define a Bayesian network with two random variables  $Z_j \in \{0, 1\}$  and  $W_j \in [q]$ , where  $Z_j$  is the parent of  $W_j$ . The conditional probabilities are as follows.

- For each  $\ell \in \{0, 1\}$ ,  $\Pr(Z_i = \ell) = \Pr(Y_i \in I_\ell)$ .
- For each  $\ell \in \{0, 1\}$  and each  $t \le q$ ,  $\Pr(W_j = t | Z_j = \ell) = \Pr(Y_j = t | Y_j \in I_\ell)$ .

In particular, if  $t \notin I_{\ell}$  then  $\Pr(W_j = t | Z_j = \ell) = 0$ .

Example 6 (Running example: alternative representation of uniformly distributed  $Y_j$ ). For the purpose of presentation, we present  $Z_j$  and  $W_j$  for  $Y_j$  that corresponds to the uniform distribution over  $\mathcal{L}(\mathcal{A})$ . We have  $\Pr(Z_j = 0) = 1/3$  and  $\Pr(Z_j = 1) = 2/3$ .

$$\Pr(W_j = 1 \succ 2 \succ 3 | Z_j = 0) = \Pr(W_j = 1 \succ 3 \succ 2 | Z_j = 0) = \frac{1}{2}, \text{ and}$$
  
$$\Pr(W_j = 2 \succ 1 \succ 3 | Z_j = 1) = \Pr(W_j = 2 \succ 3 \succ 1 | Z_j = 1)$$
  
$$= \Pr(W_j = 3 \succ 1 \succ 2 | Z_j = 1) = \Pr(W_j = 3 \succ 2 \succ 1 | Z_j = 1) = \frac{1}{4}$$

All conditional probabilities not defined above are zeros.

Applying the law of total probability, it is not hard to verify that for any  $j \leq n$ ,  $W_j$  follows the same distribution as  $Y_j$ . For any  $\vec{z} \in \{0, 1\}^n$ , we let  $\mathrm{Id}_0(\vec{z}) \subseteq [n]$  denote the indices of components of  $\vec{z}$  that equal to 0. Given  $\vec{z}$ , we define the following random variables.

- Let  $\vec{W}_{\mathrm{Id}_0(\vec{z})} = \{W_j : j \in \mathrm{Id}_0(\vec{z})\}$ . That is,  $\vec{W}_{\mathrm{Id}_0(\vec{z})}$  consists of random variables  $\{W_j : z_j = 0\}$ .
- Let  $\operatorname{Hist}(\vec{W}_{\operatorname{Id}_0(\vec{z})})$  denote the vector of the  $o + 1 = |I_0|$  random variables that correspond to the histogram of  $\vec{W}_{\operatorname{Id}_0(\vec{z})}$  restricted to  $I_0$ . Technically, the domain of every random variable in  $\vec{W}_{\operatorname{Id}_0(\vec{z})}$  is [q], but since they only receive positive probabilities on  $I_0$ , they are treated as random variables over  $I_0$  when  $\operatorname{Hist}(\vec{W}_{\operatorname{Id}_0(\vec{z})})$  is defined.
- Similarly, let  $\vec{W}_{\mathrm{Id}_1(\vec{z})} = \{W_j : j \in \mathrm{Id}_1(\vec{z})\}$  and let  $\mathrm{Hist}(\vec{W}_{\mathrm{Id}_1(\vec{z})})$  denote the vector of  $|I_1| = q o 1$  random variables that correspond to the histogram of  $\vec{W}_{\mathrm{Id}_1(\vec{z})}$ .

Example 7 (Running example:  $Id_0(\vec{z})$  and  $Id_1(\vec{z})$ ). Continuing Example 6, suppose n = 5 and  $\vec{z} = (0, 1, 1, 0, 1)$ . We have

- $Id_0(\vec{z}) = \{1, 4\}, \ \vec{W}_{Id_0(\vec{z})} = \vec{W}_{\{1,4\}} = \{W_1, W_4\}, \ and \ Hist(\vec{W}_{Id_0(\vec{z})}) \ represents \ the \ histogram \ of two \ i.i.d. \ uniform \ distributions \ over \ \{1 \succ 2 \succ 3, 1 \succ 3 \succ 2\}.$
- $Id_1(\vec{z}) = \{2, 3, 5\}, \ \vec{W}_{Id_1(\vec{z})} = \vec{W}_{\{2,3,5\}} = \{W_2, W_3, W_5\}, \ and \ Hist(\vec{W}_{Id_1(\vec{z})}) \ represents \ the \ histogram \ of \ three \ i.i.d. \ uniform \ distributions \ over \ \{2 \succ 1 \succ 3, 2 \succ 3 \succ 1, 3 \succ 1 \succ 2, 3 \succ 2 \succ 1\}.$

For any  $\vec{h}_1 \in \mathbb{Z}_{\geq 0}^{q-o-1}$ , we let  $\mathcal{H}_n^{\mathbb{Z}}|_{\vec{h}_1}$  denote the  $I_0$  components of  $\vec{h} \in \mathcal{H}_n^{\mathbb{Z}}$  whose  $I_1$  components are  $\vec{h}_1$ . Formally,

$$\mathcal{H}_n^{\mathbb{Z}}|_{\vec{h}_1} = \{\vec{h}_0 \in \mathbb{Z}_{\geq 0}^{o+1} : (\vec{h}_0, \vec{h}_1) \in \mathcal{H}_n^{\mathbb{Z}}\}$$

By definition,  $\vec{x} \in \mathcal{H}_n^{\mathbb{Z}}$  if and only if  $\vec{x}_{I_0} \in \mathcal{H}_n^{\mathbb{Z}}|_{\vec{x}_{I_1}}$ . We note that  $\mathcal{H}_n^{\mathbb{Z}}|_{\vec{h}_1}$  can contain two or more elements, because even though  $\vec{x}_{I_0}$  is completely determined by  $\vec{x}_{I_1}$  for any  $\vec{x} \in \mathcal{H}_{\leq 0}$ , this relationship may not hold for  $\vec{x} \in \mathcal{H}_n^{\mathbb{Z}}$ . Later in Claim 1 we will prove that the number of vectors in  $\mathcal{H}_n^{\mathbb{Z}}|_{\vec{h}_1}$  is bounded above by a constant that does not depend on n.

**Example 8** (**Running example:**  $\mathcal{H}_{n}^{\mathbb{Z}}|_{\vec{h}_{1}}$ ). Continuing Example 7, let  $\vec{h}_{1} = (1, 1, 1, 0)$ . Then, we have  $\mathcal{H}_{n}^{\mathbb{Z}}|_{\vec{h}_{1}} = \{(2, 0)\}$ . Notice that in this example  $|\mathcal{H}_{n}^{\mathbb{Z}}|_{\vec{h}_{1}}| \leq 1$  for all  $\vec{h}_{1}$ , because the  $\vec{b}$  components corresponding to  $\mathbf{A}^{=}$  are 0's, which means that  $\vec{h}_{0}$  is determined by  $\vec{h}_{1}$ , i.e.,  $\vec{h}_{0} = \mathbf{D} \cdot (\vec{h}_{1}, n)^{\top}$ .

It is possible that  $\mathcal{H}_{n}^{\mathbb{Z}}|_{\vec{h}_{1}} = \emptyset$ , because  $\vec{h}_{0} = \mathbf{D} \cdot (\vec{h}_{1}, n)^{\top}$  may not be a vector of non-negative integers. For example, when  $\vec{h}_{1} = (1, 2, 0, 0)$ , we have  $\vec{h}_{0} = \mathbf{D} \cdot (\vec{h}_{1}, n)^{\top} = (-1, 3)$ , which means that  $\mathcal{H}_{n}^{\mathbb{Z}}|_{\vec{h}_{1}} = \emptyset$ .

Next, we apply the law of total probability to the  $(\vec{Z}, \vec{W})$  representation of  $\vec{X}_{\vec{\pi}}$  and  $\mathcal{H}_n^{\mathbb{Z}}|_{\vec{h}_1}$ , to

obtain the following estimate on  $\Pr(\vec{X}_{\vec{\pi}} \in \mathcal{H}) = \Pr(\vec{X}_{\vec{\pi}} \in \mathcal{H}_n) = \Pr(\vec{X}_{\vec{\pi}} \in \mathcal{H}_n^{\mathbb{Z}}).$ 

$$\Pr(\vec{X}_{\vec{\pi}} \in \mathcal{H}) = \Pr(\vec{X}_{\vec{\pi}} \in \mathcal{H}_{n}^{\mathbb{Z}})$$

$$= \sum_{\vec{z} \in \{0,1\}^{n}} \Pr(\vec{Z} = \vec{z}) \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_{n}^{\mathbb{Z}} \middle| \vec{Z} = \vec{z}\right) \quad (\text{The law of total probability})$$

$$= \sum_{\vec{z} \in \{0,1\}^{n}} \Pr(\vec{Z} = \vec{z}) \Pr\left(\operatorname{Hist}(\vec{W}_{\operatorname{Id}_{0}(\vec{z})}) \in \mathcal{H}_{n}^{\mathbb{Z}}|_{\operatorname{Hist}(\vec{W}_{\operatorname{Id}_{1}(\vec{z})})} \middle| \vec{Z} = \vec{z}\right)$$

$$= \sum_{\vec{z} \in \{0,1\}^{n}} \Pr(\vec{Z} = \vec{z}) \sum_{\vec{h}_{1} \in \mathbb{Z}_{\geq 0}^{q-o-1}} \Pr\left(\operatorname{Hist}(\vec{W}_{\operatorname{Id}_{1}(\vec{z})}) = \vec{h}_{1} \middle| \vec{Z} = \vec{z}\right)$$

$$\times \Pr\left(\operatorname{Hist}(\vec{W}_{\operatorname{Id}_{0}(\vec{z})}) \in \mathcal{H}_{n}^{\mathbb{Z}}|_{\vec{h}_{1}} \middle| \vec{Z} = \vec{z}, \operatorname{Hist}(\vec{W}_{\operatorname{Id}_{1}(\vec{z})}) = \vec{h}_{1}\right) \quad (\text{The law of total probability})$$

$$= \sum_{\vec{z} \in \{0,1\}^{n}} \Pr(\vec{Z} = \vec{z}) \sum_{\vec{h}_{1} \in \mathbb{Z}_{\geq 0}^{q-o-1}} \Pr\left(\operatorname{Hist}(\vec{W}_{\operatorname{Id}_{1}(\vec{z})}) = \vec{h}_{1} \middle| \vec{Z} = \vec{z}\right) \times \Pr\left(\operatorname{Hist}(\vec{W}_{\operatorname{Id}_{0}(\vec{z})}) \in \mathcal{H}_{n}^{\mathbb{Z}}|_{\vec{h}_{1}} \middle| \vec{Z} = \vec{z}\right)$$

$$(10)$$

$$\leq \sum_{\vec{z} \in \{0,1\}^n : |\mathrm{Id}_0(\vec{z})| \ge 0.9\epsilon n} \Pr(\vec{Z} = \vec{z}) \sum_{\vec{h}_1 \in \mathbb{Z}_{\ge 0}^{q-o-1}} \Pr\left(\mathrm{Hist}(\vec{W}_{\mathrm{Id}_1(\vec{z})}) = \vec{h}_1 \mid \vec{Z} = \vec{z}\right) \\ \times \Pr\left(\mathrm{Hist}(\vec{W}_{\mathrm{Id}_0(\vec{z})}) \in \mathcal{H}_n^{\mathbb{Z}}|_{\vec{h}_1} \mid \vec{Z} = \vec{z}\right) + \Pr(|\mathrm{Id}_0(\vec{z})| < 0.9\epsilon n)$$
(11)

where we recall that  $|Id_0(\vec{z})|$  denotes the number of 0's in  $\vec{z}$ . Equation (10) holds because according to the Bayesian network structure,  $W_j$ 's are independent of each other given  $Z_j$ 's, which means that for any  $\vec{z} \in \{0,1\}^n$ ,  $Hist(\vec{W}_{Id_0(\vec{z})})$  and  $Hist(\vec{W}_{Id_1(\vec{z})})$  are independent given  $\vec{z}$ . The following example illustrates the summand in (11) when  $\vec{z} = (0, 1, 1, 0, 1)$  and  $0.9\epsilon < \frac{2}{5}$ , following the setting of Example 8.

**Example 9** (Running example: summand in (11)). Continuing Example 8, the summand in (11) becomes the following:

$$\begin{aligned} \Pr(\vec{Z} = \vec{z}) & \sum_{\vec{h}_1 \in \mathbb{Z}_{\geq 0}^{q-o-1}} \Pr\left( \text{Hist}(\vec{W}_{Id_1(\vec{z})}) = \vec{h}_1 \mid \vec{Z} = \vec{z} \right) \times \Pr\left( \text{Hist}(\vec{W}_{Id_0(\vec{z})}) \in \mathcal{H}_n^{\mathbb{Z}} \mid_{\vec{h}_1} \mid \vec{Z} = \vec{z} \right) \\ = \Pr(\vec{Z} = (0, 1, 1, 0, 1)) & \sum_{\vec{h}_1 \in \mathbb{Z}_{\geq 0}^{q-o-1}} \Pr\left( \text{Hist}(\vec{W}_{\{2,3,5\}}) = \vec{h}_1 \mid \vec{Z} = (0, 1, 1, 0, 1) \right) \\ & \times \Pr\left( \text{Hist}(\vec{W}_{\{1,4\}}) \in \mathcal{H}_n^{\mathbb{Z}} \mid_{\vec{h}_1} \mid \vec{Z} = (0, 1, 1, 0, 1) \right) \end{aligned}$$

As a more concrete example, let  $\vec{h}_1 = (1, 1, 1, 0)$  as in Example 8, we summand in the equation above becomes the following:

$$\Pr\left(\textit{Hist}(\vec{W}_{\{2,3,5\}}) = (1,1,1,0) \mid \vec{Z} = (0,1,1,0,1)\right) \times \Pr\left(\textit{Hist}(\vec{W}_{\{1,4\}}) \in \{(2,0)\} \mid \vec{Z} = (0,1,1,0,1)\right)$$

In words, it is the product of the following two terms:

- (1) the probability for the histogram of  $\vec{W}_{\{2,3,5\}}$  to be (1,1,1,0). I.e., the second, third, and fifth agents' votes are  $\{2 \succ 1 \succ 2, 2 \succ 3 \succ 1, 3 \succ 1 \succ 2\}$  in any order, and
- (2) the probability for the histogram of  $\vec{W}_{\{1,4\}}$  to be (2,0). I.e., both the first and the fourth agents vote for  $1 \succ 2 \succ 3$ .

We emphasize that this example only illustrates (11) for  $\vec{z} = (0, 1, 1, 0, 1)$  and  $\vec{h}_1 = (1, 1, 1, 0)$ . (11) requires summing over other combinations of  $\vec{z}$  and  $\vec{h}_1$ .

To upper-bound (11), we will show that given  $|\mathrm{Id}_0(\vec{z})| \ge 0.9\epsilon n$ , for any  $\vec{h}_1 \in \mathbb{Z}_{\ge 0}^{q-o-1}$ ,

$$\Pr\left(\operatorname{Hist}(\vec{W}_{\operatorname{Id}_{0}(\vec{z})}) \in \mathcal{H}_{n}^{\mathbb{Z}}|_{\vec{h}_{1}} \mid \vec{Z} = \vec{z}\right) = O((0.9\epsilon n)^{-\frac{o}{2}}) = O(n^{-\frac{o}{2}})$$
(12)

(12) follows after combining the following two parts.

• Part 1: Claim 1 below, which states that the number of integer vectors in  $\mathcal{H}_n^{\mathbb{Z}}|_{\vec{h}_1}$  is upper bounded by a constant that only depends on  $\mathcal{H}$ , which means that it does not depend on n.

Claim 1. There exists a constant  $C_{\mathcal{H}} > 0$  such that for each  $\vec{h}_1 \in \mathbb{Z}_{\geq 0}^{q-o-1}$ ,  $|\mathcal{H}_n^{\mathbb{Z}}|_{\vec{h}_1}| \leq C_{\mathcal{H}}$ .

*Proof.* We first prove an observation, which states that there exists a constant  $C^*$  that only depends on  $\mathcal{H}$ , such that for any  $\vec{x} \in \mathcal{H}_n^{\mathbb{Z}}$ , we have  $|(\vec{x}_{I_0})^{\top} - \mathbf{D} \cdot (\vec{x}_{I_1}, n)^{\top}|_{\infty} < C^*$ .

According to the V-representation of  $\mathcal{H}$ , we can write  $\vec{x} = \vec{v} + \vec{x}'$ , where  $\vec{v} \in \mathcal{V}$  is bounded, and  $\vec{x}' \in \mathcal{H}_{\leq 0}$  which means that  $(\vec{x}'_{I_0})^{\top} - \mathbf{D} \cdot (\vec{x}'_{I_1}, \vec{x}' \cdot \vec{1})^{\top} = (\vec{0})^{\top}$ . Recall that  $C_{\max}$  is the maximum  $L_{\infty}$  norm of vectors in  $\mathcal{V}$ . It follows that  $|\vec{v}|_{\infty} \leq C_{\max}$ , which means that

$$\left|\vec{v}\cdot\vec{1}\right| = \left|\vec{x}\cdot\vec{1} - \vec{x}'\cdot\vec{1}\right| = \left|n - \vec{x}'\cdot\vec{1}\right| \le qC_{\max}$$

Let C' denote the maximum absolute value of entries in **D** and let  $C^* = C_{\max} + 2qC_{\max}C'$ . We have:

$$|(\vec{x}_{I_0})^{\top} - \mathbf{D} \cdot (\vec{x}_{I_1}, n)^{\top}|_{\infty} = |(\vec{v}_{I_0} + \vec{x}'_{I_0})^{\top} - \mathbf{D} \cdot (\vec{v}_{I_1} + \vec{x}'_{I_1}, n)^{\top}|_{\infty}$$
$$= |(\vec{v}_{I_0})^{\top} - \mathbf{D} \cdot (\vec{v}_{I_1}, n - \vec{x}' \cdot \vec{1})^{\top}|_{\infty} \le C_{\max} + 2qC_{\max}C' = C^*$$

For any  $\vec{h}_0 \in \mathcal{H}_n^{\mathbb{Z}}|_{\vec{h}_1}$ , because  $(\vec{h}_0, \vec{h}_1) \in \mathcal{H}_n^{\mathbb{Z}}$ , according to the observation above, we have

$$|\left(\vec{h}_{0}\right)^{\top} - \mathbf{D} \cdot \left(\vec{h}_{1}, n\right)^{\top}|_{\infty} < C^{*}$$

Therefore,  $\mathcal{H}_{n}^{\mathbb{Z}}|_{\vec{h}_{1}}$  is contained in an  $|I_{0}|$ -dimensional cube whose edge length is  $2C^{*}$  and is centered at  $\mathbf{D} \cdot (\vec{h}_{1}, n)^{\top}$ . It is not hard to verify that the cube contains no more than  $(2C^{*}+1)^{q}$  integer points. This proves claim by letting  $C_{\mathcal{H}} = (2C^{*}+1)^{q}$ .

• Part 2: The point-wise anti-concentration bound [66, Lemma 3 in the Appendix]. For completeness, we recall the lemma in our notation below.

**Lemma'** ([66, Lemma 3 in the Appendix]). Given  $q^* \in \mathbb{N}$  and  $\epsilon > 0$ , there exists a constant  $C^* > 0$  such that for any  $n^* \in \mathbb{N}$  and strictly positive (by  $\epsilon$ ) vector  $\vec{\pi}^*$  of  $n^*$ distributions over  $[q^*]$ , and any vector  $\vec{x}^* \in \mathbb{Z}_{\geq 0}^{q^*}$ , we have  $\Pr(\vec{X}_{\vec{\pi}^*} = \vec{x}^*) < C^*(n^*)^{\frac{1-q^*}{2}}$ .

We note that the constant in  $O(n^{-\frac{o}{2}})$  in (12) only depends on  $\mathcal{H}$  (therefore q) and  $\epsilon$  but not on  $\Pi$  or n.

Then, (12) follows after applying Lemma' to (constantly many) vectors in  $\mathcal{H}_n^{\mathbb{Z}}|_{\vec{h}_1}$  (guaranteed by Claim 1), by letting  $q^* = |I_0| = o+1$  and  $n^* = |\mathrm{Id}_0(\vec{z})|$ . The next example illustrates the application of Lemma' in the setting of Example 9.

**Example 10** (Running example: Equation (12)). Continuing Example 9, recall that in this running example  $\vec{z} = (0, 1, 1, 0, 1)$  (Example 7) and  $\vec{h}_1 = (1, 1, 1, 0)$  (Example 8). Therefore, (12) becomes:

$$\Pr\left(\textit{Hist}(\vec{W}_{\{1,4\}}) \in \{(2,0)\} \mid \vec{Z} = (0,1,1,0,1)\right)$$

Then, we let  $q^* = |I_0| = 2$  and  $n^* = 2$  in Lemma' and apply it to  $\vec{x}^* = (2,0)$ , which is the only vector in  $\mathcal{H}_n^{\mathbb{Z}}|_{\vec{h}_1}$ .

Back to (11), we now upper-bound the  $\Pr(|\mathrm{Id}_0(\vec{z})| < 0.9\epsilon n)$  part in (11). Because random variables in  $\vec{Y}$  are above  $\epsilon$ , for all  $j \leq n$ ,  $Z_j$  takes 0 with probability at least  $\epsilon$ . Therefore,  $\mathbb{E}(|\mathrm{Id}_0(\vec{Z})|) \geq \epsilon n$ . By Hoeffding's inequality,  $\Pr(|\mathrm{Id}_0(\vec{Z})| < 0.9\epsilon n)$  is exponentially small in n, which is  $O(n^{-\frac{o}{2}})$  when n is sufficiently large.

Putting all together, we have:

$$\Pr(\vec{X}_{\vec{\pi}} \in \mathcal{H}) \leq \sum_{\vec{z} \in \{0,1\}^n : |\mathrm{Id}_0(\vec{z})| \geq 0.9\epsilon n} \Pr(\vec{Z} = \vec{z}) \sum_{\vec{h}_1 \in \mathbb{Z}_{\geq 0}^{q-o-1}} \Pr\left(\mathrm{Hist}(\vec{W}_{\mathrm{Id}_1(\vec{z})}) = \vec{h}_1 \mid [\vec{Z}]_{\mathrm{Id}_1(\vec{z})} = \vec{1}\right) \\ \times O(n^{-\frac{o}{2}}) + O(n^{-\frac{o}{2}}) = O(n^{-\frac{o}{2}})$$

This proves the polynomial upper bound when  $\mathcal{H}_n^{\mathbb{Z}} \neq \emptyset$  and  $\mathcal{H}_{\leq 0} \cap \operatorname{CH}(\Pi) \neq \emptyset$ , where the constant in  $O(n^{-\frac{o}{2}})$  depends on  $\mathcal{H}$  and  $\epsilon$  (but not on  $\Pi$  or n).

**Proof of the polynomial lower bound on Sup.** The proof is done in the following five steps. In **Step 1**, for any  $n \in \mathbb{N}$  that is sufficiently large, we define a non-negative integer vector  $\vec{y}^n \in \mathcal{H}_n^{\mathbb{Z}}$  and its neighborhood  $B^n \subseteq \mathcal{H}_n$  such that vectors in  $B^n$  are  $O(\sqrt{n})$  away from Cone(II). In **Step 2**, we define a vector  $\vec{\pi} \in \Pi^n$  chosen by the adversary to achieve the lower bound. In **Step 3**, we prove that  $B^n$  contains  $\Theta\left(n^{\frac{\dim(\mathcal{H}_{\leq 0})-1}{2}}\right)$  many non-negative integer vectors. In **Step 4**, we show that when preferences are generated according to  $\vec{\pi}$ , for any non-negative integer  $\vec{x} \in B^n$ , the probability for  $\vec{X}_{\vec{\pi}}$  to be  $\vec{x}$  is  $\Theta(n^{\frac{(1-q)}{2}})$ . Finally, in **Step 5** we show that the probability for  $\vec{X}_{\vec{\pi}}$  to be in  $\mathcal{H}$  is at least  $\Omega\left(n^{\frac{\dim(\mathcal{H}_{\leq 0})-1}{2}}\right) \times \Omega\left(n^{\frac{(1-q)}{2}}\right) = \Omega(n^{-\frac{o}{2}})$ .

Step 1. For any sufficiently large  $n \in \mathbb{N}$ , define a non-negative integer vector  $\vec{y}^n \in \mathcal{H}_n^{\mathbb{Z}}$ and its neighborhood  $B^n \subseteq \mathcal{H}_n$ .  $\vec{y}^n$  will be defined in Step 1.1 as an integer approximation to  $\vec{x}^{\ell} = \ell \vec{x}^* + \sqrt{\ell} \vec{x}^{@} + \vec{x}^{\#}$ , whose components are defined as follows. See Figure 7 for an illustration.

- $\vec{x}^* \in \mathcal{H}_{\leq 0} \cap CH(\Pi)$ , which may not be integral.
- $\vec{x}^{@}$  is an inner point of  $\mathcal{H}_{\leq 0}$ , which means that  $\mathbf{A}^{=} \cdot (\vec{x}^{@})^{\top} = (\vec{0})^{\top}$  and  $\mathbf{A}^{+} \cdot (\vec{x}^{@})^{\top} < (\vec{0})^{\top}$ . Note that  $\vec{x}^{@}$  may not be integral, non-negative, in  $\mathcal{H}$ , or in CH(II).
- $\vec{x}^{\#} \in \mathcal{H}$  is an integer vector, which may not be in Cone(II). For example,  $\vec{x}^{\#}$  can be any integer vector in  $\mathcal{H}_{n^{\#}}^{\mathbb{Z}}$  for the smallest  $n^{\#} \in \mathbb{N}$  such that  $\mathcal{H}_{n^{\#}}^{\mathbb{Z}} \neq \emptyset$ . Such  $n^{\#}$  exists because otherwise the 0 case of the theorem holds.

•  $\ell \in \mathbb{R}_{\geq 0}$  is a number that is used to guarantee  $\vec{y}^n \cdot \vec{1} = \vec{x}^\ell \cdot \vec{1} = n$ . That is,  $\vec{x}^\ell \cdot \vec{1} = \ell + \sqrt{\ell} (\vec{x}^{@} \cdot \vec{1}) + \vec{x}^{\#} \cdot \vec{1} = n$ , or equivalently, because  $\ell \geq 0$ , we have  $\ell = \left( -\frac{\vec{x}^{@} \cdot \vec{1}}{2} + \sqrt{n - \vec{x}^{\#} \cdot \vec{1} + (\frac{\vec{x}^{@} \cdot \vec{1}}{2})^2} \right)^2$ . We note that  $\ell$  may not be an integer.

 $\vec{x}^*$ ,  $\vec{x}^{@}$ , and  $\vec{x}^{\#}$  are arbitrarily chosen but fixed throughout the proof. Let us illustrate a choice of them in the following example.

**Example 11** (Running example:  $\vec{x}^*, \vec{x}^@, \vec{x}^\#$ , and  $\vec{x}^\ell$ ). Continuing Example 10, recall that  $\Pi$  only contains the uniform distribution. Therefore,  $\vec{x}^* = (\frac{1}{6}, \frac{1}{6}, \frac{1}{6}, \frac{1}{6}, \frac{1}{6}, \frac{1}{6})$ , which is the only distribution in  $\mathcal{H}_{\leq 0} \cap CH(\Pi)$ . Let  $\vec{x}^@ = (1.2, -0.2, 1.2, -0.2, 0, 0)$  and  $\vec{x}^\# = (2, 0, 1, 1, 1, 0)$  (where  $n^\# = \vec{x}^\# \cdot \vec{1} = 5$ ). Then,

$$\vec{x}^{l} = \ell \cdot \underbrace{(\underbrace{\frac{1}{6}, \frac{1}{6}, \frac{1}{6}, \frac{1}{6}, \frac{1}{6}, \frac{1}{6}, \frac{1}{6}}_{\vec{x}^{*}})}_{\vec{x}^{*}} + \sqrt{\ell} \cdot \underbrace{(\underbrace{1.2, -0.2, 1.2, -0.2, 0, 0}_{\vec{x}^{@}})}_{\vec{x}^{@}} + \underbrace{(\underbrace{2, 0, 1, 1, 1, 0}_{\vec{x}^{\#}})}_{\vec{x}^{\#}}$$

When n = 100,  $\ell + 2\sqrt{\ell} + 5 = 100$ , which means that  $\ell = (\sqrt{96} - 1)^2$ .

Step 1.1 Define a non-negative integer approximation  $\vec{y}^n$  to  $\vec{x}^{\ell}$ . First, we note that  $\ell$  only depends on  $\mathcal{H}$  (because of the choices of  $\vec{x}^{@}$  and  $\vec{x}^{\#}$ ) and n but not on  $\Pi$  or  $\epsilon$ . Notice that while  $\vec{x}^{\ell} \cdot \vec{1} = n$ , it may contain negative components because  $\vec{x}^{@}$  may contain negative components. Let n to be sufficiently large so that  $\vec{x}^{\ell} \in \mathbb{R}^{q}_{\geq 0}$ , which means that  $\vec{x}^{\ell} \in \mathcal{H}_{n}$ . This can be done because (i) each component of  $\vec{x}^{\ell}$  is  $\Theta(n)$ , because  $\vec{x}^{\ell}$  is largely determined by  $\ell \vec{x}^{*}$  and  $\ell = \Theta(n)$ , and (ii) each component of  $\vec{x}^{*}$  is at least  $\epsilon$ , because  $\vec{x}^{*} \in CH(\Pi)$ .

Let 
$$\mathbf{A}_n = \begin{bmatrix} \mathbf{A} \\ \vec{1} \\ -\vec{1} \\ -\mathbb{I}_q \end{bmatrix}$$
 and  $\vec{b}_n = (\vec{b}, n, -n, \underbrace{0, \dots, 0}_q)$ , where  $\mathbb{I}_q$  is the  $q \times q$  identity matrix. It follows

that  $\mathcal{H}_n$  is defined by  $\mathbf{A}_n$  and  $\vec{b}_n$ . In other words,

$$\mathcal{H}_n = \{ \vec{x} : \mathbf{A}_n \cdot \left( \vec{x} \right)^\top \le \left( \vec{b}_n \right)^\top \}$$

Let  $C_{\mathcal{H}}^1 = q\Delta(\mathbf{A}_n)$ , where  $\Delta(\mathbf{A}_n)$  is the maximum absolute value among the determinants of all square submatrices of  $\mathbf{A}_n$ . Notice that  $C_{\mathcal{H}}^1$  only depends on  $\mathbf{A}$  but not on n,  $\epsilon$ , or  $\Pi$ . It is not hard to verify that  $\vec{x}^{\ell} \in \mathcal{H}_n$  and recall that we have assumed  $\mathcal{H}_n^{\mathbb{Z}} \neq \emptyset$  as a condition for the polynomial bound on the sup part of the theorem, which means that  $\mathcal{H}_n$  contains an integer vector. Therefore, by [15, Theorem 1(i)] (where we let  $\vec{w} = \vec{0}$ ), there exists a (non-negative) integer vector  $\vec{y}^n \in \mathcal{H}_n$ such that  $|\vec{y}^n - \vec{x}^{\ell}|_{\infty} \leq C_{\mathcal{H}}^1$ . Note that by definition  $\vec{y}^n \in \mathcal{H}_n^{\mathbb{Z}}$ . For completeness, we recall [15, Theorem 1(i)] in our notation as follows.

**Theorem 1(i) in [15].** Let A be an  $L \times q$  integer matrix,  $\vec{b}$  be a q-dimensional vector, and let  $\vec{w}$  be a q-dimensional vector such that  $A \cdot (\vec{x})^{\top} \leq (\vec{b})^{\top}$  has an integer solution and  $\max\{\vec{w} \cdot \vec{x} : A \cdot (\vec{x})^{\top} \leq (\vec{b})^{\top}\}$  exists. Then, for each optimal solution  $\vec{x}_{opt}$  to  $\max\{\vec{w} \cdot \vec{x} : A \cdot (\vec{x})^{\top} \leq (\vec{b})^{\top}\}$ , there exists an optimal solution  $\vec{z}_{opt} \in \mathbb{Z}^q$  to  $\max\{\vec{w} \cdot \vec{x} : A \cdot (\vec{x})^{\top} \leq (\vec{b})^{\top}, \vec{x} \text{ is integral}\}$  with  $|\vec{x}_{opt} - \vec{z}_{opt}|_{\infty} \leq q\Delta(A)$ .
Step 1.2 Define  $\vec{y}^n$ 's neighborhood  $B^n \subseteq \mathcal{H}_n$ . For any  $\gamma > 0$ , we first define a neighborhood  $B_\gamma \subseteq \mathbb{R}^q$  of  $\vec{x}^{@}$  that consists of vectors  $\vec{x}$  such that (i)  $\mathbf{A}^{=} \cdot (\vec{x})^{\top} = (\vec{0})^{\top}$ , (ii)  $|\vec{x}_{I_1} - \vec{x}_{I_1}^{@}|_{\infty} \leq \gamma$ , and (iii)  $\vec{x} \cdot \vec{1} = \vec{x}^{@} \cdot \vec{1}$ . Formally,

$$B_{\gamma} \triangleq \{ \vec{x} : \forall i \in I_1, |x_i - x_i^{@}| \le \gamma \text{ and } (\vec{x}_{I_0})^{\top} = \mathbf{D} \cdot \left( \vec{x}_{I_1}, \vec{x}^{@} \cdot \vec{1} \right)^{\top} \}$$

Recall that  $\vec{x}^{@}$  is an inner point of  $\mathcal{H}_{\leq 0}$ , which means that  $\mathbf{A}^{=} \cdot (\vec{x}^{@})^{\top} = (\vec{0})^{\top}$ , or equivalently,  $(\vec{x}_{I_0}^{@})^{\top} = \mathbf{D} \cdot (\vec{x}_{I_1}^{@}, \vec{x}^{@} \cdot \vec{1})^{\top}$ . Therefore, for any  $\vec{x} \in B_{\gamma}$ , we have

$$\vec{x}_{I_0} - \vec{x}_{I_0}^{@} = \mathbf{D} \cdot \left[ \left( \vec{x}_{I_1}, \vec{x} \cdot \vec{1} \right)^\top - \left( \vec{x}_{I_1}^{@}, \vec{x}^{@} \cdot \vec{1} \right)^\top \right] = \mathbf{D} \cdot \left( \vec{x}_{I_1} - \vec{x}_{I_1}^{@}, 0 \right)^\top$$

Therefore, it is not hard to verify that

$$B_{\gamma} - \vec{x}^{@} = \{ \vec{\Delta} \in \mathbb{R}^{q} : \forall i \in I_{1}, |\Delta_{i}| \leq \gamma \text{ and } \left( \vec{\Delta}_{I_{0}} \right)^{\top} = \mathbf{D} \cdot \left( \vec{\Delta}_{I_{1}}, 0 \right)^{\top} \}$$
(13)

Recall that  $\vec{x}^{@}$  is an inner point of  $\mathcal{H}_{\leq 0}$ , which means that

$$\mathbf{A}^{=} \cdot \left( \vec{x}^{@} \right)^{\top} = \left( \vec{0} \right)^{\top} \text{ and } \mathbf{A}^{+} \cdot \left( \vec{x}^{@} \right)^{\top} < \left( \vec{0} \right)^{\top}$$

Therefore, there exists  $\gamma > 0$  such that for any  $\vec{x} \in B_{\gamma}$ , we have  $\mathbf{A}^{=} \cdot (\vec{x})^{\top} = (\vec{0})^{\top}$  and  $\mathbf{A}^{+} \cdot (\vec{x})^{\top} < (\vec{0})^{\top}$ . In other words, all vectors in  $B_{\gamma}$  are inner points in  $\mathcal{H}_{\leq 0}$ . It is not hard to verify that  $\dim(B_{\gamma}) = q - o - 1 = \dim(\mathcal{H}_{\leq 0}) - 1$ , where the -1 comes from the additional linear constraint  $\vec{x} \cdot \vec{1} = \vec{x}^{@} \cdot \vec{1}$ . Notice that  $B_{\gamma}$  only depends on  $\mathcal{H}_{\leq 0}$  but not on  $n, \epsilon$ , or II. Let

$$B^n \triangleq \vec{y}^n + \sqrt{\ell}(B_\gamma - \vec{x}^@) = \{\vec{y}^n + \sqrt{\ell}\vec{\Delta} : \vec{\Delta} \in B_\gamma - \vec{x}^@\}$$

Intuitively,  $B^n$  is defined by first scaling up  $(B_{\gamma} - \vec{x}^{\textcircled{0}})$  by  $\sqrt{\ell}$ , and then add it on top of  $\vec{y}^n$ . This means that at a high level  $B^n$  consists of inner points of a local space that is similar to  $\mathcal{H}_{\leq 0}$  and is centered at  $\vec{y}^n$ , plus an additional linear constraint that requires  $\vec{x} \cdot \vec{1} = n$ . Note that  $\vec{x}^{\ell}$  may not be in  $B^n$ .

**Example 12** (Running example:  $B_{\gamma}$  and  $B_n$ ). Continuing Example 11, let  $\gamma = 1$ , we have

$$B_{1} = \{ (-3x_{213} - 4x_{231} - x_{312} - 3x_{321} + 4, 2x_{213} + 3x_{231} + 2x_{321} - 2, x_{213}, x_{231}, x_{312}, x_{321}) : x_{213} \in [0.2, 2.2], x_{231} \in [-1.2, 0.8], x_{312} \in [-1, 1], x_{321} \in [-1, 1] \}$$

$$B_1 - \vec{x}^{@} = \{ (-3\Delta_{213} - 4\Delta_{231} - \Delta_{312} - 3\Delta_{321}, 2\Delta_{213} + 3\Delta_{231} + 2\Delta_{321}, \Delta_{213}, \Delta_{231}, \Delta_{312}, \Delta_{321}) : (\Delta_{213}, \Delta_{231}, \Delta_{312}, \Delta_{321}) \in [-1, 1]^4 \}$$

Let n = 100, we have  $B^{100} = \vec{y}^n + (\sqrt{96} - 1)^2 \cdot (B_1 - \vec{x}^{@})$ .

In the following three steps, we prove  $B^n \subseteq \mathcal{H}_n$  for any sufficiently large n.

• First, we prove  $B^n \subseteq \mathcal{H}$ . Because  $\vec{y}^n \in \mathcal{H}_n^{\mathbb{Z}}$ , we have  $\mathbf{A}^{=} \cdot (\vec{y}^n)^{\top} \leq (\vec{b}^{=})^{\top}$ , where  $\vec{b}^{=}$  is the subvector of  $\vec{b}$  that corresponds to  $\mathbf{A}^{=}$ . For all  $\vec{\Delta} \in B_{\gamma} - \vec{x}^{@}$ , recall from (13) that  $\vec{\Delta}_{I_0} = \mathbf{D} \cdot (\vec{\Delta}_{I_1}, 0)^{\top}$ , which means that  $\mathbf{A}^{=} \cdot (\Delta)^{\top} = (\vec{0})^{\top}$ . Therefore,

$$\mathbf{A}^{=} \cdot \left( \vec{y}^{n} + \sqrt{\ell} \vec{\Delta} \right)^{\top} = \mathbf{A}^{=} \cdot \left( \vec{y}^{n} \right)^{\top} + \mathbf{A}^{=} \cdot \left( \sqrt{\ell} \vec{\Delta} \right)^{\top} = \mathbf{A}^{=} \cdot \left( \vec{y}^{n} \right)^{\top} \le \left( \vec{b}^{=} \right)^{\top}$$

Also notice that:

$$\vec{y}^n + \sqrt{\ell}\vec{\Delta} = \vec{y}^n - \vec{x}^\ell + \vec{x}^\ell + \sqrt{\ell}\vec{\Delta} = (\vec{y}^n - \vec{x}^\ell) + \ell\vec{x}^* + \sqrt{\ell}(\vec{x}^{@} + \vec{\Delta}) + \vec{x}^{\#}$$

Therefore, we have the following bound on  $\mathbf{A}^+ \cdot \left( \vec{y}^n + \sqrt{\ell} \vec{\Delta} \right)^\top$ :

$$\mathbf{A}^{+} \cdot \left(\vec{y}^{n} + \sqrt{\ell}\vec{\Delta}\right)^{\top} = \mathbf{A}^{+} \cdot \left(\vec{y}^{n} - \vec{x}^{\ell}\right)^{\top} + \mathbf{A}^{+} \cdot \left(\ell\vec{x}^{*}\right)^{\top} + \mathbf{A}^{+} \cdot \left(\sqrt{\ell}(\vec{x}^{@} + \vec{\Delta})\right)^{\top} + \mathbf{A}^{+} \cdot \left(\vec{x}^{\#}\right)^{\top} \leq \left(O(1) \cdot \vec{1} + \vec{0} - \Omega(\sqrt{\ell}) \cdot \vec{1}\right)^{\top} + \left(\vec{b}^{+}\right)^{\top}$$
(14)

(14) follows after noticing that  $|\vec{y}^n - \vec{x}^\ell|_{\infty} = O(1), \ \vec{x}^* \in \mathcal{H}_{\leq 0}, \ \vec{x}^{@} + \vec{\Delta}$  is an inner point in  $\mathcal{H}_{\leq 0}$  (which mean that  $\mathbf{A}^+ \cdot \left(\vec{x}^{@} + \vec{\Delta}\right)^\top \leq -\Omega(1) \cdot \vec{1}$ ), and  $\vec{x}^{\#} \in \mathcal{H}$ . Therefore, when  $\ell$  is sufficiently large, we have:

$$\mathbf{A}^{+} \cdot \left( \vec{y}^{n} + \sqrt{\ell} \vec{\Delta} \right)^{\top} \leq \left( \vec{b}^{+} \right)^{\top}$$

This means that  $\vec{y}^n + \sqrt{\ell} \vec{\Delta} \in \mathcal{H}$  and therefore  $B^n \in \mathcal{H}$ .

- Second, for any  $\vec{\Delta} \in B_{\gamma} \vec{x}^{@}$  we have  $\vec{\Delta} \cdot \vec{1} = 0$ . Therefore,  $(\vec{y}^n + \sqrt{\ell}\vec{\Delta}) \cdot \vec{1} = n$ .
- Third, we prove  $B^n \subseteq \mathbb{R}^q_{\geq 0}$ . Recall that  $\vec{y}^n$  is O(1) away from  $\vec{x}^\ell$  (as constructed in Step 1.1). Each component of  $\vec{x}^\ell$  is  $\Omega(\ell)$ , because each component of  $\vec{x}^*$  is at least  $\epsilon > 0$ . Also note that for any  $\vec{\Delta} \in B_{\gamma} - \vec{x}^{(0)}$ ,  $|\vec{\Delta}|_{\infty} = O(1)$ . Therefore, each component of each vector in  $B^n$  is  $\Omega(\ell) - O(\sqrt{\ell}) - O(1)$ , which is strictly positive when  $\ell$  is sufficiently large.

Step 2. Define  $\vec{\pi} = (\pi_1, \ldots, \pi_n) \in \Pi^n$  s.t.  $\mathbb{E}(\vec{X}_{\vec{\pi}}) = \sum_{j=1}^n \pi_j$  is  $O(\sqrt{n})$  away from  $B^n$ in  $L_{\infty}$ . Because  $\vec{x}^* \in \mathcal{H}_{\leq 0} \cap CH(\Pi)$ , by Carathéodory's theorem for convex/conic hulls (see e.g., [36, p. 257]), we can write  $\vec{x}^*$  as the convex combination of  $1 \leq t \leq q$  distributions in  $\Pi$  regardless of the cardinality of  $\Pi$ , which can be infinity. Formally, let  $\vec{x}^* = \sum_{i=1}^t \alpha_i \pi_i^*$ , where for each  $i \leq t$ ,  $\alpha_i > 0$  and  $\pi_i^* \in \Pi$ , and  $\sum_{i=1}^t \alpha_i = 1$ . We note that  $\vec{x}^* \geq \epsilon \cdot \vec{1}$ , because  $\Pi$  is strictly positive (by  $\epsilon$ ). We now define  $\vec{\pi} \in \Pi^n$  and then prove that  $\vec{\pi} \cdot \vec{1}$  is  $O(\sqrt{n})$  away from  $\ell \vec{x}^* = \sum_{i=1}^t \ell \alpha_i \pi_i^*$  in  $L_{\infty}$ .

We now define  $\vec{\pi} \in \Pi^n$  and then prove that  $\vec{\pi} \cdot \vec{1}$  is  $O(\sqrt{n})$  away from  $\ell \vec{x}^* = \sum_{i=1}^t \ell \alpha_i \pi_i^*$  in  $L_{\infty}$ . Formally, for each  $i \leq t-1$ , let  $\vec{\pi}_i^{\ell}$  denote the vector of  $\beta_i = \lfloor \ell \alpha_i \rfloor$  copies of  $\pi_i^*$ . Let  $\vec{\pi}_k^{\ell}$  denote the vector of  $\beta_t = n - \sum_{i=1}^{t-1} \beta_i$  copies of  $\pi_t^*$ . It follows that for any  $i \leq t-1$ ,  $|\beta_i - \ell \alpha_i| \leq 1$ , and  $|\beta_t - \ell \alpha_t| \leq t + (\vec{y}^n \cdot \vec{1} - \ell \vec{x}^* \cdot \vec{1}) = O(\sqrt{\ell}) = O(\sqrt{n})$ , where the constant in  $O(\sqrt{n})$  depends on  $\mathcal{H}$  (because  $\vec{y}^n \cdot \vec{1} - \ell \vec{x}^* \cdot \vec{1}$  depends on  $\mathcal{H}$ ) but not on  $\Pi$  (because  $t \leq q$ ),  $\epsilon$ , or n.

Let  $\vec{\pi} = (\vec{\pi}_1^{\ell}, \dots, \vec{\pi}_t^{\ell})$ , or equivalently

$$\vec{\pi} = (\underbrace{\pi_1^*, \dots, \pi_1^*}_{\beta_1}, \underbrace{\pi_2^*, \dots, \pi_2^*}_{\beta_2}, \dots, \underbrace{\pi_t^*, \dots, \pi_t^*}_{\beta_t})$$

It follows that  $\mathbb{E}(\vec{X}_{\vec{\pi}}) = \vec{\pi} \cdot \vec{1} = \sum_{i=1}^{t} \beta_i \pi_i^*$ , which means that

$$|\mathbb{E}(\vec{X}_{\vec{\pi}}) - \ell \vec{x}^*|_{\infty} = \left| \sum_{i=1}^t (\beta_i - \ell \alpha_i) \pi_i^* \right|_{\infty} = O(\sqrt{n})$$

Recall that  $\vec{x}^{\ell} - \ell \vec{x}^* = \sqrt{\ell} \vec{x}^{@} + \vec{x}^{\#}$ . This means that  $\mathbb{E}(\vec{X}_{\vec{\pi}}) = \vec{\pi} \cdot \vec{1}$  is also  $O(\sqrt{n})$  away from  $\vec{x}^{\ell}$  in  $L_{\infty}$ . Recall that any vector in  $B^n$  is  $O(\sqrt{\ell})$  away from  $\vec{y}^n$ , which is O(1) away from  $\vec{x}^{\ell}$ . Therefore,  $\vec{\pi} \cdot \vec{1}$  is  $O(\sqrt{n})$  away from any vector in  $B^n$  in  $L_{\infty}$ , where the constant in the asymptotic bound depends on  $\mathcal{H}$  but not on  $\Pi$ ,  $\epsilon$ , or n.

Step 3.  $B^n$  contains  $\Omega\left(\sqrt{n}^{\dim(\mathcal{H}_{\leq 0})-1}\right)$ -many integer vectors. Intuitively, this is true because  $B^n$  consists of enough vectors from an neighborhood of  $\vec{y}^n$  that looks like  $\mathcal{H}_{\leq 0}$  with the additional linear constraint  $\vec{x} \cdot \vec{1} = n$ . Therefore, the  $I_1$  components of vectors in  $B^n$  can be viewed as flexible variables, each of which can take any value in an  $\Omega(\sqrt{n})$  interval, which contains  $\Omega(\sqrt{n})$  integers. Once the  $I_1$  component of a vector in  $B^n$  is given, its  $I_0$  components are more or less determined (see, e.g., Claim 1 and its proof).

More precisely, we will enumerate  $\Omega\left(\sqrt{n}^{\dim(\mathcal{H}_{\leq 0})-1}\right)$  many integer vectors in  $B^n$  of the form

$$\vec{y}^n + \left( \mathbf{D} \cdot \left( \vec{\Delta}_{I_1}, 0 \right)^\top, \vec{\Delta}_{I_1} \right), \text{ where } \vec{\Delta}_{I_1} \in \mathbb{Z}_{\geq 0}^{I_1}$$

Recall that we have assumed w.l.o.g. that  $I_0 = \{1, \ldots, o+1\}$  and  $I_1 = \{o+1, \ldots, q\}$ . Let  $\rho$  be the least common multiple of the denominators of entries in **D**. For example,  $\rho = 1$  in Example 5 because all entries in **D** are integers.

Then, we enumerate  $\vec{\Delta}_{I_1} = (\Delta_{o+2}, \dots, \Delta_q) \in \mathbb{Z}_{\geq 0}^{I_1}$  that satisfies the following two conditions.

- Condition 1. For each  $o + 2 \le j \le q$ ,  $\rho$  divides  $\Delta_j$ , and
- Condition 2. For each  $o + 2 \le j \le q$ ,  $|\Delta_j| < \frac{\gamma}{2}\sqrt{n}$ . Recall that  $\gamma > 0$  is the constant used to define  $B_{\gamma}$ .

Condition 1 guarantees that  $\vec{\Delta}_{I_0} = \mathbf{D} \cdot \left(\vec{\Delta}_{I_1}, 0\right)^\top \in \mathbb{Z}^{I_0}$ . Condition 2 guarantees that  $\vec{y}^n + \vec{\Delta} \in B^n$ when  $\frac{\gamma}{2}\sqrt{n} < \gamma\sqrt{\ell}$ , which holds for any sufficiently large *n* because  $\lim_{n\to\infty} \frac{n}{\ell} = 1$ . Notice that the total number of combinations of  $\vec{\Delta}_{I_1}$ 's that satisfy both conditions is at least  $\left(\lfloor\frac{\gamma\sqrt{n}}{\rho}\rfloor\right)^{|I_1|}$ . When *n* is sufficiently large so that  $\frac{\gamma\sqrt{n}}{\rho} > 1$  and  $\frac{\gamma}{2}\sqrt{n} < \gamma\sqrt{\ell}$ , the number of integer vectors in  $B^n$  is  $\Omega\left(\sqrt{n}^{|I_1|}\right) = \Omega\left(\sqrt{n}^{\dim(\mathcal{H}_{\leq 0})-1}\right)$ , where the constant in the asymptotic lower bound depends on  $\mathcal{H}$  but not on  $\Pi$ ,  $\epsilon$ , or *n*. This proves Step 3.

Step 4. The probability for  $\vec{X}_{\vec{\pi}}$  to be any given vector in  $B^n$  is  $\Omega\left(n^{\frac{(1-q)}{2}}\right)$ . This follows after Step 2  $(\vec{\pi} \cdot \vec{1} \text{ is } O(\sqrt{n})$  away from  $B^n$ ) and Lemma 1 below, which extends the pointwise concentration bound for i.i.d. Poisson multinomial variables [66, Lemma 4 in the Appendix] to general Poisson multinomial variables that correspond to strictly positive (but not necessarily identical) distributions.

Lemma 1 (Point-wise concentration bound for Poisson multinomial variables). For any  $q \in \mathbb{N}$ , any  $\epsilon > 0$ , and any  $\alpha > 0$ , there exists  $C_{q,\epsilon,\alpha} > 0$  such that for any  $n \in \mathbb{N}$ , any (n,q)-Poisson

multinomial random variable  $\vec{X}_{\vec{\pi}}$  where  $\vec{\pi}$  is above  $\epsilon$ , and any integer vector  $\vec{x} \in \mathbb{Z}_{\geq 0}^q$  with  $\vec{x} \cdot 1 = n$ and  $|\vec{x} - \mathbb{E}(\vec{X}_{\vec{\pi}})|_{\infty} < \alpha \sqrt{n}$ , we have:

$$\Pr(\vec{X}_{\vec{\pi}} = \vec{x}) > C_{q,\epsilon,\alpha} \cdot n^{\frac{1-q}{2}}$$

*Proof.* The proof proceeds in three steps. In **Step (i)**, we prove that it suffices to prove the lemma for a special  $\vec{\pi}$ , where at most  $2^q$  types of distributions are used. This is achieved by analyzing the following linear program given  $\epsilon > 0$ ,  $\vec{x} \in \mathbb{Z}_{\geq 0}^q$ , and  $\vec{\mu} \in \mathbb{R}_{\geq 0}^q$  are given, and the variables are  $\vec{\pi} = (\pi_1, \ldots, \pi_n)$ , where each  $\pi_j$  is a distribution over [q] that is above  $\epsilon$ .

$$\min_{\vec{\pi}} \Pr(\vec{X}_{\vec{\pi}} = \vec{x})$$
s.t. 
$$\sum_{j=1}^{n} \pi_j = \vec{\mu}$$
for each  $j \le n, \pi_j \ge \epsilon \cdot \vec{1}$  and  $\pi_j \cdot \vec{1} = 1$ 
(15)

At a high level, (15) can be viewed as an extension of ideas and techniques for Poisson binomial variables developed by Hoeffding [30] to PMVs. In **Step (ii)**, we prove the lemma for any  $\vec{\pi}'$  that consists of a constant number of different distributes (each distribution may appear multiple times in  $\vec{\pi}'$ ). This can be viewed as an extension of the point-wise concentration bound for i.i.d. Poisson multinomial variables [66, Lemma 4 in the Appendix] to PMVs of constant number of different distributions. In **Step (iii)** we combine results in Step 1 and 2 to prove the lemma.

Step (i). For any  $B \subseteq [q]$ , let  $\vec{\pi}|_B$  denote the collection (equivalently, subvector) of distributions in  $\vec{\pi}$  whose *B*-components are exactly  $\epsilon$ .

**Example 13.** For example, let  $q = 5, \epsilon = 0.1, n = 4, \ \vec{\pi} = (\pi_1, \pi_2, \pi_3, \pi_4)$ , where

$\pi_1$	$\pi_1$	$\pi_3 = \pi_4 = \pi_{uni}$
(0.25, 0.2, 0.3, 0.15, 0.1)	(0.2, 0.4, 0.15, 0.15, 0.1)	(0.2, 0.2, 0.2, 0.2, 0.2)

We have  $\vec{\pi}|_{\emptyset} = (\pi_3, \pi_4), \ \vec{\pi}|_{\{5\}} = (\pi_1, \pi_2), \ and \ for \ any \ other \ B \subseteq [5], \ \vec{\pi}|_B = \emptyset.$ 

By definition,  $\vec{\pi}$  and  $\bigcup_{B \subseteq [q]} \vec{\pi}|_B$  contain the same (multi-)set of distributions. We have the following claim about the optimal solutions to (15).

**Claim 2.** (15) has an optimal solution  $\vec{\pi}^*$  where for each  $B \subseteq [q]$  with  $\vec{\pi}^*|_B \neq \emptyset$ , all distributions in  $\vec{\pi}^*|_B$  are the same.

*Proof.* Let  $\Pi_{\epsilon}$  denote the set of all distributions over [q] that are above  $\epsilon$ . It follows that  $\Pi_{\epsilon}$  is compact and  $\Pr(\vec{X}_{\vec{\pi}} = \vec{x})$  is continuous, because it can be viewed as a polynomial in  $\vec{\pi}$ . Therefore, due to the extreme value theorem, (15) has solutions. Let  $\vec{\pi}^*$  denote an arbitrary solution with the maximum total number of probabilities in distributions in  $\vec{\pi}^*$  that equal to  $\epsilon$ , that is,

$$\vec{\pi}^* = (\pi_1^*, \dots, \pi_n^*) \in \arg\max_{\vec{\pi} \text{ is a solution to } (15)} |\{j \le n, i \le q : \pi_j(i) = \epsilon\}|$$

Suppose for the sake of contradiction the claim is not true. Then, there exists  $B \subseteq [q]$  such that  $\pi^*|_B$  contains at least two different distributions. W.l.o.g. let  $B = \{q'+1, \ldots, q\}$  and let the two distributions be  $\pi_1^*$  and  $\pi_2^*$  such that for some  $q^* \leq q'$ , we have that for each  $1 \leq i \leq q^*$ ,  $\pi_1^*(i) \neq \pi_2^*(i)$  and for each  $q^* + 1 \leq i \leq q'$ ,  $\pi_1^*(i) = \pi_2^*(i)$ . It follows that  $q^* \geq 2$ .

Let  $\vec{\pi} = (\pi_1, \ldots, \pi_n) \in \Pi_{\epsilon}^n$  denote an arbitrary vector of n distributions in  $\Pi_{\epsilon}$  (i.e.,  $\vec{\pi}$  may not be a solution to (15)) such that for every  $1 \leq i \leq q^*$ ,  $\pi_1(i) \neq \pi_2(i)$  and for each  $q^* + 1 \leq i \leq q'$ ,  $\pi_1(i) = \pi_2(i)$ . For any  $\vec{\psi} = (\psi_1, \ldots, \psi_{q^*})$  such that  $\vec{\psi} \cdot \vec{1} = 0$  and  $|\vec{\psi}|_{\infty}$  is sufficiently small, we let  $\vec{\pi}_{\vec{\psi}}$  denote the vector of distributions that is obtained from  $\vec{\pi}$  by replacing  $\pi_1$  by  $\pi_1 + (\vec{\psi}, \vec{0})$  and replacing  $\pi_2$  by  $\pi_2 - (\vec{\psi}, \vec{0})$ .

**Example 14.** Continuing Example 13, we let  $B = \{5\}$ . Then, q' = 4,  $q^* = 3$ ,  $\psi = (\psi_1, \psi_2, \psi_3)$  and

$$\vec{\pi}_{\vec{\psi}} = ((0.25 + \psi_1, 0.2 + \psi_2, 0.3 + \psi_3, 0.15, 0.1), (0.2 - \psi_1, 0.4 - \psi_2, 0.15 - \psi_3, 0.15, 0.1), \pi_3, \pi_4)$$

For any  $\vec{x} \in \mathbb{Z}_{\geq 0}^{q}$  with  $\vec{x} \cdot \vec{1} = n$  and any  $\vec{\pi}$ , we will prove that  $\Pr\left(\vec{X}_{\vec{\pi}_{\vec{\psi}}} = \vec{x}\right) - \Pr\left(\vec{X}_{\vec{\pi}} = \vec{x}\right)$  can be calculated as follows.

$$\Pr\left(\vec{X}_{\vec{\pi}_{\vec{\psi}}} = \vec{x}\right) - \Pr\left(\vec{X}_{\vec{\pi}} = \vec{x}\right)$$

$$= \sum_{i \le q^*} F_{ii}(-\psi_i^2 + \psi_i(\pi_2(i) - \pi_1(i))) + \sum_{1 \le i < t \le q^*} F_{it}(-2\psi_i\psi_t + \psi_i(\pi_2(t) - \pi_1(t)) + \psi_t(\pi_2(i) - \pi_1(i)))$$
(16)

where for any  $1 \le i \le t \le q^*$ ,  $F_{it}$  is the probability for  $\vec{X}_{\{\pi_3,\ldots,\pi_n\}}$  to be the vector that is obtained from  $\vec{x}$  by subtracting 1 from the *i*-th element and the *t*-th element (and from *i*-th element twice if i = t). Note that some  $F_{it}$ 's can be 0.

**Example 15.** Continuing Example 14, we let  $\vec{x} = (2, 1, 1, 0, 0)$ . Recall that  $\pi_3$  and  $\pi_4$  are uniform distributions. We have  $F_{11} = \frac{1}{25}$ ,  $F_{12} = F_{13} = F_{23} = \frac{2}{25}$ , and  $F_{22} = F_{33} = 0$ . Then, (16) becomes:

$$\Pr\left(\vec{X}_{\vec{\pi}_{\vec{\psi}}} = \vec{x}\right) - \Pr\left(\vec{X}_{\vec{\pi}} = \vec{x}\right) = \frac{1}{25}(-\psi_1^2 - 0.05\psi_1) + \frac{2}{25}(-2\psi_1\psi_2 + 0.2\psi_1 - 0.05\psi_2) \\ + \frac{2}{25}(-2\psi_1\psi_3 - 0.15\psi_1 - 0.05\psi_3) + \frac{2}{25}(-2\psi_2\psi_3 - 0.15\psi_2 + 0.2\psi_3)$$

Formally, (16) is proved by applying the law of total probability to the histogram of  $(\pi_3, \ldots, \pi_n)$ . For any  $1 \le i < t \le q^*$ , let  $\vec{e}_{i,t} \in \{0,1\}^q$  denote the vector that takes 1 on the *i*-th component and the *t*-th component, and takes 0 on other components. For any  $1 \le i \le q^*$ , let  $\vec{e}_{ii} \in \{0,2\}^q$  denote the vector that takes 2 on the *i*-th component and takes 0 on other components. According to the law of total probability, we have:

$$\Pr\left(\vec{X}_{\vec{\pi}} = \vec{x}\right) = \sum_{1 \le i \le t \le q^*} \Pr\left(\vec{X}_{(\pi_3, \dots, \pi_n)} = \vec{x} - \vec{e}_{it}\right) \times \Pr\left(X_{(\pi_1, \pi_2)} = \vec{e}_{it} | \vec{X}_{(\pi_3, \dots, \pi_n)} = \vec{x} - \vec{e}_{it}\right)$$
$$= \sum_{1 \le i \le t \le q^*} F_{it} \times \Pr\left(X_{(\pi_1, \pi_2)} = \vec{e}_{it} | \vec{X}_{(\pi_3, \dots, \pi_n)} = \vec{x} - \vec{e}_{it}\right)$$
$$= \sum_{1 \le i \le t \le q^*} F_{it} \times \Pr\left(X_{(\pi_1, \pi_2)} = \vec{e}_{it}\right)$$

The last equation holds because the *n* random variables are independent. A similar formula can be obtained for  $\Pr\left(\vec{X}_{\vec{\pi}_{\vec{\psi}}} = \vec{x}\right)$ . That is,

$$\Pr\left(\vec{X}_{\vec{\pi}_{\vec{\psi}}} = \vec{x}\right) = \sum_{1 \le i \le t \le q^*} F_{it} \times \Pr\left(X_{(\pi_1 + (\vec{\psi}, \vec{0}), \pi_2 - (\vec{\psi}, \vec{0}))} = \vec{e}_{it}\right)$$

Therefore,

$$\Pr\left(\vec{X}_{\vec{\pi}_{\vec{\psi}}} = \vec{x}\right) - \Pr\left(\vec{X}_{\vec{\pi}} = \vec{x}\right) = \sum_{1 \le i \le t \le q^*} F_{it} \times \left[\Pr(X_{(\pi_1 + (\vec{\psi}, \vec{0}), \pi_2 - (\vec{\psi}, \vec{0}))} = \vec{e}_{it}) - \Pr(X_{(\pi_1, \pi_2)} = \vec{e}_{it})\right]$$
(17)

Next, we calculate  $\Pr\left(X_{(\pi_1+(\vec{\psi},\vec{0}),\pi_2-(\vec{\psi},\vec{0}))}=\vec{e}_{it}\right) - \Pr\left(X_{(\pi_1,\pi_2)}=\vec{e}_{it}\right)$  for i = t and i < t, respectively.

• When i = t, we have:

$$\Pr\left(X_{(\pi_1+(\vec{\psi},\vec{0}),\pi_2-(\vec{\psi},\vec{0}))} = \vec{e}_{ii}\right) - \Pr\left(\vec{X}_{(\pi_1,\pi_2)}\right)$$
$$= (\pi_1(i) + \psi_i)(\pi_2(i) - \psi_i) - \pi_1(i)\pi_2(i)$$
$$= -\psi_i^2 + \psi_i(\pi_2(i) - \pi_1(i))$$
(18)

• When  $1 \le i < t \le q^*$ , we have:

$$\Pr\left(X_{(\pi_1+(\vec{\psi},\vec{0}),\pi_2-(\vec{\psi},\vec{0}))} = \vec{e}_{ii}\right) - \Pr\left(\vec{X}_{(\pi_1,\pi_2)}\right)$$
  
=  $[(\pi_1(i) + \psi_i)(\pi_2(t) - \psi_t) + (\pi_1(t) + \psi_t)(\pi_2(i) - \psi_i)] - [\pi_1(i)\pi_2(t) + \pi_1(t)\pi_2(i)]$   
=  $-2\psi_i\psi_t + \psi_i(\pi_2(t) - \pi_1(t)) + \psi_t(\pi_2(i) - \pi_1(i))$  (19)

(16) follows after combining (17), (18), and (19).

For convenience, we rewrite (16) in matrix form. For any  $1 \leq i < t \leq q^*$ , let  $F_{ti} = F_{it}$  and let  $\mathbf{F} = (F_{it})_{q^* \times q^*}$  denote the  $q^* \times q^*$  symmetric matrix.  $\vec{\delta} = (\pi_2(1) - \pi_1(1), \ldots, \pi_2(q^*) - \pi_1(q^*))$ . According to the definition of  $q^*$ , no component of  $\vec{\delta}$  is 0. With the matrix notation, (16) becomes

$$\Pr\left(\vec{X}_{\vec{\pi}_{\vec{\psi}}} = \vec{x}\right) - \Pr\left(\vec{X}_{\vec{\pi}} = \vec{x}\right) = -\vec{\psi} \cdot \mathbf{F} \cdot \left(\vec{\psi}\right)^{\top} + \vec{\psi} \cdot \mathbf{F} \cdot \left(\vec{\delta}\right)^{\top}$$
(20)

Example 16. Continuing Example 15, we have  $\mathbf{F} = \begin{bmatrix} \frac{1}{25} & \frac{2}{25} & \frac{2}{25} \\ \frac{2}{25} & 0 & \frac{2}{25} \\ \frac{2}{25} & \frac{2}{25} & 0 \end{bmatrix}$  and  $\vec{\delta} = (-0.05, 0.2, -0.15).$ 

Let A denote the  $(q^*-1) \times q^*$  matrix  $\begin{bmatrix} 1 & & -1 \\ & \ddots & & \vdots \\ & & 1 & -1 \end{bmatrix}$  and  $\vec{\psi'} = (\psi_1, \dots, \psi_{q^*-1})$ . Recall that

 $\vec{\psi} \cdot \vec{1} = 0$ , we have  $\psi_{q^*} = -\psi_1 - \cdots - \psi_{q^*-1}$ , which means that  $\vec{\psi} = \vec{\psi'} \cdot A$ . Because  $\pi_1$  and  $\pi_2$  are probability distributions, we have  $\vec{\delta} \cdot \vec{1} = \vec{0}$ , which means that  $\delta_{q^*} = -\delta_1 - \cdots - \delta_{q^*-1}$ . Therefore, let  $\vec{\delta'} = (\pi_2(1) - \pi_1(1), \dots, \pi_2(q^*-1) - \pi_1(q^*-1))$ , we have  $\vec{\delta} = \vec{\delta'} \cdot A$ . Let  $\mathbf{F'} = A \cdot \mathbf{F} \cdot (A)^{\top}$ . Then, (20) becomes:

$$\Pr\left(\vec{X}_{\vec{\pi}_{\vec{\psi}}} = \vec{x}\right) - \Pr\left(\vec{X}_{\vec{\pi}} = \vec{x}\right) = -\vec{\psi}' \cdot \mathbf{F}' \cdot \left(\vec{\psi}'\right)^{\top} + \vec{\psi}' \cdot \mathbf{F}' \cdot \left(\vec{\delta}'\right)^{\top}$$
(21)

Example 17. Continuing Example 16, we have  $A = \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \end{bmatrix}$ ,  $\mathbf{F}' = \begin{bmatrix} -\frac{3}{25} & -\frac{2}{25} \\ -\frac{2}{25} & -\frac{4}{25} \end{bmatrix}$  and  $\vec{\delta}' = (-0.05, 0.2)$ . Therefore,  $\mathbf{F}' \cdot \left(\vec{\delta}'\right)^{\top} = \begin{bmatrix} -\frac{1}{100} \\ -\frac{7}{250} \end{bmatrix}$  and (21) becomes:

$$\Pr\left(\vec{X}_{\vec{\pi}_{\vec{\psi}}} = \vec{x}\right) - \Pr\left(\vec{X}_{\vec{\pi}} = \vec{x}\right) = -[\psi_1, \psi_2] \cdot \begin{bmatrix} -\frac{3}{25} & -\frac{2}{25} \\ -\frac{2}{25} & -\frac{4}{25} \end{bmatrix} \cdot \begin{bmatrix} \psi_1 \\ \psi_2 \end{bmatrix} + [\psi_1, \psi_2] \cdot \begin{bmatrix} -\frac{1}{100} \\ -\frac{7}{250} \end{bmatrix}$$

Notice that  $\mathbf{F}'$  depends on both  $\vec{x}$  and  $\vec{\pi}$ . Next, we consider the case for  $\vec{\pi}^*$ , which we recall is an optimal solution to (15) and  $q^*$  is defined based on  $\vec{\pi}^*$ . We will prove that  $\mathbf{F}' \cdot \left(\vec{\delta}'\right)^\top = \left(\vec{0}\right)^\top$ , where  $\mathbf{F}'$  is the matrix corresponding to  $\vec{x}$  and  $\vec{\pi}^*$ . Suppose for the sake of contradiction that this is not true. W.l.o.g. suppose the first component of  $\mathbf{F}' \cdot \left(\vec{\delta}'\right)^\top$  is non-zero. Then, by letting  $\psi_2 = \cdots = \psi_{q^*-1} = 0$ , (21) becomes  $A\psi_1^2 + B\psi_1$  for some constants A and B with  $B \neq 0$ , which means that there exists  $\psi_1 \neq 0$  such that (21) is strictly less than zero. This contradicts the assumption that  $\vec{\pi}^*$  is an optimal solution to (15). The following example shows how to choose  $\vec{\psi}$ when  $\mathbf{F}' \cdot \left(\vec{\delta}'\right)^\top \neq \left(\vec{0}\right)^\top$ , to obtain another feasible solution with smaller objective value. Notice that in this example  $\vec{x}$  is not an optimal solution to (15).

**Example 18.** Continuing Example 17, notice that the first component of  $\mathbf{F}' \cdot \left(\vec{\delta}'\right)^{\top}$  is non-zero. Therefore, by letting  $\psi_2 = 0$ , we have

$$\Pr\left(\vec{X}_{\vec{\pi}_{\vec{\psi}}} = \vec{x}\right) - \Pr\left(\vec{X}_{\vec{\pi}} = \vec{x}\right) = \frac{3}{25}\psi_1^2 - \frac{1}{100}\psi_1$$

Let  $\psi_1 = 0.05$ , which means that  $\psi_3 = -0.05$ . It is not hard to verify that  $\vec{\pi}_{\vec{\psi}}$  is a feasible solution to (15) with a smaller objective value.

Therefore, for any  $\gamma \in \mathbb{R}$ , if we let  $\vec{\psi} = \gamma \vec{\delta}$ , then (21) becomes zero, which means that if  $\vec{\pi}^*_{\gamma \vec{\delta}}$  is strictly positive (by  $\epsilon$ ), then it is also an optimal solution to (15). Recall that all components of  $\vec{\delta}$  are non-zero. Therefore, we can start from  $\gamma = 0$  and gradually increase the value of  $\gamma$  until any of the first  $q^*$  components in  $\pi^*_1 + (\gamma \vec{\delta}, \vec{0})$  or in  $\pi^*_2 - (\gamma \vec{\delta}, \vec{0})$  becomes  $\epsilon$ . Then, it is not hard to verify that  $\vec{\pi}^*_{\gamma \vec{\delta}}$  is an optimal solution to (15) with strictly more probabilities that equal to  $\epsilon$ . This contradicts the assumption that  $\vec{\pi}^*$  contains maximum number of probabilities that equal to  $\epsilon$  among optimal solutions to (15), and therefore concludes the proof of Claim 2.

Step (ii). We prove the following special case of the lemma.

**Claim 3.** For any  $\epsilon > 0, \alpha > 0, q \in \mathbb{N}$ , and  $Q \in \mathbb{N}$ , there exist constants  $C_{q,\epsilon,\alpha,Q}$  and N such that for any set  $\Pi_Q$  of Q distributions over [q] that are strictly positive by  $\epsilon$ , any  $\vec{\pi}' \in \Pi^n_Q$ , and any integer vector  $\vec{x} \in \mathbb{Z}_{\geq 0}^q$  with  $\vec{x} \cdot 1 = n$  and  $|\vec{x} - \mathbb{E}(\vec{X}_{\vec{\pi}'})|_{\infty} < \alpha \sqrt{n}$ , we have

$$\Pr\left(\vec{X}_{\vec{\pi}'} = \vec{x}\right) > C_{q,\epsilon,\alpha,Q} \cdot n^{\frac{1-q}{2}}$$

Proof. Because  $|\Pi_Q| = Q$ , there exists  $\pi^* \in \Pi_Q$  that appears in  $\vec{\pi}'$  for at least  $\lceil \frac{n}{Q} \rceil$  times. Let  $n' = \lceil \frac{n}{Q} \rceil$  and let  $\vec{X}_1$  denote the (n', q)-PMV that corresponds to  $(\underbrace{\pi^*, \ldots, \pi^*}_{n'})$ . Let  $\vec{X}_2$  denote the (n-n', q)-PMV that corresponds to the remaining distributions in  $\vec{\pi}'$ . Recall that each distribution in  $\vec{\pi}'$  is strictly positive by  $\epsilon$ . By Hoeffding's inequality, let  $\alpha' = \frac{(1-\epsilon)^2}{2}\log(4q)$ , for each  $i \leq q$ , we have

$$\Pr\left(|[\vec{X}_2]_i - [\mathbb{E}(\vec{X}_2)]_i| > \alpha'\sqrt{n-n'}\right) \le \frac{1}{2q}$$

Therefore, by the union bound, we have:

$$\Pr\left(|\vec{X}_2 - \mathbb{E}(\vec{X}_2)|_{\infty} > \alpha'\sqrt{n-n'}\right) \le \frac{1}{2}$$

Notice that  $\vec{X}_1$  and  $\vec{X}_2$  are independent. Now we can calculate  $\Pr\left(\vec{X}_{\vec{\pi}'} = \vec{x}\right)$  by the law of total probability, by enumerating the target values for  $\vec{X}_2$ , denoted by  $\vec{x}_2$ , as follows.

$$\Pr\left(\vec{X}_{\vec{\pi}'} = \vec{x}\right)$$

$$= \sum_{\vec{x}_2 \in \mathbb{Z}_{\geq 0}^q : \vec{x}_2 \cdot \vec{1} = n - n'} \Pr\left(\vec{X}_1 = \vec{x} - \vec{x}_2 | \vec{X}_2 = \vec{x}_2\right) \cdot \Pr\left(\vec{X}_2 = \vec{x}_2\right)$$
(The law of total probability)
(22)

$$= \sum_{\vec{x}_2 \in \mathbb{Z}_{\geq 0}^q: \vec{x}_2 \cdot \vec{1} = n - n'} \Pr\left(\vec{X}_1 = \vec{x} - \vec{x}_2\right) \cdot \Pr\left(\vec{X}_2 = \vec{x}_2\right) \qquad (\vec{X}_1 \text{ and } \vec{X}_2 \text{ are independent})$$

$$\geq \sum_{\vec{x}_2 \in \mathbb{Z}_{\geq 0}^q: \vec{x}_2 \cdot \vec{1} = n - n' \text{ and } |\vec{x}_2 - \mathbb{E}(\vec{X}_2)|_{\infty} \leq \alpha' \sqrt{n - n'}} \Pr\left(\vec{X}_1 = \vec{x} - \vec{x}_2\right) \cdot \Pr\left(\vec{X}_2 = \vec{x}_2\right)$$

$$= \sum_{\vec{x}_2 \in \mathbb{Z}_{\geq 0}^q: \vec{x}_2 \cdot \vec{1} = n - n' \text{ and } |\vec{x}_2 - \mathbb{E}(\vec{X}_2)|_{\infty} \leq \alpha' \sqrt{n - n'}} \Omega\left(n^{\frac{1 - q}{2}}\right) \cdot \Pr\left(\vec{X}_2 = \vec{x}_2\right)$$

$$= \Omega\left(n^{\frac{1 - q}{2}}\right) \cdot \Pr\left(|\vec{X}_2 - \mathbb{E}(\vec{X}_2)|_{\infty} \leq \alpha' \sqrt{n - n'}\right) = \Omega\left(n^{\frac{1 - q}{2}}\right)$$
(23)

(23) follows after applying the point-wise concentration bound for i.i.d. Poisson multinomial variables [66, Lemma 4 in the Appendix] to  $\vec{X}_1$  in the following way. Recall that  $\mathbb{E}(\vec{X}_{\vec{\pi}}) = \mathbb{E}(\vec{X}_1) + \mathbb{E}(\vec{X}_2), |\vec{x} - \mathbb{E}(\vec{X}_{\vec{\pi}})|_{\infty} < \alpha \sqrt{n}$ , and  $|\vec{x}_2 - \mathbb{E}(\vec{X}_2)|_{\infty} \le \alpha' \sqrt{n - n'}$ . We have:

$$|\vec{x}_1 - \mathbb{E}(\vec{X}_1)|_{\infty} = |\vec{x} - \mathbb{E}(\vec{X}_1) - (\vec{x}_2 - \mathbb{E}(\vec{X}_2))|_{\infty} = \alpha\sqrt{n} + \le \alpha'\sqrt{n - n'} \le (\alpha Q + \alpha'(Q - 1))\sqrt{n'}$$

This completes the proof of Claim 3.

**Step (iii).** By Claim 2, there exists a vector  $\vec{\pi}^*$  of *n* distributions over [q], each of which is above  $\epsilon$ , such that (i)  $\Pr(\vec{X}_{\vec{\pi}} = \vec{x}) \ge \Pr(\vec{X}_{\vec{\pi}^*} = \vec{x})$  and  $\mathbb{E}(\vec{X}_{\vec{\pi}}) = \mathbb{E}(\vec{X}_{\vec{\pi}^*})$ , and (ii)  $\vec{\pi}^*$  consists of no more than  $Q = 2^q$  different distributions that correspond to different subsets of [q]. Note that  $\vec{\pi}^*$  may not be in  $\Pi^n$ . Lemma 1 follows after applying Claim 3 with  $\vec{\pi}' = \vec{\pi}^*$ . 

Step 5. Final calculations. Finally, combining Step 3 and 4, we have:

$$\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}\right) \ge \Pr\left(\vec{X}_{\vec{\pi}} \in B^n\right) \ge |B^n| \times \min_{\vec{x} \in B^n} \Pr\left(\vec{X}_{\vec{\pi}} = \vec{x}\right)$$
$$\ge \Omega\left(n^{\frac{\dim(\mathcal{H}_{\leq 0}) - 1}{2}}\right) \times \Omega\left(n^{\frac{(1-q)}{2}}\right) = \Omega\left(n^{-\frac{o}{2}}\right),$$

where the constants in the asymptotic bounds depend on  $\mathcal{H}$  (and therefore q) and  $\epsilon$  but not on other parts of  $\Pi$  or n. We require n to be sufficiently large to guarantee the existence of  $\vec{x}^*$  and  $\vec{x}^{@}$ , and  $\frac{\gamma\sqrt{n}}{\rho} > 1$  and  $\frac{\gamma}{2}\sqrt{n} < \gamma\sqrt{\ell}$ . This proves the polynomial lower bound on Sup.

**Proof of the exponential bounds on Inf.** The proof is similar to the proof of the exponential case in Theorem 1. The lower bound is straightforward and the upper bound is proved by choosing an arbitrary  $\pi \in CH(\Pi)$  such that  $\pi \notin \mathcal{H}_{\leq 0}$ , and define  $\vec{\pi}$  in a similar way as in the proof of the polynomial lower bound of the Sup part.

**Proof of the polynomial upper bound on Inf.** Notice that the condition for this case implies that the polynomial case for Sup holds, which means that there exists  $\vec{\pi} \in \Pi^n$  such that  $\Pr(\vec{X}_{\vec{\pi}} \in \mathcal{H}) = \Theta\left(n^{\frac{\dim(\mathcal{H}_{\leq 0})-q}{2}}\right)$ . This immediately implies an  $O\left(n^{\frac{\dim(\mathcal{H}_{\leq 0})-q}{2}}\right)$  upper bound on Inf.

**Proof of the polynomial low bound on Inf.** The proof is similar to the proof of the polynomial lower bound on the Sup part in Theorem 1, except that when defining  $\vec{y}^n$ , we let  $\vec{x}^* = \sum_{j=1}^n \pi_j / n \in CH(\Pi) \cap \mathcal{H}_{\leq 0}$ . More precisely, we have the following lemma that will be used in the proof of other propositions in this paper. Notice that the constant  $C_{\Pi,\mathcal{H}}$  in the lemma only depends on  $\Pi$  (therefore q and  $\epsilon$ ) and  $\mathcal{H}$  but not on  $\vec{\pi}$ .

**Lemma 2.** For any  $q \in \mathbb{N}$ , any closed and strictly positive  $\Pi$  over [q], and any polyhedron  $\mathcal{H}$ with integer matrix  $\mathbf{A}$ , there exists  $C_{\Pi,\mathcal{H}} > 0$  such that for any  $n \in \mathbb{N}$  with  $\mathcal{H}_n^{\mathbb{Z}} \neq \emptyset$  and any  $\vec{\pi} = (\pi_1, \ldots, \pi_n) \in \Pi^n$  with  $\sum_{j=1}^n \pi_j \in \mathcal{H}_{\leq 0}$ ,

$$\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}\right) \ge C_{\Pi, \mathcal{H}} \cdot n^{-\frac{\dim(\mathcal{H}_{\leqslant 0})}{2}}$$

Proof. For any  $n \in \mathbb{N}$  that is sufficiently large, we define a non-negative integer vector  $\vec{y}^n \in \mathcal{H}_n^{\mathbb{Z}}$  and a neighborhood  $B^n \subseteq \mathcal{H}_n$  of  $\vec{y}^n$  that is similar to Step 1 of proof for the polynomial lower bound in the Sup case of Theorem 1. The only difference is that we let  $\vec{x}^* = \sum_{j=1}^n \pi_j / n = \mathbb{E}(\vec{X}_{\vec{\pi}}) / n \in$  $CH(\Pi) \cap \mathcal{H}_{\leq 0}$ . The rest of the proof is the same as Step 3, 4, and 5 of the proof for the polynomial lower bound in the Sup case of Theorem 1.

To see that the constant  $C_{\Pi,\mathcal{H}}$  does not depend on  $\vec{\pi}$ , we first notice that there exists a constant  $\alpha_{\Pi,\mathcal{H}}$  such that  $\mathbb{E}(\vec{X}_{\vec{\pi}}) = n\vec{x}^*$  is no more than  $\alpha_{\Pi,\mathcal{H}}(\sqrt{n})$  away from  $\vec{x}^\ell = \ell \vec{x}^* + \sqrt{\ell} \vec{x}^{@} + \vec{x}^{\#}$ , which is O(1) away from  $\vec{y}^n$ . Recall that  $\vec{y}^n$  is the "center" of  $B^n$ , whose "radius" only depends on  $\ell$  and  $B_{\gamma}$  but not on  $\vec{x}^*$ . Therefore,  $\vec{y}^n$  (which is O(1) close to  $\vec{x}^\ell$  and the constant only depends on  $\mathbf{A}$  but not on  $\vec{x}^*$ ) is no more than  $\alpha'_{\Pi,\mathcal{H}}\sqrt{n}$  away from any vector in  $B^n$  for some  $\alpha'_{\Pi,\mathcal{H}} > 0$ . Finally, we note that the constant in the point-wise concentration bound (Lemma 1) does not depend on  $\vec{\pi}$ .

This completes the proof of Theorem 1.

### A.2 Proof of Theorem 2

**Theorem 2.** (Smoothed Likelihood of PMV-in-C). Given any  $q, I \in \mathbb{N}$ , any closed and strictly positive  $\Pi$  over [q], and any  $C = \bigcup_{i \in I} \mathcal{H}_i$  characterized by integer matrices, for any  $n \in \mathbb{N}$ ,

$$\sup_{\vec{\pi}\in\Pi^n} \Pr\left(\vec{X}_{\vec{\pi}}\in\mathcal{C}\right) = \begin{cases} 0 & \text{if } \alpha_n = -\infty \\ \exp(-\Theta(n)) & \text{if } -\infty < \alpha_n < 0 \\ \Theta\left(n^{\frac{\alpha_n - q}{2}}\right) & \text{otherwise } (i.e. \ \alpha_n > 0) \end{cases},$$
$$\inf_{\vec{\pi}\in\Pi^n} \Pr\left(\vec{X}_{\vec{\pi}}\in\mathcal{C}\right) = \begin{cases} 0 & \text{if } \beta_n = -\infty \\ \exp(-\Theta(n)) & \text{if } -\infty < \beta_n < 0 \\ \Theta\left(n^{\frac{\beta_n - q}{2}}\right) & \text{otherwise } (i.e. \ \beta_n > 0) \end{cases}.$$

*Proof.* For convenience, we first recall Inequality (6) in the main text.

$$\max_{i \leq I} \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i\right) \leq \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{C}\right) \leq \sum_{i \leq I} \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i\right)$$
(6)

The theorem is proved by combining (6) and applications of Theorem 1 to  $\Pi$  and  $\mathcal{H}_i$ . We first introduce some notation. Let  $\mathcal{C}_n^{\mathbb{Z}}$  denote the set of all non-negative integer vectors in  $\mathcal{C}$  whose  $L_1$  norm is n. That is,

$$\mathcal{C}_n^{\mathbb{Z}} = \bigcup_{i \leq I} \mathcal{H}_{i,n}^{\mathbb{Z}}$$

Given a distribution  $\pi$  over [q], let  $\mathcal{I}_{\mathcal{C},n}^{\pi} \subseteq \{1, \ldots, I\}$  denote the set of indices i in  $\mathcal{H}_i$  such that the weight on  $(\pi, \mathcal{H}_i)$  in the activation graph  $\mathcal{G}_{\Pi,\mathcal{C},n}$  is positive at n. Equivalently,  $\mathcal{I}_{\mathcal{C},n}^{\pi}$  can be defined as follows.

$$\mathcal{I}_{\mathcal{C},n}^{\pi} = \{ i \leq I : \mathcal{H}_{i,n}^{\mathbb{Z}} \neq \emptyset \text{ and } \pi \in \mathcal{H}_{i,\leqslant 0} \}$$

For example,  $\mathcal{I}_{\mathcal{C},n}^{\pi} = \{2\}$  in Figure 3.

Next, we let  $\Pi_{\mathcal{C},n} \subseteq \operatorname{CH}(\Pi)$  denote the distributions  $\pi$  in  $\operatorname{CH}(\Pi)$  such that  $\mathcal{I}_{\mathcal{C},n}^{\pi} \neq \emptyset$ , or equivalently,  $\dim_{\mathcal{C},n}^{\max}(\pi) > 0$ . Namely,

$$\Pi_{\mathcal{C},n} = \{ \pi \in \operatorname{CH}(\Pi) : \mathcal{I}_{\mathcal{C},n}^{\pi} \neq \emptyset \}$$

We start with the proof for the sup part.

**Proof for the sup part.** We prove the sup part by discussing the three cases (0, exponential, and polynomial) as follows.

- The 0 case of sup. This case is straightforward because  $\alpha_n = -\infty$  means that no polyhedron in  $\mathcal{C}$  is active at n, which means that  $\mathcal{C}$  does not contain any non-negative integer vector whose size is n, while any outcome of the PMV  $\vec{X}_{\vec{\pi}}$  is a non-negative integer vector whose size is n.
- The exponential case of sup. We recall that in this case  $C_n^{\mathbb{Z}} \neq \emptyset$  and  $\Pi_{\mathcal{C},n} = \emptyset$ . To prove the exponential lower bound of sup, it suffices to prove that there exists  $\vec{\pi} \in \Pi^n$  such that  $\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{C}\right) = \exp(-O(n))$ . Because  $C_n^{\mathbb{Z}} = \bigcup_{i \leq I} \mathcal{H}_{i,n}^{\mathbb{Z}} \neq \emptyset$ , there exists  $i^* \leq I$  such that  $\mathcal{H}_{i^*}$  is active at n, i.e.,  $\mathcal{H}_{i^*,n}^{\mathbb{Z}} \neq \emptyset$ . The exponential lower bound follows after applying the lower bound of (6) to an arbitrary  $\vec{\pi} \in \Pi^n$ :

$$\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{C}\right) \ge \max_{i \le I} \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i\right) \ge \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_{i^*}\right) \ge \epsilon^n = \exp(-O(n))$$

To prove the **exponential upper bound of sup**, it suffices to prove that for every  $\vec{\pi} \in \Pi^n$ , we have  $\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{C}\right) = \exp(-\Omega(n))$ . This will be proved by combining the upper bound in (6) and the 0 or exponential upper bounds on the sup part of Theorem 1 when applied to  $\Pi$  and each  $\mathcal{H}_i$ .

More precisely, because  $\Pi_{\mathcal{C},n} = \emptyset$ , for every  $\pi \in \Pi^n$ , we have  $\mathcal{I}_{\mathcal{C},n}^{\pi} = \emptyset$ . This mean that for every  $\pi \in \operatorname{CH}(\Pi)$  and every  $i \leq I$ , either  $\mathcal{H}_{i,n}^{\mathbb{Z}} = \emptyset$  or  $\pi \notin \mathcal{H}_{i,\leq 0}$  (or both hold). If the former holds, then the 0 case of Theorem 1 can be applied to  $\Pi$  and  $\mathcal{H}_i$ , which means that for any  $\vec{\pi} \in \Pi^n$ , we have  $\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i\right) = 0$ . If the former does not hold and the latter holds, then the exponential case of Theorem 1 applies to  $\Pi$  and  $\mathcal{H}_i$ , which means that for every  $\vec{\pi} \in \Pi^n$ , we have  $\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i\right) = \exp(-\Omega(n))$ . Notice that I is a constant. Therefore, following the upper bound of (6), for every  $\vec{\pi} \in \Pi^n$ , we have:

$$\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{C}\right) \leq \sum_{i \leq I} \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i\right) \leq I \cdot \exp(-\Omega(n)) = \exp(-\Omega(n))$$

• The polynomial bounds of sup. We recall that in this case  $C_n^{\mathbb{Z}} \neq \emptyset$  and  $\Pi_{\mathcal{C},n} \neq \emptyset$ . Like the exponential case, the proof is done by combining (6) and the applications of Theorem 1 to  $\Pi$  and each  $\mathcal{H}_i$ . To prove the polynomial lower bound of sup, it suffices to prove that there exists  $\vec{\pi} \in \Pi^n$  such that  $\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{C}\right) = \Omega(n^{\frac{\alpha_n-q}{2}})$ . Because  $\Pi_{\mathcal{C},n} \neq \emptyset$ , we let  $(\pi^*, i^*)$ denote the pair that achieves  $\alpha_n$ . More precisely, let

$$(\pi^*, i^*) = \arg \max_{\pi \in \Pi_{\mathcal{C}, n}} \max_{i \in \mathcal{I}_{\mathcal{C}, n}^{\pi}} \dim(\mathcal{H}_{i, \leq 0})$$

Following the definition of  $(\pi^*, i^*)$ , we have  $\mathcal{H}_{i^*,n}^{\mathbb{Z}} \neq \emptyset$ ,  $\pi^* \in \mathcal{H}_{i^*,\leqslant 0}$  (which means that  $\mathcal{H}_{i^*,\leqslant 0} \cap \operatorname{CH}(\Pi) \neq \emptyset$ ), and  $\dim(\mathcal{H}_{i^*}) = \alpha_n$ . This means that the polynomial case of the sup part of Theorem 1 holds when the theorem is applied to  $\Pi$  and  $\mathcal{H}_{i^*}$ . Therefore, there exists  $\vec{\pi} \in \Pi^n$  such that  $\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_{i^*}\right) = \Omega(n^{\frac{\dim(\mathcal{H}_{i^*})-q}{2}})$ . Notice that we cannot immediately let  $\vec{\pi} = (\pi^*, \ldots, \pi^*)$  because it is possible that  $\pi^* \notin \Pi$ . It follows after the lower bound of (6) that

$$\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{C}\right) \ge \max_{i \le I} \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i\right) \ge \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_{i^*}\right) = \Omega\left(n^{\frac{\dim(\mathcal{H}_{i^*}) - q}{2}}\right) = \Omega\left(n^{\frac{\alpha_{n-q}}{2}}\right)$$

This proves the polynomial lower bound of sup.

To prove the **polynomial upper bound of sup**, it suffices to prove that for every  $\vec{\pi} \in \Pi^n$ , we have  $\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{C}\right) = O\left(n^{\frac{\alpha_n-q}{2}}\right)$ . This will be proved by combining the upper bound in (6) and the (0, exponential, or polynomial) upper bounds on the sup part of Theorem 1 when the theorem is applied to  $\Pi$  and every  $\mathcal{H}_i$ . More precisely, for each  $i \leq I$ , applying the sup part of Theorem 1 to  $\Pi$  and  $\mathcal{H}_i$  results in the following three cases.

- The 0 case. If  $\mathcal{H}_{i,n}^{\mathbb{Z}} = \emptyset$ , then for every  $\vec{\pi} \in \Pi^n$ , we have  $\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i\right) = 0$ .
- The exponential case. If  $\mathcal{H}_{i,n}^{\mathbb{Z}} \neq \emptyset$  and  $\mathcal{H}_{i,\leqslant 0} \cap \operatorname{CH}(\Pi) = 0$ , then for every  $\vec{\pi} \in \Pi^n$ , we have  $\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i\right) = \exp(-\Omega(n))$ .
- The polynomial case. Otherwise  $(\mathcal{H}_{i,n}^{\mathbb{Z}} \neq \emptyset \text{ and } \mathcal{H}_{i,\leqslant 0} \cap \operatorname{CH}(\Pi) \neq 0)$ , for every  $\vec{\pi} \in \Pi^n$ , we have  $\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i\right) = O\left(n^{\frac{\dim(\mathcal{H}_{i,\leqslant 0})-q}{2}}\right)$ .

In the polynomial case, for any  $\pi \in \mathcal{H}_{i,\leq 0} \cap \operatorname{CH}(\Pi)$ , we have  $i \in \mathcal{I}_{\mathcal{C},n}^{\pi} \neq \emptyset$ , which means that  $\pi \in \Pi_{\mathcal{C},n}$ . This means that  $\dim(\mathcal{H}_{i,\leq 0}) \leq \alpha_n$ . Therefore, in all three cases we have  $\operatorname{Pr}\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i\right) = O\left(\frac{\alpha_n - q}{2}\right)$ . Again, recall that I is a constant. Therefore, following the upper bound in (6), for every  $\vec{\pi} \in \Pi^n$ , we have:

$$\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{C}\right) \leq \sum_{i \leq I} \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i\right) \leq I \cdot O\left(n^{\frac{\alpha_n - q}{2}}\right) = O\left(n^{\frac{\alpha_n - q}{2}}\right)$$

This proves the polynomial upper bound of sup.

**Proof for the inf part.** We now turn to the inf part by discussing the three cases (0, exponential, and polynomial) as follows. At a high level, the proofs for the upper (respectively, lower) bounds of inf are similar to the proofs for the lower (respectively, upper) bounds of sup. We include the formal proof below for completeness.

• The 0 case of inf. Like the 0 case of the sup part, this case is straightforward.

• The exponential case of inf. We recall that in this case  $C_n^{\mathbb{Z}} \neq \emptyset$  and  $\Pi_{\mathcal{C},n} \neq \operatorname{CH}(\Pi)$ . To prove the exponential lower bound of inf, it suffices to prove that for every  $\vec{\pi} \in \Pi^n$ , we have  $\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{C}\right) = \exp(-O(n))$ . Because  $C_n^{\mathbb{Z}} = \bigcup_{i \leq I} \mathcal{H}_{i,n}^{\mathbb{Z}} \neq \emptyset$ , by definition there exists  $i^* \leq I$  such that  $\mathcal{H}_{i^*,n}^{\mathbb{Z}} \neq \emptyset$ . The exponential lower bound follows after the lower bound of (6):

$$\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{C}\right) \ge \max_{i \le I} \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i\right) \ge \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_{i^*}\right) \ge \epsilon^n = \exp(-O(n))$$

To prove the **exponential upper bound of inf**, it suffices to prove that there exists  $\vec{\pi} \in \Pi^n$ such that  $\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{C}\right) = \exp(-\Omega(n))$ . Because  $\Pi_{\mathcal{C},n} \neq \operatorname{CH}(\Pi)$ , there exists  $\pi \in \operatorname{CH}(\Pi)$  such that  $\pi \notin \Pi_{\mathcal{C},n}$ , which means that  $\mathcal{I}_{\mathcal{C},n}^{\pi} = \emptyset$ . Let  $\vec{\pi} = (\pi_1, \ldots, \pi_n) \in \Pi$  denote an arbitrary vector such that  $\sum_{j=1}^n \pi_j$  is  $\Theta(1)$  away from  $n \cdot \pi$ . Because  $\mathcal{I}_{\mathcal{C},n}^{\pi} = \emptyset$ , for all  $i \leq I$ , either  $\mathcal{H}_{i,n}^{\mathbb{Z}} = \emptyset$  or  $\pi \notin \mathcal{H}_{i,\leq 0}$  (or both hold). If the former holds, then by applying the 0 upper bound on the sup part of Theorem 1 to  $\Pi$  and  $\mathcal{H}_i$ , we have  $\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i\right) = 0$ . If the former does not hold and the latter holds, then following Hoeffding's inequality and the union bound (applied to each of the q coordinate of  $\vec{X}_{\vec{\pi}}$ ), we have  $\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i\right) = \exp(-\Omega(n))$ . Therefore, following the upper bound of (6), we have:

$$\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{C}\right) \leq \sum_{i \leq I} \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i\right) \leq I \cdot \exp(-\Omega(n)) = \exp(-\Omega(n))$$

This proves the exponential upper bound of inf.

• The polynomial bounds of inf. We recall that in this case  $C_n^{\mathbb{Z}} \neq \emptyset$  and  $\Pi_{\mathcal{C},n} = \operatorname{CH}(\Pi)$ . Like in the exponential case, the proof is done by combining (6) and the applications of Theorem 1 to  $\Pi$  and each  $\mathcal{H}_i$ . To prove the **polynomial lower bound of inf**, it suffices to prove that for every  $\vec{\pi} = (\pi_1, \ldots, \pi_n) \in \Pi^n$ , we have  $\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{C}\right) = \Omega(n^{\frac{\beta_n - q}{2}})$ . Let  $\pi^* = \frac{1}{n} \sum_{i=1}^n \pi_i \in \operatorname{CH}(\Pi)$ . Because  $\Pi_{\mathcal{C},n} = \operatorname{CH}(\Pi)$ , we have  $\pi^* \in \Pi_{\mathcal{C},n}$ , which means that  $\mathcal{I}_{\Pi,n}^{\pi^*} \neq \emptyset$ . Therefore, there exists  $i^* \in \mathcal{I}_{\Pi,n}^{\pi^*}$  such that  $\dim(\mathcal{H}_{i^*,\leq 0}) \geq \beta_n$ . Additionally, recall that for all  $i \in \mathcal{I}_{\Pi,n}^{\pi^*}$  we have  $\mathcal{H}_{i^*,n}^{\mathbb{Z}} \neq \emptyset$  and  $\pi^* \in \mathcal{H}_{i^*,\leq 0}$ . Therefore, Lemma 2 can be applied to  $\vec{\pi}$  and  $\mathcal{H}_{i^*,n}^{\mathbb{Z}}$ , giving us  $\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_{i^*}\right) = \Omega\left(n^{\frac{\beta_n - q}{2}}\right)$ . It follows after the lower bound of (6) that:

$$\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{C}\right) \ge \max_{i \le I} \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_{i}\right) \ge \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_{i^{*}}\right) \ge \Omega\left(n^{\frac{\dim(\mathcal{H}_{i^{*}})-q}{2}}\right) = \Omega\left(n^{\frac{\beta_{n}-q}{2}}\right)$$

This proves the polynomial lower bound of inf.

To prove the **polynomial upper bound of inf**, it suffices to prove that there exists  $\vec{\pi} \in \Pi^n$  such that  $\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{C}\right) = O(n^{\frac{\beta_n - q}{2}})$ . Because  $\Pi_{\mathcal{C},n} = \operatorname{CH}(\Pi)$ , we define  $(\pi^*, i^*)$  to be the pair that achieves  $\beta_n$ . More precisely, let

$$(\pi^*, i^*) = \arg\min_{\pi \in \Pi_{\mathcal{C}, n}} \max_{i \in \mathcal{I}_{\mathcal{C}, n}^{\pi}} \dim(\mathcal{H}_{i, \leq 0})$$

It follows that  $\mathcal{H}_{i^*,n}^{\mathbb{Z}} \neq \emptyset$ ,  $\pi^* \in \mathcal{H}_{i^*,\leq 0}$  (which means that  $\mathcal{H}_{i^*,\leq 0} \cap \operatorname{CH}(\Pi) \neq \emptyset$ ), and dim $(\mathcal{H}_{i^*}) = \beta_n$ . Let  $\vec{\pi} = (\pi_1, \ldots, \pi_n) \in \Pi^n$  denote an arbitrary vector such that  $\sum_{j=1}^n \pi_n$  is  $\Theta(1)$  from  $n\pi^*$ . The polynomial upper bound of inf is proved by combining the upper bound in (6) and the (0, exponential, or polynomial) upper bounds on the sup part of Theorem 1 when the theorem is applied to  $\Pi$  and every  $\mathcal{H}_i$ . More precisely, for each  $i \leq I$ , applying the sup part of Theorem 1 to  $\vec{\pi}$  and  $\mathcal{H}_i$  results in the following three cases.

- The 0 case. If  $\mathcal{H}_{i,n}^{\mathbb{Z}} = \emptyset$ , then  $\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i\right) = 0$ .
- The exponential case. If  $\mathcal{H}_{i,n}^{\mathbb{Z}} \neq \emptyset$  and  $\pi^* \notin \mathcal{H}_{i,\leqslant 0}$ , then  $\mathbb{E}(\vec{X}_{\vec{\pi}})$  is  $\Theta(n)$  away from  $\mathcal{H}_{i,\leqslant 0}$ . Following Hoeffding's inequality and the union bound (applied to each of the q coordinate of  $\vec{X}_{\vec{\pi}}$ ), we have  $\Pr\left(\vec{X}_{\vec{\pi}^*} \in \mathcal{H}_i\right) = \exp(-\Omega(n))$ .
- The polynomial case. Otherwise  $(\mathcal{H}_{i,n}^{\mathbb{Z}} \neq \emptyset \text{ and } \pi^* \in \mathcal{H}_{i,\leq 0}, \text{ which means that } i \in \mathcal{I}_{\Pi,n}^{\pi^*})$ , we have

$$\Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i\right) = O\left(n^{\frac{\dim(\mathcal{H}_{i,\leq 0})-q}{2}}\right) \le O\left(n^{\frac{\beta_n-q}{2}}\right),$$

because

$$\dim(\mathcal{H}_i) \le \dim(\mathcal{H}_{i^*}) = \beta_n$$

Therefore, in all three cases above, we have  $\Pr\left(\vec{X}_{\vec{\pi}^*} \in \mathcal{H}_i\right) = O(\frac{\beta_n - q}{2})$ . Again, recall that I is a constant. Therefore, following the upper bound in (6), we have:

$$\Pr\left(\vec{X}_{\vec{\pi}^*} \in \mathcal{C}\right) \le \sum_{i \le I} \Pr\left(\vec{X}_{\vec{\pi}} \in \mathcal{H}_i\right) \le I \cdot O\left(n^{\frac{\beta_n - q}{2}}\right) = O\left(n^{\frac{\alpha_n - q}{2}}\right)$$

This proves the polynomial upper bound of inf.

# **B** Appendix for Section 4.1: Integer Positional Scoring Rules

## B.1 Proof of Theorem 3

**Theorem 3.** (Smoothed likelihood of ties: positional scoring rules). Let  $\mathcal{M} = (\Theta, \mathcal{L}(\mathcal{A}), \Pi)$  be a strictly positive and closed single-agent preference model and let  $\vec{s}$  be an integer scoring vector. For any  $2 \le k \le m$  and any  $n \in \mathbb{N}$ ,

$$\widetilde{Tie}_{\Pi}^{\max}(r_{\vec{s}},k,n) = \begin{cases} 0 & \text{if } \forall P \in \mathcal{L}(\mathcal{A})^{n}, |r_{\vec{s}}(P)| \neq k \\ \exp(-\Theta(n)) & \text{otherwise, if } \forall \pi \in CH(\Pi), |r_{\vec{s}}(\pi)| < k \\ \Theta(n^{-\frac{k-1}{2}}) & \text{otherwise} \end{cases}$$
$$\widetilde{Tie}_{\Pi}^{\min}(r_{\vec{s}},k,n) = \begin{cases} 0 & \text{if } \forall P \in \mathcal{L}(\mathcal{A})^{n}, |r_{\vec{s}}(P)| \neq k \\ \exp(-\Theta(n)) & \text{otherwise, if } \exists \pi \in CH(\Pi), |r_{\vec{s}}(\pi)| < k \\ \Theta(n^{-\frac{k-1}{2}}) & \text{otherwise} \end{cases}$$

*Proof.* The theorem is proved by modeling the set of profiles with k winners as the union of constantly many polyhedra, then applying Theorem 2. More precisely, we have the following three steps. In Step 1, for each potential winner set  $T \subseteq \mathcal{A}$ , we define a polyhedron  $\mathcal{H}^{\vec{s},T}$  that characterizes profiles whose winners are T. In Step 2, we prove properties of  $\mathcal{H}^{\vec{s},T}$ , and in particular,  $\dim(\mathcal{H}^{\vec{s},T}_{\leq 0}) = m! - |T| + 1$ . In Step 3 we formally apply Theorem 2 to  $\mathcal{C} = \bigcup_{T \subseteq \mathcal{A}: |T| = k} \mathcal{H}^{\vec{s},T}$ .

Step 1: Define  $\mathcal{H}^{\vec{s},T}$ . For any  $T \subseteq \mathcal{A}, \mathcal{H}^{\vec{s},T}$  consists of (i) equations that represent the scores of alternatives in T being equal, and (ii) inequalities that represent the score of any alternative in T being strictly larger than the score of any alternative in  $\mathcal{A} \setminus T$ . Formally, we first define a set of constraints to model the score difference between two alternatives.

**Definition 15** (Score difference vector). For any scoring vector  $\vec{s} = (s_1, \ldots, s_m)$  and any pair of different alternatives a, b, let  $Score_{a,b}^{\vec{s}}$  denote the m!-dimensional vector indexed by rankings in  $\mathcal{L}(\mathcal{A})$ : for any  $R \in \mathcal{L}(\mathcal{A})$ , the R-element of  $Score_{a,b}^{\vec{s}}$  is  $s_{j_1} - s_{j_2}$ , where  $j_1$  and  $j_2$  are the ranks of a and b in R, respectively.

Let  $\vec{x}_{\mathcal{A}} = (x_R : R \in \mathcal{L}(\mathcal{A}))$  denote the vector of m! variables, each of which represents the multiplicity of a linear order in a profile. Therefore,  $\operatorname{Score}_{a,b}^{\vec{s}} \cdot \vec{x}_{\mathcal{A}}$  represents the score difference between a and b in the profile whose histogram is  $\vec{x}_{\mathcal{A}}$ .

For any  $T \subseteq \mathcal{A}$ , we define the polyhedron  $\mathcal{H}^{\vec{s},T}$  as follows.

**Definition 16.** For any integer scoring vector  $\vec{s}$  and any  $T \subseteq A$ , we let  $\mathbf{E}^{\vec{s},T}$  denote the matrix whose row vectors are  $\{Score_{a,b}^{\vec{s}} : a \in T, b \in T, a \neq b\}$ . Let  $\mathbf{S}^{\vec{s},T}$  denote the matrix whose row vectors are  $\{Score_{a,b}^{\vec{s}} : a \notin T, b \in T\}$ . Let  $\mathbf{A}^{\vec{s},T} = \begin{bmatrix} \mathbf{E}^{\vec{s},T} \\ \mathbf{S}^{\vec{s},T} \end{bmatrix}$ ,  $\vec{b} = (\vec{0}, -\vec{1})$ , and let  $\mathcal{H}^{\vec{s},T}$  denote the corresponding polyhedron.

Step 2: Prove properties about  $\mathcal{H}^{\vec{s},T}$ . We prove the following claim about  $\mathcal{H}^{\vec{s},T}$ .

**Claim 4.** For any integer scoring vector  $\vec{s}$  and any  $T \subseteq A$ , we have:

- (i) for any integral profile P,  $Hist(P) \in \mathcal{H}^{\vec{s},T}$  if and only if  $r_{\vec{s}}(P) = T$ ;
- (ii) for any  $\vec{x} \in \mathbb{R}^q$ ,  $\pi \in \mathcal{H}_{\leq 0}^{\vec{s},T}$  if and only if  $T \subseteq r_{\vec{s}}(\vec{x})$ ;
- (*iii*) dim $(\mathcal{H}_{\leq 0}^{\vec{s},T}) = m! |T| + 1.$

*Proof.* (i) and (ii) follow after Definition 16. To prove (iii), we first note that  $\dim(\mathcal{H}_{\leq 0}^{\vec{s},T}) = m! - m!$ Rank( $\mathbf{A}^{=}$ ), where  $\mathbf{A}^{=}$  is the essential equalities of  $\mathbf{A}^{\vec{s},T}$ , according to equation (9) on page 100 in [57]. We note that  $\mathbf{A}^{=} = \mathbf{E}^{\vec{s},T}$ , because by definition  $\mathbf{E}^{\vec{s},T} \subseteq \mathbf{A}^{=}$ . To prove that no row in  $\mathbf{S}^{\vec{s},T}$ is in  $\mathbf{A}^{=}$ , it suffices to show that there exists  $\vec{x} \in \mathcal{H}_{\leq 0}^{\vec{s},T}$  such that  $\mathbf{S}^{\vec{s},T} \cdot (\vec{x})^{\top} < (\vec{0})^{\top}$ . This is proved by constructing a profile P such that  $r_{\vec{s}}(P) = T$  and then let  $\vec{x} = \text{Hist}(P)$ . We first define two cyclic permutations:  $\sigma_1$  over T and  $\sigma_2$  over  $\mathcal{A} \setminus T$ . More precisely, let

$$\sigma_1 = 1 \rightarrow 2 \rightarrow \cdots \rightarrow |T| \rightarrow 1 \text{ and } \sigma_2 = |T| + 1 \rightarrow |T| + 2 \rightarrow \cdots \rightarrow m \rightarrow |T| + 1$$

Then we let

$$P = \{\sigma_1^i(\sigma_2^j(1 \succ 2 \succ \dots \succ m)) : i \le |T|, j \le m - |T|\}$$

It is not hard to see that  $r_{\vec{s}}(P) = T$  and consequently  $\mathbf{E}^{\vec{s},T} \supseteq \mathbf{A}^=$ , which means that  $\mathbf{E}^{\vec{s},T} = \mathbf{A}^=$ . Therefore, it suffices to prove that  $\operatorname{Rank}(\mathbf{E}^{\vec{s},T}) = |T| - 1$ . We first prove that  $\operatorname{Rank}(\mathbf{E}^{\vec{s},T}) \leq T$ |T| - 1. W.l.o.g. let  $T = \{1, \ldots, k\}$ , where k = |T|. It is not hard to verify that all rows of  $\mathbf{E}^{\vec{s},T}$  can be represented by linear combinations of |T| - 1 rows  $\begin{bmatrix} \operatorname{Score}_{1,2} \\ \vdots \\ \operatorname{Score}_{1,k} \end{bmatrix}$ . This means that

 $\operatorname{Rank}(\mathbf{E}^{\vec{s},T}) \le |T| - 1.$ 

Next, we prove  $\operatorname{Rank}(\mathbf{E}^{\vec{s},T}) \geq |T| - 1$  by contradiction. Suppose for the sake of contradiction that  $\operatorname{Rank}(\mathbf{E}^{\vec{s},T}) \leq |T| - 2$ . Then, there exists  $\ell \leq k - 2$  rows of  $\mathbf{E}^{\vec{s},T}$ , denoted by  $\{\operatorname{Score}_{a_i,b_i}^{\vec{s}} : i \leq \ell\}$ , such that any  $\operatorname{Score}_{a,b}^{\vec{s}}$  is a linear combination of them. Let G denote the unweighted undirected graph over  $\{1, \ldots, k\}$  with the following  $\ell$  edges:  $\{\{a_i, b_i\} : i \leq \ell\}$ . Because G contains k-2edges and k nodes, G is not connected. W.l.o.g. suppose there is no edge in G between  $\{1, \ldots, t\}$ and  $\{t+1,\ldots,k\}$  for some  $1 \le t \le k-1$ . We now construct a profile P to show that  $\mathrm{Score}_{1,k}^{\vec{s}}$ is not a linear combination of  $\{\text{Score}_{a_i,b_i}^{\vec{s}} : i \leq \ell\}$ . We first define a linear order R and two cyclic permutations.

$$R = [1 \succ \ldots \succ t \succ k + 1 \succ \cdots \succ m \succ t + 1 \succ \cdots \succ k]$$

Let  $\sigma_1 = 1 \rightarrow \cdots \rightarrow t \rightarrow 1$  denote the cyclic permutation among  $\{1, \ldots, t\}$  and let  $\sigma_2 = t + 1 \rightarrow t$  $\dots \rightarrow k \rightarrow t+1$  denote the cyclic permutation among  $\{t+1,\dots,k\}$ . Let

$$P = \{\sigma_1^i(\sigma_2^j(R)) : 1 \le i \le t, 1 \le j \le k-t\}$$

It is not hard to verify that in P, the total scores of  $\{1, \ldots, t\}$  are the same, the scores of  $\{t + t\}$  $1, \ldots, k$  are the same, and the former is strictly larger than the latter. Therefore, for all  $i \leq \ell$ ,  $\operatorname{Score}_{a_i,b_i}^{\vec{s}} \cdot \operatorname{Hist}(P) = 0$ , but  $\operatorname{Score}_{1,k}^{\vec{s}} \cdot \operatorname{Hist}(P) > 0$ , which means that  $\operatorname{Score}_{1,k}^{\vec{s}}$  is not a linear combination of  $\{\text{Score}_{a_i,b_i}^{\vec{s}}: i \leq \ell\}$ , which is a contradiction. This means that  $\text{Rank}(\mathbf{E}^{\vec{s},T}) \geq |T| - 1$ . 

This completes the proof of Claim 4.

Step 3: Apply Theorem 2. Let  $\mathcal{C} = \bigcup_{T \subseteq \mathcal{A}: |T|=k} \mathcal{H}^{\vec{s},T}$ . It follows that for any profile P,  $|r_{\vec{s}}(P)| = k$  if and only if  $\text{Hist}(P) \in \mathcal{C}$ . Therefore, for any n and any  $\vec{\pi} \in \Pi^n$ , we have:

$$\Pr_{P \sim \vec{\pi}}(|r_{\vec{s}}(P)| = k) = \Pr(\vec{X}_{\vec{\pi}} \in \mathcal{C})$$

Recall that for all  $T_k \subseteq \mathcal{A}$  with  $|T_k| = k$ , from Claim 4 (iii) we have  $\dim(\mathcal{H}^{\vec{s},T_k}) = m! - |T_k| + 1 =$ m! - k + 1. This means that  $\alpha_n = \beta_n = m! - k + 1$  when they are non-negative (which holds for certain n).

Therefore, to prove Theorem 3, it suffices to prove that the conditions for the 0, exponential, and polynomial cases in Theorem 3 are equivalent to the conditions for the 0, exponential, and polynomial cases in Theorem 2 (applied to  $\mathcal{C}$  and  $\Pi$ ), respectively. The 0 case is straightforward. Recall from Claim 4(iii) that  $\dim(\mathcal{H}_{\leq 0}^{\vec{s},T}) = m! - |T| + 1$ . Therefore, to apply Theorem 2 to obtain Theorem 3, it suffices to prove that for any  $\pi \in CH(\Pi)$ ,  $|r_{\vec{s}}(\pi)| \geq k$  if and only if  $\pi \in \mathcal{C}_{\leq 0}$  (which is equivalent to  $\alpha_n \neq -\infty$ ). The "if" direction holds because if  $\pi \in \mathcal{C}_{\leq 0}$ , then there exists  $T \subseteq \mathcal{A}$  with |T| = k such that  $\pi \in \mathcal{H}_{\leq 0}^{\vec{s},T}$ , which means that  $T \subseteq r_{\vec{s}}(\pi)$  by Claim 4(ii). Therefore,  $|r_{\vec{s}}(\pi)| \ge k|$ . The "only if" direction holds because if  $|r_{\vec{s}}(\pi)| \geq k|$ , then let  $T \subseteq r_{\vec{s}}(\pi)$  denote an arbitrary set with |T| = k. It follows from Claim 4(ii) that  $\pi \in \mathcal{H}_{\leq 0}^{\vec{s},T}$ , which implies that  $\pi \in \mathcal{C}_{\leq 0}$ . 

#### $\mathbf{C}$ Appendix for Section 4.2: EO-Based Rules

#### C.1 Proof of Proposition 1

**Proposition 1.** For any 
$$\mathcal{A}$$
 and any  $n \ge m^4$ ,  $\{\mathrm{EO}(P) : P \in \mathcal{L}(\mathcal{A})^n\} = \begin{cases} \mathcal{O}_{\mathcal{A}} & \text{if } 2 \mid n \\ \mathcal{O}'_{\mathcal{A}} & \text{if } 2 \nmid n \end{cases}$ .

*Proof.* Let  $EO_n = \{EO(P) : P \in \mathcal{L}(\mathcal{A})^n\}$ . We first prove the  $2 \mid n$  case. By definition  $EO_n \subseteq \mathcal{O}_{\mathcal{A}}$ . We prove  $\mathcal{O}_{\mathcal{A}} \subseteq EO_n$  by explicitly constructing an *n*-profile *P* such that EO(P) = O for any  $O \in \mathcal{O}_{\mathcal{A}}$ . Let  $O = T_1 \triangleright \cdots \triangleright T_t \triangleright T_0 \triangleright T_{t+1} \triangleright \cdots \triangleright T_{2t}$  denote the tier representation of O. We define a weighted directed graph  $G_O$  such that

$$w_{G_O}(e) = \begin{cases} 2(t+1-i) & \text{if } e \in T_i \text{ for some } 1 \le i \le t \\ -2(t+1-i) & \text{if } \bar{e} \in T_i \text{ for some } 1 \le i \le t \\ 0 & \text{if } e \in T_0 \end{cases}$$

By McGarvey's theorem [42], there exists a profile  $P_O$  of no more than  $m(m+1)t < m^4$  votes such that WMG( $P_O$ ) =  $G_O$ . Let R be an arbitrary linear order and let  $\overline{R}$  denote its reverse order. Let  $P_2 = \{R, \overline{R}\}$ . It follows that WMG( $P_2$ ) is the empty graph. Let  $P = P_O + \frac{n-|P_O|}{2} \times P_2$ . It follows that |P| = n and WMG(P) = WMG( $P_O$ ) =  $G_O$ , which means that EO(P) = O. This means that  $\mathcal{O}_A \subseteq EO_n$ . Therefore, the lemma holds for  $2 \mid n$ .

Next, we prove the  $2 \nmid n$  case. For any *n*-profile *P*, the middle tier of EO(*P*) is empty because the weight on any edge must be odd. This means that EO<sub>n</sub>  $\subseteq \mathcal{O}'_{\mathcal{A}}$ . Like the  $2 \mid n$  case, we prove  $\mathcal{O}'_{\mathcal{A}} \subseteq \text{EO}_n$  by explicitly constructing an *n*-profile *P* such that EO(*P*) = *O* for any  $O \in \mathcal{O}'_{\mathcal{A}}$ . More precisely, let  $O = T_1 \triangleright \cdots \triangleright T_t \triangleright T_0 \triangleright T_{t+1} \triangleright \cdots \triangleright T_{2t}$  denote the tier representation of *O*, where  $T_0 = \emptyset$ . We define a weighted directed graph  $G_O$  such that

$$w_{G_O}(e) = \begin{cases} 2(t-i)+1 & \text{if } e \in T_i \text{ for some } 1 \le i \le t \\ -2(t-i)-1 & \text{if } \bar{e} \in T_i \text{ for some } 1 \le i \le t \end{cases}$$

By McGarvey's theorem [42], there exists a profile  $P_O$  of no more than  $m(m+1)t < m^4$  votes such that WMG( $P_O$ ) =  $G_O$ . Let  $P = P_O + \frac{n - |P_O|}{2} \times P_2$ . It follows that |P| = n and WMG(P) = WMG( $P_O$ ) =  $G_O$ , which means that EO(P) = O. This means that  $\mathcal{O}'_{\mathcal{A}} \subseteq EO_n$ . Therefore, the lemma holds for  $2 \nmid n$ .

## C.2 Formal Definition of Edge-Order-Based Rules (Section 4.2)

**Definition 17.** Given an edge-order-based rule r, any  $2 \le k \le m$ , any  $n \in \mathbb{N}$ , and any distribution  $\pi$  over  $\mathcal{L}(\mathcal{A})$ , let

$$\mathcal{O}_{r,k,n}^{\pi} = \{ O \in \mathcal{O}_{\mathcal{A}}^{n} : |r(O)| = k \text{ and } O \text{ refines } EO(\pi) \}$$

For any set of distributions  $\Pi$  over  $\mathcal{L}(\mathcal{A})$ , let  $\mathcal{O}_{r,k,n}^{\Pi} = \bigcup_{\pi \in CH(\Pi)} \mathcal{O}_{r,k,n}^{\pi}$ . When  $\mathcal{O}_{r,k,n}^{\Pi} \neq \emptyset$ , we let  $\ell_{\min} = \min_{O \in \mathcal{O}_{r,k,n}^{\Pi}} Ties(O)$ . When  $\mathcal{O}_{r,k,n}^{\pi} \neq \emptyset$  for all  $\pi \in CH(\Pi)$ , we let  $\ell_{mm} = \max_{\pi \in CH(\Pi)} \min_{O \in \mathcal{O}_{r,k,n}^{\pi}} Ties(O)$ .

## C.3 Proof of Theorem 4

**Theorem 4. (Smoothed likelihood of ties: edge-order-based rules).** Let  $\mathcal{M} = (\Theta, \mathcal{L}(\mathcal{A}), \Pi)$ be a strictly positive and closed single-agent preference model and let r be an edge-order-based rule. For any  $2 \le k \le m$  and any  $n \in \mathbb{N}$ ,

$$\begin{split} \widetilde{Tie}_{\Pi}^{\max}(r,k,n) &= \begin{cases} 0 & if \,\forall O \in \mathcal{O}_{\mathcal{A}}^{n}, |r(O)| \neq k \\ \exp(-\Theta(n)) & otherwise \, if \, \mathcal{O}_{r,k,n}^{\Pi} = \emptyset \\ \Theta\left(n^{-\frac{\ell_{\min}}{2}}\right) & otherwise \end{cases} \\ \widetilde{Tie}_{\Pi}^{\min}(r,k,n) &= \begin{cases} 0 & if \,\forall O \in \mathcal{O}_{\mathcal{A}}^{n}, |r(O)| \neq k \\ \exp(-\Theta(n)) & otherwise \, if \,\exists \pi \in CH(\Pi) \, s.t. \, \mathcal{O}_{r,k,n}^{\pi} = \emptyset \\ \Theta\left(n^{-\frac{\ell_{\min}}{2}}\right) & otherwise \end{cases}$$

*Proof.* Like in the proof of Theorem 3, the theorem is proved by modeling the set of profiles with k winners as the union of constantly many polyhedra, then applying Theorem 2. More precisely, we have the following three steps. In Step 1, for each palindromic order O, we define a polyhedron  $\mathcal{H}^O$  that characterizes the profiles whose palindromic orders are O. In Step 2, we prove properties about  $\mathcal{H}^O$ , in particular dim $(\mathcal{H}^O_{\leq 0}) = m! - \text{Ties}(O)$ . In Step 3 we formally apply Theorem 2 to  $\mathcal{C} = \bigcup_{O \in \mathcal{O}^A_{*}: |r(O)| = k} \mathcal{H}^O$ .

We first recall the definition of pairwise difference vector in [66].

**Definition 18** (Pairwise difference vector [66]). For any pair of different alternatives a, b, let  $Pair_{a,b}$  denote the m!-dimensional vector indexed by rankings in  $\mathcal{L}(\mathcal{A})$ : for any  $R \in \mathcal{L}(\mathcal{A})$ , the R-element of  $Pair_{a,b}$  is 1 if  $a \succ_R b$ ; otherwise it is -1.

Step 1: Define  $\mathcal{H}^{O}$ . For any palindromic order  $O \in \mathcal{O}_{\mathcal{A}}$ , we define a polyhedron  $\mathcal{H}^{O}$  that represents profiles whose edge order is O.

**Definition 19.** For any  $O \in \mathcal{O}_{\mathcal{A}}$ , we let  $\mathbf{E}^{O}$  denote the matrix whose row vectors are  $\{Pair_{a,b} - Pair_{c,d} : (a,b) \equiv_{O} (c,d)\}$ ; we let  $\mathbf{S}^{O}$  denote the matrix whose row vectors are  $\{Pair_{c,d} - Pair_{a,b} : (a,b) \triangleright_{O} (c,d)\}$ . Let  $\mathbf{A}^{O} = \begin{bmatrix} \mathbf{E}^{O} \\ \mathbf{S}^{O} \end{bmatrix}$ ,  $\vec{b} = (\vec{0}, -\vec{1})$ , and let  $\mathcal{H}^{O}$  denote the corresponding polyhedron.

 $\mathbf{E}^O$  contains redundant rows and is defined as in Definition 19 for notational convenience. For any  $1 \leq i \leq t$ , rows in  $\mathbf{E}^O$  that correspond to  $T_i$  are the same as the rows in  $\mathbf{E}^O$  that correspond to  $T_{2t+1-i}$ .

Step 2: Prove properties about  $\mathcal{H}^O$ . We have the following claim about  $\mathcal{H}^O$ .

**Claim 5.** For any profile P and any  $O \in \mathcal{O}_A$ , we have:

- (i)  $Hist(P) \in \mathcal{H}^O$  if and only if EO(P) = O.
- (ii)  $Hist(P) \in \mathcal{H}_{\leq 0}^O$  if and only if O refines EO(P).
- (iii)  $\dim(\mathcal{H}^O_{\leq 0}) = m! Ties(O).$

*Proof.* Part (i) and (ii) follow after the definition of  $\mathcal{H}^O$  and  $\mathcal{H}^O_{\leq 0}$ , respectively. For part (iii), let  $O = T_1 \triangleright \cdots \triangleright T_t \triangleright T_0 \triangleright T_{t+1} \triangleright \cdots \triangleright T_{2t}$  denote the tier representation of O. Let  $\mathbf{A}^=$  denote the essential equalities of  $\mathbf{A}^O$ . Again, it is not hard to verify that  $\mathbf{A}^= = \mathbf{E}^O$  due to McGarvey's theorem [42]. Therefore, it suffices to prove Rank( $\mathbf{E}^O$ ) = Ties(O).

We first note that  $\operatorname{Rank}(\mathbf{E}^O) \leq \operatorname{Ties}(O)$ , because for any  $0 \leq i \leq t$  such that  $T_i = \{e_1, \ldots, e_\ell\}$ and any profile P, the following  $\ell - 1$  equations

$$(\operatorname{Pair}_{e_1} - \operatorname{Pair}_{e_2}) \cdot \operatorname{Hist}(P) = 0, (\operatorname{Pair}_{e_1} - \operatorname{Pair}_{e_3}) \cdot \operatorname{Hist}(P) = 0, \dots, (\operatorname{Pair}_{e_1} - \operatorname{Pair}_{e_\ell}) \cdot \operatorname{Hist}(P) = 0$$

imply that all edges in  $T_i$  have the same weights in WMG(P). To prove that Rank( $\mathbf{E}^O$ )  $\geq$  Ties(O), suppose for the sake of contradiction that Rank( $\mathbf{A}^O$ ) < Ties(O). Then, there exists a tier  $T_i = \{e_1, \ldots, e_\ell\}$  where  $0 \leq i \leq t$  and no more than  $s < \ell - 1$  rows in  $\mathbf{E}^O$ , denoted by  $\mathbf{E}' = \left(\operatorname{Pair}_{e_{i_1}} - \operatorname{Pair}_{e_{j_1}}, \ldots, \operatorname{Pair}_{i_s} - \operatorname{Pair}_{j_s}\right)^{\top}$ , whose linear combinations include other rows that correspond to  $T_i$ . Let  $\mathbf{E}_i$  denote the rows of  $\mathbf{E}^O$  that correspond to  $T_i$ . By McGarvey's theorem [42], there exists a profile P such that the  $w_P(e_{i_1}) = w_P(e_{j_1}), \ldots, w_P(e_{i_s}) = w_P(e_{j_s})$ , but not all  $e_1, \ldots, e_\ell$  have the same weights in WMG(P). This means that  $\mathbf{E}' \cdot (\operatorname{Hist}(P))^{\top} = \left(\vec{0}\right)^{\top}$  but  $\mathbf{E}_i \cdot (\operatorname{Hist}(P))^\top \neq \left(\vec{0}\right)^\top$ , which means that not all rows in  $\mathbf{E}_i$  are linear combinations of rows in  $\mathbf{E}'$ , which is a contradiction. This proves that  $\operatorname{Rank}(\mathbf{E}^O) = \operatorname{Ties}(O)$ , which implies part (iii) of the claim.

**Step 3: Apply Theorem 2.** Let  $\mathcal{C} = \bigcup_{O \in \mathcal{O}_{\mathcal{A}}^n : |r(O)|=k} \mathcal{H}^O$ . It follows that for any profile P, |r(P)| = k if and only if  $\operatorname{Hist}(P) \in \mathcal{C}$ . Therefore, for any n and any  $\vec{\pi} \in \Pi^n$ , we have:

$$\Pr_{P \sim \vec{\pi}}(|r(P)| = k) = \Pr(\vec{X}_{\vec{\pi}} \in \mathcal{C})$$

Recall that for all  $O \in \mathcal{O}_{\mathcal{A}}^n$  with |r(O)| = k, from Claim 5 (iii) we have  $\dim(\mathcal{H}^O) = m! - \operatorname{Ties}(P) + 1$ . This means that  $\alpha_n = m! - \ell_{\min}$  and  $\beta_n = m! - \ell_{\min}$  when they are non-negative (which holds for certain n).

Therefore, to prove Theorem 4, it suffices to prove that the conditions for the 0, exponential, and polynomial cases in Theorem 4 are equivalent to the conditions for the 0, exponential, and polynomial cases in Theorem 2 (applied to C and  $\Pi$ ), respectively. This follows a similar reasoning as in Step 3 of the proof of Theorem 3 combined with Claim 5 (ii). This completes the proof for Theorem 4.

### C.4 Proof of Proposition 2

**Proposition 2.** (Max smoothed likelihood of ties: Copeland<sub> $\alpha$ </sub>). Let  $\mathcal{M} = (\Theta, \mathcal{L}(\mathcal{A}), \Pi)$  be a strictly positive and closed single-agent preference model with  $\pi_{uni} \in CH(\Pi)$ . Let  $l_{\alpha} = \min\{t \in \mathbb{N} : t\alpha \in \mathbb{Z}\}$ . For any  $2 \leq k \leq m$  and any  $n \in \mathbb{N}$ ,

$$\widetilde{Tie}_{\Pi}^{\max}(Cd_{\alpha},k,n) = \begin{cases} 0 & \text{if } 2 \nmid n, 2 \mid k, \text{ and } \begin{cases} k = m, \text{ or } \\ k = m - 1 \\ (1)k = m, \text{ or } \\ (2)k = m - 1 \text{ and } \alpha \geq \frac{1}{2}, \text{ or } \\ (3)k = m - 1 \text{ and } k \leq l_{\alpha}(l_{\alpha} + 1) \\ \Theta(n^{-\frac{l_{\alpha}(l_{\alpha}+1)}{4}}) & \text{if } 2 \mid n, 2 \mid k, k = m - 1, \alpha < \frac{1}{2}, \text{ and } k > l_{\alpha}(l_{\alpha} + 1) \\ \Theta(1) & \text{otherwise } (i.e., \text{ if } 2 \nmid k \text{ or } k \leq m - 2) \end{cases}$$

*Proof.* The proposition is proved by applying Theorem 4. Let  $\mathcal{G}_{\mathcal{A}}$  denote the set of all directed unweighted graphs over  $\mathcal{A}$ . For any graph  $G \in \mathcal{G}_{\mathcal{A}}$  and any pair of alternatives a, b, if there is no edge between a and b then we say that a and b are *tied* in G. Let  $\mathcal{G}'_{\mathcal{A}} \subset \mathcal{G}_{\mathcal{A}}$  denote the set of all tournament graphs over  $\mathcal{A}$ , i.e.,  $\mathcal{G}'_{\mathcal{A}}$  consists of graphs without ties.

For any profile P, let UMG(P) denote the unweighted majority graph of P, which is a graph in  $\mathcal{G}_{\mathcal{A}}$  such that there is an edge  $a \to b$  if and only if  $w_P(a, b) > 0$ . It is not hard to see that Copeland<sub> $\alpha$ </sub> can be defined over  $\mathcal{G}_{\mathcal{A}}$ .

Like Proposition 1, the set of all graphs in  $\mathcal{G}_{\mathcal{A}}$  resulted from UMGs of *n*-profiles can be characterized by  $\mathcal{G}_{\mathcal{A}}^n$  that is defined in the following claim.

**Claim 6.** For any 
$$n > m^4$$
, we have  $\mathcal{G}^n_{\mathcal{A}} = \{UMG(P) : P \in \mathcal{L}(\mathcal{A})^n\} = \begin{cases} \mathcal{G}_{\mathcal{A}} & \text{if } 2 \mid n \\ \mathcal{G}'_{\mathcal{A}} & \text{if } 2 \nmid n \end{cases}$ 

*Proof.* The claim is proved by directly applying Proposition 1 and noticing that the middle tier of a palindromic order is empty if and only if its UMG does not contain any ties.  $\Box$ 

Applying the 0 case in Theorem 4. In this part we prove that the 0 case of the proposition can be obtained from applying the 0 case in Theorem 4. By Claim 6, it suffices to prove that for any  $n > m^4$ , there exists  $G \in \mathcal{G}_A^n$  such that  $|Cd_\alpha(G)| = k$  if and only if  $2 \mid n$ , or  $2 \nmid k$ , or  $k \leq m-2$ .

- The "if" direction. We first prove the "if" direction by construction. When 2 | n, by Claim 6 we have  $\mathcal{G}_{\mathcal{A}}^n = \mathcal{G}_{\mathcal{A}}$ , and we can choose a graph G such that all alternatives in  $\{1, \ldots, k\}$  are tied to each other, and all of them are strictly preferred to other alternatives. It follows that  $\operatorname{Cd}_{\alpha}(G) = \{1, \ldots, k\}$ , which means that  $|\operatorname{Cd}_{\alpha}(G)| = k$ . When  $2 \nmid k$  or  $k \leq m-2$ , we can use the complete graphs in the  $\Theta(1)$  case as illustrated in Figure 8 (a) and (b).
- The "only if" direction. Now we prove the "only if" direction for all  $n > m^4$ . When  $2 \nmid n, 2 \mid k$ , and m = k, we have  $\mathcal{G}_{\mathcal{A}}^n = \mathcal{G}_{\mathcal{A}}'$ . Suppose for the sake of contradiction that there exists  $G \in \mathcal{G}_{\mathcal{A}}'$  such that  $|\mathrm{Cd}_{\alpha}(G)| = k$ . Notice that G is a complete graph. Therefore, the total Copeland<sub> $\alpha$ </sub> score for all alternatives is  $\binom{m}{2}$ , which means that each alternative must get  $\frac{m+1}{2} \notin \mathbb{N}$  points. This is a contradiction because the Copeland<sub> $\alpha$ </sub> score of each alternative must be an integer as G is a complete graph. When  $2 \nmid n, 2 \mid k$ , and m = k + 1, again we have  $\mathcal{G}_{\mathcal{A}}^n = \mathcal{G}_{\mathcal{A}}'$ . Suppose for the sake of contradiction that there exists  $G \in \mathcal{G}_{\mathcal{A}}'$  such that  $|\mathrm{Cd}_{\alpha}(G)| = k$ . Again, the total Copeland<sub> $\alpha$ </sub> score for all alternatives is  $\binom{m}{2}$ . Suppose each winner gets  $u \in \mathbb{N}$  points. Then, the total Copeland<sub> $\alpha$ </sub> score must be between (m-1)u and mu 1. Notice that the total Copeland<sub> $\alpha$ </sub> score for all alternatives is  $\binom{m}{2}$ . If  $u \leq \frac{m+1}{2}$ , then we have  $\binom{m}{2} > mu 1$ ; and if  $u \geq \frac{m+1}{2}$ , then  $\binom{m}{2} < (m-1)u$ . Either case leads to a contradiction.

Applying the exponential case in Theorem 4. We now prove that the exponential case of Theorem 4 do not occur. We note that  $\text{UMG}(\pi_{\text{uni}})$  is the empty graph, which means that  $\text{EO}(\pi_{\text{uni}})$  only contains the middle tier  $T_0$ . Therefore, for any palindromic order O with  $|\text{Cd}_{\alpha}(O)| = k$ , we have that O refines  $\text{EO}(\pi_{\text{uni}})$ , which means that  $\mathcal{O}_{r,k,n}^{\Pi} \neq \emptyset$ .

Applying the polynomial case in Theorem 4. The remainder of the proof focuses on characterizing  $\ell_{\min}$  in the polynomial case of Theorem 4. We first prove an alternative characterization of  $\ell_{\min}$  that will be frequently used. It states that  $\ell_{\min}$  equals to the minimum number of tied pairs in graphs in  $\mathcal{G}^n_{\mathcal{A}}$  where there are exactly k Copeland<sub> $\alpha$ </sub> winners. Recall from Claim 6 that when nis sufficiently large (>  $m^4$ ), which is the case we will focus in the rest of the proof, for any even number n,  $\mathcal{G}^n_{\mathcal{A}} = \mathcal{G}_{\mathcal{A}}$ , which is the set of all unweighted directed graphs over  $\mathcal{A}$ ; and for any odd number n,  $\mathcal{G}^n_{\mathcal{A}} = \mathcal{G}'_{\mathcal{A}}$ , which is the set of all unweighted tournament graphs over  $\mathcal{A}$ . With a little abuse of notation, for any unweighted directed graph G over  $\mathcal{A}$ , we let Ties(G) denote the number of tied pairs in G.

**Claim 7.** For any  $\Pi$  with  $\pi_{uni} \in CH(\Pi)$ , n, and k such that  $\mathcal{O}_{r,k,n}^{\Pi} \neq \emptyset$  under Copeland<sub> $\alpha$ </sub>, we have:

$$\ell_{\min} = \min\{Ties(G) : G \in \mathcal{G}_{\mathcal{A}}^n \ s.t. \ |Cd_{\alpha}(G)| = k\}$$

*Proof.* Because  $\text{EO}(\pi_{\text{uni}})$  only contains the middle tier  $T_0$ , any palindromic order refines it, which means that  $\mathcal{O}_{r,k,n}^{\Pi} = \{ O \in \mathcal{O}_{\mathcal{A}}^n : |\text{Cd}_{\alpha}(O)| = k \}.$ 

For any palindromic order O, let  $\mathrm{UMG}(O) \in \mathcal{G}_{\mathcal{A}}$  be the graph such that there is an edge  $a \to b$ if and only if  $a \to b$  is ranked before the middle tier. It is not hard to verify that  $\mathrm{Cd}_{\alpha}(O) =$  $\mathrm{Cd}_{\alpha}(\mathrm{UMG}(O))$ . Also it is not hard to verify that  $O \in \mathcal{O}_{\mathcal{A}}^n$  if and only if  $\mathrm{UMG}(O) \in \mathcal{G}_{\mathcal{A}}^n$ . Therefore, for any  $O \in \mathcal{O}_{r,k,n}^{\Pi}$  such that  $\mathrm{Ties}(O) = \ell_{\min}$ , we have  $\mathrm{UMG}(O) \in \mathcal{G}_{\mathcal{A}}^n$  and  $|\mathrm{Cd}_{\alpha}(\mathrm{UMG}(O))| = k$ . We also have  $\mathrm{Ties}(O) \geq \mathrm{Ties}(\mathrm{UMG}(O))$  because the latter corresponds to the  $T_0$  part in  $\mathrm{Ties}(O)$ . Therefore, we have  $\ell_{\min} \geq \min{\mathrm{Ties}(G) : G \in \mathcal{G}_{\mathcal{A}}^n}$  s.t.  $|\mathrm{Cd}_{\alpha}(G)| = k$ }. For any  $G^* = \arg \min_{G \in \mathcal{G}^n_{\mathcal{A}}: |\operatorname{Cd}_{\alpha}(G)| = k} \operatorname{Ties}(G)$ , we construct a palindromic order O whose  $T_0$  consists of all tied edges in  $G^*$ , and each of the other tiers contains exactly one edge. It is not hard to verify that  $\operatorname{Ties}(O) = \operatorname{Ties}(G^*)$  and  $\operatorname{UMG}(O) = G^*$ , which means that  $O \in \mathcal{O}^n_{\mathcal{A}}$  and  $|\operatorname{Cd}_{\alpha}(O)| = k$ . Therefore,  $\operatorname{Ties}(G^*) \geq \ell_{\min}$ . This proves the claim.  $\Box$ 

The  $\Theta(1)$  case:  $2 \nmid k$  or  $k \leq m-2$ . We first prove the  $\Theta(1)$  case as a warm up. Part of the proof will be reused in the proof of other cases. In fact, the  $\Theta(1)$  case covers the most number of combinations of m and k: it happens when k is odd or  $k \leq m-2$ . In light of Claim 7, it suffices to prove that in this case there exists a tournament graph G such that  $|Cd_{\alpha}(G)| = k$ , because Ties(G) = 0 and any tournament graph is in  $\mathcal{G}^n_{\mathcal{A}}$ . We will explicitly construct such G in the two subcases.

- $2 \nmid k$ . The following graph G has exactly k Copeland<sub> $\alpha$ </sub> winners  $\{1, \ldots, k\}$ , see Figure 8 (a) for an example of m = 6, k = 5. For any  $1 \leq i \leq k$  and any  $1 \leq s \leq \frac{k-1}{2}$  there is an edge  $i \rightarrow (1 + (i + s 1 \mod k))$ , for example the black edges in Figure 8 (a). For any  $1 \leq i \leq k$  and any  $k + 1 \leq j \leq m$ , there is an edge  $i \rightarrow j$ , for example the blue edges in Figure 8 (a).
- 2 | k and  $k \leq m 2$ . The following graph G has exactly k Copeland<sub> $\alpha$ </sub> winners  $\{1, \ldots, k\}$ , see Figure 8 (b) for an example of m = 6, k = 4. The edges are defined in the following steps.

Step 1: edges within  $\{1, \ldots, k\}$ . These edges are colored black in Figure 8 (b) and is the similar to the graph in Figure 8 (a), except that there are  $\frac{k}{2}$  "diagonal" edges whose directions need to be assigned because k is an even number. Formally, for any  $1 \le i \le \frac{k}{2}$  and any  $1 \le s \le \frac{k}{2}$ , there is an edge  $i \to i + s$ . For any  $\frac{k}{2} + 1 \le i \le k$  and any  $1 \le s \le \frac{k}{2} - 1$ , there is an edge  $i \to (1 + (i + s - 1 \mod k))$ .

Step 2: edges between  $\{1, \ldots, k\}$  and  $\{k + 1, \ldots, m\}$ . These edges are colored blue in Figure 8 (b). For any  $1 \le i \le \frac{k}{2}$  there is an edge  $(k + 1) \rightarrow i$ ; and for any other  $(i, j) \in \{1, \ldots, k\} \times \{k + 1, \ldots, m\}$ , there is an edge  $i \rightarrow j$ .

Step 3: edges within  $\{k+1,\ldots,m\}$ . There is an edge  $(k+2) \rightarrow (k+1)$ , colored red in Figure 8 (b). Directions of other edges are assigned arbitrarily to make G a complete graph.



Figure 8: Constructions for Copeland<sub> $\alpha$ </sub>, where the winners are  $\{1, \ldots, k\}$ .

The  $\Theta(n^{-\frac{k}{4}})$  case:  $2 \mid n, 2 \mid k$ , and  $\left\{ \begin{array}{l} (1)k = m, \mathrm{or} \\ (2)k = m - 1 \mathrm{and} \ \alpha \geq \frac{1}{2}, \mathrm{or} \\ (3)k = m - 1 \mathrm{and} \ k \leq l_{\alpha}(l_{\alpha} + 1) \end{array} \right.$  We prove that

 $\ell_{\min} = \frac{k}{2}$  by applying Claim 7 in the three subcases indicated in the statement of the proposition.

• (1)  $2 \mid n$ ,  $2 \mid k$ , and k = m. We first prove  $\ell_{\min} \leq \frac{k}{2}$ . By Claim 7, it suffices to construct an unweighted graph  $G \in \mathcal{G}_{\mathcal{A}}^n = \mathcal{G}_{\mathcal{A}}$  such that  $|Cd_{\alpha}(G)| = k$  and  $Ties(G) = \frac{k}{2}$ . In fact, the following unweighted directed graph G with  $\frac{k}{2}$  ties has k Copeland winners  $\{1, \ldots, k\}$ , see Figure 8 (c) for an example. For any  $1 \leq i \leq k$  and any  $1 \leq s \leq \frac{k}{2} - 1$ , there is an edge  $i \rightarrow (1 + (i + s - 1 \mod k))$ . There are  $\frac{k}{2}$  ties in G, namely  $\{1, \frac{k}{2} + 1\}, \{2, \frac{k}{2} + 2\}, \ldots, \{\frac{k}{2}, k\}$ .

Next, we prove  $\ell_{\min} \geq \frac{k}{2}$ . By Claim 7, it suffices to prove that it is impossible for any unweighted directed graph G with exactly k = m Copeland<sub> $\alpha$ </sub> winners to have strictly less than  $\frac{k}{2}$  ties. Suppose for the sake of contradiction that such graph exists, denoted by G'. Then, there must exist an alternative, w.l.o.g. alternative 1, that is not tied to any other alternative in G', otherwise at least  $\frac{k}{2}$  ties are necessary to "cover" all alternatives. This means that the Copeland<sub> $\alpha$ </sub> score of alternative 1 must be an integer, denoted by u, which means that the Copeland<sub> $\alpha$ </sub> score of all alternatives must be u. Therefore, the total Copeland<sub> $\alpha$ </sub> score of all alternatives is uk.

Notice that each directed each edge contributes 1 to the total Copeland<sub> $\alpha$ </sub> score and each tie contributes  $2\alpha \in [0, 2]$  to the total Copeland<sub> $\alpha$ </sub> score. Therefore, the total Copeland<sub> $\alpha$ </sub> score is between  $\binom{m}{2} - (\frac{k}{2} - 1)$  (which corresponds to the case where there are  $\frac{k}{2} - 1$  ties and  $\alpha = 0$ ) and  $\binom{m}{2} + (\frac{k}{2} - 1)$  (which corresponds to the case where there are  $\frac{k}{2} - 1$  ties and  $\alpha = 1$ ). However, if  $u \geq \frac{k}{2}$  then  $uk > \binom{m}{2} + (\frac{k}{2} - 1)$ ; and if  $u \leq \frac{k}{2} - 1$  then  $uk < \binom{m}{2} - (\frac{k}{2} - 1)$ . In either case  $uk \notin [\binom{m}{2} - (\frac{k}{2} - 1), \binom{m}{2} + (\frac{k}{2} - 1)]$ , which is a contradiction. Therefore,  $\ell_{\min} = \frac{k}{2}$ .

- (2) 2 | n, 2 | k, k = m 1, and  $\alpha \geq \frac{1}{2}$ .  $\ell_{\min} \leq \frac{k}{2}$  is proved by constructing a graph  $G \in \mathcal{G}_{\mathcal{A}}^n = \mathcal{G}_{\mathcal{A}}$  whose subgraph over  $\{1, \ldots, m-1\}$  is the same as the graph constructed in case (1) above, and then we add m-1 edges  $i \to m$  for each  $i \leq m-1$ .  $\ell_{\min} \geq \frac{k}{2}$  is proved by a similar argument as in case (1) above. More precisely, suppose for the sake of contradiction that there exists a graph G' such that  $\operatorname{Cd}_{\alpha}(G') = k = m-1$  and  $\operatorname{Ties}(G') < \frac{k}{2}$ . Then, there must exist an alternative, w.l.o.g. alternative 1, that is not tied to any other alternative in G'. This means that the Copeland<sub> $\alpha$ </sub> score of alternative 1 must be an integer, denoted by u. Because  $\alpha \geq \frac{1}{2}$ , the total Copeland<sub> $\alpha$ </sub> score is between  $\binom{m}{2}$  and  $\binom{m}{2} + \frac{k}{2} 1$ , and is also between uk and u(k+1) 1. However, if  $u \leq \frac{k}{2}$ , then  $u(k+1) 1 < \binom{m}{2} = \frac{k(k+1)}{2}$ ; and if  $u \geq \frac{k}{2} + 1$ , then  $uk > \binom{m}{2} + \frac{k}{2} 1$ . In either case  $[\binom{m}{2}, \binom{m}{2} + \frac{k}{2} 1] \cap [uk, u(k+1) 1] = \emptyset$ , which is a contradiction. This means that  $\ell_{\min} = \frac{k}{2}$ .
- (3) 2 | n, 2 | k, k = m-1, and  $k \leq l_{\alpha}(l_{\alpha}+1)$ .  $\ell_{\min} \leq \frac{k}{2}$  is proved by the same graph as in case (2) above.  $\ell_{\min} \geq \frac{k}{2}$  is proved by a similar argument as in case (1) above. More precisely, for the sake of contradiction suppose there exists a graph G' such that  $Cd_{\alpha}(G') = k = m-1$  and  $Ties(G') < \frac{k}{2}$ . Then, there must exist a winner that is not tied to any other alternative in G'. Therefore, the Copeland\_{\alpha} score of the k winners is an integer. However, in order for the Copeland\_{\alpha} score of any alternative a who is tied in G' (with any other alternative) to be an integer, a must be tied with at least  $l_{\alpha}$  alternatives. Among these  $l_{\alpha}$  alternatives, at least  $l_{\alpha} 1$  must be co-winners, each of which must be tied to at least  $l_{\alpha}$  alternatives for their total Copeland\_{\alpha} scores to be integral. It is not hard to verify that the total number of ties in G' is  $\frac{l_{\alpha}(l_{\alpha}+1)}{2} \geq \frac{k}{2}$ , which is a contradiction. This means that  $\ell_{\min} = \frac{k}{2}$ .

The  $\Theta(n^{-\frac{l_{\alpha}(l_{\alpha}+1)}{4}})$  case:  $2 \mid n, 2 \mid k, k = m - 1, \alpha < \frac{1}{2}$ , and  $k > l_{\alpha}(l_{\alpha} + 1)$ . Like the  $\Theta(n^{-\frac{k}{4}})$  case, we apply Claim 7 to prove that  $\ell_{\min} = \frac{l_{\alpha}(l_{\alpha}+1)}{2}$ . We first prove  $\ell_{\min} \leq \frac{l_{\alpha}(l_{\alpha}+1)}{2}$  by explicitly constructing an unweighted directed graph  $G \in \mathcal{G}_{\mathcal{A}}^n = \mathcal{G}_{\mathcal{A}}$  such that the Copeland<sub> $\alpha$ </sub> winners

are  $\{1, \ldots, k\}$  whose Copeland<sub> $\alpha$ </sub> scores are  $\frac{k}{2}$ . Let  $\mathcal{A} = \mathcal{A}_1 \cup \mathcal{A}_2 \cup \{m\}$ , where  $\mathcal{A}_1 = \{1, \ldots, l_{\alpha}\}$ and  $\mathcal{A}_1 = \{l_{\alpha} + 1, \ldots, m - 1\}$ . The construction of G depends on the parity of  $l_{\alpha}$ , discussed in the following two cases.

•  $l_{\alpha}$  is odd. We construct  $G \in \mathcal{G}_{\mathcal{A}}^n = \mathcal{G}_{\mathcal{A}}$  in the following three steps, illustrated in Figure 9. Note that Figure 9 is only used for the purpose of illustration but does not correspond to the  $\Theta(n^{-\frac{l_{\alpha}(l_{\alpha}+1)}{4}})$  case, because  $k = 8 < 12 = l_{\alpha}(l_{\alpha}+1)$ . A real example requires  $k \geq 14$ , which contains too many edges in G to be easily visible.

Step 1. Edges within  $A_1 \cup \{m\}$  and edges within  $A_2$ . All alternatives in  $A_1 \cup \{m\}$  are tied, i.e., there is no edge between any pair of alternatives in  $A_1 \cup \{m\}$ . It is not hard to verify that  $|A_2|$  is an odd number. Therefore, we let the subgraph of G over  $A_2$  to be the completely graph as the one shown in Figure 8 (a). After this step, the Copeland<sub> $\alpha$ </sub> score of each alternative in  $A_1 \cup \{m\}$  becomes  $\alpha l_{\alpha}$ , and the Copeland<sub> $\alpha$ </sub> score of each alternative in  $A_2$ .

Step 2. Edges between  $A_1$  and  $A_2$ . We assign all edges between  $A_1$  and  $A_2$  an direction such that each alternative in  $A_1$  has exactly  $\frac{k}{2} - \alpha l_{\alpha}$  outgoing edges, and each alternative in  $A_2$  has  $\frac{l_{\alpha}-1}{2}$  or  $\frac{l_{\alpha}+1}{2}$  outgoing edges. This can be done by assigning edges from  $A_1$  to  $A_2$  evenly across  $A_2$ . In the graph that combines edges in Step 1 and Step 2, the Copeland<sub> $\alpha$ </sub> score of each alternative in  $A_1 \cup \{m\}$  becomes exactly  $\frac{k}{2}$ , the Copeland<sub> $\alpha$ </sub> score of the remaining  $\frac{k-l_{\alpha}(l_{\alpha}+1)}{2} + \alpha l_{\alpha}^2$  alternatives in  $A_2$  becomes  $\frac{k}{2}$ , and the Copeland<sub> $\alpha$ </sub> score of the remaining  $\frac{k+l_{\alpha}(l_{\alpha}+1)}{2} - \alpha l_{\alpha}^2$  alternatives in  $A_2$  becomes  $\frac{k}{2} - 1$ .

Step 3. Edges between m and  $A_2$ . Finally, the directions of edges in this step are used to guarantee the Copeland<sub> $\alpha$ </sub> score of all alternatives in  $A_2$  is  $\frac{k}{2}$ .

Let G denote the union of edges defined in Step 1-3. It is not hard to verify that the Copeland<sub> $\alpha$ </sub> score of all alternatives in  $A_1 \cup A_2$  is  $\frac{k}{2}$ , and the Copeland<sub> $\alpha$ </sub> score of alternative m is  $\alpha l_{\alpha} + \frac{k - l_{\alpha}(l_{\alpha} + 1)}{2} + \alpha l_{\alpha}^2 < \frac{k}{2}$ , because  $\alpha < \frac{1}{2}$ . This proves  $\ell_{\min} \leq \frac{l_{\alpha}(l_{\alpha} + 1)}{2}$  when  $l_{\alpha}$  is odd.



Figure 9: An illustration of the constructions for the  $\Theta(n^{-\frac{l_{\alpha}(l_{\alpha}+1)}{4}})$  case with odd  $l_{\alpha}$ , where m = 9, k = 8,  $\alpha = \frac{1}{3}$ , and  $l_{\alpha} = 3$ . Note that this figure is only used for illustrating the three-step construction. It does not correspond to the  $\Theta(n^{-\frac{l_{\alpha}(l_{\alpha}+1)}{4}})$  case because  $k < l_{\alpha}(l_{\alpha}+1)$ . In (b) and (c), edges from right to left are colored red.

•  $l_{\alpha}$  is even. Like the  $2 \nmid l_{\alpha}$  case, we construct  $G \in \mathcal{G}_{\mathcal{A}}^n = \mathcal{G}_{\mathcal{A}}$  in the following three steps.

Step 1. Edges within  $A_1 \cup \{m\}$  and edges within  $A_2$ . All alternatives in  $A_1 \cup \{m\}$ are tied, i.e., there is no edge between any pair of alternatives in  $A_1 \cup \{m\}$ . It is not hard to verify that  $|A_2|$  is an odd number. Therefore, we let the subgraph of G over  $A_2$ to be isomorphic to the (complete) subgraph over  $\{1, \ldots, k\}$  as the one in Figure 8 (b). After this step, the Copeland<sub> $\alpha$ </sub> score of each alternative in  $A_1 \cup \{m\}$  becomes  $\alpha l_{\alpha}$ , the Copeland<sub> $\alpha$ </sub> score of  $\frac{k-l_{\alpha}}{2}$  alternative in  $A_2$  becomes  $\frac{k-l_{\alpha}}{2}$ , and the Copeland<sub> $\alpha$ </sub> score of the remaining  $\frac{k-l_{\alpha}}{2}$  alternative in  $A_2$  becomes  $\frac{k-l_{\alpha}}{2} - 1$ .

Step 2. Edges between  $A_1$  and  $A_2$ . We assign all edges between  $A_1$  and  $A_2$  and direction such that each alternative in  $A_1$  has exactly  $\frac{k}{2} - \alpha l_{\alpha}$  outgoing edges, and each alternative in  $A_2$  has  $\frac{l_{\alpha}}{2} - 1$  or  $\frac{l_{\alpha}}{2}$  outgoing edges. This can be done by assigning edges from  $A_1$  to  $A_2$  uniformly across  $A_2$ , starting with alternatives whose Copeland<sub> $\alpha$ </sub> scores are  $\frac{k-l_{\alpha}}{2}$  in Step 1. In the graph that combines edges in Step 1 and Step 2, the Copeland<sub> $\alpha$ </sub> score of each alternative in  $A_1 \cup \{m\}$  becomes exactly  $\frac{k}{2}$ , the Copeland<sub> $\alpha$ </sub> score of  $\frac{l_{\alpha}^2}{2} + \alpha l_{\alpha}$ alternatives in  $A_2$  becomes  $\frac{k}{2}$ , and the Copeland<sub> $\alpha$ </sub> score of the remaining  $k - l_{\alpha} - \frac{l_{\alpha}^2}{2} - \alpha l_{\alpha}$ (which is strictly positive because  $k > l_{\alpha}(l_{\alpha} + 1)$  and  $l_{\alpha} \ge 4$  because  $\alpha < \frac{1}{2}$ ) alternatives in  $A_2$  becomes  $\frac{k}{2} - 1$ .

Step 3. Edges between m and  $A_2$ . Finally, the directions of edges in this step are used to guarantee the Copeland<sub> $\alpha$ </sub> score of all alternatives in  $A_2$  is  $\frac{k}{2}$ .

Let G denote the union of edges defined in Step 1-3. It is not hard to verify that the Copeland<sub> $\alpha$ </sub> score of all alternatives in  $A_1 \cup A_2$  is  $\frac{k}{2}$ , and the Copeland<sub> $\alpha$ </sub> score of alternative  $m \text{ is } \frac{l_{\alpha}^2}{2} + \alpha l_{\alpha} + \alpha l_{\alpha} \leq \frac{l_{\alpha}^2}{2} + 2 \leq \frac{l_{\alpha}^2 + l_{\alpha}}{2} < \frac{k}{2}, \text{ because } l_{\alpha} \geq 4 \text{ and } k > l_{\alpha}(l_{\alpha} + 1). \text{ This proves } l_{\min} \leq \frac{l_{\alpha}(l_{\alpha} + 1)}{2} \text{ when } l_{\alpha} \text{ is even.}$ 

We now prove that  $\ell_{\min} \geq \frac{l_{\alpha}(l_{\alpha}+1)}{2}$ . The proof is similar to the proof of the  $\Theta(n^{-\frac{k}{4}})$  case (3). More precisely, for the sake of contradiction that there exists a graph G' such that  $\operatorname{Cd}_{\alpha}(G') = k = 1$ m-1 and Ties $(G') < \frac{l_{\alpha}(l_{\alpha}+1)}{2} < \frac{k}{2}$ . Then, there must exist a winner that is not tied to any other alternative in G'. Therefore, the Copeland<sub> $\alpha$ </sub> score of the k winners is an integer. However, in order for the Copeland<sub> $\alpha$ </sub> score of any alternative a who is tied in G' (with any other alternative) to be an integer, following the same argument as in the proof of the  $\Theta(n^{-\frac{k}{4}})$  case (3), we must have Ties $(G') \geq \frac{l_{\alpha}(l_{\alpha}+1)}{2}$ , which is a contradiction. This means that  $\ell_{\min} = \frac{l_{\alpha}(l_{\alpha}+1)}{2}$ 

This completes the proof for Proposition 2.

#### C.5**Proof of Proposition 3**

**Proposition 3.** (Max smoothed likelihood of ties: maximin). Let  $\mathcal{M} = (\Theta, \mathcal{L}(\mathcal{A}), \Pi)$  be a strictly positive and closed single-agent preference model where  $\pi_{uni} \in CH(\Pi)$ . For any  $2 \leq k \leq m$ and any  $n \in \mathbb{N}$ ,

$$\widetilde{Tie}_{\Pi}^{\max}(MM,k,n) = \Theta(n^{-\frac{k-1}{2}})$$

*Proof.* The proposition is proved by applying Theorem 4.

The 0 case and the exponential case. We first prove that when  $n > m^4$ , the 0 case and the exponential case of Theorem 4 do not occur. By Proposition 1 and notice that  $EO(\pi_{uni})$  only contains  $T_0$ , it suffices to prove that for any k, there exists  $O \in \mathcal{O}'_{\mathcal{A}}$  such that  $|\mathcal{M}(O)| = k$ . This is done by first defining a directed weighted graph G, and then let O = EO(G). Let G denote the following directed weighted graph over  $\mathcal{A}$ , as illustrated in Figure 10.



Figure 10: Construction for maximin where m = 5 and k = 4.

• G contains the following cycle of k edges, the weight of each of which is  $m^2 + 1$ .

$$E = \{1 \to 2, 2 \to 3, \dots, (k-1) \to k, k \to 1\}$$

- The other edges within  $\{1, \ldots k\}$  have different weights, whose parity is the same as  $m^2 + 1$  and whose absolute values are strictly smaller than  $m^2 + 1$  and strictly larger than 0.
- The absolute values of any other edge has the same parity as  $m^2 + 1$  and are strictly larger than  $m^2 + 1$ . In addition, for any  $1 \le i \le k$  and any  $k + 1 \le j \le m$ , the weight on  $i \to j$  is positive.

It is not hard to check that  $MM(G) = \{1, \ldots, k\}$ . Let O = EO(G). Because there is no tie in G, the middle tier  $T_0$  in EO(G) is empty, which means that  $O \in \mathcal{O}'_A \subseteq \mathcal{O}^n_A$ .

The polynomial case. We first prove that  $\ell_{\min} \leq k - 1$  by using O defined above for the 0 case and exponential case. There are only two tiers with more than one elements, namely the tiers corresponding to E and the inverse E, which means that Ties(O) = k - 1.

We now prove that  $\ell_{\min} \geq k-1$ . For any  $O \in \mathcal{O}_{r,k,n}^{\Pi}$ , suppose w.l.o.g. that the maximin winners are  $\{1, \ldots, k\}$ . For each  $1 \leq i \leq k$ , let  $i \to a_i$  denote an arbitrary edge that corresponds to the min score of *i*. Because  $\{1, \ldots, k\}$  have the same min score,  $\{i \to a_i : 1 \leq i \leq k\}$  must be in the same tier. If they are in  $T_0$ , then  $T_0$  contains at least 2k edges  $\{i \to a_i, a_i \to i : 1 \leq i \leq k\}$ , which means that  $\text{Ties}(O) \geq |T_0|/2 \geq k$ . If they are not in  $T_0$ , then  $\text{Ties}(O) \geq k-1$ . In either case we have  $\ell_{\min} \geq k-1$ .

Therefore,  $\ell_{\min} = k - 1$ . This completes the proof for Proposition 3.

# C.6 Proof of Proposition 4

**Proposition 4.** (Max smoothed likelihood of ties: Schulze). Let  $\mathcal{M} = (\Theta, \mathcal{L}(\mathcal{A}), \Pi)$  be a strictly positive and closed single-agent preference model with  $\pi_{uni} \in CH(\Pi)$ . For any  $2 \leq k \leq m$  and any  $n \in \mathbb{N}$ ,

$$\widetilde{Tie}_{\Pi}^{\max}(Sch,k,n) = \Theta(n^{-\frac{k-1}{2}})$$

*Proof.* The proposition is proved by applying Theorem 4.

The 0 case and the exponential case. When  $n > m^4$ , the 0 case and the exponential case of Theorem 4 do not occur. This is prove by the same graph G as in the proof of Proposition 3 that is illustrated in Figure 10.

The polynomial case.  $\ell_{\min} \leq k-1$  is proved by using O = EO(G) for the same G used in Proposition 3.

We now prove that  $\ell_{\min} \geq k - 1$ . For any  $O \in \mathcal{O}_{r,k,n}^{\Pi}$ , suppose w.l.o.g. that the Schulze winners are  $A_1 = \{1, \ldots, k\}$ . We define the following t unweighted directed graphs. For any  $1 \leq i \leq t + 1$ , let  $G_i$  denote the graph over  $\mathcal{A}$  whose edges are  $T_1 \cup \ldots \cup T_{t+1-i}$ . Let  $G_{t+1}$  denote the graph without any edges. By definition  $G_1 = \text{UMG}(O)$ . Given any unweighted directed graph G', a node a is undominated, if for any node b that there exists a directed path from b to a, there exists a directed path from a to b.

**Claim 8.** For each  $1 \leq i \leq t$ ,  $\{1, \ldots, k\}$  are undominated in  $G_i$ .

*Proof.* Suppose for the sake of contradiction that the claim is not true and w.l.o.g. alternative 1 is dominated by j in  $G_i$  for some  $1 \le i \le t$ . Let w denote the weight of the edges in  $T_{t+1-i}$ . This means that  $s[j,1] \ge w$  and in each path from 1 to j there is an edge whose weight is strictly less than w. Therefore, s[j,1] > s[1,j], which contracts the assumption that 1 is a winner.

For each  $1 \leq i \leq t$ , let  $S_i \subset 2^{A_1}$  denote the connected components in  $G_i$  that intersect  $A_1 = \{1, \ldots, k\}$  and let  $\eta_i = |S_i|$ . By Claim 8, each set S in  $S_i$  has no incoming edges from  $A \setminus S$  in  $G_i$ , otherwise  $S \cap A_1$  are dominated in  $G_i$ . If  $\eta_1 > 1$ , then we have  $|T_0| \geq 2(k-1)$ , which means that  $\text{Ties}(O) \geq k - 1$  and therefore  $\ell_{\min} \geq k - 1$ , which proves the proposition. In the rest of the proof we assume that  $\eta_1 = 1$ .

For each  $1 \leq i \leq t$ , because  $G_{i+1}$  is obtained from  $G_i$  by removing some edges, each set in  $S_{i+1}$ must be a subset of a set in  $S_i$ . If  $\eta_{i+1} > \eta_i$ , then there are at least  $\eta_{i+1} - \eta_i + 1$  sets in  $S_{i+1}$ , denoted by  $S'_{i+1}$ , that are strict subsets of some sets in  $G_i$ . For any  $S'_{i+1} \in S'_{i+1}$ , let  $S'_i \in S'_i$  be the set such that  $S'_{i+1} \subsetneq S'_i$ . By Claim 8,  $S'_{i+1}$  does not contain any incoming edges from  $\mathcal{A} \setminus S'_{i+1}$  in  $G_{i+1}$ , and in particular, no edges from  $S'_i \setminus S'_{i+1}$  to  $S'_{i+1}$  (which are in  $G_i$  and there is at least one such edge because  $S'_i$  is a connected component in  $G_i$ ) are in  $G_{i+1}$ . This means that  $T_{t+1-i}$  contains at least one incoming edge to each set in  $S'_{i+1}$ , and such sets of edges are non-overlapping because  $S'_{i+1}$  consists of non-overlapping subsets of  $\mathcal{A}$ . Therefore,  $|T_{t+1-i}| \geq \eta_{i+1} - \eta_i + 1$ . If  $\eta_{i+1} = \eta_i$ , then apparently we have  $|T_{t+1-i}| \geq \eta_{i+1} - \eta_i + 1 = 1$ . Recall that  $\eta_1 = 1$  and  $\eta_{t+1} = k$ , we have:

$$\operatorname{Ties}(O) = \sum_{i=1}^{t} (|T_{t+1-i}| - 1) + |T_0|/2 \ge \sum_{i=1}^{t} (\eta_{i+1} - \eta_i) = \eta_{t+1} - \eta_1 = k - 1$$

Again, we have  $\ell_{\min} \ge k - 1$ .

This completes the proof for Proposition 4.

### C.7 Proof of Proposition 5

**Proposition 5.** (Max smoothed likelihood of ties: ranked pairs). Let  $\mathcal{M} = (\Theta, \mathcal{L}(\mathcal{A}), \Pi)$  be a strictly positive and closed single-agent preference model with  $\pi_{uni} \in CH(\Pi)$ . For any  $2 \leq k \leq m$ and any  $n \in \mathbb{N}$ ,

$$\Omega(n^{-\frac{k-1}{2}}) \le \widetilde{Tie}_{\Pi}^{\max}(RP, k, n) \le n^{-\Omega(\frac{\log k}{\log \log k})}$$

Moreover, when  $m \ge k + 5 \lceil \log k \rceil$ ,

$$\widetilde{\mathit{Tie}}_{\Pi}^{\max}(\mathit{RP},k,n) \leq \Omega(n^{-\frac{\lceil \log k \rceil}{2}})$$

When k = 2,

$$\widetilde{Tie}_{\Pi}^{\max}(RP,2,n) = \Theta(n^{-0.5})$$

*Proof.* The proposition is proved by applying Theorem 4. It is not hard to verify that the 0 case and the exponential case of Theorem 4 do not occur when n is sufficiently large. Let  $s_k$  denote the largest integer such that  $s_k! \leq k$ . It follows that  $s_k = O(\frac{\log k}{\log \log k})$ . To prove the first inequality in the statement of the proposition, it suffices to prove that  $s_k - 1 \leq \ell_{\min} \leq k - 1$ .

We first prove that  $\ell_{\min} \leq k - 1$  by constructing a weighted directed graph G such that  $\operatorname{Ties}(\operatorname{EO}(G)) = k - 1$ ,  $|\operatorname{RP}(G)| = k$ , and  $\operatorname{EO}(G) \in \mathcal{O}'_{\mathcal{A}} \subseteq \mathcal{O}_{\mathcal{A}}$ . In fact, we can use the graph in the proof of Proposition 3 illustrated in Figure 10. The winner is determined by the tie-breaking order among edges in the cycle  $1 \rightarrow 2 \rightarrow \cdots \rightarrow k \rightarrow 1$ .

We now prove that  $\ell_{\min} \geq s_k - 1$ . Let  $O \in \mathcal{O}_{\mathcal{A}}^n$  denote the palindromic order such that  $\text{Ties}(O) = \ell_{\min}$ . The total number of ways to break ties in ranked pairs is no more than  $(\ell_{\min} + 1)!$ , and the maximum is achieved when a tier before  $T_0$  contains all l + 1 tied edges. Therefore, we must have  $(\ell_{\min} + 1)! \geq k$ , which means that  $\ell_{\min} \geq s_k - 1$ .

**The "moreover" part.** This part is proved by construction. When  $m \ge 5 \lceil \log k \rceil + k$ , we construct a weighted directed graph G such that (1)  $EO(G) \in \mathcal{O}'_{\mathcal{A}}$ , (2) |RP(G)| = k, and (3)  $Ties(EO(G)) = \lceil \log k \rceil$ . The construction is illustrated in Figure 11 (b). We let  $\mathcal{A}$  denote the union of the following three sets of alternatives.

- $\{0, \ldots, k-1\}$ . These are the ranked pairs winners. Notice that the index starts at 0 instead of 1.
- For each  $i \leq \lfloor \log k \rfloor$ , there are five alternatives  $\{a_i, b_i, c_i, d_i, f_i\}$ .
- Let A denote the remaining  $m k 5 \lceil \log k \rceil$  alternatives.

There are  $\lceil \log k \rceil$  pairs of edges with same weights in G: for any  $i \leq \lceil \log k \rceil$ ,  $w_G(a_i \rightarrow c_i) = w_G(b_i \rightarrow d_i)$ . The edges are divided into the following groups such that weights in an earlier group are higher than weights in an later group. We further require that no pair of edges have the same weight except  $a_i \rightarrow c_i$  and  $b_i \rightarrow d_i$ , and in particular, no edge has weight 0.

- Group 1. For each  $i \leq \lceil \log k \rceil$ , we have the following edges:  $c_i \to b_i$ ,  $d_i \to a_i$ ,  $c_i \to f_i$ ,  $d_i \to f_i$ . For each  $0 \leq j \leq k 1$ , there is an edge from j to each alternative in A. For each  $0 \leq j \leq k 1$  and each  $i \leq \lceil \log k \rceil$ , if the *i*-th digit from right in j's binary representation is 0, then there is an edge  $j \to a_i$ ; otherwise there is an edge  $j \to b_i$ .
- Group 2. For each  $i \leq \lceil \log k \rceil$  and each  $0 \leq j \leq k 1$ , there is an edge  $f_i \to j$ .



Figure 11: Construction for ranked pairs.  $a_i \to c_i$  and  $b_i \to d_i$  have the same weights. The figure only shows Group 1 edges.

• Group 3. All remaining edges are in Group 3.

For each  $i \leq \lceil \log k \rceil$ , if  $a_i \to c_i$  is fixed before (respectively, after)  $b_i \to d_i$ , then there is a path from each  $0 \leq j \leq k-1$  whose *i*-th digit from right in binary is 0 (respectively, 1) to  $f_i$ . Therefore, after edges in Group 1 are fixed by breaking ties between  $\{a_i \to c_i, b_i \to d_i\}$  for all  $i \leq \lceil \log k \rceil$ , there is a unique alternative  $0 \leq j \leq k-1$  that has a path to all  $f_i$ 's. Then, each alternative in  $\{0, \ldots, k-1\}$  except *j* will have an incoming edge from at least one  $f_i$ . It follows that *j* will be the ranked pairs winner. We note that each alternative in  $\{0, \ldots, k-1\}$  can be made to win by breaking ties in some way.

Therefore, in EO(G) we have  $T_0 = \emptyset$  and there are  $\lceil \log k \rceil$  tiers containing more than one edge, each of which contain exactly two edges. This means that  $\text{Ties}(\text{EO}(G)) = \lceil \log k \rceil$ , which means that  $\ell_{\min} \leq \lceil \log k \rceil$ .

When k = 2. The  $\Omega(n^{-0.5})$  lower bound has already been proved for general k. Therefore, it suffices to prove  $\ell_{\min} \ge 1$ . Suppose for the sake of contradiction that this is not true. Then, there exists a palindromic order O such that Ties(O) < 1 and |RP(O)| = 2. However, Ties(O) = 0 means that no edges are tied in WMG(P). Therefore,  $|\text{RP}(O)| = 1 \ne 2$ , which is a contradiction.

This completes the proof of Proposition 5.

# D Appendix for Section 4.3: STV and Coombs

### D.1 Formal Definitions and Statements of Results

We first formally define multi-round score-based elimination (MRSE) rules, which are generalizations of STV and Coombs.

**Definition 20.** A multi-round score-based elimination (MRSE) rule r for m alternatives is defined as a vector of m - 1 rules  $(r_2, \ldots, r_m)$  such that for each  $2 \le i \le m$ ,  $r_i$  is an integer positional scoring voting rule over i alternatives that outputs a total preorder over them according to their scores. Given a profile P over m alternatives, r(P) is chosen in m-1 steps. For each  $1 \le i \le m-1$ , in round i a loser (alternative in the lowest tier) under  $r_{m+1-i}$  is eliminated from the election. An alternative b is a winner if there is a way to break ties so that b is the remaining alternative after m-1 rounds of elimination. We now define the counterpart to palindromic orders in Section 4.2, called *PUT structures*, that contain all information needed to determine the winners for MRSE rules.

**Definition 21** (**PUT structure**). A PUT structure over  $\mathcal{A}$  is a mapping W that maps each  $B \subsetneq \mathcal{A}$  to a total pre-order over  $\mathcal{A} \setminus B$ . Let  $\mathcal{W}_{\mathcal{A}}$  denote the set of all PUT structures over  $\mathcal{A}$ .

For any MRSE rule r and any profile P, let  $PS_r(P)$  denote the PUT structure W such that for any  $B \subsetneq \mathcal{A}$ ,  $W(B) = r_{m-|B|}(P|_{\mathcal{A}\setminus B})$  is a total preorder over  $\mathcal{A}\setminus B$ , where  $P|_{\mathcal{A}\setminus B}$  is the profile over  $\mathcal{A}\setminus B$  that is obtained from P by removing all alternatives in B.

We note that  $PS_r(P)$  depends on both r and P. A PUT structure W can be equivalently represented by a *PUT graph*, where the nodes are non-empty subsets of  $\mathcal{A}$ , each node B is labeled by W(B), and there is an edge between node B and B' if  $B' = B \cup \{a\}$  and a is in the lowest tier of W(B). With a little abuse of notation, we will also use W to denote its PUT graph.

The major component of W is its subgraph whose nodes are reachable from  $\emptyset$ , denoted by MC(W). For example, a profile P and the major component of  $PS_{STV}(P)$  are shown in Figure 12. We note that the winners only depend on the major component of the PUT structure of the profile.

**Definition 22** (Tier representation and refinement of PUT structures). For any PUT structure  $W \in W_A$  and any  $B \subsetneq A$ , let  $T_1 \succ \cdots \succ T_s$  denote the tier representation of W(B), where for each  $i \leq s$ , alternatives in  $T_i$  are tied. Let  $Ties(W(B)) = \sum_{i \leq s} (|T_i| - 1)$  and let  $Ties(W) = \sum_{B \subseteq A} Ties(W(B))$ .

A PUT structure  $W_1$  refines another PUT structure  $W_2$ , if for all  $B \subsetneq A$ ,  $W_1(B)$  refines  $W_2(B)$ in the total preorder sense.

**Example 19.** Figure 12 shows a profile P and the PUT graph of  $PS_{STV}(P)$ .  $MC(PS_{STV}(P))$  is the subgraph whose nodes are  $\{\emptyset, \{1\}, \{2\}, \{1,2\}, \{1,3\}, \{2,3\}, \{1,2,4\}, \{1,3,4\}, \{2,3,4\}\}$ . Let  $W = PS_{STV}(P)$ . We have  $Ties(W(\emptyset)) = Ties(W(\{1\})) = Ties(W(\{2\})) = 1$ , and Ties(W(B)) = 0 for any other  $B \subsetneq A$ , which means that Ties(W) = 3.



Figure 12: A preference profile P, its PUT structure  $PS_{STV}(P)$  under STV and the corresponding PUT graph. The major component consists of the subgraph with the nine nodes in black.

Next, we define notation that will be used to present the theorem on the smoothed likelihood of ties under STV.

For any weak order O with tier representation  $T_1 \succ \cdots \succ T_t$ , where elements in any  $T_i$  are tied, we let GCD(O) denote the greatest common divisor of the sizes of the t tiers in O. That is,

$$\operatorname{GCD}(O) = \gcd(|T_1|, \dots, |T_t|)$$

**Example 20.** Let W denote the PUT structure in Figure 12 (b). We have  $GCD(W(\emptyset)) = GCD(3 \succ 4 \succ \{1,2\}) = gcd(1,1,2) = 1$ . As another example,  $GCD(\{3,4\} \succ \{1,2\}) = gcd(2,2) = 2$ .

The next lemma states that for any sufficiently large n, W can be represented by the PUT structure under STV or Coombs of some n-profile if and only if n is divisible by GCD(W(B)) for all  $B \subsetneq A$ .

**Lemma 3** (**PUT structures under STV and Coombs**). Let  $r \in \{STV, Coombs\}$ . For any  $m \ge 3$ , there exists  $N \in \mathbb{N}$  such that for any PUT structure W over A and any  $n \ge N$ ,

$$(\exists P \in \mathcal{L}(\mathcal{A})^n \text{ s.t. } PS_r(P) = W) \iff (\forall B \subsetneq \mathcal{A}, GCD(W(B)) \mid n)$$

The proof can be found in Appendix D.2. In light of Lemma 3, for any  $\mathcal{A}$  and any  $n \in \mathbb{N}$ , we define  $\mathcal{W}_n$  to be the PUT structures over  $\mathcal{A}$  such that for every  $B \subsetneq \mathcal{A}$ ,  $gcd(W(B)) \mid n$ .

**Definition 23.** Let  $r \in \{STV, Coombs\}$ . Given any  $\mathcal{A}$ , any  $n \in \mathbb{N}$ , any  $2 \leq k \leq m$ , and any distribution  $\pi$  over  $\mathcal{L}(\mathcal{A})$ , let

$$\mathcal{W}_{r,k,n}^{\pi} = \{ W \in \mathcal{W}_n : |r(W)| = k \text{ and } W \text{ refines } PS_r(\pi) \}$$

For any set of distributions  $\Pi$  over  $\mathcal{L}(\mathcal{A})$ , let  $\mathcal{W}_{r,k,n}^{\Pi} = \bigcup_{\pi \in CH(\Pi)} \mathcal{W}_{r,k,n}^{\pi}$ . When  $\mathcal{W}_{r,k,n}^{\Pi} \neq \emptyset$ , we let

$$w_{\min} = \min_{W \in \mathcal{W}_{r,k,n}^{\Pi}} Ties(W).$$

When  $\mathcal{W}_{r,k,n}^{\pi} \neq \emptyset$  for all  $\pi \in CH(\Pi)$ , we let

$$w_{mm} = \max_{\pi \in CH(\Pi)} \min_{W \in \mathcal{W}_{r,k,n}^{\pi}} Ties(W)$$

In words,  $\mathcal{W}_n$  consists of all PUT structures resulted from all *n*-profiles under  $r \in \{\text{STV}, \text{Coombs}\}$ .  $\mathcal{W}_{r,k,n}^{\pi}$  is the set of PUT structures W that satisfies three conditions: (1) W is the PUT structure of an *n*-profile under r; (2) there are exactly k winners in W according to r, and (3) W refines  $\text{PS}_r(\pi)$ .  $\mathcal{W}_{r,k,n}^{\Pi}$  is the union of all PUT structures corresponding to all  $\pi \in \text{CH}(\Pi)$ .  $w_{\min}$  is the minimum number of ties in PUT structures in  $\mathcal{W}_{r,k,n}^{\Pi}$ .  $w_{\min}$  is the maximum is taken for all  $\pi \in \text{CH}(\Pi)$ , and for any given  $\pi$ , the minimum is taken for all PUT structures in  $\mathcal{W}_{r,k,n}^{\pi}$ .

We note that  $w_{\min}$  and  $w_{\min}$  depend on  $r \in \{\text{STV}, \text{Coombs}\}, \Pi, k$ , and n, which are clear from the context. In fact,  $w_{\min}$  and  $w_{\min}$  correspond to  $m! - \alpha_n$  and  $m! - \beta_n$  in Theorem 2, respectively, and we have the following theorem for STV and Coombs.

**Theorem 5.** (Smoothed likelihood of ties: STV and Coombs). Let  $r \in \{STV, Coombs\}$ and let  $\mathcal{M} = (\Theta, \mathcal{L}(\mathcal{A}), \Pi)$  be a strictly positive and closed single-agent preference model. For any  $2 \leq k \leq m$  and any  $n \in \mathbb{N}$ ,

$$\widetilde{Tie}_{\Pi}^{\max}(r,k,n) = \begin{cases} 0 & \text{if } \forall W \in \mathcal{W}_n, |r(W)| \neq k \\ \exp(-\Theta(n)) & \text{otherwise if } \mathcal{W}_{r,k,n}^{\Pi} = \emptyset \\ \Theta\left(n^{-\frac{w_{\min}}{2}}\right) & \text{otherwise} \end{cases}$$
$$\widetilde{Tie}_{\Pi}^{\min}(r,k,n) = \begin{cases} 0 & \text{if } \forall W \in \mathcal{W}_n, |r(W)| \neq k \\ \exp(-\Theta(n)) & \text{otherwise if } \exists \pi \in CH(\Pi) \text{ s.t. } \mathcal{W}_{r,k,n}^{\pi} = \emptyset \\ \Theta\left(n^{-\frac{w_{\min}}{2}}\right) & \text{otherwise} \end{cases}$$

The proof can be found in Appendix D.3. Like in Section 4.2, we next characterize max smoothed likelihood of ties for STV and Coombs for distributions  $\Pi$  where  $\pi_{uni} \in CH(\Pi)$ .

**Proposition 6.** (Max smoothed likelihood of ties: STV and Coombs). Let  $r \in \{STV, Coombs\}$ and let  $\mathcal{M} = (\Theta, \mathcal{L}(\mathcal{A}), \Pi)$  be a strictly positive and closed single-agent preference model with  $\pi_{uni} \in CH(\Pi)$ . For any  $2 \leq k \leq m$  and any  $n \in \mathbb{N}$ ,

$$\widetilde{Tie}_{\Pi}^{\max}(r,k,n) = \begin{cases} \Theta(n^{-\frac{k-1}{2}}) & \text{if } \begin{cases} (1)m \ge 4, \text{ or } \\ (2)m = 3 \text{ and } k = 2, \text{ or } \\ (3)m = k = 3 \text{ and } (2 \mid n \text{ or } 3 \mid n) \\ 0 & \text{otherwise } (i.e., m = k = 3, 2 \nmid n, \text{ and } 3 \nmid n) \end{cases}$$

The proof of Proposition 6 can be found in Appendix D.4.

### D.2 Proof of Lemma 3

**Lemma 3.** (PUT structures under STV and Combs). Let  $r \in \{STV, Coombs\}$ . For any  $m \ge 3$ , there exists  $N \in \mathbb{N}$  such that for any PUT structure W over A and any  $n \ge N$ ,

$$(\exists P \in \mathcal{L}(\mathcal{A})^n \text{ s.t. } PS_r(P) = W) \iff (\forall B \subsetneq \mathcal{A}, GCD(W(B)) \mid n)$$

*Proof.* We first prove the Lemma for r = STV, then comment on how to modify the proof for Coombs.

**Proof for STV.** Notice The  $\Rightarrow$  direction for STV follows after noticing that the total plurality score of all alternatives after *B* is removed, which is *n*, is divisible by GCD(W(B)). More precisely, for any *n*-profile *P* such that  $\text{PS}_{\text{STV}}(P) = W$  and any  $B \subsetneq \mathcal{A}$ , let the tier representation of W(B) be:

$$W(B) = T_1 \succ T_2 \succ \cdots \succ T_t$$

For every  $s \leq t$ , let  $\gamma_s$  denote the plurality score of any alternative in  $T_s$  after B is removed. That is,

$$\gamma_s = \text{Score}_{\text{Plu}}(P|_{\mathcal{A}\setminus B}, a_s) \text{ for any } a_s \in T_s$$

Therefore, we have

$$n = \sum_{a \in (\mathcal{A} \setminus B)} \operatorname{Score}_{\operatorname{Plu}}(P|_{\mathcal{A} \setminus B}, a) = \sum_{1 \le s \le t} \gamma_s |T_s|,$$

which is divisible by GCD(W(B)).

The  $\Leftarrow$  direction is proved in the following steps.

Step 1 for  $\Leftarrow$ . For any  $B \subseteq A$  and any pair of alternatives  $a, b \in A \setminus B$ , in this step we define an *m*!-profile  $P_{B,a,b}$  as the building block.

**Definition 24** ( $P_{B,a,b}$ ). For any pair of different alternatives a, b and any  $B \subsetneq (\mathcal{A} \setminus \{a, b\})$ ,  $P_{B,a,b}$  is obtained from  $\mathcal{L}(\mathcal{A})$  as follows. For each  $B^*$  such that  $B \subseteq B^* \subseteq (\mathcal{A} \setminus \{a, b\})$ ,

If |B<sup>\*</sup> \B| is an even number, then we replace [B<sup>\*</sup> ≻ b ≻ a ≻ others] by [B<sup>\*</sup> ≻ a ≻ b ≻ others], where "others" represents the other alternatives, and alternatives in B<sup>\*</sup> and "others" are ranked w.r.t. the lexicographic order.

• If  $|B^* \setminus B|$  is an odd number, then we replace  $[B^* \succ a \succ b \succ others]$  by  $[B^* \succ b \succ a \succ others]$ .

An example is shown in Table 3, where only the differences between  $\mathcal{L}(\mathcal{A})$  and  $P_{\{1\},2,3}$  are shown.

$B^*$	{1}	$\{1,4\}$	$\{1, 5\}$	$\{1, 4, 5\}$
$\mathcal{L}(\mathcal{A})$	$1 \succ 3 \succ 2 \succ 4 \succ 5$	$1 \succ 4 \succ 2 \succ 3 \succ 5$	$1 \succ 5 \succ 2 \succ 3 \succ 4$	$1 \succ 4 \succ 5 \succ 3 \succ 2$
$P_{\{1\},2,3}$	$1 \succ 2 \succ 3 \succ 4 \succ 5$	$1 \succ 4 \succ 3 \succ 2 \succ 5$	$1 \succ 5 \succ 3 \succ 2 \succ 4$	$1 \succ 4 \succ 5 \succ 2 \succ 3$

Table 3: Differences between  $\mathcal{L}(\mathcal{A})$  and  $P_{\{1\},2,3}$  for  $m = 4, B = \{1\}, a = 2, and b = 3$ .

We now prove that  $P_{B,a,b}$  satisfies two properties described in the following claim. The first property states that for any  $B' \neq B$ , after B' is removed, the plurality scores of the remaining alternatives in  $P_{B,a,b}|_{A \setminus B'}$  are equal. The second property states that after B is removed, the plurality score of a is increased by 1, the plurality score of b is decreased by 1, and the plurality scores of other alternatives are the same, after B is removed from  $P_{B,a,b}$ .

Claim 9 (Properties of  $P_{B,a,b}$ ). Let  $P_{B,a,b}$  denote the profile defined in Definition 24,

(i) for any  $B' \subsetneq A$  with  $B' \neq B$  and any  $c \in A \setminus (B \cup \{a, b\})$ , we have

$$Score_{Plu}(P_{B,a,b}|_{\mathcal{A}\setminus B'}, c) = Score_{Plu}(P_{B,a,b}|_{\mathcal{A}\setminus B'}, a) = Score_{Plu}(P_{B,a,b}|_{\mathcal{A}\setminus B'}, b)$$

(ii) for any  $c \in \mathcal{A} \setminus (B \cup \{a, b\})$ , we have

$$Score_{Plu}(P_{B,a,b}|_{\mathcal{A}\backslash B}, c) = Score_{Plu}(P_{B,a,b}|_{\mathcal{A}\backslash B}, a) - 1 = Score_{Plu}(P_{B,a,b}|_{\mathcal{A}\backslash B}, b) + 1$$

*Proof.* We first verify that (i) holds after  $B' \subsetneq \mathcal{A}$  is removed, for the following cases of B'.

- If  $B \not\subseteq B'$ , then after B' is removed, at least one alternative in B is not removed. According to the definition of  $P_{B,a,b}$  (Definition 24), the plurality scores of all remaining alternatives are the same, which verifies (i).
- If  $a \in B'$ , then notice that  $P_{B,a,b}$  only switch positions of a and b in some rankings where they are adjacent. Therefore, after B' is removed, which means that a is removed, (i) holds.
- Similarly, we can show that if  $b \in B'$ , then (i) holds.

The only remaining case is  $B \subseteq B' \subseteq (\mathcal{A} \setminus \{a, b\})$ . Next, we prove that assuming  $B \subseteq B' \subseteq (\mathcal{A} \setminus \{a, b\})$ , (ii) holds if and only if B' = B, and otherwise (i) holds (i.e., when  $B \subsetneq B' \subseteq (\mathcal{A} \setminus \{a, b\})$ ).

First, for any  $c \in \mathcal{A} \setminus (B' \cup \{a, b\})$ , notice that the difference between  $P_{B,a,b}$  and  $\mathcal{L}(\mathcal{A})$  only affects the plurality scores of a and b after B' is removed. Therefore, for any B', we have

$$\operatorname{Score}_{\operatorname{Plu}}(P_{B,a,b}|_{\mathcal{A}\backslash B'},c) = \operatorname{Score}_{\operatorname{Plu}}(\mathcal{L}(\mathcal{A})|_{\mathcal{A}\backslash B'},c)$$

Next, we calculate the difference between the plurality score of a in  $P_{B,a,b}$  after B' is removed and the plurality score of a in  $\mathcal{L}(\mathcal{A})$  after B' is removed. This is done by examining the effect of switching the (adjacent) locations of a and b in some votes as described in Definition 24. More precisely, for every  $B^*$  with  $B \subseteq B^* \subseteq (\mathcal{A} \setminus \{a, b\})$ , we have the following observations.

• If  $B^* \not\subseteq B'$ , then *a* is not ranked in the top positions of  $[B^* \succ a \succ b \succ$  others] or  $[B^* \succ b \succ a \succ$  others] after *B'* is removed. Therefore, the difference between  $P_{B,a,b}$  and  $\mathcal{L}(\mathcal{A})$  in the vote that corresponds to  $B^*$  does not affect the difference in the plurality scores of *a* in  $P_{B,a,b}$  and  $\mathcal{L}(\mathcal{A})$  after *B'* is removed.

- If  $B^* \subseteq B'$  and  $|B^* \setminus B|$  is an even number, then according to the definition of  $P_{B,a,b}$  (Definition 24),  $P_{B,a,b}$  replaces  $[B^* \succ b \succ a \succ \text{others}]$  in  $\mathcal{L}(\mathcal{A})$  by  $[B^* \succ a \succ b \succ \text{others}]$ . Notice that after B' is removed, b is ranked at the top position in the former ranking and a is ranked at the top position in the latter ranking. Therefore, the plurality score of a (respectively, b) is increased (respectively, decreased) by 1 in  $P_{B,a,b}$  after B' is removed, compared to the plurality score of a (respectively, b) in  $\mathcal{L}(\mathcal{A})$ .
- Similarly, if  $B^* \subseteq B'$  and  $|B^* \setminus B|$  is an odd number, then the plurality score of a (respectively, b) is decreased (respectively, increased) by 1 in  $P_{B,a,b}$  after B' is removed, compared to the plurality score of a (respectively, b) in  $\mathcal{L}(\mathcal{A})$ .

Therefore, we have the following calculation.

$$\operatorname{Score}_{\operatorname{Plu}}(P_{B,a,b}|_{\mathcal{A}\backslash B'},a) - \operatorname{Score}_{\operatorname{Plu}}(\mathcal{L}(\mathcal{A})|_{\mathcal{A}\backslash B'},a)$$
$$= \sum_{B^*:B\subseteq B^*\subseteq B'}(-1)^{|B^*\backslash B|} = \sum_{i=0}^{|B'|-|B|}(-1)^i \binom{|B'|-|B|}{i} = \begin{cases} 1 & \text{if } B'=B\\ 0 & \text{otherwise} \end{cases}$$

Similarly, we have the following calculation for b.

$$\operatorname{Score}_{\operatorname{Plu}}(P_{B,a,b}|_{\mathcal{A}\setminus B'}, b) - \operatorname{Score}_{\operatorname{Plu}}(\mathcal{L}(\mathcal{A})|_{\mathcal{A}\setminus B'}, b) = -\sum_{B^*: B\subseteq B^*\subseteq B'} (-1)^{|B^*\setminus B|} = \begin{cases} -1 & \text{if } B' = B\\ 0 & \text{otherwise} \end{cases}$$

It follows that (ii) holds when B' = B, otherwise (i) holds. This proves Claim 9.

Step 2 for  $\Leftarrow$ . For every  $B \subsetneq \mathcal{A}$ , we use  $P_{B,a,b}$  to fine-tune the plurality scores of alternatives when B is removed, so that the weak order among them becomes W(B). For every  $0 \le n' \le m! - 1$ such that for every  $B \subsetneq \mathcal{A}$ ,  $\operatorname{GCD}(W(B)) \mid n'$ , we construct a profile  $P_{n'}$  such that  $|P_{n'}| \equiv n'$ (mod m!) and  $\operatorname{PS}_{\mathrm{STV}}(P_{n'}) = W$  in the following steps.

- Step 2.1. Let  $n' = \lfloor \frac{n}{m!} \rfloor \times m!$ . Initialize  $P_{n'}$  to be n' copies of  $[1 \succ 2 \succ \cdots \succ m]$ .
- Step 2.2. For every  $B \subsetneq \mathcal{A}$ , let  $a_B \in (\mathcal{A} \setminus B)$  denote an arbitrary alternative. For every  $b \in \mathcal{A} \setminus (B \cup \{a_B\})$ , we add  $\operatorname{Score}_{\operatorname{Plu}}(P_{B,a_B,b}|_{\mathcal{A} \setminus B}, b)$  copies of  $P_{B,a_B,b}$  to  $P_{n'}$ , which can be seen as transferring the plurality score of b after to  $a_B$ . After this step, for every  $B \subsetneq \mathcal{A}$  and every  $b \in \mathcal{A} \setminus (B \cup \{a_B\})$ , we have:

$$\operatorname{Score}_{\operatorname{Plu}}(P_{n'}|_{\mathcal{A}\setminus B}, a_B) = \operatorname{Score}_{\operatorname{Plu}}(P_{n'}|_{\mathcal{A}\setminus B}, b) + n'$$

• Step 2.3. For every  $B \subsetneq \mathcal{A}$ , we add multiple copies of  $P_{B,a,b}$  for certain combinations of (a, b) such that the plurality scores for the same tier are the same, while the order between tiers may not be consistent with W(B). Notice that before Step 2.3, the plurality scores of alternatives in each tier are already the same, except the tier that contains the distinguished alternative  $a_B$  defined in Step 2.3. Given  $B \subsetneq \mathcal{A}$ , let  $T_1 \succ T_2 \succ \cdots \succ T_t$  denote the tier representation of W(B). Following the definition of GCD(W(B)), there exists t integers  $\eta_1, \ldots, \eta_t$  such that

$$GCD(W(B)) = \sum_{s=1}^{t} \eta_s \times |T_s|$$

For every  $1 \leq s \leq t$ , and every  $b \in T_s$ , we add the following profiles to  $P_{n'}$ , where we let  $P_{B,a_B,a_B} = \emptyset$  for convenience.

- $\begin{array}{l} \mbox{ if } \eta_t \geq 0, \mbox{ then we add } \frac{\eta_t n'}{\operatorname{GCD}(W(B))} \mbox{ copies of } P_{B,a_B,b}; \\ \mbox{ if } \eta_t < 0, \mbox{ then we add } \frac{\eta_t n'}{\operatorname{GCD}(W(B))} \mbox{ copies of } P_{B,b,a_B}. \end{array}$
- Step 2.4. Finally, we will add multiple copies of  $P_{B,a,b}$  for certain combinations of (a, b) to  $P_{n'}$  so that the order between tiers are the same as in W. Formally, for every  $B \subsetneq \mathcal{A}$  such that the tier representation of W(B) is  $T_1 \succ \cdots \succ T_t$ , and every  $1 \le s \le t-1$ , we add  $C_s$  copies of  $\bigcup_{a \in T_s, b \in T_t} P_{B,a,b}$  to  $P_{n'}$ , where  $C_1, \ldots, C_t 1$  are constants to guarantee W(B). This can be achieved because the effect of  $\bigcup_{a \in T_s, b \in T_t} P_{B,a,b}$  to  $P_{n'}$  increases the plurality score of every alternative in  $T_s$  by  $|T_t|$  and reduces the plurality score of every alternative in  $T_t$  by  $|T_s|$ , after B is removed.

It follows the construction that  $|P_{n'}| \equiv n' \pmod{m!}$  and  $\operatorname{PS}_{\operatorname{STV}}(P_{n'}) = W$ .

Step 3 for  $\Leftarrow$ . Let  $N = \max_{1 \le n' \le m! = 1} P_{n'}$ . For any n > N, let  $n' = n \pmod{m!}$ , and

$$P = P_{n'} + \frac{n - n'}{m!} \times \mathcal{L}(\mathcal{A})$$

It is not hard to verify that P is an n-profile and  $PS_{STV}(P) = W$ , which proves the  $\Leftarrow$  part of Lemma 3, and therefore completes the proof of the STV part of Lemma 3.

**Proof for Coombs.** Notice that Coombs uses veto in each round, and the veto score of a ranking is the plurality score of the reverse ranking. In light of this connection, the proof is similar to the STV part and the main difference is that for Coombs, we use  $P'_{B,a,b}$  that is defined from  $P_{B,a,b}$  (Definition 24) by reversing all rankings. This completes the proof of Lemma 3.

### D.3 Proof of Theorem 5

**Theorem 5.** (Smoothed likelihood of ties: STV). Let  $\mathcal{M} = (\Theta, \mathcal{L}(\mathcal{A}), \Pi)$  be a strictly positive and closed single-agent preference model. There exists  $N \in \mathbb{N}$  such that for any  $2 \leq k \leq m$  and any n > N,

$$\begin{split} \widetilde{Tie}_{\Pi}^{\max}(STV,k,n) &= \begin{cases} 0 & \text{if } \forall W \in \mathcal{W}_n, |STV(W)| \neq k \\ \exp(-\Theta(n)) & \text{otherwise if } \mathcal{W}_{STV,k,n}^{\Pi} = \emptyset \\ \Theta\left(n^{-\frac{w_{\min}}{2}}\right) & \text{otherwise} \end{cases} \\ \widetilde{Tie}_{\Pi}^{\min}(STV,k,n) &= \begin{cases} 0 & \text{if } \forall W \in \mathcal{W}_n, |STV(W)| \neq k \\ \exp(-\Theta(n)) & \text{otherwise if } \exists \pi \in CH(\Pi) \text{ s.t. } \mathcal{W}_{STV,k,n}^{\pi} = \emptyset \\ \Theta\left(n^{-\frac{w_{\min}}{2}}\right) & \text{otherwise} \end{cases} \end{split}$$

*Proof.* Like in the proof of Theorem 3, the theorem is proved by modeling the set of profiles with k winners as the union of constantly many polyhedra, then applying Theorem 2. As in the proof of Lemma 3, we first present the proof for r = STV, then comment on how to modify the proof for Coombs.

**Proof for STV.** The proof proceeds in the following three steps. In Step 1, for each PUT structure W, we define a polyhedron  $\mathcal{H}^W$  that characterizes the profiles whose PUT structures are W. In Step 2, we prove properties about  $\mathcal{H}^W$ , in particular  $\dim(\mathcal{H}^W_{\leq 0}) = m! - \operatorname{Ties}(W)$ . In Step 3 we formally apply Theorem 2 to  $\mathcal{C} = \bigcup_{W \in \mathcal{W}_n: |\mathrm{STV}(W)| = k} \mathcal{H}^W$ .

Step 1 for STV: Define  $\mathcal{H}^W$ . We first define the vectors that will be used in the construction.

**Definition 25** (Pairwise difference vector for STV). For any pair of different alternatives a, b and any  $B \in \mathcal{A} \setminus \{a, b\}$ , we let  $Pair_{B,a,b}$  denote the vector such that for any  $R \in \mathcal{L}(\mathcal{A})$ , the R-th component of  $Pair_{B,a,b}$  is  $Score_{Plu}(R|_{\mathcal{A} \setminus B}, a) - Score_{Plu}(R|_{\mathcal{A} \setminus B}, b)$ .

In words,  $\operatorname{Pair}_{B,a,b}$  represents the difference between the plurality score of a and the plurality score of b after B is removed. We now use these vectors to construct a polyhedron that corresponds to a given PUT structure W.

**Definition 26**  $(\mathcal{H}^W)$ . Given a PUT structure W, for any  $B \subsetneq \mathcal{A}$ ,  $\mathcal{H}^W$  is characterized by the following constraints  $\mathbf{A}^W \cdot (\vec{x}_{\mathcal{A}})^\top \leq (\vec{b})^\top$ : for any pair of alternatives a  $\succ_{W(B)}$  b, there is an inequality  $\operatorname{Pair}_{B,b,a} \cdot \vec{x}_{\mathcal{A}} \leq -1$ ; and for any pair of alternatives  $a \equiv_{W_B} b$ , there is an inequality  $\operatorname{Pair}_{B,b,a} \cdot \vec{x}_{\mathcal{A}} \leq 0$ .

Step 2 for STV: Prove properties about  $\mathcal{H}^W$ . We prove the following claim about  $\mathcal{H}^W$ .

Claim 10 (Properties of  $\mathcal{H}^W$ ). For any PUT structure W, we have:

- (i) for any profile P,  $Hist(P) \in \mathcal{H}^W$  if and only if  $PS_{STV}(P) = W$ ;
- (ii) for any  $\vec{x} \in \mathbb{R}^{m!}$ ,  $\vec{x} \in \mathcal{H}_{\leq 0}^{W}$  if and only if W refines  $PS_{STV}(\vec{x})$ ;
- (iii)  $\dim(\mathcal{H}^W_{\leq 0}) = m! Ties(W).$

*Proof.* (i) and (ii) follow after the definition of  $\mathcal{H}^W$ . To prove part (iii), again, according to equation (9) on page 100 in [57], it suffices to prove that the rank of the essential equalities of  $\mathbf{A}^W$ , denoted by  $\mathbf{A}^=$ , is Ties(W). We first prove that

$$\mathbf{A}^{=} = \{ \operatorname{Pair}_{B,a,b} : \forall B \subsetneq \mathcal{A} \text{ and } a \equiv_{W(B)} b \},$$
(24)

where  $a \equiv_{W(B)} b$  means that a and b are tied in W(B). Clearly  $\mathbf{A}^{=}$  contains the right hand side of Equation (24). By Lemma 3, there exists an (m!)-profile  $P^*$  such that  $\mathrm{PS}_{\mathrm{STV}}(P^*) = W$ . Therefore,  $\mathrm{Hist}(P^*)$  is an inner point of  $\mathcal{H}^W$  in the sense that all inequalities not mentioned in the right hand side of Equation (24) are strict under  $\mathrm{Hist}(P^*)$ , which means that Equation (24) holds.

We now prove that  $\operatorname{Rank}(\mathbf{A}^{=}) = \operatorname{Ties}(W)$ . For any  $B \subsetneq \mathcal{A}$ , let  $W(B) = T_1 \succ \cdots \succ T_s$  denote the tier presentation of W(B). Let  $\mathbf{A}_B$  be the sub-matrix of  $\mathbf{A}^{=}$  that consists of  $\operatorname{Pair}_{B,b,a}$  for all pairs of alternatives  $a \equiv_{W(B)} b$ . It is not hard to check that  $\operatorname{Rank}(\mathbf{A}_B) \leq \operatorname{Ties}(W)$ , because for any  $T_i$ , each vector in { $\operatorname{Pair}_{B,a,b} : a, b \in T_i$ } can be represented as the linear combination of a subset of  $|T_i| - 1$  linear orders. This means that  $\operatorname{Rank}(\mathbf{A}^{=}) \leq \sum_{B \subseteq \mathcal{A}} \operatorname{Ties}(W(B)) = \operatorname{Ties}(W)$ .

 $\operatorname{Rank}(\mathbf{A}^{=}) \geq \operatorname{Ties}(W)$  follows after Lemma 3. More precisely, suppose for the sake of contradiction that  $\operatorname{Rank}(\mathbf{A}^{=}) < \operatorname{Ties}(W)$  and let A denote an arbitrary row basis of  $\mathbf{A}^{=}$ . Due to the pigeon hole principle, there exists  $B \subsetneq A$  such that A contains no more than  $\operatorname{Ties}(W(B))$  rows in  $\mathbf{A}_B$ , which means that there exists a tier  $T_i$  such that A contains no more than  $|T_i| - 2$  rows in  $\{\operatorname{Pair}_{B,a,b} : a, b \in T_i\}$ . This means that  $T_i$  can be partitioned into two sets  $\{a_1, \ldots, a_{t_1}\}$  and  $\{b_1, \ldots, b_{t_2}\}$  such that A does not contain  $\operatorname{Pair}_{B,a_{i_1},b_{i_2}}$  for any combination of  $1 \leq i_1 \leq t_1$  and  $1 \leq i_2 \leq t_2$ . By Lemma 3, there exists an (m!)-profile P such that

- for all  $B' \neq B$ , all alternatives are tied after B' is removed, and
- $\operatorname{PS}_r(P)(B)$  consists of two tiers: the first tier is  $T_1 \cup \cdots \cup T_{i-1}\{a_1, \ldots, a_{t_1}\}$  and the second tier is  $\{b_1, \ldots, b_{t_2}\} \cup T_{i+1} \cup \cdots \cup T_t$ .

It is not hard to check that

$$A \cdot (\operatorname{Hist}(P))^{\top} = \left(\vec{0}\right)^{\top}$$
 but  $\operatorname{Pair}_{B,a_1,b_1} \cdot \operatorname{Hist}(P) \neq 0.$ 

This means that  $\operatorname{Pair}_{B,a_1,b_1} \in \mathbf{A}^=$  is not a linear combination of rows in A, which contradicts the assumption that A is a basis of  $\mathbf{A}^=$ .

Therefore,  $\operatorname{Rank}(\mathbf{A}^{=}) = \operatorname{Ties}(W)$ . This completes the proof of Claim 10.

Step 3 for STV: Apply Theorem 2. Let  $\mathcal{C} = \bigcup_{W \in \mathcal{W}_n: |STV(W)|=k} \mathcal{H}^W$ . It follows that for any profile P, |STV(P)| = k if and only if  $\text{Hist}(P) \in \mathcal{C}$ . Therefore, for any n and any  $\vec{\pi} \in \Pi^n$ , we have:

$$\Pr_{P \sim \vec{\pi}}(|\mathrm{STV}(P)| = k) = \Pr(\vec{X}_{\vec{\pi}} \in \mathcal{C})$$

Recall that for all PUT structure W over  $\mathcal{A}$ , by Claim 10 (iii) we have  $\dim(\mathcal{H}^W) = m! - \operatorname{Ties}(P) + 1$ . This means that  $\alpha_n = m! - \ell_{\min}$  and  $\beta_n = m! - \ell_{\min}$  when  $W \in \mathcal{W}_n$ .

Therefore, to prove Theorem 5, it suffices to prove that the conditions for the 0, exponential, and polynomial cases in Theorem 5 are equivalent to the conditions for the 0, exponential, and polynomial cases in Theorem 2 (applied to C and  $\Pi$ ), respectively. This follows a similar reasoning as in Step 3 of the proof of Theorem 3 combined with Claim 10 (ii). This completes the proof of the STV part of Theorem 5.

**Proof for Coombs.** As in the proof of the Coombs part of Lemma 3, notice that Coombs uses veto in each round, and the veto score of a ranking is the plurality score of the reverse ranking. Therefore, the proof is similar to the STV part and the main difference is that for Coombs, when defining the pairwise score difference vectors (Definition 25), we use the veto rule instead of the plurality rule. This completes the proof of Lemma 3.

### D.4 Proof of Proposition 6

**Proposition 6.** (Max smoothed likelihood of ties: STV and Coombs). Let  $r \in \{STV, Coombs\}$  and let  $\mathcal{M} = (\Theta, \mathcal{L}(\mathcal{A}), \Pi)$  be a strictly positive and closed single-agent preference model with  $\pi_{uni} \in CH(\Pi)$ . For any  $2 \leq k \leq m$  and any  $n \in \mathbb{N}$ ,

$$\widetilde{Tie}_{\Pi}^{\max}(r,k,n) = \begin{cases} \Theta(n^{-\frac{k-1}{2}}) & \text{if } \begin{cases} (1)m \ge 4, \text{ or } \\ (2)m = 3 \text{ and } k = 2, \text{ or } \\ (3)m = k = 3 \text{ and } (2 \mid n \text{ or } 3 \mid n) \\ 0 & \text{ otherwise } (i.e., m = k = 3, 2 \nmid n, \text{ and } 3 \nmid n) \end{cases}$$

*Proof.* According to Theorem 5, we will prove that (1) the exponential case does not happen, and (2)  $w_{\min} = k - 1$  in the polynomial cases.

To see (1), notice that if the 0 case of Theorem 5 does not hold, then there exists  $W \in \mathcal{W}_n$ such that  $|\mathrm{STV}(W)| = k$ . Also notice that  $\pi_{\mathrm{uni}} \in \mathrm{CH}(\Pi)$  and  $\mathrm{PS}_{\mathrm{STV}}(\pi_{\mathrm{uni}})$  is the PUT structure where all alternatives are tied after any  $B \subsetneq \mathcal{A}$  is removed, which means that any PUT structure is a refinement of  $\mathrm{PS}_{\mathrm{STV}}(\pi_{\mathrm{uni}})$ . Therefore,  $\mathcal{W}_{r,k,n}^{\Pi} \neq \emptyset$ , which means that the exponential case of Theorem 5 does not hold.

To see (2), notice that for any PUT structure W, the outgoing edges of any node labeled  $B \subsetneq \mathcal{A}$ with  $|B| \leq m-2$ , the number of outgoing edges of B is Ties(W(B)) + 1. Therefore, Ties(W) is at least the total number of nodes in W that do not have outgoing edges minus one. It follows that for any PUT structure W with |STV(W)| = k, we must have  $Ties(W) \ge k - 1$ , which means that  $w_{mm} \ge k - 1$ .

In the sequel, we will prove  $w_{\rm mm} = k - 1$  by construction for the polynomial cases described in Proposition 6. More precisely, we will prove that there exists an *n*-profile *P* such that  $\text{Ties}(\text{PS}_{\text{STV}}(P)) = k - 1$  and  $|\text{STV}(\text{PS}_{\text{STV}}(P))| = k$ . In light of Lemma 3, it suffices to construct a PUT structure *W* such that

(1) 
$$\operatorname{Ties}(W) = k - 1$$
, (2)  $|\operatorname{STV}(W)| = k$ , and (3) for every  $B \subsetneq \mathcal{A}, \operatorname{GCD}(W(B)) \mid n$ 

 $\Theta(n^{-\frac{k-1}{2}})$  subcase (1):  $m \ge 4$ . When  $m \ge 4$ , we let W denote the PUT structure such that

$$W(\emptyset) = k \succ k+1 \succ \dots \succ m \succ \{1, \dots, k-1\}$$
$$W(\{1\}) = 2 \succ \dots \succ k-2 \succ k+1 \succ \dots \succ m \succ \{k-1, k\},$$

and W(B) for any other B is a linear order so that in the graph representation of W, there are k nodes without outgoing edges, whose winners are  $\{1, \ldots, k\}$ . See Figure 13 (a) for an example of m = k = 4.



Figure 13: PUT structures for STV.

It is not hard to verify that Ties(W) = k - 1 and |STV(W)| = k. Notice that for every  $B \subsetneq A$ , there exists a tier in W(B) that consists of a single alternatives, which means that GCD(W(B)) = 1 and therefore,  $W \in \mathcal{W}_n$ . This proves the proposition for the  $m \ge 4$  case.

 $\Theta(n^{-\frac{k-1}{2}})$  subcase (2): m = 3 and k = 2. The m = 3 and k = 2 case is proved by the PUT structure illustrated in Figure 13 (b).

 $\Theta(n^{-\frac{k-1}{2}})$  subcase (3): m = k = 3 and  $(2 \mid n \text{ or } 3 \mid n)$ . The case where m = k = 3 and  $2 \mid n$  is proved by the PUT structure  $W_1$  illustrated in Figure 14 (a). The case where m = k = 3 and  $3 \mid n$  is proved by the PUT structure  $W_2$  illustrated in Figure 14 (b).


Figure 14: PUT structures for STV.

**The 0 case.** When m = k = 3,  $2 \nmid n$ , and  $3 \nmid n$ , suppose for the sake of contradiction there exists an *n*-profile *P* such that |STV(P)| = 3. Let  $W = \text{PS}_{\text{STV}}(P)$ . Then, either W(B) is a tie among the three alternatives for some  $B \subsetneq \mathcal{A}$  (which means that GCD(()W(B)) = 3, and therefore  $3 \mid n$ ), or W(B) is a tie among two alternatives for some |B| = 1 (which means that GCD(()W(B)) = 2, and therefore  $2 \mid n$ ). Either case leads to a contradiction. Therefore,  $\mathcal{W}_{r,k,n}^{\Pi} = \emptyset$  in this case.  $\Box$ 

## E Experimental Results for All Rules



Figure 15: Percentage of tied profiles under IC.



Figure 16: Histograms of number of candidates and number of voters in the 307 Preflib SOC data studied in this paper.