Sinkhorn Natural Gradient for Generative Models

Zebang Shen* Zhenfu Wang[†] Alejandro Ribeiro* Hamed Hassani* *Department of Electrical and Systems Engineering [†]Department of Mathematics University of Pennsylvania

{zebang@seas,zwang423@math,aribeiro@seas,hassani@seas}.upenn.edu

Abstract

We consider the problem of minimizing a functional over a parametric family of probability measures, where the parameterization is characterized via a pushforward structure. An important application of this problem is in training generative adversarial networks. In this regard, we propose a novel Sinkhorn Natural Gradient (SiNG) algorithm which acts as a steepest descent method on the probability space endowed with the Sinkhorn divergence. We show that the Sinkhorn information matrix (SIM), a key component of SiNG, has an explicit expression and can be evaluated accurately in complexity that scales logarithmically with respect to the desired accuracy. This is in sharp contrast to existing natural gradient methods that can only be carried out approximately. Moreover, in practical applications when only Monte-Carlo type integration is available, we design an empirical estimator for SIM and provide the stability analysis. In our experiments, we quantitatively compare SiNG with state-of-the-art SGD-type solvers on generative tasks to demonstrate its efficiency and efficacy of our method.

1 Introduction

Consider the minimization of a functional \mathcal{F} over a parameterized family probability measures $\{\alpha_{\theta}\}$:

$$\min_{\theta \in \Theta} \left\{ F(\theta) := \mathcal{F}(\alpha_{\theta}) \right\},\tag{1}$$

where $\Theta \subseteq \mathbb{R}^d$ is the feasible domain of the parameter θ . We assume that the measures α_{θ} are defined over a common ground set $\mathcal{X} \subseteq \mathbb{R}^q$ with the following structure: $\alpha_{\theta} = T_{\theta \sharp} \mu$, where μ is a fixed and known measure and T_{θ} is a push-forward mapping. More specifically, μ is a simple measure on a latent space $\mathcal{Z} \subseteq \mathbb{R}^{\bar{q}}$, such as the standard Gaussian measure $\mu = \mathcal{N}(\mathbf{0}_{\bar{q}}, \mathbf{I}_{\bar{q}})$, and the parameterized map $T_{\theta} : \mathcal{Z} \to \mathcal{X}$ transforms the measure μ to α_{θ} . This type of push-forward parameterization is commonly used in deep generative models, where T_{θ} represents a neural network parametrized by weights θ [Goodfellow et al., 2014, Salimans et al., 2018, Genevay et al., 2018]. Consequently, methods to efficiently and accurately solve problem (1) are of great importance in machine learning.

The de facto solvers for problem (1) are generic nonconvex optimizers such as Stochastic Gradient Descent (SGD) and its variants, Adam [Kingma and Ba, 2014], Amsgrad [Reddi et al., 2019], RMSProp [Hinton et al.], etc. These optimization algorithms directly work on the parameter space and are agnostic to the fact that α_{θ} 's are probability measures. Consequently, SGD type solvers suffer from the complex optimization landscape induced from the neural-network mappings T_{θ} .

An alternative to SGD type methods is the natural gradient method, which is originally motivated from Information Geometry [Amari, 1998, Amari et al., 1987]. Instead of simply using the Euclidean structure of the parameter space Θ in the usual SGD, the natural gradient method endows the parameter space with a "natural" metric structure by pulling back a known metric on the probability space and then searches the steepest descent direction of $F(\theta)$ in the "curved" neighborhood of θ . In particular, the natural gradient update is invariant to reparametrization. This allows natural gradient to avoid the undesirable saddle point or local minima that are artificially created by the highly nonlinear maps T_{θ} . The classical Fisher-Rao Natural Gradient (FNG) [Amari, 1998] as well as its many variants [Martens and Grosse, 2015, Thomas et al., 2016, Song et al., 2018] endows the probability space with the KL divergence and admits update direction in closed form. However, the update rules of these methods all require the evaluation of the score function of the variable measure. Leaving aside its existence, this quantity is in general difficult to compute for push-forward measures, which limits the application of FNG type methods in the generative models. Recently, Li and Montúfar [2018] propose to replace the KL divergence in FNG by the Wasserstein distance and propose the Wasserstein Natural Gradient (WNG) algorithm. WNG shares the merit of reparameterization invariance as FNG while avoiding the requirement of the score function. However, the Wasserstein information matrix (WIM) is very difficult to compute as it does not attain a closed form expression when the dimension d of parameters is greater than 1, rendering WNG impractical.

Following the line of natural gradient, in this paper, we propose Sinkhorn Natural Gradient (SiNG), an algorithm that performs the steepest descent of the objective functional \mathcal{F} on the probability space with the Sinkhorn divergence as the underlying metric. Unlike FNG, SiNG requires only to sample from the variable measure α_{θ} . Moreover, the Sinkhorn information matrix (SIM), a key component in SiNG, can be computed in logarithmic time in contrast to WIM in WNG. Concretely, we list our contributions as follows:

- 1. We derive the Sinkhorn Natural Gradient (SiNG) update rule as the exact direction that minimizes the objective functional \mathcal{F} within the Sinkhorn ball of radius ϵ centered at the current measure. In the asymptotic case $\epsilon \rightarrow 0$, we show that the SiNG direction only depends on the Hessian of the Sinkhorn divergence and the gradient of the function F, while the effect of the Hessian of F becomes negligible. Further, we prove that SiNG is invariant to reparameterization in its continuous-time limit (i.e. using the infinitesimal step size).
- 2. We explicitly derive the expression of the Sinkhorn information matrix (SIM), i.e. the Hessian of the Sinkhorn divergence with respect to the parameter θ . We then show the SIM can be computed using logarithmic (w.r.t. the target accuracy) function operations and integrals with respect to α_{θ} .
- 3. When only Monte-Carlo integration w.r.t. α_{θ} is available, we propose to approximate SIM with its empirical counterpart (eSIM), i.e. the Hessian of the empirical Sinkhorn divergence. Further, we prove stability of eSIM. Our analysis relies on the fact that the Fréchet derivative of Sinkhorn potential with respect to the parameter θ is continuous with respect to the underlying measure μ . Such result can be of general interest.

In our experiments, we pretrain the discriminators for the celebA and cifar10 datasets. Fixing the discriminator, we compare SiNG with state-of-the-art SGD-type solvers in terms of the generator loss. The result shows the remarkable superiority of SiNG in both efficacy and efficiency.

Notation: Let $\mathcal{X} \subseteq \mathbb{R}^q$ be a compact ground set. We use $\mathcal{M}_1^+(\mathcal{X})$ to denote the space of probability measures on \mathcal{X} and use $\mathcal{C}(\mathcal{X})$ to denote the family of continuous functions mapping from \mathcal{X} to \mathbb{R} . For a function $f \in \mathcal{C}(\mathcal{X})$, we denote its L^{∞} norm by $||f||_{\infty} := \max_{x \in \mathcal{X}} |f(x)|$ and its gradient by ∇f . For a functional on general vector spaces, the Fréchet derivative is formally defined as follows. Let V and W be normed vector spaces, and $U \subseteq V$ be an open subset of V. A function $\mathcal{F} : U \to W$ is called Fréchet differentiable at $x \in U$ if there exists a bounded linear operator $A : V \to W$ such that

$$\lim_{\|h\|\to 0} \frac{\|\mathcal{F}(x+h) - \mathcal{F}(x) - Ah\|_W}{\|h\|_V} = 0.$$
(2)

If there exists such an operator A, it will be unique, so we denote $D\mathcal{F}(x) = A$ and call it the *Fréchet derivative*. From the above definition, we know that $D\mathcal{F}: U \to T(V, W)$ where T(V, W) is the family of bounded linear operators from V to W. Given $x \in U$, the linear map $D\mathcal{F}(x)$ takes one input $y \in V$ and outputs $z \in W$. This is denoted by $z = D\mathcal{F}(x)[y]$. We then define the operator norm of $D\mathcal{F}$ at x as $\|D\mathcal{F}(x)\|_{op} := \max_{h \in V} \frac{\|D\mathcal{F}(x)[h]\|_W}{\|h\|_V}$. Further, the second-order Fréchet derivative of \mathcal{F} is denoted as $D^2\mathcal{F}: U \to L^2(V \times V, W)$, where $L^2(V \times V, W)$ is the family of all continuous bilinear maps from V to W. Given $x \in U$, the bilinear map $D^2\mathcal{F}(x)$ takes two inputs $y_1, y_2 \in V$ and outputs $z \in W$. We denote this by $z = D^2\mathcal{F}(x)[y_1, y_2]$. If a function \mathcal{F} has multiple variables, we use $D_i f$ to denote the Fréchet derivative with its i^{th} variable and use $D_{ij}^2\mathcal{F}$ to denote the corresponding second-order terms. Finally, \circ denotes the composition of functions.

2 Related Work on Natural Gradient

The Fisher-Rao natural gradient (FNG) [Amari, 1998] is a now classical algorithm for the functional minimization over a class of parameterized probability measures. However, unlike SiNG, FNG as well as its many variants [Martens and Grosse, 2015, Thomas et al., 2016, Song et al., 2018] requires to evaluate the score function $\nabla_{\theta} \log p_{\theta}$ (p_{θ} denotes the p.d.f. of α_{θ}). Leaving aside its existence issue, the score function for the generative model α_{θ} is difficult to compute as it involves T_{θ}^{-1} , the inversion of the push-forward mapping, and det(JT_{θ}^{-1}), the determinant of the Jacobian of $T_{\theta}^{-1}(z)$. One can possibly recast the computation of the score function as a dual functional minimization problem over all continuous functions on \mathcal{X} [Essid et al., 2019]. However, such functional minimization problem itself is difficult to solve. As a result, FNG has limited applicability in our problem of interest.

Instead of using the KL divergence, Li and Montúfar [2018] propose to measure the distance between (discrete) probability distributions using the optimal transport and develop the Wasserstein Natural Gradient (WNG). WNG inherits FNG's merit of reparameterization invariance. However, WNG requires to compute the Wasserstein information matrix (WIM), which does not attain a closed form expression when d > 1, rendering WNG impractical [Li and Zhao, 2019, Li and Montúfar, 2020]. As a workaround, one can recast a single WNG step to a dual functional maximization problem via the Legendre duality. While itself remains challenging and can hardly be globally optimized, Li et al. [2019] simplify the dual subproblem by restricting the optimization domain to an affine space of functions (a linear combinations of several bases). Clearly, the quality of this solver depends heavily on the accuracy of this affine approximation. Alternatively, Arbel et al. [2019] restrict the dual functional optimization to a Reproducing Kernel Hilbert Space (RKHS). By adding two additional regularization terms, the simplified dual subproblem admits a closed form solution. However, in this way, the gap between the original WNG update and its kernelized version cannot be properly quantified without overstretched assumptions.

3 Preliminaries

We first introduce the entropy-regularized optimal transport distance and then its debiased version, i.e. the Sinkhorn divergence. Given two probability measures $\alpha, \beta \in \mathcal{M}_1^+(\mathcal{X})$, the entropy-regularized optimal transport distance $OT_{\gamma}(\alpha, \beta) : \mathcal{M}_1^+(\mathcal{X}) \times \mathcal{M}_1^+(\mathcal{X}) \to \mathbb{R}_+$ is defined as

$$\operatorname{OT}_{\gamma}(\alpha,\beta) = \min_{\pi \in \Pi(\alpha,\beta)} \langle c,\pi \rangle + \gamma \operatorname{KL}(\pi || \alpha \otimes \beta).$$
(3)

Here, $\gamma > 0$ is a fixed regularization parameter, $\Pi(\alpha, \beta)$ is the set of joint distributions over \mathcal{X}^2 with marginals α and β , and we use $\langle c, \pi \rangle$ to denote $\langle c, \pi \rangle = \int_{\mathcal{X}^2} c(x, y) d\pi(x, y)$. We also use $\operatorname{KL}(\pi || \alpha \otimes \beta)$ to denote the *Kullback-Leibler divergence* between the candidate transport plan π and the product measure $\alpha \otimes \beta$.

Note that $\operatorname{OT}_{\gamma}(\alpha, \beta)$ is not a valid metric as there exists $\alpha \in \mathcal{M}_{1}^{+}(\mathcal{X})$ such that $\operatorname{OT}_{\gamma}(\alpha, \alpha) \neq 0$ when $\gamma \neq 0$. To remove this bias, consider the *Sinkhorn divergence* $\mathcal{S}(\alpha, \beta) : \mathcal{M}_{1}^{+}(\mathcal{X}) \times \mathcal{M}_{1}^{+}(\mathcal{X}) \to \mathbb{R}_{+}$ introduced in Peyré et al. [2019]:

$$\mathcal{S}(\alpha,\beta) := \operatorname{OT}_{\gamma}(\alpha,\beta) - \frac{\operatorname{OT}_{\gamma}(\alpha,\alpha)}{2} - \frac{\operatorname{OT}_{\gamma}(\beta,\beta)}{2}, \tag{4}$$

which can be regarded as a *debiased version* of $OT_{\gamma}(\alpha, \beta)$. Since γ is fixed throughout this paper, we omit the subscript γ for simplicity. It has been proved that $S(\alpha, \beta)$ is nonnegative, bi-convex and metrizes the convergence in law for a compact \mathcal{X} and a Lipschitz metric *c* Peyré et al. [2019].

The Dual Formulation and Sinkhorn Potentials. The entropy-regularized optimal transport problem OT_{γ} , given in (3), is convex with respect to the joint distribution π : Its objective is a sum of a linear functional and the convex KL-divergence, and the feasible set $\Pi(\alpha, \beta)$ is convex. Consequently, there is no gap between the primal problem (3) and its Fenchel dual. Specifically, define

$$\mathcal{H}_2(f,g;\alpha,\beta) := \langle f,\alpha \rangle + \langle g,\beta \rangle - \gamma \langle \exp(\frac{1}{\gamma}(f \oplus g - c)) - 1, \alpha \otimes \beta \rangle, \tag{5}$$

where we denote $(f \oplus g)(x, y) = f(x) + g(y)$. We have

$$OT_{\gamma}(\alpha,\beta) = \max_{f,g\in\mathcal{C}(\mathcal{X})} \{\mathcal{H}_2(f,g;\alpha,\beta)\} = \langle f_{\alpha,\beta},\alpha\rangle + \langle g_{\alpha,\beta},\beta\rangle,\tag{6}$$

where $f_{\alpha,\beta}$ and $g_{\alpha,\beta}$, called the *Sinkhorn potentials* of $OT_{\gamma}(\alpha,\beta)$, are the maximizers of (6).

Training Adversarial Generative Models. We briefly describe how (1) captures the generative adversarial model (GAN): In training a GAN, the objective functional in (1) itself is defined through a maximization subproblem $\mathcal{F}(\alpha_{\theta}) = \max_{\xi \in \Xi} \mathcal{G}(\xi; \alpha_{\theta})$. Here $\xi \in \Xi \subseteq \mathbb{R}^{\overline{d}}$ is some dual adversarial variable encoding an adversarial discriminator or ground cost. For example, in the ground cost adversarial optimal transport formulation of GAN [Salimans et al., 2018, Genevay et al., 2018], we have $\mathcal{G}(\xi; \alpha_{\theta}) = \mathcal{S}_{c_{\xi}}(\alpha_{\theta}, \beta)$. Here, with a slight abuse of notation, $\mathcal{S}_{c_{\xi}}(\alpha_{\theta}, \beta)$ denotes the Sinkhorn divergence between the parameterized measure α_{θ} and a given target measure β . Notice that the symmetric ground cost c_{ξ} in $\mathcal{S}_{c_{\xi}}$ is no longer fixed to any pre-specified distance like ℓ_1 or ℓ_2 norm. Instead, c_{ξ} is encoded by a parameter ξ so that $\mathcal{S}_{c_{\xi}}$ can distinguish α_{θ} and β in an adaptive and adversarial manner. By plugging the above $\mathcal{F}(\alpha_{\theta})$ to (1), we recover the generative adversarial model proposed in [Genevay et al., 2018]:

$$\min_{\theta \in \Theta} \max_{\xi \in \Xi} \mathcal{S}_{c_{\xi}}(\alpha_{\theta}, \beta). \tag{7}$$

4 Methodology

In this section, we derive the Sinkhorn Natural Gradient (SiNG) algorithm as a steepest descent method in the probability space endowed with the Sinkhorn divergence metric. Specifically, SiNG updates the parameter θ^t by

$$\theta^{t+1} := \theta^t + \eta \cdot \mathbf{d}^t \tag{8}$$

where $\eta > 0$ is the step size and the update direction \mathbf{d}^t is obtained by solving the following problem. Recall the objective F in (1) and the Sinkhorn divergence S in (4). Let $\mathbf{d}^t = \lim_{\epsilon \to 0} \frac{\Delta \theta^t_{\epsilon}}{\sqrt{\epsilon}}$, where

$$\Delta \theta^t_{\epsilon} := \underset{\Delta \theta \in \mathbb{R}^d}{\operatorname{argmin}} F(\theta^t + \Delta \theta) \quad \text{s.t.} \quad \|\Delta \theta\| \leqslant \epsilon^{c_1}, \mathcal{S}(\alpha_{\theta^t + \Delta \theta}, \alpha_{\theta^t}) \leqslant \epsilon + \epsilon^{c_2}.$$
(9)

Here the exponent c_1 and c_2 can be arbitrary real satisfying $1 < c_2 < 1.5$, $c_1 < 0.5$ and $3c_1 - 1 \ge c_2$. Proposition 4.1 depicts a simple expression of \mathbf{d}^t . Before proceeding to derive this expression, we note that $\Delta \theta = 0$ globally minimizes the non-negative function $S(\alpha_{\theta^t + \Delta\theta}, \alpha_{\theta^t})$, which leads to the following first and second order optimality criteria:

$$\nabla_{\theta} \mathcal{S}(\alpha_{\theta}, \alpha_{\theta^{t}})_{|\theta=\theta^{t}} = 0 \quad \text{and} \quad \mathbf{H}(\theta^{t}) := \nabla_{\theta}^{2} \mathcal{S}(\alpha_{\theta}, \alpha_{\theta^{t}})_{|\theta=\theta^{t}} \ge 0.$$
(10)

This property is critical in deriving the explicit formula of the Sinkhorn natural gradient. From now on, the term $\mathbf{H}(\theta^t)$, which is a key component of SiNG, will be referred to as the *Sinkhorn information matrix (SIM)*.

Proposition 4.1. Assume that the minimum eigenvalue of $\mathbf{H}(\theta^t)$ is strictly positive (but can be arbitrary small) and that $\nabla^2_{\theta} F(\theta)$ and $\mathbf{H}(\theta)$ are continuous w.r.t. θ . The SiNG direction has the following explicit expression

$$\mathbf{d}^{t} = -\frac{\sqrt{2}}{\sqrt{\langle \mathbf{H}(\theta^{t})^{-1} \nabla_{\theta} F(\theta^{t}), \nabla_{\theta} F(\theta^{t}) \rangle}} \cdot \mathbf{H}(\theta^{t})^{-1} \nabla_{\theta} F(\theta^{t}).$$
(11)

Interestingly, the SiNG direction does not involve the Hessian of F. This is due to a Lagrangian-based argument that we sketch here. Note that the continuous assumptions on $\nabla^2_{\theta} F(\theta)$ and $\mathbf{H}(\theta)$ enable us to approximate the objective and the constraint in (9) via the second-order Taylor expansion.

Proof sketch for Proposition 4.1. The second-order Taylor expansion of the Lagrangian of (9) is

$$\bar{G}(\Delta\theta) = F(\theta^t) + \langle \nabla_{\theta}F(\theta^t), \Delta\theta \rangle + \frac{1}{2} \langle \nabla_{\theta}^2 F(\theta^t) \Delta\theta, \Delta\theta \rangle + \frac{\lambda}{2} \langle \mathbf{H}(\theta^t) \Delta\theta, \Delta\theta \rangle - \lambda\epsilon - \lambda\epsilon^{c_2},$$
(12)

where $\lambda \ge 0$ is the dual variable. Since the minimum eigenvalue of $\mathbf{H}(\theta^t)$ is strictly positive, for a sufficiently small ϵ , by taking $\lambda = \mathcal{O}(\frac{1}{\sqrt{\epsilon}})$, we have that $\mathbf{H}(\theta^t) + \frac{1}{\lambda}\nabla_{\theta}^2 F(\theta^t)$ is also positive definite. In such case, a direct computation reveals that \overline{G} is minimized at

$$\overline{\Delta\theta^*} = -\frac{1}{\lambda} \left(\mathbf{H}(\theta^t) + \frac{1}{\lambda} \nabla_{\theta}^2 F(\theta^t) \right)^{-1} \nabla_{\theta} F(\theta^t).$$
(13)

Consequently, the term involving $\nabla_{\theta}^2 F(\theta^t)$ vanishes when ϵ approaches zero and we obtain the result. The above argument is made precise in Appendix A.1. **Remark 4.1.** Note that our derivation also applies to the Fisher-Rao natural gradient or the Wasserstein natural gradient: If we replace the Sinkhorn divergence by the KL divergence (or the Wasserstein distance), the update direction $\mathbf{d}^t \simeq [\mathbf{H}(\theta^t)]^{-1} \nabla_{\theta} F(\theta^t)$ still holds, where $\mathbf{H}(\theta^t)$ is the Hessian matrix of the KL divergence (or the Wasserstein distance). This observation works for a general functional as a local metric Thomas et al. [2016] as well.

The following proposition states that SiNG is invariant to reparameterization in its continuous time limit ($\eta \rightarrow 0$). The proof is stated in Appendix A.2.

Proposition 4.2. Let Φ be an invertible and smoothly differentiable function and denote a reparameterization $\phi = \Phi(\theta)$. Define $\tilde{\mathbf{H}}(\bar{\phi}) := \nabla_{\phi}^2 S(\alpha_{\Phi^{-1}(\phi)}, \alpha_{\Phi^{-1}(\bar{\phi})})|_{\phi = \bar{\phi}}$ and $\tilde{F}(\bar{\phi}) := F(\Phi^{-1}(\bar{\phi}))$. Use $\dot{\theta}$ and $\dot{\phi}$ to denote the time derivative of θ and ϕ respectively. Consider SiNG in its continuous-time limit under these two parameterizations:

 $\dot{\theta}_s = -\mathbf{H}(\theta_s)^{-1} \nabla F(\theta_s) \text{ and } \dot{\phi}_s = -\tilde{\mathbf{H}}(\phi_s)^{-1} \nabla \tilde{F}(\phi_s) \text{ with } \phi_0 = \Phi(\theta_0).$ (14)

Then θ_s and ϕ_s are related by the equation $\phi_s = \Phi(\theta_s)$ at all time $s \ge 0$.

The SiNG direction is a "curved" negative gradient of the loss function $F(\theta)$ and the "curvature" is exactly given by the Sinkhorn Information Matrix (SIM), i.e. the Hessian $H(\theta^t) = \nabla_{\theta}^2 S(\alpha_{\theta}, \alpha_{\theta^t})|_{\theta=\theta^t}$ of the Sinkhorn divergence. An important question is whether SIM is computationally tractable. In the next section, we derive its explicit expression and describe how it can be efficiently computed. This is in sharp contrast to the Wasserstein information matrix (WIM) as in the WNG method proposed in Li and Montúfar [2018], which does not attain an explicit form for d > 1 (d is the parameter dimension).

While computing the update direction \mathbf{d}_t involves the inversion of $\mathbf{H}(\theta^t)$, it can be computed using the classical conjugate gradient algorithm, requiring only a matrix-vector product. Consequently, our Sinkhorn Natural Gradient (SiNG) admits a simple and elegant implementation based on modern auto-differential mechanisms such as PyTorch. We will elaborate this point in Appendix E.

5 Sinkhorn Information Matrix

In this section, we describe the explicit expression of the *Sinkhorn information matrix (SIM)* and show that it can be computed very efficiently using simple function operations (e.g. log and exp) and integrals with respect to α_{θ} (with complexity logarithmic in terms of the reciprocal of the target accuracy). The computability of SIM and hence SiNG is the key contribution of our paper. In the case when we can only compute the integration with respect to α_{θ} in a Monte Carlo manner, an empirical estimator of SIM (eSIM) is proposed in the next section with a delicate stability analysis.

Since $S(\cdot, \cdot)$ is a linear combination of terms like $OT_{\gamma}(\cdot, \cdot)$ -see (4), we can focus on the term $\nabla^2_{\theta}OT_{\gamma}(\alpha_{\theta}, \alpha_{\theta^t})_{|\theta=\theta^t}$ in $\mathbf{H}(\theta^t)$ and the other term $\nabla^2_{\theta}OT_{\gamma}(\alpha_{\theta}, \alpha_{\theta})_{|\theta=\theta^t}$ can be handled similarly. Having these two terms, SIM is computed as $\mathbf{H}(\theta^t) = [\nabla^2_{\theta}OT_{\gamma}(\alpha_{\theta}, \alpha_{\theta^t}) + \nabla^2_{\theta}OT_{\gamma}(\alpha_{\theta}, \alpha_{\theta})]_{|\theta=\theta^t}$.

Recall that the entropy regularized optimal transport distance OT_{γ} admits an equivalent dual concavemaximization form (6). Due to the concavity of \mathcal{H}_2 w.r.t. g in (5), the corresponding optimal $g_f = \operatorname{argmax}_{g \in \mathcal{C}(\mathcal{X})} \mathcal{H}_2(f, g; \alpha, \beta)$ can be explicitly computed for any fixed $f \in \mathcal{C}(\mathcal{X})$: Given a function $\overline{f} \in \mathcal{C}(\mathcal{X})$ and a measure $\alpha \in \mathcal{M}_1^+(\mathcal{X})$, define the Sinkhorn mapping as

$$\mathcal{A}(\bar{f},\alpha)(y) := -\gamma \log \int_{\mathcal{X}} \exp\left(-\frac{1}{\gamma}c(x,y) + \frac{1}{\gamma}\bar{f}(x)\right) \mathbf{d}\alpha(x).$$
(15)

The first-order optimality of g_f writes $g_f = \mathcal{A}(f, \alpha)$. Then, (6) can be simplified to the following problem with a single potential variable:

$$OT_{\gamma}(\alpha_{\theta},\beta) = \max_{f \in \mathcal{C}(\mathcal{X})} \left\{ \mathcal{H}_{1}(f,\theta) := \langle f, \alpha_{\theta} \rangle + \langle \mathcal{A}(f,\alpha_{\theta}), \beta \rangle \right\},$$
(16)

where we emphasize the impact of θ to \mathcal{H}_1 by writing it explicitly as a variable for \mathcal{H}_1 . Moreover, in \mathcal{H}_1 the dependence on β is dropped as β is fixed. We also denote the optimal solution to the R.H.S. of (16) by f_{θ} which is one of the Sinkhorn potentials for $OT_{\gamma}(\alpha_{\theta}, \beta)$.

The following proposition describes the explicit expression of $\nabla^2_{\theta} OT_{\gamma}(\alpha_{\theta}, \alpha_{\theta^t})_{|\theta=\theta^t}$ based on the above dual representation. The proof is provided in Appendix B.1.

Proposition 5.1. Recall the definition of the dual-variable function $\mathcal{H}_1 : \mathcal{C}(\mathcal{X}) \times \Theta \to \mathbb{R}$ in (16) and the definition of the second-order Fréchet derivative at the end of Section 1. For a parameterized push-forward measure $\alpha_{\theta} = T_{\theta \sharp} \mu$ and a fixed measure $\beta \in \mathcal{M}_1^+(\mathcal{X})$, we have

$$\nabla^2_{\theta} OT_{\gamma}(\alpha_{\theta}, \beta) = -D^2_{11} \mathcal{H}_1(f_{\theta}, \theta) \circ (Df_{\theta}, Df_{\theta}) + D^2_{22} \mathcal{H}_1(f_{\theta}, \theta), \tag{17}$$

where Df_{θ} denotes the Fréchet derivative of the Sinkhorn potential f_{θ} w.r.t. the parameter θ .

Remark 5.1 (SIM for 1*d*-Gaussian). It is in general difficult to give closed form expression of the SIM. However, in the simplest case when α_{θ} is a one-dimensional Gaussian distribution with a parameterized mean, i.e. $\alpha_{\theta} = \mathcal{N}(\mu(\theta), \sigma^2)$, SIM can be explicitly computed as $\nabla_{\theta}^2 S(\alpha_{\theta}, \beta) = 2\nabla_{\theta}^2 \mu(\theta)$ due to the closed form expression of the entropy regularized optimal transport between Gaussian measures [Janati et al., 2020].

Suppose that we have the Sinkhorn potential f_{θ} and its the Fréchet derivative Df_{θ} . Then the terms $D_{ij}^2 \mathcal{H}_1(f,\theta), i, j = 1, 2$ can all be evaluated using a constant amount of simple function operations, e.g. log and exp, since we know the explicit expression of \mathcal{H}_1 . Consequently, it is sufficient to have estimators f_{θ}^{ϵ} and g_{θ}^{ϵ} of f_{θ} and Df_{θ} respectively, such that $\|f_{\theta}^{\epsilon} - f_{\theta}\|_{\infty} \leq \epsilon$ and $\|g_{\theta}^{\epsilon} - Df_{\theta}\|_{op} \leq \epsilon$ for an arbitrary target accuracy ϵ . This is because the high accuracy approximation of f_{θ} and Df_{θ} imply the high accuracy approximation of $\nabla_{\theta}^2 OT_{\gamma}(\alpha_{\theta}, \beta)$ due to the Lipschitz continuity of the terms $D_{ij}^2 \mathcal{H}_1(f, \theta), i, j = 1, 2$. We derive these expressions and their Lipschitz continuity in Appendix B.

For the Sinkhorn Potential f_{θ} , its estimator f_{θ}^{θ} can be efficiently computed using the Sinkhorn-Knopp algorithm Sinkhorn and Knopp [1967]. We provide more details on this in Appendix B.2.

Proposition 5.2 (Computation of the Sinkhorn Potential f_{θ} – (Theorem 7.1.4 in [Lemmens and Nussbaum, 2012] and Theorem B.10 in [Luise et al., 2019]). Assume that the ground cost function c is bounded, i.e. $0 \leq c(x, y) \leq M_c, \forall x, y \in \mathcal{X}$. Denote $\lambda := \frac{\exp(M_c/\gamma) - 1}{\exp(M_c/\gamma) + 1} < 1$ and define

$$\mathcal{B}(f,\theta) := \mathcal{A}(\mathcal{A}(f,\alpha_{\theta}),\beta).$$
(18)

Then the fixed point iteration $f^{t+1} = \mathcal{B}(f^t, \theta)$ converges linearly: $\|f^{t+1} - f_{\theta}\|_{\infty} = \mathcal{O}(\lambda^t)$.

For the Fréchet derivative Df_{θ} , we construct its estimator in the following proposition.

Proposition 5.3 (Computation of the Fréchet derivative Df_{θ}). Let f_{θ}^{ϵ} be an approximation of f_{θ} such that $||f_{\theta}^{\epsilon} - f_{\theta}||_{\infty} \leq \epsilon$. Choose a large enough l, for instance $l = \lceil \log_{\lambda} \frac{1}{3} \rceil/2$. Define $\mathcal{E}(f, \theta) = \mathcal{B}(\cdots \mathcal{B}(f, \theta) \cdots, \theta)$, the l times composition of \mathcal{B} in its first variable. Then the sequence

$$g_{\theta}^{t+1} = D_1 \mathcal{E} \left(f_{\theta}^{\epsilon}, \theta \right) \circ g_{\theta}^t + D_2 \mathcal{E} \left(f_{\theta}^{\epsilon}, \theta \right) \tag{19}$$

converges linearly to a ϵ -neighborhood of Df_{θ} , i.e. $\|g_{\theta}^{t+1} - Df_{\theta}\|_{op} = \mathcal{O}(\epsilon + (\frac{2}{3})^t \|g_{\theta}^0 - Df_{\theta}\|_{op})$.

We deferred the proof to the above proposition to Appendix B.3. The high-accuracy estimators f_{θ}^{ϵ} and g_{θ}^{ϵ} derived in the above propositions can both be obtained using $\mathcal{O}(\log \frac{1}{\epsilon})$ function operations and integrals. With the expression of SIM and the two propositions discussing the efficient computation of f_{θ} and Df_{θ} , we obtain the following theorem.

Theorem 5.1 (Computability of SIM). For any given target accuracy $\epsilon > 0$, there exists an estimator $\mathbf{H}_{\epsilon}(\theta)$, such that $\|\mathbf{H}_{\epsilon}(\theta) - \mathbf{H}(\theta)\|_{op} \leq \epsilon$, and the estimator can be computed using $\mathcal{O}(\log \frac{1}{\epsilon})$ simple function operations and integrations with respect to α_{θ} .

This result shows a significantly broader applicability of SiNG than WNG, as the latter can only be used in limited situations due to the intractability of computing WIM.

6 Empirical Estimator of SIM

In the previous section, we derived an explicit expression for the Sinkhorn information matrix (SIM) and described how it can be computed efficiently. In this section, we provide an empirical estimator for SIM (eSIM) in the case where the integration w.r.t. α_{θ} can only be computed in a Monte-Carlo manner. Moreover, we prove the stability of eSIM by showing that the Fréchet derivative of the Sinkhorn potential with respect to the parameter θ is continuous with respect to the underlying measure μ , which is interesting on its own.

Recall that the parameterized measure has the structure $\alpha_{\theta} = T_{\theta \sharp} \mu$, where $\mu \in \mathcal{M}_{1}^{+}(\mathcal{Z})$ is some probability measure on the latent space $\mathcal{Z} \subseteq \mathbb{R}^{\bar{q}}$ and $T_{\theta} : \mathcal{Z} \to \mathcal{X}$ is some push-forward mapping parameterized by $\theta \in \Theta$. We use $\bar{\mu}$ to denote an empirical measure of μ with *n* Dirac measures: $\bar{\mu} = \frac{1}{n} \sum_{i=1}^{n} \delta_{z_i}$ with $z_i \stackrel{\text{iid}}{\sim} \mu$ and we use $\bar{\alpha}_{\theta}$ to denote the corresponding empirical measure of α_{θ} : $\bar{\alpha}_{\theta} = T_{\theta \sharp} \bar{\mu} = \frac{1}{n} \sum_{i=1}^{n} \delta_{T_{\theta}(z_i)}$. Based on the above definition, we propose the following empirical estimator for the Sinkhorn information matrix (eSIM)

$$\bar{\mathbf{H}}(\theta^t) = \nabla^2_{\theta} \mathcal{S}(\bar{\alpha}_{\theta}, \bar{\alpha}_{\theta^t})_{|\theta=\theta^t}.$$
(20)

The following theorem shows stability of eSIM. The proof is provided in Appendix C.

Theorem 6.1. Define the bounded Lipschitz metric of measures $d_{bl} : \mathcal{M}_1^+(\mathcal{X}) \times \mathcal{M}_1^+(\mathcal{X}) \to \mathbb{R}_+$ by

$$d_{bl}(\alpha,\beta) := \sup_{\|\xi\|_{bl} \le 1} |\langle \xi, \alpha \rangle - \langle \xi, \beta \rangle|,$$
(21)

where we denote $\|\xi\|_{bl} := \max\{\|\xi\|_{\infty}, \|\xi\|_{Lip}\}$ with $\|\xi\|_{Lip} := \max_{x,y \in \mathcal{X}} \frac{|\xi(x) - \xi(y)|}{\|x - y\|}$. Assume that the ground cost function is bounded and Lipschitz continuous. Then

$$\|\bar{\mathbf{H}}(\theta^t) - \mathbf{H}(\theta^t)\|_{op} = \mathcal{O}(d_{bl}(\mu, \bar{\mu})).$$
(22)

In the rest of this subsection, we analyze the structure of $\bar{\mathbf{H}}(\theta^t)$ and describe how it can be efficiently computed. Similar to the previous section, we focus on the term $\nabla^2_{\theta} OT_{\gamma}(\bar{\alpha}_{\theta}, \beta)$ with $\bar{\alpha}_{\theta} = \frac{1}{n} \sum_{i=1}^n \delta_{T_{\theta}(z_i)}$ and $\beta = \frac{1}{n} \sum_{i=1}^n \delta_{y_i}$ for arbitrary $y_i \in \mathcal{X}$.

First, notice that the output of the Sinkhorn mapping (15) is determined solely by the function values of the input \bar{f} at the support of α . Using $\mathbf{f} = [\mathbf{f}_1, \dots, \mathbf{f}_n] \in \mathbb{R}^n$ with $\mathbf{f}_i = \bar{f}(x_i)$ to denote the value extracted from \bar{f} on $\operatorname{supp}(\bar{\alpha})$, we define for a discrete probability measures $\bar{\alpha} = \frac{1}{n} \sum_{i=1}^n \delta_{x_i}$ the discrete Sinkhorn mapping $\bar{\mathcal{A}}(\mathbf{f}, \bar{\alpha}) : \mathbb{R}^n \times \mathcal{M}_1^+(\mathcal{X}) \to \mathcal{C}(\mathcal{X})$ as

$$\bar{\mathcal{A}}(\mathbf{f},\bar{\alpha})(y) := -\gamma \log\left(\frac{1}{n}\sum_{i=1}^{n}\exp\left(-\frac{1}{\gamma}c(x_i,y) + \frac{1}{\gamma}\mathbf{f}_i\right)\right) = \mathcal{A}(\bar{f},\bar{\alpha})(y),\tag{23}$$

where the last equality should be understood as two functions being identical. Since both $\bar{\alpha}_{\theta}$ and β in $OT_{\gamma}(\bar{\alpha}_{\theta}, \beta)$ are discrete, (16) can be reduced to

$$OT_{\gamma}(\bar{\alpha}_{\theta},\beta) = \max_{\mathbf{f}\in\mathbb{R}^n} \left\{ \bar{\mathcal{H}}_1(\mathbf{f},\theta) = \frac{1}{n} \mathbf{f}^{\top} \mathbf{1}_n + \frac{1}{n} \sum_{i=1}^n \bar{\mathcal{A}}(\mathbf{f},\bar{\alpha}_{\theta})(y_i) \right\}.$$
 (24)

Now, let \mathbf{f}_{θ} be the solution to the above problem. We can compute the first order gradient of $OT_{\gamma}(\bar{\alpha}_{\theta}, \beta)$ with respect to θ by

$$\nabla_{\theta} OT_{\gamma}(\bar{\alpha}_{\theta}, \beta) = J_{\mathbf{f}_{\theta}}^{\top} \cdot \nabla_{1} \bar{\mathcal{H}}_{1}(\mathbf{f}_{\theta}, \theta) + \nabla_{2} \bar{\mathcal{H}}_{1}(\mathbf{f}_{\theta}, \theta).$$
(25)

Here $J_{\mathbf{f}_{\theta}} = \frac{\partial \mathbf{f}_{\theta}}{\partial \theta} \in \mathbb{R}^{n \times d}$ denotes the Jacobian matrix of \mathbf{f}_{θ} with respect to θ and $\nabla_i \overline{\mathcal{H}}_1$ denotes the gradient of $\overline{\mathcal{H}}_1$ with respect to its i^{th} variable for i = 1, 2. Importantly, the optimality condition of \mathbf{f}_{θ} implies $\nabla_1 \overline{\mathcal{H}}_1(\mathbf{f}_{\theta}, \theta) = \mathbf{0}_n$. Further, we compute the second order gradient of $OT_{\gamma}(\overline{\alpha}_{\theta}, \beta)$ with respect to θ by (we omit the parameter $(\mathbf{f}_{\theta}, \theta)$ of $\overline{\mathcal{H}}_1$)

$$\nabla^{2}_{\theta} \mathrm{OT}_{\gamma}(\bar{\alpha}_{\theta},\beta) = T_{\mathbf{f}_{\theta}} \times_{1} \nabla_{1} \bar{\mathcal{H}}_{1} + J_{\mathbf{f}_{\theta}}^{\top} \cdot \nabla_{11} \bar{\mathcal{H}}_{1} \cdot J_{\mathbf{f}_{\theta}} + J_{\mathbf{f}_{\theta}}^{\top} \cdot \nabla_{12} \bar{\mathcal{H}}_{1} + \nabla_{21} \bar{\mathcal{H}}_{1}^{\top} \cdot J_{\mathbf{f}_{\theta}} + \nabla_{22} \bar{\mathcal{H}}_{1},$$
(26)

where $T_{\mathbf{f}_{\theta}} = \frac{\partial^2 \mathbf{f}_{\theta}}{\partial \theta^2} \in \mathbb{R}^{n \times d \times d}$ is a tensor denoting the second-order Jacobian matrix of \mathbf{f}_{θ} with respect to θ and \times_1 denotes the tensor product along its first dimension. Using the fact that $\nabla_1 \bar{\mathcal{H}}_1(\mathbf{f}_{\theta}, \theta) = \mathbf{0}_n$, we drop the first term and simplify $\nabla_{\theta}^2 OT_{\gamma}(\bar{\alpha}_{\theta}, \beta)$ to (again we omit the parameter $(\mathbf{f}_{\theta}, \theta) \circ \bar{\mathcal{H}}_1$)

$$\nabla^{2}_{\theta} OT_{\gamma}(\bar{\alpha}_{\theta}, \beta) = J^{\top}_{\mathbf{f}_{\theta}} \cdot \nabla_{11} \bar{\mathcal{H}}_{1} \cdot J_{\mathbf{f}_{\theta}} + J^{\top}_{\mathbf{f}_{\theta}} \cdot \nabla_{12} \bar{\mathcal{H}}_{1} + \nabla_{21} \bar{\mathcal{H}}_{1}^{\top} \cdot J_{\mathbf{f}_{\theta}} + \nabla_{22} \bar{\mathcal{H}}_{1}.$$
(27)

As we have the explicit expression of $\overline{\mathcal{H}}_1$, we can explicitly compute $\nabla_{ij}\overline{\mathcal{H}}_1$ given that we have the Sinkhorn potential \mathbf{f}_{θ} . Further, if we can compute $J_{\mathbf{f}_{\theta}}$, we are then able to compute $\nabla^2_{\theta} OT_{\gamma}(\bar{\alpha}_{\theta}, \beta)$. The following propositions can be viewed as discrete counterparts of Proposition 5.2 and Proposition 5.3 respectively. Both \mathbf{f}_{θ} and $J_{\mathbf{f}_{\theta}}$ can be well-approximated using a number of finite dimensional vector/matrix operations which is logarithmic in the desired accuracy. Besides, given these two quantities, one can easily check that $\nabla_{ij}\overline{\mathcal{H}}_1$ can be evaluated within $\mathcal{O}((n+d)^2)$ arithmetic operations. Consequently, we can compute an ϵ -accurate approximation of eSIM in time $\mathcal{O}((n+d)^2 \log \frac{1}{\epsilon})$.

Proposition 6.1 (Computation of the Sinkhorn Potential \mathbf{f}_{θ}). Assume that the ground cost function c is bounded, i.e. $0 \leq c(x, y) \leq M_c, \forall x, y \in \mathcal{X}$. Denote $\lambda := \frac{\exp(M_c/\gamma) - 1}{\exp(M_c/\gamma) + 1} < 1$ and define

$$\bar{\mathcal{B}}(\mathbf{f},\theta) := \bar{\mathcal{A}}(\mathbf{g},\beta) \text{ with } \mathbf{g} = [\bar{\mathcal{A}}(\mathbf{f},\bar{\alpha}_{\theta})(y_1),\ldots,\bar{\mathcal{A}}(\mathbf{f},\bar{\alpha}_{\theta})(y_n)] \in \mathbb{R}^n.$$
(28)

Then the fixed point iteration $\mathbf{f}^{t+1} = \bar{\mathcal{B}}(\mathbf{f}^t, \theta)$ converges linearly: $\|\mathbf{f}^{t+1} - \mathbf{f}_{\theta}\|_{\infty} = \mathcal{O}(\lambda^t)$

Proposition 6.2 (Computation of the Jacobian $J_{\mathbf{f}_{\theta}}$). Let \mathbf{f}_{ϵ} be an approximation of \mathbf{f}_{θ} such that $\|\mathbf{f}_{\epsilon} - \mathbf{f}_{\theta}\|_{\infty} \leq \epsilon$. Pick $l = [\log_{\lambda} \frac{1}{3}]/2$. Define $\overline{\mathcal{E}}(\mathbf{f}, \theta) = \overline{\mathcal{B}}(\cdots, \overline{\mathcal{B}}(\mathbf{f}, \theta) \cdots, \theta)$, the *l* times composition of $\overline{\mathcal{B}}$ in its first variable. Then the sequence of matrices

$$\mathbf{J}^{t+1} = J_1 \bar{\mathcal{E}}(\mathbf{f}_{\epsilon}, \theta) \cdot \mathbf{J}^t + J_2 \bar{\mathcal{E}}(\mathbf{f}_{\epsilon}, \theta), \tag{29}$$

converges linearly to an ϵ neighbor of $J_{\mathbf{f}_{\theta}}$: $\|\mathbf{J}^{t+1} - J_{\mathbf{f}_{\theta}}\|_{op} = \mathcal{O}(\epsilon + (\frac{2}{3})^t \|\mathbf{J}^0 - J_{\mathbf{f}_{\theta}}\|_{op})$. Here $J_i \bar{\mathcal{E}}$ denotes the Jacobian matrix of $\bar{\mathcal{E}}$ with respect to its *i*th variable.

The SiNG direction \mathbf{d}_t involves the inversion of $\bar{\mathbf{H}}(\theta^t)$. This can be (approximately) computed using the classical conjugate gradient (CG) algorithm, using only matrix-vector products. Combining eSIM and CG, we describe a simple and elegant PyTorch-based implementation for SiNG in Appendix E,

7 Experiment

In this section, we compare SiNG with other SGD-type solvers by training generative models. We did not compare with WNG Li and Montúfar [2018] since WNG can only be implemented for the case where the parameter dimension d is 1. We also tried to implement KWNG Arbel et al. [2019], which however diverges in our setting. In particular, we encounter the case when the KWNG direction has negative inner product with the euclidean gradient direction, leading to its divergence. As we discussed in the related work, the gap between KWNG and WNG cannot be quantified with reasonable assumptions, which explains our observation. In all the following experiments, we pick the push-forward map T_{θ} to be the generator network in DC-GAN [Radford et al., 2015]. For more detailed experiment settings, please see Appendix D.

7.1 Squared- ℓ_2 -norm as Ground Metric

We first consider the distribution matching problem, where our goal is to minimize the Sinkhorn divergence between the parameterized generative model $\alpha_{\theta} = T_{\theta \sharp} \mu$ and a given target distribution β ,

$$\min_{\theta \in \Theta} F(\theta) = \mathcal{S}(\alpha_{\theta}, \beta).$$
(30)

Here, T_{θ} is a neural network describing the push-forward map with its parameter summarized in θ and μ is a zero-mean isometric Gaussian distribution. In particular, the metric on the ground set \mathcal{X} is set to the vanilla squared- ℓ_2 norm, i.e. $c(x, y) = ||x - y||^2$ for $x, y \in \mathcal{X}$. Our experiment considers a specific instance of problem (30) where we take the measure β to be the distribution of the images in the CelebA dataset. We present the comparison of the generator loss (the objective value) vs time plot in right figure. The entropy regularization parameter γ is set to 0.01 for both the objective and the constraint. We can see that SiNG is much more efficient at reducing the objective value than ADAM given the same amount of time.



7.2 Squared- ℓ_2 -norm with an Additional Encoder as Ground Metric

We then consider a special case of problem (7), where the metric on the ground set \mathcal{X} is set to squared- ℓ_2 -norm with a *fixed* parameterized encoder (i.e. we fix the variable ξ in the max part of (7)): $c_{\xi}(x,y) = \|\phi_{\xi}(x) - \phi_{\xi}(y)\|^2$. Here $\phi_{\xi}(\cdot) : \mathcal{X} \to \mathbb{R}^{\hat{q}}$ is a neural network encoder that outputs an embedding of the input in a high dimensional space ($\hat{q} > q$, where we recall q is the dimension of the ground set \mathcal{X}). In particular, we set $\phi_{\xi}(\cdot)$ to be the discriminator network in DC-GAN without the last classification layer [Radford et al., 2015]. Two specific instances are considered: we take the measure β to be the distribution of the images in either the CelebA or the Cifar10 dataset. The



Figure 1: Generator losses on CelebA (left) and Cifar10 (right).

parameter ξ of the encoder ϕ is obtained in the following way: we first use SiNG to train a generative model by alternatively taking a SiNG step on θ and taking an SGD step on ξ . After sufficiently many iterations (when the generated image looks real or specifically 50 epochs), we fix the encoder ϕ_{ξ} . We then set the objective functional (1) to be $\mathcal{F}(\alpha_{\theta}) = S_{c_{\xi}}(\alpha_{\theta}, \beta)$ (see (7)), and compare SiNG and SGD-type algorithms in the minimization of \mathcal{F} under a consensus random initialization. We report the comparison in Figure 1, where we observe the significant improvement from SiNG in both accuracy and efficiency. Such phenomenon is due to the fact that SiNG is able to use geometry information by considering SIM while other method does not. Moreover, the pretrained ground cost c_{ξ} may capture some non-trivial metric structure of the images and consequently geometry-faithfully method like our SiNG can thus do better.

7.3 Training GAN with SiNG



Figure 2: Comparison of the visual quality of the images generated by Adam (left) and SiNG (right).

Finally, we showcase the the advantage of training a GAN model using SiNG over SGD-based solvers. Specifically, we consider the GAN model (7). The entropy regularization of the Sinkhorn divergence objective is set to $\gamma = 100$ as suggested in Table 2 of [Genevay et al., 2018]. The regularization for the constraint is set to $\gamma = 1$ in SiNG. We used ADAM as the optimizer for the discriminators (with step size 10^{-3} and batch size 4000). The result is reported in Figure 2. We can see that the images generated using SiNG are much more vivid than the ones obtained using SGD-based optimizers. We remark that our main goal has been to showcase that SiNG is more efficient in reducing the objective value compared to SGD-based solvers, and hence, we have used a relatively simpler DC-GAN type generator and discriminator (details given in the supplementary materials). If more sophisticated ResNet type generators and discriminators are used, the image quality can be further improved.

8 Broader Impact

We propose the Sinkhorn natural gradient (SiNG) algorithm for minimizing an objective functional over a parameterized family of generative-model type measures. While our results do not immediately lead to broader societal impacts (as they are mostly theoretical), they can lead to new potential positive impacts. SiNG admits explicit update rule which can be efficiently carried out in an exact manner under both continuous and discrete settings. Being able to exploit the geometric information provided in the Sinkhorn information matrix, we observe the remarkable advantage of SiNG over existing state-of-the-art SGD-type solvers. Such algorithm is readily applicable to many types of existing generative adversarial models and possibly helps the development of the literature.

Acknowledgment

This work is supported by NSF CPS-1837253.

References

- S.-I. Amari. Natural gradient works efficiently in learning. *Neural computation*, 10(2):251–276, 1998.
- S.-I. Amari, O. Barndorff-Nielsen, R. Kass, S. Lauritzen, C. Rao, et al. Differential geometrical theory of statistics. In *Differential geometry in statistical inference*, pages 19–94. Institute of Mathematical Statistics, 1987.
- M. Arbel, A. Gretton, W. Li, and G. Montúfar. Kernelized wasserstein natural gradient. *arXiv preprint* arXiv:1910.09652, 2019.
- M. Essid, D. F. Laefer, and E. G. Tabak. Adaptive optimal transport. *Information and Inference: A Journal of the IMA*, 8(4):789–816, 2019.
- J. Feydy, T. Séjourné, F.-X. Vialard, S.-i. Amari, A. Trouve, and G. Peyré. Interpolating between optimal transport and mmd using sinkhorn divergences. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pages 2681–2690, 2019.
- A. Genevay, G. Peyre, and M. Cuturi. Learning generative models with sinkhorn divergences. In *International Conference on Artificial Intelligence and Statistics*, pages 1608–1617, 2018.
- I. Goodfellow, J. Pouget-Abadie, M. Mirza, B. Xu, D. Warde-Farley, S. Ozair, A. Courville, and Y. Bengio. Generative adversarial nets. In *Advances in neural information processing systems*, pages 2672–2680, 2014.
- G. Hinton, N. Srivastava, and K. Swersky. Neural networks for machine learning lecture 6a overview of mini-batch gradient descent.
- H. Janati, B. Muzellec, G. Peyré, and M. Cuturi. Entropic optimal transport between (unbalanced) gaussian measures has a closed form. *arXiv preprint arXiv:2006.02572*, 2020.
- D. P. Kingma and J. Ba. Adam: A method for stochastic optimization. *arXiv preprint arXiv:1412.6980*, 2014.
- B. Lemmens and R. Nussbaum. *Nonlinear Perron-Frobenius Theory*, volume 189. Cambridge University Press, 2012.
- W. Li and G. Montúfar. Natural gradient via optimal transport. *Information Geometry*, 1(2):181–214, 2018.
- W. Li and G. Montúfar. Ricci curvature for parametric statistics via optimal transport. *Information Geometry*, pages 1–29, 2020.
- W. Li and J. Zhao. Wasserstein information matrix. arXiv preprint arXiv:1910.11248, 2019.
- W. Li, A. T. Lin, and G. Montúfar. Affine natural proximal learning. In *International Conference on Geometric Science of Information*, pages 705–714. Springer, 2019.

- G. Luise, S. Salzo, M. Pontil, and C. Ciliberto. Sinkhorn barycenters with free support via frank-wolfe algorithm. In *Advances in Neural Information Processing Systems* 32. 2019.
- J. Martens and R. Grosse. Optimizing neural networks with kronecker-factored approximate curvature. In *International conference on machine learning*, pages 2408–2417, 2015.
- G. Peyré, M. Cuturi, et al. Computational optimal transport. *Foundations and Trends*® *in Machine Learning*, 11(5-6):355–607, 2019.
- A. Radford, L. Metz, and S. Chintala. Unsupervised representation learning with deep convolutional generative adversarial networks, 2015.
- S. J. Reddi, S. Kale, and S. Kumar. On the convergence of adam and beyond. *arXiv preprint* arXiv:1904.09237, 2019.
- T. Salimans, H. Zhang, A. Radford, and D. Metaxas. Improving gans using optimal transport. In *International Conference on Learning Representations*, 2018.
- R. Sinkhorn and P. Knopp. Concerning nonnegative matrices and doubly stochastic matrices. *Pacific Journal of Mathematics*, 21(2):343–348, 1967.
- Y. Song, J. Song, and S. Ermon. Accelerating natural gradient with higher-order invariance. Proceedings of the 35th International Conference on Machine Learning, ICML 2018, Stockholmsmässan, Stockholm, 2018.
- P. Thomas, B. C. Silva, C. Dann, and E. Brunskill. Energetic natural gradient descent. In *International Conference on Machine Learning*, pages 2887–2895, 2016.

A Appendix Section for Methodology

A.1 **Proof of Proposition 4.1**

Denote the Lagrangian function by

$$G_{\lambda}(\Delta\theta) = F(\theta^{t} + \Delta\theta) + \lambda \left(S(\alpha_{\theta^{t} + \Delta\theta}, \alpha_{\theta^{t}}) - \epsilon - \epsilon^{c_{2}} \right).$$
(31)

We have the following inequality which characterize a lower bound of the solution to (9) (recall that $1 < c_2 < 1.5, c_1 < 0.5$ and $3c_1 - 1 \ge c_2$),

$$\min_{\Delta\theta\in\mathbb{R}^{d}} F(\theta^{t} + \Delta\theta) \\
\text{s.t.} \|\Delta\theta\| \leq \epsilon^{c_{1}} \\
\mathcal{S}(\alpha_{\theta^{t} + \Delta\theta}, \alpha_{\theta^{t}}) \leq \epsilon + \epsilon^{c_{2}} = \min_{\|\Delta\theta\| \leq \epsilon^{c_{1}}} \max_{\lambda \geq 0} G_{\lambda}(\Delta\theta) \geq \max_{\lambda \geq 0} \min_{\|\Delta\theta\| \leq \epsilon^{c_{1}}} G_{\lambda}(\Delta\theta). \quad (32)$$

We now focus on the R.H.S. of the above inequality. Denote the second-order Taylor expansion of the Lagrangian G_{λ} by \bar{G}_{λ} :

$$\bar{G}_{\lambda}(\Delta\theta) = F(\theta^{t}) + \langle \nabla_{\theta}F(\theta^{t}), \Delta\theta \rangle + \frac{1}{2} \langle \nabla_{\theta}^{2}F(\theta^{t})\Delta\theta, \Delta\theta \rangle + \frac{\lambda}{2} \langle \mathbf{H}(\theta^{t})\Delta\theta, \Delta\theta \rangle - \lambda\epsilon - \lambda\epsilon^{c_{2}},$$

where we used the optimality condition (10) of $S(\alpha, \alpha^t)$ so that the first-order term of $S(\alpha, \alpha^t)$ vanishes. Besides, $\mathbf{H}(\theta)$ is defined in (10). The error of such approximation can be bounded as

$$G_{\lambda}(\Delta\theta) - \bar{G}_{\lambda}(\Delta\theta) = \mathcal{O}((\lambda+1) \|\Delta\theta\|^3).$$
(33)

Further, for any fixed λ , denote $\Delta \theta_{\lambda}^* = \operatorname{argmin}_{\|\Delta \theta\| \leq \epsilon^{c_1}} G_{\lambda}(\Delta \theta)$.

We can then derive the following lower bound on the minimization subproblem of the R.H.S. of (32):

$$\max_{\lambda \ge 0} \min_{\|\Delta\theta\| \le \epsilon^{c_1}} G_{\lambda}(\Delta\theta) = \max_{\lambda \ge 0} \bar{G}_{\lambda}(\Delta\theta_{\lambda}^*) - \mathcal{O}((\lambda+1)\|\Delta\theta_{\lambda}^*\|^3)$$

$$\geq \max_{\lambda \ge 0} \bar{G}_{\lambda}(\Delta\theta_{\lambda}^*) - \mathcal{O}((\lambda+1)\epsilon^{3c_1})$$

$$\geq \max_{\lambda \ge 0} \min_{\|\Delta\theta\| \le \epsilon^{c_1}} \bar{G}_{\lambda}(\Delta\theta) - \mathcal{O}((\lambda+1)\epsilon^{3c_1}),$$

Note that for sufficiently large λ , $\mathbf{H}(\theta^t) + \frac{1}{\lambda} \nabla_{\theta}^2 F(\theta^t) > 0$ by recalling the positive definiteness of $\mathbf{H}(\theta^t)$. In this case, as a convex program, $\min_{\|\Delta\theta\| \le \epsilon^{c_1}} \bar{G}_{\lambda}(\Delta\theta)$ admits the closed form solution: Denote $\overline{\Delta\theta}_{\lambda}^* = \operatorname{argmin} \bar{G}_{\lambda}(\Delta\theta)$. We have

$$\overline{\Delta\theta_{\lambda}^{*}} = -\frac{1}{\lambda} \left(\mathbf{H}(\theta^{t}) + \frac{1}{\lambda} \nabla_{\theta}^{2} F(\theta^{t}) \right)^{-1} \nabla_{\theta} F(\theta^{t}) \text{ and } \bar{G}(\overline{\Delta\theta_{\lambda}^{*}}) = F(\theta^{t}) - \frac{\bar{a}}{2\lambda} - \lambda\epsilon - \lambda\epsilon^{c_{2}}, \quad (34)$$

where we denote $\bar{a} := \langle \left[\mathbf{H}(\theta^{t}) + \frac{1}{\lambda} \nabla_{\theta}^{2} F(\theta^{t}) \right]^{-1} \nabla_{\theta} F(\theta^{t}), \nabla_{\theta} F(\theta^{t}) \rangle > 0.$

For sufficiently small ϵ , by taking $\lambda = \sqrt{\frac{a}{2\epsilon}}$ with $a := \langle [\mathbf{H}(\theta^t)]^{-1} \nabla_{\theta} F(\theta^t), \nabla_{\theta} F(\theta^t) \rangle > 0$ (note that $\|\overline{\Delta \theta_{\lambda}^*}\| = \mathcal{O}(\sqrt{\epsilon}) < \epsilon^{c_1}$ and is hence feasible for $c_1 < 0.5$), the R.H.S. of (32) has the following lower bound (recall that we have $3c_1 - 1 \ge c_2$)

$$\max_{\lambda \ge 0} \min_{\|\Delta\theta\| \le \epsilon^{c_1}} G_{\lambda}(\Delta\theta) \ge F(\theta^t) - \left(\frac{\bar{a}}{\sqrt{2a}} + \sqrt{\frac{a}{2}}\right)\sqrt{\epsilon} - \mathcal{O}(\epsilon^{c_2 - 0.5}).$$
(35)

This result leads to the following lower bound on (9):

$$\lim_{\epsilon \to 0} \frac{F(\theta^t + \Delta \theta^t_{\epsilon}) - F(\theta^t)}{\sqrt{\epsilon}} \ge -\sqrt{2\langle [\mathbf{H}(\theta^t)]^{-1} \nabla_{\theta} F(\theta^t), \nabla_{\theta} F(\theta^t) \rangle},$$
(36)

where $\Delta \theta_{\epsilon}^{t}$ is the solution to (9). Finally, observe that the equality is achieved by taking $\Delta \theta_{\epsilon}^{t} = \sqrt{2\epsilon} (\mathbf{H}(\theta^{t}))^{-1} \nabla_{\theta} F(\theta^{t})$.

$$\frac{\sqrt{\langle [\mathbf{H}(\theta^{t})]^{-1} \nabla_{\theta} F(\theta^{t}), \nabla_{\theta} F(\theta^{t}) \rangle}}{\lim_{\epsilon \to 0} \frac{F(\theta^{t} + \Delta \theta^{t}_{\epsilon}) - F(\theta^{t})}{\sqrt{\epsilon}} = \lim_{\epsilon \to 0} \frac{1}{\sqrt{\epsilon}} \langle \nabla F(\theta^{t}), \Delta \theta^{t}_{\epsilon} \rangle = -\sqrt{2 \langle [\mathbf{H}(\theta^{t})]^{-1} \nabla_{\theta} F(\theta^{t}), \nabla_{\theta} F(\theta^{t}) \rangle},$$
(37)

and $\Delta \theta_{\epsilon}^{t}$ is feasible for sufficiently small ϵ (note that we have $\frac{1}{2} \langle \mathbf{H}(\theta^{t}) \Delta \theta_{\epsilon}^{t}, \Delta \theta_{\epsilon}^{t} \rangle = \epsilon$):

$$S(\alpha_{\theta^t + \Delta\theta^t_{\epsilon}}, \alpha_{\theta^t}) \leqslant \frac{1}{2} \langle \mathbf{H}(\theta^t) \Delta\theta^t_{\epsilon}, \Delta\theta^t_{\epsilon} \rangle + \mathcal{O}(\epsilon^{1.5}) < \epsilon + \epsilon^{c_2},$$
(38)

and $\|\Delta \theta^t_\epsilon\| = \mathcal{O}(\sqrt{\epsilon}) < \epsilon^{c_1}$ for $c_1 < 0.5$. This leads to our conclusion.

A.2 Proof of Proposition 4.2

Our goal is to show that the continuous-time limit of $\Phi(\theta_s)$ satisfies the same differential equation as ϕ_s provided that $\Phi(\theta_0) = \phi_0$. To do so, first compute the differential equation of $\Phi(\theta_s)$

$$\frac{\partial \Phi(\theta_s)}{\partial s} = \nabla_{\theta} \Phi(\theta_s) \dot{\theta}_s = -\nabla_{\theta} \Phi(\theta_s) \mathbf{H}(\theta_s)^{-1} \nabla F(\theta_s), \tag{39}$$

where $\nabla_{\theta} \Phi(\theta_s)$ is the Jacobian matrix of $\Phi(\theta)$ w.r.t. θ at $\theta = \theta_s$. We then compute the differential equation of ϕ_s (note that $\nabla_{\phi} \Phi^{-1}(\phi_s)$ is the Jacobian matrix of $\Phi^{-1}(\phi)$ w.r.t. ϕ at $\phi = \phi_s$)

$$\begin{aligned} \dot{\phi}_{s} &= -\left[\nabla_{\phi}^{2} \mathcal{S}(\alpha_{\Phi^{-1}(\phi)}, \alpha_{\Phi^{-1}(\phi_{s})})_{|\phi=\phi_{s}}\right]^{-1} \nabla_{\phi} F(\Phi^{-1}(\phi))_{|\phi=\phi_{s}} \\ &= -\left[\nabla_{\phi} \Phi^{-1}(\phi_{s})^{\top} \nabla_{\theta}^{2} \mathcal{S}(\alpha_{\theta}, \alpha_{\theta^{s}})_{|\theta=\theta_{s}} \nabla_{\phi} \Phi^{-1}(\phi_{s})\right]^{-1} \nabla_{\phi} \Phi^{-1}(\phi_{s})^{\top} \nabla F(\theta)_{|\theta=\theta_{s}} \\ &= -\left[\nabla_{\phi} \Phi^{-1}(\phi_{s})\right]^{-1} \nabla_{\theta}^{2} \mathcal{S}(\alpha_{\theta}, \alpha_{\theta^{s}})_{|\theta=\theta_{s}} \nabla F(\theta)_{|\theta=\theta_{s}} \\ &= -\nabla_{\theta} \Phi(\theta_{s}) \mathbf{H}(\theta_{s})^{-1} \nabla F(\theta_{s}) \end{aligned} \tag{41}$$
$$&= \frac{\partial \Phi(\theta_{s})}{\partial s}.$$

Here we use the following lemma in (40). We use $\Phi^{-1}(\phi_s) = \theta_s$ and the inverse function theorem $\nabla_{\theta} \Phi(\theta_s) = \left[\nabla_{\phi} \Phi^{-1}(\phi_s) \right]^{-1}$ in (41).

Lemma A.1.

$$\nabla_{\phi}^{2} \mathcal{S}(\alpha_{\Phi^{-1}(\phi)}, \alpha_{\Phi^{-1}(\phi_{s})})|_{\phi=\phi_{s}} = \nabla_{\phi} \Phi^{-1}(\phi_{s})^{\top} \nabla_{\theta}^{2} \mathcal{S}(\alpha_{\theta}, \alpha_{\theta^{s}})|_{\theta=\theta_{s}} \nabla_{\phi} \Phi^{-1}(\phi_{s})$$
(42)

Proof. This lemma can be proved with simple computations. We compute only for the terms in $\nabla^2_{\theta} OT_{\gamma}(\alpha_{\theta}, \alpha_{\theta^s})$ as example. The terms in $\nabla^2_{\theta} OT_{\gamma}(\alpha_{\theta}, \alpha_{\theta})$ can be computed similarly. Recall the expression

$$\nabla^{2}_{\theta} OT_{\gamma}(\alpha_{\theta}, \beta) = D^{2}_{11} \mathcal{H}_{1}(f_{\theta}, \theta) \circ (Df_{\theta}, Df_{\theta}) + D^{2}_{12} \mathcal{H}_{1}(f_{\theta}, \theta) \circ (Df_{\theta}, \mathcal{I}_{d}) + D^{2}_{21} \mathcal{H}_{1}(f_{\theta}, \theta) \circ (\mathcal{I}_{d}, Df_{\theta}) + D^{2}_{22} \mathcal{H}_{1}(f_{\theta}, \theta) \circ (\mathcal{I}_{d}, \mathcal{I}_{d}).$$

$$(43)$$

We compute

$$\nabla^{2}_{\phi} OT_{\gamma}(\alpha_{\Phi^{-1}(\phi)},\beta) = D^{2}_{11} \mathcal{H}_{1}(f_{\Phi^{-1}(\phi)},\Phi^{-1}(\phi)) \circ (Df_{\Phi^{-1}(\phi)} \circ J_{\Phi^{-1}}(\phi), Df_{\Phi^{-1}(\phi)} \circ J_{\Phi^{-1}}(\phi)) \\
+ D^{2}_{12} \mathcal{H}_{1}(f_{\Phi^{-1}(\phi)},\Phi^{-1}(\phi)) \circ (Df_{\Phi^{-1}(\phi)} \circ J_{\Phi^{-1}}(\phi), J_{\Phi^{-1}}(\phi)) \\
+ D^{2}_{21} \mathcal{H}_{1}(f_{\Phi^{-1}(\phi)},\Phi^{-1}(\phi)) \circ (J_{\Phi^{-1}}(\phi), Df_{\Phi^{-1}(\phi)} \circ J_{\Phi^{-1}}(\phi)) \\
+ D^{2}_{22} \mathcal{H}_{1}(f_{\Phi^{-1}(\phi)},\Phi^{-1}(\phi)) \circ (J_{\Phi^{-1}}(\phi), J_{\Phi^{-1}}(\phi)).$$
(44)

Plugging $\Phi^{-1}(\phi_s) = \theta_s$ to the above equality, we have

$$\nabla_{\phi}^{2} \mathrm{OT}_{\gamma}(\alpha_{\Phi^{-1}(\phi)},\beta)_{|\phi=\phi_{s}} = \nabla_{\phi} \Phi^{-1}(\phi_{s})^{\top} \nabla_{\theta}^{2} \mathrm{OT}_{\gamma}(\alpha_{\theta},\beta)_{|\theta=\theta_{s}} \nabla_{\phi} \Phi^{-1}(\phi_{s}).$$
(45)

B Appendix on SIM

B.1 Proof of Proposition 5.1

We will derive the explicit expression of $\nabla^2_{\theta} OT_{\gamma}(\alpha_{\theta}, \alpha_{\theta^t})|_{\theta=\theta^t}$ based on the dual representation (16). Recall the definition of the Fréchet derivative in Definition 2 and its chain rule $D(f \circ g)(x) = Df(g(x)) \circ Dg(x)$. We compute the first-order gradient by

$$\nabla_{\theta} OT_{\gamma}(\alpha_{\theta}, \beta) = \nabla_{\theta} \mathcal{H}_{1}(f_{\theta}, \theta) = \underbrace{D_{1} \mathcal{H}_{1}(f_{\theta}, \theta) \circ Df_{\theta}}_{\mathcal{G}_{1}(f_{\theta}, \theta)} + \underbrace{D_{2} \mathcal{H}_{1}(f_{\theta}, \theta)}_{\mathcal{G}_{2}(f_{\theta}, \theta)}, \tag{46}$$

where $D_i \mathcal{H}_1$ denote the Fréchet derivative of \mathcal{H}_1 with respect to its *i*th variable. Importantly, the optimality condition of (16) implies that $D_1 \mathcal{H}_1(f_{\theta}, \theta)[g] = 0, \forall g \in \mathcal{C}(\mathcal{X})$.

Further, in order to compute the second order gradient of $OT_{\gamma}(\alpha_{\theta}, \beta)$ with respect to θ , we first compute the gradient of $\mathcal{G}_i, i = 1, 2$:

$$\nabla_{\theta} \mathcal{G}_{1}(f_{\theta},\theta) = D_{1} \mathcal{H}_{1}(f_{\theta},\theta) \circ D^{2} f_{\theta} + D_{11}^{2} \mathcal{H}_{1}(f_{\theta},\theta) \circ (Df_{\theta},Df_{\theta}) + D_{12}^{2} \mathcal{H}_{2}(f_{\theta},\theta) \circ (Df_{\theta},\mathcal{I}_{d}),$$
(47)

$$\nabla_{\theta} \mathcal{G}_2(f_{\theta}, \theta) = D_{21}^2 \mathcal{H}_1(f_{\theta}, \theta) \circ (\mathcal{I}_d, Df_{\theta}) + D_{22}^2 \mathcal{H}_1(f_{\theta}, \theta) \circ (\mathcal{I}_d, \mathcal{I}_d).$$
(48)

Using the fact that $D_1\mathcal{H}_1(f_\theta, \theta)[g] = 0, \forall g \in \mathcal{C}(\mathcal{X})$, we can drop the first term in the R.H.S. of (47). Combining the above results, we have

$$\nabla^{2}_{\theta} OT_{\gamma}(\alpha_{\theta}, \beta) = D^{2}_{11} \mathcal{H}_{1}(f_{\theta}, \theta) \circ (Df_{\theta}, Df_{\theta}) + D^{2}_{12} \mathcal{H}_{1}(f_{\theta}, \theta) \circ (Df_{\theta}, \mathcal{I}_{d}) + D^{2}_{21} \mathcal{H}_{1}(f_{\theta}, \theta) \circ (\mathcal{I}_{d}, Df_{\theta}) + D^{2}_{22} \mathcal{H}_{1}(f_{\theta}, \theta) \circ (\mathcal{I}_{d}, \mathcal{I}_{d}).$$

$$(49)$$

Moreover, we can further simplify the above expression by noting that for any $g \in T(\mathbb{R}^d, \mathcal{C}(\mathcal{X}))$, i.e. any bounded linear operators from \mathbb{R}^d to $\mathcal{C}(\mathcal{X})$,

$$\nabla_{\theta} \left(D_1 \mathcal{H}_1(f_{\theta}, \theta) \circ g \right) = D_{11}^2 \mathcal{H}_1(f_{\theta}, \theta) \circ (g, Df_{\theta}) + D_{12}^2 \mathcal{H}_1(f_{\theta}, \theta) \circ (g, \mathcal{I}_d) = 0.$$
(50)

Plugging in $g = Df_{\theta}$ in the above equality we have

$$D_{11}^2 \mathcal{H}_1(f_\theta, \theta) \circ (Df_\theta, Df_\theta) = -D_{12}^2 \mathcal{H}_1(f_\theta, \theta) \circ (Df_\theta, \mathcal{I}_d).$$
(51)

Consequently we derive (we omit the identity operator $(\mathcal{I}_d, \mathcal{I}_d)$ for the second term)

$$\nabla^2_{\theta} OT_{\gamma}(\alpha_{\theta}, \beta) = -D^2_{11} \mathcal{H}_1(f_{\theta}, \theta) \circ (Df_{\theta}, Df_{\theta}) + D^2_{22} \mathcal{H}_1(f_{\theta}, \theta),$$
(52)

where we note that $D_{12}^2 \mathcal{H}_1(f_{\theta}, \theta) \circ (Df_{\theta}, \mathcal{I}_d)$ is symmetric from (51) and

$$D_{21}^{2}\mathcal{H}_{1}(f_{\theta},\theta)\circ(\mathcal{I}_{d},Df_{\theta}) = \left[D_{12}^{2}\mathcal{H}_{1}(f_{\theta},\theta)\circ(Df_{\theta},\mathcal{I}_{d})\right]^{\top} = D_{12}^{2}\mathcal{H}_{1}(f_{\theta},\theta)\circ(Df_{\theta},\mathcal{I}_{d}).$$
 (53)

These two terms can be computed explicitly and involve only simple function operations like exp and log and integration with respect to α_{θ} and β , as discussed in the following.

B.1.1 Explicit Expression of $\nabla^2_{\theta} OT_{\gamma}(\alpha_{\theta}, \beta)$

Denote $\mathbf{A}_1 = D_{11}^2 \mathcal{H}_1(f_{\theta}, \theta) \circ (Df_{\theta}, Df_{\theta})$ as the first term of (52). We note that $\mathbf{A}_1 \in \mathbb{R}^{d \times d}$ is a matrix and hence is a bilinear operator. If we can compute $h_1^\top \mathbf{A}_1 h_2$ for any two directions $h_1, h_2 \in \mathbb{R}^d$, we are able to compute entries of \mathbf{A}_1 by taking h_1 and h_2 to be the canonical bases. We compute this quantity $h_1^\top \mathbf{A}_1 h_2$ as follows.

For a fixed $y \in \mathcal{X}$, denote $\mathcal{T}_y : \mathcal{X} \times \mathcal{C}(\mathcal{X}) \to \mathbb{R}$ by

$$\mathcal{T}_y(x, f) := \exp(-c(x, y)/\gamma) \exp(f(x)/\gamma).$$

Denote $g_1 = Df_{\theta}[h_1] \in \mathcal{C}(\mathcal{X})$ for some direction $h_1 \in \mathbb{R}^d$ (recall that $Df_{\theta} \in T(\mathbb{R}^d, \mathcal{C}(\mathcal{X}))$), where T(V, W) is the family of bounded linear operators from set V to set W). Use the chain rule of Fréchet derivative to compute

$$\left(D_1 \mathcal{A}(f, \alpha_\theta)[g_1]\right)(y) = -\frac{\int_{\mathcal{X}} \mathcal{T}_y(x, f) g_1(x) \mathbf{d}\alpha_\theta(x)}{\int_{\mathcal{X}} \mathcal{T}_y(x, f) \mathbf{d}\alpha_\theta(x)}.$$
(54)

Let $h_2 \in \mathbb{R}^d$ be another direction and denote $g_2 = Df_{\theta}[h_2] \in \mathcal{C}(\mathcal{X})$. We compute

$$\frac{\left(D_{11}^{2}\mathcal{A}(f,\alpha_{\theta})[g_{1},g_{2}]\right)(y)}{\gamma\int_{\mathcal{X}}\mathcal{T}_{y}(x,f)\mathbf{d}\alpha_{\theta}(x)} - \frac{\int_{\mathcal{X}^{2}}\mathcal{T}_{y}(x,f)\mathcal{T}_{y}(x',f)g_{1}(x)g_{2}(x')\mathbf{d}\alpha_{\theta}(x)\mathbf{d}\alpha_{\theta}(x')}{\gamma\left[\int_{\mathcal{X}}\mathcal{T}_{y}(x,f)\mathbf{d}\alpha_{\theta}(x)\right]^{2}}.$$
(55)

Moreover, for any two directions $h_1, h_2 \in \mathbb{R}^d$, we compute $D_{11}^2 \mathcal{H}_1(f, \theta) [Df_\theta[h_1], Df_\theta[h_2]]$ by

$$D_{11}^{2}\mathcal{H}_{1}(f,\theta)\left[Df_{\theta}[h_{1}], Df_{\theta}[h_{2}]\right] = \int_{\mathcal{X}} \left(D_{11}^{2}\mathcal{A}(f_{\theta},\alpha_{\theta})\left[Df_{\theta}[h_{1}], Df_{\theta}[h_{2}]\right]\right)(y)\mathbf{d}\beta(y), \quad (56)$$

which by plugging in (55) yields closed a form expression with only simple function operations like exp and log and integration with respect to α_{θ} and β .

We then compute the second term of (52). Using the change-of-variable formula, we have

$$\mathcal{A}(f, T_{\theta \sharp} \mu)(y) = -\gamma \log \int_{\mathcal{Z}} \exp\left(-\frac{1}{\gamma} c(T_{\theta}(z), y) + \frac{1}{\gamma} f(T_{\theta}(z))\right) \mathbf{d}\mu(z).$$
(57)

For any $f \in C(\mathcal{X})$, the first-order Fréchet derivative of $\mathcal{H}_1(f, \theta)$ w.r.t. its second variable is given by

$$D_{2}\mathcal{H}_{1}(f,\theta) = \int_{\mathcal{Z}} \langle \nabla_{\theta} T_{\theta}(z), \nabla f(T_{\theta}(z)) \rangle d\mu(z) + \int_{\mathcal{X}} \frac{\int_{\mathcal{Z}} \mathcal{T}_{y}(T_{\theta}(z), f) \langle \nabla_{\theta} T_{\theta}(z), \nabla_{1} c(T_{\theta}(z), y) - \nabla f(T_{\theta}(z)) \rangle d\mu(z)}{\int_{\mathcal{Z}} \mathcal{T}_{y}(T_{\theta}(z), f) d\mu(z)} d\beta(y).$$

Denote $u_z(\theta, f) = \nabla_1 c (T_\theta(z), y) - \nabla f (T_\theta(z))$. The second-order Fréchet derivative is given by

$$D_{22}^{2}\mathcal{H}_{1}(f,\theta)$$

$$= \int_{\mathcal{Z}} \nabla_{\theta}^{2} T_{\theta}(z) \times_{1} \nabla f(T_{\theta}(z)) + \nabla_{\theta} T_{\theta}(z)^{\top} \nabla^{2} f(T_{\theta}(z)) \nabla_{\theta} T_{\theta}(z) d\mu(z)$$

$$+ \frac{1}{\gamma} \int_{\mathcal{X}} \frac{\int_{\mathcal{Z}} \mathcal{T}_{y}(T_{\theta}(z), f) \nabla_{\theta} T_{\theta}(z)^{\top} u_{z}(\theta, f) u_{z}(\theta, f)^{\top} \nabla_{\theta} T_{\theta}(z) d\mu(z)}{\int_{\mathcal{Z}} \mathcal{T}_{y}(T_{\theta}(z), f) \nabla_{\theta}^{2} T_{\theta}(z) \times_{1} u_{z}(\theta, f) d\mu(z)} d\beta(y)$$

$$+ \int_{\mathcal{X}} \frac{\int_{\mathcal{Z}} \mathcal{T}_{y}(T_{\theta}(z), f) \nabla_{\theta} T_{\theta}(z)^{\top} [\nabla_{11} c(T_{\theta}(z), y) - \nabla^{2} f(T_{\theta}(z))] \nabla_{\theta} T_{\theta}(z) d\mu(z)}{\int_{\mathcal{Z}} \mathcal{T}_{y}(T_{\theta}(z), f) d\mu(z)} d\beta(y)$$

$$+ \frac{1}{\gamma} \int_{\mathcal{X}} \frac{\int_{\mathcal{Z}} \mathcal{T}_{y}(T_{\theta}(z), f) \nabla_{\theta} T_{\theta}(z)^{\top} u_{z}(\theta, f) d\mu(z)}{[\int_{\mathcal{Z}} \mathcal{T}_{y}(T_{\theta}(z), f) d\mu(z)]^{2}} d\beta(y).$$
(58)

Here $\nabla_{\theta} T_{\theta}(z) \in \mathbb{R}^{q \times d}$ and $\nabla_{\theta}^2 T_{\theta}(z) \in \mathbb{R}^{q \times d \times d}$ denote the first and second order Jacobian of $T_{\theta}(z)$ w.r.t. to θ ; \times_1 denotes the tensor product along the first dimension; $\nabla f \in \mathbb{R}^q$ and $\nabla^2 f \in \mathbb{R}^{q \times q}$ denote the first and second order gradient of f w.r.t. its input; $\nabla_1 c \in \mathbb{R}^q$ and $\nabla_{11} c \in \mathbb{R}^{q \times q}$ denote the first and second order gradient of c w.r.t. its first input. By plugging in $f = f_{\theta}$, we have the explicit expression of the second term of (52).

B.2 More details in Proposition 5.2

First, we recall some existing results about the Sinkhorn potential f_{θ} .

Assumption B.1. The ground cost function c is bounded and we denote $M_c := \max_{x,y \in \mathcal{X}} c(x,y)$.

It is known that, under the above boundedness assumption on the ground cost function c, f_{θ} is a solution to the generalized DAD problem (eq. (7.4) in [Lemmens and Nussbaum, 2012]), which is the fixed point to the operator $\mathcal{B} : \mathcal{C}(\mathcal{X}) \times \Theta \to \mathcal{C}(\mathcal{X})$ defined as

$$\mathcal{B}(f,\theta) := \mathcal{A}\big(\mathcal{A}(f,\alpha_{\theta}),\beta\big). \tag{59}$$

Further, the Birkhoff-Hopf Theorem (Sections A.4 and A.7 in [Lemmens and Nussbaum, 2012]) states that $\exp(\mathcal{B}/\gamma)$ is a contraction operator under the Hilbert metric with a contraction factor λ^2 where $\lambda := \frac{\exp(M_c/\gamma) - 1}{\exp(M_c/\gamma) + 1} < 1$ (see also Theorem B.5 in [Luise et al., 2019]): For strictly positive functions $u, u' \in \mathcal{C}(\mathcal{X})$, define the Hilbert metric as

$$d_H(u, u') := \log \max_{x, y \in \mathcal{X}} \frac{u(x)u'(y)}{u'(x)u(y)}.$$
 (60)

For any measure $\alpha \in \mathcal{M}_1^+(\mathcal{X})$, we have

$$d_H(\exp(\mathcal{A}(f,\alpha_\theta)/\gamma),\exp(\mathcal{A}(f',\alpha_\theta)/\gamma)) \leq \lambda d_H(\exp(f/\gamma),\exp(f'/\gamma)).$$
(61)

Consequently, by applying the fixed point iteration

$$f^{t+1} = \mathcal{B}(f^t, \theta), \tag{62}$$

also known as the Sinkhorn-Knopp algorithm, one can compute f_{θ} in logarithmic time: $||f^{t+1} - f_{\theta}||_{\infty} = O(\lambda^t)$ (Theorem. 7.1.4 in [Lemmens and Nussbaum, 2012] and Theorem B.10 in [Luise et al., 2019]).

While the above discussion shows that the output of the Sinkhorn-Knopp algorithm well approximates the Sinkhorn potential f_{θ} , it would be useful to discuss more about the boundedness property of the sequence $\{f^t\}$ produced by the above Sinkhorn-Knopp algorithm. We first show that under bounded initialization f^0 , the entire sequence $\{f^t\}$ is bounded.

Lemma B.1. Suppose that we initialize the Sinkhorn-Knopp algorithm with $f^0 \in C(\mathcal{X})$ such that $\|f^0\|_{\infty} \leq M_c$. One has $\|f^t\|_{\infty} \leq M_c$, for $t = 1, 2, 3, \cdots$.

Proof. For $||f||_{\infty} \leq M_c$ and any measure $\alpha \in \mathcal{M}_1^+(\mathcal{X})$, we have

$$\|\mathcal{A}(f,\alpha)\|_{\infty} = \gamma \|\log \int_{\mathcal{X}} \exp\{-c(x,\cdot)/\gamma\} \exp\{f(x)/\gamma\} \mathbf{d}\alpha(x)\|_{\infty} \leq \gamma \log \exp(M_c/\gamma) \leq M_c.$$

One can then check the lemma via induction.

We then show that the sequence $\{f^t\}$ has bounded first, second and third-order gradients under the following assumptions on the ground cost function c.

Assumption B.2. The cost function c is G_c -Lipschitz continuous with respect to one of its inputs: For all $x, x' \in \mathcal{X}$,

$$|c(x,y) - c(x',y)| \le G_c ||x - x'||.$$

Assumption B.3. The gradient of the cost function c is L_c -Lipschitz continuous: for all $x, x' \in \mathcal{X}$,

$$\|\nabla_1 c(x,y) - \nabla_1 c(x',y)\| \le L_c \|x - x'\|.$$

Assumption B.4. The Hessian matrix of the cost function c is $L_{2,c}$ -Lipschitz continuous: for all $x, x' \in \mathcal{X}$,

$$\|\nabla_{11}^2 c(x,y) - \nabla_{11}^2 c(x',y)\| \le L_{2,c} \|x - x'\|.$$

Lemma B.2. Assume that the initialization $f^0 \in C(\mathcal{X})$ satisfies $||f^0||_{\infty} \leq M_c$. (*i.*) Under Assumptions B.1 and B.2, $\exists G_f$ such that $||\nabla f^t||_{2,\infty} \leq G_f, \forall t > 0$. (*ii.*) Under Assumptions B.1 - B.3, $\exists L_f$ such that $||\nabla^2 f^t(x)|| \leq L_f, \forall t > 0$. (*iii.*) Under Assumptions B.1 - B.4, $\exists L_{2,f}$ such that $||\nabla^2 f^t(x) - \nabla^2 f^t(y)||_{op} \leq L_{2,f} ||x - y||, \forall t > 0$. (*iv.*) For $||f||_{\infty} \leq M_c$, the function $\mathcal{B}(f, \theta)(x)$ is G_f -Lipschitz continuous.

Proof. We denote $k(x, y) := \exp\{-c(x, y)/\gamma\}$ in this proof.

(i) Under Assumptions B.1 and B.2, k is $[G_c/\gamma]$ -Lipschitz continuous w.r.t. its first variable. For $f \in \mathcal{C}(\mathcal{X})$ such that $||f||_{\infty} \leq M_c$, we bound

$$\begin{aligned} |\mathcal{A}(f,\alpha)(x) - \mathcal{A}(f,\alpha)(y)| &= \gamma |\log \int_{\mathcal{X}} [k(z,y) - k(z,x)] \exp\{f(z)/\gamma\} \mathbf{d}\alpha(z)| \\ &\leqslant \gamma \exp(M_c/\gamma) G_c/\gamma \|x - y\|_2 = \exp(M_c/\gamma) G_c \|x - y\|_2. \end{aligned}$$

L			
L			
	_	_	_

Using Lemma B.1, we know that $\{f^t\}$ is M_c -bounded and hence

$$\|\nabla f^{t+1}\|_{2,\infty} \leqslant G_f = \exp(2M_c/\gamma)G_c^2.$$

(ii) Under Assumption B.1, $k(x, y) \ge \exp(-M_c/\gamma)$. We compute

$$\nabla \left(\mathcal{A}(f,\alpha) \right)(x) = \frac{\int_{\mathcal{X}} k(z,x) \exp\{f(z)/\gamma\} \nabla_1 c(x,z) \mathbf{d}\alpha(z)}{\int_{\mathcal{X}} k(z,x) \exp\{f(z)/\gamma\} \mathbf{d}\alpha(z)}. \qquad \# \frac{g_1(x)}{g_2(x)}$$

Let $g_1 : \mathbb{R}^q \to \mathbb{R}^q$ and $g_2 : \mathbb{R}^q \to \mathbb{R}$ be the numerator and denominator of the above expression. If we have (a) $\|g_1\|_{2,\infty} \leq G_1$, (b) $\|g_1(x) - g_1(y)\| \leq L_1 \|x - y\|$ and (c) $\|g_2\|_{\infty} \leq G_2$, (d) $|g_2(x) - g_2(y)| \leq L_2 \|x - y\|$, (e) $g_2 \geq \overline{G}_2 > 0$, we can bound

$$\left\|\frac{g_1(x)}{g_2(x)} - \frac{g_1(y)}{g_2(y)}\right\| = \left\|\frac{g_1(x)g_2(y) - g_1(y)g_2(x)}{g_2(x)g_2(y)}\right\| \le \frac{G_2L_1 + G_1L_2}{\bar{G}_2^2} \|x - y\|,\tag{63}$$

which means that $\nabla(\mathcal{A}(f,\alpha))$ is *L*-Lipschitz continuous with $L = \frac{G_2L_1+G_1L_2}{G_2^2}$. We now prove (a)-(e).

(a)
$$\|\int_{\mathcal{X}} k(z,x) \exp\{f(z)/\gamma\} \nabla_1 c(x,z) \mathbf{d}\alpha(z)\|_{2,\infty} \leq \exp(M_c/\gamma) \cdot G_c$$
 (Assumption B.2).

(b) Note that for any two bounded and Lipschitz continuous functions $h_1 : \mathcal{X} \to \mathbb{R}$ and $h_2 : \mathcal{X} \to \mathbb{R}^q$, their product is also Lipschitz continuous:

$$\|h_1(x) \cdot h_2(x) - h_1(y) \cdot h_2(y)\| \le [|h_1|_{\infty} \cdot G_{h_2} + \|h_2\|_{2,\infty} \cdot G_{h_1}]\|x - y\|,$$
(64)

where G_{h_i} denotes the Lipschitz constant of h_i , i = 1, 2. Hence for g_1 , we have

$$\begin{split} \|g_{1}(x) - g_{1}(y)\| &\leq \exp(M_{c}/\gamma) \cdot (L_{c} + G_{c}^{2}/\gamma) \cdot \|x - y\|,\\ \text{since } k(x, y) &\leq 1, \|\nabla_{1}k(x, y)\| \leq G_{c}/\gamma, \|\nabla_{1}c(x, y)\| \leq G_{c}, \|\nabla_{11}^{2}c(x, y)\|_{op} \leq L_{c}.\\ (c) &\|\int_{\mathcal{X}} k(z, \cdot) \exp\{f(z)/\gamma\} \mathbf{d}\alpha(z)\|_{\infty} \leq \exp(M_{c}/\gamma).\\ (d) &|\int_{\mathcal{X}} [k(z, x) - k(z, y)] \exp\{f(z)/\gamma\} \mathbf{d}\alpha(z)| \leq \exp(M_{c}/\gamma) \cdot G_{c}/\gamma \cdot \|x - y\|.\\ (e) &\int_{\mathcal{X}} k(z, x) \exp\{f(z)/\gamma\} \mathbf{d}\alpha(z) \geq \exp(-2M_{c}/\gamma) > 0. \end{split}$$

Combining the above points, we prove the existence of L_f .

For (iii), compute that $\nabla^2 (A(f, \alpha))(x)$

$$= \frac{\int_{\mathcal{X}} k(z,x) \exp\{f(z)/\gamma\} \nabla_1 c(x,z) \nabla_1 c(x,z)^\top \mathbf{d}\alpha(z)}{\int_{\mathcal{X}} k(z,x) \exp\{f(z)/\gamma\} \mathbf{d}\alpha(z)}$$
#1

$$+\frac{\int_{\mathcal{X}} k(z,x) \exp\{f(z)/\gamma\} \nabla_{11}^2 c(x,z) \mathbf{d}\alpha(z)}{\int_{\mathcal{X}} k(z,x) \exp\{f(z)/\gamma\} \mathbf{d}\alpha(z)}$$
#2

$$-\frac{\int_{\mathcal{X}} k(z,x) \exp\{f(z)/\gamma\} \nabla_1 c(x,z) \mathbf{d}\alpha(z) \left[\int_{\mathcal{X}} k(z,x) \exp\{f(z)/\gamma\} \nabla_1 c(x,z) \mathbf{d}\alpha(z)\right]^{\top}}{\left[\int_{\mathcal{X}} k(z,x) \exp\{f(z)/\gamma\} \mathbf{d}\alpha(z)\right]^2}. \quad \#3$$

We now analyze #1-#3 individually.

#1 Note that for any two bounded and Lipschitz continuous functions $h_1 : \mathcal{X} \to \mathbb{R}$ and $h_2 : \mathcal{X} \to \mathbb{R}^{q \times q}$, their product is also Lipschitz continuous:

$$\|h_1(x) \cdot h_2(x) - h_1(y) \cdot h_2(y)\|_{op} \leq [|h_1|_{\infty} \cdot G_{h_2} + \|h_2\|_{op,\infty} \cdot G_{h_1}]\|x - y\|, \quad (65)$$

where G_{h_i} denotes the Lipschitz constant of $h_i, i = 1, 2$.

Take $h_1(x) = k(z', x) \exp\{f(z')/\gamma\}/\int_{\mathcal{X}} k(z, x) \exp\{f(z)/\gamma\} d\alpha(z)$. h_1 is bounded since $k(z', x) \leq 1$ and $\int_{\mathcal{X}} k(z, x) \exp\{f(z)/\gamma\} d\alpha(z) \geq \exp(-2M_c/\gamma) > 0$. h_1 is Lipschitz continuous since we additionally have k(z', x) being Lipschitz continuous (see (63)).

Take $h_2(x) = \nabla_1 c(x, z) \nabla_1 c(x, z)^{\top}$. h_2 is bounded since $\|\nabla_1 c(x, z)\| \leq G_c$ (Assumption B.2). h_2 is Lipschitz continuous due to Assumption B.3.

- #2 Following the similar argument as #1, we have the result. Note that $h_2(x) = \nabla_{11}^2 c(x, z)$ is Lipschitz continuous due to Assumption B.4.
- #3 We follow the similar argument as #1 by taking

$$h_1(x) = \frac{k(z', x) \exp\{f(z')/\gamma\}k(z', x) \exp\{f(z')/\gamma\}}{\left[\int_{\mathcal{X}} k(z, x) \exp\{f(z)/\gamma\} \mathbf{d}\alpha(z)\right]^2},$$

and taking

$$h_2(x) = \nabla_1 c(x, z) [\nabla_1 c(x, z)]^\top.$$

Combining the above points, we prove the existence of $L_{2,f}$.

(iv) As a composition of \mathcal{A} , we also have that $\mathcal{B}(f,\theta)$ is G_f -Lipschitz continuous (see G_f in (i)).

Moreover, based on the above continuity results, we can show that the first-order gradient $\nabla f_{\theta}^{\epsilon}$ (and second-order gradient $\nabla^2 f_{\theta}^{\epsilon}$) also converges to ∇f_{θ} (and $\nabla^2 f_{\theta}$) in time logarithmically depending on $1/\epsilon$.

Lemma B.3. Under Assumptions B.1-B.3, the Sinkhorn-Knopp algorithm, i.e. the fixed point iteration

$$f^{t+1} = \mathcal{B}(f^t, \theta), \tag{66}$$

computes ∇f_{θ} in logarithm time: $\|\nabla f^{t+1} - \nabla f_{\theta}\|_{2,\infty} = \epsilon$ with $t = \mathcal{O}(\log \frac{1}{\epsilon})$.

Proof. For a fix point $x \in \mathcal{X}$ and any direction $h \in \mathbb{R}^q$, we have

$$f^t(x+\eta\cdot h) - f^t(x) = \eta [\nabla f^t(x)]^\top h + \frac{\eta^2}{2} h^\top \nabla^2 f^t(x+\tilde{\eta}_1\cdot h)h,$$

where $\eta > 0$ is some constant to be determined later and $0 \le \tilde{\eta}_1 \le \eta$ is obtained from the mean value theorem. Similarly, we have for $0 \le \tilde{\eta}_2 \le \eta$

$$f_{\theta}(x+\eta\cdot h) - f_{\theta}(x) = \eta [\nabla f_{\theta}(x)]^{\top} h + \frac{\eta^2}{2} h^{\top} \nabla^2 f_{\theta}(x+\tilde{\eta}_2\cdot h)h.$$

We can then compute

$$\left| \left[\nabla f^t(x) - \nabla f_\theta(x) \right]^\top h \right| \leq \frac{2}{\eta} \| f^t - f_\theta \|_\infty + \eta L_f \| h \|^2.$$

Take $h = \nabla f^t(x) - \nabla f_{\theta}(x)$ and $\eta = \frac{2}{L_f}$. We derive from the above inequality

$$\|\nabla f^t(x) - \nabla f_\theta(x)\|^2 \leq 2L_f \|f^t - f_\theta\|_{\infty}.$$

Consequently, if we have $2L_f \|f^t - f_\theta\|_{\infty} \leq \epsilon^2$, we can prove that $\|\nabla f^t - \nabla f_\theta\|_{2,\infty} \leq \epsilon$ since x is arbitrary. This can be achieve in logarithmic time using the Sinkhorn-Knopp algorithm.

Lemma B.4. Under Assumptions B.1-B.4, the Sinkhorn-Knopp algorithm, i.e. the fixed point iteration

$$f^{t+1} = \mathcal{B}(f^t, \theta), \tag{67}$$

computes $\nabla^2 f_{\theta}$ in logarithm time: $\|\nabla^2 f^{t+1} - \nabla^2 f_{\theta}\|_{op,\infty} = \epsilon$ with $t = \mathcal{O}(\log \frac{1}{\epsilon})$.

Proof. This follows a similar argument as Lemma B.3 by noticing that the third order gradient of f^t (and f_{θ}) is bounded due to Assumption B.4.

B.3 Proof of Proposition 5.3

We now construct a sequence $\{g^t\}$ to approximate the Fréchet derivative of the Sinkhorn potential Df_{θ} such that for all $t \ge T(\epsilon)$ with some integer function $T(\epsilon)$ of the target accuracy ϵ , we have $\|g_{\theta}^t - Df_{\theta}\|_{op} \le \epsilon$. In particular, we show that such ϵ -accurate approximation can be achieved using a logarithmic amount of simple function operations and integrations with respect to α_{θ} .

For a given target accuracy $\epsilon > 0$, denote $\bar{\epsilon} = \epsilon/L_l$, where L_l is a constant defined in Lemma B.5. First, Use the Sinkhorn-Knopp algorithm to compute $f_{\theta}^{\bar{\epsilon}}$, an approximation of f_{θ} such that $\|f_{\theta}^{\bar{\epsilon}} - f_{\theta}\|_{\infty} \leq \bar{\epsilon}$. This computation can be done in $\mathcal{O}(\log \frac{1}{\epsilon})$ from Proposition 5.2.

Denote $\mathcal{E}(f,\theta) = \mathcal{B}^l(f,\theta) = \mathcal{B}(\cdots \mathcal{B}(f,\theta),\cdots,\theta)$, the *l* times composition of \mathcal{B} in its first variable. Pick $l = \lceil \log_{\lambda} \frac{1}{3} \rceil / 2$. From the contraction of \mathcal{A} under the Hilbert metric (61), we have

$$\begin{aligned} \|\mathcal{E}(f,\theta) - \mathcal{E}(f',\theta)\|_{\infty} &\leq \gamma d_{H}(\exp(\mathcal{E}(f,\theta)/\gamma), \exp(\mathcal{E}(f',\theta)/\gamma)) \\ &\leq \gamma \lambda^{2l} d_{H}(\exp(f/\gamma), \exp(f'/\gamma)) \leq 2\lambda^{2l} \|f - f'\|_{\infty} \leq \frac{2}{3} \|f - f'\|_{\infty}, \end{aligned}$$

where we use $||f - f'||_{\infty} \leq d_H(\exp(f), \exp(f')) \leq 2||f - f'||_{\infty}$ in the first and third inequalities. Consequently, $\mathcal{E}[f, \theta]$ is a contraction operator w.r.t. f under the l_{∞} norm, which is equivalent to

$$\|D^{1}\mathcal{E}(f,\theta)\|_{op} \leqslant \frac{2}{3}.$$
(68)

Now, given arbitrary initialization $g^0_{\theta} : \Theta \to T(\mathbb{R}^d, \mathcal{C}(\mathcal{X}))^1$, construct iteratively

$$g_{\theta}^{t+1} = D_1 \mathcal{E}(f_{\theta}^{\bar{\epsilon}}, \theta) \circ g_{\theta}^t + D_2 \mathcal{E}(f_{\theta}^{\bar{\epsilon}}, \theta),$$
(69)

where \circ denotes the composition of (linear) mappings. In the following, we show that

$$\|g_{\theta}^{t+1} - Df_{\theta}\|_{op} \leq 3\epsilon + (\frac{2}{3})^t \|g_{\theta}^0 - Df_{\theta}\|_{op}$$

First, note that f_{θ} is a fixed point of $\mathcal{E}(\cdot, \theta)$

$$f_{\theta} = \mathcal{E}(f_{\theta}, \theta).$$

Take the Fréchet derivative w.r.t. θ on both sides of the above equation. Using the chain rule, we compute

$$Df_{\theta} = D_1 \mathcal{E}(f_{\theta}, \theta) \circ Df_{\theta} + D_2 \mathcal{E}(f_{\theta}, \theta).$$
(70)

For any direction $h \in \mathbb{R}^d$, we bound the difference of the directional derivatives by

$$\begin{split} \|g_{\theta}^{t+1}[h] - Df_{\theta}[h]\|_{\infty} \\ &\leq \|D_{1}\mathcal{E}(f_{\theta},\theta) \big[Df_{\theta}[h] \big] - D_{1}\mathcal{E}(f_{\theta}^{\bar{\epsilon}},\theta) \big[g_{\theta}^{t}[h] \big] \|_{\infty} + \|D_{2}\mathcal{E}(f_{\theta}^{\bar{\epsilon}},\theta)[h] - D_{2}\mathcal{E}(f_{\theta},\theta)[h] \|_{\infty} \\ &\leq \frac{2}{3} \|Df_{\theta}[h] - g_{\theta}^{t}[h]\|_{\infty} + L_{l} \big(\|f_{\theta}^{\bar{\epsilon}} - f_{\theta}\|_{\infty} + \|\nabla f_{\theta}^{\bar{\epsilon}} - \nabla f_{\theta}\|_{\infty} \big) \|h\|_{\infty} \\ &\leq \frac{2}{3} \|Df_{\theta} - g_{\theta}^{t}\|_{op} \|h\|_{\infty} + \epsilon \|h\|_{\infty}, \end{split}$$

where in the second inequality we use the bound on $D_1\mathcal{E}$ in (68) and the L_l -Lipschitz continuity of $D_2\mathcal{E}$ with respect to its first argument (recall that $f_{\theta}^{\bar{\epsilon}}$ is obtained from the Sinkhorn-Knopp algorithm and hence $\|f_{\theta}^{\bar{\epsilon}}\|_{\infty} \leq M_c$ from Lemma B.1 and $\|\nabla f_{\theta}^{\bar{\epsilon}}\|_{2,\infty} \leq G_f$ from (i) of Lemma B.2). The above inequality is equivalent to

$$\|g_{\theta}^{t+1} - Df_{\theta}\|_{op} - 3\epsilon \leqslant \frac{2}{3} \left(\|Df_{\theta} - g_{\theta}^{t}\|_{op} - 3\epsilon \right) \Rightarrow \|g_{\theta}^{t+1} - Df_{\theta}\|_{op} \leqslant 3\epsilon + \left(\frac{2}{3}\right)^{t} \|g_{\theta}^{0} - Df_{\theta}\|_{op}.$$

Therefore, after $T(\epsilon) = \mathcal{O}(\log \frac{1}{\epsilon})$ iterations, we find $g_{\theta}^{T(\epsilon)}$ such that $\|g_{\theta}^{T(\epsilon)} - Df_{\theta}\|_{op} \leq 4\epsilon$. **Assumption B.5** (Boundedness of $\nabla_{\theta}T_{\theta}(x)$). There exists some $G_T > 0$ such that for any $x \in \mathcal{X}$ and $\theta \in \Theta$, $\|\nabla_{\theta}T_{\theta}(x)\|_{op} \leq G_T$.

¹Recall that $T(\mathbb{R}^d, \mathcal{C}(\mathcal{X}))$ is the family of bounded linear operators from \mathbb{R}^d to $\mathcal{C}(\mathcal{X})$

Lemma B.5 (Lipschitz continuity of $D_2\mathcal{E}$). Under Assumptions B.1 - B.3 and B.5, $D_2\mathcal{E}$ is Lipschitz continuous with respect to its first variable: For $f, f' \in \mathcal{C}(\mathcal{X})$ such that $||f||_{\infty} \leq M_c$ ($||f'||_{\infty} \leq M_c$) and $||\nabla f||_{\infty} \leq G_f$ ($||\nabla f'||_{\infty} \leq G_f$), and $\theta \in \Theta$ there exists some L_l such that

$$\|D_{2}\mathcal{E}(f,\theta) - D_{2}\mathcal{E}(f',\theta)\|_{op} \leq L_{l}(\|f - f'\|_{\infty} + \|\nabla f - \nabla f'\|_{2,\infty}).$$
(71)

Proof. Recall that $\mathcal{E}(\cdot, \theta) = \mathcal{B}^{l}(\cdot, \theta)$. Using the chain rule of Fréchet derivative, we compute

$$D_2 \mathcal{B}^l(f,\theta) = D_1 \mathcal{B} \big(\mathcal{B}^{l-1}(f,\theta), \theta \big) \circ D_2 \mathcal{B}^{l-1}(f,\theta) + D_2 \mathcal{B} \big(\mathcal{B}^{l-1}(f,\theta), \theta \big).$$
(72)

We bound the two terms on the R.H.S. individually.

Analyze the first term of (72). For a given f, use A_f and B_f to denote two linear operators depending on f. We have $||A_f \circ B_f - A_{f'} \circ B_{f'}||_{op} = \mathcal{O}(||f - f'||_{\infty} + ||\nabla f - \nabla f'||_{2,\infty})$ if both A_f and B_f are bounded, $||A_f - A_{f'}||_{op} = \mathcal{O}(||f - f'||_{\infty} + ||\nabla f - \nabla f'||_{2,\infty})$, and $||B_f - B_{f'}||_{op} = \mathcal{O}(||f - f'||_{\infty} + ||\nabla f - \nabla f'||_{2,\infty})$;

$$|A_{f} \circ B_{f} - A_{f'} \circ B_{f'}||_{op} \leq ||A_{f} \circ B_{f} - A_{f} \circ B_{f'}||_{op} + ||A_{f} \circ B_{f'} - A_{f'} \circ B_{f'}||_{op} \leq [\max_{f} ||B_{f}||_{op} \cdot L_{A} + \max_{f} ||A_{f}||_{op} \cdot L_{B}] [||f - f'||_{\infty} + ||\nabla f - \nabla f'||_{2,\infty}],$$
(73)

where L_A and L_B denote the constants of operators A_f and B_f such that

$$||A_f - A_{f'}|| \leq L_A \Big[||f - f'||_{\infty} + ||\nabla f - \nabla f'||_{2,\infty} \Big] ||B_f - B_{f'}|| \leq L_B \Big[||f - f'||_{\infty} + ||\nabla f - \nabla f'||_{2,\infty} \Big].$$

We now take

$$A_f = D_1 \mathcal{B} (\mathcal{B}^{l-1}(f,\theta),\theta)$$
 and $B_f = D_2 \mathcal{B}^{l-1}(f,\theta)$.

 $||A_f||_{op}$ is bounded from the following lemma.

Lemma B.6. $\mathcal{B}(f,\theta)$ is 1-Lipschitz continuous with respect to its first variable.

Proof. We compute that for any measure κ and any function $g \in C(\mathcal{X})$,

$$D_1 \mathcal{A}(f,\kappa)[g] = \frac{\int_{\mathcal{X}} \exp\{-\frac{1}{\gamma} (c(x,y) - f(x))\}g(x) \mathbf{d}\kappa(x)}{\int_{\mathcal{X}} \exp\{-\frac{1}{\gamma} (c(x,y) - f(x))\} \mathbf{d}\kappa(x)}.$$
(74)

Note that

$$\|D_1\mathcal{A}(f,\kappa)[g]\|_{\infty} \leq \|\frac{\int_{\mathcal{X}} \exp\{-\frac{1}{\gamma} \left(c(x,y) - f(x)\right)\} \mathbf{d}\kappa(x)}{\int_{\mathcal{X}} \exp\{-\frac{1}{\gamma} \left(c(x,y) - f(x)\right)\} \mathbf{d}\kappa(x)}\|_{\infty} \cdot \|g\|_{\infty} = \|g\|_{\infty},$$
(75)

and consequently we have $||D_1\mathcal{A}(f,\kappa)||_{op} \leq 1$. Further, since \mathcal{B} is the composition of \mathcal{A} in its first variable, we have that $||D_1\mathcal{B}(f,\theta)||_{op} \leq 1$.

 $||B_f||_{op}$ is bounded from the following lemma.

Lemma B.7. Assume that $f \in C(\mathcal{X})$ satisfies $||f||_{\infty} \leq M_c$ and $||\nabla f||_{2,\infty} \leq G_f$. Under Assumptions B.2 and B.5, $\forall l \geq 1$, $||D_2\mathcal{B}^l(f,\theta)||_{op}$ is M_l -bounded, with $M_l = l \cdot \exp(3M_c/\gamma) \cdot G_T \cdot (G_c + G_f)$.

Proof. In this proof, we denote $\tilde{\mathcal{A}}(f, \theta) := \mathcal{A}(f, \alpha_{\theta})$ to make the dependence of \mathcal{A} on θ explicit. Using the chain rule of Fréchet derivative, we compute

$$D_2 \mathcal{B}^l(f,\theta) = D_1 \mathcal{B} \big(\mathcal{B}^{l-1}(f,\theta), \theta \big) \circ D_2 \mathcal{B}^{l-1}(f,\theta) + D_2 \mathcal{B} \big(\mathcal{B}^{l-1}(f,\theta), \theta \big).$$
(76)

We will use M_l to denote the upper bound of $||D_2\mathcal{B}^l(f,\theta)||_{op}$. Consequently we have

$$M_{l} \leq \|D_{1}\mathcal{B}(\mathcal{B}^{l-1}(f,\theta),\theta)\|_{op}\|D_{2}\mathcal{B}^{l-1}(f,\theta)\|_{op} + \|D_{2}\mathcal{B}(\mathcal{B}^{l-1}(f,\theta),\theta)\|_{op} \\ \leq M_{l-1} + \|D_{2}\mathcal{B}(\mathcal{B}^{l-1}(f,\theta),\theta)\|_{op},$$

where we use Lemma B.6 in the second inequality. Recall that $\mathcal{B}(f,\theta) = \mathcal{A}(\tilde{\mathcal{A}}(f,\theta),\beta)$. Again using the chain rule of the Fréchet derivative, we compute

$$D_2\mathcal{B}(f,\theta) = D_1\mathcal{A}\big(\tilde{\mathcal{A}}(f,\theta),\beta\big) \circ D_2\tilde{\mathcal{A}}(f,\theta),\tag{77}$$

and hence

$$\|D_2\mathcal{B}(f,\theta)\|_{op} \le \|D_1\mathcal{A}\big(\tilde{\mathcal{A}}(f,\theta),\beta\big)\|_{op} \cdot \|D_2\tilde{\mathcal{A}}(f,\theta)\|_{op} \le \|D_2\tilde{\mathcal{A}}(f,\theta)\|_{op},\tag{78}$$

where we use (75) in the second inequality. We now bound $||D_2 \tilde{\mathcal{A}}(f, \theta)||_{op}$. Denote

 $\omega_y(x) := \exp(-c(x, y)/\gamma) \exp(f(x)/\gamma).$

We have $\exp(-2M_c/\gamma) \leq \omega_y(x) \leq \exp(M_c/\gamma)$ from $||f||_{\infty} \leq M_c$ and Assumption B.1. For any direction $h \in \mathbb{R}^q$ (note that $D_2 \tilde{\mathcal{A}}(f, \theta)[h] : \mathcal{X} \to \mathbb{R}$) and any $y \in \mathcal{X}$, we compute

$$\left(D_2 \tilde{\mathcal{A}}(f,\theta)[h] \right)(y) = \frac{\int_{\mathcal{X}} \omega_y(T_\theta(x)) \langle [\nabla_\theta T_\theta(x)]^\top \left[-\nabla_1 c(T_\theta(x), y) + \nabla f(T_\theta(x)) \right], h \rangle \mathbf{d}\mu(x)}{\int_{\mathcal{X}} \omega_y(T_\theta(x)) \mathbf{d}\mu(x)},$$

where $\nabla_{\theta} T_{\theta}(x)$ denotes the Jacobian matrix of $T_{\theta}(x)$ w.r.t. θ . Consequently we bound

$$\begin{aligned} \|D_2 \tilde{\mathcal{A}}(f,\theta)[h]\|_{\infty} &\leq \exp(3M_c/\gamma) \|\nabla_{\theta} T_{\theta}(x)\|_{op} \cdot [\|\nabla_1 c \big(T_{\theta}(x), y\big)\| + \|\nabla f \big(T_{\theta}(x)\big)\|] \cdot \|h\| \\ &\leq \exp(3M_c/\gamma) \cdot G_T \cdot (G_c + G_f) \|h\|, \end{aligned}$$

which implies

$$\|D_2\tilde{\mathcal{A}}(f,\theta)\|_{op} \leq \exp(3M_c/\gamma) \cdot G_T \cdot (G_c + G_f).$$
(79)

To show the Lipschitz continuity of A_f , i.e. $||A_f - A_{f'}|| \leq L_A ||f - f'||_{\infty}$, we first establish the following continuity lemmas of $D_1 \mathcal{B}(\cdot, \theta)$ and $\mathcal{B}^{l-1}(\cdot, \theta)$.

Lemma B.8. For $f \in C(\mathcal{X})$ such $||f||_{\infty} \leq M_c$, $D_1\mathcal{B}(f,\theta)$ is L-Lipschitz continuous with respect to its first variable with $L = 2L_A$.

Proof. Use the chain rule of Fréchet derivative to compute

$$D_1 \mathcal{B}(f, \theta) = \underbrace{D_1 \mathcal{A}(\mathcal{A}(f, \alpha_{\theta}), \beta)}_{U_f} \circ \underbrace{D_1 \mathcal{A}(f, \alpha_{\theta})}_{V_f}.$$
(80)

We analyze the Lipschitz continuity of $||D_1\mathcal{B}(f,\theta)||_{op}$ following the same logic as (73):

- The 1-boundedness of U_f and V_f is from Lemma B.6.
- The L_A -Lipschitz continuity of V_f is from Lemma B.11.
- The L_A -Lipschitz continuity of U_f is from Lemmas B.6 and B.11.

Consequently, we have that $D_1\mathcal{B}(f,\theta)$ is $2L_{\mathcal{A}}$ -Lipschitz continuous w.r.t. its first variable. **Lemma B.9.** $\forall l, \mathcal{B}^l(f,\theta)$ is 1-Lipschitz continuous with respect to its first variable.

Proof. Use the chain rule of Fréchet derivative to compute

$$D_1 \mathcal{B}^l(f,\theta) = D_1 \mathcal{B}\left(\mathcal{B}^{l-1}(f,\theta),\theta\right) \circ D_1 \mathcal{B}^{l-1}(f,\theta).$$
(81)

Consequently $||D_1\mathcal{B}^l(f,\theta)||_{op} \leq ||D_1\mathcal{B}(f,\theta)||_{op}^l$. Further, we have $||D_1\mathcal{B}(f,\theta)||_{op} \leq 1$ from Lemma B.6 which leads to the result.

We have that A_f is Lipschitz continuous since (i) A_f is the composition of Lipschitz continuous operators $D_1\mathcal{B}(\cdot,\theta)$ and $\mathcal{B}^{l-1}(f\cdot,\theta)$ and (ii) for $||f||_{\infty} \leq M_c$, $\forall l \geq 0$, $||\mathcal{B}^l(f,\theta)||_{\infty} \leq M_c$ (the argument is similar to Lemma B.1).

We prove $||B_f - B_{f'}|| \leq L_l [||f - f'||_{\infty} + ||\nabla f - \nabla f'||_{2,\infty}]$ via induction. The following lemma establishes the base case for $D_2\mathcal{B}(f,\theta)$ (when l = 2). Note that the boundedness of $||f||_{\infty} (||f'||_{\infty})$ and $||\nabla f||_{\infty} (||\nabla f'||_{\infty})$ remains valid after the operator \mathcal{B} (Lemma B.1 and (i) of Lemma (B.2)).

Lemma B.10. There exists constant L_1 such that for $||f||_{\infty} \leq M_c$ ($||f'||_{\infty} \leq M_c$) and $||\nabla f||_{\infty} \leq G_f$ ($||\nabla f'||_{\infty} \leq G_f$)

$$\|D_{2}\mathcal{B}(f,\theta) - D_{2}\mathcal{B}(f',\theta)\|_{op} \leq L_{1} \Big[\|f - f'\|_{\infty} + \|\nabla f - \nabla f'\|_{2,\infty}\Big].$$
(82)

Proof. In this proof, we denote $\tilde{\mathcal{A}}(f,\theta):=\mathcal{A}(f,\alpha_{\theta})$ to make the dependence of \mathcal{A} on θ explicit. Recall that $\mathcal{B}(f,\theta) = \mathcal{A}(\tilde{\mathcal{A}}(f,\theta),\beta)$. Use the chain rule of Fréchet derivative to compute

$$D_2 \mathcal{B}(f,\theta) = \underbrace{D_1 \mathcal{A}(\mathcal{A}(f,\alpha_\theta),\beta)}_{U_f} \circ \underbrace{D_2 \tilde{\mathcal{A}}(f,\theta)}_{V_f}.$$
(83)

We analyze the Lipschitz continuity of $||D_2\mathcal{B}(f,\theta)||_{op}$ following the same logic as (73):

- The 1-boundedness of U_f is from Lemma B.6.
- The $\exp(3M_c/\gamma) \cdot G_T \cdot (G_c + G_f)$ -boundedness of V_f is from (79).
- The L_A -Lipschitz continuity of U_f is from Lemmas B.6 and B.11 and the fact that for $||f||_{\infty} \leq M_c$, $||\mathcal{A}(f, \theta)||_{\infty} \leq M_c$ (the argument is similar to Lemma B.1).
- Denote

$$\mathcal{T}_y(x, f) := \exp(-c(x, y)/\gamma) \exp(f(x)/\gamma).$$

We compute

$$V_f = \frac{\int_{\mathcal{Z}} \mathcal{T}_y \big(T_\theta(z), f \big) [\nabla_\theta T_\theta(z)]^\top \big[-\nabla_1 c(T_\theta(z), y) + \nabla f \big(T_\theta(z) \big) \big] \mathbf{d} \mu(z)}{\int_{\mathcal{Z}} \mathcal{T}_y \big(T_\theta(z), f \big) \mathbf{d} \mu(z)}, \quad \# \frac{P_f}{Q_f}$$

Denote the numerator by P_f and the denominator by Q_f . Following the similar idea as (63), we show that both $||P_f||_{op}$ and $||Q_f||_{\infty}$ are bounded, Q_f is Lipschitz continuous w.r.t. f, Q_f is positive and bounded from below, and $||P_f - P_{f'}||_{op} \leq L_v[||f - f'||_{\infty} + ||\nabla f - \nabla f'||_{2,\infty}]$ for some constant L_v .

- The boundedness of $||P_f||_{op}$ is from the boundedness of f, Assumptions B.5, B.2, and the boundedness of ∇f .
- The boundedness of $||Q_f||_{\infty}$ is from the boundedness of f.
- Use DQ_f to denote the Fréchet derivative of Q_f w.r.t. f. For any function $g \in C(\mathcal{X})$,

$$DQ_f[g] = \int_{\mathcal{X}} \mathcal{T}_y(x, f) g(x) / \gamma \mathbf{d}\alpha_\theta(x), \tag{84}$$

where we recall that $\alpha_{\theta} = T_{\theta \sharp} \mu$. Further, we have $\|DQ_f[g]\|_{\infty} \leq \exp(M_c/\gamma)/\gamma \|g\|_{\infty}$, which implies the Lipschitz continuity of Q_f (for $\|f\|_{\infty} \leq M_c$).

- We prove that for $||f||_{\infty} \leq M_c$ $(||f'||_{\infty} \leq M_c)$ and $||\nabla f||_{\infty} \leq G_f$ $(||\nabla f'||_{\infty} \leq G_f)$,

$$||P_f - P_{f'}||_{op} \leq L_v [||f - f'||_{\infty} + ||\nabla f - \nabla f'||_{2,\infty}].$$

For a fixed $z \in \mathcal{Z}$, denote

$$p_f^z := \mathcal{T}_y \big(T_\theta(z), f \big) [\nabla_\theta T_\theta(z)]^\top \left[-\nabla_1 c(T_\theta(z), y) + \nabla f \big(T_\theta(z) \big) \right].$$

Note that $P_f = \int_{\mathcal{Z}} p_f^z \mathbf{d}\mu(z)$. For any direction $h \in \mathbb{R}^d$, we bound

$$\begin{split} \|p_f^z[h] - p_{f'}^z[h]\|_{op} \\ \leqslant \|D_2 \mathcal{T}_y \big(T_\theta(z), f \big) \|_{op} \|f - f'\|_{\infty} \cdot \max_y \left| [\nabla_\theta T_\theta(z)h]^\top \left[-\nabla_1 c(T_\theta(z), y) + \nabla f \big(T_\theta(z) \big) \right] \right| \\ &+ \left[\max_y \mathcal{T}_y \big(T_\theta(z), f \big) \right] \cdot \|\nabla_\theta T_\theta(z)h\| \|\nabla f \big(T_\theta(z) \big) - \nabla f' \big(T_\theta(z) \big) \| \\ \leqslant \exp(M_c/\gamma)/\gamma \cdot G_T \cdot (G_c + G_f) \cdot \|f - f'\|_{\infty} \cdot \|h\| + \exp(M_c/\gamma) \cdot G_T \cdot \|h\| \cdot \|\nabla f - \nabla f'\|_{2,\infty}. \end{split}$$

Consequently, we have that there exists a constant L_v such that

$$\|p_{f}^{z}[h] - p_{f'}^{z}[h]\|_{\infty} \leq L_{v}[\|f - f'\|_{\infty} + \|\nabla f - \nabla f'\|_{2,\infty}] \cdot \|h\|$$

The above lemma shows the base case for the induction. Now suppose that the inequality $||D_2\mathcal{B}^k(f,\theta) - D_2\mathcal{B}^k(f',\theta)||_{op} \leq L_k [||f - f'||_{\infty} + ||\nabla f - \nabla f'||_{2,\infty}]$ holds. For the case of k + 1, we compute the Fréchet derivative

$${}_{2}\mathcal{B}^{k+1}(f,\theta) = D_{1}\mathcal{B}\big(\mathcal{B}^{k}(f,\theta),\theta\big) \circ D_{2}\mathcal{B}^{k}(f,\theta) + D_{2}\mathcal{B}\big(\mathcal{B}^{k}(f,\theta),\theta\big)$$

and hence we can bound

D

$$\begin{split} \|D_{2}\mathcal{B}^{k+1}(f,\theta) - D_{2}\mathcal{B}^{k+1}(f',\theta)\|_{op} \\ &\leq \|D_{1}\mathcal{B}\big(\mathcal{B}^{k}(f,\theta),\theta\big) \circ \big(D_{2}\mathcal{B}^{k}(f,\theta) - D_{2}\mathcal{B}^{k}(f',\theta)\big)\big\|_{op} \\ &+ \|\Big(D_{1}\mathcal{B}\big(\mathcal{B}^{k}(f,\theta),\theta\big) - D_{1}\mathcal{B}\big(\mathcal{B}^{k}(f',\theta),\theta\big)\Big) \circ D_{2}\mathcal{B}^{k}(f',\theta)\|_{op} \\ &+ \|D_{2}\mathcal{B}\big(\mathcal{B}^{k}(f,\theta),\theta\big) - D_{2}\mathcal{B}\big(\mathcal{B}^{k}(f',\theta),\theta\big)\big\|_{op} \\ &\leq \|D_{2}\mathcal{B}^{k}(f,\theta) - D_{2}\mathcal{B}^{k}(f',\theta)\|_{op} \\ &+ L_{\mathcal{A}}\|\mathcal{B}^{k}(f,\theta) - \mathcal{B}^{k}(f',\theta)\|_{\infty}\|D_{2}\mathcal{B}^{k}(f',\theta)\|_{op} \\ &+ L_{1}\big[\|\mathcal{B}^{k}(f,\theta) - \mathcal{B}^{k}(f',\theta)\|_{\infty} + \|\nabla\mathcal{B}^{k}(f,\theta) - \nabla\mathcal{B}^{k}(f',\theta)\|_{2,\infty}\big] \\ &\leq L_{k}\big[\|f - f'\|_{\infty} + \|\nabla f - \nabla f'\|_{2,\infty}\big] + L_{\mathcal{A}} \cdot M_{k} \cdot \|f - f'\|_{\infty} \\ &+ L_{1}\|f - f'\|_{\infty} + L_{1}\|\nabla\mathcal{B}^{k}(f,\theta) - \nabla\mathcal{B}^{k}(f',\theta)\|_{2,\infty} \\ &\leq (L_{k} + L_{1} + L_{\mathcal{A}}\mathcal{M}_{k})\big[\|f - f'\|_{\infty} + \|\nabla f - \nabla f'\|_{2,\infty}\big] + L_{1}\|\nabla\mathcal{B}^{k}(f,\theta) - \nabla\mathcal{B}^{k}(f',\theta)\|_{2,\infty}. \tag{86}$$

Here in the third inequality, we use the induction for the first term, Lemma B.7 for the second term. Notice that $\nabla \mathcal{A}(f, \theta)$ is Lipschitz continuous w.r.t. f: Denote $k(x, y) := \exp\{-c(x, y)/\gamma\}$. For any fixed $x \in \mathcal{X}$,

$$\nabla \left(\mathcal{A}(f,\alpha) \right)(x) = \frac{\int_{\mathcal{X}} k(z,x) \exp\{f(z)/\gamma\} \nabla_1 c(x,z) \mathbf{d}\alpha(z)}{\int_{\mathcal{X}} k(z,x) \exp\{f(z)/\gamma\} \mathbf{d}\alpha(z)}, \qquad \qquad \# \frac{g_1(f)}{g_2(f)}$$

where we denote the numerator and denominator of the above expression by $g_1 : \mathcal{C}(\mathcal{X}) \to \mathbb{R}^q$ and $g_2 : \mathcal{C}(\mathcal{X}) \to \mathbb{R}$. From the boundedness of g_1 and g_2 , the Lipschitz continuity of g_1 and g_2 w.r.t. to f, and the fact that g_2 is positive and bounded away from zero, we conclude that there exists some constant $L_{\mathcal{A},f}$ such that for any $x \in \mathcal{X}$ (this follows similarly as (63))

$$\|\nabla \big(\mathcal{A}(f,\alpha)\big)(x) - \nabla \big(\mathcal{A}(f',\alpha)\big)(x)\| \leq L_{\mathcal{A},f} \|f - f'\|_{\infty}.$$
(87)

Recall that \mathcal{B}^k is the compositions of operators in the form of \mathcal{A} . Consequently, we have that

$$\|\nabla \mathcal{B}^k(f,\theta) - \nabla \mathcal{B}^k(f',\theta)\|_{2,\infty} \leq L_{\mathcal{A},f} \|f - f'\|_{\infty}$$

Plugging this result into (86), we prove that the induction holds for k + 1:

 $\|D_{2}\mathcal{B}^{k+1}(f,\theta) - D_{2}\mathcal{B}^{k+1}(f',\theta)\|_{op} \leq (L_{k} + L_{1} + L_{\mathcal{A}}M_{k} + L_{1}L_{\mathcal{A},f})[\|f - f'\|_{\infty} + \|\nabla f - \nabla f'\|_{2,\infty}].$ Consequently, for any finite *l*, we have $\|B_{f} - B_{f'}\| \leq L_{l}[\|f - f'\|_{\infty} + \|\nabla f - \nabla f'\|_{2,\infty}]$, where $L_{l} = l \cdot (L_{1} + L_{\mathcal{A}}M_{k} + L_{1}L_{\mathcal{A},f}).$

Lemma B.11. Under Assumption B.1, for $f \in C(\mathcal{X})$ such $||f||_{\infty} \leq M_c$, there exists constant $L_{\mathcal{A}}$ such that $D_1\mathcal{A}(f, \alpha_{\theta})$ is $L_{\mathcal{A}}$ -Lipschitz continuous with respect to its first variable.

Proof. Let $g \in C(\mathcal{X})$ any function. Denote $\mathcal{T}_y(x, f) := \exp(-c(x, y)/\gamma) \exp(f(x)/\gamma)$. For a fixed point $y \in \mathcal{X}$ and any function $g \in C(\mathcal{X})$, we compute that

$$\left(D_1 \mathcal{A}(f,\theta)[g]\right)(y) = \frac{\int_{\mathcal{X}} \mathcal{T}_y(x,f) g(x) \mathbf{d}\alpha_\theta(x)}{\int_{\mathcal{X}} \mathcal{T}_y(x,f) \mathbf{d}\alpha_\theta(x)}, \qquad \qquad \# \frac{g_1(f)}{g_2(f)}$$

where we denote the numerator and denominator of the above expression by $g_1 : \mathcal{C}(\mathcal{X}) \to \mathbb{R}^q$ and $g_2 : \mathcal{C}(\mathcal{X}) \to \mathbb{R}$. From the boundedness of g_1 and g_2 , the Lipschitz continuity of g_1 and g_2 w.r.t. to f, and the fact that g_2 is positive and bounded away from zero, we conclude that there exists some constant $L_{\mathcal{A}}$ such that for any $x \in \mathcal{X}$ (this follows similarly as (63)).

Analyze the second term of (72). We bound the second term of (72) using Lemma B.10:

$$\begin{split} \|D_2 \mathcal{B}(\mathcal{B}^{l-1}(f,\theta),\theta) - D_2 \mathcal{B}(\mathcal{B}^{l-1}(f',\theta),\theta)\|_{op} \\ &\leq L_1[\|\mathcal{B}^{l-1}(f,\theta) - \mathcal{B}^{l-1}(f',\theta)\|_{\infty} + \|\nabla \mathcal{B}^{l-1}(f,\theta) - \nabla \mathcal{B}^{l-1}(f',\theta)\|_{2,\infty}] \\ &\leq L_1[\|f - f'\|_{\infty} + L_{\mathcal{A},f}\|f - f'\|_{\infty}] = L_1 \cdot (1 + L_{\mathcal{A},f})\|f - f'\|_{\infty}, \end{split}$$

where we use (87) in the second inequality.

Combing the analysis for the two terms of (72), we conclude the result.

B.4 Proof of Theorem 5.1

We prove that the approximation error of $\nabla^2_{\theta} OT_{\gamma}(\alpha_{\theta}, \beta)$ using the estimated Sinkhorn potential f^{ϵ}_{θ} and the estimated Fréchet derivative g^{ϵ}_{θ} is of the order

$$\mathcal{O}(\|f_{\theta}^{\epsilon} - f_{\theta}\|_{\infty} + \|\nabla f_{\theta}^{\epsilon} - \nabla f_{\theta}\|_{2,\infty} + \|\nabla^2 f_{\theta}^{\epsilon} - \nabla^2 f_{\theta}\|_{op,\infty} + \|g_{\theta}^{\epsilon} - Df_{\theta}\|_{op}).$$

The other term $\nabla^2_{\theta} OT_{\gamma}(\alpha_{\theta}, \alpha_{\theta})$ is handled in a similar manner.

Recall the simplified expression of $\nabla^2_{\theta} OT_{\gamma}(\alpha_{\theta}, \beta)$ in (52). Given the estimator $f^{\epsilon}_{\theta}(g^{\epsilon}_{\theta})$ of $f_{\theta}(Df_{\theta})$, we need to prove the following bounds of differences in terms of the estimation accuracy: For any $h_1, h_2 \in \mathbb{R}^d$,

$$\begin{aligned} |D_{11}^{2}\mathcal{H}_{1}(f_{\theta},\theta) \Big[Df_{\theta}[h_{1}], Df_{\theta}[h_{2}] \Big] - D_{11}^{2}\mathcal{H}_{1}(f_{\theta}^{\epsilon},\theta) \Big[g_{\theta}^{\epsilon}[h_{1}], g_{\theta}^{\epsilon}[h_{2}] \Big] | \\ &= \mathcal{O}\left(\|h_{1}\| \cdot \|h_{2}\| \cdot \left(\|f_{\theta}^{\epsilon} - f_{\theta}\|_{\infty} + \|g_{\theta}^{\epsilon} - Df_{\theta}\|_{op} \right) \right), \end{aligned}$$

$$\begin{aligned} \|D^{2}\mathcal{H}_{1}(f_{0},\theta) - D^{2}\mathcal{H}_{2}(f^{\epsilon},\theta)\| \end{aligned}$$

$$(88)$$

$$\|\mathcal{D}_{22}\mathcal{H}_{1}(f_{\theta}, \theta) - \mathcal{D}_{22}\mathcal{H}_{1}(f_{\theta}, \theta)\|_{op} = \mathcal{O}\left(\|f_{\theta}^{\epsilon} - f_{\theta}\|_{\infty} + \|\nabla f_{\theta}^{\epsilon} - \nabla f_{\theta}\|_{2,\infty} + \|\nabla^{2} f_{\theta}^{\epsilon} - \nabla^{2} f_{\theta}\|_{op,\infty}\right).$$

$$(89)$$

Note that from the definition of the operator norm the first results is equivalent to the bound in the operator norm. Using Propositions 5.2 and 5.3 and Lemmas B.3, B.4, we know that we can compute the estimators f_{θ}^{ϵ} and g_{θ}^{ϵ} such that $\|f_{\theta}^{\epsilon} - f_{\theta}\|_{\infty} \leq \epsilon$, $\|\nabla f_{\theta}^{\epsilon} - \nabla f_{\theta}\|_{2,\infty} \leq \epsilon$, and $\|\nabla^2 f_{\theta}^{\epsilon} - \nabla^2 f_{\theta}\|_{op,\infty} \leq \epsilon$, and $\|g_{\theta}^{\epsilon} - Df_{\theta}\|_{op} \leq \epsilon$ in logarithm time $\mathcal{O}(\log \frac{1}{\epsilon})$. Together with (88) and (89) proved above, we can compute an ϵ -accurate estimation of $\nabla_{\theta}^2 OT_{\gamma}(\alpha_{\theta}, \beta)$ (in the operator norm) in logarithm time $\mathcal{O}(\log \frac{1}{\epsilon})$.

Bounding (88). Recall the definition of $D_{11}^2 \mathcal{H}_1(f_\theta, \theta) [Df_\theta[h_1], Df_\theta[h_2]]$ in (56). Denote

$$A_{1} = D_{11}^{2} \mathcal{A}(f_{\theta}, \alpha_{\theta}), v_{1} = Df_{\theta}[h_{1}], v_{2} = Df_{\theta}[h_{2}], A_{2} = D_{11}^{2} \mathcal{A}(f_{\theta}^{\epsilon}, \alpha_{\theta}), u_{1} = g_{\theta}^{\epsilon}[h_{1}], u_{2} = g_{\theta}^{\epsilon}[h_{2}].$$

Based on these definitions, we have

$$D_{11}^{2}\mathcal{H}_{1}(f_{\theta},\theta)\left[Df_{\theta}[h_{1}], Df_{\theta}[h_{2}]\right] = \int_{\mathcal{X}} A_{1}[v_{1},v_{2}](y)\mathbf{d}\beta(y)$$
$$D_{11}^{2}\mathcal{H}_{1}(f_{\theta}^{\epsilon},\theta)\left[g_{\theta}^{\epsilon}[h_{1}], g_{\theta}^{\epsilon}[h_{2}]\right] = \int_{\mathcal{X}} A_{2}[u_{1},u_{2}](y)\mathbf{d}\beta(y).$$

Using the triangle inequality, we have

$$||A_{1}[v_{1}, v_{2}] - A_{2}[u_{1}, u_{2}]||_{\infty}$$

$$\leq ||A_{1}[v_{1} - u_{1}, v_{2}]||_{\infty} + ||A_{1}[u_{1}, v_{2} - u_{2}]||_{\infty} + ||(A_{1} - A_{2})[u_{1}, u_{2}]||_{\infty}.$$
(90)

We bound the three terms on the R.H.S. individually.

For the first term on the R.H.S. of (90), we recall the explicit expression of $A_1[v_1, v_2](y)$ in (55) as

$$A_{1}[v_{1}, v_{2}](y) = \frac{\int_{\mathcal{X}} \mathcal{T}_{y}(x, f_{\theta}) v_{1}(x) v_{2}(x) \mathbf{d}\alpha_{\theta}(x)}{\gamma \int_{\mathcal{X}} \mathcal{T}_{y}(x, f_{\theta}) \mathbf{d}\alpha_{\theta}(x)} - \frac{\int_{\mathcal{X}^{2}} \mathcal{T}_{y}(x, f_{\theta}) \mathcal{T}_{y}(x', f_{\theta}) v_{1}(x) v_{2}(x') \mathbf{d}\alpha_{\theta}(x) \mathbf{d}\alpha_{\theta}(x')}{\gamma \left[\int_{\mathcal{X}} \mathcal{T}_{y}(x, f_{\theta}) \mathbf{d}\alpha_{\theta}(x)\right]^{2}}$$

Here we recall $\mathcal{T}_y(x, f) := \exp(-c(x, y)/\gamma) \exp(f(x)/\gamma)$. We bound using the facts that $\mathcal{T}_y(x, f_\theta)$ is bounded from above and bounded away from zero

$$|A_{1}[v_{1} - u_{1}, v_{2}](y)| \leq \left|\frac{\int_{\mathcal{X}} \mathcal{T}_{y}(x, f_{\theta}) (v_{1}(x) - u_{1}(x)) v_{2}(x) \mathbf{d}\alpha_{\theta}(x)}{\gamma \int_{\mathcal{X}} \mathcal{T}_{y}(x, f_{\theta}) \mathbf{d}\alpha_{\theta}(x)} + \left|\frac{\int_{\mathcal{X}^{2}} \mathcal{T}_{y}(x, f_{\theta}) \mathcal{T}_{y}(x', f_{\theta}) (v_{1}(x) - u_{1}(x)) v_{2}(x') \mathbf{d}\alpha_{\theta}(x) \mathbf{d}\alpha_{\theta}(x')}{\gamma \left[\int_{\mathcal{X}} \mathcal{T}_{y}(x, f_{\theta}) \mathbf{d}\alpha_{\theta}(x)\right]^{2}}\right| \\ = \mathcal{O}(\|v_{1} - u_{1}\|_{\infty} \cdot \|v_{2}\|_{\infty}).$$

Further, we have $||u_1 - v_1||_{\infty} = \mathcal{O}(||Df_{\theta} - g_{\theta}^{\epsilon}||_{op} \cdot ||h_1||)$ and $||v_1||_{\infty} = \mathcal{O}(||h_2||)$. Consequently, the first term on the R.H.S. of (90) is of order $\mathcal{O}(||Df_{\theta} - g_{\theta}^{\epsilon}||_{op} \cdot ||h_1|| \cdot ||h_2||)$.

Following the same argument, we have the second term on the R.H.S. of (90) is of order $\mathcal{O}(\|Df_{\theta} - g_{\theta}^{\epsilon}\|_{op} \cdot \|h_1\| \cdot \|h_2\|)$.

To bound the third term on the R.H.S. of (90), denote

$$A_{11}[u_1, u_2] := \frac{\int_{\mathcal{X}} \mathcal{T}_y(x, f_\theta) u_1(x) u_2(x) \mathbf{d}\alpha_\theta(x)}{\gamma \int_{\mathcal{X}} \mathcal{T}_y(x, f_\theta) \mathbf{d}\alpha_\theta(x)} \text{ and } A_{21}[u_1, u_2] := \frac{\int_{\mathcal{X}} \mathcal{T}_y(x, f_\theta^{\epsilon}) u_1(x) u_2(x) \mathbf{d}\alpha_\theta(x)}{\gamma \int_{\mathcal{X}} \mathcal{T}_y(x, f_\theta^{\epsilon}) \mathbf{d}\alpha_\theta(x)}$$

and denote

$$A_{12}[u_1, u_2] := \frac{\int_{\mathcal{X}} \mathcal{T}_y(x, f_\theta) u_1(x) \mathbf{d}\alpha_\theta(x) \int_{\mathcal{X}} \mathcal{T}_y(x', f_\theta) u_2(x') \mathbf{d}\alpha_\theta(x')}{\gamma \left[\int_{\mathcal{X}} \mathcal{T}_y(x, f_\theta) \mathbf{d}\alpha_\theta(x)\right]^2},$$

and
$$A_{22}[u_1, u_2] := \frac{\int_{\mathcal{X}} \mathcal{T}_y(x, f_\theta^{\epsilon}) u_1(x) \mathbf{d}\alpha_\theta(x) \int_{\mathcal{X}} \mathcal{T}_y(x', f_\theta^{\epsilon}) u_2(x') \mathbf{d}\alpha_\theta(x')}{\gamma \left[\int_{\mathcal{X}} \mathcal{T}_y(x, f_\theta^{\epsilon}) \mathbf{d}\alpha_\theta(x)\right]^2}.$$

We show that both $|(A_{11} - A_{21})[u_1, u_2]|$ and $|(A_{12} - A_{22})[u_1, u_2]|$ are of order $\mathcal{O}(||Df_{\theta} - g_{\theta}^{\epsilon}||_{op} \cdot ||h_1|| \cdot ||h_2||)$. $||h_1|| \cdot ||h_2||)$. This then implies $|(A_1 - A_2)[u_1, u_2]| = \mathcal{O}(||Df_{\theta} - g_{\theta}^{\epsilon}||_{op} \cdot ||h_1|| \cdot ||h_2||)$. With the argument similar to (63), we obtain that $|(A_{11} - A_{21})[u_1, u_2]| = \mathcal{O}(||Df_{\theta} - g_{\theta}^{\epsilon}||_{op} \cdot ||u_1|| \cdot ||u_2||)$ using the boundedness and Lipschitz continuity of the numerator and denominator of $A_{11}[u_1, u_2]$ w.r.t. to f_{θ} and the fact that the denominator is positive and bounded away from zero (see the discussion following (63)). Further, since both Df_{θ} and g_{θ}^{ϵ} are bounded linear operators, we have that $u_1 = \mathcal{O}(h_1)$ and $u_2 = \mathcal{O}(h_2)$. Consequently, we prove that $|(A_{11} - A_{21})[u_1, u_2]| = \mathcal{O}(||f_{\theta} - f_{\theta}^{\epsilon}||_{op} \cdot ||h_1|| \cdot ||h_2||)$. Similarly, we can prove that $|(A_{12} - A_{22})[u_1, u_2]| = \mathcal{O}(||f_{\theta} - f_{\theta}^{\epsilon}||_{op} \cdot ||h_1|| \cdot ||h_2||)$.

Altogether, we have proved (88).

Bounding (89). Recall that the expression of $D_{22}^2 \mathcal{H}_1(f, \theta)$ in (58). For a fixed $y \in \mathcal{X}$ and a fixed $z' \in \mathcal{Z}$, denote (recall that $u_z(\theta, f) = \nabla_1 c(T_\theta(z), y) - \nabla f(T_\theta(z))$)

$$\begin{split} B_{1}(f) &= \nabla_{\theta}^{2} T_{\theta}(z') \times_{1} \nabla f\left(T_{\theta}(z')\right) \\ B_{2}(f) &= \nabla_{\theta} T_{\theta}(z')^{\top} \nabla^{2} f\left(T_{\theta}(z')\right) \nabla_{\theta} T_{\theta}(z') \\ B_{3}(f) &= \frac{\int_{\mathbb{Z}} \mathcal{T}_{y}\left(T_{\theta}(z), f\right) \nabla_{\theta} T_{\theta}(z)^{\top} u_{z}(\theta, f) u_{z}(\theta, f)^{\top} \nabla_{\theta} T_{\theta}(z) \mathbf{d} \mu(z)}{\int_{\mathbb{Z}} \mathcal{T}_{y}\left(T_{\theta}(z), f\right) \nabla_{\theta}^{2} T_{\theta}(z) \times_{1} u_{z}(\theta, f) \mathbf{d} \mu(z)} \\ B_{4}(f) &= \frac{\int_{\mathbb{Z}} \mathcal{T}_{y}\left(T_{\theta}(z), f\right) \nabla_{\theta}^{2} T_{\theta}(z) \times_{1} u_{z}(\theta, f) \mathbf{d} \mu(z)}{\int_{\mathbb{Z}} \mathcal{T}_{y}\left(T_{\theta}(z), f\right) \nabla_{\theta} T_{\theta}(z)^{\top} \nabla_{11} c(T_{\theta}(z), y) \nabla_{\theta} T_{\theta}(z) \mathbf{d} \mu(z)} \\ B_{5}(f) &= \frac{\int_{\mathbb{Z}} \mathcal{T}_{y}\left(T_{\theta}(z), f\right) \nabla_{\theta} T_{\theta}(z)^{\top} \nabla^{2} f\left(T_{\theta}(z)\right) \nabla_{\theta} T_{\theta}(z) \mathbf{d} \mu(z)}{\int_{\mathbb{Z}} \mathcal{T}_{y}\left(T_{\theta}(z), f\right) \mathbf{d} \mu(z)} \\ B_{6}(f) &= -\frac{\int_{\mathbb{Z}} \mathcal{T}_{y}\left(T_{\theta}(z), f\right) \nabla_{\theta} T_{\theta}(z)^{\top} u_{z}(\theta, f) \mathbf{d} \mu(z)}{\int_{\mathbb{Z}} \mathcal{T}_{y}\left(T_{\theta}(z), f\right) \nabla_{\theta} T_{\theta}(z)^{\top} u_{z}(\theta, f) \mathbf{d} \mu(z)} \\ B_{7}(f) &= \frac{\int_{\mathbb{Z}} \mathcal{T}_{y}\left(T_{\theta}(z), f\right) \nabla_{\theta} T_{\theta}(z)^{\top} u_{z}(\theta, f) \mathbf{d} \mu(z) \left[\int_{\mathbb{Z}} \mathcal{T}_{y}\left(T_{\theta}(z), f\right) \nabla_{\theta} T_{\theta}(z)^{\top} u_{z}(\theta, f) \mathbf{d} \mu(z)\right]^{\top}}{\left[\int_{\mathbb{Z}} \mathcal{T}_{y}\left(T_{\theta}(z), f\right) \mathbf{d} \mu(z)\right]^{2}} \end{split}$$

Based on these definitions, we have

$$D_{22}^2 \mathcal{H}_1(f,\theta) = \int_{\mathcal{Z}} \sum_{i=1}^2 B_i(f) \mathbf{d}\mu(z') + \int_{\mathcal{X}} \sum_{i=3}^7 B_i(f) \mathbf{d}\beta(y).$$

We bound the above seven terms individually.

Assumption B.6. For a fixed $z \in \mathcal{Z}$ and $\theta \in \Theta$, use $\nabla_{\theta}^2 T_{\theta}(z) \in T(\mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}^q)^2$ to denote the second-order Jacobian of $T_{\theta}(z)$ w.r.t. θ . Use \times_1 to denote the tensor product along the first dimension. For any two vectors $g, g' \in \mathbb{R}^d$, we assume that

$$\|\nabla_{\theta}^2 T_{\theta}(z) \times_1 g - \nabla_{\theta}^2 T_{\theta}(z) \times_1 g'\|_{op} = \mathcal{O}(\|g - g'\|).$$
(91)

For the first term, using the boundedness of $\nabla^2_{\theta} T_{\theta}(z')$ (Assumption B.6), we have that

$$||B_1(f_\theta) - B_1(f_\theta^{\epsilon})||_{op} = \mathcal{O}(||\nabla f_\theta - \nabla f_\theta^{\epsilon}||_{2,\infty}).$$

For the second term, using the boundedness of $\nabla_{\theta} T_{\theta}(z')$, we have that

$$\|B_2(f_\theta) - B_2(f_\theta^\epsilon)\|_{op} = \mathcal{O}(\|\nabla^2 f_\theta - \nabla^2 f_\theta^\epsilon\|_{op,\infty}).$$

For the third term, note that $||u_z(\theta, f_\theta) - u_z(\theta, f_\theta^{\epsilon})|| = \mathcal{O}(||\nabla f_\theta - \nabla f_\theta^{\epsilon}||_{2,\infty})$. With the argument similar to (63), we obtain that

$$\|B_3(f_\theta) - B_3(f_\theta^\epsilon)\|_{op} = \mathcal{O}(\|f_\theta - f_\theta^\epsilon\|_{\infty} + \|\nabla f_\theta - \nabla f_\theta^\epsilon\|_{2,\infty}).$$
(92)

This is from the boundedness and Lipschitz continuity of $\mathcal{T}_y(T_\theta(z), f)$ w.r.t. to f, the boundedness and Lipschitz continuity of $u_z(\theta, f)$ w.r.t. ∇f , and the fact that $\mathcal{T}_y(T_\theta(z), f)$ is positive and bounded away from zero.

For the forth term, following the similar argument as the third term and using the boundedness of $\nabla_{\theta}^2 T_{\theta}(z)$, we have that

$$\|B_4(f_\theta) - B_4(f_\theta^{\epsilon})\|_{op} = \mathcal{O}(\|f_\theta - f_\theta^{\epsilon}\|_{\infty} + \|\nabla f_\theta - \nabla f_\theta^{\epsilon}\|_{2,\infty}).$$
(93)

For the fifth term, following the similar argument as the third term and using the boundedness of $\nabla_{\theta} T_{\theta}(z)$ and $\nabla_{11} c(T_{\theta}(z), y)$, we have that

$$\|B_5(f_\theta) - B_5(f_\theta^\epsilon)\|_{op} = \mathcal{O}(\|f_\theta - f_\theta^\epsilon\|_{\infty}).$$
(94)

For the sixth term, following the similar argument as the third term and using the boundedness of $\nabla_{\theta} T_{\theta}(z)$, we have that

$$\|B_6(f_\theta) - B_6(f_\theta^\epsilon)\|_{op} = \mathcal{O}(\|f_\theta - f_\theta^\epsilon\|_\infty + \|\nabla^2 f_\theta - \nabla^2 f_\theta^\epsilon\|_{op,\infty}).$$
(95)

For the last term, following the similar argument as the third term and using the boundedness of $\nabla_{\theta} T_{\theta}(z)$, we have that

$$\|B_7(f_\theta) - B_7(f_\theta^\epsilon)\|_{op} = \mathcal{O}(\|f_\theta - f_\theta^\epsilon\|_\infty + \|\nabla f_\theta - \nabla f_\theta^\epsilon\|_{2,\infty}).$$
(96)

Combing the above results, we obtain (89).

²Recall that T(U, W) is the family of bounded linear operators from U to W.

C eSIM appendix

C.1 Proof of Theorem 6.1

In this section, we use f_{θ}^{μ} to denote the Sinkhorn potential to $OT_{\gamma}(T_{\theta \sharp}\mu,\beta)$. This allows us to emphasize the continuity of its Fréchet derivative w.r.t. the underlying measure μ . Similarly, we write $\mathcal{B}_{\mu}(f,\theta)$ and $\mathcal{E}_{\mu}(f,\theta)$ instead of $\mathcal{B}(f,\theta)$ and $\mathcal{E}(f,\theta)$, which are used to characterize the fixed point property of the Sinkhorn potential.

To prove Theorem 6.1, we need the following lemmas.

Lemma C.1. The Sinkhorn potential f^{μ}_{θ} is Lipschitz continuous with respect to μ :

$$\|f^{\mu}_{\theta} - f^{\mu}_{\theta}\|_{\infty} = \mathcal{O}(d_{bl}(\mu, \bar{\mu})).$$
(97)

Lemma C.2. The gradient of the Sinkhorn potential f^{μ}_{θ} is Lipschitz continuous with respect to μ :

$$\nabla f^{\mu}_{\theta} - \nabla f^{\bar{\mu}}_{\theta} \|_{2,\infty} = \mathcal{O}(d_{bl}(\mu, \bar{\mu})).$$
(98)

Lemma C.3. The Hessian of the Sinkhorn potential f^{μ}_{θ} is Lipschitz continuous with respect to μ :

$$\nabla^2 f^{\mu}_{\theta} - \nabla^2 f^{\mu}_{\theta} \|_{op,\infty} = \mathcal{O}(d_{bl}(\mu,\bar{\mu})).$$
⁽⁹⁹⁾

Lemma C.4. The Fréchet derivative of the Sinkhorn potential f^{μ}_{θ} w.r.t. the parameter θ , i.e. Df^{μ}_{θ} , is Lipschitz continuous with respect to μ :

$$\|Df^{\mu}_{\theta} - Df^{\bar{\mu}}_{\theta}\|_{op} = \mathcal{O}(d_{bl}(\mu, \bar{\mu})).$$

$$(100)$$

Once we have these lemmas, we can prove 6.1 in the same way as the proof of 5.1 in Appendix B.4.

C.2 Proof of Lemma C.1

Note that from the definition of the bounded Lipschitz distance, we have

$$d_{bl}(\alpha,\bar{\alpha}) = \sup_{\|\xi\|_{bl} \leq 1} |\langle \xi, \alpha \rangle - \langle \xi, \bar{\alpha} \rangle| = \sup_{\|\xi\|_{bl} \leq 1} |\langle \xi \circ T_{\theta}, \mu \rangle - \langle \xi \circ T_{\theta}, \bar{\mu} \rangle|$$

$$\leq \sup_{\|\xi\|_{bl} \leq 1} \|\xi \circ T_{\theta}\|_{bl} \cdot d_{bl}(\mu,\bar{\mu}) \leq G_T \cdot d_{bl}(\mu,\bar{\mu}),$$
(101)

where we use $\|\xi \circ T_{\theta}\|_{lip} \leq G_T$ from Assumption B.5.

We have Lemma C.1 by combining the above results with the following lemma.

Lemma C.5. Under Assumption B.1 and Assumption B.2, the Sinkhorn potential is Lipschitz continuous with respect to the bounded Lipschitz metric: Given measures α , α' and β , we have

$$\|f_{\alpha,\beta} - f_{\alpha',\beta}\|_{\infty} \leqslant G_{bl}d_{bl}(\alpha',\alpha) \quad \text{and} \quad \|g_{\alpha,\beta} - g_{\alpha',\beta'}\|_{\infty} \leqslant G_{bl}d_{bl}(\alpha',\alpha)$$

where $G_{bl} = 2\gamma \exp(2M_c/\gamma)G'_{bl}/(1-\lambda^2)$ with $G'_{bl} = \max\{\exp(3M_c/\gamma), 2G_c \exp(3M_c/\gamma)/\gamma\}$ and $\lambda = \frac{\exp(M_c/\gamma)-1}{\exp(M_c/\gamma)+1}$.

Proof. Let (f,g) and (f',g') be the Sinkhorn potentials to $\operatorname{OT}_{\gamma}(\alpha,\beta)$ and $\operatorname{OT}_{\gamma}(\alpha',\beta)$ respectively. Denote $u := \exp(f/\gamma)$, $v := \exp(g/\gamma)$ and $u' := \exp(f'/\gamma)$, $v' := \exp(g'/\gamma)$. From Lemma C.7, u is bounded in terms of the L^{∞} norm:

$$\|u\|_{\infty} = \max_{x \in \mathcal{X}} |u(x)| = \max_{x \in \mathcal{X}} \exp(f/\gamma) \leq \exp(2M_c/\gamma),$$

which also holds for v, u', v'. Additionally, from Lemma C.8, ∇u exists and $\|\nabla u\|$ is bounded:

$$\max_{x} \|\nabla u(x)\| = \max_{x} \frac{1}{\gamma} |u(x)| \|\nabla f(x)\| \le \frac{1}{\gamma} \|u(x)\|_{\infty} \max_{x} \|\nabla f(x)\| \le \frac{G_c \exp(2M_c/\gamma)}{\gamma}$$

Define the mapping $A_{\alpha}\mu := 1/(L_{\alpha}\mu)$ with

$$L_{\alpha}\mu = \int_{\mathcal{X}} l(\cdot, y)\mu(y)\mathbf{d}\alpha(y),$$

where $l(x, y) := \exp(-c(x, y)/\gamma)$. From Assumption B.1, we have $||l||_{\infty} \leq \exp(M_c/\gamma)$ and from Assumption B.2 we have $||\nabla_x l(x, y)|| \leq \exp(M_c/\gamma)\frac{G_c}{\gamma}$. From the optimality condition of f and g, we have $v = A_{\alpha}u$ and $u = A_{\beta}v$. Similarly, $v' = A_{\alpha'}u'$ and $u' = A_{\beta}v'$. Recall the definition of the Hilbert metric in (60). Note that $d_H(\mu, \nu) = d_H(1/\mu, 1/\nu)$ if $\mu(x) > 0$ and $\nu(x) > 0$ for all $x \in \mathcal{X}$ and hence $d_H(L_{\alpha}\mu, L_{\alpha}\nu) = d_H(A_{\alpha}\mu, A_{\alpha}\nu)$. We recall the result in (61) using the above notations.

Lemma C.6 (Birkhoff-Hopf Theorem Lemmens and Nussbaum [2012], see Lemma B.4 in Luise et al. [2019]). Let $\lambda = \frac{\exp(M_c/\gamma)-1}{\exp(M_c/\gamma)+1}$ and $\alpha \in \mathcal{M}_1^+(\mathcal{X})$. Then for every $u, v \in \mathcal{C}(\mathcal{X})$, such that u(x) > 0, v(x) > 0 for all $x \in \mathcal{X}$, we have

$$d_H(L_{\alpha}u, L_{\alpha}v) \leq \lambda d_H(u, v).$$

Note that

$$\|\log\mu - \log\nu\|_{\infty} \leq d_H(\mu,\nu) = \|\log\mu - \log\nu\|_{\infty} + \|\log\nu - \log\mu\|_{\infty} \leq 2\|\log\mu - \log\nu\|_{\infty}$$

In the following, we derive upper bound for $d_H(\mu, \nu)$ and use such bound to analyze the Lipschitz continuity of the Sinkhorn potentials f and g.

Construct $\tilde{v}:=A_{\alpha}u'$. Using the triangle inequality (which holds since $v(x), v'(x), \tilde{v}(x) > 0$ for all $x \in \mathcal{X}$), we have

$$d_H(v,v') \leqslant d_H(v,\tilde{v}) + d_H(\tilde{v},v') \leqslant \lambda d_H(u,u') + d_H(\tilde{v},v'),$$

where the second inequality is due to Lemma C.6. Note that $u' = A_{\beta}v'$. Apply Lemma C.6 again to obtain

$$d_H(u, u') \le \lambda d_H(v, v').$$

Together, we obtain

$$d_H(v,v') \leq \lambda^2 d_H(v,v') + d_H(\tilde{v},v') + \lambda d_H(\tilde{u},u') \leq \lambda^2 d_H(v,v') + d_H(\tilde{v},v')$$

which leads to

$$d_H(v,v') \leqslant \frac{1}{1-\lambda^2} [d_H(\tilde{v},v')].$$

To bound $d_H(\tilde{v}, v')$, observe the following:

$$d_{H}(v',\tilde{v}) = d_{H}(L_{\alpha'}u',L_{\alpha}u') \leq 2 \|\log L_{\alpha'}u' - \log L_{\alpha}u'\|_{\infty}$$

= $2 \max_{x \in \mathcal{X}} |\nabla \log(a_{x})([L_{\alpha'}u'](x) - [L_{\alpha}u'](x))| = 2 \max_{x \in \mathcal{X}} \frac{1}{a_{x}} |[L_{\alpha'}u'](x) - [L_{\alpha}u'](x)|$
 $\leq 2 \max\{\|1/L_{\alpha'}u'\|_{\infty}, \|1/L_{\alpha}u'\|_{\infty}\}\|L_{\alpha'}u' - L_{\alpha}u'\|_{\infty},$ (102)

where $a_x \in [[L_{\alpha'}u'](x), [L_{\alpha}u'](x)]]$ in the second line is from the mean value theorem. Further, in the inequality we use $\max\{\|1/L_{\alpha}u'\|_{\infty}, \|1/L_{\alpha}u'\|_{\infty}\} = \max\{\|A_{\alpha'}u'\|_{\infty}, \|A_{\alpha}u'\|_{\infty}\} \leq \exp(2M_c/\gamma)$. Consequently, all we need to bound is the last term $\|L_{\alpha'}u' - L_{\alpha}u'\|_{\infty}$.

We first note that $\forall x \in \mathcal{X}, ||l(x, \cdot)u'(\cdot)||_{bl} < \infty$: In terms of $||\cdot||_{\infty}$

$$||l(x,\cdot)u'(\cdot)||_{\infty} \leq ||l(x,\cdot)||_{\infty} ||u'||_{\infty} \leq \exp(3M_c/\gamma) < \infty.$$

In terms of $\|\cdot\|_{lip}$, we bound

$$\begin{aligned} \|l(x,\cdot)u'(\cdot)\|_{lip} &\leqslant \|l(x,\cdot)\|_{\infty} \|u'\|_{lip} + \|l(x,\cdot)\|_{lip} \|u'\|_{\infty} \\ &\leqslant \exp(M_c/\gamma) \frac{G_c \exp(2M_c/\gamma)}{\gamma} + \exp(M_c/\gamma) \frac{G_c}{\gamma} \exp(2M_c/\gamma) = \frac{2G_c \exp(3M_c/\gamma)}{\gamma} < \infty \end{aligned}$$

Together we have $||l(x,y)u'(y)||_{bl} \leq \max\{\exp(3M_c/\gamma), \frac{2G_c \exp(3M_c/\gamma)}{\gamma}\}$. From the definition of the operator L_{α} , we have

$$\|L_{\alpha'}u' - L_{\alpha}u'\|_{\infty} = \max_{x} |\int_{\mathcal{X}} l(x,y)u'(y)\mathbf{d}\alpha'(y) - \int_{\mathcal{X}} l(x,y)u'(y)\mathbf{d}\alpha(y)| \leq \|l(x,y)u'(y)\|_{bl}d_{bl}(\alpha',\alpha).$$

All together we derive

$$d_{H}(v',v) \leqslant \frac{2\exp(2M_{c}/\gamma) \|l(x,y)u'(y)\|_{bl}}{1-\lambda^{2}} \cdot d_{bl}(\alpha',\alpha) \quad (\lambda = \frac{\exp(M_{c}/\gamma) - 1}{\exp(M_{c}/\gamma) + 1}).$$

Further, since $d_H(v', v) \ge \|\log v' - \log v\|_{\infty} = \frac{1}{\gamma} \|f' - f\|_{\infty}$, we have the result:

$$\|f' - f\|_{\infty} \leq \frac{2\gamma \exp(2M_c/\gamma) \|l(x,y)u'(y)\|_{bl}}{1 - \lambda^2} \cdot d_{bl}(\alpha',\alpha).$$

Similar argument can be made for $||g' - g||_{\infty}$.

Lemma C.7 (Boundedness of the Sinkhorn Potentials). Let (f, g) be the Sinkhorn potentials of problem (6) and assume that there exists $x_o \in \mathcal{X}$ such that $f(x_o) = 0$ (otherwise shift the pair by $f(x_o)$). Then, under Assumption B.1, $||f||_{\infty} \leq 2M_c$ and $||g||_{\infty} \leq 2M_c$.

Next, we analyze the Lipschitz continuity of the Sinkhorn potential $f_{\alpha,\beta}(x)$ with respect to the input x.

Assumption B.2 implies that $\nabla_x c(x, y)$ exists and for all $x, y \in \mathcal{X}, \|\nabla_x c(x, y)\| \leq G_c$. It further ensures the Lipschitz-continuity of the Sinkhorn potential.

Lemma C.8 (Proposition 12 of Feydy et al. [2019]). Under Assumption B.2, for a fixed pair of measures (α, β) , the corresponding Sinkhorn potential $f : \mathcal{X} \to \mathbb{R}$ is G_c -Lipschitz continuous, i.e. for $x_1, x_2 \in \mathcal{X}$

$$|f_{\alpha,\beta}(x_1) - f_{\alpha,\beta}(x_2)| \le G_c ||x_1 - x_2||.$$
(103)

Further, the gradient $\nabla f_{\alpha,\beta}$ exists at every point $x \in \mathcal{X}$, and $\|\nabla f_{\alpha,\beta}(x)\| \leq G_c, \forall x \in \mathcal{X}$.

Lemma C.9. Under Assumption B.3, for a fixed pair of measures (α, β) , the gradient of the corresponding Sinkhorn potential $f : \mathcal{X} \to \mathbb{R}$ is Lipschitz continuous,

$$\|\nabla f(x_1) - \nabla f(x_2)\| \le L_f \|x_1 - x_2\|, \tag{104}$$

where $L_f := \frac{4G_c^2}{\gamma} + L_c$.

C.3 Proof of Lemma C.2

We have Lemma C.2 by combining (101) with the following lemma.

Lemma C.10 (Lemma C.2 restated). Under Assumption B.1 and Assumption B.2, the gradient of the Sinkhorn potential is Lipschitz continuous with respect to the bounded Lipschitz metric: Given measures α , α' and β , we have

$$\|\nabla f_{\alpha,\beta} - \nabla f_{\alpha',\beta}\|_{\infty} = \mathcal{O}(d_{bl}(\alpha',\alpha))$$

Proof. From the optimality condition of the Sinkhorn potentials, one have that

$$\int_{\mathcal{X}} h_{\alpha,\beta}(x,y) \mathbf{d}\beta(y) = 1, \text{ with } h_{\alpha,\beta}(x,y) := \exp\left(\frac{1}{\gamma} \left(f_{\alpha,\beta}(x) + g_{\alpha,\beta}(y) - c(x,y)\right)\right).$$
(105)

Taking gradient w.r.t. x on both sides of the above equation, the expression of $\nabla f_{\alpha,\beta}$ writes

$$\nabla f_{\alpha,\beta}(x) = \frac{\int_{\mathcal{X}} h_{\alpha,\beta}(x,y) \nabla_x c(x,y) \mathbf{d}\beta(y)}{\int_{\mathcal{X}} h_{\alpha,\beta}(x,y) \mathbf{d}\beta(y)} = \int_{\mathcal{X}} h_{\alpha,\beta}(x,y) \nabla_x c(x,y) \mathbf{d}\beta(y).$$
(106)

We have that $\forall x, y, h_{\alpha,\beta}(x)$ is Lipschitz continuous w.r.t. α , which is due to the boundedness of $f_{\alpha,\beta}(x), g_{\alpha,\beta}(y)$ and the ground cost c, and Lemma C.1. Further, since $\|\nabla_x c(x,y)\|$ is bounded from Assumption B.2 we have the Lipschitz continuity of $\nabla f_{\alpha,\beta}$ w.r.t. α , i.e.

$$\|\nabla f_{\alpha,\beta}(x) - \nabla f_{\alpha',\beta}(x)\| = \mathcal{O}(d_{bl}(\alpha',\alpha)).$$

C.4 Proof of Lemma C.3

We have Lemma C.3 by combining (101) with the following lemma.

Lemma C.11 (Lemma C.3 restated). Under Assumptions B.1-B.3, the Hessian of the Sinkhorn potential is Lipschitz continuous with respect to the bounded Lipschitz metric: Given measures α , α' and β , we have

$$\|\nabla^2 f_{\alpha,\beta} - \nabla^2 f_{\alpha',\beta}\|_{op,\infty} = \mathcal{O}(d_{bl}(\alpha',\alpha))$$

Proof. Taking gradient w.r.t. x on both sides of (106), the expression of $\nabla^2 f_{\alpha,\beta}$ writes

$$\nabla^2 f_{\alpha,\beta}(x) = \int_{\mathcal{X}} \frac{1}{\gamma} h_{\alpha,\beta}(x,y) (\nabla f_{\alpha,\beta}(x) - \nabla_x c(x,y)) [\nabla_x c(x,y)]^\top + h_{\alpha,\beta}(x,y) \nabla^2_{xx} c(x,y) \mathbf{d}\beta(y).$$

From the boundedness of $h_{\alpha,\beta}$, $\nabla f_{\alpha,\beta}$ and $\nabla_x c$, and the Lipschitz continuity of $h_{\alpha,\beta}$ and $\nabla f_{\alpha,\beta}$ w.r.t. α , we have that the first integrand of $\nabla^2 f_{\alpha,\beta}$ is Lipschitz continuous w.r.t. α . Further, combining the boundedness of $\|\nabla^2_{xx}c(x,y)\|$ from Assumption B.3 and the Lipschitz continuity of $h_{\alpha,\beta}$ w.r.t. α , we have the Lipschitz continuity of $\nabla^2 f_{\alpha,\beta}(x)$, i.e.

$$\|\nabla^2 f_{\alpha,\beta}(x) - \nabla^2 f_{\alpha',\beta}(x)\| = \mathcal{O}(d_{bl}(\alpha',\alpha)).$$

C.5 Proof of Lemma C.4

The optimality of the Sinkhorn potential f^{μ}_{θ} can be restated as

$$f^{\mu}_{\theta} = \mathcal{B}_{\mu}(f^{\mu}_{\theta}, \theta), \tag{107}$$

where we recall the definition of \mathcal{B}_{μ} in (18)

$$\mathcal{B}_{\mu}(f,\theta) = \mathcal{A}\big(\mathcal{A}(f,T_{\theta\sharp}\mu),\beta_{\mu}\big).$$
(108)

Note that it is possible that β_{μ} depends on μ , which is the case in $OT_{\gamma}(\alpha_{\theta}, \alpha_{\theta^{t}})$ as $\beta_{\mu} = \alpha_{\theta^{t}} = T_{\theta^{t}\sharp}\mu$. Under Assumption B.1, let $\lambda = \frac{e^{M_{c}/\gamma} - 1}{e^{M_{c}/\gamma} + 1}$. By repeating the above fixed point iteration (107) $l = \lfloor \log_{\lambda} \frac{1}{3} \rfloor/2$ times, we have that

$$f^{\mu}_{\theta} = \mathcal{E}_{\mu}(f^{\mu}_{\theta}, \theta), \tag{109}$$

where $\mathcal{E}_{\mu}(f,\theta) = \mathcal{B}_{\mu}^{l}(f,\theta) = \mathcal{B}_{\mu}(\cdots \mathcal{B}_{\mu}(f,\theta)\cdots,\theta)$ is the *l* times composition of \mathcal{B}_{μ} in its first variable. We have from (68)

$$||D^1 \mathcal{E}_{\mu}(f,\theta)||_{op} \leqslant \frac{2}{3},\tag{110}$$

where we recall for a (linear) operator $C : C(\mathcal{X}) \to C(\mathcal{X}), \|C\|_{op} := \max_{f \in C(\mathcal{X})} \frac{\|Cf\|_{\infty}}{\|f\|_{\infty}}.$

Let $h \in \mathbb{R}^d$ be any direction. Taking Fréchet derivative w.r.t. θ on both sides of (109), we derive

$$Df^{\mu}_{\theta}[h] = D_1 \mathcal{E}_{\mu}(f^{\mu}_{\theta}, \theta) \left[Df^{\mu}_{\theta}[h] \right] + D_2 \mathcal{E}_{\mu}(f^{\mu}_{\theta}, \theta)[h].$$
(111)

Using the triangle inequality, we bound

$$\begin{split} \|Df_{\theta}^{\mu}[h] - Df_{\theta}^{\mu}[h]\|_{\infty} \\ &\leq \|D_{1}\mathcal{E}_{\mu}(f_{\theta}^{\mu},\theta)[Df_{\theta}^{\mu}[h]] - D_{1}\mathcal{E}_{\bar{\mu}}(f_{\theta,\bar{\mu}},\theta)[Df_{\theta}^{\bar{\mu}}[h]]\|_{\infty} \\ &+ \|D_{2}\mathcal{E}_{\mu}(f_{\theta}^{\mu},\theta)[h] - D_{2}\mathcal{E}_{\bar{\mu}}(f_{\theta}^{\bar{\mu}},\theta)[h]\|_{\infty} \\ &\leq \|D_{1}\mathcal{E}_{\mu}(f_{\theta}^{\mu},\theta)[Df_{\theta}^{\mu}[h]] - D_{1}\mathcal{E}_{\mu}(f_{\theta}^{\mu},\theta)[Df_{\theta}^{\bar{\mu}}[h]]\|_{\infty} \qquad (112) \\ &+ \|D_{1}\mathcal{E}_{\mu}(f_{\theta,\mu},\theta)[Df_{\theta}^{\bar{\mu}}[h]] - D_{1}\mathcal{E}_{\mu}(f_{\theta,\bar{\mu}},\theta)[Df_{\theta}^{\bar{\mu}}[h]]\|_{\infty} \qquad (2) \\ &+ \|D_{1}\mathcal{E}_{\mu}(f_{\theta,\bar{\mu}},\theta)[Df_{\theta}^{\bar{\mu}}[h]] - D_{1}\mathcal{E}_{\bar{\mu}}(f_{\theta,\bar{\mu}},\theta)[Df_{\theta}^{\bar{\mu}}[h]]\|_{\infty} \qquad (3) \\ &+ \|D_{2}\mathcal{E}_{\mu}(f_{\theta}^{\mu},\theta)[h] - D_{2}\mathcal{E}_{\bar{\mu}}(f_{\theta}^{\bar{\mu}},\theta)[h]\|_{\infty}. \qquad (4) \end{split}$$

The following subsections analyze (1) to (4) individually. In summary, we have

and (2), (3), (4) are all of order $\mathcal{O}(d_{bl}(\mu, \bar{\mu}) \cdot \|h\|)$. Therefore we conclude

$$\frac{1}{3} \|Df^{\mu}_{\theta}[h] - Df^{\bar{\mu}}_{\theta}[h]\|_{\infty} = \mathcal{O}(d_{bl}(\mu,\bar{\mu}) \cdot \|h\|) \Rightarrow \|Df^{\mu}_{\theta} - Df^{\bar{\mu}}_{\theta}\|_{op} = \mathcal{O}(d_{bl}(\mu,\bar{\mu})).$$
(114)

C.5.1 Bounding ①

From the linearity of $D_1 \mathcal{E}_{\mu}(f^{\mu}_{\theta}, \theta)$ and (110), we bound

$$() = \|D_1 \mathcal{E}_{\mu}(f^{\mu}_{\theta}, \theta) [Df^{\mu}_{\theta}[h] - Df^{\bar{\mu}}_{\theta}[h]]\|_{\infty}$$

$$\leq \|D_1 \mathcal{E}_{\mu}(f^{\mu}_{\theta}, \theta)\|_{op} \|Df^{\mu}_{\theta}[h] - Df^{\bar{\mu}}_{\theta}[h]\|_{\infty} \leq \frac{2}{3} \|Df^{\mu}_{\theta}[h] - Df^{\bar{\mu}}_{\theta}[h]\|_{\infty}.$$

C.5.2 Bounding ⁽²⁾

From Lemma B.8, we know that $D_1 \mathcal{B}_{\mu}(f, \theta)$ is Lipschitz continuous w.r.t. its first variable:

$$\|D_1 \mathcal{B}_{\mu}(f, \theta) - D_1 \mathcal{B}_{\mu}(f', \theta)\|_{op} = \mathcal{O}(\|f - f'\|_{\infty}).$$
(115)

Recall that $\mathcal{E}_{\mu}(f,\theta) = \mathcal{B}_{\mu}^{l}(f,\theta)$. Using the chain rule of the Fréchet derivative, we have

$$D_1 \mathcal{E}_{\mu}(f,\theta) = D_1 \mathcal{B}^l_{\mu}(f,\theta) = D_1 \mathcal{B}_{\mu} \left(\mathcal{B}^{l-1}_{\mu}(f,\theta), \theta \right) \circ D_1 \mathcal{B}^{l-1}_{\mu}(f,\theta).$$
(116)

Consequently, we can bound (2) in a recursive way: for any two functions $f, f' \in C(\mathcal{X})$

$$\begin{split} \|D_{1}\mathcal{B}_{\mu}^{l}(f,\theta) - D_{1}\mathcal{B}_{\mu}^{l}(f',\theta)\|_{op} \\ &= \|D_{1}\mathcal{B}_{\mu}\big(\mathcal{B}_{\mu}^{l-1}(f,\theta),\theta\big) \circ D_{1}\mathcal{B}_{\mu}^{l-1}(f,\theta) - D_{1}\mathcal{B}_{\mu}\big(\mathcal{B}_{\mu}^{l-1}(f',\theta),\theta\big) \circ D_{1}\mathcal{B}_{\mu}^{l-1}(f',\theta)\|_{op} \\ &\leq \|D_{1}\mathcal{B}_{\mu}\big(\mathcal{B}_{\mu}^{l-1}(f,\theta),\theta\big) \circ \big(D_{1}\mathcal{B}_{\mu}^{l-1}(f,\theta) - D_{1}\mathcal{B}_{\mu}^{l-1}(f',\theta)\big)\|_{op} \\ &+ \|\Big(D_{1}\mathcal{B}_{\mu}\big(\mathcal{B}_{\mu}^{l-1}(f,\theta),\theta\big) - D_{1}\mathcal{B}_{\mu}\big(\mathcal{B}_{\mu}^{l-1}(f',\theta),\theta\big)\Big) \circ D_{1}\mathcal{B}_{\mu}^{l-1}(f',\theta)\|_{op} \\ &\leq \|D_{1}\mathcal{B}_{\mu}\big(\mathcal{B}_{\mu}^{l-1}(f,\theta),\theta\big)\|_{op}\|D_{1}\mathcal{B}_{\mu}^{l-1}(f,\theta) - D_{1}\mathcal{B}_{\mu}^{l-1}(f',\theta)\|_{\infty} \\ &+ \mathcal{O}(\|\mathcal{B}_{\mu}^{l-1}(f,\theta) - \mathcal{B}_{\mu}^{l-1}(f',\theta)\|_{\infty} \cdot \|D_{1}\mathcal{B}_{\mu}^{l-1}(f',\theta)\|_{op}) \\ &= \mathcal{O}(\|f - f'\|_{\infty}) + \|D_{1}\mathcal{B}_{\mu}^{l-1}(f,\theta) - D_{1}\mathcal{B}_{\mu}^{l-1}(f',\theta)\|_{\infty}, \end{split}$$

where in the first inequality we use the triangle inequality, in the second inequality, we use the definition of $\|\cdot\|_{op}$ and (115), and in the last equality we use (115) and the fact that \mathcal{B}^k is Lipschitz continuous with respect its first argument for any finite k (see Lemma B.9). Besides, since f^{μ}_{θ} is continuous with respect to μ (see Lemma C.1), we have

$$\|D_1 \mathcal{B}^l(f^{\mu}_{\theta}, \theta) - D_1 \mathcal{B}^l(f^{\mu}_{\theta}, \theta)\|_{op} = \mathcal{O}(d_{bl}(\mu, \bar{\mu})).$$
(117)

We then show that $\|Df_{\theta}^{\bar{\mu}}[h]\|_{\infty} = \mathcal{O}(\|h\|_{\infty})$: Using (111), we have that

$$\|Df_{\theta}^{\bar{\mu}}[h]\|_{\infty} \leq \frac{2}{3} \|Df_{\theta}^{\bar{\mu}}[h]\|_{\infty} + \|D_{2}\mathcal{E}_{\mu}(f_{\theta}^{\mu},\theta)[h]\|_{\infty} \Rightarrow \|Df_{\theta}^{\bar{\mu}}[h]\|_{\infty} \leq 3 \|D_{2}\mathcal{E}_{\mu}(f_{\theta}^{\mu},\theta)\|_{op}\|[h]\|_{\infty}.$$

Lemma B.7 shows that $||D_2 \mathcal{E}_{\mu}(f^{\mu}_{\theta}, \theta)||_{op}$ is bounded and therefore we have

$$\|Df^{\mu}_{\theta}[h]\|_{\infty} = \mathcal{O}(\|h\|_{\infty}). \tag{118}$$

Combining the above results, we obtain

$$(2 \leq \|D_1 \mathcal{B}^l(f^{\mu}_{\theta}, \theta) - D_1 \mathcal{B}^l(f^{\bar{\mu}}_{\theta}, \theta)\|_{op} \|Df^{\bar{\mu}}_{\theta}[h]\|_{\infty} = \mathcal{O}(d_{bl}(\mu, \bar{\mu}) \cdot \|h\|_{\infty}).$$

C.5.3 Bounding ③

Denote $\omega_y(x) = \exp(-\frac{c(x,y)}{\gamma}) \exp(\bar{f}(x)/\gamma)$. Assume that $\|\bar{f}\|_{\infty} \leq M_c$ and $\|\nabla \bar{f}\|_{2,\infty} \leq G_f$. Then we have for any $y \in \mathcal{X}$,

$$\|\omega_y\|_{\infty} \leq \exp(M_c/\gamma), \quad \|\nabla\omega_y\|_{2,\infty} \leq \exp(M_c/\gamma)(G_c + G_f)/\gamma.$$
(119)

Therefore, $\|\omega_y\|_{bl} = \max\{\exp(M_c/\gamma), \exp(M_c/\gamma)(G_c + G_f)/\gamma\}$ is bounded (recall the definition of bounded Lipschitz norm in Theorem 6.1). Besides, for any $y \in \mathcal{X}$, $\omega_y(x)$ is positive and bounded away from zero

$$\omega_y(x) \ge \exp(-2M_c/\gamma). \tag{120}$$

For a fixed measure κ and $g \in \mathcal{C}(\mathcal{X})$, we compute that

$$D_1 \mathcal{A}(\bar{f},\kappa)[g] = \frac{\int_{\mathcal{X}} \omega_y(x) g(x) \mathbf{d}\kappa(x)}{\int_{\mathcal{X}} \omega_y(x) \mathbf{d}\kappa(x)}.$$
(121)

This expression allows us to bound for two measures κ and κ'

$$\begin{split} &\| \left(D_1 \mathcal{A}(\bar{f},\kappa) - D_1 \mathcal{A}(\bar{f},\kappa') \right) [g] \|_{\infty} = \| \frac{\int_{\mathcal{X}} \omega_y(x) g(x) \mathbf{d} \kappa(x)}{\int_{\mathcal{X}} \omega_y(x) \mathbf{d} \kappa(x)} - \frac{\int_{\mathcal{X}} \omega_y(x) g(x) \mathbf{d} \kappa'(x)}{\int_{\mathcal{X}} \omega_y(x) \mathbf{d} \kappa'(x)} \|_{\infty} \\ &\leqslant \| \frac{\int_{\mathcal{X}} \omega_y(x) g(x) \mathbf{d} \kappa(x)}{\int_{\mathcal{X}} \omega_y(x) \mathbf{d} \kappa(x)} - \frac{\int_{\mathcal{X}} \omega_y(x) g(x) \mathbf{d} \kappa(x)}{\int_{\mathcal{X}} \omega_y(x) \mathbf{d} \kappa'(x)} \|_{\infty} + \| \frac{\int_{\mathcal{X}} \omega_y(x) g(x) \mathbf{d} \kappa(x)}{\int_{\mathcal{X}} \omega_y(x) \mathbf{d} \kappa'(x)} - \frac{\int_{\mathcal{X}} \omega_y(x) g(x) \mathbf{d} \kappa'(x)}{\int_{\mathcal{X}} \omega_y(x) \mathbf{d} \kappa'(x)} \|_{\infty}. \end{split}$$

We now bound these two terms individually. For the first term, we have

$$\begin{split} &\|\frac{\int_{\mathcal{X}}\omega_{y}(x)g(x)\mathbf{d}\kappa(x)}{\int_{\mathcal{X}}\omega_{y}(x)\mathbf{d}\kappa(x)} - \frac{\int_{\mathcal{X}}\omega_{y}(x)g(x)\mathbf{d}\kappa(x)}{\int_{\mathcal{X}}\omega_{y}(x)\mathbf{d}\kappa'(x)}\|_{\infty} \\ &\leqslant \|\int_{\mathcal{X}}\omega_{y}(x)g(x)\mathbf{d}\kappa(x)\|_{\infty}\|\frac{\int_{\mathcal{X}}\omega_{y}(x)\left[\mathbf{d}\kappa(x) - \mathbf{d}\kappa'(x)\right]}{\int_{\mathcal{X}}\omega_{y}(x)\mathbf{d}\kappa(x)\int_{\mathcal{X}}\omega_{y}(x)\mathbf{d}\kappa'(x)}\|_{\infty} \\ &\leqslant \|\omega_{y}\|_{\infty} \cdot \|g\|_{\infty} \cdot \|\omega_{y}(x)\|_{bl} \cdot d_{bl}(\kappa,\kappa') \cdot \exp(4M_{c}/\gamma) = \mathcal{O}(\|g\|_{\infty} \cdot d_{bl}(\kappa,\kappa')), \end{split}$$

where we use (119) and (120) in the last equality. For the second term, we bound

$$\begin{split} \| \frac{\int_{\mathcal{X}} \omega_y(x) g(x) \mathbf{d} \kappa(x)}{\int_{\mathcal{X}} \omega_y(x) \mathbf{d} \kappa(x)} - \frac{\int_{\mathcal{X}} \omega_y(x) g(x) \mathbf{d} \kappa'(x)}{\int_{\mathcal{X}} \omega_y(x) \mathbf{d} \kappa(x)} \|_{\infty} \leqslant \| \frac{\int_{\mathcal{X}} \omega_y(x) g(x) [\mathbf{d} \kappa(x) - \mathbf{d} \kappa'(x)]}{\int_{\mathcal{X}} \omega_y(x) \mathbf{d} \kappa(x)} \|_{\infty} \\ \leqslant \exp(M_c/\gamma) \cdot \| \omega_y(x) \|_{bl} \cdot \|g\|_{bl} \cdot d_{bl}(\kappa, \kappa') = \mathcal{O}(\|g\|_{bl} \cdot d_{bl}(\kappa, \kappa')). \end{split}$$

Combining the above inequalities, we have

$$\| \left(D_1 \mathcal{A}(\bar{f},\kappa) - D_1 \mathcal{A}(\bar{f},\kappa') \right) [g] \|_{\infty} = \mathcal{O}(\|g\|_{bl} \cdot d_{bl}(\kappa,\kappa')).$$
(122)

Denote $\alpha = T_{\theta \sharp} \mu$ and $\bar{\alpha} = T_{\theta \sharp} \bar{\mu}$. From the chain rule of the Fréchet derivative, we compute

$$\begin{split} &\| \left(D_{1}\mathcal{B}_{\mu}(f,\theta) - D_{1}\mathcal{B}_{\bar{\mu}}(f,\theta) \right) [g] \|_{\infty} \\ &= \| \left(D_{1}\mathcal{A} \left(\mathcal{A}(f,\alpha), \beta_{\mu} \right) \circ D_{1}\mathcal{A}(f,\alpha) - D_{1}\mathcal{A} \left(\mathcal{A}(f,\bar{\alpha}), \beta_{\bar{\mu}} \right) \circ D_{1}\mathcal{A}(f,\bar{\alpha}) \right) [g] \|_{\infty} \\ &\leq \| D_{1}\mathcal{A} \left(\mathcal{A}(f,\alpha), \beta_{\mu} \right) \left[\left(D_{1}\mathcal{A}(f,\alpha) - D_{1}\mathcal{A}(f,\bar{\alpha}) \right) [g] \right] \|_{\infty} \\ &+ \| \left(D_{1}\mathcal{A} \left(\mathcal{A}(f,\alpha), \beta_{\mu} \right) - D_{1}\mathcal{A} \left(\mathcal{A}(f,\alpha), \beta_{\bar{\mu}} \right) \right) \left[D_{1}\mathcal{A}(f,\bar{\alpha}) [g] \right] \|_{\infty} \\ &+ \| \left(D_{1}\mathcal{A} \left(\mathcal{A}(f,\alpha), \beta_{\bar{\mu}} \right) - D_{1}\mathcal{A} \left(\mathcal{A}(f,\bar{\alpha}), \beta_{\bar{\mu}} \right) \right) \left[D_{1}\mathcal{A}(f,\bar{\alpha}) [g] \right] \|_{\infty} . \end{split}$$

We now bound these three terms one by one. For the first term, use (110) to derive

$$\begin{split} \|D_1\mathcal{A}\big(\mathcal{A}(f,\alpha),\beta_{\mu}\big)\big[\big(D_1\mathcal{A}(f,\alpha)-D_1\mathcal{A}(f,\bar{\alpha})\big)[g]\big]\|_{\infty} \\ &\leqslant \|D_1\mathcal{A}(f,\alpha)[g]-D_1\mathcal{A}(f,\bar{\alpha})[g]\|_{\infty}=\mathcal{O}(\|g\|_{bl}\cdot d_{bl}(\alpha,\bar{\alpha})), \end{split}$$

where we use $||D_1 \mathcal{A}(\mathcal{A}(f, \alpha), \beta_\mu)||_{op} \leq 1$ (75) and (122) in the second equality. Combining the above result with (101) gives

$$\left\| D_1 \mathcal{A} \big(\mathcal{A}(f,\alpha), \beta_\mu \big) \big[\big(D_1 \mathcal{A}(f,\alpha) - D_1 \mathcal{A}(f,\bar{\alpha}) \big) [g] \big] \right\|_{\infty} = \mathcal{O}(\|g\|_{bl} \cdot d_{bl}(\mu,\bar{\mu})).$$

For the second term, use (122) to derive

$$\begin{split} \left\| \left(D_1 \mathcal{A} \big(\mathcal{A}(f, \alpha), \beta_{\mu} \big) - D_1 \mathcal{A} \big(\mathcal{A}(f, \alpha), \beta_{\bar{\mu}} \big) \right) [D_1 \mathcal{A}(f, \bar{\alpha})[g]] \right\|_{\infty} \\ &= \mathcal{O}(\|D_1 \mathcal{A}(f, \bar{\alpha})[g]\|_{bl} \cdot d_{bl}(\beta_{\mu}, \beta_{\bar{\mu}})). \end{split}$$

We now bound $||D_1\mathcal{A}(f,\bar{\alpha})[g]||_{bl}$. From (75), we have that $||D_1\mathcal{A}(f,\bar{\alpha})[g]||_{\infty} \leq ||g||_{\infty}$. Besides, note that $D_1\mathcal{A}(f,\bar{\alpha})[g]$ is a function mapping from \mathcal{X} to \mathbb{R} and recall the expression of $D_1\mathcal{A}(f,\bar{\alpha})[g]$ in (121). To show that $D_1\mathcal{A}(f,\bar{\alpha})[g](y)$ is Lipschitz continuous w.r.t. y, we use the similar argument as (63): Under Assumption B.1 and assume that $||f||_{\infty} \leq M_c$, the numerator and denominator of (63)

are both Lipschitz continuous w.r.t. y and bounded; the denominator is positive and bounded away from zero. Consequently, we can bound for any $y \in \mathcal{X}$

$$\|\nabla_y D_1 \mathcal{A}(f, \bar{\alpha})[g](y)\| \leq 2\exp(4M_c/\gamma) \|g\|_{\infty} \cdot G_c,$$
(123)

and therefore

$$\left\| \left(D_1 \mathcal{A} \big(\mathcal{A}(f, \alpha), \beta_\mu \big) - D_1 \mathcal{A} \big(\mathcal{A}(f, \alpha), \beta_{\bar{\mu}} \big) \right) \left[D_1 \mathcal{A}(f, \bar{\alpha})[g] \right] \right\|_{\infty} = \mathcal{O}(\|g\|_{\infty} \cdot d_{bl}(\beta_\mu, \beta_{\bar{\mu}})).$$

For the third term, first note that we can use (101) and the mean value theorem to bound

$$\|\mathcal{A}(f,\alpha) - \mathcal{A}(f,\bar{\alpha})\|_{\infty} = \mathcal{O}(\max_{y\in\mathcal{X}} \|\omega_y\|_{bl} \cdot d_{bl}(\alpha,\bar{\alpha})) = \mathcal{O}(d_{bl}(\mu,\bar{\mu})).$$
(124)

Hence, we use Lemma B.11 to derive

$$\left\| \left(D_1 \mathcal{A}(\mathcal{A}(f,\alpha),\beta_{\bar{\mu}}) - D_1 \mathcal{A}(\mathcal{A}(f,\bar{\alpha}),\beta_{\bar{\mu}}) \right) \left[D_1 \mathcal{A}(f,\bar{\alpha})[g] \right] \right\|_{\infty}$$

= $\mathcal{O}(\|\mathcal{A}(f,\alpha) - \mathcal{A}(f,\bar{\alpha})\|_{\infty} \cdot \|D_1 \mathcal{A}(f,\bar{\alpha})[g]\|_{\infty}) = \mathcal{O}(\|g\|_{\infty} \cdot d_{bl}(\mu,\bar{\mu})),$

where we use (124) and the fact that $||D_1\mathcal{A}(f,\bar{\alpha})||_{op}$ is bounded in the last equality. Combing the above three results, we have

$$\| \left(D_1 \mathcal{B}_{\mu}(f,\theta) - D_1 \mathcal{B}_{\bar{\mu}}(f,\theta) \right) [g] \|_{\infty} = \mathcal{O}(\|g\|_{bl} \cdot d_{bl}(\mu,\bar{\mu})).$$
(125)

Recall that $\mathcal{E}_{\mu}(f,\theta) = \mathcal{B}_{\mu}^{l}(f,\theta)$. Using the chain rule of the Fréchet derivative, we have

$$D_1 \mathcal{E}_{\mu}(f,\theta) = D_1 \mathcal{B}^l_{\mu}(f,\theta) = D_1 \mathcal{B}_{\mu} \left(\mathcal{B}^{l-1}_{\mu}(f,\theta), \theta \right) \circ D_1 \mathcal{B}^{l-1}_{\mu}(f,\theta).$$
(126)

Denote $g = Df_{\theta}^{\overline{\mu}}[h]$. We can bound (3) in the following way:

$$\begin{split} (\mathfrak{J} &= \|D_{1}\mathcal{B}_{\mu}\big(\mathcal{B}_{\mu}^{l-1}(f,\theta),\theta\big)\big[D_{1}\mathcal{B}_{\mu}^{l-1}(f,\theta)[g]\big] - D_{1}\mathcal{B}_{\bar{\mu}}\big(\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta),\theta\big)\big[D_{1}\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta)[g]\big]\|_{\infty} \\ &\leq \|D_{1}\mathcal{B}_{\mu}\big(\mathcal{B}_{\mu}^{l-1}(f,\theta),\theta\big)\big[\big(D_{1}\mathcal{B}_{\mu}^{l-1}(f,\theta) - D_{1}\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta)\big)\big[g]\big]\|_{\infty} \\ &+ \|\Big(D_{1}\mathcal{B}_{\mu}\big(\mathcal{B}_{\mu}^{l-1}(f,\theta),\theta\big) - D_{1}\mathcal{B}_{\mu}\big(\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta),\theta\big)\Big)\Big[D_{1}\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta)[g]\big]\|_{\infty} \\ &+ \|\Big(D_{1}\mathcal{B}_{\mu}\big(\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta),\theta\big) - D_{1}\mathcal{B}_{\bar{\mu}}\big(\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta),\theta\big)\Big)\Big[D_{1}\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta)[g]\big]\|_{\infty} \\ &\leq \|D_{1}\mathcal{B}_{\mu}\big(\mathcal{B}_{\mu}^{l-1}(f,\theta),\theta\big)\|_{op}\|\big(D_{1}\mathcal{B}_{\mu}^{l-1}(f,\theta) - D_{1}\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta)\big)[g]\|_{\infty} \qquad \#1 \\ &+ \mathcal{O}(\|\mathcal{B}_{\mu}^{l-1}(f,\theta) - \mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta)\|_{\infty} \cdot \|D_{1}\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta)[g]\|_{\infty}) \qquad \#2 \end{split}$$

$$+ \mathcal{O}(\|D_1 \mathcal{B}^{l-1}_{\bar{\mu}}(f,\theta)[g]\|_{bl} \cdot d_{bl}(\mu,\bar{\mu})), \qquad \#3$$

where in the first inequality we use the triangle inequality, in the second inequality we use the definition of $\|\cdot\|_{op}$, (115) and (125). We now analyze the R.H.S. of the above inequality one by one. For the first term, use $\|D_1\mathcal{B}_{\mu}(\mathcal{B}^{l-1}_{\mu}(f,\theta),\theta)\|_{op} \leq 1$ and then use (125). We have

$$#1 \leq \| \left(D_1 \mathcal{B}_{\mu}(f,\theta) - D_1 \mathcal{B}_{\bar{\mu}}(f,\theta) \right) [g] \|_{\infty} = \mathcal{O}(\|g\|_{bl} \cdot d_{bl}(\mu,\bar{\mu})).$$

For the second term, note that \mathcal{B}^k_{μ} is the composition of the terms $\mathcal{A}(f, \alpha)$ and $\mathcal{A}(f, \beta_{\mu})$. Using a similar argument like (124), for any finite k, we have

$$|\mathcal{B}_{\mu}^{l-1}(f,\theta) - \mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta)||_{\infty} = \mathcal{O}(d_{bl}(\mu,\bar{\mu})).$$

Together with the fact that $||D_1\mathcal{B}(f,\theta)||_{op} \leq 1$, we have

$$#2 = \mathcal{O}(\|g\|_{\infty} \cdot d_{bl}(\mu, \bar{\mu})).$$

Finally, for the third term, note that \mathcal{B}_{μ} is the composition of the terms $\mathcal{A}(f, \alpha)$ and $\mathcal{A}(f, \beta_{\mu})$. Using a similar argument like (123) to bound

$$#3 = \mathcal{O}(\|g\|_{\infty} \cdot d_{bl}(\mu, \bar{\mu})).$$

Combining these three results, we have

$$\mathfrak{T} = \| (D_1 \mathcal{B}^l_{\mu}(f,\theta) - D_1 \mathcal{B}^l_{\bar{\mu}}(f,\theta)) [g] \|_{\infty} = \mathcal{O}(\|g\|_{bl} \cdot d_{bl}(\mu,\bar{\mu})).$$
(127)

We now bound $\|Df_{\theta}^{\bar{\mu}}[h]\|_{bl}$ $(g = Df_{\theta}^{\bar{\mu}}[h])$. From the fixed point definition of the Sinkhorn potential in (107), we can compute the Fréchet derivative Df_{θ}^{μ} by

$$Df^{\mu}_{\theta} = D_1 \mathcal{A} \left(\mathcal{A}(f^{\mu}_{\theta}, \alpha_{\theta}), \beta_{\mu} \right) \circ D_1 \mathcal{A}(f^{\mu}_{\theta}, \alpha_{\theta}) \circ Df^{\mu}_{\theta} + D_1 \mathcal{A} \left(\mathcal{A}(f^{\mu}_{\theta}, \alpha_{\theta}), \beta_{\mu} \right) \circ D_2 \tilde{\mathcal{A}}(f^{\mu}_{\theta}, \theta),$$
(128)

where we recall $\tilde{\mathcal{A}}(f,\theta):=\mathcal{A}(f,\alpha_{\theta})$. For any direction $h \in \mathbb{R}^d$ and any $y \in \mathcal{X}$, $Df_{\theta}^{\mu}[h]$ is a function with its gradient bounded by

$$\begin{aligned} \|\nabla_{y}Df^{\mu}_{\theta}[h](y)\| &\leq \|\nabla_{y}\left(D_{1}\mathcal{A}\left(\mathcal{A}(f^{\mu}_{\theta},\alpha_{\theta}),\beta_{\mu}\right)\left[D_{1}\mathcal{A}(f^{\mu}_{\theta},\alpha_{\theta})\left[Df^{\mu}_{\theta}[h]\right]\right]\right)(y)\| & \#1\\ &+ \|\nabla_{y}\left(D_{1}\mathcal{A}\left(\mathcal{A}(f^{\mu}_{\theta},\alpha_{\theta}),\beta_{\mu}\right)\left[D_{2}\tilde{\mathcal{A}}(f^{\mu}_{\theta},\theta)[h]\right]\right)(y)\|. & \#2\end{aligned}$$

We now bound the R.H.S. individually:

For #1, take $\bar{f} = \mathcal{A}[f, \alpha_{\theta}], \kappa = \beta_{\mu}$ and $g = D_1 \mathcal{A}(f_{\theta}^{\mu}, \alpha_{\theta}) [Df_{\theta}^{\mu}[h]]$ in (121). Using (123) and (118), we have

$$#1 = \mathcal{O}(\|g\|_{\infty}) = \mathcal{O}(\|Df^{\mu}_{\theta}[h]\|_{\infty}) = \mathcal{O}(\|h\|).$$
(129)

For #2, take $\bar{f} = \mathcal{A}[f, \alpha_{\theta}], \kappa = \beta_{\mu}$ and $g = D_2 \tilde{\mathcal{A}}(f_{\theta}^{\mu}, \theta)[h]$ in (121). Using (123) and (79), we have

$$#2 = \mathcal{O}(\|g\|_{\infty}) = \mathcal{O}(\|D_2\tilde{\mathcal{A}}(f^{\mu}_{\theta}, \theta)[h]\|_{\infty}) = \mathcal{O}(\|h\|).$$
(130)

Combining these two bounds, we have

$$\|Df^{\mu}_{\theta}[h]\|_{bl} = \mathcal{O}(\|h\|).$$
(131)

By plugging the above result to (127), we bound

$$(3) = \| (D_1 \mathcal{B}^l_{\mu}(f, \theta) - D_1 \mathcal{B}^l_{\bar{\mu}}(f, \theta)) [g] \|_{\infty} = \mathcal{O}(d_{bl}(\mu, \bar{\mu}) \cdot \|h\|).$$
(132)

C.5.4 Bounding (4)

We have from the triangle inequality

$$(\Phi \leq \| D_2 \mathcal{E}_{\mu}(f^{\mu}_{\theta}, \theta)[h] - D_2 \mathcal{E}_{\bar{\mu}}(f^{\mu}_{\theta}, \theta)[h] \|_{\infty} + \| D_2 \mathcal{E}_{\bar{\mu}}(f^{\mu}_{\theta}, \theta)[h] - D_2 \mathcal{E}_{\bar{\mu}}(f^{\bar{\mu}}_{\theta}, \theta)[h] \|_{\infty}.$$
 (133)

We analyze these two terms on the R.H.S..

For the first term of (133), use the chain rule of Fréchet derivative to compute

$$D_{2}\mathcal{E}_{\mu}(f,\theta)[h] = D_{1}\mathcal{B}_{\mu}\left(\mathcal{B}_{\mu}^{l-1}(f,\theta),\theta\right)\left[D_{2}\mathcal{B}_{\mu}^{l-1}(f,\theta)[h]\right] + D_{2}\mathcal{B}_{\mu}\left(\mathcal{B}_{\mu}^{l-1}(f,\theta),\theta\right)[h].$$
(134)

Consequently, we can bound

$$\begin{aligned} &\| \left(D_{2} \mathcal{E}_{\mu}(f,\theta) - D_{2} \mathcal{E}_{\bar{\mu}}(f,\theta) \right)[h] \|_{\infty} \\ \leqslant \| D_{1} \mathcal{B}_{\mu} \left(\mathcal{B}_{\mu}^{l-1}(f,\theta), \theta \right) \left[D_{2} \mathcal{B}_{\mu}^{l-1}(f,\theta)[h] \right] - D_{1} \mathcal{B}_{\bar{\mu}} \left(\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta), \theta \right) \left[D_{2} \mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta)[h] \right] \|_{\infty} \quad \#1 \\ &+ \| D_{2} \mathcal{B}_{\mu} \left(\mathcal{B}_{\mu}^{l-1}(f,\theta), \theta \right)[h] - D_{2} \mathcal{B}_{\bar{\mu}} \left(\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta), \theta \right)[h] \|_{\infty}. \qquad \#2 \end{aligned}$$

We analyze #1 and #2 individually.

Bounding #1. We first note that $\mathcal{A}(f, \alpha)$ is Lipschitz continuous w.r.t. α (see also (124)):

$$\|\mathcal{A}(f,\alpha) - \mathcal{A}(f,\alpha')\|_{\infty} \leq \exp(2M_c/\gamma) \cdot \|\omega_y\|_{bl} \cdot d_{bl}(\alpha,\alpha') = \mathcal{O}(d_{bl}(\alpha,\alpha')),$$
(135)

where in the equality we use (119). As \mathcal{B}^k_{μ} is the composition of \mathcal{A} , it is Lipschitz continuous with respect to μ for finite k. Note that the boundedness of $||f||_{\infty}$ and $||\nabla f||_{\infty}$ remains valid after the

operator \mathcal{B} (Lemma B.1 and (i) of Lemma (B.2)). We then bound

$$\#1 \leq \|D_{1}\mathcal{B}_{\mu}(\mathcal{B}_{\mu}^{l-1}(f,\theta),\theta)[(D_{2}\mathcal{B}_{\mu}^{l-1}(f,\theta) - D_{2}\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta))[h]]\|_{\infty}$$

$$+ \|(D_{1}\mathcal{B}_{\mu}(\mathcal{B}_{\mu}^{l-1}(f,\theta),\theta) - D_{1}\mathcal{B}_{\mu}(\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta),\theta))[D_{2}\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta)[h]]\|_{\infty}$$

$$+ \|(D_{1}\mathcal{B}_{\mu}(\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta),\theta) - D_{1}\mathcal{B}_{\bar{\mu}}(\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta),\theta))(D_{2}\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta)[h]]\|_{\infty}$$

$$\leq \|D_{1}\mathcal{B}_{\mu}(\mathcal{B}_{\mu}^{l-1}(f,\theta),\theta)\|_{op}\|D_{2}\mathcal{B}_{\mu}^{l-1}(f,\theta)[h] - D_{2}\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta)[h]\|_{\infty}$$

$$+ \mathcal{O}(\|\mathcal{B}_{\mu}^{l-1}(f,\theta) - \mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta)\|_{\infty} \cdot \|D_{2}\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta)[h]\|_{\infty})$$

$$+ \mathcal{O}(d_{bl}(\mu,\bar{\mu}) \cdot \|D_{2}\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta)[h]\|_{\infty})$$

$$\leq \|D_{2}\mathcal{B}_{\mu}^{l-1}(f,\theta)[h] - D_{2}\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta)[h]\|_{\infty})$$

where in the second inequality we use the definition of $\|\cdot\|_{op}$, (115) and (125), and in the last inequality we use the fact that $\|D_1\mathcal{B}_{\mu}(f,\theta)\|_{op} \leq 1$, \mathcal{B}^k_{μ} is Lipschitz continuous with respect to μ for finite k (see the discussion above) and that $\|D_2\mathcal{B}^{l-1}_{\mu}(f,\theta)\|_{op}$ is bounded (see Lemma B.7.

Bounding #2. To make the dependences of A on θ and μ explicit, we denote

$$\hat{\mathcal{A}}(f,\theta,\mu) = \mathcal{A}(f,T_{\theta\sharp}\mu)$$

To bound the second term, we first establish that for any $k \ge 0$, $\nabla \mathcal{B}_{\mu}^{k+1}(f, \theta)$ is Lipschitz continuous w.r.t. μ , i.e.

$$\|\nabla \mathcal{B}^{k+1}_{\mu}(f,\theta) - \nabla \mathcal{B}^{k+1}_{\bar{\mu}}(f,\theta)\|_{2,\infty} = \mathcal{O}(d_{bl}(\mu,\bar{\mu})),$$
(136)

as follows: First note that $\nabla \hat{\mathcal{A}}(f, \theta, \mu)$ is Lipschitz continuous w.r.t. μ , i.e.

$$\|\nabla \hat{\mathcal{A}}(f,\theta,\mu)(y) - \nabla \hat{\mathcal{A}}(f,\theta,\bar{\mu})(y)\| = \mathcal{O}(d_{bl}(\mu,\bar{\mu})).$$
(137)

This is because for any $y \in \mathcal{X}$ (note that $\hat{\mathcal{A}}(f, \theta, \mu)(\cdot) : \mathcal{X} \to \mathbb{R}$ is a function of y),

$$\begin{split} \|\nabla \hat{\mathcal{A}}(f,\theta,\mu)(y) - \nabla \hat{\mathcal{A}}(f,\theta,\bar{\mu})(y)\| \\ &= \|\frac{\int_{\mathcal{X}} \omega_y(x) \nabla_1 c(y,x) \mathbf{d}\alpha_\theta(x)}{\int_{\mathcal{X}} \omega_y(x) \mathbf{d}\alpha_\theta(x)} - \frac{\int_{\mathcal{X}} \omega_y(x) \nabla_1 c(y,x) \mathbf{d}\bar{\alpha}_\theta(x)}{\int_{\mathcal{X}} \omega_y(x) \mathbf{d}\bar{\alpha}_\theta(x)}\| \\ &\leq \|\frac{\int_{\mathcal{X}} \omega_y(x) \nabla_1 c(y,x) (\mathbf{d}\alpha_\theta(x) - \mathbf{d}\bar{\alpha}_\theta(x))}{\int_{\mathcal{X}} \omega_y(x) \mathbf{d}\alpha_\theta(x)}\| \\ &+ \|\int_{\mathcal{X}} \omega_y(x) \nabla_1 c(y,x) \mathbf{d}\bar{\alpha}_\theta(x)\| \cdot \|\frac{\int_{\mathcal{X}} \omega_y(x) (\mathbf{d}\alpha_\theta(x) - \mathbf{d}\bar{\alpha}_\theta(x))}{\int_{\mathcal{X}} \omega_y(x) \mathbf{d}\alpha_\theta(x) \int_{\mathcal{X}} \omega_y(x) \mathbf{d}\bar{\alpha}_\theta(x)}\| \\ &= \mathcal{O}(d_{bl}(\mu,\bar{\mu})). \end{split}$$

Here in the last equality, we use the facts that $\|\omega_y(\cdot)\nabla_1 c(y,\cdot)\|_{bl}$ and $\|\omega_y\|_{bl}$ are bounded, and $\int_{\mathcal{X}} \omega_y(x) d\alpha_\theta(x)$ is strictly positive and bounded away from zero. Recall that $\mathcal{B}_\mu(f,\theta) = \mathcal{A}(\hat{\mathcal{A}}(f,\theta,\mu),\beta_\mu)$. We can then prove (136) by bounding

Here we bound &1 using (137), the Lipschitz continuity of ∇A w.r.t. its second variable; we bound &2 using the Lipschitz continuity of $\nabla \hat{A}$ w.r.t. its first variable and (124), the Lipschitz continuity of \hat{A} w.r.t. μ ; we bound &3 using (124), the Lipschitz continuity of \hat{A} w.r.t. μ , and the fact that \mathcal{B}_{μ}^{k} is the composition of the terms $\mathcal{A}(f, \alpha)$ and $\mathcal{A}(f, \beta_{\mu})$.

We then establish that $D_2 \mathcal{B}_{\mu}(f, \theta)$ is Lipschitz continuous w.r.t. μ .

Assumption C.1. $\|\nabla_z [\nabla_\theta T_\theta(z)]\|_{op}$ is bounded

Lemma C.12. Assume that $||f||_{\infty} \leq M_c$, $||\nabla f||_{2,\infty} \leq G_f$, $||\nabla^2 f||_{op,\infty} \leq L_f$ Under Assumptions B.5, C.1 and B.1, we have

$$\|D_2\mathcal{B}_{\mu}(f,\theta) - D_2\mathcal{B}_{\bar{\mu}}(f,\theta)\|_{op} = \mathcal{O}(d_{bl}(\mu,\bar{\mu})).$$
(138)

Proof. Denote $\omega_y(x) = \exp\left(\frac{-c(x,y)+f(x)}{\gamma}\right)$ and

$$\phi_y(z) = [\nabla_\theta T_\theta(z)]^\top [-\nabla_1 c(T_\theta(z), y) + \nabla f(T_\theta(z))]$$

where $\nabla_{\theta} T_{\theta}(z)$ denotes the Jacobian matrix of $T_{\theta}(z)$ with respect to θ .

The Fréchet derivative $D_2 \hat{\mathcal{A}}(f, \theta, \mu)[h]$ can be computed by

$$D_2\hat{\mathcal{A}}(f,\theta,\mu)[h] = \frac{\int_{\mathcal{X}} \omega_y (T_\theta(z)) \langle \phi_y(z), h \rangle \mathbf{d}\mu(z)}{\int_{\mathcal{X}} \omega_y (T_\theta(z)) \mathbf{d}\mu(z)}.$$
(139)

Recall that $||f||_{\infty} \leq M_c$, $||\nabla f||_{2,\infty} \leq G_f$. Using the above expression we can bound

$$\begin{split} &\| \left(D_2 \mathcal{A}(f,\theta,\mu) - D_2 \mathcal{A}(f,\theta,\bar{\mu}) \right) [h] \|_{\infty} \\ &= \| \frac{\int_{\mathcal{X}} \omega_y (T_{\theta}(z)) \langle \phi_y(z), h \rangle d\mu(z)}{\int_{\mathcal{X}} \omega_y (T_{\theta}(z)) d\mu(z)} - \frac{\int_{\mathcal{X}} \omega_y (T_{\theta}(z)) \langle \phi_y(z), h \rangle d\bar{\mu}(x)}{\int_{\mathcal{X}} \omega_y (T_{\theta}(z)) \langle \phi_y(z), h \rangle d\mu(z)} \|_{\infty} \\ &\leq \| \frac{\int_{\mathcal{X}} \omega_y (T_{\theta}(z)) \langle \phi_y(z), h \rangle d\mu(z)}{\int_{\mathcal{X}} \omega_y (T_{\theta}(z)) d\mu(z)} - \frac{\int_{\mathcal{X}} \omega_y (T_{\theta}(z)) \langle \phi_y(z), h \rangle d\bar{\mu}(x)}{\int_{\mathcal{X}} \omega_y (T_{\theta}(z)) d\mu(z)} \|_{\infty} \\ &+ \| \frac{\int_{\mathcal{X}} \omega_y (T_{\theta}(z)) \langle \phi_y(z), h \rangle d\bar{\mu}(x)}{\int_{\mathcal{X}} \omega_y (T_{\theta}(z)) d\mu(z)} - \frac{\int_{\mathcal{X}} \omega_y (T_{\theta}(z)) \langle \phi_y(z), h \rangle d\bar{\mu}(x)}{\int_{\mathcal{X}} \omega_y (T_{\theta}(z)) d\bar{\mu}(x)} \|_{\infty} \\ &= \| \frac{\int_{\mathcal{X}} \omega_y (T_{\theta}(z)) \langle \phi_y(z), h \rangle [d\mu(z) - d\bar{\mu}(x)]}{\int_{\mathcal{X}} \omega_y (T_{\theta}(z)) d\mu(z)} \|_{\infty} \\ &+ \| \frac{\int_{\mathcal{X}} \omega_y (T_{\theta}(z)) \langle \phi_y(z), h \rangle d\bar{\mu}(x)}{\int_{\mathcal{X}} \omega_y (T_{\theta}(z)) d\mu(z)} \|_{\infty} \\ &\leq \exp(2M_c/\gamma) \cdot \| \omega_y (T_{\theta}(z)) \langle \phi_y(z), h \rangle \|_{bl} \cdot d_{bl}(\mu, \bar{\mu}) \\ &+ \exp(5M_c/\gamma) \cdot \| \phi_y \|_{\infty} \cdot \| h \|_{\infty} \cdot \| \omega_y \|_{bl} \cdot d_{bl}(\mu, \bar{\mu}). \end{split}$$

For the first term, note that $\|\omega_y(T_\theta(z))\langle \phi_y(z),h\rangle\|_{bl} \leq \|\omega_y\|_{bl} \cdot \|\phi_y\|_{bl} \cdot \|h\|_{\infty}$ and $\|\omega_y\|_{bl}$ is bounded (see (119)). We just need to bound $\|\phi_y\|_{bl}$. Under Assumption B.5 that $\|\nabla_\theta T_\theta(z)\|_{op} \leq G_T$, we clearly have that $\|\phi_y\|_{\infty}$ is bounded. For $\|\phi_y\|_{lip}$, compute that

$$\nabla_{z}\phi_{y}(z) = \nabla_{z}[\nabla_{\theta}T_{\theta}(z)] \times_{1} \left[-\nabla_{1}c(T_{\theta}(z), y) + \nabla f(T_{\theta}(z))\right] \\ + \nabla_{\theta}T_{\theta}(z)^{\top} \left[-\nabla_{11}^{2}c(T_{\theta}(z), y) + \nabla^{2}f(T_{\theta}(z))\right] \nabla_{\theta}T_{\theta}(z).$$

Recall that $\|\nabla^2 f(x)\|_{op}$ is bounded. Consequently, under Assumption C.1, we can see that $\|\nabla_z \phi_y(z)\|$ is bounded. Together, $\|\phi_y\|_{bl}$ is bounded. As a result, we have

$$\| \left(D_2 \hat{\mathcal{A}}(f,\theta,\mu) - D_2 \hat{\mathcal{A}}(f,\theta,\bar{\mu}) \right) [h] \|_{\infty} = \mathcal{O}(d_{bl}(\mu,\bar{\mu}) \cdot \|h\|).$$
(140)

Based on the above result, we can further bound

$$\begin{split} &\| \left(D_2 \mathcal{B}_{\mu}(f,\theta) - D_2 \mathcal{B}_{\bar{\mu}}(f,\theta) \right) [h] \|_{\infty} \\ &= \| \left(D_1 \mathcal{A} \left(\hat{\mathcal{A}}(f,\theta,\mu),\beta \right) \circ D_2 \hat{\mathcal{A}}(f,\theta,\mu) - D_1 \mathcal{A} \left(\hat{\mathcal{A}}(f,\theta,\bar{\mu}),\bar{\beta} \right) \circ D_2 \hat{\mathcal{A}}(f,\theta,\bar{\mu}) \right) [h] \|_{\infty} \\ &\leq \| D_1 \mathcal{A} \left(\hat{\mathcal{A}}(f,\theta,\mu),\beta \right) \left[\left(D_2 \hat{\mathcal{A}}(f,\theta,\mu) - D_2 \hat{\mathcal{A}}(f,\theta,\bar{\mu}) \right) [h] \right] \|_{\infty} \qquad \# \# 1 \end{split}$$

$$+ \left\| \left(D_1 \mathcal{A} \big(\hat{\mathcal{A}}(f, \theta, \mu), \beta \big) - D_1 \mathcal{A} \big(\hat{\mathcal{A}}(f, \theta, \mu), \bar{\beta} \big) \right) \left[D_2 \hat{\mathcal{A}}(f, \theta, \bar{\mu})[h] \right] \right\|_{\infty} \qquad \# \# 2$$

$$+ \| \left(D_1 \mathcal{A} \big(\hat{\mathcal{A}}(f, \theta, \mu), \bar{\beta} \big) - D_1 \mathcal{A} \big(\hat{\mathcal{A}}(f, \theta, \bar{\mu}), \bar{\beta} \big) \right) [D_2 \hat{\mathcal{A}}(f, \theta, \bar{\mu})[h]] \|_{\infty}. \qquad \# \# 3$$

For the first term, use $\|D_1\mathcal{A}(\hat{\mathcal{A}}(f,\theta,\mu),\beta)\|_{op} \leq 1$ (75) and (140) to bound

$$##1 \leq \|D_2\hat{\mathcal{A}}(f,\theta,\mu)[h] - D_2\hat{\mathcal{A}}(f,\theta,\bar{\mu})[h]\|_{\infty} = \mathcal{O}(d_{bl}(\mu,\bar{\mu}) \cdot \|h\|).$$

For the second term, recall the expression of $D_2 \hat{\mathcal{A}}(f, \theta, \bar{\mu})[h]$ in (139). Under Assumption B.1 and assume that $||f||_{\infty} \leq M_c$, one can see that $||D_2 \hat{\mathcal{A}}(f, \theta, \bar{\mu})[h]|_{bl} = \mathcal{O}(||h||)$. Further, use (122) and $d_{bl}(\beta, \bar{\beta}) = \mathcal{O}(d_{bl}(\mu, \bar{\mu}))$ from (101) to bound

$$##2 = \mathcal{O}(\|D_2\hat{\mathcal{A}}(f,\theta,\bar{\mu})[h]\|_{bl} \cdot d_{bl}(\beta,\bar{\beta})) = \mathcal{O}(\|h\| \cdot d_{bl}(\mu,\bar{\mu})).$$

For the third term, use Lemma B.11 to bound

$$##3 = \mathcal{O}(\|D_2\hat{\mathcal{A}}(f,\theta,\bar{\mu})[h]\|_{\infty} \cdot \|\hat{\mathcal{A}}(f,\theta,\mu) - \hat{\mathcal{A}}(f,\theta,\bar{\mu})\|_{\infty}) = \mathcal{O}(d_{bl}(\mu,\bar{\mu}) \cdot \|h\|),$$

where we use $\|D_2 \hat{\mathcal{A}}(f, \theta, \bar{\mu})[h]\|_{\infty} = \mathcal{O}(\|h\|)$ and (124). Altogether, we have

$$\|D_2 \mathcal{B}_{\mu}(f,\theta)[h] - D_2 \mathcal{B}_{\bar{\mu}}(f,\theta)[h]\|_{\infty} = \mathcal{O}(d_{bl}(\mu,\bar{\mu}) \cdot \|h\|).$$

$$(141)$$

We are now ready to bound #2.

$$\begin{aligned} \#2 &\leq \|D_{2}\mathcal{B}_{\mu}(\mathcal{B}_{\mu}^{l-1}(f,\theta),\theta)[h] - D_{2}\mathcal{B}_{\mu}(\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta),\theta)[h]\|_{\infty} \\ &+ \|D_{2}\mathcal{B}_{\mu}(\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta),\theta)[h] - D_{2}\mathcal{B}_{\bar{\mu}}(\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta),\theta)[h]\|_{\infty} \\ &= \mathcal{O}(\|\mathcal{B}_{\mu}^{l-1}(f,\theta) - \mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta)\|_{\infty} + \|\nabla\mathcal{B}_{\mu}^{l-1}(f,\theta) - \nabla\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta)\|_{2,\infty}) \\ &+ \mathcal{O}(d_{bl}(\mu,\bar{\mu}) \cdot \|h\|) \\ &= \mathcal{O}(d_{bl}(\mu,\bar{\mu}) \cdot \|h\|), \end{aligned}$$

where we use Lemma B.10 and (138) (124) in the first equality.

Combining #1 and #2. Combining the above results, we yield

$$\|D_{2}\mathcal{B}_{\mu}^{l}(f,\theta)[h] - D_{2}\mathcal{B}_{\bar{\mu}}^{l}(f,\theta)[h]\|_{\infty} \leq \|D_{2}\mathcal{B}_{\mu}^{l-1}(f,\theta)[h] - D_{2}\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta)[h]\|_{\infty} + \mathcal{O}(d_{bl}(\mu,\bar{\mu}) \cdot \|h\|_{\infty}),$$

which, via recursion, implies that (recall that $D_2 \mathcal{E}_{\mu}(f,\theta)[h] = D_2 \mathcal{B}^l_{\mu}(f,\theta)[h]$)

$$\|D_2 \mathcal{E}_{\mu}(f,\theta)[h] - D_2 \mathcal{E}_{\bar{\mu}}(f,\theta)[h]\|_{\infty} = \mathcal{O}(d_{bl}(\mu,\bar{\mu}) \cdot \|h\|).$$
(142)

To bound the second term of (133), compute the expression of $D_2 \mathcal{E}_{\bar{\mu}}(f,\theta)[h]$ via the chain rule:

$$D_{2}\mathcal{E}_{\bar{\mu}}(f,\theta)[h] = D_{1}\mathcal{B}_{\bar{\mu}}\left(\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta),\theta\right) \left[D_{2}\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta)[h]\right] + D_{2}\mathcal{B}_{\bar{\mu}}\left(\mathcal{B}_{\bar{\mu}}^{l-1}(f,\theta),\theta\right)[h].$$
(143)

Recall that $\mathcal{E}_{\bar{\mu}}(f,\theta) = \mathcal{B}_{\bar{\mu}}^{l}(f,\theta)$. We then show in an inductive manner that the second term of (133) is of order $\mathcal{O}(d_{bl}(\mu,\bar{\mu}) \cdot ||h||)$: For any finite $k \ge 1$,

$$\|D_2 \mathcal{B}^k_{\bar{\mu}}(f^{\mu}_{\theta}, \theta)[h] - D_2 \mathcal{B}^k_{\bar{\mu}}(f^{\bar{\mu}}_{\theta}, \theta)[h]\|_{\infty} = \mathcal{O}(d_{bl}(\mu, \bar{\mu}) \cdot \|h\|).$$
(144)

For the base case when l = 1, we only have the second term of (143) in $D_2 \mathcal{E}_{\bar{\mu}}(f, \theta)[h]$. Consequently, from Lemma B.10, we have

$$\begin{split} \|D_{2}\mathcal{B}_{\bar{\mu}}\big(\mathcal{B}_{\bar{\mu}}^{l-1}(f_{\theta}^{\mu},\theta),\theta\big) - D_{2}\mathcal{B}_{\bar{\mu}}\big(\mathcal{B}_{\bar{\mu}}^{l-1}(f_{\theta}^{\bar{\mu}},\theta),\theta\big)\|_{op} \\ &= \mathcal{O}(\|\mathcal{B}_{\bar{\mu}}^{l-1}(f_{\theta}^{\mu},\theta) - \mathcal{B}_{\bar{\mu}}^{l-1}(f_{\theta}^{\bar{\mu}},\theta)\|_{\infty} + \|\nabla\mathcal{B}_{\bar{\mu}}^{l-1}(f_{\theta}^{\mu},\theta) - \nabla\mathcal{B}_{\bar{\mu}}^{l-1}(f_{\theta}^{\bar{\mu}},\theta)\|_{2,\infty}) = \mathcal{O}(d_{bl}(\mu,\bar{\mu})), \end{split}$$
(145)

where we use (136) in the second equality. Now assume that for l = k the statement (144) holds. For any two function $f, f' \in C(\mathcal{X})$, we bound

Plug in $f = f_{\theta}^{\mu}$ and $f' = f_{\theta}^{\bar{\mu}}$ and use Lemmas C.1 and C.2. We prove the statement (144) holds for l = k + 1. Consequently, we have that

$$\|D_2 \mathcal{E}_{\bar{\mu}}(f^{\mu}_{\theta}, \theta)[h] - D_2 \mathcal{E}_{\bar{\mu}}(f^{\bar{\mu}}_{\theta}, \theta)[h]\|_{\infty} = \mathcal{O}(d_{bl}(\mu, \bar{\mu}) \cdot \|h\|).$$
(146)

In conclusion, we have

$$(\underline{4}) = \mathcal{O}(d_{bl}(\mu, \bar{\mu}) \cdot ||h||). \tag{147}$$

Layer (type) Conv2d-1	Output Shape [-1, 64, 32, 32]	Param # 4.800
LeakyReLU-2	[-1, 64, 32, 32]	0
Conv2d-3	[-1, 128, 16, 16]	204,800
BatchNorm2d-4	[-1, 128, 16, 16]	256
LeakyReLU-5	[-1, 128, 16, 16]	0
Conv2d-6	[-1, 256, 8, 8]	819,200
BatchNorm2d-7	[-1, 256, 8, 8]	512
LeakyReLU-8	[-1, 256, 8, 8]	0
Conv2d-9	[-1, 512, 4, 4]	3,276,800
BatchNorm2d-10	[-1, 512, 4, 4]	1,024
LeakyReLU-11	[-1, 512, 4, 4]	0

Table 1: Structure of the encoder

Table 2: Structure of the generator

Layer (type)	Output Shape	Param #
ConvTranspose2d-1	[-1, 256, 4, 4]	262,144
BatchNorm2d-2	[-1, 256, 4, 4]	512
ReLU-3	[-1, 256, 4, 4]	0
ConvTranspose2d-4	[-1, 128, 8, 8]	524,288
BatchNorm2d-5	[-1, 128, 8, 8]	256
ReLU-6	[-1, 128, 8, 8]	0
ConvTranspose2d-7	[-1, 64, 16, 16]	131,072
BatchNorm2d-8	[-1, 64, 16, 16]	128
ReLU-9	[-1, 64, 16, 16]	0
ConvTranspose2d-10	[-1, 3, 32, 32]	3,072
Tanh-11	[-1, 3, 32, 32]	0

D Experiment Details

We use the generator from DC-GAN Radford et al. [2015]. And the adversarial ground cost c_{ξ} in the form of

$$c_{\xi}(x,y) = \|\phi_{\xi}(x) - \phi_{\xi}(y)\|_{2}^{2},$$
(148)

where $\phi_{\xi} : \mathbb{R}^q \to \mathbb{R}^{\hat{q}}$ is an encoder that maps the original data point (and the generated image) to a higher dimensional space ($\hat{q} > q$). We pick ϕ_{ξ} to be an CNN with a similar structure as the discriminator of DC-GAN except that we discard the last layer which was used for classification. Specifically, the networks used are given in Table 1 and 2.

We set the step size β of SiNG to be 30 and set the maximum allow Sinkhorn divergence in each iteration to be 0.1. Note that the step size is set after the normalization in (11). For Adam, RMSprop, and AMSgrad, we set all of their initial step sizes to be $1.0 \times e^{-3}$, which is in general recommended by the GAN literature. The minibatch sizes of both the real images and the generated images for each iteration are set to 3000. We uniformly set the γ parameter in the objective (recall that $\mathcal{F}(\alpha_{\theta}) = S_{c_{\xi}}(\alpha_{\theta}, \beta)$) and the constraint to 100.

The code is in https://github.com/shenzebang/Sinkhorn_Natural_Gradient.

E PyTorch Implementation

In this section, we focus on the empirical version of SiNG, where we approximate the gradient of the function F by a minibatch stochastic gradient and approximate SIM by eSIM. In this case, all components involved in the optimization procedure can be represented by finite dimensional vectors.

It is known that the stochastic gradient admits an easy implementation in PyTorch. However, at the first sight, the computation of eSIM is quite complicated as it requires to construct two sequences f^t and g^t to estimate the Sinkhorn potential and the Fréchet derivative. As we discussed earlier, it is well known that we can solve the inversion of a p.s.d. matrix via the Conjugate Gradient (CG) method with only matrix-vector-product operations. In particular, in this case, we no longer need to explicitly form eSIM in the computer memory. Consequently, to implement the empirical version of SiNG using CG and eSIM, one can resort to the auto-differential mechanism provided by PyTorch: First, we use existing PyTorch package like geomloss³ to compute the tensor f representing the Sinkhorn potential f^{ϵ}_{θ} . Note the the sequence f^t is constructed implicitly by calling geomloss. We then use the ".detach()" function in PyTorch to maintain only the value of the f while discarding all of its "grad_fn" entries. We then enable the "autograd" mechanism is PyTorch and run several loops of Sinkhorn mapping $\mathcal{A}(f, \alpha_{\theta})$ ($\mathcal{A}(f, \alpha_{\theta t})$) so that the output tensor now records all the dependence on the parameter θ via the implicitly constructed computational graph. We can then easily compute the matrix-vector-product use the Pearlmutter's algorithm (Pearlmutter, 1994).

³https://www.kernel-operations.io/geomloss/