
CHARACTERIZATIONS OF NON-NORMALIZED DISCRETE PROBABILITY DISTRIBUTIONS AND THEIR APPLICATION IN STATISTICS

A PREPRINT

S. Betsch

Institute of Stochastics,
Karlsruhe Institute of Technology (KIT),
Germany.
Steffen.Betsch@kit.edu

B. Ebner

Institute of Stochastics,
Karlsruhe Institute of Technology (KIT),
Germany.
Bruno.Ebner@kit.edu

F. Nestmann

Institute of Stochastics,
Karlsruhe Institute of Technology (KIT),
Germany.
Franz.Nestmann2@kit.edu

October 20, 2021

ABSTRACT

From the distributional characterizations that lie at the heart of Stein's method we derive explicit formulae for the mass functions of discrete probability laws that identify those distributions. These identities are applied to develop tools for the solution of statistical problems. Our characterizations, and hence the applications built on them, do not require any knowledge about normalization constants of the probability laws. To demonstrate that our statistical methods are sound, we provide comparative simulation studies for the testing of fit to the Poisson distribution and for parameter estimation of the negative binomial family when both parameters are unknown. We also consider the problem of parameter estimation for discrete exponential-polynomial models which generally are non-normalized.

1 Introduction

In recent years, research in probability and statistics has witnessed the rise of Stein's method but also the emergence of methods to tackle the analysis and application of models which are based on non-normalized probability laws. In this work, we seek to apply findings from the research on Stein's method to contribute to the solution of testing and estimation problems involving non-normalized statistical distributions. We focus on the analysis of discrete probability laws, and how the theoretical results can be used to develop statistical methods. As such, we tie on to Betsch and Ebner (2021) who provide similar tools for continuous distributions. A rather well-known approach to the problem of parameter estimation for non-normalized continuous probability distributions is the score matching technique due to Hyvärinen (2005, 2007) (see Yu et al., 2019, for recent progress). Another approach is known as noise contrastive estimation (cf. Gutmann and Hyvärinen, 2010), but a number of 2019/20 papers indicate that the proposition and study of new tools remains an important issue, see Matsuda and Hyvärinen (2019), Uehara et al. (2020a), and Uehara et al. (2020b).

The tool box which is now known as Stein's method goes back to the work of Stein (1972) (see also Stein, 1986) who sought for an alternative proof of the central limit theorem that provides a bound on the rate of the convergence.

MSC 2020 subject classifications. Primary 62E10 Secondary 62G10, 65C05

Key words and phrases. Discrete exponential-polynomial models; goodness-of-fit tests; negative binomial distribution; non-normalized models; Stein characterizations

This inherent feature made Stein's method popular. The idea is applied in all kinds of settings, as it allows to find bounds on distributional distances between sequences of probability laws and a limit distribution, and as it often applies in the absence of stochastic independence. The application of the method to discrete distributions goes back to Chen (1975), who first derived corresponding results for the Poisson distribution, known as the Stein-Chen method. The method has since been extended to other discrete distributions, like the binomial distribution (by Ehm, 1991), the geometric distribution (by Peköz, 1996), the negative binomial distribution (by Brown and Phillips, 1999), discrete Gibbs measures (by Eichelsbacher and Reinert, 2008), and others. The foundation of the method are characterization results for the underlying probability law. While for the first distributions in consideration specific identities were used or devised, general approaches have emerged that apply to many different distributions at once. In this context, we mention the generator approach of Barbour (1988, 1990) and Götze (1991) who use time-reversible Markov processes, where the stationary distribution is the probability law of interest, to characterize that law. On the other hand, a direct derivation of characterizations is possible, and a well-known class of such identities can be found under the name of 'density approach'. For the continuous case, first ideas on the density approach came from Stein (1986) and Stein et al. (2004), and a more complete version is due to Ley and Swan (2013b). The corresponding characterizations for discrete distributions are given by Ley and Swan (2013a).

The contribution at hand is certainly not the first application of Stein's method in statistics. Indeed, similar problems in the context of non-normalized models are tackled with the use of so-called Stein discrepancies by the machine learning community, though many of them refer to the continuous setting. Let us mention some papers that explore these tools. Namely, Chwiałkowski et al. (2016), Liu et al. (2016), and Yang et al. (2018) consider the construction of tests of fit, Gorham and Mackey (2015) build measures of sample quality, and Barp et al. (2019) solve estimation problems for non-normalized models.

Our new work is based, in a strict sense, not on what is generally called Stein's method, but rather on the characterization identities we refer to above. More precisely, we take as a starting point the discrete density approach identity as provided by Ley and Swan (2013a). To sketch the idea, consider a probability mass function p on \mathbb{N}_0 as well as an \mathbb{N}_0 -valued random variable X . Subject to few regularity conditions, X is governed by p if, and only if,

$$\mathbb{E} \left[\Delta^+ f(X) + \frac{\Delta^+ p(X)}{p(X)} f(X+1) \right] = 0$$

holds for a large enough class of test functions f . Hereby Δ^+ denotes the forward difference operator. Our first contribution lies in proving that this characterization can essentially be restated as to X being governed by p if, and only if, the probability mass function ρ_X of X satisfies

$$\rho_X(k) = \mathbb{E} \left[- \frac{\Delta^+ p(X)}{p(X)} \mathbb{1}\{X \geq k\} \right], \quad k \in \mathbb{N}_0.$$

With regard to applications in statistics, this second identity is more accessible. We can, for instance, tackle the goodness-of-fit testing problem as follows. Assume we are to test whether a sample X_1, \dots, X_n of \mathbb{N}_0 -valued random variables follows one of the laws of a parametric family of distributions $\{p_\vartheta : \vartheta \in \Theta\}$, where Θ denotes the parameter space. By the above characterization, if X_1, \dots, X_n are governed by one of the p_ϑ , then the difference between

$$\hat{\rho}_n(k) = \frac{1}{n} \sum_{j=1}^n \mathbb{1}\{X_j = k\}$$

and

$$\frac{1}{n} \sum_{j=1}^n - \frac{\Delta^+ p_{\hat{\vartheta}_n}(X_j)}{p_{\hat{\vartheta}_n}(X_j)} \mathbb{1}\{X_j \geq k\}$$

ought to be small for each $k \in \mathbb{N}_0$. Here we denote by $\hat{\vartheta}_n$ an estimator of ϑ based on X_1, \dots, X_n . Thus, in line with the idea of characterization based goodness-of-fit testing, our proposal is to use

$$\sum_{k=0}^{\infty} \left(\frac{1}{n} \sum_{j=1}^n \mathbb{1}\{X_j = k\} + \frac{1}{n} \sum_{j=1}^n \frac{\Delta^+ p_{\hat{\vartheta}_n}(X_j)}{p_{\hat{\vartheta}_n}(X_j)} \mathbb{1}\{X_j \geq k\} \right)^2$$

as a test statistic for the hypothesis

$$\mathbf{H}_0 : \rho_{X_1} \in \{p_\vartheta : \vartheta \in \Theta\},$$

and to reject the hypothesis for large values of the statistic. Supposing that X_1, \dots, X_n are governed by p_{ϑ_0} for some (unknown) $\vartheta_0 \in \Theta$, the very same heuristic leads us to propose

$$\hat{\vartheta}_n = \operatorname{argmin}_{\vartheta \in \Theta} \sum_{k=0}^{\infty} \left(\frac{1}{n} \sum_{j=1}^n \mathbb{1}\{X_j = k\} + \frac{1}{n} \sum_{j=1}^n \frac{\Delta^+ p_\vartheta(X_j)}{p_\vartheta(X_j)} \mathbb{1}\{X_j \geq k\} \right)^2$$

as an estimator for the unknown ϑ_0 . The paper at hand formalizes these ideas and puts them on firm mathematical ground. We also provide examples for the theoretical results as well as for the testing and estimation methods we propose. In Section 2 we introduce basic notation and recall the density approach identity. In Section 3 we prove the new characterization result as indicated above. Section 4 contains further characterizations based on transformations of the probability mass function such as distribution functions, characteristic functions, and generating functions. In Section 5 we discuss examples. We then construct and study empirically, in Section 6, the test of fit for the Poisson distribution. In Section 7 a discrepancy measure as above leads to minimum distance estimators for the negative binomial distribution, which are put to a test in a simulation study. Section 8 deals with similar parameter estimators in the non-normalized class of discrete exponential-polynomial models.

2 The foundation: Stein characterizations for discrete distributions

We denote by $p : \mathbb{Z} \rightarrow [0, 1]$ a probability mass function (pmf) defined on the integers. We assume that the support of p , that is, $\text{spt}(p) = \{k \in \mathbb{Z} : p(k) > 0\}$ is connected, in the sense that

$$(C1) \quad \text{spt}(p) = \{L, L+1, \dots, R\}, \text{ where } L, R \in \mathbb{Z} \cup \{\pm\infty\}, L < R.$$

This prerequisite is quite usual in the context of Stein's method in the discrete setting. We further assume that

$$(C2) \quad \sup_{k \in \{L, \dots, R-1\}} \left| \frac{\Delta^+ p(k) \cdot \min\{P(k), 1 - P(k)\}}{p(k)p(k+1)} \right| < \infty,$$

where $\Delta^+ f(k) = f(k+1) - f(k)$ is the forward difference operator. Moreover, we denote by $P(k) = \sum_{\ell=L}^k p(\ell)$ the distribution function corresponding to p . Assumption (C2) is known from the continuous setting, see Lemma 13.1 of Chen et al. (2011). The supremum in (C2) runs from L to ∞ whenever $R = \infty$. In what follows, we stick to the convention of empty sums being set to 0.

Definition 2.1. Let p be a pmf that satisfies (C1) and (C2). We write \mathcal{F}_p for the class of functions $f : \{L, \dots, R\} \rightarrow \mathbb{R}$ such that

$$(a) \quad \sum_{k=L}^R |\Delta^+(p(k)f(k))| < \infty \text{ and } \sum_{k=L}^R \Delta^+(p(k)f(k)) = 0, \text{ where we put } f(R+1) = 0 \text{ if } R < \infty, \text{ as well as}$$

$$(b) \quad \sup_{k \in \{L, \dots, R\}} |\Delta^+ f(k)| < \infty, \text{ and } \sup_{k \in \{L, \dots, R\}} \left| \frac{\Delta^+ p(k)}{p(k)} f(k+1) \right| < \infty.$$

Conditions (C2) and (b) vanish completely whenever the support of p is finite. We now state the characterization theorem that is known as Stein's density approach for discrete distributions. The proof is an easy adaptation of the proof of Theorem 2.1 from Ley and Swan (2013a) taking into account the different class of test functions. We give a full proof in Appendix A to make it possible for the reader to comprehend how the assumptions come into play. Denote by $(\Omega, \mathcal{A}, \mathbb{P})$ the probability space which underlies all random quantities in this work.

Theorem 2.2 (Discrete density approach). *Let p be a pmf which satisfies (C1) and (C2), and let $X : \Omega \rightarrow \mathbb{R}$ be a random variable such that $\mathbb{P}(X \in \text{spt}(p)) > 0$. Then, $\mathbb{P}(X = k | X \in \text{spt}(p)) = p(k)$, $k \in \mathbb{Z}$, if and only if,*

$$\mathbb{E} \left[\Delta^+ f(X) + \frac{\Delta^+ p(X)}{p(X)} f(X+1) \mid X \in \text{spt}(p) \right] = 0,$$

for all $f \in \mathcal{F}_p$, where $\mathbb{E}[\cdot | \cdot]$ denotes the conditional expectation.

We use the abbreviation $X|p \sim p$ for $\mathbb{P}(X = k | X \in \text{spt}(p)) = p(k)$, $k \in \mathbb{Z}$. There exists a very similar result for continuous probability distributions. This continuous version existed first and was initiated by Stein (1986). For the fully prepared statement we refer to Ley and Swan (2011) and Ley and Swan (2013b), and for further constructions of this type of Stein operators, see Ley et al. (2017).

Remark 2.3. It follows from the proof of Theorem 2.2 that, in the case $L > -\infty$, we may assume that $f(L) = 0$ for all $f \in \mathcal{F}_p$.

3 Distributional characterizations via the probability mass function

In this section, we derive explicit distributional characterizations via the probability mass function. The whole theory can be understood as a discrete version of the results from Betsch and Ebner (2021) who established similar characterization

identities for continuous probability laws starting with a continuous version of Theorem 2.2 as stated by Ley and Swan (2013b). We make the further assumption that the expectation of p exists, that is,

(C3) $\mathbb{E}|Z| < \infty$, where Z is a discrete random variable with pmf p .

It follows from (C3) that $\mathbb{E}|\Delta^+ p(Z) \cdot Z / p(Z)| < \infty$, and hence we also have $\mathbb{E}|\Delta^+ p(Z) / p(Z)| < \infty$. We note our first result, a proof of which is given in Appendix B.

Theorem 3.1. *Let p be a pmf which satisfies (C1) – (C3) with $L > -\infty$. Let $X : \Omega \rightarrow \mathbb{R}$ be a random variable with $\mathbb{P}(X \in \text{spt}(p)) > 0$ as well as*

$$\mathbb{E} \left| \frac{\Delta^+ p(X)}{p(X)} X \cdot \mathbb{1}\{X \in \text{spt}(p)\} \right| < \infty,$$

and denote by $\rho_{X|p}(k) = \mathbb{P}(X = k | X \in \text{spt}(p))$ the pmf of X given $X \in \text{spt}(p)$. Then, $X|p \sim p$ if, and only if,

$$\rho_{X|p}(k) = \mathbb{E} \left[- \frac{\Delta^+ p(X)}{p(X)} \mathbb{1}\{X \geq k\} \mid X \in \text{spt}(p) \right], \quad k \in \mathbb{Z}, k \geq L.$$

Notice that the integrability assumption on X implies the existence of the (conditional) expectation that appears in the theorem. Even in stating Theorem 3.1 the ordering of the integers is essential. However, if p is an admissible probability mass function on some arbitrary countable set \mathbb{S} (where \mathbb{S} is endowed with the power set as a σ -field), there exists a bijection $\iota : \mathbb{S} \rightarrow \mathbb{N}_0$, which corresponds to imposing an order on the space \mathbb{S} , and Theorem 3.1 can be applied. This leads to the following corollary which allows the handling of more general state spaces.

Corollary 3.2. *Let \mathbb{S} be a countable set and $p : \mathbb{S} \rightarrow [0, 1]$ such that $\sum_{s \in \mathbb{S}} p(s) = 1$. Let $\iota : \mathbb{S} \rightarrow \{L, \dots, R\}$, with $L > -\infty$, be a bijection so that $\tilde{p} = p \circ \iota^{-1}$ satisfies (C1) – (C3). Assume that $X : \Omega \rightarrow \mathbb{S}$ is a random variable with $\mathbb{P}(X \in \text{spt}(p)) > 0$, and*

$$\mathbb{E} \left| \frac{(\Delta^+ \tilde{p})(\iota(X))}{p(X)} \iota(X) \cdot \mathbb{1}\{X \in \text{spt}(p)\} \right| < \infty.$$

Then, $X|p \sim p$ if, and only if,

$$\rho_{X|p}(k) = \mathbb{E} \left[- \frac{(\Delta^+ \tilde{p})(\iota(X))}{p(X)} \mathbb{1}\{\iota(X) \geq \iota(s)\} \mid X \in \text{spt}(p) \right], \quad s \in \mathbb{S}.$$

Any such ordering on \mathbb{S} gives a characterization result, and if $X \sim p$, the (conditional) expectation is the same for every ordering (as $\rho_{X|p}$ does not depend on ι). However, if one intends to use the converse of the characterization (with general X), the calculation of the expectation depends on the ordering, so in practice the question of choosing an efficient ordering arises. Finding an order such that the condition (C1) – (C3) are satisfied is a non-trivial endeavor. We give one example of choosing an order such that a pmf with a support that is not bounded from below can be considered. To state the result, we first need to recall that the backward difference operator Δ^- is defined by $\Delta^- f(k) = f(k) - f(k-1)$.

Corollary 3.3. *Let p be a pmf on $\{L, \dots, R\}$ which satisfies (C1) – (C3) with $R < \infty$. Let $X : \Omega \rightarrow \mathbb{R}$ be a random variable which satisfies $\mathbb{P}(X \in \text{spt}(p)) > 0$ as well as*

$$\mathbb{E} \left| \frac{\Delta^- p(X)}{p(X)} X \cdot \mathbb{1}\{X \in \text{spt}(p)\} \right| < \infty.$$

Then, $X|p \sim p$ if, and only if,

$$\rho_{X|p}(k) = \mathbb{E} \left[\frac{\Delta^- p(X)}{p(X)} \mathbb{1}\{X \leq k\} \mid X \in \text{spt}(p) \right], \quad k \in \mathbb{Z}, k \leq R.$$

The result follows from Corollary 3.2 upon choosing $\iota : \{L, \dots, R\} \rightarrow \{-R, \dots, -L\}$, $\iota(k) = -k$, and observing that

$$(\Delta^+ \tilde{p})(\iota(X)) = (\Delta^+ \tilde{p})(-X) = p(X-1) - p(X) = -\Delta^- p(X).$$

Note that Corollary 3.3 can also be obtained via a different path. With few technical changes in Definition 2.1 and Appendix A, a Δ^- -version of Theorem 2.2 can be formulated (see also Ley and Swan, 2013a). Using this result and an adaptation of the proof of Theorem 3.1 yields another proof of Corollary 3.3.

Remark 3.4. Whenever X is assumed a priori to take values in $\{L, \dots, R\}$, the conditioning on $X \in \text{spt}(p)$ can be omitted, and when $-\infty < L < R < \infty$, the integrability condition on X is trivially satisfied. As for the regularity assumptions (C1) – (C3), notice that, by Corollary 3.2, (C1) is mostly an issue of notation but no hard restriction. Whenever we deal with discrete distributions that have finite support, conditions (C2) and (C3) are trivially satisfied. In case of an infinite support, assumption (C3) is easy to interpret. It is stated to guarantee that the statement of Theorem 3.1 is consistent, as it ensures that a random variable $Z \sim p$ satisfies the integrability condition on X . A drawback in terms of the assumptions is that we cannot give a general treatment of (C2), and that this condition can sometimes be difficult to check for a given distribution. A similar condition (with identical problems) is required by Betsch and Ebner (2021) in the continuous setting. If $L > -\infty$ and $R = \infty$ then (C2) holds if, and only if,

$$\limsup_{k \rightarrow \infty} \left| \frac{\Delta^+ p(k) \cdot (1 - P(k))}{p(k) p(k+1)} \right| < \infty. \quad (1)$$

Similar thoughts apply to other choices for L and R , but this does not solve the problem in general. However, the Stolz-Cesàro theorem (see Theorem 2.7.1 of Choudary and Niculescu, 2014) provides a useful tool for checking the condition in practice, see Example 5.3.

Remark 3.5 (Non-normalized models). As we have explained in the introduction, many statistical models, primarily in machine learning and physics, are too complex for the normalization constant of the distribution to be calculable. As estimation and testing procedures (e.g. the maximum likelihood estimator) normally rely on some knowledge about this constant, they may not be applicable to such models. Thus, we want to emphasize that our explicit characterizations do not need any knowledge about the normalization constants, and neither do any of the characterizations and statistical applications presented in subsequent sections.

4 Characterizations via transformations of the probability mass function

We also obtain characterizing formulae for transformations of the pmf, like distribution functions, characteristic functions, and probability generating functions. We focus on the identities derived for mass functions on the integers but more general spaces can be treated in the lines of Corollary 3.2.

Proposition 4.1 (Distribution functions). *Let p be a pmf which satisfies (C1) – (C3) with $L > -\infty$. Let $X : \Omega \rightarrow \mathbb{R}$ be a random variable with $\mathbb{P}(X \in \text{spt}(p)) > 0$ as well as*

$$\mathbb{E} \left| \frac{\Delta^+ p(X)}{p(X)} X \cdot \mathbb{1}\{X \in \text{spt}(p)\} \right| < \infty, \quad (2)$$

and further denote by $F_{X|p}(k) = \mathbb{P}(X \leq k | X \in \text{spt}(p))$ the distribution function of X given $X \in \text{spt}(p)$. Then, $X|p \sim p$ if, and only if,

$$F_{X|p}(k) = \mathbb{E} \left[- \frac{\Delta^+ p(X)}{p(X)} (\min\{X, k\} - L + 1) \mid X \in \text{spt}(p) \right], \quad k \in \mathbb{Z}, k \geq L.$$

The proof of this proposition is provided in Appendix C. We continue to give another characterization based on the characteristic function. The proof (see Appendix D) features the inversion formula for characteristic functions. In the continuous setting, the inversion formula requires an integrability condition that is not needed in the discrete setting. If one finds a way to handle this integrability condition, similar identities are conceivable for the continuous setting of Betsch and Ebner (2021).

Proposition 4.2 (Characteristic functions). *Assume that p is a pmf which satisfies (C1) – (C3) with $L > -\infty$. Take $X : \Omega \rightarrow \mathbb{R}$ to be a random variable with $\mathbb{P}(X \in \text{spt}(p)) > 0$ such that assumption (2) holds. Denote by $\varphi_{X|p}(t) = \mathbb{E}[e^{itX} | X \in \text{spt}(p)]$, $t \in \mathbb{R}$, the characteristic function of X given $X \in \text{spt}(p)$ (where i is the complex unit). Then, $X|p \sim p$ if, and only if,*

$$\varphi_{X|p}(t) = \mathbb{E} \left[- \frac{\Delta^+ p(X)}{p(X)} \cdot \frac{e^{itL} - e^{it(X+1)}}{1 - e^{it}} \mid X \in \text{spt}(p) \right], \quad t \in \mathbb{R}.$$

We conclude this section on transformations of the pmf with a characterization via the probability generating function, thus specializing to the case $\text{spt}(p) = \mathbb{N}_0$. A proof is given in Appendix E.

Proposition 4.3 (Generating functions). *Let p be a pmf that satisfies (C1) – (C3) with $\text{spt}(p) = \mathbb{N}_0$. Let $X : \Omega \rightarrow \mathbb{N}_0$ be a discrete random variable such that*

$$\mathbb{E} \left| \frac{\Delta^+ p(X)}{p(X)} X \right| < \infty.$$

Denote by $G_X(s) = \mathbb{E}[s^X]$, $s \in [0, 1)$, the (probability) generating function of X . Then, $X \sim p$ if, and only if,

$$G_X(s) = \mathbb{E} \left[-\frac{\Delta^+ p(X)}{p(X)} \cdot \frac{1 - s^{X+1}}{1 - s} \right], \quad s \in [0, 1).$$

5 Examples

In this section, we provide examples that fit into our framework. For each distribution we indicate why (C1) – (C3) hold and we explicitly state the characterization via Theorem 3.1, though sometimes in the unconditioned formulation. The characterizations via Propositions 4.1, 4.2, and 4.3 are not stated explicitly. We start by giving three infinite support examples which are subject of our statistical applications in the subsequent sections. More precisely, we discuss the Poisson and the negative binomial distribution as well as a discrete version of the exponential-polynomial model.

Example 5.1 (Poisson distribution). The mass function of the Poisson distribution is given as $p(k) = \lambda^k \cdot e^{-\lambda} / k!$, $k \in \mathbb{N}_0$, for some rate parameter $\lambda > 0$. In this case, we obtain

$$\frac{\Delta^+ p(k)}{p(k)} = \frac{\lambda}{k+1} - 1, \quad k \in \mathbb{N}_0.$$

Conditions (C1) and (C3) are obviously true. To see that (C2) holds, note that whenever $\frac{\lambda}{k+2} < 1$, we have

$$\begin{aligned} \left| \frac{\Delta^+ p(k) \cdot (1 - P(k))}{p(k) p(k+1)} \right| &= \left| \frac{\lambda}{k+1} - 1 \right| \left| \sum_{\ell=1}^{\infty} \frac{\lambda^{\ell-1}}{(\ell+k)!} (k+1)! \right| \leq \left| \frac{\lambda}{k+1} - 1 \right| \left| \sum_{\ell=0}^{\infty} \left(\frac{\lambda}{k+2} \right)^{\ell} \right| \\ &= \left| \frac{\lambda}{k+1} - 1 \right| \cdot \frac{1}{1 - \frac{\lambda}{k+2}}, \end{aligned}$$

and therefore (1) holds which yields (C2). Theorem 3.1 implies that a random variable $X : \Omega \rightarrow \mathbb{N}_0$ with $\mathbb{E}X < \infty$ has the Poisson distribution with parameter λ if, and only if,

$$\rho_X(k) = \mathbb{E} \left[\left(1 - \frac{\lambda}{X+1} \right) \mathbb{1}_{\{X \geq k\}} \right], \quad k \in \mathbb{N}_0.$$

Example 5.2 (Negative binomial distribution). Let $p(k) = \binom{k+r-1}{k} (1-q)^k q^r$, $k \in \mathbb{N}_0$, be the probability mass function of the negative binomial distribution with parameters $r > 0$ and $q \in (0, 1)$. An important special case arises for $r = 1$, where the negative binomial distribution reduces to the geometric distribution. These laws are frequently used in the analysis of arrival times. We have

$$\frac{\Delta^+ p(k)}{p(k)} = \frac{r+k}{k+1} (1-q) - 1, \quad k \in \mathbb{N}_0.$$

Condition (C1) is trivially satisfied, and (C3) is easily verified. We prove (1) to show that (C2) is satisfied. To this end, observe that

$$\begin{aligned} \left| \frac{\Delta^+ p(k) \cdot (1 - P(k))}{p(k) p(k+1)} \right| &= \left| \frac{r+k}{k+1} (1-q) - 1 \right| \left| \sum_{\ell=0}^{\infty} \frac{(\ell+k+r)! \cdot (k+1)!}{(\ell+k+1)! \cdot (k+r)!} (1-q)^{\ell} \right| \\ &= \left| \frac{r+k}{k+1} (1-q) - 1 \right| \left| \sum_{\ell=0}^{\infty} \frac{(r+k+\ell) \cdot (r+k+\ell-1) \cdots (r+k+1)}{(k+\ell+1) \cdot (k+\ell) \cdots (k+2)} (1-q)^{\ell} \right|. \end{aligned}$$

If $r \leq 1$, the sum on the right hand-side is bounded by $\sum_{\ell=0}^{\infty} (1-q)^{\ell} = \frac{1}{q}$. If $r > 1$, let k be large enough so that $2 \cdot (r-1) / (k+2) < q / (1-q)$, and observe that

$$\begin{aligned} \sum_{\ell=0}^{\infty} \frac{(r+k+\ell) \cdots (r+k+1)}{(k+\ell+1) \cdots (k+2)} (1-q)^{\ell} &= \sum_{\ell=0}^{\infty} \left(1 + \frac{r-1}{k+\ell+1} \right) \cdot \left(1 + \frac{r-1}{k+\ell} \right) \cdots \left(1 + \frac{r-1}{k+2} \right) \cdot (1-q)^{\ell} \\ &\leq \sum_{\ell=0}^{\infty} \left(1 + \frac{1}{2} \cdot \frac{q}{1-q} \right)^{\ell} (1-q)^{\ell} = \sum_{\ell=0}^{\infty} \left(1 - \frac{q}{2} \right)^{\ell} = \frac{2}{q}, \end{aligned}$$

where the products in the sum are empty (hence equal to 1) for $\ell = 0$. In any case, (1) follows, so (C2) is valid. Theorem 3.1 states that a discrete random variable $X : \Omega \rightarrow \mathbb{N}_0$ with $\mathbb{E}X < \infty$ follows the negative binomial law with parameters r and q if, and only if,

$$\rho_X(k) = \mathbb{E} \left[\left(1 - \frac{r+X}{X+1} (1-q) \right) \mathbb{1}\{X \geq k\} \right], \quad k \in \mathbb{N}_0.$$

Note that the statement by Johnson et al. (1993) (on p. 223) that "only a few characterizations have been obtained for the negative binomial distribution" appears to still hold true. For one recent characterization related to Stein's method, we refer to Arras and Houdré (2019).

Example 5.3 (Exponential-polynomial models). We consider the following discrete exponential-polynomial parametric model given through

$$p_\vartheta(k) = C(\vartheta)^{-1} \exp(\vartheta_1 k + \dots + \vartheta_d k^d), \quad k \in \mathbb{N},$$

where

$$C(\vartheta) = \sum_{k=1}^{\infty} \exp(\vartheta_1 k + \dots + \vartheta_d k^d),$$

and $\vartheta = (\vartheta_1, \dots, \vartheta_d) \in \mathbb{R}^{d-1} \times (-\infty, 0)$. This corresponds to a discrete exponential family in the canonical form with the sufficient statistic containing monomials up to order $d \in \mathbb{N}$, with $d \geq 2$. Clearly condition (C1) is satisfied and the restriction $\vartheta_d < 0$ ensures that (C3) holds for every ϑ as well as that $C(\vartheta) < \infty$. We have

$$\frac{p_\vartheta(k+1)}{p_\vartheta(k)} = \exp(\vartheta_1 + \vartheta_2((k+1)^2 - k^2) + \dots + \vartheta_d((k+1)^d - k^d)) \rightarrow 0,$$

as $k \rightarrow \infty$, and the Stolz-Cesàro theorem (Theorem 2.7.1 of Choudary and Niculescu, 2014) yields

$$\lim_{k \rightarrow \infty} \frac{p_\vartheta(k+1)}{1 - P_\vartheta(k)} = - \lim_{k \rightarrow \infty} \frac{p_\vartheta(k+2)}{p_\vartheta(k+1)} + 1 = 1.$$

Consequently, we obtain

$$\limsup_{k \rightarrow \infty} \left| \frac{\Delta^+ p_\vartheta(k) \cdot (1 - P_\vartheta(k))}{p_\vartheta(k) p_\vartheta(k+1)} \right| \leq \limsup_{k \rightarrow \infty} \left| \frac{p_\vartheta(k+1)}{p_\vartheta(k)} - 1 \right| \cdot \left| \frac{1 - P_\vartheta(k)}{p_\vartheta(k+1)} \right| = 1$$

for every $\vartheta \in \mathbb{R}^{d-1} \times (-\infty, 0)$, so (C2) holds. Finally observe that, since $p_\vartheta(k+1)/p_\vartheta(k) < 1$ for all but finitely many $k \in \mathbb{N}$, an \mathbb{N} -valued random variable with $\mathbb{E}X < \infty$ also satisfies

$$\mathbb{E} \left| \frac{\Delta^+ p_\vartheta(X)}{p_\vartheta(X)} X \right| < \infty.$$

Theorem 3.1 yields that a random variable $X : \Omega \rightarrow \mathbb{N}$ with $\mathbb{E}X < \infty$ has the pmf p_ϑ if, and only if,

$$\rho_X(k) = \mathbb{E} \left[\left(-\exp(\vartheta_1 + \vartheta_2((X+1)^2 - X^2) + \dots + \vartheta_d((X+1)^d - X^d)) + 1 \right) \mathbb{1}\{X \geq k\} \right], \quad k \in \mathbb{N}.$$

In Section 8 we use this characterization to construct an estimation method for this type of parametric model, focusing on a two-parameter case where $d = 3$ and $\vartheta_2 = 0$ fixed.

We now take a look at the resulting characterization for the uniform and binomial distribution.

Example 5.4 (Uniform distribution). Assume that p is given through $p(k) = 1/m$, for $k = 1, \dots, m$, $m \geq 2$. Then, (C1) – (C3) are obviously satisfied (recall Remark 3.4), and we have

$$\frac{\Delta^+ p(k)}{p(k)} = 0, \quad \text{for } k = 1, \dots, m-1, \quad \text{and} \quad \frac{\Delta^+ p(m)}{p(m)} = -1.$$

By Theorem 3.1, a random variable $X : \Omega \rightarrow \mathbb{R}$ with $\mathbb{P}(X \in \{1, \dots, m\}) > 0$ satisfies $X|p \sim p$ if, and only if,

$$\rho_{X|p}(k) = \frac{\mathbb{P}(X = m)}{\mathbb{P}(X \in \{1, \dots, m\})}, \quad k = 1, \dots, m,$$

that is, if $\rho_{X|p}(k) = 1/m$, for $k = 1, \dots, m$. This result is easily derived by a direct argument, so for the uniform distribution, our characterization contains no new information. A similar behavior was observed in the continuous setting by Betsch and Ebner (2021). Note that Corollary 3.3 leads to the very same characterization as Theorem 3.1.

Example 5.5 (Binomial distribution). Let $p(k) = \binom{m}{k} q^k (1-q)^{m-k}$, for $k = 0, \dots, m$, $m \geq 1$, and some fixed $q \in (0, 1)$. Then, we have

$$\frac{\Delta^+ p(k)}{p(k)} = \frac{q}{1-q} \cdot \frac{m-k}{k+1} - 1, \quad \text{for } k = 0, \dots, m-1, \quad \text{and} \quad \frac{\Delta^+ p(m)}{p(m)} = -1.$$

By Theorem 3.1, a random variable $X : \Omega \rightarrow \mathbb{R}$ with $\mathbb{P}(X \in \{0, \dots, m\}) > 0$ satisfies $X|p \sim p$ if, and only if,

$$\rho_{X|p}(k) = \frac{\sum_{\ell=k}^{m-1} \left(1 - \frac{q}{1-q} \cdot \frac{m-\ell}{\ell+1}\right) \mathbb{P}(X = \ell) + \mathbb{P}(X = m)}{\mathbb{P}(X \in \{0, \dots, m\})}, \quad k = 0, \dots, m.$$

Moreover, we have

$$\frac{\Delta^- p(k)}{p(k)} = 1 - \frac{1-q}{q} \cdot \frac{k}{m-k+1}, \quad \text{for } k = 1, \dots, m, \quad \text{and} \quad \frac{\Delta^- p(0)}{p(0)} = 1,$$

so Corollary 3.3 yields that $X|p \sim p$ if, and only if,

$$\rho_{X|p}(k) = \frac{\sum_{\ell=1}^k \left(1 - \frac{1-q}{q} \cdot \frac{\ell}{m-\ell+1}\right) \mathbb{P}(X = \ell) + \mathbb{P}(X = 0)}{\mathbb{P}(X \in \{0, \dots, m\})}, \quad k = 0, \dots, m.$$

Thus the example of the binomial distribution shows that Theorem 3.1 and Corollary 3.3 do not always lead to the same identity in cases where both are applicable.

We conclude this section on examples by showing that discrete Gibbs measures, which describe physical systems with countably many states, also fall into our framework.

Example 5.6 (Gibbs (or Boltzmann) distribution). We distinguish in our discussion between finite and infinite support.

- Assume that a given system can have $S \in \mathbb{N}$ states, $S \geq 2$, and let $V : \{1, \dots, S\} \rightarrow \mathbb{R} \cup \{\infty\}$ be a map (called the energy function) which assigns each state its corresponding energy. Another map $N : \{1, \dots, S\} \rightarrow \mathbb{N}$ assigns to each state the number of particles the system has in the given state. Let $\mu \in \mathbb{R}$ (the chemical potential), $T > 0$ (the temperature) and denote by κ the Boltzmann constant ($\approx 1.380649 \cdot 10^{-23}$ joule per kelvin). The Gibbs distribution is given through

$$p(k) = \frac{1}{\Xi} \exp\left(\frac{\mu N(k) - V(k)}{\kappa T}\right), \quad k \in \{1, \dots, S\},$$

where

$$\Xi = \Xi(\mu, T, N, V) = \sum_{\ell=1}^S \exp\left(\frac{\mu N(\ell) - V(\ell)}{\kappa T}\right),$$

and where we assume that $V(k) < \infty$ for at least one $k \in \{1, \dots, S\}$ (which ensures $\Xi > 0$). In this setting with finitely many states, (C1) – (C3) are trivially satisfied. We obtain $\Delta^+ p(S) / p(S) = -1$, and

$$\frac{\Delta^+ p(k)}{p(k)} = \exp\left(\frac{V(k) - V(k+1) + \mu(N(k+1) - N(k))}{\kappa T}\right) - 1, \quad k = 1, \dots, S-1.$$

Theorem 3.1 yields that a random variable $X : \Omega \rightarrow \{1, \dots, S\}$ follows the Gibbs distribution as above if, and only if, for all $k = 1, \dots, S$,

$$\rho_X(k) = \sum_{\ell=k}^{S-1} \left(1 - \exp\left(\frac{V(\ell) - V(\ell+1) + \mu(N(\ell+1) - N(\ell))}{\kappa T}\right)\right) \mathbb{P}(X = \ell) + \mathbb{P}(X = S).$$

- The Gibbs distribution immediately generalizes to the case where $S = \infty$, that is, a system with infinitely many possible states. In this general setting however, one has to make further assumptions on μ , T , N , and V to ensure that $\Xi < \infty$ and that (C2) and (C3) hold. One set of assumptions that ensures $\Xi < \infty$ and (C3) is that the number of particles is fixed $N \equiv \bar{N} \in \mathbb{N}$ and that the probability of the system being in one of the states with higher index decreases sufficiently fast, or equivalently, that the energy grows fast enough. More precisely, it is sufficient to assume that there exists some $k_0 \in \mathbb{N}$ such that, for all $k \geq k_0$, we have

$V(k) \geq c \cdot k^\alpha$, for some $c > 0$ and $\alpha \geq 1$. In order for (C2) to hold, we could additionally assume that there exists some $k_1 \in \mathbb{N}$ and $c' > 0$ such that, for all $k \geq k_1$,

$$V(k) - V(k+n) \leq -c' n, \quad n \in \mathbb{N}_0,$$

as this implies

$$\left| \frac{\Delta^+ p(k) \cdot \min\{P(k), 1 - P(k)\}}{p(k)p(k+1)} \right| \leq 2 \sum_{n=0}^{\infty} \left(e^{-\frac{c'}{\kappa T}} \right)^n < \infty, \quad k \geq k_1.$$

One choice of V (to satisfy all of the conditions) is thus $V(k) = C \cdot k$ for all k larger than some fixed $k^* \in \mathbb{N}$ and some $C > 0$. The characterization via Theorem 3.1 in the case of infinitely many possible states is similar as in the finite case: We have, for any $k \in \mathbb{N}$,

$$\frac{\Delta^+ p(k)}{p(k)} = \exp\left(\frac{V(k) - V(k+1)}{\kappa T}\right) - 1.$$

A random variable $X : \Omega \rightarrow \mathbb{N}$ with $\mathbb{E}X < \infty$ follows the (infinite states) Gibbs distribution if, and only if,

$$\rho_X(k) = \mathbb{E} \left[\left(1 - \exp\left(\frac{V(X) - V(X+1)}{\kappa T}\right) \right) \mathbb{1}\{X \geq k\} \right], \quad k \in \mathbb{N}.$$

As is indicated by the names that certain quantities in the above display of the Gibbs distribution carry, the model appears in statistical mechanics, see the reprint, Gibbs (2010), of Josiah Willard Gibbs' work from 1902, or virtually any textbook on statistical mechanics. It also plays an important role in image analysis and processing, see Li (2009). A crucial observation about our characterizations is that the partition function Ξ vanishes completely.

6 Goodness-of-fit testing for the Poisson distribution

A first application of the characterization results from the previous sections is the construction of a test of fit for the Poisson distribution. Given a sample of \mathbb{N}_0 -valued independent identically distributed (i.i.d.) random variables, the problem is to test the composite hypothesis that the sample comes from some Poisson distribution $\text{Po}(\lambda)$ with an unknown rate parameter $\lambda > 0$, that is,

$$\mathbf{H}_0 : \mathbb{P}^{X_1} \in \{\text{Po}(\lambda) : \lambda > 0\}.$$

This is a classical statistical problem, well studied in the literature. Apart from Pearson's χ^2 test, see Khmaladze (2013) for recent developments, the hitherto proposed tests are based on the (conditional) empirical distribution function, see Beltrán-Beltrán and O'Reilly (2019); Frey (2012); Gürtler and Henze (2000); Henze (1996), the empirical probability generating function, see Baringhaus and Henze (1992); Puig and Weiß (2020); Rueda and O'Reilly (1999), on the integrated distribution function, see Klar (1999), on a characterization by mean distance, see Székely and Rizzo (2004), on quadratic forms of score vectors, see Inglot (2019), on Charlier polynomials, see Ledwina and Wylupek (2017), on conditional probabilities ratio, see Beltrán-Beltrán and O'Reilly (2019), and on relating first- and second-order moments, see Kyriakoussis et al. (1998). For a survey of classical procedures and a comparative simulation study see Gürtler and Henze (2000). However, the construction of new and powerful methods is still of relevance: As Nikitin (2017) stated on p.4 of his contribution that "[...] one should keep in mind that any hypothesis has to be tested with several possible criteria. The point of the matter is that with absolute confidence we can only reject it, while each new test which fails to reject the null-hypothesis gradually brings the statistician closer to the perception that this hypothesis is true".

The idea of our new method is to estimate the two quantities that appear in the characterization via Theorem 3.1 as given in Example 5.1, and to compare these empirical quantities. Based on the sample X_1, \dots, X_n , let

$$\hat{e}_n(k) = \frac{1}{n} \sum_{j=1}^n \left(1 - \frac{\hat{\lambda}_n}{X_j + 1} \right) \mathbb{1}\{X_j \geq k\}, \quad k \in \mathbb{N}_0,$$

be an estimator of the expectation that arises in the characterization, where $\hat{\lambda}_n = n^{-1} \sum_{j=1}^n X_j$ is a consistent estimator of the rate parameter. Also consider the empirical probability mass function,

$$\hat{\rho}_n(k) = \frac{1}{n} \sum_{j=1}^n \mathbb{1}\{X_j = k\}, \quad k \in \mathbb{N}_0,$$

as an estimator of ρ_{X_1} . By Theorem 3.1 (see Example 5.1), if the sample X_1, \dots, X_n comes from a Poisson distribution, the absolute difference between $\hat{e}_n(k)$ and $\hat{\rho}_n(k)$ ought to be small for every $k \in \mathbb{N}_0$. On the other hand, if the sample does not come from the Poisson law, we expect their absolute difference to be large. Based on this heuristic, we suggest to use as a test statistic the squared difference of \hat{e}_n and $\hat{\rho}_n$ summed over $k \in \mathbb{N}_0$, that is,

$$T_n^{Po} = \sum_{k=0}^{\infty} (\hat{e}_n(k) - \hat{\rho}_n(k))^2,$$

and to reject the Poisson hypothesis \mathbf{H}_0 for large values of T_n^{Po} . Note that we do not need to introduce any weight functions to make the infinite sum in the definition of T_n^{Po} converge, and observe that we choose the squared distance to obtain a finite double sum representation for T_n^{Po} , namely

$$T_n^{Po} = \frac{1}{n^2} \sum_{j, \ell=1}^n \left[\left(1 - \frac{\hat{\lambda}_n}{X_j + 1} \right) (X_\ell - 1 - \hat{\lambda}_n) \mathbb{1}\{X_j \geq X_\ell\} \right. \\ \left. + (X_j + 1 - \hat{\lambda}_n) \left(1 - \frac{\hat{\lambda}_n}{X_\ell + 1} \right) \mathbb{1}\{X_j < X_\ell\} + \mathbb{1}\{X_j = X_\ell\} \right],$$

which is a numerically stable representation, hence easily implemented in a computer. The calculation of T_n^{Po} involves only straight forward algebra and consists, mainly, of writing the squared difference of $\hat{e}_n(k)$ and $\hat{\rho}_n(k)$ as a double sum, multiplying the corresponding terms, and solving the sum over k of the indicator functions.

Remark 6.1. The major advantage in using Theorem 3.1 to construct the test is that the empirical quantities are integrable with respect to the counting measure $\sum_{k=0}^{\infty} \delta_k$. Of course, the Propositions 4.1, 4.2, and 4.3 provide similar heuristics for the construction of goodness-of-fit tests [see also Betsch and Ebner (2019), Allison et al. (2019), and Betsch and Ebner (2020) for the continuous setting], but the quantities are not necessarily integrable/summable and thus require the introduction of some weight function. What is more, it seems difficult to obtain explicit formulae as in the previous equation for T_n^{Po} , so the routines would rely on numerical integration which is computationally costly. We must therefore leave it as a problem of further research to employ these characterizations via distribution, characteristic, or generating functions in the construction of goodness-of-fit tests.

As a proof of concept, we carry out a simulation study in order to compare our new test of poissonity with established procedures. All simulations are performed in the statistical computing environment R, see R Core Team (2020). We consider the sample size $n = 50$ and the nominal level of significance is set to 0.05. Based on the methodology for asymptotic theory detailed by Henze (1996), we expect the (limit) distribution of the test statistics considered in the following to depend on the unknown parameter λ . Consequently, we use for the implementation of the tests a similar parametric bootstrap procedure as the one suggested by Gürtler and Henze (2000). For a given sample X_1, \dots, X_n and a statistic T_n , simulate an approximate critical value $c_{n,B}^*$ for a level α test procedure as follows:

- 1) Calculate $\hat{\lambda}_n(X_1, \dots, X_n)$ and generate B bootstrap samples of size n with distribution $\text{Po}(\hat{\lambda}_n)$, i.e., generate i.i.d. $\text{Po}(\hat{\lambda}_n)$ random variables $X_{j,1}^*, \dots, X_{j,n}^*, j = 1, \dots, B$.
- 2) Compute $T_{j,n}^* = T_n(X_{j,1}^*, \dots, X_{j,n}^*)$ for $j = 1, \dots, B$.
- 3) Derive the order statistics $T_{1:B}^* \leq \dots \leq T_{B:B}^*$ of $T_{1,n}^*, \dots, T_{B,n}^*$ and calculate

$$c_{n,B}^* = T_{k:B}^* + (1 - \alpha) \cdot (T_{(k+1):B}^* - T_{k:B}^*),$$

where $k = \lfloor (1 - \alpha) \cdot B \rfloor$ and $\lfloor \cdot \rfloor$ denotes the floor function.

- 4) Reject the hypothesis \mathbf{H}_0 if $T_n(X_1, \dots, X_n) > c_{n,B}^*$.

This parametric bootstrap procedure was used for all of the following procedures to generate the critical points. We consider the test of Baringhaus and Henze (1992) based on the statistic

$$\text{BH} = \frac{1}{n} \sum_{i,j=1}^n \left(\frac{\hat{\lambda}_n^2}{X_i + X_j + 1} + \frac{X_i X_j}{X_i + X_j - 1} \right) - \hat{\lambda}_n \left(n - \frac{1}{n} \left(\sum_{j=1}^n \mathbb{1}\{X_j = 0\} \right)^2 \right).$$

The mean distance test by Székely and Rizzo (2004) is based on

$$\text{SR} = n \sum_{j=0}^{\infty} \left(\widehat{M}_n(j) - P(j; \hat{\lambda}_n) \right)^2 P(j; \hat{\lambda}_n),$$

where $\widehat{M}_n(j)$ is an estimator of the CDF based on the mean distance and $P(j; \lambda)$ (resp. $p(j; \lambda)$) denote the distribution function (resp. pmf) of $\text{Po}(\lambda)$, respectively. Note that SR is implemented in the R-package *energy*, see Rizzo and Szekely (2019). The test of Rueda et al. (1991) is based on

$$\text{RU} = \frac{1}{n} \sum_{i,j=1}^n \frac{1}{X_i + X_j + 1} + \frac{n \cdot (1 - e^{-2\widehat{\lambda}_n})}{2\widehat{\lambda}_n} - 2 \sum_{i=1}^n \left(\frac{(-1)^{X_i} \cdot X_i! \cdot (1 - e^{-\widehat{\lambda}_n})}{\widehat{\lambda}_n^{X_i+1}} + \sum_{j=1}^{X_i} \frac{(-1)^{j+1} \cdot X_i!}{(X_i - j + 1)! \cdot \widehat{\lambda}_n^j} \right).$$

Note that in the original paper of Rueda et al. (1991) and in a slight handwritten correction thereof available on the internet, as well as in the work of Gürtler and Henze (2000), the explicit formula of the RU-statistic contains errors. We have corrected and numerically checked the formula given above against the integral representation used to introduce the test. The integrated distribution function based tests of Klar (1999) are defined via

$$K_1 = \sqrt{n} \left(\sum_{j=0}^M \left| \widehat{F}_n(j) - P(j; \widehat{\lambda}_n) \right| + \widehat{\lambda}_n - \sum_{j=0}^M \left(1 - P(j; \widehat{\lambda}_n) \right) \right)$$

and

$$K_2 = \sqrt{n} \sup_{1 \leq k \leq M} \left| \sum_{j=0}^{k-1} \left(\widehat{F}_n(j) - P(j; \widehat{\lambda}_n) \right) \right|,$$

where $M = \max\{X_1, \dots, X_n\}$ and \widehat{F}_n is the empirical distribution function of X_1, \dots, X_n . For representations of the Kolmogorov-Smirnov statistic and the modified Cramér-von Mises statistic, we follow the representation given by Gürtler and Henze (2000), namely

$$\text{KS} = \sqrt{n} \sup_{0 \leq k \leq M} \left| \widehat{F}_n(x) - P(k; \widehat{\lambda}_n) \right|$$

and

$$\text{CM} = n \sum_{j=0}^M \left(\widehat{F}_n(j) - P(j; \widehat{\lambda}_n) \right)^2 \cdot \frac{1}{n} \sum_{k=1}^n \mathbb{1}\{X_k = j\}.$$

The simulation study consists of the following 45 representatives of families of distributions. In order to show that all the considered testing procedures maintain the nominal level α of 5%, we consider the $\text{Po}(\lambda)$ distribution with $\lambda \in \{1, 5, 10, 30\}$. As examples for alternative distributions, we consider the discrete uniform distribution $\mathcal{U}\{0, 1, \dots, m\}$ with $m \in \{1, 2, 3, 5, 6\}$, several different instances of the binomial distribution $\text{Bin}(m, q)$, several Poisson mixtures of the form $\text{PP}(q; \vartheta_1, \vartheta_2) = q \cdot \text{Po}(\vartheta_1) + (1 - q) \cdot \text{Po}(\vartheta_2)$, a 0.9/0.1 mixture of $\text{Po}(3)$ and point mass in 0 denoted by $\text{Po}(3)\delta_0$, discrete Weibull distributions $\text{W}(\vartheta_1, \vartheta_2)$, zero-modified Poisson distributions $\text{zmPo}(\lambda, q)$, the zero-truncated Poisson distributions $\text{ztPo}(\lambda)$ with $\lambda \in \{2, 3, 5\}$, and the absolute discrete normal distribution $|\text{N}(\mu, 1)|$ with $\mu \in \{0, 2, 3\}$. Note that most distributions were generated by the packages *extraDistr*, see Wolodzko (2019), and *actuar*, see Dutang et al. (2008), and that a significant part of these distributions can also be found in the simulation study presented by Gürtler and Henze (2000). Furthermore we indicate that the chosen design of simulation parameters coincides with the study by Gürtler and Henze (2000) which facilitates the comparison to other tests of poissonity not considered here.

Every entry in Table 1 is based on 100000 repetitions and 500 bootstrap samples of size 50. All of the considered procedures maintain the significance level $\alpha = 5\%$ under the hypothesis, which supports the statement that the parametric bootstrap procedure is well calibrated. Overall the best performing tests are K_2 , BH and SR. The new test based on T_n^{Po} is competitive to the stated procedures, although it never outperforms them all at once for the considered alternatives.

Distr. / Test	T_n^{Po}	BH	SR	RU	K ₁	K ₂	KS	CM
Po(1)	5	5	5	5	5	5	5	5
Po(5)	5	5	5	5	5	5	5	5
Po(10)	5	5	5	5	5	5	5	5
Po(30)	5	5	5	5	5	5	5	5
$\mathcal{U}\{0, 1\}$	99	99	99	99	99	99	99	99
$\mathcal{U}\{0, 1, 2\}$	39	9	22	15	64	68	50	58
$\mathcal{U}\{0, 1, 2, 3\}$	46	33	20	27	61	16	45	51
$\mathcal{U}\{0, 1, 2, 3, 4, 5\}$	69	65	58	62	75	39	60	63
$\mathcal{U}\{0, 1, 2, 3, 4, 5, 6\}$	85	85	83	85	86	66	72	76
Bin(2, 0.5)	81	81	89	87	86	90	83	81
Bin(4, 0.25)	18	22	23	24	18	22	21	15
Bin(10, 0.1)	7	7	7	7	6	7	7	6
Bin(10, 0.5)	57	52	49	52	82	88	60	68
Bin(1, 0.5)	73	77	82	81	80	82	76	77
Bin(2, 2/3)	34	38	44	44	43	45	37	39
Bin(3, 0.75)	19	22	26	26	26	27	21	23
Bin(9, 0.9)	6	7	8	8	8	8	7	8
Bin(5, 0.5)	82	85	80	84	88	89	67	71
Bin(10, 2/3)	41	45	41	44	48	50	28	31
Bin(15, 0.75)	23	27	26	26	28	30	16	18
Bin(45, 0.9)	8	9	10	9	9	8	6	7
PP(0.5; 2, 5)	64	64	69	65	72	74	53	57
PP(0.5; 3, 5)	17	19	20	19	20	21	12	14
PP(0.25; 1, 5)	93	95	96	95	87	88	75	73
PP(0.05; 1, 5)	23	33	32	32	13	12	8	7
PP(0.01; 1, 5)	7	9	9	9	6	5	5	5
Po(3) δ_0	54	62	54	59	32	31	32	26
W(0.5, 1)	73	77	82	81	80	82	76	77
W(0.25, 1)	22	24	27	28	26	26	26	25
W(0.5, 2)	49	52	52	51	48	52	51	52
W(0.25, 2)	8	8	7	6	6	8	7	10
W(0.75, 2)	28	32	35	35	26	32	30	21
W(0.1, 1)	10	10	10	8	10	10	10	10
W(0.9, 3)	97	97	99	99	98	99	97	93
zmPo(1, 0.1)	91	93	90	92	81	84	90	64
zmPo(1, 0.5)	17	18	19	19	19	18	18	19
zmPo(1, 0.8)	71	72	74	74	74	74	74	74
zmPo(2, 0.1)	8	9	8	8	6	6	6	6
zmPo(3, 0.1)	24	30	24	27	13	12	11	9
ztPo(2)	93	99	83	95	38	39	56	19
ztPo(3)	12	18	18	18	9	10	7	9
ztPo(5)	4	1	1	1	5	5	5	5
N(0, 1)	45	48	48	50	42	46	47	41
N(2, 1)	44	46	59	53	54	61	42	37
N(3, 1)	88	78	94	86	96	98	85	90

 Table 1: Empirical rejection rates of the tests of poissonity (sample size $n = 50$, significance level $\alpha = 0.05$).

7 Parameter estimation in the family of negative binomial distributions

The characterizations we employ contain information about the underlying probability law and lead to empirical discrepancy measures being close to zero if the distribution generating the data is the one stated in the characterization. These measures can be used for estimation of the parameters of the considered parametric family of distributions. To illustrate this point, we propose a minimum distance estimation procedure for the family of negative binomial distributions. Our objective is to estimate the unknown parameters $q \in (0, 1)$ and $r > 0$ of a negative binomial distribution based on an i.i.d. \mathbb{N}_0 -valued sample X_1, \dots, X_n . Estimation in this particular family is not trivial, since Aragón et al. (1992) have shown the conjecture of Anscombe dating back to 1950, that the maximum likelihood equations have a unique solution if, and only if, $\bar{X}_n < S_n^2$ with $\bar{X}_n = n^{-1} \sum_{j=1}^n X_j$, the sample mean, and $S_n^2 = n^{-1} \sum_{j=1}^n (X_j - \bar{X}_n)^2$, the sample variance. However, as Johnson et al. (1993) state in their Section 8.3, so called "[...] underdispersed samples [...] will occasionally be encountered, even when a negative binomial model is appropriate." The moment estimators defined by $\tilde{q}_n = \bar{X}_n / S_n^2$ and

$$\tilde{r}_n = \frac{(\bar{X}_n)^2}{(1 - \tilde{q}_n) S_n^2} = \frac{(\bar{X}_n)^2}{S_n^2 - \bar{X}_n},$$

see display (5.49) and (5.50) of Johnson et al. (1993), perform comparably bad as the maximum likelihood estimators, since, in underdispersed samples, they lead to negative values of \tilde{r}_n or values of \tilde{q}_n that are greater than one, see the following simulation study and Figure 2.

The heuristic for our new method is similar to that of the previous section, again based on Theorem 3.1 (see also Example 5.2). Thus, we define

$$\hat{e}_n(k; r, q) = \frac{1}{n} \sum_{j=1}^n \left(1 - \frac{r + X_j}{X_j + 1} (1 - q) \right) \mathbb{1}\{X_j \geq k\}, \quad k \in \mathbb{N}_0,$$

and let $\hat{\rho}_n$ be as in the previous section. Similar to the test for the Poisson distribution, we consider the empirical discrepancy measure

$$\begin{aligned} S_n^{NB}(r, q) &= \sum_{k=0}^{\infty} (\hat{e}_n(k; r, q) - \hat{\rho}_n(k))^2 \\ &= \frac{1}{n^2} \sum_{j, \ell=1}^n \left[\left(1 - \frac{r + X_j}{X_j + 1} (1 - q) \right) \left(q(r + X_\ell) - r - 1 \right) \mathbb{1}\{X_j \geq X_\ell\} \right. \\ &\quad \left. + \left(q(r + X_j) - r + 1 \right) \left(1 - \frac{r + X_\ell}{X_\ell + 1} (1 - q) \right) \mathbb{1}\{X_j < X_\ell\} + \mathbb{1}\{X_j = X_\ell\} \right], \end{aligned}$$

where the proposed estimators for (r, q) are defined by

$$(\hat{r}_n, \hat{q}_n) = \operatorname{argmin}_{(r, q)} S_n^{NB}(r, q). \quad (3)$$

In this particular example we expect that it is possible to minimize the quadratic equation in r and q explicitly to obtain formulae for the estimators. However, in the following section we cannot hope for an explicit solution to the optimization problem and for reasons of consistency of the presentation, we use a numerical routine to find the values of the estimators in both cases. Note that similar estimators for parametric families of continuous distributions are investigated by Betsch et al. (2021).

For a comparison of the two presented methods we conduct a simulation study in R and use the `optim` routine to find the minimal values in (3). The option `method` was fixed to `L-BFGS-B`, thus choosing an implementation of the routine suggested by Byrd et al. (1995), and the maximum number of iterations to `maxit=1000`. As starting values for the optimization routine we choose independent uniformly distributed random numbers $r_s \sim \mathcal{U}(1, 3)$ and $q_s \sim \mathcal{U}(0.1, 0.9)$. For different $(r_0, q_0) \in (0, \infty) \times (0, 1)$ we simulate 200 i.i.d. samples of size $n = 100$ from a negative binomial distribution with parameters (r_0, q_0) and calculate the minimum distance estimators (\hat{r}_n, \hat{q}_n) as well as the moment estimators $(\tilde{r}_n, \tilde{q}_n)$. Then the bias of the estimation is derived by subtracting the underlying 'true' parameters (r_0, q_0) . In Figures 1 and 2 the results of the different simulations are plotted, with estimation results of the moment estimators plotted as black crosses and the results of the minimum distance estimators as red circles. It is visible in Figure 1 that for small values of q_0 , both procedures perform comparably, although the values of the moments estimators seem to scatter a little more than those of the minimum distance estimators. A completely different picture is seen in Figure 2, where values of q_0 in the neighborhood of 1 and greater values of r_0 are assumed. The moment estimators $(\tilde{r}_n, \tilde{q}_n)$ regularly

produce values which are clearly outside of the defined parameter space $(0, \infty) \times (0, 1)$ as opposed to the minimum distance estimators (\hat{r}_n, \hat{q}_n) which do not show this behavior due to the optimization constraints. Nevertheless, some convergence failures in the optimization routine did occur and they are not exclusively related to the underdispersed samples and only happen for somewhat extreme parameter configurations. We chose to visually assess the quality of estimation, since empirical versions of the bias and mean squared error are very sensitive to big discrepancies, and hence did not provide valuable information on the quality of the estimation procedures. It would be of interest to find theoretical statements for the estimators (\hat{r}_n, \hat{q}_n) such as consistency results or a central limit theorem type asymptotic distribution.




Figure 1: Biases of the minimum distance (red) and method of moments (black) estimation procedures simulated for different parameters, $(r, q) \in \{2, 5, 10\} \times \{0.05, 0.25, 0.5\}$, of the negative binomial distribution with sample size $n = 100$ and 200 repetitions of the simulations. A red circle represents the value of $(\hat{r}_n - r_0, \hat{q}_n - q_0)$, while a black cross stands for $(\hat{r}_n - r_0, \hat{q}_n - q_0)$. The assumed true value (r_0, q_0) in every subfigure is highlighted by the intersection of the dotted lines.




Figure 2: Biases of the minimum distance (red) and method of moments (black) estimation procedures simulated for different parameters, $(r, q) \in \{20, 30, 50\} \times \{0.8, 0.9, 0.95\}$, of the negative binomial distribution with sample size $n = 100$ and 200 repetitions of the simulations. A red circle represents the value of $(\hat{r}_n - r_0, \hat{q}_n - q_0)$, while a black cross stands for $(\tilde{r}_n - r_0, \tilde{q}_n - q_0)$. The assumed true value (r_0, q_0) in every subfigure is highlighted by the intersection of the dotted lines. The number of cases of underdispersed samples, the number of convergence failures of the optimization routine, and the number of cases where both occurred are stated in the plot.

8 Parameter estimation in discrete exponential-polynomial models

In this final section we present an application to a non-normalized model, namely parameter estimation in the discrete exponential-polynomial models introduced in Example 5.3. We follow Betsch et al. (2021) who apply the continuous version of our estimation method to continuous exponential-polynomial models. In their work, they compare the method with two other methods for parameter estimation in non-normalized continuous models. More specifically, they implemented the score matching approach of Hyvärinen (2007) as well as noise-contrastive estimators from Gutmann and Hyvärinen (2012). As another contribution that focuses on the continuous exponential-polynomial distribution and the corresponding parameter estimation problem, let us mention Hayakawa and Takemura (2016). In our search through the literature we have found only few methods for the parameter estimation in the discrete version of the model. As such, contrastive divergence methods based on the initial proposal by Hinton (2002) can be applied in principle though it does not avoid dealing with the normalization constant $C(\vartheta)$ and Lyu (2009) proposes a discrete version of the score matching approach but does not give details on its implementation. More recently Takenouchi and Kanamori (2017) proposed a method (that avoids any calculation or approximation of the normalization constant) based on suitable homogeneous divergences which are empirically localized. We use this latter method as a comparison to our approach.

Assume that X_1, \dots, X_n is an i.i.d. \mathbb{N} -valued sample from the exponential-polynomial model $p_{\vartheta^{(0)}}$ in Example 5.3 with some unknown parameter $\vartheta^{(0)} \in \mathbb{R}^{d-1} \times (-\infty, 0)$ (with $d \in \mathbb{N}$, $d \geq 2$, fixed and known). We seek to estimate $\vartheta^{(0)}$ based on X_1, \dots, X_n . Very similar to the previous section, we consider $\hat{\rho}_n$ as before, and put

$$\hat{e}_n(k; \vartheta) = \frac{1}{n} \sum_{j=1}^n \left(1 - \exp \left(\vartheta_1 + \vartheta_2((X_j + 1)^2 - X_j^2) + \dots + \vartheta_d((X_j + 1)^d - X_j^d) \right) \right) \mathbb{1}\{X_j \geq k\}, \quad k \in \mathbb{N}.$$

We define the empirical discrepancy measure

$$S_n^{PE}(\vartheta) = \sum_{k=0}^{\infty} (\hat{e}_n(k; \vartheta) - \hat{\rho}_n(k))^2.$$

In line with Theorem 3.1, or more precisely, Example 5.3, we propose as an estimator

$$\hat{\vartheta}_n = \operatorname{argmin}_{\vartheta} S_n^{PE}(\vartheta). \quad (4)$$

To see if this approach leads to sensible estimators, we conduct simulations in a two-parameter special case of the model. Following the continuous-case simulation setting of Betsch et al. (2021), we consider $d = 3$ but fix $\vartheta_2 = 0$, thus effectively estimating the parameters of the parametric family given through

$$p_{(\vartheta_1, \vartheta_3)}(k) = C(\vartheta_1, \vartheta_3)^{-1} \exp(\vartheta_1 k + \vartheta_3 k^3), \quad k \in \mathbb{N}, \quad \vartheta_1 \in \mathbb{R}, \quad \vartheta_3 < 0, \quad (5)$$

which, though simpler than the general case, is still a non-normalized model and thus inaccessible to explicit maximum likelihood estimation. The discrepancy measure $S_n^{PE}(\vartheta)$ can be calculated as

$$\begin{aligned} S_n^{PE}(\vartheta) = \frac{1}{n^2} \sum_{j, \ell=1}^n & \left[(E_j(\vartheta_1, \vartheta_3) - 1)(E_{\ell}(\vartheta_1, \vartheta_3) \cdot X_{\ell} - X_{\ell} + 2) \mathbb{1}\{X_j \geq X_{\ell}\} + \mathbb{1}\{X_j = X_{\ell}\} \right. \\ & \left. + (E_j(\vartheta_1, \vartheta_3) - 1)(E_{\ell}(\vartheta_1, \vartheta_3) - 1) \cdot X_j \cdot \mathbb{1}\{X_j < X_{\ell}\} \right], \end{aligned}$$

where

$$E_i(\vartheta_1, \vartheta_3) = \exp(\vartheta_1 + \vartheta_3 + 3\vartheta_3 X_i + 3\vartheta_3 X_i^2), \quad i = 1, \dots, n.$$

For a comparison, we consider the estimator proposed by Takenouchi and Kanamori (2017). For positive constants $\alpha, \alpha', \gamma > 0$, with $\alpha > \alpha'$, and $\bar{\alpha} = (\alpha + \gamma\alpha') / (1 + \gamma)$, their estimator is given as

$$\begin{aligned} \tilde{\vartheta}_n = \operatorname{argmin}_{\vartheta} & \left\{ \frac{1}{1 + \gamma} \log \left(\sum_{k \in \mathcal{Z}} \left(\frac{n_k}{n} \right)^{\alpha} q_{\vartheta}(k)^{1-\alpha} \right) + \frac{\gamma}{1 + \gamma} \log \left(\sum_{k \in \mathcal{Z}} \left(\frac{n_k}{n} \right)^{\alpha'} q_{\vartheta}(k)^{1-\alpha'} \right) \right. \\ & \left. - \log \left(\sum_{k \in \mathcal{Z}} \left(\frac{n_k}{n} \right)^{\bar{\alpha}} q_{\vartheta}(k)^{1-\bar{\alpha}} \right) \right\}, \end{aligned} \quad (6)$$

where \mathcal{Z} is the set of all values that appear in the sample X_1, \dots, X_n , the variable n_k denotes how often the value k is found in the sample, and

$$q_{\vartheta}(k) = \exp(\vartheta_1 k + \dots + \vartheta_d k^d), \quad k \in \mathbb{N}.$$




Figure 3: Simulated biases of the estimators $\hat{\vartheta}_n$ (red) and $\tilde{\vartheta}_n$ (black) in the discrete exponential-polynomial model for different parameters $(\vartheta_1^{(0)}, \vartheta_3^{(0)})$ ($n = 100; 200$ repetitions). A red circle represents the value of $(\hat{\vartheta}_{n,1} - \vartheta_1^{(0)}, \hat{\vartheta}_{n,3} - \vartheta_3^{(0)})$, while a black cross stands for $(\tilde{\vartheta}_{n,1} - \vartheta_1^{(0)}, \tilde{\vartheta}_{n,3} - \vartheta_3^{(0)})$. The assumed true value in every subfigure is highlighted by the intersection of the dotted lines.

Since Takenouchi and Kanamori (2017) do not propose a specific way of choosing the constants α , α' and γ , we use the values that appear most frequently in their simulation study and therefore set $\alpha = 1.1$, $\alpha' = 0.1$ and $\gamma = 1/9$.

As in the previous section, we use the software R for the simulation and the `optim` routine to find the minimal values in (4) and (6). Again, the option method is fixed to L-BFGS-B and the maximum number of iterations to `maxit=1000`. As starting values for the optimization we choose independent uniformly distributed random numbers $\vartheta_1^{(s)} \sim \mathcal{U}(-1, 1)$ and $\vartheta_3^{(s)} \sim \mathcal{U}(-1, 0)$. For different $(\vartheta_1^{(0)}, \vartheta_3^{(0)}) \in \mathbb{R} \times (-\infty, 0)$, we simulate 200 i.i.d. samples of size $n = 100$ from the discrete exponential-polynomial model in (5) with parameters $(\vartheta_1^{(0)}, \vartheta_3^{(0)})$ and calculate the estimators $(\hat{\vartheta}_{n,1}, \hat{\vartheta}_{n,3})$ and $(\tilde{\vartheta}_{n,1}, \tilde{\vartheta}_{n,3})$ presented in (4) and (6) respectively. The biases of the estimators are given by

subtracting the underlying 'true' parameters $(\vartheta_1^{(0)}, \vartheta_3^{(0)})$. The simulation of a discrete exponential-polynomial model is rather simple as $\vartheta_3^{(0)} < 0$ ensures that the probability $p_{(\vartheta_1^{(0)}, \vartheta_3^{(0)})}(k)$ is rapidly decreasing as k grows. From a practical point of view and minding the usual calculation accuracy, we only need to deal with a discrete distribution with finite support. In Figure 3 the results of the simulations are presented and it is visible that both procedures perform comparable and overall well. The newly proposed estimators tend to scatter more which favors the competing estimators. However, our new estimators require no (data-dependent or quick fix) choice of parameters (like α , α' and γ for the estimators of Takenouchi and Kanamori, 2017). Introducing additional parameters, for instance through suitable weight functions, is also conceivable for our method. It would certainly allow for some choice which improves the overall performance, but it also leads to a less intuitive implementation as these parameters need to be chosen in practice. Note that, as in the previous simulation, some convergence failures in the optimization routine occurred (less than ten percent per parameter configuration).

Appendix

A Proof of Theorem 2.2

The following proof is, up to technical details involving the class of test functions, due to Ley and Swan (2013a) and given here for the reader's convenience.

Assume that $X|p \sim p$. Then, for $f \in \mathcal{F}_p$,

$$\begin{aligned} \mathbb{E} \left[\Delta^+ f(X) + \frac{\Delta^+ p(X)}{p(X)} f(X+1) \mid X \in \text{spt}(p) \right] &= \sum_{k=L}^R \left(p(k) \Delta^+ f(k) + f(k+1) \Delta^+ p(k) \right) \\ &= \sum_{k=L}^R \Delta^+ (p(k) f(k)) = 0, \end{aligned}$$

using assumption (a). To prove the converse, take a discrete random variable Z with mass function p , independent of X . For $m \in \mathbb{Z}$, define $f_m : \{L, \dots, R\} \rightarrow \mathbb{R}$ via

$$f_m(k) = \frac{1}{p(k)} \sum_{\ell=L}^{k-1} (\mathbb{1}\{\ell \leq m\} - \mathbb{P}(Z \leq m)) p(\ell)$$

which satisfies

$$\Delta^+ (p(k) f_m(k)) = (\mathbb{1}\{k \leq m\} - \mathbb{P}(Z \leq m)) p(k), \quad k \in \{L, \dots, R\}.$$

Therefore,

$$\sum_{k=L}^R \left| \Delta^+ (p(k) f_m(k)) \right| \leq 2 \sum_{k=L}^R p(k) = 2 < \infty,$$

as well as

$$\sum_{k=L}^R \Delta^+ (p(k) f_m(k)) = \sum_{k=L}^R (\mathbb{1}\{k \leq m\} - \mathbb{P}(Z \leq m)) p(k) = \mathbb{P}(Z \leq m) - \mathbb{P}(Z \leq m) = 0.$$

Moreover, we have $f_m(k+1) \leq 2 \cdot P(k) / p(k+1)$ and

$$\begin{aligned} |f_m(k+1)| &= \frac{1}{p(k+1)} \left| \sum_{\ell=L}^R (\mathbb{1}\{\ell \leq m\} - \mathbb{P}(Z \leq m)) p(\ell) - \sum_{\ell=k+1}^R (\mathbb{1}\{\ell \leq m\} - \mathbb{P}(Z \leq m)) p(\ell) \right| \\ &= \frac{1}{p(k+1)} \left| \sum_{\ell=k+1}^R (\mathbb{1}\{\ell \leq m\} - \mathbb{P}(Z \leq m)) p(\ell) \right| \\ &\leq \frac{2}{p(k+1)} \cdot (1 - P(k)), \end{aligned}$$

and thus get

$$\sup_{k \in \{L, \dots, R\}} \left| \frac{\Delta^+ p(k)}{p(k)} f_m(k+1) \right| \leq 2 \cdot \sup_{k \in \{L, \dots, R-1\}} \left| \frac{\Delta^+ p(k) \cdot \min\{P(k), 1 - P(k)\}}{p(k)p(k+1)} \right| < \infty.$$

Now, notice that

$$\begin{aligned} \Delta^+ f_m(k) &= \left(\frac{1}{p(k+1)} - \frac{1}{p(k)} \right) \sum_{\ell=L}^k (\mathbb{1}\{\ell \leq m\} - \mathbb{P}(Z \leq m)) p(\ell) + \frac{1}{p(k)} (\mathbb{1}\{k \leq m\} - \mathbb{P}(Z \leq m)) p(k) \\ &= \left(1 - \frac{p(k+1)}{p(k)} \right) \frac{1}{p(k+1)} \sum_{\ell=L}^k (\mathbb{1}\{\ell \leq m\} - \mathbb{P}(Z \leq m)) p(\ell) + \mathbb{1}\{k \leq m\} - \mathbb{P}(Z \leq m) \\ &= -\frac{\Delta^+ p(k)}{p(k)} f_m(k+1) + \mathbb{1}\{k \leq m\} - \mathbb{P}(Z \leq m), \end{aligned}$$

where the calculation is valid for $k \in \{L, \dots, R-1\}$, but the equality obviously also holds for $k = R$ [using our convention $f_m(R+1) = 0$] if $R < \infty$. From this relation, we immediately get that

$$\sup_{k \in \{L, \dots, R\}} |\Delta^+ f_m(k)| < \infty,$$

so $f_m \in \mathcal{F}_p$, as well as, by the assumption in the converse implication,

$$0 = \mathbb{E} \left[\Delta^+ f(X) + \frac{\Delta^+ p(X)}{p(X)} f(X+1) \mid X \in \text{spt}(p) \right] = \mathbb{E} [\mathbb{1}\{X \leq m\} - \mathbb{P}(Z \leq m) \mid X \in \text{spt}(p)],$$

which implies the claim.

B Proof of Theorem 3.1

First assume that $X|p \sim p$. Then, for all $k \in \mathbb{Z}, k \geq L$, we have

$$\rho_{X|p}(k) = p(k) = - \sum_{\ell=k}^R (p(\ell+1) - p(\ell)) = - \sum_{\ell=k}^R \frac{\Delta^+ p(\ell)}{p(\ell)} p(\ell) = \mathbb{E} \left[- \frac{\Delta^+ p(X)}{p(X)} \mathbb{1}\{X \geq k\} \mid X \in \text{spt}(p) \right].$$

For the converse implication, assume that

$$\rho_{X|p}(k) = \mathbb{E} \left[- \frac{\Delta^+ p(X)}{p(X)} \mathbb{1}\{X \geq k\} \mid X \in \text{spt}(p) \right], \quad k \in \mathbb{Z}, k \geq L.$$

We obtain for $f \in \mathcal{F}_p$

$$\begin{aligned} \mathbb{E} [\Delta^+ f(X) \mid X \in \text{spt}(p)] &= \sum_{\ell=L}^R (\Delta^+ f(\ell)) \rho_{X|p}(\ell) = \sum_{\ell=L}^R \Delta^+ f(\ell) \cdot \mathbb{E} \left[- \frac{\Delta^+ p(X)}{p(X)} \mathbb{1}\{X \geq \ell\} \mid X \in \text{spt}(p) \right] \\ &= \mathbb{E} \left[- \frac{\Delta^+ p(X)}{p(X)} \sum_{\ell=L}^X (f(\ell+1) - f(\ell)) \mid X \in \text{spt}(p) \right] \\ &= \mathbb{E} \left[- \frac{\Delta^+ p(X)}{p(X)} f(X+1) \mid X \in \text{spt}(p) \right], \end{aligned}$$

where we use that $f(L) = 0$ by Remark 2.3, and where Fubini's theorem is applicable as

$$\begin{aligned} \sum_{\ell=L}^R \mathbb{E} \left| \Delta^+ f(\ell) \frac{\Delta^+ p(X)}{p(X)} \mathbb{1}\{X \geq \ell, X \in \text{spt}(p)\} \right| &\leq \sup_{k \in \{L, \dots, R\}} |\Delta^+ f(k)| \cdot \mathbb{E} \left| \frac{\Delta^+ p(X)}{p(X)} (X - L) \cdot \mathbb{1}\{X \in \text{spt}(p)\} \right| \\ &< \infty. \end{aligned}$$

Theorem 2.2 implies the claim.

C Proof of Proposition 4.1

If $X|p \sim p$, then Theorem 3.1 implies that, for $k \in \mathbb{Z}, k \geq L$,

$$\begin{aligned} F_{X|p}(k) &= \sum_{\ell=L}^k \rho_{X|p}(\ell) = \mathbb{E} \left[-\frac{\Delta^+ p(X)}{p(X)} \sum_{\ell=L}^k \mathbb{1}\{X \geq \ell\} \mid X \in \text{spt}(p) \right] \\ &= \mathbb{E} \left[-\frac{\Delta^+ p(X)}{p(X)} (\min\{X, k\} - L + 1) \mid X \in \text{spt}(p) \right]. \end{aligned}$$

To prove the converse, assume that

$$F_{X|p}(k) = \mathbb{E} \left[-\frac{\Delta^+ p(X)}{p(X)} (\min\{X, k\} - L + 1) \mid X \in \text{spt}(p) \right], \quad k \in \mathbb{Z}, k \geq L.$$

Then, we have

$$\begin{aligned} \rho_{X|p}(k) &= F_{X|p}(k) - F_{X|p}(k-1) = \mathbb{E} \left[-\frac{\Delta^+ p(X)}{p(X)} \left(\sum_{\ell=L}^k \mathbb{1}\{X \geq \ell\} - \sum_{\ell=L}^{k-1} \mathbb{1}\{X \geq \ell\} \right) \mid X \in \text{spt}(p) \right] \\ &= \mathbb{E} \left[-\frac{\Delta^+ p(X)}{p(X)} \mathbb{1}\{X \geq k\} \mid X \in \text{spt}(p) \right], \quad k \in \mathbb{Z}, k > L, \end{aligned}$$

and

$$\rho_{X|p}(L) = F_{X|p}(L) = \mathbb{E} \left[-\frac{\Delta^+ p(X)}{p(X)} \mathbb{1}\{X \geq L\} \mid X \in \text{spt}(p) \right],$$

so Theorem 3.1 yields $X|p \sim p$.

D Proof of Proposition 4.2

First, if $X|p \sim p$, Theorem 3.1 gives

$$\begin{aligned} \varphi_{X|p}(t) &= \sum_{\ell=L}^R e^{it\ell} \rho_{X|p}(\ell) = \mathbb{E} \left[-\frac{\Delta^+ p(X)}{p(X)} \sum_{\ell=L}^X e^{it\ell} \mid X \in \text{spt}(p) \right] \\ &= \mathbb{E} \left[-\frac{\Delta^+ p(X)}{p(X)} \cdot \frac{e^{itL} - e^{it(X+1)}}{1 - e^{it}} \mid X \in \text{spt}(p) \right], \quad t \in \mathbb{R}, \end{aligned}$$

where the use of Fubini's theorem (in case $R = \infty$) is admissible by the argument from Appendix B, and where we use the complex geometric sum in the last step. For the converse, assume that $\varphi_{X|p}$ is given through the stated formula. The inversion formula for characteristic functions applied to the atoms of $X|X \in \text{spt}(p)$ (see Theorem 2.3.2 of Cuppens, 1975) gives, for $k \in \mathbb{Z}, k \geq L$,

$$\begin{aligned} \rho_{X|p}(k) &= \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T e^{-itk} \varphi_{X|p}(t) dt \\ &= \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T e^{-itk} \mathbb{E} \left[-\frac{\Delta^+ p(X)}{p(X)} \cdot \frac{e^{itL} - e^{it(X+1)}}{1 - e^{it}} \mid X \in \text{spt}(p) \right] dt \\ &= \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T e^{-itk} \mathbb{E} \left[-\frac{\Delta^+ p(X)}{p(X)} \sum_{\ell=L}^X e^{it\ell} \mid X \in \text{spt}(p) \right] dt \\ &= \lim_{T \rightarrow \infty} \frac{1}{2T} \mathbb{E} \left[-\frac{\Delta^+ p(X)}{p(X)} \sum_{\ell=L}^X \int_{-T}^T e^{-itk} e^{it\ell} dt \mid X \in \text{spt}(p) \right] \\ &= \lim_{T \rightarrow \infty} \frac{1}{2T} \mathbb{E} \left[-\frac{\Delta^+ p(X)}{p(X)} \sum_{\ell=L}^X \left(\frac{e^{iT(\ell-k)} - e^{-iT(\ell-k)}}{i(\ell-k)} \mathbb{1}\{\ell \neq k\} + 2T \mathbb{1}\{\ell = k\} \right) \mid X \in \text{spt}(p) \right] \\ &= \lim_{T \rightarrow \infty} \mathbb{E} \left[-\frac{\Delta^+ p(X)}{p(X)} \sum_{\ell=L}^X \left(\frac{\sin(T(\ell-k))}{T(\ell-k)} \mathbb{1}\{\ell \neq k\} + \mathbb{1}\{\ell = k\} \right) \mid X \in \text{spt}(p) \right] \end{aligned}$$

$$\begin{aligned}
 &= \mathbb{E} \left[- \frac{\Delta^+ p(X)}{p(X)} \sum_{\ell=L}^X \mathbb{1}\{\ell = k\} \mid X \in \text{spt}(p) \right] \\
 &= \mathbb{E} \left[- \frac{\Delta^+ p(X)}{p(X)} \mathbb{1}\{X \geq k\} \mid X \in \text{spt}(p) \right],
 \end{aligned}$$

where the use of dominated convergence in the second to last step is easily justified (if necessary) by the integrability assumptions on X . Theorem 3.1 yields the claim.

E Proof of Proposition 4.3

The necessity part follows from Theorem 3.1 with a calculation similar to Appendix D. For the sufficiency part, assume that G_X is given through the stated formula. Then, for all $s \in [0, 1)$, we have

$$G_X(s) = \mathbb{E} \left[- \frac{\Delta^+ p(X)}{p(X)} \cdot \frac{1 - s^{X+1}}{1 - s} \right] = \sum_{\ell=0}^{\infty} s^{\ell} \mathbb{E} \left[- \frac{\Delta^+ p(X)}{p(X)} \mathbb{1}\{X \geq \ell\} \right].$$

Since $G_X(s) = \sum_{\ell=0}^{\infty} s^{\ell} \rho_X(\ell) \leq 1$, for $s \in [0, 1)$, G_X is a convergent power series in $[0, 1)$. We can thus differentiate G_X in $(0, 1)$ to obtain

$$\frac{d^k}{ds^k} G_X(s) = \sum_{\ell=k}^{\infty} \ell(\ell-1) \cdots (\ell-k+1) s^{\ell-k} \rho_X(\ell)$$

as well as

$$\frac{d^k}{ds^k} G_X(s) = \sum_{\ell=k}^{\infty} \ell(\ell-1) \cdots (\ell-k+1) s^{\ell-k} \mathbb{E} \left[- \frac{\Delta^+ p(X)}{p(X)} \mathbb{1}\{X \geq \ell\} \right].$$

We conclude that

$$k! \cdot \mathbb{E} \left[- \frac{\Delta^+ p(X)}{p(X)} \mathbb{1}\{X \geq k\} \right] = \frac{d^k}{ds^k} G_X(s) \Big|_{s=0} = k! \cdot \rho_X(k)$$

so Theorem 3.1 implies the claim.

References

- Allison, J. S., Betsch, S., Ebner, B., and Visagie, I. J. H. (2019). New weighted L^2 -type tests for the inverse Gaussian distribution. *ArXiv e-prints*, 1910.14119.
- Aragón, J., Eberly, D., and Eberly, S. (1992). Existence and uniqueness of the maximum likelihood estimator for the two-parameter negative binomial distribution. *Statistics & Probability Letters*, 15(5):375–379.
- Arras, B. and Houdré, C. (2019). *On Stein’s method for infinitely divisible laws with finite first moment*. SpringerBriefs in Probability and Mathematical Statistics. Springer International Publishing, Cham.
- Barbour, A. D. (1988). Stein’s method and Poisson process convergence. *Journal of Applied Probability*, 25(A):175–184.
- Barbour, A. D. (1990). Stein’s method for diffusion approximations. *Probability Theory and Related Fields*, 84(3):297–322.
- Baringhaus, L. and Henze, N. (1992). A goodness of fit test for the Poisson distribution based on the empirical generating function. *Statistics & Probability Letters*, 13(4):269–274.
- Barp, A., Briol, F.-X., Duncan, A. B., Girolami, M. A., and Mackey, L. W. (2019). Minimum Stein discrepancy estimators. In Wallach, H., Larochelle, H., Beygelzimer, A., d’Alché-Buc, F., Fox, E., and Garnett, R., editors, *Proceedings of the 33rd International Conference on Neural Information Processing Systems*, Advances in Neural Information Processing Systems 32, pages 12964–12976. Curran Associates, Inc.
- Beltrán-Beltrán, J. I. and O’Reilly, F. J. (2019). On goodness of fit tests for the Poisson, negative binomial and binomial distributions. *Statistical Papers*, 60(1):1–18.
- Betsch, S. and Ebner, B. (2019). A new characterization of the Gamma distribution and associated goodness-of-fit tests. *Metrika*, 82(7):779–806.

- Betsch, S. and Ebner, B. (2020). Testing normality via a distributional fixed point property in the Stein characterization. *TEST*, 29(1):105–138.
- Betsch, S. and Ebner, B. (2021). Fixed point characterizations of continuous univariate probability distributions and their applications. *Annals of the Institute of Statistical Mathematics*, 73(1):31–59.
- Betsch, S., Ebner, B., and Klar, B. (2021). Minimum L^q -distance estimators for non-normalized parametric models. *The Canadian Journal of Statistics*, 42(2):514–548.
- Brown, T. C. and Phillips, M. J. (1999). Negative binomial approximation with Stein’s method. *Methodology and Computing in Applied Probability*, 1(4):407–421.
- Byrd, R. H., Lu, P., Nocedal, J., and Zhu, C. (1995). A limited memory algorithm for bound constrained optimization. *SIAM Journal on Scientific Computing*, 16(5):1190–1208.
- Chen, L. H. Y. (1975). Poisson approximation for dependent trials. *Annals of Probability*, 3(3):534–545.
- Chen, L. H. Y., Goldstein, L., and Shao, Q.-M. (2011). *Normal approximation by Steins method*. Probability and its applications. Springer, Berlin.
- Choudary, A. D. R. and Niculescu, C. P. (2014). *Real Analysis on Intervals*. Springer India, New Delhi.
- Chwialkowski, K., Strathmann, H., and Gretton, A. (2016). A kernel test of goodness of fit. In Balcan, M. F. and Weinberger, K. Q., editors, *Proceedings of the 33rd International Conference on Machine Learning*, volume 48 of *JMLR: W&CP*, pages 2606–2615. Proceedings of Machine Learning Research.
- Cuppens, R. (1975). *Decomposition of Multivariate Probabilities*. Probability and Mathematical Statistics : A Series of Monographs and Textbooks. Academic Press, New York.
- Dutang, C., Goulet, V., and Pigeon, M. (2008). actuar: An R package for actuarial science. *Journal of Statistical Software*, 25(7).
- Ehm, W. (1991). Binomial approximation to the Poisson binomial distribution. *Statistics & Probability Letters*, 11(1):7–16.
- Eichelsbacher, P. and Reinert, G. (2008). Stein’s method for discrete Gibbs measures. *The Annals of Applied Probability*, 18(4):1588–1618.
- Frey, J. (2012). An exact Kolmogorov–Smirnov test for the Poisson distribution with unknown mean. *Journal of Statistical Computation and Simulation*, 82(7):1023–1033.
- Gibbs, J. W. (2010). *Elementary Principles in Statistical Mechanics: Developed with Especial Reference to the Rational Foundation of Thermodynamics*. Cambridge Library Collection - Mathematics. Cambridge University Press, Cambridge.
- Gorham, J. and Mackey, L. (2015). Measuring sample quality with Stein’s method. In Cortes, C., Lawrence, N. D., Lee, D. D., Sugiyama, M., and Garnett, R., editors, *Proceedings of the 28th International Conference on Neural Information Processing Systems*, Advances in Neural Information Processing Systems 28, pages 226–234. Curran Associates, Inc.
- Götze, F. (1991). On the rate of convergence in the multivariate CLT. *The Annals of Probability*, 19(2):724–739.
- Gürtler, N. and Henze, N. (2000). Recent and classical goodness-of-fit tests for the Poisson distribution. *Journal of Statistical Planning and Inference*, 90(2):207–225.
- Gutmann, M. U. and Hyvärinen, A. (2010). Noise-contrastive estimation: A new estimation principle for unnormalized statistical models. In Teh, Y. W. and Titterton, M., editors, *Proceedings of the 13th International Conference on Artificial Intelligence and Statistics (AISTATS)*, volume 9 of *JMLR: W&CP*, pages 297–304. Journal of Machine Learning Research - Proceedings Track.
- Gutmann, M. U. and Hyvärinen, A. (2012). Noise-contrastive estimation of unnormalized statistical models, with applications to natural image statistics. *Journal of Machine Learning Research*, 13(1):307–361.
- Hayakawa, J. and Takemura, A. (2016). Estimation of exponential-polynomial distribution by holonomic gradient descent. *Communications in Statistics - Theory and Methods*, 45(23):6860–6882.
- Henze, N. (1996). Empirical-distribution-function goodness-of-fit tests for discrete models. *The Canadian Journal of Statistics / La Revue Canadienne de Statistique*, 24(1):81–93.
- Hinton, G. E. (2002). Training products of experts by minimizing contrastive divergence. *Neural Computation*, 14(8):1771–1800.
- Hyvärinen, A. (2005). Estimation of non-normalized statistical models by score matching. *Journal of Machine Learning Research*, 6(24):695–709.

- Hyvärinen, A. (2007). Some extensions of score matching. *Computational Statistics & Data Analysis*, 51(5):2499–2512.
- Inglot, T. (2019). Data driven efficient score tests for Poissonity. *Probability and Mathematical Statistics*, 39(1):115–126.
- Johnson, N. L., Kotz, S., and Kemp, A. W. (1993). *Univariate discrete distributions (2nd Edition)*. Wiley Series in Probability and Mathematical Statistics. Wiley, New York.
- Khmaladze, E. (2013). Note on distribution free testing for discrete distributions. *The Annals of Statistics*, 41(6):2979–2993.
- Klar, B. (1999). Goodness-of-fit tests for discrete models based on the integrated distribution function. *Metrika*, 49(1):53–69.
- Kyriakoussis, A., Li, G., and Papadopoulos, A. (1998). On characterization and goodness-of-fit test of some discrete distribution families. *Journal of Statistical Planning and Inference*, 74(2):215–228.
- Ledwina, T. and Wylupek, G. (2017). On Charlier polynomials in testing Poissonity. *Communications in Statistics - Simulation and Computation*, 46(3):1918–1932.
- Ley, C., Reinert, G., and Swan, Y. (2017). Stein’s method for comparison of univariate distributions. *Probability Surveys*, 14:1–52.
- Ley, C. and Swan, Y. (2011). A unified approach to Stein characterizations. *ArXiv e-prints*, 1105.4925v3.
- Ley, C. and Swan, Y. (2013a). Local Pinsker inequalities via Stein’s discrete density approach. *IEEE Transactions on Information Theory*, 59(9):5584–5591.
- Ley, C. and Swan, Y. (2013b). Stein’s density approach and information inequalities. *Electronic Communications in Probability*, 18(7):1–14.
- Li, S. Z. (2009). *Markov Random Field Modeling in Image Analysis (Third Edition)*. Springer-Verlag, London.
- Liu, Q., Lee, J. D., and Jordan, M. (2016). A kernelized Stein discrepancy for goodness-of-fit tests. In Balcan, M. F. and Weinberger, K. Q., editors, *Proceedings of the 33rd International Conference on Machine Learning*, volume 48 of *JMLR: W&CP*, pages 276–284. Proceedings of Machine Learning Research.
- Lyu, S. (2009). Interpretation and generalization of score matching. In *Proceedings of the 25th Conference on Uncertainty in Artificial Intelligence*, UAI’09, pages 359–366. AUAI Press.
- Matsuda, T. and Hyvärinen, A. (2019). Estimation of non-normalized mixture models. In Chaudhuri, K. and Sugiyama, M., editors, *Proceedings of the 22nd International Conference on Artificial Intelligence and Statistics (AISTATS)*, volume 89 of *PMLR*, pages 2555–2563. Proceedings of Machine Learning Research.
- Nikitin, Y. Y. (2017). Tests based on characterizations, and their efficiencies: A survey. *Acta et Commentationes Universitatis Tartuensis de Mathematica*, 21(1):3–24.
- Peköz, E. A. (1996). Stein’s method for geometric approximation. *Journal of Applied Probability*, 33(3):707–713.
- Puig, P. and Weiß, C. H. (2020). Some goodness-of-fit tests for the Poisson distribution with applications in biodosimetry. *Computational Statistics & Data Analysis*, 144:106878.
- R Core Team (2020). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna.
- Rizzo, M. and Székely, G. (2019). *energy: E-Statistics: Multivariate Inference via the Energy of Data*. R package version 1.7-7.
- Rueda, R. and O’Reilly, F. (1999). Tests of fit for discrete distributions based on the probability generating function. *Communications in Statistics - Simulation and Computation*, 28(1):259–274.
- Rueda, R., O’Reilly, F., and Pérez-Abreu, V. (1991). Goodness of fit for the Poisson distribution based on the probability generating function. *Communications in Statistics - Theory and Methods*, 20(10):3093–3110.
- Stein, C. (1972). A bound for the error in the normal approximation to the distribution of a sum of dependent random variables. In *Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability, Volume 2: Probability Theory*, pages 583–602, Berkeley. University of California Press.
- Stein, C. (1986). *Approximate computation of expectations*. Lecture Notes - Monograph Series, Vol. 7. Institute of Mathematical Statistics, Hayward.
- Stein, C., Diaconis, P., Holmes, S., and Reinert, G. (2004). Use of exchangeable pairs in the analysis of simulations. In Diaconis, P. and Holmes, S., editors, *Stein’s Method*, volume 46 of *Lecture Notes – Monograph Series*, pages 1–25, Beachwood. Institute of Mathematical Statistics.
- Székely, G. and Rizzo, M. (2004). Mean distance test of Poisson distribution. *Statistics & Probability Letters*, 67(3):241–247.

- Takenouchi, T. and Kanamori, T. (2017). Statistical inference with unnormalized discrete models and localized homogeneous divergences. *Journal of Machine Learning Research*, 18(56):1–26.
- Uehara, M., Kanamori, T., Takenouchi, T., and Matsuda, T. (2020a). A unified statistically efficient estimation framework for unnormalized models. In Chiappa, S. and Calandra, R., editors, *Proceedings of the 23rd International Conference on Artificial Intelligence and Statistics (AISTATS)*, volume 108 of *PMLR*, pages 809–819. Proceedings of Machine Learning Research.
- Uehara, M., Matsuda, T., and Kim, J. K. (2020b). Imputation estimators for unnormalized models with missing data. In Chiappa, S. and Calandra, R., editors, *Proceedings of the 23rd International Conference on Artificial Intelligence and Statistics (AISTATS)*, volume 108 of *PMLR*, pages 831–841. Proceedings of Machine Learning Research.
- Wolodzko, T. (2019). *extraDistr: Additional Univariate and Multivariate Distributions*. R package version 1.8.11.
- Yang, J., Liu, Q., Rao, V., and Neville, J. (2018). Goodness-of-fit testing for discrete distributions via Stein discrepancy. In Dy, J. and Krause, A., editors, *Proceedings of the 35th International Conference on Machine Learning*, volume 80 of *PMLR*, pages 5561–5570. Proceedings of Machine Learning Research.
- Yu, S., Drton, M., and Shojaie, A. (2019). Generalized score matching for non-negative data. *Journal of Machine Learning Research*, 20(76):1–70.