

Generalization of Rayleigh's Curse on Parameter Estimation with Incoherent Sources

Lijun Peng¹ and Xiao-Ming Lu^{1,*}

¹*Department of Physics, Hangzhou Dianzi University, Hangzhou 310018, China*

The basic idea behind Rayleigh's criterion on resolving two incoherent optical point sources is that the overlap between the spatial modes from different sources would reduce the estimation precision for the locations of the sources, dubbed Rayleigh's curse. We generalize the concept of Rayleigh's curse to the abstract problems of quantum parameter estimation with incoherent sources. To manifest the effect of Rayleigh's curse on quantum parameter estimation, we define the curse matrix in terms of quantum Fisher information and introduce the global and local immunity to the curse accordingly. We further derive the expression for the curse matrix and give the necessary and sufficient condition on the immunity to Rayleigh's curse. For estimating the one-dimensional location parameters with a common initial state, we demonstrate that the global immunity to the curse on quantum Fisher information is impossible for more than two sources.

I. INTRODUCTION

Rayleigh's criterion on distinguishing two incoherent optical point sources determines a characteristic distance, below which resolving two incoherent sources by imaging are thought to be difficult [1]. The basic idea behind Rayleigh's criterion is that the overlap between the spot images would impede the resolution of the point sources. As noted in the Feynman lectures on physics [2, Sec. 30–4], Rayleigh's criterion on the resolution of point sources of lights is a rough idea. Instead, a meticulous method of measuring the resolution power of incoherent point sources can be achieved by resorting to statistical approaches, e.g., parameter estimation [3, 4] or hypothesis testing [5–8].

One great advantage of the parameter estimation approach as well as the hypothesis testing approach is the availability of optimizing over quantum measurements [9–11] using quantum estimation and detection theory [12, 13]. The quantum limit of the estimation precision for the separation of the two incoherent point sources has been revealed to be unexpectedly much higher than that of direct imaging in the sub-Rayleigh region [9], indicating that the overlap between quantum states may not impede the estimation of the separation, as the conventional wisdom suggests. Tsang *et al.* introduced Rayleigh's curse in Ref. [9] to dub the effect that “the positions of two incoherent sources should become harder to estimate when their radiations overlap.” and showed that the estimation precision of the separation between the two optical point sources is immune to Rayleigh's curse when optimizing quantum measurements. A lot of efforts has been denoted into further investigating and demonstrating the quantum superiority brought by optimizing measurements [14–33].

To further understand and systematically study the effect of the overlap between quantum states from different sources on the precision of parameter estimation,

we in this work generalize Rayleigh's curse to the abstract problems of parameter estimation with incoherent sources. We consider the influence of Rayleigh's curse on the quantum Fisher information (QFI) matrix [12, 13, 34], which plays the pivotal role in quantum parameter estimation. To manifest Rayleigh's curse on the QFI, we define the QFI-curse matrix by comparing the real QFI matrix and that as if the quantum states from different sources are orthogonal. We further introduce the global and local immunity to the QFI curse and derive the necessary and sufficient condition on the immunity of the QFI-curse. For estimating one-dimensional location parameters with a common initial states, we show that the global immunity is impossible for more than three incoherent sources. Moreover, for two-source cases, we derives an analytic form of the QFI-curse matrix.

II. PARAMETER ESTIMATION WITH INCOHERENT SOURCES

Let us start by supposing a set of incoherent sources, each emitting a signal that is transmitted to a common receiver (see Fig. 1 for an illustration). The signals are received in the form of quantum states. Assume that the quantum states received from each source are pure states and denote them by a set of state vectors $|\psi_j\rangle$ for $j = 1, 2, \dots, n$. Even when quantum state emitted by different sources are orthogonal, the received states may be non-orthogonal, due to some physical processes during the transmission and reception, e.g., diffraction of light, noise effects, and filtering operations. These processes are represented by an abstract channel in Fig. 1. The incoherence of the sources is manifested in the fact that the relative phases between different state vectors $|\psi_j\rangle$ are totally random. As a result, the quantum state of the receiver is described by the density operator in the form of

$$\rho = \sum_{j=1}^n w_j |\psi_j\rangle\langle\psi_j|, \quad (2.1)$$

* lxm@hdu.edu.cn; <http://xmlu.me>




FIG. 1. Incoherent source model.

where the weight w_j is the prior probability of the system being in the state $|\psi_j\rangle$.

In this work, we consider parameter estimation problems with incoherent sources. We assume that the state vectors $|\psi_j\rangle$ depend on a d -dimensional vector parameter $\theta = (\theta_1, \theta_2, \dots, \theta_d)$ and the weights w_j 's are independent of θ . The value of θ need be estimated by observing the quantum system possessed by the receiver. Particularly, we consider the cases where the number of incoherent sources is much less than the dimension of the Hilbert space for the underlying quantum system. In other words, the density operators considered in this work are of low rank. This includes the special case of infinite dimensional Hilbert space.

Quantum parameter estimation theory provides a powerful tool for pursuing the fundamental limits on the estimation error. For any unbiased estimator and any quantum measurement, the error-covariance matrix \mathcal{E} of parameter estimation, defined by $\mathcal{E}_{jk} = \mathbb{E}[(\hat{\theta}_j - \theta_j)(\hat{\theta}_k - \theta_k)]$ with $\hat{\theta}_j$ being the estimate for θ_j and \mathbb{E} the expectation over the observations, obeys the quantum Cramér-Rao bound (QCRB)

$$\mathcal{E} \geq \mathcal{F}^{-1}, \quad (2.2)$$

where \mathcal{F} is the QFI matrix [12, 13, 34–37]. The entries of the QFI matrix are defined by

$$\mathcal{F}_{\mu\nu} = \text{Re tr}(L_\mu L_\nu \rho), \quad (2.3)$$

where the symmetric logarithmic derivative (SLD) operator L_μ for θ_μ is an Hermitian operator satisfying

$$\frac{1}{2}(L_\mu \rho + \rho L_\mu) = \partial_\mu \rho \quad (2.4)$$

with $\partial_\mu := \partial/\partial\theta_\mu$ being defined for short. Despite that the QCRB is not guaranteed to be attainable in general for the joint estimation of multiple parameters, the QFI matrix still reflects lots of information about the quantum limit of estimation errors [38–44].

Gauge symmetry.— Before calculating the QFI matrix, observe that, due to the incoherent characteristic of the model, the density operator in Eq. (2.1) is invariant under the (local) gauge transformations $|\psi_j\rangle \mapsto e^{i\alpha_j} |\psi_j\rangle$ for $j = 1, 2, \dots, n$, where α_j are arbitrary real-valued functions of θ . The QFI matrix is solely determined by the state vectors $|\psi_j\rangle$ and their derivatives $\partial_\mu |\psi_j\rangle$; The latter will have additional terms under the local gauge transformations, that is,

$$\partial_\mu |\psi\rangle \mapsto \partial_\mu e^{i\alpha} |\psi\rangle = e^{i\alpha} \partial_\mu |\psi\rangle + i(\partial_\mu \alpha) e^{i\alpha} |\psi\rangle \quad (2.5)$$




FIG. 2. Classical and quantum incoherent-source models. (a) In classical model, the state from each source is a probability distribution. (b) In quantum model, the (pure) state from each source is represented by a state vector $|\psi_j\rangle$ in a Hilbert space. The infinitesimal transformation of a state vector can be characterized by its covariant derivatives $|\nabla_\mu \psi_j\rangle$ in the tangent plane. The state vectors, their covariant derivatives, and the prior probabilities of each state determine the QFI matrix.

and the inner products between the state vector and its derivative transform as

$$\langle \psi | \partial_\mu | \psi \rangle \mapsto \langle \psi | \partial_\mu | \psi \rangle + i(\partial_\mu \alpha). \quad (2.6)$$

To avoid the cumbersomeness caused by these additional terms, we use the covariant derivative $|\nabla_\mu \psi\rangle := (\mathbf{1} - |\psi\rangle\langle\psi|)\partial_\mu |\psi\rangle$ instead of the ordinary derivatives for convenience. The covariant derivatives are always orthogonal to the state vectors and changes as $|\nabla_\mu \psi\rangle \mapsto e^{i\alpha} |\nabla_\mu \psi\rangle$ under the gauge transformation $|\psi\rangle \mapsto e^{i\alpha} |\psi\rangle$. Moreover, it satisfies that $\partial_\mu |\psi\rangle\langle\psi| = |\nabla_\mu \psi\rangle\langle\psi| + |\psi\rangle\langle\nabla_\mu \psi|$ and $\partial_\mu \langle\psi|\psi\rangle = \langle\nabla_\mu \psi|\psi\rangle + \langle\psi|\nabla_\mu \psi\rangle$. The QFI matrix is totally determined by the mathematical structure of the state vectors, their covariant derivatives, and the prior probability of each state, see Fig. 2 for an illustration.

III. CURSE ON THE QFI

Now, we say that a set $\{|\psi_j\rangle\}$ of state vectors satisfies the local orthogonalization condition if

$$\langle \psi_j | \psi_k \rangle = \langle \psi_j | \nabla_\mu \psi_k \rangle = 0 \quad \forall j \neq k \quad \text{and} \quad \forall \mu. \quad (3.1)$$

In such case, it can be shown that $L_\mu = 2 \sum_j \partial_\mu |\psi_j\rangle\langle\psi_j|$ is the SLD operator for ρ with respect to the parameter θ_μ and thus the entries of the QFI matrix are simply given by

$$\tilde{\mathcal{F}}_{\mu\nu} := 4 \sum_{j=1}^n w_j \text{Re} \langle \nabla_\mu \psi_j | \nabla_\nu \psi_j \rangle. \quad (3.2)$$

This QFI matrix can be understood as the average of the QFI matrix in each state vectors with the prior probability.

QFI-curse matrix.— We shall focus on the cases where the local orthogonalization condition Eq. (3.1) is not satisfied. The QFI matrix can be formally written as

$$\mathcal{F} = \tilde{\mathcal{F}} - \mathcal{C}, \quad (3.3)$$

where $\tilde{\mathcal{F}}$ is the QFI matrix given by Eq. (3.2) and \mathcal{C} is a matrix manifesting the influence of the overlap between the state vectors on the QFI matrix. We call \mathcal{C} the QFI-curse matrix as it inclines to diminish the QFI when the local orthogonalization condition does not hold.

Assume that the state vectors $|\psi_j\rangle$ and their non-vanishing derivatives $|\nabla_\mu\psi_j\rangle$ are all linearly-independent and the prior weights w_j are all independent of θ . We show in the Appendix A that the curse matrix for the QFI can be expressed as

$$\mathcal{C}_{\mu\nu} = \text{Re tr}(\mathbf{X}_\mu \mathbf{A} \mathbf{X}_\nu), \quad (3.4)$$

where \mathbf{A} is the Gram matrix of $\{|\phi_j\rangle\}$ with $|\phi_j\rangle := \sqrt{w_j}|\psi_j\rangle$ being defined for convenience, i.e., $\mathbf{A}_{jk} := \langle\phi_j|\phi_k\rangle$, and \mathbf{X}_μ are $n \times n$ Hermitian matrices determined by

$$\frac{1}{2}(\mathbf{A} \mathbf{X}_\mu + \mathbf{X}_\mu \mathbf{A}) = \mathbf{Z}_\mu \quad (3.5)$$

with \mathbf{Z}_μ being the $n \times n$ Hermitian matrix defined by

$$[\mathbf{Z}_\mu]_{jk} := i \langle\phi_j|\nabla_\mu\phi_k\rangle - i \langle\nabla_\mu\phi_j|\phi_k\rangle. \quad (3.6)$$

Note that all diagonal entries of \mathbf{Z}_μ are zero according to the above definition, for any state vector is orthogonal to its covariant derivative. Due to Eq. (3.5), the entries of the curse matrix can also be written as

$$\mathcal{C}_{\mu\nu} = \text{tr}(\mathbf{X}_\mu \mathbf{Z}_\nu), \quad (3.7)$$

which may be more convenient for practical calculation.

To understand and even dispel Rayleigh's curse on parameter estimation with incoherent sources, we need to further analyze the QFI-curse matrix. Let v be a column vector in \mathbb{R}^d and represent a direction in the parameter space. We interpret the quantity $v^\top \mathcal{C} v$ as the QFI curse along v , where \top denotes matrix transpose. The meaning of the QFI curse along v can be understood in an intuitive way by resorting to the second form of the QCRB [12, Chapter VIII]: $v^\top \mathcal{E}^{-1} v \geq v^\top \mathcal{F} v$ for any v in \mathbb{R}^d . Through a contour plot of $v^\top \mathcal{E}^{-1} v$ as a function of v , the second form of QCRB implies that, for each positive number c , the concentration ellipsoid given by those v that are restricted by $v^\top \mathcal{E}^{-1} v = c$ always lies outside the ellipsoid whose equation is $v^\top \mathcal{F} v = c$. See Fig. 3 for an illustration.

Note that $v^\top \mathcal{C} v = \text{tr}(\mathbf{X} \mathbf{A} \mathbf{X}) \geq 0$ is satisfied for any v , where we have defined $\mathbf{X} := \sum_\mu v_\mu \mathbf{X}_\mu$. In other words, the curse matrix \mathcal{C} is positive semidefinite. This reflects the negative influence of the overlaps between state vectors on the QFI of the mixed states. In fact, the positivity of the curse matrix can be understood from the perspective of the monotonicity of QFI under quantum operations [45–47] as follows. Let $\tilde{\rho} = \sum_j w_j |\psi_j\rangle\langle\psi_j| \otimes |j\rangle\langle j|$ with $|j\rangle$ being a set of orthonormal states in an ancillary Hilbert space and independent of θ . It can be shown that the QFI matrix for $\tilde{\rho}$ is the same of $\tilde{\mathcal{F}}$ given by Eq. (3.2). The incoherent-source state ρ can be obtained by performing on $\tilde{\rho}$ the partial trace operation with respective




FIG. 3. Geometric view of the QFI-curse along a direction v . Here, we consider two-parameter estimation as an example. The contour lines of $v^\top \mathcal{E}^{-1} v$, $v^\top \mathcal{F} v$, and $v^\top \tilde{\mathcal{F}} v$ are all ellipses. For each positive number c , the ellipse given by $v^\top \mathcal{E}^{-1} v = c$ must lay outside that given by $v^\top \mathcal{F} v = c$, which in turn must lay outside that given by $v^\top \tilde{\mathcal{F}} v = c$. The QFI curse along v is the distance between the $v^\top \tilde{\mathcal{F}} v$ and $v^\top \mathcal{F} v$.

to the ancilla, which is a quantum operation and thus cannot increase the QFI according to the monotonicity of QFI. Therefore, the QFI-curse matrix \mathcal{C} must be positive semidefinite.

IV. IMMUNITY TO THE QFI CURSE

When $v^\top \mathcal{C} v$ vanishes, we say that the QFI is *immune* to Rayleigh's curse along the direction v . Moreover, we say that the immunity is *global* if it holds for a direction v that is independent of the true value of the parameters of interest; Otherwise, we say the immunity is *local*. Finding all vectors v such that $v^\top \mathcal{C} v$ vanishes is mathematically equivalent to seeking the kernel of the curse matrix.

We show in the following that the QFI is immune to Rayleigh's curse along a parameter direction v , if and only if

$$\langle\psi_j|D_v\psi_k\rangle = \langle D_v\psi_j|\psi_k\rangle \quad \forall j < k \quad (4.1)$$

are satisfied, where we $|D_v\psi\rangle := \sum_\mu v_\mu |\nabla_\mu\psi\rangle$ is the directional derivative of a state vector $|\psi\rangle$ along v . Defining $\mathbf{Z} = \sum_\mu v_\mu \mathbf{Z}_\mu$, the condition Eq. (4.1) is then equivalent to $\mathbf{Z} = \mathbf{O}$. Notice that $\mathbf{Z} = (\mathbf{A} \mathbf{X} + \mathbf{X} \mathbf{A})/2$ according to the definitions. The sufficiency of Eq. (4.1) is then evident by observing that $v^\top \mathcal{C} v = \text{tr}(\mathbf{X} \mathbf{Z})$. To prove the necessity of condition Eq. (4.1), note that $v^\top \mathcal{C} v = \text{tr}(\mathbf{X} \mathbf{A} \mathbf{X})$ vanishes if and only if $\sqrt{\mathbf{A} \mathbf{X}}$ is a zero matrix, for $\text{tr}(\mathbf{X} \mathbf{A} \mathbf{X})$ is the Hilbert-Schmidt norm of $\sqrt{\mathbf{A} \mathbf{X}}$. In such case, \mathbf{Z} must be a zero matrix due to $\mathbf{Z} = (\mathbf{A} \mathbf{X} + \mathbf{X} \mathbf{A})/2$.

QFI-curse formula.— To calculate the QFI-curse matrix, one needs to solve \mathbf{X}_μ from Eq. (3.5). This can be

done in general by invoking the eigenvalue decomposition of \mathbf{A} . Let \mathbf{P}_q be the eigen-projection of \mathbf{A} with the eigenvalue λ_q , i.e., $\mathbf{A}\mathbf{P}_q = \lambda_q\mathbf{P}_q$. It then follows from Eq. (3.5) that $\mathbf{X}_\mu = \sum_{qq'} 2/(\lambda_q + \lambda_{q'})\mathbf{P}_q\mathbf{Z}_\mu\mathbf{P}_{q'}$, implying

$$\mathcal{C}_{\mu\nu} = \sum_{qq'} \frac{2}{\lambda_q + \lambda_{q'}} \text{tr}(\mathbf{Z}_\mu\mathbf{P}_q\mathbf{Z}_\nu\mathbf{P}_{q'}). \quad (4.2)$$

For two-source problems, the QFI curse along v can be simply expressed as (see Appendix B for a detailed derivation)

$$v^\top \mathcal{C} v = 4w_1w_2 \left[|\gamma|^2 + \frac{(\text{Re } \gamma \langle \psi_2 | \psi_1 \rangle)^2}{1 - |\langle \psi_2 | \psi_1 \rangle|^2} \right] \quad (4.3)$$

with $\gamma := i \langle \psi_1 | D_v \psi_2 \rangle - i \langle D_v \psi_1 | \psi_2 \rangle$ being defined for brevity. Without loss of generality, assume that $\langle \psi_2 | \psi_1 \rangle$ is real and nonnegative, which can always be implemented by changing the phases of $|\psi_1\rangle$ and $|\psi_2\rangle$, i.e., fixing a specific gauge. Then, $\langle \psi_2 | \psi_1 \rangle$ equals the fidelity between the two state vectors and will be denoted by f . For two close sources such that the fidelity $f \approx 1$, we have $\gamma \approx 0$, for $f \rightarrow 1$ implies $|\psi_1\rangle \rightarrow |\psi_2\rangle$. To see the limit of Eq. (4.3) as $f \rightarrow 1$, we introduce a normalized vector $|\tau\rangle := (|\psi_1\rangle - |\psi_2\rangle)/\sqrt{2-2f}$ so that γ can be written as

$$\gamma = i\sqrt{2-2f}(\langle \tau | D_v \psi_2 \rangle + \langle D_v \psi_1 | \tau \rangle). \quad (4.4)$$

Notice that $|\tau\rangle$ is the unit vector representing the direction from the tip of $|\psi_2\rangle$ to that of $|\psi_1\rangle$ and thus is well defined under the limit $|\psi_1\rangle \rightarrow |\psi_2\rangle$. Substituting Eq. (4.4) into Eq. (4.3), we get

$$v^\top \mathcal{C} v = 8w_1w_2 \left[(1-f)|\tilde{\gamma}|^2 + \frac{f^2(\text{Re } \tilde{\gamma})^2}{1+f} \right], \quad (4.5)$$

where $\tilde{\gamma} := i \langle \tau | D_v \psi_2 \rangle + i \langle D_v \psi_1 | \tau \rangle$ is in general a complex number. For two close sources such that the fidelity f is close to unit, we get

$$v^\top \mathcal{C} v \xrightarrow{f \rightarrow 1} 4w_1w_2(\text{Re } \tilde{\gamma})^2. \quad (4.6)$$

This tell us that the local immunity to the QFI curse along a direction v can be achieved for two sources, if the quantity $\text{Re } \tilde{\gamma}$ goes to zero as the two state vectors become close.

V. UNITARY PARAMETER ESTIMATION

We here consider the cases where the state vectors are given by $|\psi_j\rangle = U_j |\psi\rangle$, where U_j is a family of unitary operators depending on the parameter θ of interest and $|\psi\rangle$ is the initial state in common. The covariant derivatives of the state vectors are given by

$$|\nabla_\mu \psi_j\rangle = (\mathbf{1} - |\psi_j\rangle\langle\psi_j|)(\partial_\mu U_j) |\psi\rangle = -iG_{\mu,j} |\psi_j\rangle, \quad (5.1)$$

where we have defined

$$G_{\mu,j} := i(\partial_\mu U_j)U_j^\dagger - i \langle \psi_j | (\partial_\mu U_j) U_j^\dagger | \psi_j \rangle \mathbf{1} \quad (5.2)$$

with $\mathbf{1}$ being the identity operator. Note that the operator $G_{\mu,j}$ is Hermitian due to $\partial_\mu(U_j U_j^\dagger) = (\partial_\mu U_j)U_j^\dagger + U_j(\partial_\mu U_j)^\dagger = 0$ and satisfies $\langle \psi_j | G_{\mu,j} | \psi_j \rangle = 0$. Therefore, we get

$$[\mathbf{Z}_\mu]_{jk} = \langle \psi_j | (G_{\mu,k} + G_{\mu,j}) | \psi_k \rangle. \quad (5.3)$$

The necessary and sufficient condition on the immunity to Rayleigh's curse along a direction v , i.e., Eq. (4.1), becomes

$$\sum_\mu v_\mu \langle \psi_j | (G_{\mu,k} + G_{\mu,j}) | \psi_k \rangle = 0 \quad \forall j < k. \quad (5.4)$$

We further assume that each state vector $|\psi_j\rangle$ depends on an individual parameter θ_j standing for the shifted location from a common initial state $|\psi\rangle$ by a fixed unitary transformation, i.e., $|\psi_j\rangle = \exp(-i\theta_j H) |\psi\rangle$ with H being an Hermitian operator and independent of θ . This class of problems includes the superresolution for two incoherent optical point sources [9], where the state $|\psi_j\rangle$ is for a photon's spatial degree of freedom at the image plane from the j -th source. To be specific, assuming a spatially-invariant unit-magnification imaging system, the state vectors from each source are in the form of $|\psi_j\rangle = e^{-i\theta_j H} |\psi\rangle$, where H is the momentum operator (\hbar is set to be unit) and $|\psi\rangle$ is the state of a photon's spatial degree of freedom when the point optical source is located at the origin of coordinate. In Ref. [9], Tsang *et al.* showed that the QFI about the separation, i.e., $\theta_1 - \theta_2$, is independent of its true value, provided that the two sources are equally weighted ($w_1 = w_2$) and the normalized amplitude-point-spread function for the imaging system has a constant position-independent phase so that $\langle \psi_2 | \psi_1 \rangle$, $\langle \psi_2 | \nabla_1 \psi_1 \rangle$, and $\langle \nabla_2 \psi_2 | \psi_1 \rangle$ have the same phase. In our context, it can be reinterpreted that the QFI has the global immunity to Rayleigh's curse along $v = (1/\sqrt{2}, -1/\sqrt{2})$, for the directional derivative along v is equivalent to the partial derivative with respect to the separation $(\theta_1 - \theta_2)/\sqrt{2}$ when we use the rotated parameters $(\theta_1 + \theta_2)/\sqrt{2}$ and $(\theta_1 - \theta_2)/\sqrt{2}$ as the parameters of interest. We will explain it in detail and generalize the results in what follows.

Without loss of generality, we assume that $\langle \psi | H | \psi \rangle = 0$ henceforth; Otherwise, we can take the gauge transformation $|\psi_j\rangle \mapsto \exp(i\theta_j \langle \psi | H | \psi \rangle) |\psi_j\rangle$ and replace H by $\Delta H := H - \langle \psi | H | \psi \rangle \mathbf{1}$. In such case, we have $|\nabla_\mu \psi_j\rangle = -i\delta_{\mu j} H |\psi_j\rangle$. The necessary and sufficient condition Eq. (4.1) on the immunity to the QFI curse along v then becomes

$$(v_j + v_k) \langle \psi | H e^{i(\theta_j - \theta_k)H} | \psi \rangle = 0 \quad \forall j < k. \quad (5.5)$$

Since the quantity $\langle \psi | H e^{i(\theta_j - \theta_k)H} | \psi \rangle$ in general depends on the true values of θ , the global immunity can only be

achieved when $v_j = -v_k$ for all different j and k , which is only possible for the case of two sources. For two-source cases, this global immunity along $v \propto (1, -1)$ is irrelevant to the prior weights w_j and the specific form of H and $|\psi\rangle$, implying that such a global immunity still holds for two unequal weighted sources and for point-spread functions with position-dependent phases.

The QFI-curse matrix for the model considered above can be obtained as follows. By defining a characteristic function as $g(x) := \langle \psi | e^{-ixH} | \psi \rangle$ and noting that $g'(x) = -i \langle \psi | H e^{-ixH} | \psi \rangle$, it follows from Eq. (4.3) that

$$v^\top \mathcal{C} v = 4w_1 w_2 (v_1 + v_2)^2 \Theta(\theta_2 - \theta_1)$$

$$\text{with } \Theta(x) := |g'(x)|^2 + \frac{[\text{Im } g'(x)g(x)]^2}{1 - |g(x)|^2}. \quad (5.6)$$

This form implies that the QFI-curse matrix can be simply written as

$$\mathcal{C} = 4w_1 w_2 \Theta(\theta_2 - \theta_1) \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}. \quad (5.7)$$

When we rotate the parameters θ_1 and θ_2 to $(\theta_1 + \theta_2)/\sqrt{2}$ and $(\theta_1 - \theta_2)/\sqrt{2}$, the curse matrix is diagonal as $\mathcal{C} = 8w_1 w_2 \Theta(\theta_2 - \theta_1) \text{diag}\{2, 0\}$. It is worth noting that for the case of locating two incoherent optical point sources, the QFI matrix regarding the centroid $(\theta_2 + \theta_1)/2$ and separation $\theta_2 - \theta_1$ is no longer diagonal when the two intensities are unequal [22]. However, here, it can be seen that the QFI-curse matrix is still diagonal.

For two close sources such that $|\theta_2 - \theta_1| \ll 1$, we can expand the relevant quantities with respect to $\theta_2 - \theta_1$ as follows:

$$g(\delta) = 1 - \frac{\delta^2}{2} \langle \psi | H^2 | \psi \rangle + \mathcal{O}(\delta^3), \quad (5.8)$$

$$g'(\delta) = -\delta \langle \psi | H^2 | \psi \rangle + \frac{i\delta^2}{2} \langle \psi | H^3 | \psi \rangle + \mathcal{O}(\delta^3), \quad (5.9)$$

where δ is short for $\theta_2 - \theta_1$ and we have used $g'(0) = -i \langle \psi | H | \psi \rangle = 0$ and $g''(0) = -\langle \psi | H^2 | \psi \rangle$. Therefore, for small δ , we get

$$\Theta(\delta) \approx \delta^2 \left(\langle \psi | H^2 | \psi \rangle^2 + \frac{\langle \psi | H^3 | \psi \rangle^2}{4 \langle \psi | H^2 | \psi \rangle} \right). \quad (5.10)$$

As $\delta \rightarrow 0$, we can see that the local immunity of the QFI-curse occurs along any direction v .

In short, we have shown that, for the two-incoherent-source parameter estimation model $\rho = \sum_{j=1,2} w_j |\psi_j\rangle\langle\psi_j|$ with $|\psi_j\rangle = e^{-i\theta_j H} |\psi\rangle$, the QFI is globally immune along the direction $v \propto (1, -1)$ and locally immune along all direction, to the Rayleigh's curse at the region of small δ .

VI. DISCUSSION

To summarize, we have generalized the concept of Rayleigh's curse to quantum parameter estimation with

incoherent sources. To manifest the influence of the overlap between quantum states from each sources on estimating the carried parameters, we have defined the QFI-curse matrix and introduced the global and local immunities to the QFI-curse accordingly. Using the technology of non-orthonormal basis, we have derived the computable formula for the QFI-curse matrix and also given a compact form of the QFI-curse matrix for the case of two incoherent sources. We have applied the QFI-curse approach to the estimation of one-dimensional location parameters. We believe that our approach is useful to quantum metrology with incoherent sources.

In this work, we mainly focus on the condition on the vanishing of the QFI curse along a parameter direction. It will be also interesting to investigate the condition on the maximal QFI curse along a parameter direction, which corresponds to the vanishing of QFI along a parameter direction. Bisketzi, Branford, and Datta have numerically demonstrated that in the regime of small separations, no more than two independent parameters can be effectively estimated, as the QFI matrix is no more than rank two [48]. Calculating the QFI-curse matrix for the same scenario as Ref. [48] and investigating the condition on the maximal QFI curses will help to understand the above-mentioned phenomena. We leave further explorations on this issue in the future work.

ACKNOWLEDGMENTS

We thank Chandan Datta, Stefano Pirandola, and Animesh Datta for the helpful communications and enlightening discussions. This work is supported by the National Natural Science Foundation of China (Grants No. 11805048, No. 61871162, and No. 11935012) and Zhejiang Provincial Natural Science Foundation of China (Grant No. LY18A050003).

Appendix A: Derivation of the formula for the QFI-curse matrix

We here give a detailed proof of the formula Eq. (3.4) for the QFI-curse matrix. Our approach is based on the use of non-orthonormal basis of the subspace supporting the underlying density operator and its partial derivatives with respect to the parameters of interest.

1. Non-orthonormal basis

We follow Ref. [49] to use non-orthogonal bases for expanding the relevant operators involved in calculating QFI matrix. Let $\mathcal{B} = \{|\phi_j\rangle \mid j = 1, 2, \dots, n\}$ be a set of linearly independent vectors spanning a Hilbert space \mathcal{H}_s . Any bounded operator A acting on \mathcal{H}_s can be uniquely

expressed as

$$A = \sum_{jk} R(A)_{jk} |\phi_j\rangle\langle\phi_k|, \quad (\text{A1})$$

where $R(A)_{jk}$ are the coefficients of the expansion. We call $R(A)$, the matrix with entries $R(A)_{jk}$, the R-matrix for an operator A in the basis \mathcal{B} . In this work, we will use the following properties of the R-matrices [31, 49, 50]:

1. $R(A^\dagger) = R(A)^\dagger$.
2. $R(aA+bB) = aR(A)+bR(B)$ for any bounded linear operators A and B on the subspace spanned by \mathcal{B} and any two complex numbers a and b .
3. $\text{tr} A = \text{tr}[R(A)\Omega]$, where Ω is the Gram matrix of \mathcal{B} and given by $\Omega_{jk} := \langle\phi_j|\phi_k\rangle$.
4. $R(AB) = R(A)\Omega R(B)$.

The first and second properties are obvious. The third property can be proved as

$$\begin{aligned} \text{tr}(A) &= \sum_{jk} R(A)_{jk} \text{tr}(|\phi_j\rangle\langle\phi_k|) \\ &= \sum_{jk} R(A)_{jk} \langle\phi_k|\phi_j\rangle = \text{tr}[R(A)\Omega]. \end{aligned} \quad (\text{A2})$$

The fourth property can be proved by noting that

$$\begin{aligned} AB &= \sum_{jk} R(A)_{jk} |\phi_j\rangle\langle\phi_k| \sum_{\ell m} R(B)_{\ell m} |\phi_\ell\rangle\langle\phi_m| \\ &= \sum_{jm} \left[\sum_{k\ell} R(A)_{jk} \langle\phi_k|\phi_\ell\rangle R(B)_{\ell m} \right] |\phi_j\rangle\langle\phi_m|. \end{aligned} \quad (\text{A3})$$

2. General expression for the QFI-curse matrix

To calculate the QFI of the incoherent-source density operator $\rho = \sum_j w_j |\psi_j\rangle\langle\psi_j|$, Genoni and Tufarelli in Ref. [49] used the non-orthogonal bases constituted by the state vectors $|\psi_j\rangle$ and their derivatives $\partial_\mu |\psi_j\rangle$ with respect to θ_μ . In this work, we improve their approach by using the weighted state vectors $\sqrt{w_j} |\psi_j\rangle$ and their covariant derivatives $\sqrt{w_j} |\nabla_\mu \psi_j\rangle$ instead to constitute the non-orthogonal basis. We make the following assumptions on the incoherent-source density operators: (i) w_j are independent of θ and (ii) all non-vanishing $|\nabla_\mu \psi_j\rangle$ and $|\psi_j\rangle$ are linearly independent. Let us choose $|\phi_j\rangle = \sqrt{w_j} |\psi_j\rangle$ for $j = 1, 2, \dots, n$ and take from the non-vanishing covariant derivatives as $|\phi_j\rangle$ for $j > n$. It follows that $\rho = \sum_{j=1}^n |\phi_j\rangle\langle\phi_j|$ and $\partial_\mu \rho = \sum_{j=1}^n |\nabla_\mu \phi_j\rangle\langle\phi_j| + |\phi_j\rangle\langle\nabla_\mu \phi_j|$. By partitioning the basis into $\{|\phi_j\rangle \mid j = 1, 2, \dots, n\}$ and the remainder, the R-matrices for ρ and $\partial_\mu \rho$ can be expressed in the block form

$$R(\rho) = \begin{bmatrix} \mathbb{I} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \quad \text{and} \quad R(\partial_\mu \rho) = \begin{bmatrix} \mathbf{0} & \mathbf{D}_\mu^\dagger \\ \mathbf{D}_\mu & \mathbf{0} \end{bmatrix}, \quad (\text{A4})$$

respectively, where \mathbb{I} is the identity matrix of size n .

To calculate the QFI matrix, one usually need solve the equation $(L_\mu \rho + \rho L_\mu)/2 = \partial_\mu \rho$ for the SLD operator L_μ . We here give a method to calculate the QFI without explicitly solving L_μ . One of the advantages of absorbing the weighting factors $\sqrt{w_j}$ into the basis vectors is that $R(\rho)$ is idempotent, namely, $R(\rho)^2 = R(\rho)$. Using this property and the properties of the R-matrix, it can be shown that $\mathcal{F}_{\mu\nu} = \text{Re tr}(\Psi_\mu^\dagger \Omega \Psi_\nu)$, where $\Psi_\mu := R(L_\mu \rho) = R(L_\mu) \Omega R(\rho)$. So knowing the matrices Ψ_μ and Ω is sufficient to get the QFI matrix.

To proceed, we partition other relevant operators and the Gram matrix into the same block form as Eq. (A4):

$$\Psi_\mu = \begin{bmatrix} i\mathbf{X}_\mu & \mathbf{0} \\ \mathbf{Y}_\mu & \mathbf{0} \end{bmatrix}, \quad R(L_\mu) = \begin{bmatrix} \mathbf{M}_\mu & \mathbf{N}_\mu^\dagger \\ \mathbf{N}_\mu & \mathbf{K}_\mu \end{bmatrix}, \quad \Omega = \begin{bmatrix} \mathbf{A} & \mathbf{B}^\dagger \\ \mathbf{B} & \mathbf{C} \end{bmatrix}, \quad (\text{A5})$$

where the imaginary unit before \mathbf{X}_μ is introduced for later convenience, the all-zero blocks in Ψ_μ is a result of $\Psi_\mu = R(L_\mu) \Omega R(\rho)$ and the special form of $R(\rho)$ in Eq. (A4). Note that the square matrices \mathbf{A} , \mathbf{C} , \mathbf{M}_μ , and \mathbf{K}_μ are all Hermitian. With the R-matrices, the SLD equation $(L_\mu \rho + \rho L_\mu)/2 = \partial_\mu \rho$ is equivalent to $(\Psi_\mu + \Psi_\mu^\dagger)/2 = R(\partial_\mu \rho)$. Substituting the block form of Ψ_μ in Eq. (A5) into this SLD equation and comparing block by block both sides, we get $\mathbf{X}_\mu = \mathbf{X}_\mu^\dagger$ and $\mathbf{Y}_\mu = 2\mathbf{D}_\mu$. With the block forms of Ω and Ψ_μ , it follows from $\mathcal{F}_{\mu\nu} = \text{Re tr}(\Psi_\mu^\dagger \Omega \Psi_\nu)$ that

$$\begin{aligned} \mathcal{F}_{\mu\nu} &= \text{Re tr} \begin{bmatrix} -i\mathbf{X}_\mu & 2\mathbf{D}_\mu^\dagger \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \begin{bmatrix} \mathbf{A} & \mathbf{B}^\dagger \\ \mathbf{B} & \mathbf{C} \end{bmatrix} \begin{bmatrix} i\mathbf{X}_\nu & \mathbf{0} \\ 2\mathbf{D}_\nu & \mathbf{0} \end{bmatrix} \\ &= \text{Re tr}(\mathbf{X}_\mu^\dagger \mathbf{A} \mathbf{X}_\nu + 4\mathbf{D}_\mu^\dagger \mathbf{C} \mathbf{D}_\nu) - \text{tr}(\mathbf{Z}_\mu \mathbf{X}_\nu + \mathbf{X}_\mu \mathbf{Z}_\nu), \end{aligned} \quad (\text{A6})$$

where we have defined $\mathbf{Z}_\mu := i(\mathbf{B}^\dagger \mathbf{D}_\mu - \mathbf{D}_\mu^\dagger \mathbf{B})$.

To obtain the QFI matrix, we still need to solve the Hermitian matrix \mathbf{X}_μ . Substituting Eq. (A5) into $\Psi_\mu = R(L_\mu) \Omega R(\rho)$ yields

$$i\mathbf{X}_\mu = \mathbf{M}_\mu \mathbf{A} + \mathbf{N}_\mu^\dagger \mathbf{B} \quad \text{and} \quad 2\mathbf{D}_\mu = \mathbf{N}_\mu \mathbf{A} + \mathbf{K}_\mu \mathbf{B}. \quad (\text{A7})$$

It follows from the above equalities that $\mathbf{A} \mathbf{X}_\mu + (\mathbf{A} \mathbf{X}_\mu)^\dagger = -i\mathbf{A} \mathbf{N}_\mu^\dagger \mathbf{B} + i\mathbf{B}^\dagger \mathbf{N}_\mu \mathbf{A}$ and $2\mathbf{B}^\dagger \mathbf{D}_\mu - 2\mathbf{D}_\mu^\dagger \mathbf{B} = \mathbf{B}^\dagger \mathbf{N}_\mu \mathbf{A} - \mathbf{A} \mathbf{N}_\mu^\dagger \mathbf{B}$, which together implies $(\mathbf{A} \mathbf{X}_\mu + \mathbf{X}_\mu \mathbf{A})/2 = \mathbf{Z}_\mu$. Moreover, it can be shown that

$$\text{Re tr}(\mathbf{X}_\mu \mathbf{A} \mathbf{X}_\nu) = \text{tr} \left(\mathbf{X}_\mu \frac{\mathbf{A} \mathbf{X}_\nu + \mathbf{X}_\nu \mathbf{A}}{2} \right) = \text{tr}(\mathbf{X}_\mu \mathbf{Z}_\nu) \quad (\text{A8})$$

and also $\text{Re tr}(\mathbf{X}_\mu \mathbf{A} \mathbf{X}_\nu) = \text{tr}(\mathbf{X}_\nu \mathbf{Z}_\mu)$ by interchanging the subscripts μ and ν . Therefore, the last equality in Eq. (A6) can be simplified to

$$\mathcal{F}_{\mu\nu} = 4 \text{Re tr}(\mathbf{D}_\mu^\dagger \mathbf{C} \mathbf{D}_\nu) - \text{Re tr}(\mathbf{X}_\mu \mathbf{A} \mathbf{X}_\nu). \quad (\text{A9})$$

The covariant derivatives $|\nabla_\nu \phi_j\rangle$ have two indexes ν and j , so it is convenient to use ν and j as a composite index for the matrices \mathbf{B} , \mathbf{C} , and \mathbf{D}_μ . The entries of \mathbf{D}_μ can be denoted by $[\mathbf{D}_\mu]_{\nu j, k}$. It follows from $\partial_\mu \rho =$

$\sum_{j=1}^n |\nabla_\mu \phi_j\rangle\langle\phi_j| + |\phi_j\rangle\langle\nabla_\mu \phi_j|$ that $[D_\mu]_{\nu j, k} = \delta_{\mu\nu} \delta_{jk}$. Therefore, we have

$$\begin{aligned} \text{tr}(D_\mu^\dagger C D_\nu) &= \sum_{j,k,\ell=1}^n \sum_{\alpha,\beta=1}^d [D_\mu^\dagger]_{j,\alpha k} C_{\alpha k, \beta \ell} [D_\nu]_{\beta \ell, j} \\ &= \sum_{j,k,\ell=1}^n \sum_{\alpha,\beta=1}^d \delta_{\mu\alpha} \delta_{jk} C_{\alpha k, \beta \ell} \delta_{\beta\nu} \delta_{\ell,j} \\ &= \sum_{j=1}^n C_{\mu j, \nu j} = \sum_{j=1}^n \langle \nabla_\mu \phi_j | \nabla_\nu \phi_j \rangle \end{aligned} \quad (\text{A10})$$

and

$$\begin{aligned} [Z_\mu]_{jk} &= i \sum_{\alpha,\beta=1}^d [B^\dagger]_{j,\alpha\ell} [D_\mu]_{\alpha\ell, k} - [D_\mu^\dagger]_{j,\alpha\ell} B_{\alpha\ell, k} \\ &= i(B_{\mu k, j}^* - B_{\mu j, k}) \\ &= i \langle \phi_j | \nabla_\mu \phi_k \rangle - i \langle \nabla_\mu \phi_j | \phi_k \rangle. \end{aligned} \quad (\text{A11})$$

3. Calculating the QFI-curse matrix

The last thing to get the curse matrix is solving X_μ from the equation

$$\frac{1}{2}(A X_\mu + X_\mu A) = Z_\mu. \quad (\text{A12})$$

Let P_q be the eigen-projection of A with the eigenvalue λ_q . It follows from Eq. (A12) that

$$P_q Z_\mu P_{q'} = \frac{\lambda_q + \lambda_{q'}}{2} P_q X_\mu P_{q'}, \quad (\text{A13})$$

where we have used $A P_{q'} = \lambda_{q'} P_{q'}$ and $P_q A = \lambda_q P_q$. It then follows that

$$X_\mu = \sum_{qq'} P_q X_\mu P_{q'} = \sum_{qq'} \frac{2}{\lambda_q + \lambda_{q'}} P_q Z_\mu P_{q'}. \quad (\text{A14})$$

As a result, the entries of the curse matrix can be given by

$$C_{\mu\nu} = \text{tr}(X_\mu Z_\nu) = \sum_{qq'} \frac{2}{\lambda_q + \lambda_{q'}} \text{tr}(Z_\mu P_q Z_\nu P_{q'}). \quad (\text{A15})$$

Appendix B: Two-source formula for the QFI-curse

We here consider the 2-rank density operators, where the state vectors $|\psi_j\rangle$ in the mixed states comes from two sources. For 2-rank density operators, the matrix A can always be written as

$$A = \frac{1}{2}(I + \mathbf{r} \cdot \boldsymbol{\sigma}) = \frac{1}{2} \left(I + \sum_{j=1}^3 r_j \sigma_j \right), \quad (\text{B1})$$

where $\mathbf{r} = (r_1, r_2, r_3) \in \mathbb{R}^3$ and $\boldsymbol{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$ is a vector of Pauli matrices. The eigenvalues of A are given by

$$\lambda_\pm = \frac{1 \pm r}{2} \text{ with } r := |\mathbf{r}| = \sqrt{r_1^2 + r_2^2 + r_3^2}. \quad (\text{B2})$$

The corresponding eigen-projections are given by

$$P_\pm = \frac{1}{2}(I \pm \mathbf{n} \cdot \boldsymbol{\sigma}), \quad (\text{B3})$$

where $\mathbf{n} = \mathbf{r}/r$ is a unit vector. To calculate the curse matrix, we first need to obtain Z_μ . For a given vector v in the parameter space \mathbb{R}^n , define $Z = \sum_\mu v_\mu Z_\mu$. It then follows from Eq. (A15) that the curse along the direction v can be expressed as

$$\begin{aligned} v^\top C v &= 2 \text{tr}(Z P_+ Z P_- + Z P_- Z P_+) \\ &\quad + \frac{1}{\lambda_+} \text{tr}(Z P_+ Z P_+) + \frac{1}{\lambda_-} \text{tr}(Z P_- Z P_-). \end{aligned} \quad (\text{B4})$$

Since Z_μ are Hermitian and their diagonal entries are all equal to zero, we can always express Z as $Z = \mathbf{z} \cdot \boldsymbol{\sigma}$, where \mathbf{z} is a 3-dimensional real vector with the components $z_j = \frac{1}{2} \sum_\mu v_\mu \text{tr}(Z_\mu \sigma_j)$. Using Eq. (B3) and the algebraic properties of the Pauli matrices, it is easy to show that

$$2 \text{tr}(Z P_+ Z P_- + \text{H.c.}) = \text{tr}[Z^2 - Z(\mathbf{n} \cdot \boldsymbol{\sigma})Z(\mathbf{n} \cdot \boldsymbol{\sigma})]. \quad (\text{B5})$$

Due to the identities $(\mathbf{a} \cdot \boldsymbol{\sigma})(\mathbf{b} \cdot \boldsymbol{\sigma}) = (\mathbf{a} \cdot \mathbf{b})I + i(\mathbf{a} \times \mathbf{b}) \cdot \boldsymbol{\sigma}$ and $(\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{c} \times \mathbf{d}) = (\mathbf{a} \cdot \mathbf{c})(\mathbf{b} \cdot \mathbf{d}) - (\mathbf{b} \cdot \mathbf{c})(\mathbf{a} \cdot \mathbf{d})$ for any vectors $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}$ in \mathbb{R}^3 , we get

$$\begin{aligned} \text{tr}[Z(\mathbf{n} \cdot \boldsymbol{\sigma})Z(\mathbf{n} \cdot \boldsymbol{\sigma})] &= \text{tr}[\mathbf{z} \cdot \mathbf{n} I + i(\mathbf{z} \times \mathbf{n}) \cdot \boldsymbol{\sigma}]^2 \\ &= 2(\mathbf{z} \cdot \mathbf{n})^2 - 2(\mathbf{z} \times \mathbf{n}) \cdot (\mathbf{z} \times \mathbf{n}) \\ &= 4(\mathbf{z} \cdot \mathbf{n})^2 - 2(\mathbf{z} \cdot \mathbf{z}). \end{aligned} \quad (\text{B6})$$

Since P_\pm are one-dimensional projection operators, it is easy to get

$$P_\pm Z P_\pm = \text{tr}(Z P_\pm) P_\pm = \pm(\mathbf{n} \cdot \mathbf{z}) P_\pm \quad (\text{B7})$$

and thus

$$\text{tr}[Z P_\pm Z P_\pm] = (\mathbf{n} \cdot \mathbf{z})^2. \quad (\text{B8})$$

Putting Eqs. (B4), (B5), (B6) and (B8) together, we get

$$\begin{aligned} v^\top C v &= 4\mathbf{z} \cdot \mathbf{z} + \left(\frac{1}{\lambda_+ \lambda_-} - 4 \right) (\mathbf{n} \cdot \mathbf{z})^2 \\ &= 4\mathbf{z} \cdot \mathbf{z} + \frac{4}{1 - r^2} (\mathbf{r} \cdot \mathbf{z})^2. \end{aligned} \quad (\text{B9})$$

Comparing the definition $A_{jk} = \sqrt{w_j w_k} \langle \psi_j | \psi_k \rangle$ with

$$A = \frac{1}{2}(I + \mathbf{r} \cdot \boldsymbol{\sigma}) = \frac{1}{2} \begin{bmatrix} 1 + r_3 & r_1 - ir_2 \\ r_1 + ir_2 & 1 - r_3 \end{bmatrix}, \quad (\text{B10})$$

the coefficients of the Pauli vector can be expressed as $r_1 = 2\sqrt{w_1 w_2} \operatorname{Re} \langle \psi_2 | \psi_1 \rangle$, $r_2 = 2\sqrt{w_1 w_2} \operatorname{Im} \langle \psi_2 | \psi_1 \rangle$, and $r_3 = w_1 - w_2$. It can be shown that

$$\begin{aligned} 1 - r^2 &= 1 - 4w_1 w_2 |\langle \psi_2 | \psi_1 \rangle|^2 - (w_1 - w_2)^2 \\ &= (w_1 + w_2)^2 - 4w_1 w_2 |\langle \psi_2 | \psi_1 \rangle|^2 - (w_1 - w_2)^2 \\ &= 4w_1 w_2 \left(1 - |\langle \psi_2 | \psi_1 \rangle|^2 \right), \end{aligned} \quad (\text{B11})$$

where we have used $w_1 + w_2 = 1$ in the second equality. Note that $\mathbf{z}_1 - i\mathbf{z}_2 = \mathbf{Z}_{12} = \sqrt{w_1 w_2} \gamma$ with

$$\gamma := i \langle \psi_1 | D_v \psi_2 \rangle - i \langle D_v \psi_1 | \psi_2 \rangle \quad (\text{B12})$$

being defined. So we get $\mathbf{z}_1 = \sqrt{w_1 w_2} \operatorname{Re} \gamma$ and $\mathbf{z}_2 = -\sqrt{w_1 w_2} \operatorname{Im} \gamma$. Thus, we have

$$\mathbf{z} \cdot \mathbf{z} = w_1 w_2 |\gamma|^2, \quad (\text{B13})$$

$$\begin{aligned} \mathbf{r} \cdot \mathbf{z} &= 2w_1 w_2 (\operatorname{Re} \langle \psi_2 | \psi_1 \rangle \operatorname{Re} \gamma - \operatorname{Im} \langle \psi_2 | \psi_1 \rangle \operatorname{Im} \gamma) \\ &= 2w_1 w_2 \operatorname{Re} \gamma \langle \psi_2 | \psi_1 \rangle. \end{aligned} \quad (\text{B14})$$

Substituting Eqs. (B11), (B13), and (B14) into Eq. (B9), we get Eq. (4.3)

-
- [1] F. R. S. Lord Rayleigh, Xxxi. investigations in optics, with special reference to the spectroscope, *Philosophical Magazine Series 5* **8**, 261 (1879).
- [2] R. Feynman, R. Leighton, and M. Sands, *The Feynman Lectures on Physics: Volume I* (Addison-Wesley, 1963).
- [3] S. Ram, E. S. Ward, and R. J. Ober, Beyond Rayleigh's criterion: A resolution measure with application to single-molecule microscopy, *Proc. Natl. Acad. Sci. U.S.A.* **103**, 4457 (2006).
- [4] J. Chao, E. S. Ward, and R. J. Ober, Fisher information theory for parameter estimation in single molecule microscopy: tutorial, *J. Opt. Soc. Am. A* **33**, B36 (2016).
- [5] C. Helstrom, Resolution of point sources of light as analyzed by quantum detection theory, *IEEE Trans. Inform. Theory* **19**, 389 (1973).
- [6] J. L. Harris, Resolving power and decision theory, *J. Opt. Soc. Am.* **54**, 606 (1964).
- [7] C. O. Acuna and J. Horowitz, A statistical approach to the resolution of point sources, *J. Appl. Statist.* **24**, 421 (1997).
- [8] M. Shahram and P. Milanfar, Statistical and information-theoretic analysis of resolution in imaging, *IEEE Trans. Inform. Theor.* **52**, 3411 (2006).
- [9] M. Tsang, R. Nair, and X.-M. Lu, Quantum theory of superresolution for two incoherent optical point sources, *Phys. Rev. X* **6**, 031033 (2016).
- [10] M. Tsang, Resolving starlight: a quantum perspective, *Contemp. Phys.* **60**, 279 (2019).
- [11] X.-M. Lu, H. Krovi, R. Nair, S. Guha, and J. H. Shapiro, Quantum-optimal detection of one-versus-two incoherent optical sources with arbitrary separation, *npj Quantum Information* **4**, 64 (2018), 1802.02300v1.
- [12] C. W. Helstrom, *Quantum Detection and Estimation Theory* (Academic Press, New York, 1976).
- [13] A. S. Holevo, *Probabilistic and Statistical Aspects of Quantum Theory* (North-Holland, Amsterdam, 1982).
- [14] S. Pirandola, B. R. Bardhan, T. Gehring, C. Weedbrook, and S. Lloyd, Advances in photonic quantum sensing, *Nat. Photonics* **12**, 724 (2018).
- [15] R. Nair and M. Tsang, Interferometric superlocalization of two incoherent optical point sources, *Opt. Express* **24**, 3684 (2016).
- [16] F. Yang, A. Taschilina, E. S. Moiseev, C. Simon, and A. I. Lvovsky, Far-field linear optical superresolution via heterodyne detection in a higher-order local oscillator mode, *Optica* **3**, 1148 (2016).
- [17] M. Paúr, B. Stoklasa, Z. Hradil, L. L. Sánchez-Soto, and J. Rehacek, Achieving the ultimate optical resolution, *Optica* **3**, 1144 (2016).
- [18] W.-K. Tham, H. Ferretti, and A. M. Steinberg, Beating Rayleigh's curse by imaging using phase information, *Phys. Rev. Lett.* **118**, 070801 (2017).
- [19] R. Nair and M. Tsang, Far-field superresolution of thermal electromagnetic sources at the quantum limit, *Phys. Rev. Lett.* **117**, 190801 (2016).
- [20] C. Lupo and S. Pirandola, Ultimate precision bound of quantum and subwavelength imaging, *Phys. Rev. Lett.* **117**, 190802 (2016).
- [21] J. Rehacek, M. Paúr, B. Stoklasa, Z. Hradil, and L. L. Sánchez-Soto, Optimal measurements for resolution beyond the rayleigh limit, *Opt. Lett.* **42**, 231 (2017).
- [22] J. Řeháček, Z. Hradil, B. Stoklasa, M. Paúr, J. Grover, A. Krzic, and L. L. Sánchez-Soto, Multiparameter quantum metrology of incoherent point sources: Towards realistic superresolution, *Phys. Rev. A* **96**, 062107 (2017).
- [23] S. Z. Ang, R. Nair, and M. Tsang, Quantum limit for two-dimensional resolution of two incoherent optical point sources, *Phys. Rev. A* **95**, 063847 (2017).
- [24] A. Chrostowski, R. Demkowicz-Dobrzański, M. Jarzyna, and K. Banaszek, On super-resolution imaging as a multiparameter estimation problem, *International Journal of Quantum Information* **15**, 1740005 (2017), <https://doi.org/10.1142/S0219749917400056>.
- [25] Z. Dutton, R. Kerviche, A. Ashok, and S. Guha, Attaining the quantum limit of superresolution in imaging an object's length via predetection spatial-mode sorting, *Phys. Rev. A* **99**, 033847 (2019).
- [26] M. Tsang, Quantum limit to subdiffraction incoherent optical imaging, *Phys. Rev. A* **99**, 012305 (2019).
- [27] S. Zhou and L. Jiang, Modern description of Rayleigh's criterion, *Phys. Rev. A* **99**, 013808 (2019).
- [28] M. Tsang, Semiparametric estimation for incoherent optical imaging, *Phys. Rev. Research* **1**, 033006 (2019).
- [29] J. Řeháček, M. Paúr, B. Stoklasa, D. Koutný, Z. Hradil, and L. L. Sánchez-Soto, Intensity-based axial localization at the quantum limit, *Phys. Rev. Lett.* **123**, 193601 (2019).
- [30] K. A. G. Bonsma-Fisher, W.-K. Tham, H. Ferretti, and A. M. Steinberg, Realistic sub-rayleigh imaging with phase-sensitive measurements, *New J. Phys.* **21**, 093010

- (2019).
- [31] C. Napoli, S. Piano, R. Leach, G. Adesso, and T. Tufarelli, Towards superresolution surface metrology: Quantum estimation of angular and axial separations, *Phys. Rev. Lett.* **122**, 140505 (2019).
- [32] Y. L. Len, C. Datta, M. Parniak, and K. Banaszek, Resolution limits of spatial mode demultiplexing with noisy detection, *International Journal of Quantum Information* **18**, 1941015 (2020).
- [33] C. Datta, M. Jarzyna, Y. L. Len, K. Lukanowski, J. Kołodyński, and K. Banaszek, Sub-rayleigh resolution of incoherent sources by array homodyning, [arXiv:2005.08693v1](https://arxiv.org/abs/2005.08693v1).
- [34] J. Liu, H. Yuan, X.-M. Lu, and X. Wang, Quantum Fisher information matrix and multiparameter estimation, *J. Phys. A: Math. Theor.* **53**, 023001 (2020).
- [35] C. Helstrom, Minimum mean-squared error of estimates in quantum statistics, *Phys. Lett. A* **25**, 101 (1967).
- [36] C. Helstrom, The minimum variance of estimates in quantum signal detection, *IEEE Trans. Inform. Theory* **14**, 234 (1968).
- [37] M. G. A. Paris, Quantum estimation for quantum technology, *Int. J. Quantum Inf.* **7**, 125 (2009).
- [38] V. P. Belavkin, Generalized uncertainty relations and efficient measurements in quantum systems, *Theor. Math. Phys.* **26**, 213 (1976).
- [39] X.-M. Lu and X. Wang, Incorporating Heisenberg's uncertainty principle into quantum multiparameter estimation, [arXiv:2008.08888](https://arxiv.org/abs/2008.08888).
- [40] J. A. Gross and C. M. Caves, One from many: Estimating a function of many parameters, [arXiv:2002.02898v1](https://arxiv.org/abs/2002.02898v1).
- [41] H. Xing and L. Fu, Measure of the density of quantum states in information geometry and its application in the quantum multi-parameter estimation, [arXiv:2006.00203](https://arxiv.org/abs/2006.00203).
- [42] A. Carollo, B. Spagnolo, A. A. Dubkov, and D. Valenti, On quantumness in multi-parameter quantum estimation, *J. Stat. Mech: Theory Exp.* **2019**, 094010 (2019).
- [43] A. Carollo, D. Valenti, and B. Spagnolo, Geometry of quantum phase transitions, *Phys. Rep.* **838**, 1 (2020).
- [44] J. Miyazaki, Symmetry in metrology: Universal reduction of quantum multi-parameter incompatibility, [arXiv:2010.15465](https://arxiv.org/abs/2010.15465).
- [45] D. Petz, Monotone metrics on matrix spaces, *Linear Algebra Appl.* **244**, 81 (1996).
- [46] F. Hiai and D. Petz, *Introduction to Matrix Analysis and Applications*, 1st ed., Universitext (Springer International Publishing, Cham, 2014).
- [47] X.-M. Lu, S. Yu, and C. H. Oh, Robust quantum metrological schemes based on protection of quantum Fisher information, *Nat. Commun.* **6**, 7282 (2015).
- [48] E. Bisketzi, D. Branford, and A. Datta, Quantum limits of localisation microscopy, *New Journal of Physics* **21**, 123032 (2019).
- [49] M. G. Genoni and T. Tufarelli, Non-orthogonal bases for quantum metrology, *Journal of Physics A: Mathematical and Theoretical* **52**, 434002 (2019).
- [50] M. Soriano and J. J. Palacios, Theory of projections with nonorthogonal basis sets: Partitioning techniques and effective hamiltonians, *Phys. Rev. B* **90**, 075128 (2014).