

Solutions of Heun's general equation and elliptic Darboux equation

(An earlier version in: Math. Meth. Appl. Sci., 44, 7165-7206 (2021))

Bartolomeu D. B. Figueiredo ¹

Centro Brasileiro de Pesquisas Físicas,

Rua Dr. Xavier Sigaud, 150, CEP 22290-180, Rio de Janeiro, RJ, Brasil

Abstract

New solutions for the elliptic Darboux equation are obtained as particular cases of solutions constructed for Heun's general equation. We consider two groups of power series expansions and two new groups of expansions in series of Gauss hypergeometric functions. The convergence of one group in power series is determined by means of ratio tests for infinite series, while the other groups are designed to solve problems which admit finite-series solutions. Actually, we envisage periodic quasi-exactly solvable potentials for which the stationary one-dimensional Schrödinger equation is reduced to the Darboux equation. In general, finite- and infinite-series solutions are obtained from power series expansions for Heun's equation. However, we show that the Schrödinger equation admits additional finite-series expansions in terms of hypergeometric functions for a family of associated Lamé potentials used in band theory of solids. For each finite-series solution we find as well four infinite-series expansions which are bounded and convergent for all values of the independent variable. Finally we find that it is possible to use transformations of variables in order to generate new solutions for the Darboux equation out of other expansions in series of hypergeometric functions for the Heun equation.

Contents

1	Introduction	2
2	Mathematical Preliminaries	5
2.1	Elliptic Darboux equation and the Schrödinger equation	5
2.2	The Gauss hypergeometric function	6
2.3	Some transformations of Heun's general equation	7
2.4	Three-term recurrence relations and convergence	8
3	Power-series solutions	9
3.1	Series $\mathring{H}^{(i)}(x)$ around $x = 0$	10
3.2	Series $\mathring{H}^{(i)}(x)$ around $x = 1$	14
4	Solutions in series of hypergeometric functions	15
4.1	The initial solution $\mathring{H}^{(1)}(x)$	15
4.2	The solutions $H^{(i)}(x)$ around $(a-1)x/(a-x) = 0$	17
4.3	The solutions $\mathbf{H}^{(i)}(x)$ around $x = 1$	19
5	Schrödinger equation for associated Lamé potentials	21
5.1	Finite-series eigenfunctions $(\mathring{\psi}^{(i)}, \tilde{\psi}^{(i)})$ in terms of $\text{sn}^2 u$	23
5.1.1	First type: Even solutions, period $2K$	23
5.1.2	Second type: Odd solutions, period $4K$	24
5.1.3	Third type: Even solutions, period $4K$	25

¹email:barto@cbpf.br

5.1.4	Fourth type: Odd solutions, period $2K$	26
5.2	Finite-series eigenfunction $(\mathring{\Psi}^{(i)}, \tilde{\Psi}^{(i)})$ in terms of $\text{cn}^2 u$	29
5.2.1	First type: even solutions, period $2K$	29
5.2.2	Second type: odd solutions, period $4K$	30
5.2.3	Third type: even solutions, period $4K$	30
5.2.4	Fourth type: odd solutions, period $2K$	31
5.3	Finite series of hypergeometric functions $\psi^{(i)}(u)$ when l is an integer and $m \neq -1/2$ is half an odd integer	33
5.3.1	Finite series when l is an integer and m is half an odd positive integer	34
5.3.2	Finite series when l is an integer and $m \neq -1/2$ is half an odd negative integer	38
5.4	Finite series $\Psi^{(i)}(u)$ of hypergeometric functions when m is an integer and $l \neq -1/2$ is half an odd integer	39
5.4.1	Finite series when m is an integer and l is half an odd positive integer	39
5.4.2	Finite series when m is an integer and $l \neq -1/2$ is half an odd negative integer	40
5.5	Infinite-series solutions	41
6	Conclusions	43
A	Homotopic transformations	45
B	Cases solved by hypergeometric functions	45
C	Some degenerate power-series solutions	48
D	Svartholm's and Erdelyi's solutions	51
	References	54

1 Introduction

We find new solutions for the elliptic Darboux equation from solutions of Heun's general equation since this includes the former equation as a particular case. By using known substitutions of variables for Heun's equation, we construct two groups of power series expansions and two groups of expansions in series of Gauss hypergeometric functions. Finally, we find that these groups are sufficient to solve systematically the Darboux equations which emerge from the stationary one-dimensional Schrödinger equation for some periodic quasi-exactly solvable (QES) potentials.

The elliptic form of the Darboux equation can be written as [1, 2]

$$\frac{d^2 U(u)}{du^2} + \left[h - \mu(\mu + 1)k^2 \text{sn}^2 u - \frac{\nu_1(\nu_1 + 1)}{\text{sn}^2 u} - \frac{\nu_2(\nu_2 + 1) \text{dn}^2 u}{\text{cn}^2 u} - \frac{\lambda(\lambda + 1)k^2 \text{cn}^2 u}{\text{dn}^2 u} \right] U(u) = 0, \quad (1)$$

where h , μ , ν_1 , ν_2 and λ are constants, while $\text{sn } u$, $\text{cn } u$ and $\text{dn } u$ are the Jacobian elliptic functions [3] modulus k ($0 < k^2 < 1$). Eq. (1) reduces to the associated Lamé equation when $\nu_1(\nu_1 + 1) = \nu_2(\nu_2 + 1) = 0$, and to the Lamé equation when $\nu_1(\nu_1 + 1) = \nu_2(\nu_2 + 1) = \lambda(\lambda + 1) = 0$.

The Darboux equation is a particular case of the Heun equation which, in an algebraic form [4, 5], reads

$$\frac{d^2 H(x)}{dx^2} + \left[\frac{\gamma}{x} + \frac{\delta}{x-1} + \frac{\epsilon}{x-a} \right] \frac{dH(x)}{dx} + \frac{\alpha\beta x - q}{x(x-1)(x-a)} H(x) = 0, \quad \epsilon = \alpha + \beta + 1 - \gamma - \delta, \quad (2)$$

where $a \in \mathbb{C} \setminus \{0, 1\}$, and the points $x = 0, 1, a, \infty$ are regular singularities. If $a = 0$ or $a = 1$ the above equation can be reduced to the Gauss hypergeometric equation [6]

$$z(1-z)\frac{d^2 Z(z)}{dz^2} + [c - (a+b+1)z]\frac{dZ(z)}{dz} - ab Z(z) = 0, \quad (3)$$

which has only three singular points. In fact, there are several cases in which the Heun equation is reducible to the hypergeometric equation [7, 8]; some of these are written in Appendix B because they are relevant for this article.

We select the form (2) because it is the one used for studying the substitutions of variables which keep invariant the form of the Heun equation [5]. These substitutions result from 8 homotopic transformations (no change of the independent variable) and 24 linear fractional transformations of the independent variable. As in the case of the hypergeometric equation, the substitutions can also be used to transform an initial solution into new solutions which, for the Darboux equation, can be rewritten in terms of elliptic functions. Thus, from an initial solution, the homotopic transformations generate a group containing 8 expansions. New groups can be generated by firstly applying on the initial solution a fractional transformation of the independent variable.

As initial solutions for Heun's equation we take a familiar power series around the origin of the independent variable [4, 9] and an expansion in series of Gauss hypergeometric functions. As a matter of fact, in Section 3 we consider two groups of power series whose initial solutions are the expansions having the form

$$\mathring{H}^{(1)}(x) = \sum_{n=0}^{\infty} \mathring{b}_n^{(1)} x^n \quad \text{and} \quad \mathring{H}^{(1)}(x) = \sum_{n=0}^{\infty} \mathring{b}_n^{(1)} (1-x)^n, \quad (4)$$

around the points $x = 0$ and $x = 1$. We will find that $\mathring{H}^{(1)}$ results from $\mathring{H}^{(1)}$ by means of the linear fractional transformation $x \mapsto 1-x$. Furthermore, the homotopic transformations lead to 16 expansions, $\mathring{H}^{(i)}$ and $\mathring{H}^{(i)}$ ($i = 1, 2, \dots, 8$). The mathring symbol $\mathring{}$ over $\mathring{H}^{(i)}$ and $\mathring{H}^{(i)}$ distinguishes these expansions from the expansions in series of hypergeometric functions, $H^{(i)}$ and $H^{(i)}$.

On the other side, in Section 4 we consider two groups of expansions in series of hypergeometric functions, $F(a, b; c; z)$, generated from

$$H^{(1)}(x) = \left(1 - \frac{x}{a}\right)^{-\alpha} \sum_{n=0}^{\infty} \frac{b_n^{(1)}}{\Gamma(n+\gamma)} \left[\frac{(a-1)x}{a-x}\right]^n F\left[n+\alpha, \gamma+\delta-\alpha-1; n+\gamma; \frac{(a-1)x}{a-x}\right], \quad (5)$$

$$H^{(1)}(x) = \sum_{n=0}^{\infty} \frac{b_n^{(1)}}{\Gamma(n+\delta)} (1-x)^n F(n+\alpha, \gamma+\delta-\alpha-1; n+\delta; 1-x). \quad (6)$$

The expansions (5) and (6) will be obtained from an expansion having the form

$$\bar{H}^{(1)}(x) = \sum_{n=0}^{\infty} \bar{b}_n^{(1)} x^n F(n+\alpha, \gamma+\delta-\alpha-1; n+\gamma; x). \quad (7)$$

In effect, we will find that the linear fractional substitutions $x \mapsto (a-1)x/(a-x)$ and $x \mapsto 1-x$ transform $\bar{H}^{(1)}$ into $H^{(1)}$ and $H^{(1)}$, respectively. The homotopic transformations lead once again to 16 expansions, $H^{(i)}$ and $H^{(i)}$. The arguments z of the above $F(a, b; c; z)$ are appropriate for the Darboux equation because for this case $|z| \leq 1$, as we will see. For some constraints between the parameters, the hypergeometric-function expansions reduce to power-series expansions; for example, $\bar{H}^{(1)} = \mathring{H}^{(1)}$ if $\gamma + \delta = \alpha + 1$. We will see as

well that the initial solutions, $\mathring{H}(x)$ and $\bar{H}^{(1)}(x)$, lead to known solutions for the confluent Heun equation.

The power-series and hypergeometric-function expansions are both given by one-sided infinite series ($n \geq 0$) whose coefficients obey three-term recurrence relations. These relations constitute a system of homogeneous linear equations which admits non trivial solutions only if the determinant of the corresponding matrix vanish (characteristic equation). This fact requires a free parameter in the differential equation. The absence of free parameters would demand two-sided infinite series ($-\infty < n < +\infty$) having a parameter which is not present in the differential equation, as in the case of Mathieu equation (see chapter VI of Ref. [3]).

Under certain conditions, the one-sided series truncate on the right at $n = N$, becoming finite series with $0 \leq n \leq N$. For finite series the convergence is decided by examining each term of the series, provided that the characteristic equation is satisfied. For infinite series the convergence requires some ratio test between successive terms of the series. In the present article, only the power expansions $\mathring{H}^{(i)}$ are intended to be used as infinite-series solutions; by this reason, there is no concern about ratio tests for the other expansions.

Finite-series solutions occur in quasi-exactly solvable (QES) problems of quantum mechanics [10, 11], for which one part of the energy spectrum and the corresponding eigenfunctions can be computed explicitly. In fact, a QES problem is sometimes defined as a problem which presents solutions given by finite series with coefficients satisfying three-term or higher order recurrence relations [12]. The “quasi-exactly solvable” part of the spectrum is obtained from the finite-series solutions, but there are also infinite-series solutions which are convergent and bounded, as we will see.

In Section 2 we explain how to get solutions for the Darboux equation from solutions of the Heun equation and show that, for some elliptic potentials, the Schrödinger equation reduces to the Darboux equation. In addition, we collect mathematical informations used in subsequent sections: properties of the hypergeometric functions, transformations of the Heun equation and some aspects of the three-term recurrence relations for the series coefficients.

In Section 3 we present the power series solutions $\mathring{H}^{(i)}$ and $\mathring{\mathbf{H}}^{(i)}$, while in Section 4 we construct the expansions $H^{(i)}$ and $\mathbf{H}^{(i)}$ in series in hypergeometric functions. The solutions of the four groups are generated straightforward by applying substitutions of variable on the expansion $\mathring{H}^{(1)}(x)$ written in (4), and on $\bar{H}^{(1)}(x)$ written in (7). Despite this, all the expansions are written explicitly for the purposes of applications.

The largest part of the present study is Section 5 which deals with a family of QES potentials which appear in band theory of solids. In this case, the equation reduces to an associated Lamé equation and admits finite-series solutions resulting from power series expansions and from expansions in series of hypergeometric functions. In addition to finite-series eigenfunctions (which characterize the quasi-exact solvability) we get as well infinite-series eigenfunctions, convergent and bounded for any value of the independent variable.

Final remarks and additional references are in Section 6, Appendix A gives the homotopic transformations of the Heun equation, while in Appendix B the Heun equation is reduced to hypergeometric equation when its parameters satisfy certain constraints. In Appendix C, we write degenerate solutions of the Schrödinger equation for some particular values of the parameters of the associated Lamé potential. Eventually, in Appendix D we discuss the possibility of developing new solutions for the Darboux equation by starting from expansions in series of hypergeometric functions (for the Heun equation) given by Svartholm [13] and by Erdélyi [14, 15].

2 Mathematical Preliminaries

In this section the Darboux equation is written as a particular case of the Heun equation, while the Schrödinger equation is written as a Darboux equation for four known elliptic potentials. We write also some properties of the Gauss hypergeometric functions and select the transformations of variables of the Heun equation which will be used to generate solutions appropriate for the Darboux equation. Finally we regard three-term recurrence relations for the series coefficients and state a known theorem which gives conditions sufficient for assuring real and distinct solutions for the characteristic equation of finite series.

2.1 Elliptic Darboux equation and the Schrödinger equation

The basic Jacobian elliptic functions $\text{sn}(u, k)$, $\text{cn}(u, k)$ and $\text{dn}(u, k)$ are written as $\text{sn } u$, $\text{cn } u$ and $\text{dn } u$, respectively [3, 16]. Here the parameter k , called modulus of the functions, is supposed to be a real number such that $0 < k^2 < 1$. These functions satisfy the relations

$$\text{sn}^2 u + \text{cn}^2 u = k^2 \text{sn}^2 u + \text{dn}^2 u = 1 \Rightarrow (k^2 - 1) \text{sn}^2 u = \text{cn}^2 u - \text{dn}^2 u, \quad 1 - k^2 = \text{dn}^2 u - k^2 \text{cn}^2 u. \quad (8)$$

Beside these, we will use the functions $\text{cd } u$ and $\text{sd } u$, defined by

$$\text{cd } u = \text{cn } u / \text{dn } u, \quad \text{sd } u = \text{sn } u / \text{dn } u. \quad (9)$$

Now we obtain the Darboux equation from the Heun equation (2) by taking $a = k^{-2}$ and achieving the substitutions

$$H[x(u)] = (\text{sn}^2 u)^{\frac{1-2\gamma}{4}} (\text{cn}^2 u)^{\frac{1-2\delta}{4}} (\text{dn}^2 u)^{\frac{1-2\epsilon}{4}} U(u), \quad x(u) = \text{sn}^2 u, \quad [a = k^{-2}]. \quad (10)$$

which lead to Eq. (1) expressed in terms of the parameters of the Heun equation, namely,

$$\frac{d^2 U(u)}{du^2} + \left\{ h - \left[(\alpha - \beta)^2 - \frac{1}{4} \right] k^2 \text{sn}^2 u - \left(\gamma - \frac{1}{2} \right) \left(\gamma - \frac{3}{2} \right) \frac{1}{\text{sn}^2 u} - \right. \\ \left. \left(\delta - \frac{1}{2} \right) \left(\delta - \frac{3}{2} \right) \frac{\text{dn}^2 u}{\text{cn}^2 u} - \left(\epsilon - \frac{1}{2} \right) \left(\epsilon - \frac{3}{2} \right) \frac{k^2 \text{cn}^2 u}{\text{dn}^2 u} \right\} U(u) = 0, \quad [\text{Darboux Eq.}] \quad (11)$$

where $h = (\gamma + \delta)^2 + 1 - 2\gamma - 2\delta - [4q - (\gamma + \epsilon)^2 - 1 + 2\gamma + 2\epsilon] k^2$. Therefore, solutions $U(u)$ for the Darboux equation (11) follow from solutions $H(x)$ for the Heun equation through (10). The associated Lamé equation and the Lamé equation occur when, in Eq. (11),

$$(\gamma, \delta) = \left(\frac{1}{2}, \frac{1}{2} \right), \left(\frac{1}{2}, \frac{3}{2} \right), \left(\frac{3}{2}, \frac{1}{2} \right), \left(\frac{3}{2}, \frac{3}{2} \right) \quad [\text{associated Lamé equation}]; \quad (12)$$

$$(\gamma, \delta, \epsilon) = \left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2} \right), \left(\frac{1}{2}, \frac{1}{2}, \frac{3}{2} \right), \left(\frac{1}{2}, \frac{3}{2}, \frac{1}{2} \right), \left(\frac{1}{2}, \frac{3}{2}, \frac{3}{2} \right), \\ \left(\frac{3}{2}, \frac{1}{2}, \frac{1}{2} \right), \left(\frac{3}{2}, \frac{1}{2}, \frac{3}{2} \right), \left(\frac{3}{2}, \frac{3}{2}, \frac{1}{2} \right), \left(\frac{3}{2}, \frac{3}{2}, \frac{3}{2} \right) \quad [\text{Lamé equation}]. \quad (13)$$

Next we regard the one-dimensional stationary Schrödinger equation,

$$\frac{d^2 \psi(u)}{du^2} + [\mathcal{E} - \mathcal{V}(u)] \psi(u) = 0, \quad u = \kappa x, \quad \mathcal{E} = \frac{2M}{\hbar^2 \kappa^2} E, \quad \mathcal{V}(u) = \frac{2M}{\hbar^2 \kappa^2} V(x), \quad (14)$$

for a particle with mass M and energy E – the constant κ has inverse-length dimension, \hbar is the Plank constant divided by 2π , x is the spatial coordinate and $V(x)$ is the potential. Eq.

(14) reduces to the Darboux equation (11), with $\psi(u) = U(u)$, for the following potentials $\mathcal{V}_i(u)$ ($i = 1, 2, 3, 4$). The three Ganguly's potentials [17, 18]:

$$\mathcal{V}_1(u) = (1 - k^2) \left[\frac{2}{\text{cn}^2 u} - \frac{(\mathbf{l}+2)(\mathbf{l}+3)}{\text{dn}^2 u} \right] \stackrel{\text{Eq. (8)}}{=} -2k^2 - (\mathbf{l}+2)(\mathbf{l}+3) + 2 \frac{\text{dn}^2 u}{\text{cn}^2 u} + (\mathbf{l}+2)(\mathbf{l}+3) \frac{k^2 \text{cn}^2 u}{\text{dn}^2 u}. \quad (15)$$

$$\mathcal{V}_2(u) = \frac{2}{\text{sn}^2 u} - \frac{(1-k^2)(\mathbf{l}+2)(\mathbf{l}+3)}{\text{dn}^2 u} \stackrel{(8)}{=} -(\mathbf{l}+2)(\mathbf{l}+3) + \frac{2}{\text{sn}^2 u} + (\mathbf{l}+2)(\mathbf{l}+3) \frac{k^2 \text{cn}^2 u}{\text{dn}^2 u}. \quad (16)$$

$$\mathcal{V}_3(u) = \mathbf{m}(\mathbf{m}+1) k^2 \text{sn}^2 u + \mathbf{l}(\mathbf{l}+1) \frac{k^2 \text{cn}^2 u}{\text{dn}^2 u}, \quad (17)$$

where \mathbf{l} and \mathbf{m} are constants. For the potential $\mathcal{V}_3(u)$ the Schrödinger equation becomes an associated Lamé equation which was firstly considered by Khare and Sukhatme [19, 20]. On the other side, there is the Ushveridze potential [11]

$$\begin{aligned} \mathcal{V}_4(u) = & -4 \left(\mathbf{a} - \frac{1}{4} \right) \left(\mathbf{a} - \frac{3}{4} \right) \text{dn}^2 u + 4 \left(\mathbf{b} - \frac{1}{4} \right) \left(\mathbf{b} - \frac{3}{4} \right) \frac{\text{dn}^2 u}{\text{sn}^2 u} + 4 \left(\mathbf{c} - \frac{1}{4} \right) \left(\mathbf{c} - \frac{3}{4} \right) \frac{\text{dn}^2 u}{\text{cn}^2 u} \\ & + 4 \left(\mathbf{a} + \mathbf{b} + \mathbf{c} + \mathbf{l} - \frac{1}{4} \right) \left(\mathbf{a} + \mathbf{b} + \mathbf{c} + \mathbf{l} - \frac{3}{4} \right) k^2 (k^2 - 1) \frac{\text{sn}^2 u}{\text{dn}^2 u}, \end{aligned}$$

where \mathbf{a} , \mathbf{b} , \mathbf{c} and \mathbf{l} are real constants. By rewriting \mathcal{V}_4 as

$$\begin{aligned} \mathcal{V}_4(u) \stackrel{(8)}{=} & -4 \left(\mathbf{a} - \frac{1}{4} \right) \left(\mathbf{a} - \frac{3}{4} \right) - 4 (\mathbf{a} + \mathbf{c} + \mathbf{l}) (\mathbf{a} + 2\mathbf{b} + \mathbf{c} + \mathbf{l} - 1) k^2 \\ & - 8 \left(\mathbf{b} - \frac{1}{4} \right) \left(\mathbf{b} - \frac{3}{4} \right) k^2 + 4 \left(\mathbf{a} - \frac{1}{4} \right) \left(\mathbf{a} - \frac{3}{4} \right) k^2 \text{sn}^2 u + 4 \left(\mathbf{b} - \frac{1}{4} \right) \left(\mathbf{b} - \frac{3}{4} \right) \frac{1}{\text{sn}^2 u} \\ & + 4 \left(\mathbf{c} - \frac{1}{4} \right) \left(\mathbf{c} - \frac{3}{4} \right) \frac{\text{dn}^2 u}{\text{cn}^2 u} + 4 \left(\mathbf{a} + \mathbf{b} + \mathbf{c} + \mathbf{l} - \frac{1}{4} \right) \left(\mathbf{a} + \mathbf{b} + \mathbf{c} + \mathbf{l} - \frac{3}{4} \right) \frac{k^2 \text{cn}^2 u}{\text{dn}^2 u}, \end{aligned} \quad (18)$$

the Schrödinger equation (14) once more takes the form (11) for the Darboux equation.

Thus, if we know solutions $H(x)$ for the Heun equation, the eigenfunctions $\psi(u)$ follow from

$$\psi(u) = U(u) \stackrel{(10)}{=} (\text{sn} u)^{\frac{2\gamma-1}{2}} (\text{cn} u)^{\frac{2\delta-1}{2}} (\text{dn} u)^{\frac{2\epsilon-1}{2}} H(x), \quad [a = k^{-2}, x = \text{sn}^2 u], \quad (19)$$

provided that these $\psi(u)$ are bounded for any admissible value of the variable u . The problem is QES if there are solutions $H(x)$ given by finite series, a fact which occurs when the parameters of the Schrödinger equation satisfy certain conditions. For the associated Lamé potential (17) the conditions stated in [20] are: (i) either $\mathbf{m} - \mathbf{l} \neq 0$ or $\mathbf{l} + \mathbf{m}$ is an integer, (ii) either \mathbf{l} or \mathbf{m} is an integer and the other is half an odd integer. In addition, the second case admits doubly degenerate eigenfunctions, that is, two independent eigenfunctions with the same value for the energy.

2.2 The Gauss hypergeometric function

The hypergeometric function $F(a, b; c; z)$, solution of Eq. (3), is defined by the series [6]

$$F(a, b; c; z) = F(b, a; c; z) = 1 + \frac{ab}{1! c} z + \frac{a(a+1)b(b+1)}{2! c(c+1)} z^2 + \dots = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{n! (c)_n} z^n,$$

where $(a)_n$, $(b)_n$ and $(c)_n$ are Pochhammer symbols: $(a)_0 = 1$, \dots , $(a)_n = a(a+1) \dots (a+n-1) = \Gamma(a+n)/\Gamma(a)$, etc. The above series is not defined when the parameter c is zero or negative integer. Besides this, if the parameter a or the parameter b is zero or negative integer, the series reduces to a polynomial. A infinite series $F(a, b; c; z)$ converges

absolutely for $|z| < 1$ and diverges for $|z| > 1$; by the Raabe test, the series converges on the circle $|z| = 1$ if $\operatorname{Re}(c - a - b) > 0$. Thus,

$$\begin{aligned} F(a, b; c; z) & \quad \text{converges absolutely for } |z| < 1, \text{ diverges for } |z| > 1, \\ & \quad \text{converges also on } |z| = 1 \text{ if } \operatorname{Re}(c - a - b) > 0. \end{aligned} \quad (20)$$

The following relations will be useful:

$$\begin{aligned} F(a, b; c; z) &= (1 - z)^{c-a-b} F(c - a, c - b; c; z) \quad \text{for } |z| < 1, \\ F(a, b; c; 1) &= \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)} \quad \text{if } \operatorname{Re}(c-a-b) > 0. \end{aligned} \quad (21)$$

In Section 4 we also use the relations [6]

$$\begin{aligned} \frac{d}{dz} [z^{c-1} (1 - z)^{b-c+1} F(a, b; c; z)] &= (c - 1) z^{c-2} (1 - z)^{b-c} F(a - 1, b; c - 1; z), \\ \frac{d}{dz} [(1 - z)^a F(a, b; c; z)] &= -\frac{a(c-b)}{c} (1 - z)^{a-1} F(a + 1, b; c + 1; z), \end{aligned} \quad (22)$$

which imply

$$\begin{aligned} (1 - z) \frac{d}{dz} F(a, b; c; z) &= \left(b - \frac{c-1}{z}\right) F(a, b; c; z) + \frac{c-1}{z} F(a - 1, b; c - 1; z), \\ (z - 1) \frac{d}{dz} F(a, b; c; z) &= -a F(a, b; c; z) + \frac{a(c-b)}{c} F(a + 1, b; c + 1; z). \end{aligned} \quad (23)$$

2.3 Some transformations of Heun's general equation

In 2007 Maier set down the 192 substitutions of variables which preserve the form of Heun's equation. The table 2 of his paper [5] displays the transformations of each parameter and variable of the equation. These transformations follow from the composition of 8 homotopic substitutions of the dependent variable $H(x)$ (which keep unaltered the independent variable x) with 24 linear fractional substitutions $x \mapsto y(x) = (Ax + B)/(Cx + D)$ (in general these demand a transformation of the independent variable as well). We rewrite some of these substitutions in a form convenient for applications.

We denote by $H(x) = H(q, a; \alpha, \beta, \gamma, \delta; x)$ a solution of the Heun equation (2); on the other hand, the transformations of $H(x)$ are indicated by the symbol M_i following the order in which they appear in Maier's table ($i = 1, 2, \dots, 192$). Hence, each M_i transform $H(x)$ into $M_i H(x)$ according to

$$M_i H(x) = M_i H(a, q; \alpha, \beta, \gamma, \delta; x) = f_i(x) H\left[\tilde{a}, \tilde{q}; \tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}, \tilde{\delta}; y(x)\right]. \quad (24)$$

where $f_i(x)$ and $\left[\tilde{q}, \tilde{a}; \tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}, \tilde{\delta}; y(x)\right]$ depend on the transformation considered. For example, the fractional substitutions

$$\frac{(1-a)x}{x-a}, \quad 1 - x, \quad \frac{a(x-1)}{x-a} \quad (25)$$

may be represented by M_{17} , M_{49} and M_{65} , that is,

$$M_{17} H(x) = \left(1 - \frac{x}{a}\right)^{-\alpha} H\left[1 - a, -q + \alpha\gamma; \alpha, -\beta + \gamma + \delta, \gamma, \delta; \frac{(1-a)x}{x-a}\right], \quad (26)$$

$$M_{49} H(x) = H(1 - a, -q + \alpha\beta; \alpha, \beta, \delta, \gamma; 1 - x), \quad (27)$$

$$M_{65} H(x) = \left(1 - \frac{x}{a}\right)^{-\alpha} H\left[a, q - \alpha(\beta - \delta); \alpha, -\beta + \gamma + \delta, \delta, \gamma; \frac{a(x-1)}{x-a}\right]. \quad (28)$$

As a matter of fact, for each substitution of the independent variable, there are 8 substitutions of the dependent variable on account of the homotopic transformations. These are given by the transformations $M_1 - M_4$ and $M_{25} - M_{28}$ of Maier's table but here, as in [21] and [22], they are denoted by T_i ($i = 1, \dots, 8$) according to

$$\begin{aligned} T_1 &= M_1, & T_2 &= M_{25}, & T_3 &= M_2, & T_4 &= M_{26}, \\ T_5 &= M_3, & T_6 &= M_{27}, & T_7 &= M_4, & T_8 &= M_{28}. \end{aligned} \quad (29)$$

The 8 transformations are written in Appendix A.

Notice that for the Darboux equation (11)

$$x = \text{sn}^2 u, \quad 1 - x = \text{cn}^2 u, \quad \frac{(a-1)x}{a-x} = (1 - k^2) \text{sd}^2 u, \quad \frac{a(x-1)}{x-a} = \text{cd}^2 u, \quad (30)$$

where $\text{sd} u$ and $\text{cd} u$ are the functions given in (9). From definitions [3, 16], we know that $0 \leq \text{sn}^2 u \leq 1$, $0 \leq \text{cn}^2 u \leq 1$ and $0 < \text{dn}^2 u < 1$. Further, the relations [3]

$$\begin{aligned} \text{cd} u &= \text{sn}(u + K) \quad \text{and} \quad (1 - k^2) \text{sd}^2 u = 1 - \text{cd}^2 u, \\ \text{where } K &= K(k) = \int_0^{\frac{\pi}{2}} [1 - k^2 \sin^2 \theta]^{-\frac{1}{2}} d\theta, \end{aligned} \quad (31)$$

assure that $0 \leq \text{cd}^2 u \leq 1$ and $0 \leq (1 - k^2) \text{sd}^2 u \leq 1$. Thus, the variables (30) are inside the interval $[0, 1]$ which is suitable for arguments of the hypergeometric functions, according to (20). Similarly, the variables x/a and $1 - (x/a)$ are also appropriate. As we are interested in Darboux equation, in sections III and IV we shall use the homotopic transformations and the transformations M_{17} and M_{49} given in Eqs. (26) and (27).

Therefore, in most cases we will use the substitutions of variables as a procedure to generate new solutions out of a given solution $H(x)$. However, the substitutions can be used to simply transform the Heun equation (2) into another version of itself. We will use this viewpoint in Appendix B in order to show that, for certain values of the parameters, the Heun equation (2) is transformed into an equation reducible to the hypergeometric equation (3).

2.4 Three-term recurrence relations and convergence

The following are known results which are also valid for the confluent cases of the Heun equation [21]. The three-term recurrence relations for the coefficients b_n of one-sided solutions have the form

$$\alpha_0 b_1 + \beta_0 b_0 = 0, \quad \alpha_n b_{n+1} + \beta_n b_n + \gamma_n b_{n-1} = 0 \quad (n \geq 1), \quad (32)$$

where α_n , β_n and γ_n are functions of the parameters of the equation. This system of homogeneous linear equations can be written in matrix form. Then, we see that the system has non-trivial solutions only if the determinant of a tridiagonal coefficient matrix vanishes; such condition is called 'characteristic equation' and determine the possible values of some 'free' parameter of the equation. The characteristic equation is also written as the continued fraction [23, 24]

$$\beta_0 = \frac{\alpha_0 \gamma_1}{\beta_1 -} \frac{\alpha_1 \gamma_2}{\beta_2 -} \frac{\alpha_2 \gamma_3}{\beta_3 -} \dots \quad (33)$$

A series whose coefficients satisfy three-term recurrence relations like (32) truncates on the right-hand side if $\gamma_n = 0$ for some value of $n \geq$. Precisely, if $\gamma_{N+1} = 0$, the series becomes a finite series with $0 \leq n \leq N$ [25]). So, we can write

$$\gamma_{N+1} = 0 \quad \Rightarrow \quad \text{finite series with } n \leq N. \quad (34)$$

For finite series, the characteristic equation is equivalent to the vanishing of the determinant of a $(N + 1) \times (N + 1)$ matrix. Precisely

$$\det \begin{bmatrix} \beta_0 & \alpha_0 & 0 & & \\ \gamma_1 & \beta_1 & \alpha_1 & & \\ & \ddots & \ddots & \ddots & \\ & & \gamma_{N-1} & \beta_{N-1} & \alpha_{N-1} \\ & & 0 & \gamma_N & \beta_N \end{bmatrix} = 0 \quad (35)$$

assures that there are nontrivial solutions for the coefficients b_n ($n = 0, 1, \dots, N$).

The following theorem, due to Arscott [25, 26], holds for finite series:

Let $\beta_i = \mathcal{B}_i - \Lambda$ for $i = 0, 1, \dots, N$, where Λ is an arbitrary constant. If α_i , \mathcal{B}_i and γ_i are all real and independent of Λ , and if each product $\alpha_{i-1} \gamma_i$ is positive, that is,

$$\text{if } \alpha_i, \mathcal{B}_i, \gamma_i \text{ are real and independent of } \Lambda, \text{ and if } \alpha_{i-1} \gamma_i > 0 \text{ for } i = 1, \dots, N, \quad (36)$$

then the vanishing of the determinant of the above matrix affords $N + 1$ real and distinct values for Λ ; if $\gamma_i \alpha_{i-1} \leq 0$ nothing can be said about the nature of Λ .

The previous theorem will be useful to analyse the energy spectra of finite-series wave-functions of quasi-exactly solvable problems, as the one discussed in Section 5. In these cases the characteristic equation becomes an algebraic equation of order $N + 1$ in Λ , where Λ is a linear function of the energy \mathcal{E} . For each value of \mathcal{E} , the series coefficients can be computed by a forward application of the recurrence relations, from b_0 upward to b_N ; this gives all the b_n in terms of b_0 . In this case the convergence is decided by examining each term of the series.

On the other side, the convergence of infinite-series solutions comes from ratio tests for successive terms of the series [27, 28]. In general the three-term relations (32) present two linearly independent solutions [9, 24] and, consequently, two ratios for b_{n+1}/b_n but, in order to assure the convergence the continued fraction (33), we have to choose the solution that Gautschi has called minimal (or minimal at infinity): if f_n and g_n denote such solutions and $\lim_{n \rightarrow \infty} (f_n/g_n) = 0$, then f_n is the minimal solution. Alternatives to the term *minimal* are give in a footnote on page 25 of Gautschi's paper [24]. For practical purposes, if

$$\lim_{n \rightarrow \infty} (b_{n+1}/b_n) = t_1, \quad \lim_{n \rightarrow \infty} (b_{n+1}/b_n) = t_2, \quad t_1 \neq t_2, \quad (37)$$

then a Poincaré-Perron theorem [24] assures the existence of two linearly independent solutions presenting these limits, the minimal solution being the one with smaller modulus for the limit, $|t_1|$ or $|t_2|$.

The use of a minimal solution demands a backward procedure to compute the ratios b_{n+1}/b_n , from a large value of n downward to $n = 0$. However, before this, it is necessary to solve the characteristic equation which affords relations between the parameters of the Heun equation. Some considerations concerning solutions of characteristic equation for infinite series are given in [29, 30, 31]. We just mention that solving Eq. (33) is an important part of the problem since that equation determines, for example, the energy spectra of hydrogen molecular ion (confluent Heun equation).

3 Power-series solutions

This section regards the two groups of power series solutions, $\mathring{H}^{(i)}(x)$ around $x = 0$ and $\mathring{H}^{(i)}(x)$ around $x = 1$ ($i = 1, \dots, 8$), each group being closed with respect to the homotopic transformations (A.1-A.8). The series coefficients are denoted by $\mathring{b}_n^{(i)}$ and $\mathring{b}_n^{(i)}$, respectively.

The starting-point solution $\mathring{H}^{(1)}(x)$ generates the others by means of variable transformations as follows

$$\mathring{H}^{(i)}(x) = T_i \mathring{H}^{(1)}(x), \quad \mathring{\mathbf{H}}^{(1)}(x) = M_{49} \mathring{H}^{(1)}(x), \quad \mathring{\mathbf{H}}^{(i)}(x) = T_i \mathring{\mathbf{H}}^{(1)}(x). \quad (38)$$

Thus, the present section uses only the Möbius substitution M_{49} ($x \mapsto 1 - x$) written in (27). In order to determine the convergence of $\mathring{H}^{(i)}$ for infinite series it is necessary to suppose that $|a| \neq 1$. It will be seen as well that (for a fixed value of i) both series $\mathring{H}^{(i)}$ and $\mathring{\mathbf{H}}^{(i)}$ reduce to finite series under the same circumstances since $\mathring{\gamma}_n^{(i)} = \mathring{\gamma}_n^{(i)}$ in condition (34); however, one of the series may satisfy the Arscott condition (36) while the other may not satisfy.

All solutions $\mathring{H}^{(i)}$ and $\mathring{\mathbf{H}}^{(i)}$ will be used in Section 5. The convergence of infinite series $\mathring{H}^{(i)}$ holds for minimal solutions of the recurrence relations for the coefficients $\mathring{b}_n^{(i)}$. At the end of Section 3.1, we find conditions for linear dependence between the expansions $\mathring{H}^{(i)}$ and $\mathring{H}^{(i+4)}$ for $i = 1, 2, 3, 4$. We show as well that the expansion $\mathring{H}^{(1)}$, by means of a confluence procedure, leads to a known expansion for solutions of the confluent Heun equation (CHE).

3.1 Series $\mathring{H}^{(i)}(x)$ around $x = 0$

By using the homotopic transformations T_i , the solutions around $x = 0$ are written as

$$\mathring{H}^{(1)}(x) = \sum_{n=0}^{\infty} \mathring{b}_n^{(1)} x^n, \quad \mathring{H}^{(i)}(x) = T_i \mathring{H}^{(1)}(x). \quad (39)$$

The coefficients of the recurrence relations (32) for $\mathring{b}_n^{(1)}$ are [21]

$$\begin{aligned} \mathring{\alpha}_n^{(1)} &= a(n+1)(n+\gamma), \quad \mathring{\beta}_n^{(1)} = -an(n+\gamma+\delta-1) - n(n+\alpha+\beta-\delta) - q, \\ \mathring{\gamma}_n^{(1)} &= (n+\alpha-1)(n+\beta-1). \end{aligned} \quad (40)$$

Dividing these relations by $n^2 \mathring{b}_n^{(1)}$, we get

$$\begin{aligned} a \left[1 + \frac{1+\gamma}{n} + O\left(\frac{1}{n^2}\right) \right] \frac{\mathring{b}_{n+1}^{(1)}}{\mathring{b}_n^{(1)}} - \left[a + 1 + \frac{a(\gamma+\delta-1)+\alpha+\beta-\delta}{n} + O\left(\frac{1}{n^2}\right) \right] + \\ \left[1 + \frac{\alpha+\beta-2}{n} + O\left(\frac{1}{n^2}\right) \right] \frac{\mathring{b}_{n-1}^{(1)}}{\mathring{b}_n^{(1)}} = 0. \end{aligned}$$

Then, the ratios $\mathring{b}_{n+1}^{(1)}/\mathring{b}_n^{(1)}$ for large n are

$$\frac{\mathring{b}_{n+1}^{(1)}}{\mathring{b}_n^{(1)}} \sim \frac{1}{a} \left[1 + \frac{\epsilon-2}{n} \right] \Leftrightarrow \frac{\mathring{b}_{n-1}^{(1)}}{\mathring{b}_n^{(1)}} \sim a \left[1 - \frac{\epsilon-2}{n} \right] \quad \text{and} \quad \frac{\mathring{b}_{n+1}^{(1)}}{\mathring{b}_n^{(1)}} \sim 1 + \frac{\delta-2}{n} \Leftrightarrow \frac{\mathring{b}_{n-1}^{(1)}}{\mathring{b}_n^{(1)}} \sim 1 - \frac{\delta-2}{n}.$$

Thus, $\lim_{n \rightarrow \infty} \mathring{b}_{n+1}^{(1)}/\mathring{b}_n^{(1)}$ is $\frac{1}{a}$ or 1. If $|a| \neq 1$, the two limits are distinct and by the Poincaré-Perron theorem, mentioned after Eq. (37),

$$\text{for minimal solution: } \frac{\mathring{b}_{n+1}^{(1)}}{\mathring{b}_n^{(1)}} \sim \frac{1}{a} \left[1 + \frac{\epsilon-2}{n} \right] \quad \text{if } |a| > 1, \quad \frac{\mathring{b}_{n+1}^{(1)}}{\mathring{b}_n^{(1)}} \sim 1 + \frac{\delta-2}{n} \quad \text{if } |a| < 1. \quad (41)$$

Thus, for minimal solution, the regions of convergence follow from

$$\left| \frac{\mathring{b}_{n+1}^{(1)} x^{n+1}}{\mathring{b}_n^{(1)} x^n} \right| \sim \begin{cases} |x| \left[1 + \frac{1}{n} \text{Re}(\delta-2) \right], & \text{if } |a| < 1, \\ \left| \frac{x}{a} \right| \left[1 + \frac{1}{n} \text{Re}(\epsilon-2) \right], & \text{if } |a| > 1. \end{cases} \quad (42)$$

Then, if $|a| < 1$ the solution $\mathring{H}^{\{1\}}(x)$ converges inside the circle $|x| = 1$ which contains $x = 0$ and $x = a$; by the Raabe's test [27, 28] it converges also on $|x| = 1$ if $\text{Re } \delta < 1$. On the other hand, if $|a| > 1$ the solution converges inside the circle $|x| = |a|$ which contains $x = 0$ and $x = 1$; by the Raabe's test it converges also on $|x| = |a|$ if $\text{Re } \epsilon < 1$. In sum,

$$\text{for minimal solution} \quad \begin{cases} \text{if } |a| < 1, \mathring{H}^{(1)}(x) \text{ converges for } |x| < 1 \text{ and on } |x| = 1 \text{ if } \text{Re } \delta < 1; \\ \text{if } |a| > 1, \mathring{H}^{(1)}(x) \text{ converges for } |x| < |a| \text{ and on } |x| = |a| \text{ if } \text{Re } \epsilon < 1. \end{cases}$$

Observe that Eq. (37) and the limits preceding Eqs. (41) imply that the recurrence relations also admit a non-minimal solution.

The remaining solutions $\mathring{H}^{(i)}(x)$ are generated by the homotopic transformations T_i as indicated in (39). This gives the expansions written below, where the coefficients $\mathring{\beta}_n^{(i)}$ are written as polynomials of degree 2 in n .

$$\begin{cases} \mathring{H}^{(1)}(x) = \sum_{n=0}^{\infty} \mathring{b}_n^{(1)} x^n, \\ \mathring{\alpha}_n^{(1)} = a(n + \gamma)(n + 1), \quad \mathring{\beta}_n^{(1)} = -(a + 1)n^2 - [a(\gamma + \delta - 1) + \alpha + \beta - \delta]n \\ -q, \quad \mathring{\gamma}_n^{(1)} = (n + \alpha - 1)(n + \beta - 1). \end{cases} \quad (43)$$

$$\begin{cases} \mathring{H}^{(2)}(x) = x^{1-\gamma} \sum_{n=0}^{\infty} \mathring{b}_n^{(2)} x^n, \\ \mathring{\alpha}_n^{(2)} = a(n + 2 - \gamma)(n + 1), \quad \mathring{\beta}_n^{(2)} = -(a + 1)n^2 - [a(1 - \gamma + \delta) + 1 + \epsilon - \gamma]n \\ -q - (\delta a + \epsilon)(1 - \gamma), \quad \mathring{\gamma}_n^{(2)} = (n + \alpha - \gamma)(n + \beta - \gamma). \end{cases} \quad (44)$$

$$\begin{cases} \mathring{H}^{(3)}(x) = (1 - x)^{1-\delta} \sum_{n=0}^{\infty} \mathring{b}_n^{(3)} x^n, \\ \mathring{\alpha}_n^{(3)} = a(n + \gamma)(n + 1), \quad \mathring{\beta}_n^{(3)} = -(a + 1)n^2 - [a(1 + \gamma - \delta) + \alpha + \beta - \delta]n \\ -q - a\gamma(1 - \delta), \quad \mathring{\gamma}_n^{(3)} = (n + \alpha - \delta)(n + \beta - \delta). \end{cases} \quad (45)$$

$$\begin{cases} \mathring{H}^{(4)}(x) = x^{1-\gamma}(1 - x)^{1-\delta} \sum_{n=0}^{\infty} \mathring{b}_n^{(4)} x^n, \\ \mathring{\alpha}_n^{(4)} = a(n + 2 - \gamma)(n + 1), \quad \mathring{\beta}_n^{(4)} = -(a + 1)n^2 - [a(3 - \gamma - \delta) + 1 + \epsilon - \gamma]n \\ -q - a(2 - \gamma - \delta) - \epsilon(1 - \gamma), \quad \mathring{\gamma}_n^{(4)} = (n + 1 + \alpha - \gamma - \delta)(n + 1 + \beta - \gamma - \delta). \end{cases} \quad (46)$$

$$\begin{cases} \mathring{H}^{(5)}(x) = \left[1 - \frac{x}{a}\right]^{1-\epsilon} \sum_{n=0}^{\infty} \mathring{b}_n^{(5)} x^n, \\ \mathring{\alpha}_n^{(5)} = a(n + \gamma)(n + 1), \quad \mathring{\beta}_n^{(5)} = -(a + 1)n^2 - [a(\gamma + \delta - 1) + 2\gamma + \delta - \alpha - \beta]n \\ -q - \gamma(1 - \epsilon), \quad \mathring{\gamma}_n^{(5)} = (n + \gamma + \delta - \alpha - 1)(n + \gamma + \delta - \beta - 1). \end{cases} \quad (47)$$

$$\begin{cases} \mathring{H}^{(6)}(x) = x^{1-\gamma} \left[1 - \frac{x}{a}\right]^{1-\epsilon} \sum_{n=0}^{\infty} \mathring{b}_n^{(6)} x^n, \\ \mathring{\alpha}_n^{(6)} = a(n + 2 - \gamma)(n + 1), \quad \mathring{\beta}_n^{(6)} = -(a + 1)n^2 - [a(1 - \gamma + \delta) - \alpha - \beta + \\ \delta + 2]n - q - \delta a(1 - \gamma) + \alpha + \beta - \delta - 1, \quad \mathring{\gamma}_n^{(6)} = (n - \alpha + \delta)(n - \beta + \delta). \end{cases} \quad (48)$$

$$\begin{cases} \mathring{H}^{(7)}(x) = (1-x)^{1-\delta} \left[1 - \frac{x}{a}\right]^{1-\epsilon} \sum_{n=0}^{\infty} \mathring{b}_n^{(7)} x^n, \\ \mathring{\alpha}_n^{(7)} = a(n+\gamma)(n+1), \quad \mathring{\beta}_n^{(7)} = -(a+1)n^2 - [a(1+\gamma-\delta) - \alpha - \beta + 2\gamma + \delta]n - q - \gamma a(1-\delta) - \gamma(1-\epsilon), \\ \mathring{\gamma}_n^{(7)} = (n-\alpha+\gamma)(n-\beta+\gamma). \end{cases} \quad (49)$$

$$\begin{cases} \mathring{H}^{(8)}(x) = x^{1-\gamma}(1-x)^{1-\delta} \left[1 - \frac{x}{a}\right]^{1-\epsilon} \sum_{n=0}^{\infty} \mathring{b}_n^{(8)} x^n, \\ \mathring{\alpha}_n^{(8)} = a(n+2-\gamma)(n+1), \quad \mathring{\beta}_n^{(8)} = -(a+1)n^2 - [a(3-\gamma-\delta) + 2 - \alpha - \beta + \delta]n - q - a(2-\gamma-\delta) + \alpha + \beta - \delta - 1, \\ \mathring{\gamma}_n^{(8)} = (n+1-\alpha)(n+1-\beta). \end{cases} \quad (50)$$

Since x and a are kept fixed by homotopic transformations T_i , all the solutions $\mathring{H}^{(i)}(x)$ are Heun functions which converge inside the circle $|x| = 1$ if $|a| < 1$ and inside the circle $|x| = |a|$ if $|a| > 1$; in fact, by

$$\text{by D'Alembert's test} \quad \begin{cases} \text{if } |a| < 1 \text{ } \mathring{H}^{(i)} \text{ converges for } |x| < 1 (\Rightarrow x=0 \text{ and } x=a); \\ \text{if } |a| > 1 \text{ } \mathring{H}^{(i)} \text{ converges for } |x| < |a| (\Rightarrow x=0 \text{ and } x=1). \end{cases} \quad (51)$$

In addition, by Raabe's test, the transformations (A.1-A.8) give the following conditions for convergence on $|x| = 1$ and $|x| = |a|$:

$$\text{if } |a| < 1 \quad \begin{cases} \mathring{H}^{(1)}, \mathring{H}^{(2)}, \mathring{H}^{(5)} \text{ and } \mathring{H}^{(6)} \text{ converge on } |x| = 1 \text{ if } \text{Re } \delta < 1; \\ \mathring{H}^{(3)}, \mathring{H}^{(4)}, \mathring{H}^{(7)} \text{ and } \mathring{H}^{(8)} \text{ converge on } |x| = 1 \text{ if } \text{Re } \delta > 1; \end{cases} \quad (52)$$

$$\text{if } |a| > 1 \quad \begin{cases} \mathring{H}^{(1)}, \mathring{H}^{(2)}, \mathring{H}^{(3)} \text{ and } \mathring{H}^{(4)} \text{ converge on } |x| = |a| \text{ if } \text{Re } \epsilon < 1; \\ \mathring{H}^{(5)}, \mathring{H}^{(6)}, \mathring{H}^{(7)} \text{ and } \mathring{H}^{(8)} \text{ converge on } |x| = |a| \text{ if } \text{Re } \epsilon > 1. \end{cases} \quad (53)$$

The previous restrictions on δ and ϵ hold only for infinite series since the convergence of finite series is determined by examining each term of the series. Anyway, the behaviour of the solutions at the singular points depends also on the multiplicative factors which precede the series.

Since the recurrence relations admit two solutions, Kristensson [9] used the Raabe test to determine the convergence of $\mathring{H}^{(1)}(x)$ by supposing that that $|a| > 1$. He found that

$$\text{if } |a| > 1 \text{ } \mathring{H}^{(1)} \text{ converges for } \begin{cases} |x| < 1 \text{ and at } |x| = 1 \text{ if } \text{Re } \delta < 1 \text{ (dominant solution);} \\ |x| < |a| \text{ and at } |x| = |a| \text{ if } \text{Re } \epsilon < 1 \text{ (minimal solution).} \end{cases} \quad (54)$$

Note the differences between the above treatments: we are considering only minimal solutions for both $|a| > 1$ and $|a| < 1$; in addition, we are using the homotopic transformations.

In the previous expansions we see that

$$\begin{aligned} a = 0, & \quad \Rightarrow \quad \mathring{\alpha}_n^{(i)} = 0; \\ a = -1, \quad \epsilon = \delta, \quad q = 0 & \quad \Rightarrow \quad \mathring{\beta}_n^{(1)} = \mathring{\beta}_n^{(2)} = \mathring{\beta}_n^{(7)} = \mathring{\beta}_n^{(8)} = 0; \\ a = -1, \quad \alpha + \beta = 1 + \gamma, \quad q = \gamma(1 - \delta) & \quad \Rightarrow \quad \mathring{\beta}_n^{(3)} = \mathring{\beta}_n^{(4)} = \mathring{\beta}_n^{(5)} = \mathring{\beta}_n^{(6)} = 0. \end{aligned} \quad (55)$$

For these sets of parameters, the recurrence relations (32) become two-terms relations and the Heun equation (2) can be reduced to the hypergeometric equation (3) – see Appendix B.

Finally, by considering the solutions $\mathring{H}^{(i+4)}(x)$ and $\mathring{H}^{(i)}(x)$ ($i = 1, 2, 3, 4$) we find that

$$\text{for infinite series, } \mathring{H}^{(i+4)}(x) = \mathring{H}^{(i)}(x) \quad \text{if } |x| < |a| \quad \text{and} \quad 1 - \epsilon \neq 0, 1, 2, \dots \quad (56)$$

Relations (56) results from [32]

$$(1 + y)^\kappa = \sum_{m=0}^{\infty} \frac{\Gamma(\kappa + m)}{m! \Gamma(\kappa)} y^m, \quad |y| < 1, \quad \kappa \neq 0, 1, 2, \dots \quad (57)$$

applied to the factor $[1 - x/a]^{1-\epsilon}$, and from the Cauchy product for infinite series, namely,

$$\sum_{n=0}^{\infty} f_n \sum_{m=0}^{\infty} g_m = \sum_{j=0}^{\infty} h_j, \quad \text{where} \quad h_j = \sum_{l=0}^j f_l g_{j-l}. \quad (58)$$

For instance, for $\mathring{H}^{(5)}(x)$ we have

$$\begin{aligned} \mathring{H}^{(5)}(x) &\stackrel{(47)}{=} \sum_{n=0}^{\infty} \mathring{b}_n^{(5)} x^n \left[1 - \frac{x}{a}\right]^{1-\epsilon} \stackrel{(57)}{=} \sum_{n=0}^{\infty} \mathring{b}_n^{(5)} x^n \sum_{m=0}^{\infty} \left(-\frac{1}{a}\right)^m \frac{\Gamma(1 - \epsilon + m)}{\Gamma(1 - \epsilon) m!} x^m \\ &\stackrel{(58)}{=} \sum_{j=0}^{\infty} x^j \left[\frac{1}{\Gamma(1 - \epsilon)} \sum_{l=0}^j \left(-\frac{1}{a}\right)^{j-l} \frac{\Gamma(1 - \epsilon + j - l)}{(j - l)!} \mathring{b}_l^{(5)} \right], \quad |x| < |a|, \quad 1 - \epsilon \neq 0, 1, \dots \end{aligned}$$

Thence, by taking $j = n$ and comparing the above equation with (43), we get

$$\mathring{H}^{(5)}(x) = \mathring{H}^{(1)}(x) \quad \text{with} \quad \mathring{b}_n^{(1)} = \frac{1}{\Gamma(1-\epsilon)} \sum_{l=0}^n \left(-\frac{1}{a}\right)^{n-l} \frac{\Gamma(1 - \epsilon + n - l)}{(n - l)!} \mathring{b}_l^{(5)}.$$

In fact, Eqs. (56) generalize an Ince result [33] for solutions of the Lamé equation (in which case $\gamma = \delta = \epsilon = 1/2$, $x = \text{sn}^2 u$ and $a = 1/k^2$).

Finally we set the limits of $\mathring{H}^{(1)}$ for the confluent Heun equation (CHE) which, in the form used by Arscott [21], reads

$$x(x-1) \frac{d^2 S(x)}{dx^2} + [-\gamma + (\gamma + \delta)x + \rho x(x-1)] \frac{dS(x)}{dx} + (\alpha \rho x - \sigma) S(x) = 0, \quad (59)$$

$$\rho \neq 0, \quad (CHE),$$

where ρ and σ are constants. The points $x = 0$ and $x = 1$ are regular singular points while $z = \infty$ is an irregular point. Eq. (59) results from the Heun equation (2) by the limits

$$a, \beta, q \rightarrow \infty \quad \text{such that} \quad \frac{\beta}{a} \rightarrow \frac{\epsilon}{a} \rightarrow -\rho, \quad \frac{q}{a} \rightarrow -\sigma, \quad H(x) \rightarrow S(x). \quad (60)$$

Dividing the recurrence relations by a and using the previous limits, the expansion (43) gives ($\mathring{b}_{-1}^{(1)} = 0$)

$$\begin{cases} \mathring{S}^{(1)}(x) = \sum_{n=0}^{\infty} \mathring{b}_n^{(1)} x^n, \\ (n + \gamma)(n + 1) \mathring{b}_{n+1}^{(1)} - [n^2 + (\gamma + \delta - 1 - \rho)n - \sigma] \mathring{b}_n^{(1)} - \rho(n + \alpha - 1) \mathring{b}_{n-1}^{(1)} = 0, \end{cases} \quad (61)$$

as solution to the CHE (59). From (61) we can generate other expansions by means of transformations of the CHE [22, 34]. Actually, expansion (61) includes the one used in 1935 by Baber and Hassé to obtain convergent solutions to the angular part of the two-center problem of quantum mechanics [35].

3.2 Series $\mathring{H}^{(i)}(x)$ around $x = 1$

The following power series $\mathring{H}^{(i)}(x)$ around $x = 1$ come from $\mathring{H}^{(1)}(x) = M_{49}\mathring{H}^{(i)}(x)$, $\mathring{H}^{(i)}(x) = T_i\mathring{H}^{(1)}(x)$. In the following the coefficients $\mathring{\gamma}_n^{(i)}$ are the ones of section III.A.

$$\begin{cases} \mathring{H}^{(1)}(x) = \sum_{n=0}^{\infty} \mathring{b}_n^{(1)}(1-x)^n, \\ \mathring{\alpha}_n^{(1)} = (1-a)(n+\delta)(n+1), \quad \mathring{\beta}_n^{(1)} = (a-2)n^2 + [(a-1)(\gamma+\delta-1) - \alpha - \beta \\ + \gamma]n + q - \alpha\beta, \quad \mathring{\gamma}_n^{(1)} = \mathring{\gamma}_n^{(1)}. \end{cases} \quad (62)$$

$$\begin{cases} \mathring{H}^{(2)}(x) = x^{1-\gamma} \sum_{n=0}^{\infty} \mathring{b}_n^{(2)}(1-x)^n, \\ \mathring{\alpha}_n^{(2)} = (1-a)(n+\delta)(n+1), \quad \mathring{\beta}_n^{(2)} = (a-2)n^2 + [(a-1)(1-\gamma+\delta) - \alpha \\ - \beta + \gamma]n + q - a(\gamma-1)\delta + (\gamma-1)\delta - \alpha\beta, \quad \mathring{\gamma}_n^{(2)} = \mathring{\gamma}_n^{(2)}, \end{cases} \quad (63)$$

$$\begin{cases} \mathring{H}^{(3)}(x) = (1-x)^{1-\delta} \sum_{n=0}^{\infty} \mathring{b}_n^{(3)}(1-x)^n, \\ \mathring{\alpha}_n^{(3)} = (1-a)(n+2-\delta)(n+1), \quad \mathring{\beta}_n^{(3)} = (a-2)n^2 + [(a-1)(1+\gamma-\delta) \\ - \alpha - \beta + \gamma + 2\delta - 2]n + q - a\gamma(\delta-1) - \alpha\beta + (\delta-1)(\alpha + \beta - \delta + 1), \\ \mathring{\gamma}_n^{(3)} = \mathring{\gamma}_n^{(3)}. \end{cases} \quad (64)$$

$$\begin{cases} \mathring{H}^{(4)}(x) = x^{1-\gamma}(1-x)^{1-\delta} \sum_{n=0}^{\infty} \mathring{b}_n^{(4)}(1-x)^n, \\ \mathring{\alpha}_n^{(4)} = (1-a)(n+2-\delta)(n+1), \quad \mathring{\beta}_n^{(4)} = (a-2)n^2 + [(a-1)(3-\gamma-\delta) \\ - \alpha - \beta + \gamma + 2\delta - 2]n + q - a(\gamma+\delta-2) - \alpha\beta + \gamma + \delta - 2 + (\delta-1)\epsilon, \\ \mathring{\gamma}_n^{(4)} = \mathring{\gamma}_n^{(4)}. \end{cases} \quad (65)$$

$$\begin{cases} \mathring{H}^{(5)}(x) = \left[1 - \frac{x}{a}\right]^{1-\epsilon} \sum_{n=0}^{\infty} \mathring{b}_n^{(5)}(1-x)^n, \\ \mathring{\alpha}_n^{(5)} = (1-a)(n+\delta)(n+1), \quad \mathring{\beta}_n^{(5)} = (a-2)n^2 + [(a-1)(\gamma+\delta-1) \\ + \alpha + \beta - \gamma - 2\delta]n + q - \alpha\beta + (\alpha + \beta - \gamma - \delta)\delta, \quad \mathring{\gamma}_n^{(5)} = \mathring{\gamma}_n^{(5)}. \end{cases} \quad (66)$$

$$\begin{cases} \mathring{H}^{(6)}(x) = x^{1-\gamma} \left[1 - \frac{x}{a}\right]^{1-\epsilon} \sum_{n=0}^{\infty} \mathring{b}_n^{(6)}(1-x)^n, \\ \mathring{\alpha}_n^{(6)} = (1-a)(n+\gamma)(n+1), \quad \mathring{\beta}_n^{(6)} = (a-2)n^2 + [(a-1)(1-\gamma+\delta) \\ + \alpha + \beta - \gamma - 2\delta]n + q - a(\gamma-1)\delta - \alpha\beta + (\alpha + \beta - \delta - 1)\delta, \quad \mathring{\gamma}_n^{(6)} = \mathring{\gamma}_n^{(6)}. \end{cases} \quad (67)$$

$$\begin{cases} \mathring{H}^{(7)}(x) = (1-x)^{1-\delta} \left[1 - \frac{x}{a}\right]^{1-\epsilon} \sum_{n=0}^{\infty} \mathring{b}_n^{(7)}(1-x)^n, \\ \mathring{\alpha}_n^{(7)} = (1-a)(n+2-\delta)(n+1), \quad \mathring{\beta}_n^{(7)} = (a-2)n^2 + [(a-1)(1+\gamma-\delta) \\ + \alpha + \beta - \gamma - 2]n + q - a\gamma(\delta-1) - \alpha\beta + \gamma(\delta-1) + \alpha + \beta - \gamma - 1, \\ \mathring{\gamma}_n^{(7)} = \mathring{\gamma}_n^{(7)}. \end{cases} \quad (68)$$

$$\begin{cases} \mathring{\mathbf{H}}^{(8)}(x) = x^{1-\gamma}(1-x)^{1-\delta} \left[1 - \frac{x}{a}\right]^{1-\epsilon} \sum_{n=0}^{\infty} \mathring{\mathbf{b}}_n^{(8)}(1-x)^n, \\ \mathring{\alpha}_n^{(8)} = (1-a)(n+2-\delta)(n+1), \quad \mathring{\beta}_n^{(7)} = (a-2)n^2 + [(a-1)(3-\gamma-\delta) \\ + \alpha + \beta - \gamma - 2]n + q + a(2-\gamma-\delta) - \alpha\beta + \alpha + \beta + \delta - 3, \quad \gamma_n^{(8)} = \mathring{\gamma}_n^{(8)}. \end{cases} \quad (69)$$

We find that

$$\begin{aligned} a = 1 &\Rightarrow \mathring{\alpha}_n^{(i)} = 0; \\ a = 2, \alpha + \beta + 1 = 2\gamma + \delta, q = \alpha\beta &\Rightarrow \mathring{\beta}_n^{(1)} = \mathring{\beta}_n^{(3)} = \mathring{\beta}_n^{(6)} = \mathring{\beta}_n^{(8)} = 0; \\ a = 2, \alpha + \beta = 1 + \delta, q = \alpha\beta + (\gamma - 1)\delta &\Rightarrow \mathring{\beta}_n^{(2)} = \mathring{\beta}_n^{(4)} = \mathring{\beta}_n^{(5)} = \mathring{\beta}_n^{(7)} = 0 \end{aligned} \quad (70)$$

in the expansions (62-69). Once more, for these cases, the Heun equation (2) can be reduced to the hypergeometric equation (3) – see Appendix B.

4 Solutions in series of hypergeometric functions

This section presents the two groups of solutions in series of hypergeometric functions, $F(a, b; c; z)$, which are denoted by $H^{(i)}(x)$ and $\mathbf{H}^{(i)}(x)$ ($i = 1, 2, \dots, 8$) and correspond, respectively, to the series coefficients denoted by $b_n^{(i)}$ and $\mathbf{b}_n^{(i)}$. The solutions are generated from the solution $\bar{H}^{(1)}(x)$, written in Eqs. (74a) and (74b), by means of the following transformations

$$H^{(1)}(x) = M_{17}\bar{H}^{(1)}(x), \quad H^{(i)}(x) = T_i H^{(1)}(x), \quad (71)$$

$$\mathbf{H}^{(1)}(x) = M_{49}\bar{H}^{(1)}(x), \quad \mathbf{H}^{(i)}(x) = T_i \mathbf{H}^{(1)}(x), \quad (72)$$

where M_{17} and M_{49} are the fractional transformations given in Eqs. (26) and (27), and T_i are the homotopic transformations (A.1-A.8).

In section 4.1 we construct the initial solution $\bar{H}^{(1)}(x)$. In sections 4.2 and 4.3 we write the solutions $H^{(i)}(x)$ and $\mathbf{H}^{(i)}(x)$. Formally the solutions are given by infinite series, but they are applied only for the cases in which the series become finite. By this reason, there is no discussion on convergence of the series.

To simplify the expressions of the recurrence relations, from Eq. (79) onwards, the solutions are written in terms of the functions $\tilde{F}(a, b; c; z)$ defined as

$$\tilde{F}(a, b; c; z) = \frac{1}{\Gamma(c)} F(a, b; c; z) \quad (73)$$

where Γ stands for gamma function.

4.1 The initial solution $\bar{H}^{(1)}(x)$

The solution $\bar{H}^{(1)}$ in series of hypergeometric functions $F(n + \alpha, \gamma + \delta - \alpha - 1; n + \gamma; x)$ is

$$\bar{H}^{(1)}(x) = \sum_{n=0}^{\infty} \bar{b}_n^{(1)} x^n F(n + \alpha, \gamma + \delta - \alpha - 1; n + \gamma; x). \quad (74a)$$

where the recurrence relations (32) for $\bar{b}_n^{(1)}$ take the form $\bar{\alpha}_n^{(1)} \bar{b}_{n+1}^{(1)} + \bar{\beta}_n^{(1)} \bar{b}_n^{(1)} + \bar{\gamma}_n^{(1)} \bar{b}_{n-1}^{(1)} = 0$ with $\bar{b}_{-1}^{(1)} = 0$ and

$$\begin{aligned} \bar{\alpha}_n^{(1)} &= a(n+1)(n+\gamma), \quad \bar{\beta}_n^{(1)} = -(a+1)n^2 - [a(2\alpha+1-\gamma-\delta) + \alpha + \beta - \delta]n \\ &- q - a\alpha(\alpha+1-\gamma-\delta), \quad \bar{\gamma}_n^{(1)} = \frac{(n+\alpha-1)(n+\alpha-\delta)(n+\alpha+\beta-\gamma-\delta)}{n+\gamma-1}. \end{aligned} \quad (74b)$$

The above hypergeometric functions converge for any $|x| < 1$ and on $|x| = 1$ if $\text{Re } \delta < 1$. For this reason, the solution $\bar{H}^{(1)}(x)$ will be valid only for $|x| \leq 1$ in opposition to $\dot{H}^{(1)}(x)$ which may be valid for $|x| > 1$ if $|a| > 1$ – see conditions (51). In fact, $\bar{H}^{(1)}(x) = \dot{H}^{(1)}(x)$ only if $\gamma + \delta - 1 = \alpha$.

The Heun equation (2) is equivalent to

$$(x-a) \left[x(x-1) \frac{d^2}{dx^2} + (\gamma + \delta)x \frac{d}{dx} - \gamma \frac{d}{dx} \right] H + \epsilon x(x-1) \frac{dH}{dx} + [\alpha\beta x - q] H = 0, \quad (75)$$

The solution (74a) is obtained by writing

$$H(x) = \bar{H}^{(1)}(x) = \sum_{n=0}^{\infty} \bar{b}_n^{(1)} x^n F_n^{(1)}(x). \quad \text{where} \quad (76)$$

$$F_n^{(1)}(x) = F(\rho_n, b; \sigma_n; x), \quad \rho_n = n + \alpha, \quad b = \gamma + \delta - \alpha - 1, \quad \sigma_n = n + \gamma.$$

Then,

$$\frac{dH}{dx} = \sum_{n=0}^{\infty} \bar{b}_n^{(1)} x^n \left[\frac{d}{dx} + \frac{n}{x} \right] F_n^{(1)}, \quad \frac{d^2 H}{dx^2} = \sum_{n=0}^{\infty} \bar{b}_n^{(1)} x^n \left[\frac{d^2}{dx^2} + \frac{2n}{x} \frac{d}{dx} + \frac{n(n-1)}{x^2} \right] F_n^{(1)}.$$

By inserting the previous expressions into Eq. (75) and using the hypergeometric equation

$$x(1-x) \frac{d^2 F_n^{(1)}(x)}{dx^2} + [\sigma_n - (\rho_n + b + 1)x] \frac{dF_n^{(1)}(x)}{dx} - \rho_n b F_n^{(1)}(x) = 0,$$

for $F_n(x)$, we find

$$\begin{aligned} & \sum_{n=0}^{\infty} \bar{b}_n^{(1)} x^n \left[an(1-x) \frac{dF_n^{(1)}}{dx} + (n+\epsilon)x(x-1) \frac{dF_n^{(1)}}{dx} \right] + \\ & \sum_{n=0}^{\infty} \bar{b}_n^{(1)} x^{n+1} [n(n+\gamma+\delta-1+\epsilon) + \alpha\beta - b\rho_n] F_n^{(1)} + \\ & \sum_{n=0}^{\infty} \bar{b}_n^{(1)} x^n [a\alpha(\gamma+\delta-\alpha-1) - q - an(n+\alpha) - n(n+\gamma-1+\epsilon)] F_n^{(1)} + \\ & \sum_{n=0}^{\infty} \bar{b}_n^{(1)} an(n-1+\gamma)x^{n-1} F_n^{(1)} = 0. \end{aligned} \quad (77)$$

On the other side, from the properties (23) of the hypergeometric functions we get

$$\begin{aligned} (1-x) \frac{dF_n^{(1)}}{dx} &= \left(b - \frac{\sigma_{n-1}}{x} \right) F_n^{(1)} + \frac{\sigma_{n-1}}{x} F_{n-1}^{(1)}, \quad x(x-1) \frac{dF_n^{(1)}}{dx} = \\ & -\rho_n x F_n^{(1)} + \frac{\rho_n(\sigma_n - b)}{\sigma_n} x F_{n+1}^{(1)}. \end{aligned} \quad (78)$$

By means of these equations, Eq. (77) becomes

$$\sum_{n=1}^{\infty} \bar{b}_n^{(1)} \sigma_{n-1} x^{n-1} F_{n-1}^{(1)} + \sum_{n=0}^{\infty} \bar{b}_n^{(1)} x^n \bar{\beta}_n F_n^{(1)} + \sum_{n=0}^{\infty} \bar{b}_n^{(1)} \frac{\rho_n(\sigma_n - b)(n+\epsilon)}{\sigma_n} x^{n+1} F_{n+1}^{(1)} = 0.$$

Now, by setting $n = m+1$ and $n = m-1$ in the first and last terms of the above equation, one finds

$$\begin{aligned} & \left[a\sigma_0 \bar{b}_1^{(1)} + \bar{\beta}_0 \bar{b}_0^{(1)} \right] F_0^{(1)}(x) + \\ & \sum_{n=1}^{\infty} \left[a(n+1)\sigma_n \bar{b}_{n+1}^{(1)} + \bar{\beta}_n \bar{b}_n^{(1)} + \frac{\rho_{n-1}(\sigma_{n-1} - b)(n+\epsilon-1)}{\sigma_{n-1}} \bar{b}_{n-1}^{(1)} \right] x^n F_n^{(1)}(x) = 0. \end{aligned}$$

Thence, the recurrence relations for $\bar{b}_n^{(1)}$ follow by requiring that the coefficients of the functions $x^n F_n^{(1)}(x)$ vanish.

Finally, by taking $\tilde{b}_n^{(1)} = \Gamma(n + \gamma) \bar{b}_n$, the expansion $\bar{H}^{(1)}$ is re-expressed in terms of the functions \tilde{F} defined in (73), namely,

$$\left\{ \begin{array}{l} \bar{H}^{(1)}(x) = \sum_{n=0}^{\infty} \tilde{b}_n^{(1)} x^n \tilde{F}(n + \alpha, \gamma + \delta - \alpha - 1; n + \gamma; x) \quad \text{where, in (32) for } \tilde{b}_n^{(1)}, \\ \tilde{\alpha}_n^{(1)} = a(n + 1), \quad \tilde{\beta}_n^{(1)} = -(a + 1)n^2 - [a(2\alpha + 1 - \gamma - \delta) + \alpha + \beta - \delta]n - q \\ -a\alpha(\alpha + 1 - \gamma - \delta), \quad \tilde{\gamma}_n^{(1)} = (n + \alpha - 1)(n + \alpha - \delta)(n + \alpha + \beta - \gamma - \delta). \end{array} \right. \quad (79)$$

Other solutions are given $\bar{H}^{(i)}(x) = T_i \bar{H}^{(1)}(x)$. Some of these expansions yield finite-series solutions for the problem discussed in Section 5; however, such solutions do not fulfil the Arcsott conditions (36). By this reason, we use $\bar{H}^{(1)}(x)$ to obtain the expansions $H^{(i)}(x)$ and $\mathbf{H}^{(i)}(x)$ which are used in Sections 5.3 and 5.4, respectively.

As a by-product we find that the expansion (79) admits the limits (60) for the CHE (59), leading to

$$\left\{ \begin{array}{l} \bar{S}^{(1)}(x) = \sum_{n=0}^{\infty} \tilde{b}_n^{(1)} x^n \tilde{F}(n + \alpha, \gamma + \delta - \alpha - 1; n + \gamma; x) \quad \text{with} \\ (n + 1) \tilde{b}_{n+1}^{(1)} - [n^2 + (2\alpha + 1 - \gamma - \delta - \rho)n - \sigma + \alpha(\alpha + 1 - \gamma - \delta)] \tilde{b}_n^{(1)} - \\ \rho(n + \alpha - 1)(n + \alpha - \delta) \tilde{b}_{n-1}^{(1)} = 0, \quad [\tilde{b}_{-1}^{(1)} = 0]. \end{array} \right. \quad (80)$$

This $\bar{S}^{(1)}$ is equivalent to a solution for the CHE proposed 1937 by E. Fisher [36].

4.2 The solutions $H^{(i)}(x)$ around $(a - 1)x/(a - x) = 0$

In this section $H^{(1)}(x) = M_{17} \bar{H}^{(1)}(x)$ and $H^{(i)}(x) = T_i H^{(1)}(x)$, where $\bar{H}^{(1)}$ is given in (79). The recurrence relations for the coefficients $b_n^{(i)}$ once more have the form (32).

$$\left\{ \begin{array}{l} H^{(1)}(x) = \left(1 - \frac{x}{a}\right)^{-\alpha} \sum_{n=0}^{\infty} b_n^{(1)} \left[\frac{(a-1)x}{a-x}\right]^n \tilde{F}\left[n + \alpha, \gamma + \delta - \alpha - 1; n + \gamma; \frac{(a-1)x}{a-x}\right], \\ \alpha_n^{(1)} = (1 - a)(n + 1), \quad \beta_n^{(1)} = (a - 2)n^2 + [(a - 1)(2\alpha + 1 - \gamma - \delta) - \alpha + \\ \beta - \gamma]n + q + (a - 1)\alpha(\alpha + 1 - \gamma - \delta) - \alpha\gamma, \quad \gamma_n^{(1)} = (n + \alpha - 1) \times \\ (n + \alpha - \delta)(n + \alpha - \beta). \end{array} \right. \quad (81)$$

$$\left\{ \begin{array}{l} H^{(2)}(x) = x^{1-\gamma} \left(1 - \frac{x}{a}\right)^{\gamma-\beta-1} \times \\ \sum_{n=0}^{\infty} b_n^{(2)} \left[\frac{(a-1)x}{a-x}\right]^n \tilde{F}\left[n + \beta + 1 - \gamma, \delta - \beta; n + 2 - \gamma; \frac{(a-1)x}{a-x}\right], \\ \alpha_n^{(2)} = (1 - a)(n + 1), \quad \beta_n^{(2)} = (a - 2)n^2 + [(a - 1)(2\beta + 1 - \gamma - \delta) - 2 + \\ \alpha - \beta + \gamma]n + q + a\beta(\beta + 1 - \gamma - \delta) - \beta(\beta - \delta + 1) + (\gamma - 1)(1 - \alpha + \beta), \\ \gamma_n^{(2)} = (n + \beta - \gamma)(n + \beta + 1 - \gamma - \delta)(n - \alpha + \beta) \end{array} \right. \quad (82)$$

The solutions $H^{(3)}$ and $H^{(4)}$ take the forms

$$H^{(3)}(x) = (1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{\delta-\beta-1} \times \sum_{n=0}^{\infty} b_n^{(3)} \left[\frac{(a-1)x}{a-x}\right]^n \tilde{F} \left[n + \beta + 1 - \delta, \gamma - \beta; n + \gamma; \frac{(a-1)x}{a-x}\right] \stackrel{(21)}{=} H^{(1)}(x) \Big|_{\alpha \leftrightarrow \beta}, \quad (83)$$

$$H^{(4)}(x) = x^{1-\gamma} (1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{\gamma+\delta-\alpha-2} \sum_{n=0}^{\infty} b_n^{(4)} \times \left[\frac{(a-2)x}{a-x}\right]^n \tilde{F} \left[n + 2 + \alpha - \gamma - \delta, 1 - \alpha; n + 2 - \gamma; \frac{(a-1)x}{a-x}\right] = H^{(2)}(x) \Big|_{\alpha \leftrightarrow \beta}, \quad (84)$$

where α and β must be interchanged also in the recurrence relations.

$$\left\{ \begin{aligned} H^{(5)}(x) &= (1 - \frac{x}{a})^{-\beta} \sum_{n=0}^{\infty} b_n^{(5)} \left[\frac{(a-1)x}{a-x}\right]^n \tilde{F} \left[n - \alpha + \gamma + \delta, \alpha - 1; n + \gamma; \frac{(a-1)x}{a-x}\right], \\ \alpha_n^{(5)} &= (1-a)(n+1), \quad \beta_n^{(5)} = (a-2)n^2 + [(a-1)(1-2\alpha + \gamma + \delta) + \\ &\quad \alpha - \beta - \gamma]n + q + (a-1)(\alpha-1)(\alpha-\gamma-\delta) - \beta\gamma, \\ \gamma_n^{(5)} &= (n - \alpha + \gamma + \delta - 1)(n - \alpha + \gamma)(n - \alpha + \beta). \end{aligned} \right. \quad (85)$$

$$\left\{ \begin{aligned} H^{(6)}(x) &= x^{1-\gamma} \left(1 - \frac{x}{a}\right)^{\gamma-\alpha-1} \sum_{n=0}^{\infty} b_n^{(6)} \left[\frac{(a-1)x}{a-x}\right]^n \times \\ &\quad \tilde{F} \left[n - \beta + \delta + 1, \beta - \gamma; n + 2 - \gamma; \frac{(a-1)x}{a-x}\right], \\ \alpha_n^{(6)} &= (1-a)(n+1), \quad \beta_n^{(6)} = (a-2)n^2 + [(a-1)(1-2\beta + \gamma + \delta) - \alpha + \\ &\quad \beta + \gamma - 2]n + q - \alpha + a(\beta-1)(\beta-\gamma-\delta) - (\beta-1)(\beta-\delta-1), \\ \gamma_n^{(6)} &= (n - \beta + \delta)(n - \beta + 1)(n + \alpha - \beta). \end{aligned} \right. \quad (86)$$

The solutions $H^{(7)}(x)$ and $H^{(8)}(x)$ are expressed as

$$H^{(7)}(x) = (1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{\delta-\alpha-1} \times \sum_{n=0}^{\infty} b_n^{(7)} \left[\frac{(a-1)x}{a-x}\right]^n \tilde{F} \left[n + 1 - \beta + \gamma, \beta - \delta; n + \gamma; \frac{(a-1)x}{a-x}\right] \stackrel{(21)}{=} H^{(5)}(x) \Big|_{\alpha \leftrightarrow \beta}, \quad (87)$$

$$H^{(8)}(x) = x^{1-\gamma} (1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{\gamma+\delta-\beta-2} \sum_{n=0}^{\infty} b_n^{(8)} \times \left[\frac{(a-1)x}{a-x}\right]^n \tilde{F} \left[n + 2 - \alpha, \alpha + 1 - \gamma - \delta; n + 2 - \gamma; \frac{(a-1)x}{a-x}\right] \stackrel{(21)}{=} H^{(6)}(x) \Big|_{\alpha \leftrightarrow \beta}. \quad (88)$$

In this group the hypergeometric functions converge in the region $|(a-1)x/(a-x)| < 1$ which includes only one singular point ($x=0$); in accordance with (20), a restriction on δ assure convergence also on $|(a-1)x/(a-x)| = 1$ (includes $x=1$). Precisely,

$$\text{hypergeometric functions in } H^{(i)} \text{ converge on } \left|\frac{(a-1)x}{(a-x)}\right| = 1 \text{ if } \operatorname{Re} \delta < 1. \quad (89)$$

In fact, in view of (21) there are two forms for each expansion in series of hypergeometric functions, but only one of them is suitable for $|z| = 1$. For example,

$$H^{(1)}(x) = \left(1 - \frac{x}{a}\right)^{-\alpha} \sum_{n=0}^{\infty} b_n^{(1)} \left[\frac{(a-1)x}{a-x}\right]^n \tilde{F} \left[n + \alpha, \gamma + \delta - \alpha - 1; n + \gamma; \frac{(a-1)x}{a-x}\right]$$

$$\stackrel{(21)}{=} (1-x)^{1-\delta} \left(1 - \frac{x}{a}\right)^{\delta-1-\alpha} \sum_{n=0}^{\infty} b_n^{(1)} \left[\frac{(a-1)x}{a-x}\right]^n \tilde{F} \left[n + 1 + \alpha - \delta, \gamma - \alpha; n + \gamma; \frac{(a-1)x}{a-x}\right].$$

In the first expression, the hypergeometric function converges on $x = 1$ if $\text{Re } \delta < 1$ while the factor $(1 - x/a)^{-\alpha}$ is finite because $a \neq 0, 1$. In the second expression, the hypergeometric function converges on $x = 1$ only if $\text{Re } \delta > 1$, but in this case the factor $(1-x)^{1-\delta}$ in general becomes infinite at $x = 1$. Thus, the second form is inappropriate to get the behaviour at $x = 1$.

We find that, for the expansions $H^{(i)}(x)$,

$$\begin{aligned} a = 1 & \Rightarrow \alpha_n^{(i)} = 0; \\ a = 2, \alpha + \beta + 1 = 2\gamma + \delta, q = \alpha\beta & \Rightarrow \beta_n^{(1)} = \beta_n^{(3)} = \beta_n^{(6)} = \beta_n^{(8)} = 0; \\ a = 2, \alpha + \beta = 1 + \delta, q = \alpha\beta + (\gamma - 1)\delta & \Rightarrow \beta_n^{(2)} = \beta_n^{(4)} = \beta_n^{(5)} = \beta_n^{(7)} = 0, \end{aligned} \quad (90)$$

which reproduce the conditions given in (70) for reducing the Heun equation to a hypergeometric equation.

4.3 The solutions $H^{(i)}(x)$ around $x = 1$

These arise from the solution $\bar{H}^{(1)}(x)$, given in (79), by means of the following transformations: $\mathbf{H}^{(1)}(x) = M_{49}\bar{H}^{(1)}(x)$ and $\mathbf{H}^{(i)}(x) = T_i\mathbf{H}^{(1)}(x)$.

$$\left\{ \begin{aligned} \mathbf{H}^{(1)}(x) &= \sum_{n=0}^{\infty} \mathbf{b}_n^{(1)} (1-x)^n \tilde{F}(n + \alpha, \gamma + \delta - \alpha - 1; n + \delta; 1-x), \\ \alpha_n^{(1)} &= (1-a)(n+1), \quad \beta_n^{(1)} = (a-2)n^2 + [(a-1)(2\alpha+1-\gamma-\delta) - \\ &\quad \alpha - \beta + \gamma]n + q + a\alpha(\alpha+1-\gamma-\delta) - \alpha\epsilon, \\ \gamma_n^{(1)} &= (n+\alpha-1)(n+\alpha-\gamma)(n+\alpha+\beta-\gamma-\delta). \end{aligned} \right. \quad (91)$$

$$\begin{aligned} \mathbf{H}^{(2)}(x) &= x^{1-\gamma} \sum_{n=0}^{\infty} \mathbf{b}_n^{(2)} (1-x)^n \tilde{F}(n + \beta + 1 - \gamma, \delta - \beta; n + \delta; 1-x) \\ &\stackrel{(21)}{=} \mathbf{H}^{(1)}(x) \Big|_{\alpha \leftrightarrow \beta}. \end{aligned} \quad (92)$$

$$\left\{ \begin{aligned} \mathbf{H}^{(3)}(x) &= (1-x)^{1-\delta} \sum_{n=0}^{\infty} \mathbf{b}_n^{(3)} (1-x)^n \times \\ &\quad \tilde{F}(n + 1 + \beta - \delta, \gamma - \beta; n + 2 - \delta; 1-x), \\ \alpha_n^{(3)} &= (1-a)(n+1), \quad \beta_n^{(3)} = (a-2)n^2 + [(a-1)(1+2\beta-\gamma-\delta) - \alpha - \\ &\quad \beta + \gamma + 2\delta - 2]n + q + a\beta(\beta-\gamma-\delta+1) - (\beta-\delta+1)(\alpha+\beta+1-\gamma-\delta), \\ \gamma_n^{(3)} &= (n+\beta-\delta)(n+1+\beta-\gamma-\delta)(n+\alpha+\beta-\gamma-\delta), \end{aligned} \right. \quad (93)$$

$$\begin{aligned} \mathbf{H}^{(4)}(x) &= x^{1-\gamma}(1-x)^{1-\delta} \sum_{n=0}^{\infty} \mathbf{b}_n^{(4)}(1-x)^n \times \\ &\quad \tilde{F}(n+2+\alpha-\gamma-\delta, 1-\alpha; n+2-\delta; 1-x) \stackrel{(21)}{=} \mathbf{H}^{(3)}(x) \Big|_{\alpha \leftrightarrow \beta}. \end{aligned} \quad (94)$$

$$\left\{ \begin{aligned} \mathbf{H}^{(5)}(x) &= \left(1 - \frac{x}{a}\right)^{1-\epsilon} \sum_{n=0}^{\infty} \mathbf{b}_n^{(5)}(1-x)^n \tilde{F}(n-\alpha+\gamma+\delta, \alpha-1; n+\delta; 1-x), \\ \alpha_n^{(5)} &= (1-a)(n+1), \quad \beta_n^{(5)} = (a-2)n^2 + [(a-1)(1-2\alpha+\gamma+\delta) + \alpha + \\ &\quad \beta - \gamma - 2\delta]n + q + a(\alpha - \gamma - \delta)(\alpha - 1) - (\alpha - \delta)(\alpha + \beta - \gamma - \delta - 1) - \gamma, \\ \gamma_n^{(5)} &= (n-1-\alpha+\gamma+\delta)(n-\alpha+\delta)(n-\alpha-\beta+\gamma+\delta). \end{aligned} \right. \quad (95)$$

$$\begin{aligned} \mathbf{H}^{(6)}(x) &= x^{1-\gamma} \left(1 - \frac{x}{a}\right)^{1-\epsilon} \sum_{n=0}^{\infty} \mathbf{b}_n^{(6)}(1-x)^n \tilde{F}(n+1-\beta+\delta, \beta-\gamma; n+\delta; 1-x) \\ &\stackrel{(21)}{=} \mathbf{H}^{(5)}(x) \Big|_{\alpha \leftrightarrow \beta}. \end{aligned} \quad (96)$$

$$\left\{ \begin{aligned} \mathbf{H}^{(7)}(x) &= (1-x)^{1-\delta} \left[1 - \frac{x}{a}\right]^{1-\epsilon} \sum_{n=0}^{\infty} \mathbf{b}_n^{(7)}(1-x)^n \times \\ &\quad \tilde{F}(n-\beta+\gamma+1, \beta-\delta; n+2-\delta; 1-x), \\ \alpha_n^{(7)} &= (1-a)(n+1), \quad \beta_n^{(7)} = (a-2)n^2 + [(a-1)(1-2\beta+\gamma+\delta) + \alpha + \\ &\quad \beta - \gamma - 2]n + q + a(\beta - \gamma - \delta)(\beta - 1) - (\beta - 1)(\alpha + \beta - \gamma - \delta - 1) - \gamma, \\ \gamma_n^{(7)} &= (n-\beta+\gamma)(n+1-\beta)(n-\alpha-\beta+\gamma+\delta). \end{aligned} \right. \quad (97)$$

$$\begin{aligned} \mathbf{H}^{(8)}(x) &= x^{1-\gamma}(1-x)^{1-\delta} \left[1 - \frac{x}{a}\right]^{1-\epsilon} \sum_{n=0}^{\infty} \mathbf{b}_n^{(8)}(1-x)^n \times \\ &\quad \tilde{F}(n+2-\alpha, \alpha+1-\gamma-\delta; n+2-\delta; 1-x) \stackrel{(21)}{=} \mathbf{H}^{(7)}(x) \Big|_{\alpha \leftrightarrow \beta}. \end{aligned} \quad (98)$$

The hypergeometric functions in $\mathbf{H}^{(i)}(x)$ converge in the region $|x-1| < 1$ but, by (20),

hypergeometric functions in $\mathbf{H}^{(i)}$ converge on $|x-1| = 1$ if $\text{Re } \gamma < 1$. (99)

For the above expansions, $\mathbf{H}^{(i)}(x)$,

$$\begin{aligned} a = 1 &\quad \Rightarrow \quad \alpha_n^{(i)} = 0; \\ a = 2, \quad \beta = \alpha + 1 - \delta, \quad q = \alpha\gamma &\quad \Rightarrow \quad \beta_n^{(1)} = \beta_n^{(3)} = \beta_n^{(6)} = \beta_n^{(8)} = 0; \\ a = 2, \quad \alpha = \beta + 1 - \delta, \quad q = \beta\gamma &\quad \Rightarrow \quad \beta_n^{(2)} = \beta_n^{(4)} = \beta_n^{(5)} = \beta_n^{(7)} = 0. \end{aligned} \quad (100)$$

Thence, Eqs.(32) become two-term recurrence relations for the series coefficients and the Heun equation (2) is reducible to the hypergeometric equation (3) (Appendix B).

5 Schrödinger equation for associated Lamé potentials

There are some periodic potentials for which the Schrödinger equation leads to equations of Heun family [17, 18, 19, 20, 37, 38]. Solutions for these equations in general demand numerical computations. In this section we consider only some analytical aspects concerning the solutions of the Schrödinger equation for the associated Lamé potential (17) which depends on the parameters \mathbf{m} and \mathbf{l} .

We will find that the solutions of the general Heun equation allow to solve this problem as follows:

- if $\mathbf{m} - \mathbf{l}$ or $\mathbf{l} + \mathbf{m}$ is an integer, we get finite series from the power series expansions given in Section 3; further restrictions on \mathbf{m} and \mathbf{l} assure that these solutions satisfy the Arscott criterion (36);
- if either \mathbf{l} or \mathbf{m} is an integer and the other is half an odd integer, we get finite series from the expansions in series of hypergeometric functions given in Section 4; such solutions are degenerate and satisfy the Arscott criterion only for special values of \mathbf{l} and \mathbf{m} ;
- bounded infinite-series solutions are obtained from the expansions given in section 3.1.

The finite-series solutions are valid under general restrictions on the parameters, that is, without regarding particular values for \mathbf{m} and \mathbf{l} (some specific values are discussed in Appendix C). Initially, we explain how we obtain the several types of solutions and set up some notations for the solutions.

The Schrödinger equation (14) for the associated Lamé potential (17) is

$$\frac{d^2\psi(u)}{du^2} + \left[\mathcal{E} - \mathbf{m}(\mathbf{m} + 1)k^2 \operatorname{sn}^2 u - \mathbf{l}(\mathbf{l} + 1)k^2 \frac{\operatorname{cn}^2 u}{\operatorname{dn}^2 u} \right] \psi(u) = 0, \quad (101)$$

which is the Darboux Eq. (11) with parameters ($a = 1/k^2$)

$$\alpha = \frac{\mathbf{l}-\mathbf{m}+1}{2}, \quad \beta = \frac{\mathbf{l}+\mathbf{m}+2}{2}, \quad \gamma = \delta = \frac{1}{2}, \quad q = \frac{(\mathbf{l}+1)^2}{4} - \frac{\mathcal{E}}{4k^2}, \quad (\epsilon = \mathbf{l} + \frac{3}{2}). \quad (102)$$

Thence, the eigenfunctions $\psi(u)$ follow from solutions $H(x)$ of the Heun equation through

$$\psi(u) \stackrel{\text{Eq. (19)}}{=} \operatorname{dn}^{\mathbf{l}+1} u H[x(u)], \quad 0 \leq x(u) = \operatorname{sn}^2 u \leq 1, \quad (103)$$

provided that $H(x)$ is a convergent series and $\psi(u)$ is bounded for any value of u . In addition, we impose the Arscott condition $\alpha_{n-1}\gamma_n > 0$ for finite series with two or more terms ($n \geq 1$).

The associated Lamé equation (101) remains invariant under the simultaneous substitutions

$$\mathbf{m} \mapsto -\mathbf{m} - 1, \quad \mathbf{l} \mapsto -\mathbf{l} - 1 \quad (\text{leave Eq. (101) invariant}) \quad (104)$$

which also leave invariant the solutions constructed according to (103), as we will see. Alternatively, (104) can be used to set up a new solution out of a given solution. Further, according to (30), the interchange of \mathbf{l} and \mathbf{m} in conjunction with the shift of u to $u + K$ leaves the associated Lamé equation invariant, that is,

$$u \mapsto u + K, \quad \mathbf{m} \leftrightarrow \mathbf{l} \quad (\text{leave Eq. (101) invariant}). \quad (105)$$

This substitutions may be used to get new solutions to the associated Lamé equation by means of the relations [3, 19]

$$\operatorname{sn}(u + K) = \operatorname{cd} u, \quad \operatorname{cn}(u + K) = -\sqrt{1 - k^2} \operatorname{sd} u, \quad \operatorname{dn}(u + K) = \frac{\sqrt{1 - k^2}}{\operatorname{dn} u}. \quad (106)$$

Thence, power series of $\operatorname{sn} u$ and $\operatorname{cn} u$ are transformed, respectively, in power series of $\operatorname{cd} u$ and $\operatorname{sd} u$. To obtain these series directly from solutions of the Heun equation it would be necessary to construct two additional groups by the fractional transformations M_{17} and M_{65} as we see from (30). Notice that the particular cases

$$\frac{d^2 \psi(u)}{du^2} + \left[\mathcal{E} - \mathbf{l}(\mathbf{l} + 1)k^2 \frac{\operatorname{cn}^2 u}{\operatorname{dn}^2 u} \right] \psi(u) = 0, \quad [\mathbf{m} = 0 \text{ or } \mathbf{m} = -1], \quad \text{and} \quad (107)$$

$$\frac{d^2 \psi(u)}{du^2} + [\mathcal{E} - \mathbf{m}(\mathbf{m} + 1)k^2 \operatorname{sn}^2 u] \psi(u) = 0, \quad [\mathbf{l} = 0 \text{ or } \mathbf{l} = -1 : \text{Lamé equation}], \quad (108)$$

are connected by (105). We will see that finite-series solutions which satisfy the Arscott condition for the first case do not satisfy for the second case, and vice-versa.

For comparison purposes, we write the Khare-Sukhatme notation [19, 20] for the parameters \mathbf{l} , \mathbf{m} and k^2 , namely,

$$\mathbf{l} = b, \quad \mathbf{m} = a, \quad k^2 = m : \quad \text{Khare-Sukhatme [19, 20]}. \quad (109)$$

According to (34), finite series result when $\gamma_n = 0$ for some value of n . For each power-series expansion the condition $\gamma_n = 0$ gives two possibilities for truncating the series on the right-hand side. For solutions obtained from $\mathring{H}^{(1)}$ and $\mathring{H}^{(1)}$, the two possibilities result from the fact that $(\mathring{\gamma}_n^{(1)} = \mathring{\gamma}_n^{(1)})$

$$\mathring{\gamma}_n^{(1)} = \left(n - \frac{\mathbf{m} - \mathbf{l} + 1}{2}\right) \left(n + \frac{\mathbf{l} + \mathbf{m}}{2}\right) = 0 \text{ if } \begin{cases} \mathbf{m} - \mathbf{l} = 1, 3, 5, \dots \Rightarrow 0 \leq n \leq \frac{\mathbf{m} - \mathbf{l} - 1}{2} \text{ or} \\ \mathbf{m} + \mathbf{l} = -2, -4, \dots \Rightarrow 0 \leq n \leq -\frac{\mathbf{m} + \mathbf{l} + 2}{2}. \end{cases} \quad (110)$$

For $\mathbf{m} = -1/2$, the two cases become identical. For $\mathbf{m} \neq -1/2$, if the above conditions on $\mathbf{m} - \mathbf{l}$ and $\mathbf{m} + \mathbf{l}$ are satisfied for the same values of \mathbf{m} and \mathbf{l} , thence we have to consider only the truncation which gives the small interval for n . The conditions for this will be written in relations (115c), (116c), \dots , (122c). Such conditions imply that both \mathbf{l} and \mathbf{m} are half an integer number and, consequently, they do not stand for solutions of Eqs. (107) and (108).

Solutions resulting from $\mathring{H}^{(i)}$ and $\mathring{H}^{(i)}$ are expansions in series of $\operatorname{sn}^2 u$ and $\operatorname{cn}^2 u$, respectively. We will find that finite series in terms of $\operatorname{sn}^2 u$ and $\operatorname{cn}^2 u$ cannot satisfy simultaneously the Arscott condition because this condition requires that $\mathring{\gamma}_n^{(i)} > 0$ and $\mathring{\gamma}_n^{(i)} < 0$, respectively. For the present case, the Arscott condition is important for assuring that the characteristic equation (35) yields real and distinct values for the energy \mathcal{E} .

We use the following notations for the solutions of Eq. (101):

- (1) the pairs $(\mathring{\psi}^{(i)}, \tilde{\psi}^{(i)})$ denote finite series of $\operatorname{sn}^2 u$, obtained from truncation of the powers series $\mathring{H}^{(i)}$;
- (2) the pairs $(\mathring{\Psi}^{(i)}, \tilde{\Psi}^{(i)})$ denote finite series of $\operatorname{cn}^2 u$, obtained from truncation of the powers series $\mathring{H}^{(i)}$;
- (3) $\psi^{(i)}$ ($i = 1, 2, \dots, 8$) denote finite series of hypergeometric functions obtained from expansions $H^{(i)}$ of Sec. 4.2;
- (4) $\Psi^{(i)}$ denote finite series of hypergeometric functions resulting from expansions $H^{(i)}$ of Sec. 4.3;

(5) $\mathring{\Phi}^{(i)}$ denote infinite-series solutions in terms of $\text{sn}^2 u$ whose linear dependence requires further investigation.

Each of the above items are discussed in the next subsections. We will use the following parity and periodicity properties for elliptic functions [3]:

$$\operatorname{sn}(-u) = -\operatorname{sn} u, \quad \operatorname{cn}(-u) = \operatorname{cn} u, \quad \operatorname{dn}(-u) = \operatorname{dn} u. \quad (111)$$

$$\operatorname{sn}(u + 2K) = -\operatorname{sn} u, \quad \operatorname{cn}(u + 2K) = -\operatorname{cn} u, \quad \operatorname{dn}(u + 2K) = \operatorname{dn} u. \quad (112)$$

As in the case of Lamé equation [33], these properties give four types of solutions for the associated Lamé equation: even and odd solutions with period $2K$, even and odd solutions with period $4K$. In Sections 5.1 and 5.2, solutions of each type are put together. We also take into account the special values

$$\operatorname{sn} 0 = 0, \quad \operatorname{sn} K = 1, \quad \operatorname{cn} 0 = 1, \quad \operatorname{cn} K = 0, \quad \operatorname{dn} 0 = 1, \quad \operatorname{dn} K = 1 - k^2. \quad (113)$$

5.1 Finite-series eigenfunctions $(\psi^{(i)}, \tilde{\psi}^{(i)})$ in terms of $\text{sn}^2 u$

For the associated Lamé equation (101), the power series solutions $\mathring{H}^{(i)}(x)$ lead to eight pairs of finite-series expansions $\left(\mathring{\psi}^{(i)}, \tilde{\psi}^{(i)}\right)$ by means of the truncation procedure indicated in (110). To write down the four types of expansions we observe that the pairs $\left(\mathring{\psi}^{(j+4)}, \tilde{\psi}^{(j+4)}\right)$ ($j = 1, 2, 3, 4$) can be generated from $\left(\mathring{\psi}^{(j)}, \tilde{\psi}^{(j)}\right)$ by the substitutions $\boldsymbol{l} \mapsto -\boldsymbol{l} - 1$ and $\boldsymbol{m} \mapsto -\boldsymbol{m} - 1$ which preserve the period and parity of the expansions.

The series coefficients for $\left(\mathring{\psi}^{(i)}, \tilde{\psi}^{(i)}\right)$ are denoted by $\left(\mathring{b}_n^{(i)}, \tilde{b}_n^{(i)}\right)$. However, in the recurrence relations $\mathring{\alpha}_n \mathring{b}_{n+1} + \mathring{\beta}_n \mathring{b}_n + \mathring{\gamma}_n \mathring{b}_{n-1} = 0$ and $\tilde{\alpha}_n \tilde{b}_{n+1} + \tilde{\beta}_n \tilde{b}_n + \tilde{\gamma}_n \tilde{b}_{n-1} = 0$, the coefficients are formally identical, that is,

$$\tilde{\alpha}_n^{(i)} = \mathring{\alpha}_n^{(i)}, \quad \tilde{\beta}_n^{(i)} = \mathring{\beta}_n^{(i)}, \quad \tilde{\gamma}_n^{(i)} = \mathring{\gamma}_n^{(i)}. \quad (114)$$

On the right-hand side of each expansion, inside square brackets, we write the restrictions on the parameters \mathbf{l} and \mathbf{m} that assure the Arscott condition (36).

First we write the four types of expansions, then we discuss the Arscott condition (36) and, finally, we exhibit some problems which can be solved by these expansions.

5.1.1 First type: Even solutions, period $2K$

$$\left\{ \begin{array}{ll} \psi^{(1)}(u) = \operatorname{dn}^{l+1} u \sum_{n=0}^{\frac{m-l-1}{2}} \hat{b}_n^{(1)} \operatorname{sn}^{2n} u & \text{if } m-l = 1, 3, 5, \dots, \\ & [m \leq -\frac{1}{2}, m+l < -2], \\ \tilde{\psi}^{(1)}(u) = \operatorname{dn}^{l+1} u \sum_{n=0}^{\frac{l+m+2}{2}} \tilde{b}_n^{(1)} \operatorname{sn}^{2n} u & \text{if } m+l = -2, -4, -6, \dots, \\ & [m \geq -\frac{1}{2}, m-l > 1], \end{array} \right. \quad (115a)$$

$$\begin{aligned} \alpha_n^{(1)} &= \frac{1}{k^2} \left(n + \frac{1}{2} \right) (n + 1), & \beta_n^{(1)} &= \left(\frac{1}{k^2} + 1 \right) n^2 - (l + 1)n + \\ & \frac{1}{4k^2} \mathcal{E} - \frac{1}{4} (l + 1)^2, & \gamma_n^{(1)} &= \left(n - \frac{m-l+1}{2} \right) \left(n + \frac{l+m}{2} \right); \end{aligned} \quad (115b)$$

Only one of the above expansions is valid when (from now on, M_i and N_i denote positive integers)

$$\begin{aligned} \mathbf{m} - \mathbf{l} - 1 = 2M_1 \text{ and } \mathbf{m} + \mathbf{l} + 2 = -2N_1, \quad (\Leftrightarrow \mathbf{m} = M_1 - N_1 - \tfrac{1}{2}, \\ \mathbf{l} = -M_1 - N_1 - \tfrac{3}{2}) : \psi^{(1)} \text{ if } M_1 < N_1, \quad \tilde{\psi}^{(1)} \text{ if } N_1 < M_1. \end{aligned} \quad (115c)$$

For series with two or more terms ($M_1 \geq 1$ or $N_1 \geq 1$), we see that these expansions satisfy the Arscott criterion. The same holds for the other seven cases given below: (116c), (117c), \dots , (122c).

$$\left\{ \begin{aligned} \psi^{(5)}(u) &= \operatorname{dn}^{-l} u \sum_{n=0}^{\frac{l-m-1}{2}} \mathring{b}_n^{(5)} \operatorname{sn}^{2n} u \quad \text{if} \quad \mathbf{l} - \mathbf{m} = 1, 3, 5, \dots, \\ &\quad [\mathbf{m} \geq -\tfrac{1}{2}, \mathbf{m} + \mathbf{l} > 0], \\ \tilde{\psi}^{(5)}(u) &= \operatorname{dn}^{-l} u \sum_{n=0}^{\frac{l+m}{2}} \tilde{b}_n^{(5)} \operatorname{sn}^{2n} u \quad \text{if} \quad \mathbf{m} + \mathbf{l} = 0, 2, 4, \dots, \\ &\quad [\mathbf{m} \leq -\tfrac{1}{2}, \mathbf{l} - \mathbf{m} > 1], \end{aligned} \right. \quad (116a)$$

$$\begin{aligned} \mathring{\alpha}_n^{(5)} &= \frac{1}{k^2} \left(n + \tfrac{1}{2}\right) (n+1), \quad \mathring{\beta}_n^{(5)} = -\left(\frac{1}{k^2} + 1\right) n^2 + \mathbf{l}n + \frac{1}{4k^2} \mathcal{E} - \tfrac{1}{4} \mathbf{l}^2, \\ \mathring{\gamma}_n^{(5)} &= \left(n - \frac{l-m+1}{2}\right) \left(n - \frac{l+m+2}{2}\right). \end{aligned} \quad (116b)$$

Only one expansion when

$$\begin{aligned} \mathbf{l} - \mathbf{m} - 1 = 2M_2 \text{ and } \mathbf{m} + \mathbf{l} = 2N_2, \quad (\Leftrightarrow \mathbf{m} = N_2 - M_2 - \tfrac{1}{2}, \\ \mathbf{l} = -M_2 + N_2 + \tfrac{1}{2}) : \psi^{(5)} \text{ if } M_2 < N_2, \quad \tilde{\psi}^{(5)} \text{ if } N_2 < M_2. \end{aligned} \quad (116c)$$

5.1.2 Second type: Odd solutions, period $4K$

$$\left\{ \begin{aligned} \psi^{(2)}(u) &= \operatorname{sn} u \operatorname{dn}^{l+1} u \sum_{n=0}^{\frac{m-l-2}{2}} \mathring{b}_n^{(2)} \operatorname{sn}^{2n} u \quad \text{if} \quad \mathbf{m} - \mathbf{l} = 2, 4, 6, \dots, \\ &\quad [\mathbf{m} \leq -\tfrac{1}{2}, \mathbf{m} + \mathbf{l} < -3], \\ \tilde{\psi}^{(2)}(u) &= \operatorname{sn} u \operatorname{dn}^{l+1} u \sum_{n=0}^{-\frac{m+l+3}{2}} \tilde{b}_n^{(2)} \operatorname{sn}^{2n} u \quad \text{if} \quad \mathbf{m} + \mathbf{l} = -3, -5, -7, \dots, \\ &\quad [\mathbf{m} \geq -\tfrac{1}{2}, \mathbf{m} - \mathbf{l} > 2], \end{aligned} \right. \quad (117a)$$

$$\begin{aligned} \mathring{\alpha}_n^{(2)} &= \frac{1}{k^2} \left(n + \tfrac{3}{2}\right) (n+1), \quad \mathring{\beta}_n^{(2)} = -\left(\frac{1}{k^2} + 1\right) n^2 - \left(\frac{1}{k^2} + \mathbf{l} + 2\right) n + \\ &\quad \frac{1}{4k^2} (\mathcal{E} - 1) - \tfrac{1}{4} (\mathbf{l} + 2)^2, \quad \mathring{\gamma}_n^{(2)} = \left(n - \frac{m-l}{2}\right) \left(n + \frac{l+m+1}{2}\right); \end{aligned} \quad (117b)$$

Only one expansion when

$$\begin{aligned} \mathbf{m} - \mathbf{l} - 2 = 2M_3 \text{ and } \mathbf{m} + \mathbf{l} + 3 = -2N_3, \quad (\Leftrightarrow \mathbf{m} = M_3 - N_3 - \tfrac{1}{2}, \\ \mathbf{l} = -M_3 - N_3 - \tfrac{5}{2}) : \psi^{(2)} \text{ if } M_3 < N_3, \quad \tilde{\psi}^{(2)} \text{ if } N_3 < M_3. \end{aligned} \quad (117c)$$

$$\left\{ \begin{array}{ll} \psi^{(6)}(u) = \operatorname{sn} u \operatorname{dn}^{-l} u \sum_{n=0}^{\frac{l-m-2}{2}} \mathring{b}_n^{(6)} \operatorname{sn}^{2n} u & \text{if } l-m = 2, 4, 6, \dots, \\ & [m \geq -\frac{1}{2}, m+l > 1], \\ \tilde{\psi}^{(6)}(u) = \operatorname{sn} u \operatorname{dn}^{-l} u \sum_{n=0}^{\frac{m+l-1}{2}} \tilde{b}_n^{(6)} \operatorname{sn}^{2n} u & \text{if } m+l = 1, 3, 5, \dots, \\ & [m \leq -\frac{1}{2}, l-m > 2], \end{array} \right. \quad (118a)$$

$$\begin{aligned} \mathring{\alpha}_n^{(6)} &= \frac{1}{k^2} \left(n + \frac{3}{2}\right) (n+1), & \mathring{\beta}_n^{(6)} &= -\left(\frac{1}{k^2} + 1\right) n^2 - \left(\frac{1}{k^2} - l + 1\right) n + \\ & \frac{1}{4k^2} (\mathcal{E} - 1) - \frac{1}{4} (l-1)^2, & \mathring{\gamma}_n^{(6)} &= \left(n - \frac{l-m}{2}\right) \left(n - \frac{l+m+1}{2}\right), \end{aligned} \quad (118b)$$

Only one expansion when

$$\begin{aligned} l-m-2 &= 2M_4 \text{ and } m+l-1 = 2N_4 \quad (\Leftrightarrow m = N_4 - M_4 - \frac{1}{2}, \\ l &= M_4 + N_4 + \frac{3}{2}) : \psi^{(6)} \text{ if } M_4 < N_4, \quad \tilde{\psi}^{(6)} \text{ if } N_4 < M_4. \end{aligned} \quad (118c)$$

5.1.3 Third type: Even solutions, period $4K$

$$\left\{ \begin{array}{ll} \psi^{(3)}(u) = \operatorname{cn} u \operatorname{dn}^{l+1} u \sum_{n=0}^{\frac{m-l-2}{2}} \mathring{b}_n^{(3)} \operatorname{sn}^{2n} u & \text{if } m-l = 2, 4, 6, \dots, \\ & [m \leq -\frac{1}{2}, m+l < -3] \\ \tilde{\psi}^{(3)}(u) = \operatorname{cn} u \operatorname{dn}^{l+1} u \sum_{n=0}^{-\frac{m+l+3}{2}} \tilde{b}_n^{(3)} \operatorname{sn}^{2n} u & \text{if } m+l = -3, -5, -7, \dots, \\ & [m \geq -\frac{1}{2}, m-l > 2], \end{array} \right. \quad (119a)$$

$$\begin{aligned} \mathring{\alpha}_n^{(3)} &= \frac{1}{k^2} \left(n + \frac{3}{2}\right) (n+1), & \mathring{\beta}_n^{(3)} &= -\left(\frac{1}{k^2} + 1\right) n^2 - \left(\frac{1}{k^2} + l + 1\right) n + \\ & \frac{1}{4k^2} (\mathcal{E} - 1) - \frac{1}{4} (l+1)^2, & \mathring{\gamma}_n^{(3)} &= \mathring{\gamma}_n^{(2)}. \end{aligned} \quad (119b)$$

Only one expansion when

$$\begin{aligned} m-l-2 &= 2M_3 \text{ and } m+l+3 = -2N_3 \quad (\Leftrightarrow m = M_3 - N_3 - \frac{1}{2}, \\ l &= -M_3 - N_3 - \frac{5}{2}) : \psi^{(3)} \text{ if } M_3 < N_3, \quad \tilde{\psi}^{(3)} \text{ if } N_3 < M_3. \end{aligned} \quad (119c)$$

$$\left\{ \begin{array}{ll} \psi^{(7)}(u) = \operatorname{cn} u \operatorname{dn}^{-l} u \sum_{n=0}^{\frac{l-m-2}{2}} \mathring{b}_n^{(7)} \operatorname{sn}^{2n} u & \text{if } l-m = 2, 4, 6, \dots, \\ & [m \geq -\frac{1}{2}, m+l > 1], \\ \tilde{\psi}^{(7)}(u) = \operatorname{cn} u \operatorname{dn}^{-l} u \sum_{n=0}^{\frac{m+l-1}{2}} \tilde{b}_n^{(7)} \operatorname{sn}^{2n} u & \text{if } m+l = 1, 3, 5, \dots, \\ & [m \leq -\frac{1}{2}, l-m > 2], \end{array} \right. \quad (120a)$$

$$\begin{aligned}\alpha_n^{(7)} &= \frac{1}{k^2} \left(n + \frac{3}{2}\right) (n+1), & \beta_n^{(7)} &= -\left(\frac{1}{k^2} + 1\right) n^2 - \left(\frac{1}{k^2} - l\right) n + \\ & \frac{1}{4k^2}(\mathcal{E} - 1) - \frac{1}{4}l^2, & \gamma_n^{(7)} &= \gamma_n^{(6)}.\end{aligned}\quad (120b)$$

Only one expansion when

$$\begin{aligned}l - m - 2 &= 2M_4 \quad \text{and} \quad m + l - 1 = 2N_4 \quad (\Leftrightarrow m = N_4 - M_4 - \frac{1}{2}, \\ l &= M_2 + N_4 + \frac{3}{2}) : \psi^{(7)} \text{ if } M_4 < N_4, \quad \tilde{\psi}^{(7)} \text{ if } N_4 < M_4.\end{aligned}\quad (120c)$$

5.1.4 Fourth type: Odd solutions, period $2K$

$$\left\{ \begin{aligned} \psi^{(4)}(u) &= \text{sn} u \, \text{cn} u \, \text{dn}^{l+1} u \sum_{n=0}^{\frac{m-l-3}{2}} \mathring{b}_n^{(4)} \text{sn}^{2n} u \quad \text{if} \quad m - l = 3, 5, 7, \dots, \quad ; \\ & \quad [m \leq -\frac{1}{2}, \quad m + l < -4] \\ \tilde{\psi}^{(4)}(u) &= \text{sn} u \, \text{cn} u \, \text{dn}^{l+1} u \sum_{n=0}^{-\frac{m+l+4}{2}} \tilde{b}_n^{(4)} \text{sn}^{2n} u \quad \text{if} \quad m + l = -4, -6, -8, \dots, \quad ; \\ & \quad [m \geq -\frac{1}{2}, \quad m - l > 3] \end{aligned} \right. \quad (121a)$$

$$\begin{aligned}\alpha_n^{(4)} &= \frac{1}{k^2} \left(n + \frac{3}{2}\right) (n+1), & \beta_n^{(4)} &= -\left(\frac{1}{k^2} + 1\right) n^2 - \left(\frac{2}{k^2} + l + 2\right) n + \\ & \frac{1}{4k^2}(\mathcal{E} - 4) - \frac{1}{4}(l + 2)^2, & \gamma_n^{(4)} &= \left(n - \frac{m-l-1}{2}\right) \left(n + \frac{l+m+2}{2}\right);\end{aligned}\quad (121b)$$

Only one expansion when

$$\begin{aligned}m - l - 3 &= 2M_5 \quad \text{and} \quad m + l + 4 = -2N_5, \quad (\Leftrightarrow m = M_5 - N_5 - \frac{1}{2}, \\ l &= -M_5 - N_5 - \frac{3}{2}) : \psi^{(4)} \text{ if } M_5 < N_5, \quad \tilde{\psi}^{(4)} \text{ if } N_5 < M_5.\end{aligned}\quad (121c)$$

$$\left\{ \begin{aligned} \psi^{(8)}(u) &= \text{sn} u \, \text{cn} u \, \text{dn}^{-l} u \sum_{n=0}^{\frac{l-m-3}{2}} \mathring{b}_n^{(8)} \text{sn}^{2n} u \quad \text{if} \quad l - m = 3, 5, 7, \dots, \quad ; \\ & \quad [m \geq -\frac{1}{2}, \quad m + l > 2], \\ \tilde{\psi}^{(8)}(u) &= \text{sn} u \, \text{cn} u \, \text{dn}^{-l} u \sum_{n=0}^{\frac{m+l-2}{2}} \tilde{b}_n^{(8)} \text{sn}^{2n} u \quad \text{if} \quad m + l = 2, 4, 6, \dots, \quad ; \\ & \quad [m \leq -\frac{1}{2}, \quad l - m > 3], \end{aligned} \right. \quad (122a)$$

$$\begin{aligned}\alpha_n^{(8)} &= \frac{1}{k^2} \left(n + \frac{3}{2}\right) (n+1), & \beta_n^{(5)} &= -\left(\frac{1}{k^2} + 1\right) n^2 - \left(\frac{2}{k^2} - l + 1\right) n + \\ & \frac{1}{4k^2}(\mathcal{E} - 4) - \frac{1}{4}(l - 1)^2, & \gamma_n^{(8)} &= \left(n - \frac{l-m-1}{2}\right) \left(n - \frac{l+m}{2}\right).\end{aligned}\quad (122b)$$

Only one expansion when

$$\begin{aligned}l - m - 3 &= 2M_6 \quad \text{and} \quad m + l - 2 = -2N_6 \quad (\Leftrightarrow m = N_6 - M_6 - \frac{1}{2}, \\ l &= M_6 + N_6 + \frac{5}{2}) : \psi^{(8)} \text{ if } M_6 < N_6, \quad \tilde{\psi}^{(8)} \text{ if } N_6 < M_6.\end{aligned}\quad (122c)$$

Due to the symmetry under the substitutions (104), we discuss the Arscott condition (36) only for $(\psi^{(i)}, \tilde{\psi}^{(i)})$ with $i = 1, 2, 3, 4$. As the $\alpha_n^{(i)}$ are positive, the condition $\alpha_{n-1}\gamma_n > 0$ is satisfied if $\gamma_n > 0$ for any admissible n . So, for $\psi^{(i)}$ we requires that

$$\begin{aligned}\gamma_n^{(1)} &= \left(n - \frac{m-l+1}{2}\right) \left(n + \frac{l+m}{2}\right) > 0 \text{ for } 1 \leq n \leq \frac{m-l-1}{2}, \quad [m-l = 3, 5, 7, \dots], \\ \gamma_n^{(2)} &= \gamma_n^{(3)} = \left(n - \frac{m-l}{2}\right) \left(n + \frac{l+m+1}{2}\right) > 0 \text{ for } 1 \leq n \leq \frac{m-l-2}{2}, \quad [m-l = 4, 6, 8, \dots], \\ \gamma_n^{(4)} &= \left(n - \frac{m-l-1}{2}\right) \left(n + \frac{l+m+2}{2}\right) > 0 \text{ for } 1 \leq n \leq \frac{m-l-3}{2}, \quad [m-l = 5, 7, 9, \dots].\end{aligned}$$

Thus, since the first term is negative, $\gamma_n > 0$ if the last term on the right of the above equations is also negative for each value of n . By considering the maximum value of n we find that $m < 1/2$ for the four expansions; to include all values of n , we take $m \leq -1/2$ due to (124) and similar relations for each $\gamma_n^{(i)}$. On the other side, for series with several terms, $\gamma_n^{(i)} > 0$ implies $\gamma_1^{(i)} > 0$: this imposes the following restrictions on $m + l$

$$\begin{aligned}m &\leq -\frac{1}{2} \text{ for } \psi^{(i)}, \quad m + l + 2 < 0 \text{ for } \psi^{(1)}, \quad m + l + 3 < 0 \text{ for } \psi^{(2)} \text{ and } \psi^{(3)}, \\ l + m + 4 &< 0 \text{ for } \psi^{(4)},\end{aligned}\tag{123}$$

($i = 1, 2, 3, 4$) in addition to the restrictions on $m - l$. Actually, firstly we find conditions to get $\gamma_n^{(i)} \leq 0$ by considering n as a continuous variable. For example,

$$\gamma_n^{(1)} \leq 0 \text{ if } \begin{cases} \frac{m-l+1}{2} \leq n \leq -\frac{m+l}{2} & \text{when } m \leq -\frac{1}{2}, \\ -\frac{m+l}{2} \leq n \leq \frac{m-l+1}{2} & \text{when } m \geq -\frac{1}{2}. \end{cases}\tag{124}$$

Thence, $\gamma_n^{(i)} > 0$ when n lies outside the above intervals. In particular,

$$\gamma_n^{(1)} > 0 \text{ for } n < \frac{m-l+1}{2}, \text{ with } m \leq -\frac{1}{2},$$

an interval which includes the subinterval $n \leq (m - l - 1)/2$.

On the other side, for $\tilde{\psi}^{(i)}$, we demand that ($\gamma_n^{(2)} = \gamma_n^{(3)}$)

$$\begin{aligned}\gamma_n^{(1)} &= \left(n + \frac{l+m}{2}\right) \left(n + \frac{l-m-1}{2}\right) > 0 \text{ for } 1 \leq n \leq -\frac{m+l+2}{2}, \quad [m+l = -4, -6, -8, \dots], \\ \gamma_n^{(2)} &= \left(n + \frac{l+m+1}{2}\right) \left(n + \frac{l-m}{2}\right) > 0 \text{ for } 1 \leq n \leq -\frac{m+l+3}{2}, \quad [m+l = -5, -7, -9, \dots], \\ \gamma_n^{(4)} &= \left(n + \frac{l+m+2}{2}\right) \left(n + \frac{l-m+1}{2}\right) > 0 \text{ for } 1 \leq n \leq -\frac{m+l+4}{2}, \quad [m-l = -6, -8, \dots].\end{aligned}$$

As the first term is negative, then $\gamma_n^{(i)} > 0$ if the second term on the right of these equations is also negative for each value of n . From (124) and similar relations for each $\gamma_n^{(i)}$, we find that $m \geq -1/2$. Besides this, $\gamma_1^{(i)} > 0$ restricts the values of $m - l$, namely,

$$\begin{aligned}m &\geq -\frac{1}{2} \text{ for } \tilde{\psi}^{(i)}, \quad l - m + 1 < 0 \text{ for } \tilde{\psi}^{(1)}, \quad l - m + 2 < 0 \text{ for } \tilde{\psi}^{(2)} \text{ and } \tilde{\psi}^{(3)}, \\ l - m + 3 &< 0 \text{ for } \tilde{\psi}^{(4)},\end{aligned}\tag{125}$$

($i = 1, 2, 3, 4$) in addition to the restrictions on $m + l$. Using (124) we can check, for example, that

$$\gamma_n^{(1)} > 0 \text{ for } n < -\frac{l+m}{2}, \text{ when } m \geq -\frac{1}{2},$$

where the above interval includes all the possible values for n , that is, $n \leq -(l + m + 2)/2$.

To proceed we mention some examples, Obviously, the cases (115c), (117c), \dots , (122c) are solved by finite series of $\text{sn}^2 u$. Furthermore, we find that each of these cases admits a second expansion having the same period and opposite parity. For instance, if $0 < M_1 < N_1$ in (115c), we have

$$\begin{aligned} \text{for } \mathbf{m} = M_1 - N_1 - \frac{1}{2} \\ \text{and } \mathbf{l} = -M_1 - N_1 - \frac{3}{2} : \end{aligned} \left\{ \begin{array}{l} \psi^{(1)} \text{ with } 0 \leq n \leq M_1 < N_1, \text{ even, period } 2K; \\ \psi^{(4)} \text{ with } 0 \leq n \leq (M_1 - 1), \text{ odd, period } 2K; \end{array} \right. \quad (126)$$

and so far on. In each case there are only two distinct expansions satisfying the Arscott condition.

As a second example we take $\mathbf{m} = -\frac{1}{2}$ in which case $\psi^{(i)}(u) = \tilde{\psi}^{(i)}(u)$. The eight expansions satisfy the Arscott condition with the following values for \mathbf{l} :

$$\text{even solutions, period } 2K \ (\mathbf{m} = -1/2) : \left\{ \begin{array}{l} \psi^{(1)} \text{ if } \mathbf{l} = -\frac{3}{2}, -\frac{7}{2}, -\frac{11}{2}, \dots; \\ \psi^{(5)} \text{ if } \mathbf{l} = \frac{1}{2}, \frac{5}{2}, \frac{9}{2}, \dots; \end{array} \right. \quad (127)$$

$$\text{odd solutions, period } 2K \ (\mathbf{m} = -1/2) : \left\{ \begin{array}{l} \psi^{(4)} \text{ if } \mathbf{l} = -\frac{7}{2}, -\frac{11}{2}, -\frac{15}{2}, \dots; \\ \psi^{(8)} \text{ if } \mathbf{l} = \frac{5}{2}, \frac{9}{2}, \frac{13}{2}, \dots; \end{array} \right. \quad (128)$$

$$\text{odd solutions, period } 4K \ (\mathbf{m} = -1/2) : \left\{ \begin{array}{l} \psi^{(2)} \text{ if } \mathbf{l} = -\frac{5}{2}, -\frac{9}{2}, -\frac{13}{2}, \dots; \\ \psi^{(6)} \text{ if } \mathbf{l} = \frac{3}{2}, \frac{7}{2}, \frac{11}{2}, \dots; \end{array} \right. \quad (129)$$

$$\text{even solutions, period } 4K \ (\mathbf{m} = -1/2) : \left\{ \begin{array}{l} \psi^{(3)} \text{ if } \mathbf{l} = -\frac{5}{2}, -\frac{9}{2}, -\frac{13}{2}, \dots; \\ \psi^{(7)} \text{ if } \mathbf{l} = \frac{3}{2}, \frac{7}{2}, \frac{11}{2}, \dots; \end{array} \right. \quad (130)$$

In the above expressions, there are at the most two distinct expansions for a given value of \mathbf{l} (only one solution when $\mathbf{l} = -3/2$ or $\mathbf{l} = 1/2$). Notice that the Arscott condition is applicable for series with two or more terms; further, if the condition is not fulfilled, we cannot in advance state anything on the eigenvalues of the characteristic equation.

As a last illustration, we consider solutions for Eq. (107), that is, for $\mathbf{m} = 0$ or $\mathbf{m} = -1$. There are finite-series solutions in series of $\text{sn}^{2n} u$ for any integer value of \mathbf{l} (excepting $\mathbf{l} = -1$), but only the following eight expansions satisfy the Arscott condition

$$\text{even solutions, period } 2K \ (\mathbf{m} = 0) : \left\{ \begin{array}{ll} \tilde{\psi}^{(1)} & \text{if } \mathbf{l} = -2, -4, -6, \dots; \\ \psi^{(5)} & \text{if } \mathbf{l} = 1, 3, 5, \dots; \end{array} \right. \quad (131)$$

$$\text{odd solutions, period } 2K \ (\mathbf{m} = 0) : \left\{ \begin{array}{ll} \tilde{\psi}^{(4)} & \text{if } \mathbf{l} = -4, -6, -8, \dots; \\ \psi^{(8)} & \text{if } \mathbf{l} = 3, 5, 7, \dots; \end{array} \right. \quad (132)$$

$$\text{odd solutions, period } 4K \ (\mathbf{m} = 0) : \left\{ \begin{array}{ll} \tilde{\psi}^{(2)} & \text{if } \mathbf{l} = -3, -5, -7, \dots; \\ \psi^{(6)} & \text{if } \mathbf{l} = -2, -4, -6, \dots; \end{array} \right. \quad (133)$$

$$\text{even solutions, period } 4K \ (\mathbf{m} = 0) : \left\{ \begin{array}{ll} \tilde{\psi}^{(3)} & \text{if } \mathbf{l} = -3, -5, -7, \dots; \\ \psi^{(7)} & \text{if } \mathbf{l} = -2, -4, -6, \dots. \end{array} \right. \quad (134)$$

On the other hand, for $\mathbf{m} = -1$ the remaining expansions also lead to solutions satisfying the Arscott condition; nevertheless, such solutions are identical to the above ones. For example, we find that

$$\psi^{(1)}|_{\mathbf{m}=-1} = \tilde{\psi}^{(1)}|_{\mathbf{m}=0}, \quad \tilde{\psi}^{(5)}|_{\mathbf{m}=-1} = \psi^{(5)}|_{\mathbf{m}=0}, \quad \text{and so on.}$$

From relations (131-134), we find that there are four types of finite-series solutions if $l = -4, -6, -8, \dots$, unlike the expansions of the previous examples.

5.2 Finite-series eigenfunction $\left(\mathring{\Psi}^{(i)}, \tilde{\Psi}^{(i)}\right)$ in terms of $\text{cn}^2 u$

This is similar to Section 5.1. The finite-series expansions $\left(\mathring{\Psi}^{(i)}, \tilde{\Psi}^{(i)}\right)$ now result from the power-series solutions $\mathring{H}^{(i)}(x)$ for the Heun equation, truncated as in (110). The pairs $\left(\mathring{\Psi}^{(j+4)}, \tilde{\Psi}^{(j+4)}\right)$ ($j = 1, 2, 3, 4$) can be generated from $\left(\mathring{\psi}^{(j)}, \tilde{\Psi}^{(j)}\right)$ by the substitutions $l \mapsto -l - 1$ and $m \mapsto -m - 1$. The Arscott condition is satisfied if $\mathring{\gamma}_n^{(i)} = \mathring{\gamma}_n^{(i)} < 0$, in opposition to solutions of Section 5.1.

In the recurrence relations $\mathring{\alpha}_n \mathring{b}_{n+1} + \mathring{\beta}_n \mathring{b}_n + \mathring{\gamma}_n \mathring{b}_{n-1} = 0$ and $\tilde{\alpha}_n \tilde{b}_{n+1} + \tilde{\beta}_n \tilde{b}_n + \tilde{\gamma}_n \tilde{b}_{n-1} = 0$, formally we have

$$\tilde{\alpha}_n^{(i)} = \mathring{\alpha}_n^{(i)}, \quad \tilde{\beta}_n^{(i)} = \mathring{\beta}_n^{(i)}, \quad \tilde{\gamma}_n^{(i)} = \mathring{\gamma}_n^{(i)}. \quad (135)$$

After writing the four types of expansions, we discuss the Arscott condition (36) and comment on the Lamé equation.

5.2.1 First type: even solutions, period $2K$

$$\left\{ \begin{array}{ll} \mathring{\Psi}^{(1)}(u) = \text{dn}^{l+1} u \sum_{n=0}^{\frac{m-l-1}{2}} \mathring{b}_n^{(1)} \text{cn}^{2n} u & \text{if } m-l = 1, 3, 5, \dots, \\ & [m > -\frac{1}{2}, m+l > -2], \\ \tilde{\Psi}^{(1)}(u) = \text{dn}^{l+1} u \sum_{n=0}^{-\frac{m+l+2}{2}} \tilde{b}_n^{(1)} \text{cn}^{2n} u & \text{if } m+l = -2, -4, -6, \dots, \\ & [m < -\frac{1}{2}, m-l < 1], \end{array} \right. \quad (136a)$$

$$\begin{aligned} \mathring{\alpha}_n^{(1)} &= \left(1 - \frac{1}{k^2}\right) \left(n + \frac{1}{2}\right) (n+1), & \mathring{\beta}_n^{(1)} &= \left(\frac{1}{k^2} - 2\right) n^2 - (l+1)n - \frac{\varepsilon}{4k^2} +, \\ \frac{m(m+1)-l-1}{4}, & \mathring{\gamma}_n^{(1)} &= \left(n - \frac{m-l+1}{2}\right) \left(n + \frac{l+m}{2}\right); \end{aligned} \quad (136b)$$

As in the expansions in series of $\text{sn}^2 u$, there is only one expansion when (M_1 and N_1 denote positive integers)

$$\begin{aligned} m-l-1 = 2M_1 \quad \text{and} \quad m+l+2 = -2N_1 \quad (\Leftrightarrow m = M_1 - N_1 - \frac{1}{2}, : \\ l = -M_1 - N_1 - \frac{3}{2}) : \mathring{\psi}^{(1)} \text{ if } M_1 < N_1, \quad \tilde{\psi}^{(1)} \text{ if } N_1 < M_1. \end{aligned}$$

These expansions do not satisfy the Arscott condition. Actually, the Arscott condition is not satisfied by series of $\text{cn}^2 u$ for any of the cases given in (115c), (116c), \dots , (122c). Thus, these cases will not be considered from now on.

$$\left\{ \begin{array}{ll} \mathring{\Psi}^{(5)}(u) = \frac{1}{\text{dn}^l u} \sum_{n=0}^{\frac{l-m-1}{2}} \mathring{b}_n^{(5)} \text{cn}^{2n} u & \text{if } l-m = 1, 3, 5, \dots; \\ & [m < -\frac{1}{2}, m+l < 0], \\ \tilde{\Psi}^{(5)}(u) = \frac{1}{\text{dn}^l u} \sum_{n=0}^{\frac{l+m}{2}} \tilde{b}_n^{(5)} \text{cn}^{2n} u & \text{if } l+m = 0, 2, 4, \dots; \\ & [m > -\frac{1}{2}, l-m < 1], \end{array} \right. \quad (137a)$$

$$\begin{aligned}\mathring{\alpha}_n^{(5)} &= \left(1 - \frac{1}{k^2}\right) \left(n + \frac{1}{2}\right) (n+1), & \mathring{\beta}_n^{(5)} &= \left(\frac{1}{k^2} - 2\right) n^2 + \mathbf{l}n - \frac{\varepsilon}{4k^2} + \\ & \frac{\mathbf{l} + \mathbf{m}(\mathbf{m}+1)}{4}, & \mathring{\gamma}_n^{(5)} &= \left(n - \frac{\mathbf{l} - \mathbf{m} + 1}{2}\right) \left(n - \frac{\mathbf{l} + \mathbf{m} + 2}{2}\right).\end{aligned}\quad (137b)$$

5.2.2 Second type: odd solutions, period $4K$

$$\left\{ \begin{aligned} \mathring{\Psi}^{(2)}(u) &= \operatorname{sn} u \operatorname{dn}^{l+1} u \sum_{n=0}^{\frac{\mathbf{m}-\mathbf{l}-2}{2}} \mathring{\mathbf{b}}_n^{(2)} \operatorname{cn}^{2n} u & \text{if } \mathbf{m} - \mathbf{l} = 2, 4, 6, \dots, \\ & & [\mathbf{m} > -\frac{1}{2}, \mathbf{m} + \mathbf{l} > -3], \\ \tilde{\Psi}^{(2)}(u) &= \operatorname{sn} u \operatorname{dn}^{l+1} u \sum_{n=0}^{-\frac{\mathbf{m}+\mathbf{l}+3}{2}} \tilde{\mathbf{b}}_n^{(2)} \operatorname{cn}^{2n} u & \text{if } \mathbf{m} + \mathbf{l} = -3, -5, -7, \dots, \\ & & [\mathbf{m} < -\frac{1}{2}, \mathbf{m} - \mathbf{l} < 2], \end{aligned} \right. \quad (138a)$$

$$\begin{aligned}\mathring{\alpha}_n^{(2)} &= \left(1 - \frac{1}{k^2}\right) \left(n + \frac{1}{2}\right) (n+1), & \mathring{\beta}_n^{(2)} &= \left(\frac{1}{k^2} - 2\right) n^2 + \left(\frac{1}{k^2} - \mathbf{l} - 2\right) n + \\ & \frac{1-\varepsilon}{4k^2} + \frac{\mathbf{m}(\mathbf{m}+1) - \mathbf{l} - 2}{4}, & \mathring{\gamma}_n^{(2)} &= \left(n - \frac{\mathbf{m} - \mathbf{l}}{2}\right) \left(n + \frac{\mathbf{l} + \mathbf{m} + 1}{2}\right);\end{aligned}\quad (138b)$$

$$\left\{ \begin{aligned} \mathring{\Psi}^{(6)}(u) &= \frac{\operatorname{sn} u}{\operatorname{dn}^l u} \sum_{n=0}^{\frac{\mathbf{l}-\mathbf{m}-2}{2}} \mathring{\mathbf{b}}_n^{(6)} \operatorname{cn}^{2n} u & \text{if } \mathbf{l} - \mathbf{m} = 2, 4, 6, \dots, \\ & & [\mathbf{m} < -\frac{1}{2}, \mathbf{m} + \mathbf{l} < 1], \\ \tilde{\Psi}^{(6)}(u) &= \frac{\operatorname{sn} u}{\operatorname{dn}^l u} \sum_{n=0}^{\frac{\mathbf{l}+\mathbf{m}-1}{2}} \tilde{\mathbf{b}}_n^{(6)} \operatorname{cn}^{2n} u & \text{if } \mathbf{l} + \mathbf{m} = 1, 3, 5, \dots, \\ & & [\mathbf{m} > -\frac{1}{2}, \mathbf{l} - \mathbf{m} < 2], \end{aligned} \right. \quad (139a)$$

$$\begin{aligned}\mathring{\alpha}_n^{(6)} &= \left(1 - \frac{1}{k^2}\right) \left(n + \frac{1}{2}\right) (n+1), & \mathring{\beta}_n^{(6)} &= \left(\frac{1}{k^2} - 2\right) n^2 + \left(\frac{1}{k^2} + \mathbf{l} - 1\right) n + \\ & \frac{1-\varepsilon}{4k^2} + \frac{\mathbf{m}(\mathbf{m}+1) + \mathbf{l} - 1}{4}, & \mathring{\gamma}_n^{(6)} &= \left(n - \frac{\mathbf{l} - \mathbf{m}}{2}\right) \left(n - \frac{\mathbf{l} + \mathbf{m} + 1}{2}\right).\end{aligned}\quad (139b)$$

5.2.3 Third type: even solutions, period $4K$

$$\left\{ \begin{aligned} \mathring{\Psi}^{(3)}(u) &= \operatorname{cn} u \operatorname{dn}^{l+1} u \sum_{n=0}^{\frac{\mathbf{m}-\mathbf{l}-2}{2}} \mathring{\mathbf{b}}_n^{(3)} \operatorname{cn}^{2n} u & \text{if } \mathbf{m} - \mathbf{l} = 2, 4, 6, \dots, \\ & & [\mathbf{m} > -\frac{1}{2}, \mathbf{m} + \mathbf{l} > -3], \\ \tilde{\Psi}^{(3)}(u) &= \operatorname{cn} u \operatorname{dn}^{l+1} u \sum_{n=0}^{-\frac{\mathbf{m}+\mathbf{l}+3}{2}} \tilde{\mathbf{b}}_n^{(3)} \operatorname{cn}^{2n} u & \text{if } \mathbf{m} + \mathbf{l} = -3, -5, -7, \dots, \\ & & [\mathbf{m} < -\frac{1}{2}, \mathbf{m} - \mathbf{l} < 2], \end{aligned} \right. \quad (140a)$$

$$\begin{aligned}\mathring{\alpha}_n^{(3)} &= \left(1 - \frac{1}{k^2}\right) \left(n + \frac{3}{2}\right) (n+1), & \mathring{\beta}_n^{(3)} &= \left(\frac{1}{k^2} - 2\right) n^2 + \left(\frac{1}{k^2} - \mathbf{l} - 3\right) n + \\ & \frac{1-\varepsilon}{4k^2} - \frac{1}{4}(3\mathbf{l} + 5) + \frac{1}{4}\mathbf{m}(\mathbf{m} + 1), & \mathring{\gamma}_n^{(3)} &= \left(n - \frac{\mathbf{m} - \mathbf{l}}{2}\right) \left(n + \frac{\mathbf{l} + \mathbf{m} + 1}{2}\right);\end{aligned}\quad (140b)$$

$$\left\{ \begin{array}{ll} \mathring{\Psi}^{(7)}(u) = \frac{\text{cn}u}{\text{dn}^l u} \sum_{n=0}^{\frac{l-m-2}{2}} \mathring{b}_n^{(7)} \text{cn}u^{2n} u & \text{if } \quad l - m = 2, 4, 6, \dots, \\ & [m < -\frac{1}{2}, m + l < 1], \\ \tilde{\Psi}^{(7)}(u) = \frac{\text{cn}u}{\text{dn}^l u} \sum_{n=0}^{\frac{l+m-1}{2}} \tilde{b}_n^{(7)} \text{cn}u^{2n} u & \text{if } \quad l + m = 1, 3, 5, \dots, \\ & [m > -\frac{1}{2}, l - m < 2], \end{array} \right. \quad (141a)$$

$$\begin{aligned}\mathring{\alpha}_n^{(7)} &= \left(1 - \frac{1}{k^2}\right) \left(n + \frac{3}{2}\right) (n+1), & \mathring{\beta}_n^{(7)} &= \left(\frac{1}{k^2} - 2\right) n^2 + \left(\frac{1}{k^2} + l - 2\right) n + \\ \frac{1-\mathcal{E}}{4k^2} + \frac{1}{4}(3l-2) + \frac{1}{4}\mathbf{m}(\mathbf{m}+1), & \mathring{\gamma}_n^{(7)} &= \left(n - \frac{l-m}{2}\right) \left(n - \frac{l+m+1}{2}\right).\end{aligned}\quad (141b)$$

5.2.4 Fourth type: odd solutions, period $2K$

$$\left\{ \begin{array}{l} \dot{\Psi}^{(4)}(u) = \operatorname{sn} u \operatorname{cn} u \operatorname{dn}^{l+1} u \sum_{n=0}^{\frac{m-l-3}{2}} \dot{\boldsymbol{b}}_n^{(4)} \operatorname{cn}^{2n} u \quad \text{if } m-l=3, 5, 7, \dots, \\ \qquad \qquad \qquad [m > -\tfrac{1}{2}, m+l > -4], \\ \tilde{\Psi}^{(4)}(u) = \operatorname{sn} u \operatorname{cn} u \operatorname{dn}^{l+1} u \sum_{n=0}^{-\frac{m+l+4}{2}} \tilde{\boldsymbol{b}}_n^{(4)} \operatorname{cn}^{2n} u \quad \text{if } m+l=-4, -6, -8, \dots, \\ \qquad \qquad \qquad [m < -\tfrac{1}{2}, m-l < 3], \end{array} \right. \quad (142a)$$

$$\begin{aligned} \mathring{\alpha}_n^{(4)} &= \left(1 - \frac{1}{k^2}\right) \left(n + \frac{3}{2}\right) (n+1), \quad \mathring{\beta}_n^{(4)} = \left(\frac{1}{k^2} - 2\right) n^2 + \left(\frac{1}{k^2} - l - 4\right) n + \\ &\frac{4-\mathcal{E}}{4k^2} - \frac{1}{4}(3l+8) + \frac{1}{4}\mathbf{m}(\mathbf{m}+1), \quad \mathring{\gamma}_n^{(4)} = \left(n - \frac{m-l-1}{2}\right) \left(n + \frac{l+m+2}{2}\right); \end{aligned} \quad (142b)$$

$$\left\{ \begin{array}{ll} \mathring{\Psi}^{(8)}(u) = \frac{\operatorname{sn} u \operatorname{cn} u}{\operatorname{dn}^l u} \sum_{n=0}^{\frac{l-m-3}{2}} \mathring{\mathbf{b}}_n^{(8)} \operatorname{cn}^{2n} u & \text{if } \mathbf{l} - \mathbf{m} = 3, 5, 7, \dots, \\ & [\mathbf{m} < -\frac{1}{2}, \mathbf{m} + \mathbf{l} < 2], \\ \tilde{\Psi}^{(8)}(u) = \frac{\operatorname{sn} u \operatorname{cn} u}{\operatorname{dn}^l u} \sum_{n=0}^{\frac{l+m-2}{2}} \tilde{\mathbf{b}}_n^{(8)} \operatorname{cn}^{2n} u & \text{if } \mathbf{l} + \mathbf{m} = 2, 4, 6, \dots, \\ & [\mathbf{m} > -\frac{1}{2}, \mathbf{l} - \mathbf{m} < 3], \end{array} \right. \quad (143a)$$

$$\begin{aligned}\mathring{\alpha}_n^{(8)} &= \left(1 - \frac{1}{k^2}\right) \left(n + \frac{3}{2}\right) (n+1), & \mathring{\beta}_n^{(8)} &= \left(\frac{1}{k^2} - 2\right) n^2 + \left(\frac{2}{k^2} + l - 3\right) n + \\ & \frac{4-\varepsilon}{4k^2} + \frac{3l-5}{4} + \frac{m}{4}(m+1), & \mathring{\gamma}_n^{(8)} &= \left(n - \frac{l-m-1}{2}\right) \left(n - \frac{l+m}{2}\right).\end{aligned}\quad (143b)$$

Concerning the Arscott condition (36), we notice that $(1 - 1/k^2) < 1$ implies that the coefficients $\mathring{\alpha}_n^{(i)}$ are negative. Then, the condition $\mathring{\alpha}_{n-1}\mathring{\gamma}_n > 0$ is satisfied if $\mathring{\gamma}_n < 0$ for any admissible value of n , that is,

$$\begin{aligned}\mathring{\gamma}_n^{(1)} &= \left(n - \frac{\mathbf{m}-\mathbf{l}+1}{2}\right) \left(n + \frac{\mathbf{l}+\mathbf{m}}{2}\right) < 0 \text{ for } 1 \leq n \leq \frac{\mathbf{m}-\mathbf{l}-1}{2}, [\mathbf{m} - \mathbf{l} = 3, 5, 7, \dots], \\ \mathring{\gamma}_n^{(2)} &= \mathring{\gamma}_n^{(3)} = \left(n - \frac{\mathbf{m}-\mathbf{l}}{2}\right) \left(n + \frac{\mathbf{l}+\mathbf{m}+1}{2}\right) < 0 \text{ for } 1 \leq n \leq \frac{\mathbf{m}-\mathbf{l}-2}{2}, [\mathbf{m} - \mathbf{l} = 4, 6, 8, \dots], \\ \mathring{\gamma}_n^{(4)} &= \left(n - \frac{\mathbf{m}-\mathbf{l}-1}{2}\right) \left(n + \frac{\mathbf{l}+\mathbf{m}+2}{2}\right) < 0 \text{ for } 1 \leq n \leq \frac{\mathbf{m}-\mathbf{l}-3}{2}, [\mathbf{m} - \mathbf{l} = 5, 7, 9, \dots],\end{aligned}$$

for $\mathring{\Psi}^{(i)}$. As the first term of these $\mathring{\gamma}_n^{(i)}$ is negative, $\mathring{\gamma}_n^{(i)} < 0$ if the last term is positive for each value of n . From (124) and similar relations for each $\mathring{\gamma}_n^{(i)}$, it follows that $\mathbf{m} > -1/2$. In addition, from $\mathring{\gamma}_{n=1}^{(i)} < 0$ we get restrictions on $\mathbf{m} + \mathbf{l}$, namely,

$$\begin{aligned}\mathbf{m} &> -\frac{1}{2} \text{ for } \mathring{\Psi}^{(i)}, \quad \mathbf{m} + \mathbf{l} + 2 > 0 \text{ for } \mathring{\Psi}^{(1)}, \quad \mathbf{m} + \mathbf{l} + 3 > 0 \text{ for } \mathring{\Psi}^{(2)} \text{ and } \mathring{\Psi}^{(3)}, \\ \mathbf{m} + \mathbf{l} + 4 &> 0 \text{ for } \mathring{\Psi}^{(4)}.\end{aligned}\tag{144}$$

We see, for example, that the interval

$$-\frac{\mathbf{m}+\mathbf{l}}{2} < n < \frac{\mathbf{m}-\mathbf{l}-1}{2} + 1, \quad \text{for } \mathbf{m} > -\frac{1}{2} \quad \text{and} \quad -\mathbf{m} - \mathbf{l} < 2$$

assures that $\mathring{\gamma}_n^{(1)} < 0$ for all the possible values for n .

On the other hand, for $\mathring{\Psi}^{(i)}$ we must require that $(\mathring{\gamma}_n^{(2)} = \mathring{\gamma}_n^{(3)})$

$$\begin{aligned}\mathring{\gamma}_n^{(1)} &= \left(n + \frac{\mathbf{l}+\mathbf{m}}{2}\right) \left(n + \frac{\mathbf{l}-\mathbf{m}-1}{2}\right) < 0 \text{ for } 1 \leq n \leq -\frac{\mathbf{m}+\mathbf{l}+2}{2}, [\mathbf{m} + \mathbf{l} = -4, -6, -8, \dots], \\ \mathring{\gamma}_n^{(2)} &= \left(n + \frac{\mathbf{l}+\mathbf{m}+1}{2}\right) \left(n + \frac{\mathbf{l}-\mathbf{m}}{2}\right) < 0 \text{ for } 1 \leq n \leq -\frac{\mathbf{m}+\mathbf{l}+3}{2}, [\mathbf{m} + \mathbf{l} = -5, -7, -9, \dots], \\ \mathring{\gamma}_n^{(4)} &= \left(n + \frac{\mathbf{l}+\mathbf{m}+2}{2}\right) \left(n + \frac{\mathbf{l}-\mathbf{m}+1}{2}\right) < 0 \text{ for } 1 \leq n \leq -\frac{\mathbf{m}+\mathbf{l}+4}{2}, [\mathbf{m} + \mathbf{l} = -6, -8, \dots].\end{aligned}$$

Since the first term is negative, $\mathring{\gamma}_n^{(i)} < 0$ if the last term on the right of these equations is positive. Thence, we find that the Arscott condition is assured by the following restrictions on the parameters

$$\begin{aligned}\mathbf{m} &< -\frac{1}{2} \text{ for } \mathring{\Psi}^{(i)}, \quad \mathbf{l} - \mathbf{m} + 1 > 0 \text{ for } \mathring{\Psi}^{(1)}, \quad \mathbf{l} - \mathbf{m} + 2 > 0 \text{ for } \mathring{\Psi}^{(2)} \text{ and } \mathring{\Psi}^{(3)}, \\ \mathbf{l} - \mathbf{m} + 3 &> 0 \text{ for } \mathring{\Psi}^{(4)};\end{aligned}\tag{145}$$

These conditions imply that $\mathring{\gamma}_n^{(i)} < 0$ for any admissible n . For example, by (124) and (145), we see that $\mathring{\gamma}_n^{(1)} < 0$ in the interval

$$\frac{\mathbf{m}-\mathbf{l}+1}{2} < n < -\frac{\mathbf{m}+\mathbf{l}}{2} \quad \text{for } \mathbf{l} - \mathbf{m} + 1 > 0 \quad \text{and} \quad \mathbf{m} < -\frac{1}{2}$$

which includes all the possible values of n for the solutions $\mathring{\Psi}^{(1)}$.

The eigenfunctions $(\mathring{\Psi}^{(i)}, \mathring{\Psi}^{(i)})$ do not satisfy the Arscott condition for the examples discussed in Section 5.1. However, we can check that the sixteen eigenfunctions give different types of solutions for Lamé equation (108) when $\mathbf{m} = 0, \pm 1, \pm 2, \dots$; each solution satisfies the Arscott condition. Except if $\mathbf{m} = -2$, $\mathbf{m} = -1$, $\mathbf{m} = 0$ and $\mathbf{m} = 1$, for each

value of \mathbf{m} there are four types of solutions.

$$\text{even solutions, period } 2K \ (l = 0) : \begin{cases} \mathring{\Psi}^{(1)} & \text{if } \mathbf{m} = 1, 3, 5, \dots; \\ \tilde{\Psi}^{(1)} & \text{if } \mathbf{m} = -2, -4, -6, \dots; \\ \mathring{\Psi}^{(5)} & \text{if } \mathbf{m} = -1, -3, -5, \dots; \\ \tilde{\Psi}^{(5)} & \text{if } \mathbf{m} = 0, 2, 4, \dots; \end{cases} \quad (146)$$

$$\text{odd solutions, period } 2K \ (l = 0) : \begin{cases} \mathring{\Psi}^{(4)} & \text{if } \mathbf{m} = 3, 5, 7, \dots; \\ \tilde{\Psi}^{(4)} & \text{if } \mathbf{m} = -4, -6, -8, \dots; \\ \mathring{\Psi}^{(8)} & \text{if } \mathbf{m} = -3, -5, -7, \dots; \\ \tilde{\Psi}^{(8)} & \text{if } \mathbf{m} = 2, 4, 6, \dots; \end{cases} \quad (147)$$

$$\text{odd solutions, period } 4K \ (l = 0) : \begin{cases} \mathring{\Psi}^{(2)} & \text{if } \mathbf{m} = 2, 4, 6, \dots; \\ \tilde{\Psi}^{(2)} & \text{if } \mathbf{m} = -3, -5, -7, \dots; \\ \mathring{\Psi}^{(6)} & \text{if } \mathbf{m} = -2, -4, -6, \dots; \\ \tilde{\Psi}^{(6)} & \text{if } \mathbf{m} = 1, 3, 5, \dots; \end{cases} \quad (148)$$

$$\text{even solutions, period } 4K \ (l = 0) : \begin{cases} \tilde{\Psi}^{(3)} & \text{if } \mathbf{m} = 2, 4, 6, \dots; \\ \mathring{\Psi}^{(3)} & \text{if } \mathbf{m} = -3, -5, -7, \dots; \\ \mathring{\Psi}^{(7)} & \text{if } \mathbf{m} = -2, -4, -6, \dots; \\ \tilde{\Psi}^{(7)} & \text{if } \mathbf{m} = 1, 3, 5, \dots. \end{cases} \quad (149)$$

We find the same results if we take $l = 0$ or $l = -1$. By taking $l = -1$ we simply rearrange the previous expansions. We obtain, for instance,

$$\mathring{\Psi}^{(1)} \Big|_{l=-1} = \tilde{\Psi}^{(5)} \Big|_{l=0}, \quad \tilde{\Psi}^{(1)} \Big|_{l=-1} = \mathring{\Psi}^{(5)} \Big|_{l=0}, \quad \mathring{\Psi}^{(5)} \Big|_{l=-1} = \tilde{\Psi}^{(1)} \Big|_{l=0}, \quad \tilde{\Psi}^{(5)} \Big|_{l=-1} = \mathring{\Psi}^{(1)} \Big|_{l=0}.$$

Other examples, for fixed values of \mathbf{m} and l , are presented in Appendix C.

5.3 Finite series of hypergeometric functions $\psi^{(i)}(u)$ when l is an integer and $\mathbf{m} \neq -1/2$ is half an odd integer

If \mathbf{m} is half an odd positive integer, finite-series solutions $\psi^{(i)}(u)$ with $0 \leq n \leq \mathbf{m} - 1/2$ are obtained by inserting in (103) the hypergeometric-function expansions $H^{(i)}(x)$ with $i = 1, 4, 6, 7$. In these cases, the series are finite because the factor $(n + \alpha - \beta) = (n - \mathbf{m} - 1/2)$ implies $\gamma_n^{(i)} = 0$ for $n = \mathbf{m} + 1/2$ as demanded in (34). Since $\delta = 1/2$, according to (89) the four hypergeometric functions converge when $z = 1$ or $\text{sn}^2 u = 1$, where $z = (a - 1)x/(a - x) = (1 - k^2) \text{sd}^2 u$.

If $\mathbf{m} \neq -1/2$ is half an odd negative integer, finite-series solutions $\psi^{(i)}$ with $0 \leq n \leq -\mathbf{m} - 3/2$ result from $H^{(i)}$ for $i = 2, 3, 5, 8$. Now the series terminate at $n = -\mathbf{m} - 3/2$ because $(n - \alpha + \beta) = (n + \mathbf{m} + 1/2)$ implies $\gamma_n^{(i)} = 0$ for $n = -\mathbf{m} - 1/2$ as demanded in (34). Since $\delta = 1/2$, by (89) the four solutions converge when $z = 1$ ($\text{sn}^2 u = 1$).

Eigenfunctions for negative and positive values of \mathbf{m} are connected to each other by means of the substitutions (104). In both cases (positive and negative \mathbf{m}) the solutions are degenerate in the sense that there are two independent solutions with the same energies \mathcal{E} for a given set of values for the parameters (\mathbf{m}, l, k^2) . In addition, for a given expansion the

the Arscott condition (36) is satisfied only if we impose restrictions on the values of \mathbf{l} and \mathbf{m} . However, by considering all the expansions, we find at least one expansion satisfying the condition for any allowed value of (\mathbf{m}, \mathbf{l}) .

Here and in Sec. 5.4, in order to check the parity and periodicity properties we consider the one-term expansions because in these cases the hypergeometric functions can be written in terms of elementary functions rather than in terms of infinite series. For this we use [3]

$$\begin{aligned} F(a, -a; \tfrac{1}{2}; -z^2) &= \tfrac{1}{2} \left[(\sqrt{1+z^2} + z)^{2a} + (\sqrt{1+z^2} - z)^{2a} \right], \\ F(a, 1-a; \tfrac{1}{2}; -z^2) &= \frac{1}{2\sqrt{1+z^2}} \left[(\sqrt{1+z^2} + z)^{2a-1} + (\sqrt{1+z^2} - z)^{2a-1} \right], \\ F(a, 1-a; \tfrac{3}{2}; -z^2) &= \frac{1}{(2-4a)z} \left[(\sqrt{1+z^2} + z)^{1-2a} - (\sqrt{1+z^2} - z)^{1-2a} \right]. \end{aligned} \quad (150)$$

These relations can also be applied to some two-term series. Further, we find that such parity and periodicity properties hold for any finite-series expansion.

5.3.1 Finite series when \mathbf{l} is an integer and \mathbf{m} is half an odd positive integer

First we write the eigenfunctions $\psi^{(1)}(u)$ and $\psi^{(6)}(u)$. Then, we proof the degeneracy by using the matrix form for the recurrence relations. After this, we discuss the Arscott condition and, finally, we consider the eigenfunctions $\psi^{(4)}(u)$ and $\psi^{(7)}(u)$.

$$\left\{ \begin{aligned} &\psi^{(1)}(u) \stackrel{(81,103)}{=} (\text{dn } u)^{\mathbf{m}} \sum_{n=0}^{\mathbf{m}-1/2} b_n^{(1)} [(1-k^2) \text{sd}^2 u]^n \times \\ &\tilde{F} \left[n + \frac{\mathbf{l}-\mathbf{m}+1}{2}, \frac{\mathbf{m}-\mathbf{l}-1}{2}; n + \tfrac{1}{2}; (1-k^2) \text{sd}^2 u \right], \quad \mathbf{m} = \tfrac{1}{2}, \tfrac{3}{2}, \tfrac{5}{2}, \dots, \quad [\text{even}, 4\mathbf{K}], \\ &\alpha_n^{(1)} = \left(1 - \frac{1}{k^2}\right) (n+1), \quad \beta_n^{(1)} = \left(\frac{1}{k^2} - 2\right) n^2 + \left[\left(\frac{1}{k^2} - 1\right) (\mathbf{l} - \mathbf{m} + 1) + \mathbf{m}\right] n + \\ &\frac{(\mathbf{l}+1-\mathbf{m})^2 - \mathcal{E}}{4k^2} + \frac{2\mathbf{l}\mathbf{m} - \mathbf{l} + \mathbf{m} - (\mathbf{m}-1)^2}{4}, \quad \gamma_n^{(1)} = \left(n + \frac{\mathbf{l}-\mathbf{m}-1}{2}\right) \left(n + \frac{\mathbf{l}-\mathbf{m}}{2}\right) \left(n - \mathbf{m} - \tfrac{1}{2}\right); \end{aligned} \right. \quad (151)$$

$$\left\{ \begin{aligned} &\psi^{(6)}(u) \stackrel{(86,103)}{=} \text{snu}(\text{dn } u)^{\mathbf{m}-1} \sum_{n=0}^{\mathbf{m}-1/2} b_n^{(6)} [(1-k^2) \text{sd}^2 u]^n \times \\ &\tilde{F} \left[n - \frac{\mathbf{l}+\mathbf{m}-1}{2}, \frac{\mathbf{m}+\mathbf{l}+1}{2}; n + \tfrac{3}{2}; (1-k^2) \text{sd}^2 u \right], \quad \mathbf{m} = \tfrac{1}{2}, \tfrac{3}{2}, \tfrac{5}{2}, \dots, \quad [\text{odd}, 4\mathbf{K}], \\ &\alpha_n^{(6)} = \left(1 - \frac{1}{k^2}\right) (n+1), \quad \beta_n^{(6)} = \left(\frac{1}{k^2} - 2\right) n^2 + \left[\left(1 - \frac{1}{k^2}\right) (\mathbf{l} + \mathbf{m}) + \mathbf{m} - 1\right] n + \\ &\frac{(\mathbf{l}+\mathbf{m})^2 - \mathcal{E}}{4k^2} + \frac{\mathbf{l} - 2\mathbf{l}\mathbf{m} + \mathbf{m} - (\mathbf{m}-1)^2}{4}, \quad \gamma_n^{(6)} = \left(n - \frac{\mathbf{l}+\mathbf{m}+1}{2}\right) \left(n - \frac{\mathbf{l}+\mathbf{m}}{2}\right) \left(n - \mathbf{m} - \tfrac{1}{2}\right). \end{aligned} \right. \quad (152)$$

For $\mathbf{m} = 1/2$ we obtain two independent eigenfunctions,

$$\begin{aligned} \psi^{(1)}(u) \Big|_{\mathbf{m}=1/2} &= b_0^{(1)} \sqrt{\text{dn } u} \tilde{F} \left[\tfrac{1}{2} \left(\mathbf{l} + \tfrac{1}{2}\right), -\tfrac{1}{2} \left(\mathbf{l} + \tfrac{1}{2}\right); \tfrac{1}{2}; (1-k^2) \text{sd}^2 u \right], \\ \psi^{(6)}(u) \Big|_{\mathbf{m}=1/2} &= b_0^{(6)} \frac{\text{snu}}{\sqrt{\text{dn } u}} \tilde{F} \left[-\tfrac{1}{2} \left(\mathbf{l} - \tfrac{1}{2}\right), \tfrac{1}{2} \left(\mathbf{l} + \tfrac{3}{2}\right); \tfrac{3}{2}; (1-k^2) \text{sd}^2 u \right], \end{aligned}$$

which are degenerate because both correspond to the same energy \mathcal{E} , that is,

$$\beta_0^{(1)} = 0, \quad \beta_0^{(6)} = 0 \quad \Rightarrow \quad \mathcal{E} = \left(\mathbf{l} + \tfrac{1}{2}\right)^2 + \frac{k^2}{4}, \quad [\mathbf{m} = \tfrac{1}{2}].$$

To prove the degeneracy for $\mathbf{m} \geq 3/2$, first we redefine $b_n^{(6)}$ as

$$b_n^{(6)} = \left(\frac{k^2}{1-k^2} \right)^n \Gamma \left(n - \frac{l+m-1}{2} \right) \Gamma \left(n - \frac{l+m-2}{2} \right) \hat{b}_n, \quad (153a)$$

which implies that the \hat{b}_n obey the relations $\hat{\alpha}_n \hat{b}_{n+1} + \hat{\beta}_n \hat{b}_n + \hat{\gamma}_n \hat{b}_{n-1} = 0$ ($\hat{b}_{-1} = 0$) where

$$\begin{aligned} \hat{\alpha}_n &= - \left(n - \frac{l+m-1}{2} \right) \left(n - \frac{l+m-2}{2} \right) (n+1), & \hat{\beta}_n &= \beta_n^{(6)}, \\ \hat{\gamma}_n &= - \left(1 - \frac{1}{k^2} \right) \left(n - \mathbf{m} - \frac{1}{2} \right). \end{aligned} \quad (153b)$$

We show the degeneracy by writing the relations for $b_n^{(1)}$ and \hat{b}_n in the forms $\mathbb{A} \vec{b} = 0$ and $\mathbb{B} \vec{\hat{b}} = 0$, where

$$\vec{b} = \left(b_0^{(1)}, b_1^{(1)}, \dots, b_{\mathbf{m}-\frac{1}{2}}^{(1)} \right)^t, \quad \vec{\hat{b}} = \left(\hat{b}_0, \hat{b}_1, \dots, \hat{b}_{\mathbf{m}-\frac{1}{2}} \right)^t,$$

while \mathbb{A} and \mathbb{B} denote the matrices

$$\mathbb{A} = \begin{bmatrix} \beta_0^{(1)} & \alpha_0^{(1)} & 0 & & \\ \gamma_1^{(1)} & \beta_1^{(1)} & \alpha_1^{(1)} & & \\ & \ddots & \ddots & \ddots & \\ & & \gamma_{\mathbf{m}-\frac{3}{2}}^{(1)} & \beta_{\mathbf{m}-\frac{3}{2}}^{(1)} & \alpha_{\mathbf{m}-\frac{3}{2}}^{(1)} \\ & & 0 & \gamma_{\mathbf{m}-\frac{1}{2}}^{(1)} & \beta_{\mathbf{m}-\frac{1}{2}}^{(1)} \end{bmatrix}, \quad \mathbb{B} = \begin{bmatrix} \hat{\beta}_0 & \hat{\alpha}_0 & 0 & & \\ \hat{\gamma}_1 & \hat{\beta}_1 & \hat{\alpha}_1 & & \\ & \ddots & \ddots & \ddots & \\ & & \hat{\gamma}_{\mathbf{m}-\frac{3}{2}} & \hat{\beta}_{\mathbf{m}-\frac{3}{2}} & \hat{\alpha}_{\mathbf{m}-\frac{3}{2}} \\ & & 0 & \hat{\gamma}_{\mathbf{m}-\frac{1}{2}} & \hat{\beta}_{\mathbf{m}-\frac{1}{2}} \end{bmatrix}.$$

Then (151), (152) and (153b) imply that both matrices have the same elements since

$$\mathbb{B} = \begin{bmatrix} \beta_{\mathbf{m}-\frac{1}{2}}^{(1)} & \gamma_{\mathbf{m}-\frac{1}{2}}^{(1)} & 0 & & \\ \alpha_{\mathbf{m}-\frac{3}{2}}^{(1)} & \beta_{\mathbf{m}-\frac{3}{2}}^{(1)} & \gamma_{\mathbf{m}-\frac{3}{2}}^{(1)} & & \\ & \ddots & \ddots & \ddots & \\ & & \alpha_1^{(1)} & \beta_1^{(1)} & \gamma_1^{(1)} \\ & & 0 & \alpha_0^{(1)} & \beta_0^{(1)} \end{bmatrix}. \quad (154)$$

The degeneracy results from the fact that $\det \mathbb{A} = \det \mathbb{B}$. In effect, let \mathbb{U} be a square matrix of order $(\mathbf{m} + \frac{1}{2})$, having 1's on the antidiagonal as the only nonzero elements, that is,

$$\mathbb{U} = \mathbb{U}^{-1} = \begin{bmatrix} & & 1 \\ & . & \\ 1 & & \end{bmatrix}, \quad \det \mathbb{U} = \pm 1, \quad (155)$$

where \mathbb{U}^{-1} is the matrix inverse to \mathbb{U} . Then, we find that

$$\mathbb{B} = \mathbb{U} \mathbb{A} \mathbb{U} \quad \Rightarrow \quad \det \mathbb{A} = \det \mathbb{B}, \quad (156)$$

where the last equality follows from the properties of the determinants. The above approach has already been used to get the degeneracy of wavefunctions for a problem which is ruled by a reduced confluent Heun equation [39].

Finally, in (151) and (152) $\alpha_n^{(1)} = \alpha_n^{(6)} < 0$. Thence, the Arscott condition (36) for $\mathbf{m} \geq 3/2$ requires that $\gamma_n^{(1)} < 0$ and $\gamma_n^{(6)} < 0$. From the expressions for $\gamma_n^{(1)}$ and $\gamma_n^{(6)}$ these conditions are equivalent to

$$\begin{aligned} f(n) &= \left(n + \frac{l-m-1}{2}\right) \left(n + \frac{l-m}{2}\right) > 0 \quad \text{for } \psi^{(1)}, \\ g(n) &= \left(n - \frac{l+m+1}{2}\right) \left(n - \frac{l+m}{2}\right) > 0 \quad \text{for } \psi^{(6)} \end{aligned} \quad (157)$$

for any admissible value of n , that is, $1 \leq n \leq \mathbf{m} - \frac{1}{2}$. However, these conditions are satisfied only for special values of \mathbf{l} and \mathbf{m} . As an illustration, for the Lamé equation ($\mathbf{l} = 0, \mathbf{l} = -1$), we find that

$$\begin{aligned} \text{for } \mathbf{l} = 0, \quad f(n) = g(n) > 0 \quad \text{only if } \mathbf{m} &= \frac{5}{2}, \frac{9}{2}, \frac{13}{2}, \dots; \\ \text{for } \mathbf{l} = -1 \quad (f \neq g), \quad f(n) > 0 \text{ and } g(n) > 0 \quad \text{only if } \mathbf{m} &= \frac{3}{2}, \frac{7}{2}, \frac{11}{2}, \dots \end{aligned} \quad (158)$$

So, we get two degenerate finite-series solutions for the Lamé equation when \mathbf{m} is half an odd positive integer – for $\mathbf{m} = 1/2$ see solutions after Eq. (152). These are not the finite series proposed by Ince [40].

On the other side,

$$f(n) \leq 0 \quad \text{if } \frac{m-l}{2} \leq n \leq \frac{m-l}{2} + \frac{1}{2}, \quad \text{and } g(n) \leq 0 \quad \text{if } \frac{m+l}{2} \leq n \leq \frac{m+l}{2} + \frac{1}{2}. \quad (159)$$

Then, Arscott condition holds if the above intervals are empty (no admissible n). In particular we can satisfy the condition by demanding that, for a given value of \mathbf{m} , the upper bounds of the intervals (159) are less than 1, that is, $\mathbf{m} - \mathbf{l} < 1$ and $\mathbf{m} + \mathbf{l} < 1$. Since $\mathbf{m} = 3/2, 5/2, \dots$, the Arscott condition stands if

$$\mathbf{l} > \mathbf{m} - 1 \text{ for } \psi^{(1)} \text{ (} \mathbf{l} \text{ is positive integer); } \mathbf{l} < 1 - \mathbf{m} \text{ for } \psi^{(6)} \text{ (} \mathbf{l} \text{ negative integer).} \quad (160)$$

In these cases, the degeneracy proof fails because \mathbf{l} takes different values in each expansion.

On the other hand, the degenerate eigenfunctions $\psi^{(4)}(u)$ and $\psi^{(7)}(u)$ are given by

$$\left\{ \begin{aligned} &\psi^{(4)}(u) \stackrel{(84,103)}{=} \text{sn } u (\text{dn } u)^{m-1} \sum_{n=0}^{m-1/2} b_n^{(4)} [(1-k^2) \text{sd}^2 u]^n \times \\ &\tilde{F} \left[n + \frac{l-m+2}{2}, \frac{m-l}{2}; n + \frac{3}{2}; (1-k^2) \text{sd}^2 u \right], \quad \mathbf{m} = \frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots, \quad [\text{odd}, 4K], \\ &\alpha_n^{(4)} = \left(1 - \frac{1}{k^2}\right) (n+1), \quad \beta_n^{(4)} = \left(\frac{1}{k^2} - 2\right) n^2 + \left[\left(\frac{1}{k^2} - 1\right) (\mathbf{l} - \mathbf{m} + 1) + \mathbf{m} - 1\right] n + \\ &\frac{(l+1-m)^2 - \varepsilon}{4k^2} + \frac{2lm - m^2 + 5m - 2 - l}{4}, \quad \gamma_n^{(4)} = \left(n + \frac{l-m+1}{2}\right) \left(n + \frac{l-m}{2}\right) \left(n - \mathbf{m} - \frac{1}{2}\right); \end{aligned} \right. \quad (161)$$

$$\left\{ \begin{aligned} &\psi^{(7)}(u) \stackrel{(87,103)}{=} (\text{dn } u)^m \sum_{n=0}^{m-1/2} b_n^{(7)} [(1-k^2) \text{sd}^2 u]^n \times \\ &\tilde{F} \left[n - \frac{l+m}{2}, \frac{m+l}{2}; n + \frac{1}{2}; (1-k^2) \text{sd}^2 u \right], \quad \mathbf{m} = \frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots, \quad [\text{even}, 4K], \\ &\alpha_n^{(7)} = \left(1 - \frac{1}{k^2}\right) (n+1), \quad \beta_n^{(7)} = \left(\frac{1}{k^2} - 2\right) n^2 + \left[\left(1 - \frac{1}{k^2}\right) (\mathbf{l} + \mathbf{m}) + \mathbf{m}\right] n + \\ &\frac{(l+m)^2 - \varepsilon}{4k^2} + \frac{l+m-2lm-m^2}{4}, \quad \gamma_n^{(7)} = \left(n - \frac{l+m+2}{2}\right) \left(n - \frac{l+m+1}{2}\right) \left(n - \mathbf{m} - \frac{1}{2}\right). \end{aligned} \right. \quad (162)$$

For one-term series ($\mathbf{m} = 1/2$), $\psi^{(4)}$ and $\psi^{(7)}$ are not independent respect to $\psi^{(1)}$ and $\psi^{(6)}$, since

$$\begin{aligned} \psi^{(4)}(u) = \psi^{(6)}(u) &= b_0^{(4)} \sqrt{\text{dn } u} \left[\left(\frac{\text{cn } u + \sqrt{k^2 - 1} \text{sn } u}{\text{dn } u} \right)^{l+\frac{1}{2}} - \left(\frac{\text{cn } u - \sqrt{k^2 - 1} \text{sn } u}{\text{dn } u} \right)^{l+\frac{1}{2}} \right], \quad \mathbf{m} = \frac{1}{2}, \\ \psi^{(7)}(u) = \psi^{(1)}(u) &= b_0^{(7)} \sqrt{\text{dn } u} \left[\left(\frac{\text{cn } u + \sqrt{k^2 - 1} \text{sn } u}{\text{dn } u} \right)^{l+\frac{1}{2}} + \left(\frac{\text{cn } u - \sqrt{k^2 - 1} \text{sn } u}{\text{dn } u} \right)^{l+\frac{1}{2}} \right], \quad \mathbf{m} = \frac{1}{2}, \end{aligned}$$

which are odd and even solutions, respectively, both having period $4K$. The previous expressions follow from relations (150).

To prove the degeneracy for $\mathbf{m} \geq 3/2$, first we redefine $b_n^{(7)}$ as

$$b_n^{(7)} = \left(\frac{k^2}{1-k^2} \right)^n \Gamma \left(n - \frac{l+m-1}{2} \right) \Gamma \left(n - \frac{l+m}{2} \right) \tilde{b}_n^{(7)}$$

and obtain the relations

$$- \left(n - \frac{l+m-1}{2} \right) \left(n - \frac{l+m}{2} \right) (n+1) \tilde{b}_{n+1}^{(7)} + \beta_n^{(7)} \tilde{b}_n^{(7)} + \frac{1-k^2}{k^2} \left(n - \mathbf{m} - \frac{1}{2} \right) \tilde{b}_{n-1}^{(7)} = 0.$$

Thence, if \mathbb{C} and \mathbb{D} represent the matrices corresponding to the recurrence relations for $b_n^{(4)}$ and $\tilde{b}_n^{(7)}$, we find that $\mathbb{D} = \mathbb{U}\mathbb{C}\mathbb{U}$, where \mathbb{U} is given in (155). Thus, $\psi^{(4)}$ and $\psi^{(7)}$ are degenerate because they satisfy the same characteristic equation.

For $\mathbf{m} = \frac{3}{2}$ the two-terms series $\psi^{(1)}$, $\psi^{(4)}$, $\psi^{(6)}$ and $\psi^{(7)}$ are all of them degenerate and each one corresponds to two real values, $\mathcal{E}^{(\pm)}$, for the energy \mathcal{E} (regardless of the Arscott condition). In fact, from the characteristic equations $\alpha_0^{(i)} \gamma_1^{(i)} - \beta_0^{(i)} \beta_1^{(i)} = 0$, we find

$$\mathcal{E}^{(\pm)} = \mathbf{l}^2 + \mathbf{l} + \frac{5}{2} + \frac{5}{2}k^2 \pm \sqrt{4(1-k^2)(\mathbf{l}^2 + \mathbf{l}) + 1}, \quad \mathbf{m} = \frac{3}{2}.$$

The even solutions, $\psi^{(1)}$ and $\psi^{(7)}$, can be written in term of elementary functions as

$$\begin{aligned} \psi^{(1)}(u) &= b_0^{(1)} (\text{dn } u)^{\frac{3}{2}} \left\{ \frac{1}{\Gamma(1/2)} \left[\left(\frac{\text{cn } u + \sqrt{k^2-1} \text{sn } u}{\text{dn } u} \right)^{\mathbf{l}-\frac{1}{2}} + \left(\frac{\text{cn } u - \sqrt{k^2-1} \text{sn } u}{\text{dn } u} \right)^{\mathbf{l}-\frac{1}{2}} \right] + \right. \\ &\quad \left. \frac{\sqrt{k^2-1}}{\Gamma(3/2)(1+2\mathbf{l})} \frac{\beta_0^{(1)}}{\alpha_0^{(1)}} \frac{\text{sn } u}{\text{dn } u} \left[\left(\frac{\text{cn } u + \sqrt{k^2-1} \text{sn } u}{\text{dn } u} \right)^{\mathbf{l}+\frac{1}{2}} - \left(\frac{\text{cn } u - \sqrt{k^2-1} \text{sn } u}{\text{dn } u} \right)^{\mathbf{l}+\frac{1}{2}} \right] \right\}, \quad \mathbf{m} = \frac{3}{2}, \\ \psi^{(7)}(u) &= b_0^{(7)} (\text{dn } u)^{\frac{3}{2}} \left\{ \frac{1}{\Gamma(1/2)} \left[\left(\frac{\text{cn } u + \sqrt{k^2-1} \text{sn } u}{\text{dn } u} \right)^{\mathbf{l}+\frac{3}{2}} + \left(\frac{\text{cn } u - \sqrt{k^2-1} \text{sn } u}{\text{dn } u} \right)^{\mathbf{l}+\frac{3}{2}} \right] + \right. \\ &\quad \left. \frac{\sqrt{k^2-1}}{\Gamma(3/2)(1+2\mathbf{l})} \frac{\beta_0^{(7)}}{\alpha_0^{(7)}} \frac{\text{sn } u}{\text{dn } u} \left[\left(\frac{\text{cn } u + \sqrt{k^2-1} \text{sn } u}{\text{dn } u} \right)^{\mathbf{l}+\frac{1}{2}} - \left(\frac{\text{cn } u - \sqrt{k^2-1} \text{sn } u}{\text{dn } u} \right)^{\mathbf{l}+\frac{1}{2}} \right] \right\}, \quad \mathbf{m} = \frac{3}{2}, \end{aligned}$$

where there are two expressions for $\beta_0^{(1)}$ and $\beta_0^{(7)}$ corresponding to $\mathcal{E} = \mathcal{E}^{(\pm)}$.

As $\alpha_n^{(4)} = \alpha_n^{(7)} < 0$ in (161) and (162), the Arscott condition (36) for $\mathbf{m} \geq 3/2$ requires that $\gamma_n^{(4)} < 0$ and $\gamma_n^{(7)} < 0$ or, equivalently, that

$$\begin{aligned} F(n) &= \left(n + \frac{l-m+1}{2} \right) \left(n + \frac{l-m}{2} \right) > 0 \quad \text{for } \psi^{(4)}, \\ G(n) &= \left(n - \frac{l+m+2}{2} \right) \left(n - \frac{l+m+1}{2} \right) > 0 \quad \text{for } \psi^{(7)}. \end{aligned} \quad (163)$$

These conditions are satisfied only for special values of \mathbf{l} and \mathbf{m} ; for the Lamé equation ($\mathbf{l} = 0$, $\mathbf{l} = -1$), we find that

$$\begin{aligned} \text{for } \mathbf{l} = 0, \quad F(n) > 0 \quad \text{and} \quad G(n) > 0 \quad \text{only if} \quad \mathbf{m} &= \frac{3}{2}, \frac{7}{2}, \frac{13}{2}, \dots; \\ \text{for } \mathbf{l} = -1, \quad F(n) = G(n) > 0 \quad \text{only if} \quad \mathbf{m} &= \frac{5}{2}, \frac{9}{2}, \frac{13}{2}, \dots \end{aligned} \quad (164)$$

By comparing (158) and (164), we see that for a given choice of (\mathbf{l}, \mathbf{m}) : (i) if the solutions $\psi^{(1)}$ and $\psi^{(6)}$ satisfy the Arscott condition, the solutions $\psi^{(4)}$ and $\psi^{(7)}$ do not satisfy; (ii) if the solutions $\psi^{(1)}$ and $\psi^{(6)}$ do not satisfy the Arscott condition, the solutions $\psi^{(4)}$ and $\psi^{(7)}$ satisfy the condition. Therefore, for the Lamé equation there are solutions satisfying the Arscott condition for all admissible values of (\mathbf{l}, \mathbf{m}) .

The above conclusion is not restricted to Lamé equation. Since

$$F(n) \leq 0 \text{ if } \frac{m-l-1}{2} \leq n \leq \frac{m-l}{2}, \text{ and } G(n) \leq 0 \text{ if } \frac{m+l+1}{2} \leq n \leq \frac{m+l+2}{2}, \quad (165)$$

Arcscott condition for $\psi^{(4)}$ and $\psi^{(7)}$ holds if the above intervals are empty similarly to the previous cases. As n is an integer, in the intervals (159) and (165) we can take the sign $<$ instead of \leq . Thus, for $\psi^{(4)}$ and $\psi^{(1)}$ we have, respectively,

$$\psi^{(4)}, \psi^{(1)} : F(n) \leq 0 \text{ if } \frac{m-l-1}{2} < n < \frac{m-l}{2}, \text{ and } f(n) \leq 0 \text{ if } \frac{m-l}{2} < n < \frac{m-l+1}{2}.$$

Hence, at least one of the intervals is empty for $n = 2, 3, 4, \dots$. Analogously, for $\psi^{(6)}$ and $\psi^{(7)}$ we have

$$\psi^{(6)}, \psi^{(7)} : g(n) \leq 0 \text{ if } \frac{m+l}{2} < n < \frac{m+l+1}{2}, \text{ and } G(n) \leq 0 \text{ if } \frac{m+l+1}{2} < n < \frac{m+l+2}{2}.$$

5.3.2 Finite series when l is an integer and $m \neq -1/2$ is half an odd negative integer

If $m \neq -1/2$ is half an odd negative integer, finite-series solutions $\psi^{(i)}$ with $0 \leq n \leq -m - 3/2$ result from $H^{(i)}$ for $i = 2, 3, 5, 8$. Now the series terminate at $n = -m - 3/2$ because $(n - \alpha + \beta) = (n + m + 1/2)$ implies $\gamma_n^{(i)} = 0$ for $n = -m - 1/2$ as demanded in (34). Since $\delta = 1/2$, by (89) the four solutions converge when $z = 1$ ($\text{sn}^2 u = 1$).

The eigenfunctions $\psi^{(2)}(u)$ and $\psi^{(5)}(u)$ are

$$\left\{ \begin{array}{l} \psi^{(2)}(u) \stackrel{(82,103)}{=} \frac{\text{sn } u}{\text{dn}^{m+2} u} \sum_{n=0}^{-m-3/2} b_n^{(2)} [(1-k^2) \text{sd}^2 u]^n \times \\ \tilde{F} \left[n + \frac{l+m+3}{2}, -\frac{m+l+1}{2}; n + \frac{3}{2}; (1-k^2) \text{sd}^2 u \right], \quad m = -\frac{3}{2}, -\frac{5}{2}, \dots, [\text{odd}, 4K], \\ \alpha_n^{(2)} = \left(1 - \frac{1}{k^2}\right) (n+1), \quad \beta_n^{(2)} = \left(\frac{1}{k^2} - 2\right) n^2 + \left[l - \frac{l+m+2}{k^2}\right] n + \frac{(l+m+2)^2 - \varepsilon}{4k^2} \\ -\frac{2lm+3l+m^2+7m+8}{4}, \quad \gamma_n^{(2)} = \left(n + \frac{l+m+1}{2}\right) \left(n + \frac{l+m+2}{2}\right) \left(n + m + \frac{1}{2}\right); \end{array} \right. \quad (166)$$

$$\left\{ \begin{array}{l} \psi^{(5)}(u) \stackrel{(85,103)}{=} \frac{1}{\text{dn}^{m+1} u} \sum_{n=0}^{-m-3/2} b_n^{(5)} [(1-k^2) \text{sd}^2 u]^n \times \\ \tilde{F} \left[n + \frac{m-l+1}{2}, \frac{l-m-1}{2}; n + \frac{1}{2}; (1-k^2) \text{sd}^2 u \right], \quad m = -\frac{3}{2}, -\frac{5}{2}, \dots, [\text{even}, 4K], \\ \alpha_n^{(5)} = \left(1 - \frac{1}{k^2}\right) (n+1), \quad \beta_n^{(5)} = \left(\frac{1}{k^2} - 2\right) n^2 - \left[l + \frac{m-l+1}{k^2}\right] n + \frac{(l-m-1)^2 - \varepsilon}{4k^2} \\ -\frac{(l-m-1)^2 + l+m+2 - (l+1)^2}{4}, \quad \gamma_n^{(5)} = \left(n + \frac{m-l}{2}\right) \left(n + \frac{m-l-1}{2}\right) \left(n + m + \frac{1}{2}\right). \end{array} \right. \quad (167)$$

We can verify that $\psi^{(2)}$ and $\psi^{(5)}$ result, respectively, from $\psi^{(6)}$ and $\psi^{(1)}$ by means of the substitutions (104). By this reason we do not discuss the degeneracy and the Arcscott condition.

On the other side, the eigenfunctions $\psi^{(3)}(u)$ and $\psi^{(8)}(u)$ are

$$\left\{ \begin{array}{l} \psi^{(3)}(u) \stackrel{(83,103)}{=} (\operatorname{dn} u)^{-m-1} \sum_{n=0}^{-m-3/2} b_n^{(3)} [(1-k^2) \operatorname{sd}^2 u]^n \times \\ \tilde{F} \left[n + \frac{l+m+2}{2}, -\frac{m+l+2}{2}; n + \frac{1}{2}; (1-k^2) \operatorname{sd}^2 u \right], \quad \mathbf{m} = \frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots [\text{even}, 4K], \\ \alpha_n^{(3)} = \left(1 - \frac{1}{k^2}\right) (n+1), \quad \beta_n^{(3)} = \left(\frac{1}{k^2} - 2\right) n^2 + \left[\left(\frac{1}{k^2} - 1\right) (l + \mathbf{m} + 2) - \mathbf{m} - 1\right] n \\ + \frac{(l+m+2)^2 - \varepsilon}{4k^2} - \frac{1}{4} [l + \mathbf{m} + 2 + 2(l+1)(\mathbf{m}+1) + (\mathbf{m}+1)^2], \\ \gamma_n^{(3)} = \left(n + \frac{l+m}{2}\right) \left(n + \frac{l+m+1}{2}\right) \left(n + \mathbf{m} + \frac{1}{2}\right); \end{array} \right. \quad (168)$$

$$\left\{ \begin{array}{l} \psi^{(8)}(u) \stackrel{(88,103)}{=} \operatorname{sn} u (\operatorname{dn} u)^{-m-2} \sum_{n=0}^{-m-3/2} b_n^{(8)} [(1-k^2) \operatorname{sd}^2 u]^n \times \\ \tilde{F} \left[n - \frac{l-m-2}{2}, \frac{l-m}{2}; n + \frac{3}{2}; (1-k^2) \operatorname{sd}^2 u \right], \quad \mathbf{m} = -\frac{3}{2}, -\frac{5}{2}, -\frac{7}{2}, \dots [\text{odd}, 4K], \\ \alpha_n^{(8)} = \left(1 - \frac{1}{k^2}\right) (n+1), \quad \beta_n^{(8)} = \left(\frac{1}{k^2} - 2\right) n^2 + \left[\left(1 - \frac{1}{k^2}\right) (l - \mathbf{m} - 1) - \mathbf{m} - 2\right] n + \\ \frac{(l-1-\mathbf{m})^2 - \varepsilon}{4k^2} + \frac{2(l+1)(\mathbf{m}+1) - (\mathbf{m}+1)^2 - 5\mathbf{m} - 6 + l}{4}, \\ \gamma_n^{(8)} = \left(n - \frac{l-m-1}{2}\right) \left(n - \frac{l-m}{2}\right) \left(n + \mathbf{m} + \frac{1}{2}\right). \end{array} \right. \quad (169)$$

$\psi^{(3)}$ and $\psi^{(8)}$ result, respectively, from $\psi^{(7)}$ and $\psi^{(4)}$ by means of the substitutions (104).

5.4 Finite series $\Psi^{(i)}(u)$ of hypergeometric functions when \mathbf{m} is an integer and $l \neq -1/2$ is half an odd integer

These solutions are obtained from the expansions $\mathbf{H}^{(i)}(x)$ given in Section 4.3; alternatively, they can be generated from the above solutions (Sec. 5.3) through the substitutions (105). If $l \neq -1/2$ is half an odd integer, then the factor $(n - \alpha - \beta + \gamma + \delta) = (n - l - 1/2)$ implies that four of the coefficients $\gamma_n^{(i)}$ vanish for $n = l + 1/2$. These give finite series with $0 \leq n \leq l - \frac{1}{2}$ if l is positive; according to (99) the expansions converge for all values of u . Similarly, the factor $(n + \alpha + \beta - \gamma - \delta) = (n + l + 1/2)$ implies that four expansions terminate at $n = -l - 3/2$ if $l \leq -3/2$, and by (99) the eigenfunctions are bounded for all permissible values of u .

Due to the aforementioned connection with solutions of Section 5.3, the eigenfunctions are degenerate and a given expansion satisfies the Arscott condition (36) only for special values of \mathbf{m} and l . However, once more, by considering all the expansions, we find at least one expansion satisfying the condition for any allowed value of (\mathbf{m}, l) .

5.4.1 Finite series when \mathbf{m} is an integer and l is half an odd positive integer

Now we write the degenerate eigenfunctions $\Psi^{(5)}(u)$ and $\Psi^{(7)}(u)$. By using the Eqs. (106) we find they can be obtained from $\Psi^{(1)}(u)$ and $\Psi^{(6)}(u)$.

$$\left\{ \begin{array}{l} \Psi^{(5)}(u) \stackrel{(95,103)}{=} \frac{1}{\operatorname{dn}^l u} \sum_{n=0}^{l-1/2} b_n^{(5)} \operatorname{cn}^{2n} u \tilde{F} \left(n - \frac{l-m-1}{2}, \frac{l-m-1}{2}; n + \frac{1}{2}; \operatorname{cn}^2 u \right) \\ \stackrel{(106)}{=} \Psi^{(1)}(u + K) \Big|_{l \leftrightarrow \mathbf{m}}, \quad l = \frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots [\text{even, period } 4K] \\ \alpha_n^{(5)} = \alpha_n^{(1)} \Big|_{l \leftrightarrow \mathbf{m}} = \alpha_n^{(1)}, \quad \beta_n^{(5)} = \beta_n^{(1)} \Big|_{l \leftrightarrow \mathbf{m}}, \quad \gamma_n^{(5)} = \gamma_n^{(1)} \Big|_{l \leftrightarrow \mathbf{m}}; \end{array} \right. \quad (170)$$

$$\left\{ \begin{array}{l} \Psi^{(7)}(u) \stackrel{(97,103)}{=} \frac{\text{cn } u}{\text{dn } u} \sum_{n=0}^{l-1/2} b_n^{(7)} \text{cn}^{2n} u \tilde{F}\left(n - \frac{l+m-1}{2}, \frac{l+m+1}{2}; n + \frac{3}{2}; \text{cn}^2 u\right) \\ \stackrel{(106)}{=} \Psi^{(6)}(u+K) \Big|_{l \leftrightarrow m}, \quad l = \frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots, \quad [\text{odd, period } 4K] \\ \alpha_n^{(7)} = \alpha_n^{(6)} \Big|_{l \leftrightarrow m} = \alpha_n^{(6)}, \quad \beta_n^{(7)} = \alpha_n^{(6)} \Big|_{l \leftrightarrow m}, \quad \gamma_n^{(7)} = \gamma_n^{(6)} \Big|_{l \leftrightarrow m}, \end{array} \right. \quad (171)$$

On the other side, degenerate eigenfunctions $\Psi^{(6)}(u)$ and $\Psi^{(8)}(u)$ result from $\mathbf{H}^{(6)}(u)$ and $\mathbf{H}^{(8)}(u)$. By Eqs. (106)) we see that they can also be obtained from $\Psi^{(7)}(u)$ and $\Psi^{(4)}(u)$, respectively.

$$\left\{ \begin{array}{l} \Psi^{(6)}(u) \stackrel{(96,103)}{=} \frac{1}{\text{dn } u} \sum_{n=0}^{l-1/2} b_n^{(6)} \text{cn}^{2n} u \tilde{F}\left(n - \frac{l+m}{2}, \frac{l+m}{2}; n + \frac{1}{2}; \text{cn}^2 u\right) \\ \stackrel{(106)}{=} \Psi^{(7)}(u+K) \Big|_{l \leftrightarrow m}, \quad l = \frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots, \quad [\text{even, period } 4K] \\ \alpha_n^{(6)} = \alpha_n^{(7)} \Big|_{l \leftrightarrow m} = \alpha_n^{(7)}, \quad \beta_n^{(6)} = \beta_n^{(7)} \Big|_{l \leftrightarrow m}, \quad \gamma_n^{(6)} = \gamma_n^{(7)} \Big|_{l \leftrightarrow m}; \end{array} \right. \quad (172)$$

$$\left\{ \begin{array}{l} \Psi^{(8)}(u) \stackrel{(98,103)}{=} \frac{\text{cn } u}{\text{dn } u} \sum_{n=0}^{l-1/2} b_n^{(8)} \text{cn}^{2n} u \tilde{F}\left(n - \frac{l-m-2}{2}, \frac{l-m}{2}; n + \frac{3}{2}; \text{cn}^2 u\right) \\ \stackrel{(106)}{=} \Psi^{(4)}(u+K) \Big|_{l \leftrightarrow m}, \quad l = \frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots, \quad [\text{odd, period } 4K] \\ \alpha_n^{(8)} = \alpha_n^{(4)} \Big|_{l \leftrightarrow m} = \alpha_n^{(4)}, \quad \beta_n^{(8)} = \alpha_n^{(4)} \Big|_{l \leftrightarrow m}, \quad \gamma_n^{(8)} = \gamma_n^{(4)} \Big|_{l \leftrightarrow m}, \end{array} \right. \quad (173)$$

For one-term series ($l = 1/2$), only two expansions are independent since

$$\begin{aligned} \psi^{(5)}(u) &= \Psi^{(6)}(u) = \frac{b_0^{(5)}}{\sqrt{\text{dn } u}} \left[(\text{cn } u - i \text{sn } u)^{m+\frac{1}{2}} + (\text{cn } u + i \text{sn } u)^{m+\frac{1}{2}} \right], \quad l = \frac{1}{2}, \\ \psi^{(7)}(u) &= \Psi^{(8)}(u) = \frac{b_0^{(7)}}{\sqrt{\text{dn } u}} \left[(\text{cn } u - i \text{sn } u)^{m+\frac{1}{2}} - (\text{cn } u + i \text{sn } u)^{m+\frac{1}{2}} \right], \quad l = \frac{1}{2}. \end{aligned}$$

These are even and odd solutions, respectively, both with period $4K$ and energy $\mathcal{E} = (m + \frac{1}{2})^2 + \frac{k^2}{4}$. We can also verify that, for $l = 3/2$, the four two-term series are degenerate with energies $\mathcal{E}^{(\pm)} = m^2 + m + \frac{5}{2} + \frac{5}{2}k^2 \pm \sqrt{4(1-k^2)(m^2 + m)} + 1$.

5.4.2 Finite series when m is an integer and $l \neq -1/2$ is half an odd negative integer

For the eigenfunctions $\Psi^{(1)}(u)$ and $\Psi^{(3)}(u)$ we find

$$\left\{ \begin{array}{l} \Psi^{(1)}(u) \stackrel{(91,103)}{=} \text{dn}^{l+1} u \sum_{n=0}^{-l-3/2} b_n^{(1)} \text{cn}^{2n} u \tilde{F}\left(n + \frac{l-m+1}{2}, \frac{m-l-1}{2}; n + \frac{1}{2}; \text{cn}^2 u\right) \\ \stackrel{(106)}{=} \psi^{(5)}(u+K) \Big|_{l \leftrightarrow m}, \quad l = -\frac{3}{2}, -\frac{5}{2}, -\frac{7}{2}, \dots, \quad [\text{even, period } 4K] \\ \alpha_n^{(1)} = \alpha_n^{(5)} \Big|_{l \leftrightarrow m} = \alpha_n^{(5)}, \quad \beta_n^{(1)} = \beta_n^{(5)} \Big|_{l \leftrightarrow m}, \quad \gamma_n^{(1)} = \gamma_n^{(5)} \Big|_{l \leftrightarrow m}; \end{array} \right. \quad (174)$$

$$\left\{ \begin{array}{l} \Psi^{(3)}(u) \stackrel{(93,103)}{=} \text{cn } u \text{ dn}^{l+1} u \sum_{n=0}^{-l-3/2} b_n^{(3)} \text{cn}^{2n} u \tilde{F}\left(n + \frac{l+m+3}{2}, -\frac{l+m+1}{2}; n + \frac{3}{2}; \text{cn}^2 u\right) \\ \stackrel{(106)}{=} \psi^{(2)}(u+K) \Big|_{l \leftrightarrow m}, \quad l = -\frac{3}{2}, -\frac{5}{2}, -\frac{7}{2}, \dots, \quad [\text{odd, period } 4K] \\ \alpha_n^{(3)} = \alpha_n^{(2)} \Big|_{l \leftrightarrow m} = \alpha_n^{(2)}, \quad \beta_n^{(3)} = \beta_n^{(2)} \Big|_{l \leftrightarrow m}, \quad \gamma_n^{(3)} = \gamma_n^{(2)} \Big|_{l \leftrightarrow m}. \end{array} \right. \quad (175)$$

$\Psi^{(1)}$ and $\Psi^{(3)}$ follow also from $\Psi^{(5)}$ and $\Psi^{(7)}$ by the substitutions (104).

The degenerate eigenfunctions $(\Psi^{(2)}, \Psi^{(4)})$, coming from $(\mathbf{H}^{(2)}, \mathbf{H}^{(4)})$, are given by

$$\left\{ \begin{array}{l} \Psi^{(2)}(u) \stackrel{(92,103)}{=} \text{dn}^{l+1} u \sum_{n=0}^{-l-\frac{3}{2}} b_n^{(2)} \text{cn}^{2n} u \tilde{F}\left(n + \frac{l+m+2}{2}, -\frac{m+l+2}{2}; n + \frac{1}{2}; \text{cn}^2 u\right) \\ \stackrel{(106)}{=} \psi^{(3)}(u+K) \Big|_{l \leftrightarrow m}, \quad l = -\frac{3}{2}, -\frac{5}{2}, -\frac{7}{2}, \dots, \quad [\text{even, period } 4K] \\ \alpha_n^{(2)} = \alpha_n^{(3)} \Big|_{l \leftrightarrow m} = \alpha_n^{(3)}, \quad \beta_n^{(2)} = \beta_n^{(3)} \Big|_{l \leftrightarrow m}, \quad \gamma_n^{(2)} = \gamma_n^{(3)} \Big|_{l \leftrightarrow m}; \end{array} \right. \quad (176)$$

$$\left\{ \begin{array}{l} \Psi^{(4)}(u) \stackrel{(94,103)}{=} \text{cn } u \text{ dn}^{l+1} u \sum_{n=0}^{-l-\frac{3}{2}} b_n^{(4)} \text{cn}^{2n} u \tilde{F}\left(n + \frac{l-m+2}{2}, \frac{m-l}{2}; n + \frac{3}{2}; \text{cn}^2 u\right) \\ \stackrel{(106)}{=} \psi^{(8)}(u+K) \Big|_{l \leftrightarrow m}, \quad l = -\frac{3}{2}, -\frac{5}{2}, -\frac{7}{2}, \dots, \quad [\text{odd, period } 4K] \\ \alpha_n^{(4)} = \alpha_n^{(8)} \Big|_{l \leftrightarrow m} = \alpha_n^{(8)}, \quad \beta_n^{(4)} = \beta_n^{(8)} \Big|_{l \leftrightarrow m}, \quad \gamma_n^{(4)} = \gamma_n^{(8)} \Big|_{l \leftrightarrow m}. \end{array} \right. \quad (177)$$

Notice that $\Psi^{(2)}$ and $\Psi^{(4)}$ follow from $\Psi^{(6)}$ and $\Psi^{(8)}$ by the substitutions (104). Beside this, for one-term series ($l = -3/2$): $\Psi^{(1)} = \Psi^{(2)}$ and $\Psi^{(3)} = \Psi^{(4)}$.

5.5 Infinite-series solutions

To obtain infinite-series solutions we require that $\mathring{\gamma}_n^{(i)} \neq 0$ in the power series expansions $\mathring{H}^{(i)}(x)$ written in section 3.1. This imposes restrictions on the parameters \mathbf{l} and \mathbf{m} : one condition on $\mathbf{l} - \mathbf{m}$ and other on $\mathbf{l} + \mathbf{m}$. Since $a = 1/k^2 > 1$, the relations (51) imply that the eigenfunctions are convergent and bounded for any value of the independent variable, that is, for $0 \leq x = \text{sn}^2 u \leq 1$. They are denoted by $\mathring{\Phi}^{(i)}$ while the series coefficients are denoted by $\mathring{b}_n^{(i)}$,

$$\mathring{\alpha}_n^{(i)} \mathring{b}_{n+1}^{(i)} + \mathring{\beta}_n^{(i)} \mathring{b}_n^{(i)} + \mathring{\gamma}_n^{(i)} \mathring{b}_{n-1}^{(i)} = 0, \quad \mathring{b}_{-1}^{(i)} = 0, \quad n \geq 0, \quad (178)$$

$\mathring{\alpha}_n^{(i)}$, $\mathring{\beta}_n^{(i)}$ and $\mathring{\gamma}_n^{(i)}$ being defined in Section 5.1. Below

- we write down the conditions to have infinite series;
- we show that each finite series corresponds to four infinite-series expansions (same values for \mathbf{l} and \mathbf{m}).

Therefore, for this family of quasi-exactly solvable potentials, for solutions of the Schrödinger equation we have to regard both finite-series and infinite-series wavefunctions since the infinite series are convergent and bounded for all values of the independent variable, in opposition to the case mentioned in [12].

By inserting the expansions $\mathring{H}^{(i)}(x)$ into (103), we get four types of even and odd infinite-series solutions (periods $2K$ and $4K$), namely,

$$\text{even, } 2K : \begin{cases} \mathring{\Phi}^{(1)}(u) \stackrel{(43,103)}{=} \text{dn}^{l+1}u \sum_{n=0}^{\infty} \mathring{b}_n^{(1)} \text{sn}^{2n}u, & \text{if } \frac{l-m-1}{2} \text{ and } \frac{l+m}{2} \text{ are not} \\ & \text{negative integers,} \\ \mathring{\Phi}^{(5)}(u) \stackrel{(47,103)}{=} \frac{1}{\text{dn}^l u} \sum_{n=0}^{\infty} \mathring{b}_n^{(5)} \text{sn}^{2n}u, & \text{if } \frac{l-m+1}{2} \text{ and } \frac{l+m+2}{2} \text{ are not} \\ & \text{positive integers;} \end{cases} \quad (179)$$

$$\text{odd, } 4K : \begin{cases} \mathring{\Phi}^{(2)}(u) \stackrel{(44,103)}{=} \text{sn}u \text{dn}^{l+1}u \sum_{n=0}^{\infty} \mathring{b}_n^{(2)} \text{sn}^{2n}u, & \text{if } \frac{l-m}{2} \text{ and } \frac{l+m+1}{2} \text{ are not} \\ & \text{negative integers,} \\ \mathring{\Phi}^{(6)}(u) \stackrel{(48,103)}{=} \frac{\text{sn}u}{\text{dn}^l u} \sum_{n=0}^{\infty} \mathring{b}_n^{(6)} \text{sn}^{2n}u, & \text{if } \frac{l-m}{2} \text{ and } \frac{l+m+1}{2} \text{ are not} \\ & \text{positive integers;} \end{cases} \quad (180)$$

$$\text{even, } 4K : \begin{cases} \mathring{\Phi}^{(3)}(u) \stackrel{(45,103)}{=} \text{cn}u \text{dn}^{l+1}u \sum_{n=0}^{\infty} \mathring{b}_n^{(3)} \text{sn}^{2n}u, & \text{if } \frac{l-m}{2} \text{ and } \frac{l+m+1}{2} \text{ are not} \\ & \text{negative integers,} \\ \mathring{\Phi}^{(7)}(u) \stackrel{(49,103)}{=} \frac{\text{cn}u}{\text{dn}^l u} \sum_{n=0}^{\infty} \mathring{b}_n^{(7)} \text{sn}^{2n}u, & \text{if } \frac{l-m}{2} \text{ and } \frac{l+m+1}{2} \text{ are not} \\ & \text{positive integers;} \end{cases} \quad (181)$$

$$\text{odd, } 2K : \begin{cases} \mathring{\Phi}^{(4)}(u) \stackrel{(46,103)}{=} \text{sn}u \text{cn}u \text{dn}^{l+1}u \sum_{n=0}^{\infty} \mathring{b}_n^{(4)} \text{sn}^{2n}u, & \text{if } \frac{l-m+1}{2} \text{ and } \frac{l+m+2}{2} \text{ are} \\ & \text{not negative integers,} \\ \mathring{\Phi}^{(8)}(u) \stackrel{(50,103)}{=} \frac{\text{sn}u \text{cn}u}{\text{dn}^l u} \sum_{n=0}^{\infty} \mathring{b}_n^{(8)} \text{sn}^{2n}u, & \text{if } \frac{l-m-1}{2} \text{ and } \frac{l+m}{2} \text{ are} \\ & \text{not positive integers,} \end{cases} \quad (182)$$

The expansions in each pair are related to each other by $l \mapsto -l-1$ and $m \mapsto -m-1$. From the above conditions we will see that there are four types of infinite-series solutions corresponding to a given finite-series solution.

First, consider the cases in which either l or m is an integer and the other is half an odd integer, Sections 5.3 and 5.4. Then, the eight expansions $\mathring{\Phi}^{(i)}(u)$ are given by infinite series which, however, reduce to four expansions because Eq. (56) implies that $\mathring{\Phi}^{(i+4)}(u) = \mathring{\Phi}^{(i)}(u)$ for $i = 1, 2, 3, 4$.

As examples for finite series $(\mathring{\psi}^{(i)}, \tilde{\mathring{\psi}}^{(i)})$ in terms of sn^2u , we consider the same cases given in Section 5.1. Thus, the finite-series $\mathring{\psi}^{(1)}$, given in (126), is associated with the four infinite-series expansions: $\mathring{\phi}^{(2)} = \mathring{\phi}^{(6)}$, $\mathring{\phi}^{(3)} = \mathring{\phi}^{(7)}$, $\mathring{\phi}^{(5)}$ and $\mathring{\phi}^{(8)}$. For the second example,

$\mathbf{m} = -1/2$. Then $\psi^{(j)} = \tilde{\psi}^{(j)}$ and, by using the values for \mathbf{l} written in (127-130), we find four infinite-series expansions corresponding to each of the finite series according to

$$\psi^{(i)} = \tilde{\psi}^{(i)} \mapsto \mathring{\Phi}^{(i+4)}, \quad \psi^{(i+4)} = \tilde{\psi}^{(i+4)} \mapsto \mathring{\Phi}^{(i)}, \quad (183)$$

where i assumes the four values $i = 1, 2, 3, 4$ in each case. In the third example, we take $\mathbf{m} = 0$ for Eq. (107); thence, from the values for \mathbf{l} written in (131-134), we obtain the correspondence

$$\tilde{\psi}^{(i)} \mapsto \mathring{\Phi}^{(i+4)}, \quad \tilde{\psi}^{(i+4)} \mapsto \mathring{\Phi}^{(i)}, \quad (184)$$

where i assumes the four values $i = 1, 2, 3, 4$ once more.

As an example of finite-series $(\mathring{\Psi}^{(i)}, \tilde{\Psi}^{(i)})$ in terms of $\text{cn}^2 u$, given in Section 5.2, we take the Lamé equation with $\mathbf{l} = 0$. Then, by using the parameters written in (146-149), we get the following four types of infinite-series solutions corresponding each of the 16 finite series:

$$(\mathring{\Psi}^{(1)}, \tilde{\Psi}^{(1)}), (\mathring{\Psi}^{(4)}, \tilde{\Psi}^{(4)}), (\mathring{\Psi}^{(6)}, \tilde{\Psi}^{(6)}), (\mathring{\Psi}^{(7)}, \tilde{\Psi}^{(7)}) \mapsto \mathring{\Phi}^{(2,3,5,8)}; \quad (185)$$

$$(\mathring{\Psi}^{(2)}, \tilde{\Psi}^{(2)}), (\mathring{\Psi}^{(3)}, \tilde{\Psi}^{(3)}), (\mathring{\Psi}^{(5)}, \tilde{\Psi}^{(5)}), (\mathring{\Psi}^{(8)}, \tilde{\Psi}^{(8)}) \mapsto \mathring{\Phi}^{(1,4,6,7)}. \quad (186)$$

More examples are given in Appendix C.

6 Conclusions

By treating the elliptic Darboux equation as an instance of Heun's general equation, we have applied known transformations of variables to generate new solutions for the latter equation. In Section 3 we have established two groups of power series expansions and, in Section 4, two groups of expansions in series of Gauss hypergeometric functions. Each group is constituted by eight expansions connect to each other by homotopic transformations of the dependent variable. After this, the solutions of the Heun equation can be particularized for Darboux equation and its special cases.

The relations among the eight expansions of each group require further study. In Section 3 we have seen that the solutions $\mathring{U}^{(i)}$ reduce to four expansions under certain circumstances. On the other side, Ince [33, 40] has discussed these relations for some solutions for the particular case of the Lamé equation.

The solutions we have considered may be used to solve the stationary Schrödinger equation for quasi-exactly solvable (QES) potentials. For this purpose, firstly, in section 2.1, we have seen that the Schrödinger equation for certain elliptic potentials reduces to Darboux equations. For particular values of the parameters, some finite-series solutions to each of such potentials are already known in the literature [17, 20], but finite series turn out to be systematically generated from solutions for the Heun equation without the need of specifying the parameters. This procedure provides as well suitable infinite-series eigenfunctions.

As an example, in Section 5 we have considered the Schrödinger equation for the associated Lamé potential (17) which depends on the parameters \mathbf{l} and \mathbf{m} . When either $\mathbf{m} - \mathbf{l}$ or $\mathbf{l} + \mathbf{m}$ is an integer, the solutions $\mathring{H}^{(i)}(x)$ and $\mathring{H}^{(i)}(x)$ of the Heun equation lead to finite-series eigenfunctions given by expansions in series of $\text{sn}^2 u$ and $\text{cn}^2 u$, respectively. To assure real and distinct energy eigenvalues, we use the Arscott condition (36) which, in turn, imposes further restrictions on \mathbf{l} and \mathbf{m} (such restrictions are written on the

right-hand side of the expansions given in Sections 5.1 and 5.2). Finite series in terms of $\text{sn}^2 u$ and $\text{cn}^2 u$ cannot satisfy simultaneously the Arscott condition because this condition requires that $\dot{\gamma}_n^{(i)} > 0$ and $\dot{\gamma}_n^{(i)} < 0$, respectively. This fact gives a criterion to select the appropriate solutions for the associated Lamé equation, as in the examples of Sections 5.1, 5.2 and Appendix C, where the use of each of the preceding finite-series expansions has been illustrated. From the solutions of Sections 5.1 and 5.2 we can generate new solutions by means of the substitutions (105).

On the other hand, when either \mathbf{l} or \mathbf{m} is an integer and the other is half an odd integer, we have found eigenfunctions by using the hypergeometric-function expansions $H^{(i)}(x)$ and $\mathbf{H}^{(i)}(x)$ given in Section 4. If \mathbf{l} or \mathbf{m} is a positive or a negative integer all the expansions afford finite-series solutions bounded for all values of the independent variable u . Thus, if \mathbf{l} is an integer and $\mathbf{m} \neq -1/2$ is half an odd integer we have the degenerate finite-series solutions given in Section 5.3; for $\mathbf{l} = -1$ or $\mathbf{l} = 0$, we get new solutions for the Lamé equation. If \mathbf{m} is an integer and $\mathbf{l} \neq -1/2$ is half an odd integer, we have the degenerate finite-series solutions given in Section 5.4. In fact, the solutions of Sections 5.3 and 5.4 are connected by means of the substitutions (105). The eigenfunctions are degenerate in the sense that there are two independent eigenfunctions corresponding to the same energy. The degeneracy has been shown by writing the recurrence relations in the matrix form.

However, Khare and Sukhatme [19, 20] have pointed out another type of degeneracy which is not obtained by using expansions in series of hypergeometric functions. In fact, this type of degenerate solutions comes from expansions in series of $\text{cn}^2 u$ or $\text{sn}^2 u$ for particular values for \mathbf{l} and \mathbf{m} , as explained in Appendix C.

The previous examples show how to use the solutions of the Heun equation to find finite-series solutions for the elliptic Darboux equation or, more precisely, for the associated Lamé equation. Furthermore, to each finite-series solution, in Section 5.5 we have seen that this procedure also gives four periodic infinite-series solutions which are bounded and convergent for any value of the independent variable. Hitherto, relations among these have not been determined even for the known case of the Lamé equation [40]. We note that the existence of such periodic solutions for the associated Lamé equation can also be studied by using expansions in series of trigonometric functions like the ones considered in References [40, 41, 42]. In Appendix D we explain how these solutions could be obtained as particular instances of solutions of the Heun equation (2).

Anyway, in the present case, infinite series as solutions for quasi-exactly solvable problems cannot be discarded by asserting that the wavefunctions are unbounded, as sometimes occurs [12]. Besides this, it is known that the infinite series are important in band theory of solids. Indeed, these are the only type of eigenstates admitted by a model proposed by Slater in which the Schrödinger equation reduces to a Mathieu equation [37] (this does not admit finite-series solutions).

Although we have considered only the associated Lamé potential, the solutions of the Heun equation can also be applied for the other potentials presented in Section 2.1. To finalize, we mention some supplementary issues. In the first place, for the Heun equation, there are other expansions in series of hypergeometric functions [13, 14, 15, 43] which, however, are not appropriate for the problems of Section 2.1. The same is true respecting a recent study on Darboux equation [44].

In the second place, one of the Erdélyi expansions in series of hypergeometric functions seems to constitute a suitable start-point for generating new solutions for the associated Lamé equation. To this end, from the inception we have to modify the recurrence relations for series coefficients as in Appendix D and, after that, use the homotopic transformations as well as the fractional substitutions (26-28). Similarly, by using modified recurrence relations we see that the Svartholm solution [13] leads to expansions in series of trigonometric

functions for the associated Lamé equation. However, in order to compare solutions obtained from Svartholm's and Erdelyi's expansions with the ones presented in this article, it would be necessary to find out if the finite-series expansions fulfill the Arscott condition (36).

Acknowledgement

This work was funded by Ministério da Ciência, Tecnologia e Inovações of Brazil. The author thanks Léa Jaccoud El-Jaick for computations of section 4.1 and for reading the manuscript. This work does not have any conflicts of interest.

A Homotopic transformations

The 8 homotopic transformations (29) are:

$$T_1 H(x) = H(x) = H(a, q; \alpha, \beta, \gamma, \delta; x), \quad (\text{identity}) \quad (\text{A.1})$$

$$T_2 H(x) = x^{1-\gamma} H[a, q - (\gamma - 1)(\delta a + \epsilon); \beta - \gamma + 1, \alpha - \gamma + 1, 2 - \gamma, \delta; x], \quad (\text{A.2})$$

$$T_3 H(x) = (1 - x)^{1-\delta} H[a, q - (\delta - 1)\gamma a; \beta - \delta + 1, \alpha - \delta + 1, \gamma, 2 - \delta; x], \quad (\text{A.3})$$

$$T_4 H(x) = x^{1-\gamma} (1 - x)^{1-\delta} \times \\ H[a, q - (\gamma + \delta - 2)a - (\gamma - 1)\epsilon; \alpha - \gamma - \delta + 2, \beta - \gamma - \delta + 2, 2 - \gamma, 2 - \delta; x], \quad (\text{A.4})$$

$$T_5 H(x) = \left[1 - \frac{x}{a}\right]^{1-\epsilon} H[a, q - \gamma(\alpha + \beta - \gamma - \delta); -\alpha + \gamma + \delta, -\beta + \gamma + \delta, \gamma, \delta; x], \quad (\text{A.5})$$

$$T_6 H(x) = \left[1 - \frac{x}{a}\right]^{1-\epsilon} x^{1-\gamma} \times \\ H[a, q - \delta(\gamma - 1)a - \alpha - \beta + \delta + 1; -\beta + \delta + 1, -\alpha + \delta + 1, 2 - \gamma, \delta; x], \quad (\text{A.6})$$

$$T_7 H(x) = \left[1 - \frac{x}{a}\right]^{1-\epsilon} (1 - x)^{1-\delta} \times \\ H[a, q - \gamma[(\delta - 1)a + \alpha + \beta - \gamma - \delta]; -\beta + \gamma + 1, -\alpha + \gamma + 1, \gamma, 2 - \delta; x], \quad (\text{A.7})$$

$$T_8 H(x) = \left[1 - \frac{x}{a}\right]^{1-\epsilon} x^{1-\gamma} (1 - x)^{1-\delta} \times \\ H[a, q - (\gamma + \delta - 2)a - \alpha - \beta + \delta + 1; 2 - \alpha, 2 - \beta, 2 - \gamma, 2 - \delta; x]. \quad (\text{A.8})$$

B Cases solved by hypergeometric functions

In the four groups of solutions there are cases where the recurrence relations for the series coefficients present only two terms. For such cases, by combining results given in Ref. [8] with transformations of Maier's table [5], we find that the Heun equation (2) is reduced to the hypergeometric equation (3).

In (55) we have seen that the powers expansions (43-50) present two-term recurrence relations in the following three cases:

$$a = 0; \quad a = -1, \quad \epsilon = \delta, \quad q = 0 \quad \text{and} \quad a = -1, \quad \alpha + \beta = 1 + \gamma, \quad q = \gamma(1 - \delta).$$

For $a = 0$ the Heun equation equation reads

$$x(x - 1) \frac{d^2 H}{dx^2} + [(\alpha + \beta + 1)x - (\gamma + \epsilon)] \frac{dH}{dx} + \left[\alpha\beta - \frac{q}{x}\right] H = 0, \quad (a = 0) \quad (\text{B.1})$$

and the substitution

$$H(x) = x^{k_1} F(x), \quad k_1 = \frac{1}{2} \left[1 - \gamma - \epsilon - \sqrt{(1 - \gamma - \epsilon)^2 - 4q} \right]$$

gives the hypergeometric equation

$$x(1-x) \frac{d^2 F}{dx^2} + [(2k_1 + \gamma + \epsilon) - (2k_1 + \alpha + \beta + 1)x] \frac{dF}{dx} - (k_1 + \alpha)(k_1 + \beta)F = 0.$$

For the second case ($a = -1$, $\epsilon = \delta$, $q = 0$), the Heun equation (2) reads

$$\frac{d^2 H(x)}{dx^2} + \left[\frac{\gamma}{x} + \frac{\alpha + \beta + 1 - \gamma}{2(x-1)} + \frac{\alpha + \beta + 1 - \gamma}{2(x+1)} \right] \frac{dH(x)}{dx} + \frac{\alpha\beta}{(x-1)(x+1)} H(x) = 0,$$

since $\epsilon = \delta = (\alpha + \beta + 1 - \delta)/2$. It is already known that the substitutions [8]

$$z = x^2, \quad u(z) = H(x), \quad [a = -1, \epsilon = \delta, q = 0] \quad (\text{B.2})$$

reduce the above equation to

$$z(1-z) \frac{d^2 u(z)}{dz^2} + \left[\frac{1}{2}(\gamma + 1) - \frac{1}{2}(\alpha + \beta + 2)z \right] \frac{du(z)}{dz} - \frac{\alpha\beta}{4} u(z) = 0, \quad (\text{B.3})$$

which is the hypergeometric equation (3) with $a = \alpha/2$, $b = \beta/2$ and $c = (1 + \gamma)/2$. For the third case, first we transform the Heun equation (2) by using the homotopic substitution T_3 , that is, we take

$$H(x) = (1-x)^{1-\delta} h(x), \quad [a = -1, \gamma = \alpha + \beta - 1, q = \gamma(1 - \delta)].$$

Then, the Heun equation leads to

$$\frac{d^2 h(x)}{dx^2} + \left[\frac{\alpha + \beta - 1}{x} + \frac{2 - \delta}{x-1} + \frac{2 - \delta}{x+1} \right] \frac{dh(x)}{dx} + \frac{(\alpha - \delta + 1)(\beta - \delta + 1)}{(x-1)(x+1)} h(x) = 0.$$

Now, by the substitutions [8]

$$z = x^2, \quad u(z) = h(x) \Rightarrow H(x) = (1-x)^{1-\delta} u(z), \quad (\text{B.4})$$

$$[a = -1, \gamma = \alpha + \beta - 1, q = \gamma - \gamma\delta]$$

the above equation gives the hypergeometric equation

$$z(1-z) \frac{d^2 u(z)}{dz^2} + \left[\frac{1}{2}(\alpha + \beta) - \frac{1}{2}(\alpha + \beta + 4 - 2\delta)z \right] \frac{du(z)}{dz} - \frac{(\alpha - \delta + 1)(\beta - \delta + 1)}{4} u(z) = 0, \quad (\text{B.5})$$

with parameters $a = (\alpha - \delta + 1)/2$, $b = (\beta - \delta + 1)/2$ and $c = (\alpha + \beta)/2$.

According to (70) and (90) the expansions $\mathbf{H}^{(i)}(x)$ and $H^{(i)}(x)$ have two-term recurrence relations if

$$a = 1; \quad a = 2, \quad \alpha + \beta + 1 = 2\gamma + \delta, \quad q = \alpha\beta \quad \text{and}$$

$$a = 2, \quad \alpha + \beta = 1 + \delta, \quad q = \alpha\beta + (\gamma - 1)\delta.$$

For $a = 1$ we have

$$x(x-1) \frac{d^2 H}{dx^2} + [(\alpha + \beta + 1)x - \gamma] \frac{dH}{dx} + \frac{\alpha\beta x - q}{x-1} H = 0, \quad (a = 1) \quad (\text{B.6})$$

and the change of variable

$$H(x) = (x-1)^{k_2} G(x), \quad k_2 = \frac{1}{2} \left[\gamma - \alpha - \beta - \sqrt{(\gamma - \alpha - \beta)^2 - 4\alpha\beta + 4q} \right] \quad (\text{B.7})$$

yields an hypergeometric equation, namely,

$$x(1-x) \frac{d^2 G}{dx^2} + [\gamma - (2k_2 + \alpha + \beta + 1)x] \frac{dG}{dx} - (k_2 + \alpha)(k_2 + \beta)G = 0. \quad (\text{B.8})$$

For the second case the Heun equation (2) becomes

$$\frac{d^2 H(x)}{dx^2} + \left[\frac{\gamma}{x} + \frac{\alpha + \beta + 1 - 2\gamma}{x-1} + \frac{\gamma}{x-2} \right] \frac{dH(x)}{dx} + \frac{\alpha\beta}{x(x-2)} H(x) = 0, \\ [a = 2, \alpha + \beta = 2\gamma + \delta, q = \alpha\beta].$$

It is known that the substitutions [8]

$$z = x(2-x), \quad u(z) = H(x), \quad [a = 2, \alpha + \beta = 2\gamma + \delta, q = \alpha\beta] \quad (\text{B.9})$$

transform the above equation into the hypergeometric equation (3)

$$z(1-z) \frac{d^2 u(z)}{dz^2} + \left[\gamma - \frac{1}{2}(\alpha + \beta + 2)z \right] \frac{du(z)}{dz} - \frac{\alpha\beta}{4} u(z) = 0, \quad (\text{B.10})$$

with $a = \alpha/2$, $b = \beta/2$ and $c = \gamma$. For the third case, first we transform the Heun equation (2) by using the homotopic substitution T_2 , that is, we take

$$H(x) = x^{1-\gamma} h(x), \quad [a = 2, \alpha + \beta = 1 + \delta, q = \alpha\beta + (\gamma - 1)\delta].$$

Then, the Heun equation leads to

$$\frac{d^2 h(x)}{dx^2} + \left[\frac{2-\gamma}{x} + \frac{\delta}{x-1} + \frac{2-\gamma}{x-2} \right] \frac{dh(x)}{dx} + \frac{(\alpha-\gamma+1)(\beta-\gamma+1)}{x(x-2)} h(x) = 0.$$

By the substitutions [8]

$$z = x(2-x), \quad u(z) = h(x) \Rightarrow H(x) = x^{1-\gamma} u(z), \\ [a = 2, \alpha + \beta = 1 + \delta, q = \alpha\beta + (\gamma - 1)\delta] \quad (\text{B.11})$$

we get the hypergeometric equation

$$z(1-z) \frac{d^2 u(z)}{dz^2} + \left[2 - \gamma - \frac{1}{2}(\alpha + \beta + 2 - 2\gamma)z \right] \frac{du(z)}{dz} - \frac{(\alpha-\gamma+1)(\beta-\gamma+1)}{4} u(z) = 0, \quad (\text{B.12})$$

with $a = (\alpha - \gamma + 1)/2$, $b = (\beta - \gamma + 1)/2$ and $c = 2 - \gamma$.

Finally, by (100) the expansions $\mathbf{H}^{(i)}(x)$ present two-term recurrence relations if

$$a = 2, \quad \beta = \alpha + 1 - \delta, \quad q = \alpha\gamma \quad \text{and} \quad a = 2, \quad \alpha = \beta + 1 - \delta, \quad q = \beta\gamma.$$

For the first case, in Eq. (2) we use the fractional substitution

$$H(x) = \left(1 - \frac{x}{a}\right)^{-\alpha} f(y), \quad y = \frac{(1-a)x}{x-a}, \quad [a = 2, \beta = \alpha + 1 - \delta, q = \alpha\gamma]$$

which is Maier the transformation M_{17} . Thence

$$\frac{d^2 f(y)}{dy^2} + \left[\frac{\gamma}{y} + \frac{\delta}{y-1} + \frac{\delta}{y+1} \right] \frac{df(y)}{dy} - \frac{\alpha(1 + \alpha - \gamma - 2\delta)}{(y-1)(y+1)} f(y) = 0.$$

The changes

$$z = y^2 = \left(\frac{x}{2-x}\right)^2, \quad u(z) = f(y) \Rightarrow H(x) = \left(1 - \frac{x}{2}\right)^{-\alpha} u(z), \quad (B.13)$$

$$[a = 2, \beta = \alpha + 1 - \delta, q = \alpha\gamma]$$

give the hypergeometric equation

$$z(1-z)\frac{d^2u(z)}{dz^2} + \left[\frac{1+\gamma}{2} - \frac{1+\gamma+2\delta}{2}z\right]\frac{du(z)}{dz} + \frac{\alpha(1+\alpha-\gamma-2\delta)}{4}u(z) = 0, \quad (B.14)$$

whose parameters are $a = \alpha/2$, $b = (\gamma + 2\delta - \alpha - 1)/2$ and $c = (1 + \gamma)/2$. The second case differs from the first one by the interchange $\alpha \longleftrightarrow \beta$. Thus, the changes

$$z = y^2 = \left(\frac{x}{2-x}\right)^2, \quad u(z) = f(y) \Rightarrow H(x) = \left(1 - \frac{x}{2}\right)^{-\beta} u(z), \quad (B.15)$$

$$[a = 2, \alpha = \beta + 1 - \delta, q = \beta\gamma]$$

give the hypergeometric equation

$$z(1-z)\frac{d^2u(z)}{dz^2} + \left[\frac{1+\gamma}{2} - \frac{1+\gamma+2\delta}{2}z\right]\frac{du(z)}{dz} + \frac{\beta(1+\beta-\gamma-2\delta)}{4}u(z) = 0, \quad (B.16)$$

whose parameters are $a = \beta/2$, $b = (\gamma + 2\delta - \beta - 1)/2$ and $c = (1 + \gamma)/2$.

C Some degenerate power-series solutions

We consider three cases of the Schrödinger equation (101) whose solutions are obtained as particular cases of the power-series solutions given in Section 5. For each case, we will find

- degenerate finite-series eigenfunctions which satisfy the Arscott criterion;
- four infinite-series eigenfunctions.

The cases are: (1) $\mathbf{l} = 1/2$ and $\mathbf{m} = 3/2$, (2) $\mathbf{l} = 1/2$ and $\mathbf{m} = 5/2$, (3) $\mathbf{l} = 1/2$ and $\mathbf{m} = 7/2$. For these cases the finite series have at most three terms whose coefficients are computed from the recurrence relations (32) by means of

$$b_1 = -b_0 \frac{\beta_0}{\alpha_0}; \quad b_2 = b_0 \frac{1}{\alpha_1} \left(\frac{\beta_0\beta_1}{\alpha_0} - \gamma_1 \right), \quad (C.1)$$

where b_0 is the coefficient of the first term. The characteristic equations (35) reduces to

$$\beta_0 = 0, \quad \beta_0 \beta_1 - \alpha_0 \gamma_1 = 0, \quad \beta_0 \beta_1 \beta_2 - \alpha_1 \beta_0 \gamma_2 - \alpha_0 \beta_2 \gamma_1 = 0, \quad (C.2)$$

for series with one, two and three terms respectively.

Khare and Sukhatme [19, 20] have found solutions for these cases by integrating the Schrödinger equation directly. Here, the solutions are obtained from the expansions in series of $\text{cn}^2 u$ ($\Psi^{(i)}$ and $\tilde{\Psi}^{(i)}$ written in Section 5.2); these satisfy the Arscott condition for series with two or more terms, in opposition to $\psi^{(i)}$ and $\tilde{\psi}^{(i)}$.

First case: $l = 1/2$ and $m = 3/2$

For this case, $m - l = 1$ and $m + l = 2$. Thence, (136a-143a) imply the

$$\text{finite-series expansions : } \mathring{\Psi}^{(1)} \text{ (1 term), } \tilde{\Psi}^{(5)} \text{ (2 terms), } \tilde{\Psi}^{(8)} \text{ (1 term),} \quad (\text{C.3})$$

while (179-182) give four

$$\text{infinite-series expansions : } \mathring{\Phi}^{(2)} \stackrel{(56)}{=} \mathring{\Phi}^{(6)}, \quad \mathring{\Phi}^{(3)} \stackrel{(56)}{=} \mathring{\Phi}^{(7)}, \quad \mathring{\Phi}^{(4)}, \quad \mathring{\Phi}^{(5)}. \quad (\text{C.4})$$

The one-term expansions and their respective energies read

$$\mathring{\Psi}^{(1)}(u) \stackrel{(136a)}{=} b_0^{(1)} \text{dn}^{3/2}u, \quad \mathcal{E} = \frac{9}{4}k^2; \quad (\text{C.5})$$

$$\tilde{\Psi}^{(8)}(u) \stackrel{(?)}{=} b_0^{(8)} \frac{\text{sn}u \text{cn}u}{\text{dn}^{1/2}u}, \quad \mathcal{E} = 4 + \frac{1}{4}k^2. \quad (\text{C.6})$$

The two-term expansion $\mathring{\Psi}^{(5)}(u)$ represents two solutions, but one is equivalent to the previous $\mathring{\Psi}^{(1)}(u)$, that is,

$$\begin{aligned} \tilde{\Psi}_1^{(5)}(u) &\stackrel{(137a)}{=} \frac{C_1}{\text{dn}^{1/2}u} \left(1 + \frac{k^2}{1-k^2} \text{cn}^2u \right) = \frac{C_1}{(1-k^2)\text{dn}^{1/2}u} (1 - k^2 \text{sn}^2u) \stackrel{(8)}{=} \mathring{\Psi}^{(1)}(u), \\ \tilde{\Psi}_2^{(5)}(u) &\stackrel{(137a)}{=} \frac{C_2}{\text{dn}^{1/2}u} (1 - 2\text{cn}^2u), \quad \mathcal{E} = 4 + \frac{1}{4}k^2. \end{aligned} \quad (\text{C.7})$$

Therefore, there are three independent eigensolutions. $\tilde{\Psi}^{(8)}(u)$ and $\tilde{\Psi}_2^{(5)}(u)$ are degenerate because have the same energy.

Second case: $l = 1/2$ and $m = 5/2$

Now $m - l = 2$ and $m + l = 3$. Then, proceeding as in the above case, we obtain the

$$\text{finite series : } \mathring{\Psi}^{(2)} \text{ and } \mathring{\Psi}^{(3)} \text{ (1 term), } \tilde{\Psi}^{(6)} \text{ and } \tilde{\Psi}^{(7)} \text{ (2 terms);} \quad (\text{C.8})$$

$$\text{infinite series : } \mathring{\Phi}^{(1)}, \quad \mathring{\Phi}^{(4)}, \quad \mathring{\Phi}^{(5)}, \quad \mathring{\Phi}^{(8)}. \quad (\text{C.9})$$

The one-term finite series give two eigenfunctions,

$$\mathring{\Psi}^{(2)}(u) \stackrel{(138a)}{=} b_0^{(2)} \text{sn}u \text{dn}^{3/2}u, \quad \mathcal{E} = 1 + \frac{25}{4}k^2; \quad (\text{C.10})$$

$$\mathring{\Psi}^{(3)}(u) \stackrel{(140a)}{=} b_0^{(3)} \text{cn}u \text{dn}^{3/2}u, \quad \mathcal{E} = 1 + \frac{9}{4}k^2. \quad (\text{C.11})$$

The two-term series $\tilde{\Psi}^{(6)}(u)$ represents two eigenfunctions but one of them repeats the above solution $\mathring{\Psi}^{(2)}(u)$,

$$\begin{aligned} \tilde{\Psi}_1^{(6)}(u) &\stackrel{(139a)}{=} C_1 \frac{\text{sn}u}{\text{dn}^{1/2}u} \left(1 + \frac{k^2}{1-k^2} \text{cn}^2u \right) = \mathring{\Psi}^{(2)}(u), \\ \tilde{\Psi}_2^{(6)}(u) &\stackrel{(139a)}{=} C_2 \frac{\text{sn}u}{\text{dn}^{1/2}u} (1 - 4\text{cn}^2u), \quad \mathcal{E} = 9 + \frac{1}{4}k^2. \end{aligned} \quad (\text{C.12})$$

Similarly, the two-term series $\tilde{\Psi}^{(7)}(u)$ represent two eigenfunctions but one of them repeats the above solution $\mathring{\Psi}^{(3)}(u)$,

$$\begin{aligned} \tilde{\Psi}_1^{(7)}(u) &\stackrel{(141a)}{=} D_1 \frac{\text{cn}u}{\text{dn}^{1/2}u} \left(1 + \frac{k^2}{1-k^2} \text{cn}^2u \right) = \mathring{\Psi}^{(3)}(u), \\ \tilde{\Psi}_2^{(7)}(u) &\stackrel{(141a)}{=} D_2 \frac{\text{cn}u}{\text{dn}^{1/2}u} \left(1 - \frac{4}{3} \text{cn}^2u \right), \quad \mathcal{E} = 9 + \frac{1}{4}k^2 \text{ (degenerate ver acima)}. \end{aligned} \quad (\text{C.13})$$

Therefore, in this case there are only four independent solutions, being $\tilde{\Psi}_2^{(6)}(u)$ and $\tilde{\Psi}_2^{(7)}(u)$ degenerate solutions.

Third case: $l = 1/2$ and $m = 7/2$

In this case $m - l = 3$ and $m + l = 4$. Thence we get the the following finite and infinite series:

$$\text{finite : } \mathring{\Psi}^{(1)} \text{ (2 terms), } \mathring{\Psi}^{(4)} \text{ (1 term), } \tilde{\Psi}^{(5)} \text{ (3 terms), } \tilde{\Psi}^{(8)} \text{ (2 terms);} \quad (\text{C.14})$$

$$\text{infinite : } \mathring{\Phi}^{(2)}, \quad \mathring{\Phi}^{(3)}, \quad \mathring{\Phi}^{(6)}, \quad \mathring{\Phi}^{(7)}. \quad (\text{C.15})$$

The one-term series reads

$$\mathring{\Psi}^{(4)}(u) \stackrel{(142a)}{=} \mathbf{b}_0^{(4)} \operatorname{sn} u \operatorname{cn} u \operatorname{dn}^{3/2} u, \quad \mathcal{E} = 4 + \frac{25}{4} k^2. \quad (\text{C.16})$$

The two-term expansions $\mathring{\Psi}^{(1)}(u)$ represent two eigenfunctions $\mathring{\Psi}_{\pm}^{(1)}(u)$ corresponding to the energies \mathcal{E}_{\pm} ,

$$\begin{aligned} \mathring{\Psi}_{\pm}^{(1)}(u) &\stackrel{(136a)}{=} C_{\pm} \operatorname{dn}^{3/2} u \left[1 + \frac{7k^2 - 2 \pm \sqrt{4 + 25k^4 - 4k^2}}{2(1 - k^2)} \operatorname{cn}^2 u \right], \\ \mathcal{E}_{\pm} &= 2 + \frac{29}{4} k^2 \mp \sqrt{4 + 25k^4 - 4k^2}. \end{aligned} \quad (\text{C.17})$$

The expansion $\mathring{\Psi}^{(8)}(u)$ also represents two eigenfunctions but one of these is identical to $\mathring{\Psi}^{(4)}(u)$. We find

$$\begin{aligned} \tilde{\Psi}_1^{(8)}(u) &\stackrel{(143a)}{=} D_1 \frac{\operatorname{sn} u \operatorname{cn} u}{\operatorname{dn}^{1/2} u} \left[1 + \frac{k^2}{1 - k^2} \operatorname{cn}^2 u \right] = \mathring{\Psi}^{(4)}(u), \\ \tilde{\Psi}_2^{(8)}(u) &\stackrel{(143a)}{=} D_2 \frac{\operatorname{sn} u \operatorname{cn} u}{\operatorname{dn}^{1/2} u} [1 - 2 \operatorname{cn}^2 u], \quad \mathcal{E} = 16 + \frac{1}{4} k^2. \end{aligned} \quad (\text{C.18})$$

On the other hand, as $\mathring{\Psi}^{(4)}(u)$ is a three-term expansion, it represents three eigensolutions having the form

$$\begin{aligned} \tilde{\Psi}^{(5)}(u) &\stackrel{(137a)}{=} \frac{\tilde{\mathbf{b}}_0^{(5)}}{\operatorname{dn}^{1/2} u} \left\{ 1 - \frac{2k^2}{k^2 - 1} \tilde{\beta}_0^{(5)} \operatorname{cn}^2 u + \right. \\ &\quad \left. \frac{k^2}{3(k^2 - 1)} \left[4 + \frac{2k^2}{(k^2 - 1)} \tilde{\beta}_0^{(5)} \left(\tilde{\beta}_0^{(5)} + \frac{1}{k^2} - \frac{3}{2} \right) \right] \operatorname{cn}^4 u \right\}, \end{aligned} \quad (\text{C.19})$$

where $\tilde{\beta}_0^{(5)}$ is determined from the characteristic equation

$$\begin{aligned} \left[\tilde{\beta}_0^{(5)} \right]^3 + \left(\frac{5}{k^2} - \frac{17}{2} \right) \left[\tilde{\beta}_0^{(5)} \right]^2 + \left(\frac{43k^4 - 48k^2 + 8}{2k^4} \right) \tilde{\beta}_0^{(5)} + 2 \left(1 - \frac{1}{k^2} \right) \left(\frac{4}{k^2} - 7 \right) &= 0, \\ \tilde{\beta}_0^{(5)} &= \frac{65}{16} - \frac{\mathcal{E}}{4k^2}. \end{aligned} \quad (\text{C.20})$$

The Arscott condition implies three real and distinct values for $\tilde{\beta}_0^{(5)}$. We find that, if $\mathcal{E} = 16 + k^2/4$ then $\tilde{\beta}_0^{(5)} = 4(k^2 - 1)/k^2$ and Eq. (C.20) is fulfilled. The eigenfunction takes the form

$$\tilde{\Psi}_1^{(5)}(u) = \frac{\bar{D}_1}{\operatorname{dn}^{1/2} u} (1 - 8 \operatorname{sn}^2 u \operatorname{cn}^2 u), \quad \mathcal{E} = 16 + \frac{k^2}{4} \quad (\text{C.21})$$

and in conjunction with $\tilde{\Psi}_2^{(8)}(u)$ constitute a pair of degenerate eigenfunctions as stated in [19]. In fact, the solution mentioned in [19] comes from expansions in series of $\operatorname{sn}^2 u$, namely,

$$\begin{aligned} \tilde{\psi}^{(5)}(u) &\stackrel{(116a)}{=} \frac{\tilde{b}_0^{(5)}}{\operatorname{dn}^{1/2} u} \left\{ 1 - \frac{1}{2} \left(\mathcal{E} - \frac{k^2}{4} \right) \operatorname{sn}^2 u + \right. \\ &\quad \left. \frac{k^2}{3} \left[4 + \frac{1}{8k^2} \left(\mathcal{E} - \frac{k^2}{4} \right) \left(\mathcal{E} - 4 - \frac{9}{4} k^2 \right) \right] \operatorname{sn}^4 u \right\}, \end{aligned}$$

which satisfy the characteristic equation

$$\lambda^3 - 4 \left(\frac{7}{2}k^2 + 5 \right) \lambda^2 + 16 \left(\frac{3}{2}k^4 + 16k^2 + 4 \right) \lambda - 128k^2 (3k^2 + 4) = 0, \quad \lambda = \mathcal{E} - \frac{k^2}{4}.$$

These solutions do not obey the Arscott condition, but we can see that $\tilde{\psi}^{(5)}(u) = \tilde{\Psi}^{(5)}(u)$ if $\mathcal{E} = 16 + k^2/4$.

D Svartholm's and Erdélyi's solutions

Svartholm [13] and Erdélyi [14, 15] have supposed that the coefficients of their expansions in series of hypergeometric functions satisfy recurrence relations having the form given in (32). Such conjecture has been repeated in subsequent literature [3, 9, 21, 23], including Erdélyi himself [23]. However, there are three forms for the recurrence relations, namely,

$$\alpha_0 b_1 + \beta_0 b_0 = 0, \quad \alpha_n b_{n+1} + \beta_n b_n + \gamma_n b_{n-1} = 0, \quad (n \geq 1), \quad (\text{D.1})$$

$$\begin{cases} \alpha_0 b_1 + \beta_0 b_0 = 0, & \alpha_1 b_2 + \beta_1 b_1 + [\gamma_1 + \alpha_{-1}] b_0 = 0, \\ \alpha_n b_{n+1} + \beta_n b_n + \gamma_n b_{n-1} = 0, & (n \geq 2), \end{cases} \quad (\text{D.2})$$

$$\alpha_0 b_1 + [\beta_0 + \alpha_{-1}] b_0 = 0, \quad \alpha_n b_{n+1} + \beta_n b_n + \gamma_n b_{n-1} = 0, \quad (n \geq 1) \quad (\text{D.3})$$

which reduce to (32) only if $\alpha_{-1} = 0$.

The above recurrences also occur for some expansions in series of hypergeometric functions and in series of Coulomb wave functions for solutions of confluent and double-confluent Heun equations [46]. In fact, the procedure used in Appendix A of reference [46] can be easily adapted for the expansions of Svartholm and Erdélyi.

As an example we consider the Erdélyi solutions [15]

$$H(x) = \sum_{n=0}^{\infty} b_n P_n(x) = \sum_{n=0}^{\infty} b_n F(n + \lambda, -n + \mu; \gamma; x), \quad (\text{D.4})$$

$[\gamma \neq 0, -1, -2, \dots, \quad \mu - \lambda \neq 1, 2, 3, \dots]$

where λ and μ depend on the parameters of the Heun equation and are connected by $\lambda + \mu = \gamma + \delta - 1$. The condition $\mu - \lambda \neq 1, 2, 3, \dots$ avoids two identical hypergeometric functions for different values of n . For $H(x)$ given in (D.4), Eq. (2) leads to

$$\sum_{n=0}^{\infty} b_n [\alpha_{n-1} P_{n-1}(x) + \beta_n P_n(x) + \gamma_{n+1} P_{n+1}(x)] = 0, \quad (\text{D.5})$$

which can be written in Erdélyi's notation [15] by taking $x = z$, $b_n = c_n$, $\alpha_n = M_n$, $\beta_n = L_n$ and $\gamma_n = K_n$: for example [3]

$$\alpha_n = M_n = -\frac{(n-\alpha+\lambda+1)(n-\beta+\lambda+1)(n-\gamma+\lambda+1)(n-\mu+1)}{(2n+\lambda-\mu+1)(2n+\lambda-\mu+2)}.$$

In order to get the three forms for the recurrence relations, we rewrite Eq. (D.5) as

$$\begin{aligned} & \alpha_{-1} b_0 P_{-1}(x) + [\alpha_0 b_1 + \beta_0 b_0] P_0(x) + [\alpha_1 b_2 + \beta_1 b_1 + \gamma_1 b_0] P_1(x) \\ & + \sum_{n=2}^{\infty} [\alpha_n b_{n+1} + \beta_n b_n + \gamma_n b_{n-1}] P_n(x) = 0. \end{aligned} \quad (\text{D.6})$$

After this, we specify λ and μ and equate to zero the coefficients of each independent function $P_n(x)$. Next we regard only two choices for (λ, μ) .

In the first place, for the expansion mentioned at the end of Section 6 we take $\lambda = \alpha$ and $\mu = \gamma + \delta - 1 - \alpha$. In this manner we obtain the expansion $H^{(1)}(x)$,

$$H^{(1)}(x) = \sum_{n=0}^{\infty} b_n^{(1)} F(n + \alpha, -n - \alpha - 1 + \gamma + \delta; \gamma; x), \quad (D.7)$$

$[\gamma \text{ and } 2 + 2\alpha - \gamma - \delta \neq 0, -1, -2, \dots],$

for which the recurrence relations for $b_n^{(1)}$ are given by

$$\begin{aligned} \text{Eqs. (D.1) if } 2\alpha - \gamma - \delta \neq 0, -1; & \quad \text{Eqs. (D.2) if } 2\alpha - \gamma - \delta = -1; \\ \text{Eqs. (D.3) if } 2\alpha - \gamma - \delta = 0, & \end{aligned} \quad (D.8)$$

with α_n, β_n and γ_n defined as [15]

$$\left\{ \begin{array}{ll} \alpha_n^{(1)} = -\frac{(n+1)(n+1+\alpha-\beta)(n+1+\alpha-\gamma)(n+2+\alpha-\gamma-\delta)}{(2n+2+2\alpha-\gamma-\delta)(2n+3+2\alpha-\gamma-\delta)}, & [n \geq -1] \\ \beta_n^{(1)} = \left(\frac{1}{2} - a\right) \left[n(n+2\alpha-\gamma-\delta+1) + \alpha(\alpha+1-\gamma-\delta) \right] - q + \frac{1}{2}\alpha\beta + \\ \quad \frac{1}{8}(\gamma-\delta) \left[2\alpha+2\beta-\gamma-\delta + \frac{(\gamma+\delta-2)(2\alpha-\gamma-\delta)(2\beta-\gamma-\delta)}{(2n+2\alpha-\gamma-\delta)(2n+2+2\alpha-\gamma-\delta)} \right], & [n \geq 0] \\ \gamma_n^{(1)} = -\frac{(n+2\alpha-\gamma-\delta)(n+\alpha-\delta)(n+\alpha-1)(n+\alpha+\beta-\gamma-\delta)}{(2n+2\alpha-\gamma-\delta)(2n+2\alpha-\gamma-\delta-1)}, & [n \geq 1] \end{array} \right. \quad (D.9)$$

As to the recurrence relations which are missing in Erdélyi, they result because there are terms linearly dependent in Eq. (D.6), namely,

$$\begin{aligned} \text{if } 2\alpha - \gamma - \delta = -1, & \quad P_{-1}(x) = P_1(x) = F(\alpha-1, \alpha+1; \gamma; x) \Rightarrow \text{Eqs. (D.2);} \\ \text{if } 2\alpha - \gamma - \delta = 0, & \quad P_{-1}(x) = P_0(x) = F(\alpha-1; \alpha; \gamma; x) \Rightarrow \text{Eqs. (D.3).} \end{aligned}$$

Relations (D.8) take the place of (5.2), (5.3) and (5.4) given in Erdélyi's paper [15].

In the second place, the Svartholm solution arises when we put $\lambda = \gamma + \delta - 1$ and $\mu = 0$ in (D.4). Then the Erdélyi expansion (D.4) becomes

$$\tilde{H}^{(1)}(x) = \sum_{n=0}^{\infty} \tilde{b}_n^{(1)} F(n + \gamma + \delta - 1, -n; \gamma; x), \quad (D.10)$$

$[\gamma \neq 0, -1, -2, \dots; \quad \gamma + \delta \neq 0, -1, -2, \dots]$

where the coefficients $\tilde{b}_n^{(1)}$ satisfy the recursions

$$\begin{aligned} \text{Eqs. (D.1) if } \gamma + \delta \neq 1, 2; & \quad \text{Eqs. (D.2) if } \gamma + \delta = 1; \\ \text{Eqs. (D.3) if } \gamma + \delta = 2, & \end{aligned} \quad (D.11)$$

with

$$\left\{ \begin{array}{ll} \tilde{\alpha}_n^{(1)} = -\frac{(n+1)(n+\gamma+\delta-\alpha)(n+\gamma+\delta-\beta)(n+\delta)}{(2n+\gamma+\delta)(2n+\gamma+\delta+1)}, & [n \geq -1] \\ \tilde{\beta}_n^{(1)} = \left[\frac{1}{2} - a\right] n[n + \gamma + \delta - 1] - q + \frac{1}{2}\alpha\beta + \\ \quad \frac{1}{8}(\gamma-\delta) \left[2\alpha+2\beta-\gamma-\delta + \frac{(\gamma+\delta-2)(2\alpha-\gamma-\delta)(2\beta-\gamma-\delta)}{(2n+\gamma+\delta-2)(2n+\gamma+\delta)} \right], & [n \geq 0] \\ \tilde{\gamma}_n^{(1)} = -\frac{(n+\alpha-1)(n+\beta-1)(n+\gamma+\delta-2)(n+\gamma-1)}{(2n+\gamma+\delta-3)(2n+\gamma+\delta-2)}, & [n \geq 1] \end{array} \right. \quad (D.12)$$

By the homotopic transformations we can generate eight Svartholm-type solutions, $\tilde{H}^{(i)}(x)$. These give series of trigonometric functions for the associated Lamé equation. For this, we consider the definitions of the basic elliptic functions [16]:

$$\operatorname{sn} u = \sin v, \quad \operatorname{cn} u = \cos v, \quad \operatorname{dn} u = \sqrt{1 - k^2 \sin^2 v} \text{ for } u = \int_0^v \frac{d\theta}{\sqrt{1 - k^2 \sin^2 \theta}}. \quad (\text{D.13})$$

In addition, for the associated Lamé equation ($x = \operatorname{sn}^2 u = \sin^2 v$), the relations [3]

$$\begin{aligned} F\left(-a, a; \frac{1}{2}; \sin^2 v\right) &= \cos(2av), & F\left(a, 1-a; \frac{3}{2}; \sin^2 v\right) &= \frac{\cos[(2a-1)v]}{\cos v}, \\ F\left(1-a, a; \frac{3}{2}; \sin^2 v\right) &= \frac{\sin[(2a-1)v]}{(2a-1)\sin v}, & F\left(a, 2-a; \frac{3}{2}; \sin^2 v\right) &= \frac{\sin[(2a-2)v]}{(a-1)\sin(2v)}, \end{aligned} \quad (\text{D.14})$$

allow to get the trigonometric series. For example, for $\gamma = \delta = 1/2$ we find

$$\tilde{H}^{(1)}(x) = \sum_{n=0}^{\infty} \tilde{b}_n^{(1)} F\left(-n, n; \frac{1}{2}; x\right) = \sum_{n=0}^{\infty} \tilde{b}_n^{(1)} \cos(2nv), \quad x = \operatorname{sn}^2 u = \sin^2 v, \quad (\text{D.15})$$

where the coefficients $\tilde{b}_n^{(1)}$ satisfy the relations (D.2) with

$$\begin{cases} \tilde{\alpha}_n^{(1)} = -\frac{1}{4}(n+1-\alpha)(n+1-\beta), & [n \geq -1], \\ \tilde{\beta}_n^{(1)} = \left(\frac{1}{2} - a\right)n^2 - q + \frac{1}{2}\alpha\beta, & [n \geq 0] \\ \tilde{\gamma}_n^{(1)} = -\frac{1}{4}(n+\alpha-1)(n+\beta-1), & [n \geq 1] \end{cases} \quad (\text{D.16})$$

where $a = 1/k^2$. For the Lamé equation ($\gamma = \delta = \epsilon = 1/2$), the expansions in trigonometric functions are equivalent to the solutions found by Ince in 1940 [40].

Finally we find that, by performing the limits (60) for the confluent Heun equation (59), the solutions (D.7)-(D.8) and (D.10)-(D.11) remain formally unaltered while the coefficients (D.5) and (D.12) give, respectively,

$$\begin{cases} \alpha_n^{(1)} = -\rho \frac{(n+1)(n+1+\alpha-\gamma)(n+2+\alpha-\gamma-\delta)}{(2n+2+2\alpha-\gamma-\delta)(2n+3+2\alpha-\gamma-\delta)}, & [n \geq -1], \\ \beta_n^{(1)} = -n(n+2\alpha-\gamma-\delta+1) - \alpha(\alpha+1-\gamma-\delta) + \sigma - \frac{1}{2}\alpha\rho + \\ \quad \frac{1}{4}\rho(\gamma-\delta) \left[\frac{(\gamma+\delta-2)(\gamma+\delta-2\alpha)}{(2n+2\alpha-\gamma-\delta)(2n+2+2\alpha-\gamma-\delta)} - 1 \right], & [n \geq 0], \\ \gamma_n^{(1)} = \rho \frac{(n+2\alpha-\gamma-\delta)(n+\alpha-\delta)(n+\alpha-1)}{(2n+2\alpha-\gamma-\delta)(2n+2\alpha-\gamma-\delta-1)}, & [n \geq 1]. \end{cases} \quad (\text{D.17})$$

$$\begin{cases} \tilde{\alpha}_n^{(1)} = -\rho \frac{(n+1)(n+\gamma+\delta-\alpha)(n+\delta)}{(2n+\gamma+\delta)(2n+\gamma+\delta+1)}, & [n \geq -1] \\ \tilde{\beta}_n^{(1)} = -n[n+\gamma+\delta-1] + \sigma - \frac{1}{2}\alpha\rho - \\ \quad \frac{1}{4}\rho(\gamma-\delta) \left[1 + \frac{(\gamma+\delta-2)(2\alpha-\gamma-\delta)}{(2n+\gamma+\delta-2)(2n+\gamma+\delta)} \right], & [n \geq 0] \\ \tilde{\gamma}_n^{(1)} = \rho \frac{(n+\alpha-1)(n+\gamma+\delta-2)(n+\gamma-1)}{(2n+\gamma+\delta-3)(2n+\gamma+\delta-2)}, & [n \geq 1] \end{cases} \quad (\text{D.18})$$

Note that the three forms for the recurrence relations for the CHE have already been obtained [46] independently of the solutions for the Heun equation, as aforementioned.

References

- [1] Darboux, G.: Sur une équation linéaire. C. R. Acad. Sci. 94, 1645-1648 (1882).
- [2] Humbert, P.: Fonctions de Lamé et fonctions de Mathieu. Mémorial des Sciences Mathématique, Vol. X, Gauthier-Villards (1926).
- [3] Olver, F.W.J., Lozier, D.V., Boisvert, R.F., Clark C.W. (editors): NIST Handbook of Mathematical Functions. Cambridge University Press (2010).
- [4] Heun, K.: Zur Theorie der Riemannschen Functionen zweiter Ordnung mit vier Verzweigungspunkten. Math. Ann. 33, 161-179 (1899).
- [5] Maier, R.S.: The 192 solutions of the Heun equation. Mathematics of Computation 76, 811-843 (2007).
- [6] Erdélyi, A., Magnus, W., Oberhettinger, F., Tricomi, F.G.: Higher Transcendental Functions vol. 1. McGraw-Hill, New York (1953).
- [7] Maier, R.S.: On reducing the Heun equation to the hypergeometric equation. J. Diff. Equations 213, 171-203 (2005).
- [8] Vidunas, R., Filipuk, G.: Parametric transformations between the Heun and Gauss hypergeometric functions. Funkcialaj Ekvacioj 56, 271-321 (2013).
- [9] Kristensson, G.: Second Order Differential Equations: Special Functions and Their Classification. Springer (2010).
- [10] Turbiner, A.V.: Quasi-exactly-solvable problems and $sl(2)$ algebra. Commun. Math. Phys. 118, 467-474 (1988).
- [11] Ushveridge, A.G.: Quasi-exactly solvable models in quantum mechanics. Sov. J. Part. Nucl. 20, 504-528 (1989).
- [12] Kalnins, E.G, Miller, M., Pogosyan, G.S.: Exact and quasiexact solvability of second-order superintegrable quantum systems: I. Euclidian space preliminaries. J. Math. Phys. 47, 033502 (2006).
- [13] Svartholm, N.: Die Lösung der Funchs'schen Differentialgleichung zweiter Ordnung durch Hypergeometrische Polynome. Math. Ann. 116, 413-421 (1939).
- [14] Erdélyi, A.: The Fuchsian equation of second order with four singularities. Duke Math. J. 9, 48-58 (1942).
- [15] Erdélyi, A.: Certain expansions of solutions of the Heun equation. Q. J. Math. (Oxford) 15, 62-69 (1944).
- [16] Erdélyi, A., Magnus, W., Oberhettinger, F., Tricomi, F.G.: Higher Transcendental Functions vol. 2. McGraw-Hill, New York (1953).
- [17] Ganguly, A.: Associated Lamé equation, periodic potentials and $sl(2, \mathbb{R})$. Mod. Phys. Lett. A 15, 1923-1930 (2002).
- [18] Ganguly, A.: Associated Lamé and various other new classes of elliptic potentials from $sl(2, \mathbb{R})$ and related orthogonal polynomials. J. Math. Phys. 43, 1980-1999 (2002).

- [19] Khare, A., Sukhatme, U.: New solvable and quasiexactly solvable periodic potentials. *J. Math. Phys.* 40, 5473-5494 (1999).
- [20] Khare, A., Sukhatme, U.: Some exact results for mid-band and zero band-gap states of associated Lamé potentials. *J. Math. Phys.* 42, 5652-5664 (2001).
- [21] Ronveaux, A. (editor): Heun's differential equations. Oxford University Press, New York (1995).
- [22] El-Jaick, L.J., Figueiredo, B.D.B.: Transformations of Heun's equation and its integral relations. *J. Phys. A: Math. Theor.* 44, 075204 (2011).
- [23] Erdélyi, A., Magnus, W., Oberhettinger, F., Tricomi, F.G.: Higher Transcendental Functions vol. 3. McGraw-Hill, New York (1955).
- [24] Gautschi, W.: Computational aspects of three-term recurrence relations. *SIAM Rev.* 9, 24-82 (1967).
- [25] Arscott, F.M.: Periodic Differential Equations. Pergamon Press (1964).
- [26] Arscott, F.M.: Latent roots of tri-diagonal matrices. *Edinburgh Mathematical Notes* 44, 5-7 (1961).
- [27] Whittaker, E.T., Watson, G.N.: A Course of Modern Analysis. Cambridge University Press (1954).
- [28] Knopp, K.: Infinite Sequences and Series. Dover (1956).
- [29] Liu, J.W.: Analytical solutions to the generalized spheroidal wave equation and the Green's function of one-electron diatomic molecules. *J. Math. Phys.* 33, 4026-4036 (1992).
- [30] Hodge, D.B.: Eigenvalues and eigenfunctions of the spheroidal wave equation. *J. Math. Phys.* 11, 2308-2312 (1970).
- [31] Falloon, P.E., Abbott, P.C., Wang, J.B.: Theory and computation of spheroidal wavefunctions. *J. Phys. A: Math. Gen.* 36, 5477-5495 (2003).
- [32] Gradshteyn, I.S., Ryzhik, I. M.: Table of Integrals, Series, and Products. Elsevier (2014).
- [33] Ince, E.L.: The periodic Lamé functions. *Proc. Roy. Soc. Edinburgh* 60, 47-63 (1940).
- [34] Decarreau, A., Dumont-Lepage, M.G., Maroni, P., Robert, A., Ronveaux, A.: Formes canoniques des équations confluentes de l'équation de Heun. *Ann. Soc. Sci. Brux. T92(I-II)*, 53-78 (1978).
- [35] Baber, W. G., Hassé, H.R.: The two centre problem in wave mechanics. *Proc. Cambr. Philos. Soc.* 25, 564-581 (1935).
- [36] Fisher, E.: Some differential equations involving three-term recursion formulas. *Phil. Mag.* 24, 245-256 (1937).
- [37] Slater, J.C.: A soluble problem in energy bands. *Phys. Rev.* 87, 807-835 (1951).

- [38] Matveev, V.B., Smirnov, A.O.: On the link between the Sparre equation and Darboux-Treibich-Verdier equation. *Lett. Math. Phys.* 76, 283-295 (2006).
- [39] El-Jaick, L.J., Figueiredo, B.D.B.: A limit of the confluent Heun equation and the Schrödinger equation for an inverted potential and for an electric dipole. *J. Math. Phys.* 50, 123511 (2009).
- [40] Ince, E.L.: Further investigations into the periodic Lamé functions. *Proc. Roy. Soc. Edinburgh* 60, 83-99 (1940).
- [41] Magnus, W., Winkler, S.: *Hill's Equation*. Wiley, New York (1966).
- [42] Volkmer, H.: Coexistence of periodic solutions of Ince's equation. *Analysis* 23, 97-105 (2003).
- [43] Ishkhanyan, T.A., Shahverdyan, T.A., Ishkhanyan, A. M.: Expansions of the solutions of the general Heun equation governed by two-term recurrence relations for coefficients. *Adv. High Energy Phys.* 426367 (2018).
- [44] Chiang, Y.M., A.Ching, A., Tsan, C.Y.: Symmetries of the Darboux Equation. *Kumamoto J.Math.* 31, 15-48 (2018).
- [45] Karayer, H., Demirhan, D., Büyükkiliç, F.: A particular solution of Heun equation for Hulthen and Woods-Saxon potentials. *Ann. Phys.* 526, 527-532 (2014).
- [46] Figueiredo, B.D.B.: On some solutions to generalized spheroidal wave equations and applications. *J. Phys. A: Math. Gen.* 35, 2877 (2002).