

POSITIVE ARBOREALIZATION OF POLARIZED WEINSTEIN MANIFOLDS

DANIEL ÁLVAREZ-GAVELA, YAKOV ELIASHBERG, AND DAVID NADLER

ABSTRACT. Let X be a Weinstein manifold. We show that the existence of a global field of Lagrangian planes in T^*X is equivalent to the existence of a positive arboreal skeleton for the Weinstein homotopy class of X .

CONTENTS

1. Introduction	1
2. Wc-manifolds and cotangent buildings	15
3. Symplectic neighborhoods of arboreal Lagrangians	44
4. Positive cotangent buildings	59
5. Ridgification of Lagrangians	73
6. Arborealization of skeleta	82
References	96

1. INTRODUCTION

1.1. **Arborealization program.** The symplectic topology of the cotangent bundle T^*M of a smooth manifold is determined by the smooth topology of its Lagrangian zero-section M . A general *Weinstein symplectic manifold* X (see precise definitions below), which is the symplectic counterpart of a Stein complex manifold, can be viewed as a symplectic thickening of its *skeleton*, which is a singular isotropic subcomplex $\text{Skel } X \subset X$. By analogy with the special case $X = T^*M$, $\text{Skel } X = M$, one would like to view X as the “cotangent bundle” of $\text{Skel } X$, and characterize the smooth symplectic topology of X in terms of the differential topology of $\text{Skel } X$. However, in general the singularities of $\text{Skel } X$ are too complicated to be amenable to a differential topological treatment.

In the paper [N13], the third author introduced a class of Lagrangian singularities, called *arboreal*. For arboreal singularities, the smooth topology of the singularity determines the symplectic topology of its neighborhood. In particular, it is possible to calculate the local symplectic invariants of a neighborhood of an arboreal singularity in terms of the singularity.

Date: February 15, 2022.

DA was partially supported by NSF grant DMS-1638352 and the Simons Foundation.

YE was partially supported by NSF grant DMS-1807270.

DN was partially supported by NSF grant DMS-1802373.

This was shown in [N13] for microlocal sheaves; for Fukaya categories, one can apply Lefschetz fibration calculations [Se08] to the plumbing characterization of [Sh18]. Going further, if a Weinstein manifold W has a skeleton with arboreal singularities, then its global symplectic invariants can be effectively computed knowing the smooth topology of the skeleton, see [N16] for microlocal sheaves and [GPS17] for Fukaya categories.

The paper [N13] initiated a program to determine which Weinstein manifolds admit arboreal skeleta, or as we say, *which Weinstein manifolds could be arborealized*. As evidence that this program might achieve the arborealization of a large class of Weinstein manifolds, it was shown in [N15] that germs of Whitney stratified Lagrangians can always be deformed to arboreal Lagrangians in a non-characteristic fashion, i.e. without changing their microlocal invariants.

The question of whether a Weinstein manifold can be arborealized via a homotopy of its Weinstein structure is more subtle. In two dimensions, the story is classical: generic ribbon graphs provide arboreal skeleta. In four dimensions, Starkston proved that arboreal skeleta always exist [St18]. In this paper, we establish that any Weinstein manifold admitting a polarization, i.e. a Lagrangian plane field, or equivalently a reduction of structure group of the tangent bundle from $Sp(2n)$ to $GL(n)$, can be arborealized, see Theorem 1.5 below. Moreover, our proof yields skeleta with the more specific class of *positive* arboreal singularities. Conversely, Weinstein manifolds with positive arboreal skeleta admit polarizations. On the other hand it turns out not all Weinstein manifolds can be arborealized, see the discussion in Section 1.3 below. Perhaps this is not surprising: there are homotopical obstructions to defining many symplectic invariants, and so any combinatorial route to realizing them must also encounter these obstructions.

1.2. Main results. Let (W, λ) be a $2n$ -dimensional Liouville domain. We recall the definition: W is a compact $2n$ -manifold with boundary; λ is a 1-form, called the Liouville form, such that $\omega = d\lambda$ is a symplectic form; and the corresponding vector field $Z = \omega^{-1}(\lambda)$, called the Liouville vector field, is outward transverse to ∂W .

By definition, the skeleton of a Liouville domain (W, λ) is the attractor of the negative flow of the Liouville vector field:

$$\text{Skel}(W, \lambda) = \bigcap_{t>0} Z^{-t}(W).$$

While the $2n$ -dimensional Lebesgue measure of $\text{Skel}(W, \lambda)$ is equal to 0, in general $\text{Skel}(W, \lambda)$ can be quite large if no additional conditions are imposed.

A Weinstein domain is a Liouville domain (W, λ) which admits a Morse Lyapunov function $\phi : W \rightarrow \mathbb{R}$, i.e. the critical points of ϕ are non-degenerate and Z is gradient-like for ϕ . Sometimes it is convenient to relax the Morse condition to generalized Morse or Morse-Bott. We do not consider the Lyapunov function as part of the defining data of a Weinstein domain, we merely require its existence.

The skeleton of a Weinstein domain (W, λ) is known to be the union of stable manifolds

$$\text{Skel}(W, \lambda) = \bigcup_{\lambda_p=0} S_p, \quad S_p = \{x \in W \mid \lim_{t \rightarrow \infty} Z^t(x) = p\}.$$

Each stable manifold is λ -isotropic, hence ω -isotropic and so at most half-dimensional. A result of F. Laudenbach [L92] states that if Z is Morse-Smale, i.e. its stable and unstable manifolds intersect transversely, and if moreover Z is the Euclidean gradient near critical points with respect to the coordinates which give the Morse normal form, then $\text{Skel}(W, \lambda)$ can be Whitney stratified.

However, even if Z satisfies the above conditions, in general the singularities of $\text{Skel}(W, \lambda)$ are quite complicated, and their smooth topology does not determine the symplectic topology of their neighborhoods. For example, in the simplest case when W is obtained by attaching a handle to a ball along a Legendrian sphere, the skeleton has a unique conical singular point and the symplectic topology depends on the Legendrian isotopy class of the link, not its smooth isotopy class (which is always trivial for manifolds of dimension > 4). However, for the class of arboreal singularities described in Definition 1.1 below the situation is different.

First, we introduce some auxiliary notions. A closed subset of a symplectic or contact manifold is called *isotropic* if it is stratified by isotropic submanifolds. It is called *Lagrangian* or *Legendrian* if it is isotropic and purely of the maximal possible dimension. The germ at the origin of a locally simply-connected isotropic subset $L \subset T^*\mathbb{R}^n$ of the cotangent bundle with its standard Liouville structure $\lambda = pdq$ admits a unique lift to an isotropic germ at the origin $\widehat{L} \subset J^1\mathbb{R}^n = T^*\mathbb{R}^n \times \mathbb{R}$ of the 1-jet bundle. Given an isotropic subset $\Lambda \subset S^*\mathbb{R}^n$ of the cosphere bundle, its Liouville cone $C(\Lambda) \subset T^*\mathbb{R}$, i.e. the closure of its saturation by trajectories of the Liouville vector field $Z = p\frac{\partial}{\partial p}$, is an isotropic subset.

Definition 1.1. *Arboreal Lagrangian (resp. Legendrian) singularities* form the smallest class $\text{Arb}_n^{\text{symp}}$ (resp. $\text{Arb}_n^{\text{cont}}$) of germs of closed isotropic subsets in $2n$ -dimensional symplectic (resp. $(2n+1)$ -dimensional contact) manifolds such that the following properties are satisfied:

- (i) (Invariance) $\text{Arb}_n^{\text{symp}}$ is invariant with respect to symplectomorphisms and $\text{Arb}_n^{\text{cont}}$ is invariant with respect to contactomorphisms.
- (ii) (Base case) $\text{Arb}_0^{\text{symp}}$ contains $pt = \mathbb{R}^0 \subset T^*\mathbb{R}^0 = pt$.
- (iii) (Stabilizations) If $L \subset (X, \omega)$ is in $\text{Arb}_n^{\text{symp}}$, then the product $L \times \mathbb{R} \subset (X \times T^*\mathbb{R}, \omega + dp \wedge dq)$ is in $\text{Arb}_{n+1}^{\text{symp}}$.
- (iv) (Legendrian lifts) If $L \subset T^*\mathbb{R}^n$ is in $\text{Arb}_n^{\text{symp}}$, then its Legendrian lift $\widehat{L} \subset J^1\mathbb{R}^n$ is in $\text{Arb}_n^{\text{cont}}$.
- (v) (Liouville cones) Let $\Lambda_1, \dots, \Lambda_k \subset S^*\mathbb{R}^n$ be a finite disjoint union of arboreal Legendrian germs from $\text{Arb}_{n-1}^{\text{cont}}$ centered at points $z_1, \dots, z_k \in S^*\mathbb{R}^n$. Let $\pi : S^*\mathbb{R}^n \rightarrow \mathbb{R}^n$ be the front projection. Suppose
 - $\pi(z_1) = \dots = \pi(z_k)$.
 - For any i , and smooth submanifold $Y \subset \Lambda_i$, the restriction $\pi|_Y : Y \rightarrow \mathbb{R}^n$ is an embedding (or equivalently, an immersion, since we only consider germs).
 - For any distinct i_1, \dots, i_ℓ , and any smooth submanifolds $Y_{i_1} \subset \Lambda_{i_1}, \dots, Y_{i_\ell} \subset \Lambda_{i_\ell}$, the restriction $\pi|_{Y_{i_1} \cup \dots \cup Y_{i_\ell}} : Y_{i_1} \cup \dots \cup Y_{i_\ell} \rightarrow \mathbb{R}^n$ is self-transverse.

Then the union $\mathbb{R}^n \cup C(\Lambda_1) \cup \cdots \cup C(\Lambda_k)$ of the Liouville cones with the zero-section form an arboreal Lagrangian germ from $\text{Arb}_n^{\text{symp}}$.

With the above classes defined, we can also allow boundary by additionally taking the product $L \times \mathbb{R}_{\geq 0} \subset (X \times T^*\mathbb{R}, \omega + dp \wedge dq)$ for any arboreal Lagrangian $L \subset (X, \omega)$, and similarly for arboreal Legendrians.




FIGURE 1.1. The simplest non-smooth arboreal singularity is the zero section union the positive conormal of a smooth co-oriented hypersurface.




FIGURE 1.2. This arboreal singularity consists of the zero section union the positive conormal of two smoothly intersecting co-oriented hypersurfaces.

In our paper [AGEN20a] we proved that for fixed dimension n the class of arboreal singularities contains only finitely many local models up to ambient symplectomorphism or contactomorphism. More precisely, to each member of the class $\text{Arb}_n^{\text{symp}}$ one can assign a signed rooted tree $\mathcal{T} = (T, \rho, \varepsilon)$ with at most $n + 1$ vertices. Here T is a finite acyclic graph,




FIGURE 1.3. This arboreal singularity consists of the zero section union the positive conormal of a singular co-oriented hypersurface (namely the front of the arboreal singularity of Figure 1.1).

ρ is a distinguished root vertex, and ε is a sign function on the edges of T not adjacent to ρ . This discrete data completely determines the germ:

Theorem 1.2 ([AGEN20a]). *If two arboreal singularities $L \subset (X, \omega)$, $L' \subset (X', \omega')$ have the same dimension and signed rooted tree \mathcal{T} , then there is (the germ of) a symplectomorphism $(X, \omega) \simeq (X', \omega')$ identifying L and L' .*

Within all arboreal singularities, there is the distinguished class with positive sign $\varepsilon \equiv +1$.

Definition 1.3. An *arboreal Lagrangian* L (with boundary) in a symplectic manifold (X, ω) is a piecewise smooth Lagrangian with arboreal singularities (with boundary), i.e. locally modeled on the above class (with boundary). When L is an arboreal Lagrangian whose boundary ∂L is a smooth manifold, we will say that L has *smooth boundary*.

A *positive arboreal Lagrangian* L (with boundary) in a symplectic manifold (X, ω) is a Lagrangian with positive arboreal singularities (with boundary), i.e. locally modeled on the distinguished class with positive sign $\varepsilon \equiv +1$.

Given any positive arboreal Lagrangian $L \subset (X, \omega)$, possibly with boundary, its neighborhood $U \subset X$ admits a canonical, up to homotopy, Lagrangian plane field $\xi \subset TU$. This homotopy class admits a representative ξ which is transverse to L , i.e. transverse to the closure of each smooth piece.

Definition 1.4. A *polarization* of a symplectic manifold (X, ω) is the choice of a Lagrangian plane field $\xi \subset TX$.

If (W, λ) is a Weinstein domain with positive arboreal skeleton, then (W, λ) , which retracts onto an arbitrarily small neighborhood of its skeleton, admits a polarization. The main result of this paper is that the converse also holds:




FIGURE 1.4. One can obtain a new arboreal singularity from the arboreal singularity of Figure 1.1 in two different ways depending on how it is embedded in S^*M relative to the vertical distribution $\nu = \ker(d\pi)$. One yields a positive arboreal singularity and the other does not.




FIGURE 1.5. The canonical polarization in the neighborhood of an A_2 singularity, where the positivity condition is vacuous. One could also turn the tangent field to the zero section counter-clockwise instead of clockwise. The choice is determined by what we call an orientation structure, which in this case is equivalent to a co-orientation of the origin inside the zero-section.

Theorem 1.5. *A Weinstein domain (W, λ) is homotopic to a Weinstein domain whose skeleton is positive arboreal with smooth boundary if and only if (W, λ) admits a polarization.*

Remark 1.6. The conclusion of Theorem 1.5 can be refined in several ways, some of which we sketch below. For precise formulations see Theorem 6.16 and its corollaries, which use the language of Wc-manifolds, see Definition 2.24.

- (i) In fact, for ξ a polarization of (W, λ) we can arrange it so that ξ is the canonical polarization in the neighborhood of the positive arboreal skeleton, so in particular ξ is transverse to the skeleton.




FIGURE 1.6. The canonical polarization in the neighborhood of a positive A_3 singularity, where the orientation structure is given by the co-orientation of the A_3 front.

- (ii) If so desired, the smooth boundary of the positive arboreal skeleton can be pushed out to the boundary of W , in which case it serves as a skeleton for a *Weinstein pair*, see Section 2.4. The Weinstein hypersurface in the pair is thus the ribbon of a smooth Legendrian, and can also be thought of as a stop.
- (iii) Theorem 1.5 also holds with input a Weinstein pair (W, A) instead of a Weinstein domain. In this case the conclusion is that the Weinstein hypersurface A can be enlarged by the ribbon of a smooth Legendrian so that the resulting Weinstein pair has a positive arboreal skeleton.
- (iv) Moreover, the version of Theorem 1.5 for Weinstein pairs also holds in relative form if the Weinstein hypersurface in the Weinstein pair already has a positive arboreal skeleton and the polarization restricts on it to the canonical polarization. In this case the Weinstein structure can be kept fixed near the Weinstein hypersurface.
- (v) Theorem 1.5 and its refinements hold more generally for arbitrary Weinstein manifolds, not necessarily of finite type (i.e. not completions of Weinstein domains), and the result is proved in the same way since the inductive argument stabilizes. We restrict the discussion to the finite type case for simplicity of exposition.

Existence of a polarization for a symplectic manifold (X^{2n}, ω) is equivalent to asking that the classifying map $X \rightarrow BU_n$ of the tangent bundle lifts to BO_n , where we endow TX with the homotopically unique almost complex structure compatible with ω .

Definition 1.7. A *stable polarization* of a symplectic manifold (X, ω) is the choice of a Lagrangian plane field $\xi \subset TX \oplus \mathbb{C}^d$ for some $d \geq 1$.

The existence of a stable polarization is equivalent to asking that the classifying map $W \rightarrow BU$ of the stable tangent bundle lifts to BO . Recall that a Weinstein domain (W, λ) has the homotopy type of a half-dimensional CW complex. Recall also that $BO_k \rightarrow BO_{k+1}$ is

k -connected and $BU_k \rightarrow BU_{k+1}$ is $(2k+1)$ -connected. Therefore, it follows that for Weinstein domains the existence of a polarization is equivalent to the existence of a stable polarization. From the above, one can check that many Weinstein domains admit polarizations, notably smooth complete intersections in complex affine space. Indeed, if $X \subset \mathbb{C}^N$ is a smooth complete intersection, then its normal bundle is trivial, hence its tangent bundle is stably trivial and in particular X admits a stable polarization. This proves:

Corollary 1.8. *Let $X \subset \mathbb{C}^N$ be a complete intersection. Then X is Weinstein homotopic to a Weinstein manifold whose skeleton is positive arboreal with smooth boundary.*

An arboreal skeleton with boundary comes with a natural additional orientation structure, induced from the ambient symplectic structure. We formalize this in the notion of an arboreal space and prove the following counterpart to Theorem 1.5.

Theorem 1.9. *Any compact arboreal space with boundary arises as the skeleton of a Weinstein domain (W, λ) , unique up to Weinstein homotopy.*




FIGURE 1.7. The simplest example of two non-equivalent arboreal spaces with diffeomorphic underlying arboreal Lagrangians. These are skeleta for the pair of pants and the once-punctured torus respectively. The arboreal spaces are distinguished by the orientation structure, which is additional data, and in this dimension reduces to the usual cyclic structure on ribbon graphs. We thank A. Oancea for pointing out this example.

Hence, up to Weinstein homotopy, those Weinstein domains which admit polarizations are precisely the Weinstein thickenings of positive arboreal spaces. Moreover, the relative form of Theorem 1.5 implies a uniqueness result for positive arboreal skeleta. To state it we need the notion of a Weinstein concordance, which is best defined in the language of Wc-manifolds, see Definition 2.24. Briefly, if λ_1 and λ_2 are two Weinstein structures on the same symplectic domain (W, ω) with Liouville fields Z_1, Z_2 , respectively, then a Weinstein concordance between

them is a Weinstein-type primitive λ for $\omega + du \wedge dt$ on $W \times T^*[0, 1]$ such that the Liouville field Z of λ is tangent to the boundary $W \times T_0^*[0, 1] \cup W \times T_1^*[0, 1]$ and restricts to $Z_1 + u\partial/\partial u$ and $Z_2 + u\partial/\partial u$ over $W \times T_0^*[0, 1]$ and $W \times T_1^*[0, 1]$ respectively.

Definition 1.10. A *positive arboreal concordance* is a Weinstein concordance $(W \times T^*[0, 1], \lambda)$ whose skeleton is a positive arboreal Lagrangian with boundary. We assume that the boundary of the skeleton is smooth in the interior of $W \times T^*[0, 1]$ but we allow simple corners with one face contained in $W \times T_0^*[0, 1]$ or $W \times T_1^*[0, 1]$ and the other face transverse to it.

Remark 1.11. We must allow simple corners so that for example if (W, λ) is a Weinstein manifold whose skeleton is a positive arboreal with nonempty smooth boundary, then the trivial product $(W \times T^*[0, 1], \lambda + udt)$ is a positive arboreal concordance.

The uniqueness statement then reads as follows:

Theorem 1.12. *Suppose that λ_1 and λ_2 are two homotopic Weinstein structures on W whose skeleta are positive arboreal Lagrangians with smooth boundary, transverse to polarizations ξ_1 and ξ_2 respectively. Then ξ_1 and ξ_2 are stably homotopic if and only if there is a positive arboreal concordance between λ_1 and λ_2 .*

We remark that the classification of polarizations on a Weinstein manifold up to stable homotopy is in general weaker than the classification up to homotopy.

1.3. Further development of the arborealization program. Work in progress of the authors extends the results of this paper in two directions. First, in our forthcoming work [AGEN21] we consider a 1-parametric version of the problem, by introducing a suitable notion of “arboreal homotopy of positive arboreal spaces”, i.e. the minimal sequence of combinatoral Reidemeister type moves necessary to connect two homotopic Weinstein structures with positive arboreal skeleta that have homotopic canonical polarizations. Note that substantial progress in this direction appeared in Zorn’s thesis [Z18], where local mutations of arboreal skeleta are classified.

The second direction is an extension of the arborealization program to a more general class of Weinstein manifolds. While most Weinstein manifolds considered in applications such as mirror symmetry are complete intersections and hence polarized, not all Weinstein manifolds admit polarizations. For example a cohomological obstruction to existence of a polarization is that all odd Chern classes are 2-torsion. However, this cohomological condition does not obstruct existence of a not necessarily positive arboreal skeleton.

But even allowing for non-positive arboreal skeleta, there are still obstructions to their existence. Indeed, while existence of a polarization is not a necessary condition for existence of a non-positive arboreal skeleton, there is a closely related necessary condition: existence of an $(n, n - 1)$ -polarization. This is, roughly speaking, a polarization which is allowed to degenerate to an $(n - 1)$ -dimensional isotropic subspace. More precisely, it is defined as follows.

Given a symplectic vector space (E, ω) let $L_n(E)$ denote its Lagrangian Grassmanian of non-oriented Lagrangian planes, and $I_{n-1}(E)$ denote the Grassmanian of non-oriented isotropic $(n-1)$ -dimensional subspaces. We further denote by $F_{n,n-1}(E)$ the flag manifold $\{(\lambda, \mu); \lambda \in L_n(E), \mu \in I_{n-1}(E), \mu \subset \lambda\}$. Let π_L and π_I be the tautological projections $\pi_L : F_{n,n-1}(E) \rightarrow L_n(E)$ and $\pi_I : F_{n,n-1}(E) \rightarrow I_{n-1}(E)$. Consider the double cone $C_n(E) := (F_{n,n-1} \times [0, 1] \cup L_n(E) \cup I_{n-1}(E)) / \{f \times 0 \sim \pi_L(f), f \times 1 \sim \pi_I(f)\}$. Given any symplectic vector *bundle* E , we will use the notation $C_n(E)$ for the associated cone bundle. An $(n, n-1)$ -polarization on a symplectic manifold V is by definition a *section* of the cone bundle $C_n(T(V))$.

One can show that for $n \leq 5$ any $2n$ -dimensional Weinstein manifold admits a $(n, n-1)$ -polarization, while for $n = 6$ there are obstructions (e.g. one must have $c_3(V) = c_1(V)c_2(V)$).¹ It is possible that this condition is also sufficient for existence of an arboreal skeleton. Note in particular that a Weinstein manifold of dimension four always admits an $(n, n-1)$ -polarization, so this discussion is consistent with Starkston's result [St18] that four-dimensional Weinstein manifolds always admit arboreal skeleta.

For completely arbitrary Weinstein manifolds it is not clear whether one should expect there to be a simple class of singularities which the skeleton can always be arranged to have, but if it exists such a class would have to be more general than the arboreal class. In any case, even the most optimistic hopes of constructing a reasonable skeleton for a Weinstein manifold will in all likelihood require the existence of “Maslov data”, i.e. the homotopical trivialization required to define Fukaya categories or microlocal sheaves.

1.4. Flexibility of caustics and the rigidification theorem. We discuss a toy example to illustrate our strategy of proof. Consider a Weinstein domain (W^{2n}, λ) which is obtained from the standard Darboux ball $B \subset (\mathbb{R}^{2n}, pdq - qdp)$ by attaching an index n handle, i.e. the neighborhood of a Lagrangian disk, along a Legendrian sphere $\Lambda \subset \partial B = (S^{2n-1}, \xi_{\text{std}})$. The skeleton of W is homeomorphic to the n -sphere: it consists of the Lagrangian disk union the Liouville cone on Λ with respect to the radial Liouville field on \mathbb{R}^{2n} , union the origin.

Replace the radial Liouville structure $pdq - qdp$ on \mathbb{R}^{2n} with the Morse-Bott canonical Liouville structure pdq on the cotangent bundle $T^*\mathbb{R}^n$, or rather on T^*D for $D \subset B$ a Lagrangian n -disk bounding a Legendrian unknot $\partial D \subset S^{2n-1}$ which we assume disjoint from Λ . Then the radial Liouville cone on $\Lambda \subset S^{2n-1}$ in B gets replaced with the fibrewise Liouville cone on $\Lambda \subset S^*D$ in T^*D , where S^*D denotes the cosphere bundle, see Figure 1.8.

Observe that as a result, the singularity gets spread out over the disk, i.e. instead of a cone on a point it is now a fibrewise cone over the disk. The skeleton of the new Weinstein structure on W consists of the Lagrangian handle, union the Liouville cone on Λ with respect to the canonical Liouville structure on T^*D , union the zero section D . The singularities of this new Lagrangian skeleton are thus related to the singularities of the map $\pi|_\Lambda : \Lambda \rightarrow D$, where $\pi : S^*D \rightarrow D$ is the front projection. These singularities correspond to the tangencies of Λ

¹We thank John Pardon for suggesting to us the notion of a $(n, n-1)$ -polarization, and to Søren Galatius for computations of obstructions.

with the distribution $\nu = \ker(d\pi) \subset T(S^*D)$ and are also known as caustics in the literature. For example, if $\pi|_\Lambda : \Lambda \rightarrow D$ has no caustics, then it is an immersion and so by axiom (v) of Definition 1.1, after a generic perturbation (to ensure self-transversality), the skeleton of W is arboreal.




FIGURE 1.8. The blow-up procedure for a Weinstein manifold W^{2n} with two critical points: one of index 0 and one of index n .

The idea of blowing up conical Lagrangian singularities was already present at the inception of the arborealization program, but for our purposes it is necessary to perform the blowing up procedure globally, at the level of Weinstein structures. This strategy was successfully implemented by Starkston in [St18] for the 4-dimensional case. Starkston moreover showed how to explicitly arborealize the semi-cubical cusp singularities which generically appear in the front projections of 1-dimensional Legendrians, thus leading to her result that 4-dimensional Weinstein manifolds always admit arboreal skeleta. In higher dimensions this strategy encounters the difficulty that there is no classification theorem for the generic singularities of front projections, nor can anything remotely close to a satisfactory classification be hoped for.

Nevertheless, there holds an h-principle for the simplification of caustics. The h-principle states that if there is no homotopy theoretic obstruction to the simplification of caustics, then the simplification can be achieved by means of a Legendrian isotopy (or for Lagrangians, by means of a Hamiltonian isotopy). Results in this direction first appeared in work of Entov [En99], and the full h-principle was established in work of the first author [AG18a], [AG18b]. Returning to our toy example, if we know that W admits a polarization as in the hypothesis of our main Theorem 1.5 and we further assume that the dimension of W is congruent to 2 modulo 4, then it follows from the h-principle that there exists a Legendrian isotopy which deforms Λ to a Legendrian whose front projection $\pi|_\Lambda : \Lambda \rightarrow D$ only has cusp singularities. In this case we can proceed as in [St18] to conclude that the skeleton of W can be arborealized by a Weinstein homotopy. When the dimension of W is congruent to 0 modulo 4 there is an additional homotopical subtlety to consider, even in this toy example, but we do not pursue

this further since one also encounters additional and more serious difficulties when attempting to implement the above strategy on a Weinstein manifold with a more complicated handlebody presentation. For example, one would need to know that the relevant homotopical obstruction to applying the h-principle vanishes for every handle, yet it is unclear how to arrange for this even under the global assumption of existence of a polarization.

So instead of attempting a direct application of the h-principle, we will exploit the fact that for the purposes of arborealization we can allow deformations which are more general than a Hamiltonian isotopy, namely we can deform the skeleton of a Weinstein manifold by any homotopy of the Weinstein structure. This allows the skeleton to develop genuine singularities (i.e. non-smooth points) along which its field of tangent planes jumps discontinuously. By introducing these jumps we destroy the homotopical obstruction to simplifying the singularities with the distribution $\nu = \ker(d\pi)$, and in fact we can always completely eliminate these tangencies.

This strategy was made precise in [AGEN19], where we introduced a class of singular Lagrangians, called ridgy, and proved an h-principle *without pre-conditions*, which allows one to deform any Lagrangian so that it becomes transverse to any given Lagrangian distribution, at the expense of developing ridgy singularities. This result, which we refer to as the *ridgification theorem*, uses the h-principle for the simplification of caustics as one of the key ingredients in its proof, but is better suited for the task at hand since there are no hypotheses needed for its application.




FIGURE 1.9. The ridge singularity in one dimension.

Ridgy singularities are extremely simple: they consist of the corner $\{p = |q|\} \subset \mathbb{R}^2$ together with its products and stabilizations. Moreover, ridgy singularities can be explicitly arborealized while maintaining transversality to any given Lagrangian distribution, see Figure 1.11. So our revised strategy to arborealize a Weinstein manifold W is the following: (1) blow up the singularities as before to replace Liouville cones on a point with Liouville cones on Lagrangian disks, (2) apply the ridgification theorem to get rid of all singularities of the resulting front projections, and finally (3) arborealize the ridgy Lagrangians produced by the ridgification theorem, while maintaining transversality to the relevant distributions, to obtain an arboreal skeleton for W .




FIGURE 1.10. The ridgification theorem in action: the zero section $M \subset T^*M$ gets deformed to a ridgy Lagrangian $L \subset T^*M$ which is transverse to the red distribution. Of course in this simple example a single ridge point would suffice, but the proof always produces a ridge locus which is the union of homologically trivial hypersurfaces, in this case two ridge points.



FIGURE 1.11. Arborealization of an order 1 ridge.




FIGURE 1.12. Arborealization of an order 2 ridge.

Finally, a word on how we deal with the issue of compatibility at the interaction of three or more strata of the skeleton: this is where the notion of positivity comes up in an essential way. The key fact is that if a Lagrangian distribution η on $T^*\mathbb{R}^n$ corresponds to a family of positive definite quadratic forms, then it is transverse to the conormal $T_H^*\mathbb{R}^n \subset T^*\mathbb{R}^n$ of *any* immersed hypersurface $H \subset \mathbb{R}^n$ in a neighborhood of the zero section. For the interaction of say three strata, we use the ridgification theorem to achieve transversality for two strata at a time and then conclude that transversality also holds for the third stratum *for free* using the above observation on positivity. This yields the desired compatibility. The existence of

a global polarization for W will play a key role in guiding the positivity condition globally, thus enabling the successful implementation of the above strategy.

1.5. Structure of the article. The paper is organized as follows:

In Section 2 we give the necessary preliminaries regarding Weinstein manifolds. This includes a review of notions in the literature as well as the introduction of a new framework for working with Weinstein structures. The key notions are that of a Weinstein manifold with boundary and corners, which we call a Wc-manifold, and that of a cotangent building, which provides a way of presenting a Wc-manifold in terms of successive attachments of cotangent bundles. The concept of a cotangent building gives an alternative to the standard Weinstein handlebody presentation for Weinstein manifolds, and is better suited for our purposes.

In Section 3 we prove the existence and uniqueness of symplectic neighborhoods of arboreal spaces. The existence part is an easy consequence of the stability theorem for arboreal singularities, which we proved in [AGEN20a]. The uniqueness part consists of a Darboux-Weinstein theorem for the symplectic neighborhoods of arboreal Lagrangians.

In Section 4 we discuss the notion of positivity, which is the key ingredient that allows us to control the interaction of three or more strata in the Lagrangian skeleta under consideration. We first discuss the notion of positivity for ordered tuples of Lagrangian planes, then discuss positivity for cotangent buildings and finally we relate this to the notion of positivity for arboreal spaces. We show that existence of a positive arboreal skeleton for the homotopy class of a Weinstein structure implies the existence of a polarization, proving one half of our main theorem 1.5.

In Section 5 we recall the notion of a ridgy Lagrangian from our previous work [AGEN19] and state the main result of that paper, which we call the ridgification theorem, as well as its non-integrable counterpart, which we call the formal ridgification theorem. The formal ridgification theorem also plays an important role since it holds in extension form. The main point is that by allowing for ridgy singularities we can deform any Lagrangian submanifold so that it becomes transverse to any given Lagrangian distribution, and moreover the deformation can be achieved by means of a Weinstein homotopy.

Finally, in Section 6 we show how the ridgy singularities produced by the ridgification theorem can be further deformed into arboreal singularities. Furthermore, the deformation is realized by a Weinstein homotopy and the end result is a positive arboreal Lagrangian which serves as skeleton for the deformed Weinstein structure. This proves the other half of our main theorem 1.5.

1.6. Acknowledgements. We are very grateful to Laura Starkston who collaborated with us on the initial stages of this project. We are also grateful to John Pardon who helped us to crystalize the notion of an $(n, n - 1)$ -polarization and to Søren Galatius who explained to us how to compute obstructions to its existence.

The first author thanks the Institute for Advanced Study and Princeton University for a great research environment and thanks CRM Montreal for its hospitality. The second author

thanks RIMS Kyoto and ITS ETH Zurich for their hospitality. The third author thanks MSRI for its hospitality.

Finally, we are very grateful for the support of the American Institute of Mathematics, which hosted a workshop on the arborealization program in 2018 from which this project has greatly benefited.

2. WC-MANIFOLDS AND COTANGENT BUILDINGS

In this section we recall the basic definitions concerning Liouville and Weinstein manifolds, and introduce the language of Wc-manifolds and cotangent buildings, which will be used throughout the text.

2.1. Liouville manifolds.

2.1.1. *Liouville domains.* An exact symplectic manifold (W, λ) is a compact manifold with boundary equipped with an exact symplectic form ω , together with a choice of primitive λ , i.e. $\omega = d\lambda$. Since ω is non-degenerate, the 1-form λ corresponds to a vector field Z under the contraction $\iota_Z \omega = \lambda$.

Definition 2.1. A *Liouville domain* is an exact symplectic manifold (W, λ) such that Z is outwards transverse to ∂W .

Equivalently, (W, λ) is Liouville if $\lambda|_{\partial W}$ is a contact form and the orientation defined on ∂W by the volume form $\lambda \wedge (d\lambda)^{n-1}|_{\partial W}$ coincides with the orientation of ∂W as the boundary of W , where W is itself oriented by the volume form $\omega^{\wedge n}$.

We call λ the *Liouville form* and Z the *Liouville field*. By Cartan's formula for the Lie derivative, the condition $\omega = d\lambda$ is equivalent to $\mathcal{L}_Z \omega = \omega$. This means that the vector field Z is symplectically conformally expanding, i.e. $(Z^t)_* \omega = e^t \omega$, $t \geq 0$, for Z^t the flow of Z . Equivalently, the vector field $-Z$ is symplectically conformally contracting and this viewpoint will sometimes be more natural in what follows. Note that we also have $(Z^t)_* \lambda = e^t \lambda$. The object of interest in this paper is the following.

Definition 2.2. The *skeleton* of a Liouville domain (W, λ) is the subset

$$\text{Skel}(W, \lambda) = \bigcap_{t>0} Z^{-t}(W).$$

In other words, $\text{Skel}(W, \lambda)$ is the attractor of the positive flow of the contracting field $-Z$. We will be going back and forth between Liouville domains and the closely related notion of a Liouville manifold, whose definition we recall next.

2.1.2. *Liouville manifolds.* Recall that the *symplectization* $S(Y, \xi)$ of a contact manifold (Y, ξ) equipped with a contact form α for ξ is the exact symplectic manifold $Y \times \mathbb{R}$ equipped with the primitive $\lambda = e^t \alpha$, where $t \in \mathbb{R}$. We can equivalently consider $Y \times \mathbb{R}^+$ with $\lambda = s\alpha$ for $s = e^t \in \mathbb{R}^+$ the multiplicative coordinate.

The symplectization of a contact manifold (Y, ξ) can be defined invariantly without choosing a contact form. Assuming that Y is connected and ξ is co-oriented, we define $S(Y, \xi)$ as follows. Let $N(\xi) \subset T^*Y$ be the total space of the conormal line bundle to ξ and $N_+(\xi)$ the component of $N(\xi) \setminus Y$ consisting of 1-forms defining the given co-orientation of ξ . Then $\lambda_\xi = pdq|_{N_+(\xi)}$ is a primitive of the exact symplectic form $\omega = d\lambda_\xi$ and we define $S(Y, \xi)$ as $N_+(\xi)$ endowed with this structure. A choice of a contact form α for ξ identifies $N_+(\xi)$ with $Y \times \mathbb{R}^+$ and λ_ξ with $s\alpha$. For example, the symplectization of the cosphere bundle S^*M is the complement of the zero section in T^*M .

In the other direction, the *contactization* of an exact symplectic manifold (N, μ) is the manifold $M \times \mathbb{R}$ equipped with the contact form $\mu - du$, where $u \in \mathbb{R}$. For example, the contactization of the cotangent bundle T^*M is the 1-jet bundle J^1M .

Definition 2.3. A *finite type Liouville manifold* (X, λ) consists of a (non-compact) boundaryless exact symplectic manifold such that the following conditions hold.

- (L1) The vector field Z which is ω -dual to λ is complete for $t \rightarrow \pm\infty$.
- (L2) There exists a compact domain $W \subset X$ such that Z is outward transverse to the boundary ∂W , i.e. $(W, \lambda|_W)$ is a Liouville domain, and such that the union of forward trajectories of Z starting at ∂W is equal to $X \setminus \text{Int } W$.

We call Liouville domains as in (L2) *defining*.

Definition 2.4. The *skeleton* of a finite type Liouville manifold (X, λ) is defined to be

$$\text{Skel}(X, \lambda) = \bigcup_C \bigcap_{t>0} Z^{-t}(C), \quad C \subset X \text{ is compact.}$$

Clearly, $\text{Skel}(X, \lambda) = \text{Skel}(W, \lambda)$ for any defining Liouville domain $W \subset X$. Hence the skeleton of a finite type Liouville manifold is compact. Note that for any such W we may use the Liouville flow starting at ∂W to parametrize $X \setminus \text{int}W$ as $\partial W \times [0, \infty)$. Since the Liouville field conformally expands λ we have $\lambda|_{\partial W \times [0, \infty)} = e^t(\lambda|_{\partial W})$ for $t \in [0, \infty)$ the coordinate parametrizing the Liouville flow, so $X \setminus \text{int}W$ is the positive part of the symplectization of the contact manifold $(\partial W, \lambda|_{\partial W})$.

Conversely, any Liouville domain (W, λ) can be completed to a Liouville manifold by attaching to it the positive part of the symplectization of $(\partial W, \lambda|_{\partial W})$. Explicitly:

$$X = W \bigcup_{\partial W \sim \partial W \times 0} \partial W \times [0, \infty),$$

where the Liouville form λ on W extends to $\partial W \times [0, \infty)$ as $e^t(\lambda|_{\partial W})$.

Definition 2.5. The *ideal boundary* of a finite type Liouville manifold (X, λ) is defined to be $\partial_\infty X = (X \setminus \text{Skel}(X, \lambda))/\mathbb{R}^+$, where the \mathbb{R}^+ action is given by the positive flow of Z .

Since $\lambda(Z) = 0$, the form λ descends to $\partial_\infty X$ as a contact structure ξ_∞ . A choice of a contact form α for ξ_∞ yields an isomorphism $j_\alpha : S(\partial_\infty X, \xi_\infty) = (\partial_\infty X \times \mathbb{R}, e^t\alpha) \rightarrow (X \setminus \text{Skel}(X, \lambda), \lambda)$

which maps $\partial_\infty X \times 0$ contactomorphically onto the boundary of a defining Liouville domain. We denote by $\pi_\infty : X \setminus \text{Skel}(X, \lambda) \rightarrow \partial_\infty X$ the projection along the trajectories of Z .

2.1.3. Deformations of Liouville structures.

Definition 2.6. A homotopy of finite type Liouville structures (X, λ_t) is a family λ_t , $t \in [0, 1]$, of finite type Liouville structures on a manifold X admitting a smooth family $W_t \subset X$ of defining Liouville domains.

Given such a homotopy (X, λ_t) there exists an isotopy $\phi_t : X \rightarrow X$ such that $\phi_t^* \omega_t = \omega_0$, where $\omega_t = d\lambda_t$. Moreover, one can always arrange that $\phi_t^* \lambda_t = \lambda_0 + dH_t$ for uniformly compactly supported functions H_t , see [CE12], Sections 11.1 and 11.2. In particular, it is always sufficient to consider homotopies fixing the symplectic form, and moreover changing the Liouville form by adding a compactly supported exact form. Note that for any fixed ω the space of Liouville structures on (X, ω) which are fixed at infinity is convex, hence any two such Liouville structures are canonically homotopic.

Remark 2.7. Unless explicitly stated otherwise, all Liouville structures under consideration in this paper are of finite type and hence we will stop mentioning this below.

We will often abuse terminology and say that two Liouville manifolds (X_1, λ_1) and (X_2, λ_2) are Liouville homotopic if there exists a symplectomorphism $F : (X_1, d\lambda_1) \rightarrow (X_2, d\lambda_2)$ such that $(X_1, F^* \lambda_2)$ is a Liouville manifold which is Liouville homotopic to (X_1, λ_1) . In all cases where such terminology will be abused, the symplectomorphism will be obvious and unique up to contractible choice.

If $F^* \lambda_2 = \lambda_1$ on the nose, we say that (X_1, λ_1) and (X_2, λ_2) are *Liouville isomorphic*. For (X, λ) and (Y, μ) Liouville domains or manifolds we call a smooth, proper embedding $F : (X, \lambda) \rightarrow (Y, \mu)$ a *Liouville embedding* if $F^* \mu = \lambda$.

2.1.4. *Liouville germs.* It will often be convenient to express our constructions in the language of germs.

Definition 2.8. A *Liouville germ* (\mathcal{X}, λ) is the equivalence class of defining Liouville domains for a fixed Liouville manifold (X, λ) .

Definition 2.9. A Liouville embedding of a Liouville germ (\mathcal{X}, λ) into a Liouville domain or manifold (Y, μ) is an equivalence class of Liouville embeddings $F_W : (W, \lambda|_W) \rightarrow (Y, \mu)$, where $W \subset X$ is a defining domain for (X, λ) and we identify F_{W_0} with F_{W_1} if they agree on a defining domain W_2 for (X, λ) contained in both W_0 and W_1 .

Example 2.10. The cotangent bundle $X = T^*M$ of a closed manifold M is a Liouville manifold with $\lambda = pdq$ the canonical 1-form. The Liouville field is the fibrewise radial vector field $Z = p\partial_p$ and the skeleton is the zero section $M \subset T^*M$. As a Liouville domain we can take the unit disk bundle $W = \{\|p\| \leq 1\}$ relative to any Riemannian metric on M . The ideal

boundary is the cosphere bundle S^*M , i.e the (positively) projectivized cotangent bundle, which is contactomorphic to the unit sphere bundle $\partial W = \{\|p\| = 1\}$ for any choice of Riemannian metric. We will use the notation \mathcal{T}^*M for the germ of the cotangent bundle T^*M .

2.2. Weinstein manifolds. The notion of *Weinstein manifold* was first introduced in [EG91] building on [E90] and [W91].

2.2.1. Lyapunov functions. Let (X, λ) be a Liouville manifold. While $\text{Skel}(X, \lambda)$ always has its $2n$ -dimensional Lebesgue measure equal to 0, its dimension can nonetheless be quite large if no extra conditions are imposed on the Liouville structure. For example, McDuff constructed in [MD91] a Liouville structure on $T^*\Sigma_g \setminus \Sigma_g$ for Σ_g a closed surface of genus $g > 1$ whose skeleton has codimension 1. To tame the dynamics of the Liouville flow we require existence of a Lyapunov function.

Definition 2.11. We say that $\phi : X \rightarrow \mathbb{R}$ is a *Lyapunov function* for the Liouville manifold (X, λ) if Z is gradient-like for ϕ , i.e. if there holds

$$(W1) \quad d\phi(Z) \geq \delta(\|Z\|^2 + \|d\phi\|^2) \text{ for some Riemannian metric on } X \text{ and some constant } \delta > 0.$$

Note that the space of Lyapunov functions for a given Liouville structure is convex, hence contractible as soon as it is nonempty. Consider the set of critical points $\text{Crit}(\phi) = \{d\phi = 0\}$ of the Lyapunov function, which by (W1) is the same as the zero set $\{\lambda = 0\}$ of the Liouville form. The conditions (L1) and (W1) imply that $\text{Skel}(X, \lambda)$ is the union of the Z -stable manifolds of the critical points of ϕ , i.e. points converging to $\text{Crit}(\phi)$ in forward time. However, as far as we know, we have to add some further constraints on the Lyapunov function ϕ in order to deduce any meaningful properties.

Definition 2.12. A *Morse-Weinstein manifold* is a Liouville manifold (X, λ) for which there exists a Morse Lyapunov function $\phi : X \rightarrow \mathbb{R}$.

Under this assumption it was shown in [CE12], see also [EG91, E95], that

$$(W2) \quad \text{Skel}(X, \lambda) \text{ is the union of submanifolds which are isotropic for } \lambda, \text{ and hence for } \omega.$$

Remark 2.13. Note that λ -isotropic is equivalent to ω -isotropic and Z -conic.

Without any assumptions on ϕ , is not known whether a triple (X, λ, ϕ) which satisfies (L1), (L2) and (W1) is homotopic in the class of such triples to one with a Morse Lyapunov function. However, condition (W2) holds for a larger class of taming functions, for example when ϕ is generalized Morse, see [CE12]. Moreover, it is often most natural to consider a Lyapunov function which is not Morse or even generalized Morse, such as the Morse-Bott function $\phi = \|p\|^2$ on (T^*M, pdq) .




FIGURE 2.1. The skeleton of an open Riemann surface with Morse Lyapunov function.

2.2.2. *Morse-Bott Lyapunov functions.* In this paper we will consider a version of Morse-Bott Weinstein structures which allows the critical set of the Lyapunov function to have nonempty boundary, and even corners. Our definition is a slight variation of Starkston's definition in [St18] of a Weinstein structure which is Morse-Bott with boundary.

Let $E^+ \oplus E^0 \oplus E^-$ be the decomposition of the tangent space to X at a zero of Z into generalized eigenspaces of the differential dZ with eigenvalue having positive, zero or negative real part respectively. Suppose first that a connected component C of $\text{Crit}(\phi)$ is a smooth, proper submanifold of X without boundary. Then the Morse-Bott condition on a triple (X, λ, ϕ) along the component C is the following:

(MB1) Z is non-degenerate in the directions normal to C , i.e. $TC = E^0|_C$.

When $\partial C \neq \emptyset$ we moreover demand that in a neighborhood of ∂C the triple (X, λ, ϕ) takes a special form, which we now describe. Recall the Weinstein normal form for Morse critical points in a Weinstein manifold of dimension 2. There are two possibilities depending on the index: $k = 0$ or $k = 1$.

$$(k = 1) \quad \lambda_1 = 2pdq + qdp, \quad \phi_1 = p^2 - \frac{1}{2}q^2.$$

$$(k = 0) \quad \lambda_0 = \frac{1}{2}(pdq - qdp), \quad \phi_0 = \frac{1}{4}(p^2 + q^2).$$

We also recall the standard Morse-Bott normal form corresponding to the canonical Liouville structure on $T^*\mathbb{R}$.

$$(\text{std}) \quad \lambda_{\text{std}} = pdq, \quad \phi = \frac{1}{2}p^2.$$

We construct local models for *Morse-Bott with boundary* Weinstein structures on $T^*\mathbb{R}$ with the half line $\{p = 0, q \geq 0\}$ as the critical set. The construction consists of patching up the above Morse normal forms on the half plane $\{q \leq 0\}$ with the standard Morse-Bott normal form on the half plane $\{q \geq 0\}$. Take any smooth function $\psi : \mathbb{R} \rightarrow [0, 1]$ satisfying the following properties:

- (i) $\psi = 0$ on $[0, \infty)$
- (ii) $\psi = 1$ on $(-\infty, -\varepsilon]$ for some $\varepsilon > 0$.
- (iii) $\psi < 1$ and $\psi' < 0$ on $(-\varepsilon, 0)$ for that same $\varepsilon > 0$.

In the case $k = 1$ we set $f_1(q, p) = \psi(q)pq$ and in the case $k = 0$ we set $f_0(q, p) = -\frac{1}{2}\psi(q)pq$. In both cases the patched up model is then given by $\lambda_k^{\text{MB}} = \lambda_{\text{std}} + df_k$, which has Liouville field $Z_k^{\text{MB}} = Z_{\text{std}} + X_{f_k}$ for X_{f_k} the Hamiltonian vector field of f_k . Explicitly, when $k = 1$ we have

$$Z_1^{\text{MB}} = (1 + \psi(q) + \psi'(q)q) p\partial_p - \psi(q)q\partial_q.$$

Since the coefficient which multiplies $p\partial_p$ is positive, we deduce that Z_1^{MB} admits a Lyapunov function. For example, if we set $\phi = p^2 - h(q)$ for

$$h(q) = \int_0^q \psi(t)t dt,$$

then Z_1^{MB} is the Euclidean gradient of ϕ along $p = 0$ and we have $d\phi(Z_1^{\text{MB}}) > 0$ on all of $T^*\mathbb{R}$, from which the claim follows. Similarly, one verifies that Z_0^{MB} also admits a Lyapunov function.

Definition 2.14. We call $(T^*\mathbb{R}, \lambda_k^{\text{MB}})$ the *Weinstein normal form for Morse-Bott boundary* of index $k = 0, 1$.




FIGURE 2.2. The Weinstein normal form for Morse-Bott boundary of index 1 and 0 respectively.

The space of permissible ψ is convex, hence the normal form is well defined up to a contractible choice. Note that the behavior of the Liouville flow along the half line $\{p = 0, q \leq 0\}$, which shares its boundary point $\{p = 0, q = 0\}$ with the critical set $\{p = 0, q \geq 0\}$, can be either repelling or attracting depending on whether $k = 0$ or $k = 1$. We also consider products of these normal forms. Thus we obtain Weinstein structures on $T^*\mathbb{R}^n$ where the critical set is the positive quadrant $\{p_j = 0, q_j \geq 0\}$, which not only has boundary but also has corners of all orders $\leq n$.

Definition 2.15. The *Weinstein normal form for Morse-Bott n -fold corners* $(T^*\mathbb{R}^n, \lambda_r^{\text{MB}})$ of index $0 \leq r \leq n$ is the product of $n - r$ copies of $(T^*\mathbb{R}, \lambda_0^{\text{MB}})$ and r copies of $(T^*\mathbb{R}, \lambda_1^{\text{MB}})$.

Going back to a triple (X, λ, ϕ) as before, consider a component $C \subset X$ of the critical set of ϕ such that C is a smooth, proper submanifold of X with nonempty boundary ∂C and possibly with higher order corners. By definition, if $x \in \partial_k C \subset \partial C$ is a corner of order k , then there is a collar neighborhood $U \times \mathcal{I}^k$ of x in C , where \mathcal{I} is the germ of the interval $[0, 1)$ at 0 and $U \subset \partial_k C$ is a neighborhood of x in the order k corner locus. For such a collar neighborhood we have a local symplectomorphism near x between X and $T^*U \times T^*\mathbb{R}^k \times \mathbb{C}^m$, with C corresponding to $U \times \mathcal{I}^k \times 0$. The condition we require is that for each such k -fold point $x \in \partial_k C$ there exists a collar neighborhood $U \times \mathcal{I}^k$ as above such that with respect to the splitting $T^*U \times T^*\mathbb{R}^k \times \mathbb{C}^m$ there holds:

(MB2) λ splits as a direct sum with the summand corresponding to T^*U given by the canonical Liouville form and the summand corresponding to $T^*\mathbb{R}^k$ given by the Weinstein normal form for Morse-Bott k -fold corners λ_r^{MB} for some $r \leq k$.

Definition 2.16. A *generalized Morse-Bott Weinstein manifold* is a Liouville manifold (X, λ) which admits a Lyapunov function ϕ whose critical set is a union of smooth, proper submanifolds with corners $C \subset X$ satisfying (MB1) and (MB2).

Remark 2.17. It was shown by Starkston [St18] that in the boundary repelling Morse-Bott case the intersection of the Z -stable manifold of a component C of the critical set of ϕ with a neighborhood of C is a smoothly embedded isotropic submanifold of X , even without requiring standard local models. To check that this property also holds near the boundary attracting or mixed behavior points of the Morse-Bott structures considered above is straightforward, since it suffices to consider the explicit local model.

Lemma 2.18. *Suppose that (X, λ_0, ϕ_0) is a generalized Morse-Bott Weinstein manifold. Then there exists a homotopy (λ_t, ϕ_t) , fixed outside of a compact subset, such that (λ_t, ϕ_t) is Morse-Weinstein for $t > 0$.*

Proof. We give the proof in the case where each component $C \subset \Sigma$ of the critical set of ϕ is of the critical dimension $n = \frac{1}{2} \dim(X)$, the general case is similar and in any case will not be needed for our purposes. Suppose first that $C \subset \Sigma$ has empty boundary. Since C is Lagrangian, by the Weinstein neighborhood theorem it has a neighborhood in X symplectomorphic to a neighborhood of the zero section in $(T^*C, \lambda_{\text{std}} = pdq)$. Therefore we may from the onset replace X by T^*C with the pulled back Liouville form which we still denote by λ_0 .

Let $f : C \rightarrow \mathbb{R}$ be a Morse function and let $\eta : T^*C \rightarrow [0, 1]$ be a cutoff function such that $\eta = 1$ on $\mathcal{O}p(C)$ and $\eta = 0$ outside of a slightly bigger neighborhood. Fix an auxiliary Riemannian metric on C and put $h = \eta \lambda_{\text{std}}(\nabla f)$, a function on T^*C . The restriction of the Hamiltonian vector field of the function h to C is ∇f . Hence $\phi_0 + \eta f$ is a Lyapunov function for $\lambda_0 + dh$ in a neighborhood of C and therefore on all of T^*C if f is taken sufficiently C^1 small. Moreover, for such h the Liouville form $\lambda_0 + t dh$ is Morse-Weinstein for any $t > 0$.

If ∂C is non-empty, then we need to modify the construction somewhat. Consider a slight enlargement \widehat{C} of C in X obtained by attaching an isotropic collar along ∂C so that the

Liouville field Z_0 is tangent to \widehat{C} . Then we should require that our function $f : \widehat{C} \rightarrow \mathbb{R}$ is Morse on C , compactly supported in \widehat{C} , has no critical points on ∂C and moreover has the following behavior on ∂C . Let P be the closure of a component of the order 1 corner locus $\partial_1 C = \partial C \setminus \bigcup_{k>1} \partial_k C$. If the Liouville structure is boundary attracting ($k = 1$) along P , then we demand that $df(v) > 0$ for any inwards pointing vector $v \in TC|_{\partial C}$. If the Liouville structure is boundary repelling ($k = 0$) along P , then we demand that $df(v) < 0$ for any outwards pointing vector $v \in TC|_{\partial C}$. With this restriction on f the proof proceeds just as before. \square

Remark 2.19. As we will see, if a Morse-Bott Weinstein manifold has arboreal skeleton, then one can choose the Morsifying homotopy such that $\text{Skel}(X, \lambda_t) = \text{Skel}(X, \lambda_0)$ for all $t \geq 0$.

2.2.3. Weinstein manifolds. It is easy to check that Lemma 2.18 also holds for the generalized Morse condition instead of Morse-Bott, see Proposition 9.13 in [CE12]. Therefore we will take the following as our working definition, which includes Morse, generalized Morse and Morse-Bott Weinstein structures.

Definition 2.20. A Liouville manifold (X, λ_0) is called *Weinstein* if there exists a Liouville homotopy λ_t and a family of Lyapunov functions ϕ_t for Z_t such that (X, λ_t, ϕ_t) is Morse-Weinstein for $t > 0$.

The notions of a Weinstein domain, defining Weinstein domain for a Weinstein manifold and a Weinstein germ are entirely analogous to the corresponding notions in the Liouville case. We emphasize that the Lyapunov function ϕ is not part of the structure of a Weinstein manifold, we merely require its existence. For example, Weinstein embeddings are just Liouville embeddings of Weinstein type Liouville manifolds (or domains, or germs). We do not require the embedding to pull back one Lyapunov function to the other. Similarly, a Weinstein isomorphism (resp. homotopy) is a Liouville isomorphism (resp. homotopy) of Liouville manifolds of Weinstein type.

Example 2.21. The standard Liouville structure $\lambda = pdq$ on the cotangent bundle T^*M of a closed manifold M is Weinstein. Given a Riemannian metric on M , the function $\phi = \|p\|^2$ is a Morse-Bott Lyapunov function for λ . By the procedure explained in Lemma 2.18, we can Morsify the Lyapunov function ϕ using a Morse function on M . By inspection, this procedure does not change the skeleton of T^*M , which is the zero section.

Remark 2.22. Although Definition 2.20 allows for rather degenerate Weinstein structures, in the Weinstein homotopies constructed in this paper for the purposes of arborealization of a Morse-Weinstein manifold the Weinstein structures are always of Morse-Bott type except for a finite number of times at which there is a birth/death of Morse-Bott critical components.

2.3. Wc-manifolds.

2.3.1. *Manifolds with boundary and corners.* Recall that an n -dimensional smooth manifold M has a corner of order $k \leq n$ at $x \in M$ if there is a neighborhood of x in M diffeomorphic to a neighborhood of the origin in $\mathcal{I}^k \times \mathbb{R}^{n-k}$, where \mathcal{I} denotes the germ of the interval $[0, 1)$ at 0. We denote the locus of order k corners by $\partial_k M$. The closure P of a connected component of $\partial_k M$ is called a *boundary k -face*. For $k = 1$ we will more simply call P a *boundary face*. Sometimes we will call a manifold with boundary and corners (of any order) a *bc-manifold* for short.

We always assume that each component of $\partial_k M$ is regularly embedded, so that each k -face is itself a manifold with corners. Under this assumption, if Q is a k -face, then there is an embedded collar neighborhood $Q \times \mathcal{I}^k \subset M$. We consider the germs of these collars as part of the structure. In particular, near each point $x \in \partial_k M$ we have canonical collar coordinates $x = (y, t)$, where $y \in \partial_k M$ and $t = (t_1, \dots, t_k) \in \mathcal{I}^k$. Note that in a neighborhood of x we have $\partial_k M$ cut out by $t_1 = \dots = t_k = 0$. More generally, for $j \leq k$ the components of $\partial_j M$ whose closure contains x are given by setting exactly j of the coordinates t_i equal to zero.

We demand compatibility of these collars in the sense that the remaining $n - j$ coordinates t_i give the collar structure for $\partial_j M \times \mathcal{I}^{n-j}$ near x . We call such a compatible system of collar germs a *corner structure on M* . Henceforth all manifolds with corners will be implicitly equipped with a fixed but otherwise arbitrary corner structure.




FIGURE 2.3. A corner structure on M .

Let us choose a partial ordering of the boundary faces of a manifold M with corners such that the faces at all corners are ordered in a compatible way, i.e. the order is that induced by the order of the coordinates t_i in the collar structure. We will assume this order to be part of the structure of a manifold with corners.

Given manifolds with corners W and X , we consider embeddings of W into X which satisfy the property that each j -face of W is properly embedded into some k -face of X , where $k \geq j$.

2.3.2. *Liouville manifolds with boundary and corners.* We now give an inductive definition of the notion of a Liouville manifold with corners of order $k \leq \frac{1}{2} \dim X$, starting with the base case $k = 0$ which is just our previous definition.

Definition 2.23. A Liouville manifold with corners of order k consists of an exact symplectic manifold (X^{2n}, λ) with smooth corners of order $k \leq n$ such that the following properties hold.




FIGURE 2.4. A Liouville manifold with coners W near one of its faces P with nucleus N .

- (L1) The vector field Z which is ω -dual to λ is complete for $t \rightarrow \pm\infty$.
- (L2) There exists a compact domain with corners $W \subset X$ such that Z is outwards transverse to its *vertical boundary* $\partial_v W$, which is defined to be the closure of $\partial W \cap \text{int} X$, and such that the union of forward trajectories of Z starting at $\partial_v W$ is equal to $\partial_v W \cup (X \setminus W)$.
- (L3) For $r = 1, \dots, k$, each r -face P of the *horizontal boundary* $\partial_h W$, which is defined to be $\partial W \cap \partial X$, contains a Liouville submanifold with corners $N \subset P$, called the *nucleus* of P , such that the collar neighborhood of P in X is of the form $N \times T^*\mathcal{I}^r$, where P is the product of N with the cotangent fibre over $0 \in \mathcal{I}^r$, and in this neighborhood the form λ is the direct sum of a Liouville form on N and the canonical 1-form on $T^*\mathcal{I}^r$.

It follows from the definition that the Liouville field Z is tangent to every component of $\partial_k X$, $k = 1, \dots, n$. A Liouville manifold with corners has a *horizontal boundary* $\partial_h X = \partial X$ and an *ideal vertical boundary* $\partial_\infty X = (X \setminus \text{Skel}(X, \lambda))/\mathbb{R}^+$, where just as before the action of \mathbb{R}^+ is defined by the flow of Z and the skeleton $\text{Skel}(X, \lambda)$ is defined to be the attractor of the negative flow of Z .

We call $(W, \lambda|_W)$ as in (L2) a defining Liouville domain with corners for the Liouville manifold with corners (X, λ) . Note that the horizontal boundary of W is $\partial_h W = W \cap \partial_h X$ and its vertical boundary $\partial_v W$, the closure of $\partial W \setminus \partial_h W$, is a contact manifold with corners which is naturally contactomorphic to $\partial_\infty X$ under the projection $\pi_\infty : \partial_v W \rightarrow \partial_\infty X$. Just as before, the equivalence class of defining domains for Liouville manifold with corners is called the Liouville germ of (X, λ) and is denoted by (\mathcal{X}, λ) . Notions of isomorphism and homotopy of Liouville manifolds with corners are defined exactly the same way as without corners.

2.3.3. *Wc-manifolds.* We now introduce Weinstein structures for Liouville manifolds with corners, which we call *Wc-manifolds*. We restrict our discussion to the Morse-Bott category since this is the framework in which we will work throughout this paper.

Definition 2.24. A *Wc-manifold* is a Liouville manifold with corners (X, λ) whose Liouville vector field admits a Morse-Bott Lyapunov function $\phi : X \rightarrow \mathbb{R}$, i.e. satisfying (MB1) and (MB2) in the interior of X , and such that on the collars $N \times T^*\mathcal{I}^k$ of (L3) ϕ has the form $\phi_N + \sum_{i=1}^k u_i^2$ for ϕ_N a Morse-Bott Lyapunov function on $(N, \lambda|_N)$ and u_1, \dots, u_k the momentum coordinates on $T^*\mathcal{I}^k$.




FIGURE 2.5. The prototypical example of a Wc-manifold is the cotangent bundle of a manifold M with nonempty boundary, or even corners of any order.

Defining Liouville domains for Wc-manifolds will be called defining Wc-domains, and the equivalence class of such will be called a Wc-germ. Notions of Wc-isomorphism, embedding and homotopy are the same as the corresponding notions for Liouville manifolds with corners, but demanding that all objects in sight are of Weinstein type.

Example 2.25. Let M be a smooth n -dimensional manifold with corners. Then T^*M is a Liouville manifold with corners, where the structure (L3) is induced by a corner structure on M . We call the germ \mathcal{S}^*M of T^*M a *cotangent block*. The corresponding Lagrangian distribution tangent to cotangent fibers will be denoted by ν_M . We denote by λ_M and Z_M the corresponding Liouville form and the Liouville vector field. The notation S^*M will be used for the positively projectivized cotangent bundle of M , which is the ideal boundary of \mathcal{S}^*M . For Q a k -face of M we have $P = T^*M|_Q \subset \partial_k T^*M$ a k -face of T^*M and $N = T^*Q$ is the nucleus of P . The collar coordinates near Q yield the decomposition $\mathcal{O}_p P = N \times T^*\mathcal{I}^k$. A choice of Riemannian metric on M yields defining Liouville domain with corners $W = \{\|p\| \leq \varepsilon\}$ and a Morse-Bott Lyapunov function $\phi = \|p\|^2$.

The rest of this section can be equally discussed in the Liouville and Weinstein categories. We restrict the discussion only to the Weinstein case as this is the one needed for our further applications.

2.4. Wc-hypersurfaces.

2.4.1. *Weinstein hypersurfaces.* We now turn our attention to Weinstein hypersurfaces with corners, or Wc-hypersurfaces. They were introduced (without corners) in [E18]. These are special cases of the Liouville hypersurfaces introduced by Avdek in [A12]. This notion is similar to the “stops” of Sylvan, [Sy16] and the Liouville sectors of Ganatra-Pardon-Shende, [GPS17]. Related constructions are also considered in Ekholm-Lekili’s paper [EL17]. We first recall the definition without corners.

Definition 2.26. A Weinstein hypersurface in a Weinstein manifold (X^{2n}, λ) is a Weinstein embedding $(A^{2n-2}, \lambda' = \lambda|_A) \hookrightarrow (X \setminus \text{Skel}(X, \lambda), \lambda)$ such that $\pi_\infty|_A : A \rightarrow \partial_\infty X$ is an embedding.

Note that any Weinstein hypersurface is contained in a boundary ∂W of some defining domain W . Note also that the Reeb vector field of the contact form $\lambda|_{\partial W}$ is transverse to A . Note also that the boundary ∂A is a codimension 2 contact submanifold of $(\partial W, \lambda|_{\partial W})$. By the assumption that $(A, \lambda|_A)$ is Weinstein, the skeleton $\text{Skel}(A, \lambda|_A)$ can be decomposed as a union of isotropic submanifolds, which in the top dimension $n - 1$ are Legendrian for the contact structure $\xi = \ker(\lambda|_{\partial W})$.

Lemma 2.27. *The skeleton $\text{Skel}(A, \lambda|_A)$ depends only on the projection $\pi_\infty(A) \subset \partial_\infty X$.*

Proof. Indeed, the Liouville fields corresponding to the Liouville structures λ_A and $f\lambda|_A$ for a positive function f are proportional. In fact, as computed in Lemma 12.1 of [CE12] the form $d(f\lambda)|_A$ is symplectic if and only if $k := \inf(f + df(Z)) > 0$, where Z is the expanding field for $\lambda|_A$, and in that case the restriction of the form $f\lambda$ to A is automatically Liouville, with the expanding vector field for $f\lambda|_A$ equal to $\frac{1}{k}Z$. Therefore $\text{Skel}(A, \lambda|_A)$ only depends on the contact structure ξ_∞ . \square

Note that the space of functions $f > 0$ for which $f\lambda$ is Liouville (and hence in the considered case Weinstein) is convex, therefore contractible.

Definition 2.28. We say that two Weinstein hypersurfaces A_0, A_1 in a Weinstein manifold (X, λ) are related by a *translation* if $A_1 = Z^f(A_0)$ for some function $f : A_0 \rightarrow \mathbb{R}$.

Note that given a defining domain $W \subset X$, any Weinstein hypersurface A of X can be translated to a Weinstein hypersurface contained in W . However, it may not be possible to translate it to a Weinstein hypersurface contained in ∂W . Indeed, this is equivalent to the condition $\inf(f + df(Z)) > 0$ for f the conformal factor between the contact forms corresponding to A and ∂W .

Definition 2.29. A Weinstein hypersurface germ $\mathcal{A} \subset X \setminus \text{Skel}(X, \lambda)$ is an equivalence class of Weinstein hypersurfaces $(A, \lambda' = \lambda|_A) \subset (X \setminus \text{Skel}(X, \lambda), \lambda)$ for (A, λ') defining domains for a fixed Liouville manifold, where two embeddings are equivalent if they agree on a smaller defining domain, possibly after translation.

One can think of \mathcal{A} as living in the ideal boundary $\partial_\infty X$, or one can think of \mathcal{A} as the germ of a Weinstein embedding $\mathcal{A} \rightarrow \mathcal{X} \setminus \text{Skel}(\mathcal{X}, \lambda)$. The two viewpoints are equivalent and we will go back and forth between them.

Example 2.30. An important example of a Weinstein hypersurface is the *ribbon* of a Legendrian submanifold. Let Λ be a Legendrian submanifold in the contact boundary of a Liouville domain (W, λ) . Then it admits a Darboux neighborhood $U(\Lambda) \subset \partial W$ contactomorphic to $\{\|p\|^2 + z^2 \leq \varepsilon^2\} \subset J^1(\Lambda)$ for some Riemannian metric on Λ . The strip $\Sigma(\Lambda) := U(\Lambda) \cap \{z = 0\}$ is a Weinstein hypersurface symplectomorphic to the cotangent disk bundle of Λ , whose skeleton is precisely Λ . The space of all ribbon extensions of a given Legendrian Λ is contractible. An important example to keep in mind is the case where $\mathcal{X} = \mathcal{T}^*M$ for M a smooth manifold, $W = \{\|p\| \leq 1\}$ with respect to any Riemannian metric on M and Λ is the unit conormal to a co-oriented hypersurface $H \subset M$.




FIGURE 2.6. A Legendrian $\Lambda \subset S^*M$ and its ribbon $U(\Lambda)$.

2.4.2. Weinstein pairs.

Definition 2.31. A *Weinstein pair* (X, A, λ) consists of a Weinstein manifold (X, λ) together a Weinstein hypersurface $A \subset X \setminus \text{Skel}(X, \lambda)$.

We will also call $(X, \mathcal{A}, \lambda)$ a Weinstein pair when \mathcal{A} is the germ of a Weinstein hypersurface $\mathcal{A} \subset X \setminus \text{Skel}(X, \lambda)$.

Definition 2.32. Let (X, λ) be a Liouville manifold and $Y \subset X \setminus \text{Skel}(X, \lambda)$ any subset. The *Liouville cone* on Y is the saturation of A by the Liouville trajectories:

$$\text{Cone}(Y, \lambda) = \bigcup_{t \in \mathbb{R}} Z^{-t}(Y) \subset X.$$

The *positive Liouville cone* on A is the saturation of A by the backwards Liouville trajectories:

$$\text{Cone}^+(Y, \lambda) = \bigcup_{t \geq 0} Z^{-t}(Y) \subset X.$$

Note that if Y is compact, then $\text{Cone}^+(Y, \lambda)$ has compact closure while $\text{Cone}(Y, \lambda)$ does not. The skeleton of a Weinstein pair (X, A, λ) is defined as the compact subset:

$$\text{Skel}(X, A, \lambda) = \text{Skel}(X, \lambda) \cup \text{Cone}^+(\text{Skel}(A, \lambda|_A), \lambda).$$

Note that $\text{Skel}(X, A, \lambda) \subset W$ for any defining domain W such that $A \subset \partial W$, moreover $\partial W \cap \text{Skel}(X, A, \lambda) = \text{Skel}(A, \lambda|_A)$. The notion of a pair at the level of germs is defined as follows.

Definition 2.33. A Weinstein (germ) pair $(\mathcal{X}, \mathcal{A}, \lambda)$ consists of a Weinstein germ (\mathcal{X}, λ) together with a Weinstein hypersurface germ $\mathcal{A} \subset \mathcal{X} \setminus \text{Skel}(\mathcal{X}, \lambda)$.

Note that for any representative (W, λ) of (\mathcal{X}, λ) we may take a representative A for \mathcal{A} contained in W . The skeleton of a Weinstein pair $(\mathcal{X}, \mathcal{A}, \lambda)$ is defined as the non-compact subset:

$$\text{Skel}(X, \mathcal{A}, \lambda) = \text{Skel}(X, \lambda) \cup \text{Cone}(\text{Skel}(\mathcal{A}, \lambda|_{\mathcal{A}}), \lambda).$$

Equivalently in terms of germs, we can think of $\text{Skel}(X, \mathcal{A}, \lambda)$ as the equivalence class of the compact subsets $\text{Skel}(X, A, \lambda) \subset W$ for all defining domains $(W, \lambda|_W)$ and representatives A for \mathcal{A} such that $A \subset W$.

2.4.3. *Wc-hypersurfaces.* Finally, we extend the above to include boundary and corners.

Definition 2.34. A Wc-hypersurface in a Wc-manifold (X^{2n}, λ) is an equivalence class of embeddings of Wc-germs $(\mathcal{A}^{2n-2}, \lambda' = \lambda|_{\mathcal{A}}) \hookrightarrow (X \setminus \text{Skel}(X, \lambda), \lambda)$ such that $\pi_{\infty}|_{\mathcal{A}; \mathcal{A}} \rightarrow \partial X_{\infty}$ is an embedding, with the equivalence given by translation.

All the previous discussion carried over word by word to the Wc-setting. The notion of a Weinstein pair in the Wc-category is called a Wc-pair.

Definition 2.35. A Wc-pair $(\mathcal{X}, \mathcal{A}, \lambda)$ consists of a Wc-germ (\mathcal{X}, λ) together with a Wc-hypersurface germ $\mathcal{A} \subset \mathcal{X} \setminus \text{Skel}(\mathcal{X}, \lambda)$.

2.5. Conversion between Wc-hypersurfaces and face nuclei.

2.5.1. *Nucleus-to-hypersurface conversion.* We start with a Wc-manifold (X, λ) . Let P be a boundary face of X and $N \subset P$ its nucleus. Let U be a collar neighborhood of P in X as in the definition of a Wc-manifold. So U is of the form $N \times T^*\mathcal{I}$, with $\lambda = \lambda|_N + udt$ where t is the defining coordinate for P . Attach to $N \times T^*\mathcal{I}$ the half-infinite collar $N \times T^*(-\infty, 0]$ along their common intersection and on the union consider the Weinstein structure given by the direct sum of $\lambda|_N$ and the Weinstein normal form for Morse-Bott boundary of index $k = 0$, which agrees with $\lambda_N + udt$ on $N \times T^*\mathcal{I}$. The existence of a Morse-Bott Lyapunov function is immediate from the local model. Hence the result is a Wc-manifold (X^N, λ^N) . Let \mathcal{N} be the germ of N . For $\varepsilon > 0$ note that $\mathcal{N} \times \{t = -\varepsilon\} \subset \mathcal{N} \times T^*(-\infty, 0]$ is a Wc-hypersurface for λ^N . Any two differ by a translation and all are canonically identified with \mathcal{N} . We will denote this Wc-hypersurface by $\mathcal{A}^N \subset X^N \setminus \text{Skel}(X^N, \lambda^N)$.

Note that by construction we have $\text{Skel}(X^N, \mathcal{A}^N, \lambda^N) = \text{Skel}(X, \lambda)$. The operation $(X, N) \mapsto (X^N, \mathcal{A}^N)$ only depends on the contractible choice of a collar neighborhood and the contractible choice of the Weinstein normal form for Morse-Bott boundary. Hence it produces a Wc-germ $(\mathcal{X}^N, \lambda^N)$ which is well defined up to Wc-homotopy.

Definition 2.36. The Wc-pair $(\mathcal{X}^N, \mathcal{A}^N, \lambda^N)$ is called the *nucleus-to-hypersurface* conversion of the Wc-germ (\mathcal{X}, λ) and the boundary nucleus N .




FIGURE 2.7. Converting a boundary nucleus to a Wc-hypersurface.

2.5.2. *Underlying Weinstein manifold of a Wc-manifold.* Let X be a Wc-manifold and let P be a boundary face. The resulting of converting the nucleus N of P to a Wc-hypersurface is a Wc-manifold which has one less boundary face than X . We can continue this process inductively over all faces. After forgetting the resulting Wc-hypersurfaces, we end up with a Weinstein manifold without boundary which we denote \widehat{X} . It is clear that the Weinstein homotopy class of the Weinstein manifold \widehat{X} does not depend on the order in which the boundary faces are chosen. If one wishes to, one can further deform the Weinstein structure on \widehat{X} , which at this point is generalized Morse-Bott, so that it becomes Morse-Weinstein.

Definition 2.37. The Weinstein manifold \widehat{X} obtained from a Wc-manifold X by converting all of its boundary nuclei to Wc-hypersurfaces is called the *underlying Weinstein manifold*.

Example 2.38. the underlying Weinstein manifold of the Wc-manifold $(T^*[0, 1]^n, pdq)$ is Weinstein homotopic to $(\mathbb{R}^{2n}, pdq - qdp)$.

Remark 2.39. A Wc-manifold may be thought of as a Weinstein manifold with ‘stops’. From this viewpoint, taking the underlying Weinstein manifold of a Wc-manifold is simply removing the stops.

2.5.3. *Hypersurface-to-nucleus conversion.* The converse operation to the nucleus-to-hypersurface conversion is the *hypersurface-to-nucleus* conversion, which we discuss next. Here and below we denote $U_{a,b} = \{0 < u \leq a, -b < t < b\} \subset T^*\mathbb{R}$.

Lemma 2.40. *The inclusion $(A, \lambda' = \lambda|_A) \hookrightarrow (X, \lambda)$ of a Wc-hypersurface extends, uniquely up to contractible choice, to a Liouville embedding $(A \times U_{1,\varepsilon}, \lambda' + udt) \rightarrow (X, \lambda)$, where A is identified with $A \times (u = 1, t = 0) \subset A \times U_{1,\varepsilon}$.*

Proof. Let $W \subset X$ be a defining domain such that $A \subset \partial W$ and let $A \times (-\varepsilon, \varepsilon) \subset \partial W$ be its contact surrounding, as in [E18], so that the restriction of λ to this neighborhood is of the form $\pi^* \lambda|_A + dt$, where $\pi : A \times (-\varepsilon, \varepsilon) \rightarrow A$ is the projection and $t \in (-\varepsilon, \varepsilon)$. For every $x \in X \setminus \text{Skel}(X, \lambda)$, let x^s denote the backwards flow of Z_X starting at x , where $s \in (-\infty, 0]$. For every $y \in A \times (-\varepsilon, \varepsilon)$, let $y^s \in U_{1,\varepsilon}$ denote the backwards flow of $Z_A + u \frac{\partial}{\partial u}$ starting at y , where $s \in (-\infty, 0]$. We extend the contact embedding $A \times (-\varepsilon, \varepsilon) \rightarrow \partial W$ which sends $y \mapsto x$ to a Liouville embedding $A \times U_{1,\varepsilon} \rightarrow W$ by sending $y^s \mapsto x^s$, so that by construction $Z_A + u \frac{\partial}{\partial u}$ is identified with Z_X . It is clear that the embedding is determined by the contact surrounding and its parametrization, which are unique up to contractible choice. \square

We now explain the *hypersurface-to-nucleus* conversion. Let (X, λ) be a Wc-manifold and let $\mathcal{A} \subset X \setminus \text{Skel}(X, \lambda)$ be a Wc-hypersurface. Fix defining Wc-domains A and W for \mathcal{A} and X respectively so that we have an embedding $A \rightarrow \partial W$ which extends to an embedding $A \times U_{1,\varepsilon} \rightarrow W$ as in Lemma 2.40. Consider the exact symplectic manifold which is the result of gluing W and $A \times U_{3,\varepsilon}$ along $A \times U_{1,\varepsilon}$. Consider the Weinstein structure on $U_{3,\varepsilon}$, thought of as the ε -neighborhood of the zero section in $T^*(0, 3)$, which is the restriction of the Weinstein normal form for Morse-Bott boundary of index $k = 1$ translated two units in the u direction, so that the critical set is $\{t = 0, u \geq 2\}$. The restriction of this structure to $U_{1,\varepsilon}$ is udt , hence we obtain a patched up structure on the union of W and $A \times U_{3,\varepsilon}$. Moreover, by inspection the new Liouville field points outwards at the boundary. So we may replace $U_{3,\varepsilon}$ with a smoothing $U_{3,\mu} = \{-\varepsilon < t < \varepsilon, 0 < u < \min(3, \mu(|t|))\}$ for $\mu : (0, \varepsilon) \rightarrow [1, \infty)$ a function satisfying

- (i) $\mu(x) \rightarrow \infty$ as $x \rightarrow 0$
- (ii) $\mu = 1$ on $(\varepsilon/2, \varepsilon)$
- (iii) $\mu' < 0$ on $(0, \varepsilon/2)$

and the Liouville field is still outwards pointing for the smoothed boundary. Thus we obtain a defining domain for a Wc-manifold (X^A, λ^A) . Indeed the only thing that needs to be checked is the existence of a Morse-Bott Lyapunov function, which is clear from the local models since without loss of generality we can start with a defining domain W for X such that ∂W is a regular level set of the Morse-Bott Lyapunov function ϕ for X . Note that $A \times \{t = 0, u = 3\}$ is the nucleus of the new boundary face of X^A .

By construction we have $\text{Skel}(X^A, \lambda^A) = \text{Skel}(X, A, \lambda)$, or more precisely

$$\text{Skel}(X^A, \lambda^A) = \text{Skel}(X, A, \lambda) \cup (\text{Skel}(A, \lambda|_A) \times [1, 2]).$$

In other words, the backwards λ -Liouville cone on A has a trivial collar attached to it, which does not change the topology of $\text{Skel}(X, A, \lambda)$. Note that the space of possible smoothings μ is convex, hence contractible. The choice of Morse-Bott Weinstein normal form is also contractible. Similarly the choice of collars used in the construction is contractible. Hence

we get a Wc-germ $(\mathcal{X}^A, \lambda^A)$ which is well defined up to Wc-homotopy. Finally, note that different representatives A for \mathcal{A} will also result in homotopic Wc-manifolds (X^A, λ^A) . Hence the conversion operation only depends on the Wc-hypersurface germ \mathcal{A} up to Wc-homotopy.

Definition 2.41. The Wc-germ $(\mathcal{X}^{\mathcal{A}}, \lambda^{\mathcal{A}})$ is called the *hypersurface-to-nucleus* conversion of the Wc-germ X and the Wc-hypersurface \mathcal{A} .




FIGURE 2.8. Converting a boundary nucleus to a Wc-hypersurface.

2.5.4. *Cancellation of conversions.* The hypersurface-to-nucleus conversion is inverse to the nucleus-to-hypersurface conversion at the level of Weinstein homotopy classes, as follows from a parametric Smale cancellation of Morse-Bott index 0 and 1 critical points, the existence of which is immediate from the local models. This implies the following obvious but important lemma.

Lemma 2.42. *Let $(X^{\mathcal{A}}, \lambda^{\mathcal{A}})$ be the result of hypersurface-to-nucleus conversion of a Wc-hypersurface $\mathcal{A} \subset X \setminus \text{Skel}(X, \lambda)$. Then the underlying Weinstein manifolds of (X, λ) and $(X^{\mathcal{A}}, \lambda^{\mathcal{A}})$ are Wc-homotopic.*

2.6. Gluing and splitting.

2.6.1. *Horizontal gluing.* We will now describe the basic gluing operation for Wc-manifolds. Given two Wc-manifolds (X_0, λ_0) and (X_1, λ_1) , we define their *horizontal gluing* as follows. Pick some *non-adjacent* boundary faces P_1, \dots, P_l of X_1 and Q_1, \dots, Q_l of X_0 , and for each j suppose we are given a Liouville isomorphism $\phi_j : N_j \rightarrow M_j$, where N_j and M_j are the nuclei or P_j and Q_j respectively. We have collar neighborhoods $N_j \times T^*\mathcal{I} \subset X_1$ and $M_j \times T^*\mathcal{I} \subset X_2$ and we extend ϕ_j to $\Phi_j : P_j \rightarrow Q_j$ as $\Phi_j = \phi_j \times \text{id}_{\mathbb{R}} : N_j \times \mathbb{R} \rightarrow M_j \times \mathbb{R}$.

Definition 2.43. The horizontal gluing of $(\mathcal{X}_1, \lambda_1)$ and $(\mathcal{X}_2, \lambda_2)$ along $\{\phi_j\}_j$ is defined to be the Wc-germ of the Wc-manifold $X_1 \cup X_2$ where we identify $P_j \sim Q_j$ via Φ_j .

Example 2.44. Suppose M, N are two manifolds with boundary and we have a diffeomorphism $\psi : \partial N \rightarrow \partial M$. Then ψ lifts to a Liouville isomorphism ϕ between $T^*(\partial N)$ and $T^*(\partial M)$ and we can perform the horizontal gluing $T^*N \cup_{\phi} T^*M$ which is the same as $T^*(N \cup_{\psi} M)$ for $N \cup_{\psi} M$ the smooth gluing of N and M given by ψ .




FIGURE 2.9. Horizontal gluing of two Wc-manifolds.

Note that the skeleton of the horizontal gluing of X_1 and X_2 is equal to the union of the skeleta of X_1 and X_2 .

2.6.2. *Vertical gluing.* We can also glue two Wc-manifolds X_1 and X_2 in a different way.

Definition 2.45. Suppose we have a Wc-hypersurface \mathcal{A} in \mathcal{X}_1 and a Liouville isomorphism $\phi : \mathcal{A} \rightarrow \mathcal{N}$ for \mathcal{N} the germ of the nucleus N of a boundary face of X_2 . The *vertical gluing* of \mathcal{X}_1 and \mathcal{X}_2 is defined to be the composition of two steps: a conversion a Wc-hypersurface into a boundary hypersurface, see Section 2.5 above, followed by a horizontal gluing.

Indeed, if we first convert \mathcal{A} into a boundary nucleus, we can then apply the horizontal gluing determined by ϕ as described above. The resulting Wc-germ is well defined up to Wc-isomorphism. Note that the skeleton of the vertical gluing is equal to the union of the skeleton of the pair $(\mathcal{X}_1, \mathcal{A})$ and the skeleton of \mathcal{X}_2 . We record for future reference the following slight strengthening of this observation, where we denote by λ^P the Liouville form $\lambda - d(ut)$ for t the defining coordinate of a boundary face P .

Lemma 2.46. *Let $(\mathcal{X}_i, \lambda_i)$, $i = 1, 2$ be two Wc-germs, $\mathcal{A} \subset \mathcal{X}_1 \setminus \text{Skel}(\mathcal{X}_2, \lambda_2)$ a Wc-hypersurface, N the nucleus of a boundary face P of X_1 and \mathcal{N} its germ, $\phi : \mathcal{A} \rightarrow \mathcal{N}$ a Liouville isomorphism and (\mathcal{Y}, λ) the result of vertically gluing \mathcal{X}_1 to \mathcal{X}_2 along ϕ . Then there exists a symplectic embedding of germs $\Psi : \text{int}\mathcal{X}_1 \rightarrow \mathcal{Y}$ such that the following properties hold, where we denote by $\iota : \text{int}\mathcal{X}_2 \rightarrow \mathcal{Y}$ the restriction of the inclusion to the interior of \mathcal{X}_2 .*

- (i) $\iota(\text{Skel}(\mathcal{X}_2, \lambda_2)) \cup \Psi(\text{Skel}(\mathcal{X}_1, \lambda_1) \cap \text{int}\mathcal{X}_1) = \text{Skel}(\mathcal{Y}, \lambda)$
- (ii) $\iota(\text{Skel}(\mathcal{X}_2, \lambda_2)) \cap \Psi(\text{int}\mathcal{X}_1) = \emptyset$.




FIGURE 2.10. Vertical gluing is the composition of two steps: a Wc-hypersurface to boundary nucleus conversion and a horizontal gluing.

$$(iii) \Psi^{-1}(\iota(\text{int}\mathcal{X}_2)) = \mathcal{O}p \mathcal{N} \setminus \partial\mathcal{X}_1 \subset \mathcal{X}_1 \text{ and } (\Psi)^*(\iota)_*\lambda_2 = \lambda_1^P.$$

Proof. We recall from the gluing construction that we attached $\mathcal{N} \times (-\varepsilon, \varepsilon)$ to the top of a cylinder $\mathcal{A} \times U_{3,\varepsilon} \subset Y$. Since a neighborhood $\mathcal{O}p N$ of N in X_1 is symplectomorphic to $N \times T_\varepsilon^*[0, \delta)$, we can map $\mathcal{N} \times T_\varepsilon^*(0, \delta)$ symplectomorphically to $\mathcal{A} \times U_{3,\varepsilon} \cup \mathcal{N} \times T_\varepsilon^*[0, \delta)$, relative to $\mathcal{N} \times T_\varepsilon^*(\delta/2, \delta)$ say, so that the symplectomorphism extends to the rest of $\text{int}\mathcal{X}_1$ as the identity. Moreover, the symplectomorphism can be obtained from a reparametrization of the t coordinate, hence the Liouville forms pull back as desired. \square

2.6.3. *Splitting.* The converse operation to gluing is splitting.

Definition 2.47. Given a Wc-manifold (X, λ) , a regularly embedded hypersurface $(H, \partial H) \subset (X, \partial X)$ with corners called *splitting* for X if it satisfies the following conditions:

- (i) H divides X into two parts, $X = X_+ \cup X_-$ and $X_- \cap X_+ = H$, which are manifolds with corners.
- (ii) the Liouville vector field Z of X is tangent to H .
- (iii) there exists a hypersurface with corners $(S, \partial S) \subset (H, \partial H)$ tangent to Z .
- (iv) $(S, \lambda|_S)$ is a Wc-manifold and there exists a retraction $\pi : H \rightarrow S$ such that $\lambda|_H = \pi^*(\lambda|_S)$.

Remark 2.48. That H is regularly embedded implies in particular that $\mathcal{O}p H = H \times (-\varepsilon, \varepsilon)$.

We will refer to S as the *soul* of the splitting hypersurface H and denote it by $\text{Soul}(H)$. Note that $\text{Skel}(S, \lambda|_S) = \text{Skel}(X, \lambda) \cap H$.

Example 2.49. Let M be a manifold with corners and N a regularly embedded codimension 1 submanifold with corners, so that $\mathcal{O}_p N = N \times (-\varepsilon, \varepsilon)$. Assume that N divides M into two parts: M_+ and M_- , so that $M_+ \cup M_- = M$. Then $H = T^*M|_N$ is a splitting hypersurface. Given a decomposition of the tubular neighborhood of N , $\mathcal{O}_p(N) = N \times (-\varepsilon, \varepsilon)$, we have $S := T^*N \times 0 \subset T^*M = T^*N \times T^*(-\varepsilon, \varepsilon)$ the soul of H .

The splitting hypersurface S can be used to split X into two Wc-manifolds in two ways, *horizontal* and *vertical*. For the *horizontal splitting* we view X_\pm as Wc-manifolds with an additional component H of $\partial_1 X_\pm$. The Wc-manifold S serves as the nucleus of the additional boundary component H . The *vertical splitting* consists of a horizontal splitting followed by a nucleus-to-hypersurface conversion operation, transforming the nucleus S into a Wc-hypersurface.

2.6.4. Partial horizontal gluing. One more basic operation that we will need is the creation of corners and its associated partial horizontal gluing. Let X be a Wc-manifold, P a boundary face of X and N the nucleus of P . Let $\Sigma \subset N$ be a splitting hypersurface with soul T . Then there is a deformation of the Wc-structure on X which creates an additional corner along Σ with its nucleus T .

More precisely, take a collar neighborhood $\mathcal{O}_p P = N \times T^*\mathcal{I}$, so that $\lambda = \lambda|_N + udt$ and extend it to $N \times T^*(-\varepsilon, \varepsilon)$ with the same Liouville form. Let $\Sigma \times (-\delta, \delta) \subset N$ be a tubular neighborhood of Σ in N and consider a smooth function $\mu : (0, \delta) \rightarrow [0, \infty)$ satisfying the following properties:

- (i) $\mu(x) \rightarrow \infty$ as $x \rightarrow 0$.
- (ii) $\mu(x) = 0$ on $[\delta/2, \delta)$
- (iii) $\mu'(x) < 0$ on $(0, \delta/2)$.

Set $\eta(x) = \min(\mu(x), \varepsilon/2)$ and consider the subset $\Omega \subset (\Sigma \times (-\delta, \delta)) \times T^*(-\varepsilon, \varepsilon)$ consisting of points (σ, s, t, u) such that $\eta(s) + t \geq 0$. Then the union of Ω and X along P is a new Wc-manifold which has a new corner along a parallel copy of Σ .




FIGURE 2.11. A creation of additional corners on \mathcal{T}^*M for M a closed manifold with corners is induced from a creation of additional corners on M .

Next, consider two Wc-manifolds X_0, X_1 with boundary faces P_0, P_1 and respective nuclei N_0, N_1 . Let Σ be a dividing hypersurface for N_0 with its soul T , which splits N_0 into Wc-manifolds N'_0 and N''_0 , which share a common face Σ with the nucleus T . Suppose there is a Liouville isomorphism $\phi : N_1 \rightarrow N'_0$.

Definition 2.50. The partial horizontal gluing of \mathcal{X}_1 to \mathcal{X}_0 using ϕ is the Wc-germ obtained from the following two operations:

- (a) Creating a corner along Σ , i.e. adding Σ to $\partial_2 X_0$, and replacing the face N_0 with two faces N'_0, N''_0 ;
- (b) The horizontal gluing X_0 and X_1 along ϕ .

As before, the partial horizontal gluing is well defined up to Wc-homotopy.

2.7. Wc-buildings.

2.7.1. Iterated gluing. We now introduce the iterated vertical gluing of Wc-manifolds, which will be a convenient notion for the constructions in this paper. Consider Wc-germs $(\mathcal{X}_1, \lambda_1), \dots, (\mathcal{X}_k, \lambda_k)$. We inductively define a k -level *Wc-building*, which we will denote by $(\mathcal{X}_k \rightarrow \mathcal{X}_{k-1} \rightarrow \dots \rightarrow \mathcal{X}_1)$, as the following presentation of a Wc-germ.

Definition 2.51. A 1-level Wc-building is $(\mathcal{X}_1, \lambda_1)$ itself. Suppose that the Wc-building

$$(\mathcal{Y}, \mu) := (\mathcal{X}_k \rightarrow \mathcal{X}_{k-1} \cdots \rightarrow \mathcal{X}_2)$$

is already defined. Consider a nucleus $(\mathcal{N}, \mu' := \mu|_{\mathcal{N}})$ of one of the boundary faces \mathcal{P} of (\mathcal{Y}, μ) and let ϕ be an embedding \mathcal{N} as a Wc-hypersurface in $\mathcal{X}_1 \setminus \text{Skel}(\mathcal{X}_1, \lambda_1)$. The building

$$(\mathcal{X}_k \rightarrow \mathcal{X}_{k-1} \rightarrow \dots \rightarrow \mathcal{X}_2 \rightarrow \mathcal{X}_1) := (\mathcal{X}_k \rightarrow \mathcal{X}_{k-1} \rightarrow \dots \rightarrow \mathcal{X}_2) \rightarrow \mathcal{X}_1$$

is by definition the result of a vertical gluing of (\mathcal{Y}, μ) and $(\mathcal{X}_1, \lambda_1)$ along ϕ .




FIGURE 2.12. A Wc-building.

So a Wc-building is a Wc-germ \mathcal{Y} equipped with a choice of presentation of its Wc-deformation class as the iterated vertical gluing $\mathcal{Y} = (\mathcal{X}_1 \rightarrow \dots \rightarrow \mathcal{X}_k)$.

2.7.2. *Distinguished cover of a Wc-building.* A Wc-building come equipped with a symplectic cover by the Wc-manifolds it is built out of. The relevant properties of this cover are summarized in the following lemma. Recall that given a Wc-manifold (X, λ) the Liouville form λ restricted to a neighborhood of a face P has the form $\lambda := \lambda|_N + udt$, where N is the nucleus of P and t is the defining coordinate for the face P . We recall the notation λ^P for the form $\lambda|_N - tdu = \lambda - d(ut)$ on $\mathcal{O}p N$.

Lemma 2.52. *Suppose (\mathcal{X}, λ) is presented as a k -level building $(\mathcal{X}_k \rightarrow \mathcal{X}_{k-1} \rightarrow \cdots \rightarrow \mathcal{X}_1)$. Then there exist symplectic embeddings $\Phi_j : \text{int}\mathcal{X}_j \rightarrow \mathcal{X}$, $j = 1, \dots, k$, such that*

- (i) $\bigcup_{j=1}^k (\Phi_j(\text{Skel}(\mathcal{X}_j, \lambda_j) \cap \text{int}\mathcal{X}_j)) = \text{Skel}(\mathcal{X}, \lambda)$;
- (ii) $\Phi_j(\text{int}\mathcal{X}_j) \cap \Phi_i(\text{Skel}(\mathcal{X}_i, \lambda_i) \cap \text{int}\mathcal{X}_i) = \emptyset$ if $j > i$;
- (iii) If $\Phi_j(\text{int}\mathcal{X}_j) \cap \Phi_i(\text{int}\mathcal{X}_i) \neq \emptyset$ for $j > i$ then there exists a nucleus \mathcal{N} of a boundary face of \mathcal{X}_j such that $\Phi_j^{-1}(\Phi_i(\text{int}\mathcal{X}_i)) = \mathcal{O}p \mathcal{N} \cap \text{int}\mathcal{X}_j$ and $\Phi_j^*(\Phi_i)_*\lambda_i = \lambda_j^P$.




FIGURE 2.13. Presenting a Wc-building as the union of Wc-manifolds.

Proof. We inductively construct the maps $\Phi_{i,j} : \mathcal{X}_i \rightarrow (\mathcal{X}_k \rightarrow \cdots \rightarrow \mathcal{X}_j)$, $i \geq j$, as follows, at the end of the inductive process we will set $\Phi_i = \Phi_{i,1}$. The map $\Phi_{k,k}$ is just the inclusion on \mathcal{X}_1 into \mathcal{X} . Suppose we have defined maps $\Phi_{i,j}$ for $i \geq j$. Recall that $\mathcal{Y} = (\mathcal{X}_1 \rightarrow \cdots \rightarrow \mathcal{X}_j)$ is vertically glued to \mathcal{X}_{j-1} by a Liouville isomorphism between the nucleus \mathcal{N} of a boundary face \mathcal{P} of \mathcal{Y} and a Wc-hypersurface \mathcal{A} in \mathcal{X}_{j-1} . Let $\Psi : \mathcal{Y} \rightarrow (\mathcal{X}_1 \rightarrow \cdots \rightarrow \mathcal{X}_j \rightarrow \mathcal{X}_{j-1})$ be the symplectic embedding from Lemma 2.46, set $\Phi_{i,j-1} = \Psi \circ \Phi_{i,j}$ and set $\Phi_{j-1,j-1}$ to be the restriction of the inclusion $\mathcal{X}_{j-1} \hookrightarrow (\mathcal{X}_1 \rightarrow \cdots \rightarrow \mathcal{X}_j \rightarrow \mathcal{X}_{j-1})$ to the interior of \mathcal{X}_{j-1} . The desired properties for the resulting Φ_i follow from the properties stated in Lemma 2.46. \square

Remark 2.53. Note that on the overlap $\Phi_{i_1}(\mathcal{X}_{i_1}) \cap \cdots \cap \Phi_{i_k}(\mathcal{X}_{i_k}) \subset \mathcal{X}$ the pushed-forward Liouville fields Z_{i_1}, \dots, Z_{i_k} coming from $\mathcal{X}_{i_1}, \dots, \mathcal{X}_{i_k}$ all commute with each other.

2.7.3. *Cotangent buildings.* Finally, we introduce a particular class of Wc-buildings which will be particularly relevant for our purposes.

Definition 2.54. A Wc-building $(\mathcal{X}_k \rightarrow \cdots \rightarrow \mathcal{X}_1)$ for which each Wc-germ consists of a cotangent block \mathcal{T}^*M_j is called a *cotangent building*.

We will later show that any finite type Weinstein manifold can be realized as the underlying Weinstein manifold of a cotangent building.

2.8. Legendrian submanifolds.

2.8.1. Legendrians in cotangent bundles. Consider first a cotangent bundle T^*M , where M an n -dimensional compact manifold with corners. Let Λ be an $(n - 1)$ -dimensional manifold with corners.

Definition 2.55. An embedding $\Lambda \rightarrow T^*M \setminus M$ is called *Legendrian* if its projection to S^*M is a Legendrian embedding.

With every Legendrian $\Lambda \subset T^*M$ we associate its Lagrangian Liouville cone $L := \text{Cone}(\Lambda, pdq)$ formed by forward and backwards trajectories of the Liouville field Z . Note that L is embedded.

The splitting $P \times \mathcal{I}^k$ near a k -face P of M provide canonical splittings

$$(1) \quad T^*M|_{\mathcal{O}_p P} = T^*P \times T^*\mathcal{I}^k$$

We will call a Legendrian embedding $\phi : L \rightarrow \mathcal{T}^*M \setminus M$ *adapted* to the block T^*M if $f(\mathcal{O}_p \partial L) \subset \mathcal{O}_p \partial M$ and for each stratum $P \subset \partial_k M$ we have $L \cap \mathcal{O}_p P = L_k \times \mathcal{I}^k$ for a Legendrian $L_k \subset \mathcal{T}^*P$.

By conical Lagrangians $L \subset T^*M \setminus M$ we will always mean Lagrangian Liouville cones over Legendrians.

Denote by $\pi : T^*M \rightarrow M$ the cotangent bundle projection.

Definition 2.56. An adapted Legendrian Λ is called *regular* if $\pi|_{\Lambda} : \Lambda \rightarrow M$ is an immersion with transverse self-intersections.




FIGURE 2.14. The Legendrian on the left is regular, while the one on the right is not.

2.8.2. *Legendrians in Wc-manifolds.* Let (X, λ) be a $2n$ -dimensional Wc-manifold.

Definition 2.57. An $(n-1)$ -dimensional submanifold $\Lambda \subset X \setminus \text{Skel}(X, \lambda)$ is called *Legendrian* if it projects to $\partial_\infty X$ as an embedded Legendrian submanifold of $\partial_\infty X$. Its Liouville cone $L = \text{Cone}(\Lambda, \lambda)$ is called a *conical Lagrangian*.

We say that a Legendrian Λ is *adapted* to the Wc-structure of (X, λ) if $\partial_k \Lambda \subset \partial_k X$ and moreover in a neighborhood of any corner $P \subset \partial_k X$ we have $\Lambda = \Lambda_N \times \mathcal{I}^k \subset P \times T^* \mathcal{I}^k$, where Λ_N is a Legendrian in the nucleus N of the corner P .




FIGURE 2.15. An adapted Legendrian Λ in a Wc-manifold W .

Example 2.58. Let M be a manifold and $N \subset M$ a co-oriented hypersurface. Then any submanifold $\Lambda \subset T^*M \setminus M$ which 1-1 projects to the conormal lift of N in S^*M (and is never tangent to the Liouville flow) is Legendrian. If N is adapted for the corner structure of M , then one can construct a Legendrian lift Λ which is adapted for the Wc-structure of T^*M .

We can also think of Legendrians in T^*M as equivalence classes of Legendrian embeddings into defining domains for T^*M , where the equivalence is given by translation along the Liouville flow. This gives us the notion of a Legendrian in a cotangent block \mathcal{F}^*M . Similarly, we have the notion of a Legendrian in a Wc-germ \mathcal{X} . For a Legendrian Λ in a defining domain $(W, \lambda|_W)$ for (\mathcal{X}, λ) , we only take its positive Liouville cone $\text{Cone}^+(\Lambda, \lambda)$, which is contained in $W \setminus \text{Skel}(W, \lambda)$. One can think of the equivalence class of all positive Liouville cones over all Legendrians equivalent to Λ as the whole Liouville cone $\text{Cone}(\Lambda, \lambda) \subset X \setminus \text{Skel}(X, \lambda)$.

2.9. Gluing and splitting of Legendrians.

2.9.1. Gluing Legendrians.

Lemma 2.59. Let (\mathcal{X}, λ) be a Wc-germ, $\Lambda \subset \mathcal{X} \setminus (\text{Skel}(\mathcal{X}, \lambda))$ an adapted Legendrian and L the λ -Liouville cone over Λ . Let $(\mathcal{X}^N, \lambda^N)$ be the result of a nucleus-to-hypersurface conversion of the nucleus N of one of the boundary faces of X . Then $\Lambda \subset \mathcal{X}^N$ remains Legendrian and its λ^N -Liouville cone coincides with L .

Proof. In a neighborhood of the face P of X whose nucleus N we convert to a boundary face we have $\mathcal{O}_p P = T^*N \times T^*\mathcal{I}$ and $\Lambda = \Lambda_N \times \mathcal{I}$. The conversion changes $\lambda = \lambda_N + udt$ only by the differential of a function of (u, t) , moreover the resulting Liouville field on $T^*\mathcal{I}$ is tangent to \mathcal{I} (and outwards pointing at the boundary). Hence Λ remains Legendrian and its Liouville cone is unchanged. \square

Lemma 2.60. *Let $\mathcal{X}_0, \mathcal{X}_1$ be two Wc-germs, P_0, P_1 two of their boundary faces and N_0, N_1 the nuclei of P_0, P_1 . Let $\Lambda_0 \subset \mathcal{X}_0 \setminus \text{Skel}(\mathcal{X}_0, \lambda_0)$ and $\Lambda_1 \subset \mathcal{X}_1 \setminus \text{Skel}(\mathcal{X}_1, \lambda_1)$ be two adapted Legendrians. Denote $\Lambda'_i := \Lambda_i \cap N_i$, $i = 0, 1$. Let $\phi : N_0 \rightarrow N_1$ be a Liouville isomorphism such that $\phi(\Lambda'_0) = \Lambda'_1$. Then the horizontally glued Wc-manifold $\mathcal{X}_{01} = \mathcal{X}_0 \cup_\phi \mathcal{X}_1$ contains a glued Legendrian*

$$\Lambda_{01} = \Lambda_0 \cup_{\Lambda'_1 = \phi|_{\Lambda'_0}} \Lambda_1.$$

Proof. This is obvious by definition of adapted. \square

2.9.2. *Splitting Legendrians.* Let X be a Wc-manifold and $P \subset X$ a splitting hypersurface with the soul S . A Legendrian $\Lambda \subset X \subset \text{Skel}(X, \lambda)$ is called in *in splitting position* if Λ is transverse to P and $\Lambda \cap P \subset S$.

Lemma 2.61. *Let X be a Wc-manifold, $P \subset X$ a splitting hypersurface and consider a Legendrian $\Lambda \subset X \setminus \text{Skel}(X, \lambda)$ which is in splitting position. Then under a horizontal splitting of X we can arrange it so that Λ splits into Legendrians with corners $\Lambda_\pm \subset X_\pm \setminus \text{Skel}(X_\pm, \lambda_\pm)$.*

Proof. It suffices to take a neighborhood of P in \mathcal{X} so that Λ is of product form with respect to the neighborhood, which is clearly possible. \square

Let us recall that given a contact manifold (M, ξ) a hypersurface $N \subset M$ is called *convex* if it admits a tranverse to it contact vector field v . Consider a split a tubular neighborhood $U_\varepsilon := N \times (-\varepsilon, \varepsilon)$ of a contact hypersurface N such that the contact vector field v coincides with $\frac{\partial}{\partial t}$, where t is the coordinate corresponding to the second factor. The contact structure ξ on U_ε can be then defined by a contact form $\alpha = fdt + \lambda$, where $f : N \rightarrow \mathbb{R}$ is a smooth function and λ is a 1-form on N . The set $Q = \{f = 0\} = \{x \in N; v(x) \in \xi\}$ is called the *dividing set*. It is always a regular level set of the function f . Denote $N_+ := \{f > 0\}, N_- = \{f < 0\}$. We also denote

$$\widehat{Q} := Q \times (-\varepsilon, \varepsilon) = \bigcup_{t \in (-\varepsilon, \varepsilon)} v^t(Q)$$

and call it the *dividing wall*. The contact condition implies that $\beta := \lambda|_Q$ is a contact form on Q and that the form $\lambda_f := \frac{1}{f}\lambda$ is Liouville on $N \setminus Q = N_+ \cup N_-$. The corresponding Liouville fields Z_\pm are complete of both halves N_+ and N_- and for any point $x \in N_\pm \setminus \text{Skel}(N_\pm)$ there exists a limit $\lim_{s \rightarrow \infty} Z_\pm^s(x) \in Q$. We say that the pair (N, v) , where N is a convex hypersurface and v a transverse vector field, is of *Weinstein type*, if (N_\pm, λ_f) is Weinstein.

Lemma 2.62. *Let (M, ξ) be a contact manifold, $N \subset (M, \xi)$ a Weinstein type convex hypersurface and $\Lambda \subset (M, \xi)$ a Legendrian transverse to N . Then there exists a contact isotopy $\phi_s : M \rightarrow M$, $t \in [0, 1]$ such that*

- (i) ϕ_s fixes Q ;
- (ii) ϕ_s leaves N invariant;
- (iii) there exists $\varepsilon > 0$ such that $\phi_1(\Lambda \cap U_\varepsilon) \subset Q_\varepsilon$;
- (iv) ϕ_s is supported in a neighborhood $U' \ni U_\varepsilon$.

Proof. Note that a contact vector field $Y := t \frac{\partial}{\partial t} + Z_\pm$ on $N_\pm \times (-\varepsilon, \varepsilon)$ smoothly extends to $N \times (-\varepsilon, \varepsilon)$ as equal to $t \frac{\partial}{\partial t}$ on $Q \times (-\varepsilon, \varepsilon)$. Let us choose ε sufficiently small so that the coordinate $t|_{\Lambda \cap U_\varepsilon}$ has no critical points. Denote $\Gamma := \Lambda \cap N$. A general position argument allows us to C^∞ -perturb Λ so that $\Gamma \cap (\text{Skel}(N_+) \cup \text{Skel}(N_-)) = \emptyset$ and the projection of $\pi : \Gamma \rightarrow Q$ along Liouville trajectories of N_\pm is an embedding. Note that $\lambda|_\Gamma = 0$, and hence $\pi(\Gamma) \subset Q$ is Legendrian for the contact structure $\ker \beta$ on Q . Let us parameterize $\Lambda \cap U_\varepsilon$ by an embedding $h_0 : \Gamma \times (-\varepsilon, \varepsilon) \rightarrow \Lambda \cap U_\varepsilon$ such that $h_0(\Gamma \times t) \subset N \times t$. Define a Legendrian isotopy $h_s : \Gamma \times (-\varepsilon, \varepsilon) \rightarrow \Lambda \cap U_\varepsilon$, $s \in [0, \infty)$ by the formula

$$h_s(x, t) = Y^s(h_0(x, te^{-s})).$$

Let us observe that $h_s(x, t) = (g_{s,t}(x), t)$, with $g_{s,0}(x)$ flowing a λ -isotropic submanifold Γ to $\pi(\Gamma)$ along trajectories of Z_\pm . The flow is fixed on Q . Let us analyze the isotopy in a neighborhood of Q . The form λ_f on $\mathcal{O}p Q$ can be written as $\frac{1}{u}\beta + dt$ for the coordinate $u = f$ on $\mathcal{O}p Q$, and hence, $Y = t \frac{\partial}{\partial t} - u \frac{\partial}{\partial u}$. The flow Y^t is given by $Y^s(x, u, t) = (x, e^{-s}u, e^st)$. Assuming that T is large enough so that $h_T(\Gamma \times (-\varepsilon, \varepsilon)) \subset \mathcal{O}p Q \times (-\varepsilon, \varepsilon)$ we get $h_{T+s}(x, u, t) = (x, e^{-s}x, t) \rightarrow (x, 0, t)$ as $s \rightarrow \infty$. Hence by reparameterizing the isotopy to the interval $[0, 1]$ and extending it to the end-point 1 as $(x, 0, 1)$. we get a smooth Legendrian isotopy $\phi_s : \Gamma \times (-\varepsilon, \varepsilon)$ such that $\phi_1(\Gamma \times (-\varepsilon, \varepsilon)) \subset Q \times (-\varepsilon, \varepsilon)$. The isotopy ϕ_s can be realized by an ambient contact isotopy which keeps N invariant. By cutting off this isotopy outside a smaller neighborhood of N we get the required Legendrian isotopy of Λ . \square

Lemma 2.63. *Consider a Wc-manifold (X, λ) with a dividing hypersurface $P \subset X$ and its soul S . Let $\Lambda_0 \subset X \setminus \text{Skel}(X, \lambda)$ be any Legendrian. Then there exists a Legendrian isotopy $\Lambda_t \subset X \setminus \text{Skel}(X, \lambda)$, $t \in [0, 1]$, such that Λ_1 is in a splitting position.*

Proof. We can realize $\partial_\infty X$ as a smooth boundary of a neighborhood of $\text{Skel}(X, \lambda)$ and view $\Lambda, \partial_\infty S$ and $\partial_\infty P$ as submanifolds of $\partial_\infty X$. Then the boundary $\partial_\infty P \subset \partial_\infty X$ is a convex hypersurface in the contact manifold $(\partial_\infty X, \lambda)$, and its dividing set $\partial_\infty S$ divides $\partial_\infty P$ into two copies S_\pm of S . Therefore the conclusion follows from the previous lemma. \square

2.9.3. Legendrians in Wc-buildings. Let (\mathcal{X}, λ) be a Wc-germ with a structure of a k -block building $\mathcal{X}_k \rightarrow \cdots \rightarrow \mathcal{X}_1$. We say that a Legendrian $\Lambda \subset \mathcal{X} \setminus \text{Skel}(\mathcal{X}, \lambda)$ is *compatible with the building structure* if its Liouville cone L is invariant with respect to all Liouville fields Z_i , $i = 1, \dots, k$, which we can think of as living in \mathcal{X} via Lemma 2.52.

Lemma 2.64. *Let (X, λ) be presented as a Weinstein building $\mathcal{X}_k \rightarrow \cdots \rightarrow \mathcal{X}_1$. Let $\Lambda \subset \mathcal{X} \setminus \text{Skel}(\mathcal{X}, \lambda)$ be a Legendrian. Then there exists a Legendrian isotopy Λ_t starting from $\Lambda_0 = \Lambda$ such that Λ_1 is compatible with the building, i.e. its Z -Liouville cone is invariant with respect to all Liouville fields Z_i where they are defined.*

Proof. By definition we have filtration by Wc-manifolds $\mathcal{X} \supset \mathcal{X}_{\geq 2}, \dots, \supset \mathcal{X}_{\geq k-1} \supset \mathcal{X}_k$, where $\mathcal{X}_{\geq j} = \mathcal{X}_k \rightarrow \cdots \rightarrow \mathcal{X}_j$. Note that we equivalently view \mathcal{X} as a 2-block building $\mathcal{X}_{\geq 2} \rightarrow \mathcal{X}_1$. By definition the building $\mathcal{X}_{\geq 2} \rightarrow \mathcal{X}_1$ can be split into the blocks \mathcal{X}_1 and $\mathcal{X}_{\geq 2}$ along a splitting hypersurface H which can be defined as follows. Recall that $\mathcal{X}_{\geq 2} \rightarrow \mathcal{X}_1$ is obtained from \mathcal{X}_1 and $\mathcal{X}_{\geq 2}$ by taking the nucleus N of one of the boundary faces of $\mathcal{X}_{\geq 2}$, realizing it as Wc-hypersurface in \mathcal{X}_1 , converting it to a nucleus of a boundary face and then performing the horizontal gluing along the face H of \mathcal{X}_1 and the converted face of $\mathcal{X}_{\geq 2}$. The collar neighborhood $\mathcal{O}p H$ of the face H in \mathcal{X}_1 is split as $\mathcal{O}p H = H \times \mathcal{I}$. The inverse splitting can be done by using the $H \times \varepsilon$ as a splitting surface. Note that making the Legendrian Λ compatible with the building $\mathcal{X}_{\geq 2} \rightarrow \mathcal{X}_1$ is equivalent to moving it to a splitting position with respect to the splitting hypersurface $H \times \varepsilon$. Hence, using Lemma 2.64 we can move Λ by a Legendrian isotopy to make it compatible with the building $\mathcal{X}_{\geq 2} \rightarrow \mathcal{X}_1$. The splitting defines a Legendrian $\Lambda_{\geq 2} \subset \mathcal{X}_{\geq 2} \setminus \text{Skel}(\mathcal{X}_{\geq 2}, \lambda)$. We continue by applying again 2.64 we move $\Lambda_{\geq 2}$ by a Legendrian isotopy to make it compatible with the building $\mathcal{X}_{\geq 2} \rightarrow \mathcal{X}_2$. Continuing the process we prove the lemma. \square

2.10. Weinstein handlebodies revisited.

2.10.1. *Rounded handles.* Consider an n -ball $D^n \subset S^{n-k} \times D^k$ and denote

$$K_k^n := (S^{n-k} \times D^k) \setminus \text{Int } D^n.$$

Thus, ∂K_k^n is manifold with two boundary components $\partial_- K_k^n = S^{n-k} \times \partial D^{k-1}$ and $\partial_+ K_k^n := S^{n-1}$.

Consider a Weinstein manifold with boundary $G_k^n := T^*K_k^n$. It has two boundary faces M_- and M_+ with nuclei $P_- := T^*(S^{n-k} \times \partial D^k)$ and $P_+ := T^*S^{n-1}$.

Definition 2.65. We call G_k^n a *rounded subcritical Weinstein handle*.

Remark 2.66. Our rounded handles G_k^n have nothing to do with *round* handles of index k .

Lemma 2.67. *Let W^{2n} be Weinstein domain and $\phi : \partial D^k \times D^{n-k} \rightarrow \partial W$ a Legendrian embedding, $k \leq n$. Let W' be the Weinstein domain obtained by a subcritical index k Weinstein handle attachment along the isotropic sphere $\phi(\partial D^k \times 0)$ with the framing given by the Legendrian extension. Suppose that the embedding ϕ is extended to a Legendrian embedding $\Phi : \partial D^k \times S^{n-k} \rightarrow \partial W$. Let $\Sigma = T^*(\partial D^k \times S^{n-k})$ be the Weinstein ribbon of this embedding. Then the Weinstein structure obtained by the vertical gluing of the rounded handle G_k^n along Σ is homotopic to the Weinstein structure on W' .*

Proof. The vertical gluing consists of the conversion of Σ to a boundary nucleus and the horizontal gluing to G_k^n along the new nucleus. The conversion is achieved by taking the union of W with a collar $T^*(\partial D^k \times S^{n-k} \times [0, 1])$ and deforming the Liouville field of W near Σ so that the result is a Weinstein manifold with boundary. Hence the vertical gluing is the union of W with the cotangent bundle of $\widehat{K}_k^n = (\partial D^k \times S^{n-k} \times [0, 1]) \cup (K_n^k \times 1)$. Let $L := D^k \times D^{n-k}$ be attached to $\partial D^k \times S^{n-k}$ along $\partial D^k \times D^{n-k}$, i.e. the image of ϕ . Then \widehat{K}_k^n is diffeomorphic to $T_k^n := (\partial D^k \times S^{n-k} \times [0, 1]) \cup (L \times 1)$ after smoothing corners. The underlying Weinstein structure of the vertically glued Weinstein manifold is given along the boundary of $T^*\widehat{K}_k^n$ by the Weinstein normal form for Morse-Bott boundary of index 0. Before smoothing, on $T^*T_k^n$ this corresponds to the Weinstein normal form for Morse-Bott boundary, also of index 0. By a Weinstein homotopy we can deform the structure to cancel the critical points on $T^*(\partial D^k \times S^{n-k} \times [0, 1])$ along the $[0, 1]$ direction, similar to a parametric Smale cancellation, where we think of $\partial D^k \times S^{n-k}$ as the parameter space. We are left with the Weinstein normal form for Morse-Bott boundary and corners on $T^*L \simeq T^*(D^k \times D^{n-k})$ which has index 1 on the boundary face $\partial D^k \times D^{n-k}$ and index 0 on the boundary face $D^k \times \partial D^{n-k}$. This Weinstein structure is evidently homotopic to the standard Morse-Weinstein normal form of index k on $T^*(D^k \times D^{n-k})$, which is by definition the Weinstein surgery of index k . \square




FIGURE 2.16. Attaching a subcritical isotropic handle is the same as attaching a rounded handle.

2.10.2. *Weinstein manifolds as cotangent buildings.* Recall that a cotangent building is a Wc-building $W = (B_k \rightarrow \cdots \rightarrow B_1)$ consisting of cotangent blocks $B_i := \mathcal{T}^*M_i$.

Proposition 2.68. *Any Weinstein manifold germ (\mathcal{X}, λ) admits the structure of a cotangent building with blocks $B_j = \mathcal{T}^*M_j$, where each M_j is diffeomorphic to either a disc D^n or one of the manifolds K_j^n , $j = 1, \dots, n-1$ with corners added to its boundary.*

Proof. Consider a handle decomposition $H_k \rightarrow H_{k-1} \rightarrow \dots \rightarrow H_1$ of a Weinstein germ (\mathcal{X}, λ) . If $k = 0$ then we just replace the first index 0 handle by T^*D^n . Suppose that the Weinstein structure on the handlebody $W_{\leq k} := H_{k-1} \rightarrow \dots \rightarrow H_1$ is already compatible with a structure of a cotangent building. If the index of the handle H_k is equal m for $0 < m < n$ we use Lemma 2.67 to replace the handle H_k by the rounded handle $G_m^n = T^*K_m^n$. If $m = n$ we keep it as is equal to T^*D^n . In both cases we denote by T^*M_k the block we need to attach with the nucleus of the attaching face N equal to $T^*\partial D^n$ in the latter case and $\partial_- G_m^n = T^*(S^m \times S^{n-m})$ in the former one. For the attaching, N is realized as a Weinstein hypersurface in $W^{<k} \setminus \text{Skel}(W^{<k})$. We denote by Λ the core Legendrian of that hypersurface. Let L be the $(Z_{\leq k})$ -cone of Λ . Using Lemma 2.64 we deform Λ via a Legendrian isotopy to make the cone L invariant with respect to the contracting fields on the blocks B_1, \dots, B_k . Note that this automatically gives $\widehat{M}_k := L \cup M^k$ a new structure of a Wc-manifold with additional corners. The neighborhood of \widehat{M}_k then gets the structure of a cotangent block $B_k = \mathcal{T}^*(\widehat{M}_k)$ and W can be obtained from the vertical gluing of B_k to $W_{\leq k}$. \square




FIGURE 2.17. If $B_2 \rightarrow B_1$ already has the structure of a cotangent building and we wish to extend this structure to a third block B_3 , we need to make sure the Legendrian Λ along whose ribbon the block B_3 will be attached is adapted, i.e. its Liouville cone is invariant under the Liouville flows of the blocks B_1 and B_2 .

More generally, we have the following result:

Proposition 2.69. *Any Wc-manifold admits the structure of a cotangent building.*

To prove this we will inductively apply the following lemma.

Lemma 2.70. *Given any Wc-pair $(\mathcal{X}, \mathcal{A}, \lambda)$, where \mathcal{X} and \mathcal{A} are Wc-germs which admit the structure of a cotangent building, the hypersurface-to-nucleus conversion $(\mathcal{X}^{\mathcal{A}}, \lambda^{\mathcal{A}})$ also admits the structure of a cotangent building.*

Proof. Suppose that (\mathcal{X}, λ) is a Wc-manifold which admits the structure of a cotangent building $B_k \rightarrow \cdots \rightarrow B_1$ and $\mathcal{A} \subset \mathcal{X} \setminus \text{Skel}(\mathcal{X}, \lambda)$ is a Wc-hypersurface which also admits the structure of a cotangent building $B'_r \rightarrow \cdots \rightarrow B'_1$. Then it is straightforward to verify that the hypersurface-to-nucleus conversion $(\mathcal{X}^{\mathcal{A}}, \lambda^{\mathcal{A}})$ inherits an induced structure of a cotangent building $(B'_r \times \mathcal{T}^*[0, 1]) \rightarrow \cdots \rightarrow (B'_1 \times \mathcal{T}^*[0, 1]) \rightarrow B_k \rightarrow \cdots \rightarrow B_1$. \square

Proof of Proposition 2.69. Given any Wc-manifold (\mathcal{X}, λ) , we may convert all of its boundary nuclei to Wc-hypersurfaces to obtain its underlying Weinstein manifold \widehat{X} , and \mathcal{X} can be recovered from \widehat{X} by converting back the Wc-hypersurfaces to face nuclei. By Proposition 2.68 we know that \widehat{X} admits the structure of a cotangent building, and we may assume by induction on dimension that the Wc-hypersurfaces also admit the structure of a cotangent building. Hence it follows from Lemma 2.70 that \mathcal{X} admits the structure of a cotangent building. \square

3. SYMPLECTIC NEIGHBORHOODS OF ARBOREAL LAGRANGIANS

In this section we introduce the notion of an arboreal space and prove that any arboreal space has a unique symplectic thickening.

3.1. Arboreal spaces.

3.1.1. *Pre-Ac spaces.* In [AGEN20a] we associated to each signed rooted tree $\mathcal{T} = (T, \rho, \varepsilon)$, an arboreal Lagrangian $L_{\mathcal{T}} \subset T^*\mathbb{R}^{n(\mathcal{T})}$ which is a union of smooth pieces, $L_{\mathcal{T}} = \bigcup_{\alpha \in v(\mathcal{T})} L_{\alpha}$.

For fixed $c \geq 0$, and $m \geq n(\mathcal{T}) + c$, set $d = m - n(\mathcal{T}) + c$, and consider the product

$$L(\mathcal{T}, m, c) := L_{\mathcal{T}} \times \mathbb{R}^d \times \mathbb{R}_{\geq 0}^c$$

We also have the smooth pieces $L(\mathcal{T}, m, c)_{\alpha} := L_{\alpha} \times \mathbb{R}^d \times \mathbb{R}_{\geq 0}^c$.

We equip $L(\mathcal{T}, m, c)$ with the structure sheaf $\mathcal{O}(\mathcal{T}, m, c)$ of functions restricted from smooth functions on $T^*\mathbb{R}^{n(\mathcal{T})} \times \mathbb{R}^d \times \mathbb{R}_{\geq 0}^c$ by the inclusion $L(\mathcal{T}, m, c) \hookrightarrow T^*\mathbb{R}^{n(\mathcal{T})} \times \mathbb{R}^d \times \mathbb{R}_{\geq 0}^c$.

Definition 3.1. An m -dimensional *pre-arboreal space with corners* (*pre-Ac space*) (X, \mathcal{O}) is a locally ringed Hausdorff topological space locally modeled on

$$(L(\mathcal{T}, m, c), \mathcal{O}(\mathcal{T}, m, c))$$

for varying signed rooted trees \mathcal{T} , and $c \leq m - n(\mathcal{T})$.

A *smooth sheet* $Y \rightarrow X$ of a pre-Ac space (X, \mathcal{O}) is a finite proper map from a manifold with corners such that the map is an embedding on the interior of Y , and locally the map identifies the local connected components of Y with smooth pieces

$$L(\mathcal{T}, m, c)_{\alpha} \subset L(\mathcal{T}, m, c)$$

for varying $\alpha \in v(\mathcal{T})$.

In this paper we shall only need to consider the cases where $c = 0$ or $c = 1$. When $c, d = 0$, we write $(L_{\mathcal{T}}, \mathcal{O}_{\mathcal{T}})$ in place of $(L(\mathcal{T}, m, c), \mathcal{O}(\mathcal{T}, m, c))$.




FIGURE 3.1. We can take X to be the interval $[-1, 1]$ with 1 glued to 0 as an \mathcal{A}_2 -singularity. Then $[-1, 1] \rightarrow X$ is a smooth sheet. We will introduce the notion of Ac-building to avoid the unnecessary complications of this kind of gluing.

Remark 3.2. Given a pre-Ac space (X, \mathcal{O}) , we can speak about a closed embedding into a smooth manifold (M, C_M^∞) . These are embeddings with closed image $X \rightarrow M$ with a surjective map of sheaves of local rings $C_M^\infty|_X \rightarrow \mathcal{O}$.

Remark 3.3. One can relax the definition of a pre-arboreal space by choosing the alternative structure sheaf of continuous real-valued functions on $L(\mathcal{T}, m, c)$ whose restrictions to each smooth piece $L(\mathcal{T}, m, c)_\alpha$, for $\alpha \in v(\mathcal{T})$, are smooth. One can show that every such “weak” pre-arboreal space is diffeomorphic to a strong one, and any weak diffeomorphism between strong pre-arboreals is isotopic to a strong one.

3.1.2. *Ac spaces.* In [AGEN20a] we proved that any germ of an arboreal Lagrangian is symplectomorphic to exactly one of the canonical models $L_{\mathcal{T}} \subset T^*\mathbb{R}^{n(\mathcal{T})}$, hence in particular isomorphic as the germ of a locally ringed Hausdorff topological space to the corresponding $(L_{\mathcal{T}}, \mathcal{O}_{\mathcal{T}})$. Going the other way, if we are given a local model $(L_{\mathcal{T}}, \mathcal{O}_{\mathcal{T}})$ it is not possible to reconstruct the symplectic geometry of $L_{\mathcal{T}} \subset T^*\mathbb{R}^{n(\mathcal{T})}$ unless we keep track of additional structure. This is due to the fact that not every automorphism of $(L_{\mathcal{T}}, \mathcal{O}_{\mathcal{T}})$ can be lifted to an automorphism of $L_{\mathcal{T}} \subset T^*\mathbb{R}^{n(\mathcal{T})}$, i.e. a germ of a symplectomorphism preserving $L_{\mathcal{T}}$. We now explain the additional structure that we need to keep track of.

For a signed rooted tree $\mathcal{T} = (T, \rho, \varepsilon)$, recall if we remove the root $\mathcal{T} \setminus \rho$ what remains is a disjoint union of rooted trees with signs along their edges. Let us write $\mathcal{T}' = (T, \rho, \varepsilon')$ for the signed rooted tree where we negate all of the signs along the edges in any collection of components of $\mathcal{T} \setminus \rho$. Then negating corresponding coordinates in our local models provides an isomorphism of pre-Ac spaces

$$(L_{\mathcal{T}}, \mathcal{O}_{\mathcal{T}}) \simeq (L_{\mathcal{T}'}, \mathcal{O}_{\mathcal{T}'})$$

that does not extend to an ambient symplectomorphism unless $\mathcal{T} = \mathcal{T}'$. Additionally, in the case of the tree \mathcal{A}_2 with a single unsigned edge, the nontrivial involution of the pre-Ac space $(L_{\mathcal{A}_2}, \mathcal{O}_{\mathcal{A}_2})$ that reverses the orientation on the smooth base piece $L_\rho \subset L_{\mathcal{A}_2}$ can only be extended to an ambient anti-symplectomorphism. We will see in Lemma 3.10 below these are the only ambiguities.

We introduce here additional structure to overcome the above ambiguities and hence enable us to reconstruct $L_{\mathcal{T}} \subset T^*\mathbb{R}^{n(\mathcal{T})}$ from $(L_{\mathcal{T}}, \mathcal{O}_{\mathcal{T}})$. Let us first mention some simple data we could add. Given $(L_{\mathcal{T}}, \mathcal{O}_{\mathcal{T}})$, consider the base smooth piece $L_{\rho} \subset L_{\mathcal{T}}$, and the hypersurface $H = \bigcup_{\alpha \neq \rho} L_{\alpha} \cap L_{\rho} \subset L_{\rho}$. Then a coorientation of H provides an embedding $L_{\mathcal{T}} \subset T^*\mathbb{R}^{n(\mathcal{T})}$ as the corresponding positive conormal of H . In particular, there is the specific choice of coorientation that gives the canonical embedding $L_{\mathcal{T}} \subset T^*\mathbb{R}^{n(\mathcal{T})}$ of the local model.

To go forward, we will introduce a version of the above coorientation data that can be easily glued in a global setting. For a signed rooted tree $\mathcal{T} = (T, \rho, \varepsilon)$, and vertex $\alpha \in v(\mathcal{T})$, consider the orientation bundle

$$\kappa(\mathcal{T}, m, c)_{\alpha} := \wedge^m TL(\mathcal{T}, m, c)_{\alpha}$$

Let $K(\mathcal{T}, m, c)$, denote the space of real line bundles $\kappa \rightarrow L(\mathcal{T}, m, c)$ equipped with identifications

$$\kappa|_{L(\mathcal{T}, m, c)_{\alpha}} \simeq \kappa(\mathcal{T}, m, c)_{\alpha} \quad \alpha \in v(\mathcal{T})$$

Note that given such a κ , we can extract from it a coorientation of the hypersurface $H = \bigcup_{\alpha \neq \rho} L(\mathcal{T}, m, c)_{\alpha} \cap L(\mathcal{T}, m, c)_{\rho}$ in the base smooth piece $L(\mathcal{T}, m, c)_{\rho}$. Namely, we can take the coorientation of H to be that glued under κ with the positive Liouville direction.

Lemma 3.4. *Each connected component of $K(\mathcal{T}, m, c)$ is contractible, and $\pi_0(K(\mathcal{T}, m, c))$ is a $(\mathbb{Z}/2\mathbb{Z})^{e(\mathcal{T})}$ -torsor.*

Proof. Suppose the lemma is proved for signed rooted trees with fewer vertices than \mathcal{T} .

Fix $r > 0$, let $B_r = \{|p| = r\} \subset T^*(\mathbb{R}^{m-c} \times \mathbb{R}_{\geq 0}^c)$ denote the radius r open ball bundle, and $N_r = B_r \cap L(\mathcal{T}, m, c)$ the corresponding open neighborhood of the zero-section $L(\mathcal{T}, m, c)_{\rho} \simeq \mathbb{R}^{m-c} \times \mathbb{R}_{\geq 0}^c$. Note the Liouville flow of $T^*\mathbb{R}^m$ deformation retracts N_r back to the zero-section.

Consider the Čech cover of $L(\mathcal{T}, m, c)$ given by the open N_r along with the disjoint opens

$$L(\mathcal{T}_i, m-1, c) \times \mathbb{R}_{>0} \quad i \in I$$

where $I \subset v(T)$ denotes the set of vertices adjacent to the root ρ , or equivalently the set of edges adjacent to ρ , \mathcal{T}_i is the signed rooted tree given by the connected component of $\mathcal{T} \setminus \rho$ containing $i \in I$, and the last factor $\mathbb{R}_{>0}$ of each open is given by Liouville flow. Note each intersection $N_r \cap L(\mathcal{T}_i, m-1, c) \times \mathbb{R}_{>0}$, for $i \in I$, is homeomorphic to $L(\mathcal{T}_i, m-1, c) \times (0, r)$, in particular connected.

We see that to give a line bundle on $L(\mathcal{T}, m, c)$, given its restriction to the above opens, is to give a sign ± 1 on each intersection $N_r \cap L(\mathcal{T}_i, m-1, c) \times \mathbb{R}_{>0}$, for $i \in I$. This inductively establishes the lemma. \square

We will now use the symplectic geometry to specify a distinguished element of $\kappa(\mathcal{T}, m, c) \in K(\mathcal{T}, m, c)$, i.e. a trivialization of the $(\mathbb{Z}/2\mathbb{Z})^{e(\mathcal{T})}$ -torsor of the lemma

$$K(\mathcal{T}, m, c) \simeq (\mathbb{Z}/2\mathbb{Z})^{e(\mathcal{T})}$$

Given the canonical model $L(\mathcal{T}, m, c) \subset T^*\mathbb{R}^m$, we have a canonical coorientation of the hypersurface $H = \bigcup_{\alpha \neq \rho} L(\mathcal{T}, m, c)_\alpha \cap L(\mathcal{T}, m, c)_\rho$ in the base smooth piece $L(\mathcal{T}, m, c)_\rho$. To identify the orientation bundles of the smooth pieces along this intersection it suffices to specify that *the positive codirection to the co-oriented front is glued to the positive Liouville direction*. Note this choice is invariant under symplectomorphism.

Definition 3.5. The *neutral element* $\kappa(\mathcal{T}, m, c) \in K(\mathcal{T}, m, c)$ is given by the above inductively specified gluing.

Remark 3.6. Let us mention a characterization (we will not use) of when an arboreal model is positive. Assume for simplicity \mathcal{T} has a single vertex adjacent to the root. Then the Legendrian at infinity $L(\mathcal{T}, m, c)^\infty \subset S^*\mathbb{R}^m$ projects homeomorphically to its front $H \subset \mathbb{R}^m$, and the front has a natural “tangent” bundle given by the orthogonal to the coorientation. In the case of positive \mathcal{T} , the restriction of $\kappa(\mathcal{T}, m, c)$ to $L(\mathcal{T}, m, c)^\infty$ coincides with the top exterior power of the “tangent” bundle of the front under the above homeomorphism.

Definition 3.7. An *orientation structure* κ for a pre-Ac space (X, \mathcal{O}) is a real line bundle $\kappa \rightarrow X$ equipped with identifications $\kappa|_Y \simeq \kappa_Y$ for each smooth sheet $Y \rightarrow X$, locally modeled on the neutral elements $\kappa(\mathcal{T}, m, c) \in K(\mathcal{T}, m, c)$.

Definition 3.8. An m -dimensional *arboreal space with corners (Ac space)* (X, \mathcal{O}, κ) is an m -dimensional pre-arboreal space (X, \mathcal{O}) equipped with an orientation structure κ .

Remark 3.9. When $m = 1$, one can compare the notion of orientation structure with ribbon structure for a trivalent graph. At each trivalent vertex we have a “T” singularity and in this case an orientation structure is equivalent to a cyclic orientation of the three half-edges incident to the vertex. In particular, the orientation structure rigidifies the pre-Ac space $(L_{A_2}, \mathcal{O}_{A_2})$.




FIGURE 3.2. For the A_2 singularity there are two possible orientation structures, which can be realized by the above embeddings as arboreal spaces into $(\mathbb{R}^2, dp \wedge dq)$. Since the zero section is fixed, the choice of an orientation structure is naturally equivalent to the choice of a cyclic structure on the three half-edges incident to the vertex. Note that the choice of an orientation structure is also equivalent to a choice of co-orientation for a point inside a line.

Recall from [AGEN20a] that an arboreal Lagrangian L in a symplectic manifold (M, ω) is locally symplectomorphic to a standard model. Since this model has an orientation structure given by its neutral element, and this element is invariant under symplectomorphism, we conclude that any arboreal Lagrangian has an orientation structure induced by the ambient symplectic structure. Hence *every arboreal Lagrangian has the structure of an arboreal space*.

Finally, let us return to the discussion at the beginning of this section, and observe that, locally, orientation structures are nothing more than choices of coorientations.

Lemma 3.10. *Let $\mathcal{T} = (T, \rho, \varepsilon)$ be a signed rooted tree.*

There are precisely 2^k isomorphism classes of orientation structures on the pre-Ac space $(L_{\mathcal{T}}, \mathcal{O}_{\mathcal{T}})$, where k is the number of edges adjacent to the root ρ .

Proof. By construction, an orientation structure on the local model $(L_{\mathcal{T}}, \mathcal{O}_{\mathcal{T}})$ must result by pullback of the neutral orientation structure under an isomorphism $(L_{\mathcal{T}}, \mathcal{O}_{\mathcal{T}}) \simeq (L_{\mathcal{T}'}, \mathcal{O}_{\mathcal{T}'})$ for some \mathcal{T}' . Moreover, note that $(L_{\mathcal{T}}, \mathcal{O}_{\mathcal{T}})$ is determined by the hypersurface $H = \bigcup_{\alpha \neq \rho} L_{\alpha} \cap L_{\rho} \subset L_{\rho} \simeq \mathbb{R}^{n(\mathcal{T})}$, and so we must have an isomorphism $(\mathbb{R}^{n(\mathcal{T})}, H) \simeq (\mathbb{R}^{n(\mathcal{T}')} , H')$ of hypersurfaces. So we must have $\mathcal{T}' = (T, \rho, \varepsilon')$ where ε' results from ε by negating all of the signs along the edges in some collection of components of $\mathcal{T} \setminus \rho$. These choices correspond to the 2^k choices of coorientations of the hypersurface H . Each such choice provides a distinct Lagrangian realization $L_{\mathcal{T}} \subset T^*\mathbb{R}^{n(\mathcal{T})}$ with corresponding orientation structure. \square

Remark 3.11. Note therefore that the data of an orientation structure on an arboreal singularity is equivalent to a fully signed rooted tree, i.e. a signed rooted tree in the previous sense together with additional signs on each of the edges of \mathcal{T} adjacent to the root, so that now every edge in \mathcal{T} has a sign. We will however not use this notion of a fully signed rooted tree and will always use the term signed rooted tree for rooted trees with signs on those edges not adjacent to the root.

3.2. Arboreal Darboux-Weinstein neighborhood theorem.

3.2.1. *Lagrangian subsets.* Let L be a path connected and locally simply-connected closed subset in a symplectic manifold (M, ω) . We call L a *Lagrangian subset* if each point of L has a Darboux neighborhood U such that $\oint_{\gamma} pdq = 0$ for every loop γ in $L \cap U$. Clearly, any arboreal Lagrangian $L \subset M$ is a Lagrangian subset. If f is (the germ at a point of) a diffeomorphism $M \rightarrow M$, then $f(L)$ is (the germ of) a Lagrangian subset if and only if the image under f of each smooth piece of L is Lagrangian.

If $L \subset M$ is a pre-Ac space embedded as a Lagrangian subset in a symplectic manifold (M, ω) , then L inherits an orientation structure as follows. Returning to the inductive description of orientation structures given in Lemma 3.4, along a smooth Lagrangian piece $P \subset L$, given a Lagrangian half-plane H with codimension 1 clean intersection with P , we specify that the inward normal direction ν to $P \cap H$ in H is glued to the normal direction to $P \cap H$ in P which pairs positively with ν under ω . With this structure, any pre-Ac space embedded as a Lagrangian subset gets an induced structure of an Ac space.

Note the above orientation structure on L as a Lagrangian subset agrees with the previously defined orientation structure on L as an arboreal Lagrangian, i.e. the one defined using local models.

3.2.2. *Darboux-Weinstein theorem.* The main result of this section is the following:

Theorem 3.12. *Let V be a manifold with two symplectic forms ω_0, ω_1 . Suppose $L \subset V$ is a compact arboreal space embedded as a Lagrangian subset for both symplectic forms. Let κ_0, κ_1 be the induced orientation structures on L .*

If the orientation structures are isomorphic $\kappa_0 \simeq \kappa_1$, then there exists a diffeomorphism $\varphi : V \rightarrow V$, preserving L as a subset, such that $\varphi^\omega_1 = \omega_0$. If ω_0 and ω_1 coincide on $\mathcal{O}p A$ for a closed subset $A \subset L$ then φ can be chosen to be the identity on $\mathcal{O}p A$.*

Corollary 3.13. *Any arboreal space embeds into a symplectic manifold as an arboreal Lagrangian such that the orientation structure induced from the ambient symplectic structure is isomorphic to the given orientation structure.*

Proof. By definition, any point of an arboreal space has a model neighborhood. On overlaps, these local symplectic structures have the same orientation structure, and hence according to Theorem 3.12 they are diffeomorphic via a diffeomorphism preserving the arboreal. Moreover, the relative form of that theorem allows us to construct a symplectic neighborhood inductively extending it over neighborhoods of strata of certain stratification. \square

In Theorem 3.27, proved below in Section 3.27 below, we find a Weinstein structure on a symplectic neighborhood of an arboreal space L for which L serves as the skeleton. Moreover, Theorem 3.27 ensures uniqueness of this structure up to Weinstein homotopy fixing the skeleton.

Theorem 3.12 is a corollary of the following two propositions.

Proposition 3.14. *Let V be a manifold with two symplectic forms ω_0, ω_1 . Suppose $L \subset V$ is a compact arboreal space which is a Lagrangian subset for both symplectic forms. Let κ_0, κ_1 be the induced orientation structures on L . If the orientation structures are isomorphic $\kappa_0 \simeq \kappa_1$, then there exists a diffeomorphism ϕ of $\mathcal{O}p(L)$ preserving L pointwise such that the linear interpolation $\omega_t = (1-t)\omega_0 + t\phi^*\omega_1$ is a family of symplectic forms on $\mathcal{O}p(L)$ so that L is a Lagrangian subset for each ω_t . If ω_0 and ω_1 coincide on $\mathcal{O}p A$ for a closed subset $A \subset X$ then ϕ can be chosen to be the identity on $\mathcal{O}p A$.*

Proposition 3.15. *Let $L \subset V$ be an arboreal subspace which is a Lagrangian subset for a family ω_t , $t \in [0, 1]$, of symplectic forms on L . Then there exists a diffeotopy $h_t : \mathcal{O}p(L) \rightarrow \mathcal{O}p(L)$, $t \in [0, 1]$, such that $h_t(L) = L$ and $h_t^*\omega_t = \omega_0$. If $\omega_t = \omega_0$ on $\mathcal{O}p A$ for a closed subset $A \subset L$ then the isotopy h_t can be chosen fixed on $\mathcal{O}p A$.*

Indeed, to prove Theorem 3.12 we can apply Proposition 3.15 to the output of Proposition 3.14. We will prove Propositions 3.14 and 3.15 in the next two subsections.

Remark 3.16. As the proof of Proposition 3.14 will show, the conclusion holds more generally for two symplectic manifolds (V_i, ω_i) , $i = 1, 2$, together with embeddings of a compact arboreal space L onto a Lagrangian subset of V_i , $i = 1, 2$. In particular it follows that the smooth topology of a neighborhood of a Lagrangian compact arboreal space in a symplectic manifold is uniquely determined by the compact arboreal space.

3.2.3. Proof of Proposition 3.14. Let us revisit the notion of an orientation structure. Consider a full flag $F = (0 = V_0 \subset V_1 \subset \cdots \subset V_{2n} = V)$ of subspaces in a $2n$ -dimensional vector space V , $\dim V_j = j$. Let $\Omega(F)$ be the space of all symplectic forms ω on V such that V_j is isotropic for $j \leq n$, V_j is coisotropic for $j \geq n$, and $V_j^{\perp\omega} = V_{2n-j}$ for $j \leq n$.

We begin with following elementary assertion.

Lemma 3.17. *The connected components of $\Omega(F)$ are determined by orientations of the symplectic quotients V_{2n-j}/V_j , for $j = 0, \dots, n-1$. Any sequence of orientations can occur and each connected component is convex.*

Proof. Represent ω as a matrix with respect to a basis u_1, \dots, u_{2n} of V such that $V_j = \text{span}(u_1, \dots, u_j)$. It has a block form with four n by n blocks, the two diagonal blocks being zero and the two off-diagonal blocks being A and $-A$. Moreover, A is a triangular matrix and its diagonal elements must be non-zero. Their signs naturally correspond to orientations of the symplectic quotients. \square

We use the notation of [AGEN20a] for the explicit construction of the extended local model

$${}^nL = \bigcup_{i=0}^n {}^nL_i \subset T^*\mathbb{R}^n$$

in terms of equations in the canonical coordinates $x_0, \dots, x_{n-1}, p_0, \dots, p_{n-1}$. Introduce the flag $F({}^nL) = (0 = V_0 \subset V_1 \subset \cdots \subset V_{2n} = T^*\mathbb{R}^n)$ given by the assignments

$$V_i = T_0(\bigcap_{j=0}^{n-i} {}^nL_j) = \text{Span}(\partial_{x_{n-1}}, \dots, \partial_{x_{n-i}}) \quad i \leq n$$

$$V_{2n-i} = \Theta_0(\bigcup_{j=0}^{n-i} {}^nL_j) = \text{Span}(\partial_{x_{n-1}}, \dots, \partial_{x_0}, \partial_{p_0}, \dots, \partial_{p_{2n-i}}) \quad i \leq n$$

Here $\bigcap_{j=0}^{n-i} {}^nL_j$ is a submanifold of $T^*\mathbb{R}^n$, in fact a linear subspace of ${}^nL_0 = \mathbb{R}^n$, and $T_0(\bigcap_{j=0}^{n-i} {}^nL_j)$ denotes its tangent space at the origin; $\bigcup_{j=0}^{n-i} {}^nL_j$ is not a submanifold, but has a tangent sheaf in the sense of algebraic geometry, and $\Theta_0(\bigcup_{j=0}^{n-i} {}^nL_j) = (\mathcal{I}_0/\mathcal{I}_0^2)^\vee$ denotes its fiber at the origin. Note these characterizations imply $F({}^nL)$ is preserved by diffeomorphisms of $T^*\mathbb{R}^n$ preserving nL as a subset hence preserving each nL_i as a subset, i.e. $F({}^nL)$ only depends on the sheaf of smooth functions on nL . Note also V_i is isotropic for $i \leq n$, V_i is coisotropic for $i \geq n$, and $V_i^{\perp\omega} = V_{2n-i}$ for $j \leq n$ for any symplectic form on \mathbb{R}^{2n} such that ${}^nL \subset \mathbb{R}^{2n}$ is Lagrangian.

Recall that the local model for an \mathcal{A}_{n+1} -arboreal Lagrangian is a subset of the extended local model: $L_{\mathcal{A}_{n+1}} \subset {}^nL$. We have the following reformulation of orientation structures for Ac-space structures on the germ of $L_{\mathcal{A}_{n+1}}$ at its central point:

Lemma 3.18. *Let ω_0 be a symplectic form on \mathbb{R}^{2n} for which $L_{\mathcal{A}_{n+1}}$ is a Lagrangian subset.*

If ω_1 is another such form, then the orientation structures κ_0, κ_1 induced by ω_0, ω_1 respectively on $L_{\mathcal{A}_{n+1}}$ are isomorphic if and only if ω_0 and ω_1 lie in the same connected component of $\Omega(F({}^nL))$.

Proof. Let ω be a symplectic structure on \mathbb{R}^{2n} such that nL is Lagrangian. The sign of the leading diagonal element of the matrix A in the proof of Lemma 3.17 applied to ω is precisely the pairing used to define the orientation structure on nL as a Lagrangian subset of $(\mathbb{R}^{2n}, \omega)$. The other signs are uniquely determined by quadratic information. For example, for $L_{\mathcal{A}_3} \subset T^*\mathbb{R}^2$ the third stratum is the positive conormal to $\{x_0 = x_1^2 \mid x_1 \geq 0\}$. Therefore, the vectors $2x_1\partial/\partial x_1 + \partial/\partial x_0$ and $\partial/\partial p_1 + 2x_1\partial/\partial p_0$ should span a Lagrangian subspace. Their symplectic product is $2x_1(\lambda - \mu)$ where λ is the first diagonal term of the matrix A and μ is the second diagonal term of the matrix A . So in fact $\lambda = \mu$. The general case can be proved similarly by induction. The conclusion is that the connected component of $\Omega(F({}^nL))$ is uniquely determined by the orientation structure, which in this case is just a single sign. \square

For more general trees \mathcal{T} , one can apply the above argument to each factor $T^*\mathbb{R}^n$ in $T^*\mathbb{R}^{n(\mathcal{T})}$ corresponding to each \mathcal{A}_n -subtree of \mathcal{T} with the same root ρ and deduce the general version:

Lemma 3.19. *Let \mathcal{T} be a signed rooted tree and ω_0 a symplectic form on $\mathbb{R}^{2n(\mathcal{T})}$ for which $L_{\mathcal{T}}$ is Lagrangian.*

If ω_1 is another such form, then the orientation structures κ_0, κ_1 induced by ω_0, ω_1 respectively on $L_{\mathcal{T}}$ are isomorphic if and only if for any \mathcal{A}_{m+1} -type subtree $\mathcal{A}_{m+1} \subset \mathcal{T}$ with the same root ρ , the restrictions of ω_0 and ω_1 to the factor \mathbb{R}^{2m} of $\mathbb{R}^{2n(\mathcal{T})}$ corresponding to the \mathcal{A}_{m+1} -subtree lie in the same connected component of $\Omega(F({}^mL))$.

So we have proved Proposition 3.14 for the germ of the central point of the pre-Ac space $L(\mathcal{T}, m)$ with $m = n(\mathcal{T})$, i.e. not stabilized. In the case $d = m - n(\mathcal{T}) > 0$, the situation changes as follows. Given a symplectic form ω_0 on \mathbb{R}^{2m} for which the standard model $L_{\mathcal{T}} \times \mathbb{R}^d \subset \mathbb{R}^{2n(\mathcal{T})} \times \mathbb{R}^{2d}$ is Lagrangian, consider $E \subset \mathbb{R}^{2m}$, the $(2m - d)$ -dimensional coisotropic subspace of \mathbb{R}^{2m} which is symplectically orthogonal to the isotropic $F = 0 \times \mathbb{R}^d$. Since $L_{\mathcal{T}} \times \mathbb{R}^d$ is assumed to be Lagrangian, note that F is spanned by those directions tangent to the smooth part of $L_{\mathcal{T}} \times \mathbb{R}^d$ and E is spanned by those directions tangent to $L_{\mathcal{T}} \times \mathbb{R}^d$ at 0, hence both are independent of the symplectic form ω_0 .

The symplectic reduction of $(E, \omega_0|_E)$ is a $2(m - d)$ -dimensional symplectic vector space V_0 . The quotient \mathbb{R}^{2m}/E is canonically identified via the fixed symplectic form ω_0 with the dual of $0 \times \mathbb{R}^d$. Therefore there exists a canonical symplectic isomorphism Φ between \mathbb{R}^{2m} and $V_0 \oplus (F \oplus F^*)$.

Now suppose that we have two such symplectic forms ω_0 and ω_1 on \mathbb{R}^{2m} for which $L_{\mathcal{T}} \times \mathbb{R}^d$ is Lagrangian.

Lemma 3.20. *Consider $\mathcal{S} = \mathcal{S}(\omega_0, \omega_1)$, the space of linear isomorphisms $\Psi : \mathbb{R}^{2m} \rightarrow \mathbb{R}^{2m}$ such that:*

- (i) $\Psi|_{L_{\mathcal{T}} \times \mathbb{R}^d} = \text{Id}_{L_{\mathcal{T}} \times \mathbb{R}^d}$.
- (ii) $\Psi^* \omega_1$ and ω_0 agree on $F \oplus (\mathbb{R}^{2m}/E)$.

Then \mathcal{S} is nonempty and contractible.

Example 3.21. For the smooth arboreal singularity, \mathcal{S} consists of a single element.

Proof. For the existence, write $\mathbb{R}^{2m} = \mathbb{R}^{2n(\mathcal{T})} \times \mathbb{R}^d \times \mathbb{R}^d$. Consider a linear isomorphism $\Psi : \mathbb{R}^{2m} \rightarrow \mathbb{R}^{2m}$ which is the identity on the first two factors. This induces a linear isomorphism between the V_0 and V_1 corresponding to ω_0 and ω_1 . The further condition that Ψ respects the identification $V_0 \oplus (F \oplus F^*) \simeq V_1 \oplus (F \oplus F^*)$ uniquely determines a linear isomorphism Ψ satisfying (i) and (ii).

For the contractibility, fix $\Psi_0 \in \mathcal{S}$. For any other $\Psi_1 \in \mathcal{S}$, we will write the matrix for $\Psi_1^{-1} \Psi_0$ in block form with respect to the decomposition $\mathbb{R}^{2m} = \mathbb{R}^{2n(\mathcal{T})} \times \mathbb{R}^d \times \mathbb{R}^d$, with the canonical basis. The matrix will have block form, with the diagonal factors the identity, and with all off-diagonal blocks zero except for the $(3, 2)$ block, which is an arbitrary symmetric matrix A . The space of such is contractible, so this completes the proof. \square

Lemma 3.22. *Let ω_0, ω_1 be symplectic forms on \mathbb{R}^{2m} such that $L_{\mathcal{T}} \times \mathbb{R}^d$ is Lagrangian and which induce isomorphic orientation structures. Let $\Psi \in \mathcal{S}$. Then $(1-t)\omega_0 + t\Psi^* \omega_1$ is symplectic for all $0 \leq t \leq 1$.*

Proof. On \mathbb{R}^{2m}/V_0 , the symplectic forms induced by ω_0 and $\Psi^* \omega_1$ agree. On V_0 the problem is reduced to the non-stabilized case as discussed above: the homotopy $(1-t)\omega_0 + t\Psi^* \omega_1$ will be through symplectic forms whenever the induced orientation structures agree. Indeed, the argument is unaffected by the symplectic isomorphism Ψ since it is the identity on $L_{\mathcal{T}} \times \mathbb{R}^d$. \square

Now we are ready to prove Proposition 3.14. Let $L \subset V$ be a compact arboreal space which is Lagrangian for two symplectic forms ω_0 and ω_1 . By Lemmas 3.20 and 3.22 we may construct a family of linear isomorphisms $\Psi_x : T_x V \rightarrow T_x V$, $x \in L$, such that Ψ_x is the identity on directions tangent to L at x , and such that $(1-t)\omega_0(x) + t\Psi_x^* \omega_1(x)$ is symplectic for all $x \in X$ and $0 \leq t \leq 1$. We may then integrate the family Ψ_x to a diffeomorphism $\Psi : \mathcal{O}p(L) \rightarrow \mathcal{O}p(L)$ such that $\Psi|_L = \text{Id}_L$ and such that $(1-t)\omega_0 + t\Psi^* \omega_1$ is a homotopy of symplectic forms on $\mathcal{O}p(L)$. This completes the proof.

3.2.4. Proof of Proposition 3.15.

Lemma 3.23. *Let $L \subset T^*M$ be a Lagrangian subset which is the union $M \cup C(\Lambda)$ where $C(\Lambda)$ is the Liouville cone over a Legendrian Λ . Let λ be a 1-form on T^*M such that $\lambda|_M = 0$ and $d\lambda = 0$ on L . Then there exists a smooth function $H : T^*M \rightarrow \mathbb{R}$ vanishing on M together with its differential dH , and such that λ coincides with dH on vectors tangent to L .*

Proof. For any point $p \in T_q^*M$ denote by $\gamma_{p,q}$ the path $t \mapsto (tp, q)$, $t \in [0, 1]$. Define the function H by the formula

$$H(p, q) = \int_{\gamma_{p,q}} \lambda.$$

For $(p, q) \in L$ the path $\gamma_{p,q}$ is contained in L , and hence on tangent vectors to L the 1 forms dH and λ coincide. \square

Lemma 3.24. *Let $L \subset T^*M$ be a Lagrangian subset which is the union $M \cup C(\Lambda)$. Let $A \subset M$ be a closed subset and ω_t be a family of symplectic forms on T^*M such that $\omega_0 = d(pdq)$, $\omega_t|_{\mathcal{O}_p A} = \omega_0$, and L is ω_t -Lagrangian for all t . Then there exists an isotopy $f_t : T^*M \rightarrow T^*M$ which is fixed on $M \cup \mathcal{O}_p A$, leaves L invariant and such that $f_t^* \omega_t = \omega_0$.*

Proof. There exists a family of Liouville forms λ_t for ω_t such that $\lambda_0 = pdq$, $\lambda_t|_{T^*M|_M} = 0$ and $\lambda_t|_{\mathcal{O}_p A} = pdq|_{\mathcal{O}_p A}$. By Lemma 3.23 there exists a family of functions H_t on T^*M such that $dH_t|_L = \lambda_t|_L$, $H_t|_{M \cup \mathcal{O}_p A} = 0$ and $dH_t|_{T^*M|_M} = 0$. Define $\mu_t := \lambda_t - dH_t$. Then $\mu_0 = pdq$, $\mu_t|_{T^*M|_M} = 0$, $\mu_t|_{\mathcal{O}_p A} = pdq|_{\mathcal{O}_p A}$ and $\mu_t|_L = 0$. Then Moser's homotopical method yields an isotopy $f_t : T^*M \rightarrow T^*M$ which is fixed on $M \cup \mathcal{O}_p A$, leaves L invariant and such that $f_t^* \omega_t = \omega_0 = d(pdq)$. \square

Note that any point of an arboreal space which is realized as a Lagrangian subset of a symplectic manifold X has a neighborhood which has the conical form of Lemma 3.24. Hence, we can take a finite covering of X by such neighborhoods and successively apply Lemma 3.24 to extend the isotopy. This concludes the proof of Proposition 3.15, and with it the proof of Theorem 3.12.

3.3. Weinstein thickening of an Ac-building.

3.3.1. *Ac-buildings.* Similar to the notion of a Liouville or a Weinstein manifold with corners, by a *symplectic manifold with corners* we mean a possibly non-compact manifold with corners, such that near each corner point it is symplectomorphic to $\mathbb{R}^{2m} \times T^*\mathbb{R}_+^{n-m}$.

An *arboreal Lagrangian with boundary with corners*, or *ac-Lagrangian* for short, is a closed subset of a symplectic manifold W with corners such that the germ of (W, L) at a point $x \in L$ is symplectomorphic to the germ of $(\mathbb{R}^{2n-2c} \times T^*\mathbb{R}_+^c, L_{\mathcal{T}} \times \mathbb{R}_+^c)$ at the origin for some signed rooted tree \mathcal{T} and an integer $c \geq 0$. The c is called the multiplicity of the corner. It follows from the definition that corners of L are contained in the corresponding corners of W . As in the case of manifolds with boundary with corners, we denote by $\partial_j L$ the strata of corner points of order $j \geq 1$, and write $\partial L = \bigcup_j \partial_j L$. The closure of each component of $\partial_j L$ is itself an ac-Lagrangian in a symplectic submanifold of W of codimension $2k$. The closures of components of $\partial_1 L$ are called *boundary faces* of L .

It will also be useful to consider the notion of an *Ac-building*, a special kind of Ac-space whose definition is analogous to that of a cotangent building as a special kind of Wc-manifold. Recall that an Ac-space L is the union $L = \bigcup_j L_j$ of smooth pieces L_j , and each point of L belongs to the interior of exactly one L_j .

Definition 3.25. An *Ac-building* is an Ac-space L whose smooth pieces can be ordered L_1, \dots, L_k so that:

- (i) Each $L_{>j} = \bigcup_{i>j} L_i$ is an Ac-space with boundary and corners.

- (ii) $L_{>j-1}$ is obtained from $L_{>j}$ by gluing a boundary component P_j of $L_{>j}$ to L_j along a map $P_j \rightarrow L_j$.

Lemma 3.26. *Let $K = \bigcup_j L_j$ be an Ac-building. Then the gluing maps $P_j \rightarrow L_j$ can be canonically lifted to embeddings $P_j \rightarrow T^*L_j \setminus L_j$ onto arboreal Legendrians.*

Proof. This follows from the characterization of orientation structures in terms of co-orientations of fronts, as discussed above. \square

Theorem 3.27. *Suppose (W, ω) is a $2m$ -dimensional symplectic manifold, and let $L \subset W$ be an Ac-blding. Then*

- (i) *a neighborhood of L admits a Weinstein structure with L as its skeleton,*
- (ii) *the germ near L of such a structure is unique up to Weinstein homotopy with fixed skeleton L .*

An assumption that L is an Ac-building rather than a general Ac-space simplifies the proof, but is not necessary. We will not need the more general result and will not prove it in this paper.

3.3.2. Existence.

Proof of (i). Consider the building presentation $L_k \rightarrow L_{k-1} \rightarrow \cdots \rightarrow L_1$ of L . We will inductively construct a cotangent building structure with L as its skeleton. Using Theorem 3.12 this structure can be transported to $\mathcal{O}p L \subset W$.

We argue by induction in the number k of blocks. Suppose that we already constructed a cotangent building $W_{>1} := B_k \rightarrow \cdots \rightarrow B_2$ with the skeleton $L_{>1}$ which consists of blocks $B_j = \mathcal{T}^*L_j$. By Lemma 3.26 there exists a boundary face Y of $L_{>1}$ such that L is obtained by attaching $L_{>1}$ to L_1 using a front embedding $Y \rightarrow L_1$, i.e. an embedding which factors as $Y \xrightarrow{\phi} T^*L_1 \setminus L_1$, where ϕ as an embedding onto a Legendrian arboreal. The Ac-space Y serves as the nucleus Q of a boundary face of $W_{>1}$. The embedding ϕ extends to an embedding $\Phi : Q \rightarrow T^*L_1 \setminus L_1$ as a Weinstein hypersurface. The required building L can be now obtained by the vertical gluing of $W_{>1}$ to \mathcal{T}^*L_1 . \square

3.3.3. *Liouville fields on cotangent bundles.* Before proving the second part of Theorem 3.27 we need to review some facts about Liouville fields on cotangent bundles, see also [CE12, Sect. 12.3].

Let Y be an n -dimensional bc-manifold, T^*Y its cotangent bundle with the standard Liouville form $\lambda_{\text{st}} = pdq$ and symplectic form $\omega_{\text{st}} = d(pdq)$, and Liouville field $Z_{\text{st}} = p \frac{\partial}{\partial p}$. Any other Liouville form λ has the form $\lambda_{\text{st}} - dH$ for a function $H : T^*M \rightarrow \mathbb{R}$. Respectively, the Liouville field Z of λ is given by $Z = Z_{\text{st}} + X_H$, where X_H is the Hamiltonian field of H .

We will always suppose in this section that Z is tangent to Y . This is equivalent to the condition $H|_Y = 0$. Denote $Z' := Z|_Y$, $H' = H|_Y$.

Lemma 3.28. *Let $\lambda = pdq - dH$ a Liouville form on T^*Y with $H|_Y = 0$. Let $a \in Y$ be a zero of $Z' = Z|_Y$. Choose local canonical coordinates $q_1, \dots, q_n, p_1, \dots, p_n$ centered at a . Write $H = \frac{1}{2} \sum_{i,j=1}^n a_{ij} p_i p_j + \sum_{i,j=1}^n b_{ij} p_i q_j + o(|p|^2 + |q|^2)$. Then*

$$Z' = \sum_{i,j=1}^n b_{ij} q_j \frac{\partial}{\partial q_j} + O(|q|^2), Z = \left(\sum_{i,j=1}^n b_{ij} q_j + \sum_{i,j=1}^n a_{ij} p_j \right) \frac{\partial}{\partial q_i} + \sum_{i,j=1}^n (\delta_{ij} - b_{ij}) p_i \frac{\partial}{\partial p_j} + O(|p|^2 + |q|^2).$$

The matrix of $d_a Z'$ is equal to B and the matrix of $d_a Z$ in the basis $\frac{\partial}{\partial q}, \frac{\partial}{\partial p}$ has the form

$$\begin{pmatrix} B & A \\ 0 & I - B^T \end{pmatrix}$$

where $B = (b_{ij})$, $A = (a_{ij})$.

Proof. Direct computation. \square

Definition 3.29. A zero $a \in Y$ of Z' is called *transversely non-degenerate* if the matrix $I - B^T$ has no pure imaginary eigenvalues.

Lemma 3.30. *Let (W, Z) be a Liouville structure with an arboreal skeleton X . Let $a \in X$ be a zero of Z and Y the smooth piece corresponding to the root of the arboreal singularity at the point a . Suppose that a is transversely non-degenerate with respect to Y . Denote by Z' the restriction $Z|_Y$. Then the eigenvalues λ_j of the differential $d_a Z'$ have real parts < 1 .*

Proof. By assumption Z is tangent to Y , and hence Z can be written near the zero a as

$$Z = \left(\sum_{i,j=1}^n b_{ij} q_j + \sum_{i,j=1}^n a_{ij} p_j \right) \frac{\partial}{\partial q_i} + \sum_{i,j=1}^n (\delta_{ij} - b_{ij}) p_i \frac{\partial}{\partial p_j} + O(|p|^2 + |q|^2).$$

The transverse non-degeneracy condition means the matrix $C := I - B^T = (\delta_{ij} - b_{ji})$ does not have pure imaginary eigenvalues. If $\operatorname{Re} \lambda < 0$ for one of eigenvalues of C , then the stable manifold of a must contain a curve tangent to the corresponding eigenvector. But the arboreal X cannot contain any (two-sided) curves transverse to the root smooth piece Y . \square

Lemma 3.31. *Let Y be a compact manifold, v a vector field, and $\ell : Y \rightarrow \mathbb{R}_+$ a positive Lyapunov function for v . Let A be the set of critical points of ℓ . Then for any neighborhood $U \supset A$ and any constant $C > 0$ there exists a function $\theta : \mathbb{R} \rightarrow \mathbb{R}$ with $\theta' > 0$ such that on $Y \setminus U$ there holds:*

$$(2) \quad d(\theta \circ \ell)(v) > C \theta \circ \ell$$

Proof. Let $c_0 < \dots < c_k$ be critical values of Y . Choose a sufficiently small $\sigma > 0$ and denote $B_j := \{c_j - \sigma \leq \ell \leq c_j + \sigma\}$, $C_j := \{c_j + \sigma \leq \ell \leq c_{j+1} - \sigma\}$. Note that each C_j is a trivial cobordism bounded by regular level sets $\partial_- C_j = \{\ell = c_j + \sigma\}$ and $\partial_+ C_j = \{\ell = c_{j+1} - \sigma\}$. Denote $\delta_0 := [c_0, c_0 + \sigma]$, $\Delta_0 := [c_0 + \sigma, c_1 - \sigma]$, $\delta_1 := [c_1 - \sigma, c_1 + \sigma]$, $\Delta_1 := [c_1 + \sigma, c_2 -$

$\sigma], \dots, \Delta_{k-1} := [c_{k-1} + \sigma, c_k - \sigma], \delta_k := [c_k - \sigma, c_k]$. Define a diffeomorphism $\psi_j : \partial_- C_j \times \Delta_j \rightarrow C_j$ by sending vertical intervals to the corresponding trajectories of v parameterized by ℓ , so that we have $\ell(x, u) = u$, $x \in \partial_- C_j, u \in \Delta_j$. Denote $h(x, u) := d_{x,u}\ell(v)$. We have $d(\theta \circ \ell)(v) = \theta'(u)h(x, u)$. We will successively define θ on intervals. Assuming σ so small that $B_0 \subset U$ we define $\theta(t) = t$ on δ_0 . Identifying C_0 with $\partial_- C_0 \times \Delta_0$ via ψ_0 , the inequality (2) takes the form

$$\theta'(u)h(x, u) > C\theta(u)$$

or

$$\frac{d \ln \theta(u)}{du} > \frac{C}{h(x, u)}.$$

Choose a smooth positive function $\tilde{h}(u) < \min_x h(x, u)$. Solving the equation

$$\frac{d \ln \theta(u)}{du} = \frac{C}{\tilde{h}(u)}.$$

we get $\theta(s) = \theta(c_0 + \sigma)e^{H(u)}$, where $H(u) = \int_0^u \tilde{h}(s) ds$. Note that we can assume that all the trajectories of v in $B_j \setminus U$ begin at $\{\ell = c_j - \varepsilon\}$ and end at $\{\ell = c_j + \varepsilon\}$, i.e. there exists a diffeomorphism $(\partial_- B_j \setminus U) \times \delta_j \rightarrow B_j \setminus U$. Hence we can define θ on $[c_0, c_k]$, by repeating the same argument successively to $\delta_0, \Delta_0, \delta_1, \Delta_1, \dots, \delta_k$. \square

Lemma 3.32. *Let $Z = p \frac{\partial}{\partial p} + X_H$ be a Liouville field on T^*Y . Suppose the vector field $Z' = Z|_Y$ is gradient like for some function $\ell : Y \rightarrow \mathbb{R}$. Suppose for all zeroes $a \in Y$ of the vector field Z' the eigenvalues of the differential $d_a Z'$ have real part < 1 . Then there exists a Riemannian metric on Y such that $Z|p|^2 \geq \varepsilon|p|^2$ on a neighborhood of Y .*

Proof. Let us start with any background metric on Y . Recall that the vector field Z' is gradient like for a function $\ell : Y \rightarrow \mathbb{R}$, which can be assumed positive. According to Lemma 3.32, near each zero we have

$$Z = \left(\sum_{i,j=1}^n b_{ij} q_j + \sum_{i,j=1}^n a_{ij} p_j \right) \frac{\partial}{\partial q_i} + \sum (\delta_{ij} - b_{ij}) p_i \frac{\partial}{\partial p_j} + O(|p|^2 + |q|^2).$$

By assumption, all eigenvalues of the matrix B have real parts < 1 , and hence the matrix $I - B^T$ has eigenvalues with the real part > 0 , which implies

$$(3) \quad Z \cdot |p|^2 > \varepsilon |p|^2$$

on a neighborhood U of the compact set of all zeroes of Z .

Let \widehat{Z} be the Hamiltonian extension of Z' to T^*Y with the Hamiltonian $p(Z'(q))$. Note that $Z = \widehat{Z} + \widetilde{Z}$, where $\widetilde{Z} = O(|p|)$. Hence $\widetilde{Z} \cdot |p|^2 = O(|p|^2)$. We also note that $\widehat{Z} \cdot |p|^2 = O(|p|^2)$. Hence, $Z \cdot |p|^2 = O(|p|^2)$. Denote $\Delta_1 := \max(|\frac{Z \cdot |p|^2}{|p|^2}|)$. Given any point $a \in Y \setminus U$ we choose local coordinates centered at a in a neighborhood U_a and write $|p|^2 = \sum c_{ij}(q) p_i p_j$.

Denote

$$\Delta_2 := \max_{a \in Y \setminus U} \max_{i,j; q \in U_a} |Z' \cdot c_{ij}(q)|.$$

Set $K = \max(\Delta_1, \Delta_2)$. According to Lemma 3.31 there exists a function $\theta : \mathbb{R} \rightarrow [1, \infty]$ such that the function $\theta \circ \ell$ satisfies on $Y \setminus U$ the condition

$$Z' \cdot (\theta \circ \ell) > 3K \theta \circ \ell.$$

Let us pull-back the function ℓ to T^*Y via the projection $T^*Y \rightarrow Y$. We will keep the notation ℓ for the extended function. Note that $\widehat{Z}(p, q) \cdot \ell = Z'(q) \cdot \ell$ and $\widetilde{Z} \cdot \ell = O(|p|)$.

Then we have

$$|Z \cdot (\theta \circ \ell(q))|p|^2 \geq (3K\theta \circ \ell(q) - 2K\theta \circ \ell(q)) |p|^2 + \frac{\partial}{\partial p} \cdot |p|^2 + o(|p|^2) \geq \sigma |p|^2, \quad \sigma > 0,$$

if $|p|$ is small enough.

We need to check that the conformal scaling did not destroy the estimate on U .

$$\begin{aligned} Z(\theta \circ \ell |p|^2) &\geq \varepsilon \theta \circ \ell |p|^2 + (Z \cdot \theta \circ \ell) |p|^2 \\ &= \varepsilon \theta \circ \ell |p|^2 + (\widehat{Z} \cdot \theta \circ \ell) |p|^2 + (\widetilde{Z} \cdot \theta \circ \ell) |p|^2. \end{aligned}$$

But

$$\widehat{Z} \cdot \theta \circ \ell(p, q) = \widehat{Z}' \cdot \ell(q) \geq 0,$$

while

$$|\widetilde{Z} \cdot (\theta \circ \ell)| \leq C|p|.$$

Hence, we get

$$Z \cdot (\theta \circ \ell |p|^2) \geq \varepsilon \theta \circ \ell |p|^2 - C|p|^3,$$

and for $|p| < \frac{1}{2C}$ we get $Z \cdot (\theta \circ \ell |p|^2) \geq \varepsilon \theta \circ \ell |p|^2$ for a reduced ε . Hence, rescaling the metric $|p|^2$ with the conformal factor $\theta \circ \ell$ and possibly reducing the size of the neighborhood of Y we get the required metric which satisfies the inequality $Z|p|^2 \geq \varepsilon |p|^2$. \square

Lemma 3.33. *Let Z, ℓ and the norm $|\cdot|$ be as in Lemma 3.32 We pull-back ℓ to T^*Y via the projection $T^*Y \rightarrow Y$ and will keep the notation ℓ for the extended function. Then there exists a constant $K > 0$ such that the function $\widehat{\phi} := \ell + K|p|^2$ is a Lyapunov function for Z on $\mathcal{O}_p Y$.*

Proof. It is sufficient to check the Lyapunov condition only in a neighborhood of the zero locus A of Z . Indeed, elsewhere it is true by assumption along Y , and hence, by openness of the taming condition in its neighborhood. As in the proof of Lemma 3.32 we extend Z' to T^*Y as a Hamiltonian vector field \widehat{Z} with the Hamiltonian function $\zeta(p, q) = p(Z'(q))$. Denote $\widetilde{Z} := Z - \widehat{Z} - \frac{\partial}{\partial p}$. We also assume that the function ℓ is extended to T^*Y by pulling it back from Y by the projection $T^*Y \rightarrow Y$. We have

$$Z \cdot \widehat{\phi} = KZ \cdot |p|^2 + Z \cdot \ell \geq K\varepsilon |p|^2 + \widehat{Z} \cdot \ell + \widetilde{Z} \cdot \ell.$$

Furthermore,

$$\widehat{Z}(p, q) \cdot \ell = Z'(q) \cdot \ell \geq \varepsilon_1(|d\ell|^2 + |Z'(q)|^2),$$

as ℓ is Lyapunov for Z' .

We also have

$$|\tilde{Z} \cdot \ell| \leq C_1 |d\ell| |p|, \quad |\widehat{Z}(p, q) - Z'(q)| \leq C_2 |d\ell(q)| |p|.$$

Thus,

$$\begin{aligned} d\widehat{\phi}(Z) &\geq K\varepsilon |p|^2 + \varepsilon_2 (|Z'(q)|^2 + |d\ell(q)|^2) - C_3 |d\ell| |p| \\ &\geq \varepsilon_3 (|p|^2 + |d\ell(q)|^2 + |Z'(q)|^2), \end{aligned}$$

if K is chosen large enough. We also have

$$\|Z\|^2 \leq C_2 (|p|^2 + |Z'(q)|^2) \quad \text{and} \quad |d\widehat{\phi}|^2 \leq C_3 (K^2 |p|^2 + |d\ell(q)|^2).$$

Hence,

$$d\widehat{\phi}(Z) > \varepsilon_4 (|Z|^2 + |d\phi|^2).$$

□

3.3.4. Deformations.

Proof of Theorem 3.27(ii). We will construct a Weinstein homotopy between the given Weinstein structure $(W, \omega, \lambda, Z, \phi)$ and the cotangent building structure constructed in the proof of (i). Again, we will argue by induction in the number k of blocks. The inductive argument will simultaneously establish the base case of the induction, which is simpler.

According to Lemmas 3.30 and 3.32 we have $Z|p|^2 > \varepsilon |p|^2$ on $U \setminus L_1$ for a neighborhood $U \supset L_1$. Choose $\delta > 0$ to have $\{|p| \leq 6\delta\} \subset U$. Denote $U_\delta := \{\delta \leq \phi \leq 6\delta\}$,

Assuming $\phi > 0$ and choosing a sufficiently large C we can find a Lyapunov function ψ for Z such that

- $\psi = C_1 |p|^2$ on $\{2\delta < |p| < 5\delta\}$;
- $\psi = C_2 \phi$ on $W \setminus \{|p| > 5\delta\}$;
- $\psi = \phi$ on $\{|p| < \delta\}$;
- $d\psi(Z) > 0$ on $U_\delta \setminus \{2\delta < |p| < 5\delta\}$.

We can apply the same argument to the function $\widehat{\phi} = \ell + K|p|^2$ constructed in Lemma 3.33 to get a function $\widehat{\psi}$ which coincide with $\widehat{\phi}$ on $\{|p| < \delta\}$ and with $C_1 |p|^2$ on $\{2\delta < |p| < 3\delta\}$.

A convex combination of Lyapunov functions for Z is again a Lyapunov function for Z , and hence there is a fixed on $\{|p| \geq 2\delta\}$ deformation of ψ to a Lyapunov function which is equal to $\widehat{\psi}$ on $\{|p| < \delta\}$. We will keep the notation ψ for the deformed function.

Recall that $\lambda|_{\mathcal{O}_p L_1} = pdq - dH$ for a function H vanishing on $L \cap \mathcal{O}_p L_1$. Take a function $\theta : [0, 6\delta] \rightarrow \mathbb{R}_+$ which is equal to 0 on $[0, 2\delta] \cup [5\delta, 6\delta]$ and equal to 1 on $[3\delta, 4\delta]$. Define a family of functions $H_t : U_\delta \rightarrow \mathbb{R}$ by the formula $H_t := (1 - t\theta(|p|))H$. Thus, $H_0 = H$, $H_1 = 0$ on $\{3\delta < |p| < 4\delta\}$ and $H_t = H$ on $\{0 \leq |p| \leq 2\delta\} \cup \{5\delta \leq |p| \leq 6\delta\}$ for all $t \in [0, 1]$. Denote by v the Hamiltonian field of H , by v_t the Hamiltonian field of H_t and by Z_t the Liouville field $v_t + p \frac{\partial}{\partial p}$. Then $v_t = v$ on $\{0 \leq |p| \leq 2\delta\} \cup \{5\delta \leq |p| \leq 6\delta\}$ and $v_t = (1 - t\theta(|p|))v + H\theta'(|p|)w$ on $\{3\delta < |p| < 4\delta\}$, where we denoted by w the Hamiltonian vector field of the Hamiltonian

$|p|^2$. Let us observe that the Liouville field Z_t is tangent to the arboreal L . Then we have $v_t \cdot \psi = v \cdot \psi$ on $\{0 \leq |p| \leq 2\delta\} \cup \{5\delta \leq |p| \leq 6\delta\}$ and

$$v_t \cdot |p|^2 = (1 - t\theta(|p|))v \cdot |p|^2 + H\theta'(|p|)w \cdot |p|^2 = (1 - t\theta(|p|))v \cdot |p|^2$$

on $\{3\delta < |p| < 4\delta\}$. On the other hand, on $\{3\delta < |p| < 4\delta\}$ we have

$$Z_t \cdot |p|^2 = (1 - t\theta(|p|))Z \cdot |p|^2 + 2t\theta'(|p|)|p|^2 > 0.$$

Thus, Z_t is gradient like for ψ for all $t \in [0, 1]$.

According to Lemma 3.32 the vector field Z also satisfies the condition $Z \cdot |p|^2 \geq \varepsilon|p|^2$. Hence, the vector field $p\frac{\partial}{\partial p} + v_1$ is gradient like for the function $|p|^2 + \psi$. Define Z_t for $t \in [1, 2]$ on $\{|p| \leq 4\delta\}$ by the formula

$$Z_t := p\frac{\partial}{\partial p} + (2-t)v_1.$$

We claim that the Liouville field Z_t is gradient like for the function $\psi_t := |p|^2 + (2-t)\psi$. Away from the 0-section both vector fields, $(2-t)Z_1$ and $(t-1)p\frac{\partial}{\partial p}$ are gradient like for $|p|^2$, and hence, so is $Z_t = (2-t)Z_1 + (t-1)p\frac{\partial}{\partial p}$. Hence, it is sufficient to check the Lyapunov condition in an arbitrary small neighborhood of the 0-section.

Note that we have $p\frac{\partial}{\partial p} \cdot \ell = 0$. Hence,

$$\begin{aligned} Z_t \cdot \psi_t &= ((t-1)p\frac{\partial}{\partial p} + (2-t)Z_1) \cdot (|p|^2 + (2-t)\psi) \\ &= (t-1)K(t)|p|^2 + (2-t)^2Z_1 \cdot \psi \geq 2(t-1)K(t)|p|^2 + (2-t)^2\varepsilon(|Z_1|^2 + |d\psi|^2). \end{aligned}$$

Furthermore, $|Z_t|^2 \leq 2((2-t)^2|Z_1|^2 + (t-1)^2|p|^2)$ and $|d\psi_t|^2 \leq 2(|p|^2 + (2-t)^2|d\psi|^2)$. Therefore,

$$Z_t \cdot \psi_t \geq \varepsilon_1(|Z_t|^2 + |d\psi_t|^2).$$

We also note that the Liouville field Z_t is tangent to the arboreal L for all t .

All functions ψ_t are proportional to $|p|^2$ on $\{3\delta < |p| < 4\delta\}$, and hence can be extended as proportional to ψ Lyapunov functions for Z_t for all $t \in [0, 2]$.

Finally, we observe that Z_2 is equal to $Z_{st} = p\frac{\partial}{\partial p}$ on $\{|p| \leq 4\delta\}$, and therefore one can use splitting procedure, as it is described in Section 2.6 to split the Weinstein manifold along the hypersurface $\{|p| = 4\}$ into the block T^*L_1 and a Wc-manifold with the skeleton $L_{>1}$. Hence the induction hypothesis completes the proof. \square

4. POSITIVE COTANGENT BUILDINGS

In this section we discuss the positivity relation, first for Lagrangian planes and then for cotangent buildings.

4.1. The positivity relation.

4.1.1. *Polarizations.* Let (V, ω) be a symplectic vector space and let $\tau, \nu \subset V$ be transverse linear Lagrangian subspaces. We will refer to the pair (τ, ν) as a *polarization* of V , and call τ

and ν the respective *horizontal* and *vertical* spaces of the polarization. A polarization (τ, ν) provides a canonical linear isomorphism $\nu \simeq \tau^*$ given by $\nu \ni v \mapsto \iota(v)\omega|_{\tau} \in \tau^*$, and linear symplectomorphisms $V \simeq \tau \oplus \nu \simeq T^*\tau$, where we define $T^*\tau = \tau \oplus \tau^*$.

Let $L \subset V$ be a Lagrangian subspace transverse to ν . Via the identification $V \simeq T^*\tau$, we can regard L as the graph of the differential of a quadratic form denoted by $L^{(\tau, \nu)} : \tau \rightarrow \mathbb{R}$. By construction, this differential is the composition of the canonical maps

$$dL^{(\tau, \nu)} : \tau \simeq V/\nu \simeq L \rightarrow V \rightarrow V/\tau \simeq \nu.$$

Definition 4.1. Let (V, ω) be a symplectic vector space with polarization (τ, ν) . Let $L \subset V$ be a Lagrangian subspace transverse to the vertical space ν . We write

$$L \underset{\nu}{\succ} \tau \quad (\text{resp. } L \underset{\nu}{\prec} \tau)$$

when the quadratic form $L^{(\tau, \nu)} : \tau \rightarrow \mathbb{R}$ is positive (resp. negative) definite.




FIGURE 4.1. The above Lagrangians L_3 and L_4 in \mathbb{R}^2 are given by positive-definite and negative-definite quadratic forms respectively, hence we have $L_3 \underset{L_2}{\succ} L_1$ and $L_4 \underset{L_2}{\prec} L_1$.

Note that $L \underset{\nu}{\succ} \tau$ or $L \underset{\nu}{\prec} \tau$ implies that L and τ are transverse.

One can check positivity by Hamiltonian reduction to smaller symplectic vector spaces, in particular two-dimensional symplectic planes. Given a coisotropic $W \subset V$, and any subspace $L \subset V$, denote the symplectic reduction by $[L]^W = (L \cap W)/W^\perp \subset W/W^\perp = [V]^W$. Note that if L is Lagrangian then $[L]^W$ is Lagrangian in $[V]^W$ even when the intersection $L \cap W$ is not transverse.

Lemma 4.2. Let $\Lambda(V)$ denote the Lagrangian Grassmanian of a symplectic space V . Then the map $\pi_W : \Lambda(V) \rightarrow \Lambda([V]^W)$ defined by the formula $\pi_W(L) = [L]^W$, $L \in \Lambda(V)$, is continuous.

Proof. This is clear. □

Lemma 4.3. Let V be a symplectic vector space of dimension $2n > 0$, (τ, ν) a polarization of V and $L \subset V$ a linear Lagrangian subspace. The following conditions are equivalent:

(i) *Positivity:*

$$L \succ_{\nu} \tau.$$

(ii) *Positivity restricted to subspaces:*

$$[L]^W \succ_{[\nu]^W} [\tau]^W,$$

for all coisotropics $W \subset V$ containing ν with $\dim W > n$.

(iii) *Positivity restricted to lines:*

$$[L]^W \succ_{[\nu]^W} [\tau]^W,$$

for all coisotropics $W \subset V$ containing ν with $\dim W = n + 1$.

(iv) *Positivity of all reductions:*

$$[L]^W \succ_{[\nu]^W} [\tau]^W,$$

for all coisotropics $W \subset V$.

(v) *Sylvester's criterion:*

$$\wedge^{\text{top}} [L]^{W_i} \succ_{\wedge^{\text{top}} [\nu]^{W_i}} \wedge^{\text{top}} [\tau]^{W_i},$$

for any flag of coisotropics

$$\nu \subset W_{n+1} \subset \cdots \subset W_{2n-1} \subset W_{2n} = V$$

with $\dim W_i = i$.

Proof. Equivalence of (i) with (ii) (resp. (iii)) means that a quadratic form is positive definite if and only if it is positive definite on all subspaces (resp. on all lines), which is clear. Clearly (iv) implies (iii), and the converse implication follows from Lemma 4.2 together with the two fact that the positivity condition $L \succ_{\nu} \tau$ implies that $[\nu]^W, [L]^W$ and $[\tau]^W$ are pairwise transverse. Finally, (v) is a reformulation of Sylvester's criterion that a quadratic form is positive definite if and only if its leading principal minors have positive determinant. \square

4.1.2. *Cyclic symmetries.* Note that given a polarization (τ, ν) of V , we also have the polarization (ν, τ) . Additionally, if $L \subset V$ is transverse to ν , we can regard (L, ν) as a polarization.

Lemma 4.4. *The following properties hold.*

(i) *Duality of polarization:* $L \succ_{\nu} \tau \iff L \prec_{\tau} \nu$.

(ii) *Exchange of graphs:* $L \succ_{\nu} \tau \iff \tau \prec_{\nu} L$.

(iii) *Transitivity:* suppose $K \subset V$ is another Lagrangian subspace transverse to ν . Then

$$K \succ_{\nu} L, L \succ_{\nu} \tau \implies K \succ_{\nu} \tau$$

Proof. Let ω denote the symplectic form of V . For (i), by construction, $L^{(\tau,\nu)}(X) = \omega(dL^{(\tau,\nu)}X, X)$, $X \in \tau$. Since $dL^{(\tau,\nu)} = (dL^{(\nu,\tau)})^{-1}$, for $Y \in \nu$ we have

$$L^{(\nu,\tau)}(Y) = \omega((dL^{(\tau,\nu)})^{-1}Y, Y) = -\omega(Y, (dL^{(\tau,\nu)})^{-1}Y) = -L^{(\tau,\nu)}(dL^{(\tau,\nu)}Y).$$

Hence if $L^{(\tau,\nu)}$ is positive definite, then $L^{(\nu,\tau)}$ is negative definite and vice versa. Next, let $K \subset V$ be a Lagrangian transverse to ν . Note that $K^{(\tau,\nu)} = K^{(L,\nu)} + L^{(\tau,\nu)}$ under the isomorphism $\tau \simeq V/\nu \simeq L$. Both (ii) and (iii) follow immediately (for (ii) take $K = \tau$ so that $K^{(\tau,\nu)} = 0$ and hence $\tau^{(L,\nu)} = -L^{(\tau,\nu)}$). \square

Lemma 4.5. *Suppose $L_1, L_2, L_3 \subset V$ are Lagrangian subspaces transverse to ν . Then for any permutation $\sigma \in \Sigma_3$ we have*

$$L_1 \underset{L_3}{\succ} L_2 \iff L_{\sigma(1)} \underset{L_{\sigma(3)}}{\overset{\sigma}{\succ}} L_{\sigma(2)}$$

where $\overset{\sigma}{\succ}$ denotes \succ for σ an even composition of transpositions, and \prec for σ an odd composition of transpositions.

Proof. The case of $\sigma = (23)$ (resp. $\sigma = (12)$) is part (i) (resp. (ii)) of Lemma 4.7. These permutations generate. \square

One can interpret Lemma 4.7, and in turn Lemma 4.5, as saying the ternary relation \succ provides a *partial cyclic order* on triples of Lagrangian subspaces.

Definition 4.6. We say an ordered list $L_1, \dots, L_m \subset V$ of Lagrangian planes is \succ -*cyclically ordered* (resp. \prec -*cyclically ordered*) when they are pairwise transverse and satisfy

$$L_{i_2} \underset{L_{i_1}}{\succ} L_{i_3} \quad (\text{resp. } L_{i_2} \underset{L_{i_1}}{\prec} L_{i_3})$$

whenever $i_1, i_2, i_3 \in \{1, \dots, m\}$ are cyclically ordered in \mathbb{Z}/m . We will also write $L_1 \succ L_2 \succ \dots \succ L_m$ (resp. $L_1 \prec L_2 \prec \dots \prec L_m$).

We record the following useful assertions for future reference. The assertions and their proofs hold in any partial cyclic order.

Lemma 4.7. *The positivity relation satisfies the following properties.*

- (i) *Suppose $L_1 \underset{\nu}{\prec} L_2$ and either $\tau \underset{\nu}{\prec} L_1$ or $L_2 \underset{\nu}{\prec} \tau$. Then $L_1 \underset{\tau}{\prec} L_2$.*
- (ii) *Suppose $L_3 \underset{L_4}{\succ} L_2 \underset{L_4}{\succ} L_1$. Then $L_2 \underset{L_3}{\succ} L_1$. If additionally $L \underset{L_3}{\succ} L_2$ then $L \underset{L_4}{\succ} L_1$.*
- (iii) *If $L_- \underset{L_2}{\succ} L_1$ and $L_+ \underset{L_1}{\succ} L_2$, then $L_- \underset{L_2}{\succ} L_+$.*

Proof. (i) By cyclic symmetry applied to $L_1 \underset{\nu}{\prec} L_2$, we have $L_2 \underset{L_1}{\prec} \nu$ and $\nu \underset{L_2}{\prec} L_1$. Suppose $\tau \underset{\nu}{\prec} L_1$. Then by cyclic symmetry, we have $\nu \underset{L_1}{\prec} \tau$ and hence by transitivity $L_2 \underset{L_1}{\prec} \tau$. By transpositional symmetry, we conclude $L_2 \underset{\tau}{\succ} L_1$.

Similarly, suppose $L_2 \underset{\nu}{\prec} \tau$. Then by cyclic symmetry, we have $\tau \underset{L_2}{\prec} \nu$ and hence by transitivity $\tau \underset{L_1}{\prec} L_1$. By transpositional symmetry, we conclude $L_1 \underset{\tau}{\prec} L_2$.




FIGURE 4.2. A cyclic ordering on a tuple (L_1, \dots, L_n) can be thought of as an embedding into S^1 preserving the cyclic order of all triples.

(ii) For the first assertion, $L_3 \succ_{L_4} L_2$ (resp. $L_2 \succ_{L_4} L_1$) implies $L_3 \prec_{L_2} L_4$ (resp. $L_4 \prec_{L_2} L_1$) by transpositional symmetry. Hence by transitivity we have $L_3 \prec_{L_2} L_1$, and so by transpositional symmetry $L_2 \succ_{L_3} L_1$.

For the second, suppose L satisfies $L \succ_{L_3} L_2$. By the previous part, we have $L_2 \succ_{L_3} L_1$, hence $L \succ_{L_3} L_1$ by cyclic symmetry. By transpositional symmetry, we then have $L \prec_{L_1} L_3$. By assumption, we have $L_3 \succ_{L_4} L_1$ and hence by transpositional symmetry $L_3 \prec_{L_1} L_4$ so by transitivity $L \prec_{L_1} L_4$. Finally, by transpositional symmetry, we obtain $L \succ_{L_4} L_1$.

(iii) By cyclic symmetry, we have $L_1 \succ_{L_2} L_+$, and hence by transitivity $L_- \succ_{L_2} L_+$. \square

4.1.3. *Positive zones.* It is useful to reformulate positivity more geometrically.

Definition 4.8. Let (V, ω) be a symplectic vector space with polarization (τ, ν) . Let $\Lambda(V)$ be the Lagrangian Grassmannian of V . Denote by $C(\tau, \nu) \subset \Lambda(V)$ the subset consisting of those Lagrangians $L \subset V$ transverse to ν and satisfying $L \succ_{\nu} \tau$. We refer to $C(\tau, \nu)$ as a *positive zone*, and its closure, denoted by $\overline{C}(\tau, \nu)$, as a *closed positive zone*.

Remark 4.9. As justified by part (ii) of the following lemma, we will regard any single Lagrangian $L \in \Lambda(V)$ itself as a (degenerate) closed positive zone.

Lemma 4.10. *The positive zone $C(\tau, \nu) \subset \text{Gr}(V)$ satisfies the following properties.*

- (i) $C(\tau, \nu)$ (resp. $\overline{C}(\tau, \nu)$) is non-empty, open (resp. closed), and geodesically convex for any homogenous metric on $\text{Gr}(V)$, hence contractible.
- (ii) If τ approaches ν along a geodesic in $\Lambda(V)$, then $C(\tau, \nu)$ and $\overline{C}(\tau, \nu)$ limit to ν itself.




FIGURE 4.3. The *positive zone* of a polarization (τ, ν)

Proof. (i) is immediate from usual statements about quadratic forms. For (ii), we can put everything into a standard position so that we are contracting positive definite quadratic forms towards zero. \square

Lemma 4.11. *The following properties hold.*

- (i) Suppose $L_2 \in C(L_1, \nu)$ and either $L_1 \in C(\tau, \nu)$ or $L_2 \in C(\tau, \nu)$. Then $L_2 \in C(L_2, \tau)$.
- (ii) Suppose $L_3 \in C(L_2, L_4)$, $L_2 \in C(L_1, L_4)$. Then $L_2 \in C(L_1, L_3)$ and $C(L_2, L_3) \subset C(L_1, L_4)$.
- (iii) If $L_- \in C(L_1, L_2)$ and $L_+ \in C(L_2, L_1)$, then $L_- \in C(L_+, L_2)$.

Proof. This follows immediately from Lemma 4.7 by unwinding the definition. \square

Lemma 4.12. *Let $\mathcal{L} \subset \Lambda(V)$ be a non-empty set of Lagrangian planes each transverse to a fixed Lagrangian plane $L \subset V$. The set $\mathcal{S}(\mathcal{L}, L) \subset \Lambda(V)$ of Lagrangian planes $L_- \subset V$ such that $\mathcal{L} \subset C(L_-, L)$ is convex. If \mathcal{L} is compact, then $\mathcal{S}(\mathcal{L}, L)$ is non-empty, hence contractible.*

Proof. The condition $T \in C(L_-, L)$ is equivalent to $L_- \in C(L, T)$, and thus

$$\mathcal{S}(L, \mathcal{L}) = \bigcap_{T \in \mathcal{L}} C(L, T).$$

Fix a Lagrangian plane $H \subset V$ transverse to L . Then we can identify Lagrangian planes $T \subset V$ transverse to L with quadratic forms Q_T on H , and specifically, the set of Lagrangian planes $C(L, T) \subset \Lambda(V)$ with the convex set of those quadratic forms Q on H satisfying $Q < Q_T$. Hence $\mathcal{S}(L, \mathcal{L})$ is an intersection of convex sets so itself convex.

Finally, when \mathcal{L} is compact, the set of quadratic forms Q_T , for $T \in \mathcal{L}$, is bounded, hence we may choose a quadratic form Q_- such that $Q_- < Q_T$, for all $T \in \mathcal{L}$. Thus $\mathcal{S}(L, \mathcal{L})$ is non-empty. \square

Lemma 4.13. *Let $W \subset V$ be a coisotropic subspace. For any polarization (τ, ν) of V , the reduction map π_W projects the positive zone $C(\tau, \nu) \subset \Lambda(V)$ to the positive zone $C([\tau]^W, [\nu]^W) \subset \Lambda([V]^W)$.*

Proof. For any $L \in \Lambda(V)$ transverse to ν , the pre-composition of the quadratic form $([L]^W)^{([\tau]^W, [\nu]^W)}$ with the reduction map $\tau \rightarrow [\tau]^W$ is equal to the quadratic form $L^{(\tau, \nu)}$. Since $\tau \rightarrow [\tau]^W$ is surjective, the lemma follows. \square

4.2. Positivity of polarizations.

4.2.1. Polarized Legendrians. Recall that a Legendrian embedding $\Lambda \subset \mathcal{T}^*M \setminus M$ is said to be *adapted* to the block \mathcal{T}^*M if $\mathcal{O}_p \partial \Lambda \subset \mathcal{O}_p \partial M$ and for each stratum $P \subset \partial_k M$ we have $\Lambda \cap (\mathcal{O}_p P = \mathcal{T}^*P \times \mathcal{T}^*\mathcal{I}^k) = \Lambda_k \times \mathcal{I}^k$ for a Legendrian $\Lambda_k \subset \mathcal{T}^*P$. Recall also that an adapted Legendrian Λ is called *regular* if $\pi|_\Lambda : \Lambda \rightarrow M$ is an immersion with transverse self-intersections, where $\pi : \mathcal{T}^*M \rightarrow M$ is the cotangent projection.

A *polarization* of a Lagrangian submanifold L of a symplectic manifold (M, ω) is a field $\eta \subset TM|_L$ of Lagrangian planes transverse to L . A polarization of a Legendrian Λ in a contact manifold (V, ξ) is a Lagrangian field $\tau \subset \xi|_\Lambda$ transverse to Λ (recall that ξ has a well-defined conformal symplectic structure; hence, it makes sense to talk about Lagrangian fields in ξ).

Definition 4.14. A polarization of a Legendrian $\Lambda \subset \mathcal{T}^*M \setminus M$ is a field of Lagrangian planes $\mu \subset \mathcal{T}^*M|_\Lambda$ which projects to a polarization of the projection of Λ to $S^\infty M$.

Note that μ has to be tangent to the Liouville field Z .

In particular, the Lagrangian distribution ν_M which projects to the Legendrian distribution ℓ_M tangent to the spherical Legendrian fibration of $S^\infty M$, defines a canonical polarization of any *regular* Legendrian in $\mathcal{T}^*M \setminus M$. We will refer to this polarization as *tautological*.

The space of polarizations of a given Legendrian is contractible. Any given polarization of a Legendrian admits a ribbon, whose tangent planes to fibers along the 0-section form the given polarization. The space of germs of ribbons for a given polarization is contractible. Hence the space of germs of ribbons for a given Legendrian is contractible.

Definition 4.15. A polarization μ of a *regular* Legendrian $L \subset \mathcal{T}^*M \setminus M$ is called *positive* if $\ell_M \in C(TL, \mu)$ at any point of L , where ℓ_M is the tautological polarization.

4.2.2. Positivity implies transversality of conormals.

Definition 4.16. Let (W, λ) be a Wc-manifold. A *polarization* of W is a global field of Lagrangian planes $\eta \subset TW$.

Let T^*M be the cotangent bundle of a compact manifold with corners, $\tau = TM$ the tangent field along M and $\nu = \ker(d\pi)$ the vertical field, where $\pi : T^*M \rightarrow M$ is the cotangent bundle projection.

Lemma 4.17. *Let $L \subset \mathcal{T}^*M \setminus M$ be the Lagrangian cone over a regular Legendrian, and η a polarization of \mathcal{T}^*M whose restriction to the 0-section M is positive with respect to the polarization (τ, ν) . Then η is transverse to the cone L on a sufficiently small neighborhood of $M \subset \mathcal{T}^*M$.*

The statement is a corollary of the following linear algebra lemma.

Lemma 4.18. *Let Q be a quadratic form and $L_Q = \{q = Ap\} \subset T^*\mathbb{R}^n$ the corresponding linear Lagrangian. If Q is positive or negative definite then for any co-oriented hyperplane $H \subset \mathbb{R}^n$ through the origin, its conormal $T_H^*\mathbb{R}^n \subset T^*\mathbb{R}^n$ is transverse to L_Q . Conversely, if L_Q is transverse to conormal $T_H^*\mathbb{R}^n$ for all co-oriented hyperplanes $H \subset \mathbb{R}^n$ through the origin, then Q is positive or negative definite.*

Proof. Indeed, the condition $(p, q) \in L_Q \cap \tau^*$ and $(p, q) \neq 0$ is equivalent to $Q(p) = 0$, □

Proof of Lemma 4.17. Indeed, the Liouville cone is the conormal of the immersed front projection of Λ , and hence we can apply Lemma 4.18 in a sufficiently small neighborhood of the 0-section. □




FIGURE 4.4. For a non positive-definite quadratic form such as $\eta = x^2 - y^2$, the conormal $T_H^*\mathbb{R}^2$ of a smooth hypersurface $H \subset \mathbb{R}^2$ will be tangent to η whenever TH is tangent to the light-cone $\{x = y\}$.

4.3. Positive cotangent buildings.

4.3.1. *Reductions.* Let W be a cotangent building with blocks $B_j = \mathcal{T}^*M_j, j \in J$. The notions we introduce in this section will depend on sizes of blocks, so we deviate here from the point of view of germs and take B_j to be a fixed defining domain for T^*M_j . Set $B_j^\circ = B_j \setminus M_j$, for $j \in J$. For each $i \in J$, consider the line bundle $\text{Span}(Z_i) \subset TB_i^\circ$ generated by Z_i , let

$\zeta_i = \text{Span}(Z_i)^\perp / \text{Span}(Z_i) \rightarrow B_i^\circ$ denote the corresponding symplectic normal bundle, and consider the natural reduction diagram

$$TB_i^\circ \xleftarrow{\mathbf{i}} \text{Span}(Z_i)^\perp \xrightarrow{\mathbf{p}} \zeta_i$$

Given a linear subspace $\nu \subset TB_i^\circ$, let $[\nu]^i := [\nu]^{\text{Span}(Z_i)^\perp} \subset \zeta_i$ be the reduction along the above correspondence

$$[\nu]^i = \mathbf{p}(\mathbf{i}^{-1}(\nu)) = ((\nu + \text{Span}(Z_i)) \cap \text{Span}(Z_i)^\perp) / \text{Span}(Z_i).$$

We will refer to $[\nu]^i$ as the i -reduction of ν . More generally, given a multi-index $I = (i_1 < \dots < i_m) \subset J$ (which could be empty), consider the natural reduction diagram

$$T(B_{i_1}^\circ \cap \dots \cap B_{i_m}^\circ) \xleftarrow{\mathbf{i}} \text{Span}(Z_{i_1}, \dots, Z_{i_m})^\perp \xrightarrow{\mathbf{p}} \zeta_I$$

$$\zeta_I = \text{Span}(Z_{i_1}, \dots, Z_{i_m})^\perp / \text{Span}(Z_{i_1}, \dots, Z_{i_m}).$$

Given a linear subspace $\nu \subset TB_i^\circ$, we refer to $[\nu]^I := [\nu]^{\text{Span}(Z_{i_1}, \dots, Z_{i_m})^\perp} = \mathbf{p}(\mathbf{i}^{-1}(\nu)) \subset \zeta_I$ as the I -reduction of ν . If $I = \emptyset$, then $[\nu]^I = \nu$.

4.3.2. Positivity. Let $W = \bigcup_j B_j$ be a cotangent building. Since the sizes of the blocks B_j are fixed, we can assign a *type* to any point of $\text{Skel}(W) = \bigcup_j \mathring{M}_j$ as follows:

Definition 4.19. We say that a point $a \in \mathring{M}_i$ is of *type* $I = (i_1 < \dots < i_m)$, where $i_m < i$, if $a \in B_{i_1} \cap \dots \cap B_{i_m}$ and $a \notin B_j$, for $j \notin I$ and $j < i$.

Note that we allow the type I to be empty. For any multi-index $I = (i_1 < \dots < i_m) \subset J$ and integer s , where $1 \leq s \leq m$, denote $I(s) = (i_s < \dots < i_t) \subset I$.

Definition 4.20. A cotangent building $W = \bigcup_{j=1}^k B_j$ is *positive* if for any point $a \in \mathring{M}_j$ of type $I = (i_1 < \dots < i_m)$, $I \neq \emptyset$ and any $s \leq m$ the tuple of Lagrangian planes

$$[T_a M_j]^{I(s)}, [\nu_j(a)]^{I(s)}, [\nu_{i_m}(a)]^{I(s)}, \dots, [\nu_{i_s}(a)]^{I(s)}$$

is \prec -cyclically ordered.

Definition 4.21. Let W be a cotangent building W with the notation as above. A Lagrangian distribution η on W is called *positive* if it is transverse to all the 0-sections M_i and at any point $a \in \mathring{M}_i$ of type I (which could be empty), we have $[\nu_i(a)]^I \in C([T_a M_i]^I, [\eta(a)]^I)$.

4.3.3. Existence and uniqueness of positive distributions. Positive distributions exist and are unique up to a contractible choice on any positive cotangent building. Before we prove this we need two lemmas.

Lemma 4.22. Let $B = \mathcal{T}^*M$ be a cotangent block and τ any Lagrangian distribution extending TM . Let $\Sigma \subset \mathcal{T}^*M$ be a conormal of a co-oriented hypersurface $H \subset M$. For any Lagrangian plane $T \in T_a B$, $a \in B \setminus M$ denote by $\bar{T} := [T]^{\text{Span}(Z)^\perp}$ the reduction with respect




FIGURE 4.5. The positivity condition for the interaction of two blocks.

to the conormal of the Liouville field Z . Then the angle between $\overline{T\Sigma(a)}$ and $\overline{\tau(a)}$ converges to 0 as $a \rightarrow M$.

Proof. The statement is local so we can assume that $M = \mathbb{R}^n$, $H = \{q_n = 0\}$, $a = 0$ and $\Sigma = \{q_n = 0, p_j = 0, j < n\}$. Moreover, it is sufficient to prove the lemma for a single particular extension τ , and hence we can choose $\tau = \text{Span}(\partial_{q_1}, \dots, \partial_{q_n})$. The Liouville field is $Z = \sum_i p_i \partial_{p_i}$, hence $Z|_{\Sigma} = p_n \partial_{p_n}$. Therefore $\text{Span}(Z)^\perp|_{\Sigma} = \text{Span}(\partial_{q_1}, \dots, \partial_{q_{n-1}}, \partial_{p_1}, \dots, \partial_{p_n})$, $T\Sigma \cap Z^\perp|_{\Sigma} = \text{Span}(\partial_{q_1}, \dots, \partial_{q_{n-1}}, \partial_{p_n})$ and $\tau \cap Z^\perp|_{\Sigma} = \text{Span}(\partial_{q_1}, \dots, \partial_{q_{n-1}})$ so in fact $\overline{T\Sigma}$ and $\overline{\tau}$ are identical along Σ . \square

Lemma 4.23. *Let $B_1 = \mathcal{T}^*M_1$ be the base block of a positive cotangent building W and η a distribution on $\mathcal{O}p M_1$ such that at any point $a \in M_1$ the triple $T_a M_1, \nu_1(a), \eta(a)$ is cyclically \prec -ordered. Then η is positive for W on $\mathcal{O}p M_1$.*

Proof. Take any extension τ of the distribution TM_1 to $B_1 = \mathcal{T}^*M_1$. For any point $a \in B_1 \cap \dot{M}_j$ of type $I = (i_1 = 1 < i_2 < \dots < i_k), i_k < j$, the tuple

$$[T_a M_j]^I, [\nu_j(a)]^I, \dots, [\nu_1]^I$$

is cyclically \prec -ordered. On the other hand, if the block B_1 is sufficiently thin then the triple

$$[\tau(a)]^1, [\nu_1(a)]^1, [\eta(a)]^i$$

is also cyclically \prec -ordered. Hence Lemma 4.22 implies that

$$[T_a M_j]^1, [\nu_1(a)]^1, [\eta(a)]^i$$

is cyclically \prec -ordered as well. Therefore,

$$[T_a M_j]^I, [\nu_j(a)]^I, \dots, [\nu_1]^I, [\eta(a)]^I$$

is cyclically \prec -ordered, i.e. η is positive for W on B_1 . \square

Let us denote by $\text{Pos}(W)$ the space of positive distributions on a positive cotangent building $W = \bigcup_j B_j$.

Proposition 4.24. *For any positive cotangent building W the space $\text{Pos}(W)$ is non-empty and (weakly) contractible.*

Proof. We will show that $\text{Pos}(W) \neq \emptyset$ using the convexity of the space of positive definite quadratic forms. The weak contractibility claim is a parametric version of the same argument. We argue by induction on the number of blocks.

For $W_{\leq 1} = \mathcal{T}^*M_1$ a positive distribution from $\text{Pos}(W)$ restricted to M_1 can be viewed as a field of positive definite quadratic forms on the canonical polarization ν_1 . Take any such field η and its arbitrary extension to \mathcal{T}^*M_1 . According to Lemma 4.23 the distribution η satisfies the positivity condition for W if the block B_1 is chosen sufficiently thin.

Suppose that we have already constructed η on $W_{< m} = \bigcup_{j < m} B_j$ and let us consider the next block $B_m = \mathcal{T}^*M_m$. Recall that M_m is a manifold with corners. Its corners are enumerated by the types of points in their neighborhood, i.e. if $b \in \partial_k M_m$ then points $a \in \mathring{M}_m$ in a sufficiently small neighborhood of b have a fixed type $I = (i_1 < \dots < i_k)$, where $i_k < m$. The Liouville fields Z_{i_j} of blocks $B_{i_j}, j = 1, \dots, k$, are tangent to \mathring{M}_m and yield the canonical splitting $\mathcal{O}p_{M_m} b = \mathcal{O}p_{\partial_k M_m} \times \mathcal{I}^k$. Consider the restriction of η to $\mathring{M}_m \cap W_{< m}$. It is transverse to M_m and can therefore be viewed as a field of quadratic forms on ν_m . The positivity condition means that:

(†) for any point $a \in \mathring{M}_m$ of type I , the quadratic form $\eta(a)$ is positive definite on the subspace of $\nu_m(a)$ dual to $\text{Span}(Z_{i_1}(a), \dots, Z_{i_k}(a))$.

Recall that the notion of type depends on the thickness of the blocks B_1, \dots, B_{m-1} . Let us take slightly thinner blocks $B'_i \Subset B_i$ and choose a Lagrangian field along \mathring{M}_m that

- (i) coincides on $\mathring{M}_m \cap (W'_{< m} := \bigcup_1^{m-1} B'_i)$ with the restriction of η from $W'_{< m}$,
- (ii) is given by a field of positive definite quadratic forms on ν_m .

Note in particular that (ii) implies that condition (†) holds on $W_{< m} \cap \mathring{M}_m$. Then choosing any extension to $\mathcal{O}p_{\mathring{M}_m}$, and choosing the block B_m sufficiently thin, we get the required positive distribution on $W_{\leq m}$. \square

4.4. Positive cotangent buildings and positive arboreals. We prove in this section that the skeleton of a positive cotangent building is generically a positive arboreal, where throughout this section *generically* means after a C^∞ -small perturbation of the building structure. This corresponds to a C^∞ -small perturbation of the underlying Weinstein structure. Conversely, if a cotangent building has a positive arboreal skeleton, we show the building structure can be adjusted to be made positive without changing the skeleton.

4.4.1. *From positive buildings to positive arboreals.* The key definition is the following:

Definition 4.25. Given an arboreal Lagrangian L in a symplectic manifold X , a Lagrangian distribution η along L is called *positive with respect to L* if the following condition is satisfied. Take any singular point $a \in L$. Let T be the tangent space to the root Lagrangian at a , and T' a tangent plane to any other smooth piece adjacent to a . Then $T' \in C(T, \eta(a))$.

We will later have use for the following reduced variant:

Definition 4.26. Let $L \subset X$ be a positive arboreal Lagrangian and Z a non-zero Liouville vector field tangent to L . A Lagrangian distribution μ along L is called *reduced positive with respect to (L, Z)* if for any singular point $a \in L$ the following condition is satisfied. Let T be the tangent space to the root Lagrangian at a , and T' a tangent plane to any other smooth piece adjacent to a . Then $[T']^\zeta \in C([T]^\zeta, [\eta(a)]^\zeta)$, where $\zeta = \text{Span}(Z)^\perp$.

Lemma 4.27. *Let W be a W -complex $B_k \rightarrow \cdots \rightarrow B_0$ and denote by ν_j the polarization of the block B_j , $j = 0, \dots, k$.*

- (i) *If the skeleton $\text{Skel}(W)$ is arboreal, then for each point $a \in \mathring{B}_j \cap M_i$, $0 \leq j < i \leq k$, the Lagrangian distribution $[\nu_j]^j$ is transverse to $[TM_j]^j$. Conversely, if for each point $a \in \mathring{B}_j \cap M_i$, $0 \leq j < i \leq k$, the Lagrangian distribution $[\nu_j]^j$ is transverse to $[TM_j]^j$ then $\text{Skel}(W)$ is generically arboreal.*
- (ii) *If the skeleton $\text{Skel}(W)$ is positive arboreal, then for each point $a \in \mathring{B}_j \cap M_i \cap M_\ell$, $0 \leq j < i < \ell \leq k$, the triple $[T_a M_\ell]^N, [T_a M_i]^N, [\nu_j]^N$, where $N := \text{Span}(T_a M_i, T_a M_\ell)$, is \leftarrow -cyclically ordered. Conversely, if for each point $a \in \mathring{B}_j \cap M_i \cap M_\ell$, $0 \leq j < i < \ell \leq k$, the triple $[T_a M_\ell]^N, [T_a M_i]^N, [\nu_j]^N$ is \leftarrow -cyclically ordered, then $\text{Skel}(W)$ is generically positive arboreal.*

Proof. The top block B_k is attached to B_{k-1} along a ribbon of a smooth Legendrian, and the transversality condition implies that the front projection to M_{k-1} is an immersion. Generically it has transverse self-intersections, and hence $\text{Skel}(W_{\geq k-1})$ is arboreal in the case (i) and positive arboreal in the case (ii). Continuing by induction we will assume that $\text{Skel}(W_{> j})$ is arboreal. Then the skeleton of the attaching hypersurface V_j of $W_{> j}$ to B_j is arboreal as well. By the axiom on transverse Liouville cones in the definition of arboreal singularities, the transversality condition then implies that generically $\text{Skel}(W_{\geq j})$ is arboreal. In the case (ii) the skeleton is positive arboreal, as can be seen directly from inspection of the normal forms given in [AGEN20a]. The converse direction is straightforward. \square

Corollary 4.28. *The skeleton of a positive cotangent building is generically a positive arboreal.*

4.4.2. *More on polarizations.* We will need the following lemma from linear algebra.

Lemma 4.29. *Let τ, ν denote 0-section and the cotangent fiber in $T^*\mathbb{R}^n$. Given a collection of hyperplanes $\Pi = \{\Pi_1, \dots, \Pi_k\} \subset \mathbb{R}^n$ denote by ν_1, \dots, ν_k their Lagrangian conormals, and*

set $P_j := \text{Span}(\tau, \nu_j)$. Denote by $\mathcal{P}(\Pi)$ the space of Lagrangian planes η transverse to τ such that the triples of lines $([\tau]^{P_j}, [\eta]^{P_j}, [\nu]^{P_j})$ are \prec -cyclically ordered for all $j = 1, \dots, k$. Denote by \mathcal{P}_+ the subspace of $\mathcal{P}(\Pi)$ consisting of those Lagrangian planes such that the triple (τ, ν, η) is \prec -cyclically ordered. Then $\mathcal{P}(\Pi)$ and $\mathcal{P}_+ \subset \mathcal{P}(\Pi)$ are non-empty convex subsets of the affine space of Lagrangian planes transverse to τ .

Proof. Any Lagrangian plane transverse to τ can be viewed as a quadratic form on ν . The subspace \mathcal{P}_+ consists of positive definite quadratic forms, while $\mathcal{P}(\Pi)$ consists of quadratic forms which take positive values on the covectors $A_j \in \nu = \tau^*$ which define the hyperplanes Π_i , $j = 1, \dots, k$. Both spaces are non-empty and convex. \square

Below in this section we view polarizations of a Lagrangian as transverse Lagrangian foliations on its neighborhood. Similarly, we assume that polarizations of a Legendrian Λ are extended to its ribbon, i.e. a given embedding of $\mathcal{S}^*\Lambda$ as a Weinstein hypersurface.

Lemma 4.30. *Let us fix a reference polarization ν of a Lagrangian $L_0 \subset (X, \omega)$. Then the space $\text{Pol}(L_0)$ of all polarizations on L_0 can be viewed as the space of germs along L_0 of symplectomorphisms $T^*L_0 \rightarrow T^*L_0$ which fix L_0 , or alternatively as the space of germs along L_0 of functions on T^*L_0 which vanish on L_0 together with their differential.*

Proof. For any polarization $\eta \in \text{Pol}(L_0)$ there exists a unique symplectomorphism of T^*L_0 which fixes L_0 and which sends ν onto η as foliations. Each function on H on T^*L_0 vanishing on L_0 together with its differential generates a symplectomorphism fixing L_0 as the time 1 map of its Hamiltonian flow. Conversely, each symplectomorphism germ ϕ fixing L_0 defines the required function H by the conditions $\phi^*pdq = pdq + dH$, $H|_{L_0} = 0$. \square

Let $L = C(\Lambda) \cup L_0$ be an arboreal Lagrangian, where $\Lambda \subset \partial_\infty T^*L_0$ is an arboreal Legendrian. A polarization η is called *positive* for L if for any point $\lambda \in L \cap \overline{C(\Lambda)}$ and any Lagrangian plane $T \in T_\lambda(T^*L_0)$ tangent to one of smooth pieces of L different from L_0 the triple $([T_\lambda L_0]^N, [T]^N, [\eta(\lambda)]^N)$ is \prec -cyclically ordered, where $N = \text{Span}(T_\lambda L_0, T)$. Such a polarization is automatically transverse to L on $\mathcal{O}p L_0$ (comp. Lemma 4.17).

Lemma 4.31. *Let $L = C(\Lambda) \cup L_0$ be an arboreal Lagrangian, where $\Lambda \subset \partial_\infty T^*L_0$ is an arboreal Legendrian and η a positive polarization of L_0 . Denote by $\text{Pol}(L, \eta)$ the space of polarizations μ of L_0 for which the corresponding Liouville field is tangent to the cone $C(\Lambda)$ and the triple $(T_\lambda L, \mu(\lambda), \eta(\lambda))$ is \prec -cyclically ordered for any $\lambda \in L_0$. Then $\text{Pol}(L, \eta)$ is non-empty and convex.*

Proof. Viewing $\mu \in \text{Pol}(L, \eta)$ as a function on T^*L_0 the condition $\mu \in \text{Pol}(L, \eta)$ means that μ vanishes on L and that the quadratic part of $\eta - \mu$ along L_0 is positive definite on cotangent fibers. Hence, the convexity claim for $\text{Pol}(L, \eta)$ is straightforward. Let us show that the space $\text{Pol}(L, \eta)$ is non-empty. Take any polarization η' such that the triple $(T_\lambda L_0, \nu(\lambda_0), \eta'(\lambda))$ is \prec -cyclically ordered for all $\lambda \in L_0$. In particular, ζ' is transverse to L on $\mathcal{O}p L_0$.

The tangent cone to $\overline{L \setminus L_0}$ at a point $\lambda \in L_0$ is the union of Lagrangian conormals $\nu_{j,\lambda}$ to a collection $T_{j,\lambda}$ of transverse hyperplanes $\Pi_{j,\lambda} \subset T_\lambda L_0$, $j = 1, \dots, k$. Using Lemma 4.29 we can find a homotopy η_t connecting $\eta_0 = \eta$ and $\eta_1 = \eta'$ such that the triple $([T_\lambda]^{N_{j,\lambda}}, [\nu]^{N_{j,\lambda}}, [\eta_t]^{N_{j,\lambda}})$ is \prec -cyclically ordered for all $0 \leq t \leq 1$. Here we denoted $N_{j,\lambda} := \text{Span}(T_\lambda L_0, \nu_{j,\lambda})$.

In particular, η_t is transverse to the germ of L along L_0 for all $t \in [0, 1]$. There exists a Hamiltonian isotopy ψ_t defined on $\mathcal{O}p L_0$ such that $\psi_t|_{L_0} = \text{Id}$ and $\psi_t(\eta_t) = \eta$, $t \in [0, 1]$. Denote $L^t := \psi_t(L)$. Using a parametric version of the stability theorem which keeps track of polarizations (this is Corollary 3.14 in [AGEN20a]) we can construct for each point $\lambda \in L$ a Hamiltonian isotopy $\phi_{t,\lambda}$ on $\mathcal{O}p \lambda$ such that

- (i) $\phi_{t,\lambda}|_{L_0 \cap \mathcal{O}p \lambda} = \text{Id}$;
- (ii) $\phi_t(\eta|_{\mathcal{O}p \lambda}) = \eta|_{\mathcal{O}p \lambda}$;
- (iii) $\phi_{t,\lambda}(L^t \cap \mathcal{O}p \lambda) = L \cap \mathcal{O}p \lambda$.

Then the polarization $\mu_\lambda := \phi_{1,\lambda} \circ \psi_1(\mu)$ belongs to the space $\text{Pol}(L \cap \mathcal{O}p \lambda, \eta|_{\mathcal{O}p \lambda})$. In view of compactness of L_0 we can cover L_0 by finitely many balls U_j , $j = 1, \dots, N$, centered at the points λ_j such that $\mu_{\lambda_j} \in \text{Pol}(L \cap U_j, \eta|_{U_j})$. Given a partition of unity $\sum_1^N \theta_j = 1$ subordinate to the covering, we define the required element of $\text{Pol}(L, \eta)$ by the formula $\mu = \sum_1^N \theta_j \mu_{\lambda_j}$. \square

4.4.3. From positive arboreals to positive buildings.

Proposition 4.32. *If the skeleton of a cotangent building is positive arboreal, then the cotangent building structure can be deformed to be positive without changing the skeleton. Similarly, if η is a Lagrangian distribution which is positive for the skeleton, then it can be made positive for the building without changing the skeleton.*

Proof. We argue by induction on blocks beginning with the bottom block B_1 . For a cotangent building consisting of a single block $B_1 = \mathcal{T}^* M_1$ the positivity condition is vacuous and to make η positive we just need to choose the polarization ν_1 , so that along M_1 we have $\nu_1 \in C(TM_1, \eta)$.

Suppose we have already constructed a positive building structure on $W_{<j} = \bigcup_{i < j} B_i$ such that $\eta|_{W_{<j}}$ is positive. Let $L = \text{Skel}(W)$ and denote $L_j := L \cap (B_j = T^* M_j)$. Suppose the block B_j intersects with blocks B_{j_1}, \dots, B_{j_m} , $1 \leq j_1 < \dots < j_m < j$, and let P_i be the face of M_j such that B_j is attached to B_{j_i} along a ribbon $V_i = \mathcal{T}^* P_i \subset B_{j_i} \setminus M_{j_i}$. The arboreal Lagrangian L_j intersects the ribbon V_i along an arboreal Legendrian Λ_i containing P_i . Denote by μ_i the polarization of P_i induced by the polarization ν_j of M_j and by $\bar{\nu}_{j_i}$ the polarization obtained by reducing the polarization ν_{j_i} of B_{j_i} with respect to the symplectic conormal of the Liouville field Z_{j_i} .

Arguing by induction beginning with $i = m$, we first use Lemma 4.31 to deform μ_m to a polarization μ'_m of P_m such that its Liouville field is tangent to Λ_j and the triple $(TP_m, \mu'_m, \bar{\nu}_{j_m})$ is \prec -cyclically ordered. We extend μ'_m to B_j as a polarization such that its Liouville field is tangent to L_j , which we will still denote by ν_j , forgetting the previous polarization of M_j . Note that on $\mathcal{O}p P_m \cap P_{m-1}$ the reduced triple $([TP_{m-1}]^{\zeta_m}, [\mu'_{m-1}]^{\zeta_{i_m}}, [\nu_{j_{m-1}}]^{\zeta_{i_m}})$ is \prec -ordered,

thanks to the positivity of the building $W_{<j}$. Applying Lemma 4.31 again, we can deform the polarization μ_{m-1} to a polarization μ'_{m-1} whose Liouville field is tangent to Λ_{j-1} , which coincides with $\bar{\nu}_j := [\nu_j]^{\zeta_{i_{m-1}}}$ on a neighborhood $U \supset P_m \cap P_{m-1}$, and such that the triple $(TP_{m-1}, \mu'_{m-1}, \bar{\nu}_{j_{m-1}})$ is \prec -cyclically ordered outside U . The extension claim follows from convexity: one first constructs μ'_{m-1} satisfying the positivity condition in the complement of a neighborhood $U' \Subset U$. Then, using a cut-off function θ equal to 1 on U' and 0 outside U , one redefines μ_{m-1} as $(1 - \theta)\mu'_{m-1} + \theta\bar{\nu}_j$. Again we extend to a polarization of B_j such that its Liouville field is tangent to L_j and we still call the new polarization ν_j . Continuing this process, we obtain a polarization ν_j on B_j making the building $W_{\leq j}$ positive. Finally, to ensure that η is positive along M_j we can further deform ν_j via a homotopy fixed on $W_{<j} \cap B_j$ by final application of Lemma 4.31. \square

5. RIDGIFICATION OF LAGRANGIANS

In this section we recall the ridgification theorem, as well as its formal analogue, and describe the canonical Wc-structure in the neighborhood of a ridgy Lagrangian.

5.1. Geomorphology of Lagrangian ridges.

5.1.1. *Ridgy Lagrangians.* As a stepping stone towards arboreal skeleta it will be useful to consider a class of singular Lagrangian and Legendrian submanifolds, called *ridgy*, which were introduced in [AGEN19]. Let us recall their definition. In the standard symplectic $(\mathbb{R}^2, dx \wedge dy)$ consider the subset $R = R_{1,2} = \{xy = 0, x \geq 0, y \geq 0\}$.

Definition 5.1. The *model ridge* $R_{k,n} \subset \mathbb{R}^{2n}$ of order k is the product $R_{k,n} = R^k \times \mathbb{R}^{n-k} \subset (T^*\mathbb{R})^k \times T^*\mathbb{R}^{n-k} = T^*\mathbb{R}^n$.

Example 5.2. The order n model ridge $R_{n,n} \subset T^*\mathbb{R}^n$ is the union to all the inner conormals of the faces of a quadrant in \mathbb{R}^n , hence is the union of the 2^n linear Lagrangians $\{p_j = q_k = 0, q_j, p_k \geq 0, j \in I, k \notin I\}$, where $I \subset \{1, \dots, n\}$.

Definition 5.3. An n -dimensional *ridgy Lagrangian* in a symplectic manifold (X, ω) is a closed subset $L \subset X$ which is covered by open neighborhoods U_i such that each $(U_i, U_i \cap L)$ is symplectomorphic to some $(B, B \cap R_{k,n})$, $0 \leq k \leq n$.

A ridgy Lagrangian L admits a natural stratification according to the order of ridges, so that the top dimensional stratum is the smooth part of the ridgy Lagrangian. Any ridgy Lagrangian can be viewed as the limit as $\varepsilon \rightarrow 0$ of a family of smooth Lagrangians $L(\varepsilon)$ homeomorphic to $L = L(0)$; they are obtained by using the model $R(\varepsilon) = \{xy = \varepsilon, x \geq 0, y \geq 0\}$ instead of R .

Definition 5.4. Given a contact manifold (Y, ξ) , a ridgy Legendrian $\Lambda \subset \xi$ is defined as a ridgy Lagrangian in a Weinstein hypersurface $\Sigma \subset Y$.

Note that $R_{k,n} \subset \mathbb{R}^{2n}$ is invariant with respect to the radial contracting vector field $Z_n = \sum_{j=1}^n q_j \partial / \partial q_j - p_j \partial / \partial p_j$. Hence its link $\partial R_{k,n} := R_{k,n} \cap S^{2n-1}$ in the standard contact sphere




FIGURE 5.1. A 2-dimensional ridgy Lagrangian has order 1 ridges along a union of separating curves, which intersect in order 2 ridges along a discrete set of points.

is an $(n - 1)$ -dimensional ridgy Legendrian. Similarly, $R_{k,n} \subset \mathbb{R}^{2k} \times T^*\mathbb{R}^{n-k}$ is invariant with respect to the contracting field $Z_k - \sum_{j=1}^{n-k} p_j \partial / \partial p_j$, where $\sum_{j=1}^{n-k} p_j \partial / \partial p_j$ is the canonical Liouville field for the factor $T^*\mathbb{R}^{n-k}$.




FIGURE 5.2. The 2-fold model ridge $R^2 = R \times R \subset T^*\mathbb{R} \times T^*\mathbb{R}$ is symplectomorphic to the union in $T^*\mathbb{R}^2$ of the first quadrant in \mathbb{R}^2 , the inner conormals to the positive x and y axes and the quarter-conormal to the origin lying in the first quadrant.

5.1.2. *The formal ridgification theorem.* First, we recall from [AGEN19] the notion of a tectonic field, which is the formal (i.e. non-integrable) analogue of a ridgy Lagrangian. Let M be a manifold with corners and $B = \mathcal{T}^*M$ the corresponding cotangent block. Recall that a Lagrangian plane field η along M which is transverse to the vertical distribution ν can be viewed as a field of quadratic forms on TM . To simplify the notation we will use the same symbol to denote the Lagrangian distribution and the corresponding field of quadratic forms.

By a *dividing hypersurface* $N \subset M$ we mean a properly embedded, co-oriented codimension 1 submanifold with corners such that $(N, \partial N) \subset (M, \partial M)$ is homologically trivial. A

collection of dividing hypersurfaces $\{N_j\}_j$ is said to be in *in general position* if each N_j is transverse to the intersection $N_{i_1} \cap \cdots \cap N_{i_k}$ of any subset of the other hypersurfaces.

Given a collection of dividing hypersurfaces $N_1, \dots, N_k \subset M$ in general position, a *tectonic field* λ with *faults* along $\{N_j\}_j$ is a collection of smooth graphical Lagrangian plane fields $\{\sigma_Q\}_Q$ defined over the closures \overline{Q} of the connected components Q of $M \setminus \bigcup_{j=1}^k N_j$, such that for every $j = 1, \dots, k$ there exists a field ℓ_j of non-vanishing 1-forms along N_j for which the following conditions are satisfied:

- (i) For each component P of $N_i \setminus \bigcup_{j \neq i} N_j$ adjacent to components Q_{\pm} of $M \setminus \bigcup_j N_j$ we have that the difference $\lambda_{Q_+} - \lambda_{Q_-}$ is the rank 1 quadratic form $\eta_i = \ell_i^2$, where the co-orientation of N_j points into Q_+ ;
- (ii) Along each intersection $N_{j_1} \cap \cdots \cap N_{j_m}$ the hyperplane fields $\tau_{j_s} = \ker(\eta_{j_s})$, $s = 1, \dots, m$, are transverse to all possible intersections of the τ_{j_r} , $r \neq s$.

A tectonic field η is called ε -small if it deviates from TM by an angle $< \varepsilon$, where a fixed but arbitrary Riemannian metric is understood. A tectonic field λ is called *collared* if for each k -face F of M the tectonic field λ splits as a product of a tectonic field on F and the trivial (tangent) field in the collar directions. In terms of quadratic forms, this means that the quadratic form is the sum of a quadratic form on TF and the zero form in the collar directions. In particular this gives the notion of a smooth Lagrangian plane field which is collared with respect to a corner structure. We can now state the formal version of the rigidification theorem.

Theorem 5.5 ([AGEN19]). *Let M be a manifold with corners and η any Lagrangian plane field in \mathcal{T}^*M . Then there exists a tectonic field λ transverse to η .*

Remark 5.6. The formal transversalization theorem holds in several variants, which will be needed for our applications below,.

- (i) C^0 -control: for any fixed $\varepsilon > 0$ we may demand that λ is ε -small.
- (ii) Relative form: if for a compact set $A \subset M$ we are already given an ε -small tectonic field λ_0 over $\mathcal{O}p A$ transverse to η , then λ can be chosen equal to λ_0 over $\mathcal{O}p A$. In this case the C^0 -control means that $\lambda - \lambda_0$ can be made ε -small.
- (iii) There is also an extension formulation as follows: if $K_1, K_2 \subset M$ are disjoint compact subsets and ζ is any tectonic field, then we may find a λ satisfying the conclusion of the theorem on $\mathcal{O}p(K_1)$ and such that $\lambda = \zeta$ on $\mathcal{O}p(K_2)$. In this case the C^0 -control means that $\lambda - \zeta$ can be made ε -small.
- (iv) The relative and extension forms can be combined if we assume that $\lambda_0 = \zeta$ on $\mathcal{O}p(A \cap K_2)$.
- (v) Collared version: if we assume that in the neighborhood $F \times \mathcal{I}^k$ of a k -face F the Lagrangian field μ splits as the product of $\mu_F \subset T^*F$ and the cotangent (vertical) distribution on $T^*\mathcal{I}^k$, then λ can be taken to be collared. The collared version holds with C^0 -control and with the relative and extension forms.

A tectonic field is called *aligned* if for each fault N_i we have $\ker(\eta_i) = TN_i$, where η_i is the rank 1 form giving the discontinuity of the tectonic field along N_j . Given a Lagrangian plane field η in \mathcal{T}^*M it is in general not possible to find an aligned tectonic field which is transverse to η , but it is possible to find an aligned tectonic field which is transverse to a Lagrangian field homotopic to η . More precisely, we have:

Theorem 5.7 ([AGEN19]). *Let M be a manifold with corners and η any Lagrangian field in \mathcal{T}^*M . Then there exists an aligned tectonic field λ transverse to a Lagrangian field $\hat{\eta}$ homotopic to η .*

Remark 5.8. The aligned formal transversalization theorem holds in several variants, which will be needed for our applications below.

- (i) C^0 -control: for any fixed $\varepsilon > 0$ we may demand that λ is ε -small.
- (ii) Relative form: if for a compact set $A \subset M$ we are already given an ε -small aligned tectonic field λ_0 over $Op A$ transverse to η , then λ can be chosen equal to λ_0 over $Op A$. Moreover we may assume that the homotopy between $\hat{\eta}$ and η is constant on $Op(A)$. In this case the C^0 -control means that $\lambda - \lambda_0$ can be made ε -small.
- (iii) There is also an extension formulation as follows: if $K_1, K_2 \subset M$ are disjoint compact subsets and ζ is any aligned tectonic field, then we may find a λ satisfying the conclusion of the theorem on $Op(K_1)$ and such that $\lambda = \zeta$ on $Op(K_2)$. In this case the C^0 -control means that $\lambda - \zeta$ can be made ε -small.
- (iv) The relative and extension forms can be combined if we assume that $\lambda_0 = \zeta$ on $Op(A \cap K_2)$.
- (v) Collared version: if we assume that in the neighborhood $F \times \mathcal{I}^k$ of a k -face F the Lagrangian field μ splits as the product of $\mu_F \subset T^*F$ and the cotangent (vertical) distribution on $T^*\mathcal{I}^k$, then λ can be taken to be collared and we may moreover assume the homotopy between $\hat{\eta}$ and η to be through collared distributions. The collared version holds with C^0 -control and with the relative and extension forms.

5.1.3. *The ridgification theorem.* Given a tectonic field λ in \mathcal{T}^*M , a ridgy Lagrangian $L \subset \mathcal{T}^*M$ is called δ -close to λ if for any point $a \in L$ and any tangent plane T to L at a there exists a non-fault point $b \in M$, δ -close to a such that the angle between $\lambda(b)$ and the plane T parallel transported to b along a geodesic is $< \delta$. As before, a fixed but otherwise arbitrary Riemannian metric on M is understood.

A ridgy Lagrangian in $L \subset \mathcal{T}^*M$ is called *collared* if for each collar $Q \times \mathcal{I}^k \subset M$, where Q is a k -face, we have $L \cap T^*(Q \times \mathcal{I}^k) = L_Q \times \mathcal{I}^k$ for $L_Q \subset T^*Q$ a ridgy Lagrangian and $\mathcal{I}^k \subset T^*\mathcal{I}^k$ the zero section.

Remark 5.9. If $L \subset \mathcal{T}^*M$ is a collared ridgy Lagrangian, then it is adapted for the We-structure of \mathcal{T}^*M inherited from the collar structure of M .

The ridgy Lagrangians we will consider are of a special type: they are all obtained from the zero section $M \subset \mathcal{T}^*M$ by means of a ridgy isotopy, which is defined as follows.

- (i) Let $N_1, \dots, N_m \subset M$ be co-oriented separating hypersurfaces defined by equations $\phi_j = 0$ for some C^∞ -functions $\phi_j : M \rightarrow \mathbb{R}$ without critical points on N_j . We assume that the N_j are co-oriented by the outward transversals to the domains $\{\phi_j \leq 0\}$. Denote $\phi_j^+ = \max(\phi_j, 0)$ and choose a cut-off function θ_j which is equal to 1 on N_j and to 0 outside a neighborhood of N_j . Define a function $\Phi : M \rightarrow \mathbb{R}$ (which is C^1 and piecewise C^∞) by the formula

$$\Phi := \sum_{j=1}^m \theta_j (\phi_j^+)^2.$$

An *earthquake isotopy* with faults N_j is defined as a family of Lagrangians $L_t \subset \mathcal{T}^*M$ given by the homotopy of generating functions $t\Phi$, i.e. $L_t = \{p = td\Phi\}$, $t \geq 0$.

- (ii) A *ridgy isotopy* is an earthquake isotopy followed by an ambient Hamiltonian isotopy.




FIGURE 5.3. An earthquake isotopy.

Note that $L_0 = M$ and the earthquake isotopy can be realized by an ambient Hamiltonian homotopy beginning from any $t > 0$. We can now state the integrable version of the ridgification theorem.

Theorem 5.10 ([AGEN19]). *For any Lagrangian field η in \mathcal{T}^*M there exists a ridgy isotopy of M to a ridgy Lagrangian L which is transverse to η .*

Remark 5.11. The ridgification theorem also holds in several variants:

- (i) C^0 -control: we may assume that the ridgy isotopy is C^0 -close to the identity.
- (ii) Relative form: if η is transverse to M on $\mathcal{O}p A$ for $A \subset M$ a closed subset, then we may assume that the ridgy isotopy is constant on $\mathcal{O}p A$.
- (iii) Collared version: if we assume that in the neighborhood $F \times \mathcal{I}^k$ of a k -face F the Lagrangian field μ splits as the product of $\mu_F \subset T^*F$ and the cotangent (vertical) distribution on $T^*\mathcal{I}^k$, then we may demand that the ridgy isotopy is collared.

The proof of Theorem 5.10 is given in [AGEN19]. The argument is roughly the following: first one constructs an aligned tectonic field λ transverse to η using Theorem 5.7. Then one integrates λ to a ridgy Lagrangian which is transverse to η along its ridge locus using the refined holonomic approximation [AG18a]. In the complement of the ridge locus there is no homotopical obstruction to removing the tangencies so one can achieve transversality

using the wrinkling theorem for Lagrangian submanifolds [AG18b]. Finally, one exchanges the wrinkles for order 1 ridges to complete the proof.

For our applications in the present article it will be useful to first inductively apply the holonomic approximation step in extension form and only wrinkle at the end, so we quote here Lemma 4.8 from [AGEN19] which we will use below.

Lemma 5.12 ([AGEN19]). *Let $\Lambda \subset T^*L$ be a ridgy Lagrangian, $R \subset \Lambda$ its ridge locus and γ a Lagrangian field. Suppose that γ is homotopic to a Lagrangian field $\hat{\gamma}$ which is transverse to Λ . Then there exists a Hamiltonian isotopy Λ_t of Λ and a Lagrangian field $\tilde{\gamma}$ such that:*

- (i) $\Lambda_1 \pitchfork \gamma$ on $Op(R)$.
- (ii) $\Lambda_1 \pitchfork \tilde{\gamma}$ everywhere.
- (iii) $\tilde{\gamma}$ is homotopic to γ by a homotopy fixed on $Op(R)$.

Remark 5.13. We have the following variants:

- (i) Relative version: if $A \subset \Lambda$ is a closed subset and the homotopy between γ and $\hat{\gamma}$ is fixed on $Op(A)$, then the Hamiltonian isotopy and the homotopy between $\tilde{\gamma}$ and γ can both be chosen to be fixed on $Op(A)$.
- (ii) Extension form: if $K_1, K_2 \subset L$ are closed subsets we may find Λ_t and $\tilde{\gamma}$ such that the conclusions of the lemma are satisfied on $Op(K_1)$, the ridgy isotopy is constant on $Op(K_2)$ and the homotopy between $\tilde{\gamma}$ and γ is constant on $Op(K_2)$. Furthermore, we can combine this extension version with the relative version.
- (iii) Collared version: if we assume that in the neighborhood $F \times \mathcal{I}^k$ of a k -face F the Lagrangian field γ splits as the product of $\gamma_F \subset T^*F$ and the cotangent (vertical) distribution on $T^*\mathcal{I}^k$ and we moreover assume the homotopy between γ and $\hat{\gamma}$ is through such distributions, then we may assume that Λ_t is collared and the homotopy between $\tilde{\gamma}$ and γ is collared.

We also record for future reference the intermediate step in which one integrates an aligned tectonic field, before applying holonomic approximation.

Lemma 5.14 ([AGEN19]). *Let γ be a Lagrangian field on T^*L and let λ be an aligned tectonic field on L which is transverse to a Lagrangian field homotopic to γ . Then there exists a ridgy isotopy L_t such that L_1 is transverse to a Lagrangian field homotopic to γ on a neighborhood of its ridge locus.*

As before, this holds in C^0 -small, relative, extension and collared versions.

5.2. Wc-structures for ridgy Lagrangians.

5.2.1. *Wc-building structure associated with a ridgy Lagrangian.* Next we discuss a Darboux/Weinstein type theorem for the symplectic structure in the neighborhood of a ridgy Lagrangian. Denote by L_k the locus of k -dimensional locus of $(n - k)$ -fold ridges in L . This stratifies L as a union of smooth, relatively open submanifolds $L = L_0 \cup \dots \cup L_n$.

Given a manifold with corners M we denote by $M^{\Delta,j}$ the manifold with corners obtained from M by truncating all corners of dimension $\leq j$. Thus, each i -face P from $\partial_i M$ for $i < j$ yields a 1-face $P^{\blacktriangle,i} := P^{\Delta,i} \times \Delta^{j-i} \subset \partial_1 M^{\Delta,j}$, where we denote by Δ_{j-i} an open $(j-i)$ -dimensional simplex. Denote by \mathcal{R}^{2k} the germ of the standard symplectic \mathbb{R}^{2k} at the origin.

Lemma 5.15. *Let L be a ridgy Lagrangian in a symplectic manifold (X, ω) such that for each $k \geq 1$ the symplectic normal bundle to the order k ridge locus is trivial. Then a neighborhood of L admits a structure of a Wc-building $W = (U_0 \rightarrow \cdots \rightarrow U_n)$, where*

$$U_j := \bigcup_Q \mathcal{T}^* Q^{\Delta,j-1} \times \mathcal{R}^{2j}$$

and the union is taken over all components Q of L_j , such that the following properties hold:

- (i) *The inclusion of W into X is a symplectic embedding.*
- (ii) *The skeleton of W is L .*
- (iii) *The Weinstein structure underlying W is Weinstein homotopic to the cotangent bundle structure $\mathcal{T}^*L(\varepsilon)$ for a smoothing $L(\varepsilon)$ of L .*




FIGURE 5.4. The Wc-building structure in the neighborhood of a ridgy Lagrangian

Remark 5.16. The hypothesis on the symplectic normal bundles is included only for the sake of simplicity. It is obviously satisfied by the ridgy Lagrangians produced by the ridgification theorem and these are the only ridgy Lagrangians we care about for our applications. In general one should replace $\mathcal{T}^*Q^{\Delta,j-1} \times \mathcal{R}^{2j}$ by the appropriate symplectic bundle over $\mathcal{T}^*Q^{\Delta,j-1}$ and the result is proved in the same way. Note that in either case the Wc-building structure is *not* a cotangent building.

Proof. Recall that L_k denotes the locus of k -dimensional locus of $(n-k)$ -fold ridges in L . Denote by U_0 the union of so small closed disjoint balls B_j centered at the points of L_0 that $L \cap B_j$ is invariant with respect to the radial contracting vector field inherited from the local model. Denote $L_j^{>0} := L_j \setminus U_n$. Choose a small (in a similar sense) closed tubular neighborhood U_1 of $\bar{L}_1^{>0}$. Set $L_j^{>1} = L_j^{>0} \setminus U_1$ for $j > 1$.

Continuing this process we define $U_j \supset \bar{L}_j^{>j-1}$ for $j = 0, 1, \dots, n$. Note that $\bar{L}_j^{>j-1}$ is a manifold with corners of dimension j . Note also that U_j can be presented as a symplectic

fibration $\pi_j : U_j \rightarrow \mathcal{S}^* \bar{L}_j^{>j-1}$. The restriction $\pi_j|_{\bar{L}_j^{j-1}}$ contains a subfibration $L \cap U_j \rightarrow \bar{L}_j^{j-1}$ with the fiber $R_{n-j, n-j} = R^{n-j}j$.

The structural group of the fibration π_j reduces to the discrete group of symmetries of the model ridge R^{n-j} . These symmetries commute with the action of the radial contracting field Z_j , and hence U_j admits a global contracting field Z_j^{j-1} which restricts to the radial field Z_j on fibers over \bar{L}_j^{j-1} and which vanishes along \bar{L}_j^{j-1} . This makes $(U_j, Z_j^{>j-1})$ a Wc-manifold.

The required Wc-structure on $\mathcal{O}p L$ can now be constructed by successive vertical gluing. Starting with (U_0, Z_0) we vertically attach $(U_1, Z_1^{>0})$ using a Wc-hypersurface U_0 , namely the ribbon of $\partial_\infty U_0 \cap L$, and the similar hypersurface in ∂U_1 which plays the role of the nucleus of one of the boundaries of ∂U_1 . Continuing this process we obtained the required Wc-structure.

Properties (i) and (ii) are immediate from the construction and for (iii) the homotopy between the constructed Wc-structure and $\mathcal{S}^*L(\varepsilon)$ can be obtained by repeating the whole construction for the smoothed $L(\varepsilon)$ with the stratification inherited from the order of ridges in L . Finally the description of the U_j given in the statement of the lemma follows from the usual Darboux/Weinstein theorem for isotropic submanifolds with trivial symplectic normal bundles. \square

5.2.2. Ridgy Lagrangians in Wc-buildings. Let $W = \bigcup_j B_j$ be a cotangent building, with ν_j the vertical distribution for B_j . Given a Liouville cone L over an adapted Legendrian in $W \setminus \text{Skel}(W)$ we say that L is *reduced transverse* to all ν_i defined near its corners if for each tangent plane T to $L \cap B_{i_1} \cap \cdots \cap B_{i_m}$ at a point $a \in B_{i_1} \cap \cdots \cap B_{i_m}$ the reduction $[T]^{\zeta_I} \subset \zeta_I$ is transverse to the reductions $[\nu_{i_s}]^{\zeta_I}$, where $I = (i_1 < \cdots < i_m)$.

Proposition 5.17. *Let \widehat{W} be a symplectic manifold and $W \subset \widehat{W}$ a symplectic embedding of a positive cotangent building. Let $L \subset \widehat{W} \setminus \text{Skel}(W)$ be a Lagrangian with corners such that $L \cap W$ is a Liouville cone over an adapted Legendrian in $W \setminus \text{Skel}(W)$. Let ν_{-1} be a positive distribution which is extended to \widehat{W} . Then there exists a C^0 -small ridgy isotopy which keeps $L \cap W$ adapted and deforms L to a ridgy Lagrangian \widehat{L} which is transverse to ν_{-1} and reduced transverse to all the corresponding distributions ν_i defined near its corners.*

Proof. Consider L as the 0-section of a block \mathcal{S}^*L . Let P_1 be a face of L which is a Legendrian in one of the blocks B_k . The distribution ν_k is adapted for this block structure and is invariant (and tangent) to the contracting field Z_{P_1} adapted to this block. We apply the collared version of the formal ridgification theorem 5.5 to deform the horizontal distribution on $\mathcal{O}p(P_1) = P_1 \times \mathcal{I}$ to an ε -small collared tectonic field σ'_1 in $\mathcal{S}^*P \times \mathcal{S}^*\mathcal{I}$, achieving transversality to the product of $[\nu_k]^{\zeta_1}$ and the vertical distribution on $T^*\mathcal{I}$. Note in particular that σ'_1 is reduced transverse to ν_k , i.e. $[\sigma'_1]^{\zeta_1}$ is transverse to $[\nu_k]^{\zeta_1}$. Furthermore, by the extension form of the formal transversalization theorem we may assume that $\sigma'_1 = 0$ outside of a neighborhood of P_1 in L .

Note that according to Lemma 4.17, for any other index j we have that $[\sigma_1]^{\zeta_j}$ is transverse to $[\nu_j]^{\zeta_j}$, or in other words, that σ_1 is reduced transverse to ν_j . Using the extension form of

the formal rigidification theorem 5.5 we can further inductively deform the tectonic field σ_1 into $\sigma_2, \dots, \sigma_k$ which is ε -small, collared and reduced transverse to ν_{k-1}, \dots, ν_1 and finally construct an ε -small collared tectonic field σ which is in addition is transverse to ν_{-1} .

To conclude the proof we want to apply the rigidification theorem to integrate the tectonic field σ . It will be easier to work one boundary face of L at a time. We first note that by the same argument as before we can inductively use Theorem 5.7 instead of Theorem 5.5 to replace σ with an ε -small, collared, aligned tectonic field $\hat{\sigma}$, which is no longer reduced transverse to the ν_i or transverse to ν_{-1} but has the property that there exists a family of symplectic bundle isomorphisms $\Phi_t : T(T^*L) \rightarrow T(T^*L)$ such that $\hat{\sigma}$ is reduced transverse to each $\Phi_1(\nu_j)$ and transverse to $\Phi_1(\nu_{-1})$. Then we integrate $\hat{\sigma}$ to a ridgy Lagrangian \hat{L} using Lemma 5.14 which still has the property that it is reduced transverse to each $\Phi_1(\nu_j)$ and transverse to $\Phi_1(\nu_{-1})$.

Now, starting with P_1 , we may use Lemma 5.12 to first deform \hat{L} along its ridge locus in a neighborhood of P_1 so that it becomes reduced transverse to ν_k in a neighborhood of the ridge locus. The deformation is achieved by a Hamiltonian isotopy which is C^0 -close to the identity. Moreover, by the collared version of that lemma we may assume that the resulting ridgy Lagrangian, which now and it what follows we abuse notation and still denote by \hat{L} , is still collared.

As before, Lemma 4.17 assures us that \hat{L} is reduced transverse to ν_j near the boundary of the next face P_2 in a neighborhood of its ridge locus, and our previous application of Lemma 5.12 guaranteed that the homotopical condition necessary to keep applying holonomic approximation is still satisfied, even in its collared form. Therefore we can use Lemma 5.12 once more to deform \hat{L} along its ridge locus in a neighborhood of P_2 so that it becomes reduced transverse to ν_j in a neighborhood of its ridge locus and we can continue the process until we have achieved reduced transversality to ν_{k-1}, \dots, ν_1 along the ridge locus in a neighborhood of the boundary for a collared \hat{L} . The deformation is still C^0 -small. Moreover, again by Lemma 4.17 we have that \hat{L} is transverse to ν_{-1} near its boundary, and ν_{-1} is homotopic to a Lagrangian field transverse to \hat{L} by a homotopy fixed near the boundary.

At this point we have a ridgy Lagrangian $\hat{L} \subset T^*L$ which is reduced transverse to the ν_i and which is collared with respect to the cotangent bundle structure T^*L . However, the intersection of \hat{L} with W is no longer a Liouville cone over an adapted Legendrian, so we need to fix this. Since all the above steps can be performed by C^0 -small perturbations, recall that we may assume that \hat{L} is in an ε -neighborhood of the zero section in T^*L . Therefore, by a straightforward cutoff at the level of generating functions one may interpolate between our old \hat{L} and a new \hat{L} whose intersection with W is a cone over an adapted Legendrian, while maintaining the reduced transversality to the ν_{k-1}, \dots, ν_1 . Moreover, the new \hat{L} still has the property that it is transverse to a Lagrangian field homotopic to ν_{-1} , with the homotopy fixed in a neighborhood of its boundary.

By the same argument as before, we may use Lemma 5.12 to further deform \widehat{L} to obtain a ridgy Lagrangian which is collared, reduced transverse to ν_{k-1}, \dots, ν_1 and transverse to ν_{-1} on a neighborhood of the ridge locus, and such that ν_{-1} is homotopic to a Lagrangian field transverse to \widehat{L} via a homotopy fixed in a neighborhood of the boundary of \widehat{L} and fixed in a neighborhood of its ridge locus. Hence we may apply the wrinkling theorem [AG18b] in its relative form to further deform \widehat{L} relative to its boundary and ridge locus to achieve transversality to ν_{-1} everywhere by the introduction of wrinkles, which can easily be exchanged for order 1 ridges, see for example Corollary 4.13 of [AGEN20a], or Figure 5.5 below. \square




FIGURE 5.5. The wrinkling theorem allows us to achieve transversality by the introduction of cuspidal singularities, but these can be easily replaced by order 1 ridges by simply replacing the unique local models.

6. ARBOREALIZATION OF SKELETA

In this final section we complete the proof of our main result: a Weinstein manifold admitting a polarization can be deformed to have a positive arboreal skeleton.

6.1. Immersions into arboreals. In this section we will present an inductive scheme to arborealize a ridgy Lagrangian. The most direct way of doing so produces an arboreal Lagrangian with boundary, i.e. whose boundary is itself arboreal. However, we can do slightly better by inductively capping off the new boundary components that arise so that the end result is an arboreal Lagrangian with smooth boundary. We therefore begin with a preliminary discussion which describes the capping process.

6.1.1. Capping arboreal Legendrians.

Lemma 6.1. *Let M be a smooth manifold, possibly with boundary and corners, and $\Lambda \subset (\mathcal{T}^*M \setminus M) \cap \mathring{\mathcal{T}}^*M$ an arboreal Legendrian with smooth boundary $\partial\Lambda$. Suppose that Λ is positive and that it is also positive with respect to the vertical polarization ν of \mathcal{T}^*M . Let $\widehat{\Lambda} \supset \Lambda$ be an arboreal space without boundary such that $\widehat{\Lambda} \setminus \Lambda$ is a manifold. Suppose that there exists an immersion $h : \widehat{\Lambda} \setminus \Lambda \rightarrow \mathring{M}$, transverse to $\pi(\Lambda)$, extending the co-oriented front projection $\pi : \mathcal{O}_p \partial\Lambda \rightarrow \mathring{M}$. Let $H \subset \mathcal{T}^*M \setminus M$ be the Legendrian positive conormal lift of the immersion h . Then the closed arboreal Legendrian $\widetilde{\Lambda} := \Lambda \cup H \subset \mathcal{T}^*M \setminus M$ is positive arboreal and is positive with respect to ν .*

Proof. Indeed, we just added to Λ a smooth Legendrian transverse to ν . \square

Remark 6.2. To be more precise, we should say that there exists a Legendrian lift $H \subset \mathcal{T}^*M \setminus M$ of h such that the conclusion of the lemma holds, or equivalently we could work in the idealized boundary S^*M from the onset, where the lift H is unique.

6.1.2. *Genuine transversality.* Let η be a Lagrangian distribution in a symplectic manifold W . We introduce a notion of transversality for piecewise smooth Lagrangians which can be intuitively thought of as “transversality even after smoothing”.

Definition 6.3. A piecewise smooth Lagrangian $L \subset W$ is called *genuinely transverse* to η if for each point $a \in L$ there exist Darboux coordinates (p, q) near a , with $\eta(a)$ is tangent to the cotangent fiber $q = 0$ at the origin 0 , such that the Lagrangian $\mathcal{O}p a \subset L$ is generated by a C^1 -function $H(q)$.

Remark 6.4. The ridgy Lagrangians produced by the ridgification theorem are transverse to the distribution used as input in the theorem, but in general are not genuinely transverse.

Similarly, a piece-wise smooth Lagrangian map $f : L \rightarrow W$, i.e. not necessarily an embedding, is said to be genuinely transverse to a Lagrangian distribution η if for every point $a \in L$ there exists a neighborhood $\mathcal{O}p a \subset L$ such that $f|_{\mathcal{O}p L}$ is a piecewise smooth Lagrangian embedding and $f(\mathcal{O}p L)$ is genuinely transverse to η . Note that any piece-wise smooth Lagrangian which is genuinely transverse to η can be approximated by a family of Lagrangians L_t , $t > 0$, which are transverse to η and such that $L_t \xrightarrow{C^0} L$ as $t \rightarrow 0$, where convergence is smooth where L is smooth. Indeed, this follows from the corresponding approximation property for C^1 -smooth generating functions.

Definition 6.5. Let L be an arboreal space and M a smooth manifold of the same dimension n . A map $f : M \rightarrow L$ is called an *immersion* if there exists a stratification $M = M_0 \cup M_1 \cup \dots \cup M_n$ by manifolds M_j of dimension $(n - j)$ such that f is an immersion on each stratum.

Recall that as an arboreal space, L is equipped with an orientation structure κ , which is a line bundle equipped with identifications with the orientation line bundles of the smooth pieces and compatibility at the singularities determined by symplectic geometry. The pullback $f^*\kappa$ is a line bundle on M , with fixed identifications with the orientation bundle $\wedge^n TM_0$ for P on M_0 .

Definition 6.6. An immersion $f : M \rightarrow L$ is *oriented* if $f^*\kappa$ is isomorphic to the orientation bundle of M relative to the fixed identification on M_0 .

The condition that $f : M \rightarrow L$ is oriented can be understood as follows. Let A be a component of M_1 and P, Q be components of M_0 adjacent to A . At a point $f(x) \in L$ in the closure of both $f(P)$ and $f(Q)$, the singularity of L is modeled on signed rooted tree \mathcal{T} . If $f(P)$ is contained in a smooth piece of L closer to the root ρ of \mathcal{T} than $f(Q)$ then the outward

boundary coorientation of $A \subset \partial P$ coincides with the coorientation of $f(A)$ in P . See also Remark 3.6.

Let $L \subset W$ be a positive arboreal Lagrangian and η a positive distribution for L . Then the symplectic vector bundle $TW|_L \rightarrow L$ is isomorphic to $\eta \oplus \eta^*$. We can realize η^* as a Lagrangian plane field on $TW|_L$ which is transverse to η and the space of such is contractible.

Lemma 6.7. *Let $L \subset W$ be a positive arboreal Lagrangian and η a positive distribution for L . For any oriented immersion $f : N \rightarrow L$ and any point $x \in N$, the germ of $f(L)$ at $f(x)$ is graphical with respect to the polarization (η^*, η) .*

Proof. We first check the condition along M_1 . Let $x \in M_1$, so $f(x) \in L$ is an A_2 type singularity. There are two possibilities for the germ of $f : M \rightarrow L$ near x , excluding the trivial case where $f(\mathcal{O}_p(x))$ stays in the smooth part of L , i.e. the zero section piece corresponding to the root ρ of the signed rooted tree \mathcal{T} which classified the singularity of L at $f(x)$. The first possibility is that $f(\mathcal{O}_p(x))$ is the conormal of a cooriented hyperplane together with the half-plane lying in the direction of the coorientation. The second possibility is to take the other half-plane. By inspection of the local model, the former is graphical with respect to (η^*, η) and the latter is not. On the other hand, the former gives an oriented immersion while the latter does not. The condition on M_k , $k > 1$ can be verified inductively using the above argument, or more directly one can inspect the explicit local model, in which the polarization η can also be taken to be canonical as proved in [AGEN20a]. \square

Proposition 6.8. *Let $L \subset W$ be a n -dimensional positive arboreal Lagrangian and η a positive distribution for L . Then for any oriented immersion $f : N \rightarrow L$ of an n -dimensional manifold N the piecewise smooth immersion $N \rightarrow L \hookrightarrow W$ is genuinely transverse to η .*

Proof. Near each point $x \in N$ the germ of $f(L)$ is graphical with respect to (η^*, η) so near x we can generate $f(L)$ by a piecewise smooth function on η^* . Moreover, the generating function can be smoothed and the result is still transverse to η since it remains graphical with respect to (η^*, η) . To achieve a global smoothing of L one can patch up the local smoothings with a partition of unity. \square

6.1.3. *Capping arboreal Lagrangians.* Let L be an arboreal space and Λ a smooth component of its boundary ∂L . We call the component Λ *bounding* if there exists a manifold M with $\partial M = \Lambda$ and an oriented immersion $M \rightarrow L$ extending the inclusion $\Lambda \hookrightarrow L$.

Lemma 6.9. *Let $\Lambda \subset \mathcal{T}^*M \setminus M$ be a positive arboreal Legendrian which is positive with respect to the vertical distribution ν . Suppose that the boundary of Λ is smooth and bounding. Then there exists a closed arboreal Legendrian $\widehat{\Lambda} \subset \mathcal{T}^*M \setminus M$ which is transverse to ν and satisfies the following properties, where we denote by $\widehat{L} \subset \mathcal{T}^*M$ the arboreal Lagrangian formed by the union of the Liouville cone of $\widehat{\Lambda}$ and the 0-section.*

- (i) $\Lambda' := \widehat{\Lambda} \setminus \Lambda$ is smooth;




FIGURE 6.1. On the left: the upper immersion is oriented, while the lower immersion is not. On the right: the upper smoothed Lagrangian is transverse to η , while the lower one is not.

- (ii) $\widehat{\Lambda}$ bounds in \widehat{L} an immersed submanifold which is genuinely transverse to any Lagrangian distribution η in \mathcal{T}^*M such that TM, ν, η are \prec -ordered.

Proof. Let $f : N \rightarrow \Lambda$ be an immersion bounding $\partial\Lambda$ in Λ . Denote by $\pi : \mathcal{T}^*M \rightarrow M$ the front projection. Then the image $\Sigma := \pi \circ f(N) \subset M$ is a C^1 -smooth immersed cooriented hypersurface. Hence it has a C^∞ -smooth, fixed on $\partial\Sigma$ push-off Σ' in the direction of the co-orientation, which together with Σ bounds an immersed domain $\Omega \subset M$ with a C^1 -smooth boundary $\partial\Omega = \Sigma \cup \Sigma'$. Let Λ' be the conormal lift of Σ' . Then $\widehat{\Lambda} := \Lambda \cup \Lambda'$ is the required Legendrian, where property (ii) follows from Proposition 6.8, \square




FIGURE 6.2. Closing up a positive Legendrian whose boundary is smooth and bounding.

6.2. Cones over arboreals. We now discuss the key lemma which will be used for the inductive arborealization of ridgy Lagrangians.

6.2.1. Arborealization of radial cones on arboreals.

Definition 6.10. An arboreal Lagrangian in a Wc-manifold W is called *asymptotically conical* if it coincides outside of a neighborhood of $\text{Skel}(W)$ with a Liouville cone over a Legendrian.

Note that the boundary of an asymptotically conical arboreal Lagrangian L is also asymptotically conical. We call a smooth component $C \subset \partial L$ *bounding* if there exists a possibly non-compact manifold N with $\partial N = C$ and an asymptotically conical immersion $f : N \rightarrow L$ bounding C .

Lemma 6.11. *Let Λ be a positive arboreal Legendrian in S^{2n-1} endowed with the standard contact structure. Let $L \subset \mathbb{R}^{2n}$ be the Lagrangian cone over Λ centered at 0, i.e. with respect to the radial Liouville structure on the unit ball $B \subset \mathbb{R}^{2n}$. Let (τ, ν) be a polarization of \mathbb{R}^{2n} such that every Lagrangian plane T tangent to L at 0 satisfies the condition $T^{\tau, \nu} > 0$. Let $L_0 = \tau(0)$ be the Lagrangian plane through the origin, and $\Lambda_0 \subset S^{2n-1}$ its Legendrian link. Suppose that each boundary component of Λ is bounding. Then there exist:*

- (i) a closed arboreal Legendrian $\widehat{\Lambda} \supset \Lambda$ such that $\Lambda' := \widehat{\Lambda} \setminus \Lambda$ is smooth and such that every Lagrangian plane T tangent to the Liouville cone L' over Λ' satisfies the condition $T^{\tau, \nu} > 0$;
- (ii) an asymptotically conical arboreal Lagrangian \widehat{L} in \mathbb{R}^{2n} endowed with the standard Liouville structure such that:
 - \widehat{L} coincides at infinity with the Lagrangian cone over $\Lambda \cup \Lambda_0$;
 - $\widehat{L} \supset \widehat{\Lambda}$ and $\widehat{L} \supset L_0$;
 - \widehat{L} has smooth boundary and every boundary component is bounding;
 - \widehat{L} is transverse to any Lagrangian distribution $\eta \in C(\nu, \tau)$, i.e. negative with respect to (τ, ν) .
- (iii) a compactly supported homotopy $\lambda_t = \lambda_0 + df_t$ of $\lambda_0 = pdq$ such that the Wc-manifold obtained from $(\mathbb{R}^{2n}, \lambda_1)$ by converting the ribbons of the Legendrians $\Lambda \cup \Lambda_0$ into nuclei of boundary faces has the structure of a positive cotangent building with skeleton \widehat{L} .





FIGURE 6.3. The goal of Lemma 6.11 when $n = 1$.


 FIGURE 6.4. The goal of Lemma 6.11 when $n = 2$.

Proof. (i) Let p, q be canonical coordinates for the polarization (τ, ν) . Note that the function $H = pq$ restricts to $L \setminus 0$ as a positive function which satisfies the condition $H|_L \geq c|p|^2$ for some $c > 0$. Take the function $G = \max(H, \frac{1}{2}c|p|^2)$. Then $\{H = \varepsilon\} \cap L = \{G = \varepsilon\} \cap L$ for any $\varepsilon > 0$. The function G has star-shaped level sets with corners away from L . Hence, it can be smoothed to a function with the same property keeping the restriction to L equal to $pq|_L$. We abuse notation and denote the smoothed function by the same symbol G . Note that in particular we have

$$(pdq = \frac{1}{2}(pdq - qdp) + d(pq))|_{L \cap \{G = \varepsilon\}} = 0,$$

i.e. $L \cap \{G = \varepsilon\}$ is Legendrian in the level set $G = \varepsilon$ for both contact structures, defined by the Liouville forms pdq and $\frac{1}{2}(pdq - qdp)$. Consider the Liouville form

$$\lambda = \frac{1}{2}(pdq - qdp) + \frac{1}{2}d(\alpha(G)) = \frac{1}{2}(pdq - qdp) + \frac{1}{2}\alpha'(G)d(pq),$$

where $\alpha : \mathbb{R}_+ \rightarrow [0, 1]$ a monotone cut-off function which is equal to the identity on $[0, \varepsilon]$ and to 0 on $[\tilde{\varepsilon} := \varepsilon + \delta, \infty)$ for $\delta \ll \varepsilon$. Then $\lambda = pdq$ in $\{G < \varepsilon\}$ and $\lambda = \frac{1}{2}d(pdq - qdp)$ in $\{G \geq \tilde{\varepsilon}\}$. Let $\tilde{L} = \tilde{L}_{\varepsilon, \delta}$ be the saturation of $\Lambda_{\tilde{\varepsilon}} = L \cap \{G = \tilde{\varepsilon}\}$ by the forward and backward Liouville trajectories of the Liouville field of the form λ .

As $\delta \rightarrow 0$, note that $\tilde{L}_{\varepsilon, \delta}$ converges to the union of the forward pdq -cone and the backwards $\frac{1}{2}(pdq - qdp)$ cone over Λ_{ε} . Hence for small δ a plane \tilde{T} tangent to \tilde{L} near $\tilde{\Lambda}_{\varepsilon}$ is close to a

plane spanned by a tangent plane η to $\tilde{\Lambda}_\varepsilon$ and a convex linear combination $(1-t)R + tV$, $0 \leq t \leq 1$ of the contracting vector fields $R = -\frac{1}{2}(p\frac{\partial}{\partial p} + q\frac{\partial}{\partial q})$ and $V = -p\frac{\partial}{\partial p}$. When $t = 0$ the plane \tilde{T} coincides with a plane T tangent to L , and hence corresponds to a positive definite quadratic form Q . Increasing t corresponds to adding to Q the quadratic form $\frac{t}{1-t}\ell^2$, where the hyperplane $\{\ell = 0\}$ is the front projection of η . On the other hand, the distribution ν is given by our assumption by a negative definite quadratic form P . The intersection of $\tilde{T} \cap \nu$ corresponds to critical points of the quadratic form $Q + \frac{t}{1-t}\ell^2 - P$ which is positive definite. This assures the transversality of ν to \tilde{L} .

Recall that each boundary component $C = \partial\Lambda$ is bounding. Let $f : N \rightarrow \Lambda$, where $\partial N = C$, be the bounding immersion. Let Z^t be the Liouville flow of the Liouville vector field dual to λ . Consider the conical immersion $F : N \times \mathbb{R} \rightarrow \mathbb{R}^{2n}$ given by

$$F(x, t) = Z^t(x), \quad x \in N, t \in \mathbb{R}.$$

Then the limit

$$F_\infty(\cdot) = \lim_{t \rightarrow -\infty} F(\cdot, t)$$

defines a cooriented C^1 -smooth immersion $F_\infty : N \rightarrow L_0$. Let $F'_\infty : N \rightarrow L_0$ be a C^∞ -smooth normal push-off of N in the direction of its co-orientation. By modifying F'_∞ near $\partial N = C$ we can arrange that the immersion F'_∞ together with F' define a C^1 -smooth immersion $\hat{F} := F_\infty \cup F'_\infty : \hat{N} := N \cup_C N \rightarrow L_0$ of the closed manifold \hat{N} obtained by gluing two copies of N along the boundary C . The manifold \hat{N} bounds a manifold M , namely the product $N \times [0, 1]$ with a smoothed boundary. Note that we can arrange that the immersion \hat{F} extends to M . There exists a Legendrian embedding $\phi : N \rightarrow S^{2n-1}$ such that $\phi(\partial N) = C$ and such that

$$\lim_{t \rightarrow \infty} Z^{-t} \circ \phi = F'_\infty.$$

Set $\check{N} := \phi(N)$ and denote by \check{L} the backward λ -Liouville cone of \check{N} . Then $\hat{L} := \tilde{L} \cup \check{L} \cup L_0$ is the required arboreal Lagrangian with boundary. Its boundary (which can be smoothed) is the union of Λ_0 and the forward Liouville cone of C glued along C to \check{N} .

The required Liouville form λ_1 can then be constructed as follows: starting with λ , convert the ribbon of $\hat{\Lambda}$ to a boundary nucleus and then back to a Wc-hypersurface. This has the effect of adding the backwards Liouville cone of $\hat{\Lambda}$ to the skeleton. Then, up to smoothing, further conversion of the ribbons of $\Lambda \cup \Lambda_0$ to boundary nucleus will produce a Wc-structure with skeleton equal to \hat{L} . The positive cotangent building structure is obtained by stabilizing the positive cotangent building structure for the ribbon of Λ and adding the block $\mathcal{T}^*\Lambda_0$. \square

6.2.2. *Parametric version.* Lemma 6.11 also holds in a parametric form. Namely:

Lemma 6.12. *Let $\Lambda^z \subset S^{2n-1}$ be a family of positive arboreal Legendrians in S^{2n-1} endowed with the standard contact structure, parametrized by $z \in Z$ for Z a compact manifold. Let $L^z \subset \mathbb{R}^{2n}$ be the Lagrangian cone over Λ^z centered at 0, i.e. with respect to the radial Liouville structure on the unit ball $B \subset \mathbb{R}^{2n}$. Let (τ_z, ν_z) be a family of polarizations of \mathbb{R}^{2n} such that*

every Lagrangian plane T_z tangent to L^z at 0 satisfies the condition $T_z^{\tau_z, \nu_z} > 0$. Let $L_0^z = \tau_z(0)$ be the Lagrangian plane through the origin, and $\Lambda_0^z \subset S^{2n-1}$ its Legendrian link. Suppose that each boundary component of Λ^z is bounding and moreover that the bounding manifolds N^z form a fibre bundle over Z . Then there exist:

- (i) a family of closed arboreal Legendrians $\widehat{\Lambda}^z \supset \Lambda^z$ such that $\Lambda'^z = \widehat{\Lambda}^z \setminus \Lambda^z$ is smooth and such that every Lagrangian plane T_z tangent to the Liouville cone L'^z over Λ'^z satisfies the condition $T_z^{\tau_z, \nu_z} > 0$;
- (ii) a family of asymptotically conical arboreal Lagrangians \widehat{L}^z in \mathbb{R}^{2n} endowed with the standard Liouville structure such that:
 - \widehat{L}^z coincides at infinity with the Lagrangian cone over $\Lambda^z \cup \Lambda_0^z$;
 - $\widehat{L}^z \supset \widehat{\Lambda}^z$ and $\widehat{L}^z \supset L_0^z$;
 - \widehat{L}^z has smooth boundary and every boundary component is bounding, moreover with bounding manifolds forming a fibre bundle over Z ;
 - \widehat{L}^z is transverse to any family of Lagrangian distributions $\eta_z \in C(\nu_z, \tau_z)$, i.e. negative with respect to (τ_z, ν_z) .
- (iii) a family of compactly supported homotopies $\lambda_t^z = \lambda_0 + df_t^z$ of $\lambda_0 = pdq$ such that the Wc-manifold obtained from $(\mathbb{R}^{2n}, \lambda_1^z)$ by converting the ribbons of the Legendrians $\Lambda^z \cup \Lambda_0^z$ into nuclei of boundary faces is a positive cotangent building with skeleton \widehat{L}^z .

Proof. The proof proceeds exactly as in the non-parametric case by adding a parameter everywhere. \square

6.3. From ridgy to arboreal.

6.3.1. *Recap on reduced positivity.* We recall the Definition 4.26 of reduced positivity. Let $L \subset X$ be a positive arboreal Lagrangian and Z a non-zero Liouville vector field tangent to L . A Lagrangian distribution μ along L is called reduced positive with respect to (L, Z) if for any singular point $a \in L$ the following condition is satisfied. Let T be the tangent space to the root Lagrangian at a , and T' a tangent plane to any other smooth piece adjacent to a . Then $[T']^\zeta \in C([T]^\zeta, [\eta(a)]^\zeta)$, where $\zeta = \text{Span}(Z)^\perp$.

6.3.2. *The distribution ν_∞ .* We consider in this section the following setup. Let M be a bc-manifold and $X = \mathcal{T}^*M$. Suppose the boundary faces of M are ordered: $\partial_1 M = P_1 \cup \dots \cup P_k$. Let u_j be defining coordinates near faces P_j . Suppose we are given the following objects for each $j = 1, \dots, k$:

- a neighborhood $V_j \supset P_j$;
- the canonical Liouville fields Z_j on \mathcal{T}^*P_j and the Liouville field $\widehat{Z}_j := u_j \frac{\partial}{\partial u_j} + Z_j$, on V_j .
- a Lagrangian distribution ν_j on V_j , which is tangent to \widehat{Z}_j and invariant with respect to its negative flow.

Suppose, in addition, we are given

- a ridgy Lagrangian $L \subset \mathcal{T}^*M$ which on $V_j \cap L$ is tangent to \widehat{Z}_j ;
- a Lagrangian distribution ν_{-1} on \mathcal{T}^*M which is transverse to L and all distributions ν_j .

Suppose that the following conditions are satisfied:

- for any multi-index $I = \{i_1 < \dots < i_\ell\}$ we have

$$[\nu_{i_\ell}(a)]^{I'} \prec \dots \prec [\nu_{i_1}(a)]^{I'} \prec [\nu_{-1}(a)]^{I'} \text{ on } \bigcap_1^\ell V_{i_j} \setminus \bigcup_{j \notin I} V_j;$$

where $I' = \{i_1 < \dots < i_{\ell-1}\}$.

- for any point $a \in L \cap V_j$ and any tangent plane T to L at a we have

$$[T]^j \prec [\nu_j]^j \prec [\nu_{-1}]^j.$$




FIGURE 6.5. The condition on $[\nu_{-1}]^1$.




FIGURE 6.6. The condition on a Lagrangian plane T tangent to L .

Lemma 6.13. *There exist neighborhoods $V_j \ni V'_j \supset P_j$, $j = 1, \dots, k$, and a distribution ν_∞ on \mathcal{T}^*M which satisfies the following conditions:*

- (i) $\nu_\infty = \nu_m$ on $V'_m \setminus \bigcup_{j>m} V_j$, $m = 1, \dots, k$;
 (ii) for any $1 \leq i < j \leq k$ and any point $a \in V'_i \cap (V_j \setminus V'_j)$ we have

$$[\nu_\infty(a)]^I \in C([\nu_i(a)]^I, [\nu_j(a)]^I),$$

where I is the multi-index $\{i_0 = i < i_1 < \dots < i_\ell\}$, $i_\ell < j$, such that

$$a \in \bigcap_1^\ell V_{i_s} \setminus \bigcup_{m \notin I, m < j} V_m.$$

- (iii) ν_∞ is transverse to ν_{-1} and for any point $a \in L \setminus \bigcup_1^k V_j$ and any tangent plane T to L at a we have $T \prec \nu_\infty(a) \prec \nu_{-1}(a)$.




FIGURE 6.7. The conclusion on $[\nu_\infty]^1$.




FIGURE 6.8. The conclusion on a Lagrangian plane T tangent to L .

Proof. Define the required distribution as equal to ν_m on $V'_m \setminus \bigcup_{j>m} V_j$, $m = 1, \dots, k$, and any distribution on $\mathcal{T}^*M \setminus \bigcup_1^k V_j$ which is transverse to ν_{-1} and satisfies condition (iii) along L . Next, we successively extend ν_∞ to V_k : first to $(V_k \setminus V'_k) \cap V'_{k-1}$ to satisfy the condition

$[\nu_\infty]^{k-1} \in C([\nu_k]^{k-1}, [\nu_{k-1}]^{k-1})$, then continuing the process by extending to $V_k \setminus (V'_k \cup V'_{k-1}) \cap V'_{k-2}$ to satisfy the condition $[\nu_\infty]^{k-1} \in C([\nu_k]^{k-2, k-1}, [\nu_{k-1}]^{k-2, k-1})$, etc. Next, we similarly successively extend ν_∞ to $V_{k-1} \setminus (V'_{k-1} \cap V_k)$, $V_{k-2} \setminus (V'_{k-2} \cup V_{k-1} \cup V_k)$, \dots , $V_1 \setminus (V'_1 \cup V_2 \cup \dots \cup V_k)$. Each extension is possible because it amounts to an extension of a section of a fibration with open convex (and therefore contractible) fiber. \square

6.3.3. Arborealization of ridges. Recall that for an isotropic submanifold C of a symplectic manifold, its *symplectic normal bundle*, is defined as $(TC)^\perp/TC$. Denote by $\pi_C = [\cdot]^C$ the reduction map. We consider a ridgy Lagrangian $L \subset \mathcal{T}^*M$ with the notation as in Section 6.3.2 above. Consider the stratification $L = L_0 \cup L_1 \cup \dots \cup L_n$, where L_j is the j -dimensional locus of order $(n-j)$ ridges. Thus the top-dimensional stratum L_n is the smooth locus of L .

For each component C_j of L_j , we denote by $N(C_j)$ the $2j$ -dimensional symplectic normal bundle to \overline{C}_j . A Lagrangian tangent plane T to L at a point of \overline{C}_j reduces to a Lagrangian plane $[T]^{C_j} := \pi_{C_j}(T) \subset N(C_j)$. If moreover $a \in C_{j-1} \subset \overline{C}_j$ for C_{j-1} a component of L_{j-1} and T a Lagrangian plane at a , then the plane $[T]^{C_{j-1}}$ further reduces to $[T]^{C_{j-1}}$.

Note that if L is tangent to \widehat{Z}_j on V_j , $j = 1, \dots, k$, then so are all $L_i, i = 1, \dots, k$. In particular, if $i < \ell$ we have $L_i \cap V_{i_1} \cap \dots \cap V_{i_\ell} = \emptyset$.

Let $U_n \rightarrow \dots \rightarrow U_0$ be the canonical Wc-structure associated to a ridgy Lagrangian L .

Definition 6.14. An *arborealization* of the ridgy Lagrangian L is the structure of a cotangent building $B_m \rightarrow \dots \rightarrow B_0$ on a neighborhood of L whose skeleton is arboreal and whose underlying Weinstein manifold is homotopic to that of $U_n \rightarrow \dots \rightarrow U_0$.

We say that the arborealization is positive when the resulting arboreal Lagrangian is positive, which implies $B_m \rightarrow \dots \rightarrow B_0$ is a positive complex.

In the next proposition we continue using the notation and assumptions introduced in Section 6.3.2.

Proposition 6.15. *The ridgy Lagrangian L can be arborealized to a positive arboreal Lagrangian \widehat{L} that is positive with respect to ν_{-1} , and moreover, over V'_j is invariant with respect to the negative flow of \widehat{Z}_j and reduced positive with respect to ν_j , for $j = 1, \dots, k$.*

Proof. Any Lagrangian distribution can be viewed as a field of quadratic forms with respect to the polarization (ν_∞, ν_{-1}) . Fix a field of positive definite quadratic forms Q . Denote by η_t the Lagrangian distribution generated by $-tQ$, and by ζ_t the distribution generated by $\frac{1}{t}Q$. Choose $\varepsilon > 0$ so small that the following conditions are satisfied:

- $\nu_j \prec \zeta_\varepsilon \prec \nu_{-1}$ on V_j , $j = 1, \dots, k$;
- for any point $a \in L \cap V_j$ and any tangent plane T to L at a we have

$$[T]^j \prec [\eta_\varepsilon]^j \prec [\nu_\infty]^j.$$

For each component C_j of L_j define Lagrangian distributions in the bundle $N(C_j)$ over \overline{C}_j , $j = 0, \dots, n$.

$$\begin{aligned}\underline{s}_j &:= [\eta_{\frac{\varepsilon}{j+1}}|_{C_j}]^{C_j}, \\ \overline{s}_j &:= [\zeta_{\frac{\varepsilon}{n-j+1}}|_{C_j}]^{C_j},\end{aligned}$$

Note that if $C_{j-1} \subset \overline{C}_j$ then

$$[\underline{s}_{j-1}]^{C_j} \prec \underline{s}_j|_{C_{j-1}} \prec \overline{s}_j|_{C_{j-1}} \prec [\overline{s}_{j-1}]^{C_j}.$$

We will refer to the system of distributions $\underline{s}_j, \overline{s}_j$ as a *frame of arborealization* for L .




FIGURE 6.9. A frame of arborealization is nested as illustrated in the figure.

Consider the Wc-building structure $U_n \rightarrow \dots \rightarrow U_0$ on a neighborhood of L where the germ \mathcal{U}_j is decomposed as $\bigcup_Q \mathcal{F}^* Q^{\Delta, j-1} \times \mathcal{R}^{2(n-j)}$. The Wc-manifold W is obtained by successively gluing U_n to U_{n-1} , the result of that to U_{n-2} , the result of that to U_{n-3} , etc. In this process we get a sequence of Wc-manifolds $W^{>j} = \bigcup_{i>j} U_i$ with skeleta $L^{>j} = L \setminus \bigcup_{i \geq j} U_i$. We will be proving the proposition by induction for $W^{>n-j}$, $j = 1, \dots, n+1$. We will also include into the induction hypothesis the following additional property. Recall that $W^{>j}$ is attached vertically to U_j along a Wc-hypersurface Σ (which is the nucleus of the attaching face) whose skeleton is fibered over $L_j^{\Delta, j-1}$ with the fiber which is the link ∂R^{n-j} of the ridgy singularity R^{n-j} . We will require that the arborealization \widehat{L}^j intersects the attaching face in an arboreal skeleton of Σ which is fibered over $L_j^{\Delta, j-1}$, whose fiber is the arborealization of the link ∂R^{n-j} .

The base of induction $j = 0$ is trivial because L_n in this case is smooth. Suppose that we already arborealized the ridgy Lagrangian $L^{>j}$ to $\widehat{L}^{>j}$ and deformed correspondingly the Weinstein structure on $W^{>j}$. i.e. made $\widehat{L}^{>j}$ the skeleton of the new structure so that, by inductive assumption, (i) the arboreal $\widehat{L}^{>j}$ has a smooth boundary $\partial \widehat{L}^{>j}$ which bounds an immersed n -dimensional manifold $A \subset \widehat{L}^{>j}$, (ii) $\widehat{L}^{>j}$ intersects the nucleus Σ in an arborealization of the skeleton of Σ and (iii) the skeleton is Legendrian for U_j . We also observe

that the condition that each boundary component is bounding is inherited by the skeleton of Σ . Hence, using the fiberwise polarization $(\tau = \bar{s}_j, \nu = \underline{s}_j)$ we can apply Lemma 6.12 to arborealize the fiberwise Lagrangian cone over the arboreal Legendrian.

Note that by our construction, the distribution ν_{-1} is positive for the constructed arboreal \widehat{L} , and ν_j are reduced positive on V'_j . \square




FIGURE 6.10. In the case $n = 2$ the inductive procedure has 2 steps. The figure illustrates the moment between the two steps: one has arborealized the order 1 ridges and it remains to arborealize a radial Liouville cone over an arboreal Legendrian with smooth boundary in a sphere around point corresponding to a order 2-ridge.

6.4. **Conclusion of the proof.** We are now ready to prove our main result.

6.4.1. *Proof of the main theorem.*

Theorem 6.16. *Let (X, λ) be a Wc-manifold and $\nu \subset TW$ a Lagrangian plane field. Then there exists a Weinstein homotopy λ_t of $\lambda_0 = \lambda$ and a Wc-hypersurface $A \subset X \setminus \text{Skel}(X, \lambda_1)$ such that the result of converting A to a boundary nucleus yields a Wc-manifold (X, λ_1^A) which admits a structure of a positive cotangent building with a minimal distribution ν_{-1} equal to ν . In particular, $\text{Skel}(X, \lambda_1^A)$ is a positive arboreal Lagrangian (with boundary) transverse to ν . Moreover, we can arrange it so that $\text{Skel}(A, \lambda)$ is smooth, i.e. $\text{Skel}(X, \lambda_1^A)$ has smooth boundary.*

Proof. We choose a presentation of (X, λ) as a cotangent building $(B_k \rightarrow \cdots \rightarrow B_0)$, $B_i = \mathcal{T}^*M_i$, see Proposition 2.69. Without loss of generality we may assume that the restriction of ν to B_0 serves as ν_{-1} , i.e. satisfies the condition $\nu_0 \in C(TM_0, \nu_{-1})$ over M_0 . Indeed, M_0 is a disk, $B_0 = T^*M$ is a ball and we are free to change polarization if desired. The cotangent block B_1 is attached to B_0 along \mathcal{T}^*P , where P is a boundary face of M_1 .

Consider the block B_1 and the restrictions of the distributions ν_0 and ν to B_1 . Applying Theorem 5.10 we can find a ridgy isotopy of M_1 to a ridgy Lagrangian M'_1 transverse to ν , reduced transverse to ν_0 , invariant with respect to the negative flow of Z_0 , and such that at each point $a \in M'_1$ and any tangent plane T at a we have $[T]^0 \prec [\nu_0]^0 \prec [\nu_{-1}]^0$.

Moreover, this deformation can be realized by a deformation of skeleta of Weinstein structures of B_1 . Next, we apply to the ridgy Lagrangian M'_1 Proposition 6.15 to further deform the Weinstein structure on B_1 to a structure of a cotangent building, which is positive with respect to ν_{-1} and reduced positive with respect to ν_0 .

This yields a building structure on $B_0 \cup B_1$ with positive skeleton, positive with respect to ν_{-1} . Using Proposition 4.32 we can deform the building structure on this building to make it positive and positive for ν_{-1} .

Before we proceed with the next block, we note that the Lagrangian $\mathring{M}_2 \cap W^{(1)}$ is no longer conical for the constructed Wc-structure of $W^{(1)}$. So we apply Lemma 2.64 to arrange for this property to hold. Once this is achieved we can proceed as before to produce a Wc-manifold $(W^{(2)}, \lambda^{(2)})$ with a positive W-complex structure for which ν is a minimal distribution and whose skeleton $L^{(2)}$ is a positive arboreal Lagrangian with smooth boundary. Arguing by induction in the same way over the rest of the attaching blocks we conclude the proof. \square

6.4.2. *Variants of the main theorem.* Next we state our main theorem for Weinstein pairs:

Theorem 6.17. *Let (X, λ) be a Wc-manifold, $A_0 \subset X \setminus \text{Skel}(X, \lambda)$ a Wc-hypersurface and $\nu \subset TW$ a Lagrangian plane field. Then there exists a Weinstein homotopy λ_t of $\lambda_0 = \lambda$ and a Wc-hypersurface $A_1 \subset X \setminus \text{Skel}(X, \lambda_1)$ disjoint from A_0 such that the result of converting $A = A_0 \cup A_1$ to a boundary nucleus yields a Wc-manifold (X, λ_1^A) which admits a structure of a positive cotangent building with a minimal distribution ν_{-1} equal to ν . In particular, $\text{Skel}(X, \lambda_1^A)$ is a positive arboreal Lagrangian (with boundary) transverse to ν . Moreover, we can arrange it so that $\text{Skel}(A_1, \lambda)$ is smooth, i.e. $\text{Skel}(X, \lambda_1^A)$ has smooth boundary away from A_0 .*

Proof. The same proof works applied to the cotangent building structure associated to a Weinstein pair. \square

Finally, we state our main theorem in relative version:

Theorem 6.18. *Let (X, λ) be a Wc-manifold, $A_0 \subset X \setminus \text{Skel}(X, \lambda)$ a Wc-hypersurface and $\nu \subset TW$ a Lagrangian plane field. Suppose that A_0 is endowed with the structure of a positive cotangent building with minimal distribution equal to the reduction of $\nu|_{A_0}$. Assume moreover that ν is transverse to the Liouville field Z of X near A_0 . Then there exists a Weinstein homotopy λ_t of $\lambda_0 = \lambda$, fixed near A_0 , and a Wc-hypersurface $A_1 \subset X \setminus \text{Skel}(X, \lambda_1)$ disjoint from A_0 such that the result of converting $A = A_0 \cup A_1$ to a boundary nucleus yields a Wc-manifold (X, λ_1^A) which admits a structure of a positive cotangent building with a minimal distribution ν_{-1} equal to ν . In particular, $\text{Skel}(X, \lambda_1^A)$ is a positive arboreal Lagrangian (with boundary) transverse to ν and $\text{Skel}(X, \lambda_1^A) \cap A_0 = \text{Skel}(A_0, \lambda)$. Moreover, we can arrange it so that $\text{Skel}(A_1, \lambda)$ is smooth, i.e. $\text{Skel}(X, \lambda_1^A)$ has smooth boundary away from A_0 .*

Proof. All the intermediary steps hold in relative form, so one may apply the same proof to the cotangent building structure associated to the Weinstein pair keeping everything fixed near A_0 at every stage of the argument. \square

Theorem 6.16 implies our main Theorem 1.5 as stated in the Section 1.2, simply by taking the underlying Weinstein manifold of the Wc-manifold (X, λ_1^A) , i.e. converting A back to a Wc-hypersurface. Similarly, Theorems 6.17 and 6.18 imply the variants outlined in the remarks below the statement. For the concordance statement of Theorem 1.12 simply apply Theorem 6.18 to the Wc-pair obtained from $W \times T^*[0, 1]$ by converting the face nuclei $A_0 = (W \times 0) \cup (W \times 1)$ to Wc-hypersurfaces, where the Liouville form is $\lambda = \lambda_t + udt$ for λ_t the homotopy between λ_0 and λ_1 .

REFERENCES

- [AG18a] D. Álvarez-Gavela, *Refinements of the holonomic approximation lemma*, Algebraic & Geometric Topology 18 (2018) 2265–2303. [11](#), [77](#)
- [AG18b] D. Álvarez-Gavela, *The simplification of singularities of Lagrangian and Legendrian fronts*, Inventiones Mathematicae, 214(2) (2018) 641–737. [11](#), [78](#), [82](#)
- [AGEN20a] D. Alvarez-Gavela, Y. Eliashberg, and D. Nadler, *Arboreal models and their stability*, arXiv:2101.04272 [4](#), [5](#), [14](#), [44](#), [45](#), [48](#), [50](#), [70](#), [72](#), [82](#), [84](#)
- [AGEN19] D. Alvarez-Gavela, Y. Eliashberg, and D. Nadler, *Geomorphology of Lagrangian ridges*, arXiv:1912.03439 [12](#), [14](#), [73](#), [74](#), [75](#), [76](#), [77](#), [78](#)
- [AGEN21] D. Alvarez-Gavela, Y. Eliashberg, and D. Nadler, *Reidemeister moves for positive arboreal skeleta*, in preparation. [9](#)
- [A12] R. Avdek, *Liouville hypersurfaces and connect sum cobordisms*, arXiv:1204.3145. [26](#)
- [CE12] K. Cieliebak and Y. Eliashberg, *From Stein to Weinstein and Back – Symplectic Geometry of Affine Complex Manifolds*, Colloquium Publications Vol. 59, Amer. Math. Soc. (2012). [17](#), [18](#), [22](#), [26](#), [54](#)
- [EL17] T. Ekholm and Y. Lekili, *Duality between Lagrangian and Legendrian invariants*, arXiv:1701.01284. [26](#)
- [E95] Y. Eliashberg, *Symplectic geometry of plurisubharmonic functions*, in “Gauge Theory and Symplectic Geometry”, Vol. 488 of the series NATO ASI Series, 49–67. [18](#)
- [E90] Y. Eliashberg, *Topological characterization of Stein manifolds of dimension > 2* , Internat. J. Math. **1**(1990), no. 1, 29–46. [18](#)
- [E18] Y. Eliashberg, *Weinstein manifolds revisited*, Proc. of Symp. in Pure Math., Amer. Math. Soc., **99**(2018), 59–82. [26](#), [30](#)
- [EG91] Y. Eliashberg, M Gromov, *Convex symplectic manifolds*, Proc. Symp. Pure Math., **52**(1991), Amer. Math. Soc., Providence, RI, 135–162. [18](#)
- [En99] M. Entov, *Surgery on Lagrangian and Legendrian Singularities*, Geometric and Functional Analysis, 9(2) (1999) 298–352. [5](#) [11](#)
- [GPS17] S. Ganatra, J. Pardon and V. Shende, *Covariantly functorial Floer theory on Liouville sectors*, arXiv:1706.03152. [2](#), [26](#)
- [L92] F. Laudenbach, *On the Thom–Smale complex*, an Appendix to Bismut-Zhang, *An extension of a Theorem by Cheeger and Müller*, Astérisque 205(1992). [3](#)
- [MD91] D. McDuff, *Symplectic manifolds with contact type boundaries*, Invent. Math. **103**(1991), 651–671. [18](#)
- [N13] D. Nadler, *Arboreal Singularities*, arXiv:1309.4122. [1](#), [2](#)
- [N15] D. Nadler, *Non-characteristic expansion of Legendrian singularities*, arXiv:1507.01513. [2](#)
- [N16] D. Nadler, *Wrapped microlocal sheaves on pairs of pants*, arXiv:1604.00114. [2](#)

- [Se08] P. Seidel *Fukaya categories and Picard-Lefschetz theory*, Zurich Lectures in Advanced Mathematics, European Mathematical Society (EMS), Zürich, 2008. [2](#)
- [Sh18] V. Shende, *Arboreal singularities from Lefschetz fibrations*, arXiv:1809.10359. [2](#)
- [Sy16] Z. Sylvan, *On partially wrapped Fukaya categories*, arXiv: 1604:02540v2. [26](#)
- [St18] L. Starkston, *Arboreal Singularities in Weinstein Skeleta*, *Selecta Mathematica*, 24, 4105–4140 (2018) [2](#), [10](#), [11](#), [19](#), [21](#)
- [W91] A. Weinstein, *Contact surgery and symplectic handlebodies*, *Hokkaido Math. J.* **20**(1991), 241–251. [18](#)
- [Z18] A. Zorn, *A Combinatorial Model of Lagrangian Skeleta*, Ph.D. thesis, UC Berkeley, 2018. [9](#)

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA, 02139
Email address: dgavela@mit.edu

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, STANFORD, CA 94305
Email address: eliash@stanford.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, BERKELEY, CA 94720-3840
Email address: nadler@math.berkeley.edu