

RIGID INDECOMPOSABLE MODULES IN GRASSMANNIAN CLUSTER CATEGORIES

KARIN BAUR, DUSKO BOGDANIC, ANA GARCIA ELSENER, AND JIAN-RONG LI

ABSTRACT. The coordinate ring of the Grassmannian variety of k -dimensional subspaces in \mathbb{C}^n has a cluster algebra structure with Plücker relations giving rise to exchange relations. In this paper, we study indecomposable modules of the corresponding Grassmannian cluster categories $\text{CM}(B_{k,n})$. Jensen, King, and Su have associated a Kac-Moody root system $J_{k,n}$ to $\text{CM}(B_{k,n})$ and shown that in the finite types, rigid indecomposable modules correspond to roots. In general, the link between the category $\text{CM}(B_{k,n})$ and the root system $J_{k,n}$ remains mysterious and it is an open question whether indecomposables always give roots. In this paper, we provide evidence for this association in the infinite types: we show that every indecomposable rank 2 module corresponds to a root of the associated root system. We also show that indecomposable rank 3 modules in $\text{CM}(B_{3,n})$ all give rise to roots of $J_{3,n}$. For the rank 3 modules in $\text{CM}(B_{3,n})$ corresponding to real roots, we show that their underlying profiles are cyclic permutations of a certain canonical one. We also characterize the rank 3 modules in $\text{CM}(B_{3,n})$ corresponding to imaginary roots. By proving that there are exactly 225 profiles of rigid indecomposable rank 3 modules in $\text{CM}(B_{3,9})$ we confirm the link between the Grassmannian cluster category and the associated root system in this case. We conjecture that the profile of any rigid indecomposable module in $\text{CM}(B_{k,n})$ corresponding to a real root is a cyclic permutation of a canonical profile.

CONTENTS

1. Introduction	2
Acknowledgments	5
2. Grassmannian cluster categories	5
2.1. Cohen-Macaulay modules	5
2.2. Subspace configurations	7
2.3. Auslander-Reiten translations and shifts	9
2.4. The root system $J_{k,n}$	9
3. Canonical profiles and interlacing property	10
4. Rank 2 rigid indecomposable modules in $\text{CM}(B_{k,n})$	13
5. Subspace configurations of rank 3 modules	17
5.1. Necessary conditions for indecomposability	17
5.2. Rank 3 indecomposable modules in $\text{CM}(B_{3,n})$	18
6. Auslander-Reiten quiver	23
6.1. Profiles of Auslander-Reiten translates	23
6.2. Auslander-Reiten sequences	24
6.3. Rank 3 rigid indecomposable modules in $\text{CM}(B_{3,9})$	26

6.4. Proof of Theorem 6.7	28
6.5. Tubes with rank 1 modules	29
6.6. Rigid rank 3 modules in $\mathcal{R}_{\geq 2}$	29
6.7. Rigid rank 3 modules in $\mathcal{R}_{\geq 3}$	30
References	34

1. INTRODUCTION

Consider the homogeneous coordinate ring $\mathbb{C}[\mathrm{Gr}(2, n)]$ of the Grassmannian of 2-dimensional subspaces of \mathbb{C}^n . This is one of the key initial examples of Fomin and Zelevinsky's theory of cluster algebras, [8, §12.2]: the cluster variables are the Plücker coordinates, the exchange relations arise from the short Plücker relations, and clusters are in bijection with triangulations of a convex n -gon. Scott then proved in [19] that this cluster structure can be generalized to the coordinate ring $\mathbb{C}[\mathrm{Gr}(k, n)]$, where additional cluster variables appear (in general, infinitely many) and more exchange relations. This has sparked a lot of research activities in cluster theory, e.g. [20, 10, 12, 11, 18, 4, 13, 17, 9].

In particular, Jensen, King and Su showed in [13] that the category $\mathrm{CM}(B_{k,n})$ of Cohen-Macaulay modules over a quotient $B_{k,n}$ of a preprojective algebra of affine type A provides an additive categorification of Scott's cluster algebra structure. The category $\mathrm{CM}(B_{k,n})$ is called the Grassmannian cluster category. They also show that there is a cluster character on this category, sending rigid indecomposable objects to cluster variables ([13, Section 9]). Without loss of generality, we will assume $1 \leq k \leq n/2$ from now on.

Through this categorification, the classification of rigid indecomposable modules in $\mathrm{CM}(B_{k,n})$ (i.e. indecomposable modules M with $\mathrm{Ext}^1(M, M) = 0$) becomes an important tool towards characterising cluster variables in $\mathbb{C}[\mathrm{Gr}(k, n)]$ as well as in the classification of real prime modules of quantum affine algebras of type A , [12, 15].

In this paper, we study indecomposable modules of $\mathrm{CM}(B_{k,n})$ with the goal of providing an understanding of the associated cluster algebras.

Among the indecomposable modules are the rank 1 modules which are known to be in bijection with k -subsets of $\{1, 2, \dots, n\}$. These are the building blocks of the category as any module in $\mathrm{CM}(B_{k,n})$ can be filtered by rank 1 modules (see Section 2.1). Using this, in [13, Section 8], a map is defined from indecomposable modules of $\mathrm{CM}(B_{k,n})$ to a root lattice by associating a module with its class in the Grothendieck group and identifying the latter with a root lattice (see Section 2.4). Let $J_{k,n}$ be the graph with nodes $1, 2, \dots, n-1$ on a line and node n attached to node k , see Figure 1. If $k = 2$ or $k = 3$ and $n \in \{6, 7, 8\}$, $J_{k,n}$ is a Dynkin diagram, in general, it gives rise to a Kac-Moody algebra. In the Dynkin cases, the categories $\mathrm{CM}(B_{k,n})$ have only finitely many indecomposable objects and they are known to correspond to positive real roots for the associated root system of $J_{k,n}$, [13, Section 2].




FIGURE 1. The diagram of the root system $J_{k,n}$ associated with $\text{Gr}(k, n)$, we write β for α_n .

In general, it is a very difficult problem to describe the structure of the category or to classify indecomposable modules in $\text{CM}(B_{k,n})$. In contrast to the finite case, it is not clear how the correspondence between modules in the Grassmannian cluster categories and the root system arises. However, Jensen, King and Su [13] suspect that the classes of rigid indecomposable modules indeed are roots for $J_{k,n}$. Evidence for this was given in the small rank cases for certain infinite cases in [2]. The authors give a construction of rank 2 modules via short exact sequences, and find conditions that filtration factors of these modules have to fulfill. They also show that $\text{CM}(B_{k,n})$ has at most $2 \binom{n}{6} \binom{n-6}{k-3}$ (profiles of) rigid indecomposable rank 2 modules which correspond to real roots. Here, we also consider the imaginary roots and show in Theorem 4.7 that there are at most

$$N_{k,n} = \sum_{r=3}^k \left(\frac{2r}{3} \cdot p_1(r) + 2r \cdot p_2(r) + 4r \cdot p_3(r) \right) \cdot \binom{n}{2r} \binom{n-2r}{k-r},$$

profiles of rigid rank 2 indecomposable modules in $\text{CM}(B_{k,n})$, where $p_i(r)$ is the number of partitions $r = r_1 + r_2 + r_3$ such that $r_1, r_2, r_3 \in \mathbb{Z}_{\geq 1}$ and $|\{r_1, r_2, r_3\}| = i$.

We conjecture that $N_{k,n}$ in fact is equal to the number of rigid indecomposable rank 2 modules in $\text{CM}(B_{k,n})$ (Conjecture 4.8).

In fact, any indecomposable rank 2 module corresponds to a root of $J_{k,n}$ and for $k = 3$, we show that all rank 3 modules in $\text{CM}(B_{3,n})$ map to roots for $J_{3,n}$.

Theorem 1 (Lemma 4.6 and Theorem 5.6). (1) *Every indecomposable rank 2 module in $\text{CM}(B_{k,n})$ corresponds to a root for $J_{k,n}$.*

(2) *Every indecomposable module of rank 3 in $\text{CM}(B_{3,n})$ corresponds to a root for $J_{3,n}$.*

Recall that the modules in $\text{CM}(B_{k,n})$ can be filtered by rank 1 modules which in turn correspond to k -subsets of $[n] = \{1, \dots, n\}$ (see Section 2.1). The **profile** of a module is the collection of k -subsets of a filtration by rank 1 modules ([13, §6]). If the filtration of M has rank 1 factors I_1, \dots, I_m , for some $m > 0$, where the rank 1 module I_m is a submodule of M , we write $M = I_1 | I_2 | \dots | I_m$. We also write P_M for the profile of M .

We denote by $\mathcal{P}_{k,n}$ the set of profiles of indecomposable modules in $\text{CM}(B_{k,n})$. Its elements have m rows with k entries in $[n]$.

Let $P \in \mathcal{P}_{k,n}$ be a profile with m rows. Write $P = (P_{ij})_{ij}$, $1 \leq i \leq m$, $1 \leq j \leq k$, for the profile P where the entries in every row are written in increasing order. Then P is called **weakly column decreasing** if for every $j \in [k]$ and every $i \in [m-1]$, $P_{i,j} \geq P_{i+1,j}$. We call P **canonical**, if P is weakly column decreasing and if, in addition, $P_{m,j} \geq P_{1,j-1}$ for all $j \in [2, k]$. We write $C_{k,n}^{\text{re}}$ to denote the set of all canonical

profiles in $\mathcal{P}_{k,n}$ such that the corresponding module gives rise to a real root for $J_{k,n}$ (see Section 2.4).

With this notion, we are able to prove the following.

Theorem 2 (Theorem 5.7). *If M is an indecomposable rank 3 module in $\text{CM}(B_{3,n})$ such that M corresponds to a real root for $J_{3,n}$, then the profile P_M is a cyclic permutation of a canonical profile (i.e. a cyclic permutation of the rows of the profile of M is canonical).*

We expect that Theorem 2 is true for all indecomposable modules corresponding to real roots.

Conjecture 3 (Conjecture 5.8). *If $M \in \text{CM}(B_{k,n})$ is rigid indecomposable and corresponds to a real root for $J_{k,n}$, then P_M is a cyclic permutation of a canonical profile.*

We have the following result about modules corresponding to imaginary roots.

Theorem 4 (Theorem 5.9). *Suppose that the indecomposable module $M \in \text{CM}(B_{3,n})$ corresponds to an imaginary root of $J_{3,n}$. Then P_M is one of the following:*

$$\begin{aligned} & \underline{i_1 i_5 i_7} \quad \underline{i_2 i_6 i_8} \quad \underline{i_3 i_7 i_9} \quad \underline{i_1 i_4 i_8} \quad \underline{i_2 i_5 i_9} \quad \underline{i_1 i_3 i_6} \quad \underline{i_2 i_4 i_7} \quad \underline{i_3 i_5 i_8} \quad \underline{i_4 i_6 i_9} \quad \underline{i_1 i_4 i_7} \quad \underline{i_2 i_5 i_8} \quad \underline{i_3 i_6 i_9} \\ & \underline{i_3 i_6 i_9}, \underline{i_1 i_4 i_7}, \underline{i_2 i_5 i_8}, \underline{i_3 i_6 i_9}, \underline{i_1 i_4 i_7}, \underline{i_2 i_5 i_8}, \underline{i_3 i_6 i_9}, \underline{i_1 i_4 i_7}, \underline{i_2 i_5 i_8}, \underline{i_3 i_6 i_9}, \underline{i_1 i_4 i_7}, \underline{i_2 i_5 i_8}, \\ & \underline{i_2 i_4 i_8} \quad \underline{i_3 i_5 i_9} \quad \underline{i_1 i_4 i_6} \quad \underline{i_2 i_5 i_7} \quad \underline{i_3 i_6 i_8} \quad \underline{i_4 i_7 i_9} \quad \underline{i_1 i_5 i_8} \quad \underline{i_2 i_6 i_9} \quad \underline{i_1 i_3 i_7} \quad \underline{i_2 i_5 i_8} \quad \underline{i_3 i_6 i_9} \quad \underline{i_1 i_4 i_7} \end{aligned}$$

where $1 \leq i_1 < i_2 < \dots < i_9 \leq n$.

Note that for $n = 9$, there are exactly 12 indecomposable rank 3 modules corresponding to an imaginary root. The first 9 in the list are rigid and the last three are non-rigid; we show that in this case, there are exactly 225 rigid indecomposable rank 3 modules corresponding to roots of $J_{3,9}$.

For arbitrary $n > 9$, we expect that modules such as the last three in the list of Theorem 4 are always non-rigid.

A module $M' \in \text{CM}(B_{k,n})$ is said to be a shift of the module $M \in \text{CM}(B_{k,n})$ if the profile of M' is obtained from the profile P of M by adding a fixed number (mod n) to every entry of P (Definition 2.6).

Theorem 5 (Theorem 6.7 and Corollary 6.8). *Consider the category $\text{CM}(B_{3,9})$. Every rigid indecomposable rank 3 module maps to a root of $J_{3,9}$. Among the rigid indecomposable rank 3 modules, 216 correspond to a real root and the profile of each of them is a cyclic permutation of a canonical one. Furthermore, there are 9 modules mapping to*

157

an imaginary root and their profiles are all a shift of 369.

248

Theorem 5 is proved by studying the tubes of the Auslander-Reiten quiver of $\text{CM}(B_{3,9})$ containing rigid rank 3 modules.

We also prove the following result and its dual version (Theorem 6.5). If I is a k -subset of $[n]$, we write L_I for the corresponding rank 1 module. If I and J are two k -subsets, we write $L_I | L_J$ for the rank 2 module with submodule L_J and quotient L_I .




FIGURE 2. The graph C and the quiver Q_C , $n = 6$.

Theorem 6 (Theorem 6.3). *We consider the category $\text{CM}(B_{3,n})$. Let $I = \{i_1, i_2, i_3\}$ be a 3-subset where $i_j < i_{j+1} - 1$ for $j = 1, 2, 3$ (reducing modulo n). Let $X = \{i_1 + 1, i_2 + 1, i_3 + 1\}$ and $Y = \{i_1 + 2, i_2 + 2, i_3 + 2\}$. Then there is an Auslander-Reiten sequence*

$$L_I \hookrightarrow M \twoheadrightarrow \frac{L_X}{L_Y}$$

where M is a rank 3 module. If M is indecomposable, its profile is $X \mid I \mid Y$.

The paper is organized as follows. In Section 2, we recall the definition of key results about Grassmannian cluster categories. In Section 3, we study profiles and show that a weakly column decreasing profile is canonical if and only if any two rows of its profile are interlacing. In Section 4, we give an upper bound for the number of indecomposable rank 2 modules $\text{CM}(B_{k,n})$ corresponding to roots for $J_{k,n}$. In Section 5, we concentrate on rank 3 modules and study the subspace configurations. In Section 6, we provide short exact sequences to describe the structure of the Auslander-Reiten quiver for $\text{CM}(B_{3,n})$. We use this to determine the number of rigid indecomposable rank 3 modules in the infinite case $\text{CM}(B_{3,9})$.

Acknowledgments. We thank Alastair King, Matthew Pressland, and Andrei Smolensky for helpful discussions. K. B. was supported by a Royal Society Wolfson Fellowship. She is currently on leave from the University of Graz. D.B. was supported by the Austrian Science Fund Project Number P29807-35. J.-R.L. was supported by the Austrian Science Fund (FWF): M 2633-N32 Meitner Program.

2. GRASSMANNIAN CLUSTER CATEGORIES

In this section, we recall the definition of the Grassmannian cluster categories from [13] and some of their properties, see also [2]. Let $n \geq 4$ and recall that we always assume $1 \leq k \leq \frac{n}{2}$.

2.1. Cohen-Macaulay modules. Denote by $C = (C_0, C_1)$ the circular graph with vertex set $C_0 = \mathbb{Z}_n$ clockwise around the circle, and with the edge set $C_1 = \mathbb{Z}_n$, with edge i joining vertices $i - 1$ and i , cf. Figure 2. Denote by Q_C the quiver with the same vertex set C_0 and with arrows $x_i : i - 1 \rightarrow i$, $y_i : i \rightarrow i - 1$ for every $i \in C_0$, cf. Figure 2.

Denote by $B_{k,n}$ the quotient of the complete path algebra $\widehat{\mathbb{C}Q_C}$ by the ideal generated by the $2n$ relations $xy = yx$, $x^k = y^{n-k}$, where x, y are arrows of the form x_i, y_j for appropriate i, j (two relations for each vertex of Q_C).

The center Z of $B_{k,n}$ is the ring of formal power series $\mathbb{C}[[t]]$, where $t = \sum_{i=1}^n x_i y_i$. A (maximal) Cohen-Macaulay B -module is given by a representation $\{M_i : i \in C_0\}$ of Q_C , where each M_i is a free Z -module of the same rank (cf. [13, Section 3]).

Definition 2.1 ([13, Definition 3.5]). For any $B_{k,n}$ -module M and K the field of fractions of Z , the **rank** of M , denoted by $\text{rk}(M)$, is defined to be $\text{rk}(M) = \text{len}(M \otimes_Z K)$.

Jensen, King, and Su proved that the category $\text{CM}(B_{k,n})$ is an additive categorification of the cluster algebra structure on $\mathbb{C}[\text{Gr}(k, n)]$.

The category $\text{CM}(B_{k,n})$ is exact and Frobenius with projective-injective objects given by the $B_{k,n}$ projective modules, and it has an Auslander-Reiten quiver ([13, Remark 3.3]). We denote by $\tau(M)$ the Auslander-Reiten translation of M and by $\tau^{-1}(M)$ the inverse Auslander-Reiten translation of M . The category $\text{CM}(B_{k,n})$ is periodic under τ . For more details the reader can see [2, Section 3], and for further details on periodicity see [7], [16].

Remark 2.2. A module M in $\text{CM}(B_{k,n})$ is **rigid** if $\text{Ext}_{\text{CM}}^1(M, M) = 0$. Since $\text{CM}(B_{k,n})$ is Frobenius, [13, Corollary 3.7] we have

$$\text{Ext}_{\text{CM}}^i(M, N) = \underline{\text{Hom}}_{\text{CM}}(M, \Omega^{-i}N),$$

where $\underline{\text{Hom}}_{\text{CM}}(M, \Omega^{-i}N)$ is the stable space of $\text{Hom}_{\text{CM}}(M, \Omega^{-i}N)$. By construction, Ω^{-1} is the shift for the triangulated category $\underline{\text{CM}}(B_{k,n})$ (underline indicating the stable category). Note that $\tau = \Omega^{-1}$ as the category $\underline{\text{CM}}(B_{k,n})$ is 2-Calabi Yau. It follows that a non-projective module M in $\text{CM}(B_{k,n})$ is rigid if and only if $\tau(M)$ is rigid, as τ is an autoequivalence for the triangulated category $\underline{\text{CM}}(B_{k,n})$.

A special class of objects of $\text{C}(B_{k,n})$ are the rank 1 modules which are known to be rigid.

Definition 2.3 ([13, Definition 5.1]). For any k -subset I of C_1 , a rank 1 module L_I in $\text{CM}(B_{k,n})$ is defined by

$$L_I = (U_i, i \in C_0; x_i, y_i, i \in C_1),$$

where $U_i = \mathbb{C}[[t]]$, $i \in C_0$, and

$$\begin{aligned} x_i : U_{i-1} &\rightarrow U_i \text{ is given by multiplication by } 1 \text{ if } i \in I, \text{ and by } t \text{ if } i \notin I, \\ y_i : U_i &\rightarrow U_{i-1} \text{ is given by multiplication by } t \text{ if } i \in I, \text{ and by } 1 \text{ if } i \notin I. \end{aligned}$$

By [13, Proposition 5.2], every rank 1 module is isomorphic to L_I for some k -subset I of $[n]$. So there is a bijection between the rank 1 modules in $\text{CM}(B_{k,n})$ with the k -subsets of $[n]$ and the cluster variables of $\mathbb{C}[\text{Gr}(k, n)]$ which are Plücker coordinates.

It is convenient to represent the module L_I by a lattice diagram, see Figure 3. The spaces U_0, \dots, U_n are represented by columns from left to right and U_0 and U_n are identified. The vertices in each column correspond to the natural monomial \mathbb{C} -basis of $\mathbb{C}[t]$. The column corresponding to U_{i+1} is displaced half a step vertically downwards (resp. upwards) in relation to U_i if $i+1 \in I$ (resp. $i+1 \notin I$).




FIGURE 3. The lattice diagram of $L_{\{1,5,6\}}$ in $\text{CM}(B_{3,9})$ with its rim (blue arrows, with labels $x_1, y_2, y_3, y_4, x_5, x_6, y_7, y_8, y_9$).

The upper boundary of the lattice diagram of L_I is called the **rim of L_I** (we usually omit the arrows when drawing the rim). The k subset $I \subset [n]$ of the rank 1 module L_I can be read off as the set of labels on the edges going down to the right which are on the rim of L_I , i.e. the labels of the x_i 's appearing in the rim. For simplicity, we usually do not draw the labels of the rim in the figures.

We recall the notion of peaks and valleys of rank 1 modules from [1].

Definition 2.4. If I is a k -subset of $[n]$, the set of **peaks of L_I** is the set $\{i \mid i \notin I, i+1 \in I\}$. The set of **valleys of L_I** is $\{i \mid i \in I, i+1 \notin I\}$.

The rank 1 modules can be viewed as building blocks for the category as every module in $\text{CM}(B_{k,n})$ has a filtration with factors which are rank 1 modules (cf. [13, Proposition 6.6]). Let M be a rank m module in $\text{CM}(B_{k,n})$ with factors L_{I_1}, \dots, L_{I_m} in its filtration,

where L_{I_m} is a submodule of M . We write $M = L_{I_1}|L_{I_2}| \cdots |L_{I_m}$ or $M = \frac{L_{I_1}}{\vdots} / \frac{L_{I_m}}$. The

ordered collection of k -subsets I_1, \dots, I_m in the filtration of M is called the **profile** of

M , denoted P_M . We write $P_M = \frac{I_1}{\vdots} / \frac{I_m}$ or $P_M = I_1| \cdots | I_m$ if M has a filtration having

factors L_{I_1}, \dots, L_{I_m} (in this order). We sometimes write $M = P_M$ to indicate that M is a module with profile P_M . Note that in general, such a filtration is not unique, but in case M is rigid, the filtration is unique in the sense that it gives a canonical ordered set of rank 1 composition factors ([13, Lemma 6.2]).

2.2. Subspace configurations. Here, we follow the set-up of [13, Section 6]. Let M be a module in $\text{CM}(B_{k,n})$ and consider a filtration of M . We draw the lattice diagrams of all the rank 1 modules appearing in the filtration, in the filtration order. The rims of the filtration factors in this picture, together with the multiplicities of the elements in the monomial bases are called the set of **contours of M** . For indecomposable modules, contours are close-packed in the sense that there can be no “walk” (unoriented path) going around between two layers of the contours of M . The result is a lattice diagram where the multiplicities of the elements of the monomial basis are equal to the rank of M below the lowest contour and equal to 1 between the top two contours. For the representation M , connected regions with constant multiplicity in the lattice diagram




FIGURE 4. The subspace configuration (B) corresponds to the graph of contours (A). The subspace configuration (B) simplifies to (C).

correspond to isomorphic vector spaces, e.g. we can consider the region below the lowest contour to be $\mathbb{C}^{\text{rank } M}$. This view-point gives a subspace configuration associated with M , drawn as a poset, with the convention that an edge $i - j$ with $i < j$ denotes an inclusion of an i -dimensional space into a j -dimensional space. With slight abuse of notation, we also call the poset of a subspace configuration a subspace configuration. The poset (or subspace configuration) is a graph where the labels on the vertices are viewed as multiplicities, they are elements of $\{1, 2, \dots, \text{rank } M\}$, the multiplicity rank M only appears in one vertex.

The poset can thus be viewed as a representation for the underlying graph, with the orientations on the edges coming from the inclusions in M , i.e. pointing from the smaller to the larger numbers.

Example 2.5. Let M be a module in $\text{CM}(B_{3,9})$ with profile $169 \mid 147 \mid 358$. Figure 4 (A) shows the contours of M . In (A), below the first rim, every region has an upper boundary which is a part of the rim of the quotient M by the rank 1 modules above it. In (B), we show the poset of the subspace configuration of M . This graph can be simplified to (C) as we explain below.

If M is indecomposable, the subspace configuration is indecomposable, i.e. the poset diagram is an indecomposable representation for the oriented poset graph: if the poset diagram is a direct sum of two non-zero representations for the poset graph, this induces a direct sum decomposition of M in $\text{CM}(B_{k,n})$.

To see whether the subspace configuration is indecomposable, it is convenient to simplify the poset diagrams.

The poset diagrams of subspace configurations can be simplified as follows. A subspace can be identified as the intersection of several subspaces it maps into and may be omitted: for example, in Figure 4 (B), each of the 1-dimensional subspaces have to map in two different 2-dimensional subspaces of the 3-dimensional space at the bottom. In each case, these two different 2-dimensional subspaces intersect in a subspace of dimension 1. Hence this 1-dimensional subspace mapping into them is already determined and can be omitted from the diagram.

If after applying simplifications, the underlying graph of the poset is the graph of a Dynkin diagram we know that if the poset diagram is indecomposable, it has to be a positive root for it.

Under the above rules, the subspace configuration in Figure 4 (B) from Example 2.5 simplifies to (C). This is of Dynkin type D_4 and the simplified poset is not a root for D_4 , hence the poset diagram is not indecomposable.

2.3. Auslander-Reiten translations and shifts. By the symmetry of the algebra $B_{k,n}$, adding a fixed number to every element in every k -subset of the profile of a module and changing the linear maps in the representation accordingly preserves indecomposability and rigidity.

Definition 2.6. Let M and M' be indecomposable modules in $\text{CM}(B_{k,n})$, with profiles P and P' . If there is a number $a \in [n]$ such that P' is obtained by adding a to each entry of P (reducing modulo n), we say that M' is a **shift of M by a** and that the profile P' is a **shift of P (by a)**.

Lemma 2.7. *Let M and $M' \in \text{CM}(B_{k,n})$ be indecomposable and let M' be a shift of M . Then M is rigid if and only if M' is rigid.*

If the Auslander-Reiten translation leaves a module invariant, it cannot be rigid:

Lemma 2.8. *Let M be a module in $\underline{\text{CM}}(B_{k,n})$. If $\tau^{-1}(M) = M$, then M is non-rigid.*

Proof. If $M = \tau^{-1}(M)$, then there is an almost split sequence

$$M \rightarrow E \rightarrow M$$

in $\text{CM}(B_{k,n})$. This sequence does not split, so M is not rigid. \square

Lemma 2.9. *For any indecomposable module M in $\underline{\text{CM}}(B_{k,n})$, if the profile of $\tau^{-1}(M)$ is the same as the profile of M , then M is non-rigid.*

Proof. Suppose that $\tau^{-1}(M)$ and M have the same profile. If M is rigid, then $\tau^{-1}(M) = M$ since rigid indecomposable modules are uniquely determined by their profiles. By Lemma 2.8, M is non-rigid. This is a contradiction. Therefore M is non-rigid. \square

2.4. The root system $J_{k,n}$. Recall the graph $J_{k,n}$ with vertices $1, 2, \dots, n-1$ on a line and an additional vertex n attached to vertex k , see Figure 1. This graph gives rise to a Kac-Moody root system (we also call this root system $J_{k,n}$). It is of Dynkin type D_n for $k = 2$ and of Dynkin type E_6, E_7 and E_8 , respectively, for $k = 3$ and $n = 6, 7$ and 8 , respectively. The root lattice of the Kac-Moody algebra of $J_{k,n}$ can be identified with the lattice $\mathbb{Z}^n(k) := \{x = (x_i)_i \in \mathbb{Z}^n \mid k \text{ divides } \sum_i x_i\}$, equipped with the quadratic form $q(x) = \sum_i x_i^2 + \frac{2-k}{k^2}(\sum_i x_i)^2$. The roots of this root system satisfy $q(x) \leq 2$, among them, real roots have $q(x) = 2$ and imaginary ones have $q(x) < 2$.

Let e_1, \dots, e_n be the standard basis vector of \mathbb{R}^n . Define $\alpha_i = -e_i + e_{i+1}$ for $i \in [n-1]$ and $\beta = e_1 + e_2 + \dots + e_k$. Then the set $\{\alpha_1, \dots, \alpha_{n-1}, \beta\}$ is a system of simple roots for the root system $J_{k,n}$.

Let us write $\alpha_n = \beta$ and denote by (\cdot, \cdot) the standard Euclidean inner product on \mathbb{R}^n . We have $(\alpha_i, \alpha_i) = 2$ for all i and

$$(\alpha_i, \alpha_j) = \begin{cases} -1, & \text{if } |i-j| = 1, \\ 0, & \text{otherwise,} \end{cases} \quad (\beta, \alpha_i) = \begin{cases} -1, & \text{if } i = k, \\ 0, & \text{otherwise,} \end{cases}$$

Denote by s_α the reflection about the hyperplane perpendicular to a root α of $J_{k,n}$ and we write $s_i = s_{\alpha_i}$ with $i \in [n-1]$ and $s_\beta = s_{\alpha_n}$. For any root α of $J_{k,n}$ and $v \in \mathbb{R}^n$, $s_\alpha(v) = v - 2\frac{(v, \alpha)}{(\alpha, \alpha)}\alpha$. It is customary to abbreviate the scalar in front of α and to write $\langle v, \alpha^\vee \rangle := 2\frac{(v, \alpha)}{(\alpha, \alpha)}$ as we will do later.

The **Weyl group** W of $J_{k,n}$ is the group generated by the simple reflections. We have that, for $i \in [n-1]$,

$$\begin{aligned} s_i\left(\sum_{j=1}^n x_j e_j\right) &= \sum_{j=1}^n x_j e_j - 2 \frac{(\sum_{j=1}^n x_j e_j, \alpha_i)}{(\alpha_i, \alpha_i)} \alpha_i \\ &= \sum_{j=1}^n x_j e_j - (x_{i+1} - x_i)(e_{i+1} - e_i) = \sum_{j \in [n] \setminus \{i, i+1\}} x_j e_j + x_{i+1} e_i + x_i e_{i+1}. \end{aligned}$$

Therefore $s_i(x_1, \dots, x_n)$ is the exchange of entries x_i and x_{i+1} , $i \in [n-1]$. Similarly,

$$s_\beta(x_1, \dots, x_n) = (x_1 + r, \dots, x_k + r, x_{k+1}, \dots, x_n),$$

where $r = x_{k+1} + \dots + x_n - 2\frac{1}{k} \sum_{i=1}^n x_i$.

Jensen, King and Su define in [13, Sections 2,8] a map from the modules of $\text{CM}(B_{k,n})$ to the root lattice of $J_{k,n}$ by associating a module with its class in the Grothendieck group of $\text{CM}(B_{k,n})$ (the Grothendieck group is identified with the root lattice of $J_{k,n}$). This works as follows. Let M be a module and $P = P_M$ its profile. For $i = 1, \dots, n$, let x_i be the multiplicity of i in the union of the k -subsets in the profile P . Then let $x(M) = x(P_M) = (x_1, x_2, \dots, x_n) \in \mathbb{Z}^n$ be the vector of the multiplicities. By definition, $k \mid \sum_i x_i$, so $x(M) \in \mathbb{Z}^n(k)$ and there is an element in $\text{Span}_{\mathbb{R}}\{\beta, \alpha_1, \dots, \alpha_{n-1}\}$ corresponding to $x(M)$. We denote this element by $\varphi(M)$ and define $x(\varphi(M)) = x(M)$. We say that $M \in \text{CM}(B_{k,n})$ corresponds to a real (respectively, imaginary) root if $\varphi(M)$ is a real (respectively, imaginary) root of $J_{k,n}$. This terminology is motivated by the fact that in the finite types, all modules map to roots and by our results on indecomposable rank 2 and rank 3 modules.

Example 2.10. Let M be a module in $\text{CM}(B_{3,9})$ whose profile is $P_M = \frac{359}{\frac{258}{147}} \frac{146}{146}$. Then

$$x(M) = (2, 1, 1, 2, 2, 1, 1, 1, 1), \quad q(M) = 2, \quad \text{and} \quad \varphi(M) = 4\beta + 2\alpha_1 + 5\alpha_2 + 8\alpha_3 + 6\alpha_4 + 4\alpha_5 + 3\alpha_6 + 2\alpha_7 + \alpha_8.$$

3. CANONICAL PROFILES AND INTERLACING PROPERTY

Modules in $\text{CM}(B_{k,n})$ corresponding to real roots are closely related to canonical profiles which we will discuss in this section. In particular, we will prove that a weakly column decreasing profile is canonical if and only if any two rows of the profile are interlacing.

Definition 3.1. We say that two sets J_1, J_2 of integers are r -interlacing if $|J_1 \setminus J_2| = |J_2 \setminus J_1| = r$ and $J_1 \setminus J_2 = \{a_1, a_2, \dots, a_r\}$, $J_2 \setminus J_1 = \{b_1, b_2, \dots, b_r\}$, $a_i < a_{i+1}$, $b_i < b_{i+1}$, $i \in [r-1]$, either $a_1 < b_1 < a_2 < b_2 < \dots < b_{r-1} < a_r < b_r$ or $b_1 < a_1 < b_2 < a_2 < \dots < a_{r-1} < b_r < a_r$. In particular, if $J_1 = J_2$, the sets are 0-interlacing. We call J_1, J_2 interlacing if they are i -interlacing for some $i \in \mathbb{Z}_{\geq 0}$.

A profile $P \in \mathcal{P}_{k,n}$ with k -subsets J_1, \dots, J_m is called **interlacing** if for any $i \neq j$, J_i and J_j are interlacing.

The profile P_M from Example 2.10 is interlacing, e.g. the first row and the second row of P_M are 2-interlacing, the third row and the fourth row of P_M are 1-interlacing.

Remark 3.2. The definition of i -interlacing we use in this paper is called tightly i -interlacing in [2].

Remark 3.3. Observe that there are profiles where any two successive rows and the first and the last row are interlacing but where the profile is not interlacing. An example is

the profile $\begin{array}{cccc} 1 & 8 & 15 & 20 \\ \hline 2 & 11 & 18 & 23 \\ 5 & 17 & 21 & 30 \\ \hline 4 & 13 & 19 & 29 \end{array}$ where rows 1 and 3 are not interlacing.

Recall from the introduction that if $P \in \mathcal{P}_{k,n}$ is a profile with m rows, we write $P = (P_{ij})$, $1 \leq i \leq m$, $1 \leq j \leq k$, for the profile P with all rows written in increasing order.

Definition 3.4. Let P be a profile in $\mathcal{P}_{k,n}$ with m rows.

- (1) P is called **weakly column decreasing** if for every $j \in [k]$ and for every $i \in [m-1]$, we have $P_{i,j} \geq P_{i+1,j}$.
- (2) P is called **canonical**, if it is weakly column decreasing and if for all $j \in [2, k]$, $P_{m,j} \geq P_{1,j-1}$.
- (3) If P is canonical and $q(P) = 2$, we say it is **real**. We write $C_{k,n}^{\text{re}} \subset \mathcal{P}_{k,n}$ for the set of all canonical profiles P with $q(P) = 2$.

Example 3.5. The profile $P = \begin{array}{c} 359 \\ \hline 258 \\ \hline 147 \\ \hline 146 \end{array}$ of $\mathcal{P}_{3,9}$ is canonical and $P \in C_{3,9}^{\text{re}}$.

Definition 3.6. Let P be a profile in $\mathcal{P}_{k,n}$ with m rows. Then a profile P' is a **cyclic permutation** of P if P' is obtained by cyclically permuting the rows of P .

Proposition 3.7. *Let $P \in \mathcal{P}_{k,n}$ be a weakly column decreasing profile. Then P is canonical if and only if any two rows of P are interlacing.*

Proof. For $m = 1$ there is nothing to show. So assume $m \geq 2$.

(\Rightarrow) Suppose that P is canonical and P has two or more rows. Write $P = (P_{ij})_{1 \leq i \leq m, 1 \leq j \leq k}$, $P_{ij} < P_{i,j+1}$, $i \in [m]$, $j \in [k-1]$. Consider any two rows

$$\begin{array}{ccc} P_{i_1,1} & \cdots & P_{i_1,k} \\ P_{i_2,1} & \cdots & P_{i_2,k} \end{array}$$

with $i_1 < i_2$. Let $I_i := \{P_{i_j,1}, \dots, P_{i_j,k}\}$ for $j = 1, 2$. These are both k -subsets of $[n]$ as P is a profile. Note that $|I_2 \setminus I_1| = |I_1 \setminus I_2| =: \ell$ for some $\ell \geq 0$.

To check the interlacing property, we consider $\tilde{I}_i := I_i \setminus (I_1 \cap I_2)$ for $i = 1, 2$. So we can write $\tilde{I}_1 = \{P_{i_1,j_1}, P_{i_1,j_2}, \dots, P_{i_1,j_\ell}\}$ and $\tilde{I}_2 = \{P_{i_2,j'_1}, P_{i_2,j'_2}, \dots, P_{i_2,j'_\ell}\}$ for $1 \leq j_1 < j_2 < \dots < j_\ell \leq k$ and $1 \leq j'_1 < j'_2 < \dots < j'_\ell \leq k$. Since P is canonical, $j'_r \leq j_r$ for $r = 1, \dots, \ell$ and $j_r < j'_{r+1}$ for $r = 1, \dots, \ell - 1$.

The ℓ -subsets \tilde{I}_1 and \tilde{I}_2 are disjoint subsets by construction. We need to see that they are ℓ -interlacing, i.e. that $P_{i_2,j'_1} < P_{i_1,j_1} < P_{i_2,j'_2} < \dots < P_{i_1,j_\ell}$.

In case $j'_r = j_r$, we have $P_{i_2, j'_r} \leq P_{i_1, j_r}$ by the weakly column decreasing condition and $P_{i_1, j_r} \leq P_{i_2, j'_{r+1}}$ since P is canonical. And since \tilde{I}_1 and \tilde{I}_2 are disjoint, both inequalities have to be strict.

So assume that $j'_r < j_r$. Then $P_{i_1, j'_r} < P_{i_1, j_r}$. By the weakly column decreasing condition, we have $P_{i_2, j'_r} \leq P_{i_1, j'_r}$. Therefore $P_{i_2, j'_r} < P_{i_1, j_r}$. Furthermore, $P_{i_1, j_r} \leq P_{i_2, j'_{r+1}}$ as P is canonical and so, since \tilde{I}_1 and \tilde{I}_2 are disjoint, we have $P_{i_1, j_r} < P_{i_2, j'_{r+1}}$.

(\Leftarrow) Let $P \in \mathcal{P}_{k,n}$ with $m \geq 2$ rows. Write $P = (P_{ij})_{1 \leq i \leq m, 1 \leq j \leq k}$, $P_{ij} < P_{i, j+1}$, $i \in [m]$, $j \in [k-1]$. For some $1 \leq j < k$, consider the four elements in rows 1 and m of P and in the two successive columns $j, j+1$:

$$P_{1,j} \quad P_{1,j+1}$$

$$P_{m,j} \quad P_{m,j+1}.$$

We have to show that $P_{1,j} \leq P_{m,j+1}$. Denote by I_1, I_m the k -subsets of the first and last row respectively.

a) If $P_{1,j} = P_{m,j}$, then we have $P_{1,j} < P_{m,j+1}$ since $P_{1,j} = P_{m,j} < P_{m,j+1}$.

b) Let $P_{1,j+1} = P_{m,j+1}$. Then $P_{1,j} < P_{m,j+1}$ as $P_{m,j+1} = P_{1,j+1} > P_{1,j}$.

So we have to consider the cases where $P_{1,j} \neq P_{m,j}$ and $P_{1,j+1} \neq P_{m,j+1}$. Since P is weakly column decreasing, this means that $P_{1,j} > P_{m,j}$ and $P_{1,j+1} > P_{m,j+1}$.

c) Let $P_{1,j} > P_{m,j}$ and $P_{1,j+1} > P_{m,j+1}$. Suppose that $P_{1,j} > P_{m,j+1}$. If $j = 1$, then the first row and the last row are not interlacing. This is a contradiction. Therefore $j > 1$. Now we have a set $\{P_{m,j}, P_{m,j+1}\}$ in which every element is smaller than $P_{1,j}$. We define this set $\{P_{m,j}, P_{m,j+1}\}$ to be the **active set** and the element $P_{1,j}$ the **active maximum**. We now check the elements $P_{1,j-1}, P_{1,j-2}, \dots, P_{1,1}$ one by one as follows: we compare each of these elements with the smallest element in the active set and with further elements of row m . Denote the minimum in the active set by A . In the beginning, we have $A = P_{m,j}$.

- If $P_{1,j-1} < A$ and $P_{1,j-1} \neq P_{m,j-1}$, then the elements in the active set $\{P_{m,j}, P_{m,j+1}\}$ lie between two consecutive numbers $P_{1,j-1}, P_{1,j}$ in $I_1 \setminus I_m$. This contradicts the assumption that the first and the last row are interlacing.
- If $P_{1,j-1} < A$ and $P_{1,j-1} = P_{m,j-1}$, we keep the active set and the active maximum and continue with $P_{1,j-2}$.
- If $P_{1,j-1} = A$, then we have $P_{m,j-1} < A = P_{1,j-1}$. We replace the active set by removing A and adding $P_{m,j-1}$. As $P_{1,j}$ is still larger than the elements of the new active set, we keep the active maximum. We then continue with $P_{1,j-2}$.
- If $P_{1,j-1} > A$ and $P_{1,j-1} \notin I_1 \cap I_m$, then $P_{m,j-1} < P_{m,j} = A < P_{1,j-1}$. We replace the active set by $\{P_{m,j-1}, P_{m,j}\} = \{P_{m,j-1}, A\}$ and replace the active maximum by $P_{1,j-1}$. We then continue with $P_{1,j-2}$.
- If $P_{1,j-1} > A$ and $P_{1,j-1} \in I_1 \cap I_m$, then $P_{m,j-1} < P_{m,j} = A < P_{1,j-1} < P_{1,j}$. We change the active set by removing $P_{1,j-1}$ (if it is contained in it - otherwise we do not remove anything from the active set) and by adding $P_{m,j-1}$. We keep the active maximum and continue with $P_{1,j-2}$.

Continuing this procedure, the active set always has two or more elements. By construction, every element in the active set is smaller than the active maximum. We




FIGURE 5. Two rims forming two boxes.

have one of the following situations which both contradict the assumption that I_1, I_m are interlacing:

- At some step, the elements in the active set lie between two consecutive elements in $I_1 \setminus I_m$.
- In the last step, after we check $P_{1,1}$, the elements in the active set are smaller than the active maximum element and this active maximum is the smallest element in $I_1 \setminus I_m$.

We thus get $P_{1,j} \leq P_{m,j+1}$ as claimed. \square

4. RANK 2 RIGID INDECOMPOSABLE MODULES IN $\text{CM}(B_{k,n})$

In this section, we give an upper bound for the number of indecomposable modules of rank 2 corresponding to roots in $\text{CM}(B_{k,n})$.

By [2], every rank 2 rigid indecomposable module in $\text{CM}(B_{k,n})$ corresponding to a real root is of the form $L_I \mid L_J$ where I and J are 3-interlacing. From that we can deduce that there are at most $2 \binom{n}{6} \binom{n-6}{k-3}$ such modules (as they come in pairs).

Recall from Subsection 2.2 that we can draw modules as lattice diagrams or as collections of rims, as in Figure 4. For simplicity, we often do not write the numbers (the dimensions of the vector spaces) in the regions of the lattice diagrams.

By abuse of notation, we will call a profile or the lattice diagram of a rank m module a **profile of rank m** or a **lattice diagram of rank m** .

Consider a lattice diagram of rank 2: it has two rims which may meet several times (in particular, they will do so if the module is indecomposable). Similarly, if we consider a lattice diagram of higher rank, any two successive rims may meet several times.

Definition 4.1. Consider a rank m module with filtration $L_{I_1} \mid L_{I_2} \mid \cdots \mid L_{I_m}$. Let R_1 and R_2 be the rims of L_{I_j} and $L_{I_{j+1}}$ for some $1 \leq j < m$.

We call the non-empty regions formed by the two rims, between any two successive meeting points (reducing modulo n , if necessary) the **boxes between** the two rims. In particular, we say that R_1 and R_2 **form n boxes** if there are n boxes between them.

We define the **size (or right-size) of a box** to be the sum of the sizes of the intervals in the part of I_j corresponding to this box.

The **co-size (or left-size) of a box** is the sum of the sizes of the intervals of I_j^c corresponding to this box.

Example 4.2. The two rims shown in Figure 5 form two boxes. The first box has size and co-size 2 and the second has size and co-size 1.




FIGURE 6. The module $\frac{135}{248}$ is a partial shift of M . Only the coloured parts of the graph are shifted.

In Figure 6 (a), there are three boxes. Two of them have size and co-size 1, one box has size 1 and co-size 3.

The map from indecomposable modules to elements of the root system does not depend on the order of the factors in the filtration, it only depends on the elements of the k -subsets in its profile. We thus give this set a name. Let P be the profile of an indecomposable module. The **content** of P , $\text{con}(P)$, is the multi-set consisting of the elements of all k -subsets in P , for example,

$$\text{con} \begin{pmatrix} \underline{1233} \\ \underline{4556} \\ \underline{7889} \end{pmatrix} = \{1, 2, 3, 3, 4, 5, 5, 6, 7, 8, 8, 9\}.$$

If we have a rank 2 module $M = I_1 \mid I_2$, adding a constant number to all entries in the involved k -subsets keeps some properties of the module (e.g. indecomposability, rigidity) but changes others, e.g. the content of P_M (i.e. the labels appearing in $I \cup J$) will change in general. We introduce an operation which will keep the content as we will need to study different indecomposable modules which give rise to the same root.

To define this operation, let $P = I_1 \mid I_2$ be a rank 2 profile in $\mathcal{P}_{k,n}$. Let $n' = n - |I_1^c \cap I_2^c| - |I_1 \cap I_2|$ and $\psi : [n] \setminus ((I_1^c \cap I_2^c) \cup (I_1 \cap I_2)) \rightarrow [n']$ be the bijection respecting the order inside these sets.

Definition 4.3. Let $P = I_1 \mid I_2 \in \mathcal{P}_{k,n}$ be of rank 2, let $c \in [n]$. For $j = 1, 2$, define I'_j as the set $\psi^{-1}(\psi(I_j) + c \pmod{n'})$. Then $P' := I'_1 \mid I'_2$ is called a **partial shift** of P .

The elements of $I_1 \cap I_2$ and the elements of $I_1^c \cap I_2^c$ are fixed under such a partial shift. So for elements in $I_1 \cap I_2$ the shift does not do anything.

Example 4.4. Let $P = I_1 \mid I_2$, $I_1 = \{2, 4, 8\}$ and $I_2 = \{1, 3, 5\}$. Then $I_1^c \cap I_2^c = \{6, 7\}$. Then the partial shift of P by $c = 1$ is $P' = I'_1 \mid I'_2$ for $I'_1 = \{1, 3, 5\}$, $I'_2 = \{2, 4, 8\}$. See Figure 6.

The following is immediate from the definition of partial shift.

Lemma 4.5. Let P be a profile in $\mathcal{P}_{k,n}$ and let P' be a partial shift of P . Then $\text{con}(P') = \text{con}(P)$. In particular, the root of P is the same as the root of P' . If P is of rank 2 and if the k -subsets are r -interlacing, then the k -subsets of P' are also r -interlacing.

Recall that we always assume $k \leq n/2$.

Lemma 4.6. *Let $M = L_I | L_J$ be an indecomposable rank 2 module in $\text{CM}(B_{k,n})$. Then the rims of L_I and L_J form at least three boxes. Furthermore, $q(M) \in \{2, 0, -2, \dots, 8 - 2k\}$ and $\varphi(M)$ is a root of $J_{k,n}$.*

Proof. That the two rims form at least three boxes follows from the subspace configurations (cf. Remark 3.2 in [2]): if there is no box (i.e. $I = J$), the poset diagram is a single vertex (i.e. of Dynkin type A_1) with multiplicity 2. If there is one box, the poset diagram is $1 \rightarrow 2$, a type A_2 diagram and if it has two boxes, the poset diagram is $1 \rightarrow 2 \leftarrow 1$, of type A_3 . In type A , the indecomposable representations have multiplicities ≤ 1 on all vertices and the 1's form a connected path in the diagram. So none of these is indecomposable.

Let $x(M) = x(P_M) = (x_1, \dots, x_n)$ be the vector of the multiplicities (Section 2.4).

Since M has rank 2, the sum $\sum_{i=1}^n x_i$ is equal to $2k$ and $0 \leq x_i \leq 2$ for all i . Also, since I and J form at least three boxes, we have $|I \setminus J| = |J \setminus I| \geq 3$ and so the number of x_i 's which are equal to 2 is at most $k - 3$. But then $\sum x_i^2 \leq 4(k - 3) + 6$.

Denote by a (resp. b) the number of 2's (resp. 1's) in $\{x_i : i \in [n]\}$. Note that b is even and $b \geq 6$ since there are at least three boxes. Then $\sum_i x_i^2 = 4a + b \leq 4k - 6$ and $\sum_i x_i = 2a + b = 2k$. Therefore $0 \leq a \leq k - 3$, $b = 2k - 2a$. Hence $q(M) = \sum_{i=1}^n x_i^2 - 4(k - 2) \in \{2, 0, -2, \dots, 8 - 2k\}$.

First we consider the case when $q(M) = 2$. That is, $a = k - 3$ and $b = 6$. We denote $x(M) = x(\varphi(M)) = x = (x_1, \dots, x_n)$. Up to Weyl group action, we may assume that $x(M) = (2, \dots, 2, 1, \dots, 1, 0, \dots, 0)$, where the number of 2's is $k - 3$ and the number of 1's is 6. Therefore

$$\varphi(M) = 2\beta + \alpha_{k-2} + 2\alpha_{k-1} + 3\alpha_k + 2\alpha_{k+1} + \alpha_{k+2}.$$

We have that $s_k s_{k+1} s_{k+2} s_{k-1} s_k s_{k+1} s_{k-2} s_{k-1} s_k s_\beta(\varphi(M)) = \beta$ which is a simple root. Therefore $\varphi(M)$ is a real root of $J_{k,n}$ by definition, [14, Section 5.1].

Now we consider the case when $q(M) < 2$. That is $a < k - 3$ and $b = 2k - 2a$. Denote by $Q^+ = \bigoplus_{i=1}^n \mathbb{Z}_{\geq 0} \alpha_i$ the positive part of the root lattice of $J_{k,n}$ where we write α_n for β for the moment. For $\alpha \in Q^+$, denote by $\text{supp}(\alpha)$ the support of α , i.e. the subdiagram of $J_{k,n}$ corresponding to the simple roots having non-zero coefficients in α . Let $K = \{\alpha \in Q^+ \setminus \{0\} \mid \langle \alpha, \alpha_i^\vee \rangle \leq 0, i \in [n] \text{ and } \text{supp}(\alpha) \text{ is connected}\}$. By [14, Theorem 5.4], the set of all positive imaginary roots of $J_{k,n}$ is equal to $\cup_{w \in W} w(K)$. Up to Weyl group action, we may assume that $x(M) = (2, \dots, 2, 1, \dots, 1, 0, \dots, 0)$, where the number of 2's is a and the number of 1's is b . Therefore

$$\varphi(M) = 2\beta + \alpha_{a+1} + 2\alpha_{a+2} + \dots + (k - a)\alpha_k + (k - a - 1)\alpha_{k+1} + \dots + 2\alpha_{a+b-2} + \alpha_{a+b-1}.$$

We have that $\langle \varphi(M), \alpha_a^\vee \rangle = \langle \varphi(M), \alpha_{a+b}^\vee \rangle = -1$, $\langle \varphi(M), \alpha_i^\vee \rangle = 0$ for $i \in [n-1] \setminus \{a, a+b\}$ and $\langle \varphi(M), \beta^\vee \rangle = 4 - (k - a) \leq 0$ since $a < k - 3$. Moreover, $\text{supp}(\varphi(M))$ is connected. Therefore $\varphi(M)$ is an imaginary root. \square

Note that if $a = 0$, $b = 2k$ and $q(M) = 8 - 2k$, whereas if $a = k - 3$, we have $b = 6$ and $q(M) = 2$. So when $k = 3$, all indecomposable rank 2 modules have $q(M) = 2$ and for $k > 3$, we can find rank 2 modules with the lower bound $q(M) = 8 - 2k$ by taking two k -interlacing subsets.

Theorem 4.7. *Consider rank 2 modules in $\text{CM}(B_{k,n})$.*

(1) *Let $M = L_I | L_J$ be rigid indecomposable. Then the rims of L_I and L_J form three*

boxes.

(2) The number of profiles of rigid indecomposable rank 2 modules in $\text{CM}(B_{k,n})$ is at most equal to

$$N_{k,n} = \sum_{r=3}^k \left(\frac{2r}{3} \cdot p_1(r) + 2r \cdot p_2(r) + 4r \cdot p_3(r) \right) \cdot \binom{n}{2r} \binom{n-2r}{k-r},$$

where $p_i(r)$ is the number of partitions $r = r_1 + r_2 + r_3$ such that $r_1, r_2, r_3 \in \mathbb{Z}_{\geq 1}$ and $|\{r_1, r_2, r_3\}| = i$.

Proof. Note that there are no indecomposable rank 2 modules for $k < 3$ and $N_{k,n} = 0$ in this case. So we can assume $k \geq 3$.

Part (1) follows from Lemma 4.6 and since by [3, Theorem 1.2], a rigid module can have at most 3 boxes.

For (2), we determine how many profiles with 2 rows and such that the module is indecomposable can arise for any given content. This will be the number $N_{k,r}$. To show this, we consider how many different profiles can arise from a given one through partial shifts and reordering of boxes. So assume I and J are the two rows of a rank 2 profile (of a rigid indecomposable module).

As all elements of $I \cap J$ and $I^c \cap J^c$ are fixed under partial shifts, it is enough to consider their reduced version, $I' := I \setminus I \cap J$ and $J' := J \setminus I \cap J$, for the set $\{1, 2, \dots, n\} \setminus (I^c \cap J^c) \cup (I \cap J)$. Note that the number of boxes for I and J is the same as the number of boxes for I' and J' . So it is enough to consider boxes for I' and J' .

So assume now that the rims of I' and J' form three boxes. We write M' for the induced rank 2 module. We have that $|I' \cup J'| = 2r$ for $r = k - |I \cap J|$, $r \in [3, k]$. Let r_1, r_2, r_3 be the sizes of the three boxes (ordered such that the box of size r_1 starts with the smallest $i \in \{1, 2, \dots, n\} \setminus (I^c \cap J^c)$). If $|\{r_1, r_2, r_3\}| = 1$ then $3 \mid r$ and the partial shifts yield $2r/3$ different modules with the same content. If $|\{r_1, r_2, r_3\}| = 2$, the partial shifts yield $2r$ different modules with the same content. If $|\{r_1, r_2, r_3\}| = 3$, the partial shifts yield $2r$ additional different modules with the same content. Reordering the boxes so that the sizes are r_1, r_3, r_2 (i.e. interchanging two boxes) yields $2r$ additional modules with the same content (as the sizes are all different). So $4r$ different modules with the same content. Summing up over all possible sizes of $I \cap J$, we get $N_{k,n}$ as an upper bound.

Now by [3, Theorem 1.1], for every such I, J there exists an indecomposable module, hence such a profile. This completes the proof. \square

Conjecture 4.8. *The number $N_{k,n}$ in Theorem 4.7 is equal to the number of rigid indecomposable rank 2 modules in $\text{CM}(B_{k,n})$.*

Note that in finite types the conjecture is true, the Auslander-Reiten quivers of these types are in Figures 10-12 of [13]. We also point out if one can show that $\text{CM}(B_{k,n})$ is tubular, then the conjecture is true, as then all indecomposable rank 2 modules are rigid. In particular, this means that for $(k, n) = (3, 9)$ and $(k, n) = (4, 8)$, the conjecture is true, as the category $\text{CM}(B_{k,n})$ is known to be tubular in these two cases, see [2, §3.5].

Example 4.9. In case $k = 4$ and $n = 8$, the formula gives

$$N_{4,8} = \sum_{r=3}^4 \left(\frac{2r}{3} \cdot p_1(r) + 2r \cdot p_2(r) + 4r \cdot p_3(r) \right) \cdot \binom{8}{2r} \binom{8-2r}{4-r},$$

The only possibilities to form three boxes are $r = 3$ with $r_1 = r_2 = r_3 = 1$ and $r = 4$ with $r_1 = r_2 = 1, r_3 = 2$. So $p_1(3) = 1, p_2(4) = 1$ and all other $p_i(r)$ are 0. The formula gives $N_{4,8} = 2p_1(3) \binom{8}{6} \binom{2}{1} + 8p_1(4) \binom{8}{8} \binom{0}{0} = 2 \cdot 28 \cdot 2 + 8 = 120$, which is equal to the number of rigid indecomposables in this case, cf. [2, Section 7]. For $r = 3$, all these modules correspond to real roots and the 8 modules with imaginary roots arise from $r = 4$.

5. SUBSPACE CONFIGURATIONS OF RANK 3 MODULES

In this section, we use the subspace configurations to derive necessary conditions for indecomposability (Subsection 5.1) and then use these to study the roots associated with indecomposable rank 3 modules (Subsection 5.2). We automatically have $k \geq 3$, as there are no indecomposable higher rank modules for $k \leq 2$. To find these necessary conditions, we consider the number of boxes between the two pairs of successive rims of M ; the number of valleys of the upper rim is an upper bound for the number of boxes.

Note that if a module M is indecomposable, any two successive rims of M are “closely packed” in the sense that there cannot be a walk between the two rims, [13, Section 6].

5.1. Necessary conditions for indecomposability. The first case is when $I = J$, then there are no boxes between the two rims of $L_I = L_J$.

Lemma 5.1. *Let $M = L_I|L_J|L_K \in \text{CM}(B_{k,n})$ be indecomposable. If $I = J$, then the rims of L_J and L_K form at least four boxes.*

Proof. Suppose that $M = L_I|L_J|L_K \in \text{CM}(B_{k,n})$ is indecomposable, with $I = J$. In this case, the subspace configuration of M looks like a star. It has one vertex with multiplicity 3 and $m \geq 0$ vertices with 2, each with an arrow towards 3. Such a subspace configuration is decomposable for $m \leq 3$. \square

Let I and J be k -subsets. Instead of writing that boxes are formed by the rims of L_I and L_J , we will also say that the boxes are formed by L_I and L_J or by I and J .

Lemma 5.2. *Let $M = L_I|L_J|L_K \in \text{CM}(B_{k,n})$ be indecomposable.*

(1) *If L_J and L_K form up to three boxes, then L_I, L_J form at least two boxes or the subspace configuration of M is (G) in Figure 7.*

(2) *If L_I, L_J form one box, then L_J, L_K form at least four boxes or the subspace configuration of M is (G) in Figure 7.*

Proof. Suppose that $M = L_I|L_J|L_K \in \text{CM}(B_{k,n})$ is indecomposable and L_J, L_K form at most three boxes. By Lemma 5.1, $I \neq J$, so L_I, L_J form at least one box. All possible subspace configurations of M when I and J form one box are in Figure 7. The only indecomposable one among them is (G). This proves (1).

For the same reason, if L_I, L_J form one box, then L_J, L_K form at least four boxes or the subspace configuration of M is (G). \square




FIGURE 7. All subspace configurations for $L_I|L_J|L_K$, where L_I, L_J form one box and L_J, L_K form at most three boxes.




FIGURE 8. All possible subspace configurations of $L_I|L_J|L_K$, L_I, L_J form two boxes and L_J, L_K form at most two boxes.

Lemma 5.3. *Let $M = L_I|L_J|L_K \in \text{CM}(B_{k,n})$ be indecomposable, and assume that L_I and L_J form two boxes. Then L_J, L_K form at least three boxes or the subspace configuration of M is (K) from Figure 8.*

Proof. Suppose that $M = L_I|L_J|L_K \in \text{CM}(B_{k,n})$ is indecomposable and that L_I, L_J form two boxes. Suppose that L_J, L_K form at most two boxes. All subspace configurations for such M where L_J and L_K form at most two boxes are in Figure 8. The only indecomposable one of them is (K). This proves the claim. \square

5.2. Rank 3 indecomposable modules in $\text{CM}(B_{3,n})$. If we restrict to $k = 3$, we can say even more about the profile of a rank 3 indecomposable module, as we will show now.

Lemma 5.4. *Suppose that $M = L_I|L_J|L_K \in \text{CM}(B_{3,n})$ is indecomposable. Then the subspace configuration of M cannot be (G) of Figure 7 or (K) of Figure 8.*

Proof. The subspace configurations (G) of Figure 7 and (K) of Figure 8 both correspond to a partition of a k -subset into four non-trivial parts, hence only occur for $k \geq 4$. \square

Lemma 5.5. *If $M = L_I|L_J|L_K \in \text{CM}(B_{3,n})$ is indecomposable, I and J are interlacing.*

Proof. Suppose that $M = L_I|L_J|L_K \in \text{CM}(B_{3,n})$ is indecomposable. Since $k = 3$, J and K form at most three boxes. Suppose that I and J are not interlacing, then they form at most two boxes (as $k = 3$). By Lemma 5.2, I and J then form exactly two boxes (as the subspace configuration of M cannot be Figure 7 (G) by Lemma 5.4).

Now we use Lemma 5.3 to see that J and K form at least three boxes (as the subspace configuration cannot be Figure 8 (K) by Lemma 5.4). As $k = 3$, J and K form exactly three boxes.

Since $k = 3$, one of the boxes has size 1 and one has size 2. Since I and J are not interlacing, the box of size 2 must be a 2×2 square. But then the rim of L_J has exactly two valleys, and since the number of valleys is an upper bound for the number of boxes, J and K cannot form three boxes. A contradiction. \square

Theorem 5.6. *Let $M \in \text{CM}(B_{3,n})$ be indecomposable of rank 3. Then $q(M) \in \{0, 2\}$ and $\varphi(M)$ is a real or imaginary root of $J_{3,n}$.*

Proof. Suppose that $M = L_I|L_J|L_K$ is an indecomposable module of rank 3 in $\text{CM}(B_{3,n})$. Then $n \geq 9$. It suffices to show that the content of M consists of 8 or 9 different numbers. If this is the case, we get, up to reordering the x_i , that the vector of multiplicities of M is $x(M) = (2, 1, 1, 1, 1, 1, 1, 0, \dots, 0) \in \mathbb{Z}^n$ in the first case, with $q(M) = 2$ and $x(M) = (1, 1, 1, 1, 1, 1, 1, 1, 0, \dots, 0) \in \mathbb{Z}^n$ in the second case, with $q(M) = 0$.

We consider the boxes formed by the rims. Note that since $k = 3$, any two successive rims form at most three boxes. We go through these cases.

- If $I = J$, M decomposes by Lemma 5.1, a contradiction.
- Suppose next that I and J form one box. Since J and K form at most three boxes, Lemma 5.2 tells that the subspace configuration of M is (G) in Figure 7. This is not possible by Lemma 5.4.
- Assume now that I and J form two boxes. Then using Lemma 5.3 and Lemma 5.4 we find that J and K have to form three boxes (each of size 1). In particular, they are 3-interlacing, $|J \cup K| = 6$ and $J \cap K = \emptyset$.

By Lemma 5.5, I and J are interlacing and since we have two boxes, I and J are 2-interlacing. So $|I \cap J| = 1$. This implies that $I \cup J \cup K$ has at least 7 different elements.

If $|I \cup J \cup K| = 7$, I and K have one element in common. This implies that in the subspace configuration, there is a 2 mapping into a 3 (from the common element of I and J). There is also a 1 mapping into two different 2's. Using the reduction from Section 2.2, we can remove this 1. The resulting diagram has two vertices with a 2 and each with a single edge mapping into the 3. Such a subspace configuration is never indecomposable.

So we get $|I \cup J \cup K| \geq 8$ as claimed.

- Suppose now that L_I, L_J form three boxes. In particular, as $k = 3$, they are 3-interlacing, $|I \cup J| = 6$ and $I \cap J = \emptyset$.

The subspace configurations for the cases where J and K form at most one box are not indecomposable, see Figure 9. So J and K form at least two boxes.

If $|I \cup J \cup K| = 6$, we have $K \subset I \cup J$ and since J and K form at least two boxes, $|J \cap K| \leq 1$. So either $K = I$ or two elements of K are in I ,




FIGURE 9. Subspace configurations for I, J forming three boxes and (A) $J = K$ or (B) J and K form one box

one is in J . In the first case, the subspace configuration consists of three 1's included simultaneously in two 2's. So they can be removed by the reduction of Section 2.2. This leaves three 2's mapping into one 3, not an indecomposable configuration. In the second case, we have two 1's mapping simultaneously into two 2's, they can be removed. The resulting subspace configuration contains two 2's with a single edge into the 3. Again, this is not indecomposable.

If $|I \cup J \cup K| = 7$, two elements of K are in $I \cup J$. These can be both in I or one in I and one in J . If they are both in I , we have, as before, two 1's in the subspace configuration mapping into two 2's each. Removing them leaves two 2's, each with only one edge to the 3. This is not an indecomposable subspace configuration.

If one is in I and one in J , we can also reduce two 1's mapping into two 2's each and get a subspace configuration which is not indecomposable.

Therefore $I \cup J \cup K$ has at least 8 different numbers.

We now prove that $\varphi(M)$ is a real or imaginary root in $J_{k,n}$. We will use a similar method as in the proof of Lemma 4.6.

First we consider the case when $q(M) = 2$. Up to Weyl group action, we may assume that $x(M) = (2, 1, 1, 1, 1, 1, 1, 1, 0, \dots, 0)$. Then

$$\varphi(M) = 3\beta + \alpha_1 + 3\alpha_2 + 5\alpha_3 + 4\alpha_4 + 3\alpha_5 + 2\alpha_6 + \alpha_7.$$

We have that

$$s_3 s_4 s_5 s_2 s_3 s_4 s_1 s_2 s_3 s_\beta s_6 s_7 s_5 s_6 s_4 s_5 s_3 s_4 s_2 s_3 s_\beta(\varphi(M)) = \beta$$

which is a simple root. Therefore $\varphi(M)$ is a real root of $J_{3,n}$.

Now we consider the case when $q(M) < 2$. Up to Weyl group action, we may assume that $x(M) = (1, 1, 1, 1, 1, 1, 1, 1, 0, \dots, 0)$. Therefore

$$\varphi(M) = 3\beta + 2\alpha_1 + 4\alpha_2 + 6\alpha_3 + 5\alpha_4 + 4\alpha_5 + 3\alpha_6 + 2\alpha_7 + \alpha_8.$$

We have that $\langle \varphi(M), \alpha_9^\vee \rangle = -1$, $\langle \varphi(M), \alpha_i^\vee \rangle = 0$ for $i \in [n] \setminus \{9\}$ (we denote $\beta = \alpha_n$). Moreover, $\text{supp}(\varphi(M))$ is connected. By [14, Theorem 5.4], $\varphi(M)$ is thus an imaginary root of $J_{3,n}$. □

Recall from Definition 3.4 that a profile $P = (P_{ij}) \in \mathcal{P}_{k,n}$ is said to be canonical if its entries are weakly column decreasing and if for all $j \geq 2$, $P_{m,j} \geq P_{1,j-1}$. The

set of canonical profiles with real root is denoted by $C_{k,n}^{\text{re}}$. Recall also that a cyclic permutation of a profile is obtained by permuting its rows cyclically (Definition 3.6).

Theorem 5.7. *Let $M \in \text{CM}(B_{3,n})$ be an indecomposable module of rank 3. If $q(M) = 2$, then P_M is a cyclic permutation of some P in $C_{3,n}^{\text{re}}$.*

Proof. Write $M = L_I|L_J|L_K$ with 3-subsets I, J, K . Recall that $k \geq 3$ as M has rank 3. We already know from the proof of Theorem 5.6 that $x(M) = (2, 1, \dots, 1, 0, \dots, 0)$, up to reordering the entries (the entry 1 appearing exactly 7 times).

In particular, I, J and K are pairwise different. If any two of them are interlacing, they have to be 2-interlacing or 3-interlacing (if they were 1-interlacing, $x(M)$ would contain at least two entries equal to 2).

By Lemma 5.5, I and J are interlacing. Assume first they are 2-interlacing. Then the rims of L_I and L_J form two boxes. By Lemma 5.3, L_J and L_K have to form three boxes, as otherwise, the subspace configuration of M would be the one in Figure 8 (K), which does not occur for $k = 3$ (Lemma 5.4). So J and K are 3-interlacing.

Let $\{i_1, \dots, i_9\}$ be the entries of the profile P_M of M . Without loss of generality we can assume $i_1 = i_2 < i_3 < \dots < i_9$ as the other cases are analogous.

Since I and J have the element i_1 in common, I and J are 2-interlacing and J and K are 3-interlacing, the only possible choices for the rank 1 modules are the following:

- (a) $L_I = i_1i_5i_8, L_J = i_1i_4i_7, L_K = i_3i_6i_9,$
- (b) $L_I = i_1i_5i_9, L_J = i_1i_4i_7, L_K = i_3i_6i_8,$
- (c) $L_I = i_1i_6i_8, L_J = i_1i_4i_7, L_K = i_3i_5i_9,$
- (d) $L_I = i_1i_6i_9, L_J = i_1i_4i_7, L_K = i_3i_5i_8,$
- (e) $L_I = i_1i_4i_7, L_J = i_1i_5i_8, L_K = i_3i_6i_9,$
- (f) $L_I = i_1i_3i_7, L_J = i_1i_5i_8, L_K = i_4i_6i_9,$
- (g) $L_I = i_1i_4i_6, L_J = i_1i_5i_8, L_K = i_3i_7i_9,$
- (h) $L_I = i_1i_3i_6, L_J = i_1i_5i_8, L_K = i_4i_7i_9.$

The only indecomposable subspace configuration among these is the one in (a) as one can check.

$$i_3i_6i_9$$

If M is as in (a), the profile P_M is a cyclic permutation of the canonical profile $i_1i_5i_8$.

$$i_1i_4i_7$$

Assume now that I and J are 3-interlacing. Then J and K are interlacing as otherwise, the subspace configuration would be three leaves with 1 mapping into a vertex with a 2 and from there one edge into a vertex with 3. Since $x(M)$ has only one entry 2, we must have that either I, K are 2-interlacing with J, K 3-interlacing, case (i), or that I, K are 3-interlacing with J, K 2-interlacing, case (ii).

As before, we assume without loss of generality that the entries of $I \cup J \cup K$ are of the form $i_1 = i_2 < i_3 < \dots < i_9$. Then the only indecomposable subspace configurations

are $\frac{i_1 i_4 i_7}{i_3 i_6 i_9}$ for (i), $\frac{i_1 i_5 i_8}{i_1 i_4 i_7}$ for (ii). \square

We expect that Theorem 5.7 is true for any indecomposable module in $\text{CM}(B_{k,n})$, in arbitrary rank.

Conjecture 5.8. *Let $M \in \text{CM}(B_{k,n})$ be rigid indecomposable and assume that $\varphi(M)$ is a real root in $J_{k,n}$. Then P_M is a cyclic permutation of a real canonical profile.*

Theorem 5.9. *Let $M \in \text{CM}(B_{3,n})$ be indecomposable of rank 3 and assume that $\varphi(M)$ is imaginary. Then the profile P_M of M is one of the following:*

$$\begin{aligned} & \frac{i_1 i_5 i_7}{i_3 i_6 i_9}, \frac{i_2 i_6 i_8}{i_3 i_7 i_9}, \frac{i_3 i_7 i_9}{i_1 i_4 i_8}, \frac{i_2 i_5 i_9}{i_1 i_3 i_6}, \frac{i_1 i_3 i_6}{i_2 i_4 i_7}, \frac{i_2 i_4 i_7}{i_3 i_5 i_8}, \frac{i_3 i_5 i_8}{i_4 i_6 i_9}, \frac{i_4 i_6 i_9}{i_1 i_4 i_7}, \frac{i_1 i_4 i_7}{i_2 i_5 i_8}, \frac{i_2 i_5 i_8}{i_3 i_6 i_9}, \\ & \frac{i_3 i_6 i_9}{i_1 i_4 i_7}, \frac{i_1 i_4 i_7}{i_2 i_5 i_8}, \frac{i_2 i_5 i_8}{i_3 i_6 i_9}, \frac{i_3 i_6 i_9}{i_1 i_4 i_7}, \frac{i_1 i_4 i_7}{i_2 i_5 i_8}, \frac{i_2 i_5 i_8}{i_3 i_6 i_9}, \frac{i_3 i_6 i_9}{i_1 i_4 i_7}, \frac{i_1 i_4 i_7}{i_2 i_5 i_8}, \frac{i_2 i_5 i_8}{i_3 i_6 i_9}, \\ & \frac{i_2 i_4 i_8}{i_3 i_5 i_9}, \frac{i_3 i_5 i_9}{i_1 i_4 i_6}, \frac{i_1 i_4 i_6}{i_2 i_5 i_7}, \frac{i_2 i_5 i_7}{i_3 i_6 i_8}, \frac{i_3 i_6 i_8}{i_4 i_7 i_9}, \frac{i_4 i_7 i_9}{i_1 i_5 i_8}, \frac{i_1 i_5 i_8}{i_2 i_6 i_9}, \frac{i_2 i_6 i_9}{i_1 i_3 i_7}, \frac{i_1 i_3 i_7}{i_2 i_5 i_8}, \frac{i_2 i_5 i_8}{i_3 i_6 i_9}, \frac{i_3 i_6 i_9}{i_1 i_4 i_7} \end{aligned}$$

where $i_1 < \dots < i_9 \in \mathbb{Z}_{\geq 1}$.

Proof. Denote by $i_1 \leq \dots \leq i_9$ the entries of P_M . Since $\varphi(M)$ is imaginary, we have $q(M) = 0$ and the vector $x(M)$ of multiplicities of M is $(1, 1, \dots, 1, 0, \dots, 0)$, up to reordering the x_i , with 1 occurring nine times, as seen in the proof of Theorem 5.6. Therefore I, J, K are pairwise different and if any two of them interlace, they are 3-interlacing.

By Lemma 5.5, I and J are thus 3-interlacing. If L_J, L_K are not interlacing, then $k = 3$ implies that L_J, L_K form one box. But then the subspace configuration of M is a graph with 3 leaves with label 1, mapping into a vertex with label 2 and the latter into a vertex with label 3, not indecomposable. So both I, J and J, K are 3-interlacing.

Assume that $i_1 \in I$. Then the three 3-subsets have to be as follows:

- (A) $I = i_1 i_3 i_6, J = i_2 i_5 i_8, K = i_4 i_7 i_9,$
- (B) $I = i_1 i_3 i_7, J = i_2 i_5 i_8, K = i_4 i_6 i_9,$
- (C) $I = i_1 i_4 i_6, J = i_2 i_5 i_8, K = i_3 i_7 i_9,$
- (D) $I = i_1 i_4 i_7, J = i_2 i_5 i_8, K = i_3 i_6 i_9,$
- (E) $I = i_1 i_4 i_7, J = i_3 i_6 i_9, K = i_2 i_5 i_8,$
- (F) $I = i_1 i_4 i_8, J = i_3 i_6 i_9, K = i_2 i_5 i_7,$
- (G) $I = i_1 i_5 i_7, J = i_3 i_6 i_9, K = i_2 i_4 i_8,$
- (H) $I = i_1 i_5 i_8, J = i_3 i_6 i_9, K = i_2 i_4 i_7.$

Only modules (A), (E), (F), (G) have indecomposable subspace configurations, as one can check.

Similarly, if i_1 appears in the second row of P_M , then P_M is one of the following:

$$\begin{array}{cccc} \underline{i_2 i_6 i_8} & \underline{i_2 i_5 i_9} & \underline{i_3 i_5 i_8} & \underline{i_2 i_5 i_8} \\ \underline{i_1 i_4 i_7}, & \underline{i_1 i_4 i_7}, & \underline{i_1 i_4 i_7}, & \underline{i_1 i_4 i_7}. \\ \underline{i_3 i_5 i_9} & \underline{i_3 i_6 i_8} & \underline{i_2 i_6 i_9} & \underline{i_3 i_6 i_9} \end{array}$$

If i_1 appears in the third row of P_M , then P_M is one of the following:

$$\begin{array}{cccc} \underline{i_3 i_7 i_9} & \underline{i_2 i_4 i_7} & \underline{i_4 i_6 i_9} & \underline{i_3 i_6 i_9} \\ \underline{i_2 i_5 i_8}, & \underline{i_3 i_6 i_9}, & \underline{i_2 i_5 i_8}, & \underline{i_2 i_5 i_8}. \\ \underline{i_1 i_4 i_6} & \underline{i_1 i_5 i_8} & \underline{i_1 i_3 i_7} & \underline{i_1 i_4 i_7} \end{array}$$

□

Recall that if we add a fixed number to every element of a profile P , reducing modulo n , the result is called a shift of P (Definition 2.6).

Corollary 5.10. *In $\text{CM}(B_{3,9})$, the profile of an indecomposable rank 3 module with imaginary root has to be a shift of one of the following two:*

$$\begin{array}{cc} \underline{157} & \underline{147} \\ \underline{369}, & \underline{369}. \\ 248 & 258 \end{array}$$

We will see in Proposition 6.9 that modules with the second profile are not rigid.

6. AUSLANDER-REITEN QUIVER

The goal of this section is to characterise part of the Grassmannian cluster category $\text{CM}(B_{3,n})$. In particular, we will give all Auslander-Reiten sequences where the end terms are a rank 1 and a rank 2 module respectively. We will use these results to show that there are exactly 225 rigid indecomposable rank 3 modules in the case $n = 9$.

6.1. Profiles of Auslander-Reiten translates. We will need to determine profiles of (inverses of) Auslander-Reiten translates of indecomposable modules. This is done as follows.

Let M be a rigid non-projective indecomposable module in $\text{CM}(B_{k,n})$ with filtration $M = L_{I_1} | L_{I_2} | \dots | L_{I_s}$. Recall that a peak of a rank 1 module L_I is an element $i \notin I$ such that $i + 1 \in I$ (Definition 2.4).

For $j \in [n]$ let $P_j = L_{\{j+1, j+2, \dots, j+k\}}$ be the projective rank 1 module in $\text{CM}(B_{k,n})$ with peak at j .

Remark 6.1. Let $M \in \underline{\text{CM}}(B_{k,n})$ be rigid indecomposable. Recall that $\tau^{-1}(M)$ is also rigid (Remark 2.2). To find the profile of $\tau^{-1}(M)$ we use a strategy to compute the first syzygy of M , which is the same as $\tau^{-1}(M)$. In practice, already finding the minimal projective cover of M is difficult. We will only need to do this in small ranks (ranks 1, 2, 3) where it is still feasible.

Let $\bigoplus_{j \in U} P_j$ be the minimal projective cover of M , where U is a multiset with entries in $\{0, 1, \dots, n-1\}$. We consider the lattice diagrams of $\bigoplus_{j \in U} P_j$ and of M . Then the dimensions in the corresponding lattice diagram of the syzygy of M , i.e. of $\tau^{-1}(M)$, are

given as $\dim \bigoplus_{j \in U} P_j - \dim M$. We will call this infinite tuple the **dimension lattice** of $\tau^{-1}(M)$. Assume that $\tau^{-1}(M)$ has filtration $L_{J_1} | L_{J_2} | \dots | L_{J_t}$.

From the dimension lattice $\dim \bigoplus_{j \in U} P_j - \dim M$ we can find the filtration factors of $\tau^{-1}(M)$ iteratively: Draw the lattice diagram for the dimension lattice of $\tau^{-1}(M)$.

The set J_1 is consisted of the $i \in [n]$ such that the arrow x_i is in the top rim of the lattice diagram. The set J_2 is consisted of the $i \in [n]$ such that x_i is in the top rim of the lattice diagram of $\tau^{-1}(M)/L_{J_1}$, etc.

Example 6.2. For $I = \{i, j, j+1, \dots, j+k-2\}$, Remark 6.1 gives $\tau^{-1}(L_I) = L_J$ with $J = \{i+1, i+2, \dots, i+k-1, j+k-1\}$.

6.2. Auslander-Reiten sequences. Observe that if L_I has m peaks, then by Remark 2.2, $\tau^{-1}(L_I)$ is also a rigid indecomposable module and since the rank is additive on short exact sequences, this module has rank $m-1$. We also recall that in the case where L_I has 2 peaks and where the k -subset I consists of two intervals, one with a single element, Auslander-Reiten sequences with these end terms have been determined in [2, Theorem 3.12]. Here, we give a more general result for Auslander-Reiten sequences in $\text{CM}(B_{3,n})$ with end terms a rank 1 and a rank 2 module.

Theorem 6.3. *Let L_I be a rank 1 module in $\text{CM}(B_{3,n})$, where $I = \{i_1, i_2, i_3\}$ with $i_1 < i_2 < i_3 < i_1$ in the cyclic order is a 3-subset with three peaks. Then the Auslander-Reiten sequence starting at L_I is as follows:*

$$L_I \hookrightarrow M \twoheadrightarrow \frac{L_X}{L_Y},$$

where $X = \{i_1+1, i_2+1, i_3+1\}$ and $Y = \{i_1+2, i_2+2, i_3+2\}$.

If the middle term M is indecomposable, then $P_M = X|I|Y$.

Proof. Let L_I be a rank 1 module in $\text{CM}(B_{3,n})$, where $I = \{i_1, i_2, i_3\}$ is a 3-subset with three peaks. Note that this implies $n \geq 6$. Using Remark 6.1 we find $\tau^{-1}(L_I) = \frac{L_X}{L_Y}$, where $X = \{i_1+1, i_2+1, i_3+1\}$, $Y = \{i_1+2, i_2+2, i_3+2\}$. Therefore, the Auslander-Reiten sequence starting at L_I is $L_I \rightarrow M \rightarrow \frac{L_X}{L_Y}$, where M is a rank 3 module.

We now show that if M is indecomposable, its profile is as claimed. For that, we will study the content of M and use it to find candidates for M .

So suppose that M is indecomposable. This implies $n \geq 8$, since there are no rank 3 indecomposables for $n = 6, 7$. The content $\text{con}(M)$ of M is the union of the contents of the end terms L_I and $L_X|L_Y$, so $\text{con}(M) = \{i_1, i_2, i_3, i_1+1, i_2+1, i_3+1, i_1+2, i_2+2, i_3+2\}$. By the proof of Theorem 5.6, this content consists of 8 or 9 different elements. If these numbers are all different, we have $i_1+2 < i_2$, $i_2+2 < i_3$ and $i_3+2 < i_1$. In this case, $q(M) = 0$, i.e. M corresponds to an imaginary root. By Theorem 5.9, there are 12 possibilities for the profile P_M of M . The projective cover of the sum $L_I \oplus \frac{L_X}{L_Y}$ of the end terms has to contain the projective cover of the middle term as a summand. Comparing with these 12 profiles, we see that only the module with filtration $L_X|L_I|L_Y$ works, as claimed.

If the content of M has exactly 8 different numbers, then we have one of the following situations:

- (i) $i_1 + 2 = i_2$, $i_2 + 2 < i_3$ and $i_3 + 2 < i_1$,
- (ii) $i_1 + 2 < i_2$, $i_2 + 2 = i_3$ and $i_3 + 2 < i_1$,
- (iii) $i_1 + 2 < i_2$, $i_2 + 2 < i_3$ and $i_3 + 2 = i_1$.

In this case, $q(M) = 2$, i.e. M corresponds to a real root of $J_{3,n}$. By Theorem 5.7, the corresponding profile P_M is one of the three cyclic permutations of a canonical one.

In (i), the canonical profile is $\frac{i_2}{i_1+1} \frac{i_2+2}{i_2+1} \frac{i_3+2}{i_3+1}$. Comparing projective covers, as above, we find that P_M is in fact equal to this, which is $X|I|Y$ as claimed. Similarly, for (ii) and (iii), where the profiles of M are the following two:

$$\frac{i_1+1}{i_1+2} \frac{i_2+1}{i_3} \frac{i_3+1}{i_3+2}, \quad \frac{i_1+1}{i_1} \frac{i_2+1}{i_3+2} \frac{i_3+1}{i_3+2}$$

Therefore P_M is $X|I|Y$ as claimed, in all cases. \square

Lemma 6.4. *Let $I = \{i, i+2, i+4\}$ (entries taken modulo n). Let $L_I \hookrightarrow M \rightarrow L_X|L_Y$ be as in Theorem 6.3, with $X = \{i+1, i+3, i+5\}$ and $Y = \{i+2, i+4, i+6\}$. Assume that $M = L_A \oplus N$ with $A = \{i+1, i+2, i+4\}$. The module N is indecomposable if and only if $n \geq 7$ and in this case, $N = L_B|L_Y$ for $B = I \cup X \setminus A = \{i, i+3, i+5\}$.*


Proof. If $n = 6$, the Auslander-Reiten quiver is well-known, see [13, Figure 10] and the only two 3-subsets with three peaks are $\{1, 3, 5\}$ and $\{2, 4, 6\}$. For $I = \{1, 3, 5\}$, we have $X = \{2, 4, 6\}$, $Y = \{1, 3, 5\}$, $A = \{2, 3, 5\}$ and $N = L_{\{1,4,5\}} \oplus L_{\{1,3,6\}}$.

So assume $n \geq 7$. If N is indecomposable, we are in case (A) of Figure 10. For N to be indecomposable, the two 3-subsets of the filtration of N have to be 3-interlacing. Thus the module N must be either $L_B|L_Y$ or $L_Y|L_B$ with $B = \{i, i+3, i+5\}$. Comparing the projective covers shows that $N = L_B|L_Y$.

If $N = C \oplus E$ is a direct sum of two rank 1 modules, the Auslander-Reiten quiver locally is as in (B) of Figure 10 with modules C', C'', E', E'' , and where E', E'' may be trivial modules. We have $\text{con}(C) \cup \text{con}(E) = \{i, i+2, i+3, i+4, i+5, i+6\}$. These six elements are all different since $n \geq 7$. The six entries have to be partitioned into two 3-subsets J_1, J_2 , with $C = L_{J_1}$ and $E = L_{J_2}$ in a way that C and E combined have the four peaks $\{i+1, i+3, i+5, i+6\}$. There are three ways to do so: (i) $\{i, i+2, i+6\} \cup \{i+3, i+4, i+5\}$, (ii) $\{i, i+4, i+5\} \cup \{i+2, i+3, i+6\}$, or (iii) $\{i, i+3, i+4\} \cup \{i+2, i+5, i+6\}$. Only in case (iii), the direct sum $C \oplus E$ has the peaks $\{i, i+2, i+3, i+5\}$ which are needed in order to match the projective covers. So assume

$$C = L_{\{i, i+3, i+4\}} \quad \text{and} \quad E = L_{\{i+2, i+5, i+6\}}.$$

We then find $E'' = \tau^{-1}(E) = L_{\{i+3, i+4, i+7\}}$, cf. Example 6.2. Since the content of $L_X|L_Y$ has to be contained in $\text{con}(E) \cup \text{con}(E'')$, this would imply $i+7 \equiv i+1$ modulo n , i.e. $n = 6$, a contradiction to the assumptions. \square

FIGURE 10. Part of the Auslander-Reiten quiver of $\text{CM}(B_{3,n})$.

We also have dual versions of Theorem 6.3 and of Lemma 6.4. We leave out their proofs as they are analogous to the above.

Theorem 6.5. *Let L_I be a rank 1 module in $\text{CM}(B_{3,n})$, $n \in \mathbb{Z}_{\geq 7}$, where $I = \{i_1, i_2, i_3\}$ ($i_1 < i_2 < i_3 < i_1$ in the cyclic order) is a 3-subset with three peaks. Then the Auslander-Reiten sequence ending at L_I is*

$$\frac{L_X}{L_Y} \hookrightarrow M \twoheadrightarrow L_I,$$

for $X = \{i_1 - 2, i_2 - 2, i_3 - 2\}$ and $Y = \{i_1 - 1, i_2 - 1, i_3 - 1\}$.

If M is indecomposable then $P_M = X|I|Y$.

Lemma 6.6. *In the situation of Theorem 6.5, if $I = \{i, i + 2, i + 4\}$ and $M = L_A \oplus N$ for $A = \{1, i + 2, i + 3\}$, then N is indecomposable if and only if $n \geq 7$ and in this case, $N = L_X|L_B$ for $B = \{i - 1, i + 1, i + 4\} = I \cup Y \setminus A$.*

6.3. Rank 3 rigid indecomposable modules in $\text{CM}(B_{3,9})$. We now characterise the rigid indecomposable rank 3 modules of $\text{CM}(B_{3,9})$.

Theorem 6.7. *Let M be a rigid indecomposable rank 3 module in $\text{CM}(B_{3,9})$. Then either $q(M) = 2$ and M is a cyclic permutation of a real canonical profile, or $q(M) = 0$ and the profile of M is a shift of $\frac{157}{369}$. Furthermore, every cyclic permutation of a real canonical profile and every shift of the above profile yields a rigid indecomposable module.*

There are 72 real canonical profiles of rank 3, listed in Table 1. Thus the following is immediate.

Corollary 6.8. *In $\text{CM}(B_{3,9})$, there are 225 rigid indecomposable rank 3 modules. Out of them, 216 correspond to a real root and 9 to an imaginary root.*

Proposition 6.9. *If M is an indecomposable module and if its profile is a cyclic permutation of the profile $\frac{369}{258}$, then M is not rigid.*

Proof. If M is indecomposable, with the above profile, then $\tau^{-1}(M)$ has the same profile. So M is not rigid by Lemma 2.9. The claim then follows from Lemma 2.7 since any cyclic permutation of one of the above profiles is also equal to a shift of this profile. \square

<u>258</u>	<u>259</u>	<u>258</u>	<u>358</u>	<u>259</u>	<u>259</u>	<u>359</u>	<u>259</u>	<u>358</u>	<u>259</u>	<u>359</u>	<u>359</u>	<u>358</u>	<u>269</u>	<u>259</u>	<u>368</u>	<u>359</u>	<u>359</u>
<u>147</u>	<u>147</u>	<u>247</u>	<u>247</u>	<u>247</u>	<u>148</u>	<u>247</u>	<u>248</u>	<u>247</u>	<u>148</u>	<u>248</u>	<u>247</u>	<u>257</u>	<u>148</u>	<u>248</u>	<u>257</u>	<u>257</u>	<u>248</u>
136	136	136	136	136	136	136	136	146	137	136	146	146	137	137	146	146	146
<u>359</u>	<u>269</u>	<u>269</u>	<u>369</u>	<u>359</u>	<u>369</u>	<u>269</u>	<u>368</u>	<u>359</u>	<u>269</u>	<u>369</u>	<u>369</u>	<u>369</u>	<u>369</u>	<u>369</u>	<u>368</u>	<u>359</u>	<u>269</u>
<u>248</u>	<u>248</u>	<u>158</u>	<u>257</u>	<u>258</u>	<u>248</u>	<u>258</u>	<u>257</u>	<u>248</u>	<u>158</u>	<u>258</u>	<u>258</u>	<u>257</u>	<u>248</u>	<u>158</u>	<u>258</u>	<u>258</u>	<u>258</u>
137	137	137	146	146	137	137	147	147	147	146	137	147	147	147	147	147	147
<u>469</u>	<u>379</u>	<u>369</u>	<u>369</u>	<u>369</u>	<u>369</u>	<u>369</u>	<u>369</u>	<u>469</u>	<u>379</u>	<u>369</u>	<u>469</u>	<u>369</u>	<u>379</u>	<u>369</u>	<u>469</u>	<u>469</u>	<u>469</u>
<u>258</u>	<u>258</u>	<u>358</u>	<u>268</u>	<u>259</u>	<u>258</u>	<u>258</u>	<u>258</u>	<u>358</u>	<u>268</u>	<u>358</u>	<u>258</u>	<u>268</u>	<u>258</u>	<u>259</u>	<u>358</u>	<u>358</u>	<u>268</u>
147	147	147	147	147	247	157	148	147	147	247	157	157	148	148	247	157	157
<u>379</u>	<u>379</u>	<u>379</u>	<u>479</u>	<u>469</u>	<u>379</u>	<u>469</u>	<u>379</u>	<u>479</u>	<u>469</u>	<u>479</u>	<u>379</u>	<u>479</u>	<u>479</u>	<u>479</u>	<u>479</u>	<u>479</u>	<u>479</u>
<u>268</u>	<u>268</u>	<u>259</u>	<u>268</u>	<u>368</u>	<u>269</u>	<u>358</u>	<u>268</u>	<u>368</u>	<u>368</u>	<u>268</u>	<u>269</u>	<u>368</u>	<u>368</u>	<u>269</u>	<u>369</u>	<u>368</u>	<u>369</u>
157	148	148	157	157	148	257	158	157	257	158	158	257	158	158	158	258	258

TABLE 1. The canonical profiles of rank 3 in $C_{3,9}^{re}$.

<u>469</u>	<u>369</u>	<u>379</u>	<u>359</u>	<u>358</u>	<u>469</u>	<u>469</u>	<u>269</u>	<u>359</u>
<u>258</u> ;	<u>258</u> ,	<u>258</u> ,	<u>247</u> ,	<u>257</u> ,	<u>358</u> ,	<u>268</u> ,	<u>158</u> ,	<u>257</u> .
<u>137</u>	<u>137</u>	<u>147</u>	<u>146</u>	<u>469</u>	<u>157</u>	<u>157</u>	<u>137</u>	<u>146</u>

TABLE 2. Candidates for rigid indecomposable rank 3 modules in $CM(B_{3,9})$, cf. Figures 11–14. The first one is imaginary.

Our strategy to prove Theorem 6.7 is as follows: we will show that for each of the predicted profiles there exists a rigid indecomposable module of rank 3. These are the three cyclic permutations of the 72 canonical real profiles (Theorem 5.7) and the nine shifts of the profile of the imaginary root (Corollary 5.10 and Proposition 6.9), i.e. of $\frac{157}{369}$.

The 72 real rank 3 profiles are listed in Table 1. They give 216 candidates for rigid indecomposable rank 3 modules. By Lemma 2.7, it is enough to consider them up to a shift (up to adding a constant number to each entry in the profile). Since the content of every such profile has seven distinct numbers and one number appearing twice, all 9 shifts of a given profile are different. So up to a shift, we only have to consider 24 candidates. These candidates are listed in Tables 2, 3 and 4. The first entry in Table 2 is the representative of the 9 imaginary profiles (which are all shifts of each other).

We will show that each of these candidates appears in the region of rigid objects in a tube of the Auslander-Reiten quiver of $CM(B_{3,9})$.

$$\frac{359}{247}, \frac{369}{248}, \frac{469}{358}, \frac{359}{248}.$$

$$\frac{136}{136}, \frac{137}{137}, \frac{147}{147}, \frac{136}{136}.$$

TABLE 3. Candidates for rigid indecomposable rank 3 modules in $\text{CM}(B_{3,9})$, cf. Figures 15–18.

$$\frac{269}{158}, \frac{369}{158}, \frac{359}{258}, \frac{269}{258}, \frac{469}{358}, \frac{369}{358}, \frac{469}{258}, \frac{258}{147}, \frac{369}{268}, \frac{379}{269}, \frac{368}{257}, \frac{358}{247}.$$

$$\frac{147}{147}, \frac{147}{147}, \frac{146}{146}, \frac{147}{147}, \frac{257}{257}, \frac{147}{147}, \frac{157}{157}, \frac{136}{136}, \frac{157}{157}, \frac{158}{158}, \frac{146}{146}, \frac{136}{136}.$$

TABLE 4. Candidates for rigid indecomposable rank 3 modules in $\text{CM}(B_{3,9})$, cf. Figures 19–28.

6.4. Proof of Theorem 6.7. The category $\text{CM}(B_{3,9})$ is known to be tubular ([2, Section 3.5]): it means that when we study the indecomposable objects and the irreducible maps between them, they form tubes which are bounded on one end. Consider a tube in the Auslander-Reiten quiver of the category $\text{CM}(B_{3,9})$. This has the shape of an infinite cylinder, bounded on one end. In general, on each level (row), it has the same number of vertices which correspond to the indecomposable objects of the category. The only exceptions are the tubes which contain projective-injective indecomposables at the end, such a row only has half of the entries of the other rows in the tube. We will consider the row with projective-injective objects as row 0. Then row 1 is called the **mouth** of the tube. The **rank** of a tube is its width. It is a fact that the indecomposable objects in rows $1, 2, \dots, r-1$ (and in row 0, when present) are all rigid (see e.g. [5]).

So what we will prove is that for every candidate in the Tables 2, 3 and 4 there exists a rank 3 module in the rigid region in a tube of the Auslander-Reiten quiver of $\text{CM}(B_{3,9})$. Note that the rank of a module is additive on short exact sequences. This means that in any tube, the lowest rank modules are at the mouth and that this rank grows when moving away from the mouth.

It will thus be enough to consider the set of tubes which have rank 1 modules in their mouth, the set of tubes which have rigid rank 2 modules in their mouth but no rank 1 modules, and the set of tubes which have rigid rank 3 modules in their mouth but no rank 2 modules. We call these sets of tubes $\mathcal{R}_{\geq 1}$, $\mathcal{R}_{\geq 2}$ and $\mathcal{R}_{\geq 3}$, respectively.

The tubes in the sets $\mathcal{R}_{\geq 1}$ and $\mathcal{R}_{\geq 2}$ have been determined in [2]: the tubes of the subfigures (A), (B), (C) and (D) of Figure 6 in [2] all belong to $\mathcal{R}_{\geq 1}$. There are 81 different rigid indecomposable modules of rank 3 in such tubes.

The tubes of the subfigures (G), (H), (I) and (J) of Figure 6 in [2] belong to $\mathcal{R}_{\geq 2}$; these tubes contain 36 different rigid indecomposable rank 3 modules.

We will give more details here. For $\mathcal{R}_{\geq 1}$, we will give the filtrations of representatives of these rank 3 modules in Figures 11–14 explicitly and explain how they can be computed. For $\mathcal{R}_{\geq 2}$, we give the filtrations of representatives of these rank 3 modules in Figures 15–18.

If we can prove that the tubes in $\mathcal{R}_{\geq 3}$ contain 108 rigid indecomposable rank 3 modules, we are done. Since all rank 3 modules are at the mouths of the tubes of $\mathcal{R}_{\geq 3}$, it is enough to compute the first τ -orbit (the mouth) in each tube of rank > 1 . For this, we first have to determine the modules for the candidates of Tables 2, 3 and 4 in the tubes of $\mathcal{R}_{\geq 1}$ and of $\mathcal{R}_{\geq 2}$: they are the ones remaining after finding the rank 3 modules in $\mathcal{R}_{\geq 1}$ and $\mathcal{R}_{\geq 2}$.

We give all the remaining representatives in Figures 19–28. In total, there are 108 rigid indecomposable rank 3 modules in such tubes, adding up to 225 rigid indecomposable rank 3 modules in $\text{CM}(B_{3,9})$ as claimed and this will finish the proof.

In all the figures, we indicate the representatives of the profiles (for Tables 2, 3, 4) by writing them in blue.

6.5. Tubes with rank 1 modules. There are four types of tubes in $\mathcal{R}_{\geq 1}$, shown in Figures 11–14. We explain how to obtain Figure 11, determining the profiles of the rank 3 modules works similarly in the other cases.

Consider the Auslander-Reiten sequence $M \rightarrow M' \oplus N' \rightarrow N$ in Figure 11 with

$$M = \frac{359}{246}, \quad M' = \frac{135}{246} \quad \text{and} \quad N = \frac{135}{247}.$$

Since the category is tubular, N' is indecomposable, of rank 2. It is rigid as it is in the rigid range in this tube. By Theorem 4.7 (1), the two 3-subsets of N' form three boxes, i.e. are 3-interlacing. So the module N' is either $L_{359}|L_{247}$ or it is $L_{247}|L_{359}$. Comparing the projective covers of $M \oplus N$ and $M' \oplus N'$ shows that $N' = L_{359}|L_{247}$. Then we compute $\tau^{-1}(N')$ and the rest of this row using Remark 6.1.

To find the profiles of the rank 3 modules in row 5, we consider the Auslander-Reiten sequence $M \rightarrow M' \oplus N' \rightarrow N$ for

$$M = \frac{359}{247}, \quad M' = \frac{135}{247} \quad \text{and} \quad N = \frac{136}{479}.$$

Similarly as before, N' is rigid indecomposable, it is of rank 3. The content of N' is $\{2, 3, 4, 5, 6, 7, 8, 9, 9\}$, so $q(N') = 2$, i.e. N' corresponds to a real root. By Theorem 5.7, $P_{N'}$ is a cyclic permutation of a canonical profile. The only possible choices of N' are the following:

$$\frac{369}{258}, \quad \frac{479}{369} \quad \text{and} \quad \frac{258}{479}.$$

Comparing the projective covers, we find that $N' = L_{369}|L_{258}|L_{479}$. Then we use Remark 6.1 to find $\tau^{-1}N'$ and so on.

In Figures 11, 12, 13 and 14, we have chosen representatives of the rigid rank 3 modules up to a shift. They are coloured in blue.

6.6. Rigid rank 3 modules in $\mathcal{R}_{\geq 2}$. In this section, we have the rigid rank 3 modules from tubes in $\mathcal{R}_{\geq 2}$, as shown in Figures 15–18. Part of these tubes appear in [2, Figure 6]. Here, we have determined the rows with rigid rank 3 modules.




FIGURE 11. The tube containing L_{124} .




FIGURE 12. The tube containing L_{256} .




FIGURE 13. The tube containing L_{268} .

6.7. **Rigid rank 3 modules in $\mathcal{R}_{\geq 3}$.** For the tubes in Figures 19–28, we only need to establish the first row, using the strategy from Remark 6.1, starting with the remaining candidates for rank 3 modules from Tables 2, 3 and 4.




FIGURE 14. The tube containing L_{158} .




FIGURE 15. The tube containing $L_{258}|L_{146}$.




FIGURE 16. The tube containing $L_{268}|L_{157}$.




FIGURE 17. The tube containing $L_{368}|L_{257}$.









FIGURE 18. The tube containing $L_{268}|L_{147}$.

FIGURE 19. The tube containing $L_{269}|L_{158}|L_{147}$.FIGURE 20. The tube containing $L_{369}|L_{158}|L_{147}$.FIGURE 21. The tube containing $L_{269}|L_{258}|L_{147}$.FIGURE 22. The tube containing $L_{469}|L_{358}|L_{257}$.FIGURE 23. The tube containing $L_{369}|L_{358}|L_{147}$.




FIGURE 24. The tube containing $L_{258}|L_{147}|L_{136}$.




FIGURE 25. The tube containing $L_{369}|L_{268}|L_{157}$.




FIGURE 26. The tube containing $L_{379}|L_{269}|L_{158}$.




FIGURE 27. The tube containing $L_{368}|L_{257}|L_{146}$.




FIGURE 28. The tube containing $L_{358}|L_{247}|L_{136}$.

REFERENCES

- [1] K. Baur, D. Bogdanic, *Extensions between Cohen–Macaulay modules of Grassmannian cluster categories*, Journal of Algebraic Combinatorics, 1–36, 2016.
- [2] K. Baur, D. Bogdanic, A. Garcia Elsenner, *Cluster categories from Grassmannians and root combinatorics*, Nagoya Mathematical Journal, 1–33, doi:10.1017/nmj.2019.14.
- [3] K. Baur, D. Bogdanic, J.-R. Li, *Construction of rank 2 indecomposable modules for Grassmannian cluster categories*, in preparation.
- [4] K. Baur, A. King, B.R. Marsh (previously R.J. Marsh), *Dimer models and cluster categories of Grassmannian*, Proc. Lond. Math. Soc. (3) **113**(2) (2016), 213–260.
- [5] K. Baur, B.R. Marsh, *A geometric model of tube categories*, J. Algebra **362** (2012), 178–191.
- [6] R. W. Carter, *Lie algebras of finite and affine type*, Cambridge Studies in Advanced Mathematics, **96**, Cambridge University Press, Cambridge, 2005.
- [7] L. Demonet and X. Luo, *Ice quivers with potential associated with triangulations and Cohen–Macaulay modules over orders*, Trans. Amer. Math. Soc., **368** (6) (2016), 4257–4293.
- [8] S. Fomin, A. Zelevinsky, *Cluster algebras II: Finite type classification*, Invent. Math. **154** (2003) 63–121.
- [9] C. Fraser, *Braid group symmetries of Grassmannian cluster algebras*, Selecta Math. (N.S.) **26** (2020), no. 2, Paper No. 17, 51 pp.
- [10] C. Geiss, B. Leclerc, and J. Schröer. Preprojective algebras and cluster algebras. Trends in representation theory of algebras and related topics, (2008) 253–283.
- [11] M. Gekhtman, M. Shapiro, A. Stolin, A. Vainshtein, *Poisson structures compatible with the cluster algebra structure in Grassmannians*, Lett. Math. Phys. **100** (2012), no. 2, 139–150.
- [12] D. Hernandez and B. Leclerc, *Cluster algebras and quantum affine algebras*, Duke Math. J. **154** (2010), no. 2, 265–341.
- [13] B. Jensen, A. King, X.P. Su, *A categorification of Grassmannian cluster algebras*, Proc. Lond. Math. Soc. (3) **113** (2016), no. 2, 185–212.
- [14] V. G. Kac, *Infinite-dimensional Lie algebras*, Third edition, Cambridge University Press, Cambridge, 1990.
- [15] S.-J. Kang, M. Kashiwara, M. Kim, S.-j. Oh, *Monoidal categorification of cluster algebras*, J. Amer. Math. Soc. **31** (2018), no. 2, 349–426.
- [16] B. Keller, *The periodicity conjecture for pairs of Dynkin diagrams*, Ann. of Math. (2), **177**(1) (2013), 111–170.
- [17] B.R. Marsh, K. Rietsch, *The B-model connection and mirror symmetry for Grassmannians*, Adv. Math. **366** (2020), 107027, 131 pp.
- [18] G. Muller, D. Speyer. Cluster algebras of Grassmannians are locally acyclic. *Proceedings of the American Mathematical Society*, **144**(8), (2016)3267–3281.
- [19] J. Scott, *Grassmannians and Cluster Algebras*, Proc. Lond. Math. Soc. (2) **92** (2006), 345–380. doi:10.1112/S0024611505015571
- [20] D. Speyer and L. Williams. The tropical totally positive Grassmannian. *Journal of Algebraic Combinatorics*, **22**(2), (2005) 189–210.