

COMPLEX STRUCTURES ON NILPOTENT LIE ALGEBRAS WITH ONE-DIMENSIONAL CENTER

ADELA LATORRE, LUIS UGARTE, AND RAQUEL VILLACAMPA

ABSTRACT. We classify the nilpotent Lie algebras of real dimension eight and minimal center that admit a complex structure. Furthermore, for every such nilpotent Lie algebra \mathfrak{g} , we describe the space of complex structures on \mathfrak{g} up to isomorphism.

CONTENTS

1. Introduction	1
2. Complex structures on nilpotent Lie algebras	3
3. Classification of $S_n\mathbb{N}$ complex structures on 8-dimensional NLAs	6
3.1. The families I and II	7
3.2. Study of Family I	9
3.2.1. Classification up to equivalence	9
3.2.2. Ascending central series	14
3.3. Study of Family II	15
3.3.1. Classification up to equivalence	15
3.3.2. Ascending central series	20
4. Classification of NLAs underlying Family I	20
4.1. NLAs with $\dim \mathfrak{g}_2 \geq 4$	22
4.2. NLAs with $\dim \mathfrak{g}_2 = 3$ and nilpotency step 3	23
4.3. NLAs with $\dim \mathfrak{g}_2 = 3$ and nilpotency step 4	24
5. Classification of NLAs underlying Family II	33
5.1. 4-step NLAs	33
5.2. 5-step NLAs	34
Appendix A. Real Lie algebras and complex structures in Family I	36
Appendix B. Real Lie algebras and complex structures in Family II	38
Acknowledgments	39
References	39

1. INTRODUCTION

In the last decades, the study of complex nilmanifolds has proved to be very useful in the understanding of several aspects of compact complex manifolds. By a complex nilmanifold we mean a compact quotient of a simply connected nilpotent Lie group G endowed with a *left invariant* complex structure J (namely, defined on the Lie algebra \mathfrak{g} of G) by a cocompact discrete subgroup Γ . Hence, the study of nilpotent Lie algebras \mathfrak{g} with complex structures J constitutes a crucial step in the construction of complex nilmanifolds.

Some recent results where complex nilmanifolds play an important role can be found in [2]–[5], [7], [8], [10]–[13], [22], [25]–[29] and the references therein. These include cohomological aspects of compact complex manifolds (Dolbeault, Bott-Chern, Aeppli cohomologies and Frölicher spectral sequence), existence of different classes of Hermitian metrics (as SKT, locally conformal Kähler or balanced metrics, among others), as well as the behaviour of complex properties under holomorphic deformations. It is

2010 *Mathematics Subject Classification*. Primary 17B30; Secondary 53C30, 53C15.

worth to note that in the previous results the complex structures J on the Lie algebras \mathfrak{g} underlying the nilmanifolds usually satisfy $Z(\mathfrak{g}) \cap J(Z(\mathfrak{g})) \neq \{0\}$, being $Z(\mathfrak{g})$ the center of \mathfrak{g} . These complex structures are known as *quasi-nilpotent* (see Definition 2.1) and include complex-parallelizable as well as abelian complex structures. By [20, Section 2], any \mathfrak{g} with quasi-nilpotent complex structure can be constructed as a certain extension of a lower dimensional nilpotent Lie algebra endowed with a complex structure. Therefore, the ‘essentially new’ complex structures J that arise in each even real dimension are those for which the only J -invariant subspace of the center $Z(\mathfrak{g})$ is the trivial one. These are called *strongly non-nilpotent*, or *SnN* for short, complex structures.

Although the complex geometry of nilmanifolds endowed with SnN complex structures still remains to be studied in general, some partial results have been obtained. For instance, in [19, 21] several families of such complex nilmanifolds are constructed, allowing to prove the existence of infinitely many real homotopy types of nilmanifolds with balanced Hermitian metrics and generalized complex structures of any type, respectively. Moreover, nilmanifolds endowed with SnN complex structures admitting neutral Calabi-Yau metrics with interesting deformation properties are given in [18].

A first step to investigate the properties of nilpotent Lie algebras with SnN complex structure is carried out in [20]. Several algebraic constraints to their existence in terms of the ascending central series of \mathfrak{g} can be found in [20, Section 3]. For instance, the nilpotency step s of the Lie algebra \mathfrak{g} satisfies $s \geq 3$, and the dimension of the center of \mathfrak{g} is bounded by $\dim Z(\mathfrak{g}) \leq n - 3$, where $2n = \dim \mathfrak{g} \geq 8$. Somehow, the existence of SnN complex structures J on \mathfrak{g} seems to require a large nilpotency step and a small center, which gives the idea that these J ’s might be very twisted.

It is known that SnN complex structures do not exist in real dimension ≤ 4 , and that there are only two non-isomorphic 6-dimensional nilpotent Lie algebras admitting such structures: \mathfrak{h}_{19}^- and \mathfrak{h}_{26}^+ in the notation of [30] (see also [6]). However, there is no classification in higher (even) dimensions. For this reason, our goal in this paper is to provide the complete list of 8-dimensional nilpotent Lie algebras \mathfrak{g} that admit SnN complex structures up to isomorphism. We recall that real nilpotent Lie algebras are classified only up to dimension 7. So, our method will consist in two steps: in the first one we will obtain a classification of SnN complex structures up to equivalence; then, we will achieve the classification of real Lie algebras admitting such structures. Notice that, by the previous upper bound for $Z(\mathfrak{g})$, these Lie algebras must have 1-dimensional center.

Let \mathfrak{g} be an 8-dimensional nilpotent Lie algebra, NLA for short, with 1-dimensional center. Suppose that \mathfrak{g} has a complex structure J . Starting from the generic complex structure equations for (\mathfrak{g}, J) given in [20] (see Proposition 2.8), we first collect them into two classes, named Family I and Family II. This allows us to arrive at an appropriate reduction in Section 3 that is suitable for their classification up to equivalence. We recall that similar results were obtained on 6-dimensional nilpotent Lie algebras in the case of abelian complex structures [1] and of any other type of complex structure [6]. Also complex parallelizable structures up to complex dimension seven are classified in [16] (see also [24]). Our first main result in this paper is the following:

Theorem 1.1. (Classification of complex structures) *Let J be a complex structure on an 8-dimensional NLA \mathfrak{g} with 1-dimensional center. Then, there exists a basis of $(1, 0)$ -forms $\{\omega^k\}_{k=1}^4$ in terms of which the complex structure equations of (\mathfrak{g}, J) are one (and only one) of the following:*

$$\bullet \text{ (Family I)} \quad \begin{cases} d\omega^1 = 0, \\ d\omega^2 = \varepsilon \omega^{1\bar{1}}, \\ d\omega^3 = \omega^{14} + \omega^{1\bar{4}} + a\omega^{2\bar{1}} + i\delta\varepsilon b\omega^{1\bar{2}}, \\ d\omega^4 = i\nu\omega^{1\bar{1}} + b\omega^{2\bar{2}} + i\delta(\omega^{13} - \omega^{3\bar{1}}), \end{cases}$$

where $\delta = \pm 1$, $a \in \mathbb{R}^{\geq 0}$, $b \in \mathbb{R}$ with $(a, b) \neq (0, 0)$, and the tuple (ε, ν, a, b) takes the following values:

$$(0, 0, 0, 1), (0, 0, 1, 0), (0, 0, 1, 1), (0, 1, 0, b/|b|), (0, 1, 1, b), (1, 0, 0, 1), (1, 0, 1, |b|) \text{ or } (1, 1, a, b).$$

$$\bullet \text{ (Family II)} \quad \begin{cases} d\omega^1 = 0, \\ d\omega^2 = \omega^{14} + \omega^{1\bar{4}}, \\ d\omega^3 = a\omega^{1\bar{1}} + \varepsilon(\omega^{12} + \omega^{1\bar{2}} - \omega^{2\bar{1}}) + i\mu(\omega^{24} + \omega^{2\bar{4}}), \\ d\omega^4 = i\nu\omega^{1\bar{1}} - \mu\omega^{2\bar{2}} + ib(\omega^{12} - \omega^{2\bar{1}}) + i(\omega^{13} - \omega^{3\bar{1}}), \end{cases}$$

where $a, b \in \mathbb{R}$, and the tuple $(\varepsilon, \mu, \nu, a, b)$ is equal to

$$(1, 1, 0, a, b), (1, 0, 1, a, b), (1, 0, 0, 0, b), (1, 0, 0, 1, b), (0, 1, 0, 0, 0) \text{ or } (0, 1, 0, 1, 0).$$

The proof of this theorem is given in Section 3.2 (for Family I) and Section 3.3 (for Family II). Furthermore, we also provide the relation between the different values of the tuples in Theorem 1.1 and the structure of the ascending central series of the Lie algebra \mathfrak{g} (see Table 1 and Table 2 in Theorem 3.1). This is a crucial step for the classification of 8-dimensional nilpotent Lie algebras with 1-dimensional center admitting a complex structure, which is the second main result of this paper:

Theorem 1.2. (Classification of nilpotent Lie algebras) *Let \mathfrak{g} be an 8-dimensional NLA with 1-dimensional center. Then, \mathfrak{g} has a complex structure if and only if it is isomorphic to one (and only one) in the following list:*

$$\begin{aligned} \mathfrak{g}_1^\gamma &= (0^5, 13 + 15 + 24, 14 - 23 + 25, 16 + 27 + \gamma \cdot 34), \text{ where } \gamma \in \{0, 1\}, \\ \mathfrak{g}_2^\alpha &= (0^4, 12, 13 + 15 + 24, 14 - 23 + 25, 16 + 27 + \alpha \cdot 34), \text{ where } \alpha \in \mathbb{R}, \\ \mathfrak{g}_3^\gamma &= (0^4, 12, 13 + \gamma \cdot 15 + 25, 15 + 24 + \gamma \cdot 25, 16 + 27), \text{ where } \gamma \in \{0, 1\}, \\ \mathfrak{g}_4^{\alpha, \beta} &= (0^4, 12, 15 + (\alpha + 1) \cdot 24, (\alpha - 1) \cdot 14 - 23 + (\beta - 1) \cdot 25, 16 + 27 + 34 - 2 \cdot 45), \\ &\text{where } (\alpha, \beta) \in \mathbb{R}^* \times \mathbb{R}^{>0} \text{ or } \mathbb{R}^{>0} \times \{0\}, \\ \mathfrak{g}_5 &= (0^4, 2 \cdot 12, 14 - 23, 13 + 24, 16 + 27 + 35), \\ \mathfrak{g}_6 &= (0^4, 2 \cdot 12, 14 + 15 - 23, 13 + 24 + 25, 16 + 27 + 35), \\ \mathfrak{g}_7 &= (0^5, 15, 25, 16 + 27 + 34), \\ \mathfrak{g}_8 &= (0^4, 12, 15, 25, 16 + 27 + 34), \\ \mathfrak{g}_9^\gamma &= (0^3, 13, 23, 35, \gamma \cdot 12 - 34, 16 + 27 + 45), \text{ where } \gamma \in \{0, 1\}, \\ \mathfrak{g}_{10}^\gamma &= (0^3, 13, 23, 14 + 25, 15 + 24, 16 + \gamma \cdot 25 + 27), \text{ where } \gamma \in \{0, 1\}, \\ \mathfrak{g}_{11}^{\alpha, \beta} &= (0^3, 13, 23, 14 + 25 - 35, \alpha \cdot 12 + 15 + 24 + 34, 16 + 27 - 45 - \beta(2 \cdot 25 + 35)), \\ &\text{where } (\alpha, \beta) = (0, 0), (1, 0) \text{ or } (\alpha \geq 0, 1), \\ \mathfrak{g}_{12}^\gamma &= (0^2, 12, 13, 23, 14 + 25, 15 + 24, 16 + 27 + \gamma \cdot 25), \text{ where } \gamma \in \{0, 1\}. \end{aligned}$$

For the description of the nilpotent Lie algebras in Theorem 1.2 we use the standard abbreviated notation (see Notation 4.3 for details). We note that the first eight families of Lie algebras, i.e. $\mathfrak{g}_1^\gamma, \dots, \mathfrak{g}_8$, are those having complex structures in Family I, whereas the complex structures on $\mathfrak{g}_9^\gamma, \dots, \mathfrak{g}_{12}^\gamma$ belong to Family II. The list above is ordered according to the ascending type of the nilpotent Lie algebras. The proof of Theorem 1.2 can be found in Section 4 and Section 5 (see Theorem 4.1 for the Lie algebras underlying Family I and Theorem 4.1 for those underlying Family II).

The precise relation between Theorem 1.1 and Theorem 1.2 is explained in the Appendices A and B. Given any set of complex structure equations belonging to Family I or Family II in Theorem 1.1, we show in Appendix A and Appendix B, respectively, which is the corresponding underlying real nilpotent Lie algebra in Theorem 1.2. Conversely, given a real nilpotent Lie algebra \mathfrak{g} in Theorem 1.2, the information provided in the appendices allows to construct the space of complex structures on \mathfrak{g} up to equivalence.

2. COMPLEX STRUCTURES ON NILPOTENT LIE ALGEBRAS

In this section we recall some results about real nilpotent Lie algebras (NLA for short) endowed with complex structures, paying special attention to real dimension 8. In particular, we summarize our previous results in [20].

Let \mathfrak{g} be a real Lie algebra of dimension $2n$. Its *ascending central series* $\{\mathfrak{g}_k\}_{k \geq 0}$ is given by $\mathfrak{g}_0 = \{0\}$, and

$$\mathfrak{g}_k = \{X \in \mathfrak{g} \mid [X, \mathfrak{g}] \subseteq \mathfrak{g}_{k-1}\},$$

for any $k \geq 1$. In particular, $\mathfrak{g}_1 = Z(\mathfrak{g})$ is the center of \mathfrak{g} . A Lie algebra \mathfrak{g} is said to be *nilpotent* if there is an integer $s \geq 1$ such that $\mathfrak{g}_k = \mathfrak{g}$, for every $k \geq s$. In such case, the smallest integer s satisfying the previous condition is called *nilpotency step* of \mathfrak{g} , and the Lie algebra is said to be *s-step nilpotent*. Thus, any nilpotent Lie algebra \mathfrak{g} has an associated s -tuple

$$(m_1, \dots, m_{s-1}, m_s) := (\dim \mathfrak{g}_1, \dots, \dim \mathfrak{g}_{s-1}, \dim \mathfrak{g}_s)$$

which strictly increases, i.e. $0 < m_1 < \dots < m_{s-1} < m_s = 2n$. We will say that (m_1, \dots, m_s) is the *ascending type* of \mathfrak{g} .

Obviously, NLAs with different ascending type are non-isomorphic. However, the converse is only true up to real dimension 4. Indeed, there are three non-isomorphic 4-dimensional NLAs whose ascending types are (4), (2, 4), and (1, 2, 4). In contrast, there exist four non-isomorphic 6-dimensional NLAs with the same ascending type (2, 6), for instance.

Let J be a complex structure on an NLA \mathfrak{g} , that is, an endomorphism $J: \mathfrak{g} \rightarrow \mathfrak{g}$ fulfilling $J^2 = -\text{Id}$ and the integrability condition

$$(1) \quad N_J(X, Y) := [X, Y] + J[JX, Y] + J[X, JY] - [JX, JY] = 0,$$

for all $X, Y \in \mathfrak{g}$. Observe that the terms \mathfrak{g}_k in the ascending central series may not be invariant under J . For this reason, a new series $\{\mathfrak{a}_k(J)\}_k$ adapted to the complex structure J is introduced in [9]:

$$\begin{cases} \mathfrak{a}_0(J) = \{0\}, \text{ and} \\ \mathfrak{a}_k(J) = \{X \in \mathfrak{g} \mid [X, \mathfrak{g}] \subseteq \mathfrak{a}_{k-1}(J) \text{ and } [JX, \mathfrak{g}] \subseteq \mathfrak{a}_{k-1}(J)\}, \text{ for } k \geq 1. \end{cases}$$

This series $\{\mathfrak{a}_k(J)\}_k$ is called the *ascending J -compatible series* of \mathfrak{g} . Observe that every $\mathfrak{a}_k(J) \subseteq \mathfrak{g}_k$ is an even-dimensional J -invariant ideal of \mathfrak{g} , and $\mathfrak{a}_1(J)$ is indeed the largest subspace of the center \mathfrak{g}_1 which is J -invariant.

Depending on the behaviour of the series $\{\mathfrak{a}_k(J)\}_k$, complex structures on NLAs can be classified into different types:

Definition 2.1. [9, 20] A complex structure J on a nilpotent Lie algebra \mathfrak{g} is said to be

- (i) *strongly non-nilpotent*, or *SnN* for short, if $\mathfrak{a}_1(J) = \{0\}$;
- (ii) *quasi-nilpotent*, if $\mathfrak{a}_1(J) \neq \{0\}$; moreover, J is called
 - (ii.1) *nilpotent*, if there exists an integer $t > 0$ such that $\mathfrak{a}_t(J) = \mathfrak{g}$,
 - (ii.2) *weakly non-nilpotent*, if there is an integer $t > 0$ satisfying $\mathfrak{a}_l(J) = \mathfrak{a}_t(J)$, for every $l \geq t$, and $\mathfrak{a}_t(J) \neq \mathfrak{g}$.

Remark 2.2. Some algebraic constraints to the existence of complex structures J on \mathfrak{g} are studied in [23] in terms of the *descending* central series of \mathfrak{g} . These constraints imply some estimates on the nilpotency step s ; in particular, \mathfrak{g} cannot be filiform [17]. We also recall that quasi-filiform Lie algebras of dimension ≥ 8 do not admit any complex structure [15]. Hence, $s \leq 2n - 3$ for any nilpotent Lie algebra \mathfrak{g} of dimension $2n \geq 8$ endowed with a complex structure.

Remark 2.3. In the recent paper [14], the structure of Lie algebras endowed with a *maximal nilpotent J* is studied. Such J 's are defined as those nilpotent complex structures for which $t = n$, where t is the smallest integer such that $\mathfrak{a}_t(J) = \mathfrak{g}$ in the ascending J -compatible series of \mathfrak{g} .

One can see that quasi-nilpotent complex structures on NLAs of a given dimension can be constructed from other complex structures defined on (strictly) lower dimensional NLAs (see [20, Section 2] for details). Therefore, the essentially new complex structures that arise in each even real dimension are those of strongly non-nilpotent type. That is to say, SnN complex structures constitute the fundamental piece to fully understand complex geometry on nilpotent Lie algebras.

In real dimension 4 it is well known that SnN complex structures do not exist, whereas in dimension 6 one has the following result:

Theorem 2.4. [6] *Let \mathfrak{g} be an NLA of real dimension 6. If \mathfrak{g} admits an SnN complex structure, then its ascending type is $(\dim \mathfrak{g}_k)_k = (1, 3, 6)$ or $(1, 3, 4, 6)$.*

In fact, all the pairs (\mathfrak{g}, J) with $\dim \mathfrak{g} = 6$ and $\mathfrak{a}_1(J) = \{0\}$ have been classified by means of their complex structure equations [6]. It should be noted that only two NLAs of this dimension admit SnN complex structures, namely, \mathfrak{h}_{19}^- and \mathfrak{h}_{26}^+ in the notation of [30].

Concerning higher dimensions, [20] provides several general restrictions on the terms of the ascending central series of NLAs admitting SnN complex structures. Among them, we highlight the following one:

Theorem 2.5. [20, Theorem 3.11] *Let (\mathfrak{g}, J) be a $2n$ -dimensional nilpotent Lie algebra, with $n \geq 4$, endowed with a strongly non-nilpotent complex structure J . Then, $1 \leq \dim \mathfrak{g}_1 \leq n - 3$.*

From Definition 2.1 one can clearly deduce that any complex structure on an NLA with 1-dimensional center has SnN type. Thanks to Theorem 2.5, the converse also holds in eight dimensions:

Corollary 2.6. *Let \mathfrak{g} be an 8-dimensional NLA admitting a complex structure J . The following properties are equivalent:*

- the center of \mathfrak{g} has dimension 1;
- the complex structure J is strongly non-nilpotent.

Moreover, a structural result in the spirit of Theorem 2.4 is proved in [20] for dimension 8:

Theorem 2.7. [20, Theorem 4.1] *Let \mathfrak{g} be an NLA of real dimension 8. If \mathfrak{g} admits an SnN complex structure, then its ascending type is $(\dim \mathfrak{g}_k)_k = (1, 3, 8)$, $(1, 3, 5, 8)$, $(1, 3, 6, 8)$, $(1, 3, 5, 6, 8)$, $(1, 4, 8)$, $(1, 4, 6, 8)$, $(1, 5, 8)$, or $(1, 5, 6, 8)$.*

Furthermore, the complex structure equations for all the pairs (\mathfrak{g}, J) with $\dim \mathfrak{g} = 8$ and $\mathfrak{a}_1(J) = \{0\}$ are parametrized in [20]. We summarize these results in the proposition below. However, we first need to recall some basic concepts. Let $\mathfrak{g}_{\mathbb{C}}^*$ denote the dual of the complexification $\mathfrak{g}_{\mathbb{C}}$ of \mathfrak{g} . Then, there is a natural bigraduation induced on $\bigwedge^* \mathfrak{g}_{\mathbb{C}}^* = \bigoplus_{p,q} \bigwedge_J^{p,q}(\mathfrak{g}^*)$, where the spaces $\bigwedge_J^{1,0}(\mathfrak{g}^*)$ and $\bigwedge_J^{0,1}(\mathfrak{g}^*)$ are, respectively, the eigenspaces of the eigenvalues $\pm i$ of J as an endomorphism of $\mathfrak{g}_{\mathbb{C}}^*$. For simplicity, we will simply denote each $\bigwedge_J^{p,q}(\mathfrak{g}^*)$ by $\mathfrak{g}_J^{p,q}$. Since J is a complex structure, if $d: \bigwedge^* \mathfrak{g}_{\mathbb{C}}^* \rightarrow \bigwedge^{*+1} \mathfrak{g}_{\mathbb{C}}^*$ is the extension to the complexified exterior algebra of the usual Chevalley-Eilenberg differential, we have that $\pi_{0,2} \circ d|_{\mathfrak{g}_J^{1,0}} \equiv 0$, being $\pi_{0,2}: \bigwedge^2 \mathfrak{g}_{\mathbb{C}}^* \rightarrow \mathfrak{g}_J^{0,2}$ the canonical projection. Given any basis $\{\omega^k\}_{k=1}^4$ of $\mathfrak{g}_J^{1,0}$, from now on we will denote by ω^{jk} , resp. $\omega^{j\bar{k}}$, the wedge product $\omega^j \wedge \omega^k$, resp. $\omega^j \wedge \omega^{\bar{k}}$, where $\omega^{\bar{k}}$ indicates the complex conjugate of ω^k .

Proposition 2.8. [20, Propositions 4.12, 4.13, and 4.14] *Let J be an SnN complex structure on an 8-dimensional NLA \mathfrak{g} . Then:*

- If $\dim \mathfrak{g}_2 = 3$, then the pair (\mathfrak{g}, J) is parametrized by equations of the form

$$(2) \quad \begin{cases} d\omega^1 = 0, \\ d\omega^2 = A\omega^{1\bar{1}} - B(\omega^{14} - \omega^{1\bar{4}}), \\ d\omega^3 = C\omega^{12} - E(\omega^{14} - \omega^{1\bar{4}}) + F\omega^{1\bar{1}} + D\omega^{1\bar{2}} - H(\omega^{24} - \omega^{2\bar{4}}) + G\omega^{2\bar{1}} + K\omega^{2\bar{2}}, \\ d\omega^4 = L\omega^{1\bar{1}} + M\omega^{1\bar{2}} + N\omega^{1\bar{3}} - \bar{M}\omega^{2\bar{1}} + i s \omega^{2\bar{2}} + P\omega^{2\bar{3}} - \bar{N}\omega^{3\bar{1}} - \bar{P}\omega^{3\bar{2}}, \end{cases}$$

where the coefficients are complex numbers with $s \in \mathbb{R}$ that must satisfy the Jacobi identity. Moreover:

- (i) if $(\dim \mathfrak{g}_k)_k = (1, 3, 8)$, then one can take $A = B = 0$ and $\Re L = 0$;
- (ii) if $(\dim \mathfrak{g}_k)_k = (1, 3, 6, 8)$, it is possible to assume $B = H = K = P = 0$;
- (iii) if $(\dim \mathfrak{g}_k)_k = (1, 3, 5, 8)$, we can consider $K = P = 0$ and $\Re L = 0$;
- (iv) if $(\dim \mathfrak{g}_k)_k = (1, 3, 5, 6, 8)$, then $H = K = P = s = 0$ and $\Re L \neq 0$.

- If $\dim \mathfrak{g}_2 = 4$, then the pair (\mathfrak{g}, J) is parametrized by equations of the form

$$(3) \quad \begin{cases} d\omega^1 = 0, \\ d\omega^2 = A\omega^{1\bar{1}}, \\ d\omega^3 = -D(\omega^{12} - \omega^{1\bar{2}}) - E(\omega^{14} - \omega^{1\bar{4}}) + F\omega^{1\bar{1}}, \\ d\omega^4 = L\omega^{1\bar{1}} + M\omega^{1\bar{2}} + N\omega^{1\bar{3}} - \bar{M}\omega^{2\bar{1}} + i s \omega^{2\bar{2}} - \bar{N}\omega^{3\bar{1}}, \end{cases}$$

where the coefficients are complex numbers with $s \in \mathbb{R}$ that must satisfy the Jacobi identity. Furthermore:

- (i) if $(\dim \mathfrak{g}_k)_k = (1, 4, 8)$, we can take $(\Re A, \Re L) = (0, 0)$;
- (ii) if $(\dim \mathfrak{g}_k)_k = (1, 4, 6, 8)$, then $(\Re A, \Re L) \neq (0, 0)$.

- If $\dim \mathfrak{g}_2 = 5$, then the pair (\mathfrak{g}, J) is parametrized by equations of the form

$$(4) \quad \begin{cases} d\omega^1 = 0, \\ d\omega^2 = A\omega^{1\bar{1}} - B(\omega^{14} - \omega^{1\bar{4}}), \\ d\omega^3 = F\omega^{1\bar{1}} - E(\omega^{14} - \omega^{1\bar{4}}), \\ d\omega^4 = L\omega^{1\bar{1}} + M\omega^{1\bar{2}} + N\omega^{1\bar{3}} - \bar{M}\omega^{2\bar{1}} + i s\omega^{2\bar{2}} + P\omega^{2\bar{3}} - \bar{N}\omega^{3\bar{1}} - \bar{P}\omega^{3\bar{2}} + i t\omega^{3\bar{3}}, \end{cases}$$

where the coefficients are complex numbers with $s, t \in \mathbb{R}$ that must satisfy the Jacobi identity. Moreover:

- (i) $(\dim \mathfrak{g}_k)_k = (1, 5, 8)$ gives $\Re L = 0$;
- (ii) $(\dim \mathfrak{g}_k)_k = (1, 5, 6, 8)$ requires $\Re L \neq 0$.

3. CLASSIFICATION OF SNN COMPLEX STRUCTURES ON 8-DIMENSIONAL NLAs

In this section, we classify SnN complex structures on 8-dimensional NLAs up to equivalence, refining equations (2), (3) and (4).

Let (\mathfrak{g}, J) and (\mathfrak{g}', J') be Lie algebras, each endowed with a complex structure. They are said to be *equivalent* if there is an isomorphism of Lie algebras $F: \mathfrak{g}' \rightarrow \mathfrak{g}$ such that $J' = F^{-1} \circ J \circ F$. That is, if there exists a \mathbb{C} -linear isomorphism $F^*: \mathfrak{g}_J^{1,0} \rightarrow \mathfrak{g}_{J'}^{1,0}$ such that $d_{\mathfrak{g}'} \circ F^* = F^* \circ d_{\mathfrak{g}}$, where $d_{\mathfrak{g}}$ and $d_{\mathfrak{g}'}$ are the (extended) Chevalley-Eilenberg differentials of \mathfrak{g} and \mathfrak{g}' , respectively. When there is no risk of confusion, we will denote by d the differentials $d_{\mathfrak{g}}$ and $d_{\mathfrak{g}'}$.

The main goal of this section is to prove the classification result below. Recall that by Corollary 2.6, a complex structure on an 8-dimensional NLA is SnN if and only if the center of the NLA is 1-dimensional.

Theorem 3.1. *Let J be a complex structure on an 8-dimensional NLA \mathfrak{g} with 1-dimensional center. Then, there exists a basis of $(1, 0)$ -forms $\{\omega^k\}_{k=1}^4$ in terms of which the complex structure equations of (\mathfrak{g}, J) are, up to equivalence, one of the following:*

- (i) if the ascending type of \mathfrak{g} is $(\dim \mathfrak{g}_k)_k = (1, 3, 8)$, $(1, 3, 6, 8)$, $(1, 4, 8)$, $(1, 4, 6, 8)$, $(1, 5, 8)$, or $(1, 5, 6, 8)$, then

$$(5) \quad \text{Family I: } \begin{cases} d\omega^1 = 0, \\ d\omega^2 = \varepsilon\omega^{1\bar{1}}, \\ d\omega^3 = \omega^{14} + \omega^{1\bar{4}} + a\omega^{2\bar{1}} + i\delta\varepsilon b\omega^{1\bar{2}}, \\ d\omega^4 = i\nu\omega^{1\bar{1}} + b\omega^{2\bar{2}} + i\delta(\omega^{1\bar{3}} - \omega^{3\bar{1}}), \end{cases}$$

where $\delta = \pm 1$, $\varepsilon, \nu \in \{0, 1\}$, $a \in \mathbb{R}^{\geq 0}$, $b \in \mathbb{R}$ with $(a, b) \neq (0, 0)$, and the parameters are related to the ascending type of \mathfrak{g} by Table 1.

- (ii) if the ascending type is $(\dim \mathfrak{g}_k)_k = (1, 3, 5, 8)$ or $(1, 3, 5, 6, 8)$, then

$$(6) \quad \text{Family II: } \begin{cases} d\omega^1 = 0, \\ d\omega^2 = \omega^{14} + \omega^{1\bar{4}}, \\ d\omega^3 = a\omega^{1\bar{1}} + \varepsilon(\omega^{12} + \omega^{1\bar{2}} - \omega^{2\bar{1}}) + i\mu(\omega^{24} + \omega^{2\bar{4}}), \\ d\omega^4 = i\nu\omega^{1\bar{1}} - \mu\omega^{2\bar{2}} + i b(\omega^{1\bar{2}} - \omega^{2\bar{1}}) + i(\omega^{1\bar{3}} - \omega^{3\bar{1}}), \end{cases}$$

where $a, b \in \mathbb{R}$, $\varepsilon, \mu, \nu \in \{0, 1\}$ with $\mu\nu = 0$ and $(\varepsilon, \mu) \neq (0, 0)$, and the parameters are related to the ascending type of \mathfrak{g} by Table 2.

The rest of this section is devoted to the proof of this theorem. We start with the complex structure equations (2), (3), and (4) given in Proposition 2.8. After a preliminary analysis, we reduce our classification problem to the study of two sets of structure equations, called Family I and Family II (see Section 3.1). Then, we independently study the two families, simplifying each of them and reaching (5)

$(\dim \mathfrak{g}_k)_k$	ε	ν	a	b	δ
(1, 3, 8)	0	0	1	0	± 1
				1	
(1, 3, 6, 8)	1	0	1	$b \geq 0$	± 1
	0	1		$b \in \mathbb{R}$	
	1	1	$a > 0$		
(1, 4, 8)	1	1	0	2δ	± 1
(1, 4, 6, 8)	1	0	0	1	± 1
		1		$b \in \mathbb{R} - \{0, 2\delta\}$	
(1, 5, 8)	0	0	0	1	± 1
(1, 5, 6, 8)	0	1	0	-1	± 1
				1	

TABLE 1. Complex structures in Family I up to equivalence

$(\dim \mathfrak{g}_k)_k$	ε	μ	ν	a	b
(1, 3, 5, 8)	0	1	0	0, 1	0
	1	0	0		$b \in \mathbb{R}$
	1	1	0	$a \in \mathbb{R}$	
(1, 3, 5, 6, 8)	1	0	1	$a \in \mathbb{R}$	$b \in \mathbb{R}$

TABLE 2. Complex structures in Family II up to equivalence

and (6) in Theorem 3.1, and provide their classification up to equivalence, that we have summarized in Tables 1 and 2 (Sections 3.2 and 3.3).

3.1. The families I and II. In order to get the reduced complex structure equations (5) and (6), let us first observe that we can gather (2), (3), and (4) in a more general expression, namely:

$$(7) \quad \begin{cases} d\omega^1 = 0, \\ d\omega^2 = A\omega^{1\bar{1}} - B(\omega^{14} - \omega^{1\bar{4}}), \\ d\omega^3 = F\omega^{1\bar{1}} + K\omega^{2\bar{2}} + C\omega^{12} + D\omega^{1\bar{2}} + G\omega^{2\bar{1}} - E(\omega^{14} - \omega^{1\bar{4}}) - H(\omega^{24} - \omega^{2\bar{4}}), \\ d\omega^4 = L\omega^{1\bar{1}} + is\omega^{2\bar{2}} + it\omega^{3\bar{3}} + (M\omega^{1\bar{2}} - \bar{M}\omega^{2\bar{1}}) + (N\omega^{1\bar{3}} - \bar{N}\omega^{3\bar{1}}) + (P\omega^{2\bar{3}} - \bar{P}\omega^{3\bar{2}}). \end{cases}$$

Indeed, by Proposition 2.8 any SnN complex structure J on an 8-dimensional NLA \mathfrak{g} is covered by these equations. Here, the coefficients $A, B, C, D, E, F, G, H, K, L, M, N, P$ are complex numbers and $s, t \in \mathbb{R}$, in such a way that they satisfy the Jacobi identity and the condition $\dim \mathfrak{g}_1 = 1$ (imposed by the existence of an SnN complex structure).

The Jacobi identity, i.e. $d^2\omega^k = 0$ for $1 \leq k \leq 4$, is equivalent to the following equations:

$$(8) \quad \begin{aligned} AH - BG + \bar{B}D &= 0, & AK = BK &= 0, & tH = tK = tC &= 0, \\ K\bar{N} - P\bar{C} - \bar{P}G &= 0, & H\Re L &= 0, & tD = tG &= 0, \\ isA - F\bar{P} - N\bar{C} + \bar{N}D &= 0, & \Re(P\bar{H}) &= 0, & itE + BP &= 0, \\ isB - E\bar{P} - N\bar{H} &= 0, & \Re(M\bar{B} + N\bar{E}) &= 0, & itF + AP &= 0. \end{aligned}$$

For the condition on the center, let us denote by $\{Z_k\}_{k=1}^4$ the dual basis to $\{\omega^k\}_{k=1}^4$. Using the well-known formula $d\alpha(X, Y) = -\alpha([X, Y])$, for any $\alpha \in \mathfrak{g}^*$ and $X, Y \in \mathfrak{g}$, and its extension to the

complexification, it is easy to check from (7) that

$$[X, Z_4 + \bar{Z}_4] = 0$$

for any $X \in \mathfrak{g}$. Since the center of \mathfrak{g} is 1-dimensional, necessarily $\mathfrak{g}_1 = \langle \Re Z_4 \rangle$. Furthermore, it is clear from equations (7) that the vanishing of the tuples (B, E, H) , (N, P, t) , or (C, D, G, H, K, M, P, s) , imply $\Im Z_4 \in \mathfrak{g}_1$, $\langle \Re Z_3, \Im Z_3 \rangle \subset \mathfrak{g}_1$, or $\langle \Re Z_2, \Im Z_2 \rangle \subset \mathfrak{g}_1$, respectively, which would give a contradiction to $\dim \mathfrak{g}_1 = 1$. Hence, the following conditions must be satisfied:

$$(9) \quad (B, E, H) \neq (0, 0, 0), \quad (N, P, t) \neq (0, 0, 0), \quad (C, D, G, H, K, M, P, s) \neq (0, \dots, 0).$$

We will study (7) bearing in mind the conditions (8) and (9). As noticed above, for the classification up to equivalence, we can consider \mathbb{C} -linear isomorphisms $F^* : \mathfrak{g}_J^{1,0} \rightarrow \mathfrak{g}'_{J'}^{1,0}$ commuting with the differentials, i.e. $d \circ F^* = F^* \circ d$. When an equivalence exists, we will construct such an isomorphism by means of an appropriate (and explicit) change of $(1, 0)$ -basis.

Lemma 3.2. *In the equations (7) above, one can suppose that $t = 0$.*

Proof. We will show that if $t \neq 0$, then there is another $(1, 0)$ -basis $\{\tau^k\}_{k=1}^4$ satisfying equations of the same form as (7) in which the corresponding coefficient t_τ is zero.

Let us then suppose that $t \neq 0$ in equations (7). By (8), we get $C = D = G = H = K = 0$. Hence, conditions (8) and (9) reduce to

$$(10) \quad \begin{aligned} isA - F\bar{P} &= 0, & itE + BP &= 0, & \Re(M\bar{B} + N\bar{E}) &= 0, & (N, P, t) &\neq (0, 0, 0), \\ isB - E\bar{P} &= 0, & itF + AP &= 0, & (B, E) &\neq (0, 0), & (M, P, s) &\neq (0, 0, 0). \end{aligned}$$

We first observe that $B \neq 0$, as otherwise the Jacobi condition $itE = 0$ would imply $E = 0$, which gives a contradiction to $(B, E) \neq (0, 0)$. Bearing this in mind, we now consider two cases.

On the one hand, if $E = 0$ then (10) implies $F = P = s = 0$. Taking $\tau^k = \omega^k$ for $k = 1, 4$, $\tau^2 = \omega^3$ and $\tau^3 = \omega^2$, we directly get equations of the form (7) with $t_\tau = 0$.

On the other hand, for $E \neq 0$ we consider the $(1, 0)$ -basis $\{\tau^k\}_{k=1}^4$ defined by

$$\tau^k = \omega^k, \quad k = 1, 3, 4, \quad \tau^2 = E\omega^2 - B\omega^3.$$

Then, the structure equations in terms of $\{\tau^k\}_{k=1}^4$ are again of the form (7). Using (10) it can be directly seen that the coefficient t_τ of $\tau^{3\bar{3}}$ in the equation $d\tau^4$ satisfies

$$t_\tau = is \frac{|B|^2}{|E|^2} + it + \frac{BP}{E} - \frac{\bar{B}\bar{P}}{\bar{E}} = \frac{\bar{B}}{|E|^2}(isB - E\bar{P}) + \frac{1}{E}(itE + BP) = 0.$$

□

Lemma 3.3. *In the equations (7), in addition to $t = 0$, we can also set $P = K = 0$.*

Proof. By Lemma 3.2 we can assume $t = 0$ in (7). Let us suppose that $P \neq 0$ and show the existence of a $(1, 0)$ -basis $\{\tau^k\}_{k=1}^4$ satisfying equations of the form (7) for which the corresponding coefficient P_τ is zero, in addition to $t_\tau = 0$.

If we take into account the value $t = 0$ and the choice $P \neq 0$ in (8), we immediately get $A = B = 0$. Hence, (8) and (9) are simplified to the following conditions:

$$(11) \quad \begin{aligned} K\bar{N} - P\bar{C} - \bar{P}G &= 0, & H\Re L &= 0, & (E, H) &\neq (0, 0), \\ F\bar{P} + N\bar{C} - \bar{N}D &= 0, & \Re(P\bar{H}) &= 0, & (N, P) &\neq (0, 0), \\ E\bar{P} + N\bar{H} &= 0, & \Re(N\bar{E}) &= 0, & (C, D, G, H, K, M, P, s) &\neq (0, \dots, 0). \end{aligned}$$

Since $E = -N\bar{H}/\bar{P}$, the condition $(E, H) \neq (0, 0)$ in (11) implies $H \neq 0$. In turn, this gives $\Re L = 0$, again by (11). We consider the $(1, 0)$ -basis $\{\tau^k\}_{k=1}^4$ defined by

$$\tau^1 = N\omega^1 + P\omega^2, \quad \tau^2 = \omega^1, \quad \tau^k = \omega^k, \quad k = 3, 4.$$

Using $\Re L = 0$, a direct calculation shows that the structure equations in terms of $\{\tau^k\}_{k=1}^4$ are again of the form (7), with coefficients $t_\tau = 0$ and $P_\tau = 0$ in the equation $d\tau^4$.

Finally, now that we have $P = t = 0$, it suffices to use the second equation of (8) to obtain $K = 0$, as $N \neq 0$ by (9). □

Taking into account Lemmas 3.2 and 3.3, we have:

Proposition 3.4. *Let J be a strongly non-nilpotent complex structure on an 8-dimensional NLA \mathfrak{g} . Then, there exists a basis of $(1,0)$ -forms $\{\omega^k\}_{k=1}^4$ in terms of which the complex structure equations of (\mathfrak{g}, J) are of the form:*

$$(12) \quad \begin{cases} d\omega^1 = 0, \\ d\omega^2 = A\omega^{1\bar{1}} - B(\omega^{14} - \omega^{1\bar{4}}), \\ d\omega^3 = F\omega^{1\bar{1}} + C\omega^{12} + D\omega^{1\bar{2}} + G\omega^{2\bar{1}} - E(\omega^{14} - \omega^{1\bar{4}}) - H(\omega^{24} - \omega^{2\bar{4}}), \\ d\omega^4 = L\omega^{1\bar{1}} + is\omega^{2\bar{2}} + (M\omega^{12} - \bar{M}\omega^{2\bar{1}}) + (N\omega^{13} - \bar{N}\omega^{3\bar{1}}), \end{cases}$$

where the coefficients $A, B, C, D, E, F, G, H, L, M, N$ are complex numbers and $s \in \mathbb{R}$, in such a way that they fulfill the following conditions:

$$(13) \quad \begin{aligned} AH - BG + \bar{B}D = 0, \quad isB - N\bar{H} = 0, \quad H\Re L = 0, \quad (B, E, H) \neq (0, 0, 0), \\ \Re(M\bar{B} + N\bar{E}) = 0, \quad isA - N\bar{C} + \bar{N}D = 0, \quad N \neq 0, \quad (C, D, G, H, M, s) \neq (0, \dots, 0). \end{aligned}$$

From now on, we will divide our study depending on the vanishing of the coefficient B . Indeed, in the next sections we will see that this choice gives a partition of the set of complex structures into two disjoint families. To simplify the procedure, we introduce the following terminology:

Definition 3.5. An SnN complex structure J belongs to *Family I*, resp. *Family II*, if there exists a $(1,0)$ -basis satisfying the equations (12) with $B = 0$, resp. $B \neq 0$.

In Sections 3.2 and 3.3 we separately study each one of these families in order to prove parts (i) and (ii) of Theorem 3.1.

3.2. Study of Family I. We here accomplish the study up to equivalence of those complex structures belonging to Family I. More precisely, in Section 3.2.1 we first prove that equations (5) in Theorem 3.1 parametrize all the complex structures in Family I. Furthermore, we classify them up to equivalence. Then, in Section 3.2.2, we compute the ascending central series of the underlying 8-dimensional nilpotent Lie algebras, reaching Table 1.

3.2.1. Classification up to equivalence. The starting point of this section is Proposition 3.4, where we choose $B = 0$ according to Definition 3.5. After some technical lemmas we will arrive at equations (5).

Lemma 3.6. *Let J be a complex structure in Family I. Then, there exists a basis of $(1,0)$ -forms $\{\omega^k\}_{k=1}^4$ such that*

$$(14) \quad \begin{cases} d\omega^1 = 0, \\ d\omega^2 = A\omega^{1\bar{1}}, \\ d\omega^3 = D\omega^{12} + G\omega^{2\bar{1}} - E(\omega^{14} - \omega^{1\bar{4}}), \\ d\omega^4 = L\omega^{1\bar{1}} + is\omega^{2\bar{2}} + (N\omega^{13} - \bar{N}\omega^{3\bar{1}}), \end{cases}$$

where the coefficients $A, D, E, G, L, N \in \mathbb{C}$ and $s \in \mathbb{R}$ satisfy

$$(15) \quad isA + \bar{N}D = 0, \quad \Re(N\bar{E}) = 0, \quad NE \neq 0, \quad (D, G, s) \neq (0, 0, 0).$$

Proof. We first observe that if we impose $B = 0$ in (13), then we are forced to consider $H = 0$. Consequently, equations (13) become

$$isA - N\bar{C} + \bar{N}D = 0, \quad \Re(N\bar{E}) = 0, \quad NE \neq 0, \quad (C, D, G, M, s) \neq (0, 0, 0, 0, 0).$$

In order to get equations (14), it suffices to show that the coefficients C, F , and M in (12) can be set equal to zero. This can be done by defining the new $(1,0)$ -basis $\{\tau^k\}_{k=1}^4$ as follows:

$$\tau^1 = \omega^1, \quad \tau^2 = \omega^2, \quad \tau^3 = \omega^3 + \frac{\bar{M}}{N}\omega^2, \quad \tau^4 = \omega^4 - \frac{C}{E}\omega^2 + \frac{M\bar{A} + N\bar{F}}{N\bar{E}}\omega^1.$$

Indeed, in terms of $\{\tau^k\}_{k=1}^4$ the complex structure equations are of the form (12) with new coefficients $A_\tau, \dots, N_\tau \in \mathbb{C}$ and $s_\tau \in \mathbb{R}$ satisfying

$$B_\tau = C_\tau = H_\tau = F_\tau = M_\tau = 0.$$

Renaming the basis and the coefficients, we directly obtain the equations (14). Notice that the conditions (13) reduce to (15). \square

Lemma 3.7. *Let J be a complex structure in Family I. Then, there exists a basis of $(1,0)$ -forms $\{\omega^k\}_{k=1}^4$ such that*

$$\begin{cases} d\omega^1 = 0, \\ d\omega^2 = \varepsilon \omega^{1\bar{1}}, \\ d\omega^3 = \omega^{14} + \omega^{1\bar{4}} + i\delta \varepsilon b \omega^{1\bar{2}} + G \omega^{2\bar{1}}, \\ d\omega^4 = L \omega^{1\bar{1}} + b \omega^{2\bar{2}} + i\delta (\omega^{1\bar{3}} - \omega^{3\bar{1}}), \end{cases}$$

where $\varepsilon \in \{0, 1\}$, $\delta = \pm 1$, $G, L \in \mathbb{C}$, $b \in \mathbb{R}$ satisfying $(G, b) \neq (0, 0)$.

Proof. Since J is in Family I, by Lemma 3.6 there is a basis $\{\omega^k\}_{k=1}^4$ satisfying the equations (14) together with the conditions (15). We consider the $(1,0)$ -basis $\{\tau^k\}_{k=1}^4$ defined by

$$\tau^k = \lambda_k \omega^k, \text{ for } 1 \leq k \leq 4,$$

where

$$\lambda_1 = |\Im(N\bar{E})|^{1/2}, \quad \lambda_2 = \begin{cases} 1, & \text{if } A = 0, \\ \frac{|\Im(N\bar{E})|}{A}, & \text{if } A \neq 0, \end{cases} \quad \lambda_3 = -i \frac{|\Im(N\bar{E})|^{1/2}}{E}, \quad \lambda_4 = i.$$

It suffices to rename the basis and the coefficients in the corresponding structure equations in order to get the desired result. Here, we simply note that the value $\varepsilon = 0$ (resp. $\varepsilon = 1$) in the statement of the lemma comes from the case $A = 0$ (resp. $A \neq 0$). Moreover, $\delta = \pm 1$ where the value of the sign \pm precisely corresponds to $\text{sign}(\Im(N\bar{E}))$. The coefficient in $\omega^{1\bar{2}}$ comes from the Jacobi identity, concretely from the condition $d^2\omega^4 = 0$. \square

Using the previous lemma, in the following result we arrive at the desired reduced structure equations (5) of Theorem 3.1 for complex structures in the Family I.

Proposition 3.8. *Every 8-dimensional nilpotent Lie algebra \mathfrak{g} endowed with a complex structure J in Family I admits a basis of $(1,0)$ -forms satisfying the structure equations*

$$\begin{cases} d\omega^1 = 0, \\ d\omega^2 = \varepsilon \omega^{1\bar{1}}, \\ d\omega^3 = \omega^{14} + \omega^{1\bar{4}} + a \omega^{2\bar{1}} + i\delta \varepsilon b \omega^{1\bar{2}}, \\ d\omega^4 = i\nu \omega^{1\bar{1}} + b \omega^{2\bar{2}} + i\delta (\omega^{1\bar{3}} - \omega^{3\bar{1}}), \end{cases}$$

where $\varepsilon, \nu \in \{0, 1\}$, $\delta = \pm 1$, and $a, b \in \mathbb{R}$ with $a \geq 0$ and $(a, b) \neq (0, 0)$.

Proof. Consider the complex structure equations in Lemma 3.7 in terms of a $(1,0)$ -basis $\{\sigma^k\}_{k=1}^4$ with coefficients $(\varepsilon_\sigma, \delta_\sigma, b_\sigma, G_\sigma, L_\sigma)$. We first normalize the coefficient L_σ by applying the change of basis

$$\tau^1 = \sigma^1, \quad \tau^2 = \sigma^2, \quad \tau^3 = \lambda \left(\sigma^3 + \frac{i \Re L_\sigma}{2\delta_\sigma} \sigma^1 \right), \quad \tau^4 = \lambda \sigma^4,$$

where $\lambda \in \mathbb{R}^*$ is defined by

$$\lambda = \begin{cases} 1, & \text{if } \Im L_\sigma = 0, \\ \frac{1}{\Im L_\sigma}, & \text{if } \Im L_\sigma \neq 0. \end{cases}$$

The new structure equations still follow Lemma 3.7, but now with coefficients $\varepsilon_\tau = \varepsilon_\sigma$, $G_\tau = \lambda G_\sigma$, $b_\tau = \lambda b_\sigma$, $\delta_\tau = \delta_\sigma$ and $L_\tau = \lambda(L_\sigma - \Re L_\sigma) = i\lambda \Im L_\sigma = i\nu \in \{0, i\}$, in terms of the $(1,0)$ -basis $\{\tau^k\}_{k=1}^4$.

Now, writing the complex coefficient G_τ as $G_\tau = |G_\tau|e^{i\alpha}$ for some $\alpha \in [0, 2\pi)$, we define a new $(1,0)$ -basis $\{\omega^i\}_{i=1}^4$ as follows:

$$\omega^1 = e^{-i\alpha/2} \tau^1, \quad \omega^2 = \tau^2, \quad \omega^3 = e^{-i\alpha/2} \tau^3, \quad \omega^4 = \tau^4.$$

This concludes the proof, simply denoting $a = |G_\tau| \geq 0$. \square

After having reduced the complex structure equations of Family I to (5), we must finish the study of equivalences between the different parameters in our equations.

Let J and J' two complex structures on an NLA \mathfrak{g} parametrized by Family I. Let $\{\omega^k\}_{k=1}^4$ and $\{\omega'^k\}_{k=1}^4$ be basis of $\mathfrak{g}_J^{1,0}$ and $\mathfrak{g}_{J'}^{1,0}$, respectively. Then, (\mathfrak{g}, J) , resp. (\mathfrak{g}, J') , satisfies the complex structure equations given in Proposition 3.8 with parameters $(\varepsilon, \nu, \delta, a, b)$, resp. $(\varepsilon', \nu', \delta', a', b')$. Any isomorphism F between these pairs, as presented at the beginning of Section 3, is defined by

$$(16) \quad F(\omega'^i) = \sum_{j=1}^4 \lambda_j^i \omega^j, \quad \text{for each } 1 \leq i \leq 4,$$

and satisfies the conditions

$$(17) \quad d(F(\omega'^i)) = F(d\omega'^i),$$

where the matrix $\Lambda = (\lambda_j^i)_{i,j=1,\dots,4}$ belongs to $\text{GL}(4, \mathbb{C})$. Obviously, we are interested in those isomorphisms F providing equivalence relations between different J and J' , in the sense that $(\varepsilon, \nu, \delta, a, b) \neq (\varepsilon', \nu', \delta', a', b')$. To simplify our discussion, we will make use of the following notation:

Notation 3.9. We will denote $[d(F(\omega'^k)) - F(d\omega'^k)]_{ij}$ the coefficient for ω^{ij} in the expression $d(F(\omega'^k)) - F(d\omega'^k)$. Similarly, for the coefficient of $\omega^{i\bar{j}}$.

The following result reduce the general expression of the isomorphism (16).

Lemma 3.10. The forms $F(\omega'^i) \in \mathfrak{g}_J^{1,0}$ satisfy the conditions:

$$(18) \quad F(\omega'^1) \wedge \omega^1 = 0, \quad F(\omega'^2) \wedge \omega^{12} = 0, \quad F(\omega'^3) \wedge \omega^{123} = 0, \quad F(\omega'^4) \wedge \omega^{14} = 0.$$

In particular, the matrix $\Lambda = (\lambda_j^i)_{i,j=1,\dots,4}$ that determines F is triangular and thus

$$\prod_{i=1}^4 \lambda_i^i = \det \Lambda \neq 0.$$

Proof. A direct calculation of the conditions (17) for $i = 1, 2$ shows that

$$\lambda_3^1 = \lambda_4^1 = \lambda_3^2 = \lambda_4^2 = 0.$$

Consequently, $F(\omega'^1), F(\omega'^2) \in \langle \omega^1, \omega^2 \rangle$. In particular, $F(\omega'^2) \wedge \omega^{12} = 0$, as stated in the lemma. Moreover, we notice that Λ becomes a block triangular matrix, in such a way that $\det \Lambda = \det (\lambda_j^i)_{i,j=1,2} \cdot \det (\lambda_j^i)_{i,j=3,4} \neq 0$.

If we now compute (17) for $i = 3$, then one obtains

$$[d(F(\omega'^3)) - F(d\omega'^3)]_{3\bar{1}} = -i \delta \lambda_4^3.$$

Since this expression must be equal to zero, one immediately has $\lambda_4^3 = 0$. Thus, $F(\omega'^3) \wedge \omega^{123} = 0$ as required. Moreover, $0 \neq \det (\lambda_j^i)_{i,j=3,4} = \lambda_3^3 \lambda_4^4$ and necessarily $\lambda_1^1 \neq 0$, since these three parameters are related by the annihilation of

$$[d(F(\omega'^3)) - F(d\omega'^3)]_{14} = \lambda_3^3 - \lambda_1^1 \lambda_4^4.$$

As a consequence of $\lambda_1^1, \lambda_3^3, \lambda_4^4$ being non-zero, the annihilation of

$$[d(F(\omega'^3)) - F(d\omega'^3)]_{13} = -\lambda_1^1 \lambda_3^4, \quad [d(F(\omega'^3)) - F(d\omega'^3)]_{24} = -\lambda_2^2 \lambda_4^4$$

gives $\lambda_2^2 = \lambda_3^4 = 0$. In particular, we conclude $F(\omega'^1) \wedge \omega^1 = 0$. Finally, from

$$0 = [d(F(\omega'^3)) - F(d\omega'^3)]_{12} = -\lambda_1^1 \lambda_2^4,$$

we obtain $\lambda_2^4 = 0$, i.e. $F(\omega'^4) \wedge \omega^{14} = 0$. \square

As a consequence, a first relation between the tuples $(\varepsilon, \nu, \delta, a, b)$ and $(\varepsilon', \nu', \delta', a', b')$ is attained:

Proposition 3.11. If the complex structures J and J' are equivalent, then

$$\varepsilon' = \varepsilon, \quad \nu' = \nu, \quad \delta' = \delta.$$

Moreover, there exists an isomorphism (16) satisfying Lemma 3.10 and

$$\lambda_1^1 = e^{i\theta}, \quad \lambda_4^4 = \lambda \in \mathbb{R}^*, \quad \lambda_3^3 = \lambda e^{i\theta}, \quad \nu(1 - \lambda) = 0, \quad \varepsilon(1 - \lambda_2^2) = 0,$$

where $\theta \in [0, 2\pi)$.

Proof. We first observe that Lemma 3.10 must hold in order to have an isomorphism F relating the complex structures J and J' . Taking this as a starting point, let us recalculate the conditions (17) for each $1 \leq i \leq 4$.

One can easily check that $F(d\omega'^i) = d(F(\omega'^i))$ is trivially fulfilled for $i = 1$. For $i = 2$, one simply has:

$$(19) \quad 0 = d(F(\omega'^2)) - F(d\omega'^2) = (\varepsilon \lambda_2^2 - \varepsilon' |\lambda_1^1|^2) \omega^{1\bar{1}}.$$

Observe the following. If we suppose $\varepsilon = 0$ then $\varepsilon' = 0$, as Lemma 3.10 states $\lambda_1^1 \neq 0$. Similarly, if $\varepsilon = 1$ then $\lambda_2^2 = \varepsilon' |\lambda_1^1|^2 \neq 0$, and the only possibility is taking $\varepsilon' = 1$. As a consequence, we conclude

$$\varepsilon' = \varepsilon.$$

For $i = 3$, we highlight the following terms:

$$[d(F(\omega'^3)) - F(d\omega'^3)]_{14} = \lambda_3^3 - \lambda_1^1 \lambda_4^4, \quad [d(F(\omega'^3)) - F(d\omega'^3)]_{1\bar{4}} = \lambda_3^3 - \lambda_1^1 \bar{\lambda}_4^4.$$

The annihilation of the previous coefficients give

$$(20) \quad \lambda_4^4 = \lambda \in \mathbb{R}^*, \quad \lambda_3^3 = \lambda \lambda_1^1.$$

Moreover, for $i = 4$ and taking into account (20), we get

$$0 = [d(F(\omega'^4)) - F(d\omega'^4)]_{13} = i \lambda (\delta - \delta' |\lambda_1^1|^2).$$

Since $\lambda, \lambda_1^1 \neq 0$ and $\delta, \delta' \in \{-1, 1\}$, one necessarily has

$$\delta' = \delta, \quad |\lambda_1^1|^2 = 1.$$

In particular, we can set $\lambda_1^1 = e^{i\theta}$, for some $\theta \in [0, 2\pi)$. Finally,

$$0 = [d(F(\omega'^4)) - F(d\omega'^4)]_{1\bar{1}} = i(\nu \lambda - \nu') - (b' |\lambda_1^2|^2 + 2\delta \Im(\lambda_1^3 e^{-i\theta})).$$

The imaginary part of the previous equation implies that either $\nu' = \nu = 0$ or $\nu' = \nu = 1$ with $\lambda = 1$, since $\lambda \neq 0$ and $\nu, \nu' \in \{0, 1\}$. Notice that this is equivalent to writing:

$$\nu' = \nu, \quad \nu(1 - \lambda) = 0.$$

This gives the desired results. Note that the expression $\varepsilon(1 - \lambda_2^2) = 0$ comes from rewriting (19). \square

From now on, in order to determine the space of complex structures up to equivalence, we can fix parameters ε, ν, δ and simply identify J and J' with the pairs (a, b) and (a', b') , respectively. Recall that $(a, b), (a', b') \neq (0, 0)$ and $a, a' \geq 0$.

Proposition 3.12. *The complex structures J and J' are equivalent if and only if there exists an isomorphism given by:*

$$(21) \quad \omega'^1 = e^{i\theta} \omega^1, \quad \omega'^2 = \lambda_2^2 \omega^2, \quad \omega'^3 = \lambda e^{i\theta} \omega^3, \quad \omega'^4 = \lambda \omega^4,$$

where $\theta \in [0, 2\pi)$, $\lambda_2^2 \in \mathbb{C}^*$, $\lambda \in \mathbb{R}^*$ and

$$(22) \quad \Im(\lambda_2^2 e^{-2i\theta}) = 0, \quad \nu(1 - \lambda) = 0, \quad \varepsilon(1 - \lambda_2^2) = 0.$$

Moreover, the parameters (a, b) and (a', b') that respectively determine J and J' are related by

$$(23) \quad a' = a \frac{\lambda}{\lambda_2^2 e^{-2i\theta}}, \quad b' = b \frac{\lambda}{|\lambda_2^2|^2},$$

Proof. According to previous results, if J and J' are equivalent, then there exists an isomorphism of type (16) satisfying Proposition 3.11. We must ensure that F fulfills (17) for each $1 \leq i \leq 4$.

First, one checks that the desired conditions are equivalent to the following equations:

$$\begin{aligned}
0 &= [d(F(\omega'^3)) - F(d\omega'^3)]_{1\bar{1}} = \varepsilon(\lambda_2^3 - i\delta b' \bar{\lambda}_1^2 e^{i\theta}) - \bar{\lambda}_1^4 e^{i\theta} - a' \lambda_1^2 e^{-i\theta}, \\
0 &= [d(F(\omega'^3)) - F(d\omega'^3)]_{1\bar{2}} = i\delta \varepsilon e^{i\theta} (b\lambda - b' \bar{\lambda}_2^2), \\
0 &= [d(F(\omega'^3)) - F(d\omega'^3)]_{2\bar{1}} = a\lambda e^{i\theta} - a' \lambda_2^2 e^{-i\theta}, \\
0 &= [d(F(\omega'^4)) - F(d\omega'^4)]_{1\bar{1}} = -2\delta \Im(\lambda_1^3 e^{-i\theta}) - b' |\lambda_1^2|^2, \\
0 &= [d(F(\omega'^4)) - F(d\omega'^4)]_{1\bar{2}} = -i\delta \bar{\lambda}_2^3 e^{i\theta} - b' \lambda_1^2 \bar{\lambda}_1^2, \\
0 &= [d(F(\omega'^4)) - F(d\omega'^4)]_{2\bar{2}} = b\lambda - b' |\lambda_2^2|^2.
\end{aligned}$$

Now, notice that the pairs (a, b) and (a', b') , which determine the complex structures J and J' , are related by $[d(F(\omega'^3)) - F(d\omega'^3)]_{2\bar{1}}$ and $[d(F(\omega'^4)) - F(d\omega'^4)]_{2\bar{2}}$. Hence, a' and b' are determined by these expressions, obtaining (23). In particular, the equivalence between the complex structures J and J' only depends on the parameters θ , λ , and λ_2^2 . Hence, the parameters λ_j^i for $i \neq j$ do not affect the relation (23), and they can be chosen to be zero. This solves the remaining equations and gives (21).

Finally, the first expression in (22) comes from imposing $a' \in \mathbb{R}$ in (23) whereas the other two are a direct consequence of Proposition 3.11. \square

Finally we can set the main result about equivalences of complex structures in Family I:

Theorem 3.13. *Up to equivalence, the complex structures defined by*

$$\begin{cases} d\omega^1 = 0, \\ d\omega^2 = \varepsilon \omega^{1\bar{1}}, \\ d\omega^3 = \omega^{14} + \omega^{1\bar{4}} + a\omega^{2\bar{1}} + i\delta \varepsilon b \omega^{1\bar{2}}, \\ d\omega^4 = i\nu \omega^{1\bar{1}} + b\omega^{2\bar{2}} + i\delta(\omega^{13} - \omega^{3\bar{1}}), \end{cases}$$

where $\delta = \pm 1$, $\varepsilon, \nu \in \{0, 1\}$, $a \geq 0$, $b \in \mathbb{R}$ with $(a, b) \neq (0, 0)$, are classified as follows:

- (i) $(\varepsilon, \nu, a, b) = (0, 0, 0, 1), (0, 0, 1, 0), (0, 0, 1, 1)$;
- (ii) $(\varepsilon, \nu, a, b) = (0, 1, 0, b/|b|), (0, 1, 1, b)$;
- (iii) $(\varepsilon, \nu, a, b) = (1, 0, 0, 1), (1, 0, 1, |b|)$;
- (iv) $(\varepsilon, \nu, a, b) = (1, 1, a, b)$.

Proof. Let us study different cases depending on the values of the pair (ε, ν) . Recall that the conditions given in Proposition 3.12 must be satisfied, namely

$$a' = a \frac{\lambda}{\lambda_2^2 e^{-2i\theta}}, \quad b' = b \frac{\lambda}{|\lambda_2^2|^2},$$

$$\text{where } \Im(\lambda_2^2 e^{-2i\theta}) = 0, \quad \nu(1 - \lambda) = 0, \quad \varepsilon(1 - \lambda_2^2) = 0.$$

These expressions will give us the desired equivalences between (a, b) and (a', b') , thus between our complex structures.

- (i) $(\varepsilon, \nu) = (0, 0)$: There are no restrictions on λ and λ_2^2 , so it is possible to normalize a and/or b (i.e. take $a' = 1$ or $b' = 1$) when they are non-zero. In particular, this is easy for $ab = 0$. For $ab \neq 0$, one can take $\theta = 0$, $\lambda = b/a^2$, $\lambda_2^2 = b/a$.
- (ii) $(\varepsilon, \nu) = (0, 1)$: In this case $\lambda = 1$ and $\lambda_2^2 \in \mathbb{C}^*$ is a free parameter. When a is non-zero, we can normalize it. If $a = 0$, we can choose $\lambda_2^2 = \sqrt{|b|}$, and thus $b' = \pm 1$.
- (iii) $(\varepsilon, \nu) = (1, 0)$: From the expressions above we get $\lambda_2^2 = 1$ and $\lambda \in \mathbb{R}^*$ free, so $a' = a\lambda e^{2i\theta}$, $b' = b\lambda$. Observe that $e^{2i\theta}$ is a real number, so the only possible choices are $\theta = 0$ or $\theta = \pi/2$. If $a = 0$ one can normalize b' , and if $a > 0$ we can take $a' = 1$ and $b' \geq 0$. In fact, in the last case it suffices to consider $e^{2i\theta} = b/|b|$ and $\lambda = b/a|b|$.
- (iv) $(\varepsilon, \nu) = (1, 1)$: We are forced to impose $\lambda = \lambda_2^2 = 1$ and thus $a' = ae^{2i\theta}$ and $b' = b$. Since $a, a' \geq 0$, necessarily $e^{2i\theta} = 1$ and $a' = a$.

\square

To complete the proof of Theorem 3.1 (i), it suffices to study the ascending type of the Lie algebras underlying Family I.

3.2.2. *Ascending central series.* This section is devoted to explicitly compute the ascending central series of the nilpotent Lie algebras underlying Family I, taking as starting point the structure equations in Proposition 3.8.

Let $\{Z_k\}_{k=1}^4$ be the dual basis to $\{\omega^k\}_{k=1}^4$. In terms of this basis, a generic (real) element $X \in \mathfrak{g}$ is given by

$$(24) \quad X = \sum_{i=1}^4 \alpha_i Z_i + \sum_{i=1}^4 \bar{\alpha}_i \bar{Z}_i,$$

where $\alpha_i \in \mathbb{C}$, and \bar{Z}_i is the conjugate of Z_i .

From the structure equations in Proposition 3.8, it follows that the brackets $[X, Z_k]$, for $1 \leq k \leq 4$, are given by

$$(25) \quad \begin{aligned} [X, Z_1] &= \varepsilon \bar{\alpha}_1 (Z_2 - \bar{Z}_2) + (\alpha_4 + \bar{\alpha}_4 + i \delta \varepsilon b \bar{\alpha}_2) Z_3 - a \bar{\alpha}_2 \bar{Z}_3 \\ &\quad + i \nu \bar{\alpha}_1 (Z_4 + \bar{Z}_4) + i \delta \bar{\alpha}_3 (Z_4 - \bar{Z}_4), \\ [X, Z_2] &= a \bar{\alpha}_1 Z_3 + i \delta \varepsilon b \bar{\alpha}_1 \bar{Z}_3 + b \bar{\alpha}_2 (Z_4 - \bar{Z}_4), \\ [X, Z_3] &= -i \delta \bar{\alpha}_1 (Z_4 - \bar{Z}_4), \\ [X, Z_4] &= -\alpha_1 Z_3 - \bar{\alpha}_1 \bar{Z}_3. \end{aligned}$$

Note that since X is real, the bracket $[X, \bar{Z}_k]$ is just the conjugate of $[X, Z_k]$, for $1 \leq k \leq 4$. Recall that $\varepsilon, \nu \in \{0, 1\}$, $\delta = \pm 1$, and $a, b \in \mathbb{R}$ with $a \geq 0$ and $(a, b) \neq (0, 0)$.

Due to Corollary 2.6 and $[X, Z_4 - \bar{Z}_4] = 0$ for every $X \in \mathfrak{g}$, one directly concludes

$$\mathfrak{g}_1 = \langle \Im Z_4 \rangle.$$

Observe that this is consistent with the discussion about the center in page 7, as the change of basis in the proof of Lemma 3.7 switches the real and imaginary parts of ω^4 , thus of Z_4 .

Lemma 3.14. *In the conditions above, the term \mathfrak{g}_2 in the ascending central series is given by:*

- (i) for $a \neq 0$: $\mathfrak{g}_2 = \langle \Re Z_3, \Im Z_3, \Im Z_4 \rangle$ and so $\dim \mathfrak{g}_2 = 3$;
- (ii) for $a = 0$ and $\varepsilon = 1$: $\mathfrak{g}_2 = \langle 2\delta \Im Z_2 + b \Re Z_4, \Re Z_3, \Im Z_3, \Im Z_4 \rangle$ and $\dim \mathfrak{g}_2 = 4$;
- (iii) for $a = \varepsilon = 0$: $\mathfrak{g}_2 = \langle \Re Z_2, \Im Z_2, \Re Z_3, \Im Z_3, \Im Z_4 \rangle$ and $\dim \mathfrak{g}_2 = 5$.

Proof. Let X be a generic element given by (24). Then, X belongs to the term \mathfrak{g}_2 in the ascending central series if and only if $[X, Z_k] \in \mathfrak{g}_1$, for every $1 \leq k \leq 4$. Since $\mathfrak{g}_1 = \langle \Im Z_4 \rangle$, bearing in mind (25) we get that $X \in \mathfrak{g}_2$ if and only if

$$\alpha_1 = 0, \quad a \alpha_2 = 0, \quad \alpha_4 + \bar{\alpha}_4 - i \delta \varepsilon b \alpha_2 = 0.$$

In particular, one directly has $\Re Z_3, \Im Z_3 \in \mathfrak{g}_2$. Now, to solve the previous system we must consider the following three different cases:

- If $a \neq 0$, then $\alpha_2 = 0$ and $\alpha_4 = i \alpha$ with $\alpha \in \mathbb{R}$. Hence, $X = \alpha_3 Z_3 + \bar{\alpha}_3 \bar{Z}_3 + i \alpha (Z_4 - \bar{Z}_4)$ and we get (i).
- Case $a = 0, \varepsilon = 1$. Since $b \neq 0$, we get $\alpha_2 = -\frac{i\delta}{b}(\alpha_4 + \bar{\alpha}_4)$, being $\alpha_4 \in \mathbb{C}$ a free parameter. Hence,

$$\begin{aligned} X &= -\frac{i\delta}{b}(\alpha_4 + \bar{\alpha}_4)(Z_2 - \bar{Z}_2) + \alpha_3 Z_3 + \bar{\alpha}_3 \bar{Z}_3 + \alpha_4 Z_4 + \bar{\alpha}_4 \bar{Z}_4 \\ &= -\frac{i\delta}{b}(\alpha_4 + \bar{\alpha}_4)(Z_2 - \bar{Z}_2) + \alpha_3 Z_3 + \bar{\alpha}_3 \bar{Z}_3 + \frac{1}{2}(\alpha_4 + \bar{\alpha}_4)(Z_4 + \bar{Z}_4) + \frac{1}{2}(\alpha_4 - \bar{\alpha}_4)(Z_4 - \bar{Z}_4) \\ &= -\frac{1}{2b}(\alpha_4 + \bar{\alpha}_4) (2\delta i (Z_2 - \bar{Z}_2) - b(Z_4 + \bar{Z}_4)) + \alpha_3 Z_3 + \bar{\alpha}_3 \bar{Z}_3 + \frac{1}{2}(\alpha_4 - \bar{\alpha}_4)(Z_4 - \bar{Z}_4), \end{aligned}$$

which proves (ii).

- Finally, in case $a = \varepsilon = 0$ the only remaining equation is $\alpha_4 + \bar{\alpha}_4 = 0$, i.e. $\alpha_4 = i \alpha$ with $\alpha \in \mathbb{R}$. Thus, $X = \alpha_2 Z_2 + \bar{\alpha}_2 \bar{Z}_2 + \alpha_3 Z_3 + \bar{\alpha}_3 \bar{Z}_3 + i \alpha (Z_4 - \bar{Z}_4)$, and (iii) is proved. \square

Proposition 3.15. *Let \mathfrak{g} be an 8-dimensional NLA endowed with a complex structure J in Family I with equations given in Proposition 3.8. Then, the ascending type of \mathfrak{g} is as follows:*

- (i) if $a \neq 0$ and
 - (i.1) $(\varepsilon, \nu) = (0, 0)$, then $(\dim \mathfrak{g}_k)_k = (1, 3, 8)$;

- (i.2) $(\varepsilon, \nu) \neq (0, 0)$, then $(\dim \mathfrak{g}_k)_k = (1, 3, 6, 8)$;
- (ii) if $a = 0$, $\varepsilon = 1$ and
 - (ii.1) $(\nu, b) = (1, 2\delta)$, then $(\dim \mathfrak{g}_k)_k = (1, 4, 8)$;
 - (ii.2) $(\nu, b) \neq (1, 2\delta)$, then $(\dim \mathfrak{g}_k)_k = (1, 4, 6, 8)$;
- (iii) if $a = \varepsilon = 0$ and
 - (iii.1) $\nu = 0$, then $(\dim \mathfrak{g}_k)_k = (1, 5, 8)$;
 - (iii.2) $\nu = 1$, then $(\dim \mathfrak{g}_k)_k = (1, 5, 6, 8)$.

Proof. A generic element X given by (24) belongs to the term \mathfrak{g}_3 in the ascending central series if and only if $[X, Z_k] \in \mathfrak{g}_2$, for every $1 \leq k \leq 4$. From (25) it follows that this happens for $k = 2, 3$ and 4 , since $\langle \Re Z_3, \Im Z_3, \Im Z_4 \rangle \subseteq \mathfrak{g}_2$ by Lemma 3.14. Hence, we must focus on the bracket $[X, Z_1]$, which can be rewritten as

$$[X, Z_1] = 2i\bar{\alpha}_1 (\varepsilon \Im Z_2 + \nu \Re Z_4) + \Upsilon,$$

for some $\Upsilon \in \mathfrak{g}_2$. Therefore, $X \in \mathfrak{g}_3$ if and only if one of the following cases holds:

- $\varepsilon \Im Z_2 + \nu \Re Z_4 \in \mathfrak{g}_2$, which implies that any $X \in \mathfrak{g}_3$, i.e. $\mathfrak{g}_3 = \mathfrak{g}$ and so $\dim \mathfrak{g}_3 = 8$. This case occurs when $\varepsilon = \nu = 0$, i.e. in cases (i.1) and (iii.1), and by Lemma 3.14 (ii) when $2\delta\nu - b\varepsilon = 0$. Indeed, in the latter case the element $\varepsilon \Im Z_2 + \nu \Re Z_4$ is a multiple of $2\delta \Im Z_2 + b \Re Z_4$, so (ii.1) is proved.
- $\varepsilon \Im Z_2 + \nu \Re Z_4 \notin \mathfrak{g}_2$ and $\alpha_1 = 0$, i.e. $\mathfrak{g}_3 = \langle \Re Z_2, \Im Z_2, \Re Z_3, \Im Z_3, \Re Z_4, \Im Z_4 \rangle$ and so $\dim \mathfrak{g}_3 = 6$. In this case any X belongs to \mathfrak{g}_4 , i.e. $\mathfrak{g}_4 = \mathfrak{g}$, getting the remaining cases (i.2), (ii.2) and (iii.2).

□

Combining the previous result with Theorem 3.13, one obtains part (i) of Theorem 3.1.

3.3. Study of Family II. In Section 3.3.1 we arrive at the reduced equations (6) in Theorem 3.1. We also complete the classification of complex structures up to equivalence. In Section 3.3.2, we study the ascending type of the 8-dimensional nilpotent Lie algebras that admit a complex structure in Family II, reaching Table 2. This shows that Families I and II do not overlap.

3.3.1. Classification up to equivalence. Here our starting point is Proposition 3.4, now taking $B \neq 0$ due to Definition 3.5.

Lemma 3.16. *Let J be a complex structure in Family II. Then, there exists a basis of $(1, 0)$ -forms $\{\omega^k\}_{k=1}^4$ such that*

$$(26) \quad \begin{cases} d\omega^1 = 0, \\ d\omega^2 = \omega^{14} + \omega^{1\bar{4}}, \\ d\omega^3 = F\omega^{1\bar{1}} + C\omega^{12} + \bar{C}(\omega^{1\bar{2}} - \omega^{2\bar{1}}) - is(\omega^{24} + \omega^{2\bar{4}}), \\ d\omega^4 = L\omega^{1\bar{1}} + s\omega^{2\bar{2}} + ib(\omega^{1\bar{2}} - \omega^{2\bar{1}}) + i(\omega^{1\bar{3}} - \omega^{3\bar{1}}), \end{cases}$$

where $C, F, L \in \mathbb{C}$ and $b, s \in \mathbb{R}$ satisfy that $s \Im L = 0$ and $(C, b, s) \neq (0, 0, 0)$.

Proof. Since $B \neq 0$ by hypothesis and $N \neq 0$ by the conditions (13), we can define the following $(1, 0)$ -basis

$$\tau^1 = \omega^1, \quad \tau^2 = -\frac{i}{B}\omega^2, \quad \tau^3 = \bar{N}\omega^3 - \frac{E\bar{N}}{B}\omega^2, \quad \tau^4 = i\omega^4 + i\frac{\bar{A}}{B}\omega^1.$$

With respect to $\{\tau^k\}_{k=1}^4$, the complex structure equations (12) in Proposition 3.4 become

$$\begin{cases} d\tau^1 = 0, \\ d\tau^2 = \tau^{14} + \tau^{1\bar{4}}, \\ d\tau^3 = F_\tau \tau^{1\bar{1}} + C_\tau \tau^{12} + D_\tau \tau^{1\bar{2}} + G_\tau \tau^{2\bar{1}} - H_\tau (\tau^{24} + \tau^{2\bar{4}}), \\ d\tau^4 = L_\tau \tau^{1\bar{1}} + s_\tau \tau^{2\bar{2}} + (M_\tau \tau^{1\bar{2}} + \bar{M}_\tau \tau^{2\bar{1}}) + i(\tau^{1\bar{3}} - \tau^{3\bar{1}}), \end{cases}$$

where the coefficients $F_\tau, C_\tau, D_\tau, G_\tau, H_\tau, L_\tau, M_\tau \in \mathbb{C}$ and $s_\tau \in \mathbb{R}$ are expressed in terms of the original ones as follows:

$$\begin{aligned} C_\tau &= \frac{iB\bar{N}}{B} (C\bar{B} - H\bar{A}), & D_\tau &= -i\bar{B}\bar{N}D, & G_\tau &= i\bar{N}(BG - AH), \\ H_\tau &= BH\bar{N}, & M_\tau &= M\bar{B} + N\bar{E}, & s_\tau &= -s|B|^2. \end{aligned}$$

Note that, in order to get the result, it suffices to check that

$$D_\tau = -G_\tau = \bar{C}_\tau, \quad H_\tau = i s_\tau, \quad M_\tau \in i\mathbb{R}.$$

Using the first condition in (13), it is straightforward to see that $G_\tau = i\bar{N}\bar{B}D = -D_\tau$. Now, from the third equation in the first column in (13) we have $is = \frac{N\bar{H}}{B}$, and replacing this value in the second equation of the same column, we get

$$N(A\bar{H} - B\bar{C}) + B\bar{N}D = 0.$$

By conjugating this expression, it is easy to arrive at $C_\tau = iBN\bar{D} = \bar{D}_\tau$. In addition, from the third equation in the first column in (13) we also have $H_\tau = -is|B|^2 = i s_\tau$, and from the second condition in the second column we obtain $M_\tau \in i\mathbb{R}$.

Renaming the basis and the coefficients, we directly get the equations (26). Finally, the Jacobi identity is equivalent to the equality $s \Im L = 0$, and the condition $(C, b, s) \neq (0, 0, 0)$ is necessary for the dimension of the center of \mathfrak{g} to be equal to 1. This concludes the proof of the lemma. \square

Lemma 3.17. *Let J be a complex structure in Family II. Then, there exists a basis of $(1, 0)$ -forms $\{\omega^k\}_{k=1}^4$ such that*

$$(27) \quad \begin{cases} d\omega^1 = 0, \\ d\omega^2 = \omega^{14} + \omega^{1\bar{4}}, \\ d\omega^3 = F\omega^{1\bar{1}} + \varepsilon(\omega^{12} + \omega^{1\bar{2}} - \omega^{2\bar{1}}) - is(\omega^{24} + \omega^{2\bar{4}}), \\ d\omega^4 = i\nu\omega^{11} + s\omega^{2\bar{2}} + ib(\omega^{12} - \omega^{2\bar{1}}) + i(\omega^{13} - \omega^{3\bar{1}}), \end{cases}$$

where $\varepsilon \in \{0, 1\}$, $F \in \mathbb{C}$ and $b, s, \nu \in \mathbb{R}$. Moreover, one has $s\nu = 0$ and $(\varepsilon, b, s) \neq (0, 0, 0)$.

Proof. Since J is in Family II, by Lemma 3.16 there is a basis $\{\omega^k\}_{k=1}^4$ satisfying the equations (26). Hence, writing $C = |C|e^{i\beta}$ for some $\beta \in [0, 2\pi)$, one can define a new $(1, 0)$ -basis as follows:

$$\tau^1 = \lambda\omega^1, \quad \tau^2 = \lambda\omega^2, \quad \tau^3 = \frac{1}{\lambda} \left(\omega^3 + \frac{i\Re L}{2} \omega^1 \right), \quad \tau^4 = \omega^4, \quad \text{where } \lambda = \begin{cases} 1, & \text{if } C = 0, \\ |C|^{1/3} e^{i\beta}, & \text{if } C \neq 0. \end{cases}$$

In terms of $\{\tau^k\}_{k=1}^4$ we obtain equations of the form (26) with new coefficients $C_\tau \in \{0, 1\}$, $F_\tau \in \mathbb{C}$, $L_\tau = i\nu_\tau \in i\mathbb{R}$, and $b_\tau, s_\tau \in \mathbb{R}$. Renaming the basis and the coefficients, we arrive at equations (27) in the statement. The condition $s\nu = 0$ comes from the Jacobi identity, concretely from $d^2\omega^3 = 0$, whereas $(\varepsilon, b, s) \neq (0, 0, 0)$ is a direct consequence of Lemma 3.16. \square

Lemma 3.18. *In Lemma 3.17, one can take $(s, \nu) \in \{(0, 0), (0, 1), (-1, 0)\}$.*

Proof. Observe that Lemma 3.17 states that the parameters $s, \nu \in \mathbb{R}$ must satisfy the condition $s\nu = 0$. Thus, we can study the following cases:

- Let us suppose $s = 0$. If $\nu \neq 0$, then we can define the $(1, 0)$ -basis

$$\tau^1 = \omega^1, \quad \tau^k = \frac{1}{\nu} \omega^k, \quad k = 2, 3, 4.$$

It suffices to rename the previous forms and the new coefficients in order to get (27) with $\nu = 1$. Therefore, for $s = 0$ one concludes that $\nu \in \{0, 1\}$.

- If $s \neq 0$, then $\nu = 0$ and we consider

$$\tau^1 = \omega^1, \quad \tau^k = -s\omega^k, \quad k = 2, 3, 4.$$

We reach (27) with $s = -1$.

This gives the desired reduction of the parameters s, ν . \square

Using the previous lemma, in the following result we arrive at the desired reduced structure equations (6) of Theorem 3.1 for complex structures in the Family II.

Proposition 3.19. *Every 8-dimensional nilpotent Lie algebra \mathfrak{g} endowed with a complex structure J in Family II admits a basis of $(1, 0)$ -forms satisfying the structure equations*

$$\begin{cases} d\omega^1 = 0, \\ d\omega^2 = \omega^{14} + \omega^{1\bar{4}}, \\ d\omega^3 = a\omega^{1\bar{1}} + \varepsilon(\omega^{12} + \omega^{1\bar{2}} - \omega^{2\bar{1}}) + i\mu(\omega^{24} + \omega^{2\bar{4}}), \\ d\omega^4 = i\nu\omega^{1\bar{1}} - \mu\omega^{2\bar{2}} + ib(\omega^{1\bar{2}} - \omega^{2\bar{1}}) + i(\omega^{1\bar{3}} - \omega^{3\bar{1}}), \end{cases}$$

where $\varepsilon, \mu, \nu \in \{0, 1\}$ such that $\mu\nu = 0$, $(\varepsilon, \mu) \neq (0, 0)$, and $a, b \in \mathbb{R}$.

Proof. Since J is in Family II, Lemmas 3.17 and 3.18 hold. Renaming $s = -\mu$, it suffices to see that the parameter F in (27) can be chosen to be a real number. Next we distinguish two cases depending on the value of ε :

- Assume that $\varepsilon = 0$. If we write $F = |F|e^{i\alpha}$, for some $\alpha \in [0, 2\pi)$, the result comes after applying the following change of basis:

$$\tau^k = e^{-i\alpha}\omega^k, \quad k = 1, 2, 3, \quad \tau^4 = \omega^4.$$

- Let $\varepsilon = 1$. In this case, the result is a consequence of defining a new $(1, 0)$ -basis as follows:

$$\tau^1 = \omega^1, \quad \tau^2 = \omega^2 - \frac{i\Im F}{2}\omega^1, \quad \tau^3 = \omega^3 + \frac{\mu\Im F}{2}\omega^2 + \frac{i\Im F(4b - 3\mu\Im F)}{8}\omega^1, \quad \tau^4 = \omega^4.$$

Finally, observe that if $(\varepsilon, \mu) = (0, 0)$ then it is easy to see that the dimension of the center of \mathfrak{g} would be greater than 1, which is a contradiction to Corollary 2.6. Therefore, $(\varepsilon, \mu) \neq (0, 0)$. \square

After having reduced our complex structure equations to (6), we must analyze the existence of equivalences in terms of the parameters appearing in the equations.

Let J and J' be two complex structures on an NLA \mathfrak{g} parametrized by Family II. Then, the $(1, 0)$ -basis $\{\omega^k\}_{k=1}^4$ (resp. $\{\omega'^k\}_{k=1}^4$) of (\mathfrak{g}, J) (resp. (\mathfrak{g}, J')) satisfies the complex structure equations given in Proposition 3.19 with parameters $(\varepsilon, \mu, \nu, a, b)$ (resp. $(\varepsilon', \mu', \nu', a', b')$). Similarly to Lemma 3.10, for Family II the isomorphism F given by (16) can be simplified as follows:

Lemma 3.20. *The forms $F(\omega'^k) \in \mathfrak{g}_J^{1,0}$ satisfy the conditions:*

$$F(\omega'^1) \wedge \omega^1 = 0, \quad F(\omega'^2) \wedge \omega^{12} = 0, \quad F(\omega'^3) \wedge \omega^{123} = 0, \quad F(\omega'^4) \wedge \omega^4 = 0.$$

In particular, the matrix $\Lambda = (\lambda_j^i)_{i,j=1,\dots,4}$ that determines F is triangular and thus

$$\prod_{i=1}^4 \lambda_i^i = \det \Lambda \neq 0.$$

Proof. The result comes straightforward by imposing $F(d\omega'^i) = d(F(\omega'^i))$ for $i = 1, 2, 3$, taking into account that $(\varepsilon, \mu) \neq (0, 0)$ and $\det \Lambda \neq 0$. \square

This allows us to find a first relation between the tuples $(\varepsilon, \mu, \nu, a, b)$ and $(\varepsilon', \mu', \nu', a', b')$.

Lemma 3.21. *In the conditions above, if there exists an equivalence between J and J' , then*

$$\varepsilon = \varepsilon', \quad \mu = \mu', \quad \nu = \nu'.$$

Proof. Observe that Lemma 3.20 must hold in order to have an isomorphism F relating J and J' . Imposing $F(d\omega'^i) = d(F(\omega'^i))$ for $i = 1, 2$, we get:

$$(28) \quad \lambda_4^4 = \lambda \in \mathbb{R}^*, \quad \lambda_2^2 = \lambda \lambda_1^1.$$

In addition, one has

$$[d(F(\omega'^3)) - F(d\omega'^3)]_{12} = \varepsilon \lambda_3^3 - \varepsilon' \lambda (\lambda_1^1)^2, \quad [d(F(\omega'^3)) - F(d\omega'^3)]_{24} = -i(\mu' \lambda^2 \lambda_1^1 - \mu \lambda_3^3).$$

Since $\lambda, \lambda_1^1, \lambda_3^3$ are non-zero numbers and $\varepsilon, \varepsilon', \mu, \mu' \in \{0, 1\}$, one can deduce that

$$\varepsilon = \varepsilon' \quad \mu = \mu'.$$

Finally, thanks to

$$0 = \Im \left([d(F(\omega'^4)) - F(d\omega'^4)]_{1\bar{1}} \right) = \lambda \nu - \nu' |\lambda_1^1|^2$$

we reach $\nu = \nu'$, because $\lambda, \lambda_1^1 \neq 0$ and $\nu, \nu' \in \{0, 1\}$. \square

Taking into account Lemma 3.21, we have $\varepsilon = \varepsilon', \mu = \mu', \nu = \nu'$, so it suffices to study the relation between the pairs (a, b) and (a', b') to analyze the equivalences between the complex structures J and J' .

Proposition 3.22. *If the complex structures J and J' are equivalent, then one of the following cases holds:*

- (i) if $(\varepsilon, \mu, \nu) = (1, 0, 0)$, then $a' = \lambda a$ and $b' = b$, with $\lambda \in \mathbb{R}^*$;
- (ii) if $(\varepsilon, \mu, \nu) = (0, 1, 0)$, then $a' = a/\kappa^5$ and $b' = 1/\kappa^2 (b + 2\kappa \Im \lambda_1^2)$, with $\kappa \in \mathbb{R}^*$;
- (iii) otherwise, $a' = a$ and $b' = b$.

Proof. Due to Lemma 3.20 and the condition (28), one obtains from the vanishing of $[d(F(\sigma^3)) - F(d\sigma^3)]_{14}$ that

$$\lambda_2^3 = i \mu \lambda \lambda_1^2.$$

In addition,

$$0 = [d(F(\sigma^3)) - F(d\sigma^3)]_{12} = \varepsilon (\lambda_3^3 - \lambda (\lambda_1^1)^2), \quad 0 = [d(F(\sigma^4)) - F(d\sigma^4)]_{3\bar{1}} = -i (\lambda - \lambda_3^3 \bar{\lambda}_1^1).$$

From the second equality, we deduce $\lambda_3^3 = \lambda/\bar{\lambda}_1^1$, in such a way that the first equation becomes

$$(29) \quad \varepsilon \lambda \lambda_1^1 \left(\frac{1}{|\lambda_1^1|^2} - \lambda_1^1 \right) = 0.$$

Recall that $\lambda, \lambda_1^1 \neq 0$. For $\varepsilon = 0$ one observes that the previous expression holds trivially, whereas in the case $\varepsilon = 1$ one deduces that $\lambda_1^1 = 1$. As a consequence, equation (29) is equivalent to

$$\varepsilon (1 - \lambda_1^1) = 0.$$

Now, using the last equality and equaling to zero the coefficients

$$[d(F(\sigma^3)) - F(d\sigma^3)]_{1\bar{1}}, \quad [d(F(\sigma^4)) - F(d\sigma^4)]_{1\bar{1}}, \quad [d(F(\sigma^4)) - F(d\sigma^4)]_{1\bar{2}}, \quad [d(F(\sigma^4)) - F(d\sigma^4)]_{2\bar{2}},$$

we find the following system of four equations

$$(30) \quad \begin{cases} \varepsilon (1 - \lambda_1^1) = 0, & \mu (1 - \lambda |\lambda_1^1|^2) = 0, \\ \nu (\lambda - |\lambda_1^1|^2) = 0, & \mu |\lambda_1^2|^2 - 2b' \Im (\lambda_1^2 \bar{\lambda}_1^1) - 2 \Im (\lambda_1^3 \bar{\lambda}_1^1) = 0, \end{cases}$$

together with the expressions

$$(31) \quad a' = \frac{1}{|\lambda_1^1|^2} \left(a \frac{\lambda}{\bar{\lambda}_1^1} + 2i \varepsilon \Im (\lambda_1^2 \bar{\lambda}_1^1) \right) \quad \text{and} \quad b' = \frac{1}{|\lambda_1^1|^2} (b + 2\mu \Im (\lambda_1^2 \bar{\lambda}_1^1)).$$

Recall that in Family II, $(\varepsilon, \mu) \neq (0, 0)$ and $\mu\nu = 0$ (see Proposition 3.19). Hence, there are four possible triples (ε, μ, ν) , namely: $(0, 1, 0)$, $(1, 0, 0)$, $(1, 0, 1)$, $(1, 1, 0)$.

We start with the last two triples. Observe that $(\varepsilon, \mu, \nu) = (1, 0, 1)$ and $(1, 1, 0)$ satisfy $\varepsilon = 1$ and $(\nu, \mu) \neq (0, 0)$. Hence, the first three equations in (30) imply that $\lambda_1^1 = 1$ and $\lambda = 1$. Substituting these values in (31), we get

$$a' = a + 2i \Im \lambda_1^2, \quad b' = b + 2\mu \Im \lambda_1^2.$$

Since a' is a real number, $\Im \lambda_1^2 = 0$ and thus $a' = a, b' = b$. This gives us part (iii) of the statement.

Let us now find the desired relations for the remaining cases.

First, we consider the values $(\varepsilon, \mu, \nu) = (1, 0, 0)$. From (30), we get that $\lambda_1^1 = 1$ and λ is a free parameter. Furthermore,

$$a' = \lambda a + 2i \Im \lambda_1^2, \quad b' = b.$$

We must choose λ_1^2 satisfying $\Im \lambda_1^2 = 0$ to ensure that $a' \in \mathbb{R}$. The last equation in (30) then gives us $\Im \lambda_1^3 = 0$. In particular, we get part (i) of the proposition.

Finally, we consider $(\varepsilon, \mu, \nu) = (0, 1, 0)$. From (30), one immediately sees that $\lambda = \frac{1}{|\lambda_1^1|^2}$, and only the last equation remains unsolved. Moreover, the expressions (31) turn into

$$a' = \frac{a}{|\lambda_1^1|^4 \lambda_1^1} \quad \text{and} \quad b' = \frac{1}{|\lambda_1^1|^2} (b + 2 \Im(\lambda_1^2 \bar{\lambda}_1^1)).$$

Just observe that we can choose any $\lambda_1^1 = \kappa \in \mathbb{R}^*$ to get $a' = \frac{a}{\kappa^5} \in \mathbb{R}$. After fixing any λ_1^2 we can also obtain a value for λ_1^3 from the last equation in (30). This gives us the equivalence in part (ii) of the statement. \square

We notice that different values of (a, b) and (a', b') are related by either the parameter λ or the parameters κ, λ_1^2 . Therefore, these will provide the equivalences between different complex structures.

Remark 3.23. *As a consequence of the previous proof, any isomorphism F relating J and J' follows (16) and satisfies Lemma 3.20 together with*

$$\lambda_4^4 = \lambda \in \mathbb{R}^*, \quad \lambda_2^2 = \lambda \lambda_1^1, \quad \lambda_3^3 = \frac{\lambda}{\lambda_1^1}, \quad \lambda_2^3 = i \mu \lambda \lambda_1^2,$$

and

$$\begin{cases} \lambda_1^1 = 1, & \Im \lambda_1^2 = \Im \lambda_1^3 = 0, & \text{if } (\varepsilon, \mu, \nu) = (1, 0, 0), \\ \lambda_1^1 = \kappa \in \mathbb{R}^*, & \lambda = \frac{1}{\kappa^2}, \quad \Im \lambda_1^3 = \frac{1}{\kappa} \left(\frac{1}{2} |\lambda_1^2|^2 - \frac{b}{\kappa} \Im \lambda_1^2 - 2 (\Im \lambda_1^2)^2 \right), & \text{if } (\varepsilon, \mu, \nu) = (0, 1, 0). \end{cases}$$

It can be seen that these values make condition (17) hold trivially for every $1 \leq i \leq 4$.

Finally we give the main result about equivalences of complex structures in Family II:

Theorem 3.24. *Up to equivalence, the complex structures*

$$\begin{cases} d\omega^1 = 0, \\ d\omega^2 = \omega^{14} + \omega^{1\bar{4}}, \\ d\omega^3 = a \omega^{1\bar{1}} + \varepsilon (\omega^{12} + \omega^{1\bar{2}} - \omega^{2\bar{1}}) + i \mu (\omega^{24} + \omega^{2\bar{4}}), \\ d\omega^4 = i \nu \omega^{1\bar{1}} - \mu \omega^{2\bar{2}} + i b (\omega^{1\bar{2}} - \omega^{2\bar{1}}) + i (\omega^{1\bar{3}} - \omega^{3\bar{1}}), \end{cases}$$

are determined by the tuples $(\varepsilon, \mu, \nu, a, b)$, where

- (i) $(\varepsilon, \mu, \nu, a, b) = (1, 1, 0, a, b)$;
- (ii) $(\varepsilon, \mu, \nu, a, b) = (1, 0, 1, a, b)$;
- (iii) $(\varepsilon, \mu, \nu, a, b) = (1, 0, 0, 0, b), (1, 0, 0, 1, b)$;
- (iv) $(\varepsilon, \mu, \nu, a, b) = (0, 1, 0, 0, 0), (0, 1, 0, 1, 0)$.

Proof. We first observe that parts (i) and (ii) of the theorem come straightforward from Proposition 3.22. Moreover, this proposition also gives us the relation between the pairs (a, b) and (a', b') when there is an equivalence between the complex structures J and J' .

For $(\varepsilon, \mu, \nu) = (1, 0, 0)$, observe that one has

$$a' = \lambda a, \quad b' = b \quad \text{with} \quad \lambda \in \mathbb{R}^*.$$

It suffices to set the values $\lambda = \begin{cases} \frac{1}{a}, & \text{if } a \neq 0, \\ 1, & \text{if } a = 0, \end{cases}$ and $\lambda_1^2 = \lambda_1^3 = 0$ in Remark 3.23 to obtain part (iii) of our theorem.

For $(\varepsilon, \mu, \nu) = (0, 1, 0)$, we have

$$a' = \frac{a}{\kappa^5} \quad \text{and} \quad b' = \frac{1}{\kappa^2} (b + 2 \kappa \Im \lambda_1^2), \quad \text{with} \quad \kappa \in \mathbb{R}^*.$$

One can take

$$\begin{cases} \lambda = \frac{1}{a^{2/5}}, & \kappa = a^{1/5}, & \text{if } a \neq 0, \\ \lambda = 1, & \kappa = 1, & \text{if } a = 0, \end{cases} \quad \text{and} \quad \lambda_1^2 = -\frac{ib}{2\kappa}, \quad \lambda_1^3 = \frac{ib^2}{8\kappa^3},$$

in Remark 3.23 to get $b' = 0$ and $a' \in \{0, 1\}$ in this case. This concludes our proof. \square

3.3.2. *Ascending central series.* Here, we explicitly compute the ascending central series of the nilpotent Lie algebras underlying Family II.

Proposition 3.25. *Let \mathfrak{g} be an 8-dimensional NLA endowed with a complex structure J in Family II with equations given in Proposition 3.19. Then, the ascending type of \mathfrak{g} is one of the following:*

- (i) $(\dim \mathfrak{g}_k)_k = (1, 3, 5, 8)$ if and only if $\nu = 0$;
- (ii) $(\dim \mathfrak{g}_k)_k = (1, 3, 5, 6, 8)$ if and only if $\nu = 1$.

Proof. Let us consider a generic element $X \in \mathfrak{g}$ given by (24). Using the structure equations in Proposition 3.19, one has the following brackets:

$$\begin{aligned}
 (32) \quad [X, Z_1] &= (\alpha_4 + \bar{\alpha}_4)Z_2 + (a\bar{\alpha}_1 + \varepsilon(\alpha_2 + \bar{\alpha}_2))Z_3 - (a\bar{\alpha}_1 - \varepsilon\bar{\alpha}_2)\bar{Z}_3 \\
 &\quad + i\nu\bar{\alpha}_1(Z_4 + \bar{Z}_4) + i(b\bar{\alpha}_2 + \bar{\alpha}_3)(Z_4 - \bar{Z}_4), \\
 [X, Z_2] &= -(\varepsilon(\alpha_1 + \bar{\alpha}_1) - i\mu(\alpha_4 + \bar{\alpha}_4))Z_3 - \varepsilon\bar{\alpha}_1\bar{Z}_3 - (ib\bar{\alpha}_1 + \mu\bar{\alpha}_2)(Z_4 - \bar{Z}_4), \\
 [X, Z_3] &= -i\bar{\alpha}_1(Z_4 - \bar{Z}_4), \\
 [X, Z_4] &= -\alpha_1 Z_2 - \bar{\alpha}_1 \bar{Z}_2 - i\mu\alpha_2 Z_3 + i\mu\bar{\alpha}_2 \bar{Z}_3.
 \end{aligned}$$

Since $[X, Z_4 - \bar{Z}_4] = 0$ for all $X \in \mathfrak{g}$, it is straightforward that $\mathfrak{g}_1 = \langle \Im Z_4 \rangle$, due to Corollary 2.6. To establish \mathfrak{g}_2 we must impose that X satisfies $[X, Z_k] \in \mathfrak{g}_1$ for any k . From (32), this is equivalent to

$$\alpha_1 = 0, \quad \varepsilon\alpha_2 = 0, \quad \mu\alpha_2 = 0, \quad \alpha_4 + \bar{\alpha}_4 = 0.$$

Since $(\varepsilon, \mu) \neq (0, 0)$, we get $\alpha_1 = \alpha_2 = 0$ and $\alpha_4 = i\alpha$. Therefore, $\dim \mathfrak{g}_2 = 3$ and

$$\mathfrak{g}_2 = \langle \Re Z_3, \Im Z_3, \Im Z_4 \rangle.$$

Using a similar argument for \mathfrak{g}_3 one concludes that $\alpha_1 = 0$ and $\alpha_4 = i\alpha$, so $\dim \mathfrak{g}_3 = 5$ and

$$\mathfrak{g}_3 = \langle \Re Z_2, \Im Z_2, \Re Z_3, \Im Z_3, \Im Z_4 \rangle.$$

Finally, the conditions for $X \in \mathfrak{g}_4$ reduce to $\nu\alpha_1 = 0$. It is then clear that $\nu = 0$ implies $\mathfrak{g}_4 = \mathfrak{g}$, whereas $\nu = 1$ leads to a 5-step nilpotent Lie algebra of ascending type $(1, 3, 5, 6, 8)$. \square

The combination of this result and Theorem 3.24 gives part (ii) of Theorem 3.1.

Furthermore, Propositions 3.15 and 3.25 show that the NLAs underlying Families I and II do not share the same ascending type. Since two complex structures can only be equivalent if they are defined on the same NLA, we conclude the following:

Corollary 3.26. *There are no equivalences between complex structures parametrized by Family I and complex structures parametrized by Family II.*

As a consequence, our two families provide a partition of the space of SnN complex structures on 8-dimensional NLAs. This completes the proof of Theorem 3.1.

4. CLASSIFICATION OF NLAs UNDERLYING FAMILY I

In this section we determine the real Lie algebras underlying Family I. In particular, we classify up to isomorphism the 8-dimensional NLAs endowed with an SnN complex structure defined by Family I. Recall that this family is given by the complex structure equations (5) with parameters in Table 1. The concrete correspondence between (5) and the real structure equations of the Lie algebras can be found in Appendix A. In the end, we are able to prove the following result:

Theorem 4.1. *The non-isomorphic real Lie algebras underlying Family I are*

$$\begin{aligned}
\mathfrak{n}_1^\gamma &= (0^5, 13 + 15 + 24, 14 - 23 + 25, 16 + 27 + \gamma \cdot 34), \text{ where } \gamma \in \{0, 1\}, \\
\mathfrak{n}_2^\alpha &= (0^4, 12, 13 + 15 + 24, 14 - 23 + 25, 16 + 27 + \alpha \cdot 34), \text{ with } \alpha \in \mathbb{R}, \\
\mathfrak{n}_3^\gamma &= (0^4, 12, 13 + 25 + \gamma \cdot 15, 15 + 24 + \gamma \cdot 25, 16 + 27), \text{ where } \gamma \in \{0, 1\}, \\
\mathfrak{n}_4^{\eta, \vartheta} &= (0^4, 12, 15 + (1 + \eta) \cdot 24, (\eta - 1) \cdot 14 - 23 + (\vartheta - 1) \cdot 25, 16 + 27 + 34 - 2 \cdot 45), \\
&\text{ where } (\eta, \vartheta) \in \mathbb{R}^* \times \mathbb{R}^{>0} \text{ or } \mathbb{R}^{>0} \times \{0\}, \\
\mathfrak{n}_5 &= (0^4, 2 \cdot 12, 14 - 23, 13 + 24, 16 + 27 + 35), \\
\mathfrak{n}_6 &= (0^4, 2 \cdot 12, 14 + 15 - 23, 13 + 24 + 25, 16 + 27 + 35), \\
\mathfrak{n}_7 &= (0^5, 15, 25, 16 + 27 + 34), \\
\mathfrak{n}_8 &= (0^4, 12, 15, 25, 16 + 27 + 34).
\end{aligned}$$

Moreover, their ascending types are listed in Table 3.

NLA	\mathfrak{n}_1^γ	\mathfrak{n}_2^α	\mathfrak{n}_3^γ	$\mathfrak{n}_4^{\eta, \vartheta}$	\mathfrak{n}_5	\mathfrak{n}_6	\mathfrak{n}_7	\mathfrak{n}_8
$(\dim \mathfrak{g}_k)_k$	(1, 3, 8)	(1, 3, 6, 8)		(1, 4, 8)	(1, 4, 6, 8)	(1, 5, 8)	(1, 5, 6, 8)	

TABLE 3. Ascending type of real Lie algebras underlying Family I

Remark 4.2. *This result proves the first part of Theorem 1.2. Concretely, the relation between the Lie algebras \mathfrak{g} 's and \mathfrak{n} 's is the following: $\mathfrak{g}_1^\gamma = \mathfrak{n}_1^\gamma$, $\mathfrak{g}_2^\alpha = \mathfrak{n}_2^\alpha$, $\mathfrak{g}_3^\gamma = \mathfrak{n}_3^\gamma$, $\mathfrak{g}_4^{\alpha, \beta} = \mathfrak{n}_4^{\eta, \vartheta}$, and $\mathfrak{g}_k = \mathfrak{n}_k$ for $k = 5, \dots, 8$.*

Notation 4.3. *For the description of the structure of the Lie algebras in Theorem 4.1 we are using the following abbreviated notation. For instance,*

$$\mathfrak{n}_1^\gamma = (0^5, 13 + 15 + 24, 14 - 23 + 25, 16 + 27 + \gamma \cdot 34)$$

means that there is a basis $\{e^j\}_{j=1}^8$ of the dual $(\mathfrak{n}_1^\gamma)^*$ of the nilpotent Lie algebra \mathfrak{n}_1^γ satisfying $de^1 = \dots = de^5 = 0$, $de^6 = e^1 \wedge e^3 + e^1 \wedge e^5 + e^2 \wedge e^4$, $de^7 = e^1 \wedge e^4 - e^2 \wedge e^3 + e^2 \wedge e^5$, and $de^8 = e^1 \wedge e^6 + e^2 \wedge e^7 + \gamma e^3 \wedge e^4$.

We first recall that NLAs having different ascending types are non-isomorphic. Hence, to prove our result, it suffices to analyze the NLAs underlying Family I within each of the different ascending types in Table 1.

Along our discussion, we will sometimes make use of the following invariants associated to NLAs:

- Descending type $(\dim \mathfrak{g}^k)_k$: Recall that the *descending central series* $\{\mathfrak{g}^k\}_{k \geq 0}$ of a Lie algebra \mathfrak{g} is defined by $\mathfrak{g}^0 = \mathfrak{g}$ and $\mathfrak{g}^k = [\mathfrak{g}^{k-1}, \mathfrak{g}]$, for any $k \geq 1$. When \mathfrak{g} is an s -step NLA, then $\mathfrak{g}_s = \{0\}$ and we can associate an s -tuple to \mathfrak{g} , namely,

$$(m^1, \dots, m^{s-1}, m^s) := (\dim \mathfrak{g}^1, \dots, \dim \mathfrak{g}^{s-1}, \dim \mathfrak{g}^s)$$

which strictly decreases, i.e. $2n > m^1 > \dots > m^{s-1} > m^s = 0$. We will say that $(\dim \mathfrak{g}^k)_k = (m^1, \dots, m^s)$ is the *descending type* of \mathfrak{g} .

- Betti numbers $b_k(\mathfrak{g})$: The *Chevalley-Eilenberg cohomology groups* of an m -dimensional Lie algebra \mathfrak{g} are naturally defined by

$$H_{\text{dR}}^k(\mathfrak{g}; \mathbb{R}) = \frac{\ker\{d : \bigwedge^k(\mathfrak{g}^*) \longrightarrow \bigwedge^{k+1}(\mathfrak{g}^*)\}}{\text{im}\{d : \bigwedge^{k-1}(\mathfrak{g}^*) \longrightarrow \bigwedge^k(\mathfrak{g}^*)\}},$$

where $1 \leq k \leq m$. The dimensions of these groups are known as the Betti numbers, i.e., $b_k(\mathfrak{g}) = \dim H_{\text{dR}}^k(\mathfrak{g}; \mathbb{R})$.

- Number of functionally independent generalized Casimir operators $n_I(\mathfrak{g})$: Let \mathfrak{g} be an m -dimensional real Lie algebra with basis $\{x_k\}_{k=1}^m$ and $[x_i, x_j] = \sum_{k=1}^m c_{ij}^k x_k$. The vectors

$$(33) \quad \hat{X}_k = \sum_{i,j=1}^m c_{ki}^j x_j \frac{\partial}{\partial x_i}$$

generate a basis of the coadjoint representation of \mathfrak{g} . One can construct a matrix C by rows from the coefficients of these vectors, and then $n_I(\mathfrak{g}) = m - \text{rank } C$. For further details, we refer to [31].

Let us now start the proof of Theorem 4.1. To clarify the presentation, we divide the discussion into different sections depending on the ascending type of the underlying NLAs.

4.1. NLAs with $\dim \mathfrak{g}_2 \geq 4$. In this section we study isomorphisms of NLAs for each one of the ascending types in Table 1 satisfying $\dim \mathfrak{g}_2 \geq 4$. We first focus on the case $(\dim \mathfrak{g}_k)_k = (1, 4, 8)$.

Proposition 4.4. *If \mathfrak{g} is an 8-dimensional NLA with ascending type $(1, 4, 8)$ admitting a complex structure, then $\mathfrak{g} \cong \mathfrak{n}_5$.*

Proof. Let \mathfrak{g} be an 8-dimensional NLA with ascending type $(1, 4, 8)$ admitting a complex structure J . By Theorem 3.1, the pair (\mathfrak{g}, J) is determined by the complex structure equations (5) with parameters $\varepsilon = \nu = 1$, $a = 0$, $b = 2\delta$, and $\delta = \pm 1$, i.e.,

$$d\omega^1 = 0, \quad d\omega^2 = \omega^{1\bar{1}}, \quad d\omega^3 = \omega^{14} + \omega^{1\bar{4}} + 2i\omega^{1\bar{2}}, \quad d\omega^4 = i\omega^{1\bar{1}} + 2\delta\omega^{2\bar{2}} + i\delta(\omega^{1\bar{3}} - \omega^{3\bar{1}}).$$

Considering the real basis $\{e^i\}_{i=1}^8$ given by:

$$\omega^1 = \delta e^1 - i e^2, \quad \omega^2 = -\frac{\delta}{2}(e^3 - 2i e^5), \quad \omega^3 = \delta e^6 - i e^7, \quad \omega^4 = 2\delta \left(\frac{\delta}{4} e^4 - \frac{1}{2} e^5 + i e^8 \right),$$

one directly attains the structure equations for \mathfrak{n}_5 in Theorem 4.1. \square

Observe that, according to Table 1, there are only two non-isomorphic complex structures for $(\dim \mathfrak{g}_k)_k = (1, 4, 8)$. Consequently, we have:

Corollary 4.5. *The Lie algebra \mathfrak{n}_5 admits two complex structures up to equivalence.*

We next move to the study of the Lie algebras underlying Family I with ascending type $(1, 4, 6, 8)$.

Proposition 4.6. *If \mathfrak{g} is an 8-dimensional NLA with ascending type $(1, 4, 6, 8)$ admitting a complex structure, then $\mathfrak{g} \cong \mathfrak{n}_6$.*

Proof. Let \mathfrak{g} be an 8-dimensional NLA with ascending type $(1, 4, 6, 8)$ that admits a complex structure J . By Theorem 3.1, the pair (\mathfrak{g}, J) is given by the complex structure equations (5) where $\varepsilon = 1$, $(\nu, b) = (0, 1)$ or $(1, b)$ with $b \neq 0$, 2δ , $a = 0$, and $\delta = \pm 1$, i.e.,

$$d\omega^1 = 0, \quad d\omega^2 = \omega^{1\bar{1}}, \quad d\omega^3 = \omega^{14} + \omega^{1\bar{4}} + i\delta b \omega^{1\bar{2}}, \quad d\omega^4 = i\nu \omega^{1\bar{1}} + b\omega^{2\bar{2}} + i\delta(\omega^{1\bar{3}} - \omega^{3\bar{1}}).$$

Defining the real basis $\{e^i\}_{i=1}^8$ as

$$\begin{aligned} \omega^1 &= \delta e^1 - i e^2, & \omega^3 &= (b - 2\delta\nu)(\delta e^6 - i e^7), \\ \omega^2 &= \left(\frac{2\delta\nu}{b} - 1 \right) e^3 + i\delta e^5, & \omega^4 &= \left(\frac{b}{2} - \delta\nu \right) e^4 - \delta\nu e^5 + 2\delta(b - 2\delta\nu) i e^8, \end{aligned}$$

one obtains the structure equations for the Lie algebra \mathfrak{n}_6 in Theorem 4.1. \square

Moreover, we deduce the following:

Corollary 4.7. *There exists an infinite family of non-equivalent complex structures on the Lie algebra \mathfrak{n}_6 .*

We now consider the NLAs underlying Family I with ascending type $(1, 5, 8)$.

Proposition 4.8. *If \mathfrak{g} is an 8-dimensional NLA with ascending type $(1, 5, 8)$ admitting a complex structure, then $\mathfrak{g} \cong \mathfrak{n}_7$.*

Proof. Let \mathfrak{g} be an 8-dimensional NLA with ascending type $(1, 5, 8)$ endowed with a complex structure J . Applying Theorem 3.1, we know that the pair (\mathfrak{g}, J) has complex structure equations (5) with parameters $\varepsilon = \nu = a = 0$, $b = 1$, and $\delta = \pm 1$:

$$d\omega^1 = d\omega^2 = 0, \quad d\omega^3 = \omega^{14} + \omega^{1\bar{4}}, \quad d\omega^4 = \omega^{2\bar{2}} + i\delta(\omega^{1\bar{3}} - \omega^{3\bar{1}}).$$

One directly obtains the structure equations for \mathfrak{n}_7 in Theorem 4.1 defining the real basis $\{e^i\}_{i=1}^8$ as follows:

$$\omega^1 = \delta e^1 - i e^2, \quad \omega^2 = -e^3 + i \delta e^4, \quad \omega^3 = \delta e^6 - i e^7, \quad \omega^4 = \frac{1}{2}e^5 + 2\delta i e^8.$$

□

This leads to the following result:

Corollary 4.9. *The Lie algebra \mathfrak{n}_7 has, up to equivalence, only two complex structures.*

We finally study the Lie algebras with ascending type $(1, 5, 6, 8)$, which are the remaining ones for which $\dim \mathfrak{g}_2 \geq 4$.

Proposition 4.10. *If \mathfrak{g} is an 8-dimensional NLA with ascending type $(1, 5, 6, 8)$ admitting a complex structure, then $\mathfrak{g} \cong \mathfrak{n}_8$.*

Proof. Let \mathfrak{g} be an 8-dimensional NLA with ascending type $(1, 5, 6, 8)$ endowed with a complex structure J . Applying Theorem 3.1, the complex structure equations of (\mathfrak{g}, J) follow (5) with parameters $\varepsilon = 0$, $\nu = 1$, $a = 0$, $b \in \{-1, 1\}$, and $\delta = \pm 1$, namely,

$$d\omega^1 = d\omega^2 = 0, \quad d\omega^3 = \omega^{14} + \omega^{1\bar{4}}, \quad d\omega^4 = i\omega^{1\bar{1}} + b\omega^{2\bar{2}} + i\delta(\omega^{1\bar{3}} - \omega^{3\bar{1}}).$$

If we define the real basis $\{e^i\}_{i=1}^8$ as

$$\omega^1 = \delta e^1 - i e^2, \quad \omega^2 = -(b e^3 + 4 i e^4), \quad \omega^3 = -4(e^6 - i \delta e^7), \quad \omega^4 = -2(\delta e^5 + 4 i e^8),$$

then we directly reach the structure equations for \mathfrak{n}_8 in Theorem 4.1. □

As a consequence, we can ensure the following:

Corollary 4.11. *There are exactly four non-equivalent complex structures on the Lie algebra \mathfrak{n}_8 .*

This finishes the study of the non-isomorphic NLAs underlying Family I that satisfy $\dim \mathfrak{g}_2 \geq 4$.

4.2. NLAs with $\dim \mathfrak{g}_2 = 3$ and nilpotency step 3. We now focus on those NLAs underlying Family I whose ascending type is $(1, 3, 8)$. Following Table 1, this leads us to consider the complex structure equations (5) with parameters $\varepsilon = \nu = 0$, $a = 1$, $b \in \{0, 1\}$ and $\delta \in \{-1, 1\}$, namely,

$$(34) \quad d\omega^1 = d\omega^2 = 0, \quad d\omega^3 = \omega^{14} + \omega^{1\bar{4}} + \omega^{2\bar{1}}, \quad d\omega^4 = b\omega^{2\bar{2}} + i\delta(\omega^{1\bar{3}} - \omega^{3\bar{1}}).$$

Observe that one finds four different complex structures within this set of equations. We next see that there are two algebras underlying the family above:

Proposition 4.12. *If \mathfrak{g} is an 8-dimensional NLA with ascending type $(1, 3, 8)$ admitting a complex structure, then $\mathfrak{g} \cong \mathfrak{n}_1^\gamma$.*

Proof. From the $(1, 0)$ -basis $\{\omega^k\}_{k=1}^4$ determined by (34), define a real basis $\{e^k\}_{k=1}^8$ as follows:

$$\omega^1 = \delta e^1 - i e^2, \quad \omega^2 = -e^3 + i \delta e^4, \quad \omega^3 = \delta e^6 - i e^7, \quad \omega^4 = \frac{1}{2}e^5 + 2\delta i e^8.$$

One then directly reaches \mathfrak{n}_1^γ , with $\gamma = b \in \{0, 1\}$, following Theorem 4.1. □

It remains to show that the Lie algebras \mathfrak{n}_1^0 and \mathfrak{n}_1^1 are not isomorphic. Although one can check that their descending and ascending central series coincide, the result comes as a direct consequence of their Casimir invariants. More precisely:

Lemma 4.13. *Let $\mathfrak{n}_1^\gamma = (0^5, 13 + 15 + 24, 14 - 23 + 25, 16 + 27 + \gamma \cdot 34)$, with $\gamma \in \{0, 1\}$. Then, $n_I(\mathfrak{n}_1^0) = 4$ and $n_I(\mathfrak{n}_1^1) = 2$. Therefore, \mathfrak{n}_1^0 and \mathfrak{n}_1^1 are not isomorphic.*

Proof. Using the equations of \mathfrak{n}_1^γ and the well-known formula $de(X, Y) = -e([X, Y])$, for $e \in \mathfrak{g}^*$ and $X, Y \in \mathfrak{g}$, it is easy to see that the matrix C constructed from the coefficients of the vectors (33) is

$$C = \begin{pmatrix} 0 & 0 & -x_6 & -x_7 & -x_6 & -x_8 & 0 & 0 \\ 0 & 0 & x_7 & -x_6 & -x_7 & 0 & -x_8 & 0 \\ x_6 & -x_7 & 0 & -\gamma x_8 & 0 & 0 & 0 & 0 \\ x_7 & x_6 & \gamma x_8 & 0 & 0 & 0 & 0 & 0 \\ x_6 & x_7 & 0 & 0 & 0 & 0 & 0 & 0 \\ x_8 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & x_8 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

One can check that the minors of orders 8 and 7 are all equal to zero. For those of order 6, one obtains the following expressions:

$$\gamma^2 x_7^2 x_8^4, \quad -\gamma^2 x_6 x_7 x_8^4, \quad -\gamma^2 x_7 x_8^5, \quad \gamma^2 x_6^2 x_8^4, \quad \gamma^2 x_6 x_8^5, \quad \gamma^2 x_8^6.$$

Consequently, if $\gamma = 1$ then $\text{rank } C = 6$ and $n_I = 2$ for the algebra \mathfrak{n}_1 . However, if $\gamma = 0$ then all the previous expressions vanish, and it is possible to see that $\text{rank } C = 4$, thus $n_I = 4$ for \mathfrak{n}_0 . This gives our result. \square

This concludes the proof of Theorem 4.1 for \mathfrak{n}_1^γ , with $\gamma \in \{0, 1\}$. In particular, notice that there are exactly two NLAs with ascending type $(1, 3, 8)$ admitting complex structures. In fact, the following holds:

Corollary 4.14. *There are exactly two non-equivalent complex structures on each of the Lie algebras \mathfrak{n}_1^γ , where $\gamma \in \{0, 1\}$.*

4.3. NLAs with $\dim \mathfrak{g}_2 = 3$ and nilpotency step 4. We finally consider those NLAs underlying Family I whose ascending type is $(1, 3, 6, 8)$. According to Table 1, there are several possible values for the parameters in the complex structure equations (5):

	ε	ν	a	b
$(1, 3, 6, 8)$	1	0	1	$b \geq 0$
	0	1		$b \in \mathbb{R}$
	1	1	$a > 0$	

TABLE 4. Lie algebras for case $(1, 3, 6, 8)$

We separate different cases depending on the values given in Table 4.

Lemma 4.15. *Let \mathfrak{g} be an NLA with ascending type $(1, 3, 6, 8)$ admitting a complex structure J such that (\mathfrak{g}, J) is parametrized, up to equivalence, by Family I with $\varepsilon = 0$. Then $\mathfrak{g} \cong \mathfrak{n}_2^\alpha$, with $\alpha \in \mathbb{R}$.*

Proof. First notice that the choice $\varepsilon = 0$ implies $\nu = a = 1$ and $b \in \mathbb{R}$ (see Table 4). It suffices to define the real basis

$$\omega^1 = \delta e^1 - i e^2, \quad \omega^2 = 4(\delta e^3 - i e^4), \quad \omega^3 = -4(e^6 - i \delta e^7), \quad \omega^4 = -2(\delta e^5 + 4i e^8),$$

and then one immediately reaches the algebra \mathfrak{n}_2^α in Theorem 4.1 with $\alpha = -4\delta b \in \mathbb{R}$. \square

Lemma 4.16. *Let \mathfrak{g} be an NLA with ascending type $(1, 3, 6, 8)$ admitting a complex structure J such that (\mathfrak{g}, J) is parametrized, up to equivalence, by Family I with $\varepsilon = 1$ and $b = 0$. Then $\mathfrak{g} \cong \mathfrak{n}_3^\gamma$, with $\gamma \in \{0, 1\}$, or $\mathfrak{g} \cong \mathfrak{n}_2^0$.*

Proof. From the $(1, 0)$ -basis $\{\omega^k\}_{k=1}^4$ determined by (34) with parameters $\varepsilon = 1$ and $b = 0$, we define a real basis $\{e^k\}_{k=1}^8$ by:

$$\begin{aligned} \omega^1 &= \frac{1}{\sqrt{2}} (\delta e^1 - i e^2), & \omega^3 &= \frac{a}{\sqrt{2}} (\delta e^6 - i e^7), \\ \omega^2 &= -\frac{1}{2} (e^3 - e^4) + i \delta e^5, & \omega^4 &= \frac{a}{4} (e^3 + e^4) + \delta (-\nu e^5 + i a e^8). \end{aligned}$$

One then finds the structure equations of \mathfrak{n}_3^γ given in Theorem 4.1, namely,

$$(35) \quad \begin{aligned} de^1 = de^2 = de^3 = de^4 = 0, \quad de^5 = e^{12}, \quad de^6 = e^{13} + e^{25} + \gamma e^{15}, \\ de^7 = e^{15} + e^{24} + \gamma e^{25}, \quad de^8 = e^{16} + e^{27}, \end{aligned}$$

where $\gamma = -\frac{2\delta\nu}{a} \in \mathbb{R}$ (observe $a \neq 0$). We next reduce the values of γ to 0 or 1, as stated in the lemma.

First, we observe that $\mathfrak{n}_3^\gamma \cong \mathfrak{n}_3^{-\gamma}$. Indeed, for any \mathfrak{n}_3^γ with structure equations (35), it suffices to consider a new basis $\{v^k\}_{k=1}^8$ with $v^k = -e^k$ for $k = 2, 5, 7$, and $v^k = e^k$ for $k \neq 2, 5, 7$, to check the isomorphism above. As a consequence, we can assume $\gamma \geq 0$ in (35).

Next, we notice that $\mathfrak{n}_3^\gamma \cong \mathfrak{n}_3^0$, for each $\gamma \in (0, 1)$. Indeed, for a nilpotent Lie algebra defined by (35) with $\gamma \in (0, 1)$, one can check that the following basis gives the desired isomorphism:

$$\begin{aligned} v^1 &= \frac{\sqrt{1-\gamma^2}}{2\sqrt{3}} \left((1 - \sqrt{1-\gamma^2}) e^1 + \gamma e^2 \right), \\ v^2 &= \frac{-(2 - \sqrt{3})\sqrt{1-\gamma^2}}{2\sqrt{3}} \left((1 + \sqrt{1-\gamma^2}) e^1 + \gamma e^2 \right), \\ v^3 &= -\frac{(7 - 4\sqrt{3})\sqrt{1-\gamma^2}}{12} \left(\gamma^2 e^3 + (2 - \gamma^2 + 2\sqrt{1-\gamma^2}) e^4 \right), \\ v^4 &= -\frac{\sqrt{1-\gamma^2}}{12} \left(\gamma^2 e^3 + (2 - \gamma^2 - 2\sqrt{1-\gamma^2}) e^4 \right), \\ v^5 &= -\frac{\gamma(2 - \sqrt{3})\sqrt{1-\gamma^2}}{12} \left(\gamma(e^3 + e^4) - 2(1 - \gamma^2)e^5 \right), \\ v^6 &= \frac{\gamma(1 - \gamma^2)^{3/2}(7 - 4\sqrt{3})}{12\sqrt{3}} \left(\gamma e^6 - (1 + \sqrt{1-\gamma^2}) e^7 \right), \\ v^7 &= \frac{\gamma(1 - \gamma^2)^{3/2}(2 - \sqrt{3})}{12\sqrt{3}} \left(\gamma e^6 - (1 - \sqrt{1-\gamma^2}) e^7 \right), \\ v^8 &= -\frac{\gamma^2(1 - \gamma^2)^{5/2}(7 - 4\sqrt{3})}{36} e^8. \end{aligned}$$

Finally, we remark that $\mathfrak{n}_3^\gamma \cong \mathfrak{n}_2^0$, for $\gamma \in (1, \infty)$, where \mathfrak{n}_2^0 belongs to the family found in Lemma 4.15. To prove this, it suffices to consider (35) with $\gamma > 1$ and construct the following basis:

$$\begin{aligned} v^1 &= \sqrt{6(\gamma^2 - 1)}(e^1 + \gamma e^2), & v^5 &= 3\gamma\sqrt{\gamma^2 - 1}(\gamma e^3 + \gamma e^4 + 2(\gamma^2 - 1)e^5), \\ v^2 &= -\sqrt{6}(\gamma^2 - 1)e^1, & v^6 &= 6\sqrt{6}\gamma(\gamma^2 - 1)^2 e^7, \\ v^3 &= -3\sqrt{\gamma^2 - 1}(\gamma^2 e^3 - (\gamma^2 - 2)e^4), & v^7 &= -6\sqrt{6}\gamma(\gamma^2 - 1)^{3/2}(\gamma e^6 - e^7), \\ v^4 &= 6(\gamma^2 - 1)e^4, & v^8 &= 36\gamma^2(\gamma^2 - 1)^{5/2}e^8. \end{aligned}$$

The previous discussion leaves $\gamma = 0$ and $\gamma = 1$ as the only values to be considered. \square

Lemma 4.17. *Let \mathfrak{g} be an NLA with ascending type $(1, 3, 6, 8)$ admitting a complex structure J such that (\mathfrak{g}, J) is parametrized, up to equivalence, by Family I with $\varepsilon = 1$ and $b \neq 0$. Then $\mathfrak{g} \cong \mathfrak{n}_4^{\eta, \vartheta}$, with $(\eta, \vartheta) \in \mathbb{R}^* \times \mathbb{R}^{>0}$ or $(\eta, \vartheta) \in \mathbb{R}^{>0} \times \{0\}$.*

Proof. Let us observe that the pair (\mathfrak{g}, J) is determined by the complex structure equations (34) with parameters $\varepsilon = 1$ and $b \neq 0$. Since this implies $a > 0$ (see Table 4), we can define a real basis $\{\tilde{e}^k\}_{k=1}^8$ for \mathfrak{g} as follows:

$$\omega^1 = \delta \tilde{e}^1 - i \tilde{e}^2, \quad \omega^2 = -\tilde{e}^3 + i \delta \tilde{e}^5, \quad \omega^3 = a(\delta \tilde{e}^6 - i \tilde{e}^7), \quad \omega^4 = \left(\frac{a}{2} \tilde{e}^4 - \nu \delta \tilde{e}^5\right) + 2\delta a i \tilde{e}^8.$$

Then, \mathfrak{g} is determined by the real structure equations

$$(36) \quad \begin{aligned} d\tilde{e}^1 = d\tilde{e}^2 = d\tilde{e}^3 = d\tilde{e}^4 = 0, \quad d\tilde{e}^5 = 2\tilde{e}^{12}, \quad d\tilde{e}^6 = \tilde{e}^{13} + \tilde{e}^{14} + \tilde{e}^{25} + \vartheta \tilde{e}^{15} - \rho \tilde{e}^{23}, \\ d\tilde{e}^7 = \tilde{e}^{15} - \tilde{e}^{23} + \tilde{e}^{24} + \rho \tilde{e}^{13} + \vartheta \tilde{e}^{25}, \quad d\tilde{e}^8 = \tilde{e}^{16} + \tilde{e}^{27} + \rho \tilde{e}^{35}, \end{aligned}$$

where $\rho = \frac{b}{a}$ and $\vartheta = \frac{1}{a}(b - 2\nu\delta)$. Notice that $\rho \neq 0$ since $b \neq 0$ by hypothesis.

Taking, for instance, $v^i = -\tilde{e}^i$ for $i = 2, 5, 7$ and $v^i = \tilde{e}^i$ for $i \neq 2, 5, 7$ in (36), one checks that the parameters (ϑ, ρ) and $(-\vartheta, -\rho)$ give rise to isomorphic Lie algebras. As a consequence, we can assume that $\vartheta \geq 0$ and let $\rho \in \mathbb{R} \setminus \{0\}$ in (36).

If we now consider the basis

$$\begin{aligned} e^1 &= -\sqrt{1+\vartheta}(\tilde{e}^1 + \tilde{e}^2), & e^3 &= \frac{-2}{\sqrt{1+\vartheta}}\tilde{e}^4, & e^5 &= \sqrt{1+\vartheta}\tilde{e}^5 + \frac{1}{\sqrt{1+\vartheta}}\tilde{e}^4, & e^7 &= \sqrt{1+\vartheta}(\tilde{e}^6 - \tilde{e}^7), \\ e^2 &= -\tilde{e}^2 + \tilde{e}^1, & e^4 &= -\rho\tilde{e}^3, & e^6 &= -\tilde{e}^6 - \tilde{e}^7, & e^8 &= 2\sqrt{1+\vartheta}\tilde{e}^8, \end{aligned}$$

then we attain the structure equations for the Lie algebra $\mathfrak{n}_4^{\eta, \vartheta}$ given in Theorem 4.1, with $\eta = \frac{1}{\rho} \in \mathbb{R} \setminus \{0\}$ and $\vartheta \geq 0$. To finish the proof, it remains to see that the case $\vartheta = 0$ implies $\eta > 0$. This can be done defining the following basis for $\vartheta = 0$ and $\eta < 0$:

$$v^1 = e^2, \quad v^2 = e^1, \quad v^3 = e^3, \quad v^4 = -e^4, \quad v^5 = -e^3 - e^5, \quad v^6 = e^7, \quad v^7 = e^6, \quad v^8 = e^8,$$

which shows that $\mathfrak{n}_4^{\eta, 0} \cong \mathfrak{n}_4^{-\eta, 0}$. □

To complete the proof of Theorem 4.1, it remains to show the following:

- the one-parameter family \mathfrak{n}_2^α , $\alpha \in \mathbb{R}$, contains no pairwise isomorphic NLAs;
- \mathfrak{n}_3^0 and \mathfrak{n}_3^1 are not isomorphic;
- the two-parameter family $\mathfrak{n}_4^{\eta, \vartheta}$, $(\eta, \vartheta) \in \mathbb{R}^* \times \mathbb{R}^{>0}$ or $\mathbb{R}^{>0} \times \{0\}$, has no pairwise isomorphic NLAs;
- if \mathfrak{g} and \mathfrak{g}' are two NLAs with ascending type $(1, 3, 6, 8)$ that belong to two different families of the ones given above, then $\mathfrak{g} \not\cong \mathfrak{g}'$.

In Table 5, we provide the descending type of the NLAs in the families \mathfrak{n}_2^α , \mathfrak{n}_3^γ , and $\mathfrak{n}_4^{\eta, \vartheta}$, as well as their number of functionally independent Casimir operators.

\mathfrak{g}	real parameter(s)	$\dim\{\mathfrak{g}^k\}_k$	$n_I(\mathfrak{g})$
\mathfrak{n}_2^α	$\alpha = 0$	$(4, 3, 1, 0)$	4
	$\alpha \neq 0$		2
\mathfrak{n}_3^γ	$\gamma = 0$	$(4, 3, 1, 0)$	4
	$\gamma = 1$	$(4, 2, 1, 0)$	
$\mathfrak{n}_4^{\eta, \vartheta}$	$\eta \neq 0, \vartheta = 1$	$(4, 2, 1, 0)$	2
	$\eta \neq 0, \vartheta \in (0, 1) \cup (1, \infty)$	$(4, 3, 1, 0)$	
	$\eta > 0, \vartheta = 0$		

TABLE 5. Some invariants for the NLAs with ascending central series $(1, 3, 6, 8)$

A direct consequence of these invariants is that some of the NLAs in the previous three families cannot be isomorphic (see Table 6 to check which invariant can be used in each case; the symbol “=” indicates that the NLAs under study are exactly the same, whereas the roman numbers mean that the two invariants coincide).

		\mathfrak{n}_2^α		\mathfrak{n}_3^γ		$\mathfrak{n}_4^{\eta, \vartheta}$		
		$\alpha = 0$	$\alpha \neq 0$	$\gamma = 0$	$\gamma = 1$	$\eta \in \mathbb{R} \setminus \{0\},$ $\vartheta = 1$	$\eta \in \mathbb{R} \setminus \{0\},$ $\vartheta \in (0, 1) \cup (1, \infty)$	$\eta > 0,$ $\vartheta = 0$
\mathfrak{n}_2^α	$\alpha = 0$	=	$n_I(\mathfrak{g})$	(i)	$\dim\{\mathfrak{g}^k\}_k$	$\dim\{\mathfrak{g}^k\}_k / n_I(\mathfrak{g})$	$n_I(\mathfrak{g})$	$n_I(\mathfrak{g})$
	$\alpha \neq 0$			(ii)	$n_I(\mathfrak{g})$	$n_I(\mathfrak{g})$	$\dim\{\mathfrak{g}^k\}_k$	(iii)
\mathfrak{n}_3^γ	$\gamma = 0$			=	$\dim\{\mathfrak{g}^k\}_k$	$\dim\{\mathfrak{g}^k\}_k / n_I(\mathfrak{g})$	$n_I(\mathfrak{g})$	$n_I(\mathfrak{g})$
	$\gamma = 1$				=	$n_I(\mathfrak{g})$	$\dim\{\mathfrak{g}^k\}_k / n_I(\mathfrak{g})$	$\dim\{\mathfrak{g}^k\}_k / n_I(\mathfrak{g})$
$\mathfrak{n}_4^{\eta, \vartheta}$	$\eta \neq 0,$ $\vartheta = 1$					(iv)	$\dim\{\mathfrak{g}^k\}_k$	$\dim\{\mathfrak{g}^k\}_k$
	$\eta \neq 0,$ $\vartheta \in (0, 1) \cup (1, \infty)$							(v)
	$\eta > 0,$ $\vartheta = 0$							

TABLE 6. Invariants for distinguishing NLAs with ascending central series (1, 3, 6, 8)

Consequently, to prove Theorem 4.1 it suffices to show the result below:

Proposition 4.18. *The following pairs of Lie algebras are not isomorphic:*

- (i) \mathfrak{n}_2^0 and \mathfrak{n}_3^0 .
- (ii) \mathfrak{n}_2^α and $\mathfrak{n}_2^{\alpha'}$ whenever $\alpha \neq \alpha'$ and $\alpha\alpha' \neq 0$.
- (iii) \mathfrak{n}_2^α and $\mathfrak{n}_4^{\eta, \vartheta}$ whenever $\alpha \neq 0$ and $\vartheta \neq 1$.
- (iv) $\mathfrak{n}_4^{\eta, 1}$ and $\mathfrak{n}_4^{\eta', 1}$ whenever $\eta \neq \eta'$.
- (v) $\mathfrak{n}_4^{\eta, \vartheta}$ and $\mathfrak{n}_4^{\eta', \vartheta'}$ whenever $(\eta, \vartheta) \neq (\eta', \vartheta')$ and $\vartheta, \vartheta' \neq 1$.

In order to study the five cases above, we will directly analyze the existence of isomorphisms between any two of the previous Lie algebras. For this we will consider bases of 1-forms adapted to the ascending central series and to the derived algebra, that is, for each Lie algebra $\mathfrak{g} \in \{\mathfrak{n}_2^\alpha, \mathfrak{n}_3^0, \mathfrak{n}_4^{\eta, \vartheta}\}$ we consider a basis $\{e^i\}_{i=1}^8$ of \mathfrak{g}^* satisfying the structure equations

$$\begin{aligned}
(37) \quad \mathfrak{n}_2^\alpha &= (0^4, 12, 13 + 15 + 24, 14 - 23 + 25, 16 + 27 + \alpha \cdot 34), \\
\mathfrak{n}_3^0 &= (0^4, 12, 13 + 25, 15 + 24, 16 + 27), \\
\mathfrak{n}_4^{\eta, \vartheta} &= (0^4, 12, 15 + (1 + \eta) \cdot 24, (\eta - 1) \cdot 14 - 23 + (\vartheta - 1) \cdot 25, 16 + 27 + 34 - 2 \cdot 45),
\end{aligned}$$

where $\alpha \in \mathbb{R}$, and $(\eta, \vartheta) \in \mathbb{R}^* \times \mathbb{R}^{>0}$ or $\mathbb{R}^{>0} \times \{0\}$.

Lemma 4.19. *Let \mathfrak{g} and \mathfrak{g}' be any of the Lie algebras \mathfrak{n}_2^α , \mathfrak{n}_3^0 , or $\mathfrak{n}_4^{\eta, \vartheta}$. Let $\{e^i\}_{i=1}^8$ and $\{e'^i\}_{i=1}^8$ be respective bases for \mathfrak{g}^* and \mathfrak{g}'^* satisfying (37). If $f: \mathfrak{g} \rightarrow \mathfrak{g}'$ is an isomorphism of Lie algebras, then the dual map $f^*: \mathfrak{g}'^* \rightarrow \mathfrak{g}^*$ satisfies*

$$\begin{aligned}
(38) \quad f^*(e'^i) \wedge e^{12} &= 0, \quad \text{for } i = 1, 2, & f^*(e'^5) \wedge e^{12345} &= 0, \\
f^*(e'^i) \wedge e^{1234} &= 0, \quad \text{for } i = 3, 4, & f^*(e'^i) \wedge e^{1234567} &= 0, \quad \text{for } i = 6, 7.
\end{aligned}$$

Proof. More generally, let $f: \mathfrak{g} \rightarrow \mathfrak{g}'$ be an isomorphism of two m -dimensional Lie algebras. Consider an ideal $\{0\} \neq \mathfrak{a} \subset \mathfrak{g}$, and let $\mathfrak{a}' = f(\mathfrak{a}) \subset \mathfrak{g}'$ be the corresponding ideal in \mathfrak{g}' . Let $\{x_{r+1}, \dots, x_m\}$ and $\{x'_{r+1}, \dots, x'_m\}$ be any bases for \mathfrak{a} and \mathfrak{a}' , respectively. We complete them up to respective basis $\{x_1, \dots, x_r, x_{r+1}, \dots, x_m\}$ and $\{x'_1, \dots, x'_r, x'_{r+1}, \dots, x'_m\}$ for \mathfrak{g} and \mathfrak{g}' . Denote the dual bases of \mathfrak{g}^* and \mathfrak{g}'^* , respectively, by $\{x^i\}_{i=1}^m$ and $\{x'^i\}_{i=1}^m$. In these conditions, it is proved in [21, Lemma 4.2] that the dual map $f^*: \mathfrak{g}'^* \rightarrow \mathfrak{g}^*$ satisfies

$$(39) \quad f^*(x'^i) \wedge x^1 \wedge \dots \wedge x^r = 0, \quad \text{for all } i = 1, \dots, r.$$

We will apply this result to different choices of \mathfrak{a} . Denote by $\{e_i\}_{i=1}^8$ (resp. $\{e'_i\}_{i=1}^8$) the basis for \mathfrak{g} (resp. \mathfrak{g}') dual to a basis $\{e^i\}_{i=1}^8$ (resp. $\{e'^i\}_{i=1}^8$) of \mathfrak{g}^* (resp. \mathfrak{g}'^*) which satisfies (37). The ascending

central series of \mathfrak{g} and \mathfrak{g}' are

$$\mathfrak{g}_1 = \langle e_8 \rangle \subset \mathfrak{g}_2 = \langle e_6, e_7, e_8 \rangle \subset \mathfrak{g}_3 = \langle e_3, e_4, e_5, e_6, e_7, e_8 \rangle,$$

and

$$\mathfrak{g}'_1 = \langle e'_8 \rangle \subset \mathfrak{g}'_2 = \langle e'_6, e'_7, e'_8 \rangle \subset \mathfrak{g}'_3 = \langle e'_3, e'_4, e'_5, e'_6, e'_7, e'_8 \rangle.$$

Since $f(\mathfrak{g}_k) = \mathfrak{g}'_k$ for any Lie algebra isomorphism $f: \mathfrak{g} \rightarrow \mathfrak{g}'$, by (39) applied to $\mathfrak{a} = \mathfrak{g}_k$ for $k = 1, 2, 3$ one finds (38) for $i = 1, 2, 5, 6$ and 7.

Moreover, the derived algebras of \mathfrak{g} and \mathfrak{g}' are

$$[\mathfrak{g}, \mathfrak{g}] = \langle e_5, e_6, e_7, e_8 \rangle, \quad [\mathfrak{g}', \mathfrak{g}'] = \langle e'_5, e'_6, e'_7, e'_8 \rangle.$$

Hence, applying (39) to $\mathfrak{a} = [\mathfrak{g}, \mathfrak{g}]$ we get (38) for $i = 3, 4$. \square

We next show how this lemma simplifies the construction of possible isomorphisms among our algebras. Let $f: \mathfrak{g} \rightarrow \mathfrak{g}'$ be an homomorphism of Lie algebras. Its dual map $f^*: \mathfrak{g}'^* \rightarrow \mathfrak{g}^*$ naturally extends to a map $F: \bigwedge^* \mathfrak{g}'^* \rightarrow \bigwedge^* \mathfrak{g}^*$ that commutes with the differentials, i.e. $F \circ d = d \circ F$. If $\{e^i\}_{i=1}^8$ and $\{e'^i\}_{i=1}^8$ are any bases for \mathfrak{g}^* and \mathfrak{g}'^* , respectively, then any Lie algebra isomorphism is defined by

$$(40) \quad F(e'^i) = \sum_{j=1}^8 \lambda_j^i e^j, \quad i = 1, \dots, 8,$$

and satisfies the conditions

$$(41) \quad F(de'^i) = d(F(e'^i)), \quad \text{for each } 1 \leq i \leq 8,$$

where the matrix $\Lambda = (\lambda_j^i)_{i,j=1,\dots,8}$ belongs to $\text{GL}(8, \mathbb{R})$.

Now, taking bases as in Lemma 4.19, we have that the matrix $\Lambda \in \text{GL}(8, \mathbb{R})$ is simplified according to (38) as follows:

$$(42) \quad \lambda_j^i = 0, \quad \text{for } \begin{cases} 1 \leq i \leq 2 \text{ and } 3 \leq j \leq 8, \\ 3 \leq i \leq 5 \text{ and } 6 \leq j \leq 8, \end{cases} \quad \text{and also } \lambda_5^3 = \lambda_5^4 = \lambda_8^6 = \lambda_8^7 = 0.$$

This makes (41) hold for every $1 \leq k \leq 4$. Furthermore, the conditions above imply that Λ is a block triangular matrix, so it satisfies

$$(43) \quad 0 \neq \det \Lambda = \det(\lambda_j^i)_{i,j=1,2} \cdot \det(\lambda_j^i)_{i,j=3,4} \cdot \lambda_5^5 \cdot \det(\lambda_j^i)_{i,j=6,7} \cdot \lambda_8^8.$$

In particular, observe that $\lambda_5^5 \neq 0$ and $\lambda_8^8 \neq 0$.

From now on, and similarly to Notation 3.9, we will use the following

Notation 4.20. We will denote by $[d(F(e'^k)) - F(de'^k)]_{ij}$ the coefficient for e'^{ij} in the expression $d(F(e'^k)) - F(de'^k)$.

Notice that from the structure equations (37) and bearing in mind (42), the vanishing of the term $[d(F(e'^5)) - F(de'^5)]_{12}$ yields

$$(44) \quad \lambda_5^5 = \lambda_1^1 \lambda_2^2 - \lambda_2^1 \lambda_1^2 \quad (\neq 0).$$

Therefore, the condition (41) reduces to $i = 6, 7, 8$.

We sum up the previous considerations in the following:

Lemma 4.21. Let \mathfrak{g} and \mathfrak{g}' be any of the Lie algebras \mathfrak{n}_2^α , \mathfrak{n}_3^0 , or $\mathfrak{n}_4^{\eta,\vartheta}$. If they are isomorphic, then in terms of respective bases for \mathfrak{g}^* and \mathfrak{g}'^* satisfying (37), any isomorphism between \mathfrak{g} and \mathfrak{g}' is defined by (40) satisfying (42)–(44) together with the conditions

$$[d(F(e'^k)) - F(de'^k)]_{ij} = 0,$$

for $k = 6, 7, 8$ and $1 \leq i < j \leq 8$.

We will proceed as follows. First, we will show part (i), completing in this way the results involving \mathfrak{n}_3^γ . Then, we will gather families \mathfrak{n}_2^α and $\mathfrak{n}_4^{\eta,\vartheta}$ in a single set of structure equations. This will allow us to see that there are no isomorphisms between an NLA in family \mathfrak{n}_2^α and an NLA in family $\mathfrak{n}_4^{\eta,\vartheta}$, thus proving (iii). Finally, we will prove (ii), (iv), and (v).

Proof of Proposition 4.18, part (i). Suppose \mathfrak{n}_2^0 and \mathfrak{n}_3^0 are isomorphic Lie algebras. Let $\{e^k\}_{k=1}^8$ and $\{e'^k\}_{k=1}^8$ be bases of \mathfrak{n}_2^0 and \mathfrak{n}_3^0 , respectively, in terms of which the structure equations are

$$\mathfrak{n}_2^0 : \begin{cases} de^1 = de^2 = de^3 = de^4 = 0, \\ de^5 = e^{12}, \\ de^6 = e^{13} + e^{15} + e^{24}, \\ de^7 = e^{14} - e^{23} + e^{25}, \\ de^8 = e^{16} + e^{27}. \end{cases} \quad \mathfrak{n}_3^0 : \begin{cases} de'^1 = de'^2 = de'^3 = de'^4 = 0, \\ de'^5 = e'^{12}, \\ de'^6 = e'^{13} + e'^{25}, \\ de'^7 = e'^{15} + e'^{24}, \\ de'^8 = e'^{16} + e'^{27}. \end{cases}$$

Then, we can apply Lemma 4.21 for these bases. From the annihilation of $[d(F(e'^k)) - F(de'^k)]_{15}$ and $[d(F(e'^k)) - F(de'^k)]_{25}$ for $k = 6, 7$, one gets:

$$\lambda_6^6 = \lambda_1^2 \lambda_5^5, \quad \lambda_6^7 = \lambda_2^2 \lambda_5^5, \quad \lambda_7^6 = \lambda_1^1 \lambda_5^5, \quad \lambda_7^7 = \lambda_2^1 \lambda_5^5,$$

where we recall that λ_5^5 follows (44). Furthermore, since $[d(F(e'^8)) - F(de'^8)]_{17}$ vanishes, one reaches the equation $\lambda_1^1 \lambda_7^6 + \lambda_1^2 \lambda_7^7 = 0$. Considering the values of λ_7^6 and λ_7^7 given above, this equation turns into

$$(45) \quad \lambda_1^1 \lambda_2^2 + \lambda_2^1 \lambda_1^2 = 0.$$

Similarly, we now notice that

$$\begin{aligned} 0 &= [d(F(e'^8)) - F(de'^8)]_{16} = \lambda_8^8 - \lambda_1^1 \lambda_6^6 - \lambda_1^2 \lambda_6^7 = \lambda_8^8 - 2\lambda_1^1 \lambda_1^2 \lambda_5^5, \\ 0 &= [d(F(e'^8)) - F(de'^8)]_{27} = \lambda_8^8 - \lambda_2^1 \lambda_7^6 - \lambda_2^2 \lambda_7^7 = \lambda_8^8 - 2\lambda_2^1 \lambda_2^2 \lambda_5^5. \end{aligned}$$

Equalling these two expressions, we deduce the condition

$$(46) \quad \lambda_1^1 \lambda_1^2 - \lambda_2^1 \lambda_2^2 = 0.$$

Let us consider the system of equations determined by (45) and (46):

$$\begin{pmatrix} \lambda_2^2 & \lambda_1^2 \\ \lambda_1^2 & -\lambda_2^2 \end{pmatrix} \begin{pmatrix} \lambda_1^1 \\ \lambda_2^1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

Observe that this system can be solved if and only if $(\lambda_2^1, \lambda_2^2) = (0, 0)$ or $(\lambda_1^1, \lambda_2^1) = (0, 0)$. However, both cases lead to $\lambda_5^5 = 0$, contradicting (44). Consequently, the Lie algebras \mathfrak{n}_2^0 and \mathfrak{n}_3^0 are non-isomorphic.

Proof of Proposition 4.18, parts (ii)–(v). As we have already mentioned, the first step is to gather the structure equations of families \mathfrak{n}_2^α and $\mathfrak{n}_4^{\eta,\vartheta}$ in a single set of equations. This can be simply done by considering

$$(47) \quad \mathfrak{h}^{\mathbf{a}} : \begin{cases} de^1 = de^2 = de^3 = de^4 = 0, \\ de^5 = e^{12}, \\ de^6 = A_{13} e^{13} + e^{15} + A_{24} e^{24}, \\ de^7 = B_{14} e^{14} - e^{23} + B_{25} e^{25}, \\ de^8 = e^{16} + e^{27} + C_{34} e^{34} + C_{45} e^{45}, \end{cases}$$

with

$$\mathbf{a} = (A_{13}, A_{24}, B_{14}, B_{25}, C_{34}, C_{45}) = \begin{cases} (1, 1, 1, 1, \alpha, 0), & \text{for } \mathfrak{n}_2^\alpha, \\ (0, 1 + \eta, \eta - 1, \vartheta - 1, 1, -2), & \text{for } \mathfrak{n}_4^{\eta,\vartheta}. \end{cases}$$

Notice that we are interested in the case $\alpha \in \mathbb{R}^*$. Moreover, recall $(\eta, \vartheta) \in \mathbb{R}^* \times \mathbb{R}^{>0}$ or $\mathbb{R}^{>0} \times \{0\}$.

Let us now consider two NLAs $\mathfrak{h}^{\mathbf{a}}$ and $\mathfrak{h}^{\mathbf{a}'}$ with respective bases $\{e^k\}_{k=1}^8$ and $\{e'^k\}_{k=1}^8$ in terms of which the structure equations follow (47), with

- $\mathbf{a} = (A_{13}, A_{24}, B_{14}, B_{25}, C_{34}, C_{45})$ and possible parameters $(\alpha, \eta, \vartheta)$,

- $\mathbf{a}' = (A'_{13}, A'_{24}, B'_{14}, B'_{25}, C'_{34}, C'_{45})$ and parameters $(\alpha', \eta', \vartheta')$.

Suppose $\mathfrak{h}^{\mathbf{a}}$ and $\mathfrak{h}^{\mathbf{a}'}$ are isomorphic. Then, we can apply Lemma 4.21 for bases satisfying (47).

We first show part (iii) of Proposition 4.18. Indeed, we can prove this more general result:

Lemma 4.22. *An NLA in family \mathfrak{n}_2^α , where $\alpha \in \mathbb{R}$, and an NLA in $\mathfrak{n}_4^{\eta, \vartheta}$, with $(\eta, \vartheta) \in \mathbb{R}^* \times \mathbb{R}^{>0}$ or $\mathbb{R}^{>0} \times \{0\}$, cannot be isomorphic.*

Proof. Under the conditions above, one first observes that $[d(F(e'^8)) - F(de'^8)]_{45} = 0$ gives

$$C_{45}\lambda_8^8 = C'_{45}\lambda_4^4\lambda_5^5.$$

Now, it suffices to remark that $C'_{45} = 0$ implies $C_{45} = 0$, since $\lambda_8^8 \neq 0$. Consequently, if there is an isomorphism F between $\mathfrak{h}^{\mathbf{a}}$ and $\mathfrak{h}^{\mathbf{a}'}$, then our two algebras must belong to the same family \mathfrak{n}_2^α or $\mathfrak{n}_4^{\eta, \vartheta}$. \square

Therefore, from now on we can assume that both $\mathfrak{h}^{\mathbf{a}}$ and $\mathfrak{h}^{\mathbf{a}'}$ belong to the same family. We first compute the expressions:

$$(48) \quad \begin{aligned} [d(F(e'^6)) - F(de'^6)]_{15} &= \lambda_6^6 - \lambda_1^1\lambda_5^5, & [d(F(e'^8)) - F(de'^8)]_{16} &= \lambda_8^8 - \lambda_1^1\lambda_6^6 - \lambda_1^2\lambda_6^7, \\ [d(F(e'^6)) - F(de'^6)]_{25} &= B_{25}\lambda_7^6 - \lambda_2^2\lambda_5^5, & [d(F(e'^8)) - F(de'^8)]_{35} &= C'_{45}\lambda_3^4\lambda_5^5, \\ [d(F(e'^7)) - F(de'^7)]_{15} &= \lambda_7^7 - B'_{25}\lambda_1^2\lambda_5^5, & [d(F(e'^8)) - F(de'^8)]_{45} &= C_{45}\lambda_8^8 - C'_{45}\lambda_4^4\lambda_5^5, \\ [d(F(e'^7)) - F(de'^7)]_{25} &= B_{25}\lambda_7^7 - B'_{25}\lambda_2^2\lambda_5^5, \end{aligned}$$

They will allow us to prove part (iv) of Proposition 4.18. In particular, we show the following:

Lemma 4.23. *The Lie algebras $\mathfrak{n}_4^{\eta, 1}$ and $\mathfrak{n}_4^{\eta', 1}$ are isomorphic if and only if $\eta = \eta'$.*

Proof. Let us respectively identify $\mathfrak{n}_4^{\eta, 1}$ and $\mathfrak{n}_4^{\eta', 1}$ with the NLAs $\mathfrak{h}^{\mathbf{a}}$ and $\mathfrak{h}^{\mathbf{a}'}$ determined by (47) with $\mathbf{a} = (0, 1 + \eta, \eta - 1, 0, 1, -2)$ and $\mathbf{a}' = (0, 1 + \eta', \eta' - 1, 0, 1, -2)$. We first notice that under these assumptions the conditions (48) simplify. In particular, since they all must be equal to zero, it suffices to consider (44) to find:

$$\lambda_2^1 = \lambda_3^4 = \lambda_6^7 = 0, \quad \lambda_4^4 = (\lambda_1^1)^2, \quad \lambda_5^5 = \lambda_1^1\lambda_2^2, \quad \lambda_6^6 = (\lambda_1^1)^2\lambda_2^2, \quad \lambda_8^8 = (\lambda_1^1)^3\lambda_2^2,$$

where $\lambda_1^1\lambda_2^2 \neq 0$. Moreover, we have

$$0 = [d(F(e'^6)) - F(de'^6)]_{24} = (1 + \eta)\lambda_6^6 - (1 + \eta')\lambda_2^2\lambda_4^4 - \lambda_2^1\lambda_4^5,$$

so the result comes straightforward simply replacing the values above and solving the corresponding equation. \square

Only parts (ii) and (v) of Proposition 4.18 remain to be proved. Although we will show these two results separately, it will be useful to have the following expressions:

$$(49) \quad \begin{aligned} [d(F(e'^6)) - F(de'^6)]_{13} &= A_{13}\lambda_6^6 - \lambda_1^1(\lambda_3^5 + A'_{13}\lambda_3^3) - A'_{24}\lambda_1^2\lambda_3^4, \\ [d(F(e'^6)) - F(de'^6)]_{23} &= -\lambda_7^6 - \lambda_2^2(\lambda_3^5 + A'_{13}\lambda_3^3) - A'_{24}\lambda_2^2\lambda_3^4, \\ [d(F(e'^6)) - F(de'^6)]_{24} &= A_{24}\lambda_6^6 - \lambda_2^1(\lambda_4^5 + A'_{13}\lambda_4^3) - A'_{24}\lambda_2^2\lambda_4^4, \\ [d(F(e'^7)) - F(de'^7)]_{13} &= A_{13}\lambda_7^7 + \lambda_1^2(\lambda_3^3 - B'_{25}\lambda_3^5) - B'_{14}\lambda_1^1\lambda_3^4, \\ [d(F(e'^7)) - F(de'^7)]_{23} &= -\lambda_7^7 + \lambda_2^2(\lambda_3^3 - B'_{25}\lambda_3^5) - B'_{14}\lambda_2^1\lambda_3^4, \\ [d(F(e'^7)) - F(de'^7)]_{24} &= A_{24}\lambda_7^7 + \lambda_2^2(\lambda_4^3 - B'_{25}\lambda_4^5) - B'_{14}\lambda_2^1\lambda_4^4, \\ [d(F(e'^8)) - F(de'^8)]_{34} &= C_{34}\lambda_8^8 + \lambda_3^4(C'_{34}\lambda_4^3 - C'_{45}\lambda_4^5) - \lambda_4^4(C'_{34}\lambda_3^3 - C'_{45}\lambda_3^5). \end{aligned}$$

The annihilation of these expressions and those in (48) will allow us to obtain the desired results. More precisely, if one focus on the family $\mathfrak{n}_4^{\eta, \vartheta}$ for $\vartheta \neq 1$, one attains the following lemma:

Lemma 4.24. *The Lie algebras $\mathfrak{n}_4^{\eta, \vartheta}$ and $\mathfrak{n}_4^{\eta', \vartheta'}$, with $(\eta, \vartheta), (\eta', \vartheta') \in \mathbb{R}^* \times \mathbb{R}^{>0} \setminus \{1\}$ or $\mathbb{R}^{>0} \times \{0\}$, are isomorphic if and only if $\eta = \eta'$ and $\vartheta = \vartheta'$.*

Proof. Let us respectively identify $\mathfrak{n}_4^{\eta, \vartheta}$ and $\mathfrak{n}_4^{\eta', \vartheta'}$ with the NLAs $\mathfrak{h}^{\mathbf{a}}$ and $\mathfrak{h}^{\mathbf{a}'}$ determined by (47) with $\mathbf{a} = (0, 1 + \eta, \eta - 1, \vartheta - 1, 1, -2)$ and $\mathbf{a}' = (0, 1 + \eta', \eta' - 1, \vartheta' - 1, 1, -2)$; namely,

$$\begin{aligned} A_{13} &= 0, & A_{24} &= 1 + \eta, & B_{14} &= \eta - 1, & B_{25} &= \vartheta - 1, & C_{34} &= 1, & C_{45} &= -2, \\ A'_{13} &= 0, & A'_{24} &= 1 + \eta', & B'_{14} &= \eta' - 1, & B'_{25} &= \vartheta' - 1, & C'_{34} &= 1, & C'_{45} &= -2. \end{aligned}$$

We first observe that equalling to zero the expressions $[d(F(e'^8)) - F(de'^8)]_{34}$, $[d(F(e'^8)) - F(de'^8)]_{35}$ and $[d(F(e'^8)) - F(de'^8)]_{45}$ given in (48) and (49), and bearing in mind (43), one obtains:

$$(50) \quad \lambda_3^3 = \lambda_5^5 - 2\lambda_3^5, \quad \lambda_3^4 = 0, \quad \lambda_4^4 = \frac{\lambda_8^8}{\lambda_5^5}.$$

In addition, annihilating the first expression in (49), one gets:

$$\lambda_1^1 \lambda_3^5 = 0.$$

This naturally leads to distinguish two cases:

- Suppose $\lambda_3^5 = 0$. Then, we observe that $\lambda_3^3 = \lambda_5^5$ due to (50). Since $[d(F(e'^6)) - F(de'^6)]_{23}$, $[d(F(e'^7)) - F(de'^7)]_{13}$ and $[d(F(e'^7)) - F(de'^7)]_{23}$ in (49) must be zero, we reach

$$\lambda_1^2 = \lambda_7^6 = 0, \quad \lambda_7^7 = \lambda_2^2 \lambda_5^5.$$

Note that this implies $\lambda_5^5 = \lambda_1^1 \lambda_2^2 (\neq 0)$ from (44). Bearing all these values in mind, the vanishing of $[d(F(e'^6)) - F(de'^6)]_{15}$, $[d(F(e'^6)) - F(de'^6)]_{25}$ and $[d(F(e'^7)) - F(de'^7)]_{25}$ in (48) gives:

$$\lambda_2^1 = 0, \quad \lambda_6^6 = (\lambda_1^1)^2 \lambda_2^2, \quad \vartheta = \vartheta'.$$

In particular, we see that the existence of an isomorphism F of this type (namely, with $\lambda_3^5 = 0$) requires $\vartheta' = \vartheta$. Let us see that also $\eta' = \eta$. Indeed, it suffices to solve λ_8^8 from $[d(F(e'^8)) - F(de'^8)]_{16} = 0$, where the left hand side is determined by (48), and then replace it in (50) to obtain:

$$\lambda_4^4 = (\lambda_1^1)^2, \quad \lambda_8^8 = (\lambda_1^1)^3 \lambda_2^2.$$

The result comes from (49), simply solving $[d(F(e'^6)) - F(de'^6)]_{24} = 0$.

- Take $\lambda_3^5 \neq 0$. Consequently, $\lambda_1^1 = 0$ and also $\lambda_5^5 = -\lambda_2^1 \lambda_1^2 (\neq 0)$ from (44). Now, using (50) when equalling to zero the expressions of $[d(F(e'^7)) - F(de'^7)]_{13}$, $[d(F(e'^7)) - F(de'^7)]_{15}$ and $[d(F(e'^7)) - F(de'^7)]_{25}$ computed in (48) and (49), one finds:

$$\lambda_3^3 = \frac{\vartheta' - 1}{\vartheta' + 1} \lambda_5^5, \quad \lambda_3^5 = \frac{\lambda_5^5}{\vartheta' + 1}, \quad \lambda_6^6 = (\vartheta' - 1) \lambda_1^2 \lambda_5^5, \quad \lambda_7^7 = \frac{\vartheta' - 1}{\vartheta' - 1} \lambda_2^2 \lambda_5^5.$$

(Recall $\vartheta, \vartheta' \in [0, 1) \cup (1, \infty)$.) These values together with the vanishing of $[d(F(e'^6)) - F(de'^6)]_{23}$, $[d(F(e'^7)) - F(de'^7)]_{23}$ and $[d(F(e'^8)) - F(de'^8)]_{16}$, determined in (48) and (49), give rise to

$$\lambda_2^2 = 0, \quad \lambda_7^6 = -\frac{\lambda_2^1 \lambda_5^5}{1 + \vartheta'}, \quad \lambda_8^8 = (\vartheta' - 1) (\lambda_1^2)^2 \lambda_5^5.$$

Let us remark that this implies $\lambda_4^4 = (\vartheta' - 1) (\lambda_1^2)^2$ due to (50). Finally, considering the vanishing of $[d(F(e'^6)) - F(de'^6)]_{25}$ and $[d(F(e'^7)) - F(de'^7)]_{24}$, one reaches the equations:

$$\vartheta + \vartheta' = 0, \quad \eta + \eta' = 0.$$

Since $\vartheta, \vartheta' \geq 0$, one needs $\vartheta = \vartheta' = 0$. However, this implies $\eta, \eta' > 0$, which contradicts the second equation. Hence, there are no valid isomorphisms in this case.

This finishes the proof of the lemma. \square

The lemma below proves part (ii) of Proposition 4.18, thus completing the proof of Theorem 4.1.

Lemma 4.25. *The Lie algebras \mathfrak{n}_2^α and $\mathfrak{n}_2^{\alpha'}$, with $\alpha, \alpha' \in \mathbb{R} \setminus \{0\}$, are isomorphic if and only if $\alpha = \alpha'$.*

Proof. We respectively identify \mathbf{n}_2^α and $\mathbf{n}_2^{\alpha'}$ with the NLAs $\mathfrak{h}^{\mathbf{a}}$ and $\mathfrak{h}^{\mathbf{a}'}$ given by (47) with $\mathbf{a} = (1, 1, 1, 1, \alpha, 0)$ and $\mathbf{a}' = (1, 1, 1, 1, \alpha', 0)$; namely,

$$\begin{aligned} A_{13} = A_{24} = B_{14} = B_{25} = 1, & \quad C_{34} = \alpha, & \quad C_{45} = 0, \\ A'_{13} = A'_{24} = B'_{14} = B'_{25} = 1, & \quad C'_{34} = \alpha', & \quad C'_{45} = 0. \end{aligned}$$

First observe that one can solve the following parameters from (48):

$$(51) \quad \lambda_6^6 = \lambda_1^1 \lambda_5^5, \quad \lambda_7^6 = \lambda_2^1 \lambda_5^5, \quad \lambda_6^7 = \lambda_1^2 \lambda_5^5, \quad \lambda_7^7 = \lambda_2^2 \lambda_5^5, \quad \lambda_8^8 = \lambda_5^5 ((\lambda_1^1)^2 + (\lambda_1^2)^2),$$

where $\lambda_5^5 \neq 0$ comes determined by (44). Since the following equation must hold

$$(52) \quad 0 = [d(F(e'^8)) - F(de'^8)]_{26} = -\lambda_5^5 (\lambda_1^1 \lambda_2^1 + \lambda_1^2 \lambda_2^2),$$

we are led to distinguish two cases.

- Let us assume $\lambda_2^1 = 0$. From (44), (51) and (52), one indeed has:

$$\lambda_2^1 = \lambda_1^2 = \lambda_6^7 = \lambda_7^6 = 0, \quad \lambda_5^5 = \lambda_1^1 \lambda_2^2, \quad \lambda_6^6 = (\lambda_1^1)^2 \lambda_2^2, \quad \lambda_7^7 = \lambda_1^1 (\lambda_2^2)^2, \quad \lambda_8^8 = (\lambda_1^1)^3 \lambda_2^2,$$

with $\lambda_1^1 \lambda_2^2 \neq 0$. These values greatly simplify the expressions of $[d(F(e'^6)) - F(de'^6)]_{13}$, $[d(F(e'^6)) - F(de'^6)]_{23}$, $[d(F(e'^6)) - F(de'^6)]_{24}$ and $[d(F(e'^7)) - F(de'^7)]_{23}$ given in (49). Since they must be equal to zero, one obtains

$$\lambda_3^4 = \lambda_3^5 = 0, \quad \lambda_3^3 = \lambda_1^1 \lambda_2^2, \quad \lambda_4^4 = (\lambda_1^1)^2.$$

If we now replace all these values in the equation $[d(F(e'^8)) - F(de'^8)]_{34} = 0$, whose left hand side can be found in (49), then we get

$$(\alpha - \alpha') (\lambda_1^1)^3 \lambda_2^2 = 0.$$

As $\lambda_1^1 \lambda_2^2 \neq 0$, this implies $\alpha = \alpha'$.

- Let $\lambda_2^1 \neq 0$. Solving λ_1^1 from (52) and taking into account (44) and (51), we compute

$$0 = [d(F(e'^8)) - F(de'^8)]_{27} = \frac{(\lambda_2^2)^2 + (\lambda_2^1)^2}{(\lambda_2^1)^2} ((\lambda_1^2)^2 - (\lambda_2^1)^2).$$

Notice that this enables to solve $\lambda_1^2 = \delta \lambda_2^1$, where $\delta \in \{-1, 1\}$. In fact, from (44), (51) and (52), we get

$$(53) \quad \begin{aligned} \lambda_1^1 &= -\delta \lambda_2^2, \quad \lambda_1^2 = \delta \lambda_2^1, \quad \lambda_6^6 = -\delta \lambda_2^2 \lambda_5^5, \quad \lambda_7^6 = \lambda_2^1 \lambda_5^5, \quad \lambda_6^7 = \delta \lambda_2^1 \lambda_5^5, \quad \lambda_7^7 = \lambda_2^2 \lambda_5^5, \\ \lambda_8^8 &= \lambda_5^5 ((\lambda_2^1)^2 + (\lambda_2^2)^2), \quad \text{where } \delta \in \{-1, 1\} \text{ and } \lambda_5^5 = -\delta ((\lambda_2^1)^2 + (\lambda_2^2)^2). \end{aligned}$$

Replacing these values in $[d(F(e'^6)) - F(de'^6)]_{13}$, $[d(F(e'^6)) - F(de'^6)]_{23}$ and $[d(F(e'^7)) - F(de'^7)]_{24}$ determined by (49), one finds

$$(54) \quad \lambda_3^3 = -\lambda_3^5 + \delta ((\lambda_2^1)^2 - (\lambda_2^2)^2), \quad \lambda_3^4 = 2\delta \lambda_2^1 \lambda_2^2, \quad \lambda_4^4 = \frac{\lambda_2^2}{\lambda_2^1} (\lambda_4^3 - \lambda_4^5) - (\lambda_2^1)^2 - (\lambda_2^2)^2,$$

as the three expressions above must vanish. Moreover, from (49) we also have

$$0 = [d(F(e'^7)) - F(de'^7)]_{23} = -2\lambda_2^2 \lambda_3^5,$$

and two new cases arise.

- Assume $\lambda_2^2 = 0$. Then, the expressions (53) and (54) simplify, and (49) gives us

$$\begin{aligned} 0 &= [d(F(e'^7)) - F(de'^7)]_{13} = -2\delta \lambda_2^1 \lambda_3^5, \\ 0 &= [d(F(e'^8)) - F(de'^8)]_{34} = -(\lambda_2^1)^2 (\alpha' \lambda_3^5 + \delta (\alpha - \alpha') (\lambda_2^1)^2). \end{aligned}$$

By hypothesis $\lambda_2^1 \neq 0$, so the first equations leads to $\lambda_3^5 = 0$, and the second one allows us to conclude $\alpha = \alpha'$.

– If $\lambda_2^2 \neq 0$ then $\lambda_3^5 = 0$, which simplifies (53) and (54). We then compute:

$$0 = [d(F(e'^6)) - F(de'^6)]_{14} = 2\delta\lambda_2^2\lambda_4^5,$$

$$0 = [d(F(e'^7)) - F(de'^7)]_{14} = \frac{\delta((\lambda_2^1)^2 + (\lambda_2^2)^2)}{\lambda_2^1}(\lambda_4^3 - \lambda_4^5 - 2\lambda_2^1\lambda_2^2).$$

One can solve $\lambda_4^5 = 0$ from the first equation and $\lambda_4^3 = 2\lambda_2^1\lambda_2^2$ from the the second one. Finally, the condition $[d(F(e'^8)) - F(de'^8)]_{34} = 0$, where the left hand side is determined by (49), becomes

$$\delta(\alpha - \alpha')((\lambda_2^1)^2 + (\lambda_2^2)^2)^2 = 0.$$

This necessarily requires $\alpha = \alpha'$.

Hence, we can conclude that there only exists an isomorphism between \mathfrak{n}_2^α and $\mathfrak{n}_2^{\alpha'}$ when $\alpha = \alpha'$. This finishes the proof of our lemma. \square

5. CLASSIFICATION OF NLAs UNDERLYING FAMILY II

In this section we classify the real nilpotent Lie algebras that underlie Family II (see equations (6) and parameters in Table 2). More precisely, we determine up to isomorphism the 8-dimensional NLAs that admit complex structures given by Family II. The concrete correspondence between the complex and the real structure equations can be found in Appendix B. In particular, we obtain the following result:

Theorem 5.1. *The non-isomorphic real Lie algebras underlying Family II are:*

$$\mathfrak{m}_1^\gamma = (0^3, 13, 23, 35, \gamma \cdot 12 - 34, 16 + 27 + 45),$$

$$\mathfrak{m}_2^\gamma = (0^3, 13, 23, 14 + 25, 15 + 24, 16 + \gamma \cdot 25 + 27),$$

$$\mathfrak{m}_3^{\alpha,\beta} = (0^3, 13, 23, 14 + 25 - 35, \alpha \cdot 12 + 15 + 24 + 34, 16 + 27 - 45 - \beta(2 \cdot 25 + 35)),$$

$$\mathfrak{m}_4^\gamma = (0^2, 12, 13, 23, 14 + 25, 15 + 24, 16 + \gamma \cdot 25 + 27),$$

where $\gamma \in \{0, 1\}$ and $(\alpha, \beta) = (0, 0), (1, 0)$ or $(\alpha \geq 0, 1)$. Moreover, their ascending types are listed in Table 7.

NLA	\mathfrak{m}_1^γ	\mathfrak{m}_2^γ	$\mathfrak{m}_3^{\alpha,\beta}$	\mathfrak{m}_4^γ
$(\dim \mathfrak{g}_k)_k$	(1, 3, 5, 8)			(1, 3, 5, 6, 8)

TABLE 7. Ascending type of real Lie algebras underlying Family II

Remark 5.2. *This result proves the second part of Theorem 1.2, thus completing its proof. The relation between the Lie algebras \mathfrak{g} 's and \mathfrak{m} 's is the following: $\mathfrak{g}_9^\gamma = \mathfrak{m}_1^\gamma$, $\mathfrak{g}_{10}^\gamma = \mathfrak{m}_2^\gamma$, $\mathfrak{g}_{11}^{\alpha,\beta} = \mathfrak{m}_3^{\alpha,\beta}$ and $\mathfrak{g}_{12}^\gamma = \mathfrak{m}_4^\gamma$. Recall that the real Lie algebras are described according to Notation 4.3.*

In order to prove Theorem 5.1, we develop a similar study to the one made for Family I, dividing the discussion into different sections depending on the ascending type of the underlying NLAs.

5.1. 4-step NLAs. Let us note that the ascending type of those 4-step NLAs underlying Family II is precisely (1, 3, 5, 8). Hence, we must consider the complex structure equations (6) with the appropriate values of the tuple $(\varepsilon, \mu, \nu, a, b)$ given in Table 2.

Proposition 5.3. *Let (\mathfrak{g}, J) be a 4-step NLA admitting a complex structure parametrized, up to equivalence, by Family II with $(\varepsilon, \mu, \nu, a, b) = (0, 1, 0, \gamma, 0)$, where $\gamma \in \{0, 1\}$. Then, $\mathfrak{g} \cong \mathfrak{m}_1^\gamma$. Moreover, \mathfrak{m}_1^0 and \mathfrak{m}_1^1 are not isomorphic.*

Proof. First observe that we obtain the equations of \mathfrak{m}_1^γ from (6) just considering $\{e^i\}_{i=1}^8$ given by

$$\omega^1 = -\frac{1}{8}(e^1 + ie^2), \quad \omega^2 = \frac{1}{16}(e^4 + ie^5), \quad \omega^3 = -\frac{1}{32}(e^6 + ie^7), \quad \omega^4 = -\frac{1}{128}(32e^3 - ie^8).$$

The second part of the result follows since the value $\gamma = 0$ gives four decomposable d -exact 2-forms, while the value $\gamma = 1$ only gives three. \square

Proposition 5.4. *Let (\mathfrak{g}, J) be a 4-step NLA admitting a complex structure parametrized, up to equivalence, by Family II with $(\varepsilon, \mu, \nu, a, b) = (1, 0, 0, a, b)$, where $a \in \{0, 1\}$ and $b \in \mathbb{R}$. Then, $\mathfrak{g} \cong \mathfrak{m}_2^\gamma$, with $\gamma = 0$ for $b = 0$ and $\gamma = 1$ otherwise. Moreover, \mathfrak{m}_2^0 and \mathfrak{m}_2^1 are not isomorphic.*

Proof. The equations of \mathfrak{m}_2^γ can be attained from (6) defining $\{e^i\}_{i=1}^8$ as follows:

$$(55) \quad \begin{aligned} \omega^1 &= \frac{1}{2\eta} \left(-\frac{e^1}{\sqrt{3}} + i e^2 \right), & \omega^3 &= \frac{1}{12\eta^4} (-\sqrt{3} e^6 + e^7), \\ \omega^2 &= \frac{1}{2\eta^3} \left(\frac{e^4}{3} - i \left(\frac{e^5}{\sqrt{3}} - 2a\eta^2 e^2 \right) \right), & \omega^4 &= \frac{1}{6\eta^5} \left(-\sqrt{3}\eta^3 e^3 + i \left(\frac{e^8}{2} - \frac{b\eta}{\sqrt{3}} e^6 \right) \right), \end{aligned}$$

where $\eta = 1$ for $b = 0$ and $\eta = \frac{-\sqrt{3}}{4b}$ otherwise.

Finally, observe that the second Betti numbers of \mathfrak{m}_2^0 and \mathfrak{m}_2^1 do not coincide. Indeed, it can be seen that

$$H_{\text{dR}}^2(\mathfrak{m}_2^0) = \langle [e^{12}], [e^{25}], [e^{34}], [e^{35}], [e^{17} + e^{26}], [e^{38} - e^{46} - e^{57}] \rangle,$$

$$H_{\text{dR}}^2(\mathfrak{m}_2^1) = \langle [e^{12}], [e^{25}], [e^{34}], [e^{35}], [e^{17} + e^{26}] \rangle.$$

□

Remark 5.5. *For the remaining case $(\varepsilon, \mu, \nu, a, b) = (1, 1, 0, a, b)$, it suffices to define*

$$\omega^1 = \frac{1}{2}(-e^1 + i e^2), \quad \omega^2 = \frac{1}{2}(e^4 - i e^5), \quad \omega^3 = \frac{1}{4}(-e^6 + i e^7), \quad \omega^4 = \frac{1}{4}(-2e^3 + i e^8),$$

to get the family of nilpotent Lie algebras studied by the authors in [19]. Hence, it is known that the underlying real Lie algebras are $\mathfrak{m}_3^{\alpha, \beta}$. Indeed, the cited paper proves that the algebras $\mathfrak{m}_3^{\alpha, \beta}$ are non-isomorphic for different values of α, β satisfying $(\alpha, \beta) = (0, 0), (1, 0)$ or $(\alpha \geq 0, 1)$.

To finish the study for the ascending type $(1, 3, 5, 8)$ one needs to prove that there are no isomorphisms between \mathfrak{m}_i^\bullet and \mathfrak{m}_j^\bullet for $i \neq j$, $i, j \in \{1, 2, 3\}$. Although the descending type of these three families is exactly the same, namely $(8, 5, 3, 1)$, one can make use of the number of functionally independent Casimir invariants n_I and the second Betti number b_2 . In fact, these two invariants allow us to conclude that there are no isomorphisms between any two NLAs belonging to two different families (see Table 8 below).

\mathfrak{g}	real parameter(s)	$b_2(\mathfrak{g})$	$n_I(\mathfrak{g})$
\mathfrak{m}_1^γ	$\gamma \in \{0, 1\}$	6	2
\mathfrak{m}_2^γ	$\gamma = 0$	6	4
	$\gamma = 1$	5	
$\mathfrak{m}_3^{\alpha, \beta}$	$(\alpha, \beta) = (0, 0), (1, 0),$ or $(\alpha \geq 0, 1)$	4	2

TABLE 8. Some invariants for the NLAs with ascending type $(1, 3, 5, 8)$

5.2. 5-step NLAs. According to Table 2, this case corresponds to the parameters $(\varepsilon, \mu, \nu, a, b) = (1, 0, 1, a, b)$. If we first apply (55) to the complex structure equations (6), we directly obtain the real Lie algebras \mathfrak{m}_4^γ , $\gamma \in \{0, 1\}$, given in Theorem 5.1. One now needs to show that \mathfrak{m}_4^0 and \mathfrak{m}_4^1 are not isomorphic. For this purpose we make use of the following result, which can be proved similarly to Lemma 4.19:

Lemma 5.6. *Let $\mathfrak{g} = \mathfrak{m}_4^0$ and $\mathfrak{g}' = \mathfrak{m}_4^1$. Let $\{e^i\}_{i=1}^8$ and $\{e'^i\}_{i=1}^8$ be respective bases for \mathfrak{g}^* and \mathfrak{g}'^* satisfying the corresponding structure equations in Theorem 5.1. If $f: \mathfrak{g} \rightarrow \mathfrak{g}'$ is an isomorphism of Lie algebras, then the dual map $f^*: \mathfrak{g}'^* \rightarrow \mathfrak{g}^*$ satisfies*

$$\begin{aligned} f^*(e'^i) \wedge e^{12} &= 0, \quad \text{for } i = 1, 2, & f^*(e'^i) \wedge e^{12345} &= 0, \quad \text{for } i = 4, 5. \\ f^*(e'^3) \wedge e^{123} &= 0, & f^*(e'^i) \wedge e^{1234567} &= 0, \quad \text{for } i = 6, 7. \end{aligned}$$

This allows us to show the following proposition, which completes the proof of Theorem 5.1.

Proposition 5.7. *The Lie algebras \mathfrak{m}_4^0 and \mathfrak{m}_4^1 are not isomorphic.*

Proof. Let $\{e^i\}_{i=1}^8$ and $\{e'^i\}_{i=1}^8$ be bases of $(\mathfrak{m}_4^0)^*$ and $(\mathfrak{m}_4^1)^*$, respectively, satisfying the corresponding structure equations in Theorem 5.1. Consider a Lie algebra isomorphism F between \mathfrak{m}_4^0 and \mathfrak{m}_4^1 given by (40). Using Notation 4.20, observe that the conditions (41) are equivalent to

$$[d(F(e'^k)) - F(de'^k)]_{ij} = 0, \text{ for every } 1 \leq k \leq 8 \text{ and } 1 \leq i < j \leq 8.$$

We first notice that, as a consequence of Lemma 5.6, one has

$$\lambda_j^i = 0, \text{ for } \begin{cases} 1 \leq i \leq 2 \text{ and } 3 \leq j \leq 8, \\ 3 \leq i \leq 5 \text{ and } 6 \leq j \leq 8, \end{cases} \text{ and also } \lambda_4^3 = \lambda_5^3 = \lambda_8^6 = \lambda_8^7 = 0.$$

Moreover, to ensure $\Lambda = (\lambda_j^i)_{i,j=1,\dots,8} \in \text{GL}(8, \mathbb{R})$ one needs $\lambda_3^3 \lambda_8^8 \neq 0$. We next show that:

$$(56) \quad \lambda_2^2 = \epsilon \lambda_1^1, \quad \lambda_1^2 = \epsilon \lambda_2^1, \quad \text{where } \epsilon = \pm 1.$$

Let us observe that $[d(F(e'^3)) - F(de'^3)]_{12} = 0$ together with $[d(F(e'^k)) - F(de'^k)]_{13} = 0$ and $[d(F(e'^k)) - F(de'^k)]_{23} = 0$, for $k = 4, 5$, give the following expressions:

$$(57) \quad \lambda_3^3 = \lambda_1^1 \lambda_2^2 - \lambda_2^1 \lambda_1^2 (\neq 0), \quad \lambda_4^4 = \lambda_1^1 \lambda_3^3, \quad \lambda_5^5 = \lambda_2^1 \lambda_3^3, \quad \lambda_4^5 = \lambda_1^1 \lambda_3^3, \quad \lambda_5^5 = \lambda_2^1 \lambda_3^3.$$

Furthermore, thanks to $[d(F(e'^k)) - F(de'^k)]_{14} = 0$ for $k = 6, 7$ one can solve

$$(58) \quad \lambda_6^6 = \lambda_1^1 \lambda_4^4 + \lambda_1^2 \lambda_5^5, \quad \lambda_7^7 = \lambda_1^1 \lambda_4^4 + \lambda_1^2 \lambda_5^5.$$

Considering these values and those in (57), from $[d(F(e'^k)) - F(de'^k)]_{25} = 0$, with $k = 6, 7$, one gets the system of equations

$$(59) \quad (\lambda_1^1)^2 - (\lambda_2^1)^2 + (\lambda_2^2)^2 - (\lambda_2^1)^2 = 0, \quad \lambda_1^1 \lambda_1^2 - \lambda_2^1 \lambda_2^2 = 0.$$

If we suppose $\lambda_1^1 = 0$, then (56) follows directly as a consequence of $\lambda_3^3 = -\lambda_2^1 \lambda_1^2 \neq 0$. Otherwise, one can solve $\lambda_1^2 = \frac{\lambda_2^1 \lambda_2^2}{\lambda_1^1}$ using the second equation in (59), and then

$$\lambda_3^3 = \frac{\lambda_2^2}{\lambda_1^1} ((\lambda_1^1)^2 - (\lambda_2^1)^2).$$

In particular, one observes that $\lambda_2^2 \neq 0$ and $|\lambda_2^1| \neq |\lambda_1^1|$. If we now substitute the value of λ_1^2 in the first equation of (59), a quartic equation in λ_1^1 arises. Its only valid solutions are $\lambda_1^1 = \epsilon \lambda_2^2$, where $\epsilon = \pm 1$, thus $\lambda_2^1 = \epsilon \lambda_1^1$ and we get (56).

As a consequence of the values found for λ_1^2 and λ_2^2 , one should note that (57) and (58) become:

$$\begin{aligned} \lambda_3^3 &= \epsilon ((\lambda_1^1)^2 - (\lambda_2^1)^2), & \lambda_4^4 &= \epsilon \lambda_5^5 = \lambda_1^1 \lambda_3^3, & \lambda_4^5 &= \epsilon \lambda_5^5 = \lambda_2^1 \lambda_3^3, \\ \lambda_6^6 &= \lambda_3^3 ((\lambda_1^1)^2 + (\lambda_2^1)^2), & \lambda_7^7 &= 2\epsilon \lambda_1^1 \lambda_2^1 \lambda_3^3. \end{aligned}$$

Then, one has

$$0 = [d(F(e'^8)) - F(de'^8)]_{26} = -(\lambda_2^1 \lambda_6^6 + \lambda_2^2 \lambda_7^7) = -\lambda_2^1 \lambda_3^3 (3(\lambda_1^1)^2 + (\lambda_2^1)^2).$$

Since $\lambda_3^3 \neq 0$, we are forced to take

$$\lambda_2^1 = \lambda_2^2 = 0.$$

Therefore, from $[d(F(e'^6)) - F(de'^6)]_{13} = 0$ and $[d(F(e'^7)) - F(de'^7)]_{23} = 0$, we solve

$$\lambda_4^6 = \lambda_1^1 \lambda_3^4, \quad \lambda_5^7 = \epsilon \lambda_1^1 \lambda_3^4,$$

and then we get

$$\begin{aligned} 0 &= [d(F(e'^8)) - F(de'^8)]_{14} = \lambda_6^8 - \lambda_1^1 \lambda_4^6 = \lambda_6^8 - (\lambda_1^1)^2 \lambda_3^4, \\ 0 &= [d(F(e'^8)) - F(de'^8)]_{25} = \lambda_6^8 - \lambda_2^2 (\lambda_5^5 + \lambda_5^7) = \lambda_6^8 - (\lambda_1^1)^2 (\lambda_3^3 + \lambda_3^4). \end{aligned}$$

If we solve λ_6^8 from the first equation above and then replace it in the second one, it suffices to recall that $\lambda_3^3 = \epsilon (\lambda_1^1)^2$ to conclude $\lambda_1^1 = 0$. However, this is not possible. \square

APPENDIX A. REAL LIE ALGEBRAS AND COMPLEX STRUCTURES IN FAMILY I

Ascending type	(ε, ν, a, b)	Real basis $\{e^k\}_{k=1}^8$	NLA
(1, 3, 8)	$(0, 0, 1, b),$ $b \in \{0, 1\}$	$\omega^1 = \delta e^1 - i e^2, \quad \omega^2 = -e^3 + \delta i e^4,$ $\omega^3 = \delta e^6 - i e^7, \quad \omega^4 = \frac{1}{2} e^5 + 2 \delta i e^8.$	\mathfrak{g}_1^b
(1, 3, 6, 8)	$(0, 1, 1, b)$	$\omega^1 = \delta e^1 - i e^2, \quad \omega^2 = 4(\delta e^3 - i e^4),$ $\omega^3 = -4(e^6 - \delta i e^7), \quad \omega^4 = -2(\delta e^5 + 4 i e^8).$	$\mathfrak{g}_2^{-4\delta b}$
	$(1, 1, a, 0)$ $0 < a < 2$	See (*)	\mathfrak{g}_2^0
	$(1, 1, 2, 0)$	$\omega^1 = \frac{\delta}{\sqrt{2}}(e^1 + i e^2),$ $\omega^2 = -\frac{1}{2}(e^3 - e^4) - i e^5,$ $\omega^3 = \sqrt{2}\delta(e^6 + i e^7),$ $\omega^4 = \frac{1}{2}(e^3 + e^4) + e^5 + 2 \delta i e^8.$	\mathfrak{g}_3^1
	$(1, 1, a, 0),$ $a > 2$	See (**)	\mathfrak{g}_3^0
	$(1, 0, 1, 0)$	$\omega^1 = \frac{1}{\sqrt{2}}(\delta e^1 - i e^2),$ $\omega^2 = -\frac{1}{2}(e^3 - e^4) + i \delta e^5,$ $\omega^3 = \frac{1}{\sqrt{2}}(\delta e^6 - i e^7), \quad \omega^4 = \frac{1}{4}(e^3 + e^4) + i \delta e^8.$	
	$(1, 0, 1, b)$ $b > 0$	See (***)	$\mathfrak{g}_4^{\frac{a s}{b}, \frac{ b-2\delta\nu }{a}}$
	$(1, 1, a, b)$ $a > 0, b \neq 0, 2\delta$		
	$(1, 1, a, 2\delta)$ $a > 0$	$\omega^1 = -\frac{1}{2}(e^1 - e^2 - i(e^1 + e^2)),$ $\omega^2 = \frac{a}{2}e^4 + i\left(\frac{e^3}{2} + e^5\right),$ $\omega^3 = -\frac{a}{2}(e^6 - e^7 - i(e^6 + e^7)),$ $\omega^4 = -\left(\frac{a+2}{4}e^3 + e^5\right) + i\delta a e^8.$	$\mathfrak{g}_4^{\frac{a}{2}, 0}$
(1, 4, 8)	$(1, 1, 0, 2\delta)$	$\omega^1 = \delta e^1 - i e^2, \quad \omega^2 = -\frac{\delta}{2}(e^3 - 2 i e^5),$ $\omega^3 = \delta e^6 - i e^7, \quad \omega^4 = 2\delta\left(\frac{\delta}{4}e^4 - \frac{1}{2}e^5 + i e^8\right).$	\mathfrak{g}_5
(1, 4, 6, 8)	$(1, 0, 0, 1)$	$\omega^1 = \delta e^1 - i e^2,$ $\omega^2 = \left(\frac{2\delta\nu}{b} - 1\right)e^3 + i \delta e^5,$	\mathfrak{g}_6
	$(1, 1, 0, b)$ $b \neq 0, 2\delta$	$\omega^3 = (b - 2\delta\nu)(\delta e^6 - i e^7),$ $\omega^4 = \left(\frac{b}{2} - \delta\nu\right)e^4 - \delta\nu e^5 + 2\delta(b - 2\delta\nu)i e^8.$	
(1, 5, 8)	$(0, 0, 0, 1)$	$\omega^1 = \delta e^1 - i e^2, \quad \omega^2 = -e^3 + \delta i e^4,$ $\omega^3 = \delta e^6 - i e^7, \quad \omega^4 = \frac{1}{2}e^5 + 2 \delta i e^8.$	\mathfrak{g}_7
(1, 5, 6, 8)	$(0, 1, 0, b)$ $b = \pm 1$	$\omega^1 = \delta e^1 - i e^2, \quad \omega^2 = -(b e^3 + 4 i e^4),$ $\omega^3 = -4(e^6 - i \delta e^7), \quad \omega^4 = -2(\delta e^5 + 4 i e^8).$	\mathfrak{g}_8

Here, $\delta = \pm 1$ and $s = \text{sign}(b - 2\nu\delta)$.

(*) Case $(\varepsilon, \nu, a, b) = (1, 1, a, 0)$ with $0 < a < 2$:

$$\begin{aligned}\Re \omega^1 &= \frac{-\delta a^2}{2\sqrt{3}(4-a^2)} e^2, & \Re \omega^3 &= \frac{\delta a^6}{48\sqrt{3}(4-a^2)^2} \left(a e^6 - \sqrt{4-a^2} e^7 \right), \\ \Im \omega^1 &= \frac{\delta a^2}{4\sqrt{3}(4-a^2)} \left(\sqrt{4-a^2} e^1 + a e^2 \right), & \Im \omega^3 &= \frac{\delta a^6}{24\sqrt{3}(4-a^2)^2} e^6, \\ \Re \omega^2 &= \frac{a^3}{24(4-a^2)} \left(\sqrt{4-a^2} e^3 + a e^4 \right), & \Re \omega^4 &= \frac{a^4}{48(4-a^2)^{3/2}} \left(a^2 e^3 - a \sqrt{4-a^2} e^4 + 4 e^5 \right), \\ \Im \omega^2 &= \frac{-a^3}{12(4-a^2)^{3/2}} \left(a e^3 - \sqrt{4-a^2} e^4 + a e^5 \right), & \Im \omega^4 &= \frac{\delta a^8}{144(4-a^2)^{5/2}} e^8.\end{aligned}$$

(**) Case $(\varepsilon, \nu, a, b) = (1, 1, a, 0)$ with $a > 2$:

$$\begin{aligned}\Re \omega^1 &= -\frac{\delta a^2}{(a^2-4)} \sqrt{\frac{3}{2}} \left(e^1 + \frac{1}{2-\sqrt{3}} e^2 \right), \\ \Im \omega^1 &= \frac{\delta a^2}{2(a^2-4)} \sqrt{\frac{3}{2}} \left((a + \sqrt{a^2-4}) e^1 + \frac{a - \sqrt{a^2-4}}{2-\sqrt{3}} e^2 \right), \\ \Re \omega^2 &= -\frac{3a^3}{4(a^2-4)} \left(\frac{a - \sqrt{a^2-4}}{(2-\sqrt{3})^2} e^3 - (a + \sqrt{a^2-4}) e^4 \right), \\ \Im \omega^2 &= \frac{3a^3}{2(a^2-4)^{3/2}} \left(\frac{a - \sqrt{a^2-4}}{(2-\sqrt{3})^2} e^3 + (a + \sqrt{a^2-4}) e^4 - \frac{2a}{2-\sqrt{3}} e^5 \right), \\ \Re \omega^3 &= \frac{-\delta a^6}{(a^2-4)^2(2-\sqrt{3})} \left(\frac{3}{2} \right)^{3/2} \left(\frac{a - \sqrt{a^2-4}}{2-\sqrt{3}} e^6 - (a + \sqrt{a^2-4}) e^7 \right), \\ \Im \omega^3 &= \frac{-3\delta a^6}{(2-\sqrt{3})(a^2-4)^2} \sqrt{\frac{3}{2}} \left(\frac{1}{2-\sqrt{3}} e^6 - e^7 \right), \\ \Re \omega^4 &= -\frac{3a^5}{8(a^2-4)^{3/2}} \left(\frac{a - \sqrt{a^2-4}}{(2-\sqrt{3})^2} e^3 + (a + \sqrt{a^2-4}) e^4 - \frac{8}{a(2-\sqrt{3})} e^5 \right), \\ \Im \omega^4 &= \frac{-9\delta a^8}{(2-\sqrt{3})^2(a^2-4)^{5/2}} e^8.\end{aligned}$$

(***) Cases $(\varepsilon, \nu, a, b) = (1, 0, 1, b)$ with $b > 0$, and $(\varepsilon, \nu, a, b) = (1, 1, a, b)$ with $a > 0$ and $b \neq 0, 2\delta$:

$$\begin{aligned}\Re \omega^1 &= -\frac{\delta}{2} \left(\sqrt{\frac{a}{a+s(b-2\nu\delta)}} e^1 - e^2 \right), & \Re \omega^3 &= -\frac{a\delta}{2} \left(e^6 - \sqrt{\frac{a}{a+s(b-2\nu\delta)}} e^7 \right), \\ \Im \omega^1 &= \frac{s}{2} \left(\sqrt{\frac{a}{a+s(b-2\nu\delta)}} e^1 + e^2 \right), & \Im \omega^3 &= \frac{sa}{2} \left(e^6 + \sqrt{\frac{a}{a+s(b-2\nu\delta)}} e^7 \right), \\ \Re \omega^2 &= \frac{sa}{b} e^4, & \Re \omega^4 &= -\sqrt{\frac{a}{a+s(b-2\nu\delta)}} \left(\frac{a+sb}{4} e^3 + \delta\nu s e^5 \right), \\ \Im \omega^2 &= \delta s \sqrt{\frac{a}{a+s(b-2\nu\delta)}} \left(\frac{e^3}{2} + e^5 \right), & \Im \omega^4 &= \delta a \sqrt{\frac{a}{a+s(b-2\nu\delta)}} e^8,\end{aligned}$$

where $s = \text{sign}(b - 2\nu\delta)$.

APPENDIX B. REAL LIE ALGEBRAS AND COMPLEX STRUCTURES IN FAMILY II

Ascending type	$(\varepsilon, \mu, \nu, a, b)$	Real basis $\{e^k\}_{k=1}^8$	NLA
(1, 3, 5, 8)	(0, 1, 0, 0, 0)	$\omega^1 = -\frac{1}{8}(e^1 + i e^2),$ $\omega^2 = \frac{1}{16}(e^4 + i e^5),$	\mathfrak{g}_9^0
	(0, 1, 0, 1, 0)	$\omega^3 = -\frac{1}{32}(e^6 + i e^7),$ $\omega^4 = -\frac{1}{128}(32 e^3 - i e^8).$	\mathfrak{g}_9^1
	(1, 0, 0, a, 0) $a \in \{0, 1\}$	$\omega^1 = \frac{1}{2\eta} \left(-\frac{e^1}{\sqrt{3}} + i e^2 \right),$ $\omega^2 = \frac{1}{2\eta^3} \left(\frac{e^4}{3} - i \left(\frac{e^5}{\sqrt{3}} - 2\gamma\eta^2 e^2 \right) \right),$	\mathfrak{g}_{10}^0
	(1, 0, 0, a, b) $a \in \{0, 1\}, b \neq 0$	$\omega^3 = \frac{1}{12\eta^4} (-\sqrt{3} e^6 + i e^7),$ $\omega^4 = \frac{1}{6\eta^5} \left(-\sqrt{3} \eta^3 e^3 + i \left(\frac{e^8}{2} - \frac{b\eta}{\sqrt{3}} e^6 \right) \right).$	\mathfrak{g}_{10}^1
	(1, 1, 0, 0, 0)	$\omega^1 = \frac{1}{2}(-e^1 + i\sqrt{3}e^2),$ $\omega^2 = \frac{\sqrt{3}}{4}(e^4 - i\sqrt{3}e^5),$ $\omega^3 = \frac{3}{8}(-\sqrt{3}e^6 + ie^7),$ $\omega^4 = \frac{\sqrt{3}}{4}(-e^3 + \frac{3i}{2}e^8).$	$\mathfrak{g}_{11}^{0,0}$
	(1, 1, 0, a, 0) $a \neq 0$	$\omega^1 = \frac{2a}{\sqrt{3}}(-e^1 + i\sqrt{3}e^2),$ $\omega^2 = \frac{4a^2}{\sqrt{3}}(e^4 - i\sqrt{3}e^5),$ $\omega^3 = \frac{8a^3}{\sqrt{3}}(-\sqrt{3}e^6 + ie^7),$ $\omega^4 = a \left(-e^3 + \frac{32ia^3}{\sqrt{3}} e^8 \right).$	$\mathfrak{g}_{11}^{1,0}$
(1, 1, 0, a, b) $b \neq 0$	$\omega^1 = -\frac{b}{3}(e^1 - i s_a s_b \sqrt{3} e^2),$ $\omega^2 = \frac{b^2}{3\sqrt{3}}(s_a s_b e^4 - i\sqrt{3} e^5),$ $\omega^3 = -\frac{b^3}{9}(\sqrt{3} s_a s_b e^6 - i e^7),$ $\omega^4 = -\frac{b s_a s_b}{2\sqrt{3}} \left(e^3 + \frac{4ib^3}{9} (e^6 - e^8) \right).$	$\mathfrak{g}_{11}^{\frac{2\sqrt{3} a }{ b }, 1}$	
(1, 3, 5, 6, 8)	(1, 0, 1, a, 0)	$\omega^1 = \frac{1}{2\eta} \left(-\frac{e^1}{\sqrt{3}} + i e^2 \right),$ $\omega^2 = \frac{1}{2\eta^3} \left(\frac{e^4}{3} - i \left(\frac{e^5}{\sqrt{3}} - 2a\eta^2 e^2 \right) \right),$	\mathfrak{g}_{12}^0
	(1, 0, 1, a, b) $b \neq 0$	$\omega^3 = \frac{1}{12\eta^4} (-\sqrt{3} e^6 + i e^7),$ $\omega^4 = \frac{1}{6\eta^5} \left(-\sqrt{3} \eta^3 e^3 + i \left(\frac{e^8}{2} - \frac{b\eta}{\sqrt{3}} e^6 \right) \right).$	\mathfrak{g}_{12}^1

$$\text{where } \eta = \begin{cases} 1, & b = 0, \\ \frac{-\sqrt{3}}{4b}, & b \neq 0, \end{cases} \quad \text{and } s_x = \begin{cases} 1, & x > 0, \\ -1, & x < 0. \end{cases}$$

ACKNOWLEDGMENTS

This work has been partially supported by the projects MTM2017-85649-P (AEI/FEDER, UE), and E22-17R “Algebra y Geometría” (Gobierno de Aragón/FEDER).

REFERENCES

- [1] A. Andrada, M.L. Barberis, I. Dotti, Classification of abelian complex structures on 6-dimensional Lie algebras, *J. Lond. Math. Soc. (2)* **83** (2011), no. 1, 232–255. Corrigendum: *J. Lond. Math. Soc. (2)* **87** (2013), no. 1, 319–320.
- [2] D. Angella, *Cohomological aspects in complex non-Kähler geometry*, Lecture Notes in Math. 2095, Springer 2014.
- [3] M.L. Barberis, I.G. Dotti, M. Verbitsky, Canonical bundles of complex nilmanifolds, with applications to hypercomplex geometry, *Math. Res. Lett.* **16** (2009), no. 2, 331–347.
- [4] G. Bazzoni, Vaisman nilmanifolds, *Bull. London Math. Soc.* **49** (2017), no. 5, 824–830.
- [5] L. Bigalke, S. Rollenske, Erratum to: The Frölicher spectral sequence can be arbitrarily non-degenerate, *Math. Ann.* **358** (2014), no. 3-4, 1119–1123.
- [6] M. Ceballos, A. Otal, L. Ugarte, R. Villacampa, Invariant complex structures on 6-nilmanifolds: classification, Frölicher spectral sequence and special Hermitian metrics, *J. Geom. Anal.* **26** (2016), no. 1, 252–286.
- [7] S. Console, A. Fino, Dolbeault cohomology of compact nilmanifolds, *Transform. Groups* **6** (2001), 111–124.
- [8] S. Console, A. Fino, Y.S. Poon, Stability of Abelian complex structures, *Internat. J. Math.* **17** (2006), no. 4, 401–416.
- [9] L.A. Cordero, M. Fernández, A. Gray, L. Ugarte, Compact nilmanifolds with nilpotent complex structures: Dolbeault cohomology, *Trans. Amer. Math. Soc.* **352** (2000), no. 12, 5405–5433.
- [10] I.G. Dotti, A. Fino, Hypercomplex eight-dimensional nilpotent Lie groups, *J. Pure Appl. Algebra* **184** (2003), no. 1, 41–57.
- [11] A. Fino, M. Parton, S. Salamon, Families of strong KT structures in six dimensions, *Comment. Math. Helv.* **79** (2004), 317–340.
- [12] A. Fino, L. Vezzoni, On the existence of balanced and SKT metrics on nilmanifolds, *Proc. Amer. Math. Soc.* **144** (2016), 2455–2459.
- [13] A. Fino, L. Vezzoni, A correction to “Tamed symplectic forms and strong Kähler with torsion metrics”, *J. Symplectic Geom.* **17** (2019), no. 4, 1079–1081.
- [14] Q. Gao, Q. Zhao, F. Zheng, Maximal nilpotent complex structures, arXiv:2005.13886 [math.DG].
- [15] L. García Vergnolle, E. Remm, Complex structures on quasi-filiform Lie algebras, *J. Lie Theory* **19** (2009), no. 2, 251–265.
- [16] M. Gong, Classification of nilpotent Lie algebras of dimension 7 (over algebraically closed fields and \mathbb{R}), *PhD. thesis*, University of Waterloo, 1998.
- [17] M. Goze, E. Remm, Non existence of complex structures on filiform Lie algebras, *Comm. Algebra* **30** (2002), no. 8, 3777–3788.
- [18] A. Latorre, L. Ugarte, On the stability of compact pseudo-Kähler and neutral Calabi-Yau manifolds, *J. Math. Pures Appl.*, doi.org/10.1016/j.matpur.2020.09.001.
- [19] A. Latorre, L. Ugarte, R. Villacampa, A family of complex nilmanifolds with infinitely many real homotopy types, *Complex Manifolds* **5** (2018), 89–102.
- [20] A. Latorre, L. Ugarte, R. Villacampa, The ascending central series of nilpotent Lie algebras with complex structure, *Trans. Amer. Math. Soc.* **372** (2019), no. 6, 3867–3903.
- [21] A. Latorre, L. Ugarte, R. Villacampa, On the real homotopy type of generalized complex nilmanifolds, *Mathematics* **8** (2020), no. 9, 1562, 12 pp.
- [22] C. Maclaughlin, H. Pedersen, Y.S. Poon, S. Salamon, Deformation of 2-step nilmanifolds with abelian complex structures, *J. Lond. Math. Soc.* **73** (2006), 173–193.
- [23] D.V. Millionshchikov, Complex structures on nilpotent Lie algebras and descending central series, *Rendiconti Seminario Matematico Univ. Pol. Torino* **74** (2016), no. 1, 163–182.
- [24] I. Nakamura, Complex parallelisable manifolds and their small deformations, *J. Differential Geom.* **10** (1975), no. 1, 85–112.
- [25] L. Ornea, A. Otiman, M. Stanciu, Compatibility between non-Kähler structures on complex (nil)manifolds, arXiv:2003.10708 [math.DG].
- [26] A. Otal, L. Ugarte, R. Villacampa, Hermitian metrics on compact complex manifolds and their deformation limits, Special metrics and group actions in geometry, 269–290, Springer INdAM Ser., 23, Springer, Cham, 2017.
- [27] S. Rollenske, Geometry of nilmanifolds with left-invariant complex structure and deformations in the large, *Proc. Lond. Math. Soc.* **99** (2009), 425–460.
- [28] S. Rollenske, The Kuranishi space of complex parallelisable nilmanifolds, *J. Eur. Math. Soc.* **13** (2011), no. 3, 513–531.
- [29] S. Rollenske, A. Tomassini, X. Wang, Vertical-horizontal decomposition of Laplacians and cohomologies of manifolds with trivial tangent bundles, *Ann. Mat. Pura Appl. (4)* **199** (2020), no. 3, 833–862.
- [30] S.M. Salamon, Complex structures on nilpotent Lie algebras, *J. Pure Appl. Algebra* **157** (2001), no. 2-3, 311–333.
- [31] L. Šnobl, P. Winternitz, Classification and Identification of Lie Algebras, CRM Monogr. Ser., vol. 33, Amer. Math. Soc., Providence, RI, 2014.

(A. Latorre) DEPARTAMENTO DE MATEMÁTICA APLICADA, UNIVERSIDAD POLITÉCNICA DE MADRID, C/ JOSÉ ANTONIO NOVAIS 10, 28040 MADRID, SPAIN

Email address: `adela.latorre@upm.es`

(L. Ugarte) DEPARTAMENTO DE MATEMÁTICAS - I.U.M.A., UNIVERSIDAD DE ZARAGOZA, CAMPUS PLAZA SAN FRANCISCO, 50009 ZARAGOZA, SPAIN

Email address: `ugarte@unizar.es`

(R. Villacampa) CENTRO UNIVERSITARIO DE LA DEFENSA - I.U.M.A., ACADEMIA GENERAL MILITAR, CRTA. DE HUESCA S/N. 50090 ZARAGOZA, SPAIN

Email address: `raquelvg@unizar.es`