MATHEMATICAL COMPARISON OF CLASSICAL AND QUANTUM MECHANISMS IN OPTIMIZATION UNDER LOCAL DIFFERENTIAL PRIVACY

YUUYA YOSHIDA

ABSTRACT. Let $\varepsilon > 0$. An *n*-tuple $(p_i)_{i=1}^n$ of probability vectors is called ε -differentially private $(\varepsilon$ -DP) if $e^{\varepsilon}p_j - p_i$ has no negative entries for all i, j = 1, ..., n. An *n*-tuple $(\rho_i)_{i=1}^n$ of density matrices is called classical-quantum ε -differentially private (CQ ε -DP) if $e^{\varepsilon}\rho_j - \rho_i$ is positive semi-definite for all i, j = 1, ..., n. Denote by $C_n(\varepsilon)$ the set of all ε -DP *n*-tuples, and by $CQ_n(\varepsilon)$ the set of all CQ ε -DP *n*-tuples. By considering optimization problems under local differential privacy, we define the subset $EC_n(\varepsilon)$ of $CQ_n(\varepsilon)$ that is essentially classical. Roughly speaking, an element in $EC_n(\varepsilon)$ is the image of $(p_i)_{i=1}^n \in C_n(\varepsilon)$ by a completely positive and trace-preserving linear map (CPTP map). In a preceding study, it is known that $EC_2(\varepsilon) = CQ_2(\varepsilon)$. In this paper, we show that $EC_n(\varepsilon) \neq CQ_n(\varepsilon)$ for every $n \geq 3$, and estimate the difference between $EC_n(\varepsilon)$ in a certain manner.

1. INTRODUCTION

In data analysis, data analysts need to know only some statistical information about private data while protecting the private data. They hope to maximally utilize private data under some privacy protection. In general, protection and utilization of private data have a trade-off relation, and researchers optimize the trade-off relation [7–9, 13, 14, 17].

As a way protecting private data, Warner [15] proposed randomized response in 1965, in which private data X are converted to other data Y subject to a conditional probability distribution $\mathbb{P}_{Y|X}$, and the data Y is released instead of X. Since a data analyst collects only randomized data Y, private data X are protected.

Differential privacy.—However, private data are not always protected in the above way. For instance, if Y is always equal to X, then it is clear that private data are not protected. To enforce protection of data, we impose the following condition on the

²⁰¹⁰ Mathematics Subject Classification. Primary 81P45, Secondary 68R01 68R05 62B10.

Key words and phrases. differential privacy, randomized response, quantum state, optimization, data processing inequality.

conditional probability distribution $\mathbb{P}_{Y|X}$:

(1.1) $\forall x, x', \ \mathbb{P}_{Y|X}(\cdot|x') \le e^{\varepsilon} \mathbb{P}_{Y|X}(\cdot|x),$

where $\varepsilon > 0$ is a constant, and the inequality is entrywise. This condition is called ε -differential privacy (ε -DP) [4–6]. Differential privacy (DP) was introduced by Dwork et al. [6] and Dwork [5] in the global privacy context (the case when a company or government releases users' data partially for machine learning). After that, DP was also introduced by Duchi et al. [4] in the local privacy context (the case when data providers do not trust a data analyst). Definition (1.1) is that in the local privacy context.

DP has been studied intensively by using classical probability theory. However, there are only a few studies of quantum versions of DP [1–3, 16, 18] to the best of our knowledge. In this paper, we define a quantum version of DP and investigate its mathematical aspects when *n*-ary data X are converted to quantum states ρ depending on X (i.e., classical-quantum setting) in the local privacy context.

To define a quantum version of DP, we consider an *n*-tuple of quantum states $(\rho_x)_{x=1}^n$, where x and ρ_x correspond to an input classical state and its output quantum state, respectively. Now, a data analyst needs to measure a quantum state by a measurement $(M_y)_{y=1}^m$ in order to obtain some information about the quantum state. Hence, following the classical definition of DP, we impose the following condition on the quantum states ρ_x :

(1.2)
$$\forall (M_y)_{y=1}^m \text{ POVM}, \text{ the c.p.d. } \mathbb{P}(y|x) = \text{Tr } \rho_x M_y \text{ satisfies } \varepsilon\text{-DP},$$

where "c.p.d." is an abbreviation of "conditional probability distribution"; a quantum state and a POVM are briefly explained below.

- A quantum state is defined as a *density matrix*, i.e., a positive semi-definite matrix with trace one.
- An *m*-tuple $(M_i)_{i=1}^m$ of positive semi-definite matrices is called a *positive-operator-valued measure (POVM)* if the sum of all M_i is equal to the identity matrix. A POVM is regarded as a measurement in quantum information theory.
- Given a quantum state ρ and a measurement $(M_i)_{i=1}^m$, the probability of obtaining each outcome $i = 1, \ldots, m$ is Tr ρM_i .

Condition (1.2) is called *classical-quantum* ε -*differential privacy* (CQ ε -DP) [16], and is equivalent to the following one:

$$\forall x, x', \ \rho_{x'} \le e^{\varepsilon} \rho_x,$$

where for Hermitian matrices H and H' the inequality $H \leq H'$ means for H' - H to be positive semi-definite. The definition of CQ ε -DP is a simple extension of the classical

 $\mathbf{2}$

one, because (1.1) can be written as

$$\forall x, x', \ p_{x'} \le e^{\varepsilon} p_x$$

if replacing the probability distributions $\mathbb{P}_{Y|X}(\cdot|x)$ with probability vectors p_x , where for probability vectors p and p' the inequality $p \leq p'$ means for p'-p to be non-negative. From now on, we use an *n*-tuple $(p_i)_{i=1}^n$ of probability vectors instead of $(\mathbb{P}_{Y|X}(\cdot|x))_{x=1}^n$.

We summarize the above definitions.

Definition 1.1 (Classical ε -DP [4] and classical-quantum ε -DP [16]). Let $\varepsilon > 0$ be a real number and $n \geq 2$ be an integer. An n-tuple $(p_i)_{i=1}^n$ of probability vectors is called ε -differentially private (ε -DP) if $p_i \leq e^{\varepsilon}p_j$ for all i, j = 1, ..., n. An n-tuple $(\rho_i)_{i=1}^n$ of density matrices is called classical-quantum ε -differentially private (CQ ε -DP) if $\rho_i \leq e^{\varepsilon}\rho_j$ for all i, j = 1, ..., n. Also, define the sets $C_n^{(d)}(\varepsilon)$, $CQ_n^{(d)}(\varepsilon)$, $C_n(\varepsilon)$ and $CQ_n(\varepsilon)$ as

$$C_n^{(d)}(\varepsilon) = \{\varepsilon \text{-}DP \ (p_i)_{i=1}^n : all \ p_i \ are \ probability \ vectors \ in \ \mathbb{R}^d\} \quad (d \ge 2),$$

$$CQ_n^{(d)}(\varepsilon) = \{CQ \ \varepsilon \text{-}DP \ (\rho_i)_{i=1}^n : all \ \rho_i \ are \ density \ matrices \ on \ \mathbb{C}^d\} \quad (d \ge 2),$$

$$C_n(\varepsilon) = \bigcup_{d \ge 2} C_n^{(d)}(\varepsilon), \quad CQ_n(\varepsilon) = \bigcup_{d \ge 2} CQ_n^{(d)}(\varepsilon).$$

If $(\rho_i)_{i=1}^n$ is CQ ε -DP, then all ρ_i have the same support, i.e., all the ranges of ρ_i are equal to one another. Hence, we often implicitly assume that all ρ_i have full rank if $(\rho_i)_{i=1}^n$ is CQ ε -DP.

Embedding classical states into quantum ones.—Next, let us consider a subset of $CQ_n(\varepsilon)$ that corresponds to $C_n(\varepsilon)$. For a probability vector $p = (p(i))_{i=1}^d \in \mathbb{R}^d$, define diag(p) as the diagonal matrix with diagonal entries $p(1), \ldots, p(d)$, which is a density matrix on \mathbb{C}^d . Since a quantum (resp. classical) state is a density matrix (resp. probability vector), the mapping diag (\cdot) is an embedding from the set of classical states into the set of quantum ones. Using the mapping diag (\cdot) , we obtain the set

$$\operatorname{diag}(\mathcal{C}_n(\varepsilon)) \coloneqq \{ (\operatorname{diag}(p_i))_{i=1}^n : (p_i)_{i=1}^n \in \mathcal{C}_n(\varepsilon) \}$$

that corresponds to $C_n(\varepsilon)$.

Essentially classical elements.—The set diag($C_n(\varepsilon)$) is much smaller than $CQ_n(\varepsilon)$, but actually, there is a set larger than diag($C_n(\varepsilon)$) that is "essentially classical". To describe such a set, we consider two optimization problems: one is the classical case

$$S_n^{\mathbb{C}}(\varepsilon; \Phi) = \sup_{\substack{(p_i)_{i=1}^n \in \mathbb{C}_n(\varepsilon)\\ \text{Privacy protection}}} \underbrace{\Phi(\operatorname{diag}(p_1), \dots, \operatorname{diag}(p_n))}_{\text{Utility}}$$

which is often considered in information-theoretic studies of DP [7–9, 13, 14, 16]; the other is the quantum case

$$S_n^{\text{CQ}}(\varepsilon; \Phi) = \sup_{\substack{(\rho_i)_{i=1}^n \in \text{CQ}_n(\varepsilon)\\ \text{Privacy protection}}} \underbrace{\Phi(\rho_1, \dots, \rho_n)}_{\text{Utility}}.$$

The above Φ is a real-valued function of *n* density matrices that represents the utility of private data, and the conditions $(p_i)_{i=1}^n \in C_n(\varepsilon)$ and $(\rho_i)_{i=1}^n \in CQ_n(\varepsilon)$ represent the privacy protection. Since the data analyst's purpose is to maximally utilize private data under the privacy protection, we arrive at the above optimization problems.

Now, we want to define a subset of $CQ_n(\varepsilon)$ that is "essentially classical". For this purpose, assume that the objective function Φ must satisfy monotonicity for *completely* positive and trace-preserving linear maps (CPTP maps); for the definition of CPTP maps, see Appendix.

Definition 1.2 (Monotonicity for CPTP maps). A real-valued function Φ of n density matrices is called monotone for CPTP maps if

$$\Phi(\Lambda(\rho_1),\ldots,\Lambda(\rho_n)) \le \Phi(\rho_1,\ldots,\rho_n)$$

for all density matrices ρ_1, \ldots, ρ_n and CPTP maps Λ . This inequality is called the data processing inequality (or information processing inequality).

Since a CPTP map is regarded as a quantum operation in quantum information theory, information-theoretic quantities usually satisfy monotonicity for CPTP maps. For example, quantum relative entropy, symmetric logarithmic derivative (SLD) Fisher information, Kubo–Mori–Bogoljubov (KMB) Fisher information, right logarithmic derivative (RLD) Fisher information, and trace distance satisfy monotonicity for CPTP maps [10, Theorems 5.7 and 6.2], [11, Theorem 6.7 and Lemma 6.9].

By monotonicity for CPTP maps, it follows that

$$\sup_{\substack{(p_i)_{i=1}^n \in C_n(\varepsilon)\\ \Lambda \text{ CPTP map}}} \Phi\left(\Lambda(\operatorname{diag}(p_1)), \ldots, \Lambda(\operatorname{diag}(p_n))\right) \leq S_n^{\mathcal{C}}(\varepsilon; \Phi).$$

Moreover, the opposite inequality also holds since the identity mapping on Herm(d) is a CPTP map, where Herm(d) denotes the set of all Hermitian matrices on \mathbb{C}^d . This fact leads us to the following definition.

Definition 1.3 (Essentially classical element). Let $\varepsilon > 0$ be a real number and $n \ge 2$ be an integer. We say that $(\rho_i)_{i=1}^n \in CQ_n(\varepsilon)$ is essentially classical if there exist $(p_i)_{i=1}^n \in C_n(\varepsilon)$ and a CPTP map Λ such that $\Lambda(\operatorname{diag}(p_i)) = \rho_i$ for all $i = 1, \ldots, n$. We denote by $EC_n(\varepsilon)$ the set of all essentially classical elements in $CQ_n(\varepsilon)$. Although an element in $EC_n(\varepsilon)$ consists of quantum states, the equality

$$S_n^{\mathrm{EC}}(\varepsilon; \Phi) = S_n^{\mathrm{C}}(\varepsilon; \Phi)$$

holds, where $S_n^{\text{EC}}(\varepsilon; \Phi)$ is defined in the same way as $S_n^{\text{CQ}}(\varepsilon; \Phi)$. Hence, the comparison of $S_n^{\text{C}}(\varepsilon; \Phi)$ and $S_n^{\text{CQ}}(\varepsilon; \Phi)$ is the same as that of $S_n^{\text{EC}}(\varepsilon; \Phi)$ and $S_n^{\text{CQ}}(\varepsilon; \Phi)$.

Comparison of $EC_n(\varepsilon)$ and $CQ_n(\varepsilon)$.—Although the set $EC_n(\varepsilon)$ is a subset of $CQ_n(\varepsilon)$, we are interested in whether they are equal to each other or not. If $EC_n(\varepsilon)$ is equal to $CQ_n(\varepsilon)$, then $S_n^{C}(\varepsilon; \Phi) = S_n^{EC}(\varepsilon; \Phi) = S_n^{CQ}(\varepsilon; \Phi)$, i.e., $CQ \varepsilon$ -DP mechanisms have no quantum advantage in optimization. In this perspective, it is important to compare $EC_n(\varepsilon)$ with $CQ_n(\varepsilon)$. The following fact follows from [16, Theorem 1].

Proposition 1.4. For all $\varepsilon > 0$, $EC_2(\varepsilon) = CQ_2(\varepsilon)$.

By Proposition 1.4, it follows that $S_2^{C}(\varepsilon; \Phi) = S_2^{EC}(\varepsilon; \Phi) = S_2^{CQ}(\varepsilon; \Phi)$. However, it turns out that $EC_n(\varepsilon) \neq CQ_n(\varepsilon)$ for every $n \geq 3$ (Corollary 1.8). Hence, using the following definition, we investigate the difference between $EC_n(\varepsilon)$ and $CQ_n(\varepsilon)$.

Definition 1.5. For $\varepsilon > 0$ and $n \ge 2$, define the set $\mathcal{E}_n(\varepsilon)$ as

$$\mathcal{E}_n(\varepsilon) = \{ \varepsilon' > 0 : \mathrm{CQ}_n(\varepsilon) \subset \mathrm{EC}_n(\varepsilon') \}.$$

Actually, $\mathcal{E}_n(\varepsilon)$ is non-empty (Theorem 1.6). Since $\mathrm{EC}_n(\varepsilon)$ is monotonically increasing in $\varepsilon > 0$, the set $\mathcal{E}_n(\varepsilon)$ is an interval that is not bounded above, i.e., of the form $[\varepsilon_{\inf}, \infty)$ or $(\varepsilon_{\inf}, \infty)$. Since $\mathrm{EC}_n(\varepsilon)$ is a subset of $\mathrm{CQ}_n(\varepsilon)$, and since $\mathrm{CQ}_n(\varepsilon)$ is strictly increasing in $\varepsilon > 0$, every $\varepsilon' \in \mathcal{E}_n(\varepsilon)$ is greater than or equal to ε . Therefore, Proposition 1.4 implies that $\mathcal{E}_2(\varepsilon) = [\varepsilon, \infty)$. We estimate the infimum of $\mathcal{E}_n(\varepsilon)$.

Main results.—In this paper, we show the following theorems.

Theorem 1.6. For all $\varepsilon > 0$ and $n \ge 2$, there exists $\varepsilon' \in \mathcal{E}_n(\varepsilon)$ such that $e^{\varepsilon'} - 1 = (n-1)(e^{\varepsilon} - 1)$.

Theorem 1.7. For all $n \geq 2$, $\varepsilon > 0$ and $\varepsilon' \in \mathcal{E}_n(\varepsilon)$,

$$\frac{e^{\varepsilon'}-1}{e^{\varepsilon}-1} \ge F_n(e^{\varepsilon}-1),$$

where F_n is defined as follows:

$$g_n(t) = \begin{cases} \frac{2}{t-1} (\sqrt{(n-1)(n-t)} + n - t) & 1 < t \le n, \\ \infty & t = 1, \end{cases}$$

$$a_{n,t}(x) = t \left(\frac{n-t}{n-1} (x+2)^2 - x^2 \right) & (1 \le t \le n, \ x \ge 0), \end{cases}$$

$$f_{n,k}(x) = \frac{(n+2k)x + \sqrt{(n+2k)^2x^2 + 8na_{n,k}(x)}}{2a_{n,k}(x)} & (1 \le k \le n/2, \ 0 \le x < g_n(k)), \end{cases}$$

$$F_n(x) = \min\{f_{n,k}(x) : 1 \le k \le n/2 \ \text{with} \ x < g_n(k)\} \quad (x \ge 0). \end{cases}$$

For several properties of the functions g_n , $a_{n,t}$, $f_{n,k}$ and F_n , see Lemmas 4.1 and 4.2.

Theorem 1.6 with n = 2 implies Proposition 1.4. Moreover, by Theorems 1.6 and 1.7, the infimum $\varepsilon_{\inf} = \varepsilon_{\inf}(n, \varepsilon) = \inf \mathcal{E}_n(\varepsilon)$ satisfies that

(1.3)
$$F_n(e^{\varepsilon} - 1) \le \frac{e^{\varepsilon_{\inf}} - 1}{e^{\varepsilon} - 1} \le n - 1.$$

Lemma 4.2, which is proved in Section 4, yields that

- for all $x \ge 0$, $F_2(x) = 1$;
- for all $n \ge 3$, F_n is strictly increasing;
- for all $n \ge 2$,

$$F_n(0) = \sqrt{\frac{n(n-1)}{2\lfloor n/2 \rfloor \lceil n/2 \rceil}}$$
 and $\lim_{x \to \infty} F_n(x) = \frac{n+2}{4}$,

where $\lfloor x \rfloor$ (resp. $\lceil x \rceil$) denotes the greatest (resp. least) integer $\leq x$ (resp. $\geq x$) for a real number x.

Since $F_n(x) > F_n(0) > 1$ for all $n \ge 3$ and x > 0, we obtain the following corollary.

Corollary 1.8. For all $\varepsilon > 0$ and $n \ge 3$, $\text{EC}_n(\varepsilon) \neq \text{CQ}_n(\varepsilon)$.

Since we give a concrete objective function Φ such that $S_n^{\rm C}(\varepsilon; \Phi) = S_n^{\rm EC}(\varepsilon; \Phi) < S_n^{\rm CQ}(\varepsilon; \Phi)$ for every $n \geq 3$ (Theorem 2.2), Corollary 1.8 also follows from Theorem 2.2 (although Theorem 1.7 is proved by using Theorem 2.2). Theorem 2.2 implies a sufficient condition for a CQ ε -DP *n*-tuple not to lie in EC_n(ε) (Corollary 2.3). Using Corollary 2.3, we construct CQ ε -DP *n*-tuples that do not lie in EC_n(ε) (Section 6).

We mention a relation among this paper and existing studies briefly. Ref. [16] handles the classical-quantum setting as well as this paper, but Refs. [1–3,18] consider the case when input and output states are quantum. The definition of CQ ε -DP can be regarded as a special case of quantum DP [3], but [3] does not include our results. Supplement on the set $EC_n(\varepsilon)$.—Actually, the set $EC_n(\varepsilon)$ can be written without CPTP maps.

Proposition 1.9. For all $\varepsilon > 0$ and $n \ge 2$,

(1.4)
$$\operatorname{EC}_{n}(\varepsilon) = \left\{ \left(\sum_{k} p_{i}(k) \sigma_{k} \right)_{i=1}^{n} : (p_{i})_{i=1}^{n} \in \operatorname{C}_{n}(\varepsilon), \text{ density matrices } \sigma_{k} \right\},$$

where the above sum is taken all over k = 1, ..., d if d is the dimension of the vector space that $p_1, ..., p_n$ inhabit.

Proposition 1.9 can easily be checked; see Appendix. Although we have defined the set $EC_n(\varepsilon)$ with CPTP maps, the same set is obtained even if replacing CPTP maps with positive and trace-preserving linear maps (PTP maps). That is, complete positivity is unnecessary, and positivity suffices in Definition 1.3. However, we have used CPTP maps in Definition 1.3 because CPTP maps are more natural in quantum information theory than PTP maps, and monotonicity for CPTP maps is used in Section 2.

2. Another main result

In this section, we state another main result (Theorem 2.2), which is proved in Section 5. Theorem 2.2 asserts that a certain objective function Φ satisfies that $S_n^{\rm C}(\varepsilon; \Phi) = S_n^{\rm EC}(\varepsilon; \Phi) < S_n^{\rm CQ}(\varepsilon; \Phi)$ for all $\varepsilon > 0$ and $n \ge 3$. The objective function Φ in Theorem 2.2 is constructed by using the RLD Fisher information of a one-parameter family.

Definition 2.1 (RLD Fisher information [10, p. 260]). For density matrices ρ and σ with full rank, we denote the RLD Fisher information of the one-parameter family $((1 - \theta)\rho + \theta\sigma)_{\theta \in [0,1]}$ at the point θ as

$$J_{\theta}(\rho,\sigma) = \operatorname{Tr}(\sigma-\rho)^2((1-\theta)\rho + \theta\sigma)^{-1}.$$

For probability vectors p and q, we set $J_{\theta}(p,q) = J_{\theta}(\operatorname{diag}(p),\operatorname{diag}(q))$.

The function J_{θ} satisfies monotonicity for CPTP maps (see also Definition 1.2). If $(\rho_i)_{i=1}^n$ is CQ ε -DP, we may assume that all ρ_i have full rank (see Section 1), and hence, we can consider the value $J_{\theta}(\rho_i, \rho_j)$ for all $i, j = 1, \ldots, n$. Also, for probability vectors p and q, the value $J_{\theta}(p,q)$ is the Fisher information in the classical sense. From now on, we denote by $\operatorname{avg}_{i\neq j} \alpha_{i,j}$ the arithmetic mean of real numbers $\alpha_{i,j}, i \neq j$.

Theorem 2.2. For real numbers $\theta \in [0, 1]$ and $\varepsilon > 0$ and an integer $n \ge 2$, we define the suprema $M_n^{\mathrm{C}}(\varepsilon; J_{\theta})$, $M_n^{\mathrm{EC}}(\varepsilon; J_{\theta})$ and $M_n^{\mathrm{CQ}}(\varepsilon; J_{\theta})$ as

$$M_n^{\mathcal{C}}(\varepsilon; J_{\theta}) = \sup_{\substack{(p_i)_{i=1}^n \in \mathcal{C}_n(\varepsilon) \\ (\rho_i)_{i=1}^n \in \mathcal{X}_n(\varepsilon)}} \min_{\substack{i \neq j}} J_{\theta}(p_i, p_j),$$
$$M_n^{\mathcal{X}}(\varepsilon; J_{\theta}) = \sup_{\substack{(\rho_i)_{i=1}^n \in \mathcal{X}_n(\varepsilon) \\ i \neq j}} \min_{\substack{i \neq j}} J_{\theta}(\rho_i, \rho_j) \quad (\mathcal{X} = \mathrm{EC}, \mathrm{CQ})$$

Then, for all $\theta \in [0,1]$, $\varepsilon > 0$ and $n \ge 2$, we have $M_n^{CQ}(\varepsilon; J_\theta) = M_2^{C}(\varepsilon; J_\theta)$ and

$$M_n^{\text{EC}}(\varepsilon; J_{\theta}) = M_n^{\text{C}}(\varepsilon; J_{\theta}) = \sup_{\substack{(p_i)_{i=1}^n \in \mathcal{C}_n(\varepsilon) \\ i \neq j}} \arg J_{\theta}(p_i, p_j)$$
$$= \frac{f_{\theta}(e^{\varepsilon}, 1) + f_{\theta}(1, e^{\varepsilon})}{n - 1} \max_{1 \le k \le n/2} \frac{k(n - k)}{ke^{\varepsilon} + n - k},$$

where $f_{\theta}(\alpha, \beta) \coloneqq (\alpha - \beta)^2 / ((1 - \theta)\alpha + \theta\beta)$ for $\alpha, \beta > 0$. Moreover, $M_n^{\text{EC}}(\varepsilon; J_{\theta}) < M_n^{\text{CQ}}(\varepsilon; J_{\theta})$ for all $\theta \in [0, 1]$, $\varepsilon > 0$ and $n \ge 3$.

If we set $\Phi(\rho_1, \ldots, \rho_n) = \min_{i \neq j} J_{\theta}(\rho_i, \rho_j)$, then $S_n^{\mathcal{X}}(\varepsilon; \Phi) = M_n^{\mathcal{X}}(\varepsilon; J_{\theta})$ for $\mathcal{X} = \mathcal{C}, \mathcal{EC}, \mathcal{CQ}$. Hence, Theorem 2.2 gives us a concrete objective function Φ such that $S_n^{\mathcal{C}}(\varepsilon; \Phi) = S_n^{\mathcal{EC}}(\varepsilon; \Phi) < S_n^{\mathcal{CQ}}(\varepsilon; \Phi)$ for all $\varepsilon > 0$ and $n \geq 3$. Moreover, Theorem 2.2 implies Corollary 1.8 and the following corollary immediately.

Corollary 2.3. Let $\varepsilon > 0$ be a real number and $n \ge 3$ be an integer. If $(\rho_i)_{i=1}^n \in CQ_n(\varepsilon)$ satisfies that $M_n^C(\varepsilon; J_\theta) < \arg_{i \ne j} J_\theta(\rho_i, \rho_j)$ for some $\theta \in [0, 1]$, then $(\rho_i)_{i=1}^n$ does not lie in $EC_n(\varepsilon)$.

3. Proof of Theorem 1.6

Denote by I_d the identity matrix of order d, and by $\mathbf{1}_d$ the column vector $[1, \ldots, 1]^\top \in \mathbb{R}^d$. We often use the bra-ket notation: for $u \in \mathbb{C}^d$, $|u\rangle$ and $\langle u|$ denote the column vector u and its conjugate transpose, respectively. Hence, $\langle \cdot | \cdot \rangle$ is the standard Hermitian inner product on \mathbb{C}^d , and $|u\rangle\langle u|$ is a rank-one orthogonal projection for every unit vector $u \in \mathbb{C}^d$.

To prove Theorem 1.6, we begin with the following preliminary lemma.

Lemma 3.1. If density matrices ρ_1, \ldots, ρ_n on \mathbb{C}^d are orthogonal to each other, i.e., $\operatorname{Tr} \rho_i \rho_j = 0$ for all $i \neq j$, then for all density matrices $\sigma_1, \ldots, \sigma_n$ on $\mathbb{C}^{d'}$ there exists a CPTP map Λ such that $\Lambda(\rho_i) = \sigma_i$ for all $i = 1, \ldots, n$.

Proof. Let ρ_1, \ldots, ρ_n be density matrices that are orthogonal to each other. For $i = 1, \ldots, n$, take the orthogonal projection P_i onto the support of ρ_i . Put $P_0 = I_d - \sum_{i=1}^n P_i$. Then P_0, P_1, \ldots, P_n are also orthogonal to each other. Defining $\Lambda(X) =$

 $\sum_{i=0}^{n} (\operatorname{Tr} X P_{i}) \sigma_{i} \text{ for } X \in \operatorname{Herm}(d), \text{ we find that } \Lambda \text{ is a CPTP map satisfying that } \Lambda(\rho_{i}) = \sigma_{i} \text{ for all } i = 1, \ldots, n.$

Proof of Theorem 1.6. Let $\varepsilon > 0$ be a real number and $n \ge 2$ be an integer. Define $\varepsilon' > 0$ as $e^{\varepsilon'} - 1 = (n-1)(e^{\varepsilon} - 1)$, and the *n*-tuple $(p_i)_{i=1}^n$ of probability vectors in \mathbb{R}^n as

$$[p_1,\ldots,p_n] = \frac{(e^{\varepsilon'}-1)I_n + \mathbf{1}_n\mathbf{1}_n^{\top}}{e^{\varepsilon'}+n-1} = \frac{1}{e^{\varepsilon'}+n-1} \begin{bmatrix} e^{\varepsilon'} & 1 & \cdots & 1\\ 1 & e^{\varepsilon'} & \ddots & 1\\ \vdots & \ddots & \ddots & 1\\ 1 & \cdots & 1 & e^{\varepsilon'} \end{bmatrix}.$$

Then $(p_i)_{i=1}^n$ lies in $C_n(\varepsilon')$. Moreover, for every $k = 1, \ldots, n$,

(3.1)
$$(e^{\varepsilon'} + n - 2)p_k - \sum_{i \neq k} p_i = \frac{(e^{\varepsilon'} + n - 2)e^{\varepsilon'} - (n - 1)}{e^{\varepsilon'} + n - 1}e_k = (e^{\varepsilon'} - 1)e_k,$$

where $(e_i)_{i=1}^n$ denotes the standard basis of \mathbb{R}^n .

Now, let $(\rho_i)_{i=1}^n$ be CQ ε -DP. We show that $(\rho_i)_{i=1}^n$ lies in EC_n (ε') . By the definitions of CQ ε -DP and ε' , for every $k = 1, \ldots, n$,

$$(e^{\varepsilon'} + n - 2)\rho_k - \sum_{i \neq k} \rho_i \ge (e^{\varepsilon'} + n - 2)\rho_k - \sum_{i \neq k} e^{\varepsilon}\rho_k$$
$$= (e^{\varepsilon'} + n - 2 - (n - 1)e^{\varepsilon})\rho_k = 0$$

Thus, the left-hand side can be rewritten as

$$(e^{\varepsilon'} + n - 2)\rho_k - \sum_{i \neq k} \rho_i = (e^{\varepsilon'} - 1)\sigma_k,$$

with density matrices $\sigma_1, \ldots, \sigma_n$. By Lemma 3.1, there exists a CPTP map Λ such that $\Lambda(|e_k\rangle\langle e_k|) = \sigma_k$ for every $k = 1, \ldots, n$. This and (3.1) yield that for every $k = 1, \ldots, n$,

$$(e^{\varepsilon'} + n - 2)\Lambda(\operatorname{diag}(p_k)) - \sum_{i \neq k} \Lambda(\operatorname{diag}(p_i)) = (e^{\varepsilon'} - 1)\Lambda(\operatorname{diag}(|e_k\rangle\langle e_k|)) = (e^{\varepsilon'} - 1)\sigma_k$$
$$= (e^{\varepsilon'} + n - 2)\rho_k - \sum_{i \neq k} \rho_i.$$

Solving the above simultaneous equations, we obtain $\Lambda(\operatorname{diag}(p_k)) = \rho_k$ for every $k = 1, \ldots, n$. This implies that $(\rho_i)_{i=1}^n$ lies in $\operatorname{EC}_n(\varepsilon')$.

4. Proof of Theorem 1.7

In this section, assuming Theorem 2.2, we prove Theorem 1.7. First, we begin with several lemmas on the functions g_n , $a_{n,t}$, $f_{n,k}$ and F_n in Theorem 1.7. These lemmas are necessary to prove Theorem 1.7.

Lemma 4.1. For g_n and $a_{n,t}$ in Theorem 1.7, the following facts hold.

- (1) For all $n \geq 2$, g_n is strictly decreasing.
- (2) For all $n \ge 2$, $1 \le t < n$ and $x \ge 0$, the inequality $x < g_n(t)$ is equivalent to $a_{n,t}(x) > 0$.
- (3) For all $n \ge 2$, 1 < t < n and $x \ge 0$, the equality $x = g_n(t)$ is equivalent to $a_{n,t}(x) = 0$.

Proof. Proof of fact 1. Trivial.

Proof of fact 2. If t = 1, the assertion is trivial. For all $n \ge 2$, 1 < t < n and $x \ge 0$, we have

$$a_{n,t}(x) > 0 \iff \sqrt{\frac{n-t}{n-1}}(x+2) > x \iff 2\sqrt{\frac{n-t}{n-1}} > \left(1 - \sqrt{\frac{n-t}{n-1}}\right)x$$
$$\iff 2\sqrt{\frac{n-t}{n-1}}\left(1 + \sqrt{\frac{n-t}{n-1}}\right) > \left(1 - \frac{n-t}{n-1}\right)x = \frac{t-1}{n-1}x$$
$$\iff x < \frac{2}{t-1}(\sqrt{(n-1)(n-t)} + n - t) = g_n(t).$$

Therefore, fact 2 holds.

Proof of fact 3. Due to 1 < t < n, fact 3 can be proved in a similar way to fact 2. \Box

Lemma 4.2. For g_n , $a_{n,t}$, $f_{n,k}$ and F_n in Theorem 1.7, the following facts hold. (1) For all $n \ge 2$, $1 \le k \le n/2$ and $0 \le x < g_n(k)$, the quadratic equation

(4.1)
$$\frac{k(n-k)}{n-1}(x+2)^2y^2 = (xy+2)(kxy+n)$$

for y has the unique positive solution $y = f_{n,k}(x)$, and the other solution is negative.

- (2) For all $x \ge 0$, $F_2(x) = 1$.
- (3) For all $n \ge 3$ and $1 \le k \le n/2$, $f_{n,k}(x)$ is strictly increasing in $0 \le x < g_n(k)$.
- (4) For all $n \geq 3$, F_n is strictly increasing.
- (5) For all $n \geq 2$,

$$F_n(0) = \sqrt{\frac{n(n-1)}{2\lfloor n/2 \rfloor \lceil n/2 \rceil}} \quad and \quad \lim_{x \to \infty} F_n(x) = \frac{n+2}{4}.$$

11

(6) For all $n \ge 3$ and $1 < t \le n/2$, the quadratic equation

$$p_{n,t}(x) \coloneqq a_{n,t}(x)(n+2)^2/4 - (n+2t)(n+2)x - 8n = 0$$

for x has a unique positive solution $x = G_n(t)$, and the other solution is negative. Moreover, $G_n(t) < g_n(t)$ for all $n \ge 3$ and $1 < t \le n/2$.

(7) For all $n \ge 3$ and $x \ge 0$,

$$F_n(x) = \min\{f_{n,k}(x) : 1 \le k \le n/2 \text{ with } x < G_n(k)\},\$$

where $G_n(1) \coloneqq \infty$.

(8) For all $n \ge 4$ and $2 \le k \le n/2$, $G_n(k) < 2n - 6$.

Proof. Proof of fact 1. Let $n \ge 2$ and $1 \le k \le n/2$ be integers, and $0 \le x < g_n(k)$ be a real number. Eq. (4.1) can be rewritten as

(4.1)
$$\iff k \left(\frac{n-k}{n-1} (x+2)^2 - x^2 \right) y^2 - (n+2k)xy - 2n = 0$$

(4.2) $\iff a_{n,k}(x)y^2 - (n+2k)xy - 2n = 0.$

Since $a_{n,k}(x) > 0$ by fact 2 of Lemma 4.1, fact 1 follows.

Proof of fact 2. Since $F_2(x) = f_{2,1}(x)$, fact 2 follows from the definition of $f_{2,1}$.

Proof of fact 3. Let $n \ge 3$ and $1 \le k \le n/2$ be integers. We show that $f_{n,k}$ is strictly increasing. Put $y = f_{n,k}(x)$ with $0 \le x < g_n(k)$. Differentiating both sides in (4.2) with respect to x, we have

$$a'_{n,k}(x)y^2 + 2a_{n,k}(x)yy' - (n+2k)y - (n+2k)xy' = 0,$$

which yields that

(4.3)
$$(2a_{n,k}(x)y - (n+2k)x)y^{-1}y' = n + 2k - a'_{n,k}(x)y$$

Since $y > (n + 2k)x/a_{n,k}(x)$, if the right-hand side in (4.3) is positive, then y' is also positive. Write the right-hand side in (4.3) as h(x). Since

(4.4)
$$f_{n,k}(0) = \frac{\sqrt{8na_{n,k}(0)}}{2a_{n,k}(0)} = \sqrt{\frac{2n}{a_{n,k}(0)}} = \sqrt{\frac{n(n-1)}{2k(n-k)}},$$

it turns out that

$$h(0) = n + 2k - a'_{n,k}(0)f_{n,k}(0) = n + 2k - \frac{4k(n-k)}{n-1}\sqrt{\frac{n(n-1)}{2k(n-k)}}$$
$$= n + 2k - \sqrt{\frac{8nk(n-k)}{n-1}} = (\sqrt{n} - \sqrt{2k})^2 + \sqrt{8nk}\left(1 - \sqrt{\frac{n-k}{n-1}}\right) > 0,$$

where the assumption $n \geq 3$ has been used to obtain the last inequality. We show that h(x) > 0 by contradiction. Suppose that there exists $0 < x_1 < g_n(k)$ such that $h(x_1) \leq 0$. By the continuity of h, we can take the minimum value $0 < x_2 \leq x_1$ such that $h(x_2) \leq 0$. It is clear that $h(x_2) = 0$ (if not so, then x_2 would not be the minimum). This implies that $f'_{n,k}(x_2) = 0$, since h(x) is the right-hand side in (4.3). Also, we find that $h(x_2 - \delta) > 0$ for all $0 < \delta < x_2$, and thus,

$$h'(x_2) = \lim_{\delta \to +0} \frac{h(x_2 - \delta) - h(x_2)}{-\delta} = \lim_{\delta \to +0} \frac{h(x_2 - \delta)}{-\delta} \le 0.$$

However,

$$h'(x_2) = \frac{2k(k-1)}{n-1} f_{n,k}(x_2) > 0$$

since $f'_{n,k}(x_2) = 0$, $f_{n,k}(x_2) > 0$, and

$$h'(x) = -a''_{n,k}(x)y - a'_{n,k}(x)y' = \frac{2k(k-1)}{n-1}y - a'_{n,k}(x)y'$$

By this contradiction, we conclude that h(x) > 0 and thus, y' > 0. Therefore, $f_{n,k}$ is strictly increasing.

Proof of fact 4. Let $n \ge 3$ be an integer. We show that F_n is strictly increasing. Let $x_2 > x_1 > 0$. For i = 1, 2, define the integer n_i as

$$n_i = \max\{1 \le k \le n/2 : x_i < g_n(k)\} = \max\{1 \le k \le n/2 : k < g_n^{-1}(x_i)\},\$$

where g_n^{-1} is the inverse function of g_n . By fact 1 of Lemma 4.1, $g_n^{-1}(x_2) < g_n^{-1}(x_1)$ and thus, $n_2 \leq n_1$. By this and fact 3,

$$F_n(x_1) = \min_{1 \le k \le n_1} f_{n,k}(x_1) < \min_{1 \le k \le n_2} f_{n,k}(x_2) = F_n(x_2).$$

Therefore, F_n is strictly increasing.

Proof of fact 5. Let $n \ge 2$ be an integer. From (4.4), it follows that

$$F_n(0) = \min_{1 \le k \le n/2} \sqrt{\frac{n(n-1)}{2k(n-k)}} = \sqrt{\frac{n(n-1)}{2\lfloor n/2 \rfloor \lceil n/2 \rceil}}$$

Since $x \ge g_n(k)$ for all $2 \le k \le n/2$ and $x \ge g_n(2)$, and since $a_{n,1}(x) = 4(x+1)$, we obtain

$$\lim_{x \to \infty} F_n(x) = \lim_{x \to \infty} f_{n,1}(x) = \lim_{x \to \infty} \frac{(n+2)x + \sqrt{(n+2)^2 x^2 + 8na_{n,1}(x)}}{2a_{n,1}(x)} = \frac{n+2}{4}.$$

Proof of fact 6. Let $n \ge 3$ be an integer and $1 < t \le n/2$ be a real number. Then

$$p_{n,t}(0) = a_{n,t}(0)(n+2)^2/4 - 8n = \frac{t(n-t)}{n-1}(n+2)^2 - 8n$$

> $(n+2)^2 - 8n = (n-2)^2 > 0.$

Also, the leading coefficient of $p_{n,t}(x)$ is $t(1-t)(n+2)^2/4(n-1) < 0$. Therefore, the quadratic equation $p_{n,t}(x) = 0$ has a unique positive solution $x = G_n(t)$, and the other solution is negative. By fact 3 of Lemma 4.1, we have $a_{n,t}(g_n(t)) = 0$ and thus, $p_{n,t}(g_n(t)) < 0$. From this and $g_n(t) > 0$, it follows that $G_n(t) < g_n(t)$.

Proof of fact 7. Let $n \ge 3$ be an integer. By facts 4 and 5, $F_n(x) < (n+2)/4$ for all $x \ge 0$. Thus, for all $x \ge 0$,

(4.5)
$$F_n(x) = \min\{f_{n,k}(x) : 1 \le k \le n/2 \text{ with } x < g_n(k) \text{ and } f_{n,k}(x) < (n+2)/4\}.$$

For all $2 \le k \le n/2$ and $0 \le x < g_n(k)$, we have

$$f_{n,k}(x) < (n+2)/4 \iff a_{n,k}(x) \left(\frac{n+2}{4}\right)^2 - (n+2k)x \cdot \frac{n+2}{4} - 2n > 0$$
$$\iff p_{n,k}(x) > 0 \iff x < G_n(k),$$

where (a) is derived from fact 1, (4.2), and $a_{n,k}(x) > 0$ (fact 2 of Lemma 4.1); (b) is derived from fact 6 (since the leading coefficient of $p_{n,k}(x)$ is $k(1-k)(n+2)^2/4(n-1) < 0$). The above equivalence of $f_{n,k}(x) < (n+2)/4$ and $x < G_n(k)$ is also true for k = 1, since $f_{n,1}(x) < f_{n,1}(+\infty) = (n+2)/4$ (see the definition of $f_{n,1}$ and fact 3) and $G_n(1) = \infty$. Therefore, the right-hand side in (4.5) is equal to

$$\min\{f_{n,k}(x) : 1 \le k \le n/2 \text{ with } x < G_n(k)\}.$$

Proof of fact 8. Let $n \ge 4$ and $2 \le k \le n/2$ be integers. If $k \ge 4$, then fact 6 of Lemma 4.2 and fact 1 of Lemma 4.1 imply that

$$G_n(k) < g_n(k) \le g_n(4) = \frac{2}{3}(\sqrt{(n-1)(n-4)} + n - 4) < \frac{2}{3}(n-2+n-4) < 2n-6.$$

Consider the remaining cases k = 2, 3. Recalling the definition of $p_{n,t}$, we have

$$p_{n,k}(2n-6) = a_{n,k}(2n-6)(n+2)^2/4 - (n+2k)(n+2)(2n-6) - 8n$$

Since

$$a_{n,2}(2n-6) = 8\left(\frac{n-2}{n-1}(n-2)^2 - (n-3)^2\right)$$

= $8\left(\frac{n-2}{n-1}(n-1)(n-3) + \frac{n-2}{n-1} - (n-3)^2\right)$
= $8\left((n-2)(n-3) + 1 - \frac{1}{n-1} - (n-3)^2\right) = 8\left(n-2 - \frac{1}{n-1}\right)$

and

$$a_{n,3}(2n-6) = 12\left(\frac{n-3}{n-1}(n-2)^2 - (n-3)^2\right)$$
$$= 12(n-3)\left(\frac{(n-2)^2}{n-1} - (n-3)\right) = \frac{12(n-3)}{n-1},$$

we have

$$p_{n,2}(2n-6) = 2\left(n-2-\frac{1}{n-1}\right)(n+2)^2 - 2(n+4)(n+2)(n-3) - 8n$$

= $2(n+2)\left((n-2)(n+2) - (n+4)(n-3)\right) - \frac{2(n+2)^2}{n-1} - 8n$
= $2(n+2)(8-n) - 2 \cdot \frac{(n-1)(n+5) + 9}{n-1} - 8n$
< $2(n+2)(8-n) - 2(n+5) - 8n$
= $2(n+2)(4-n) - 2(n-3) < 0$

and

$$p_{n,3}(2n-6) = \frac{3(n-3)}{n-1}(n+2)^2 - 2(n+6)(n+2)(n-3) - 8n$$
$$= (n-3)(n+2)\left(\frac{3(n+2)}{n-1} - 2(n+6)\right) - 8n$$
$$\leq (n-3)(n+2)\left(6 - 2(n+3)\right) - 8n$$
$$\leq (n-3)(n+2) \cdot (-2)(n+3) - 8n < 0.$$

Since the leading coefficient of $p_{n,k}(x)$ is $k(1-k)(n+2)^2/4(n-1) < 0$ for k = 2, 3, it follows from fact 6 and 2n-6 > 0 that $G_n(2) < 2n-6$ and $G_n(3) < 2n-6$.

Next, assuming Theorem 2.2, we prove Theorem 1.7.

Proof of Theorem 1.7. For n = 2, the assertion follows from fact 2 of Lemma 4.2 (see also the sentences below Definition 1.5). Let $n \ge 3$ be an integer, $\varepsilon > 0$ be a real number, and $\varepsilon_1 > 0$ satisfy $e^{\varepsilon_1} - 1 < (e^{\varepsilon} - 1)F_n(e^{\varepsilon} - 1)$. We show that $M_n^{\text{CQ}}(\varepsilon; J_{1/2}) > M_n^{\text{EC}}(\varepsilon_1; J_{1/2})$. Put $x = e^{\varepsilon} - 1$ and $y = (e^{\varepsilon_1} - 1)/x < F_n(x)$. By Theorem 2.2,

$$M_n^{CQ}(\varepsilon; J_{1/2}) = M_2^{C}(\varepsilon; J_{1/2}) = \frac{4(e^{\varepsilon} - 1)^2}{e^{\varepsilon} + 1} \cdot \frac{1}{e^{\varepsilon} + 1} = \frac{4x^2}{(x+2)^2},$$
$$M_n^{EC}(\varepsilon_1; J_{1/2}) = M_n^{C}(\varepsilon_1; J_{1/2}) = \frac{4(e^{\varepsilon_1} - 1)^2}{e^{\varepsilon_1} + 1} \cdot \frac{1}{n-1} \max_{1 \le k \le n/2} \frac{k(n-k)}{ke^{\varepsilon_1} + n-k}$$
$$= \frac{4(xy)^2}{(n-1)(xy+2)} \max_{1 \le k \le n/2} \frac{k(n-k)}{kxy+n}.$$

Hence, it suffices to show that for every $1 \le k \le n/2$,

$$\frac{1}{(x+2)^2} > \frac{y^2}{(n-1)(xy+2)} \cdot \frac{k(n-k)}{kxy+n}$$

This inequality is equivalent to

(4.6)
$$h(y) \coloneqq a_{n,k}(x)y^2 - (n+2k)xy - 2n$$
$$= \frac{k(n-k)}{n-1}(x+2)^2y^2 - (xy+2)(kxy+n) < 0$$

(see also (4.1) and (4.2)). If $a_{n,k}(x) \leq 0$, then inequality (4.6) is trivial. If $a_{n,k}(x) > 0$, then inequality (4.6) holds by the inequality $0 < y < F_n(x) \le f_{n,k}(x)$ and fact 1 of Lemma 4.2. Therefore, $M_n^{CQ}(\varepsilon; J_{1/2}) > M_n^{EC}(\varepsilon_1; J_{1/2})$. Let $\varepsilon' \in \mathcal{E}_n(\varepsilon)$. Then $M_n^{EC}(\varepsilon'; J_{1/2}) \ge M_n^{CQ}(\varepsilon; J_{1/2}) > M_n^{EC}(\varepsilon_1; J_{1/2})$. Since $EC_n(\varepsilon)$ is monotonically increasing in $\varepsilon > 0$, so is $M_n^{EC}(\varepsilon; J_{1/2})$. Thus, $\varepsilon' > \varepsilon_1$, i.e.,

$$\frac{e^{\varepsilon'}-1}{e^{\varepsilon}-1} > \frac{e^{\varepsilon_1}-1}{e^{\varepsilon}-1}$$

Since $\varepsilon_1 > 0$ is arbitrary as long as $e^{\varepsilon_1} - 1 < (e^{\varepsilon} - 1)F_n(e^{\varepsilon} - 1)$, we obtain

$$\frac{e^{\varepsilon'}-1}{e^{\varepsilon}-1} \ge F_n(e^{\varepsilon}-1).$$

Theorem 1.7 and Lemma 4.2 yield the following corollaries.

Corollary 4.3. For all $n \geq 3$, $\varepsilon > 0$ and $\varepsilon' \in \mathcal{E}_n(\varepsilon)$,

$$\frac{e^{\varepsilon'}-1}{e^{\varepsilon}-1} > \sqrt{\frac{n(n-1)}{2\lfloor n/2 \rfloor \lceil n/2 \rceil}}.$$

Proof. The assertion follows from Theorem 1.7, facts 4 and 5 of Lemma 4.2 immediately. \Box

Corollary 4.4. For all $n \geq 3$, $\varepsilon > 0$ with $e^{\varepsilon} \geq 2n - 5$, and $\varepsilon' \in \mathcal{E}_n(\varepsilon)$,

$$\frac{e^{\varepsilon'}-1}{e^{\varepsilon}-1} \ge f_{n,1}(e^{\varepsilon}-1) > \frac{n+2}{4}(1-e^{-\varepsilon}),$$

where $f_{n,1}$ is defined in Theorem 1.7.

Proof. If n = 3, the assertion follows from Theorem 1.7 and $F_3(x) = f_{3,1}(x)$. If $n \ge 4$, the assertion follows from Theorem 1.7, facts 7 and 8 of Lemma 4.2.

5. Proof of Theorem 2.2

In this section, we prove Theorem 2.2. First, let us begin with the classical optimization, for which we need the following definition and lemma [14, Theorem 4].

Definition 5.1 (Sublinear function). We say that a function $\phi: (0, \infty)^n \to \mathbb{R}$ is sublinear if $\phi(x+y) \leq \phi(x) + \phi(y)$ and $\phi(\alpha x) = \alpha \phi(x)$ for all $x, y \in (0, \infty)^n$ and $\alpha > 0$.

Lemma 5.2. Let $\Phi_{\mathcal{C}}$ be a real-valued function of n probability vectors with the following condition: there exists a sublinear function $\phi: (0, \infty)^n \to \mathbb{R}$ such that

(5.1)
$$\Phi_{\mathcal{C}}(p_1, \dots, p_n) = \sum_{p_1(k), \dots, p_n(k) > 0} \phi(p_1(k), \dots, p_n(k)),$$

where the above sum is taken all over k with $p_1(k), \ldots, p_n(k) > 0$. Then, for all $\varepsilon > 0$ and $n \ge 2$,

$$\sup_{(p_i)_{i=1}^n \in \mathcal{C}_n(\varepsilon)} \Phi_{\mathcal{C}}(p_1, \dots, p_n) \\ = \max\left\{\sum_{v \in \mathcal{S}_n(\varepsilon)} \phi(v(1), \dots, v(n))\alpha_v : \begin{array}{l} \sum_{v \in \mathcal{S}_n(\varepsilon)} \alpha_v v = \mathbf{1}_n, \\ \forall v \in \mathcal{S}_n(\varepsilon), \ \alpha_v \ge 0 \end{array}\right\},$$

where $\mathcal{S}_n(\varepsilon) \coloneqq \{1, e^{\varepsilon}\}^n$.

Many information-theoretic quantities can be expressed as (5.1). Such examples are relative entropy, Fisher information, total variation distance. Especially, $J_{\theta}(p,q)$ is expressed as

$$J_{\theta}(p,q) = \sum_{p(k),q(k)>0} f_{\theta}(p(k),q(k)),$$

where the above sum is taken all over k with p(k), q(k) > 0, and the function f_{θ} defined in Theorem 2.2 is sublinear. We now prove the following lemma by using Lemma 5.2. **Lemma 5.3.** Let $\psi: (0,\infty)^2 \to \mathbb{R}$ be a sublinear function with $\psi(1,1) = 0$, and Ψ be the function $\Psi(p,q) = \sum_{p(k),q(k)>0} \psi(p(k),q(k))$ of two probability vectors. Then, for all $\varepsilon > 0$ and $n \ge 2$,

$$M_n^{\mathcal{C}}(\varepsilon; \Psi) \coloneqq \sup_{\substack{(p_i)_{i=1}^n \in \mathcal{C}_n(\varepsilon) \\ n-1}} \min_{\substack{i \neq j \\ i \neq j}} \Psi(p_i, p_j) = \sup_{\substack{(p_i)_{i=1}^n \in \mathcal{C}_n(\varepsilon) \\ i \neq j}} \operatorname{avg}_{i \neq j} \Psi(p_i, p_j)$$
$$= \frac{\psi(e^{\varepsilon}, 1) + \psi(1, e^{\varepsilon})}{n-1} \max_{1 \leq k \leq n/2} \frac{k(n-k)}{ke^{\varepsilon} + n-k}.$$

Proof. Let $\varepsilon > 0$ be a real number and $n \ge 2$ be an integer. The following inequality holds:

(5.2)
$$M_n^{\mathcal{C}}(\varepsilon; \Psi) \le \sup_{(p_i)_{i=1}^n \in \mathcal{C}_n(\varepsilon)} \arg_{i \neq j} \Psi(p_i, p_j).$$

Recall the definition $\mathcal{S}_n(\varepsilon) = \{1, e^{\varepsilon}\}^n$. Set

$$\Phi_{\mathcal{C}}(p_1,\ldots,p_n) = \sum_{i \neq j} \Psi(p_i,p_j) \text{ and } \phi(x) = \sum_{i \neq j} \psi(x(i),x(j))$$

for $x \in (0, \infty)^n$. Lemma 5.2 yields that

(5.3)
$$\sup_{\substack{(p_i)_{i=1}^n \in \mathcal{C}_n(\varepsilon) \\ i \neq j}} \sum_{i \neq j} \Psi(p_i, p_j) \\ = \max\left\{\sum_{v \in \mathcal{S}_n(\varepsilon)} \sum_{i \neq j} \psi(v(i), v(j)) \alpha_v : \begin{array}{l} \sum_{v \in \mathcal{S}_n(\varepsilon)} \alpha_v v = \mathbf{1}_n, \\ \forall v \in \mathcal{S}_n(\varepsilon), \ \alpha_v \ge 0 \end{array}\right\}.$$

Consider the partition of $S_n(\varepsilon)$ into the n+1 subsets

$$\mathcal{S}_{n,k}(\varepsilon) \coloneqq \{ v \in \mathcal{S}_n(\varepsilon) : \text{the number of } i \text{ with } v(i) = e^{\varepsilon} \text{ is } k \} \quad (k = 0, 1, \dots, n).$$

If $v \in S_{n,k}(\varepsilon)$, then $\sum_{i \neq j} \psi(v(i), v(j)) = (\psi(e^{\varepsilon}, 1) + \psi(1, e^{\varepsilon}))k(n-k)$ due to the assumption $\psi(1, 1) = 0$. Thus, for every $k = 0, 1, \ldots, n$, we have

$$\sum_{v \in S_n(\varepsilon)} \sum_{i \neq j} \psi(v(i), v(j)) \alpha_v = \sum_{k=0}^n \sum_{v \in S_{n,k}(\varepsilon)} \sum_{i \neq j} \psi(v(i), v(j)) \alpha_v$$
$$= \sum_{k=0}^n \left(\psi(e^{\varepsilon}, 1) + \psi(1, e^{\varepsilon}) \right) k(n-k) \beta_k,$$

where $\beta_k = \sum_{v \in S_{n,k}(\varepsilon)} \alpha_v$. Since the equality $\sum_{v \in S_n(\varepsilon)} \alpha_v v = \mathbf{1}_n$ yields

$$\sum_{k=0}^{n} (ke^{\varepsilon} + n - k)\beta_k = \sum_{v \in \mathcal{S}_n(\varepsilon)} \alpha_v \left\langle \mathbf{1}_n | v \right\rangle = \left\langle \mathbf{1}_n | \mathbf{1}_n \right\rangle = n,$$

the right-hand side in (5.3) is bounded above by

$$\max\left\{\sum_{k=0}^{n} \left(\psi(e^{\varepsilon}, 1) + \psi(1, e^{\varepsilon})\right) k(n-k)\beta_{k} : \begin{array}{l} \sum_{k=0}^{n} (ke^{\varepsilon} + n - k)\beta_{k} = n, \\ \beta_{0}, \dots, \beta_{n} \ge 0 \end{array}\right\}$$
$$= \left(\psi(e^{\varepsilon}, 1) + \psi(1, e^{\varepsilon})\right) n \max_{0 \le k \le n} \frac{k(n-k)}{ke^{\varepsilon} + n - k}$$
$$(5.4) \qquad = \left(\psi(e^{\varepsilon}, 1) + \psi(1, e^{\varepsilon})\right) n \max_{1 \le k \le n/2} \frac{k(n-k)}{ke^{\varepsilon} + n - k}.$$

From (5.2), (5.3) and (5.4), it follows that

(5.5)
$$M_n^{\mathcal{C}}(\varepsilon;\Psi) \le \sup_{(p_i)_{i=1}^n \in \mathcal{C}_n(\varepsilon)} \arg_{i \ne j} \Psi(p_i, p_j) \le \frac{\psi(e^{\varepsilon}, 1) + \psi(1, e^{\varepsilon})}{n-1} \max_{1 \le k \le n/2} \frac{k(n-k)}{ke^{\varepsilon} + n-k}.$$

Fix an arbitrary integer $1 \leq k \leq n/2$. Let *d* be the number of elements in $S_{n,k}(\varepsilon)$, i.e., $d = \binom{n}{k}$. Then the vector space $\mathbb{R}^{S_{n,k}(\varepsilon)}$ is isomorphic to \mathbb{R}^d . Define the probability vectors $p_1, \ldots, p_n \in \mathbb{R}^{S_{n,k}(\varepsilon)}$ as

$$p_i(v) = \frac{v(i)}{\binom{n-1}{k-1}e^{\varepsilon} + \binom{n-1}{k}} \quad (v \in \mathcal{S}_{n,k}(\varepsilon); \ i = 1, \dots, n).$$

Then $(p_i)_{i=1}^n$ is ε -DP, and moreover,

$$\min_{i \neq j} \Psi(p_i, p_j) = \min_{i \neq j} \frac{1}{\binom{n-1}{k-1}e^{\varepsilon} + \binom{n-1}{k}} \sum_{v \in \mathcal{S}_{n,k}(\varepsilon)} \psi(v(i), v(j))$$
$$= \frac{1}{\binom{n-1}{k-1}e^{\varepsilon} + \binom{n-1}{k}} \cdot \left(\psi(e^{\varepsilon}, 1) + \psi(1, e^{\varepsilon})\right) \binom{n-2}{k-1}$$
$$= \frac{\psi(e^{\varepsilon}, 1) + \psi(1, e^{\varepsilon})}{((n-1)/(n-k))e^{\varepsilon} + (n-1)/k} = \frac{\psi(e^{\varepsilon}, 1) + \psi(1, e^{\varepsilon})}{n-1} \cdot \frac{k(n-k)}{ke^{\varepsilon} + n-k}.$$

Since $1 \le k \le n/2$ is arbitrary, the inequalities in (5.5) turn to equality.

Next, we consider the quantum optimization, for which we need the following lemmas.

Lemma 5.4. Let $n \ge 2$ be an integer and $c \in [0, 1]$ be a real number. There exists an n-tuple $(u_i)_{i=1}^n$ of unit vectors in \mathbb{R}^n such that $\langle u_i | u_j \rangle = c$ for all $i \ne j$.

Proof. Since the matrix $A = (1 - c)I_n + c |\mathbf{1}_n\rangle\langle \mathbf{1}_n|$ is positive semi-definite and consists of real numbers, there exists a real square matrix B of order n such that $A = B^{\top}B$. The column vectors $u_1, \ldots, u_n \in \mathbb{R}^n$ of B satisfy that $\langle u_i | u_j \rangle = 1$ if i = j and $\langle u_i | u_j \rangle = c$ if $i \neq j$.

18

Lemma 5.5. Let $c \in [0,1)$ and $\varepsilon, t > 0$ be real numbers, $(u_i)_{i=1}^n$ be the n-tuple in Lemma 5.4, and ρ_1, \ldots, ρ_n be the density matrices defined as

$$\rho_i = \frac{1}{n+t} (I_n + t |u_i\rangle \langle u_i|) \quad (i = 1, \dots, n).$$

If $D = (e^{\varepsilon} - 1)^2 + 4(1 - c^2)e^{\varepsilon}$ and

$$0 < t \le t_{\max} \coloneqq \frac{2(e^{\varepsilon} - 1)}{\sqrt{D} + 1 - e^{\varepsilon}}$$

then $(\rho_i)_{i=1}^n$ is CQ ε -DP.

Proof. Let $i \neq j$ be positive integers less than or equal to n. We show that

(5.6)
$$|u_i\rangle\langle u_i| - e^{\varepsilon} |u_j\rangle\langle u_j| \le \frac{1 - e^{\varepsilon} + \sqrt{D}}{2}I_n$$

Take an orthonormal system $(e_i)_{i=1}^2$ of \mathbb{C}^n such that $u_i = e_1$, $u_j = \alpha e_1 + \beta e_2$, $\alpha = c$ and $\beta = \sqrt{1 - \alpha^2}$. Then the matrix $|u_i\rangle\langle u_i| - e^{\varepsilon} |u_j\rangle\langle u_j|$ can be expressed as a square matrix of order 2:

(5.7)
$$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} - e^{\varepsilon} \begin{bmatrix} \alpha^2 & \alpha\beta \\ \alpha\beta & \beta^2 \end{bmatrix} = \begin{bmatrix} 1 - e^{\varepsilon}\alpha^2 & -e^{\varepsilon}\alpha\beta \\ -e^{\varepsilon}\alpha\beta & -e^{\varepsilon}\beta^2 \end{bmatrix} =: A.$$

Since $\operatorname{Tr} A = 1 - e^{\varepsilon}$, det $A = -e^{\varepsilon}\beta^2$ and $D = (\operatorname{Tr} A)^2 - 4 \operatorname{det} A$, the greatest eigenvalue of A is equal to $(1 - e^{\varepsilon} + \sqrt{D})/2$. Therefore, inequality (5.6) holds. Consequently, we obtain

$$t(|u_i\rangle\langle u_i| - e^{\varepsilon} |u_j\rangle\langle u_j|) \le t_{\max} \frac{1 - e^{\varepsilon} + \sqrt{D}}{2} I_n = (e^{\varepsilon} - 1)I_n$$

for all $0 < t \le t_{\text{max}}$. This implies $\rho_i \le e^{\varepsilon} \rho_j$.

Recalling the definition of $M_n^{\mathrm{X}}(\varepsilon; J_{\theta})$, we have the monotonicity

(5.8)
$$M_2^{\mathcal{X}}(\varepsilon; J_\theta) \ge M_3^{\mathcal{X}}(\varepsilon; J_\theta) \ge \cdots$$

for X = C, EC, CQ. This monotonicity is used below.

Lemma 5.6. For all $\theta \in [0, 1]$, $\varepsilon > 0$ and $n \ge 2$, $M_n^{CQ}(\varepsilon; J_\theta) = M_2^{C}(\varepsilon; J_\theta)$.

Proof. Let $\theta \in [0, 1]$, $c \in [0, 1)$ and $\varepsilon, t > 0$ be real numbers, and $n \ge 2$ be an integer. Take a CQ ε -DP *n*-tuple $(\rho_i)_{i=1}^n$ in Lemma 5.5, i.e.,

$$\rho_i = \frac{1}{n+t} (I_n + t |u_i\rangle \langle u_i|), \quad u_i \in \mathbb{R}^n \quad (i = 1, \dots, n),$$

 $\langle u_i | u_j \rangle = 1$ if i = j and $\langle u_i | u_j \rangle = c$ if $i \neq j$. Fix $i \neq j$ arbitrarily. The matrix $|u_i \rangle \langle u_i | - |u_j \rangle \langle u_j |$ can be expressed as a square matrix of order 2 in the same way as (5.7):

$$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} - \begin{bmatrix} \alpha^2 & \alpha\beta \\ \alpha\beta & \beta^2 \end{bmatrix} = \begin{bmatrix} \beta^2 & -\alpha\beta \\ -\alpha\beta & -\beta^2 \end{bmatrix} = -\beta \begin{bmatrix} -\beta & \alpha \\ \alpha & \beta \end{bmatrix},$$

where $\alpha = c$ and $\beta = \sqrt{1 - \alpha^2}$. Moreover, $(|u_i\rangle \langle u_i| - |u_j\rangle \langle u_j|)^2$ is expressed as

$$\beta^2 \begin{bmatrix} -\beta & \alpha \\ \alpha & \beta \end{bmatrix}^2 = \beta^2 I_2.$$

Denote by λ_1 and λ_2 two eigenvalues of the matrix

$$A \coloneqq I_2 + t \left((1-\theta) \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + \theta \begin{bmatrix} \alpha^2 & \alpha\beta \\ \alpha\beta & \beta^2 \end{bmatrix} \right) = \begin{bmatrix} 1 + t(1-\theta\beta^2) & t\theta\alpha\beta \\ t\theta\alpha\beta & 1 + t\theta\beta^2 \end{bmatrix}.$$

It follows that

$$J_{\theta}(\rho_i, \rho_j) = \frac{t^2}{n+t} \operatorname{Tr}(|u_i\rangle\langle u_i| - |u_j\rangle\langle u_j|)^2 \left(I_n + t\left((1-\theta)|u_i\rangle\langle u_i| + \theta|u_j\rangle\langle u_j|\right)\right)^{-1}$$
$$= \frac{t^2}{n+t}\beta^2 \operatorname{Tr} A^{-1} = \frac{t^2}{n+t}\beta^2(1/\lambda_1 + 1/\lambda_2) = \frac{t^2}{n+t}\beta^2\frac{\lambda_1 + \lambda_2}{\lambda_1\lambda_2} = \frac{t^2}{n+t}\beta^2\frac{\operatorname{Tr} A}{\det A}.$$

Since $\operatorname{Tr} A = 2 + t$ and

$$\det A = (1 + t(1 - \theta\beta^2))(1 + t\theta\beta^2) - (t\theta\alpha\beta)^2 = 1 + t + t^2(1 - \theta\beta^2)\theta\beta^2 - (t\theta\beta)^2\alpha^2 = 1 + t + t^2\theta\beta^2 - (t\theta\beta)^2 = 1 + t + t^2\theta(1 - \theta)\beta^2 = 1 + t + t^2\theta(1 - \theta)(1 - c^2),$$

we have

(5.9)
$$J_{\theta}(\rho_i, \rho_j) = \frac{t^2}{n+t} \beta^2 \frac{\operatorname{Tr} A}{\det A} = \frac{t^2}{n+t} \cdot (1-c^2) \cdot \frac{2+t}{1+t+t^2\theta(1-\theta)(1-c^2)}.$$

Finally, putting $t = t_{\max}$, we take the limit $c \to 1 - 0$. Set $s = e^{\varepsilon} - 1 > 0$. Then $D = s^2 + 4(1 - c^2)e^{\varepsilon}$ and

$$t_{\max} = \frac{2s}{\sqrt{D} - s} = \frac{2s(\sqrt{D} + s)}{D - s^2} = \frac{s(\sqrt{D} + s)}{2(1 - c^2)e^{\varepsilon}}.$$

Thus, the positive number $t = t_{\text{max}}$ diverges to $+\infty$ as $c \to 1-0$, and moreover,

$$\lim_{c \to 1-0} \frac{t}{n+t} = 1, \quad \lim_{c \to 1-0} (1-c^2)t = \frac{s^2}{e^{\varepsilon}},$$
$$\lim_{c \to 1-0} \frac{2+t}{1+t+t^2\theta(1-\theta)(1-c^2)} = \lim_{c \to 1-0} \frac{t}{t+t^2\theta(1-\theta)(1-c^2)}$$
$$= \lim_{c \to 1-0} \frac{1}{1+t\theta(1-\theta)(1-c^2)} = \frac{1}{1+\theta(1-\theta)s^2/e^{\varepsilon}} = \frac{e^{\varepsilon}}{e^{\varepsilon}+\theta(1-\theta)s^2}.$$

Since Proposition 1.4 and Lemma 5.3 yield that

$$M_2^{CQ}(\varepsilon; J_\theta) = M_2^C(\varepsilon; J_\theta) = \frac{f_\theta(e^\varepsilon, 1) + f_\theta(1, e^\varepsilon)}{e^\varepsilon + 1}$$
$$= \frac{1}{e^\varepsilon + 1} \cdot \frac{(e^\varepsilon - 1)^2(e^\varepsilon + 1)}{((1 - \theta)e^\varepsilon + \theta)(\theta e^\varepsilon + 1 - \theta)} = \frac{s^2}{((1 - \theta)s + 1)(\theta s + 1)},$$

it turns out that

$$M_2^{\mathcal{C}}(\varepsilon; J_{\theta}) = M_2^{\mathcal{C}\mathcal{Q}}(\varepsilon; J_{\theta}) \ge M_n^{\mathcal{C}\mathcal{Q}}(\varepsilon; J_{\theta}) \ge \lim_{c \to 1-0} J_{\theta}(\rho_1, \rho_2)$$
$$= \lim_{c \to 1-0} \frac{t^2}{n+t} \cdot (1-c^2) \cdot \frac{2+t}{1+t+t^2\theta(1-\theta)(1-c^2)}$$
$$= \frac{s^2}{e^{\varepsilon}} \cdot \frac{e^{\varepsilon}}{e^{\varepsilon} + \theta(1-\theta)s^2} = \frac{s^2}{e^{\varepsilon} + \theta(1-\theta)s^2} = \frac{s^2}{((1-\theta)s+1)(\theta s+1)}$$
$$= M_2^{\mathcal{C}}(\varepsilon; J_{\theta}),$$

where inequality (5.8) has been used to obtain the first inequality.

Proof of Theorem 2.2. The former assertion follows from Lemmas 5.3 and 5.6 immediately. Let us show the latter assertion. Let $\theta \in [0, 1]$ and $\varepsilon > 0$ be real numbers, and $n \ge 3$ be an integer. Since Lemmas 5.3, 5.6 and inequality (5.8) imply that

$$M_n^{\mathcal{C}}(\varepsilon; J_{\theta}) \le M_3^{\mathcal{C}}(\varepsilon; J_{\theta}) = \frac{f_{\theta}(e^{\varepsilon}, 1) + f_{\theta}(1, e^{\varepsilon})}{2} \cdot \frac{2}{e^{\varepsilon} + 2}$$
$$= \frac{e^{\varepsilon} + 1}{e^{\varepsilon} + 2} M_2^{\mathcal{C}}(\varepsilon; J_{\theta}) < M_2^{\mathcal{C}}(\varepsilon; J_{\theta}) = M_n^{\mathcal{CQ}}(\varepsilon; J_{\theta}),$$

we obtain the latter assertion.

6. Concrete CQ ε -DP *n*-tuples that do not lie in EC_n(ε)

Using Corollary 2.3, we construct CQ ε -DP *n*-tuples that do not lie in EC_n(ε). In this section, we use the following lemmas instead of Lemmas 5.4 and 5.5.

Lemma 6.1. Let $d \ge 2$ be an integer and $c \in [1/d, 1]$ be a real number. There exists a (d+1)-tuple $(u_i)_{i=1}^{d+1}$ of unit vectors in \mathbb{C}^d such that $|\langle u_i|u_j\rangle| = c$ for all $i \ne j$.

Proof. For $z \in \mathbb{C}$, define the Hermitian matrix $A(z) = (\alpha_{i,j})$ of order d+1 as $\alpha_{i,j} = 1$ if i = j and $\alpha_{i,j} = z$ if i < j. Denote by eig A(z) the set of all eigenvalues of A(z). Then eig $A(c) = \{1 + dc, 1 - c\}$, eig $A(-c) = \{1 - dc, 1 + c\}$, min eig $A(c) = 1 - c \ge 0$ and min eig $A(-c) = 1 - dc \le 0$. Since the minimum eigenvalue of A(z) can be expressed as

min eig
$$A(z) = \min_{\|u\|=1} \langle u | A(z) | u \rangle$$
,

it follows that

$$\left|\min \operatorname{eig} A(z) - \min \operatorname{eig} A(z')\right| \le \left\|A(z) - A(z')\right\|$$

for all $z, z' \in \mathbb{C}$, where $\|\cdot\|$ denotes the operator norm. This shows that min eig A(z) is continuous in z. Thus, the intermediate value theorem implies that min eig $A(z_0) = 0$ for some $z_0 \in \mathbb{C}$ of magnitude c. Therefore, there exists a $d \times (d+1)$ complex matrix B such that $A(z_0) = B^*B$. The column vectors $u_1, \ldots, u_{d+1} \in \mathbb{C}^d$ of B satisfy that $\langle u_i | u_j \rangle = 1$ if i = j and $\langle u_i | u_j \rangle = z_0$ if i < j.

Lemma 6.2. Let $c \in [1/d, 1)$ and $\varepsilon, t > 0$ be real numbers, $(u_i)_{i=1}^{d+1}$ be the (d+1)-tuple in Lemma 6.1, and $\rho_1, \ldots, \rho_{d+1}$ be the density matrices defined as

$$\rho_i = \frac{1}{d+t} (I_d + t |u_i\rangle \langle u_i|) \quad (i = 1, \dots, d+1).$$

If $D = (e^{\varepsilon} - 1)^2 + 4(1 - c^2)e^{\varepsilon}$ and

$$0 < t \le t_{\max} \coloneqq \frac{2(e^{\varepsilon} - 1)}{\sqrt{D} + 1 - e^{\varepsilon}},$$

then $(\rho_i)_{i=1}^{d+1}$ is CQ ε -DP.

Proof. See the proof of Lemma 5.5.

Theorem 6.3. Let $(\rho_i)_{i=1}^3$ be a CQ ε -DP 3-tuple in Lemma 6.2 with d = 2 and $t = t_{\max}$. Then $(\rho_i)_{i=1}^3$ does not lie in EC₃ (ε) .

Proof. Set $s = e^{\varepsilon} - 1 > 0$. Then $D = s^2 + 4(1 - c^2)(s + 1)$ and $t = t_{\max} = 2s/(\sqrt{D} - s)$. Let $i \neq j$ be positive integers less than or equal to 3. We show that $M_3^{\rm C}(\varepsilon; J_{1/2}) < J_{1/2}(\rho_i, \rho_j)$. First, Theorem 2.2 implies that

$$M_3^{\rm C}(\varepsilon; J_{1/2}) = \frac{4(e^{\varepsilon} - 1)^2}{2(e^{\varepsilon} + 1)} \cdot \frac{2}{e^{\varepsilon} + 2} = \frac{4s^2}{(s+2)(s+3)}$$

Also, it follows form the same calculation as (5.9) that

$$J_{1/2}(\rho_i, \rho_j) = \frac{t^2}{2+t} \cdot (1-c^2) \cdot \frac{2+t}{1+t+(t/2)^2(1-c^2)} = \frac{(1-c^2)t^2}{1+t+(t/2)^2(1-c^2)},$$

where we must replace n in (5.9) with the dimension d = 2. Thus,

$$M_{3}^{C}(\varepsilon; J_{1/2}) < J_{1/2}(\rho_{i}, \rho_{j}) \iff \frac{4s^{2}}{(s+2)(s+3)} < \frac{(1-c^{2})t^{2}}{1+t+(t/2)^{2}(1-c^{2})}$$
$$\iff \frac{(s+2)(s+3)}{s^{2}} > \frac{4}{1-c^{2}} \cdot \frac{1+t+(t/2)^{2}(1-c^{2})}{t^{2}}$$
$$\iff 1 + \frac{5s+6}{s^{2}} > \frac{4}{1-c^{2}} \left(\frac{1+t}{t^{2}} + \frac{1-c^{2}}{4}\right) \iff \frac{5s+6}{s^{2}} > \frac{4}{1-c^{2}} \cdot \frac{1+t}{t^{2}}$$

Recalling that $D = s^2 + 4(1 - c^2)(s + 1)$ and $t = t_{\text{max}} = \frac{2s}{\sqrt{D-s}}$, we have

$$\frac{t^2}{1+t} = t - 1 + \frac{1}{1+t} = \frac{2s}{\sqrt{D} - s} - 1 + \frac{\sqrt{D} - s}{\sqrt{D} + s} = \frac{2s}{\sqrt{D} - s} + \frac{-2s}{\sqrt{D} + s}$$
$$= \frac{4s^2}{D - s^2} = \frac{s^2}{(1 - c^2)(s + 1)}.$$

Therefore,

$$M_3^{\rm C}(\varepsilon; J_{1/2}) < J_{1/2}(\rho_i, \rho_j) \iff \frac{5s+6}{s^2} > \frac{4(s+1)}{s^2}.$$

Since the right inequality always holds, so does the left inequality. By Corollary 2.3, $(\rho_i)_{i=1}^3$ does not lie in EC₃(ε).

Corollary 6.4. Let $n \geq 3$ be an integer, and $(\rho_i)_{i=1}^3$ be a $CQ \in DP 3$ -tuple in Lemma 6.2 with d = 2 and $t = t_{\max}$. Then every $(\sigma_i)_{i=1}^n \in CQ_n(\varepsilon)$ with $\sigma_i = \rho_i$, i = 1, 2, 3, does not lie in $EC_n(\varepsilon)$.

Proof. If $(\sigma_i)_{i=1}^n \in EC_n(\varepsilon)$, then $(\sigma_i)_{i=1}^3 \in EC_3(\varepsilon)$. Thus, the assertion follows from Theorem 6.3 and Definition 1.3.

7. Conclusion

We have investigated the difference between the sets $\text{EC}_n(\varepsilon)$ and $\text{CQ}_n(\varepsilon)$, which is represented by the infimum $\varepsilon_{\inf} = \varepsilon_{\inf}(n, \varepsilon) = \inf \mathcal{E}_n(\varepsilon)$. This infimum has the upper and lower bounds as (1.3). The lower bound can probably be improved, but we do not know whether the upper bound can be improved or not. It is desirable to find tighter bounds for ε_{\inf} .

Although we have not fixed the dimension d of the finite-dimensional vector spaces \mathbb{C}^d and \mathbb{R}^d , it is also important to study the case when d is fixed. For instance, it is an

interesting problem to find extreme points of $CQ_n^{(d)}(\varepsilon)$. For the classical case, Holohan et al. [12] studied extreme points of $C_n^{(d)}(\varepsilon)$.

We have used Lemma 6.1 to construct CQ ε -DP *n*-tuples that do not lie in EC_n(ε). Instead of Lemma 6.1, one might use symmetric, informationally complete, positiveoperator-valued measures (SIC-POVMs). In this case, one can probably prove that the CQ ε -DP d^2 -tuple (ρ_i) $_{i=1}^{d^2}$ of density matrices on \mathbb{C}^d constructed by a SIC-POVM does not lie in EC_{d²}(ε). However, we can prove this statement only for large $\varepsilon > 0$ if using Corollary 2.3. Hence, one needs an alternative criterion instead of Corollary 2.3 to prove the above statement.

Acknowledgments

Partial contents of this paper are also used in the author's doctoral thesis. The author is grateful to Prof. François Le Gall and Prof. Yoshimichi Ueda for giving me some advice on a part of this paper (precisely, the doctoral thesis). The author was supported by JSPS KAKENHI Grant Number JP19J20161.

Appendix

In this appendix, we discuss linear mappings from $\operatorname{Herm}(d)$ into $\operatorname{Herm}(d')$ which are used in quantum information theory. Recall that $\operatorname{Herm}(d)$ is the set of all Hermitian matrices on \mathbb{C}^d . Denote by $\operatorname{PSD}(d)$ the set of all positive semi-definite matrices on \mathbb{C}^d . First, let us begin with several basic terms (see also a textbook in quantum information theory, e.g., [10,11]). For two linear mapping Λ_i , i = 1, 2, from $\operatorname{Herm}(d_i)$ into $\operatorname{Herm}(d'_i)$, the tensor product $\Lambda_1 \otimes \Lambda_2$ is a linear mapping from $\operatorname{Herm}(d_1) \otimes \operatorname{Herm}(d_2)$ into $\operatorname{Herm}(d'_1) \otimes$ $\operatorname{Herm}(d'_2)$. Since $\operatorname{Herm}(d_1) \otimes \operatorname{Herm}(d_2)$ can be regarded as $\operatorname{Herm}(d_1d_2)$, the tensor product $\Lambda_1 \otimes \Lambda_2$ is also a linear mapping from $\operatorname{Herm}(d_1d_2)$ into $\operatorname{Herm}(d'_1d'_2)$. Let id_d be the identity mapping on $\operatorname{Herm}(d)$.

- A linear mapping Λ from Herm(d) into Herm(d') is called *positive* if Λ(PSD(d)) ⊂ PSD(d').
- A linear mapping Λ from $\operatorname{Herm}(d)$ into $\operatorname{Herm}(d')$ is called *completely positive* if $\Lambda \otimes \operatorname{id}_k$ is positive for every integer $k \geq 2$.
- A linear mapping Λ from $\operatorname{Herm}(d)$ into $\operatorname{Herm}(d')$ is called *trace-preserving* if $\operatorname{Tr} \Lambda(X) = \operatorname{Tr} X$ for every $X \in \operatorname{Herm}(d)$.
- A linear mapping Λ from $\operatorname{Herm}(d)$ into $\operatorname{Herm}(d')$ is called *CPTP* if Λ is completely positive and trace-preserving.

In quantum information theory, a quantum channel is a CPTP map.

Example (Entanglement breaking channel). The linear mapping Λ below is a CPTP map called *entanglement breaking channel*. Let $\sigma_1, \ldots, \sigma_m$ be density matrices on $\mathbb{C}^{d'}$

and $(M_k)_{k=1}^m$ be a POVM, i.e., $M_1, \ldots, M_m \ge 0$ and $\sum_{k=1}^m M_k = I_d$. For example, $(|e_k\rangle\langle e_k|)_{k=1}^d$ is a POVM, where $(e_k)_{k=1}^d$ denotes the standard basis of \mathbb{C}^d . Define the linear mapping Λ from $\operatorname{Herm}(d)$ into $\operatorname{Herm}(d')$ as $\Lambda(X) = \sum_{k=1}^m (\operatorname{Tr} M_k X) \sigma_k$. It can easily be checked that Λ is a CPTP map. This fact is used implicitly in this section.

Next, we prove Proposition 1.9.

Proof of Proposition 1.9. First, assume that $(\rho_i)_{i=1}^n$ lies in $\mathrm{EC}_n(\varepsilon)$, i.e., there exist $(p_i)_{i=1}^n \in \mathrm{C}_n(\varepsilon)$ and a CPTP map Λ such that $\rho_i = \Lambda(\mathrm{diag}(p_i))$ for all $i = 1, \ldots, n$. Denote by $(e_k)_{k=1}^d$ the standard basis of \mathbb{C}^d , where d is the dimension of the vector space that p_1, \ldots, p_n inhabit. Then $\rho_i = \sum_{k=1}^d p_i(k)\Lambda(|e_k\rangle\langle e_k|)$ for all $i = 1, \ldots, n$. Since all $\Lambda(|e_k\rangle\langle e_k|)$ are density matrices, $(\rho_i)_{i=1}^n$ lies in the right-hand side of (1.4).

Conversely, assume that $(\rho_i)_{i=1}^n$ lies in the right-hand side of (1.4): there exist $(p_i)_{i=1}^n \in \mathcal{C}_n(\varepsilon)$ and density matrices σ_k such that $\rho_i = \sum_{k=1}^d p_i(k)\sigma_k$ for all $i = 1, \ldots, n$, where d is the dimension of the vector space that p_1, \ldots, p_n inhabit. Define the CPTP map Λ as $\Lambda(X) = \sum_{k=1}^d \langle e_k | X | e_k \rangle \sigma_k$. Then $\Lambda(\operatorname{diag}(p_i)) = \rho_i$ for all $i = 1, \ldots, n$. Therefore, $(\rho_i)_{i=1}^n$ lies in $\operatorname{EC}_n(\varepsilon)$.

References

- S. Aaronson and G. N. Rothblum. Gentle measurement of quantum states and differential privacy. In STOC'19—Proceedings of the 51st Annual ACM SIGACT Symposium on Theory of Computing, pages 322–333. ACM, New York, 2019.
- [2] Y. Du, M.-H. Hsieh, T. Liu, D. Tao, and N. Liu. Quantum noise protects quantum classifiers against adversaries. *Phys. Rev. Research*, 3(2):023153, 18, 2021.
- [3] Y. Du, M.-H. Hsieh, T. Liu, S. You, and D. Tao. Quantum differentially private sparse regression learning. preprint, available at https://arxiv.org/abs/2007.11921, 2020.
- [4] J. C. Duchi, M. I. Jordan, and M. J. Wainwright. Local privacy and statistical minimax rates. In 2013 IEEE 54th Annual Symposium on Foundations of Computer Science—FOCS 2013, pages 429–438. IEEE Computer Soc., Los Alamitos, CA, 2013.
- [5] C. Dwork. Differential privacy. In Automata, languages and programming. Part II, volume 4052 of Lecture Notes in Comput. Sci., pages 1–12. Springer, Berlin, 2006.
- [6] C. Dwork, F. McSherry, K. Nissim, and A. Smith. Calibrating noise to sensitivity in private data analysis. In *Theory of cryptography*, volume 3876 of *Lecture Notes in Comput. Sci.*, pages 265–284. Springer, Berlin, 2006.
- [7] Q. Geng, P. Kairouz, S. Oh, and P. Viswanath. The staircase mechanism in differential privacy. IEEE J. Sel. Topics Signal Process., 9(7):1176–1184, 2015.
- [8] Q. Geng and P. Viswanath. The optimal noise-adding mechanism in differential privacy. *IEEE Trans. Inform. Theory*, 62(2):925–951, 2016.
- Q. Geng and P. Viswanath. Optimal noise adding mechanisms for approximate differential privacy. IEEE Trans. Inform. Theory, 62(2):952–969, 2016.
- [10] M. Hayashi. Quantum Information Theory: Mathematical Foundation, Second Edition. Springer, Berlin, Heidelberg, 2017.

- [11] M. Hayashi, S. Ishizaka, A. Kawachi, G. Kimura, and T. Ogawa. Introduction to Quantum Information Science. Springer, Berlin, Heidelberg, 2015.
- [12] N. Holohan, D. J. Leith, and O. Mason. Extreme points of the local differential privacy polytope. *Linear Algebra Appl.*, 534:78–96, 2017.
- [13] N. Holohan, D. J. Leith, and O. Mason. Optimal differentially private mechanisms for randomised response. *IEEE Trans. Inf. Forensics Secur.*, 12(11):2726–2735, 2017.
- [14] P. Kairouz, S. Oh, and P. Viswanath. Extremal mechanisms for local differential privacy. J. Mach. Learn. Res., 17:Paper No. 17, 51, 2016.
- [15] S. L. Warner. Randomized response: a survey technique for eliminating evasive answer bias. J. Amer. Statist. Assoc., 60(309):63-69, 1965.
- [16] Y. Yoshida and M. Hayashi. Classical mechanism is optimal in classical-quantum differentially private mechanisms. In 2020 IEEE International Symposium on Information Theory (ISIT), pages 1973–1977. 2020.
- [17] Y. Yoshida, M.-H. Yung, and M. Hayashi. Optimal mechanism for randomized responses under universally composable security measure. In 2019 IEEE International Symposium on Information Theory (ISIT), pages 547–551. 2019.
- [18] L. Zhou and M. Ying. Differential privacy in quantum computation. In 2017 IEEE 30th Computer Security Foundations Symposium (CSF), pages 249–262. 2017.

Yuuya Yoshida, Graduate School of Mathematics, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, 464-8602, Japan

Email address: m17043e@math.nagoya-u.ac.jp