On a cylinder freely floating in oblique waves

Nikolay Kuznetsov

Laboratory for Mathematical Modelling of Wave Phenomena, Institute for Problems in Mechanical Engineering, Russian Academy of Sciences, V.O., Bol'shoy pr. 61, St. Petersburg 199178, Russian Federation E-mail address: nikolay.g.kuznetsov@gmail.com

Abstract

The coupled motion is investigated for a mechanical system consisting of water and a body freely floating in it. Water occupies either a half-space or a layer of constant depth into which an infinitely long surface-piercing cylinder is immersed, thus allowing us to study the so-called oblique waves. Under the assumption that the motion is of small amplitude near equilibrium and describes time-harmonic oscillations, the phenomenon's linear setting reduces to a spectral problem with the radian frequency as the spectral parameter. If the radiation condition holds, then the total energy is finite and the equipartition of kinetic and potential energy holds for the whole system. On this basis, it is proved that no wave modes are trapped under some restrictions on their frequencies; in the case when a symmetric cylinder has two immersed parts restrictions are imposed on the type of mode as well.

1 Introduction

This paper continues the author's studies dealing with the motion of a mechanical system consisting of a water layer of constant depth and a rigid body freely floating in it; the initial note [1] was written on the occasion of V. M. Babich's 80th birthday, who recently turned 90. It was F. John [2], who proposed the linear problem describing the coupled motion of water bounded from above by the atmosphere and a partially immersed body. The latter floats freely according to Archimedes' law being unaffected by all external forces (for example due to constraints on its motion) except for gravity. The motion of water is assumed to be irrotational (viscosity is neglected as well as the surface tension), whereas the motion of the whole system is supposed to be of small amplitude near equilibrium; this allows us to use a linear model.

In the framework of the linear theory of water waves, two- and three-dimensional statements of the problem are possible; the one considered here is two-dimensional which is essential for formulation of restrictions to be imposed when investigating the question of uniqueness. The latter is of paramount importance (it is the first one on the list of important open problems in the survey [3]), because there are examples of nonuniqueness; see, for example, [1]. For this reason restrictions on the frequency range and on the body's shape are required. It should be noted that such restrictions are unnecessary for similar acoustical problems; see, for example, the monograph [4].

The original John's formulation of the floating body problem is rather cumbersome because he did not use matrices to express the equations of body's motion (the matrix form of these equations described below demonstrates their simple structure; see also [5], [6] and [7]). Anyway, the problem was neglected by researches during 60 years after publication of the article [8], in which the uniqueness theorem was proved for the threedimensional problem under the assumptions that the so-called John condition holds for the body (it is described below for two-dimensional geometry) and the frequency of oscillations is sufficiently large.

On the other hand, the problem of time-harmonic oscillations of water in the presence of fixed rigid bodies attracted much attention in the second half of the 20th century; initially, the case of a single body had been investigated in the article [8] mentioned above. Numerous results about this problem were presented in detail in the summarising monograph [9], which also contains extensive literature. In particular, the first non-uniqueness example due to M. McIver [10] was generalized in [9]. Namely, examples of non-trivial solutions were constructed for the two-dimensional homogeneous problem with an arbitrary number of fixed surface-piercing bodies (only two bodies were considered in [10]); the so-called inverse procedure was applied for this purpose. The presence of multiple bodies violates the original John's condition according to which only a single surfacepiercing body is admissible in the two-dimensional problem. From the hydrodynamic viewpoint, these non-trivial solutions describe trapped modes, that is, free oscillations of water having final energy. These results were developed further in the article [11], in which the uniqueness theorem due to John and McIver's non-uniqueness example were extended to the case of infinitely long, surface-piercing cylinders in oblique waves. The problem for fixed cylinders considered in [11] is a particular case of that formulated below in \S 2. Another approach to the question of uniqueness in the problem with fixed cylinders was proposed in [12].

2 Statement of the problem

Let an infinitely long surface-piercing cylinder of uniform cross-section float freely in water which is either infinitely deep or bounded from below by a horizontal rigid bottom. The Cartesian coordinate system (x, y) is chosen in a plane orthogonal to the cylinder's generators (directed along the z-axis), so that the y-axis is directed upwards, whereas the mean free surface of the water intersects this plane along the x-axis. Thus, the cross-section W of the water domain is a subset of $\mathbb{R}^2_- = \{x \in \mathbb{R}, y < 0\}$. Let \hat{B} denote the bounded two-dimensional domain whose closure is the cross-section of the cylinder's equilibrium position; we suppose that $\hat{B} \setminus \overline{\mathbb{R}^2_-}$ —the part of the body located above the water surface—is a nonempty domain, whereas the immersed part $B = \hat{B} \cap \mathbb{R}^2_-$ is the




Figure 1: A definition sketch of the cylinder's cross-section \widehat{B} with two immersed parts; they are denoted by B_{-} and B_{+} , and their wetted boundaries are S_{-} and S_{+} respectively. The cross-section F of the free surface of the water consists of three parts; two of them lying on the x-axis outside |x| > a are denoted by F_{∞} and the third one F_{0} is between x = -b and x = +b. Furthermore, $W = W_{0} \cup W_{\infty}$, where W_{0} is the part of the water domain located in the vertical strip under F_{0} and W_{∞} is its complement. The equation of the horizontal bottom is y = -h.

union of a finite number of domains. Thus, $D = \widehat{B} \cap \partial \mathbb{R}^2_-$ consists of the same number of nonempty intervals of the *x*-axis; see Fig. 1 for the case of two immersed parts, where

$$D = \{ x \in (-a, -b) \cup (b, a), y = 0 \}.$$

Notice that $W = \mathbb{R}^2_{-} \setminus \overline{B}$, when the water has infinite depth, or

$$W = \{x \in \mathbb{R}, \ -h < y < 0\} \setminus \overline{B}, \quad \text{where } h > b_0 = \sup_{(x,y) \in B} |y|,$$

when the water has the constant finite depth h (see Fig. 1). The cross-section of the bottom is denoted by $H = \{x \in \mathbb{R}, y = -h\}$ in the last case. Furthermore, W is assumed to be a Lipschitz domain, and so the unit normal **n** pointing to the exterior of W is defined almost everywhere on ∂W . Finally, we denote by $S = \partial \widehat{B} \cap \mathbb{R}^2_-$ the wetted contour (the number of its components is equal to the number of immersed domains), whereas $F = \partial \mathbb{R}^2_- \setminus \overline{D}$ is the free surface at rest.

To describe the small-amplitude coupled motion of the water/cylinder system it is standard to apply the linear setting, in which case the first-order approximation of unknowns is used. These are the velocity potential $\Phi(x, y, z; t)$ and the vector column $\mathbf{q}(t)$ describing the motion of the cylinder; its three components are as follows:

• q_1 and q_2 are the displacements of the centre of mass in the horizontal and vertical directions respectively from its rest position $(x^{(0)}, y^{(0)})$;

• q_3 is the angle of rotation about the axis that goes through the centre of mass orthogonally to the (x, y)-plane (the angle is measured from the x- to the y-axis).

We omit relations governing the time-dependent behaviour (details can be found in [1]) and turn to the description of time-harmonic oscillations of the coupled water/cylinder system in the presence of oblique waves for which purpose the following ansatz

$$\left(\Phi(x, y, z; t), \mathbf{q}(t)\right) = \operatorname{Re}\left\{\left(\operatorname{e}^{\operatorname{i}(kz - \omega t)}\varphi(x, y), \operatorname{ie}^{-\omega t}\mathbf{z}\right)\right\}$$
(1)

is applied. Here $\omega > 0$ is the radian frequency of oscillations, to which the wavenumber $\nu = \omega^2/g$ corresponds; g > 0 is the acceleration due to gravity that acts in the direction opposite to the *y*-axis. Furthermore, $k \in [0, \nu)$ is the prescribed wavenumber component that the wave train has parallel to the generators of the cylinder; $\varphi \in H^1_{loc}(W)$ is a complex-valued function and $\mathbf{z} \in \mathbb{C}^3$. Thus, k/ν is the sine of the angle between the wave crests and the plane normal to the generators; waves are called oblique when k > 0.

To be specific, we consider the case of infinitely deep W first. In the absence of incident waves, we obtain the following problem for (φ, \mathbf{z}) :

$$(\nabla^2 - k^2)\varphi = 0 \quad \text{in } W, \tag{2}$$

$$\partial_y \varphi - \nu \varphi = 0 \quad \text{on } F, \tag{3}$$

$$\partial_{\mathbf{n}}\varphi = \omega \mathbf{N}^{\mathsf{T}} \mathbf{z} \left(= \omega \sum_{1}^{3} N_{j} z_{j}\right) \quad \text{on } S,$$
(4)

$$\nabla \varphi \to 0 \quad \text{as } y \to -\infty,$$
 (5)

$$\omega^2 \mathbf{E} \mathbf{z} = -\omega \int_S \varphi \, \mathbf{N} \, \mathrm{d}s + g \, \mathbf{K} \mathbf{z}. \tag{6}$$

Here $\nabla = (\partial_x, \partial_y)$ is the spatial gradient, whereas $\mathbf{N} = (N_1, N_2, N_3)^{\mathsf{T}}$ (the operation $^{\mathsf{T}}$ transforms a vector row into a vector column and vice versa), where $(N_1, N_2)^{\mathsf{T}} = \mathbf{n}$, $N_3 = (x - x^{(0)}, y - y^{(0)})^{\mathsf{T}} \times \mathbf{n}$ and \times stands for the vector product. In the equations of the body motion (6), the 3×3 matrices are as follows:

$$\mathbf{E} = \begin{pmatrix} I^{M} & 0 & 0\\ 0 & I^{M} & 0\\ 0 & 0 & I_{2}^{M} \end{pmatrix} \quad \text{and} \quad \mathbf{K} = \begin{pmatrix} 0 & 0 & 0\\ 0 & I^{D} & I_{x}^{D}\\ 0 & I_{x}^{D} & I_{xx}^{D} + I_{y}^{B} \end{pmatrix}.$$
(7)

The positive elements of the mass/inertia matrix \mathbf{E} are as follows:

$$\begin{split} I^M &= \rho_0^{-1} \int_{\widehat{B}} \rho(x, y) \, \mathrm{d}x \mathrm{d}y \,, \\ I_2^M &= \rho_0^{-1} \int_{\widehat{B}} \rho(x, y) \Big[\left(x - x^{(0)} \right)^2 + \left(y - y^{(0)} \right)^2 \Big] \mathrm{d}x \mathrm{d}y \,. \end{split}$$

Here $\rho(x, y) \ge 0$ is the density distribution within the body and $\rho_0 > 0$ is the constant density of water. On the right-hand side of relation (6), we have forces and their moments: the first term is due to the hydrodynamic pressure and the second one is related to the buoyancy (see, for example, [2]). The non-zero elements of the matrix **K** are

$$I^{D} = \int_{D} dx > 0, \quad I^{D}_{x} = \int_{D} (x - x^{(0)}) dx,$$
$$I^{D}_{xx} = \int_{D} (x - x^{(0)})^{2} dx > 0, \quad I^{B}_{y} = \int_{B} (y - y^{(0)}) dx dy.$$

It should be noted that the matrix \mathbf{K} is symmetric.

In relations (3), (4) and (6), ω is a spectral parameter which is sought together with the eigenvector (φ , \mathbf{z}). Since W is a Lipschitz domain and $\varphi \in H^1_{loc}(W)$, relations (2)–(4) are understood in the sense of the following integral identity:

$$\int_{W} \nabla \varphi \nabla \psi \, \mathrm{d}x \mathrm{d}y = \nu \int_{F} \varphi \, \psi \, \mathrm{d}x + \omega \int_{S} \psi \, \mathbf{N}^{\mathsf{T}} \mathbf{z} \, \mathrm{d}s. \tag{8}$$

It must hold for an arbitrary smooth ψ having a compact support in \overline{W} .

Together with (5) the following condition

$$\int_{W \cap \{|x|=b\}} \left| \partial_{|x|} \varphi - \mathrm{i}\ell\varphi \right|^2 \mathrm{d}s = o(1) \quad \text{as } b \to \infty, \quad \ell = (\nu^2 - k^2)^{1/2}, \tag{9}$$

specifies the behaviour of φ at infinity; (5) means that the velocity field decays with depth, whereas (9) yields that the potential given by formula (1) describes outgoing waves. This radiation condition is similar to that used in [8], where the problem was considered for two-dimensional water waves in the presence of a fixed obstacle.

The relations listed above must be augmented by the following subsidiary conditions concerning the equilibrium position:

• Archimedes' law, $I^M = \int_B dx dy$ (the mass of the displaced liquid is equal to that of the body);

• $\int_B (x - x^{(0)}) dx dy = 0$ (the centre of buoyancy lies on the same vertical line as the centre of mass);

• The matrix **K** is positive semi-definite; moreover, the 2×2 matrix **K**' that stands in the lower right corner of **K** is positive definite (see [2]).

The last of these requirements yields stability of the body equilibrium position, which follows from the results formulated, for example, in the paper [2, § 2.4]. The stability is understood in the classical sense, that is, an instantaneous infinitesimal disturbance causes the position changes which remain infinitesimal, except for purely horizontal drift, for all subsequent times.

In conclusion of this section, we note that relations (5) and (9) must be amended in the case when W has finite depth. Namely, the no flow condition

$$\partial_u \varphi = 0 \quad \text{on } H \tag{10}$$

replaces (5), whereas ℓ must be changed to ℓ_0 in (9), where ℓ_0 is the unique positive root of $\ell_0 \tanh(\ell_0 h) = \ell$.

3 Equipartition of energy

It is known (see, for example, $[9, \S 2.2.1]$), that a potential, satisfying relations (2), (3), (5) and (9), has an asymptotic representation at infinity of the same type as Green's function. Namely, if W has infinite depth, then

$$\varphi(x,y) = A_{\pm}(y) e^{i\ell|x|} + r_{\pm}(x,y),$$

$$|r_{\pm}|^2, |\nabla r_{\pm}| = O([x^2 + y^2]^{-1}) \text{ as } x^2 + y^2 \to \infty,$$
 (11)

and the following equality holds

$$\ell \int_{-\infty}^{0} \left(|A_{+}(y)|^{2} + |A_{-}(y)|^{2} \right) \mathrm{d}y = -\mathrm{Im} \int_{S} \overline{\varphi} \,\partial_{\mathbf{n}} \varphi \,\mathrm{d}s.$$
(12)

Assuming that (φ, \mathbf{z}) is a solution of the problem (2)–(6) and (9), we rearrange the last formula using the coupling conditions (4) and (6). First, transposing the complex conjugate of equation (6), we obtain

$$\omega^{2} \left(\mathbf{E} \overline{\mathbf{z}} \right)^{\mathsf{T}} = -\omega \int_{S} \overline{\varphi} \mathbf{N}^{\mathsf{T}} \, \mathrm{d}s + g \left(\mathbf{K} \overline{\mathbf{z}} \right)^{\mathsf{T}}$$

This relation and condition (4) yield that the inner product of both sides with \mathbf{z} can be written in the form

$$\omega^2 \,\overline{\mathbf{z}}^\mathsf{T} \mathbf{E} \mathbf{z} - g \overline{\mathbf{z}}^\mathsf{T} \mathbf{K} \mathbf{z} = -\int_S \overline{\varphi} \,\partial_\mathbf{n} \varphi \,\mathrm{d}s. \tag{13}$$

Second, substituting this equality into (12), we obtain

$$\ell \int_{-\infty}^{0} \left(|A_{+}(y)|^{2} + |A_{-}(y)|^{2} \right) \mathrm{d}y = \mathrm{Im} \Big\{ \omega^{2} \, \overline{\mathbf{z}}^{\mathsf{T}} \mathbf{E} \mathbf{z} - g \overline{\mathbf{z}}^{\mathsf{T}} \mathbf{K} \mathbf{z} \Big\}.$$
(14)

In the same way as in [6], this yields the following assertion about the kinetic and potential energy of the water motion.

Proposition 1. Let (φ, \mathbf{z}) be a solution of problem (2)–(6) and (9), then

$$\int_{W} \left(|\nabla \varphi|^2 + k^2 |\varphi|^2 \right) \mathrm{d}x \,\mathrm{d}y < \infty \quad and \quad \nu \int_{F} |\varphi|^2 \,\mathrm{d}x < \infty \,, \tag{15}$$

that is, $\varphi \in H^1(W)$. Moreover, the following equality holds:

$$\int_{W} \left(|\nabla \varphi|^{2} + k^{2} |\varphi|^{2} \right) \mathrm{d}x \,\mathrm{d}y + \omega^{2} \overline{\mathbf{z}}^{\mathsf{T}} \mathbf{E} \mathbf{z} = \nu \int_{F} |\varphi|^{2} \,\mathrm{d}x + g \,\overline{\mathbf{z}}^{\mathsf{T}} \mathbf{K} \mathbf{z}.$$
(16)

Here the kinetic energy of the water/body system stands on the left-hand side, whereas we have the potential energy of this coupled motion on the right-hand side. Thus the last formula generalises the energy equipartition equality valid when a fixed body is immersed into water. Indeed, \mathbf{z} vanishes for such a body, and (16) turns into the well-known equality (see, for example, formula (4.99) in [9]).

Proposition 1 shows that if (φ, \mathbf{z}) is a solution of problem (2)–(6) and (9) with complex-valued components, then its real and imaginary parts separately satisfy this problem. This allows us to consider (φ, \mathbf{z}) as an element of the real product space $H^1(W) \times \mathbb{R}^3$ in what follows.

Definition 1. Let the subsidiary conditions concerning the equilibrium position (see § 2) hold for the freely floating body \widehat{B} . A non-trivial real solution $(\varphi, \mathbf{z}) \in H^1(W) \times \mathbb{R}^3$ of problem (8) and (6) is called a *mode trapped* by this body, whereas the corresponding value of ω is referred to as a *trapping frequency*.

In the case of finite depth, the remainder in formula (11) has the following behaviour uniformly in $y \in [-h, 0]$:

$$|r_{\pm}(x,y)|, |\nabla r_{\pm}(x,y)| = O(|x|^{-1}) \text{ as } |x| \to \infty,$$

whereas formula (12) holds with ℓ changed to ℓ_0 . Therefore, Proposition 1 is true for problem (2)–(6) and (9) with condition (5) replaced by (10) and ℓ changed to ℓ_0 in (9). Definition 1 remains unchanged for the finite depth case.

4 On the absence of trapped modes

In order to determine conditions on the domain \widehat{B} and on the frequency ω guaranteeing the absence of non-trivial solutions $(\varphi, \mathbf{z}) \in H^1(W) \times \mathbb{R}^3$, we write (16) as follows:

$$\mathbf{z}^{\mathsf{T}}(\omega^{2}\mathbf{E} - g\,\mathbf{K})\mathbf{z} = \nu \int_{F} |\varphi|^{2}\,\mathrm{d}x - \int_{W} |\nabla\varphi|^{2}\,\mathrm{d}x\mathrm{d}y.$$
(17)

It is clear that the left-hand side is non-negative for a nonzero \mathbf{z} provided $\omega^2 \geq \lambda_0$; the latter is the largest λ satisfying det $(\lambda \mathbf{E} - g\mathbf{K}) = 0$. Let the equilibrium position be stable for the cylinder, whose cross-section is \widehat{B} (see § 2 for conditions of stability), it is convenient to say that property Ω holds for ω if $\omega^2 \geq \lambda_0$. Thus, we arrive at the following.

Proposition 2. Let the cylinder's cross-section be \widehat{B} and let property Ω hold for ω . If the inequality

$$\nu \int_{F} |\varphi|^2 \,\mathrm{d}x < \int_{W} \left(|\nabla \varphi|^2 + k^2 |\varphi|^2 \right) \,\mathrm{d}x \mathrm{d}y \,, \tag{18}$$

holds for every non-trivial $\varphi \in H^1(W)$ satisfying (8) and (6), then ω is not a trapping frequency.

Indeed, for a non-trivial (φ, \mathbf{z}) inequality (18) is incompatible with property Ω .

4.1 Cylinders with single immersed part

For the sake of simplicity we suppose that W has infinite depth. We recall that John's condition requires a simply connected domain $\widehat{B} \cap \mathbb{R}^2_-$ to belong to the strip lying between the verticals going through the points, where the contour $\partial \widehat{B}$ intersects the x-axis.

Theorem 1. Let W have infinite depth and let the domain $\widehat{B} \cap \mathbb{R}^2_{-}$ be simply connected and satisfy John's condition. Let also the subsidiary conditions concerning the equilibrium position (see § 2) hold. Then inequality (18) holds for a non-trivial solution $(\varphi, \mathbf{z}) \in H^1(W) \times \mathbb{R}^3$ of problem (8) and (6).

Proof. Since $\widehat{B} \cap \mathbb{R}^2_{-}$ is simply connected, D consists of a single interval, and so the intersection of the free surface and the (x, y)-plane is $F = F_{\infty}$ (see Fig. 1); moreover,

 $F = F_+ \cup F_-$, where F_+ (F_-) is the ray lying on the x-axis to the right (left) of D. Let us prove the inequality

$$\nu \int_{F_{\pm}} |\varphi|^2 \,\mathrm{d}x < \int_{W_{\pm}} \left(|\nabla \varphi|^2 + k^2 |\varphi|^2 \right) \,\mathrm{d}x \mathrm{d}y \,, \tag{19}$$

where $W_{\pm} \subset W$ is the subdomain lying strictly under F_{\pm} ; in view of John's condition these subdomains are well defined.

Following John let us define

$$a^{(\pm)}(x) = \int_{-\infty}^{0} \varphi(x, y) e^{\nu y} dy$$
 on F_{\pm} . (20)

Differentiating this function twice and using equation (2), we obtain

$$a_{xx}^{(\pm)} = k^2 a^{(\pm)} - \int_{-\infty}^{0} \varphi_{yy}(x, y) e^{\nu y} dy.$$

Integrating by parts twice and taking into account conditions (3) and (5), we conclude that $a_{xx}^{(\pm)} + \ell^2 a^{(\pm)} = 0$ on F_{\pm} because $k^2 - \nu^2 = -\ell^2$.

Since $\varphi \in H^1(W)$, we have that $\lim_{|x|\to\infty} a^{(\pm)}(x) = 0$, and so $a^{(\pm)} \equiv 0$ on F_{\pm} . Now, integrating by parts in (20), we see that

$$\varphi(x,0) = \int_{-\infty}^0 \varphi_y(x,y) \,\mathrm{e}^{\nu y} \,\mathrm{d}y \quad \text{for } (x,0) \in F_{\pm} \,.$$

Squaring both sides, applying the Schwarz inequality to the integral and integrating over F_{\pm} , we obtain

$$\nu \int_{F_{\pm}} |\phi^{(\pm)}(x,0)|^2 \,\mathrm{d}x \le \frac{1}{2} \int_{W_{\pm}} |\varphi_y|^2 \,\mathrm{d}x \mathrm{d}y < \int_{W_{\pm}} |\nabla \varphi|^2 \,\mathrm{d}x \mathrm{d}y;$$

here the coefficient 1/2 results from integration of $e^{2\nu y}$. The last inequality is stronger than (19), and so it implies (18) because $F = F_+ \cup F_-$. \Box

Propositions 1 and 2, and Theorem 1 yield.

Corollary 1. Let W have infinite depth and let the domain $\widehat{B} \cap \mathbb{R}^2_{-}$ be simply connected and satisfy John's condition. Let also the subsidiary conditions concerning the equilibrium position (see § 2) hold. If property Ω holds for ω , then problem (2)–(6) and (9) has only a trivial solution for this ω .

Analogous assertion is true when the water domain has finite depth.

4.2 Cylinders with two immersed parts

In the case of a cylinder with two immersed parts (see Fig. 1), we are going to apply the trick based on John's condition which was used in § 4.1. This is possible if the cylinder's cross-section is symmetric about a vertical axis (the y-axis in Fig. 1), the density distribution ρ is also symmetric within the cylinder and each of the domains B_+ and B_- is simply connected and satisfies John's condition, that is, $\beta \geq \pi/2$ (see Fig. 1). Let also the subsidiary conditions concerning the equilibrium position (see § 2) hold. Moreover, we restrict the cylinder's admissible motions to sway, heave and roll, and consider separately the classes of velocity potentials consisting of odd and even functions of x.

4.2.1 Sway motion

In this case $\mathbf{z} = (z_1, 0, 0)^{\mathsf{T}}, z_1 \in \mathbb{R}$, and so the boundary condition (4) takes the form

$$\partial_{\mathbf{n}}\varphi = \omega N_1 z_1 \quad \text{on } S \,, \tag{21}$$

and system (6) splits. Its first equation is as follows

$$\omega I^M z_1 = -\int_S \varphi N_1 \,\mathrm{d}s\,,\tag{22}$$

whereas the second and third equations turn into the orthogonality conditions:

$$\int_{S} \varphi N_j \,\mathrm{d}s = 0 \,, \quad j = 2, 3. \tag{23}$$

Thus, we have to determine frequency intervals for which the equality

$$I^{M}(\omega z_{1})^{2} = \nu \int_{F} |\varphi|^{2} \,\mathrm{d}x - \int_{W} \left(|\nabla \varphi|^{2} + k^{2} |\varphi|^{2} \right) \,\mathrm{d}x \mathrm{d}y \tag{24}$$

obtained from (17) cannot hold for a non-trivial φ .

Let us assume that the domains B_+ and B_- are simply connected and each satisfies John's condition, that is, $\beta \ge \pi/2$ (see Fig. 1). Repeating literally considerations used in § 4.1, we arrive at the inequality:

$$\nu \int_{F_{\infty}} |\varphi|^2 \,\mathrm{d}x < \int_{W_{\infty}} \left(|\nabla \varphi|^2 + k^2 |\varphi|^2 \right) \,\mathrm{d}x \mathrm{d}y \,. \tag{25}$$

It remains to estimate $\nu \int_{F_0} |\varphi|^2 dx$, and for this purpose we consider the function

$$a(x) = \int_{-\infty}^{0} \varphi(x, y) e^{\nu y} dy, \qquad (26)$$

which, in view of John's condition, is well defined for $x \in F_0$. In the same way as in § 4.1, we obtain that it satisfies the equation $a_{xx} + \ell^2 a = 0$ on F_0 .

Assuming that $\varphi(x, y)$ is odd in x, we get that $a(x) = C \sin \ell x$, which we substitute into (26) and differentiate. After squaring both obtained equalities, we apply the Schwarz

inequality to the integrals. In this way, we find that for every $(x, 0) \in F_0$:

$$2\nu C^{2} \sin^{2} \ell x \leq \int_{-\infty}^{0} |\varphi(x,y)|^{2} \,\mathrm{d}y\,, \qquad (27)$$

$$2\nu \ell^2 C^2 \cos^2 \ell x \le \int_{-\infty}^0 |\varphi_x(x,y)|^2 \,\mathrm{d}y \,. \tag{28}$$

Furthermore, integration by parts in (26) yields

$$\varphi(x,0) = -\nu C \sin \ell x + \int_{-\infty}^{0} \varphi_y(x,y) e^{\nu y} dy$$

for every $(x, 0) \in F_0$, and so

$$\nu |\varphi(x,0)|^2 \le 2\nu^3 C^2 \sin^2 \ell x + \int_{-\infty}^0 |\varphi_y(x,y)|^2 \,\mathrm{d}y \,. \tag{29}$$

Integrating this over F_0 and using (27), we obtain

$$\nu \int_{F_0} |\varphi(x,0)|^2 \, \mathrm{d}x \le 2\nu (k^2 + \ell^2) C^2 \int_{F_0} \sin^2 \ell x \, \mathrm{d}x + \int_{W_0} |\varphi_y|^2 \, \mathrm{d}x \mathrm{d}y \\
\le k^2 \int_{W_0} |\varphi|^2 \, \mathrm{d}x \mathrm{d}y + \int_{W_0} |\varphi_y|^2 \, \mathrm{d}x \mathrm{d}y + 2\nu \ell^2 C^2 \int_{F_0} \sin^2 \ell x \, \mathrm{d}x \,. \quad (30)$$

Let us say that property Ω_{-} holds for ω if the inequalities

$$\pi m \le \ell \, b \le \pi (2m+1)/2$$

are valid for some m = 0, 1, ...; we recall that $\ell = (\nu^2 - k^2)^{1/2}$, $\nu = \omega^2/g$ and 2b is the spacing between B_+ and B_- .

Since property Ω_{-} is equivalent to the inequality

$$\int_0^b \sin^2 \ell x \, \mathrm{d}x \le \int_0^b \cos^2 \ell x \, \mathrm{d}x \,,$$

we can estimate the last term in (30) with the help of (28) provided this property holds. In this way, we arrive at the estimate

$$\nu \int_{F_0} |\varphi|^2 \, \mathrm{d}x \le \int_{W_0} \left(|\nabla \varphi|^2 + k^2 |\varphi|^2 \right) \, \mathrm{d}x \mathrm{d}y \,,$$

which combined with (25) leads to a contradiction with equality (24), unless (φ, \mathbf{z}) is trivial. This completes the proof of the following.

Proposition 3. Let W have infinite depth and let the domain \widehat{B} be symmetric about the y-axis and such that $\widehat{B} \cap \mathbb{R}^2_{-}$ is the union of two simply connected domains each satisfying John's condition. Let also the subsidiary conditions concerning the equilibrium position (see § 2) hold.

If property Ω_{-} holds for ω , then problem (2)–(6) and (9) has only a trivial solution (φ, \mathbf{z}) for this ω provided φ is odd in x and $\mathbf{z} = (z_1, 0, 0)^{\mathsf{T}}$.

Let the motion of water in the presence of a symmetric cylinder floating freely (see Fig. 1) be described by φ even in x. Then equation (22) implies that $z_1 = 0$, which means that such a potential does not comport with the free sway motion of a symmetric cylinder. However, there are examples of motionless symmetric cylinders with two immersed parts that trap modes at particular frequencies; see [11], §§ 5 and 6.

Now, equality (24) turns into

$$\nu \int_{F} |\varphi|^2 \,\mathrm{d}x = \int_{W} \left(|\nabla \varphi|^2 + k^2 |\varphi|^2 \right) \,\mathrm{d}x \mathrm{d}y \,, \tag{31}$$

and so the uniqueness theorem proved in § 3 of [11], where the case of fixed B_- and B_+ was discussed, is applicable to the problem under consideration here. The corresponding assertion formulated below is analogous to Proposition 3, but involves property Ω_+ instead of Ω_- ; namely, property Ω_+ holds for ω if the inequalities

$$\pi(2m+1)/2 \le \ell b \le \pi(m+1)$$
, where $\ell = (\nu^2 - k^2)^{1/2}$ and $\nu = \omega^2/g$,

take place for some $m = 0, 1, \ldots$ Thus, these properties are of the same kind; either of them is valid when ℓb belongs to one interval in a sequence; these sequences for Ω_+ and Ω_- are complementary and their intervals have common endpoints.

Proposition 4. Let W have infinite depth and let the domain \hat{B} satisfy the assumptions of Proposition 3. If property Ω_+ holds for ω , then problem (2)–(6) and (9) has only a trivial solution (φ, \mathbf{z}) for this ω provided φ is even in x and $\mathbf{z} = (z_1, 0, 0)^{\mathsf{T}}$.

Comparing Propositions 3 and 4 with the uniqueness theorem proved in [11], we see that for a cylinder with two immersed parts the frequency intervals, where the potential (either odd or even in x) is trivial, are the same irrespective whether the cylinder floats freely or is fixed and depend only on the potential's parity.

4.2.2 Heave motion

In this case $\mathbf{z} = (0, z_2, 0)^{\mathsf{T}}, z_2 \in \mathbb{R}$, and so the boundary condition (4) takes the form

$$\partial_{\mathbf{n}}\varphi = \omega N_2 z_2 \quad \text{on } S \,, \tag{32}$$

and system (6) splits. Its second equation is as follows

$$\left(\omega^2 I^M - g I^D\right) z_2 = -\omega \int_S \varphi \, N_2 \, \mathrm{d}s \,, \tag{33}$$

whereas the second and third equations turn into the orthogonality conditions

$$\int_{S} \varphi N_1 \,\mathrm{d}s = 0 \,, \quad \int_{S} \varphi N_3 \,\mathrm{d}s = 0 \,. \tag{34}$$

The last one is a consequence of the equality $I_x^D = 0$, which follows from the symmetry about the *y*-axis; indeed, $x^{(0)} = 0$ in this case. Furthermore, relation (17) takes the form:

$$\left(\omega^2 I^M - gI^D\right) z_2^2 = \nu \int_F |\varphi|^2 \,\mathrm{d}x - \int_W \left(|\nabla \varphi|^2 + k^2 |\varphi|^2\right) \,\mathrm{d}x \mathrm{d}y \,. \tag{35}$$

Assuming that $\varphi(x, y)$ is odd in x, we see that the right-hand side of (33) vanishes because the integrand has the same parity as φ . Then the last equality turns into (31), and so we are in a position to refer to the considerations used in [11], § 3. In this way, we arrive at the following assertion (it is similar to Propositions 3 and 4) for the present problem.

Proposition 5. Let W have infinite depth and let the domain \widehat{B} satisfy the assumptions of Proposition 3. If property Ω_{-} holds for ω , then problem (2)–(6) and (9) has only a trivial solution (φ, \mathbf{z}) for this ω provided φ is odd in x and $\mathbf{z} = (0, z_2, 0)^{\mathsf{T}}$.

Proof. According to considerations in [11], § 3, property Ω_{-} guarantees that φ (satisfying (31) and odd in x) is trivial. Then the boundary condition (32) yields that \mathbf{z} is also trivial. \Box

Let us turn to the case when $\varphi(x, y)$ is even in x and apply the approach used in the proof of Proposition 3 for determining the frequency intervals when equality (35) cannot hold if φ is non-trivial. Considerations based on property Ω_+ show that the right-hand side of (35) is strictly negative when this property is valid. On the other hand, the left-hand side of this equality is nonnegative if $\omega^2 \geq gI^D/I^M$; this yields the following.

Proposition 6. Let W have infinite depth and let the domain \widehat{B} satisfy the assumptions of Proposition 3. If property Ω_+ holds for $\omega \ge \sqrt{gI^D/I^M}$, then problem (2)–(6) and (9) has only a trivial solution (φ, \mathbf{z}) for this ω provided φ is even in x and $\mathbf{z} = (0, z_2, 0)^{\mathsf{T}}$.

The essential distinction between this assertion and Proposition 5 is as follows. Along with property Ω_+ , the assumptions of Proposition 6 include the inequality $\omega^2 \geq gI^D/I^M$, whereas no condition other than property Ω_- is imposed in Proposition 5.

4.2.3 Roll motion

In this case $\mathbf{z} = (0, 0, z_3)^{\mathsf{T}}, z_3 \in \mathbb{R}$, and so the boundary condition (4) takes the form

$$\partial_{\mathbf{n}}\varphi = \omega N_3 z_3 \quad \text{on } S \,, \tag{36}$$

and system (6) splits. The first and second equations turn into the orthogonality conditions

$$\int_{S} \varphi N_1 \,\mathrm{d}s = 0 \,, \quad \int_{S} \varphi N_2 \,\mathrm{d}s = 0 \,. \tag{37}$$

Again, the last one is a consequence of the equality $I_x^D = 0$, which follows from symmetry of \hat{B} about the *y*-axis. The third equation is as follows

$$\left[\omega^2 I_2^M - g\left(I_{xx}^D + I_y^B\right)\right] z_3 = -\omega \int_S \varphi N_3 \,\mathrm{d}s\,,\tag{38}$$

where $N_3(x,y) = xN_2 - (y - y^{(0)})N_1$ and $N_3(x,y) = -N_3(-x,y)$ for $(\pm x, y) \in S_{\pm}$ in view of symmetry. Now, relation (17) takes the form:

$$\left[\omega^{2}I_{2}^{M} - g\left(I_{xx}^{D} + I_{y}^{B}\right)\right]z_{3}^{2} = \nu \int_{F} |\varphi|^{2} \,\mathrm{d}x - \int_{W} \left(|\nabla\varphi|^{2} + k^{2}|\varphi|^{2}\right) \,\mathrm{d}x \,\mathrm{d}y \,. \tag{39}$$

Assuming that $\varphi(x, y)$ is even in x, we see that the right-hand side of (38) vanishes (cf. § 4.2.2), and the last equality turns into (31). Again we are in a position to refer to the considerations used in [11], § 3, which lead to.

Proposition 7. Let W have infinite depth and let the domain \hat{B} satisfy the assumptions of Proposition 3. If property Ω_+ holds for ω , then problem (2)–(6) and (9) has only a trivial solution (φ, \mathbf{z}) for this ω provided φ is even in x and $\mathbf{z} = (0, 0, z_3)^{\mathsf{T}}$.

Comparing this assertion and Proposition 4, we see that for a freely floating symmetric cylinder with two immersed parts property Ω_+ guarantees uniqueness at all frequencies satisfying it provided φ is even in x and the cylinder executes either sway or roll motion.

In the case when $\varphi(x, y)$ is odd in x, we again apply the approach used for the proof of Proposition 6. Since $I_x^D = 0$ in view of symmetry of \hat{B} , the assumption that \mathbf{K}' is a positive definite matrix (see § 2) yields the inequality $I_{xx}^D + I_y^B > 0$. Thus we obtain.

Proposition 8. Let W have infinite depth and let the domain \widehat{B} satisfy the assumptions of Proposition 3. If property Ω_{-} holds for

$$\omega \ge \sqrt{g \left(I_{xx}^D + I_y^B \right) / I_2^M} \,, \tag{40}$$

then problem (2)–(6) and (9) has only a trivial solution (φ, \mathbf{z}) for this ω provided φ is odd in x and $\mathbf{z} = (0, 0, z_3)^{\mathsf{T}}$.

As in the case of Propositions 5 and 6, the distinction between Propositions 7 and 8 is that the latter one includes the extra inequality (40).

In conclusion of this section, it should be mentioned that there are analogues of Propositions 3–8 for water of finite depth; their formulations are similar to those above.

5 Discussion

First, the obtained results guarantee uniqueness of a solution in proper classes of functions for the problem of scattering of obliquely incoming plane waves by an infinitely long cylinder floating freely. This problem is left for future research.

Second, equations (33) and (38) demonstrate that one may expect the existence of trapped modes for some particular bodies in heave and roll motion. To find these bodies and the corresponding trapped modes is another interesting problem for future research. A hint to its solution (at least for k = 0) can be found in the article [7].

References

- N. Kuznetsov, On the problem of time-harmonic water waves in the presence of a freely-floating structure. — St Petersburg Math. J. 22 (2011), 985–995.
- [2] F. John, On the motion of floating bodies, I. Comm. Pure Appl. Math. 2 (1949), 13–57.

- [3] F. Ursell, Some unsolved and unfinished problems in the theory of waves. Wave Asymptotics, eds. P.A. Martin, G.R. Wickham, Cambridge, Cambridge University Press (1992), 220–244.
- [4] D. Colton, R. Kress, Integral Equation Methods in Scattering Theory. Philadelphia, SIAM (2013).
- [5] N. Kuznetsov, O. Motygin, On the coupled time-harmonic motion of water and a body freely floating in it. — J. Fluid Mech. 679 (2011), 616–627.
- [6] N. Kuznetsov, O. Motygin, On the coupled time-harmonic motion of deep water and a freely floating body: trapped modes and uniqueness theorems. — J. Fluid Mech. 703 (2012), 142–162.
- [7] N. Kuznetsov, O. Motygin, Freely floating structures trapping time-harmonic water waves. — Quart. J. Mech. Appl. Math. 68 (2015), 173–193.
- [8] F. John, On the motion of floating bodies, II. Comm. Pure Appl. Math. 3 (1950), 45–101.
- [9] N. Kuznetsov, V. Maz'ya, B. Vainberg, *Linear Water Waves: A Mathematical Approach.* Cambridge, Cambridge University Press (2002).
- [10] M. McIver, An example of non-uniqueness in the two-dimensional linear water-wave problem. — J. Fluid Mech. **315** (1996), 257–266.
- [11] N. Kuznetsov, R. Porter, D. V. Evans, M. J. Simon, Uniqueness and trapped modes for surface-piercing cylinders in oblique waves. — J. Fluid Mech. 365 (1998), 351– 368.
- [12] N. Kuznetsov, Uniqueness in the problem of an obstacle in oblique waves. C. R. Mecanique. 331 (2003), 183–188.
- [13] N. Kuznetsov, O. Motygin, On the coupled time-harmonic motion of water and a body freely floating in it. — J. Fluid Mech. 679 (2011), 616–627.