CLASSES OF ALGEBRAS AND CLOSURE OPERATIONS

I. S. GUTIERREZ, ANSELMO TORRESBLANCA-BADILLO, AND DAVID A. TOWERS

In memory of Prof. Dr. Klaus Doerk

ABSTRACT. The calculus of classes and closure operations has proved to be a useful tool in group theory and has led to a deep theory in the study of finite soluble groups. More recently, parallel theories have started to be developed in various varieties of algebras, such as Lie, Leibniz and Malcev algebras. This paper seeks to investigate the extent to which these later theories can be generalised to the variety of all non-associative algebras.

1. INTRODUCTION

In the early 1960s Philip Hall introduced a systematic use of closure operations into group theory (see [8], [9]). This was followed by the development of the concept of a formation by Gaschütz in 1963. Since then, these concepts have contributed significantly to the theory of finite soluble groups, as is expounded in [6] and [1].

In recent years, closure operators have been more widely applied in various mathematical systems. In particular, a parallel theory has been developed in the context of a universal algebra (see [11], [12], [13], for instance), and in various varieties of algebras, such as Lie algebras (see [3], [2], [14], for example), Leibniz algebras (see [5] and [4], for example) and Malcev algebras ([15]). It is the purpose of this paper to investigate the extent to which some of these last three sets of results can be generalised to the variety of all non-associative algebras.

In section 2 we introduce the concepts of classes of algebras and of closure operations and present their basic properties. In section 3 we introduce a partial order on closure operations, define the join of a set of closure operations and show that the join of various such operations are again closure operations. In the fourth section the concepts of homomorph, formation, saturation, Schunck class and Fitting class are presented, and some equivalent conditions for a non-empty homomorph and for a non-empty formation to be saturated are proved. The final section is devoted to conclusions and ideas for further work.

Throughout this work, unless otherwise stated, all of the algebras considered are assumed to be non-associative algebras over a field K.

Date: January 12, 2022.

²⁰¹⁰ Mathematics Subject Classification. Primary 68R05; Secondary 05C69.

Key words and phrases. Non-associative algebras, finite-dimensional algebra, solvable algebras, nilpotent algebras, supersolvable algebras, classes of algebras, formations, Schunck classes, closure operations.

2 I. S. GUTIERREZ, ANSELMO TORRESBLANCA-BADILLO, AND DAVID A. TOWERS

An algebra is said to be simple if it has only the trivial ideals 0 and A and it is not the one-dimensional algebra. The core of a subalgebra H of A with respect to A, denoted by $\operatorname{Core}_A(H)$, is defined to be the largest ideal of A contained in H. An algebra is called primitive if it has a core-free maximal subalgebra.

If $S \subseteq A$, then the ideal of A generated by S or the ideal closure of S in A is defined to be the unique smallest ideal of A containing S. That is, the intersection of all ideals of A which contain S, and it is denoted like in group theory by S^A . It is straightforward to check that

(1.1)
$$S^{A} = \left\{ \sum_{s \in S} \left(\sum_{k=1}^{n_{s}} a_{k} s y_{k} \right) \mid n_{s} \in \mathbb{N}, \ x_{k}, y_{k} \in A \right\}.$$

A subspace I is called a subideal of A if there exist a chain of subalgebras

(1.2)
$$I = I_0 < I_1 < \dots < I_n = A,$$

where I_j is an ideal of I_{j+1} for each $0 \le j \le n-1$.

The Frattini subalgebra of A, denoted by F(A), is defined as the intersection of the maximal subalgebras of A; if A has no maximal subalgebras then it is defined to be A. E. I. Marshall [10] proved that for finite-dimensional Lie algebras over an algebraically closed field of characteristic zero the Frattini subalgebra is an ideal and D. A. Towers [16] showed that the restriction of algebraic closure can be removed to obtain that it is a characteristic ideal. However, in general, the Frattini subalgebra need not be an ideal, so we define the Frattini ideal, $\Phi(A)$, to be equal to $\operatorname{Core}_A(F(A))$.

Set $A^1 = A^{(1)} = A$, and then for $n \ge 1$ define inductively

(1.3)
$$A^{n+1} = \sum_{i+j=n+1} A^i A^j$$

and

(1.4)
$$A^{(n+1)} = A^{(n)}A^{(n)} = (A^{(n)})^2.$$

The subset A^n is called the *n*-th power of A. The chain of subsets

(1.5)
$$A^1 \supseteq A^2 \supseteq \cdots \supseteq A^n \supseteq \cdots$$

is a chain of ideals of A. A non-associative algebra A is called **nilpotent** if $A^n = 0$ for some n and **solvable** if $A^{(r)} = 0$ for some r. The smallest natural number n (respectively r) with this property is called the **nilpotency index** (respectively **solvability index**) of A. Note that the elements of the chain

(1.6)
$$A^{(1)} \supseteq A^{(2)} \supseteq \cdots \supseteq A^{(n)} \supseteq \cdots$$

beyond $A^{(2)}$, are not, generally speaking, ideals in A.

Motivated by the concept of supersolubility in group theory a similar concept has been introduced in Lie algebras, where normal subgroups with cyclic factors were replaced by ideals with one dimensional factors. In general, we say that an algebra A is called **supersolvable** if it is solvable and has a flag of ideals

$$0 = A_0 \le A_1 \le \ldots \le A_n = A$$

where dim $A_i = i$ for $0 \le i \le n$. For a general non-associative algebra A, the added condition that A be solvable is needed, as the one-dimensional algebra spanned by a single idempotent has such a flag, trivially, but is not solvable.

If A is an algebra over a field K, an A-bimodule is a vector space M over K with bilinear mappings

 $A \times M \to M, (x, m) \mapsto xm \text{ and } M \times A \to M, (m, x) \mapsto mx.$

The split null extension of A by $M, A \rtimes M$, is the algebra over K with underlying vector space $A \oplus M$ and multiplication

$$(x+m)(y+n) = xy + (xn+my)$$
 for all $x, y \in A, m, n \in M$.

The regular bimodule, Reg(A), is the underlying vector space of A considered as a bimodule with mx and xm as the multiplication in A.

2. Classes of Algebras and Closure Operations

A class of algebras \mathfrak{X} is a collection of algebras with the property that if $A \in \mathfrak{X}$, then every algebra isomorphic to A belongs to \mathfrak{X} . The algebras which belong to a class \mathfrak{X} are called \mathfrak{X} -algebras.

Following K. Doerk and T. O. Hawkes [6] and A. Ballester-Bolinches and L. Ezquerro [1], we denote the empty class of algebras by \emptyset and we use Gothic font when a single capital letter denotes a class of algebras.

The following are examples of specific classes of algebras:

- \emptyset denotes the empty class of algebras
- \mathfrak{A} denotes the class of all abelian algebras
- ${\mathfrak N}$ –denotes the class of all nilpotent algebras
- \mathfrak{N}_c denotes the class of all nilpotent algebras of index $\leq c, c \in \mathbb{N}$
- \mathfrak{S} denotes the class of all solvable algebras
- \mathfrak{U} denotes the class of all supersolvable algebras
- \mathfrak{E} denotes the class of all finite dimensional algebras.

Algebra classes are partially ordered by inclusion and we write $\mathfrak{X} \subseteq \mathfrak{Y}$ to denote that the class \mathfrak{X} is a subclass of the class \mathfrak{Y} .

A map which sends a class of algebras to a class of algebras is called a class map. Among class maps are the so-called **closure operations**, which play a central role in studying of algebra classes.

A class map c assigning to each class of algebras \mathfrak{X} a class of algebras $c\mathfrak{X}$ is called an **operation**, if it holds the following conditions:

- (C1) $\mathsf{c}\emptyset = \emptyset$ and
- (C2) $\mathfrak{X} \subseteq c\mathfrak{X} \subseteq c\mathfrak{Y}$, providing $\mathfrak{X} \subseteq \mathfrak{Y}$. (We say c is expanding and monotonic respectively).

An operation c is called a closure operation, if it is idempotent, that is, if

(C3) $c\mathfrak{X} = c(c\mathfrak{X})$, for all class \mathfrak{X} .

Definition 2.1. Let \mathfrak{X} be a class of algebras. Then the following are important examples of class maps:

$$\begin{split} \mathbf{s} & \mathbf{\mathfrak{X}} = (A \mid A \leq K, \text{ for some } K \in \mathbf{\mathfrak{X}}) \\ \mathbf{s}_n & \mathbf{\mathfrak{X}} = (A \mid A \text{ a subideal of } K \text{ for some } K \in \mathbf{\mathfrak{X}}) \\ & \bar{\mathbf{s}}_n & \mathbf{\mathfrak{X}} = (A \mid K \text{ a subideal of } A, \text{ for some } K \in \mathbf{\mathfrak{X}}) \\ & \mathbf{D}_0 & \mathbf{\mathfrak{X}} = (A \mid A = \bigoplus_{j=1}^r I_j \text{ with each } I_j \in \mathbf{\mathfrak{X}}) \\ & \mathbf{Q} & \mathbf{\mathfrak{X}} = (A \mid \exists H \in \mathbf{\mathfrak{X}} \text{ and } \exists \varphi : H \longrightarrow A \text{ an epimorphism}) \\ & \mathbf{R}_0 & \mathbf{\mathfrak{X}} = (A \mid \exists I_1, \dots, I_r \trianglelefteq A \text{ with } A/I_j \in \mathbf{\mathfrak{X}} \text{ and } \bigcap_{j=1}^r I_j = 0) \\ & \mathbf{R} & \mathbf{\mathfrak{X}} = (A \mid \exists (I_j)_{j \in J}, I_j \trianglelefteq A \text{ with } A/I_j \in \mathbf{\mathfrak{X}} \text{ for all } j \in J \text{ and } \bigcap_{j \in J} I_j = 0) \\ & \mathbf{N}_0 & \mathbf{\mathfrak{X}} = (A \mid \exists K_1, \dots, K_r \text{ subideals of } A \text{ with } K_i \in \mathbf{\mathfrak{X}} \text{ and } A = \sum_{i=1}^r K_i) \\ & \mathbf{N} & \mathbf{\mathfrak{X}} = (A \mid \exists (I_j)_{j \in J}, I_j \text{ a subideal of } A \text{ with } I_j \in \mathbf{\mathfrak{X}} \text{ for all } j \in J \text{ and } A = \sum_{j \in J} I_j) \\ & \mathbf{E}_\Phi & \mathbf{\mathfrak{X}} = (A \mid \exists N \le A \text{ with } N \le \Phi(A) \text{ and } A/N \in \mathbf{\mathfrak{X}}) \\ & \mathbf{P} & \mathbf{\mathfrak{X}} = (A \mid \mathbf{Q}(A) \cap \mathbf{\mathfrak{P}} \subseteq \mathbf{\mathfrak{X}}). \end{split}$$

Note that, if \mathfrak{X} is a class of algebras, then $\mathfrak{s}\mathfrak{X}$ is the class of all subalgebras of \mathfrak{X} -algebras, and $\mathfrak{Q}\mathfrak{X}$ is the class of all quotients of \mathfrak{X} -algebras.

A class \mathfrak{X} is called c-closed if $\mathfrak{X} = c\mathfrak{X}$. If c is a closure operation, then is $c\mathfrak{X}$ a c-closed class, for all classes \mathfrak{X} . It is clear from (C1) and (C2) that the classes \emptyset and \mathfrak{E} are c-closed for any operation c.

In this terminology $s\mathfrak{X} = \mathfrak{X}$ means that \mathfrak{X} is closed under subalgebras, $Q\mathfrak{X} = \mathfrak{X}$ indicate that \mathfrak{X} is quotient-closed, whereas $D_0\mathfrak{X} = \mathfrak{X}$ implies that \mathfrak{X} is closed under forming direct sums.

Definition 2.2. If \mathfrak{X} and \mathfrak{Y} are two classes of algebras, we define their product $\mathfrak{X}\mathfrak{Y}$ as follows:

(2.1)
$$\mathfrak{X}\mathfrak{Y} = (A \mid \exists N \leq A, N \in \mathfrak{X} with A/N \in \mathfrak{Y})$$

Algebras in the class $\mathfrak{X}\mathfrak{Y}$ are called \mathfrak{X} -by- \mathfrak{Y} algebras. If $\mathfrak{X} = \emptyset$ or $\mathfrak{Y} = \emptyset$, then we define $\mathfrak{X}\mathfrak{Y} = \emptyset$.

For powers of a class, we set $\mathfrak{X}^0 = (1)$, and for $n \in \mathbb{N}$ make the inductive definition $\mathfrak{X}^n = (\mathfrak{X}^{n-1})\mathfrak{X}$. An algebra in \mathfrak{X}^2 is called **meta-X**.

Remark 2.3. This product is neither associative nor commutative in general.

We will find conditions on \mathfrak{X} and \mathfrak{Y} such that $\mathfrak{X}(\mathfrak{YJ}) = (\mathfrak{XY})\mathfrak{Z}$ does hold.

Lemma 2.4. With the exception of P, all the class maps in definition 2.1 are closure operations.

Proof. It is very easy to verify that all the class maps are expanding and monotonic. We show now that all this class maps are idempotent. Notice that, if c is an expanding class map, then $c\mathfrak{X} \subseteq c(c\mathfrak{X})$, for all class \mathfrak{X} . Therefore it is sufficient to demonstrate that $c(c\mathfrak{X}) \subseteq c\mathfrak{X}$, with c the above defined class maps.

- (1) Operator s. Let $A \in s(s\mathfrak{X})$. Then there exists $H \in s\mathfrak{X}$ such that $A \leq H$. Similarly, there is an algebra $V \in \mathfrak{X}$ with $H \leq V$. Then we have $A \leq H \leq V$, with $V \in \mathfrak{X}$; that is $A \in s\mathfrak{X}$.
- (2) Operator s_n . Let $A \in s_n(s_n\mathfrak{X})$. Then there exist $H \in s_n\mathfrak{X}$ and $K \in \mathfrak{X}$ such that A is a subideal of H and H is a subideal of K. This proves that A is a subideal of K with $K \in \mathfrak{X}$; that is, $A \in s_n\mathfrak{X}$.
- (3) Operator \bar{s}_n . Let $A \in \bar{s}_n(\bar{s}_n \mathfrak{X})$. Then there exist $H \in \bar{s}_n \mathfrak{X}$ and $V \in \mathfrak{X}$ such that V is a subideal of H and H is a subideal of A. This proves that $A \in \bar{s}_n \mathfrak{X}$.
- (4) Operator D_0 . Let $A \in \mathsf{D}_0(\mathsf{D}_0\mathfrak{X})$. Then $A = \bigoplus_{i=1}^r K_i$ with $K_i \in \mathsf{D}_0\mathfrak{X}$, $i = 1, \ldots, r$. Therefore each algebra K_i has the form $K_i = \bigoplus_{j=1}^{r_i} N_{ij}$ with $N_{ij} \in \mathfrak{X}$. Since $\bigoplus_{i=1}^r \bigoplus_{j=1}^{r_i} N_{ij} = \bigoplus_{i=1}^r N_i = A$, it follows that $A \in \mathsf{D}_0\mathfrak{X}$.
- (5) Operator Q. Let $A \in Q(Q\mathfrak{X})$. Then there exist algebras $H \in Q\mathfrak{X}$ and $V \in \mathfrak{X}$ and epimorphisms $\varphi : H \longrightarrow A$ and $\psi : V \longrightarrow H$. Then we have an epimorphism $\varphi \psi : V \longrightarrow A$, with $V \in \mathfrak{X}$; that is $A \in Q\mathfrak{X}$.
- (6) Operator R₀. Let $A \in \mathsf{R}_0(\mathsf{R}_0\mathfrak{X})$. Then A has ideals N_1, \ldots, N_r such that $A/N_i \in \mathsf{R}_0\mathfrak{X}$ and $\bigcap_{i=1}^r N_i = 0$. Therefore each algebra A/N_i has ideals K_{ij}/N_i , $j = 1, \ldots, r_i$, such that $A/K_{ij} \cong (A/N_i)/(K_{ij}/N_i)$ and $\bigcap_{j=1}^{r_i} K_{ij} = N_i$. Since $\bigcap_{i=1}^r \bigcap_{j=1}^{r_i} K_{ij} = \bigcap_{i=1}^r N_i = 0$, it follows that $A \in \mathsf{R}_0\mathfrak{X}$.
- (7) Operator R. Is similar to (6).
- (8) Operator N₀. Let $A \in N_0(N_0\mathfrak{X})$. Then A has subideals K_1, \ldots, K_r such that $K_i \in N_0\mathfrak{X}$ and $A = \sum_{i=1}^r K_i$. Therefore each subalgebra K_i has subideals $N_{ij}, j = 1, \ldots, r_i$ with $N_{ij} \in \mathfrak{X}$ and $K_i = \sum_{j=1}^{r_i} N_{ij}$. Since $\sum_{i=1}^r \sum_{j=1}^{r_i} N_{ij} = \sum_{i=1}^r N_i = A$, it follows that $A \in N_0\mathfrak{X}$.
- (9) $\overline{\text{Operator N}}$. Is similar to (8).
- (10) Operator \mathbf{E}_{Φ} . Let $A \in \mathbf{E}_{\Phi}(\mathbf{E}_{\Phi}\mathfrak{X})$. Then A has an ideal $N \leq \Phi(A)$ such that $A/N \in \mathbf{E}_{\Phi}\mathfrak{X}$. Consequently A/N possesses an ideal $K/N \leq \Phi(A/N)$ such that $A/K \cong (A/N)/(K/N) \in \mathfrak{X}$. Since $\Phi(A/N) = \Phi(A)/N$, we have $K \leq \Phi(A)$. Therefore $A \in \mathbf{E}_{\Phi}\mathfrak{X}$.

Remark 2.5. Let c be a closure operation. If $\{\mathfrak{X}_j\}_{j\in J}$ is a family of c-closed algebras classes, it follows immediately that the class $\bigcap_{i\in J}\mathfrak{X}_j$ is also c-closed.

A product of operations is defined as follows:

Definition 2.6. Let c_1 and c_2 be operations. The product c_1c_2 is defined by composition thus $(c_1c_2)\mathfrak{X} = c_1(c_2\mathfrak{X})$, for all classes \mathfrak{X} .

Remark 2.7. If c_1 and c_2 are closure operations, then c_1c_2 need not be, in general, a closure operation. For example, if we define the class maps E, and L as follows:

$$\mathsf{E}\mathfrak{X} = (A \mid \exists \ 0 = I_0 \trianglelefteq I_1 \trianglelefteq \cdots \trianglelefteq I_n \trianglelefteq A \text{ with } I_{j+1}/I_j \in \mathfrak{X} \text{ for all } j = 0, \dots, n-1)$$

 $\mathsf{L}\mathfrak{X} = (A \mid \forall S \subset A, \ S \ finite \ \exists H \leq A, \ with \ H \in \mathfrak{X} \ and \ S \subseteq H)$

It is easy to show that they are closure operations, and in context of infinitedimensional Lie algebras, $EL\mathfrak{A} = E\mathfrak{A}$, whereas $ELEL\mathfrak{A} = ELE\mathfrak{A}$ contains non-solvable algebras, for instance the direct sum of a set of solvable algebras of unbounded derived length.

Remark 2.8. Let c be a closure operation. Conditions (C2) and (C3) imply that $c\mathfrak{X}$ is the uniquely determined, smallest c-closed class that contains the class \mathfrak{X} .

Remark 2.9. Let c be a closure operation. If $\{\mathfrak{X}_j\}_{j\in J}$ is a family of c-closed algebras classes it follows immediately that the class $\bigcap_{i\in J}\mathfrak{X}_j$ is also c-closed.

Some elementary closure properties of class product are discussed now.

Lemma 2.10. Given algebras A_1, A_2, H and epimorphisms $\mu_i : A_i \longrightarrow H$, let be $N_i := \ker(\mu_i)$. Then

(2.2)
$$A_1 \land A_2 := \{ (a_1, a_2) \mid a_i \in A_i, \ \mu_1(a_1) = \mu_2(a_2) \}$$

is a subalgebra of $A_1 \oplus A_2$. Furthermore, there exists epimorphisms $\alpha_i : A_1 \land A_2 \longrightarrow A_i$ (i = 1, 2) with

$$\ker(\alpha_1) = \{(0, a_2) \mid a_2 \in N_2\} \cong N_2$$

and

$$\ker(\alpha_2) = \{ (a_1, 0) \mid a_1 \in N_1 \} \cong N_1.$$

In addition there is an epimorphism $\mu: A_1 \land A_2 \longrightarrow H$ with

$$\ker(\mu) = \ker(\alpha_1) \oplus \ker(\alpha_2)$$

Proof. It is clear that $A_1 \land A_2$ is a subalgebra of $A_1 \oplus A_2$. Let $\pi_i : A_1 \oplus A_2 \longrightarrow A_i$ the homomorphism defined by

$$\pi_i(a_1, a_2) := a_i.$$

Then the restriction of π_i on $A_1 \downarrow A_2$ is an epimorphism α_i with

$$\ker(\alpha_1) = \{ (a_1, a_2) \mid a_i \in A_i, \ \mu_1(a_1) = \mu_2(a_2), \ a_1 = 0 \}$$
$$= \{ (0, a_2) \mid a_i \in A_i, \ 0 = \mu_1(a_1) = \mu_2(a_2) \}$$
$$= \{ (0, a_2) \mid a_2 \in \ker(\mu_2) \}$$
$$= 0 \oplus N_2.$$

On the other hand $\mu(a_1, a_2) := \mu_1(a_1) = \mu_2(a_2)$ defines an epimorphism μ from $A_1 \downarrow A_2$ to H. Evidently $\ker(\mu) = \ker(\alpha_1) \oplus \ker(\alpha_2)$.

Lemma 2.11. If \mathfrak{X} and \mathfrak{Y} are classes of algebras, then $\mathfrak{Y}(\mathfrak{QX}) \subseteq \mathfrak{Q}(\mathfrak{YX})$.

Proof. Let $A \in \mathfrak{Y}(q\mathfrak{X})$. Then A has an ideal $I \in \mathfrak{Y}$ with $A/I \in q\mathfrak{X}$. Thus there exists an algebra $H \in \mathfrak{X}$ and an epimorphism $\mu_1 : H \longrightarrow A/I$. Let $\mu_2 : A \longrightarrow A/I$ be the canonical epimorphism. Consider now the subalgebra

(2.3)
$$S := H \land A := \{(h, l) \mid h \in H, \ l \in A, \ \mu_1(h) = \mu_2(l) = l + I\}$$

of the direct sum $H \oplus A$. Using Lemma 2.10 it follows that there exist epimorphisms $\alpha_1 : S \longrightarrow H$ and $\alpha_2 : S \longrightarrow A$ with

$$\ker(\alpha_1) = \{(0,i) \mid i \in I\} = 0 \oplus I \cong I \in \mathfrak{Y}, \text{ and}$$
$$\ker(\alpha_2) = \{(h,0) \mid h \in \ker(\mu_1)\} = \ker(\mu_1) \oplus 0 \cong \ker(\mu_1).$$

Then $S/\ker(\alpha_1) \cong H \in \mathfrak{X}$ and we have that $S \in \mathfrak{Y}\mathfrak{X}$.

On the other hand $S/\ker(\alpha_2) \cong A$, and therefore $A \in Q(\mathfrak{YX})$.

Lemma 2.12. Let c be any of the closure operations s, q or s_n . If \mathfrak{X} and \mathfrak{Y} are c-closed classes of algebras, then \mathfrak{XY} is also a c-closed class.

Proof. Operator s. Suppose $\mathfrak{X} = s\mathfrak{X}$ and $\mathfrak{Y} = s\mathfrak{Y}$, and let $A \in \mathfrak{X}\mathfrak{Y}$. Then A has an ideal $I \in \mathfrak{X}$ with $A/I \in \mathfrak{Y}$. Let $V \leq A$. We must show that $V \in \mathfrak{X}\mathfrak{Y}$. To this end, consider the quotient algebra (V + I)/I and notice that $(V + I)/I \leq A/I \in \mathfrak{Y} = s\mathfrak{Y}$. Hence $(V + I)/I \in \mathfrak{Y}$. Using the second isomorphism theorem we have that $(V + I)/I \cong V/(V \cap I) \in \mathfrak{Y}$. Furthermore, $V \cap I \leq I \in \mathfrak{X} = s\mathfrak{X}$ and from here it follows that $V \in \mathfrak{X}\mathfrak{Y}$.

Operator \mathfrak{Q} . Suppose $\mathfrak{X} = \mathfrak{Q}\mathfrak{X}$ and $\mathfrak{Y} = \mathfrak{Q}\mathfrak{Y}$, and let $A \in \mathfrak{X}\mathfrak{Y}$. Then A has an ideal $I \in \mathfrak{X}$ with $A/I \in \mathfrak{Y}$. Let K be an ideal of A. We show that $A/K \in \mathfrak{X}\mathfrak{Y}$. To do that, consider the ideal (I + K)/K of A/K. Certainly we have $(I + K)/K \cong I/(I \cap K) \in \mathfrak{Q}\mathfrak{X} = \mathfrak{X}$. Additionally,

$$\frac{A/K}{(I+K)/K}\cong \frac{A}{I+K}\cong \frac{A/I}{(I+K)/I}\in \mathsf{Q}(A/I)\subseteq \mathsf{Q}\mathfrak{Y}=\mathfrak{Y}.$$

Consequently, $A/K \in \mathfrak{XY}$.

Operator s_n . Suppose that $\mathfrak{X} = s_n \mathfrak{X}$ and $\mathfrak{Y} = s_n \mathfrak{Y}$. Now we prove that $s_n(\mathfrak{XY}) \subseteq \mathfrak{XY}$. To this end, we show that if K is a subideal of $A \in \mathfrak{XY}$, then $K \in \mathfrak{XY}$. Let I be an ideal of A with $I \in \mathfrak{X}$ and $A/I \in \mathfrak{Y}$. Since $K \cap I$ is a subideal of $I \in \mathfrak{X}$ we have $K \cap I \in s_n \mathfrak{X} = \mathfrak{X}$; also $K/(K \cap I) \cong (K+I)/I \trianglelefteq A/I \in \mathfrak{Y}$. Therefore $K/(K \cap I) \in s_n \mathfrak{Y} = \mathfrak{Y}$. Hence $K \in \mathfrak{XY}$.

Lemma 2.13. Let \mathfrak{X} , \mathfrak{Y} and \mathfrak{Z} are classes of algebras. Then each of the following two conditions is sufficient to guarantee the associativity of the product of classes.

(1) $\mathfrak{X} = \mathsf{N}_0 \mathfrak{X}$ and $\mathfrak{Y} = \mathsf{Q} \mathfrak{Y}$. (2) $\mathfrak{X} = \bar{\mathsf{s}}_n \mathfrak{X}$ and $\mathfrak{Y} = \mathsf{Q} \mathfrak{Y}$

Proof. It is obvious that $\mathfrak{X}(\mathfrak{YJ}) \subseteq (\mathfrak{XY})\mathfrak{Z}$. Then it is enough to prove that $(\mathfrak{XY})\mathfrak{Z} \subseteq \mathfrak{X}(\mathfrak{YJ})$.

(1) Let A be an algebra belonging to the class $(\mathfrak{XY})\mathfrak{Z}$. Then A has a chain of subalgebras

$$(2.4) 0 \le K \le I \le A,$$

such that $K \in \mathfrak{X}$, $I/K \in \mathfrak{Y}$ and $A/I \in \mathfrak{Z}$. Let K^A be the ideal closure of K in A. Then, by using (1.1), we have $K^A = \sum_{k \in K} (k)$ and it follows that K^A is an ideal of A, which belongs to $\mathbb{N}_0 \mathfrak{X} = \mathfrak{X}$. Furthermore, the quotient algebra A/K^A has an ideal $I/K^A \cong (I/K)/(K^A/K) \in \mathbb{Q}(I/K) \subseteq \mathbb{Q}\mathfrak{Y} = \mathfrak{Y}$, and $(A/A^K)/(I/K^A) \cong A/I \in \mathfrak{Z}$. Thus $A/K^A \in \mathfrak{Y}\mathfrak{Z}$, and we have proved that $A \in \mathfrak{X}(\mathfrak{Y}\mathfrak{Z})$, as desired.

(2) Let us suppose again that A has a chain (2.4) of subalgebras with the stated properties, and let K^A be the ideal closure of K in A. Since $K \in \mathfrak{X}$ and $K \trianglelefteq \trianglelefteq K^A$ we have that $K^A \in \bar{\mathbf{s}}_n \mathfrak{X} = \mathfrak{X}$. The rest is similar to the argument in (1).

3. A partial order on closure operations

Definition 3.1. Let c_1 and c_2 be class maps, in particular, closure operations. We say that c_1 is contained in c_2 , and write $c_1 \leq c_2$ if $c_1 \mathfrak{X} \subseteq c_2 \mathfrak{X}$, for every algebras class \mathfrak{X} .

Remarks 3.2. (1) It is straightforward to verify that \leq is a relation of partial order on the set of class maps and hence on the set of closure operations.

- (2) If c_1 and c_2 are closure operations, $c_1 \leq c_2$ if and only if c_2 -closure invariably implies c_1 -closure.
- (3) From the definition, it is obvious that $D_0 \leq R_0$, $s_n \leq s$ and $D_0 \leq N_0$.

Lemma 3.3. Let c_1 and c_2 be closure operations. Then $c_1 \leq c_2$ if and only if every c_2 -closed class is c_1 -closed.

Proof. Let us suppose that $c_1 \leq c_2$, and $\mathfrak{X} = c_2\mathfrak{X}$. Then $c_1\mathfrak{X} \subseteq c_2\mathfrak{X} = \mathfrak{X}$. Owing to the fact that c_1 is expanding, we have $\mathfrak{X} \subseteq c_1\mathfrak{X}$, and therefore $\mathfrak{X} = c_1\mathfrak{X}$.

Reciprocally, assume that every c_2 -closed class is also c_1 -closed, and let \mathfrak{X} be any class of algebras. Since c_2 is a closure operation, we have $\mathfrak{X} \subseteq c_2 \mathfrak{X}$ and $c_2 \mathfrak{X}$ is c_2 -closed. Seeing that c_1 is monotonic, it follows by assumption that $c_1 \mathfrak{X} \subseteq c_1(c_2 \mathfrak{X}) = c_2 \mathfrak{X}$. Thus $c_1 \leq c_2$.

Definition 3.4 (The join of a set of closure operations). Let $\{c_j \mid j \in J\}$ be a set of closure operations. We define the join $c := \langle c_j \mid j \in J \rangle$ by

(3.1)
$$\mathsf{c}\mathfrak{X} := \{ \mathfrak{Y} \mid \mathfrak{X} \subseteq \mathfrak{Y} = \mathsf{c}_j \mathfrak{Y} \text{ for all } j \in J \}.$$

for any algebras class \mathfrak{X} .

Lemma 3.5. Let $\{c_j \mid j \in J\}$ be a set of closure operations and let $c = \langle c_j \mid j \in J \rangle$ be their join. Then

- (1) the class map c is a closure operation;
- (2) if \mathfrak{X} is a class of algebras, then $\mathfrak{C}\mathfrak{X}$ is the uniquely determined, smallest class containing \mathfrak{X} which is simultaneously c_j -closed for all $j \in J$;
- (3) in the partial order on the set of closure operations defined above the join c is the least upper bound of the set {c_j | j ∈ J}.
- *Proof.* (1) From the definition of c follows immediately that it is expanding and monotonic. To prove the last property note that, for any class \mathfrak{X} , we have

$$\{\mathfrak{Y} \mid \mathfrak{X} \subseteq \mathfrak{Y} = \mathsf{c}_j \mathfrak{Y} \text{ for all } j \in J\} = \{\mathfrak{Y} \mid \mathsf{c}\mathfrak{X} \subseteq \mathfrak{Y} = \mathsf{c}_j \mathfrak{Y} \text{ for all } j \in J\},\$$

and therefore $c(c\mathfrak{X}) = c\mathfrak{X}$. Then c is also idempotent.

- (2) It follows easily from the definition that if c is a closure operation and $\{\mathfrak{X}_j \mid j \in J\}$ is a set of c-closed classes then $\bigcap_{j \in J} \mathfrak{X}_j$ is c-closed. Thus $c\mathfrak{X}$ is a c_j -closed class for all $j \in J$. On the other hand, any c_j -closed class \mathfrak{Y} for all $j \in J$, that contain \mathfrak{X} certainly contains $c\mathfrak{X}$ by definition, and therefore Statement (2) is true.
- (3) Let \mathfrak{X} be a class of algebras. Since $\mathfrak{X} \subseteq c\mathfrak{X}$ we have for all $j \in J$

(3.2)
$$c_j \mathfrak{X} \subseteq c_j(c\mathfrak{X}) = c\mathfrak{X};$$

that is, $c_j \leq c$ for all $j \in J$, and therefore c is an upper bound of the set $\{c_j \mid j \in J\}$. Let now A be any upper bound for the set $\{c_j \mid j \in J\}$ and let \mathfrak{X} be any A-closed class. By Lemma 3.3 we have that \mathfrak{X} is a c_j -closed class for all $j \in J$. Then by (2) follows that

$$(3.3) C\mathfrak{X} \subseteq \mathfrak{X} = \mathsf{A}\mathfrak{X}.$$

This shows Statement (3).

Lemma 3.6. Let c_1, \ldots, c_n be closure operations, and let \mathfrak{X} be a class of algebras. Then $\mathfrak{X} = c_1 \cdots c_n \mathfrak{X}$ if and only if $\mathfrak{X} = \langle c_1 \cdots c_n \rangle \mathfrak{X}$.

Proof. Let suppose that $\mathfrak{X} = c_1 \cdots c_n \mathfrak{X}$ and let $j \in \{1, \ldots, n\}$. By Lemma 3.5 (2) it is sufficient to prove that $\mathfrak{X} = c_j \mathfrak{X}$. Since c_{j+1}, \ldots, c_n are expanding and monotonic, we have $\mathfrak{X} \subseteq c_{j+1} \cdots c_n \mathfrak{X}$, and hence $c_j \mathfrak{X} \subseteq c_j c_{j+1} \cdots c_n \mathfrak{X}$. On the other hand, c_1, \ldots, c_{j-1} are also expanding and monotonic, and then

(3.4)
$$\mathsf{c}_{j}\mathfrak{X} \subseteq \mathsf{c}_{1}\cdots\mathsf{c}_{j-1}(\mathsf{c}_{j}\mathfrak{X}) \subseteq \mathsf{c}_{1}\cdots\mathsf{c}_{j-1}(\mathsf{c}_{j}\mathsf{c}_{j+1}\cdots\mathsf{c}_{n}\mathfrak{X}).$$

Hence $c_j \mathfrak{X} \subseteq c_1 \cdots c_n \mathfrak{X} = \mathfrak{X}$, and consequently $\mathfrak{X} = c_j \mathfrak{X}$ as required.

Conversely, assume that $\mathfrak{X} = \langle c_1 \cdots c_n \rangle \mathfrak{X}$. Then it holds that $c_j \mathfrak{X} = \mathfrak{X}$ for all $j \in \{1, \ldots, n\}$, and $\mathfrak{X} = c_1 \cdots c_n \mathfrak{X}$.

Lemma 3.7. Let c_1 and c_2 be closure operations. Then any two of the following statements are equivalent:

- (1) The class map c_1c_2 is a closure operation;
- $(2) \mathsf{c}_2 \mathsf{c}_1 \leq \mathsf{c}_1 \mathsf{c}_2$
- (3) $c_1c_2 = \langle c_1, c_2 \rangle$.

Proof. (1) \Rightarrow (2). Let \mathfrak{X} be any class of algebras, and suppose that c_1c_2 is a closure operation. From the expanding and monotonic properties of c_1 and c_2 we have

$$(3.5) c_2 c_1 \mathfrak{X} \subseteq c_2 c_1 (c_2 \mathfrak{X}) \subseteq c_1 (c_2 c_1 (c_2 \mathfrak{X})) = (c_1 c_2)^2 \mathfrak{X} = (c_1 c_2) \mathfrak{X}.$$

Then we have $c_2c_1 \leq c_1c_2$.

 $(2) \Rightarrow (3)$. Let \mathfrak{X} be any class of algebras, and let $\mathfrak{X} \subseteq \mathfrak{Y} = c_1 \mathfrak{Y} = c_2 \mathfrak{Y}$. Then $(c_1c_2)\mathfrak{X} \subseteq (c_1c_2)\mathfrak{Y} = c_1(c_2\mathfrak{Y}) = c_1\mathfrak{Y} = \mathfrak{Y}$. This show that $(c_1c_2)\mathfrak{X} \subseteq \langle c_1, c_2 \rangle \mathfrak{X}$. On the other hand, it is clear that $(c_1c_2)\mathfrak{X}$ is a c_1 -closed class containing \mathfrak{X} , and since

$$\mathsf{c}_2(\mathsf{c}_1\mathsf{c}_2)\mathfrak{X} = (\mathsf{c}_2\mathsf{c}_1)\mathsf{c}_2\mathfrak{X} = (\mathsf{c}_1\mathsf{c}_2)\mathsf{c}_2\mathfrak{X} = (\mathsf{c}_1\mathsf{c}_2)\mathfrak{X},$$

the class $(c_1c_2)\mathfrak{X}$ is a c_2 -closed class. Therefore $\langle c_1, c_2 \rangle \mathfrak{X} \subseteq (c_1c_2)\mathfrak{X}$, and (3) holds.

 $(3) \Rightarrow (1)$. This follows from the fact that $\langle c_1, c_2 \rangle$ is a closure operation. \Box

Lemma 3.8. The class maps products sD_0 , $E_{\Phi}D_0$ and $E_{\Phi}Q$ are closure operations.

Proof. Let \mathfrak{X} be any class of algebras.

- 10 I. S. GUTIERREZ, ANSELMO TORRESBLANCA-BADILLO, AND DAVID A. TOWERS
 - (1) Let $A \in D_0 \mathfrak{sX}$. Then there exist algebras $A_1, \ldots, A_n \in \mathfrak{X}$ with subalgebras H_1, \ldots, H_n , respectively such that

$$A \cong \bigoplus_{j=1}^n H_j.$$

Since $\bigoplus_{j=1}^{n} H_j$ can be identified with the obvious subgroup of $\bigoplus_{j=1}^{n} A_j \in D_0 \mathfrak{X}$ we conclude that $A \in sD_0 \mathfrak{X}$. This proves that $D_0 s \leq sD_0$ and then by using Lemma 3.7 the map sD_0 is a closure operation.

(2) Let $A \in \mathsf{D}_0\mathsf{E}_\Phi\mathfrak{X}$. Then there exist algebras $A_1, \ldots, A_n \in \mathsf{E}_\Phi\mathfrak{X}$ satisfying

$$A \cong \bigoplus_{j=1}^n A_j.$$

For all $j \in \{1, ..., n\}$ we have that there exist $I_j \leq A_j$ with $I_j \leq \Phi(A_j)$ and $A_j/I_j \in \mathfrak{X}$. Since $\Phi(A) = \bigoplus_{j=1}^n \Phi(A_j)$, by [16, Theorem 4.8], and $A/\bigoplus_{j=1}^n I_j \cong \bigoplus_{j=1}^n (A_j/I_j) \in \mathsf{D}_0\mathfrak{X}$, we have that $A/\bigoplus_{j=1}^n I_j \in \mathsf{D}_0\mathfrak{X}$. On the other hand, $I := \bigoplus_{j=1}^n I_j \leq \bigoplus_{j=1}^n \Phi(A_j) = \Phi(A)$, and therefore $A/I \in$ $\mathsf{D}_0\mathfrak{X}$ and $A \in \mathsf{E}_\Phi\mathsf{D}_0\mathfrak{X}$. Thus $\mathsf{D}_0\mathsf{E}_\Phi \leq \mathsf{E}_\Phi\mathsf{D}_0$ and we use Lemma 3.7 to conclude.

(3) Let $A \in \mathsf{QE}_{\Phi}\mathfrak{X}$. Then $A \cong H/N$, where $N \trianglelefteq H$ and H has an ideal I such that $I \le \Phi(H)$ and $H/I \in \mathfrak{X}$. Thus

(3.6)
$$\frac{H/N}{(I+N)/N} \cong \frac{H}{I+N} \cong \frac{H/I}{(I+N)/N} \in Q\mathfrak{X}.$$

Furthermore,

(3.7)
$$(I+N)/N \le (\Phi(H)+N)/N \le \Phi(H/N).$$

Following (3.6) and (3.7) we have $A \in E_{\Phi}Q\mathfrak{X}$. Then $QE_{\Phi} \leq E_{\Phi}Q$ and the assertion is true.

Definition 3.9. Let A_1, \ldots, A_r be algebras, and let $D := \bigoplus_{i=1}^r A_i$ be their direct sum.

(1) The map $\pi_i: D \longrightarrow A_i$ defined by

$$\pi(a_1,\ldots,a_r):=a_i$$

- is called the projection of D onto the *i*-th component.
- (2) A subalgebra U of D is called subdirect if $\pi_i(U) = A_i$, for each $i = 1, \ldots, r$.

Definition 3.10. A subdirect product is a subalgebra A of a direct product $\prod_{i \in I} A_i$ such that every induced projection (the composite $p_j s : A \longrightarrow A_j$ of a projection $p_j : \prod_{i \in I} A_i \longrightarrow A_j$ with the subalgebra inclusion $s : A \longrightarrow \prod_{i \in I} A_i$) is surjective.

Lemma 3.11. Let I_1, \ldots, I_n be ideals of an algebra A. Then the function $f : A \longrightarrow \bigoplus_{i=1}^n (A/I_j)$ defined by

$$f(x) = (x + I_1, \dots, x + I_n)$$

is a algebra homomorphism; its kernel is $\bigcap_{j=1}^{n} I_j$ and its image f(A) is subdirect. In particular, $A / \bigcap_{j=1}^{n} I_j$ is isomorphic with a subdirect subalgebra of $\bigoplus_{j=1}^{n} (A/I_j)$. *Proof.* It is clear that f is a homomorphism. Moreover, $f(x) = (I_1, \ldots, I_n)$, if and only if $x \in I_j$, for all $j \in \{1, \ldots, n\}$, and therefore $\ker(f) = \bigcap_{j=1}^n I_j$. Finally, $\pi_j(f(A)) = \{x + I_j \mid x \in A\} = A/I_j$, whence f(A) is subdirect. The final assertion follows from the isomorphism theorem. \Box

Lemma 3.12. Let \mathfrak{X} be a class of algebras.

- (1) An algebra A belongs to $R_0 \mathfrak{X}$ if and only if A is isomorphic with a subdirect subalgebra of a direct sum of a finite set of \mathfrak{X} -algebras.
- (2) The class maps product QR_0 is a closure operation.
- *Proof.* (1) Let $A \in \mathbb{R}_0 \mathfrak{X}$. Then there exist ideals I_1, \ldots, I_n of A such that $A/I_j \in \mathfrak{X}$, for all $j \in \{1, \ldots, n\}$ and $\bigcap_{j=1}^n I_j = 0$. By using Lemma 3.11 we have that A is isomorphic with a subdirect subalgebra of $\bigoplus_{j=1}^n (A/I_j)$.

Reciprocally, if $f : A \longrightarrow \bigoplus_{j=1}^{n} I_j$ is a monomorphism with f(A) subdirect and each $I_j \in \mathfrak{X}$. We consider now the following functions:



Since f(A) is subdirect we have for all $j \in \{1, ..., n\}$ that

(3.8)
$$\operatorname{Im}(\pi_j f) = \pi_j(f(A)) = I_j.$$

We define $N_j := \ker(\pi_j f)$. Then $A/N_j \cong I_j \in \mathfrak{X}$ and $\bigcap_{j=1}^n I_j = \ker(f) = 0$. Thus $A \in \mathsf{R}_0 \mathfrak{X}$.

(2) Let $A \in \mathsf{R}_0 Q \mathfrak{X}$. Then, using (1), there exist algebras $H_1, \ldots, H_n \in \mathfrak{X}$ and ideals $I_j \leq H_j$, for $j \in \{1, \ldots, n\}$, such that $A \leq \bigoplus_{j=1}^n (H_j/I_j)$ and $\pi_j(A) = H_j/I_j$, for all $j \in \{1, \ldots, n\}$. Let α denote the natural homomorphism from $\bigoplus_{j=1}^n H_j$ onto $\bigoplus_{j=1}^n (H_j/I_j)$ and define H as the inverse image of A under α . To prove that H is subdirect in $\bigoplus_{j=1}^n H_j$ we consider the following maps:

Note that

$$H_j/I_j \cong \tilde{\pi}_j(A) = \tilde{\pi}_j(\alpha(H)) = \gamma_j(\pi_j(H)) = \pi_j(H)/I_j.$$

Then, $\pi_j(H) = H_j$ for all $j \in \{1, \ldots, n\}$ and this shows that $H \in \mathsf{R}_0\mathfrak{X}$. Moreover, $A \cong H/(H \cap \ker(\alpha)) \in \mathsf{QR}_0\mathfrak{X}$. Thus $\mathsf{R}_0\mathsf{Q} \leq \mathsf{QR}_0$, and so the map QR_0 is a closure operation. 12 I. S. GUTIERREZ, ANSELMO TORRESBLANCA-BADILLO, AND DAVID A. TOWERS

Remark 3.13. If S is a set of algebras, we use (S) to denote the smallest class of algebras containing S, and when $S = \{A\}$, a singleton, we write (A) rather than $(\{A\})$.

We end this section by describing unary closure operations and a useful exponential notation for this maps.

Definition 3.14. A closure operation c is called unary if $c\mathfrak{X} = \bigcup \{c(A) \mid A \in \mathfrak{X}\}$ for all classes \mathfrak{X} . If c is a unary closure operation and \mathfrak{X} a class of algebras, we define

$$\mathfrak{X}^{\mathsf{C}} = (A \mid \mathsf{c}(A) \subseteq \mathfrak{X}).$$

- **Remarks 3.15.** (1) The maps Q, s and s_n are examples of unary operations whereas D_0 , R_0 and N_0 are not.
 - (2) It follows from definition of unary operation that $\mathfrak{X}^{\mathsf{C}}$ is the unique largest c -closed class contained in \mathfrak{X} . In particular holds $\mathsf{c}(\mathfrak{X}^{\mathsf{C}}) = \mathfrak{X}^{\mathsf{C}}$.

Lemma 3.16. Let c_1 and c_2 be closure operations and \mathfrak{X} a c_2 -closed class of algebras. If c_1 is unary and $c_1c_2 \leq c_2c_1$, then \mathfrak{X}^{C_1} is a c_2 -closed class.

Proof. We have $c_1c_2\mathfrak{X}^{\mathsf{C}_1} \subseteq c_2c_1\mathfrak{X}^{\mathsf{C}_1} = c_2\mathfrak{X}^{\mathsf{C}_1} \subseteq c_2\mathfrak{X} = \mathfrak{X}$, and therefore $c_2\mathfrak{X}^{\mathsf{C}_1}$ is a c_1 -closed subclass of \mathfrak{X} . Consequently $c_2\mathfrak{X}^{\mathsf{C}_1} \subseteq \mathfrak{X}^{\mathsf{C}_1}$, and it follows that $\mathfrak{X}^{\mathsf{C}_1}$ is c_2 -closed.

4. Classes defined by closure properties

Schunck classes, formations, varieties, and Fitting classes are example of classes of algebras which satisfy certain closure properties. This classes are the main subject of this section.

Definition 4.1. (1) A non-empty Q-closed class \mathfrak{H} is called a homomorph. That is, \mathfrak{H} contains, along with an algebra A, all epimorphic images of A.

- (2) A homomorph which is also R₀-closed is called a formation. By lemma 3.12 (2) a formation is precisely a QR₀-closed class, and classes which are simultaneously closed under s, Q, and D₀ are formations. Furthermore, (Q, R₀)X = QR₀X is the formation generated by the class X.
- (3) A homomorph which is also R-closed is called a variety.
- (4) A homomorph which is also P-closed is called a Schunck class.
- (5) A non-empty class \mathfrak{X} is called a Fitting class if and only if, $\langle s_n, N_0 \rangle \mathfrak{X} = \mathfrak{X}$.

Lemma 4.2. Let \mathfrak{X} be an R-closed class of algebras and L an algebra. Then the set

$$\mathscr{S}(\mathfrak{X},L) := \{I \mid I \leq L, \ L/I \in \mathfrak{X}, \}$$

partially ordered by inclusion, has a unique minimal element, denoted by $L^{\mathfrak{X}}$ and called the \mathfrak{X} -residual of L. If \mathfrak{X} is a formation and $\varphi : L \longrightarrow \varphi(L)$ is an epimorphism, then $\varphi(L^{\mathfrak{X}}) = \varphi(L)^{\mathfrak{X}}$.

Proof. Let $R := \bigcap \{I \mid I \in \mathscr{S}(\mathfrak{X}, L)\}$. It is clear that R is an ideal of L. We consider now the set

$$\mathcal{T} := \{ I/R \mid I \in \mathscr{S}(\mathfrak{X}, L) \}.$$

It is verified that $\bigcap \mathcal{T} = 0$ and $(L/R)/(I/R) \in \mathfrak{X}$. Therefore $L/R \in \mathfrak{R}\mathfrak{X} = \mathfrak{X}$ and we have $R \in \mathscr{S}(\mathfrak{X}, L)$. Then R is the desired smallest element of $\mathscr{S}(\mathfrak{X}, L)$, which is denoted by $L^{\mathfrak{X}}$.

Let $\varphi : L \longrightarrow \varphi(L)$ be an epimorphism. Let $R := L^{\mathfrak{X}}$, $T := \varphi(L)^{\mathfrak{X}}$ and $I := \varphi^{-1}(T)$, the inverse image of T in L. Then by using the isomorphism theorem we have

$$L/I \cong \varphi(L)/T \in \mathfrak{X},$$

and therefore $L^{\mathfrak{X}} \leq I$ and so $\varphi(L^{\mathfrak{X}}) \leq \varphi(I) = \varphi(\varphi^{-1}(T)) = T = \varphi(L)^{\mathfrak{X}}$. Conversely, $\varphi(L)/\varphi(L^{\mathfrak{X}}) \in \mathsf{q}(L/L^{\mathfrak{X}}) \subseteq \mathsf{q}\mathfrak{X} = \mathfrak{X}$, and then $T \leq \varphi(L^{\mathfrak{X}})$. Thus $\varphi(L)^{\mathfrak{X}} = \varphi(L^{\mathfrak{X}})$. \Box

Definition 4.3. Let \mathfrak{X} be a homomorph. We say that a subalgebra B of A is an \mathfrak{X} -projector if

(1)
$$B \in \mathfrak{X}$$
, and
(2) $B \leq C \leq A$, $C_0 \leq C$, $C/C_0 \in \mathfrak{X}$ implies that $B + C_0 = C$.

We denote the set of all \mathfrak{X} -projectors of A by $Proj_{\mathfrak{X}}(A)$ and write $B \in Proj_{\mathfrak{X}}(A)$ for B is an \mathfrak{X} -projector of A. In the following we assume that all algebras A are finite dimensional and solvable; that is, $A \in \mathfrak{E} \cap \mathfrak{S}$.

Lemma 4.4. If $B \in Proj_{\mathfrak{X}}(A)$, then B is maximal in the set $\{C \leq A \mid C \in \mathfrak{X}\}$; in particular, if $A \in \mathfrak{X}$, then B = A.

Proof. If $B \leq C \leq A$ and $C \in \mathfrak{X}$, put $C_0 = 0$ in Definition 4.3 (2).

Lemma 4.5. (1) If $B \in Proj_{\mathfrak{X}}(A)$, and $B \leq C \leq A$, then $B \in Proj_{\mathfrak{X}}(C)$. (2) If $B \in Proj_{\mathfrak{X}}(A)$ and $I \leq A$, then $B + I/I \in Proj_{\mathfrak{X}}(A/I)$.

Proof. These are clear.

Lemma 4.6. Suppose that \mathfrak{X} is a homomorph which contains a non-zero algebra. Then the one-dimensional solvable algebra is in \mathfrak{X} , and $B \in \operatorname{Proj}_{\mathfrak{X}}(A)$, $A \neq 0$ implies that $B \neq 0$

Proof. Suppose that $A \in Proj_{\mathfrak{X}}(A)$, $A \neq 0$. Since A is solvable there exists $A_1 \trianglelefteq A$ such that $\dim(A/A_1) = 1$. If $B \in Proj_{\mathfrak{X}}(A)$ and $A \neq 0$, then $B \neq 0$, by Lemma 4.4.

Lemma 4.7. Suppose that $B \in Proj_{\mathfrak{X}}(A)$ and $0 \neq B \leq U \leq A$. If $U \leq V \leq A$, then U = V.

Proof. Suppose that $U \neq V$. There is a subalgebra W of V such that dim W/U = 1, since A is solvable. Then $W/U \in \mathfrak{X}$, by Lemma 4.6, and B + U = W, putting W = C, $U = C_0$ in Definition 4.3 (2). But $B \leq U$, so U = W, a contradiction. \Box

Lemma 4.8. Suppose that $B \leq A$, $U/B \in Proj_{\mathfrak{X}}(A/B)$ and $C \in Proj_{\mathfrak{X}}(U)$. Then $C \in Proj_{\mathfrak{X}}(A)$.

Proof. We use induction on dim A. The result is trivial if dim A = 0. Suppose that $C \leq D \leq A$, $D_0 \leq D$ and $D/D_0 \in \mathfrak{X}$. Then $D + B \geq C + B = U$, by Lemma 4.4, and $U/B \in Proj_{\mathfrak{X}}(C + B/B)$, by Lemma 4.5 (2). But $C + B/B \cong C/C \cap B$ and so $U \cap C/B \cap C \in Proj_{\mathfrak{X}}(C/B \cap C)$.

If D < A, then, since $C \in Proj_{\mathfrak{X}}(U \cap D)$, $C \in Proj_{\mathfrak{X}}(D)$ by induction and $C + D_0 = D$. Thus, $C \in Proj_{\mathfrak{X}}(A)$.

So suppose that D = A. Since $A/(D_0 + B) \in \mathfrak{X}$, $U/B + (D_0 + B)/B = A/B$ and so $U + B + D_0 = A$. Since $U \ge B$, $U + D_0 = A$. Hence $U/D_0 \cap U \cong A/D_0 \in \mathfrak{X}$. It follows that $C + (D_0 \cap U) = U$ and $C + D_0 = U + D_0 = A$.

We will call an ideal B of A abelian if $B^2 = 0$. Note that, if A has an abelian ideal, then it will have a minimal abelian ideal. However, in general, a solvable algebra may not have an abelian ideal, as we will see later.

Lemma 4.9. Let \mathfrak{X} be a homomorph and let B be a minimal abelian ideal of A. Suppose that $A/B \in \mathfrak{X}$, $A \notin \mathfrak{X}$ and $C \in Proj_{\mathfrak{X}}(A)$. Then C complements B in A.

Proof. We have that A = B + C and $C \neq A$, from Definition 4.3 (2). Moreover, $B \cap C$ is an abelian ideal of A, since B is abelian. Hence $B \cap C = 0$.

Lemma 4.10. Let \mathfrak{X} be a formation and let B be a minimal abelian ideal of A. Suppose that $A/B \in \mathfrak{X}$, $A \notin \mathfrak{X}$ and that C complements B in A. Then $C \in \mathfrak{X}$.

Proof. We have that $C \cong A/B \in \mathfrak{X}$ and C is a maximal subalgebra of A, so it remains to prove that, if C_0 is an ideal of A and $A/C_0 \in \mathfrak{X}$, then $A = C + C_0$. Since \mathfrak{X} is a formation, $A/B \cap C_0 \in \mathfrak{X}$. But B is a minimal ideal of A, so $B \cap C_0 = 0$ or B. The former is ruled out by the fact that $A \notin \mathfrak{X}$, so $B \subseteq C_0$ and $C + C_0 \supseteq C + B = A$. \Box

Definition 4.11. If B is an abelian ideal of A we can consider B as an A/Bbimodule by defining

 $B \times A/B \rightarrow B, (b, a + B) \mapsto ba \text{ and } A/B \times B \rightarrow B, (a + B, b) \mapsto ab.$

Using this we can define the split null extension, $B \rtimes A/B$, of B by A/B in the usual way. Then a homomorph \mathfrak{X} is said to be **split** if, whenever $A \in \mathfrak{X}$ and B is an abelian ideal of A, the split null extension of B by A/B, also belongs to \mathfrak{X} .

Lemma 4.12. Let \mathfrak{X} be a formation. Then \mathfrak{X} is split.

Proof. Suppose that $A \in \mathfrak{X}$ and that B is an abelian ideal of A. Let C be isomorphic to B as an A-bimodule, and form the split null extension $X = C \rtimes A$. Then $Y = B \oplus C$ is a direct sum of two isomorphic X-bimodules. Let D be the diagonal subalgebra of Y. Then D is an ideal of X. Also

 $A + D = A + B + C = X, A \cap D = 0$ and $X/D \cong X/C \cong A \in \mathfrak{X}$.

Thus $X = X/C \cap D \in \mathfrak{X}$ and so $X/B \in \mathfrak{X}$. But X/B is the split null extension of $B + C/B \cong B$ by A/B.

Definition 4.13. A homomorph \mathfrak{X} is saturated if all solvable algebras which have an abelian ideal have \mathfrak{X} -projectors.

Theorem 4.14. Let $\mathfrak{X} \subseteq \mathfrak{S}$ be a non-empty homomorph. Then the following are equivalent

- (1) \mathfrak{X} is saturated; and
- (2) if $A \notin \mathfrak{X}$ and A has a minimal abelian ideal B with $A/B \in \mathfrak{X}$, then $Proj_{\mathfrak{X}}(A) \neq \emptyset$.

Proof. $(1) \Rightarrow (2)$ is clear.

 $(2) \Rightarrow (1)$: Suppose that (ii) holds. We need to show that, if A is solvable and has an abelian ideal, $Proj_{\mathfrak{X}}(A) \neq \emptyset$. Let A be a minimal counter-example. Then there exists a subalgebra $S \subseteq A$ such that $B \subseteq S$, $S/B \in Proj_{\mathfrak{X}}(A/B)$.

If $S \neq A$ then there exists $C \in Proj_{\mathfrak{X}}(S)$, by the minimality of A, and $B \in Proj_{\mathfrak{X}}(A)$, by Lemma 4.5. If S = A, then $A/B \in \mathfrak{X}$. Either $A \in \mathfrak{X}$ and $A \in Proj_{\mathfrak{X}}(A)$ or $A \notin \mathfrak{X}$ and $Proj_{\mathfrak{X}}(A) \neq \emptyset$ by hypothesis. \Box

Corollary 4.15. Let \mathfrak{X} be a non-empty formation. Then the following are equivalent:

- (1) \mathfrak{X} is saturated; and
- (2) if $A \notin \mathfrak{X}$, B is a minimal abelian ideal of A and $A/B \in \mathfrak{X}$, then there is a complement to B in A.

Proof. Suppose that $A \notin \mathfrak{X}$, B is a minimal abelian ideal of A and $A/B \in \mathfrak{X}$. By Lemmas 4.9 and 4.10, $C \in Proj_{\mathfrak{X}}(A)$ if and only if C complements B in A.

Lemma 4.16. Suppose that \mathfrak{X} is a non-empty formation and $A/\Phi(A) \in \mathfrak{X}$ implies that $A \in \mathfrak{X}$. Then \mathfrak{X} is saturated.

Proof. We must show that, if $A \notin \mathfrak{X}$, B is a minimal abelian ideal of A and $A/B \in \mathfrak{X}$, then there is a complement C to B in A, by Corollary 4.15. If $B \subseteq \Phi(A)$ then $A/\Phi(A) \in \mathfrak{X}$, so $A \in \mathfrak{X}$, a contradiction. Hence $B \not\subseteq \Phi(A)$, and so there is a maximal subalgebra C of A with $B \not\subseteq C$. Then A = B + C and $B \cap C = 0$. \Box

5. Conclusions and future work

In the above, we have seen that the basic language and results from the theory of classes of groups and closure operations can be applied successfully in this general setting. But to develop the theories of formations, Schunck classes and Fitting theory appears to require extra restrictions on the algebras under consideration. Difficulties arise because the algebras in the derived series, $A^{(n)}$ are not necessarily ideals unless n = 2 and minimal ideals of solvable algebras are not necessarily abelian. As a result, the natural examples of saturated formations and Schunck classes, such as nilpotent algebras, are no longer valid unless we add extra restrictions to the definitions, as the following example shows.

Example 5.1. Let A be the three-dimensional algebra over a field K spanned by x, y, z with products $x^2 = z$, yz = z and zx = y, all other products being zero. Then $A^{(2)} = Ky + Kz$, $A^{(3)} = Kz$, $A^{(4)} = 0$, so this is a solvable algebra. However, $A^{(2)}$ is a minimal ideal of A and is not abelian; indeed, it is not even nilpotent. The

only proper subalgebras of A are Ky, Kz and $A^{(2)}$, and the only nilpotent ones are one-dimensional.

Then Ky and Kz are the only possible \mathfrak{N} -projectors. But both are subalgebras of A, $A^{(2)}$ is an ideal of A and $A/A^{(2)} \in \mathfrak{N}$, and $Ky + A^{(2)} = Kz + A^{(2)} = A^{(2)} \neq A$. Hence A has no \mathfrak{N} -projectors. Clearly \mathfrak{N} is a formation, but would not saturated if we required that all solvable algebras have \mathfrak{N} -projectors. Notice also that $\phi(A) = A^{(2)}$ is not nilpotent.

So, the next step is to find a class of non-associative algebras for which the theory can be progressed beyond this point. We would, of course, wish such a class to include the important varieties for which an advanced structure theory exists and for which such a theory has already been developed, such as Lie, Leibniz and Malcev algebras.

References

- 1. A. Ballester-Bolinches and L. M. Ezquerro. Classes of finite groups. Springer, 2006.
- 2. D. W. Barnes. On the theory of soluble Lie algebras, Math. Zeit. 106 (1968), 343-354.
- D.W. Barnes. Saturated formations of soluble Lie algebras in characteristic zero, Archiv. der Math. 30 (1978), 477-480.
- D. W. Barnes. On locally defined formations of soluble Lie and Leibniz Algebras, Bull. Aust. Math. Soc. 86 (2012), 322–326 doi:10.1017/S0004972711003443.
- D. W. Barnes. Schunck Classes of Soluble Leibniz Algebras, Communications in Algebra, 41:11, (2013) 4046-4065, DOI: 10.1080/00927872.2012.700978.
- 6. K. Doerk and T. Hawkes. Finite soluble groups. W de Gruyter, 1990.
- I. S. Gutierrez, and M. Navarro. Clases de álgebras de Lie y subálgebras de Cartan. Revista Colombiana de Matemáticas, vol 42, 2008.
- P. Hall. On the finiteness of certain soluble groups, Proc. London Math. Soc.9(3) (1959). 595–622.
- 9. P. Hall. On non-strictly simple groups, Proc. Cambridge Philos.Soc.59 (1963). 531-553.
- 10. E. I. Marshall. The Frattini subalgebra of a Lie algebra. J. London Math. Soc. 42 (1967), 416-22.
- L. A. Shemetkov, On the product of formations of algebraic systems. Algeb. Logika, 23, No. 6 (1984), 721-729.
- L.A. Shemetkov and A. N. Skiba. Formations of Algebraic Systems [in Russian], Nauka, Moscow (1989).
- 13. A.N. Skiba. Algebra of Formations [in Russian], Belarus. Navuka, Minsk (1997).
- E. L. Stitzinger. On saturated formations of solvable Lie algebras, Pacific J. Math. 47 no. 2 (1973), 531-538.
- 15. E. L. Stitzinger. Supersolvable Malcev algebras, J. Algebra 103 (1986), 69-79.
- 16. D. A. Towers. A Frattini theory for algebras. Proc. London Math. Soc., 27: 3 (1973), 440-462.
- K. A. Zhevlakov, A. M. Slin'ko, I. P. Shestakov, and A. I. Shirshov, Rings that are nearly associative, Academic Press, 1982.

Department of Mathematics and Statistics, Universidad del Norte, Km 5 via a Puerto Colombia, Barranquilla - Colombia

 $Current \; address:$ Department of Mathematics and Statistics, Universidad del Norte, Km5via a Puerto Colombia, Barranquilla - Colombia

 $Email \ address: \verb"isgutier@uninorte.edu.co"$

Department of Mathematics and Statistics, Universidad del Norte, Km 5 via a Puerto Colombia, Barranquilla - Colombia

Current address: Department of Mathematics and Statistics, Universidad del Norte, Km 5 via a Puerto Colombia, Barranquilla - Colombia

 $Email \ address: \verb"atorresblanca@uninorte.edu.co"$

DEPARTMENT OF MATHEMATICS AND STATISTICS, LANCASTER UNIVERSITY, , LANCASTER - UK

 $Current \; address:$ Department of Mathematics and Statistics, Lancaster University, , Lancaster - UK

 $Email \ address: \texttt{d.towers@lancaster.ac.uk}$