

# $\mathbb{Z}_2$ -graded polynomial identities for the Jordan algebra of $2 \times 2$ upper triangular matrices

Dimas José Gonçalves  
 Universidade Federal de São Carlos (UFSCar)  
 Departamento de Matemática  
 13565-905 São Carlos, SP, Brasil  
 e-mail: dimasjg@ufscar.br

Mateus Eduardo Salomão  
 Universidade Federal de São Carlos (UFSCar)  
 Departamento de Matemática  
 13565-905 São Carlos, SP, Brasil  
 e-mail: m.salomaoutfpr@gmail.com

November 24, 2020

**Keywords:** Upper triangular matrix algebra, Jordan algebra, Graded polynomial identities.

**2010 AMS MSC Classification:** 16R10, 17C05.

## Abstract

Let  $K$  be a field (finite or infinite) of  $\text{char}(K) \neq 2$  and let  $UT_n = UT_n(K)$  be the  $n \times n$  upper triangular matrix algebra over  $K$ . If  $\cdot$  is the usual product on  $UT_n$  then with the new product  $a \circ b = (1/2)(a \cdot b + b \cdot a)$  we have that  $UT_n$  is a Jordan algebra, denoted by  $UJ_n = UJ_n(K)$ . In this paper, we describe the set of all  $\mathbb{Z}_2$ -graded polynomial identities of  $UJ_2$  with any nontrivial  $\mathbb{Z}_2$ -grading. Moreover, we describe a linear basis for the corresponding relatively free  $\mathbb{Z}_2$ -graded algebra.

## 1 Introduction

Let  $K$  be a field and let  $UT_n = UT_n(K)$  be the algebra of  $n \times n$  upper triangular matrices over  $K$ . This algebra plays an important role in PI-Theory, and its polynomial identities were described in [9, 10].

With respect to gradings, let  $G$  be any group. The  $G$ -gradings on  $UT_n$  were described as follows: In [12], it was proved that every  $G$ -grading on  $UT_n$  is

isomorphic to an elementary  $G$ -grading ; in [2], the elementary  $G$ -gradings were classified. In [2], it was proved that two  $G$ -gradings on  $UT_n$  are isomorphic if and only if they satisfy the same  $G$ -graded polynomial identities. Moreover, the set of all  $G$ -graded polynomial identities of  $UT_n$  was described as follows: in [2] when  $K$  is an infinite field, in [4] when  $K$  is a finite field.

From now on, we assume  $\text{char}(K) \neq 2$ . Denote by  $UJ_n = UJ_n(K)$  the vector space  $UT_n$  with a new product  $\circ$  given by

$$u \circ v = (1/2)(u \cdot v + v \cdot u)$$

where  $u, v \in UT_n$ . Then  $UJ_n$  is a Jordan algebra.

In [6], the polynomial identities of  $UJ_2$  were described when  $K$  is an infinite field of  $\text{char}(K) \neq 2, 3$ . It is an open problem to describe the polynomial identities of  $UJ_n$  when  $n \geq 3$ .

With respect to gradings, let  $G$  be any group. All  $\mathbb{Z}_2$ -gradings on  $UJ_2$  were described in [6]. After, in [7] it was proved that if  $K$  is infinite, then every  $G$ -grading on  $UJ_n$  is, up to a graded isomorphism, either elementary or MT (mirror type). Moreover, in [7] the authors proved that two  $G$ -gradings on  $UJ_n$  are isomorphic if and only if they satisfy the same  $G$ -graded polynomial identities.

In [6], Koshlukov and Martino described the set of all  $\mathbb{Z}_2$ -graded polynomial identities of  $UJ_2$  when  $K$  is any field of characteristic 0 and, as a consequence, in [1] it was proved that the variety of Jordan algebras generated by  $UJ_2$  endowed with any  $G$ -grading has the Specht property.

Since almost nothing is known concerning polynomial identities of Jordan algebras, and motivated by the results above, given any nontrivial  $\mathbb{Z}_2$ -grading on  $UJ_2$  we describe the set of its  $\mathbb{Z}_2$ -graded polynomial identities when  $K$  is any field (finite or infinite). Moreover, we describe a linear basis for the corresponding relatively free  $\mathbb{Z}_2$ -graded algebra when  $K$  is any field (finite or infinite). In order to obtain our descriptions we use some ideas from the paper [6].

We draw the reader's attention to the fact that gradings and graded polynomial identities for  $UT_n$  were studied in another context too: Lie algebra. See, for example, [8].

## 2 Preliminaries

Throughout this section,  $K$  is a field (finite or infinite) with  $\text{char}(K) \neq 2$ .

Let  $UT_n = UT_n(K)$  be the  $n \times n$  upper triangular matrix algebra over  $K$ . On this algebra we consider the usual product  $\cdot$  and so it is an associative algebra. Denote by  $UJ_n = UJ_n(K)$  the vector space  $UT_n$  with a new product  $\circ$  given by

$$u \circ v = (1/2)(u \cdot v + v \cdot u)$$

where  $u, v \in UT_n$ . Then  $UJ_n$  is a Jordan algebra.

Let  $e_{ij}$  be the matrix unit in  $UJ_n$  whose  $(i, j)$ th entry equals 1 and all other entries equal 0. We denote

$$1 = e_{11} + e_{22}, \quad a = e_{11} - e_{22} \quad \text{and} \quad b = e_{12}.$$

Note that

$$a \circ a = 1 \text{ and } a \circ b = b \circ b = 0.$$

For convenience, if  $u, v \in UJ_n$  we will write  $u \circ v = uv$ .

Let  $\mathbb{Z}_2 = \{0, 1\}$  and denote by

$$UJ_2 = (UJ_2)_0 \oplus (UJ_2)_1$$

a  $\mathbb{Z}_2$ -grading on  $UJ_2$ . Thus  $(UJ_2)_i \circ (UJ_2)_j \subseteq (UJ_2)_{i+j}$  for all  $i, j \in \mathbb{Z}_2$ .

The next proposition was proved in [6].

**Proposition 2.1.** The following decompositions  $UJ_2 = (UJ_2)_0 \oplus (UJ_2)_1$  are  $\mathbb{Z}_2$ -gradings on  $UJ_2(K)$  :

1. The associative grading:  $(UJ_2)_0 = K \oplus Kb, (UJ_2)_1 = Ka$ ;
2. The scalar grading:  $(UJ_2)_0 = K, (UJ_2)_1 = Ka \oplus Kb$ ;
3. The classical grading:  $(UJ_2)_0 = K \oplus Ka, (UJ_2)_1 = Kb$ ;
4. The trivial grading:  $(UJ_2)_0 = UJ_2, (UJ_2)_1 = 0$ ;

where we identify  $K$  with the scalar matrices in  $UJ_2$ . The four gradings are pairwise nonisomorphic. They are, up to a  $\mathbb{Z}_2$ -graded isomorphism, the only  $\mathbb{Z}_2$ -gradings on  $UJ_2$ .

*Proof.* See [6, Lemma 2, Lemma 3, Proposition 4]. □

Let  $Y = \{y_1, y_2, \dots\}$  and  $Z = \{z_1, z_2, \dots\}$  be disjoint infinite sets. If  $X = Y \cup Z$ , denote by  $J(X)$  the free  $\mathbb{Z}_2$ -graded Jordan algebra, freely generated by  $X$  over  $K$ . We remember that  $J(X)$  is the free Jordan algebra, freely generated by  $X$  over  $K$ , and

$$J(X) = (J(X))_0 \oplus (J(X))_1$$

is its  $\mathbb{Z}_2$ -grading where  $\|y_i\| = 0, \|z_i\| = 1$  for all  $i$ , and if  $u, v$  are monomials in  $J(X)$  then

$$\|uv\| = \|u\| + \|v\|.$$

Note that the notation  $\|\cdot\|$  means the homogeneous degree, that is, if  $f \in (J(X))_0$  then  $\|f\| = 0$  and  $f$  is called even; if  $f \in (J(X))_1$  then  $\|f\| = 1$  and  $f$  is called odd.

We remember that a  $T_{\mathbb{Z}_2}$ -ideal of  $J(X)$  is an ideal of  $J(X)$  closed under all  $\mathbb{Z}_2$ -graded endomorphisms of  $J(X)$ . If  $W \subseteq J(X)$  we denote by  $\langle W \rangle^{T_{\mathbb{Z}_2}}$  the  $T_{\mathbb{Z}_2}$ -ideal of  $J(X)$  generated by  $W$ , that is, the minimal  $T_{\mathbb{Z}_2}$ -ideal of  $J(X)$  containing  $W$ . If  $f \in \langle W \rangle^{T_{\mathbb{Z}_2}}$  we say that  $f$  is consequence of the polynomials in  $W$ .

By using similar arguments as in [3, Proposition 4.2.3] we state the following:

**Proposition 2.2.** Let  $K$  be a field with  $|K|$  elements. Let  $f \in J(X)$ ,  $w \in X$  and

$$f = \sum_{i=0}^{d_w} f^{(i)}$$

where  $f^{(i)}$  is the homogeneous component of  $f$  with  $\deg_w f^{(i)} = i$ . If  $d_w < |K|$  then

$$\langle f \rangle^{T_{\mathbb{Z}_2}} = \langle f^{(0)}, f^{(1)}, \dots, f^{(d_w)} \rangle^{T_{\mathbb{Z}_2}}.$$

Let  $K[x_1, \dots, x_n]$  be the free commutative algebra, freely generated by  $x_1, \dots, x_n$  over  $K$ . The next lemma is consequence of [3, Proposition 4.2.3].

**Lemma 2.3.** Let  $K$  be a field with  $|K| \geq q$ . Given  $f \in K[x_1, \dots, x_n]$  write

$$f(x_1, \dots, x_n) = \sum_{d_1=0}^{q-1} \dots \sum_{d_n=0}^{q-1} \lambda_{(d_1, \dots, d_n)} x_1^{d_1} \dots x_n^{d_n},$$

where  $\lambda_{(d_1, \dots, d_n)} \in K$ . If  $f(\alpha_1, \dots, \alpha_n) = 0$  for all  $\alpha_1, \dots, \alpha_n \in K$  then  $\lambda_{(d_1, \dots, d_n)} = 0$  for all  $(d_1, \dots, d_n)$ .

If  $UJ_2 = (UJ_2)_0 \oplus (UJ_2)_1$  is a  $\mathbb{Z}_2$ -grading and  $f(y_1, \dots, y_s, z_1, \dots, z_n) \in J(X)$ , we remember that  $f$  is a  $\mathbb{Z}_2$ -graded polynomial identity for  $UJ_2$  if

$$f(Y_1, \dots, Y_s, Z_1, \dots, Z_n) = 0$$

for all  $Y_1, \dots, Y_s \in (UJ_2)_0$  and  $Z_1, \dots, Z_n \in (UJ_2)_1$ . The set of all  $\mathbb{Z}_2$ -graded polynomial identities of  $UJ_2$ , denoted by  $Id$ , is a  $T_{\mathbb{Z}_2}$ -ideal of  $J(X)$ . If the grading is the trivial, then  $Id$  was described in [6, Theorem 19] when  $K$  is infinite with  $\text{char}(K) \neq 2, 3$ . We want to describe  $Id$  when the grading is nontrivial. In this case, by Proposition 2.1, it is sufficient to consider the associative, scalar and classical gradings only.

If  $u, v, w \in J(X)$  we denote by  $(u, v, w)$  the associator, that is,

$$(u, v, w) = (uv)w - u(vw).$$

Note that

$$(w, v, u) = -(u, v, w) \text{ and } (v, u, w) = (u, v, w) - (u, w, v). \quad (1)$$

If  $f_1, f_2, \dots, f_n \in J(X)$  we use the following convention:

$$f_1 f_2 \dots f_{n-1} f_n = (f_1 f_2 \dots f_{n-1}) f_n.$$

If  $u, v, c, d \in J(X)$  then

$$uvcd + udcv + vdcu = (uv)(cd) + (uc)(vd) + (ud)(vc). \quad (2)$$

In fact, the identity (2) is true for any Jordan algebra (see [5, Chapter I.7]). Moreover, renaming and comparing two expressions from (2), we may also obtain

$$uvcd + udcv + vdcu = wvdc + uc dv + vcd u. \quad (3)$$

### 3 The associative grading

Let  $T_{\text{Ass}}(UJ_2)$  be the set of all  $\mathbb{Z}_2$ -graded polynomial identities for  $UJ_2$  with the associative grading. In this section we will describe  $T_{\text{Ass}}(UJ_2)$  for any field  $K$  of  $\text{char}(K) = p \neq 2$ .

We remember that

$$UJ_2 = (UJ_2)_0 \oplus (UJ_2)_1,$$

where

$$(UJ_2)_0 = \text{span}\{e_{11} + e_{22}, e_{12}\} \text{ and } (UJ_2)_1 = \text{span}\{e_{11} - e_{22}\}.$$

**Lemma 3.1.** The polynomials

$$(y_1, y_2, y_3), (z_1, y_1, y_2), (z_1, y_1, z_2), (z_1, z_2, z_3) \text{ and } (z_1 z_2, z_3, z_4)$$

belong to  $T_{\text{Ass}}(UJ_2)$ .

*Proof.* The proof consists of a direct verification.  $\square$

**Notation 3.2.** Let  $I$  be the  $T_{\mathbb{Z}_2}$ -ideal of  $J(X)$  generated by the polynomials in Lemma 3.1.

Define the equivalence relation  $\equiv$  on  $J(X)$  as follows: if  $f, g \in J(X)$  then

$$f \equiv g \Leftrightarrow f + I = g + I.$$

**Lemma 3.3.** The polynomials

$$(z_1, y_1, y_2), (y_1, z_1, y_2) \text{ and } (y_1, y_2, z_1)$$

belong to  $I$ .

*Proof.* By definition we have  $(z_1, y_1, y_2) \in I$ . By (1),

$$(y_1, z_1, y_2) = (z_1, y_1, y_2) - (z_1, y_2, y_1) \in I \text{ and } (y_1, y_2, z_1) = -(z_1, y_2, y_1) \in I$$

as desired.  $\square$

**Lemma 3.4.** The polynomials

$$(z_1 z_2, x_1, x_2), (x_1, z_1 z_2, x_2) \text{ and } (x_1, x_2, z_1 z_2)$$

belong to  $I$ , where  $x_1$  and  $x_2$  are any variables in  $Y \cup Z$ .

*Proof.* By Lemma 3.1 and Lemma 3.3 we obtain

$$(z_1 z_2, x_1, x_2) \in \langle (y_1, y_2, y_3), (y_1, z_1, y_2), (y_1, y_2, z_1), (z_1 z_2, z_3, z_4) \rangle^{T_{\mathbb{Z}_2}} \subseteq I.$$

By  $(z_1 z_2, x_1, x_2) \in I$  and (1) we obtain  $(x_1, z_1 z_2, x_2), (x_1, x_2, z_1 z_2) \in I$ .  $\square$

**Lemma 3.5.** The polynomial

$$((y_1y_2)z_1)z_2 - ((y_1z_1)z_2)y_2 - ((y_2z_1)z_2)y_1 + (z_1z_2)(y_1y_2)$$

belongs to  $I$ .

*Proof.* Let  $f = ((y_1y_2)z_1)z_2 - ((y_1z_1)z_2)y_2 - ((y_2z_1)z_2)y_1 + (z_1z_2)(y_1y_2)$ . If  $u = y_1$ ,  $v = z_1$ ,  $c = z_2$  and  $d = y_2$  in (2) we obtain

$$\begin{aligned} f &= ((y_1y_2)z_1)z_2 + ((y_1y_2)z_2)z_1 - (z_1y_1)(z_2y_2) - (z_2y_1)(z_1y_2) \\ &= ((y_1y_2)z_1)z_2 + ((y_1y_2)z_2)z_1 - ((z_1y_1)y_2)z_2 - ((z_2y_1)y_2)z_1 + (z_1y_1, y_2, z_2) + (z_2y_1, y_2, z_1) \\ &= (y_2, y_1, z_1)z_2 + (y_2, y_1, z_2)z_1 + (z_1y_1, y_2, z_2) + (z_2y_1, y_2, z_1). \end{aligned}$$

By Lemma 3.3 and  $(z_1, y_1, z_2) \in I$  we conclude that  $f \in I$ .  $\square$

**Lemma 3.6.** The subalgebras of  $J(X)/I$  generated by the sets

$$Y + I = \{y + I : y \in Y\} \quad \text{and} \quad Z + I = \{z + I : z \in Z\}$$

are commutative and associative.

*Proof.* Let  $A_Y$  and  $A_Z$  be the subalgebras of  $J(X)/I$  generated by the sets  $Y + I$  and  $Z + I$ , respectively. The algebra  $J(X)$  is commutative, thus  $A_Y$  and  $A_Z$  are commutative too.

Since  $(y_1, y_2, y_3) \in I$  we have that  $A_Y$  is associative.

Finally, let  $f_1, f_2, f_3$  polynomials in the variables  $z_1, z_2, \dots$ . Note that  $f_i$  is a sum of even and odd elements. We shall prove that  $f = (f_1, f_2, f_3) \in I$ . In this case, it is sufficient to suppose that each  $f_i$  is either even or odd. The next table shows that  $f$  is consequence of  $g \in I$ .

$f_1$	$f_2$	$f_3$	$g$
Even	Even	Even	$(y_1, y_2, y_3)$
Even	Even	Odd	$(y_1, y_2, z_1)$
Even	Odd	Even	$(y_1, z_1, y_2)$
Even	Odd	Odd	$(z_1z_2, x_1, x_2), (y_1, y_2, z_1)$ and $(y_1, y_2, y_3)$
Odd	Even	Even	$(z_1, y_1, y_2)$
Odd	Even	Odd	$(z_1, y_1, z_2)$
Odd	Odd	Even	$(z_1z_2, x_1, x_2), (y_1, y_2, z_1)$ and $(y_1, y_2, y_3)$
Odd	Odd	Odd	$(z_1, z_2, z_3)$

We will prove the fourth case. Let  $f_1, f_2, f_3$  be monomials even, odd, odd respectively. We will show by induction on  $\deg(f_1)$  that  $f = (f_1, f_2, f_3) \in I$ . If  $\deg(f_1) = 2$  then  $f$  is consequence of  $(z_1z_2, x_1, x_2) \in I$ . Suppose  $\deg(f_1) \geq 4$  and write  $f_1 = f'_1 f''_1$  where  $f'_1$  and  $f''_1$  are monomials with degree  $< \deg(f_1)$ . If  $f'_1$  and  $f''_1$  are odd, then  $f$  is consequence of  $(z_1z_2, x_1, x_2) \in I$ . If  $f'_1$  and  $f''_1$  are even we obtain the equivalences below as follows: by  $(y_1, y_2, z_1) \in I$  we obtain  $(\Delta 1)$ ; by induction hypothesis we obtain  $(\Delta 2)$ ; by  $(y_1, y_2, y_3) \in I$  we obtain  $(\Delta 3)$ .

$$\begin{aligned} \overbrace{(f'_1 f''_1) f_2}^{(\Delta 1)} f_3 &\equiv \overbrace{(f'_1 (f''_1 f_2))}^{(\Delta 2)} f_3 &\equiv \overbrace{f'_1 ((f''_1 f_2) f_3)}^{(\Delta 2)} &\equiv \overbrace{f'_1 (f''_1 (f_2 f_3))}^{(\Delta 3)} &\equiv (f'_1 f''_1) (f_2 f_3). \end{aligned}$$

Thus  $(f'_1 f''_1, f_2, f_3) \equiv 0$  as desired.

Since  $(f_1, f_2, f_3) = -(f_3, f_2, f_1)$  we obtain the seventh case. The other cases are trivial. Therefore the subalgebra  $A_Z$  is associative.  $\square$

**Lemma 3.7.** Let  $f, g \in J(X)$  and  $Z' = z_1 z_2 \cdots z_s$ , where  $s$  is even. Then

$$(f Z')g \equiv (fg)Z'.$$

*Proof.* Denote  $Z'' = z_1 z_2 \cdots z_{s-1}$ . By Lemma 3.4 we have

$$(f Z')g \equiv (f(Z'' z_s))g \equiv f((Z'' z_s)g) \equiv f(g(Z'' z_s)) \equiv (fg)(Z'' z_s) \equiv (fg)Z'.$$

The lemma is proved.  $\square$

Denote by  $Sym(s)$  the symmetric group of  $\{1, \dots, s\}$ .

**Lemma 3.8.** If  $s$  is odd and  $\sigma \in Sym(s)$ , then

$$(y_1 z_{\sigma(1)})(z_{\sigma(2)} \cdots z_{\sigma(s)}) \equiv (y_1 z_1)(z_2 \cdots z_s).$$

*Proof.* By Lemma 3.7 and Lemma 3.6 we have

$$\begin{aligned} (y_1 z_{\sigma(1)})(z_{\sigma(2)} \cdots z_{\sigma(s)}) &\equiv y_1(z_{\sigma(1)}(z_{\sigma(2)} \cdots z_{\sigma(s)})) \equiv y_1(z_1(z_2 \cdots z_s)) \\ &\equiv (y_1 z_1)(z_2 \cdots z_s). \end{aligned}$$

The proof is complete.  $\square$

**Lemma 3.9.** If  $s$  is even and  $\sigma \in Sym(s)$ , then

$$(((y_1 z_{\sigma(1)})z_{\sigma(2)})y_2)(z_{\sigma(3)} \cdots z_{\sigma(s)}) \equiv (((y_1 z_1)z_2)y_2)(z_3 \cdots z_s).$$

*Proof.* Suppose  $s = 2$ . Since  $(z_1, y_1, z_2) \in I$ , we obtain

$$(((y_1 z_1)z_2)y_2) \equiv ((z_1 y_1)z_2)y_2 \equiv (z_1(y_1 z_2))y_2 \equiv ((y_1 z_2)z_1)y_2.$$

If  $s \geq 4$ , denote  $Z' = z_{\sigma(3)} \cdots z_{\sigma(s)}$ . By Lemma 3.7 we have

$$(((y_1 z_{\sigma(1)})z_{\sigma(2)})y_2)Z' \equiv ((y_1 z_{\sigma(1)})(z_{\sigma(2)}Z'))y_2$$

and also

$$(((y_1 z_{\sigma(1)})z_{\sigma(2)})y_2)Z' \equiv ((y_1(z_{\sigma(1)}Z'))z_{\sigma(2)})y_2.$$

Now we use the Lemma 3.6 to order the variables  $z_1, \dots, z_s$ .  $\square$

**Lemma 3.10.** Let  $S$  be the subset of  $J(X)$  formed by all polynomials

- (a)  $Y'Z'$ ,
- (b)  $(Y'z_{j_1})Z'$ ,
- (c)  $((y_i z_{j_1})z_{j_2})Y'Z'$ ,

where  $Y' = y_{i_1} \cdots y_{i_r}$  with  $r \geq 0$  and  $i_1 \leq \dots \leq i_r$ ;  $Z' = z_{l_1} \cdots z_{l_s}$  with  $s \geq 0$  even and  $j_1 \leq j_2 \leq l_1 \leq \dots \leq l_s$ . Then the quotient vector space  $J(X)/I$  is spanned by the set of all elements  $g + I$  where  $g \in S$ .

*Proof.* Let  $A$ ,  $B$  and  $C$  be the sets of all elements  $g + I$  where  $g$  is in (a), (b) and (c), respectively. Denote  $D = A \cup B \cup C$ .

**Claim 1.** If  $Y' = y_{k_1} y_{k_2} \cdots y_{k_m}$ ,  $Y'' = y_{b_1} y_{b_2} \cdots y_{b_t}$  and  $Z' = z_{l_1} \cdots z_{l_s}$  with  $s \geq 0$  even then  $((Y' z_{j_1}) z_{j_2}) Y'' Z' + I \in \text{span} D$ .

*Proof of the Claim 1.* The proof is by induction on  $m$ . If  $m = 0$ , by Lemmas 3.7 and 3.6 we obtain  $((z_{j_1} z_{j_2}) Y'' Z' + I \in A \subset \text{span} D$ . If  $m = 1$ , by Lemmas 3.6 and 3.9 we obtain  $((y_{k_1} z_{j_1}) z_{j_2}) Y'' Z' + I \in C \subset \text{span} D$ .

Suppose  $m \geq 2$ . By  $(y_1, y_2, y_3) \equiv 0$  and Lemma 3.5 we have

$$\begin{aligned} ((Y' z_{j_1}) z_{j_2}) Y'' Z' &\equiv (((y_{k_1} y_{k_2} \cdots y_{k_m}) z_{j_1}) z_{j_2}) Y'' Z' \\ &\equiv (((((y_{k_1} \cdots y_{k_{m-1}}) z_{j_1}) z_{j_2}) y_{k_m}) Y'' Z' \\ &\quad + (((y_{k_m} z_{j_1}) z_{j_2}) (y_{k_1} \cdots y_{k_{m-1}})) Y'' Z' \\ &\quad - ((z_{j_1} z_{j_2}) (y_{k_1} y_{k_2} \cdots y_{k_m})) Y'' Z' \\ &\equiv (((y_{k_1} \cdots y_{k_{m-1}}) z_{j_1}) z_{j_2}) (y_{k_m} Y'' Z' \\ &\quad + ((y_{k_m} z_{j_1}) z_{j_2}) ((y_{k_1} \cdots y_{k_{m-1}}) Y'')) Z' \\ &\quad - ((y_{k_1} y_{k_2} \cdots y_{k_m}) Y'') ((z_{j_1} z_{j_2}) Z'). \end{aligned}$$

By Lemmas 3.6 and 3.9 it follows that  $((y_{k_m} z_{j_1}) z_{j_2}) ((y_{k_1} \cdots y_{k_{m-1}}) Y'') Z' + I \in C \subset \text{span} D$  and  $((y_{k_1} y_{k_2} \cdots y_{k_m}) Y'') ((z_{j_1} z_{j_2}) Z' + I \in A \subset \text{span} D$ . By induction,  $((y_{k_1} \cdots y_{k_{m-1}}) z_{j_1}) z_{j_2}) (y_{k_m} Y'') Z' + I \in \text{span} D$ . The Claim 1 is proved.

Now, if  $f$  is a monomial in  $J(X)$ , we shall prove by induction on  $\deg(f)$  that  $f + I \in \text{span} D$ .

The cases  $\deg(f) = 1$  and  $\deg(f) = 2$  are trivial.

Suppose  $\deg(f) \geq 3$  and write  $f = gh$  where  $g, h \in J(X)$  are monomials with degree  $< \deg(f)$ . By induction hypothesis it follows that  $g + I$  and  $h + I$  belong to  $D$ . We have six cases to consider:

1.  $g + I$  and  $h + I$  belong to  $A$ .

In this case,  $g + I = Y' Z' + I$  and  $h + I = Y'' Z'' + I$ . By Lemma 3.7 we have

$$f \equiv (Y' Z')(Y'' Z'') \equiv ((Y' Z') Y'') Z'' \equiv ((Y' Y'') Z') Z'' \equiv (Y' Y'')(Z' Z'').$$

By Lemma 3.6 it follows that  $f + I \in A \subset \text{span} D$ .

2.  $g + I$  belongs to  $A$  and  $h + I$  belongs to  $B$ .

In this case,  $g + I = Y' Z' + I$  and  $h + I = (Y'' z_{j_1}) Z'' + I$ . By Lemmas 3.7 and 3.3 we have

$$f \equiv (Y' Z')((Y'' z_{j_1}) Z'') \equiv (Y'(Y'' z_{j_1}))(Z' Z'') \equiv ((Y' Y'') z_{j_1})(Z' Z'').$$

By Lemmas 3.6 and 3.8 it follows that  $f + I \in B \subset \text{span} D$ .



3.  $g + I$  belongs to  $A$  and  $h + I$  belongs to  $C$ .

In this case,  $g + I = Y'Z' + I$  and  $h + I = (((y_i z_{j_1}) z_{j_2}) Y'') Z'' + I$ . By Lemma 3.7 and  $(y_1, y_2, y_3) \in I$  we have

$$\begin{aligned} f &\equiv (Y'Z')(((y_i z_{j_1}) z_{j_2}) Y'') Z'' \equiv (Y'(((y_i z_{j_1}) z_{j_2}) Y''))(Z'Z'') \\ &\equiv (((y_i z_{j_1}) z_{j_2}) (Y'Y''))(Z'Z''). \end{aligned}$$

By Lemmas 3.6 and 3.9 it follows that  $f + I \in C \subset \text{span}D$ .

4.  $g + I$  and  $h + I$  belong to  $B$ .

In this case,  $g + I = (Y'z_{j_1})Z' + I$  and  $h + I = (Y''z_{j_2})Z'' + I$ . By Lemma 3.7, by  $(z_1, y_1, z_2) \in I$  and Lemma 3.3 we have

$$\begin{aligned} f &\equiv ((Y'z_{j_1})Z')((Y''z_{j_2})Z'') \equiv ((Y'z_{j_1})(Y''z_{j_2}))(Z'Z'') \\ &\equiv (((Y'z_{j_1})Y'')z_{j_2})(Z'Z'') \equiv (((Y'Y'')z_{j_1})z_{j_2})(Z'Z''). \end{aligned}$$

By Claim 1 it follows that  $f + I \in \text{span}D$ .

5.  $g + I$  belongs to  $B$  and  $h + I$  belongs to  $C$ .

In this case,  $g + I = (Y'z_{j_1})Z' + I$  and  $h + I = (((y_i z_{j_2}) z_{j_3}) Y'') Z'' + I$ . We have the following congruences:

$$\begin{aligned} f &\equiv ((Y'z_{j_1})Z')(((y_i z_{j_2}) z_{j_3}) Y'') Z'' \equiv [(Y'z_{j_1})(((y_i z_{j_2}) z_{j_3}) Y'')](Z'Z'') \\ &\equiv [(((Y'z_{j_1})Y'')((y_i z_{j_2}) z_{j_3}))](Z'Z'') \equiv [(((Y'Y'')z_{j_1})((y_i z_{j_2}) z_{j_3}))](Z'Z'') \\ &\equiv [(Y'Y'')(z_{j_1}((y_i z_{j_2}) z_{j_3}))](Z'Z'') \equiv [(Y'Y'')((z_{j_1} z_{j_3})(y_i z_{j_2}))](Z'Z'') \\ &\equiv [(Y'Y'')(y_i z_{j_2})][(z_{j_1} z_{j_3})(Z'Z'')] \equiv [((Y'Y'')y_i)z_{j_2}][(z_{j_1} z_{j_3})(Z'Z'')]. \end{aligned}$$

By Lemmas 3.6 and 3.8 it follows that  $f + I \in B \subset \text{span}D$ .

6.  $g + I$  and  $h + I$  belong to  $C$ .

In this case,  $g + I = (((y_i z_{j_1}) z_{j_2}) Y') Z' + I$  and  $h + I = (((y_j z_{j_3}) z_{j_4}) Y'') Z'' + I$ . We have the following congruences:

$$\begin{aligned} f &\equiv (((y_i z_{j_1}) z_{j_2}) Y') Z' (((y_j z_{j_3}) z_{j_4}) Y'') Z'' \\ &\equiv [(((y_i z_{j_1}) z_{j_2}) ((y_j z_{j_3}) z_{j_4})) (Y'Y'')](Z'Z'') \\ &\equiv [(((y_i z_{j_1}) (y_j z_{j_3})) (z_{j_2} z_{j_4})) (Y'Y'')](Z'Z'') \\ &\equiv [(((y_i z_{j_1}) (y_j z_{j_3})) (Y'Y''))][(z_{j_2} z_{j_4})(Z'Z'')] \\ &\equiv [(((y_i y_j) z_{j_1}) z_{j_3}) (Y'Y'')][(z_{j_2} z_{j_4})(Z'Z'')]. \end{aligned}$$

By Claim 1 it follows that  $f + I \in \text{span}D$ .

The proof is complete.  $\square$

### 3.1 The associative grading, when $K$ is an infinite field

In this subsection we describe the  $\mathbb{Z}_2$ -graded identities for  $UJ_2$  with the associative grading when  $K$  is infinite.

**Theorem 3.11.** If  $K$  is an infinite field of  $\text{char}(K) \neq 2$  then  $I = T_{\text{Ass}}(UJ_2)$ , that is,  $T_{\text{Ass}}(UJ_2)$  is generated, as a  $T_{\mathbb{Z}_2}$ -ideal, by the polynomials in Lemma 3.1. Moreover, the set in Lemma 3.10 is a basis for the quotient vector space  $J(X)/I$ .

*Proof.* By Lemma 3.1 we have  $I \subseteq T_{\text{Ass}}(UJ_2)$ .

Let  $S$  be the set in Lemma 3.10 and write  $\overline{S} = \{g + T_{\text{Ass}}(UJ_2) : g \in S\}$ . Since  $I \subseteq T_{\text{Ass}}(UJ_2)$  we have by Lemma 3.10 that  $J(X)/T_{\text{Ass}}(UJ_2) = \text{span}\overline{S}$ .

We shall prove that  $\overline{S}$  is a linearly independent set. Let

$$f = \sum_{g \in S} \lambda_g g \in T_{\text{Ass}}(UJ_2), \quad \lambda_g \in K.$$

Since  $K$  is an infinite field, by Proposition 2.2 every multihomogeneous component of  $f$  belongs to  $T_{\text{Ass}}(UJ_2)$ . Thus it is sufficient to suppose the three cases below:

$$\begin{aligned} f &= \lambda(y_1^{k_1} \dots y_r^{k_r})(z_1^{t_1} \dots z_s^{t_s}) \\ &\quad + \sum_{i=1}^r \lambda_i(((y_i z_1) z_2)(y_1^{k_1} \dots y_i^{k_i-1} \dots y_r^{k_r}))(z_1^{t_1-1} z_2^{t_2-1} z_3^{t_3} \dots z_s^{t_s}) \end{aligned}$$

or

$$\begin{aligned} f &= \lambda(y_1^{k_1} \dots y_r^{k_r})(z_1^{t_1} \dots z_s^{t_s}) \\ &\quad + \sum_{i=1}^r \lambda_i(((y_i z_1) z_1)(y_1^{k_1} \dots y_i^{k_i-1} \dots y_r^{k_r}))(z_1^{t_1-2} z_2^{t_2} z_3^{t_3} \dots z_s^{t_s}) \end{aligned}$$

or

$$f = \lambda(y_1^{k_1} \dots y_r^{k_r} z_1)(z_1^{t_1} \dots z_s^{t_s})$$

where  $t_1 + \dots + t_s$  is even. We shall prove that  $\lambda = \lambda_i = 0$  for all  $i$ . Denote

$$1 = e_{11} + e_{22}, \quad a = e_{11} - e_{22} \quad \text{and} \quad b = e_{12}.$$

In the first and second cases, let  $Y_i = \alpha_i 1 + \beta_i b$  and  $Z_i = \gamma_i a$ , where  $\alpha_i, \beta_i, \gamma_i \in K$ .

We have

$$f(Y_1, \dots, Y_r, Z_1, \dots, Z_s) = \begin{bmatrix} A & B \\ 0 & A \end{bmatrix} = 0,$$

where

$$\begin{aligned} A &= \left( \lambda + \sum_{i=1}^r \lambda_i \right) \alpha_1^{k_1} \dots \alpha_r^{k_r} \gamma_1^{t_1} \dots \gamma_s^{t_s} = 0; \\ B &= \sum_{i=1}^r \left( \lambda k_i + \lambda_i(k_i - 1) + \sum_{\substack{j=1 \\ j \neq i}}^r \lambda_j k_j \right) \alpha_1^{k_1} \dots \alpha_i^{k_i-1} \dots \alpha_r^{k_r} \beta_i \gamma_1^{t_1} \dots \gamma_s^{t_s} = 0. \end{aligned}$$

Since  $\alpha_i, \beta_i, \gamma_i$  are any elements of  $K$ , and  $K$  is infinite, we have by Lemma 2.3 that

$$\lambda + \sum_{i=1}^r \lambda_i = 0 \quad (4)$$

and also

$$\lambda k_i + \lambda_i(k_i - 1) + \sum_{\substack{j=1 \\ j \neq i}}^r \lambda_j k_i = 0$$

for all  $i = 1, \dots, r$  that is

$$\lambda k_i + \lambda_1 k_i + \dots + \lambda_i(k_i - 1) + \dots + \lambda_r k_i = 0 \quad (5)$$

for all  $i = 1, \dots, r$ . By the equalities (4) and (5) we obtain the system

$$\begin{cases} \lambda + \lambda_1 + \lambda_2 + \dots + \lambda_r = 0 \\ \lambda k_1 + \lambda_1(k_1 - 1) + \lambda_2 k_1 + \dots + \lambda_r k_1 = 0 \\ \lambda k_2 + \lambda_1 k_2 + \lambda_2(k_2 - 1) + \dots + \lambda_r k_2 = 0 \\ \vdots \\ \lambda k_r + \lambda_1 k_r + \lambda_2 k_r + \dots + \lambda_r(k_r - 1) = 0 \end{cases}$$

with augmented matrix

$$\begin{bmatrix} 1 & 1 & 1 & \dots & 1 & 0 \\ k_1 & k_1 - 1 & k_1 & \dots & k_1 & 0 \\ k_2 & k_2 & k_2 - 1 & \dots & k_2 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ k_r & k_r & k_r & \dots & k_r - 1 & 0 \end{bmatrix} \sim \begin{bmatrix} 1 & 1 & 1 & \dots & 1 & 0 \\ 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 1 & 0 \end{bmatrix}.$$

Thus,  $\lambda = \lambda_i = 0$  for all  $i = 1, \dots, r$ .

In the third case, that is,

$$f(y_1, \dots, y_r, z_1, \dots, z_s) = \lambda(y_1^{k_1} \dots y_r^{k_r} z_1)(z_1^{t_1} \dots z_s^{t_s})$$

where  $t_1 + \dots + t_s$  is even, let  $Y_i = 1$  and  $Z_i = a$  for all  $i$ . Then

$$f(Y_1, \dots, Y_r, Z_1, \dots, Z_s) = \lambda a = 0$$

and so  $\lambda = 0$ .

Therefore, the set  $\overline{S}$  is a basis for the quotient vector space  $J(X)/T_{\text{Ass}}(UJ_2)$ . Moreover, since  $I \subseteq T_{\text{Ass}}(UJ_2)$ , by Lemma 3.10 we have  $I = T_{\text{Ass}}(UJ_2)$ .  $\square$

### 3.2 The associative grading, when $K$ is a finite field

In this subsection we describe the  $\mathbb{Z}_2$ -graded identities for  $UJ_2$  with the associative grading when  $K$  is finite. Throughout this subsection,  $K$  is a finite field with  $|K| = q$  elements and  $\text{char}(K) \neq 2$ .

Since  $(K - \{0\}, \cdot)$  is a group with  $q - 1$  elements it follows that  $x^{q-1} = 1$  for all  $x \in K - \{0\}$ . Therefore,  $x^q = x$  for all  $x \in K$ .

A direct consequence of this fact is the following lemma.

**Lemma 3.12.** The polynomials

$$(y_1^q - y_1)(y_2^q - y_2), \quad z_1^q - z_1 \quad \text{and} \quad (y_1^q - y_1)z_1$$

belong to  $T_{\text{Ass}}(UJ_2)$ .

**Notation 3.13.** Let  $I'$  be the  $T_{\mathbb{Z}_2}$ -ideal of  $J(X)$  generated by the polynomials in Lemmas 3.1 and 3.12.

**Lemma 3.14.** The polynomial  $(y_1^q, x_1, x_2)$  belongs to  $I'$ , where  $x_1$  and  $x_2$  are any variables in  $Y \cup Z$ .

*Proof.* Since  $(y_1, y_2, y_3), (y_1, z_1, y_2), (y_1, y_2, z_1) \in I \subseteq I'$  (see definition of  $I$  and Lemma 3.3) it follows that  $(y_1^q, x_1, x_2) \in I'$  when  $x_1 \in Y$  or  $x_2 \in Y$ .

We will prove that  $(y_1^q, z_1, z_2) \in I'$ .

**Claim 1.**  $(y_1^n, z_1, z_2) + I' = n[(y_1 z_1)z_2]y_1^{n-1} - n[(z_1 z_2)y_1^n] + I'$  for all  $n \geq 1$ .

*Proof of the claim.* The case  $n = 1$  is trivial. Suppose  $n \geq 2$ . By Lemmas 3.5 and 3.7,

$$\begin{aligned} (y_1^n, z_1, z_2) + I' &= (y_1^n z_1)z_2 - y_1^n(z_1 z_2) + I' \\ &= ((y_1^{n-1} z_1)z_2)y_1 + ((y_1 z_1)z_2)y_1^{n-1} - (z_1 z_2)y_1^n - y_1^n(z_1 z_2) + I' \\ &= (y_1^{n-1}, z_1, z_2)y_1 + ((y_1 z_1)z_2)y_1^{n-1} - (z_1 z_2)y_1^n + I'. \end{aligned}$$

Now we apply the induction hypothesis on the first summand and  $(y_1, y_2, y_3) \in I'$  to conclude the proof of the claim.

In particular, if  $n = q$  then

$$(y_1^q, z_1, z_2) + I' = q[(y_1 z_1)z_2]y_1^{q-1} - q[(z_1 z_2)y_1^q] + I' = I'$$

and the proof is complete.  $\square$

**Lemma 3.15.** The following equality is valid:

$$((y_i z_1)z_2)y_j^q + I' = ((y_i z_1)z_2)y_j + ((y_j z_1)z_2)y_i - (z_1 z_2)(y_i y_j) + I'.$$

*Proof.* Let  $g = ((y_i z_1)z_2)y_j^q$ . By Lemma 3.14 we have

$$g + I' = (y_i z_1)(z_2 y_j^q) + I'.$$

Since  $(y_j^q - y_j)z \in I'$  we obtain

$$y_j^q z + I' = y_j z + I'.$$

We use this equality,  $(z_1, y_1, z_2) \in I'$ ,  $(z_1, y_1, y_2) \in I'$  and Lemma 3.5 to obtain

$$\begin{aligned} g + I' &= (y_i z_1)(y_j z_2) + I' = ((y_i z_1)y_j)z_2 + I' = ((y_i y_j)z_1)z_2 + I' \\ &= ((y_i z_1)z_2)y_j + ((y_j z_1)z_2)y_i - (z_1 z_2)(y_i y_j) + I' \end{aligned}$$

as desired.  $\square$

Denote by  $\Lambda_n$  the set of all elements  $(s_1, \dots, s_n) \in \mathbb{Z}^n$  such that:

- a)  $0 \leq s_1, \dots, s_n < 2q$ ;
- b) If  $s_i \geq q$  for some  $i$ , then  $s_j < q$  for all  $j \neq i$ .

**Lemma 3.16.** The quotient vector space  $J(X)/I'$  is spanned by the set of all polynomials  $g + I'$  such that

- (a)  $g = \overline{Y}$  or
- (b)  $g = Y'_1 Z'_1$  or
- (c)  $g = (Y'_1 z_j) Z'_2$  or
- (d)  $g = (((y_i z_j) z_l) Y'_1) Z'_3$

where

- $\overline{Y} = y_1^{k_1} \dots y_r^{k_r}$  with  $(k_1, \dots, k_r) \in \Lambda_r$  and  $r \geq 1$ ;
- $Y'_1 = y_1^{k_1} \dots y_r^{k_r}$  with  $0 \leq k_1, \dots, k_r < q$  and  $r \geq 1$ ;
- $Z'_1 = z_1^{t_1} \dots z_s^{t_s}$  with  $0 \leq t_1, \dots, t_s < q$ ,  $s \geq 1$  and  $t_1 + \dots + t_s > 0$  even;
- $Z'_2 = z_j^{t_j} z_{j+1}^{t_{j+1}} \dots z_s^{t_s}$  with  $j \geq 1$ ,  $s \geq 1$ ,  $0 \leq t_j < q - 1$ ,  $0 \leq t_{j+1}, \dots, t_s < q$  and  $t_j + \dots + t_s \geq 0$  even;
- $Z'_3 = z_l^{t_l} z_{l+1}^{t_{l+1}} \dots z_s^{t_s}$  with  $1 \leq j \leq l$ ,  $s \geq 1$ ,  $0 \leq t_{l+1}, \dots, t_s < q$  and  $t_l + t_{l+1} + \dots + t_s \geq 0$  even. Moreover, if  $j < l$  then  $0 \leq t_l < q - 1$ . If  $j = l$  then  $0 \leq t_l < q - 2$ .

*Proof.* Firstly, with respect to  $\overline{Y}, Y'_1, Z'_1, Z'_2$  and  $Z'_3$ , these polynomials are well defined. See Lemma 3.6.

Let  $A, B, C$  and  $D$  be the sets of all elements  $g + I'$  where  $g$  is in (a), (b), (c) and (d) respectively. Denote  $E = A \cup B \cup C \cup D$ . Let  $f$  be a monomial in  $J(X)$ , we shall prove that  $f + I' \in \text{span}E$ .

Since  $I \subseteq I'$  we have by Lemma 3.10 that the quotient vector space  $J(X)/I'$  is spanned by the set of all polynomials:

- (a')  $Y' Z' + I'$ ,
- (b')  $(Y' z_j) Z'' + I'$ ,
- (c')  $(((y_i z_j) z_l) Y') Z'' + I'$ ,

where  $Y' = y_1^{k_1} y_2^{k_2} \dots y_r^{k_r}$ ;  $0 \leq k_1, \dots, k_r$ ;  $r \geq 1$ ;  $Z' = z_1^{l_1} z_2^{l_2} \dots z_s^{l_s}$ ;  $0 \leq l_1, l_2, \dots, l_s$ ;  $s \geq 1$ ;  $l_1 + l_2 + \dots + l_s$  even;  $Z'' = z_l^{t_l} z_{l+1}^{t_{l+1}} \dots z_s^{t_s}$ ;  $0 \leq t_l, t_{l+1}, \dots, t_s$ ;  $1 \leq j \leq l$ ;  $s \geq 1$ ;  $t_l + t_{l+1} + \dots + t_s$  even.

Let  $A', B'$  and  $C'$  be the sets of all elements in (a'), (b') and (c'), respectively. We shall prove that  $A' \cup B' \cup C' \subseteq \text{span}E$ .

*Case 1.*  $f + I' \in A'$ .

In this case,  $f + I' = (y_1^{k_1} \cdots y_r^{k_r})(z_1^{l_1} \cdots z_s^{l_s}) + I'$ . Since  $(y_i^q - y_i)(y_j^q - y_j) \in I'$  we obtain

$$y_i^q y_j^q + I' = y_i^q y_j + y_i y_j^q - y_i y_j + I' \text{ and } y_i^{2q} + I' = 2y_i^{q+1} - y_i^2 + I'.$$

By using several times these two equalities and Lemma 3.6, we can suppose  $(k_1, \dots, k_r) \in \Lambda_r$ .

If  $l_1 = \dots = l_s = 0$  then  $f + I' \in A \subset \text{span}E$ .

Suppose  $l_i \neq 0$  for some  $i$ . Since  $(z_i^q - z_i) \in I'$  we obtain

$$z_i^q + I' = z_i + I'. \quad (6)$$

By using several times this equality, we can suppose  $0 \leq l_1, \dots, l_s < q$ . We have two cases to consider:  $0 \leq k_1, \dots, k_r < q$  or  $k_m \geq q$  for some  $m$ . In the first case,  $f + I' \in B \subset \text{span}E$ . In the second case, write  $k_m = q + u_m$ , where  $0 \leq u_m < q$ . Let

$$j_1 = \min\{j \mid l_j \geq 1\}.$$

If  $l_{j_1} - 1 \geq 1$  denote  $j_2 = j_1$ ; otherwise denote

$$j_2 = \min\{j \mid l_j \geq 1; j \neq j_1\}.$$

By Lemmas 3.6 and 3.7, we have

$$f + I' = [(y_m^q(z_{j_1} z_{j_2})) \underbrace{(y_1^{k_1} \cdots y_m^{u_m} \cdots y_r^{k_r})}_{Y'_1}] \underbrace{(z_{j_1}^{l_{j_1}-1} z_{j_2}^{l_{j_2}-1} \cdots z_s^{l_s})}_{Z'_3} + I'.$$

Since  $(y_m^q - y_m)z \in I'$  we obtain

$$y_m^q z + I' = y_m z + I'. \quad (7)$$

By using this equality and Lemma 3.14 we obtain

$$f + I' = [((y_m^q z_{j_1}) z_{j_2}) Y'_1] Z'_3 + I' = [((y_m z_{j_1}) z_{j_2}) Y'_1] Z'_3 + I' \in D \subset \text{span}E.$$

This case is finished.

*Case 2.*  $f + I' \in B'$ .

In this case,  $f + I' = ((y_1^{k_1} \cdots y_r^{k_r}) z_j)(z_j^{t_j} z_{j+1}^{t_{j+1}} \cdots z_s^{t_s}) + I'$ . By Lemma 3.3 and (7), we can suppose  $0 \leq k_1, \dots, k_r < q$ . By (6), we can suppose  $0 \leq t_j, t_{j+1}, \dots, t_s < q$ . If  $t_j < q - 1$  then  $f + I' \in C \subset \text{span}E$ . If  $t_j = q - 1$ , by Lemma 3.6, Lemma 3.7 and (6) we obtain

$$\begin{aligned} f + I' &= \underbrace{((y_1^{k_1} \cdots y_r^{k_r}) z_j)}_{Y'_1} \underbrace{(z_j^{q-1} z_{j+1}^{t_{j+1}} \cdots z_s^{t_s})}_{Z'_2} + I' = (Y'_1 z_j)(z_j^{q-1} Z'_2) + I' \\ &= (Y'_1 z_j^q) Z'_2 + I' = (Y'_1 z_j) Z'_2 + I' \in C \subset \text{span}E. \end{aligned}$$

Case 3.  $f + I' \in C'$ .

In this case,  $f + I' = (((y_i z_j) z_l) \underbrace{(y_1^{k_1} \dots y_r^{k_r})}_{Y'_1} \underbrace{(z_l^{t_l} z_{l+1}^{t_{l+1}} \dots z_s^{t_s})}_{Z'_3}) + I'$ . By Lemma

3.14, Lemma 3.15 and Case 1, we can suppose  $0 \leq k_1, \dots, k_r < q$ . By (6), we can suppose  $0 \leq t_l, \dots, t_s < q$  too. We have two cases to consider:

(3.a)  $j < l$ .

In this case,  $f + I' = (((y_i z_j) z_l) Y'_1) (z_l^{t_l} Z'_3) + I'$ . If  $t_l = q - 1$ , by Lemma 3.7 and (6) we have

$$f + I' = (((y_i z_j) z_l^q) Y'_1) Z'_3 + I' = (((y_i z_j) z_l) Y'_1) Z'_3 + I'.$$

Thus, we can suppose  $0 \leq t_l < q - 1$ . In this case,  $f + I' \in D \subset \text{span}E$ .

(3.b)  $j = l$ .

In this case,  $f + I' = (((y_i z_l) z_l) Y'_1) (z_l^{t_l} Z'_3) + I'$ . If  $t_l = q - 1$ , we can use the same argument as in Case 3.a. If  $t_l = q - 2$  then there exists  $k > l$  such that  $t_k \geq 1$ , because  $q - 2$  is odd. Write  $f + I' = (((y_i z_l) z_l) Y'_1) ((z_l^{q-2} z_k) Z''_3) + I'$ . By Lemma 3.7 and (6) we obtain

$$\begin{aligned} f + I' &= (((y_i z_l) (z_l^{q-1} z_k)) Y'_1) Z''_3 + I' \\ &= (((y_i z_l^q) z_k) Y'_1) Z''_3 + I' = (((y_i z_l) z_k) Y'_1) Z''_3 + I' \in D \subset \text{span}E. \end{aligned}$$

Thus, we can suppose  $0 \leq t_l < q - 2$ . In this case,  $f + I' \in D \subset \text{span}E$ .

The lemma is proved.  $\square$

**Theorem 3.17.** If  $K$  is a finite field with  $|K| = q$  elements and  $\text{char}(K) \neq 2$  then  $I' = T_{\text{Ass}}(UJ_2)$ , that is,  $T_{\text{Ass}}(UJ_2)$  is generated, as a  $T_{\mathbb{Z}_2}$ -ideal, by the polynomials in Lemmas 3.1 and 3.12. Moreover, the set in Lemma 3.16 is a basis for the quotient vector space  $J(X)/I'$ .

*Proof.* By Lemmas 3.1 and 3.12 we have  $I' \subseteq T_{\text{Ass}}(UJ_2)$ .

Denote by  $S'$  the set of all polynomials  $g$  in Lemma 3.16 - item (a). Denote by  $S''$  the set of all polynomials  $g$  in Lemma 3.16 - items (b), (c), (d). Let  $S = S' \cup S''$  and  $\overline{S} = \{g + T_{\text{Ass}}(UJ_2) : g \in S\}$ . Since  $I' \subseteq T_{\text{Ass}}(UJ_2)$ , by Lemma 3.16 it follows that  $J(X)/T_{\text{Ass}}(UJ_2) = \text{span}\overline{S}$ .

We shall prove that  $\overline{S}$  is a linearly independent set. Let

$$f(y_1, \dots, y_r, z_1, \dots, z_s) = \sum_{g \in S} \lambda_g g \in T_{\text{Ass}}(UJ_2), \quad \lambda_g \in K.$$

In particular,

$$h = f(y_1, \dots, y_r, 0, \dots, 0) = \sum_{g \in S'} \lambda_g g = \sum_{k \in \Lambda_r} \lambda_k y_1^{k_1} \dots y_r^{k_r} \in T_{\text{Ass}}(UJ_2), \quad k = (k_1, \dots, k_r).$$

Let  $*$  be the involution on the associative algebra  $UT_2$  defined by:

$$\begin{pmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{pmatrix}^* = \begin{pmatrix} a_{22} & a_{12} \\ 0 & a_{11} \end{pmatrix}.$$

Note that the symmetric elements of  $UT_2$  form a vector subspace with basis  $\{e_{11} + e_{22}, e_{12}\}$ . Moreover, if  $u, v \in (UJ_2)_0$  then

$$u \circ v = u \cdot v$$

where  $\cdot$  is the usual product of  $UT_2$ . Thus  $h = h(y_1, \dots, y_r)$  is a  $*$ -polynomial identity for  $UT_2$  if  $y_1, y_2, \dots$  are symmetric variables. By [11, Lemma 5.8], we obtain  $\lambda_k = 0$  for all  $k \in \Lambda_r$ .

In particular,

$$f(y_1, \dots, y_r, z_1, \dots, z_s) = \sum_{g \in S''} \lambda_g g.$$

Write

$$f = \sum_t f_t, \quad t = (t_1, \dots, t_s),$$

where  $f_t$  is multihomogeneous with respect to variables  $z_1, \dots, z_s$  and  $\deg_{z_i} f_t = t_i$  for all  $i$ . Since  $|K| = q$  and  $\deg_{z_i} f < q$  for all  $i$ , by Proposition 2.2 we have  $f_t \in T_{\text{Ass}}(UJ_2)$  for all  $t$  and

$$\begin{aligned} f_t &= \sum_k \lambda_k (y_1^{k_1} \dots y_r^{k_r}) (z_1^{t_1} \dots z_s^{t_s}) \\ &\quad + \sum_{i=1}^r \sum_k \lambda_{(i,k)} (((y_i z_1) z_2) (y_1^{k_1} \dots y_i^{k_i} \dots y_r^{k_r})) (z_1^{t_1-1} z_2^{t_2-1} z_3^{t_3} \dots z_s^{t_s}) \end{aligned} \quad (8)$$

or

$$\begin{aligned} f_t &= \sum_k \lambda_k (y_1^{k_1} \dots y_r^{k_r}) (z_1^{t_1} \dots z_s^{t_s}) \\ &\quad + \sum_{i=1}^r \sum_k \lambda_{(i,k)} (((y_i z_1) z_1) (y_1^{k_1} \dots y_i^{k_i} \dots y_r^{k_r})) (z_1^{t_1-2} z_2^{t_2} \dots z_s^{t_s}) \end{aligned} \quad (9)$$

or

$$f_t = \sum_k \lambda_k ((y_1^{k_1} \dots y_r^{k_r}) z_1) (z_1^{t_1-1} z_2^{t_2} \dots z_s^{t_s}) \quad (10)$$

where  $k = (k_1, \dots, k_r)$ ,  $0 \leq k_j < q$  for all  $j$ . We shall prove that  $\lambda_k = \lambda_{(i,k)} = 0$  for all  $k, i$ .

Suppose  $f_t$  as in (8). Let  $Y_i = \alpha_i 1 + \beta_i e_{12}$  and  $Z_i = e_{11} - e_{22}$ , where  $\alpha_i, \beta_i \in K$ . We have

$$f_t(Y_1, \dots, Y_r, Z_1, \dots, Z_s) = \begin{bmatrix} A & B \\ 0 & A \end{bmatrix} = 0,$$

where



$$\begin{aligned}
A &= \sum_k \lambda_k \alpha_1^{k_1} \cdots \alpha_r^{k_r} + \sum_{i=1}^r \sum_{\substack{k \\ k_i < q-1}} \lambda_{(i,k)} \alpha_1^{k_1} \cdots \alpha_i^{k_i+1} \cdots \alpha_r^{k_r} + \sum_{i=1}^r \sum_{\substack{k \\ k_i = q-1}} \lambda_{(i,k)} \alpha_1^{k_1} \cdots \alpha_i^q \cdots \alpha_r^{k_r}; \\
B &= \sum_k \lambda_k \left( \sum_{j=1}^r k_j \alpha_1^{k_1} \cdots \alpha_j^{k_j-1} \beta_j \cdots \alpha_r^{k_r} \right) \\
&\quad + \sum_{i=1}^r \sum_{\substack{k \\ k_i < q-1}} \lambda_{(i,k)} \left( k_i \alpha_1^{k_1} \cdots \alpha_i^{k_i} \beta_i \cdots \alpha_r^{k_r} + \sum_{\substack{j=1 \\ j \neq i}}^r k_j \alpha_1^{k_1} \cdots \alpha_i^{k_i+1} \cdots \alpha_j^{k_j-1} \beta_j \cdots \alpha_r^{k_r} \right) \\
&\quad + \sum_{i=1}^r \sum_{\substack{k \\ k_i = q-1}} \lambda_{(i,k)} \left( (q-1) \alpha_1^{k_1} \cdots \alpha_i^{q-1} \beta_i \cdots \alpha_r^{k_r} + \sum_{\substack{j=1 \\ j \neq i}}^r k_j \alpha_1^{k_1} \cdots \alpha_i^q \cdots \alpha_j^{k_j-1} \beta_j \cdots \alpha_r^{k_r} \right).
\end{aligned}$$

Since  $B = 0$  for all  $\alpha_1, \dots, \alpha_r, \beta_1, \dots, \beta_r \in K$  and  $\deg_{\beta_i} B = 1 < q$  for all  $i$ , it follows that every homogeneous component of  $B$  with respect to  $\beta_i$  of degree 1 is zero too. Thus,  $B = B_1 + \dots + B_r$  where

$$\begin{aligned}
B_i &= \sum_k \lambda_k k_i \alpha_1^{k_1} \cdots \alpha_i^{k_i-1} \beta_i \cdots \alpha_r^{k_r} + \sum_{\substack{k \\ k_i < q-1}} \lambda_{(i,k)} k_i \alpha_1^{k_1} \cdots \alpha_i^{k_i} \beta_i \cdots \alpha_r^{k_r} \\
&\quad + \sum_{\substack{l=1 \\ l \neq i}}^r \sum_{\substack{k \\ k_l < q-1}} \lambda_{(l,k)} k_i \alpha_1^{k_1} \cdots \alpha_l^{k_l+1} \cdots \alpha_i^{k_i-1} \beta_i \cdots \alpha_r^{k_r} + \sum_{\substack{k \\ k_i = q-1}} \lambda_{(i,k)} (q-1) \alpha_1^{k_1} \cdots \alpha_i^{q-1} \beta_i \cdots \alpha_r^{k_r} \\
&\quad + \sum_{\substack{l=1 \\ l \neq i}}^r \sum_{\substack{k \\ k_l = q-1}} \lambda_{(l,k)} k_i \alpha_1^{k_1} \cdots \alpha_l^q \cdots \alpha_i^{k_i-1} \beta_i \cdots \alpha_r^{k_r} = 0.
\end{aligned}$$

and  $B_1 = \dots = B_r = 0$ . Since  $B_i = 0$  for all  $\alpha_1, \dots, \alpha_r, \beta_i \in K$  and  $\deg_{\alpha_i} B_i < q$  it follows that every homogeneous component of  $B_i$  with respect to  $\alpha_i$  is zero too. Thus,  $B_i = B_{i,0} + B_{i,1} + \dots + B_{i,q-1}$  where  $\deg_{\alpha_i} B_{i,j} = j$ , and

$$B_{i,q-1} = \sum_{\substack{k \\ k_i = q-1}} \lambda_{(i,k)} (q-1) \alpha_1^{k_1} \cdots \alpha_i^{q-1} \beta_i \cdots \alpha_r^{k_r} = 0.$$

Since  $B_{i,q-1}$  is a polynomial identity for  $K$  and  $\deg_{\alpha_j} B_{i,q-1} < q$  for all  $j$ , by Lemma 2.3 we obtain  $\lambda_{(i,k)} = 0$  for all  $i$  and  $k = (k_1, \dots, k_{i-1}, q-1, k_{i+1}, \dots, k_r)$ .

In particular, by (8) we have

$$\begin{aligned}
f_t &= \sum_k \lambda_k y_1^{k_1} \cdots y_r^{k_r} z_1^{t_1} \cdots z_s^{t_s} \\
&\quad + \sum_{i=1}^r \sum_{\substack{k \\ k_i < q-1}} \lambda_{(i,k)} (((y_i z_1) z_2) (y_1^{k_1} \cdots y_i^{k_i} \cdots y_r^{k_r})) (z_1^{t_1-1} z_2^{t_2-1} z_3^{t_3} \cdots z_s^{t_s}).
\end{aligned}$$

Since  $\deg_{y_j} f_t < q$  and  $\deg_{z_j} f_t < q$  for all  $j$ , every multihomogeneous component of  $f_t$  belongs to  $T_{\text{Ass}}(UJ_2)$ . Now, using similar arguments as in Theorem 3.11 it follows that  $\lambda_k = \lambda_{(i,k)} = 0$  for all  $i, k$  as desired.

The second case (9) is analogous to the first (8).  
Now, let  $f_t$  as in (10), that is,

$$f_t = \sum_k \lambda_k((y_1^{k_1} \cdots y_r^{k_r}) z_1)(z_1^{t_1-1} \cdots z_s^{t_s}).$$

Since  $\deg_{y_j} f_t < q$  for all  $j$ , it follows that

$$f_{t,k}(y_1, \dots, y_r, z_1, \dots, z_s) = \lambda_k((y_1^{k_1} \cdots y_r^{k_r}) z_1)(z_1^{t_1-1} \cdots z_s^{t_s}) \in T_{\text{Ass}}(UJ_2)$$

for all  $k$ . Thus, if  $Y_i = 1$  and  $Z_i = e_{11} - e_{22}$  then

$$0 = f_{t,k}(Y_1, \dots, Y_r, Z_1, \dots, Z_s) = \lambda_k(e_{11} - e_{22})$$

and so  $\lambda_k = 0$  for all  $k$ .

Therefore, the set  $\bar{S}$  is a basis for the quotient vector space  $J(X)/T_{\text{Ass}}(UJ_2)$ .  
Moreover, since  $I' \subseteq T_{\text{Ass}}(UJ_2)$ , by Lemma 3.16 we have  $I' = T_{\text{Ass}}(UJ_2)$ .  $\square$

## 4 The scalar grading

Let  $T_{\text{Sca}}(UJ_2)$  be the set of all  $\mathbb{Z}_2$ -graded polynomial identities for  $UJ_2$  with the scalar grading. In this section we will describe  $T_{\text{Sca}}(UJ_2)$  for any field  $K$  of  $\text{char}(K) = p \neq 2$ .

We remember that

$$UJ_2 = (UJ_2)_0 \oplus (UJ_2)_1,$$

where

$$(UJ_2)_0 = \text{span}\{e_{11} + e_{22}\} \text{ and } (UJ_2)_1 = \text{span}\{e_{11} - e_{22}, e_{12}\}.$$

**Lemma 4.1.** The polynomials

$$(y_1, y_2, y_3), (z_1, y_1, y_2), (y_1, z_1, z_2) \text{ and } z_1(z_2, z_3, z_4)$$

belong to  $T_{\text{Sca}}(UJ_2)$ .

*Proof.* The proof consists of a direct verification.  $\square$

**Notation 4.2.** Let  $I$  be the  $T_{\mathbb{Z}_2}$ -ideal of  $J(X)$  generated by the polynomials in Lemma 4.1.

Define the equivalence relation  $\equiv$  on  $J(X)$  as follows: if  $f, g \in J(X)$  then

$$f \equiv g \Leftrightarrow f + I = g + I.$$

**Lemma 4.3.** Let  $x_1, x_2, x_3 \in Y \cup Z$ .

- (a) If  $x_i \in Y$  for some  $1 \leq i \leq 3$ , then  $(x_1, x_2, x_3) \in I$ .
- (b)  $z_1(x_1, x_2, x_3) \in I$ .

*Proof.* By  $(y_1, y_2, y_3), (z_1, y_1, y_2), (y_1, z_1, z_2) \in I$  and (1) we prove (a). By  $z_1(z_2, z_3, z_4) \in I$  and (a) we prove (b).  $\square$

**Lemma 4.4.** If  $s$  is even and  $\sigma \in \text{Sym}(s)$  then

$$(z_{\sigma(1)}z_{\sigma(2)})(z_{\sigma(3)}z_{\sigma(4)})\cdots(z_{\sigma(s-1)}z_{\sigma(s)}) \equiv (z_1z_2)(z_3z_4)\cdots(z_{s-1}z_s).$$

*Proof.* Since  $(z_{\sigma(i)}z_{\sigma(i+1)})$  is an even polynomial, by Lemma 4.3 - (a) it is sufficient to prove  $(z_1z_2)(z_3z_4) \equiv (z_1z_3)(z_2z_4)$ . By Lemma 4.3 we have

$$\begin{aligned} (z_1z_2)(z_3z_4) &\equiv z_1(z_2(z_3z_4)) \equiv z_1((z_2z_3)z_4) \equiv z_1((z_3z_2)z_4) \\ &\equiv z_1(z_3(z_2z_4)) \equiv (z_1z_3)(z_2z_4). \end{aligned}$$

The proof is complete.  $\square$

**Lemma 4.5.** Let  $S$  be the subset of  $J(X)$  formed by all polynomials

- (a)  $Y'Z'$  and
- (b)  $Y'(z_{i_0}Z')$ ,

where  $Y' = y_1^{k_1}\cdots y_r^{k_r}$ ;  $k_i \geq 0$  for all  $i$ ;  $r \geq 0$ ;  $Z' = (z_{i_1}z_{j_1})(z_{i_2}z_{j_2})\cdots(z_{i_t}z_{j_t})$ ;  $i_1 \leq j_1 \leq i_2 \leq j_2 \leq \cdots \leq i_t \leq j_t$ ;  $t \geq 0$ ;  $i_0 \geq 1$ . Then the quotient vector space  $J(X)/I$  is spanned by the set of all elements  $g + I$  where  $g \in S$ .

*Proof.* Let  $A$  and  $B$  be the sets of all elements  $g + I$  where  $g$  is in (a) and (b), respectively. Denote  $C = A \cup B$ . If  $f(y_1, \dots, y_r, z_1, \dots, z_s)$  is a monomial in  $J(X)$ , we shall prove that  $f + I \in \text{span}C$ .

By Lemma 4.3 - (a) we have

$$f \equiv y_1^{k_1}\cdots y_r^{k_r}g(z_1, \dots, z_s), \quad (11)$$

where  $g(z_1, \dots, z_s)$  is a monomial in the variables  $z_1, z_2, \dots, z_s$ .

Let  $A_Z$  be the subalgebra of  $J(X)/I$  generated by the set  $\{z + I : z \in Z\}$ .

**Claim 1.** The vector space  $A_Z$  is spanned by all elements  $Z'' + I$  and  $z_{i_0}Z'' + I$  where  $Z'' = (z_{i_1}z_{j_1})(z_{i_2}z_{j_2})\cdots(z_{i_t}z_{j_t})$  and  $t \geq 0$ .

*Proof of the Claim 1.* Let  $h \in A_Z$  be a monomial. We will prove the result by induction on  $\deg(h) = n$ . For  $n = 1, 2, 3$  is trivial. Suppose  $n \geq 4$  and write  $h = h_1h_2$  where  $\deg(h_1), \deg(h_2) < n$ . We use the induction hypothesis on  $h_1$  and  $h_2$ , and by Lemma 4.3 - (a) we have the desired.

By (11), Claim 1 and Lemma 4.4 we finish the proof of the lemma.  $\square$

## 4.1 The scalar grading, when $K$ is an infinite field

In this subsection we describe the  $\mathbb{Z}_2$ -graded identities for  $UJ_2$  with the scalar grading when  $K$  is infinite.

**Theorem 4.6.** If  $K$  is an infinite field of  $\text{char}(K) \neq 2$  then  $I = T_{\text{Sca}}(UJ_2)$ , that is,  $T_{\text{Sca}}(UJ_2)$  is generated, as a  $T_{\mathbb{Z}_2}$ -ideal, by the polynomials in Lemma 4.1. Moreover, the set in Lemma 4.5 is a basis for the quotient vector space  $J(X)/I$ .

*Proof.* If  $g = g(z_1, \dots, z_n) = z_{i_0}(z_{i_1}z_{j_1})(z_{i_2}z_{j_2})\cdots(z_{i_t}z_{j_t})$  where  $i_1 \leq j_1 \leq i_2 \leq j_2 \leq \dots \leq i_t \leq j_t$ , and

$$d = (\deg_{z_1} g, \deg_{z_2} g, \dots, \deg_{z_n} g),$$

we denote  $g = g_{(z_{i_0}, d)}$ .

By Lemma 4.1 we have  $I \subseteq T_{\text{Sca}}(UJ_2)$ .

Let  $S$  be the subset in Lemma 4.5 and write  $\overline{S} = \{g + T_{\text{Sca}}(UJ_2) : g \in S\}$ . Since  $I \subseteq T_{\text{Sca}}(UJ_2)$  we have by Lemma 4.5 that  $J(X)/T_{\text{Sca}}(UJ_2) = \text{span}\overline{S}$ .

We shall prove that  $\overline{S}$  is a linearly independent set. Let

$$f(y_1, \dots, y_r, z_1, \dots, z_n) = \sum_{g \in S} \lambda_g g \in T_{\text{Sca}}(UJ_2), \lambda_g \in K.$$

Since  $K$  is an infinite field, every multihomogeneous component of  $f$  belongs to  $T_{\text{Sca}}(UJ_2)$ . Thus it is sufficient to suppose

$$f = \lambda y_1^{k_1} \cdots y_r^{k_r} (z_{i_1}z_{j_1})(z_{i_2}z_{j_2})\cdots(z_{i_t}z_{j_t}) \quad \text{or} \quad f = \sum_{i_0=1}^n \lambda_{i_0} y_1^{k_1} \cdots y_r^{k_r} g_{(z_{i_0}, d)},$$

where  $i_1 \leq j_1 \leq i_2 \leq j_2 \leq \dots \leq i_t \leq j_t$ ;  $t \geq 0$ . We have to prove that  $\lambda = \lambda_{i_0} = 0$  for all  $i_0$ . Denote

$$1 = e_{11} + e_{22}, \quad a = e_{11} - e_{22} \quad \text{and} \quad b = e_{12}.$$

In the first case, let  $Y_i = 1$  and  $Z_i = a$  for all  $i$ . Then

$$f(Y_1, \dots, Y_r, Z_1, \dots, Z_n) = \lambda 1 = 0$$

and so  $\lambda = 0$ .

In the second case, let  $Y_i = 1$  for all  $i$ ,  $Z_{i_0} = a + b$ ,  $Z_i = a$  for all  $i \neq i_0$ . Since  $ab = b^2 = 0$  we obtain

$$f(Y_1, \dots, Y_r, Z_1, \dots, Z_n) = (\lambda_1 + \dots + \lambda_n)a + \lambda_{i_0}b = 0$$

and so  $\lambda_{i_0} = 0$  as desired.

Therefore, the set  $\overline{S}$  is a basis for the quotient vector space  $J(X)/T_{\text{Sca}}(UJ_2)$ . Moreover, since  $I \subseteq T_{\text{Sca}}(UJ_2)$ , by Lemma 4.5 we have  $I = T_{\text{Sca}}(UJ_2)$ .  $\square$

**Remark 4.7.** There is a missing identity in the statement of [6, Proposition 8]. But that missing identity was used in the proof of [6, Proposition 8]. Here we give the complete list of these identities.

## 4.2 The scalar grading, when $K$ is a finite field

In this subsection we describe the  $\mathbb{Z}_2$ -graded identities for  $UJ_2$  with the scalar grading when  $K$  is finite. Throughout this subsection,  $K$  is a finite field with  $|K| = q$  elements and  $\text{char}(K) \neq 2$ .

**Lemma 4.8.** If  $u = \alpha a + \beta b$ , where  $\alpha, \beta \in K$ ,  $a = e_{11} - e_{22}$  and  $b = e_{12}$ , then

- i)  $u^{2n} = \alpha^{2n}1$ ,
- ii)  $u^{2n-1} = \alpha^{2n-1}a + \alpha^{2n-2}\beta b$ ,

for all  $n \in \mathbb{N}$ . In particular,  $u^q = \alpha a + \alpha^{q-1}\beta b$ .

*Proof.* By using induction on  $n$  we can prove i) and ii). Now, since  $p \neq 2$  we have  $p$  odd. Thus  $q$  is odd and

$$u^q = \alpha^q a + \alpha^{q-1}\beta b = \alpha a + \alpha^{q-1}\beta b.$$

The lemma is proved. □

**Lemma 4.9.** The polynomials

$$y_1^q - y_1 \text{ and } (z_1^q - z_1)z_2$$

belong to  $T_{\text{Sca}}(UJ_2)$ .

*Proof.* We will check the last polynomial only. Since  $ab = b^2 = 0$ , we have by Lemma 4.8 that  $(z_1^q - z_1)z_2 \in T_{\text{Sca}}(UJ_2)$ . □

**Notation 4.10.** Let  $I'$  be the  $T_{\mathbb{Z}_2}$ -ideal of  $J(X)$  generated by the polynomials in Lemmas 4.1 and 4.9.

**Lemma 4.11.** Let  $\widehat{S}$  be the subset of  $J(X)$  formed by all polynomials

(a)  $Y_1'Z_1'$  and

(b)  $Y_1'(z_{i_0}Z_1')$ ,

where  $Y_1' = y_1^{k_1} \dots y_r^{k_r}$ ;  $0 \leq k_i < q$  for all  $i$ ;  $r \geq 0$ ;  $Z_1' = (z_{i_1}z_{j_1})(z_{i_2}z_{j_2}) \dots (z_{i_t}z_{j_t})$ ;  $i_1 \leq j_1 \leq i_2 \leq j_2 \leq \dots \leq i_t \leq j_t$ ;  $t \geq 0$ ;  $0 \leq \deg_{z_k}(Z_1') < q$  for all  $k$ ;  $i_0 \geq 1$ . Then the quotient vector space  $J(X)/I'$  is spanned by the set of all elements  $g + I'$  where  $g \in \widehat{S}$ .

*Proof.* Let  $A$  and  $B$  be the sets of all elements  $g + I'$  where  $g$  is in (a) and (b), respectively. Denote  $C = A \cup B$ . If  $f(y_1, \dots, y_r, z_1, \dots, z_s)$  is a monomial in  $J(X)$ , we shall prove that  $f + I' \in \text{span}C$ .

Since  $I \subseteq I'$  we have by Lemma 4.5 that the quotient vector space  $J(X)/I'$  is spanned by the set of all polynomials:

(a')  $Y'Z' + I'$ ,

(b')  $Y'(z_{i_0}Z') + I'$ ,

where  $Y' = y_1^{k_1} \dots y_r^{k_r}$ ;  $k_i \geq 0$  for all  $i$ ;  $r \geq 0$ ;  $Z' = (z_{i_1}z_{j_1})(z_{i_2}z_{j_2}) \dots (z_{i_t}z_{j_t})$ ;  $i_1 \leq j_1 \leq i_2 \leq j_2 \leq \dots \leq i_t \leq j_t$ ;  $t \geq 0$ ;  $i_0 \geq 1$ .

Let  $A'$  and  $B'$  be the sets of all elements in (a') and (b'), respectively. We shall prove that  $A' \cup B' \subseteq \text{span}C$ .

*Case 1.*  $f + I' \in A'$ .

In this case,  $f + I' = (y_1^{k_1} \cdots y_r^{k_r})(z_{i_1} z_{j_1})(z_{i_2} z_{j_2}) \cdots (z_{i_t} z_{j_t}) + I'$ . Since  $(y_i^q - y_i) \in I'$  we can suppose  $0 \leq k_1, \dots, k_r < q$ .

Now,

$$\underbrace{(z_k z_k) \cdots (z_k z_k)}_{(q-1)/2 \text{ times}} (z_k z_l) = z_k^{q-1} (z_k z_l)$$

and  $z_k^{q-1} (z_k z_l) + I' = (z_k^{q-1} z_k) z_l + I' = (z_k^q) z_l + I' = z_k z_l + I'$  (see Lemma 4.9). Thus, we can suppose  $0 \leq \deg_{z_k} Z' < q$  for all  $k$  and so  $f + I' \in A \subseteq \text{span} C$ .

*Case 2.*  $f + I' \in B'$ .

We can use analogous argument to prove  $f + I' \in B \subseteq \text{span} C$ .

The lemma is proved.  $\square$

**Theorem 4.12.** If  $K$  is a finite field with  $|K| = q$  elements and  $\text{char}(K) \neq 2$  then  $I' = T_{\text{Sca}}(UJ_2)$ , that is,  $T_{\text{Sca}}(UJ_2)$  is generated, as a  $T_{\mathbb{Z}_2}$ -ideal, by the polynomials in Lemmas 4.1 and 4.9. Moreover, the set in Lemma 4.11 is a basis for the quotient vector space  $J(X)/I'$ .

*Proof.* By Lemmas 4.1 and 4.9 we have  $I' \subseteq T_{\text{Sca}}(UJ_2)$ .

Consider the subset  $\widehat{S}$  in Lemma 4.11 and write  $\overline{S} = \{g + T_{\text{Sca}}(UJ_2) : g \in \widehat{S}\}$ . Since  $I' \subseteq T_{\text{Sca}}(UJ_2)$  we have by Lemma 4.11 that  $J(X)/T_{\text{Sca}}(UJ_2) = \text{span} \overline{S}$ . We shall prove that  $\overline{S}$  is a linearly independent set.

If  $k = (k_1, \dots, k_r)$  we write  $y_1^{k_1} \cdots y_r^{k_r} = y^{[k]}$ . If  $g = (z_{i_1} z_{j_1}) \cdots (z_{i_t} z_{j_t})$ ,  $\deg_{z_i} g = d_i$  and  $d = (d_1, \dots, d_n)$  then we write  $g = z^{[d]}$ . Let

$$f(y_1, \dots, y_r, z_1, \dots, z_n) = \sum_{g \in \overline{S}} \lambda_g g \in T_{\text{Sca}}(UJ_2),$$

where  $\lambda_g \in K$ . We will prove that  $\lambda_g = 0$  for all  $g$ . Since  $\deg_{y_i} f < q$  for all  $i$  we can suppose

$$f = \lambda y^{[k]} + \sum_d \lambda_d y^{[k]} z^{[d]} + \sum_{i=1}^n \sum_d \lambda_{(i,d)} y^{[k]} (z_i z^{[d]}).$$

By replacing  $y_i$  and  $z_i$  by the matrices 1 and 0 respectively, for all  $i$ , we obtain  $0 = f(1, \dots, 1, 0, \dots, 0) = \lambda \cdot 1$  and so  $\lambda = 0$ . Now, since  $T_{\text{Sca}}(UJ_2)$  is a  $\mathbb{Z}_2$ -graded ideal of  $J(X)$  it follows that  $f_1, f_2 \in T_{\text{Sca}}(UJ_2)$  where

$$f_1 = \sum_d \lambda_d y^{[k]} z^{[d]} \quad \text{and} \quad f_2 = \sum_{i=1}^n \sum_d \lambda_{(i,d)} y^{[k]} (z_i z^{[d]}).$$

Since  $\deg_{z_i} f_1 < q$  for all  $i$  it follows that every  $g_d = \lambda_d y^{[k]} z^{[d]} \in T_{\text{Sca}}(UJ_2)$ . By replacing  $y_i$  and  $z_i$  by the matrices 1 and  $a = e_{11} - e_{22}$  respectively, for all  $i$ , we obtain  $0 = g_d(1, \dots, 1, a, \dots, a) = \lambda_d \cdot 1$  and so  $\lambda_d = 0$ .

Denote  $a = e_{11} - e_{22}$ ,  $b = e_{12}$  and  $Z_i = \alpha_i a + \beta_i b$ , where  $\alpha_i, \beta_i \in K$ . We have

$$f_2(1, \dots, 1, Z_1, \dots, Z_n) = \alpha a + \beta b = 0, \quad \text{where} \quad \beta = \sum_{i=1}^n \sum_d \lambda_{(i,d)} \beta_i \alpha_1^{d_1} \cdots \alpha_n^{d_n}$$

and  $\alpha \in K$ . Since  $\alpha_i, \beta_i$  are any elements of  $K$  for all  $i$ ,  $\beta = 0$ ,  $\deg_{\alpha_i} \beta < q$  and  $\deg_{\beta_i} \beta < q$  it follows that  $\lambda_{(i,d)} = 0$  for all  $i, d$ . See Lemma 2.3.

Therefore, the set  $\bar{S}$  is a basis for the quotient vector space  $J(X)/T_{\text{Sca}}(UJ_2)$ . Moreover, since  $I' \subseteq T_{\text{Sca}}(UJ_2)$ , by Lemma 4.11 we have  $I' = T_{\text{Sca}}(UJ_2)$  and the theorem is proved.  $\square$

## 5 The classical grading

Let  $T_{\text{Cla}}(UJ_2)$  be the set of all  $\mathbb{Z}_2$ -graded polynomial identities for  $UJ_2$  with the classical grading. In this section we will describe  $T_{\text{Cla}}(UJ_2)$  for any field  $K$  of  $\text{char}(K) = p \neq 2$ .

We remember that

$$UJ_2 = (UJ_2)_0 \oplus (UJ_2)_1,$$

where

$$(UJ_2)_0 = \text{span}\{e_{11} + e_{22}, e_{11} - e_{22}\} \text{ and } (UJ_2)_1 = \text{span}\{e_{12}\}.$$

**Lemma 5.1.** The polynomials

$$(y_1, y_2, y_3), \quad z_1 z_2 \text{ and } (y_1, z_1, y_2)$$

belong to  $T_{\text{Cla}}(UJ_2)$ .

*Proof.* The proof is a direct verification.  $\square$

**Notation 5.2.** Let  $I$  be the  $T_{\mathbb{Z}_2}$ -ideal of  $J(X)$  generated by the polynomials in Lemma 5.1.

Define the equivalence relation  $\equiv$  on  $J(X)$  as follows: if  $f, g \in J(X)$  then

$$f \equiv g \Leftrightarrow f + I = g + I.$$

**Lemma 5.3.** The polynomial

$$y_1(y_2(y_3 z_1)) - \frac{1}{2}(y_1(z_1(y_2 y_3)) + y_2(z_1(y_1 y_3)) + y_3(z_1(y_1 y_2)) - z_1(y_1(y_2 y_3)))$$

belongs to  $I$ .

*Proof.* Let

$$f = 2y_1(y_2(y_3 z_1)) - y_1(z_1(y_2 y_3)) - y_2(z_1(y_1 y_3)) - y_3(z_1(y_1 y_2)) + z_1(y_1(y_2 y_3)).$$

If  $u = y_2$ ,  $v = y_3$ ,  $c = z_1$  and  $d = y_1$  in (3) we obtain

$$\begin{aligned} f &= 2((y_3 z_1) y_2) y_1 - ((z_1 y_2) y_1) y_3 - ((z_1 y_3) y_1) y_2 \\ &= (y_3, z_1, y_2) y_1 + (y_2, z_1 y_3, y_1) + (y_3, z_1 y_2, y_1). \end{aligned}$$

Since  $(y_1, z_1, y_2) \in I$  it follows that  $f \in I$  and so

$$y_1(y_2(y_3 z_1)) - \frac{1}{2}(y_1(z_1(y_2 y_3)) + y_2(z_1(y_1 y_3)) + y_3(z_1(y_1 y_2)) - z_1(y_1(y_2 y_3))) = \frac{1}{2} f \in I.$$

$\square$

**Lemma 5.4.** The subalgebra of  $J(X)/I$  generated by the set

$$Y + I = \{y + I : y \in Y\}$$

is commutative and associative.

*Proof.* Let  $A_Y$  be the subalgebra of  $J(X)/I$  generated by the set  $Y + I$ . The algebra  $J(X)$  is commutative, thus  $A_Y$  is commutative too. Since  $(y_1, y_2, y_3) \in I$  we have that  $A_Y$  is associative, and the lemma is proved.  $\square$

**Notation 5.5.** If  $Y' = y_{i_1} \cdots y_{i_r}$  and  $Y'' = y_{j_1} \cdots y_{j_s}$  are monomials in  $J(Y)$  such that  $r, s \geq 0$ ,  $i_1 \leq i_2 \leq \dots \leq i_r$  and  $j_1 \leq j_2 \leq \dots \leq j_s$ , we will write  $Y' < Y''$  if

- i)  $r < s$  or
- ii)  $r = s$  and  $i_1 = j_1, i_2 = j_2, \dots, i_l = j_l, i_{l+1} < j_{l+1}$  for some  $l$ .

**Lemma 5.6.** Let  $S$  be the subset of  $J(X)$  formed by all polynomials

- (a)  $Y'$  and
- (b)  $Y'(z_i Y'')$ ,

where  $Y' = y_{i_1} \cdots y_{i_r}$ ;  $r \geq 0$ ;  $i_1 \leq \dots \leq i_r$ ;  $Y'' = y_{j_1} \cdots y_{j_s}$ ;  $s \geq 0$ ;  $j_1 \leq \dots \leq j_s$ ;  $Y' \leq Y''$ . Then the quotient vector space  $J(X)/I$  is spanned by the set of all elements  $g + I$  where  $g \in S$ .

*Proof.* Let  $A$  and  $B$  be the sets of all elements  $g + I$  where  $g$  is in (a) and (b), respectively. Denote  $C = A \cup B$ . If  $f$  is a monomial in  $J(X)$ , we shall prove by induction on  $\deg(f)$  that  $f + I \in \text{span}C$ .

The cases  $\deg(f) = 1$  and  $\deg(f) = 2$  are trivial.

Suppose  $\deg(f) \geq 3$  and write  $f = gh$  where  $g, h \in J(X)$  are monomials with  $\deg g < \deg f$ . By induction hypothesis it follows that  $g + I$  and  $h + I$  belong to  $C$ . Since  $z_1 z_2 \in I$  it is sufficient to consider two cases:

1.  $g + I$  and  $h + I$  belong to  $A$ .

In this case,  $g + I = Y' + I$ ,  $h + I = Y'_1 + I$  and  $f \equiv Y'Y'_1$ . By Lemma 5.4 it follows that  $f + I \in A \subset \text{span}C$ .

2.  $g + I$  belongs to  $A$  and  $h + I$  belongs to  $B$ .

In this case,  $g + I = Y' + I$  and  $h + I = Y'_1(z_i Y'_2) + I$ . By Lemma 5.3 we obtain

$$\begin{aligned} f &\equiv Y'(Y'_1(z_i Y'_2)) \\ &\equiv \frac{1}{2} (Y'(z_i(Y'_1 Y'_2)) + Y'_1(z_i(Y' Y'_2)) + Y'_2(z_i(Y' Y'_1)) - z_i(Y'(Y'_1 Y'_2))). \end{aligned} \tag{12}$$

We will show that  $f + I \in \text{span}B$ . Firstly, we use Lemma 5.4 to order the variables in  $Y'_1 Y'_2$ ,  $Y' Y'_2$ ,  $Y' Y'_1$ ,  $Y'(Y'_1 Y'_2)$  appearing in (12). Now, if  $Y_3 > Y_4$  are monomials in  $J(Y)$  then by  $(y_1, z_1, y_2) \in I$  we have  $Y_3(z_i Y_4) \equiv Y_4(z_i Y_3)$ . If necessary we can use this to prove that the summands in (12) are in  $B$ . Thus  $f + I \in \text{span}B \subset \text{span}C$ .



The lemma is proved.  $\square$

## 5.1 The classical grading, when $K$ is an infinite field

In this subsection we describe the  $\mathbb{Z}_2$ -graded identities for  $UJ_2$  with the classical grading when  $K$  is infinite.

**Theorem 5.7.** If  $K$  is an infinite field of  $\text{char}(K) \neq 2$  then  $I = T_{\text{Cla}}(UJ_2)$ , that is,  $T_{\text{Cla}}(UJ_2)$  is generated, as a  $T_{\mathbb{Z}_2}$ -ideal, by the polynomials in Lemma 5.1. Moreover, the set in Lemma 5.6 is a basis for the quotient vector space  $J(X)/I$ .

*Proof.* By Lemma 5.1 we have  $I \subseteq T_{\text{Cla}}(UJ_2)$ .

Let  $S$  be the set in Lemma 5.6. Write  $\bar{S} = \{g + T_{\text{Cla}}(UJ_2) : g \in S\}$ . Since  $I \subseteq T_{\text{Cla}}(UJ_2)$  we have by Lemma 5.6 that  $J(X)/T_{\text{Cla}}(UJ_2) = \text{span}\bar{S}$ .

We shall prove that  $\bar{S}$  is a linearly independent set. Let

$$f(y_1, \dots, y_r, z_1, \dots, z_n) = \sum_{g \in S} \lambda_g g \in T_{\text{Cla}}(UJ_2), \quad \lambda_g \in K.$$

Since  $K$  is an infinite field, every multihomogeneous component of  $f$  belongs to  $T_{\text{Cla}}(UJ_2)$ .

Thus it is sufficient to suppose

$$f(y_1, \dots, y_r) = \lambda y_1^{k_1} \dots y_r^{k_r} \text{ or } f(y_1, \dots, y_r, z_j) = \sum_l \lambda_l y_1^{k_1-l_1} \dots y_r^{k_r-l_r} (z_j (y_1^{l_1} \dots y_r^{l_r}))$$

where  $l = (l_1, \dots, l_r)$ ,  $0 \leq l_i \leq k_i$  for all  $i$ , and  $y_1^{k_1-l_1} \dots y_r^{k_r-l_r} \leq y_1^{l_1} \dots y_r^{l_r}$ . We shall prove that  $\lambda = \lambda_l = 0$  for all  $l$ .

In the first case,  $f(1, \dots, 1) = \lambda \cdot 1 = 0$  and so  $\lambda = 0$ .

In the second case, let  $Y_i = \alpha_i e_{11} + \beta_i e_{22}$  for all  $i$  and let  $Z_j = e_{12}$ , where  $\alpha_i, \beta_i \in K$ . Remember that  $(UJ_2)_0 = \text{span}\{e_{11}, e_{22}\}$ . We have

$$f(Y_1, \dots, Y_r, Z_j) = u e_{12} = 0 \text{ where}$$

$$u = (1/4) \sum_l \lambda_l (\alpha_1^{k_1} \dots \alpha_r^{k_r} + \alpha_1^{k_1-l_1} \dots \alpha_r^{k_r-l_r} \beta_1^{l_1} \dots \beta_r^{l_r} + \alpha_1^{l_1} \dots \alpha_r^{l_r} \beta_1^{k_1-l_1} \dots \beta_r^{k_r-l_r} + \beta_1^{k_1} \dots \beta_r^{k_r}).$$

Since  $y_1^{k_1-l_1} \dots y_r^{k_r-l_r} \leq y_1^{l_1} \dots y_r^{l_r}$ , the coefficient of  $\alpha_1^{k_1-l_1} \dots \alpha_r^{k_r-l_r} \beta_1^{l_1} \dots \beta_r^{l_r}$  in  $u$  is  $(1/4)\lambda_l$  if  $l = (l_1, \dots, l_r) \neq k = (k_1, \dots, k_r)$ . Since  $K$  is infinite and  $u = 0$  for all  $\alpha_i, \beta_i \in K$  it follows that  $\lambda_l = 0$  for all  $l \neq k$ . Now

$$u = (1/2)\lambda_k (\alpha_1^{k_1} \dots \alpha_r^{k_r} + \beta_1^{k_1} \dots \beta_r^{k_r})$$

and with analogous argument we have  $\lambda_k = 0$  too.

Therefore, the set  $\bar{S}$  is a basis for the quotient vector space  $J(X)/T_{\text{Cla}}(UJ_2)$ . Moreover, since  $I \subseteq T_{\text{Cla}}(UJ_2)$ , by Lemma 5.6 we have  $I = T_{\text{Cla}}(UJ_2)$ .  $\square$

## 5.2 The classical grading, when $K$ is a finite field

Throughout this subsection,  $K$  is a finite field with  $|K| = q$  elements and  $\text{char}(K) \neq 2$ .

**Lemma 5.8.** The polynomial  $y_1^q - y_1$  belongs to  $T_{\text{Cla}}(UJ_2)$ .

*Proof.* In fact, given  $Y_1 \in (UJ_2)_0$  we have  $Y_1 = \alpha e_{11} + \beta e_{22}$  for some  $\alpha, \beta \in K$ . Since  $|K| = q$  we obtain  $Y_1^q = \alpha^q e_{11} + \beta^q e_{22} = \alpha e_{11} + \beta e_{22} = Y_1$  as desired.  $\square$

**Notation 5.9.** Let  $I'$  be the  $T_{\mathbb{Z}_2}$ -ideal of  $J(X)$  generated by the polynomials in Lemmas 5.1 and 5.8.

**Lemma 5.10.** Let  $\widehat{S}$  be the subset of  $J(X)$  formed by all polynomials

- (a)  $Y'_1$  and
- (b)  $Y'_1(z_i Y'_2)$ ,

where  $Y'_1 = y_1^{k_1} \cdots y_r^{k_r}$ ;  $0 \leq k_1, \dots, k_r < q$ ;  $Y'_2 = y_1^{l_1} \cdots y_r^{l_r}$ ;  $0 \leq l_1, \dots, l_r < q$ ;  $Y'_1 \leq Y'_2$ ;  $r \geq 1$ ;  $z_i \in Z$ . Then the quotient vector space  $J(X)/I'$  is spanned by the set of all elements  $g + I'$  where  $g \in \widehat{S}$ .

*Proof.* Let  $A$  and  $B$  be the sets of all elements  $g + I'$  where  $g$  is in (a) and (b), respectively. Denote  $C = A \cup B$ . Let  $f$  be a monomial in  $J(X)$ , we shall prove that  $f + I' \in \text{span}C$ .

Since  $I \subseteq I'$  we have by Lemma 5.6 that the quotient vector space  $J(X)/I'$  is spanned by the set of all polynomials:

- (a')  $Y' + I'$ ,
- (b')  $Y'(z_i Y'') + I'$ ,

where  $Y' = y_1^{k_1} \cdots y_r^{k_r}$ ;  $0 \leq k_1, \dots, k_r < q$ ;  $Y'' = y_1^{l_1} \cdots y_r^{l_r}$ ;  $0 \leq l_1, \dots, l_r < q$ ;  $Y' \leq Y''$ ;  $r \geq 0$ ;  $z_i \in Z$ . Let  $A'$  and  $B'$  be the sets of all elements in (a') and (b'), respectively. We shall prove that  $A' \cup B' \subseteq \text{span}C$ .

*Case 1.*  $f + I' \in A'$ .

In this case,  $f + I' = y_1^{k_1} \cdots y_r^{k_r} + I'$ . Since  $y_i^q - y_i \in I'$  we can suppose  $0 \leq k_1, \dots, k_r < q$ . Thus,  $f + I' \in A \subset \text{span}C$ .

*Case 2.*  $f + I' \in B'$ .

In this case,  $f + I' = (y_1^{k_1} \cdots y_r^{k_r})(z_i(y_1^{l_1} \cdots y_r^{l_r})) + I'$ . As in Case 1, we can suppose  $0 \leq k_1, \dots, k_r < q$  and  $0 \leq l_1, \dots, l_r < q$ . Since  $(y_1, z_i, y_2) \in I'$  we obtain

$$(y_1^{k_1} \cdots y_r^{k_r})(z_i(y_1^{l_1} \cdots y_r^{l_r})) + I' = (y_1^{l_1} \cdots y_r^{l_r})(z_i(y_1^{k_1} \cdots y_r^{k_r})) + I'.$$

Thus, we can suppose  $y_1^{k_1} \cdots y_r^{k_r} \leq y_1^{l_1} \cdots y_r^{l_r}$ , and consequently  $f + I' \in B \subset \text{span}C$  as desired. The lemma is proved.  $\square$

**Theorem 5.11.** If  $K$  is a finite field with  $|K| = q$  elements and  $\text{char}(K) \neq 2$  then  $I' = T_{\text{Cla}}(UJ_2)$ , that is,  $T_{\text{Cla}}(UJ_2)$  is generated, as a  $T_{\mathbb{Z}_2}$ -ideal, by the polynomials in Lemmas 5.1 and 5.8. Moreover, the set in Lemma 5.10 is a basis for the quotient vector space  $J(X)/I'$ .

*Proof.* By Lemmas 5.1 and 5.8 we have  $I' \subseteq T_{\text{Cla}}(UJ_2)$ .

Consider the set  $\widehat{S}$  in Lemma 5.10 and write  $\overline{S} = \{g + T_{\text{Cla}}(UJ_2) : g \in \widehat{S}\}$ . Since  $I' \subseteq T_{\text{Cla}}(UJ_2)$ , by Lemma 5.10 it follows that  $J(X)/T_{\text{Cla}}(UJ_2) = \text{span}\overline{S}$ .

We shall prove that  $\overline{S}$  is a linearly independent set. Let

$$f(y_1, \dots, y_r, z_1, \dots, z_n) = \sum_{g \in \overline{S}} \lambda_g g \in T_{\text{Cla}}(UJ_2), \quad \lambda_g \in K.$$

In particular,

$$h = f(y_1, \dots, y_r, 0, \dots, 0) = \sum_k \lambda_k y_1^{k_1} \dots y_r^{k_r} \in T_{\text{Cla}}(UJ_2),$$

where  $k = (k_1, \dots, k_r)$ ,  $0 \leq k_i < q$  for all  $i$ . Since  $|K| = q$  and  $\deg_{y_i} h < q$  for all  $i$ , we have

$$h_k(y_1, \dots, y_r) = \lambda_k y_1^{k_1} \dots y_r^{k_r} \in T_{\text{Cla}}(UJ_2)$$

for all  $k$ . Thus,  $h_k(1, \dots, 1) = \lambda_k \cdot 1 = 0$  and so  $\lambda_k = 0$ .

Now, we have

$$f(y_1, \dots, y_r, z_1, \dots, z_n) = \sum_{i=1}^n \sum_{(l,m)} \lambda_{(l,m)} y_1^{l_1} \dots y_r^{l_r} (z_i(y_1^{m_1} \dots y_r^{m_r})) \in T_{\text{Cla}}(UJ_2),$$

where  $l = (l_1, \dots, l_r)$ ,  $0 \leq l_j < q$  for all  $j$ ,  $m = (m_1, \dots, m_r)$ ,  $0 \leq m_j < q$  for all  $j$ , and  $y_1^{l_1} \dots y_r^{l_r} \leq y_1^{m_1} \dots y_r^{m_r}$ . Since  $f(y_1, \dots, y_r, 0, \dots, 0, z_i, 0, \dots, 0) \in T_{\text{Cla}}(UJ_2)$  we can suppose

$$f = f(y_1, \dots, y_r, z_i) = \sum_{(l,m)} \lambda_{(l,m)} y_1^{l_1} \dots y_r^{l_r} (z_i(y_1^{m_1} \dots y_r^{m_r})) \in T_{\text{Cla}}(UJ_2).$$

Let  $Y_j = \alpha_j e_{11} + \beta_j e_{22}$  for all  $j$ , and  $Z_i = e_{12}$ , where  $\alpha_j, \beta_j \in K$ . We have  $f(Y_1, \dots, Y_r, Z_i) = u e_{12} = 0$  where

$$u = (1/4) \sum_{(l,m)} \lambda_{(l,m)} (\alpha_1^{l_1+m_1} \dots \alpha_r^{l_r+m_r} + \alpha_1^{l_1} \dots \alpha_r^{l_r} \beta_1^{m_1} \dots \beta_r^{m_r} + \alpha_1^{m_1} \dots \alpha_r^{m_r} \beta_1^{l_1} \dots \beta_r^{l_r} + \beta_1^{l_1+m_1} \dots \beta_r^{l_r+m_r}).$$

Since  $\alpha_i^q = \alpha_i$  and  $\beta_i^q = \beta_i$  we can write

$$u = \sum_{(l,m)} \eta_{(l,m)} \alpha_1^{l_1} \dots \alpha_r^{l_r} \beta_1^{m_1} \dots \beta_r^{m_r} = 0$$

where  $l = (l_1, \dots, l_r)$ ,  $0 \leq l_i < q$  for all  $i$ ,  $m = (m_1, \dots, m_r)$ ,  $0 \leq m_i < q$  for all  $i$ ,  $\eta_{(l,m)} \in K$ . In particular,  $\eta_{(l,m)} = 0$  for all  $(l,m)$ . Now, if  $l \neq (0, \dots, 0)$  and  $y_1^{l_1} \dots y_r^{l_r} \leq y_1^{m_1} \dots y_r^{m_r}$  we have  $(1/4)\lambda_{(l,m)} = \eta_{(l,m)} = 0$  and so  $\lambda_{(l,m)} = 0$ . In particular,

$$u = (1/2) \sum_{(l,m), l=(0,\dots,0)} \lambda_{(l,m)} (\alpha_1^{m_1} \dots \alpha_r^{m_r} + \beta_1^{m_1} \dots \beta_r^{m_r}) = 0.$$

Since  $0 \leq m_i < q$  for all  $i$  we obtain  $\lambda_{(l,m)} = 0$  if  $l = (0, \dots, 0)$ .

Therefore, the set  $\overline{S}$  is a basis for the quotient vector space  $J(X)/T_{\text{Cla}}(UJ_2)$ . Moreover, since  $I' \subseteq T_{\text{Cla}}(UJ_2)$ , by Lemma 5.10 we have  $I' = T_{\text{Cla}}(UJ_2)$ .  $\square$

## Funding

Mateus Eduardo Salomão was supported by Ph.D. grant from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES). Dimas José Gonçalves was partially supported by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) grant No. 2018/23690-6, and by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) grant No. 406401/2016-0.

## References

- [1] L. Centrone, F. Martino, M. S. Souza. Specht property for some varieties of Jordan algebras of almost polynomial growth. *J. Algebra* 521 (2019) 137–165.
- [2] O. M. Di Vincenzo, P. Koshlukov, A. Valenti. Gradings on the algebra of upper triangular matrices and their graded identities. *J. Algebra* 275 (2004), no. 2, 550–566.
- [3] V. Drensky. Free algebras and PI-algebras. Graduate course in algebra. Springer-Verlag Singapore, Singapore, 2000. xii+271 pp. ISBN: 981-4021-48-2.
- [4] D. J. Gonçalves, E. Riva. Graded polynomial identities for the upper triangular matrix algebra over a finite field. *J. Algebra* 559 (2020), 625–645.
- [5] N. Jacobson. Structure and representations of Jordan algebras, in: Amer. Math. Soc. Colloq. Publ., vol. 39, Amer. Math. Soc, Providence, RI, 1968.
- [6] P. Koshlukov, F. Martino. Polynomial identities for the Jordan algebra of upper triangular matrices of order 2. *J. Pure Appl. Algebra* 216 (2012), no. 11, 2524–2532.
- [7] P. Koshlukov, F. Y. Yasumura. Group gradings on the Jordan algebra of upper triangular matrices. *Linear Algebra Appl.* 534 (2017), 1–12.
- [8] P. Koshlukov, F. Yukihide. Group gradings on the Lie algebra of upper triangular matrices. *J. Algebra* 477 (2017), 294–311.
- [9] Y. N. Maltsev. A basis for the identities of the algebra of upper triangular matrices, *Algebra Logika* 10 (1971) 393–400 (in Russian). Translation: *Algebra Logic* 10 (1971) 242–247.

- [10] P.N. Siderov. A basis for identities of an algebra of triangular matrices over an arbitrary field. *Pliska Stud. Math. Bulgar.* 2 (1981) 143–152 (in Russian).
- [11] R. I. Q. Urure, D. J. Gonçalves. Identities with involution for  $2 \times 2$  upper triangular matrices algebra over a finite field. *Linear Algebra Appl.* 544 (2018), 223–253.
- [12] A. Valenti, M. V. Zaicev. Group gradings on upper triangular matrices. *Arch. Math.* 89 (1) (2007), 33-40.