

APPROXIMATION AND LOCALIZED POLYNOMIAL FRAME ON CONIC DOMAINS

YUAN XU

ABSTRACT. Highly localized kernels constructed by orthogonal polynomials have been fundamental in the recent development of approximation and computational analysis on the unit sphere, unit ball, and several other regular domains. In this work, we first study homogeneous spaces that are assumed to contain highly localized kernels and establish a framework for approximation and localized tight frames in such spaces, which extends recent works on bounded regular domains. We then show that the framework is applicable to homogeneous spaces defined on bounded conic domains, which consists of conic surfaces and the solid domains bounded by such surfaces and hyperplanes. The highly localized kernels on conic domains require precise estimates that rely on recently discovered addition formulas for orthogonal polynomials with respect to special weight functions on each domain and an intrinsic distance that takes into account the boundary of the domain, the latter is not comparable to the Euclidean distance at around the apex of the cone.

The main results provide construction of semi-discrete localized tight frame in weighted L^2 norm and characterization of best approximation by polynomials on conic domains. The latter is achieved by using a K -functional, defined via the differential operator that has orthogonal polynomials as eigenfunctions, as well as a modulus of smoothness defined via a multiplier operator that is equivalent to the K -functional. Several intermediate results are of interest in their own right, including the Marcinkiewicz-Zygmund inequalities, positive cubature rules, Christoffel functions, and Bernstein type inequalities. Moreover, although the highly localizable kernels hold only for special families of weight functions on each domain, many intermediate results are shown to hold for doubling weights defined via the intrinsic distance on the domain.

CONTENTS

1. Introduction	2
1.1. Background	2
1.2. New contributions	4
1.3. Organization and convention	7
2. Homogeneous spaces with highly localized kernels	7
2.1. Localizable homogeneous spaces	8
2.2. Examples of localizable homogeneous spaces	12
2.3. An auxiliary maximal function	15
2.4. Marcinkiewicz-Zygmund inequality	17
2.5. Christoffel function for doubling weight	19

Date: December 12, 2021.

2010 Mathematics Subject Classification. 41A10, 41A63, 42C10, 42C40.

Key words and phrases. Approximation, conic domain, Fourier orthogonal series, homogeneous spaces, localized kernel, orthogonal polynomials, polynomial frame.

The author is partially supported by Simons Foundation Grant #849676.

2.6. Positive cubature rules	22
2.7. Tight polynomial frames	23
3. Polynomial Approximation on homogeneous spaces	25
3.1. Addition formula and differential operator	26
3.2. Convolution structure	27
3.3. Characterization of best approximation	30
3.4. Near best approximation operator	31
3.5. Bernstein and Nikolskii inequalities	32
3.6. Proof of the main results	35
4. Homogeneous space on conic surfaces	37
4.1. Distance on the surface of the cone	37
4.2. A family of doubling weights	40
4.3. Orthogonal polynomials on the conic surface	42
4.4. Highly localized kernels	44
4.5. Maximal ε -separated sets and MZ inequality	50
4.6. Cubature rules and localized tight frames	52
4.7. Characterization of best approximation	54
5. Homogeneous space on solid cones	56
5.1. Distance on solid cone	56
5.2. A family of doubling weights	58
5.3. Orthogonal polynomials on the solid cone	59
5.4. Highly localized kernels	60
5.5. Maximal ε -separated sets and MZ inequality	66
5.6. Cubature rules and localized tight frames	68
5.7. Characterization of best approximation	70
References	71

1. INTRODUCTION

In recent years highly localized kernels constructed via orthogonal polynomials have become important tools in approximation theory, both computational and theoretical harmonic analysis, and functional analysis. We start with an introduction on the background of this development in the first subsection, describe our main results in the second subsection and state the organization of the work in the third subsection.

1.1. Background. Let Ω be a set in \mathbb{R}^d , either an algebraic surface or a domain with non-empty interior, and let ϖ be a nonnegative weight function defined on Ω , normalized with unit integral with respect to the Lebesgue measure dx on Ω , so that the bilinear form

$$\langle f, g \rangle_{\varpi} = \int_{\Omega} f(x)g(x)\varpi(x)dx$$

is a well defined inner product on the space of polynomials restricted to Ω . Let $\mathcal{V}_n^d(\varpi)$ be the space of orthogonal polynomials of degree n with respect to this inner product. Assume that ϖ is regular so that the orthogonal decomposition

$$L^2(\Omega, \varpi) = \bigoplus_{n=0}^{\infty} \mathcal{V}_n^d(\varpi)$$

holds. Let $P_n(\varpi; \cdot, \cdot)$ be the reproducing kernel of the space $\mathcal{V}_n^d(\varpi)$. The projection operator $\text{proj}_n : L^2(\Omega, \varpi) \mapsto \mathcal{V}_n^d(\varpi)$ can be written as

$$(\text{proj}_n f)(x) = \int_{\Omega} P_n(\varpi; x, y) f(y) \varpi(y) dy, \quad f \in L^2(\Omega, \varpi).$$

If \hat{a} is a cut-off function, defined as a compactly supported function in $C^\infty(\mathbb{R}_+)$ and $\hat{a}(t) = 1$ for t near zero, then our highly localized kernels are of the form

$$L_n(\varpi; x, y) := \sum_{j=0}^{\infty} \hat{a}\left(\frac{j}{n}\right) P_j(\varpi; x, y).$$

For orthogonal systems on several regular domains, the kernel $L_n(\varpi; x, y)$ for a suitable \hat{a} satisfies a *localization principle*, meaning that the kernel decays at rates faster than any inverse polynomial rate away from the main diagonal $y = x$ in $\Omega \times \Omega$ with respect to an intrinsic distance in Ω .

To get a sense of the highly localized estimate, we specify to two cases. The first one is the unit sphere \mathbb{S}^{d-1} of \mathbb{R}^d with $\varpi(x)dx = d\sigma$ is the surface measure, for which the kernel $L_n(\cdot, \cdot) = L_n(\varpi; \cdot, \cdot)$ is highly localized in the sense that [34], for every $\kappa > 0$,

$$(1.1) \quad |L_n(x, y)| \leq c_\kappa \frac{n^{d-1}}{(1 + n \, \text{d}_{\mathbb{S}}(x, y))^\kappa}, \quad x, y \in \mathbb{S}^{d-1},$$

where $\text{d}_{\mathbb{S}}$ denotes the geodesic distance

$$(1.2) \quad \text{d}_{\mathbb{S}}(x, y) = \arccos \langle x, y \rangle, \quad x, y \in \mathbb{S}^{d-1}.$$

For the unit ball, the classical weight function is $\varpi_\mu(x) = (1 - \|x\|^2)^{\mu - \frac{1}{2}}$, $\mu > -\frac{1}{2}$. For $\mu \geq 0$, the kernel $L_n(\varpi_\mu; \cdot, \cdot)$ is highly localized in the sense that [38]

$$(1.3) \quad |L_n(\varpi_\mu; x, y)| \leq c_\kappa \frac{n^d}{\sqrt{\varpi_\mu(n; x)} \sqrt{\varpi_\mu(n; y)} (1 + n \, \text{d}_{\mathbb{B}}(x, y))^\kappa}, \quad x, y \in \mathbb{B}^d,$$

where $\varpi_\mu(n; x) = (1 - \|x\|^2 + n^{-2})^\mu$ and $\text{d}_{\mathbb{B}}$ is the distance on \mathbb{B}^d defined by

$$(1.4) \quad \text{d}_{\mathbb{B}}(x, y) = \arccos \left(\langle x, y \rangle + \sqrt{1 - \|x\|^2} \sqrt{1 - \|y\|^2} \right).$$

By choosing κ large, the kernel decays away from $y = x$ by (1.1) and (1.3) faster than any polynomial rate. Furthermore, under a technical restriction on the cut-off function, both estimates can be improved to sub-exponential rate [28]. The proof of these estimates relies on closed-form formula of the reproducing kernel $P_n(\varpi; \cdot, \cdot)$ on the unit sphere and the unit ball; see the next subsection.

Let us denote the integral operator with $L_n(\varpi)$ as its kernel by $L_n * f$; that is,

$$(1.5) \quad L_n * f(x) := \int_{\Omega} f(y) L_n(\varpi; x, y) \varpi(y) dy.$$

If \hat{a} has the support $[0, 2]$ and satisfies $\hat{a}(t) = 1$ for $0 \leq t \leq 1$, then the operator $L_n * f$ is a polynomial of degree $2n$ and it reproduces polynomials of degree n . Furthermore, in many cases, it provides the near best polynomial approximation to $f \in L^p(\Omega, \varpi)$ in the sense that

$$\|L_n * f - f\|_{L^p(\Omega, \varpi)} \leq c \inf_{\deg g \leq n} \|f - g\|_{L^p(\Omega, \varpi)}, \quad 1 \leq p \leq \infty.$$

The rapid decay of the kernel makes it a powerful tool for studying polynomial approximation on Ω . Moreover, the operator $L_n * f$ plays an important role in the characterization of best approximation, which is a central problem in approximation

theory. For the unit sphere, for example, it is used in the classical result that characterizes the error of best polynomial approximation by a modulus of smoothness defined via spherical means or by its equivalent K -functional defined via the Laplace-Beltrami operator on the sphere. The theory has a long history that spans from [4, 36] to [42] with contributions from many researchers in between. It has also been extended, more recently, to weight approximation on the sphere, the unit ball, and the simplex in [52].

The highly localized kernels can also be used to construct localized frames via a semi-continuous Calderón type decomposition based on the decomposition,

$$f = \sum_{j=0}^{\infty} L_{2^j} * L_{2^j} * f, \quad f \in L^2(\Omega, \varpi),$$

which holds if the cut-off function \hat{a} satisfies $\sum_{k=0}^{\infty} |\hat{a}(2^{-k}t)|^2 = 1$. To define the frame elements, the integrals in the right-hand side of the above expansion need to be appropriately discretized by a cubature rule, established with the help of the highly localized kernels, which states that

$$\int_{\Omega} f(x) dx = \sum_{\xi \in \Xi_j} \lambda_{\xi} f(\xi), \quad f \in \Pi_{2^j}(\Omega),$$

where $\lambda_{\xi} > 0$, Ξ_j is a finite subset in Ω , and $\Pi_n(\Omega)$ denotes the space of polynomials, restricted on Ω , of degree at most n . The discretization leads to the tight frame $\{\psi_{\xi}\}_{\xi \in \Xi}$ of $L^2(\Omega, \varpi)$, where Ξ is a discrete set of well separated, with respect to the intrinsic distance, points in Ω , and the frame elements ψ_{ξ} , also called *needlets*, are of the form

$$\psi_{\xi}(x) = \sqrt{\lambda_{\xi}} L_{2^j}(x, \xi), \quad \xi \in \Xi,$$

which is highly localized with its center at ξ since λ_{ξ} can be quantized and $L_{2^j}(x, \xi)$ is highly localized. The tight frame means that, for all $f \in L^2(\Omega, \varpi)$,

$$f = \sum_{\xi \in \Xi} \langle f, \psi_{\xi} \rangle_{\varpi} \psi_{\xi} \quad \text{and} \quad \int_{\Omega} |f(x)|^2 \varpi(x) dx = \sum_{\xi \in \Xi} |\langle f, \psi_{\xi} \rangle|^2.$$

Because of their rapid decay, needlets can be used for decomposition of spaces of functions and distributions in various settings, including L^p , Sobolev, Besov, and Triebel-Lizorkin spaces.

The above narrative has been carried out in the setting of the regular domains. The localized kernels are most heavily studied and used when Ω is the unit sphere \mathbb{S}^{d-1} of \mathbb{R}^d , for which our outline of the localized frames is initiated in [34, 35]. They are further explored in, for example, [12, 28], on the sphere, and also studied on the interval $[-1, 1]$ with the Jacobi weight [5, 24, 37], the unit ball with the Gegenbauer weight [25, 38], and the simplex with the multivariable Jacobi weight [28] as well as for \mathbb{R}_+^d with the product Laguerre weight and \mathbb{R}^d with the Hermite weight [15, 18, 20, 39] and for product domains [29]. The subexponential decay of the kernel was established in [28], following earlier work in [16]. The needlets on the sphere and the ball have found applications in computational harmonics analysis, mathematical physics and statistics; see, for example, [2, 3, 26, 27, 31, 49] and references in them.

1.2. New contributions. Our starting point is the recent study of orthogonal structures on conic domains of \mathbb{R}^{d+1} in [53, 54], which shows that many properties of spherical harmonics and classical orthogonal polynomials on the unit ball have analogs on conic domains. Most importantly, there are families of orthogonal polynomials on

conic domains that possess properties akin to the addition formula and the Laplace-Beltrami operator for the spherical harmonics, the former provides a closed-form formula for the reproducing kernels and the latter is a second order differential operator that has orthogonal polynomials as eigenfunctions. This paves the way for carrying out the narrative outlined in the previous subsection on conic domains, which has been a topic largely untouched hitherto. In the present paper, we will deal with two types of conic domains and they are standardized as:

- (1) conic surface \mathbb{V}_0^{d+1} defined by

$$\mathbb{V}_0^{d+1} := \{(x, t) \in \mathbb{R}^{d+1} : \|x\| = t, 0 \leq t \leq 1, x \in \mathbb{R}^d\};$$

- (2) solid cone \mathbb{V}^{d+1} bounded by \mathbb{V}_0^{d+1} and the hyperplane $t = 1$.

Two other conic domains, double hyperbolic surface and solid hyperboloid, will be dealt with elsewhere because of the page limitation. On each domain, a closed-form formula of the reproducing kernels is uncovered for a family of weight functions, which allows us to study highly localized kernels and carries out the program described in Subsection 1.1, and a differential operator is defined that has orthogonal polynomials as eigenfunctions, which leads to a K -functional that is used to characterize the best approximation by polynomials on the domain.

Instead of dealing with each case individually, we will develop a unified theory for all homogeneous spaces that contain highly localized kernels that we describe in the first two subsections below. The unified theory is applicable to conic domains, which is described in the third and the fourth subsections.

1.2.1. Localizable homogeneous space. A homogeneous space is a measure space $(\Omega, d\mu, d)$, where $d\mu$ is a nonnegative doubling measure with respect to the metric $d(\cdot, \cdot)$ on Ω . When $d\mu = w(x)dx$, where w is a doubling weight function on Ω , we denote the homogeneous space by (Ω, w, d) . For example, the space $(\mathbb{S}^{d-1}, d\sigma, d_{\mathbb{S}})$ is a homogeneous space, so is $(\mathbb{B}^d, d\mu, d_{\mathbb{B}})$ for $\mu > -\frac{1}{2}$.

We call a homogeneous space (Ω, ϖ, d) *localizable* if the orthogonal polynomials with respect to ϖ on Ω admit highly localized kernels, which requires a fast decay estimate on $|L_n(\varpi; x, y)|$ as well as on $|L_n(\varpi; x_1, y) - L_n(\varpi; x_2, y)|$; see Subsection 2.1 for precise definition. Our first main result is to show that the narrative on the localized tight frame in the previous subsection can be established in a localizable homogeneous space. This provides a unified theory that is applicable to conical domains to be studied in later sections. Furthermore, it allows us to establish several intermediate results that are of interest in their own right for doubling weight functions, including Marcinkiewicz-Zygmund inequalities and positive cubature rules.

The development generalizes and follows closely recent studies on regular domains, especially those on the unit sphere and the unit ball. We follow the approach on the unit sphere in [7, 12], which is formulated in the homogeneous space on the sphere. While some of the results for the unit sphere can be extended easily, others need to be dealt with more carefully. Amongst various reasons, we mention that, on the unit sphere, it is the Lebesgue measure $d\sigma$ that admits the highly localized kernels. It is, however, not the case in general. For example, the highly localized kernels on the conic surface \mathbb{V}_0^{d+1} are established for the weight function $w_{-1, \gamma}(t) = t^{-1}(1-t)^\gamma$, which does not include the Lebesgue measure. While spherical caps of the same radius on \mathbb{S}^{d-1} have the same constant surface area, this is not the case on \mathbb{V}_0^{d+1} , where we need to deal with conical caps defined via the intrinsic distance $d_{\mathbb{V}_0}$, defined in (1.6) below,

and their volumes are measured by integrals over the caps against the weight function $w_{-1,\gamma}$.

1.2.2. Best approximation by polynomials. The characterization of best approximation is at the core of approximation theory (cf. [17]). The classical approach on the unit sphere relies essentially on multiplier operators of the Fourier harmonic series and the convolution structure arising from the addition formula. We provide a general framework on localized homogeneous space by assuming that the reproducing kernel $P_n(\varpi; \cdot, \cdot)$ satisfies an addition formula of a specific form satisfied by all known cases, including those on conic domains, and that there is a second order differential operator that has orthogonal polynomials as eigenfunctions. The former allows us to introduce a convolution structure that can be used to define a modulus of smoothness via a multiplier operator, and the latter is used to define a K -functional. The two quantities are then shown to be equivalent and either one can be used to characterize the error of the best polynomial approximation on the domain. The framework generalizes the classical result on the unit sphere and is applicable to other regular domains satisfying our assumptions, which are met by all four types of conic domains.

1.2.3. Conic surface \mathbb{V}_0^{d+1} and cone \mathbb{V}^{d+1} . On the conic surface \mathbb{V}_0^{d+1} , the orthogonal structure is studied in [53] for the weight function $w_{\beta,\gamma}(t) = t^\beta(1-t)^\gamma$ for $\beta > -d$ and $\gamma > -1$. For our study we need an intrinsic metric on the conic surface. In contrast to the unit sphere with its geodesic distance $d_S(\cdot, \cdot)$ that depend only on relative positions of the points, the conic surface \mathbb{V}_0^{d+1} has a boundary at $t = 1$ and an apex point at $t = 0$ that its metric needs to take into account. The distance that we will use is defined by

$$(1.6) \quad d_{V_0}((x, t), (y, s)) = \arccos \left[\sqrt{\frac{\langle x, y \rangle + ts}{2}} + \sqrt{1-t}\sqrt{1-s} \right].$$

Using this distance function and the closed-form formula of the reproducing kernel proved in [53], we shall show that $w_{\beta,\gamma}$ is a doubling weight and that the space $(\Omega, w_{-1,\gamma}, d_{V_0})$ is a localizable homogeneous space. These results are established through delicate estimates, partly because the distance $d_{V_0}(\cdot, \cdot)$ is not comparable to the Euclidean distance at around the apex of \mathbb{V}_0^{d+1} . With the highly localized kernel, we can then carry out our program and establish positive cubature rules, and use them to establish the highly localized frame. Furthermore, using the second order differential operator that has orthogonal polynomials with respect to $w_{-1,\gamma}$ as eigenfunctions, which is an analog of the Laplace-Beltrami operator in the unit sphere, we can define a K -functional and use it to establish a characterization of the best approximation by polynomials in the weighted L^p norm on \mathbb{V}_0^{d+1} .

Our study on the cone \mathbb{V}^{d+1} follows along the similar line. The distance function is defined by identifying \mathbb{V}^{d+1} with a subset of \mathbb{V}_0^{d+2} . The orthogonal structure is studied in [53] for the weight function $W_{\beta,\gamma,\mu}(x, t) = t^\beta(1-t)^\gamma(t^2 - \|x\|^2)^{\mu-\frac{1}{2}}$, which has a more involved close form formula for its reproducing kernels. While part of the estimate can be carried out for the kernel associated with $W_{0,\gamma,\mu}$, the localizable homogeneous space is established for $W_{0,\gamma,0}$, which nevertheless allows us to complete our program on the solid cone \mathbb{V}^{d+1} .

Several remarks are in order over the above description of our main results. First of all, analysis on conical domains have not been seriously studied in the literature as far as we are aware, and it does pose new phenomena and new challenges. In

particular, while the framework for localizable homogeneous space is applicable to conic domains, the verification of assertions under which the framework holds possess considerable difficulty in each case. Second, there are many intermediate results that are of interests. For example, to quantize the coefficients in positive cubature rule, we need bounds on the Christoffel functions

$$\lambda_n(\mathbf{w}; x) = \inf_{\substack{g(x)=1 \\ g \in \Pi_n(\Omega)}} \int_{\Omega} |g(x)|^2 \mathbf{w}(x) dx,$$

which are of interest in their own right. See, for example, [6, 14, 22, 23, 40, 41, 50] and their references for recent works on Christoffel functions in several variables. While the lower bound of the Christoffel function on a homogenous space can be derived using highly localized kernels when \mathbf{w} admits such kernels, we will establish the upper bound for all doubling weight under the assumption that certain fast decaying polynomials exist on Ω . The latter requires the construction of such polynomials in each conic domain. Third, there have been several recent works that deal with multivariate approximation by polynomials or Marcinkiewicz-Zygmund inequalities and positive cubature rules on either polyhedra domain or C^2 domain [8, 9, 47, 48]. They do not cover conical domains and our approach is different and relies on a specific orthogonal structure on the domain, akin to those on the unit sphere and the unit ball.

Finally, we should mention the important book *Analysis on Symmetric Cones* [19] and the literature around it. While the topic works in the theory of Euclidean Jordan algebras and lies in a more abstract setting, our study requires very specific structures of Fourier orthogonal series and is far less abstract. We have not been able to discern a connection between the two topics duo to our lack of background in Jordan algebras. It would be of great interest if a connection could be identified.

1.3. Organization and convention. The paper is organized as follows. The localizable homogeneous space will be defined and studied in the next section, which consists of results discussed in Subsection 1.2.1. The best approximation by polynomials in the homogeneous space is discussed in the third section, which consists of the results described in Subsection 1.2.2. The results on the conic surface and solid cone will be discussed in Sections 4 and 5, respectively. Each section will contain several subsections and its organization will be described in the preamble of the section.

Throughout this paper, we will denote by $L^p(\Omega, \mathbf{w})$ the weighted L^p space with respect to the weight function \mathbf{w} defined on the domain Ω for $1 \leq p \leq \infty$. Its norm will be denote by $\|\cdot\|_{p, \mathbf{w}}$ for $1 \leq p \leq \infty$ with the understanding that the space is $C(\Omega)$ with the uniform norm when $p = \infty$.

For conical domains, in an attempt to distinguish conic surfaces and solid conic bodies, we shall denote the operator on the surface in sans serif font, such as \mathbf{P}_n and \mathbf{L}_n , and the operator on the solid domains in bold font, such as \mathbf{P}_n and \mathbf{L}_n .

Finally, we shall use c, c', c_1, c_2 etc. to denote positive constants that depend on fixed parameters and their values may change from line to line. Furthermore, we shall write $A \sim B$ if $c'A \leq B \leq cA$.

2. HOMOGENEOUS SPACES WITH HIGHLY LOCALIZED KERNELS

Much of recent work on regular domains, such as the sphere, the ball, and the regular simplex falls in the framework of homogeneous space. What distinguishes these regular domains is the existence of highly localized kernels constructed via orthogonal polynomials. From the prior work on these regular domains and what we will prove

for the conic domains latter sections, it becomes clear that the desired estimates for the highly localized kernels share a common formation. In this section, we work with homogeneous spaces that are assumed to contain highly localized kernels and carry out our analysis on such spaces.

The first subsection contains basics on homogeneous space and orthogonal polynomials, and it contains three assertions that define highly localized kernels precisely. Three examples are given in the second subsection, which also serves as a review of the Jacobi polynomials, spherical harmonics, and orthogonal polynomials on the ball that will be used later. An auxiliary maximal function is introduced in the third subsection and, with the help of highly localized kernels for ϖ , is shown to be bounded in weighted L^p space with respect to a doubling weight. Using this maximal function, Marcinkiewicz-Zygmund inequalities are established in the fourth subsection for all doubling weights. The fifth subsection contains the fourth assertion that is used to obtain an upper bound for the Christoffel function. The latter is needed in the sixth subsection, where the positive cubature rules are established. With these preparations, the localized tight frame is defined and studied in the seventh subsection.

The development of this section follows and generalizes the study on the unit sphere. Some parts of the work follow those on the sphere with little extra effort but by no means all. Since the geodesic distance on the sphere and the surface measure on the unit sphere will be replaced by a distance function and a weight function that offer no specific geometric information, we will provide complete proofs in the most part and will be brief only when the proof follows that on the unit sphere fairly straightforwardly.

2.1. Localizable homogeneous spaces.

2.1.1. Homogeneous spaces. A homogeneous space is a measure space $(\Omega, \mu, \mathbf{d})$ with a positive measure μ and a metric \mathbf{d} such that all open balls

$$B(x, r) = \{y \in \Omega : \mathbf{d}(x, y) < r\},$$

are measurable and μ is a regular measure satisfying the doubling property

$$\mu(B(x, 2r)) \leq c\mu(B(x, r)), \quad \forall x \in \Omega, \quad \forall r > 0,$$

where c is independent of x and r . Such a measure μ is called a doubling measure.

Let Ω be a domain in \mathbb{R}^d . We assume that Ω is equipped with an intrinsic distance \mathbf{d} , so that the Lebesgue measure \mathbf{m} on Ω is a doubling measure and, for each $\varepsilon > 0$, $\min_{x \in \Omega} \mathbf{m}(B(x, \varepsilon)) \geq c_\varepsilon > 0$. Let \mathbf{w} be a nonnegative integrable function defined on Ω . For a given set $E \subset \Omega$, we define

$$\mathbf{w}(E) = \int_E \mathbf{w}(x) \mathbf{d}\mathbf{m}(x).$$

The weight function \mathbf{w} is called a doubling weight if it satisfies the doubling condition: there exists a constant $L > 0$ such that

$$\mathbf{w}(B(x, 2r)) \leq L\mathbf{w}(B(x, r)), \quad \forall x \in \Omega, \quad r \in (0, r_0),$$

where r_0 is the largest positive number such that $B(x, r) \subset \Omega$. If \mathbf{w} is a doubling weight, then $\mathbf{d}\mu = \mathbf{w}(x) \mathbf{d}\mathbf{m}$ is a doubling measure. We are particularly interested in the homogeneous space with its measure so defined, which we denote by $(\Omega, \mathbf{w}, \mathbf{d})$.

For a doubling weight \mathbf{w} on Ω with respect to \mathbf{d} , we denote by $L(\mathbf{w})$ the least constant L for the doubling condition $\mathbf{w}(B(x, 2r)) \leq L\mathbf{w}(B(x, r))$. Let $\alpha(\mathbf{w})$ be a

positive number such that

$$(2.1) \quad \sup_{B(x,r) \subset \Omega} \frac{w(B(x, 2^m r))}{w(B(x, r))} \leq c_{L(w)} 2^{\alpha(w)m}, \quad m = 1, 2, \dots$$

Then $\alpha(w) \leq \log_2 L(w)$, as can be seen by iteration. We call $L(w)$ the doubling constant and $\alpha(w)$ the doubling index.

Lemma 2.1. *Let w be a doubling weight on Ω .*

(i) *If $0 < r < t$ and $x \in \Omega$, then*

$$w(B(x, t)) \leq c_{L(w)} \left(\frac{t}{r} \right)^{\alpha(w)} w(B(x, r)).$$

(ii) *For $x, y \in \Omega$ and $n = 1, 2, \dots$,*

$$w(B(x, n^{-1})) \leq c_{L(w)} (1 + nd(x, y))^{\alpha(w)} w(B(y, n^{-1})).$$

Proof. Assume $2^{m-1} \leq t/r \leq 2^m$, (i) follows immediately from the definition of $\alpha(w)$. By the triangle inequality of the distance function, $w(B(x, n^{-1})) \leq w(B(y, d(x, y) + n^{-1}))$, applying (i) to bound it by $w(B(y, n^{-1}))$ proves (ii). \square

Many fundamental results that hold for the Euclidean space can be extended to homogeneous spaces [44]. For example, the Hardy-Littlewood maximal function is defined, for a locally integrable function f , by

$$(2.2) \quad M_w f(x) := \sup_{r>0} \frac{1}{w(B(x, r))} \int_{B(x, r)} |f(y)| w(y) dy, \quad x \in \Omega,$$

which possesses the usual properties of the maximal function [44, p. 13]. For example, for every $f \in L^p(\Omega, w)$,

$$(2.3) \quad \|M_w f\|_{p,w} \leq c_p \|f\|_{p,w}, \quad 1 < p \leq \infty,$$

where $\|f\|_{p,w}$ denotes the L^p norm with respect to the measure $w(x)dx$.

We study homogeneous spaces that contain highly localized kernels, to be defined below. Such spaces are known to hold for certain weight functions on several regular domains. We adopt the convention of denoting the weight function that admits highly localized kernels by ϖ and we also assume that ϖ is normalized by $\int_{\Omega} \varpi(x) dx = 1$. The kernels are defined via orthogonal polynomials with respect to the inner product $\langle \cdot, \cdot \rangle_{\varpi}$ defined in terms of ϖ in (2.4).

2.1.2. Orthogonal polynomials. Let $\Pi(\Omega)$ denote the space of polynomials restricted to Ω and let $\Pi_n(\Omega)$ denote its subspace of polynomials of degree at most n . If the interior of Ω is open, then $\Pi(\Omega)$ contains all polynomials of degree n in d variables and we write $\Pi_n = \Pi_n(\Omega)$. If Ω is an algebraic surface, such as the sphere \mathbb{S}^{d-1} , then $\Pi_n(\Omega)$ contains all polynomials restricted to the surface.

Let ϖ be a normalized weight function on Ω . Then the bilinear form $\langle \cdot, \cdot \rangle_{\varpi}$

$$(2.4) \quad \langle f, g \rangle_{\varpi} := \int_{\Omega} f(x)g(x)\varpi(x)dx$$

is a well defined inner product on $\Pi_n(\Omega)$. Let $\mathcal{V}_n^d(\Omega, \varpi)$ be the space of orthogonal polynomials of degree n with respect to this inner product. We are particularly interested in the case when Ω is either a quadratic surface, such as the unit sphere \mathbb{S}^{d-1} or

the conical surface \mathbb{V}_0^d , for which

$$(2.5) \quad \dim \mathcal{V}_n^d(\Omega, \varpi) = \binom{n+d-2}{n} + \binom{n+d-3}{n-1}, \quad n = 1, 2, 3, \dots,$$

where we assume $\binom{n}{k} = 0$ if $k < 0$, or a solid domain bounded by a quadratic surface, such as the unit ball \mathbb{B}^d and the solid cone \mathbb{V}^d , for which

$$(2.6) \quad \dim \mathcal{V}_n^d(\Omega, \varpi) = \binom{n+d-1}{n}, \quad n = 0, 1, 2, \dots,$$

The reproducing kernel of $\mathcal{V}_n^d(\Omega, \varpi)$, denoted by $P_n(\varpi; \cdot, \cdot)$, is uniquely determined by

$$\int_{\Omega} f(y) P_n(\varpi; x, y) \varpi(y) \mathrm{d}\mathbf{m}(y) = f(x), \quad \forall f \in \mathcal{V}_n^d(\Omega, \varpi).$$

If $\{P_{\nu,n} : 1 \leq \nu \leq \dim \mathcal{V}_n^d(\Omega, \varpi)\}$ is an orthogonal basis of $\mathcal{V}_n^d(\Omega, \varpi)$, then

$$(2.7) \quad P_n(\varpi; x, y) = \sum_{\nu=1}^{\dim \mathcal{V}_n^d(\Omega, \varpi)} \frac{P_{\nu,n}(x) P_{\nu,n}(y)}{\langle P_{\nu,n}, P_{\nu,n} \rangle_{\varpi}}.$$

The kernel plays an essential role in the study of the Fourier orthogonal series, since it is the kernel of the orthogonal projection operator $\mathrm{proj}_n : L^2(\Omega, \varpi) \mapsto \mathcal{V}_n^d(\Omega, \varpi)$; more precisely,

$$(2.8) \quad \mathrm{proj}_n(\varpi; f, x) = \int_{\Omega} f(y) P_n(\varpi; x, y) \varpi(y) \mathrm{d}\mathbf{m}(y), \quad f \in L^2(\Omega, \varpi).$$

The Fourier orthogonal series of $f \in L^2(\Omega, \varpi)$ is defined by

$$(2.9) \quad f = \sum_{n=0}^{\infty} \mathrm{proj}_n(\varpi; f) = \sum_{n=0}^{\infty} \sum_{\nu=1}^{\dim \mathcal{V}_n^d(\Omega, \varpi)} \hat{f}_{\nu,n} P_{\nu,n}, \quad \hat{f}_{\nu,n} = \frac{\langle f, P_{\nu,n} \rangle_{\varpi}}{\langle P_{\nu,n}, P_{\nu,n} \rangle_{\varpi}}.$$

For these definitions and orthogonal polynomials of several variables in general, we refer to [14].

Our highly localized kernels are defined by a sampling of the kernels of the Fourier series through a smooth cut-off function, which we define first.

Definition 2.2. A nonnegative function $\hat{a} \in C^\infty(\mathbb{R})$ is said to be admissible if it obeys either one of the conditions

- (a) $\mathrm{supp} \hat{a} \subset [0, 2]$ and $\hat{a}(t) = 1$, $t \in [0, 1]$; or
- (b) $\mathrm{supp} \hat{a} \subset [1/2, 2]$.

Let \hat{a} be an admissible cut-off function. For $n = 0, 1, 2, \dots$, we define the kernel function $L_n(\varpi)$ on $\Omega \times \Omega$ by

$$(2.10) \quad L_n(\varpi; x, y) = \sum_{k=0}^{\infty} \hat{a}\left(\frac{k}{n}\right) P_n(\varpi; x, y), \quad x, y \in \Omega,$$

Since \hat{a} is supported on $[0, 2]$, this is a kernel of polynomials of degree at most $2n$ in either x or y variable.

2.1.3. Localizable homogeneous spaces. The main purpose of this section is to study homogeneous spaces $(\Omega, \varpi, \mathbf{d})$ for which the kernels $L_n(\varpi; \cdot, \cdot)$ are assumed to satisfy the following assertions.

Definition 2.3. *The kernels $L_n(\varpi; \cdot, \cdot)$, $n = 1, 2, \dots$, are called highly localized if they satisfy the following assertions:*

Assertion 1. *For $\kappa > 0$ and $x, y \in \Omega$,*

$$|L_n(\varpi; x, y)| \leq c_\kappa \frac{1}{\sqrt{\varpi(B(x, n^{-1}))} \sqrt{\varpi(B(y, n^{-1}))} (1 + n\mathbf{d}(x, y))^\kappa}.$$

Assertion 2. *For $0 < \delta \leq \delta_0$ with some $\delta_0 < 1$ and $x_1 \in B(x_2, \frac{\delta}{n})$,*

$$|L_n(\varpi; x_1, y) - L_n(\varpi; x_2, y)| \leq c_\kappa \frac{n\mathbf{d}(x_1, x_2)}{\sqrt{\varpi(B(x_1, n^{-1}))} \sqrt{\varpi(B(x_2, n^{-1}))} (1 + n\mathbf{d}(x_2, y))^\kappa}.$$

Assertion 3. *For sufficient large $\kappa > 0$,*

$$\int_{\Omega} \frac{\varpi(y)}{\varpi(B(y, n^{-1})) (1 + n\mathbf{d}(x, y))^\kappa} \mathbf{d}m(y) \leq c.$$

The third assertion affirms the sharpness of the first two assertions. It implies the following more general inequality.

Lemma 2.4. *Let ϖ be a doubling weight that satisfies Assertion 3 with $\kappa > 0$. For $0 < p < \infty$, let $\tau = \kappa - \frac{p}{2}|\alpha(\varpi)|1 - \frac{p}{2}| > 0$. Then, for $x \in \Omega$,*

$$(2.11) \quad \int_{\Omega} \frac{\varpi(y)}{\varpi(B(y, n^{-1}))^{\frac{p}{2}} (1 + n\mathbf{d}(x, y))^\tau} \mathbf{d}m(y) \leq c \varpi(B(x, n^{-1}))^{1 - \frac{p}{2}}.$$

Proof. Denote the left-hand side of (2.11) by $J_{p,\tau}$. When $p = 2$, the inequality (2.11) is precisely the Assertion 3. The case $p \neq 2$ can be deduced from the case $d = 2$. Indeed, by Lemma 2.1, it is easy to see that,

$$(2.12) \quad (1 + n\mathbf{d}(x, y))^{-\alpha(\varpi)} \leq \frac{\varpi(B(y, n^{-1}))}{\varpi(B(x, n^{-1}))} \leq (1 + n\mathbf{d}(x, y))^{\alpha(\varpi)}.$$

For $p \neq 2$, using the right-hand side inequality if $0 < p < 2$ and the left-hand side inequality if $p > 2$, we obtain

$$\varpi(B(y, n^{-1}))^{\frac{p}{2}} \geq \frac{\varpi(B(y, n^{-1})) \varpi(B(x, n^{-1}))^{\frac{p}{2}-1}}{(1 + n\mathbf{d}(x, y))^{\alpha(\varpi)|1 - \frac{p}{2}|}},$$

which gives $J_{p,\tau} \leq \varpi(B(x, n^{-1}))^{1 - \frac{p}{2}} J_{2,\kappa}$. Hence, (2.11) for $p \neq 2$ follows from the case $p = 2$. The proof is completed. \square

Corollary 2.5. *For $0 < p < \infty$ and $x \in \Omega$, the highly localized kernel satisfies*

$$(2.13) \quad \int_{\Omega} |L_n(\varpi; x, y)|^p \varpi(y) \mathbf{d}m(y) \leq c [\varpi(B(x, n^{-1}))]^{1-p}.$$

In particular, for $p = 1$, it ensures that the Assertion 1 is sharp in the sense that the estimate guarantees the boundedness of $|L_n(\varpi; x, y)|$ and it ensures that Assertion 2 is sharp in the same sense.

Definition 2.6. *The homogeneous space $(\Omega, \varpi, \mathbf{d})$ is called localizable if ϖ is a doubling weight that admits highly localized kernels. For convenience, we also say that the domain Ω , or the weight ϖ , admits a localizable homogeneous space without specifying ϖ , or Ω .*

Throughout the rest of this section, we assume that the weight function ϖ admits a localizable homogeneous space on Ω .

2.2. Examples of localizable homogeneous spaces. In this subsection, we give three examples of localizable homogeneous spaces: the interval $[-1, 1]$ with the Jacobi weight, the unit sphere \mathbb{S}^{d-1} with the surface measure, the unit ball \mathbb{B}^d with its classical weight function. These cases have been thoroughly studied fairly recently in the literature, and they motivate our definition of the localizable homogeneous spaces. Moreover, their corresponding orthogonal polynomials serve as the building blocks for orthogonal polynomials on the conic domains.

2.2.1. The interval $[-1, 1]$ with the Jacobi weight. For $\alpha, \beta > -1$, the Jacobi weight function is defined by

$$w_{\alpha, \beta}(t) := (1-t)^\alpha(1+t)^\beta, \quad -1 < t < 1.$$

Its normalization constant $c'_{\alpha, \beta}$, defined by $c'_{\alpha, \beta} \int_{-1}^1 w_{\alpha, \beta}(x) dx = 1$, is given by

$$(2.14) \quad c'_{\alpha, \beta} = \frac{1}{2^{\alpha+\beta+1}} c_{\alpha, \beta} \quad \text{with} \quad c_{\alpha, \beta} := \frac{\Gamma(\alpha + \beta + 2)}{\Gamma(\alpha + 1)\Gamma(\beta + 1)}.$$

The intrinsic distance of the interval $[-1, 1]$ is defined by

$$(2.15) \quad d_{[-1, 1]}(t, s) = \arccos \left(ts + \sqrt{1-t^2} \sqrt{1-s^2} \right),$$

which is the projection of the distance $|\theta - \phi|$ on the upper half of the unit circle if we set $t = \cos \theta$ and $s = \cos \phi$. The space $([-1, 1], w_{\alpha, \beta}, d_{[-1, 1]})$ is a localizable homogeneous space if $\alpha, \beta \geq -\frac{1}{2}$.

The orthogonal polynomials with respect to $w_{\alpha, \beta}$ are the Jacobi polynomials $P_n^{(\alpha, \beta)}$, which are given by the hypergeometric function as [46, (4.21.2)]

$$P_n^{(\alpha, \beta)}(x) = \frac{(\alpha + 1)_n}{n!} {}_2F_1 \left(\begin{matrix} -n, n + \alpha + \beta + 1 \\ \alpha + 1 \end{matrix}; \frac{1-x}{2} \right).$$

These polynomials satisfy the orthogonal relations

$$c'_{\alpha, \beta} \int_{-1}^1 P_n^{(\alpha, \beta)}(x) P_m^{(\alpha, \beta)}(x) w_{\alpha, \beta}(x) dx = h_n^{(\alpha, \beta)} \delta_{m, n},$$

where $h_n^{(\alpha, \beta)}$ is the square of the L^2 norm that satisfies

$$h_n^{(\alpha, \beta)} = \frac{(\alpha + 1)_n (\beta + 1)_n (\alpha + \beta + n + 1)}{n! (\alpha + \beta + 2)_n (\alpha + \beta + 2n + 1)}.$$

The Jacobi polynomials are eigenfunctions of a second order differential operator:

$$[(1-x)\partial^2 - (\alpha - \beta + (\alpha + \beta + 2)x)\partial] u = -n(n + \alpha + \beta + 1)u,$$

where $\partial f = f'$ and $u = P_n^{(\alpha, \beta)}$ for $n = 0, 1, 2, \dots$. The highly localized kernel for the Jacobi polynomials will be denoted by

$$(2.16) \quad L_n^{(\alpha, \beta)}(x, y) = \sum_{j=0}^{\infty} \hat{a} \left(\frac{j}{n} \right) \frac{P_j^{(\alpha, \beta)}(x) P_j^{(\alpha, \beta)}(y)}{h_j^{(\alpha, \beta)}},$$

where \widehat{a} is an admissible cutoff function. This kernel decays rapidly in terms of the distance $d_{[-1,1]}(\cdot, \cdot)$. Let the ball $B(x, r)$ be defined with respect to this distance. Then the Jacobi weight is a doubling weight and satisfies,

$$w_{\alpha,\beta}(B(x, n^{-1})) \sim n^{-1} w_{\alpha,\beta}(n; t),$$

where $w_{\alpha,\beta}(n; t)$ is defined by

$$w_{\alpha,\beta}(n; t) = (1 - t + n^{-2})^{\alpha+\frac{1}{2}} (1 + t + n^{-2})^{\beta+\frac{1}{2}}.$$

The following estimate of the localized kernel can be found in [37].

Theorem 2.7. *Let $\alpha, \beta \geq -1/2$ and $0 < \varepsilon \leq 1$. Then, for any $\kappa > 0$ and $n \geq 1$,*

$$|L_n^{(\alpha,\beta)}(t, s)| \leq \frac{cn}{\sqrt{w_{\alpha,\beta}(n; t)} \sqrt{w_{\alpha,\beta}(n; s)}} (1 + nd_{[-1,1]}(t, s))^{-\kappa},$$

This verifies Assertion 1 of the Definition 2.3. The Assertions 2 and 3 can be found in [37, 24], respectively. The proof of the estimate relies on the estimate for the special case

$$L_n^{(\alpha,\beta)}(t) = L_n^{(\alpha,\beta)}(t, 1), \quad -1 \leq t \leq 1.$$

If $s = 1$ and $t = \cos \theta$, then $d_{[-1,1]}(t, 1) = \theta \sim \sin \theta/2 \sim \sqrt{1-t}$ and we obtain [5],

$$(2.17) \quad \left| L_n^{(\alpha,\beta)}(t) \right| \leq c \frac{n^{2\alpha+2}}{(1 + n\sqrt{1-t})^\kappa}, \quad -1 \leq t \leq 1,$$

which is used to establish the estimate in Theorem 2.7. There is also a more general estimate stated under weaker assumption on the cut-off function [5] and [12, Theorem 2.6.7], which will be useful in later development.

Theorem 2.8. *Let ℓ be a positive integer and let η be a function that satisfy, $\eta \in C^{3\ell-1}(\mathbb{R})$, $\text{supp } \eta \subset [0, 2]$ and $\eta^{(j)}(0) = 0$ for $j = 0, 1, 2, \dots, 3\ell-2$. Then, for $\alpha \geq \beta \geq -\frac{1}{2}$, $t \in [-1, 1]$ and $n \in \mathbb{N}$,*

$$(2.18) \quad \left| \frac{d^m}{dt^m} L_n^{(\alpha,\beta)}(t) \right| \leq c_{\ell,m,\alpha} \left\| \eta^{(3\ell-1)} \right\|_\infty \frac{n^{2\alpha+2m+2}}{(1 + n\sqrt{1-t})^\ell}, \quad m = 0, 1, 2, \dots$$

The Jacobi polynomials with equal parameters are the Gegenbauer polynomials C_n^λ , which are of particular interest for our study. These polynomials are orthogonal with respect to the the weight function

$$w_\lambda(x) = (1 - x^2)^{\lambda-\frac{1}{2}}, \quad -1 < x < 1, \quad \lambda > -\frac{1}{2}.$$

More precisely, the polynomials C_n^λ satisfy the orthogonal relation

$$(2.19) \quad c_\lambda \int_{-1}^1 C_n^\lambda(x) C_m^\lambda(x) w_\lambda(x) dx = h_n^\lambda \delta_{n,m},$$

where the normalization constant c_λ of w_λ and the norm square h_n^λ are given by

$$(2.20) \quad c_\lambda = \frac{\Gamma(\lambda+1)}{\Gamma(\frac{1}{2})\Gamma(\lambda+\frac{1}{2})} \quad \text{and} \quad h_n^\lambda = \frac{\lambda}{n+\lambda} C_n^\lambda(1) = \frac{\lambda}{n+\lambda} \frac{(2\lambda)_n}{n!}.$$

The polynomial C_n^λ is a constant multiple of the Jacobi polynomial $P_n^{(\lambda-\frac{1}{2}, \lambda-\frac{1}{2})}$ and it is also related to the Jacobi polynomial by a quadratic transform [46, (4.7.1)]

$$(2.21) \quad C_{2n}^\lambda(x) = \frac{(\lambda)_n}{(\frac{1}{2})_n} P_n^{(\lambda-\frac{1}{2}, \lambda-\frac{1}{2})}(2x^2 - 1).$$

2.2.2. *The unit sphere \mathbb{S}^{d-1} .* With the unit weight function, or the Lebesgue measure $d\sigma$, the space $(\mathbb{S}^{d-1}, d\sigma, d_{\mathbb{S}})$ is a localizable homogeneous space, where the metric $d_{\mathbb{S}} = d_{\mathbb{S}^{d-1}}$ is the geodesic distance defined in (1.2). The orthogonal polynomials are spherical harmonics. Let \mathcal{P}_n^d denote the space of homogeneous polynomials of degree n in d variables. A spherical harmonics Y of degree n is an element of \mathcal{P}_n^d that satisfies $\Delta Y = 0$, where Δ is the Laplace operator of \mathbb{R}^d . If $Y \in \mathcal{P}_n^d$, then $Y(x) = \|x\|^n Y(x')$, $x' = x/\|x\| \in \mathbb{S}^{d-1}$, so that Y is determined by its restriction on the unit sphere.

Let $\mathcal{H}_n(\mathbb{S}^{d-1})$ denote the space of spherical harmonics of degree n . Its dimension $\dim \mathcal{H}_n(\mathbb{S}^{d-1})$ is given by (2.5). Spherical harmonics of different degrees are orthogonal on the sphere. For $n \in \mathbb{N}_0$ let $\{Y_\ell^n : 1 \leq \ell \leq \dim \mathcal{H}_n(\mathbb{S}^{d-1})\}$ be an orthonormal basis of $\mathcal{H}_n(\mathbb{S}^{d-1})$; then

$$\frac{1}{\omega_d} \int_{\mathbb{S}^{d-1}} Y_\ell^n(\xi) Y_{\ell'}^m(\xi) d\sigma(\xi) = \delta_{\ell, \ell'} \delta_{m, n},$$

where ω_d denotes the surface area $\omega_d = 2\pi^{\frac{d}{2}}/\Gamma(\frac{d}{2})$ of \mathbb{S}^{d-1} . Let $P_n(\cdot, \cdot)$ denote the reproducing kernel of the space $\mathcal{H}_n(\mathbb{S}^{d-1})$. Among many properties of the spherical harmonics (see, for example, [12, 45]), one characteristic property is a closed-formula for this kernel [12, (1.2.3) and (1.2.7)],

$$(2.22) \quad P_n(x, y) = \sum_{\ell=1}^{\dim \mathcal{H}_n(\mathbb{S}^{d-1})} Y_\ell^n(x) Y_\ell^n(y) = Z_n^{\frac{d-2}{2}}(\langle x, y \rangle), \quad x, y \in \mathbb{S}^{d-1},$$

where Z_n^λ is defined in terms of the Gegenbauer polynomial by

$$(2.23) \quad Z_n^\lambda(t) = \frac{n + \lambda}{\lambda} C_n^\lambda(t), \quad \lambda = \frac{d-2}{2}.$$

Because of the second equal sign, the identity (2.22) is often called the *addition formula* of spherical harmonics. The function in its right-hand side is often called zonal harmonic, since it depends only on $\langle x, y \rangle$. This formula allows us to use (2.17) to derive the estimate (1.1), which is Assertion 1, for the highly localized kernel

$$L_n(x, y) = \sum_{k=0}^{\infty} \hat{a}\left(\frac{k}{n}\right) P_k(x, y), \quad x, y \in \mathbb{S}^{d-1}.$$

The other two assertions also hold; see, for example, [12, Thm. 6.5] and [12, Cor. 2.6.6].

Another characteristic property of the spherical harmonics is that they are eigenfunctions of the Laplace-Beltrami operator Δ_0 on the sphere, which is the restriction of the Laplace operator Δ on the unit sphere; see, for example, [12, Section 1.4] for its explicit expression. More precisely, it is known [12, (1.4.9)] that

$$(2.24) \quad \Delta_0 Y = -n(n + d - 2)Y, \quad Y \in \mathcal{H}_n^d,$$

so that the eigenvalues depend only on n . This relation, as well as the operator Δ_0 , plays an important role in the approximation theory on the unit sphere.

2.2.3. *The unit ball \mathbb{B}^d with classical weight function.* The classical weight function on the unit ball \mathbb{B}^d of \mathbb{R}^d is defined by

$$(2.25) \quad W_\mu(x) = (1 - \|x\|)^{\mu - \frac{1}{2}}, \quad x \in \mathbb{B}^d, \quad \mu > -\frac{1}{2}.$$

Its normalization constant is $b_\mu^\mathbb{B} = \Gamma(\mu + \frac{d+1}{2})/(\pi^{\frac{d}{2}}\Gamma(\mu + \frac{1}{2}))$. For the ball $B(x, r)$ defined via the distance function $\mathbf{d}_\mathbb{B}$ in (1.4) on the unit ball [38, Lemma 5.1],

$$W_\mu(B(x, r)) \sim r^d \left(\sqrt{1 - \|x\|^2} + r \right)^{2\mu}, \quad 0 < r \leq 1,$$

so that W_μ is a doubling weight. The space $(\mathbb{B}^d, W_\mu, \mathbf{d}_\mathbb{B})$ is a localizable homogeneous space for $\mu \geq 0$, where $\mathbf{d}_\mathbb{B}(\cdot, \cdot)$ is the distance defined in (1.4) on the unit ball.

Let $\mathcal{V}_n^d(\mathbb{B}^d, W_\mu)$ be the space of orthogonal polynomials of degree n with respect to W_μ . An orthogonal basis of $\mathcal{V}_n^d(\mathbb{B}^d, W_\mu)$ can be given explicitly in terms of the Jacobi polynomials and spherical harmonics. For $0 \leq m \leq n/2$, let $\{Y_\ell^{n-2m} : 1 \leq \ell \leq \dim \mathcal{H}_{n-2m}(\mathbb{S}^{d-1})\}$ be an orthonormal basis of \mathcal{H}_{n-2m}^d . Define [14, (5.2.4)]

$$(2.26) \quad P_{\ell, m}^n(x) = P_m^{(\mu - \frac{1}{2}, n - 2m + \frac{d-2}{2})}(2\|x\|^2 - 1) Y_\ell^{n-2m}(x).$$

Then $\{P_{\ell, m}^n : 0 \leq m \leq n/2, 1 \leq \ell \leq \dim \mathcal{H}_{n-2m}(\mathbb{S}^{d-1})\}$ is an orthogonal basis of $\mathcal{V}_n^d(W_\mu)$. Let $\mathbf{P}_n(W_\mu; \cdot, \cdot)$ denote the reproducing kernel of the space $\mathcal{V}_n(\mathbb{B}^d, W_\mu)$. This kernel satisfies an analog of the addition formula [51]: for $x, y \in \mathbb{B}^d$,

$$(2.27) \quad \mathbf{P}_n(W_\mu; x, y) = c_{\mu - \frac{1}{2}} \int_{-1}^1 Z_n^{\mu + \frac{d-1}{2}} \left(\langle x, y \rangle + u \sqrt{1 - \|x\|^2} \sqrt{1 - \|y\|^2} \right) \times (1 - u^2)^{\mu-1} du,$$

where $\mu > 0$ and it holds for $\mu = 0$ under the limit

$$(2.28) \quad \lim_{\mu \rightarrow 0+} c_{\mu - \frac{1}{2}} \int_{-1}^1 f(t)(1 - t^2)^{\mu-1} du = \frac{f(1) + f(-1)}{2}.$$

This is an analog of the addition formula for the unit ball and it allows us to use (2.17) to derive the estimate (1.3), which is Assertion 1, for the highly localized kernel now denoted by

$$\mathbf{L}_n(W_\mu; x, y) = \sum_{k=0}^{\infty} \widehat{a} \left(\frac{k}{n} \right) \mathbf{P}_k(W_\mu; x, y).$$

All three assertions were proved in [38]. For the first one, see also [12, Thm. 11.5.3].

The orthogonal polynomials with respect to W_μ on the unit ball are eigenfunctions of a second order differential operator [14, (5.2.3)]:

$$(2.29) \quad \left(\Delta - \langle x, \nabla \rangle^2 - (2\mu + d - 1) \langle x, \nabla \rangle \right) u = -n(n + 2\mu + d - 1)u$$

for all $u \in \mathcal{V}_n(\mathbb{B}^d, W_\mu)$. The differential operator in the left-hand side is the analog of the Laplace-Beltrami operator on the sphere.

2.3. An auxiliary maximal function. Some of the polynomial inequalities in the latter sections will be established for norms defined via a doubling weight. The study of approximation and polynomial inequality with doubling weights was pioneered in [32] for the interval $[-1, 1]$ and developed subsequently in [7] for the unit sphere \mathbb{S}^{d-1} . The main tool for the latter is an auxiliary maximal function. We define its analog on the domain Ω that admits a localizable homogeneous space.

Definition 2.9. For $\beta > 0$, $n \in \mathbb{N}$ and $f \in C(\Omega)$, a maximal function $f_{\beta, n}^*$ is defined by

$$f_{\beta, n}^*(x) := \max_{y \in \Omega} |f(y)| (1 + n \mathbf{d}(x, y))^{-\beta}, \quad x \in \Omega.$$

The development below follows closely [12, Sect. 5.2 and 5.3] but with a somewhat more streamlined proof.

We clearly have $|f(x)| \leq f_{\beta,n}^*(x)$ for all $x \in \Omega$. The maximal function $f_{\beta,n}^*$ has an upper bound in the Hardy-Littlewood maximum function M_μ defined in (2.2). To see this, we first prove a lemma. For $f \in C(\Omega)$ and $r > 0$, we define

$$\text{osc}(f)(x, r) := \sup_{x_1, x_2 \in B(x, r)} |f(x_1) - f(x_2)|, \quad x \in \Omega.$$

Lemma 2.10. *If $f \in \Pi_n(\Omega)$ and $0 < \delta \leq 1$, then for any $\beta > 0$,*

$$\text{osc}(f)\left(x, \frac{\delta}{n}\right) \leq c_\beta \delta f_{\beta,n}^*(x), \quad x \in \Omega,$$

where the constant c_β depends only on d and β when β is large.

Proof. We use the highly localized kernel L_n with an admissible cut-off function of type (a). For $\delta > 0$ and $x, y \in \Omega$, set

$$A_{n,\delta}(x, y) := \sup_{z \in B(x, \frac{\delta}{n})} |L_n(\varpi; x, y) - L_n(\varpi; z, y)|.$$

Let $L_n(\varpi) * f$ be defined as in (1.5). Then $L_n(\varpi) * f = f$ for $f \in \Pi_n(\Omega)$. If $x_1, x_2 \in B(x, r)$, then by the triangle inequality

$$|L_n(\varpi; x_1, y) - L_n(\varpi; x_2, y)| \leq 2A_{n,\delta}(x, y).$$

Hence, it follows by (1.5) that

$$\begin{aligned} \text{osc}(f)\left(x, \frac{\delta}{n}\right) &\leq 2 \int_{\Omega} |f(y)| A_{n,\delta}(x, y) \varpi(y) \, d\mathbf{m}(y) \\ &\leq 2f_{\beta,n}^*(x) \int_{\Omega} (1 + n\mathbf{d}(x, y))^\beta A_{n,\delta}(x, y) \, d\mathbf{m}(y) \\ &\leq c_\beta \delta f_{\beta,n}^*(x), \end{aligned}$$

where the last step follows from Assertion 1 and Assertion 3. \square

Theorem 2.11. *Let w be a doubling weight. For $f \in \Pi_n(\Omega)$, $\beta > 0$ and setting $\gamma = \alpha(w)/\beta$, then*

$$(2.30) \quad f_{\beta,n}^*(x) \leq c_{\beta,L(w)} \left(M_w(|f|^\gamma)(x) \right)^{1/\gamma}, \quad x \in \Omega.$$

Proof. We apply Lemma 2.10 with $c_\beta \delta = \frac{1}{4}$ to obtain, by $(a+b)^\gamma \leq 2^\gamma(a^\gamma + b^\gamma)$,

$$\frac{|f(y)|^\gamma}{(1 + n\mathbf{d}(x, y))^{\alpha(w)}} \leq 2^\gamma \frac{\min_{z \in B(y, \frac{\delta}{n})} |f(z)|^\gamma}{(1 + n\mathbf{d}(x, y))^{\alpha(w)}} + \frac{1}{2^\gamma} (f_{\beta,n}^*(x))^\gamma,$$

Taking maximum over $x \in \Omega$, the left-hand side becomes $(f_{\beta,n}^*(x))^\gamma$ using $\alpha(w) = \beta\gamma$, so that the inequality can be rearranged to give

$$(2.31) \quad (f_{\beta,n}^*(x))^\gamma \leq c(1 + n\mathbf{d}(x, y))^{-\alpha(w)} \min_{z \in B(y, \frac{\delta}{n})} |f(z)|^\gamma,$$

where $c = 2^\gamma/(1 - 2^{-\gamma})$. We now estimate $\min_{z \in B(y, \frac{\delta}{n})} |f(z)|^\gamma$, following the proof in the case of the unit sphere [12, Thm. 5.2.2].

Let $\theta = \max\{\frac{\delta}{n}, \mathbf{d}(x, y)\}$. Then $B(y, \frac{\delta}{n}) \subset B(x, 2\theta) \subset B(y, 3\theta)$. By the doubling condition for w and (ii) of Lemma 2.1,

$$w(B(y, \frac{\delta}{n})) \geq \frac{1}{c_{L(w)}} \left(\frac{3\theta n}{\delta} \right)^{-\alpha(w)} w(B(y, 3\theta)) \geq \frac{1}{c_{L(w)}} \left(\frac{3\theta n}{\delta} \right)^{-\alpha(w)} w(B(x, 2\theta)),$$

which implies, together with $n\theta \leq 1 + nd(x, y)$, the estimate

$$\begin{aligned} \min_{z \in B(y, \frac{\delta}{n})} |f(z)|^\gamma &\leq \frac{1}{w(B(y, \frac{\delta}{n}))} \int_{B(y, \frac{\delta}{n})} |f(z)|^\gamma w(z) \, dm(z) \\ &\leq c_{L(w)} \left(\frac{3\theta n}{\delta} \right)^{\alpha(w)} \frac{1}{w(B(x, 2\theta))} \int_{B(x, 2\theta)} |f(z)|^\gamma w(z) \, dm(z) \\ &\leq c_{L(w)} \left(\frac{3}{\delta} \right)^{\alpha(w)} (1 + nd(x, y))^{\alpha(w)} M_w(|f|^\gamma)(x). \end{aligned}$$

Together with (2.31), this completes the proof. \square

Let $\|\cdot\|_{p,w}$ denote the norm of $L^p(\Omega, w)$ and we adopt this notation for $0 < p < 1$, even though it is no longer a norm. The L^p boundedness of $f_{\beta,n}^*$ follows from that of $M_\mu(|f|^\gamma)$ for $\gamma p > 1$ or $\beta > \alpha(w)/p$.

Corollary 2.12. *If $0 < p \leq \infty$, $f \in \Pi_n(\Omega)$ and $\beta > \alpha(w)/p$, then*

$$\|f\|_{p,w} \leq \|f_{\beta,n}^*\|_{p,w} \leq c \|f\|_{p,w},$$

where c depends also on $L(w)$ and β when β is either large or close to $\alpha(w)/p$.

2.4. Marcinkiewicz-Zygmund inequality. These inequalities are between the L^p norm and a discrete L^p norm defined with respect to a well-distributed set of points. We start with the definition of well-separated sets of points.

Definition 2.13. *Let Ξ be a discrete set in Ω .*

- (a) *A finite collection of subsets $\{S_z : z \in \Xi\}$ is called a partition of Ω if $S_z^\circ \cap S_y^\circ = \emptyset$ when $z \neq y$ and $\Omega = \bigcup_{z \in \Xi} S_z$.*
- (b) *Let $\varepsilon > 0$. A discrete subset Ξ of Ω is called ε -separated if $d(x, y) \geq \varepsilon$ for every two distinct points $x, y \in \Xi$.*
- (c) *Ξ is called maximal if there is a constant $c_d > 1$ such that*

$$(2.32) \quad 1 \leq \sum_{z \in \Xi} \chi_{B(z, \varepsilon)}(x) \leq c_d, \quad \forall x \in \Omega,$$

where χ_E denotes the characteristic function of the set E .

Evidently (2.32) implies $\Omega = \bigcup_{z \in \Xi} B(z, \varepsilon)$. Moreover, it implies that the cardinality $\#\Xi$ of Ξ satisfies

$$c_1 \frac{m(\Omega)}{\max_{z \in \Xi} m(B(z, \varepsilon))} \leq \#\Xi \leq c_2 \frac{m(\Omega)}{\min_{z \in \Xi} m(B(z, \varepsilon))}.$$

For the unit sphere, all balls of the same radius have the same volume; that is, $\sigma(B(x, r)) = \sigma(B(0, 1))r^{d-1}$ for all $x \in \mathbb{S}^{d-1}$, where σ is the surface measure on \mathbb{S}^{d-1} . This allows us to deduce (2.32) from $\Omega = \bigcup_{x \in \Xi} B(x, \varepsilon)$ and the cardinality of the maximal ε -separated set $\Xi_{\mathbb{S}}$ on the sphere \mathbb{S}^{d-1} satisfies $c'_d \varepsilon^{-d+1} \leq \#\Xi_{\mathbb{S}} \leq c_d \varepsilon^{-d+1}$; see, for example, [12, p. 114].

As we shall show in the next section that the volume of the ball $B(x, r)$ on the conic surface depends on the position of x . In particular, we no longer have $\varpi(B(x, \varepsilon)) \sim \varpi(B(y, \varepsilon))$ for all x, y in the domain in general.

Theorem 2.14. *Let w be a doubling weight on Ω . Let $\varepsilon = \frac{\delta}{n}$ for $n = 1, 2, \dots$ and $\delta \in (0, 1)$. If Ξ is a maximal ε -separated subset of Ω , then for all $f \in \Pi_n(\Omega)$ and $0 < p < \infty$,*

$$(2.33) \quad \left(\sum_{y \in \Xi} |\text{osc}(f)(y, \frac{\delta}{n})|^p w(B(y, \frac{\delta}{n})) \right)^{\frac{1}{p}} \leq c_p \delta \|f\|_{p,w},$$

where c_p depends on p , when p is close to 0, and on d and $L(w)$.

Proof. If $y \in B(x, \frac{\delta}{n})$, then $f_{2\alpha(w)/p,n}^*(y) \sim f_{2\alpha(w)/p,n}^*(x)$. Hence, it follows from Lemma 2.10 that

$$\begin{aligned} \sum_{y \in \Xi} |\text{osc}(f)(y, \frac{\delta}{n})|^p w(B(y, \frac{\delta}{n})) &\leq c(c_p \delta)^p \sum_{y \in \Xi} \int_{B(y, \frac{\delta}{n})} \left(f_{2\alpha(w)/p,n}^*(x) \right)^p w(x) dm(x) \\ &\leq (c_p \delta)^p \int_{\Omega} \left(f_{2\alpha(w)/p,n}^*(x) \right)^p w(x) dm(x). \end{aligned}$$

The last integral is bounded by $\|f\|_{p,w}^p$ by Corollary 2.12. \square

We now prove the Marcinkiewicz-Zygmund inequality for polynomials.

Theorem 2.15. *Let w be a doubling weight on Ω . Let Ξ be a maximal $\frac{\delta}{n}$ -separated subset of Ω and let $\delta > 0$ and $\delta \leq 1$.*

(i) *For all $0 < p < \infty$ and $f \in \Pi_m(\Omega)$ with $n \leq m \leq cn$,*

$$(2.34) \quad \left(\sum_{z \in \Xi} \left(\max_{x \in B(z, \frac{\delta}{n})} |f(x)|^p \right) w(B(z, \frac{\delta}{n})) \right)^{\frac{1}{p}} \leq c_{w,p} \|f\|_{p,w},$$

where $c_{w,p}$ depends on $L(w)$ and on p when p is close to 0.

(ii) *For $0 < r < 1$, there is a $\delta_r > 0$ such that for $\delta \leq \delta_r$, $r \leq p < \infty$ and $f \in \Pi_n(\Omega)$,*

$$(2.35) \quad \|f\|_{p,w} \leq c_{w,r} \left(\sum_{z \in \Xi} \left(\min_{x \in B(z, \frac{\delta}{n})} |f(x)|^p \right) w(B(z, \frac{\delta}{n})) \right)^{\frac{1}{p}},$$

where $c_{w,r}$ depends only on $L(w)$ and on r when r is close to 0. Furthermore,

$$(2.36) \quad \|f\|_{\infty} \leq c \max_{z \in \Xi} |f(z)|.$$

Proof. For every $y \in \Xi$, choose $z_y \in B(y, \frac{\delta}{n})$ such that $|f(z_y)| = \sup_{x \in B(y, \frac{\delta}{n})} |f(x)|$. For $f \in \Pi_m(\Omega)$, we have $|f(z_y)| \leq f_{\beta,m}^*(z_y) \leq c f_{\beta,n}^*(y)$. Hence,

$$\begin{aligned} \sum_{z \in \Xi} \left(\max_{x \in B(z, \frac{\delta}{n})} |f(x)|^p \right) w(B(z, \frac{\delta}{n})) &\leq c \sum_{z \in \Xi} (f_{\beta,n}^*(y))^p w(B(z, \frac{\delta}{n})) \\ &\leq c \int_{\Omega} (f_{\beta,n}^*(x))^p w(x) dm(x), \end{aligned}$$

from which (i) follows from the bound in Corollary 2.12. We now prove (ii). Since Ξ is maximal $\frac{\delta}{n}$ -separated,

$$\begin{aligned} \|f\|_{p,w}^p &\leq \sum_{z \in \Xi} \int_{B(z, \frac{\delta}{n})} |f(x)|^p w(x) dm(x) \\ &\leq 2^p \sum_{z \in \Xi} |\text{osc}(f)(z, \frac{\delta}{n})|^p w(B(z, \frac{\delta}{n})) + 2^p \sum_{z \in \Xi} \left(\min_{x \in B(z, \frac{\delta}{n})} |f(x)|^p \right) w(B(z, \frac{\delta}{n})). \end{aligned}$$

The first term in the right-hand side is bounded by $(2c_r\delta)^p \|f\|_{p,w}^p$ by Lemma 2.10. Hence, choosing $\delta_r = 1/(4c_r)$ so that $(2c_r\delta)^p \leq 2^{-p}$ for $\delta \leq \delta_r$, we conclude that

$$(1 - 2^{-p}) \|f\|_{p,w}^p \leq 2^p \sum_{z \in \Xi} \left(\min_{x \in B(z, \frac{\delta}{n})} |f(x)|^p \right) w(B(z, \frac{\delta}{n})).$$

Taking the power of $1/p$ proves (2.35). In particular, since the constant in (2.35) is independent of p , it readily implies (2.36). This completes the proof. \square

For the unit sphere without weight and the unit ball with the classical weight, the MZ inequality was established in [5, 33, 34, 38]. For the unit sphere with the doubling weight, it was established in [7].

2.5. Christoffel function for doubling weight. Let w be a doubling weight on Ω . The Christoffel function $\lambda_n(w; \cdot)$ is defined by

$$(2.37) \quad \lambda_n(w; x) := \inf_{\substack{g(x)=1 \\ g \in \Pi_n(\Omega)}} \int_{\Omega} |g(x)|^2 w(x) dm(x).$$

It is known that the Christoffel function is closely related to the kernel $K_n(w; \cdot, \cdot)$ of the partial sum operator

$$K_n(w; x, y) = \sum_{k=0}^n P_k(w; x, y), \quad x, y \in \Omega.$$

More precisely, it is known [14, Theorem 3.6.6.] that, for $n = 0, 1, 2, \dots$,

$$(2.38) \quad \lambda_n(w; x) = \frac{1}{K_n(w; x, x)}, \quad x \in \Omega.$$

The Christoffel function encodes essential information on weighted approximation by polynomials on the domain Ω . It will also be useful in the study of cubature rules in the next subsection, for which we need an upper bound of $\lambda_n(w; x)$. We establish this bound for all doubling weights on the localizable homogeneous space under the assumption that there exist fast decaying polynomials on Ω . More precisely, we make the following assertion.

Assertion 4. *Let Ω be compact. For each $x \in \Omega$, there is a nonnegative polynomial T_x of degree at most n that satisfies*

- (1) $T_x(x) = 1$, $T_x(y) \geq \delta > 0$ for $y \in B(x, \frac{1}{n})$ for some δ independent of n , and, for each $\gamma > 1$,

$$0 \leq T_x(y) \leq c_\gamma (1 + nd(x, y))^{-\gamma}, \quad y \in \Omega;$$

- (2) *there is a polynomial q_n such that $q_n(x)T_x(y)$ is a polynomial of degree at most rn , for some positive integer r , in x -variable and $c_1 \leq q_n(x) \leq c_2$ for $x \in \Omega$ for some positive numbers c_1 and c_2 .*

The fast decaying polynomials of one variable are studied in [30], see also [43], which leads to fast decaying polynomials on the unit sphere by the addition formula (2.22). For the unit ball, such polynomials are constructed in [38]. We use these polynomials to derive an upper bound for the Christoffel function. First, we need a lemma.

Lemma 2.16. *Assume Assertion 4. Let $\alpha > 0$ be a positive number and let $x \in \Omega$ be fixed. For a doubling weight \mathbf{w} on Ω , there is a polynomial Q_x of degree n such that, for all $y \in \Omega$,*

$$(2.39) \quad c_1(1 + nd(x, y))^\alpha \mathbf{w}(B(y, \frac{1}{n})) \leq Q_x(y) \leq c_2(1 + nd(x, y))^\alpha \mathbf{w}(B(y, \frac{1}{n})),$$

where c_1 and c_2 are positive constant independent of n , x and y .

Proof. Let T_x and q be polynomials in Assertion 4 of degree at most $\lfloor \frac{n}{2} \rfloor$. For fixed x , we define Q_x by

$$Q_x(y) = q_n(y) \int_{\Omega} T_y(v) (1 + nd(x, v))^\alpha \mathbf{w}(v) dm(v), \quad y \in \Omega.$$

Since $q_n(y)T_y(v)$ is a polynomial of degree at most n in y , it is easy to see that Q_x is a polynomial of degree at most n . Using Assertion 4 with $\gamma > \alpha + 1$ and using the triangle inequality $1 + nd(x, v) \leq (1 + nd(x, y))(1 + nd(y, v))$, we obtain

$$\begin{aligned} Q_x(y) &\leq c(1 + nd(x, y))^\alpha \int_{\Omega} \frac{1}{(1 + nd(y, v))^{\gamma-\alpha}} \mathbf{w}(v) dm(v) \\ &\leq c(1 + nd(x, y))^\alpha \mathbf{w}(B(y, \frac{1}{n})), \end{aligned}$$

since, by the doubling property of \mathbf{m} and choosing γ sufficiently large,

$$\begin{aligned} (2.40) \quad \int_{\Omega} \frac{\mathbf{w}(v) dm(v)}{(1 + nd(y, v))^{\gamma-\alpha}} &\leq \sum_{k=1}^{\infty} \int_{B(y, \frac{2^k}{n}) \setminus B(y, \frac{2^{k-1}}{n})} \frac{\mathbf{w}(v) dm(v)}{(1 + nd(y, v))^{\gamma-\alpha}} \\ &\leq c \sum_{k=1}^{\infty} \frac{1}{(1 + 2^{k-1})^{\gamma-\alpha}} \mathbf{w}(B(y, \frac{2^k}{n})) \\ &\leq c \mathbf{w}(B(y, \frac{1}{n})) \sum_{k=1}^{\infty} \frac{L(\mathbf{w})^{k-1}}{(1 + 2^{k-1})^{\gamma-\alpha}} \\ &\leq c \mathbf{w}(B(y, \frac{1}{n})). \end{aligned}$$

This establishes the upper bound of Q_x . In the other direction, we use $q_n(y) \geq c_1$, $T_y(v) \geq \delta > 0$ for $v \in B(y, \frac{1}{n})$ and $T_y(v) \geq 0$ to obtain

$$\begin{aligned} Q_x(y) &\geq (1 + nd(x, y))^\alpha \int_{\Omega} T_y(v) (1 + nd(y, v))^{-\alpha} \mathbf{w}(v) dm(v) \\ &\geq c\delta(1 + nd(x, y))^\alpha \int_{B(y, \frac{1}{n})} (1 + nd(y, v))^{-\alpha} \mathbf{w}(v) dm(v) \\ &\geq c\delta(1 + nd(x, y))^\alpha \mathbf{w}(B(y, \frac{1}{n})), \end{aligned}$$

since the last integral is trivially bounded below by $c\mathbf{w}(B(y, \frac{1}{n}))$. This completes the proof. \square

Proposition 2.17. *Assume Assertion 4 holds. Let \mathbf{w} be a doubling weight on the domain Ω . Then*

$$\lambda_n(\mathbf{w}; x) \leq c \mathbf{w}(B(x, \frac{1}{n})), \quad x \in \Omega.$$

Proof. Let $m = \lfloor \frac{n}{3} \rfloor$. Let $T_x(y)$ be the polynomial of degree m in Assertion 4. Let Ξ be a maximal $\frac{\delta}{n}$ -separated set in Ω with $\delta \leq \delta_0$ as in Theorem 2.15. By the inequality

(2.35) with $p = 1$ and (ii) of Lemma 2.1 for both \mathbf{m} and \mathbf{w} , we obtain

$$\begin{aligned} \int_{\Omega} [T_x(y)]^2 \mathbf{w}(y) d\mathbf{m}(y) &\leq c \sum_{z \in \Xi} [T_x(z)]^2 \mathbf{w}(B(z, \frac{\delta}{n})) \\ &\leq c \frac{\mathbf{w}(B(x, \frac{\delta}{n}))}{[\mathbf{m}(B(x, \frac{\delta}{n}))]^2} \sum_{z \in \Xi} [T_x(z)]^2 (1 + nd(x, z))^\alpha [\mathbf{m}(B(z, \frac{\delta}{n}))]^2, \end{aligned}$$

where $\alpha = \alpha(\mathbf{w}) + 2\alpha(\mathbf{m})$. Let Q_x be the polynomial of degree m that satisfies (2.39) with \mathbf{m} in place of \mathbf{w} . Using the lower bound of Q_x in (2.39) and applying the inequality (2.34) with $p = 1$ on the polynomial $[T_x(z)]^2 Q_y(z)$ of degree n with respect to the measure $d\mathbf{m}$, we obtain

$$\begin{aligned} \int_{\Omega} [T_x(y)]^2 \mathbf{w}(y) d\mathbf{m}(y) &\leq c \frac{\mathbf{w}(B(x, \frac{\delta}{n}))}{[\mathbf{m}(B(x, \frac{\delta}{n}))]^2} \sum_{z \in \Xi} [T_x(z)]^2 Q_x(z) \mathbf{m}(B(z, \frac{\delta}{n})) \\ &\leq c \frac{\mathbf{w}(B(x, \frac{\delta}{n}))}{[\mathbf{m}(B(x, \frac{\delta}{n}))]^2} \int_{\Omega} [T_x(v)]^2 Q_x(v) d\mathbf{m}(v). \end{aligned}$$

Using now the upper bound of Q_x in (2.39), again with \mathbf{m} in place of \mathbf{w} , and the upper bound of $T_x(v)$ in Assertion 4, and (ii) of Lemma 2.1 for the measure $d\mathbf{m}$, we further deduce that the last integral is bounded by

$$\begin{aligned} \int_{\Omega} [T_x(v)]^2 Q_x(v) d\mathbf{m}(v) &\leq c \int_{\Omega} \frac{\mathbf{m}(B(v, \frac{\delta}{n}))}{(1 + nd(x, v))^{2\gamma - \alpha}} d\mathbf{m}(v) \\ &\leq c \mathbf{m}(B(x, \frac{\delta}{n})) \int_{\Omega} \frac{1}{(1 + nd(x, z))^{2\gamma - \alpha - \alpha(\mathbf{m})}} d\mathbf{m}(v) \\ &\leq c [\mathbf{m}(B(x, \frac{\delta}{n}))]^2, \end{aligned}$$

where the last step follows from (2.40). Together, from the last two displayed inequalities follows that

$$\int_{\Omega} [T_x(y)]^2 \mathbf{w}(y) d\mathbf{m}(y) \leq c \mathbf{w}(B(x, \frac{\delta}{n})).$$

Since $T_x(x) = 1$, this completes the proof by (2.37). \square

Proposition 2.18. *If \mathbf{w} is a weight function for which Assertion 1 holds, then*

$$\lambda_n(\mathbf{w}; x) \geq c \mathbf{w}(B(x, \frac{1}{n})), \quad x \in \Omega.$$

Proof. Let $L_n(\mathbf{w}; \cdot, \cdot)$ be the localized kernel defined with a cut-off function of type (a). It follows immediately that $K_n(\mathbf{w}; x, x) \leq L_n(\mathbf{w}; x, x)$. Hence, if \mathbf{w} admits Assertion 1, then

$$K_n(\mathbf{w}; x, x) \leq L_n(\mathbf{w}; x, x) \leq \frac{c}{\mathbf{w}(B(x, \frac{1}{n}))},$$

which is equivalent to the lower bound $\lambda_n(\mathbf{w}; x)$ by (2.38). \square

Together, the last two propositions give both upper and lower bounds for the Christoffel functions. Such bounds are known for regular domains, see for example [14, 23, 50] and the references therein, and it has been established recently for fairly general convex domains [22, 40] and planar domains with piecewise boundary [41]. None of the previous results, however, apply to conic domains that will come out as special cases of the above theorem.

2.6. Positive cubature rules. Let \mathbf{w} be a doubling weight on Ω . A cubature rule of degree n for \mathbf{w} is a finite linear combination of point evaluations that satisfies

$$\int_{\Omega} f(x) \mathbf{w}(x) d\mathbf{m}(x) = \sum_{k=1}^N \lambda_k f(x_k), \quad \forall f \in \Pi_n(\Omega),$$

where $\lambda_k \in \mathbb{R}$ and $x_k \in \Omega$. When $\lambda_k > 0$ for all k , the cubature rule is called positive. Given a maximal ε -separated subset Ξ on Ω , we establish the existence of a positive cubature rule based on the points in Ξ , which plays an essential role for discretizing our near best approximation $L_n * f$.

The proof will use the Farkas lemma (cf. [12, Lemma 6.3.2]) stated below.

Lemma 2.19. *Let V be a finite dimensional real Hilbert space with inner product $\langle \cdot, \cdot \rangle$. Then for any points a^1, a^2, \dots, a^m and $\zeta \in V$, exactly of the following two systems has a solution:*

- (1) $\sum_{j=1}^m \mu_j a^j = \zeta, \quad 0 \leq \mu_1, \mu_2, \dots, \mu_m \in \mathbb{R}.$
- (2) $\langle a^j, x \rangle \geq 0, j = 1, 2, \dots, m$ and $\langle \zeta, x \rangle < 0$ for some $x \in V$.

We use the lemma to prove the existence of the positive cubature rules. For the unit sphere and the unit ball, this was established in [33] and later in [5, 34, 38]. We follow the proof in [12, Theorem 6.3.3] for the doubling weight on the unit sphere.

Theorem 2.20. *Let \mathbf{w} be a doubling weight on Ω . Let Ξ be a maximum $\frac{\delta}{n}$ -separated subset of Ω . There is a $\delta_0 > 0$ such that for $0 < \delta < \delta_0$ there exist positive numbers $\lambda_z, z \in \Xi$, so that*

$$(2.41) \quad \int_{\Omega} f(x) \mathbf{w}(x) d\mathbf{m}(x) = \sum_{z \in \Xi} \lambda_z f(z), \quad \forall f \in \Pi_n(\Omega).$$

Moreover, $\lambda_z \geq c_1 \mathbf{w}(B(z, \frac{\delta}{n}))$ and, if Assertion 4 holds, then $\lambda_z \sim \mathbf{w}(B(z, \frac{\delta}{n}))$.

Proof. Let ϖ be the weight function that admits highly localized kernels on Ω . We use Lemma 2.19 with the inner product defined by $\langle \cdot, \cdot \rangle_{\varpi}$. Recall that $K_n(\varpi; \cdot, \cdot)$ denotes the reproducing kernel of $\Pi_n(\Omega)$. Let $\{z_k : 1 \leq k \leq N\}$ be an enumeration of Ξ . We define the functions a^j and ζ by

$$\begin{aligned} a^j(x) &= K_n(\varpi; x, z_j), \quad 1 \leq j \leq N, \\ \zeta(x) &= \int_{\Omega} K_n(\varpi; x, y) \mathbf{w}(y) d\mathbf{m}(y) - \frac{1}{2c_d} \sum_{j=1}^N K_n(\varpi; x, z_j) \mathbf{w}(B(z_j, \frac{\delta}{n})), \end{aligned}$$

where c_d is the constant in (2.32). For every $f \in \Pi_n(\Omega)$, the reproducing property of $K_n(\varpi; \cdot, \cdot)$ implies that

$$\begin{aligned} \langle f, a^j \rangle_{\varpi} &= f(z_j), \quad 1 \leq j \leq N, \\ \langle f, \zeta \rangle_{\varpi} &= \int_{\Omega} f(y) \mathbf{w}(y) d\mathbf{m}(y) - \frac{1}{2c_0} \sum_{j=1}^N f(z_j) \mathbf{w}(B(z_j, \frac{\delta}{n})). \end{aligned}$$

Assume $\langle f, a^j \rangle_{\varpi} \geq 0$ for all j , so that f is nonnegative on Ξ . Then, by (2.32),

$$\begin{aligned} \int_{\Omega} f(x) \mathbf{w}(x) d\mathbf{m}(x) &= \sum_{j=1}^N \int_{B(z_j, \frac{\delta}{n})} f(x) \frac{\mathbf{w}(x)}{\sum_{z \in \Xi} \chi_{B(y, \frac{\delta}{n})}(x)} d\mathbf{m}(x) \\ &\geq \frac{1}{c_d} \sum_{j=1}^N f(z_j) \int_{B(z_j, \frac{\delta}{n})} \mathbf{w}(x) d\mathbf{m}(x) - \sum_{j=1}^N \int_{B(z_j, \frac{\delta}{n})} \text{osc}(f)(x, \frac{\delta}{n}) \mathbf{w}(x) d\mathbf{m}(x) \\ &\geq \frac{1}{c_d} \sum_{j=1}^N f(z_j) \mathbf{w}(B(z_j, \frac{\delta}{n})) - c_1 \delta \int_{\Omega} |f(x)| \mathbf{w}(x) d\mathbf{m}(x) \end{aligned}$$

by (2.33). Hence, using the Marcinkiewicz-Zygmund inequality in (ii) of Theorem 2.15 with $p = 1$, we conclude that

$$\langle f, \zeta \rangle_{\varpi} \geq ((2c_d)^{-1} - c_{\mathbf{w},1} c_1 \delta) \sum_{j=1}^N f(z_j) \mathbf{w}(B(z_j, \frac{\delta}{n})) \geq 0$$

if we choose $\delta \leq (2c_d)^{-1}/(c_{\mathbf{w},1} c_1)$. Consequently, (2) of Farkas lemma does not hold so that (1) must hold, which shows that there exist $\mu_j \geq 0$ such that

$$\int_{\Omega} f(x) \mathbf{w}(x) d\mathbf{m}(x) - \frac{1}{2c_0} \sum_{j=1}^N f(z_j) \mathbf{w}(B(z_j, \frac{\delta}{n})) = \sum_{j=1}^N \mu_j f(z_j).$$

This establishes the cubature rule (2.41) with $\lambda_{z_j} = (2c_0)^{-1} \mathbf{w}(B(z_j, \frac{\delta}{n})) + \mu_j$.

It follows immediately that $\lambda_z \geq c \mathbf{w}(B(z, \frac{\delta}{n}))$ for all $z \in \Xi$. To prove the upper bound of this inequality, we set $m = \lfloor \frac{n}{2} \rfloor$ and use the reproducing kernel $K_m(\mathbf{w}; \cdot, \cdot)$ of $\Pi_n(\Omega)$ with respect to \mathbf{w} . Since $K_m(\mathbf{w}; z, z) > 0$, we define a polynomial

$$q_z(x) = \frac{K_m(\mathbf{w}; x, z)}{K_m(\mathbf{w}; z, z)}, \quad z \in \Xi.$$

For each z , the polynomial q_z^2 is a polynomial of degree at most n and $q_z(z) = 1$. Hence, by (2.41),

$$\lambda_z \leq \sum_{y \in \Xi} \lambda_y [q_z(y)]^2 = \int_{\Omega} [q_z(x)]^2 \mathbf{w}(x) d\mathbf{m}(x) = \int_{\Omega} \frac{K_m(\mathbf{w}; x, z)^2}{K_m(\mathbf{w}; z, z)^2} \mathbf{w}(x) d\mathbf{m}(x).$$

Hence, using the reproducing property of the kernel $K_m(\mathbf{w}; \cdot, \cdot)$,

$$\int_{\Omega} K_m(\mathbf{w}; x, z)^2 \mathbf{w}(x) d\mathbf{m}(x) = K_m(\mathbf{w}; z, z),$$

we conclude that $\lambda_z \leq c[K_m(\mathbf{w}; z, z)]^{-1} = \lambda_n(\mathbf{w}; z)$. Hence, $\lambda_z \leq c \mathbf{w}(B(z, \frac{\delta}{n}))$ by Proposition 2.17. \square

2.7. Tight polynomial frames. The highly localized kernels are powerful tools when they exist. We use them to study approximation behavior of the integral operator $L_n(\varpi) * f$ that has $L_n(\varpi; \cdot, \cdot)$ as its kernel, or more precisely,

$$(2.42) \quad L_n(\varpi) * f(x) = \int_{\Omega} f(y) L_n(\varpi; x, y) \varpi(y) d\mathbf{m}(y).$$

The fast decaying of the kernel indicates that $L_n(\varpi) * f$ provides a good approximation to the function f . Its approximation property will be studied in the next section. For

now, we observe that $L_n(\varpi) * f$ is a polynomial of degree at most $2n$ by the property of the cut-off function.

We now use the operator $L_n(\varpi) * f$ and the positive cubature rules based on the discrete set Ξ to construct tight polynomial frames in $L^2(\Omega, \varpi)$. The construction is fairly standard by now, so is the proof of the tight frame. See, for example, [12, 34, 37, 38] for the case of the interval, the unit sphere and the unit ball. We recall the procedure and will be brief in proof.

Let (Ω, ϖ, d) be a localizable homogeneous space. For the highly localized kernels $L_n(\varpi; x, y)$, we assume its cut-off function \hat{a} is of type (b) and satisfies

$$(2.43) \quad \begin{aligned} \hat{a}(t) &\geq \rho > 0, & \text{if } t \in [3/5, 5/3], \\ [\hat{a}(t)]^2 + [\hat{a}(2t)]^2 &= 1, & \text{if } t \in [1/2, 1]. \end{aligned}$$

If g is a nonnegative even function in $C^\infty(\mathbb{R})$, so that $\text{supp } g = [-1, 1]$, $g(0) = 1$ and $|g(t)|^2 + |g(t+1)|^2 = 1$ on $[-1, 0]$, then $\hat{a}(t) := g(\log_2 t)$ has the desired properties. The last assumption on \hat{a} implies

$$(2.44) \quad \sum_{j=0}^{\infty} \left[\hat{a}\left(\frac{t}{2^j}\right) \right]^2 = 1, \quad t \in [1, \infty).$$

Let w be the doubling weight that admits Assertion 4. Then the cubature rules in Theorem 2.20 are well established for w . For $j = 0, 1, \dots$, let $\varepsilon_j = \frac{\delta}{2^j}$ and let Ξ_j be a maximal ε_j -separated subset in Ω , where δ is chosen so that the cubature rule in Theorem 2.20 holds; that is, there are $\lambda_{z,j} > 0$ for $z \in \Xi_j$ such that

$$(2.45) \quad \int_{\Omega} f(x) w(x) dm(x) = \sum_{z \in \Xi_j} \Xi_{z,j} f(z), \quad f \in \Pi_{2^j}(\Omega);$$

moreover, $\lambda_{z,j} \sim w(B(z, 2^{-j}))$ since we assume that Assertion 4 holds for w . In terms of the kernel $L_n(w; \cdot, \cdot)$ defined via the cut-off function \hat{a} , we define

$$F_0(x, z) := 1, \quad \text{and} \quad F_j(x, z) := L_{2^{j-1}}(w; x, z), \quad j = 1, 2, 3, \dots$$

Accordingly, we define $F_j * f = L_{2^{j-1}} * f$, that is,

$$F_j * f(x) := \int_{\Omega} f(y) F_j(x, y) w(y) dm(y), \quad j = 0, 1, 2, \dots$$

Now, for $z \in \Xi_j$ and $j = 1, 2, \dots$, we define our frame elements by

$$\psi_{z,j}(x) := \sqrt{\lambda_{z,j}} F_j(x, z).$$

Then $\Psi := \{\psi_{z,j} : z \in \Xi_j, 1 \leq j \leq \infty\}$ is a frame system.

By the orthogonality of the reproducing kernel, it follows readily that

$$F_j * F_j * f = \sum_{k=1}^{2^j} \left| \hat{a}\left(\frac{k}{2^{j-1}}\right) \right|^2 \text{proj}_k(w; f).$$

Hence, the following semi-discrete Calderón type decomposition follows from (2.44),

$$(2.46) \quad f = \sum_{k=0}^{\infty} \text{proj}_k(w; f) = \sum_{j=0}^{\infty} F_j * F_j * f, \quad f \in L^2(\Omega, w).$$

Furthermore, the system Ψ is a tight frame in $L^2(\Omega, w)$ norm.

Theorem 2.21. *Assume that Ω admits a localizable homogenous space. Let \mathbf{w} be a doubling weight satisfying Assertion 4. If $f \in L^2(\Omega, \mathbf{w})$, then*

$$(2.47) \quad f = \sum_{j=0}^{\infty} \sum_{z \in \Xi_j} \langle f, \psi_{z,j} \rangle_{\mathbf{w}} \psi_{z,j} \quad \text{in } L^2(\Omega, \mathbf{w})$$

and

$$(2.48) \quad \|f\|_{2,\mathbf{w}} = \left(\sum_{j=0}^{\infty} \sum_{z \in \Xi_j} |\langle f, \psi_{z,j} \rangle_{\mathbf{w}}|^2 \right)^{1/2}.$$

Proof. Since $F_j(x, \cdot)F_j(\cdot, y)$ is a polynomial of degree 2^j , we can apply the cubature rule (2.45) to discretize $F_j * F_j$, which gives

$$F_j * F_j(x, y) = \int_{\Omega} F_j(x, z) F_j(z, y) \mathbf{w}(z) \mathrm{d}\mathbf{m}(z) = \sum_{z \in \Xi_j} \psi_{z,j}(x) \psi_{z,j}(y),$$

from which the identity (2.47) follows from (2.46) right away. Furthermore, let $S_R f$ be the sum $S_R f = \sum_{j=0}^R \sum_{z \in \Xi_j} \langle f, \psi_{z,j} \rangle_{\mathbf{w}} \psi_{z,j}$. Then, it follows immediately

$$\langle f, S_R f \rangle_{\mathbf{w}} = \sum_{j=0}^R \sum_{z \in \Xi_j} |\langle f, \psi_{z,j} \rangle_{\mathbf{w}}|^2.$$

Taking the limit $R \rightarrow \infty$ we obtain the tight frame identity (2.48). \square

For the weight function ϖ that admits the highly localized kernels, the frame element $\psi_{z,j}$ has near exponential rate of decay away from its center with respect to the distance $\mathbf{d}(\cdot, \cdot)$ on Ω .

Theorem 2.22. *Let $(\Omega, \varpi, \mathbf{d})$ be a localizable homogeneous space. There, there is a constant $c_{\kappa} > 0$ depending only on κ, d, ϖ and $\hat{\mathbf{a}}$ such that for $z \in \Xi_j, j = 0, 1, \dots$,*

$$(2.49) \quad |\psi_{z,j}(x)| \leq c_{\kappa} \frac{1}{\sqrt{\varpi(B(z, 2^{-j}))}(1 + 2^j \mathbf{d}(x, z))^{\kappa}}, \quad x \in \Omega.$$

This follows readily from the fast decaying of the kernel and $\lambda_{z,j} \sim \mathbf{w}(B(z, 2^{-j}))$. It shows that the frame Ψ is highly localized, which makes it a powerful tool for various applications, such as decompositions of functions spaces and computational harmonic analysis, on more specific domains.

3. POLYNOMIAL APPROXIMATION ON HOMOGENEOUS SPACES

We consider approximation by polynomials in the space $L^p(\Omega, \varpi)$ when the weight function ϖ posses two additional properties that are analogs of two characteristics of spherical harmonics: addition formula for the reproducing kernels and a differential operator that has orthogonal polynomials as eigenfunctions. With appropriate weight functions, these properties are shared by orthogonal polynomials on the unit sphere, on the unit ball, and on the simplex. The addition formula premises that the reproducing kernels possess a one-dimensional character, which leads to a convolution structure that allows us to reduce much of the study of the Fourier orthogonal series to that of the Fourier-Jacobi series of one variable. The differential operator that has orthogonal polynomials as eigenfunctions leads to a K -functional. Together they provide us with tools for characterizing the best approximation by polynomials, which is a central problem in approximation theory. The goal of this section is to develop this framework,

which extends the results on the unit sphere and the unit ball and it is applicable on the conic domains.

The two characteristic properties are defined and discussed in the first subsection. The first one is used to define the convolution structure that leads to the definition of the modulus of smoothness in the second subsection. The second one is used to define a K -functional in the third subsection, where the main results on the characterization of the best polynomial approximation are stated and discussed. The near-best approximation operator, used to provide a direct estimate, is studied in the fourth subsection with the help of the differential operator, where some of the results are established with norms defined via a doubling weight. The Bernstein inequality, essential for proving the inverse estimate, is proved in the fifth subsection, together with the Nikolskii inequality. Finally, the proof of the main results is given in the seventh subsection.

3.1. Addition formula and differential operator. Let ϖ be a weight function defined on Ω . For studying approximation by polynomials, we require two more properties on the orthogonal structure of $L^2(\Omega, \varpi)$. These properties are analogs to the two characteristics of spherical harmonics on the unit sphere: the Laplace-Beltrami operator (2.24) and the addition formula (2.22), which serve as the quintessential examples in the study [53, 54] as well as in this subsection. These properties are shared by several other domains, including the unit ball, the simplex, and conic domains. Below we assume these properties and use them to carry out our subsequent study.

The first property that we assume is an analog of the Laplace-Beltrami operator. Recall that $\mathcal{V}_n(\Omega, \varpi)$ denote the space of orthogonal polynomials of degree n with respect to ϖ on Ω .

Definition 3.1. Let ϖ be a weight function on Ω . We denote by \mathfrak{D}_ϖ the second order derivation operator that has orthogonal polynomials with respect to ϖ as eigenfunctions; more precisely,

$$(3.1) \quad \mathfrak{D}_\varpi Y = -\mu(n)Y, \quad \forall Y \in \mathcal{V}_n(\Omega, \varpi),$$

where μ is a nonnegative quadratic polynomial.

For the unit sphere \mathbb{S}^{d-1} , the operator \mathfrak{D}_ϖ is the Laplace-Beltrami operator given by (2.24). For the unit ball \mathbb{B}^d , \mathfrak{D}_ϖ is the operator given by (2.29). In both cases, the operator is a differential operator. For the unit sphere and the unit ball with reflection invariant weight functions, \mathfrak{D}_ϖ is a differential-difference operator, called Dunkl Laplacian [13, 14]. For $d = 2$ and Ω is a domain with non-empty interior, the existence of a second order differential operator that satisfies (3.1) was characterized in [21]. There are essentially five cases with positive weight functions on regular domains but only two are compact domains: disk and triangle. The characterization for $d > 2$ remains an open problem. The operators for the conic domains are the recent additions, discovered in [53, 54] and will be recalled in later sections.

The second property that we assume is the addition formula, which gives a closed-form formula for the reproducing kernel $P_n(\varpi; \cdot, \cdot)$ of $\mathcal{V}_n(\Omega, \varpi)$.

Definition 3.2. Let ϖ be a weight function on Ω . The reproducing kernel $P_n(\varpi; \cdot, \cdot)$ is said to satisfy an addition formula if, for some $\alpha \geq \beta \geq -\frac{1}{2}$,

$$P_n(\varpi; x, y) = \int_{[-1,1]^m} Z_n^{(\alpha,\beta)}(\xi(x, y; u)) d\tau(u), \quad Z_n^{(\alpha,\beta)}(t) = \frac{P_n^{(\alpha,\beta)}(1)P_n^{(\alpha,\beta)}(t)}{h_n^{(\alpha,\beta)}},$$

where m is a positive integer; $\xi(x, y; u)$ is a function of $u \in [-1, 1]^m$, symmetric in x and y , and $\xi(x, y; u) \in [-1, 1]$; moreover, $d\tau$ is a probability measure on $[-1, 1]^m$, which can degenerate to have a finite support.

The classical addition formula is the one for the unit sphere \mathbb{S}^{d-1} , given in (2.22), which is the degenerate case, with $\alpha = \beta = \frac{d-3}{2}$. The one for the classical weight ϖ_μ on the unit ball \mathbb{B}^d is given by (2.27), which has $m = 1$ and $\alpha = \beta = \mu + \frac{d-2}{2}$. Similar formula holds for reflection invariant weight functions on the unit sphere and on the unit ball (cf. [14, p. 221 and p. 265]). Furthermore, for the simplex in \mathbb{R}^d (cf. [14, p. 275]) and the conic domains \mathbb{V}_0^{d+1} and \mathbb{V}^{d+1} , the polynomial $Z_n^{(\alpha, \beta)}$ is given by Z_{2n}^λ for some $\lambda \geq 0$, so that, by the quadratic transform (2.21), it is given by the Jacobi polynomial with $\alpha = \lambda - \frac{1}{2}$ and $\beta = -\frac{1}{2}$.

The addition formula premises a one-dimensional structure for the reproducing kernel. In all known cases, the estimates of the highly localized kernel $L_n(\varpi; \cdot, \cdot)$ are established with the help of this formula, since the addition formula implies

$$(3.2) \quad L_n(\varpi; x, y) = \int_{[-1, 1]^m} L_n^{(\alpha, \beta)}(\xi(x, y; u)) d\tau(u),$$

where the function $t \mapsto L_n^{(\alpha, \beta)}(t) = L_n^{(\alpha, \beta)}(t, 1)$ is the kernel defined in (2.16) for the Jacobi polynomials.

In the rest of this section, we assume that the orthogonal polynomials with respect to ϖ possess both the derivation operator and the addition formula. We do not, however, require that ϖ admits highly localized kernels that satisfy Assertions 1–3. In fact, some of our results hold if ϖ satisfies Assertions 1 and 3, but need not satisfy Assertion 2.

3.2. Convolution structure. Making use of the one-dimensional structure premised by the addition formula, we define a convolution operator on Ω .

Definition 3.3. Assume ϖ admits the addition formula. For $f \in L^1(\Omega, \varpi)$ and $g \in L^1([-1, 1], w_{\alpha, \beta})$, we define the convolution of f and g by

$$(f *_{\varpi} g)(x) := \int_{\Omega} f(y) T^{(\alpha, \beta)} g(x, y) \varpi(y) d\mathbf{m}(y),$$

where the operator $g \mapsto T^{(\alpha, \beta)} g$ is defined by

$$T^{(\alpha, \beta)} g(x, y) := \int_{[-1, 1]^m} g(\xi(x, y; u)) d\tau(u).$$

The addition formula shows that the projection operator $\text{proj}_n(\varpi; f)$ and the reproducing kernel $P_n(\varpi; \cdot, \cdot)$ satisfy

$$(3.3) \quad \text{proj}_n(\varpi; f, x) = f *_{\varpi} Z_n^{(\alpha, \beta)} \quad \text{and} \quad P_n(\varpi; x, y) = T^{(\alpha, \beta)} \left(Z_n^{(\alpha, \beta)} \right).$$

The operator $T^{(\alpha, \beta)}$ is defined in [54] with a more generic orthogonal polynomial in place of $P_n^{(\alpha, \beta)}$, where the following boundedness of the operator is established [54, Lemma 6.3].

Lemma 3.4. Let $g \in L^1([-1, 1], w_{\alpha, \beta})$. Then, for each $Q_n \in \mathcal{V}_n(\Omega, \varpi)$,

$$(3.4) \quad \int_{\Omega} T^{(\alpha, \beta)} g(x, y) Q_n(y) \varpi(y) dy = \Lambda_n^{(\alpha, \beta)}(g) Q_n(x),$$

where

$$\Lambda_n^{(\alpha,\beta)}(g) = c_{\alpha,\beta} \int_{-1}^1 g(t) R_n^{(\alpha,\beta)}(t) w_{\alpha,\beta}(t) dt \quad \text{with} \quad R_n^{(\alpha,\beta)}(t) := \frac{P_n^{(\alpha,\beta)}(t)}{P_n^{(\alpha,\beta)}(1)}$$

Furthermore, for $1 \leq p \leq \infty$ and $x \in \Omega$,

$$(3.5) \quad \left\| T^{(\alpha,\beta)} g(x, \cdot) \right\|_{L^p(\Omega, \varpi)} \leq \|g\|_{L^p([-1,1], w_{\alpha,\beta})}.$$

The convolution operator $f *_{\varpi} g$ is also defined in [54] and shown to satisfy the usual Young's inequality.

Theorem 3.5. *Let $p, q, r \geq 1$ and $p^{-1} = r^{-1} + q^{-1} - 1$. For $f \in L^q(\Omega, \varpi)$ and $g \in L^r([-1, 1]; w_{\alpha,\beta})$,*

$$(3.6) \quad \|f *_{\varpi} g\|_{L^p(\Omega, \varpi)} \leq \|f\|_{L^q(\Omega, \varpi)} \|g\|_{L^r([-1,1]; \varpi)}.$$

As an application of these results, we consider the Cesàro (C, δ) means $S_n^{\delta}(\varpi; f)$ of the Fourier orthogonal series with respect to ϖ . For $\delta > 0$, the operator $S_n^{\delta}(\varpi)$ is defined by

$$S_n^{\delta}(\varpi; f) := \frac{1}{\binom{n+\delta}{n}} \sum_{k=0}^n \binom{n-k+\delta}{n-k} \text{proj}_k(\varpi; f).$$

It can be written as an integral operator with the kernel $K_n^{\delta}(\varpi; \cdot, \cdot)$ being the (C, δ) mean of the reproducing kernel $P_n(\varpi; \cdot, \cdot)$. By the addition formula, the kernel $K_n^{\delta}(\varpi; \cdot, \cdot)$ can be written as

$$K_n^{\delta}(\varpi; x, y) = \int_{[-1,1]^m} k_n^{(\alpha,\beta),\delta}(\xi(x, y; u), 1) d\tau(u),$$

where $k_n^{(\alpha,\beta),\delta}(s, t)$ denotes the kernel of the (C, δ) mean of the Fourier-Jacobi series. In particular, using the positivity of the kernel $k_n^{(\alpha,\beta),\delta}[1]$ and the boundedness of the kernel [46, Theorem 9.1.3], we obtain the following theorem.

Theorem 3.6. *Let ϖ be a weight function that admits the addition formula. The Cesàro (C, δ) means of the Fourier orthogonal series with respect to ϖ satisfy*

- (1) *If $\delta \geq \alpha + \beta + 2$, then $S_n^{\delta}(\varpi; f)$ is a nonnegative operator;*
- (2) *If $\delta > \alpha + \frac{1}{2}$, then for $n = 0, 1, 2, \dots$,*

$$\|S_n^{\delta}(\varpi; f)\|_{p, \varpi} \leq \|f\|_{p, \varpi}, \quad 1 \leq p \leq \infty.$$

To further explore the one-dimensional structure that leads to the definition of the convolution, we define an operator $S_{\theta, \varpi} f$ as follows.

Definition 3.7. *Assume the addition formula holds for the weight function ϖ . For $0 \leq \theta \leq \pi$, the translation operator $S_{\theta, \varpi}$ is defined by*

$$(3.7) \quad \text{proj}_n(\varpi; S_{\theta, \varpi} f) = R_n^{(\alpha,\beta)}(\cos \theta) \text{proj}_n(\varpi; f), \quad n = 0, 1, 2, \dots$$

As a corollary of Theorem 3.6, a function $f \in L^1(\Omega, \varpi)$ is uniquely determined by its orthogonal projections $\text{proj}_n(\varpi; f)$, $n \geq 0$. Hence, the operator $S_{\theta, \varpi}$ is well defined for all $f \in L^1(\Omega, \varpi)$. For the unit sphere with the Lebesgue measure $d\sigma$, the operator S_{θ} is an integral operator given by [12, (2.16)]

$$S_{\theta, d\sigma} f(x) = \frac{1}{\omega_{d-1}(\sin \theta)^{d-1}} \int_{c(x, \theta)} f(y) d\ell(y),$$

where $c(x, \theta) = \{y : \langle x, y \rangle = \cos \theta\}$ is the spherical cap and $d\ell$ denotes the Lebesgue measure on the cap. For the unit ball \mathbb{B}^d with the classical weight W_μ , the operator S_{θ, W_μ} can also be written as an explicit integral operator [12, Theorem 11.2.6].

Proposition 3.8. *The operator $S_{\theta, \varpi}$ satisfies the following properties:*

(i) *For $f \in L^2(\Omega, \varpi)$ and $g \in L^1([-1, 1], w_{\alpha, \beta})$,*

$$(f *_{\varpi} g)(x) = c_{\alpha, \beta} \int_0^\pi S_{\theta, \varpi} f(x) g(\cos \theta) w_{\alpha, \beta}(\cos \theta) \sin \theta d\theta.$$

(ii) *$S_{\theta, \varpi} f$ preserves positivity; that is, $S_{\theta, \varpi} f \geq 0$ if $f \geq 0$.*

(iii) *For $f \in L^p(\varpi; \Omega)$, if $1 \leq p \leq \infty$, or $f \in C(\Omega)$ if $p = \infty$,*

$$\|S_{\theta, \varpi} f\|_{p, \varpi} \leq \|f\|_{p, \varpi} \quad \text{and} \quad \lim_{\theta \rightarrow 0} \|S_{\theta, \varpi} f - f\|_{p, \varpi} = 0.$$

Proof. From the Fubini theorem and (3.4), it is easy to see that

$$\begin{aligned} \text{proj}_n(\varpi; f *_{\varpi} g) &= \Lambda_n^{(\alpha, \beta)}(g) \text{proj}_n(\varpi; f) \\ &= c_{\alpha, \beta} \int_0^\pi \text{proj}_n(\varpi; S_{\theta, \varpi} f) g(\cos \theta) w_{\alpha, \beta}(\cos \theta) \sin \theta d\theta, \end{aligned}$$

from which (i) follows. To prove (ii), we let g_n be a non-negative function such that $g_n(\cos \theta)$ is supported on $[-\frac{1}{n}, \frac{1}{n}]$ and $\int_0^\pi g_n(\cos \theta) w_{\alpha, \beta}(\cos \theta) \sin \theta d\theta = 1$. Then, using the expression in (i), $f *_{\varpi} g_n$ converges to $S_{\theta, \varpi} f$, which proves (ii). Moreover, by (3.6), $\|f *_{\varpi} g_n\|_{p, \varpi} \leq \|f\|_{p, \varpi}$, so that by the Fatou lemma, $\|S_{\theta, \varpi} f\|_{p, \varpi} \leq \|f\|_{p, \varpi}$. Finally, if $f_n = S_n^\delta(\varpi; f)$ with $\delta \geq \alpha + \beta + 2$, then $f_n \geq 0$ and $f_n \rightarrow f$ in $L^p(\Omega, \varpi)$; furthermore, by (3.7), $S_{\theta, \varpi} f_n$ converges to f_n for $\theta \rightarrow 0$; consequently, by the triangle inequality, $\|S_{\theta, \varpi} f - f\|_{p, \varpi}$ converges to 0 when $\theta \rightarrow 0$. This completes the proof. \square

The operator $S_{\theta, \varpi}$ can be used to define a modulus of smoothness. For $r > 0$, we defined the r -th difference operator

$$\Delta_{\theta, \varpi}^r = (I - S_{\theta, \varpi})^{r/2} = \sum_{n=0}^{\infty} (-1)^n \binom{r/2}{n} (S_{\theta, \varpi})^n,$$

where I denote the identity operator, in the distribution sense, by

$$(3.8) \quad \text{proj}_n(\varpi; \Delta_{\theta, \varpi}^r f) = \left(1 - R_n^{(\alpha, \beta)}(\cos \theta)\right)^{r/2} \text{proj}_n(\varpi; f), \quad n = 0, 1, 2, \dots$$

Definition 3.9. *Let $r > 0$ and $0 < \theta < \pi$. For $f \in L^p(\Omega, \varpi)$ and $1 \leq p < \infty$ or $f \in C(\Omega)$ and $p = \infty$, the weighted r th order modulus of smoothness is defined by*

$$(3.9) \quad \omega_r(f, t)_{p, \varpi} := \sup_{0 < \theta \leq t} \|\Delta_{\theta, \varpi}^r f\|_{p, \varpi}, \quad 0 < t < \pi.$$

For the unit sphere with the Lebesgue measure, the definition of this modulus of smoothness is classical. For weighed approximation on the unit sphere, the unit ball, and the simplex, it is given in [52]. The proof of the following lemma is standard, see [12, Proposition 10.1.2] for example, and will be omitted.

Proposition 3.10. *Let $f \in L^p(\Omega, \varpi)$ if $1 \leq p < \infty$ and $f \in C(\Omega)$ if $p = \infty$. Then*

- (1) $\omega_r(f, t)_{p, \varpi} \leq 2^{r+2} \|f\|_{p, \varpi}$;
- (2) $\omega_r(f, t)_{p, \varpi} \rightarrow 0$ if $t \rightarrow 0+$;
- (3) $\omega_r(f, t)_{p, \varpi}$ is monotone nondecreasing on $(0, \pi)$;
- (4) $\omega_r(f + g, t)_{p, \varpi} \leq \omega_r(f, t)_{p, \varpi} + \omega_r(g, t)_{p, \varpi}$;

(5) For $0 < s < r$,

$$\omega_r(f, t)_{p, \varpi} \leq 2^{(r-s)+2} \omega_s(f, t)_{p, \varpi}.$$

3.3. Characterization of best approximation. Let w be a doubling weight on Ω . For $f \in L^p(\Omega, w)$, we denote by $E_n(f)_{p, w}$ the best approximation to f from $\Pi_n(\Omega)$ in the norm $\|\cdot\|_{p, w}$ of $L^p(\Omega, w)$; that is,

$$E_n(f)_{p, w} := \inf_{g \in \Pi_n(\Omega)} \|f - g\|_{p, w}, \quad 1 \leq p \leq \infty.$$

A central problem of approximation theory is to character this quantity by the smoothness of the functions, usually in terms of a modulus of smoothness or a K -functional, which are often equivalent. In this subsection, we state our main results on the characterization of the best approximation.

The modulus of smoothness is already defined. We now define the K -functional via the differential operator \mathfrak{D}_ϖ in (3.1). Since the operator $-\mathfrak{D}_\varpi$ has nonnegative eigenvalues, it is a non-negative operator. A function $f \in L^p(\Omega; \varpi)$ belongs to the Sobolev space $W_p^r(\Omega; \varpi)$ if there is a function $g \in L^p(\Omega; \varpi)$, which we denote by $(-\mathfrak{D}_\varpi)^{\frac{r}{2}} f$, such that

$$(3.10) \quad \text{proj}_n(\varpi; (-\mathfrak{D}_\varpi)^{\frac{r}{2}} f) = \mu(n)^{\frac{r}{2}} \text{proj}_n(\varpi; f),$$

where we assume that $f, g \in C(\Omega)$ when $p = \infty$. The fractional differential operator $(-\mathfrak{D}_\varpi)^{\frac{r}{2}} f$ is a linear operator on the space $W_p^r(\Omega; \varpi)$ defined by (3.1).

Let w be a doubling weight. We denote by $W_p^r(\Omega, w)$ the Sobolev space that consists of $f \in L^p(\Omega, w)$ such that $(-\mathfrak{D}_\varpi)^{\frac{r}{2}} f \in L^p(\Omega, w)$. The space is well defined as we shall see later in Theorem 3.17. We define our K -functional as follows.

Definition 3.11. Let $r > 0$ and $1 \leq p \leq \infty$. The r -th K -functional of $f \in L^p(\Omega, w)$ is defined by

$$K_r(f, t)_{p, w} := \inf_{g \in W_p^r(\Omega, w)} \left\{ \|f - g\|_{p, w} + t^r \|(-\mathfrak{D}_\varpi)^{\frac{r}{2}} g\|_{p, w} \right\}.$$

We first state of our characterization theorem in terms of the K -functional, which contains two parts, the first part is the direct estimate or the Jackson inequality, and the second part is the inverse estimate. For the first part, we need another assertion to ensure that a Bernstein inequality holds. This is Assertion 5 and it will be stated in Subsection 3.5.

Theorem 3.12. Let ϖ be a weight function that admits Assertions 1, 3 and 5. Let $f \in L^p(\Omega, \varpi)$ if $1 \leq p < \infty$ and $f \in C(\Omega)$ if $p = \infty$. Then for $r > 0$ and $n = 1, 2, \dots$,

(i) the direct estimate

$$E_n(f)_{p, \varpi} \leq c K_r(f; n^{-1})_{p, \varpi};$$

(ii) the inverse estimate

$$K_r(f; n^{-1})_{p, \varpi} \leq c n^{-r} \sum_{k=0}^n (k+1)^{r-1} E_k(f)_{p, \varpi}.$$

It is worth mentioning that the direct estimate holds for the weight norm of ϖ , whereas the inverse estimate holds for all doubling weight. We can also replace the above characterization by the modulus of smoothness $\varpi_r(f; t)_{p, \varpi}$, since the two quantities will be shown to be equivalent.

Theorem 3.13. *Let ϖ be a weight function that admits Assertions 1, 3 and 5. Let $f \in L^p(\Omega, \varpi)$ if $1 \leq p < \infty$ and $f \in C(\Omega)$ if $p = \infty$. If $0 < \theta \leq \pi/2$ and $r > 0$, then*

$$(3.11) \quad c_1 K_r(f; \theta)_{p, \varpi} \leq \omega_r(f; \theta)_{p, \varpi} \leq c_2 K_r(f; \theta)_{p, \varpi}.$$

For the unit sphere and the unit ball, these theorems are known to hold when ϖ being reflection invariant weight functions. For the unit sphere with the Lebesgue measure, the characterization of best approximation has a long history. Starting from [4, 36] for $r = 2$, the problem was studied by a number of authors and finally completed in [42]; we refer to [12, p. 102 and p. 263] for historical notes. These results are extended to the weighted case, with reflection invariant weight functions, on the unit sphere, the unit ball, and the simplex, in [52].

Another pair of K -functional and modulus of smoothness was used to characterize the best approximation for the unit sphere with the Lebesgue measure in [11]. The approach appears to be very much domain-specific and relies on the geometric and the differential structure on the unit sphere.

We will give the proof of these two theorems in Subsection 3.6, after proving several preliminary results that are of interest in their own right.

3.4. Near best approximation operator. In this subsection we study the approximation property of the operator $L_n(\varpi) * f$, defined in (2.42), that has the highly localized kernel as its kernel. By (3.2), the operator can be written in terms of the convolution operator by

$$L_n(\varpi) * f = f *_{\varpi} L_n^{(\alpha, \beta)}, \quad n = 0, 1, 2, \dots$$

We now show that $L_n(\varpi) * f$ provides near-best approximation to f .

Theorem 3.14. *Let \hat{a} be admissible of type (a). Assume ϖ admits Assertions 1 and 3. Then the operator $L_n * f$ satisfies*

- (a) $L_n(\varpi) * f$ is a polynomial of degree at most $2n$;
- (b) $L_n(\varpi) * g = g$ for any polynomial $g \in \Pi_n(\Omega)$ of degree n ;
- (c) For $f \in L^p(\Omega, \varpi)$, $1 \leq p \leq \infty$,

$$\|L_n(\varpi) * f\|_{p, \varpi} \leq c \|f\|_{p, \varpi} \quad \text{and} \quad \|L_n(\varpi) * f - f\|_{p, \varpi} \leq c E_n(f)_{p, \varpi}.$$

Proof. The first two properties are immediate consequences of the definition. For the third property, in the case $p = 1$ we obtain

$$\|L_n(\varpi) * f\|_{1, \varpi} \leq c \|f\|_{1, \varpi} \max_{x \in \Omega} \int_{\Omega} |L_n(\varpi; x, y)| \varpi(y) dm(y).$$

By Assertion 1 and (2.11) with $p = 1$ and a large τ , the integral in the right-hand side is bound by

$$\max_{x \in \Omega} \int_{\Omega} \frac{\varpi(y) dm(y)}{\sqrt{\varpi(B(y, n^{-1}))} \sqrt{\varpi(B(x, n^{-1}))} (1 + nd(x, y))^{\tau}} \leq c,$$

which shows that $\|L_n(\varpi) * f\|_{p, \varpi}$ is bounded for $p = 1$. The same inequality also shows that the boundedness holds for $p = \infty$. The case $1 < p < \infty$ follows from the Riesz-Thorin interpolation theorem. By item (b), the boundedness of $\|L_n(\varpi) * f\|_{p, \varpi}$ implies, by the triangle inequality,

$$\|L_n(\varpi) * f - f\|_{p, \varpi} \leq \|L_n(\varpi) * (f - g)\|_{p, \varpi} + \|f - g\|_{p, \varpi} \leq c \|f - g\|_{p, \varpi}$$

for any polynomial $g \in \Pi_n(\Omega)$. Taking infimum over g completes the proof. \square

Because of property (c), we call the operator $L_n(\varpi) * f$ near best approximation polynomial. Such operators have been used extensively in approximation theory and computational harmonic analysis on the unit sphere and the unit ball (see [12], for example), and in various other domains.

Theorem 3.15. *Let the assumption be the same as in the previous theorem. For $f \in W_p^r(\Omega, \varpi)$, $1 \leq p \leq \infty$,*

$$(3.12) \quad \|f - L_n(\varpi) * f\|_{p, \varpi} \leq cn^{-r} \|(-\mathfrak{D}_\varpi)^{\frac{r}{2}} f\|_{p, \varpi}, \quad n = 1, 2, \dots$$

Proof. Let ℓ be a positive integer. Without loss of generality, we can assume $n > \ell$. Since $L_n(\varpi) * f = f$ if f is a polynomial of degree n , we can write

$$\begin{aligned} f - L_n(\varpi) * f &= \sum_{k=n+1}^{\infty} \left(1 - \widehat{a}\left(\frac{k}{n}\right)\right) \text{proj}_k(\varpi; f) \\ &= \sum_{k=n+1}^{\infty} \left(1 - \widehat{a}\left(\frac{k}{n}\right)\right) \mu(k)^{-\frac{r}{2}} \text{proj}_k(\varpi; F), \end{aligned}$$

where $F = (-\mathfrak{D}_\varpi)^{\frac{r}{2}} f$. Summation by parts $\ell + 1$ times, we obtain

$$f - L_n(\varpi) * f = \sum_{k=n+1}^{\infty} b_k \text{proj}_k(\varpi; F) = \sum_{k=n+1}^{\infty} (\Delta^{\ell+1} b_k) A_k^\ell S_k^\ell(\varpi; F),$$

where $b_k = (1 - \widehat{a}(\frac{k}{n}))\mu(k)^{-\frac{r}{2}}$, $A_k^\ell = \binom{k+\ell}{k} \sim k^\ell$ and $S_k^\ell(\varpi; F)$ denotes the k -th Cesàro (C, ℓ) mean of the Fourier orthogonal series with respect to ϖ . Since \widehat{a} is in C^∞ and its support is $[0, 2]$, it is easy to see that $|\Delta^{\ell+1} b_k| \leq ck^{-r-\ell-1}$. Hence, it follows that

$$\sum_{k=n+1}^{\infty} |\Delta^{\ell+1} b_k| A_k^\ell \leq c \sum_{k=n+1}^{\infty} k^{-r-\ell-1} k^\ell \leq cn^r.$$

Since the choice of ℓ implies that $\|S_k^\ell(\varpi; F)\|_{p, \varpi} \leq \|F\|_{p, \varpi}$ by the convergence of the Cesàro means, we obtain

$$\|f - L_n(\varpi) * f\|_{p, \varpi} \leq cn^{-r} \|F\|_{p, \varpi},$$

which is what we need to prove. \square

3.5. Bernstein and Nikolskii inequalities. The Bernstein inequality is essential for studying approximation by polynomials. We establish such an inequality for the deviation operator \mathfrak{D}_ϖ defined by (3.1). First we prove a proposition that is of independent interest.

Proposition 3.16. *Let $(\Omega, \varpi, \mathbf{d})$ be a localizable homogeneous space and assume Assertion 4 holds. Let $G_n(\cdot, \cdot) : \Omega \times \Omega \mapsto \mathbb{R}$ be a kernel such that $G_n(x, y) = G_n(y, x)$ for all $x, y \in \Omega$ and G_n is a polynomial of degree n in either of its variables. Let $T : f \mapsto Tf$ be the operator defined by*

$$Tf(x) = \int_{\Omega} f(y) G_n(x, y) \varpi(y) \mathbf{d}\mathbf{m}(y).$$

If f is a polynomial of degree n , then for $1 \leq p \leq \infty$ and any doubling weight \mathbf{w} ,

$$(3.13) \quad \|Tf\|_{p, \mathbf{w}} \leq c \|f\|_{p, \mathbf{w}} \max_{z \in \Omega} \int_{\Omega} |G_n(x, z)| (1 + n\mathbf{d}(z, x))^{\alpha(\mathbf{w}) + \alpha(\varpi)} \varpi(x) \mathbf{d}\mathbf{m}(x).$$

Proof. Since T is a linear integral operator, if f is a polynomial of degree n then Tf is a polynomial of the same degree. Let Ξ be an ε -separated set of Ω for $\varepsilon = \frac{\delta}{2n}$ so that the Marcinkiewicz-Zygmund inequalities in Theorem 2.15 hold for all polynomials of degree $2n$. The inequality is trivial for $p = \infty$ since it holds without $(1 + nd(z, x))^{\alpha(\mathbf{w})}$ term. For $1 \leq p < \infty$, by (2.35),

$$\|Tf\|_{p, \mathbf{w}}^p \leq c \sum_{z \in \Xi} |Tf(z)|^p \mathbf{w} \left(B(z, \frac{\delta}{2n}) \right).$$

Since $f(y)G_n(z, y)$ is a polynomial of degree $2n$ in y variable, applying (2.35) again, we obtain

$$|Tf(z)| \leq c \sum_{u \in \Xi} |f(u)| |G_n(u, z)| \varpi \left(B(u, \frac{\delta}{2n}) \right).$$

Assume $1 \leq p < \infty$. By the Hölder's inequality,

$$\begin{aligned} |Tf(z)|^p &\leq c \sum_{u \in \Xi} |f(u)|^p |G_n(z, u)| \varpi \left(B(u, \frac{\delta}{2n}) \right) \left(\sum_{u \in \Xi} |G_n(z, u)| \varpi \left(B(u, \frac{\delta}{2n}) \right) \right)^{\frac{p}{q}} \\ &\leq c \sum_{u \in \Xi} |f(u)|^p |G_n(z, u)| \varpi \left(B(u, \frac{\delta}{2n}) \right) \|G_n(z, \cdot)\|_1^{\frac{p}{q}}, \end{aligned}$$

where we used (2.34) in the second step. Using (ii) of Lemma 2.1 for both ϖ and \mathbf{w} , we see that

$$\begin{aligned} \sum_{z \in \Xi} |Tf(z)|^p \mathbf{w} \left(B(z, \frac{\delta}{2n}) \right) &\leq c \max_{z \in \Xi} \|G_n(z, \cdot)\|_1^{\frac{p}{q}} \\ &\quad \times \sum_{u \in \Xi} |f(u)|^p \mathbf{w} \left(B(u, \frac{\delta}{2n}) \right) \sum_{z \in \Xi} |G_n(z, u)| (1 + nd(z, u))^{\alpha(\mathbf{w}) + \alpha(\varpi)} \varpi \left(B(z, \frac{\delta}{2n}) \right). \end{aligned}$$

By Lemma 2.16, there is a nonnegative polynomial Q_u of degree n that satisfies (2.39) with $\alpha = \alpha(\mathbf{w}) + \alpha(\varpi)$, which allows us to apply (2.34) on the polynomial $z \mapsto G(z, u)Q_u(z)$ to show that the last sum in the right-hand side is bounded by

$$\begin{aligned} c \sum_{z \in \Xi} |G_n(z, u)| Q_u(z) \varpi \left(B(z, \frac{\delta}{2n}) \right) &\leq c \int_{\Omega} |G_n(x, u)| Q_u(x) \varpi(x) \mathrm{d}m(x) \\ &\leq c \int_{\Omega} |G_n(x, u)| (1 + nd(x, u))^{\alpha(\mathbf{w}) + \alpha(\varpi)} \varpi(x) \mathrm{d}m(x). \end{aligned}$$

Putting these together and using (2.34) and (2.35), we have proved that

$$\begin{aligned} \|Tf\|_{p, \mathbf{w}}^p &\leq c \|f\|_{p, \mathbf{w}}^p \max_{z \in \Xi} \|G_n(z, \cdot)\|_1^{\frac{p}{q}} \\ &\quad \times \max_{u \in \Xi} \int_{\Omega} |G_n(x, u)| (1 + nd(x, u))^{\alpha(\mathbf{w}) + \alpha(\varpi)} \varpi(x) \mathrm{d}m(x) \\ &\leq c \|f\|_{p, \mathbf{w}}^p \left(\max_{u \in \Xi} \int_{\Omega} |G_n(x, u)| (1 + nd(x, u))^{\alpha(\mathbf{w}) + \alpha(\varpi)} \varpi(x) \mathrm{d}m(x) \right)^{1 + \frac{p}{q}}, \end{aligned}$$

where we have used the assumption that G_n is symmetric in its variables. Since $1 + \frac{p}{q} = p$, this proves the stated inequality. \square

For $r > 0$, we denote by $L_n^{(r)}(\varpi; \cdot, \cdot)$ the kernel defined by

$$(3.14) \quad L_n^{(r)}(\varpi; x, y) = \sum_{n=0}^{\infty} \hat{a} \left(\frac{k}{n} \right) [\mu(k)]^{r/2} P_k(\varpi; x, y),$$

which is the kernel $\mathfrak{D}_{\varpi}^{r/2} L_n(x, y)$ with $\mathfrak{D}_{\varpi}^{r/2}$ applying on x variable. Our Bernstein inequality is proved under the following assumption on the decaying of this kernel.

Assertion 5. For $r > 0$ and $\kappa > 0$, the kernel $L_n^{(r)}(\varpi)$ satisfies, for $x, y \in \Omega$,

$$\left| L_n^{(r)}(\varpi; x, y) \right| \leq c_{\kappa} \frac{n^r}{\sqrt{\varpi(B(x, n^{-1}))} \sqrt{\varpi(B(y, n^{-1}))}} (1 + nd(x, y))^{-\kappa}.$$

Theorem 3.17. Let ϖ be a weight functions that admits Assertions 1–5. Let \mathbf{w} be a doubling weight on Ω . If $r > 0$, $1 \leq p \leq \infty$ and $f \in \Pi_n(\Omega)$, then

$$\|(-\mathfrak{D}_{\varpi})^{\frac{r}{2}} f\|_{p, \mathbf{w}} \leq cn^r \|f\|_{p, \mathbf{w}}.$$

Proof. Since $L_n(\varpi) * f$ reproduces polynomials of degree n , we can write

$$(-\mathfrak{D}_{\varpi})^{\frac{r}{2}} f = (-\mathfrak{D}_{\varpi})^{\frac{r}{2}} (L_n(\varpi) * f) = \int_{\Omega} f(y) L_n^{(r)}(\varpi; x, y) \varpi(y) dm(y),$$

where $L_n^{(r)}(\varpi)$ is defined by (3.14). Applying Proposition 3.16, we conclude that

$$\|(-\mathfrak{D}_{\varpi})^{\frac{r}{2}} f\|_{p, \mathbf{w}} \leq c \|f\|_{p, \mathbf{w}} \max_{x \in \Omega} \int_{\Omega} \left| L_n^{(r)}(\varpi; x, y) \right| (1 + nd(x, y))^{\alpha(\mathbf{w}) + \alpha(\varpi)} \varpi(y) dm(y).$$

By Assertion 5, the integral in the right-hand side is bounded by

$$c_{\kappa} n^r \int_{\Omega} \frac{\varpi(s)}{\sqrt{\varpi(B(x, n^{-1}))} \sqrt{\varpi(B(y, n^{-1}))}} (1 + nd(x, y))^{\kappa - \alpha(\mathbf{w}) - \alpha(\varpi)} dm(y) \leq cn^r$$

using (2.11) with $p = 1$ and $\kappa > \alpha(\mathbf{w}) + \frac{5}{4}\alpha(\varpi)$. \square

For the Laplace-Beltrami operator on the unit sphere, this Bernstein inequality is classical if \mathbf{w} is the Lebesgue measure, whereas the version with the doubling weight was proved in [7]. For the unit ball, the inequality was established in [52] for the classical weight function with a different proof that applies to reflection invariant weight functions on the unit sphere and the unit ball.

We can also prove a Nikolskii type inequality for the doubling weight on Ω that admits a localizable homogeneous space.

Theorem 3.18. Let \mathbf{w} be a doubling weight on Ω . If $0 < p < q \leq \infty$ and $f \in \Pi_n(\Omega)$, then

$$(3.15) \quad \|f\|_{q, \mathbf{w}} \leq cn^{\left(\frac{1}{p} - \frac{1}{q}\right)\alpha(\mathbf{w})} \|f\|_{p, \mathbf{w}}.$$

Proof. The proof is another one that follows as in the case of the unit sphere. The main work lies in proving the case $q = \infty$, for which we choose a maximal $\frac{\delta}{n}$ -separated subset of Ω , so that

$$\begin{aligned} \|f\|_{\infty} &\leq c \max_{z \in \Xi} |f(z)| \leq c \left(\min_{z \in \Xi} B\left(z, \frac{\delta}{n}\right) \right)^{-\frac{1}{p}} \left(\sum_{z \in \Xi} \mathbf{w}\left(B\left(z, \frac{\delta}{n}\right)\right) |f(z)|^p \right)^{\frac{1}{p}} \\ &\leq c \|f\|_{p, \mathbf{w}} \max_{z \in \Xi} \left(B\left(z, \frac{1}{n}\right) \right)^{-\frac{1}{p}}. \end{aligned}$$

Since Ω is compact, there is a positive number $\rho_0 > 0$ such that, for each $z \in \Xi$, $1 = \mathbf{w}(\Omega) = \mathbf{w}(B(z, \rho))$ for some positive number $\rho \leq \rho_0$. Let m be a positive integer such that $2^{m-1} \leq \rho n \leq 2^m$. Then, by (i) of Lemma 2.1,

$$1 = \mathbf{w}(B(z, \rho)) \leq c_{L(\mathbf{w})} 2^{m\alpha(\mathbf{w})} \mathbf{w}\left(B\left(z, \frac{1}{n}\right)\right) \leq cn^{\alpha(\mathbf{w})} \mathbf{w}\left(B\left(z, \frac{1}{n}\right)\right),$$

which implies that $w(B(z, \frac{1}{n})) \leq cn^{-\alpha(w)}$. Combing the two inequalities, we conclude that $\|f\|_\infty \leq c\|f\|_{p,w} n^{\frac{1}{p}\alpha(w)}$. This proves (3.15) for $q = \infty$.

The case $q < \infty$ reduces to that of $q = \infty$ since, using (3.15) for $q = \infty$,

$$\|f\|_{q,w}^q \leq \|f\|_\infty^{q-p} \|f\|_{q,w}^p \leq c \left(\|f\|_{p,w} n^{\frac{1}{p}\alpha(w)} \right)^{q-p} \|f\|_{q,w}^p = cn^{q(\frac{1}{p}-\frac{1}{q})\alpha(w)} \|f\|_{p,\sigma}^q,$$

which is (3.15) for $q < \infty$. \square

For the unit sphere, this inequality was established in [10]. The above proof uses essentially the same argument as seen in [12, Theorem 5.5.1]. For the unit ball with the classical weight, the inequality was proved in [25].

3.6. Proof of the main results. We are now ready to prove Theorems 3.12 and 3.13. The proof follows the same procedure used in [42, 52], which is summarized in [12, section 10]. Since most technical parts are essentially the same, we shall be brief.

Proof of Theorem 3.12.

To prove the direct estimate, we use Theorems 3.14 and 3.15 to obtain

$$\begin{aligned} \|f - L_n(\varpi; f)\|_{p,\varpi} &\leq 2\|f - g\|_{p,\varpi} + \|g - L_n(\varpi; g)\|_{p,\varpi} \\ &\leq 2\|f - g\|_{p,\varpi} + cn^{-r} \|(-\mathfrak{D}_\varpi)^{\frac{r}{2}} g\|_{p,\varpi}. \end{aligned}$$

Taking infimum over g proves (i).

To prove the inverse estimate, we choose m such that $2^{m-1} \leq n < 2^m$. Let us set $L_{2^{-1}}(\varpi) * f = 0$. Then, choosing $g = L_{2^m}(\varpi) * f$, we obtain by Theorem 3.14,

$$\begin{aligned} K_r(f, n^{-1})_{p,w} &\leq \|f - L_{2^m}(\varpi) * f\|_{p,w} + 2^{-(m-1)r} \|(-\mathfrak{D}_\varpi)^{\frac{r}{2}} L_{2^m}(\varpi) * f\|_{p,w} \\ &\leq cE_n(f)_{p,w} + 2^{-(m-1)r} \sum_{j=0}^m \|(-\mathfrak{D}_\varpi)^{\frac{r}{2}} [L_{2^j}(\varpi) * f - L_{2^{j-1}}(\varpi) * f]\|_{p,w}. \end{aligned}$$

Applying the Bernstein inequality Theorem 3.17 and using the triangle inequality with Theorem 3.14, we conclude that

$$\begin{aligned} K_r(f, n^{-1})_{p,\varpi} &\leq cE_n(f)_{p,\varpi} + 2^{-mr} \sum_{j=0}^m 2^{-jr} \|L_{2^j}(\varpi) * f - L_{2^{j-1}}(\varpi) * f\|_{p,w} \\ &\leq c2^{-mr} \sum_{j=0}^m 2^{jr} E_{2^{j-1}}(f)_{p,\varpi} \leq cn^{-r} \sum_{k=0}^m (k+1)^{r-1} E_k(f)_{p,\varpi}. \end{aligned}$$

This completes the proof. \square

The proof of the equivalence of the K -functional and the modulus of smoothness relies on two technical lemmas. Recall that \hat{a} is a cut-off function and $\mu(j)$ is the eigenvalue in (3.1).

Lemma 3.19. *Let $\alpha \geq \beta \geq -\frac{1}{2}$. For $r > 0$, $0 \leq \theta \leq 3n^{-1}$, and any $\ell \in \mathbb{N}$,*

$$(3.16) \quad \sum_{j=0}^{2n} \left| \Delta^{\ell+1} \left[\left(\frac{1 - R_j^{(\alpha,\beta)}(\cos \theta)}{\mu(j)\theta^2} \right)^r \hat{a}\left(\frac{j}{n}\right) \right] \right| (j+1)^\ell \leq c,$$

and

$$(3.17) \quad \sum_{j=0}^{2n} \left| \Delta^{\ell+1} \left[\left(\frac{\mu(j)\theta^2}{1 - R_j^{(\alpha,\beta)}(\cos \theta)} \right)^r \hat{a}\left(\frac{j}{n}\right) \right] \right| (j+1)^\ell \leq c,$$

where the difference $\triangle^{\ell+1}$ is acting on j and c depends only on α, β, ℓ, r .

Lemma 3.20. *Let $\alpha \geq \beta \geq -\frac{1}{2}$ and let $\ell = \lceil \alpha \rceil$. If $r > 0$, $k^{-1} \leq \theta \leq \frac{\pi}{2}$, $j \in \mathbb{N}_0$ and $0 \leq j \leq \ell + 1$, then for any $m \in \mathbb{N}$,*

$$(3.18) \quad \left| \triangle^j \left(\frac{(1 - (1 - R_k^{(\alpha, \beta)}(\cos \theta))^r)^{m+\ell+1}}{(1 - R_k^{(\alpha, \beta)}(\cos \theta))^r} \right) \right| \leq c_{m,r} (k\theta)^{-m\alpha} \theta^j.$$

For $\alpha = \beta = \lambda$, these two lemma are Lemma 10.4.3 and Lemma 10.4.4 in [12]. The proof relies on properties of the Jacobi polynomials and holds readily for $\alpha \geq \beta \geq -\frac{1}{2}$.

Proof of Theorem 3.13.

Choose n such that $\theta^{-1} \leq n \leq 3\theta^{-1}$. Let g be a polynomial in $\Pi_n(\Omega)$. Then so is $(I - S_{\theta, \varpi})^{r/2} g$ in $\Pi_n(\Omega)$ by (3.7). Let \hat{a} be a cut-off function of type (a), so that $L_n(\varpi)$ reproduces polynomials of degree n . Using (3.10), it follows readily that

$$(I - S_{\theta, \varpi})^{r/2} g = \theta^r \sum_{j=0}^{2n} \left(\frac{1 - R_j^{(\alpha, \beta)}(\cos \theta)}{\mu(j)\theta^2} \right)^{r/2} \hat{a}\left(\frac{j}{n}\right) \text{proj}_j \left(\varpi; (-\mathfrak{D}\varpi)^{r/2} g \right).$$

Following the proof of Theorem 3.15 to take a summation by parts and use the boundedness of the Cesàro means, we can use (3.16) to obtain

$$(3.19) \quad \left\| (I - S_{\theta, \varpi})^{r/2} g \right\|_{p, \varpi} \leq cn^{-r} \left\| (-\mathfrak{D}\varpi)^{r/2} g \right\|_{p, \varpi}.$$

Using this inequality with $g = L_{\lfloor n/2 \rfloor}(\varpi) * f$ and the Jackson estimate in Theorem 3.12, we obtain

$$\begin{aligned} \left\| (I - S_{\theta, \varpi})^{r/2} f \right\|_{p, \varpi} &\leq c \left\| f - L_{\lfloor \frac{n}{2} \rfloor}(\varpi) * f \right\|_{p, \varpi} + \left\| (I - S_{\theta, \varpi})^{r/2} L_{\lfloor \frac{n}{2} \rfloor}(\varpi) * f \right\|_{p, \varpi} \\ &\leq cK_r(f; n^{-1})_{p, \varpi} + cn^{-r} \left\| (-\mathfrak{D})^{r/2} L_{\lfloor \frac{n}{2} \rfloor}(\varpi) * f \right\|_{p, \varpi}. \end{aligned}$$

By the triangle inequality and applying the Bernstein inequality on $L_n(\varpi) * (f - g)$, we obtain

$$\begin{aligned} n^{-r} \left\| (-\mathfrak{D})^{r/2} L_{\lfloor \frac{n}{2} \rfloor}(\varpi) * f \right\|_{p, \varpi} &\leq c \|f - g\|_{p, \varpi} + n^{-r} \left\| (-\mathfrak{D})^{r/2} L_{\lfloor \frac{n}{2} \rfloor}(\varpi) * g \right\|_{p, \varpi} \\ &\leq c \left(\|f - g\|_{p, \varpi} + n^{-r} \left\| (-\mathfrak{D})^{r/2} g \right\|_{p, \varpi} \right), \end{aligned}$$

where we used, by (3.1), $(-\mathfrak{D})^{r/2} L_{\lfloor \frac{n}{2} \rfloor}(\varpi) * g = L_{\lfloor \frac{n}{2} \rfloor}(\varpi) * (-\mathfrak{D})^{r/2} g$ in the second step. Hence, taking infimum over g , it follows from the above two inequalities and $\theta \sim n^{-1}$ that

$$\left\| (I - S_{\theta, \varpi})^{r/2} f \right\|_{p, \varpi} \leq cK_r(f, \theta)_{p, \varpi},$$

from which the right-hand inequality of (3.11) follows.

In the other direction, we can follow the proof of inequality (3.19) and use (3.17) instead of (3.16), to establish

$$n^{-r} \left\| (-\mathfrak{D}\varpi)^{r/2} L_n(\varpi) * f \right\|_{p, \varpi} \leq c \left\| (I - S_{\theta, \varpi})^{r/2} f \right\|_{p, \varpi}.$$

Hence, in order to prove the left-hand inequality of (3.11), it suffices to prove that

$$(3.20) \quad \|f - L_n(\varpi) * f\|_{p, \varpi} \leq c \left\| (I - S_{\theta, \varpi})^{r/2} f \right\|_{p, \varpi}.$$

Using $(1-r)^{-1} = \sum_{i=0}^{m+\ell} r^i + r^{m+\ell}(1-r)^{-1}$ with $r = 1 - (1 - R_j^{(\alpha,\beta)}(\cos\theta))^{r/2}$, one can write, as seen in [12, p. 253], that

$$\begin{aligned} f - L_n(\varpi) * f &= \sum_{i=0}^{m+\ell} (I - L_n(\varpi)) * \left(I - (I - S_{\theta,\varpi})^{r/2} \right)^i F \\ &\quad + \sum_{j=n}^{\infty} \left(1 - \hat{a} \left(\frac{j}{n} \right) \right) \frac{(1 - (1 - R_j^{(\alpha,\beta)}(\cos\theta))^{r/2})^{m+\ell+1} F}{(1 - R_j^{(\alpha,\beta)}(\cos\theta))^{r/2}} \text{proj}_j(\varpi; F), \end{aligned}$$

where $F = (I - S_{\theta,\varpi})^{r/2} f$. While the first term is bounded by $\|F\|_{p,\varpi}$ by the boundedness of $L_n(\varpi) * f$ and $(I - (I - S_{\theta,\varpi})^{r/2})F$, the second one can be shown to be bounded by $\|F\|_{p,\varpi}$ by using (3.18) with the summation by parts and the boundedness of Cesàro means that we have used several times. This proves (3.20) and completes the proof. \square

4. HOMOGENEOUS SPACE ON CONIC SURFACES

In this section we work in the setting of homogeneous space on the conic surface

$$\mathbb{V}_0^{d+1} = \{(x, t) : \|x\| = t, x \in \mathbb{R}^d, 0 \leq t \leq 1\}.$$

We shall verify that the framework in the previous two sections is applicable on this domain for the weight function $t^{-1}(1-t)^\gamma$, which has a singularity at the apex. The verification is highly non-trivial because of new phenomena and obstacles encountered. While the structure of orthogonal polynomials on the conic surface shares characteristic features of spherical harmonics, the conic surface is markedly different from that of the unit sphere because of its apex and its boundary.

Our first task is to understand, in the first subsection, the intrinsic distance function on the conic surface, which turns out to be incomparable to the Euclidean distance around the apex. In the second subsection, we show that the Jacobi weight function $w_{\beta,\gamma}(t) = t^\beta(1-t)^\gamma$ is a doubling weight with respect to the intrinsic distance. The orthogonal structure with respect to the Jacobi weight is reviewed in the third subsection, which is used to verify Assertions 1-3 of the highly localized kernels for the weight $w_{-1,\gamma}$ in the fourth subsection. Construction of ε -separated set of \mathbb{V}_0^{d+1} is provided in the fifth subsection and used to state the Marcinkiewicz-Zygmund inequality. Assertion 4 is verified in the sixth subsection, which ensures that the positive cubature rules and the tight localized frames can both be stated for the conic surface. In the seventh subsection, Assertion 5 is verified and the characterization of the best approximation by polynomials is stated.

4.1. Distance on the surface of the cone. Our first task is to define an appropriate distance function on the surface of the cone. Unlike the sphere, the surface \mathbb{V}_0^{d+1} has a boundary at $t = 1$ and a singularity at the apex $t = 0$. Our distance function should measure the distance between points near the boundary or the apex and the distance between interior points differently. This is a well-known phenomenon as we have already seen for the interval $[-1, 1]$. More generally, the distance on the interval $[a, b]$ is given by the change of variables $x \in [a, b] \mapsto y \in [-1, 1]$,

$$d_{[a,b]}(x_1, x_2) = \frac{b-a}{2} \arccos \left(y_1 y_2 + \sqrt{1-y_1^2} \sqrt{1-y_2^2} \right),$$

where $y_i = -1 + 2\frac{x_i - a}{b - a}$. In particular, the distance function for $[0, 1]$ is given by

$$\begin{aligned} d_{[0,1]}(x_1, x_2) &= \frac{1}{2} \arccos \left((2x_1 - 1)(2y_1 - 1) + 4\sqrt{x_1(1-x_1)}\sqrt{y_1(1-y_1)} \right) \\ &= \arccos \left(\sqrt{x_1 x_2} + \sqrt{(1-x_1)(1-x_2)} \right), \end{aligned}$$

where the second identity follows from $\arccos(\alpha) = \frac{1}{2} \arccos(2\alpha^2 - 1)$. In particular, setting $x_i = \cos^2 \frac{\theta_i}{2} \in [0, 1]$, $0 \leq \theta_i \leq \pi$, we obtain $d_{[0,1]}(x_1, x_2) = \frac{1}{2}|\theta_1 - \theta_2|$.

Definition 4.1. For (x, t) and (y, s) on \mathbb{V}_0^{d+1} , define

$$(4.1) \quad d_{\mathbb{V}_0}((x, t), (y, s)) := \arccos \left(\sqrt{\frac{\langle x, y \rangle + ts}{2}} + \sqrt{1-t}\sqrt{1-s} \right).$$

Proposition 4.2. The function $d_{\mathbb{V}_0^{d+1}}(\cdot, \cdot) = d_{\mathbb{V}_0}(\cdot, \cdot)$ defines a distance on the surface of the cone \mathbb{V}_0^{d+1} .

Proof. Evidently $d_{\mathbb{V}_0}(\cdot, \cdot)$ is symmetric. Since $\|x\| = t$ and $\|y\| = s$, it follows readily that $0 \leq \sqrt{\frac{\langle x, y \rangle + ts}{2}} + \sqrt{1-t}\sqrt{1-s} \leq 1$, so that $d_{\mathbb{V}_0}((x, t), (y, s)) \geq 0$ and, furthermore, $d_{\mathbb{V}_0}((x, t), (x, t)) = \arccos 1 = 0$. Hence, we only need to prove that it satisfies the triangle inequality.

For $d = 2$, we write $(x, t) = (x_1, x_2, t) \in \mathbb{V}_0^3$ with $x_1^2 + x_2^2 = t^2$. For $(x, t), (y, s) \in \mathbb{V}_0^3$, it is easy to verify the identity

$$\left(\sqrt{(t+x_1)(s+y_1)} + \text{sign}(x_2 y_2) \sqrt{(t-x_1)(s-y_1)} \right)^2 = \frac{ts + x_1 y_1 + x_2 y_2}{2}.$$

Hence, setting $z_{x,t} = (\sqrt{t+x_1}, \text{sign}(x_2)\sqrt{t-x_1}, \sqrt{1-t})$ and define $z_{y,s}$ similarly, then they are elements in \mathbb{S}^2 and it follows that

$$d_{\mathbb{V}_0}((x, t), (y, s)) = \arccos(\langle z_{x,t}, z_{y,s} \rangle) = d_{\mathbb{S}^2}(z_{x,t}, z_{y,s}),$$

where $d_{\mathbb{S}^2}(\cdot, \cdot)$ is the geodesic distance of the unit sphere \mathbb{S}^2 . In particular, $d_{\mathbb{V}_0}(\cdot, \cdot)$ satisfies the triangle inequality. For $d > 2$, given three distinct point in \mathbb{V}_0^{d+1} , written as $(t_i \xi_i, t_i)$, $1 \leq i \leq 3$, where $\xi \in \mathbb{S}^{d-1}$, we can find a rotation in \mathbb{R}^d so that $\xi_i = (\eta_i, 0)$ with $\eta_i \in \mathbb{S}^2$. Hence, the triangle inequality for $d > 2$ follows from the triangle inequality for $d = 2$. This completes the proof. \square

The distance function $d_{\mathbb{V}_0}(\cdot, \cdot)$ is closely related to the distance function $d_{[0,1]}(\cdot, \cdot)$ of the interval $[0, 1]$ and the geodesic distance $d_{\mathbb{S}}(\cdot, \cdot)$ of the unit sphere \mathbb{S}^{d-1} .

Proposition 4.3. For $d \geq 2$ and $(x, t), (y, s) \in \mathbb{V}_0^{d+1}$, write $x = t\xi$ and $y = s\eta$ with $\xi, \eta \in \mathbb{S}^{d-1}$. Then

$$(4.2) \quad 1 - \cos d_{\mathbb{V}_0}((x, t), (y, s)) = 1 - \cos d_{[0,1]}(t, s) + \sqrt{t}\sqrt{s} [1 - \cos(\frac{1}{2}d_{\mathbb{S}}(\xi, \eta))].$$

In particular,

$$(4.3) \quad c_1 d_{\mathbb{V}_0}((x, t), (y, s)) \leq d_{[0,1]}(t, s) + (ts)^{\frac{1}{4}} d_{\mathbb{S}}(\xi, \eta) \leq c_2 d_{\mathbb{V}_0}((x, t), (y, s)).$$

Proof. Using $\arccos(\alpha) = \frac{1}{2} \arccos(2\alpha^2 - 1)$, we deduce

$$\sqrt{\frac{ts + \langle x, y \rangle}{2}} = \sqrt{ts} \sqrt{\frac{1 + \langle \xi, \eta \rangle}{2}} = \sqrt{ts} \cos\left(\frac{1}{2} \arccos \langle \xi, \eta \rangle\right).$$

Consequently, in terms of the geodesic distance on the unit sphere, we can write

$$\mathbf{d}_{\mathbb{V}_0}((x, t), (y, s)) = \arccos \left[\sqrt{t}\sqrt{s} \cos \left(\frac{1}{2} \mathbf{d}_{\mathbb{S}}(\xi, \eta) \right) + \sqrt{1-t}\sqrt{1-s} \right].$$

In particular, it follows that

$$1 - \cos \mathbf{d}_{\mathbb{V}_0}((x, t), (y, s)) = 1 - \sqrt{t}\sqrt{s} - \sqrt{1-t}\sqrt{1-s} + \sqrt{t}\sqrt{s} \left[1 - \cos \left(\frac{1}{2} \mathbf{d}_{\mathbb{S}}(\xi, \eta) \right) \right],$$

which is the identity (4.2). From this identity, (4.3) follows from $1 - \cos \theta = 2 \sin^2 \frac{\theta}{2}$, $\frac{1}{\pi} \theta \leq \sin \frac{\theta}{2} \leq \frac{\theta}{2}$ for $0 \leq \theta \leq \pi$, and $(a+b)^2/2 \leq a^2 + b^2 \leq (a+b)^2$ for $a, b \geq 0$. \square

The line segment from the apex to a point $(\xi, 1)$, $\xi \in \mathbb{S}^{d-1}$, on the top boundary of the cone \mathbb{V}_0^{d+1} can be parametrized by $l_\xi = \{(t\xi, t) : 0 \leq t \leq 1\}$. For two points $(t\xi, t)$ and $(s\xi, s)$ on l_ξ , the identity (4.2) shows that

$$\mathbf{d}_{\mathbb{V}_0}((t\xi, t), (s\xi, s)) = \frac{|\theta - \phi|}{2} = \mathbf{d}_{[0,1]}(t, s)$$

if $t = \cos^2 \frac{\theta}{2}$ and $s = \cos^2 \frac{\phi}{2}$. Moreover, the top boundary of \mathbb{V}_0^{d+1} is the unit sphere \mathbb{S}^{d-1} , or $\{(\xi, 1) : \xi \in \mathbb{S}^{d-1}\}$. For $(\xi_1, 1)$ and $(\xi_2, 1)$ on this boundary, (4.2) gives

$$\mathbf{d}_{\mathbb{V}_0}((\xi_1, 1), (\xi_2, 1)) = \frac{1}{2} \mathbf{d}_{\mathbb{S}}(\xi_1, \xi_2),$$

or half of the geodesic distance on the unit sphere.

Remark 4.1. It is well-known that the geodesic distance $\mathbf{d}_{\mathbb{S}}(\cdot, \cdot)$ of \mathbb{S}^{d-1} is proportional to the Euclidean distance; that is, $\|\xi - \eta\| \sim \mathbf{d}_{\mathbb{S}}(\xi, \eta)$ for all $\xi, \eta \in \mathbb{S}^{d-1}$. For the surface of the cone \mathbb{V}_0^{d+1} , however, this is no longer true when the points are near the apex. Indeed, for $(x, t), (y, s) \in \mathbb{V}_0^{d+1}$ with $x = t\xi$, $y = s\eta$, $\xi, \eta \in \mathbb{S}^{d-1}$,

$$\|(x, t) - (y, s)\|^2 = 2(t-s)^2 + 2ts(1 - \langle \xi, \eta \rangle) = 2(t-s)^2 + 2ts(1 - \cos \mathbf{d}_{\mathbb{S}}(\xi, \eta)).$$

In particular, if $t = s$, we see that $\|(t\xi, t) - (t\eta, t)\| \sim t \mathbf{d}_{\mathbb{S}}(\xi, \eta)$, whereas we have $\mathbf{d}_{\mathbb{V}_0}((t\xi, t), (t\eta, t)) \sim \sqrt{t} \mathbf{d}_{\mathbb{S}}(\xi, \eta)$ by (4.3). Hence the two distances are not compatible when t is small.

We will also need the following lemma in the estimate of the kernels.

Lemma 4.4. For $(x, t), (y, s) \in \mathbb{V}_0^{d+1}$,

$$|\sqrt{t} - \sqrt{s}| \leq \mathbf{d}_{\mathbb{V}_0}((x, t), (y, s)) \quad \text{and} \quad |\sqrt{1-t} - \sqrt{1-s}| \leq \mathbf{d}_{\mathbb{V}_0}((x, t), (y, s)).$$

Proof. Let $t = \cos^2 \frac{\theta}{2}$ and $s = \cos^2 \frac{\phi}{2}$, $0 \leq \theta, \phi \leq \pi$. Since $|\langle x, y \rangle| \leq ts$, we obtain

$$\cos \mathbf{d}_{\mathbb{V}_0}((x, t), (y, s)) \leq \sqrt{ts} + \sqrt{1-t}\sqrt{1-s} = \cos \frac{\theta - \phi}{2},$$

so that $\mathbf{d}_{\mathbb{V}_0}((x, t), (y, s)) \geq \frac{1}{2}|\theta - \phi|$. Elementary trigonometric identities shows that

$$|\sqrt{t} - \sqrt{s}| = \left| \cos \frac{\theta}{2} - \cos \frac{\phi}{2} \right| \leq 2 \sin \frac{|\theta - \phi|}{4} \leq \frac{|\theta - \phi|}{2} \leq \mathbf{d}_{\mathbb{V}_0}((x, t), (y, s)).$$

The inequality for $|\sqrt{1-t} - \sqrt{1-s}|$ follows from $|\sin \frac{\theta}{2} - \sin \frac{\phi}{2}| \leq 2 \sin \frac{|\theta - \phi|}{4}$. \square

4.2. A family of doubling weights. For the conic surface, balls are conic caps. For $r > 0$ and (x, t) on \mathbb{V}_0^{d+1} , we denote the conic cap centered at (x, t) with radius r by

$$\mathbf{c}((x, t), r) := \{(y, s) \in \mathbb{V}_0^{d+1} : d_{\mathbb{V}_0}((x, t), (y, s)) \leq r\}.$$

A weight function w is a doubling weight if it satisfies

$$w(\mathbf{c}((x, t), 2r)) \leq L w(\mathbf{c}((x, t), r)), \quad r > 0.$$

In comparison with the spherical cap on \mathbb{S}^{d-1} , the geometry of $\mathbf{c}(x, t)$ is more complicated. Denote the surface measure on \mathbb{S}^{d-1} by $d\sigma_{\mathbb{S}}$.

Lemma 4.5. *For $r > 0$, $t, s \in [0, 1]$, define $\tau_r(t, s) = (\cos r - \sqrt{1-t}\sqrt{1-s})/\sqrt{ts}$ and $\theta_r(t, s) = \arccos \tau_r(t, s)$. Then, for $(x, t) \in \mathbb{V}_0^{d+1}$ with $x = t\xi$, $\xi \in \mathbb{S}^{d-1}$,*

$$w(\mathbf{c}((x, t), r)) = \int_{d_{[0,1]}(t,s) \leq r} s^{d-1} \int_{d_{\mathbb{S}}(\xi, \eta) \leq \frac{1}{2}\theta_r(t,s)} w(s\eta, s) d\sigma_{\mathbb{S}}(\eta) ds.$$

Proof. From $d_{\mathbb{V}_0}((x, t), (y, s)) \leq r$, we obtain $d_{[0,1]}(t, s) \leq r$ by (4.2) and, with $x = t\xi$ and $y = s\eta$, it follows from (4.1) that

$$d_{\mathbb{S}}(\xi, \eta) \leq \arccos(2[\tau_r(t, s)]^2 - 1) = \frac{1}{2} \arccos \tau_r(t, s) = \frac{1}{2} \theta_r(t, s).$$

Hence, the stated identity follows from $d\sigma(y, s) = s^{d-1} d\sigma_{\mathbb{S}}(\eta) ds$. \square

For $\beta > -d$ and $\gamma > -1$, consider the Jacobi weight function defined on the cone

$$w_{\beta, \gamma}(t) = t^{\beta}(1-t)^{\gamma}, \quad 0 < t < 1.$$

Let $b_{\beta, \gamma}$ be the normalization constant so that $b_{\beta, \gamma} w_{\beta, \gamma}$ has unit integral on \mathbb{V}_0^{d+1} . Setting $y = s\eta$, $\eta \in \mathbb{S}^{d-1}$, then

$$b_{\beta, \gamma}^{-1} = \int_0^1 s^{d+\beta-1} (1-s)^{\gamma} ds \int_{\mathbb{S}^{d-1}} d\sigma_{\mathbb{S}^{d-1}}(\xi) = \omega_d \frac{\Gamma(\beta+d)\Gamma(\gamma+1)}{\Gamma(\beta+\gamma+d+1)},$$

where ω_d is the surface area of \mathbb{S}^{d-1} .

Proposition 4.6. *Let $r > 0$ and $(x, t) \in \mathbb{V}_0^{d+1}$. Then for $\beta > -d$ and $\gamma > -1$,*

$$(4.4) \quad \begin{aligned} w_{\beta, \gamma}(\mathbf{c}((x, t), r)) &:= b_{\beta, \gamma} \int_{\mathbf{c}((x, t), r)} w_{\beta, \gamma}(s) d\sigma(y, s) \\ &\sim r^d (t + r^2)^{\beta + \frac{d}{2}} (1 - t + r^2)^{\gamma + \frac{1}{2}}. \end{aligned}$$

In particular, $w_{\beta, \gamma}$ is a doubling weight and the doubling index $\alpha(w_{\beta, \gamma})$, defined in (2.1), is give by $\alpha(w_{\beta, \gamma}) = d + 2 \max\{0, \beta + \frac{d}{2}\} + 2 \max\{0, \gamma + \frac{1}{2}\}$.

Proof. Without loss of generality, we assume r is bounded by a small positive number $r \leq \delta$; for example, $\delta = \frac{\pi}{12}$ will do. By rotation symmetry, we could choose $x = te_1$, where $e_1 = (1, 0, \dots, 0)$. Then, by Lemma 4.5,

$$w_{\beta, \gamma}(\mathbf{c}((x, t), r)) = \omega_{d-1} \int_{d_{[0,1]}(t,s) \leq r} s^{d-1} w_{\beta, \gamma}(s) \int_0^{\frac{1}{2}\theta_r(t,s)} (\sin \theta)^{d-2} d\theta ds,$$

where we have used the identity (cf. [12, (A.5.1)])

$$(4.5) \quad \int_{\mathbb{S}^{d-1}} g(\langle \xi, \eta \rangle) d\sigma(\eta) = \omega_{d-1} \int_0^{\pi} g(\cos \theta) (\sin \theta)^{d-2} d\theta$$

with ω_{d-1} being the surface area of \mathbb{S}^{d-2} . Since $\theta \sim \sin \theta \sim \sqrt{1 - \cos \theta}$, it follows that

$$(4.6) \quad \mathbf{w}_{\beta, \gamma}(\mathbf{c}((x, t), r)) \sim \int_{\mathbf{d}_{[0,1]}(t, s) \leq r} s^{d-1} \mathbf{w}_{\beta, \gamma}(s) (1 - \tau_r(t, s))^{\frac{d-1}{2}} ds.$$

By its definition, $\tau_r(t, s) \geq 0$ and, furthermore, $\tau_r(t, s) \leq 1$ since we can write

$$(4.7) \quad 1 - \tau_r(t, s) = \frac{\cos \mathbf{d}_{[0,1]}(t, s) - \cos r}{\sqrt{t} \sqrt{s}}.$$

We need to consider three cases.

Case 1. Assume $3r^2 \leq t \leq 1 - 3r^2$. By Lemma 4.4, this implies that $s \sim t + r^2$ and $1 - s \sim 1 - t + r^2$, which allows us to conclude that

$$\begin{aligned} \mathbf{w}_{\beta, \gamma}(\mathbf{c}((x, t), r)) &\sim (t + r^2)^{\beta + \frac{d}{2}} (1 - t + r^2)^{\gamma + \frac{1}{2}} \\ &\times \int_{\mathbf{d}_{[0,1]}(t, s) \leq r} (\cos(\mathbf{d}_{[0,1]}(t, s)) - \cos r)^{\frac{d-1}{2}} \frac{ds}{\sqrt{s(1-s)}}. \end{aligned}$$

Setting $t = \sin^2 \frac{\theta}{2}$ and $s = \sin^2 \frac{\phi}{2}$ so that $\mathbf{d}_{[0,1]}(t, s) = |\theta - \phi|/2$ and the last integral is easily seen to be

$$\int_{|\theta - \phi| \leq 2r} (\cos \frac{\theta - \phi}{2} - \cos r)^{\frac{d-1}{2}} d\phi = c \int_{|\zeta| \leq 2r} (\sin \frac{\zeta - r}{2} \sin \frac{\zeta + r}{2})^{\frac{d-1}{2}} d\zeta \sim r^d.$$

This completes the proof of the first case.

Case 2. $0 \leq t \leq 3r^2$. For $(y, s) \in \mathbf{c}((x, t), r)$, we also have $\sqrt{s} \leq \sqrt{t} + r \leq (1 + \sqrt{3})r$ by Lemma 4.4, which shows, in particular, that $s \leq c(t + r^2)$. Evidently $1 - s \sim 1 - t \sim 1$ in this case. Furthermore, let $t = \sin^2 \frac{\theta}{2}$ and $s = \sin^2 \frac{\phi}{2}$; then

$$|s - t| = \left| \sin \frac{\theta - \phi}{2} \sin \frac{\theta + \phi}{2} \right| \leq cr^2$$

since $|\theta - \phi| \leq 2\mathbf{d}_{[0,1]}(t, s) \leq r$ and $\theta \leq cr$. Now, we have a trivial upper bound $1 - \tau_r(t, s) \leq 2$, which leads to, by (4.6),

$$\mathbf{w}_{\beta, \gamma}(\mathbf{c}((x, t), r)) \leq c \int_{\mathbf{d}_{[0,1]}(t, s) \leq r/2} s^{d+\beta-1} ds \sim \int_0^{r^2} \phi^{d+\beta-1} d\phi \sim r^{2\beta+2d},$$

which proves the upper bound in (4.4). For the lower bound, we consider a subset of $\mathbf{c}((x, t), r)$ with $\mathbf{d}_{[0,1]}(t, s) \leq r/2$. Using the upper bound of s and t , we then deduce

$$1 - \tau_r(t, s) = \frac{\cos \mathbf{d}_{[0,1]}(t, s) - \cos r}{\sqrt{t} \sqrt{s}} \geq \frac{\cos \frac{r}{2} - \cos r}{(3 + \sqrt{3})r^2} \geq \frac{2}{\pi^2},$$

where in the last step we have used the monotonicity of the function over $0 \leq r \leq \pi/12$, which shows then

$$\mathbf{w}_{\beta, \gamma}(\mathbf{c}((x, t), r)) \geq c \int_{\mathbf{d}_{[0,1]}(t, s) \leq r/2} s^{d+\beta-1} ds \sim r^{2\beta+2d}.$$

This completes the proof of this second case.

Case 3. $1 - t \leq 3r^2$. In this case, we clearly have $s \sim t \sim 1$ for $(y, s) \in \mathbf{c}((x, t), r)$. Since $\mathbf{d}_{[0,1]}(t, s) = \mathbf{d}_{[0,1]}(1 - t, 1 - s)$, changing variable $s \mapsto 1 - s$ in (4.6), we obtain

$$\mathbf{w}_{\beta, \gamma}(\mathbf{c}((x, t), r)) \sim c \int_{\mathbf{d}_{[0,1]}(t, s) \leq r} (1 - s)^\gamma ds = c \int_{\mathbf{d}_{[0,1]}(1-t, s) \leq r} s^\gamma ds,$$

where the last integral can be estimated as in Case 2. This completes the proof of (4.4). \square

It is worthwhile to mention that the proof relies on the geometry of $\mathbf{c}((x, t), r)$ when $t \leq cr^2$, which we describe in the following remark.

Remark 4.2. By (4.3), $(y, s) \in \mathbf{c}((x, t), r)$ is equivalent to

$$2 \sin^2 \frac{\mathbf{d}_{\mathbb{S}}(\xi, \eta)}{2} = \frac{\cos d_{[0,1]}(t, s) - \cos d_{\mathbb{V}_0}((x, t), (y, s))}{\sqrt{t}\sqrt{s}} \leq \frac{\cos d_{[0,1]}(t, s) - \cos r}{\sqrt{t}\sqrt{s}}.$$

If $t \leq cr^2$, then the proof of the Case 2 shows, by $\sin \theta \leq \theta$, that the above inequality holds whenever $\mathbf{d}_{\mathbb{S}}(\xi, \eta) \leq \frac{4}{\pi^2}$. In other words, for any s that satisfies $d_{[0,1]}(t, s) \leq r$, the set $\mathbf{c}((x, t), r)$ contains a large spherical cap $\{\eta : \mathbf{d}_{\mathbb{S}}(\xi, \eta) \leq \frac{4}{\pi^2}\}$.

Corollary 4.7. *For $d \geq 2$, $\beta > -d$ and $\gamma > -1$, the space $(\mathbb{V}_0^{d+1}, \mathbf{w}_{\beta, \gamma}, \mathbf{d}_{\mathbb{V}_0})$ is a homogeneous space.*

For convenience, we will introduce and use the function $\mathbf{w}_{\beta, \gamma, d}(n; t)$ defined by

$$(4.8) \quad \mathbf{w}_{\beta, \gamma, d}(n; t) := n^d \mathbf{w}_{\beta, \gamma}(\mathbf{c}((x, t), n^{-1})) = (t + n^{-2})^{\beta + \frac{d}{2}} (1 - t + n^{-2})^{\gamma + \frac{1}{2}}.$$

For any ξ in the unit sphere \mathbb{S}^{m-1} , a spherical cap $\mathbf{c}(\xi, r)$ has $\sigma(\mathbf{c}(\xi, r)) = cr^{m-1}$ so that, for a doubling weight w on the unit sphere \mathbb{S}^{m-1} , the function $w(n; \xi) = n^{m-1} \sigma(\mathbf{c}(\xi, \frac{1}{n}))$ is an approximation to $w(\xi)$ by the Lebesgue differentiation theorem. The function $\mathbf{w}_{\beta, \gamma, d}(n; t)$ in (4.8), however, is not an approximation to $\mathbf{w}_{\beta, \gamma}(t)$ on the conic surface since for the Lebesgue measure $d\sigma$ on \mathbb{V}_0^{d+1} ,

$$\sigma(\mathbf{c}((x, t), n^{-1})) \sim n^{-d} (t + n^{-2})^{\frac{d}{2}} (1 - t + n^{-2})^{\frac{1}{2}}$$

by (4.4), whereas it is the ratio

$$\mathbf{w}_n(x, t) = \frac{\mathbf{w}(\mathbf{c}((x, t), n^{-1}))}{\sigma(\mathbf{c}((x, t), n^{-1}))}$$

that provides an approximation to $\mathbf{w}(t)$ on \mathbb{V}_0^{d+1} .

For the unit sphere, it can be verified that $w_n(\xi) = w(n; \xi)$ is a doubling weight on \mathbb{S}^{m-1} whenever w is, and this property has been used to show, for example, that $\|f\|_{p, w} \sim \|f\|_{p, w_n}$ for all polynomials of degree at most n . For a doubling measure \mathbf{w} on the conic surface, it is not clear, however, if \mathbf{w}_n defined above is itself a doubling weight on \mathbb{V}_0^{d+1} .

4.3. Orthogonal polynomials on the conic surface. Orthogonal structure with respect to $\mathbf{w}_{\beta, \gamma}$ on the conic surface was studied in [53]. For $\beta > -d$ and $\gamma > -1$, define the inner product

$$\langle f, g \rangle_{\mathbf{w}} = \mathbf{b}_{\beta, \gamma} \int_{\mathbb{V}_0^{d+1}} f(x, t) g(x, t) \mathbf{w}_{\beta, \gamma} d\sigma(x, t),$$

where $d\sigma$ denote the surface measure on \mathbb{V}_0^{d+1} , which is well-defined for all polynomials restricted on the conic surface. Let $\mathcal{V}_n(\mathbb{V}_0^{d+1}, \mathbf{w}_{\beta, \gamma})$ be the space of orthogonal polynomials of degree n . Since \mathbb{V}_0^{d+1} is a quadratic surface in \mathbb{R}^{d+1} , the dimension of the space $\mathcal{V}_n(\mathbb{V}_0^{d+1}, \mathbf{w}_{\beta, \gamma})$ is the same as the dimension of the spherical harmonics of degree n on \mathbb{S}^d ; that is, $\dim \mathcal{V}_0(\mathbb{V}_0^{d+1}, \mathbf{w}_{\beta, \gamma}) = 1$ and

$$\dim \mathcal{V}_n(\mathbb{V}_0^{d+1}, \mathbf{w}_{\beta, \gamma}) = \binom{n+d-1}{n} + \binom{n+d-2}{n-1}, \quad n = 1, 2, 3, \dots$$

Furthermore, let $\Pi_n(\mathbb{V}_0^{d+1})$ denote the space of polynomials of degree at most n restricted on \mathbb{V}_0^{d+1} , then it is the union of $\mathcal{V}_n(\mathbb{V}_0^{d+1}, \mathbf{w}_{\beta, \gamma})$ and

$$\dim \Pi_n(\mathbb{V}_0^{d+1}) = \binom{n+d}{n} + \binom{n+d-1}{n-1}.$$

An orthogonal basis of $\mathcal{V}_n(\mathbb{V}_0^{d+1}, \varphi_{\beta, \gamma})$ can be given in terms of the Jacobi polynomials and spherical harmonics. Let $\mathcal{H}_m(\mathbb{S}^{d-1})$ be the space of spherical harmonics of degree m in d variables. Let $\{Y_\ell^m : 1 \leq \ell \leq \dim \mathcal{H}_m(\mathbb{S}^{d-1})\}$ denote an orthonormal basis of $\mathcal{H}_m(\mathbb{S}^{d-1})$. Then the polynomials

$$S_{m, \ell}^n(x, t) = P_{n-m}^{(2m+\beta+d-1, \gamma)}(1-2t)Y_\ell^m(x), \quad 0 \leq m \leq n, \quad 1 \leq \ell \leq \dim \mathcal{H}_m(\mathbb{S}^{d-1}),$$

consist of an orthogonal basis of $\mathcal{V}_n(\mathbb{V}_0^{d+1}, \varphi_{\beta, \gamma})$. More precisely,

$$\langle S_{m, \ell}^n, S_{m', \ell'}^{n'} \rangle_{\mathbf{w}_{\beta, \gamma}} = H_{m, n}^{\beta, \gamma} \delta_{n, n'} \delta_{m, m'} \delta_{\ell, \ell'}$$

where the norm $H_{m, n}^{\beta, \gamma}$ of $S_{m, \ell}^n$ is given by

$$H_{m, n}^{\beta, \gamma} = \frac{c_{\beta+d-1, \gamma}}{c_{2m+\beta+d-1, \gamma}} h_{n-m}^{(2m+\beta+d-1, \gamma)},$$

where $c_{\alpha, \beta}$ is the normalization constant in (2.14) and $h_m^{(\alpha, \beta)}$ is the norm square of the Jacobi polynomial. We call $S_{m, \ell}^n$ the Jacobi polynomials on the conic surface. The reproducing kernel of the space $\mathcal{V}_n(\mathbb{V}_0^{d+1}, \mathbf{w}_{\beta, \gamma})$ is denoted by $P_n(\mathbf{w}_{\beta, \gamma}; \cdot, \cdot)$, which can be written in terms of the above basis,

$$P_n(\mathbf{w}_{\beta, \gamma}; (x, t), (y, s)) = \sum_{m=0}^n \sum_{k=1}^{\dim \mathcal{H}_m^d} \frac{S_{m, k}^n(x, t) S_{m, k}^n(y, s)}{H_{m, n}^{\beta, \gamma}},$$

and it is the kernel of the orthogonal projection operator $\text{proj}_n(\mathbf{w}_{\beta, \gamma}) : L^2(\mathbb{V}_0^{d+1}, \mathbf{w}_{\beta, \gamma}) \rightarrow \mathcal{V}_n(\mathbb{V}_0^{d+1}, \varphi_{\beta, \gamma})$,

$$\text{proj}_n(\mathbf{w}_{\beta, \gamma}; f) = \int_{\mathbb{V}_0^{d+1}} f(y, s) P_n(\mathbf{w}_{\beta, \gamma}; \cdot, (y, s)) \mathbf{w}_{\beta, \gamma}(s) d\sigma(y, s).$$

The case $\beta = -1$ turns out to be the most interesting case, for which there is a second order differential operator that has orthogonal polynomials as eigenfunctions, akin the Laplace-Beltrami operator for the spherical harmonics.

Theorem 4.8. *Let $\Delta_0^{(\xi)}$ denote the Laplace-Beltrami operator in $\xi \in \mathbb{S}^{d-1}$. Define*

$$\Delta_{0, \gamma} := \left(t(1-t)\partial_t^2 + (d-1-(d+\gamma)t)\partial_t + t^{-1}\Delta_0^{(\xi)} \right)$$

for $\gamma > -1$. Then the polynomials in $\mathcal{V}_n(\mathbb{V}_0^{d+1}, \mathbf{w}_{-1, \gamma})$ are eigenfunctions of $\Delta_{0, \gamma}$,

$$\Delta_{0, \gamma} u = -n(n+\gamma+d-1)u, \quad \forall u \in \mathcal{V}_n(\mathbb{V}_0^{d+1}, \mathbf{w}_{-1, \gamma}).$$

The reproducing kernel enjoys an addition formula that is a mixture of the addition formula for the spherical harmonics and the Jacobi polynomials. The formula has the most elegant form when $\beta = -1$, which is stated below.

Theorem 4.9. *Let $d \geq 2$ and $\gamma \geq -\frac{1}{2}$. Then, for $(x, t), (y, s) \in \mathbb{V}_0^{d+1}$,*

$$(4.9) \quad \begin{aligned} P_n(\mathbf{w}_{-1,\gamma}; (x, t), (y, s)) &= b_{\gamma,d} \int_{[-1,1]^2} Z_{2n}^{\gamma+d-1}(\zeta(x, t, y, s; v)) \\ &\quad \times (1 - v_1^2)^{\frac{d-4}{2}} (1 - v_2^2)^{\gamma-\frac{1}{2}} dv, \end{aligned}$$

where $b_{\gamma,d}$ is a constant so that $P_0(\mathbf{w}_{-1,\gamma}; (x, t), (y, s)) = 1$ and

$$\zeta(x, t, y, s; v) = v_1 \sqrt{\frac{st + \langle x, y \rangle}{2}} + v_2 \sqrt{1-t} \sqrt{1-s};$$

moreover, the identity holds under the limit (2.28) when $\gamma = -\frac{1}{2}$ and/or $d = 2$.

In particular, the orthogonal structure for the weight function $\mathbf{w}_{-1,\gamma}$ on the conic surface satisfies both characteristic properties specified in Definition 3.1 and Definition 3.2. We show that $\mathbf{w}_{-1,\gamma}$ admits highly localized kernels in the next subsection.

4.4. Highly localized kernels. Let \hat{a} be an admissible cut-off function. For $(x, t), (y, s) \in \mathbb{V}_0^{d+1}$, define the kernel $L_n(\mathbf{w}_{-1,\gamma})$ by

$$L_n(\mathbf{w}_{-1,\gamma}; (x, t), (y, s)) = \sum_{j=0}^{\infty} \hat{a}\left(\frac{j}{n}\right) P_j(\mathbf{w}_{-1,\gamma}; (x, t), (y, s)).$$

We show that this kernel is highly localized. It is worthwhile to point out that $\mathbf{w}_{-1,\gamma}$ does not include the constant weight function, or the Lebesgue measure on the conic surface.

We shall need $\mathbf{w}_{-1,\gamma}(\mathbf{c}((x, t), r))$ with $r = n^{-2}$. Following the notation (4.8), we define

$$(4.10) \quad \mathbf{w}_{\gamma,d}(n; t) = (1 - t + n^{-2})^{\gamma+\frac{1}{2}} (t + n^{-2})^{\frac{d-2}{2}}.$$

Theorem 4.10. *Let $d \geq 2$ and $\gamma \geq -\frac{1}{2}$. Let \hat{a} be an admissible cutoff function. Then for any $\kappa > 0$ and $(x, t), (y, s) \in \mathbb{V}_0^{d+1}$,*

$$|L_n(\mathbf{w}_{-1,\gamma}; (x, t), (y, s))| \leq \frac{c_\kappa n^d}{\sqrt{\mathbf{w}_{\gamma,d}(n; t)} \sqrt{\mathbf{w}_{\gamma,d}(n; s)}} (1 + n d_{\mathbb{V}_0}((x, t), (y, s)))^{-\kappa}.$$

Proof. We only prove the case when $d > 2$ and $\gamma > -\frac{1}{2}$. The remaining cases $\gamma = -\frac{1}{2}$ and/or $d = 2$ follow similarly and are easier. Using the addition formula for the reproducing kernel and, by the quadratic transform (2.21),

$$(4.11) \quad Z_{2n}^\lambda(x) = \frac{P_n^{(\lambda-\frac{1}{2}, -\frac{1}{2})}(1) P_n^{(\lambda-\frac{1}{2}, -\frac{1}{2})}(2x^2 - 1)}{h_n^{(\lambda-\frac{1}{2}, -\frac{1}{2})}},$$

we can write $L(\mathbf{w}_{\beta,\gamma})$ in terms of the kernel $L_n^{(\lambda-\frac{1}{2}, -\frac{1}{2})}$ of the Jacobi polynomials, where $\lambda = \gamma + d - 1$. Then

$$(4.12) \quad \begin{aligned} L_n(\mathbf{w}_{-1,\gamma}; (x, t), (y, s)) &= c_\gamma \int_{[-1,1]^2} L_n^{(\lambda-\frac{1}{2}, -\frac{1}{2})}(2\zeta(x, t, y, s; v)^2 - 1) \\ &\quad \times (1 - v_1^2)^{\frac{d-2}{2}-1} (1 - v_2^2)^{\gamma-\frac{1}{2}} dv. \end{aligned}$$

Let $\theta(x, t, y, s; v) = \arccos(2\zeta(x, t, y, s; v)^2 - 1)$. Then

$$1 - \zeta(x, t, y, s; v)^2 = \frac{1}{2}(1 - \cos \theta(x, t, y, s; v)) = \sin^2 \frac{\theta(x, t, y, s; v)}{2} \sim \theta(x, t, y, s; v)^2.$$

We apply the estimate (2.17) for $L_n^{\alpha,\beta}$ with $\alpha = \lambda - 1/2$, $\beta = -1/2$ to obtain

$$|\mathbb{L}_n(\mathbf{w}_{-1,\gamma}; (x, t), (y, s))| \leq cn^{2\lambda+1} \int_{[-1,1]^2} \frac{1}{\left(1 + n\sqrt{1 - \zeta(x, t, y, s; v)}\right)^{\kappa+2\gamma+d+1}} \times (1 - v_1^2)^{\frac{d-2}{2}-1} (1 - v_2^2)^{\gamma-\frac{1}{2}} dv.$$

From its definition, it is easy to verify that $|\zeta(x, t, y, s; v)| \leq 1$ and

$$1 - \zeta(x, t, y, s; v) = 1 - \cos \mathbf{d}_{\mathbb{V}_0}((x, t), (y, s)) + \sqrt{\frac{\langle x, y \rangle + ts}{2}}(1 - v_1) + \sqrt{1 - s}\sqrt{1 - t}(1 - v_2).$$

Consequently, since $ts + \langle x, y \rangle \geq 0$, it follows that

$$\begin{aligned} 1 - \zeta(x, t, y, s; v) &\geq 1 - \cos \mathbf{d}_{\mathbb{V}_0}((x, t), (y, s)) \\ &= 2 \sin^2 \frac{\mathbf{d}_{\mathbb{V}_0}((x, t), (y, s))}{2} \geq \frac{2}{\pi^2} [\mathbf{d}_{\mathbb{V}_0}((x, t), (y, s))]^2. \end{aligned}$$

Applying this inequality, we obtain the estimate

$$\begin{aligned} |\mathbb{L}_n(\mathbf{w}_{-1,\gamma}; (x, t), (y, s))| &\leq cn^{2\lambda+1} \frac{1}{(1 + n\mathbf{d}_{\mathbb{V}_0}((x, t), (y, s)))^\kappa} \\ &\times b_{\gamma,d} \int_{[-1,1]^2} \frac{(1 - v_1^2)^{\frac{d-2}{2}-1} (1 - v_2^2)^{\gamma-\frac{1}{2}}}{\left(1 + n\sqrt{1 - \zeta(x, t, y, s; v)}\right)^{2\gamma+d+1}} dv, \end{aligned}$$

where $b_{\gamma,d} = c_{\gamma-\frac{1}{2}, \gamma-\frac{1}{2}} c_{\frac{d-4}{2}, \frac{d-4}{2}}$. The estimate of the last expression is the crux of the proof and it is contained in the lemma below. \square

Lemma 4.11. *Let $d \geq 2$ and $\gamma > -\frac{1}{2}$. Then, for $\beta \geq 2\gamma + d + 1$,*

$$\begin{aligned} b_{\gamma,d} \int_{[-1,1]^2} \frac{(1 - v_1)^{\frac{d-2}{2}-1} (1 - v_2)^{\gamma-\frac{1}{2}}}{\left(1 + n\sqrt{1 - \zeta(x, t, y, s; v)}\right)^\beta} dv \\ \leq \frac{cn^{-(2\gamma+d-1)}}{\sqrt{\mathbf{w}_{\gamma,d}(n; t)} \sqrt{\mathbf{w}_{\gamma,d}(n; s)} (1 + n\mathbf{d}_{\mathbb{V}_0}((x, t), (y, s)))^{\beta-3\gamma-\frac{3d+1}{2}}}. \end{aligned}$$

Proof. Using the lower bound of $1 - |\zeta(x, t, y, s; v)|$ in the proof of the previous theorem, the left-hand side of the stated inequality has the upper bound

$$\frac{c}{(1 + n\mathbf{d}_{\mathbb{V}_0}((x, t), (y, s)))^{\beta-2\gamma-d-1}} \int_{[-1,1]^2} \frac{(1 - v_1)^{\frac{d-2}{2}-1} (1 - v_2)^{\gamma-\frac{1}{2}}}{\left(1 + n\sqrt{1 - \zeta(x, t, y, s; v)}\right)^{2\gamma+d+1}} dv.$$

Denote the above integral by $I(x, t, y, s)$. To complete the proof, we need to show that

$$\frac{I(x, t, y, s)}{(1 + n\mathbf{d}_{\mathbb{V}_0}((x, t), (y, s)))^{\gamma+\frac{d-1}{2}}} \leq \frac{cn^{-(2\gamma+d-1)}}{\sqrt{\mathbf{w}_{\gamma}(n; t)} \sqrt{\mathbf{w}_{\gamma}(n; s)}}.$$

We need a lower bound for $1 - \zeta$. Let $\alpha = \sqrt{\frac{\langle x, y \rangle + ts}{2}}$ and $\beta = \sqrt{1 - t}\sqrt{1 - s}$. Then

$$\begin{aligned} 1 - \zeta(x, t, y, s; v) &= 1 - \alpha + (1 - v_1)\alpha - v_2\beta \geq 1 - \alpha + \frac{1}{2}(1 - v_1)\alpha - v_2\beta \\ &= 1 - \frac{1 + v_1}{2}\alpha - \frac{1 - v_1}{2}\sqrt{ts} + \frac{1 - v_1}{2}\sqrt{ts} - v_2\beta. \end{aligned}$$

Using $\alpha \leq \sqrt{ts}$ and $1 - \sqrt{ts} - \sqrt{1-t}\sqrt{1-s} \geq 0$, we then obtain

$$(4.13) \quad \begin{aligned} 1 - \zeta(x, t, y, s; v) &\geq 1 - \sqrt{ts} - \beta + \frac{1}{2}(1 - v_1)\sqrt{ts} + (1 - v_2)\beta \\ &\geq \frac{1}{2}(1 - v_1)\sqrt{t}\sqrt{s} + (1 - v_2)\sqrt{1-s}\sqrt{1-t}. \end{aligned}$$

Using this inequality in $I(x, t, y, s)$ and then making a change of variable $v_1 \mapsto 2u_1 - 1$ and $v_2 \mapsto 2u_2 - 1$, we obtain

$$I(x, t, y, s) \leq c \int_{[0,1]^2} \frac{(1 - v_1)^{\frac{d-2}{2}-1} (1 - v_2)^{\gamma-\frac{1}{2}}}{(1 + n\sqrt{1 - (1 - v_1)\sqrt{t}\sqrt{s} - 2(1 - v_2)\sqrt{1-s}\sqrt{1-t}})^{2\gamma+d+1}} dv.$$

This integral can be estimated by using the inequality [12, (13.5.8)]

$$(4.14) \quad \int_0^1 \frac{(1-t)^{a-1} dt}{(1 + n\sqrt{B + A(1-t)})^b} \leq c \frac{n^{-2a}}{A^a (1 + n\sqrt{B})^{b-2a-1}},$$

which holds for $A > 0$, $B \geq 0$, $a > 0$ and $b \geq 2a + 1$. Applying this inequality with $A = \sqrt{1-t}\sqrt{1-s}$, $a = \gamma + \frac{1}{2}$, we obtain

$$I(x, t, y, s) \leq \frac{cn^{-2\gamma-1}}{(\sqrt{1-t}\sqrt{1-s})^{\gamma+\frac{1}{2}}} \int_0^1 \frac{(1 - v_1^2)^{\frac{d-2}{2}-1}}{\left(1 + n\sqrt{\sqrt{t}\sqrt{s}(1 - v_1)}\right)^{d-1}} dv_1.$$

Applying (4.14) one more time with $B = 0$, $a = \frac{d-2}{2}$ and $b = d - 1$, we see that the last integral is bounded by $cn^{-(d-2)}$, hence,

$$\begin{aligned} I(x, t, y, s) &\leq \frac{cn^{-(2\gamma+d-1)}}{(\sqrt{1-t}\sqrt{1-s})^{\gamma+\frac{1}{2}} (\sqrt{t}\sqrt{s})^{\frac{d-2}{2}}} \\ &\leq \frac{cn^{-(2\gamma+d-1)}}{(\sqrt{1-t}\sqrt{1-s} + n^{-2})^{\gamma+\frac{1}{2}} (\sqrt{t}\sqrt{s} + n^{-2})^{\frac{d-2}{2}}}, \end{aligned}$$

where the second inequality follows since $I(x, t, y, s) \leq 1$ holds trivially by the choice of $b_{\gamma,d}$. Now, using the elementary identity [12, (11.5.13)]

$$(4.15) \quad (a + n^{-1})(b + n^{-1}) \leq 3(ab + n^{-2})(1 + n|b - a|)$$

with $a = \sqrt{t}$ and $b = \sqrt{s}$ as well as with $a = \sqrt{1-t}$ and $b = \sqrt{1-s}$, we obtain

$$I(x, t, y, s) \leq \frac{cn^{-(2\gamma+d-1)} (1 + n|\sqrt{t} - \sqrt{s}|)^{\frac{d-2}{2}} (1 + n|\sqrt{1-t} - \sqrt{1-s}|)^{\gamma+\frac{1}{2}}}{\sqrt{\mathbf{w}_{\gamma,d}(n; t)} \sqrt{\mathbf{w}_{\gamma,d}(n; s)}}.$$

Finally, by Lemma 4.4, we conclude that

$$I(x, t, y, s) \leq \frac{cn^{-(2\gamma+d-1)} (1 + n\mathbf{d}_{V_0}((x, t), (y, s)))^{\gamma+\frac{d-1}{2}}}{\sqrt{\mathbf{w}_{\gamma,d}(n; t)} \sqrt{\mathbf{w}_{\gamma,d}(n; s)}},$$

which is what we need to complete the proof. \square

The following corollary, following immediately from (4.13), will be used later.

Corollary 4.12. *Let $d \geq 2$ and $\gamma > -\frac{1}{2}$. Then, for $\beta \geq 2\gamma + d + 1$,*

$$\begin{aligned} b_{\gamma,d} \int_{[-1,1]^2} \frac{(1-v_1)^{\frac{d-2}{2}-1} (1-v_2)^{\gamma-\frac{1}{2}}}{\left(1 + n\sqrt{1-(1-v_1)\sqrt{t}\sqrt{s}-(1-v_2)\sqrt{1-s}\sqrt{1-t}}\right)^{2\gamma+d+1}} dv \\ \leq \frac{cn^{-(2\gamma+d-1)} (1 + nd_{[0,1]}((x,t), (y,s)))^{\gamma+\frac{d-1}{2}}}{\sqrt{\mathbf{w}_{\gamma,d}(n;t)} \sqrt{\mathbf{w}_{\gamma,d}(n;s)}}. \end{aligned}$$

By (4.8), we have shown that $\mathbf{w}_{-1,\gamma}$ admits Assertion 1 of the highly localized kernel. Our next result shows that it also admits Assertion 2.

Theorem 4.13. *Let \hat{a} be an admissible cutoff function. Let $d \geq 2$ and $\gamma \geq -\frac{1}{2}$. For $(x_i, t_i), (y, s) \in \mathbb{V}_0^{d+1}$ and $(x_1, t_1) \in \mathbf{c}((x_2, t_2), c^*n^{-1})$ with c^* small and for $\kappa > 0$,*

$$(4.16) \quad \begin{aligned} & |\mathbf{L}_n(\mathbf{w}_{-1,\gamma}; (x_1, t_1), (y, s)) - \mathbf{L}_n(\mathbf{w}_{-1,\gamma}; (x_2, t_2), (y, s))| \\ & \leq c_\kappa \frac{n^{d+1} \mathbf{d}_{\mathbb{V}_0}((x_1, t_1), (x_2, t_2))}{\sqrt{\mathbf{w}_{\gamma,d}(n;s)} \sqrt{\mathbf{w}_{\gamma,d}(n;t_2)} (1 + nd_{\mathbb{V}_0}((y, s), (x_2, t_2)))^\kappa}. \end{aligned}$$

Proof. Denote the left-hand side of (4.17) by K . Let $\partial L(u) = L'(u)$. Using the integral expression (4.12) of $\mathbf{L}_n(\mathbf{w}_{-1,\gamma})$, we obtain

$$(4.17) \quad \begin{aligned} K & \leq 2 \int_{[-1,1]^2} \left\| \partial L_n^{\lambda-\frac{1}{2}, -\frac{1}{2}}(2(\cdot)^2 - 1) \right\|_{L^\infty(I_v)} |\zeta_1(v)^2 - \zeta_2(v)^2| \\ & \quad \times (1-v_1^2)^{\frac{d-2}{2}-1} (1-v_2^2)^{\gamma-\frac{1}{2}} dv, \end{aligned}$$

where $\zeta_i(v) = \zeta(x_i, t_i, y, s; v)$, and I_v is the interval with end points $\zeta_1(v)$ and $\zeta_2(v)$. Since $|\zeta(\cdot)| \leq 1$, $|\zeta_1(v)^2 - \zeta_2(v)^2| \leq 2|\zeta_1(v) - \zeta_2(v)|$. We claim that

$$(4.18) \quad |\zeta_1(v) - \zeta_2(v)| \leq c \mathbf{d}_{\mathbb{V}_0}((x_1, t_1), (x_2, t_2)) (\Sigma_1 + \Sigma_2(v_1) + \Sigma_3(v_2)),$$

where

$$\begin{aligned} \Sigma_1 &= \mathbf{d}_{\mathbb{V}_0}((x_i, t_i), (y, s)) + \mathbf{d}_{\mathbb{V}_0}((x_1, t_1), (x_2, t_2)), \\ \Sigma_2(v_1) &= (1-v_1)\sqrt{s}, \\ \Sigma_3(v_2) &= (1-v_2)\sqrt{1-s}. \end{aligned}$$

To see this, we use (4.2) and writing $x_i = t_i \xi_i$ and $y = s\eta$ to obtain

$$\begin{aligned} \zeta_1(v) - \zeta_2(v) &= \cos \mathbf{d}_{\mathbb{V}_0}((x_1, t_1), (y, s)) - \cos \mathbf{d}_{\mathbb{V}_0}((x_2, t_2), (y, s)) \\ &+ (1-v_1) \left(\sqrt{t_2 s} \cos \frac{\mathbf{d}_{\mathbb{S}}(\xi_2, \eta)}{2} - \sqrt{t_1 s} \cos \frac{\mathbf{d}_{\mathbb{S}}(\xi_1, \eta)}{2} \right) \\ &+ (1-v_2) (\sqrt{1-t_2} - \sqrt{1-t_1}) \sqrt{1-s}. \end{aligned}$$

Denote temporarily $\alpha_i = \mathbf{d}_{\mathbb{V}_0}((x_i, t_i), (y, s))$ for $i = 1, 2$. Hence, using the identity

$$\begin{aligned} \cos \alpha_1 - \cos \alpha_2 &= 2 \sin \frac{\alpha_1 - \alpha_2}{2} \sin \frac{\alpha_1 + \alpha_2}{2} \\ &= 2 \sin \frac{\alpha_1 - \alpha_2}{2} \left(2 \sin \frac{\alpha_i}{2} + \sin \frac{|\alpha_1 - \alpha_2|}{2} \right), \end{aligned}$$

it follows readily that

$$|\cos \alpha_1 - \cos \alpha_2| \leq |\alpha_1 - \alpha_2| (|\alpha_1| + \frac{1}{2}|\alpha_1 - \alpha_2|).$$

By the triangle inequality of $\mathbf{d}_{\mathbb{V}_0}$, $|\alpha_1 - \alpha_2| \leq \mathbf{d}_{\mathbb{V}_0}((x_1, t_1), (x_2, t_2))$, this gives the estimate for the Σ_1 term. Moreover, assuming $t_2 \geq t_1$, for example, and applying similar argument for $\mathbf{d}_{\mathbb{S}}$, we then obtain, using Lemma 4.4 and (4.3), that

$$\begin{aligned} & \left| \sqrt{t_1} \cos \frac{\mathbf{d}_{\mathbb{S}}(\xi_1, \eta)}{2} - \sqrt{t_2} \cos \frac{\mathbf{d}_{\mathbb{S}}(\xi_2, \eta)}{2} \right| \\ & \leq |\sqrt{t_2} - \sqrt{t_1}| + \sqrt{t_1} \left| \cos \frac{\mathbf{d}_{\mathbb{S}}(\xi_1, \eta)}{2} - \cos \frac{\mathbf{d}_{\mathbb{S}}(\xi_2, \eta)}{2} \right| \\ & \leq \mathbf{d}_{\mathbb{V}_0}((x_1, t_1), (x_2, t_2)) + \frac{1}{2} (t_1 t_2)^{\frac{1}{4}} \mathbf{d}_{\mathbb{S}}(\xi_1, \xi_2) \\ & \leq c \mathbf{d}_{\mathbb{V}_0}((x_1, t_1), (x_2, t_2)), \end{aligned}$$

which verifies the $\Sigma_2(v_1)$ term. The third term with $\Sigma_3(v_2)$ follows from using Lemma 4.4 one more time. This verifies the claim (4.18).

Since $\max_{r \in I_v} |1 + n\sqrt{1-r}|^{-\sigma}$ is attained at one of the end points of the interval, it follows from (2.18) with $m = 1$.

$$\| \partial L_n^{\lambda-\frac{1}{2}, -\frac{1}{2}} (2(\cdot)^2 - 1) \|_{L^\infty(I_v)} \leq c \left[\frac{n^{2\lambda+3}}{(1 + n\sqrt{1-\zeta_1(v)^2})^\kappa} + \frac{n^{2\lambda+3}}{(1 + n\sqrt{1-\zeta_2(v)^2})^\kappa} \right].$$

Consequently, we see that K is bounded by a sum of integrals

$$\begin{aligned} K & \leq c \mathbf{d}_{\mathbb{V}_0}((x_1, t_1), (x_2, t_2)) \int_{[-1,1]^2} \left[\frac{n^{2\lambda+3}}{(1 + n\sqrt{1-\zeta_1(v)^2})^\kappa} + \frac{n^{2\lambda+3}}{(1 + n\sqrt{1-\zeta_2(v)^2})^\kappa} \right] \\ & \quad \times (\Sigma_1 + \Sigma_2(v) + \Sigma_3(v)) (1 - v_1^2)^{\frac{d-2}{2}-1} (1 - v_2^2)^{\gamma-\frac{1}{2}} dv. \end{aligned}$$

Since $(x_1, t_1) \in \mathbf{c}((x_2, t_2), c^* n^{-1})$, Σ_1 is bounded by $\Sigma_1 \leq c n^{-1} (1 + n \mathbf{d}_{\mathbb{V}_0}((x_i, t_i), (y, s)))$. Hence, we obtain from Lemma 4.11 that

$$\begin{aligned} & \int_{[-1,1]^2} \frac{n^{2\lambda+3}}{(1 + n\sqrt{1-\zeta_i(v)^2})^\kappa} \Sigma_1 (1 - v_1^2)^{\frac{d-2}{2}-1} (1 - v_2^2)^{\gamma-\frac{1}{2}} dv \\ & \leq c \frac{n^{d+1}}{\sqrt{\mathbf{w}_{\gamma,d}(n; s)} \sqrt{\mathbf{w}_{\gamma,d}(n; t_i)} (1 + n \mathbf{d}_{\mathbb{V}_0}((y, s), (x_i, t_i)))^{\kappa(\gamma, d)-1}}, \end{aligned}$$

where $\kappa(\gamma, d) = \kappa - 3\gamma - \frac{3d+1}{2}$ for either $i = 1$ or $i = 2$. Since $\mathbf{w}_{\gamma,d}(n, t_1) \sim \mathbf{w}_{\gamma,d}(n, t_2)$ and $\mathbf{d}_{\mathbb{V}_0}((x_1, t_1), (y, s)) + n^{-1} \sim \mathbf{d}_{\mathbb{V}_0}((x_2, t_2), (y, s)) + n^{-1}$ by Lemma 4.4, we can replace (x_1, t_1) in the right-hand side by (x_2, t_2) . This shows that the integral containing Σ_1 has the desired estimate.

For the remaining integrals, the same consideration shows that we only need to consider those containing $\zeta_2(v)$. For the integral that contains $\Sigma_2(v_1) = (1 - v_1)\sqrt{s}$, the factor $(1 - v_1)$ increases the power of the weight to $(1 - v_1)^{\frac{d}{2}}$, so that we can apply Lemma 4.11 with $\frac{d-2}{2}$ replaced by $\frac{d}{2}$, which leads to

$$\begin{aligned} & \int_{-1}^1 \frac{n^{2\lambda+3}}{(1 + n\sqrt{1-\zeta_2(v)^2})^\kappa} \Sigma_2(v_1) (1 - v_1^2)^{\frac{d-2}{2}-1} (1 - v_2^2)^{\gamma-\frac{1}{2}} dv \\ & \leq c \frac{n^{d+1} n^{-1} \sqrt{s}}{\sqrt{\mathbf{w}_{\gamma,d+1}(n; s)} \sqrt{\mathbf{w}_{\gamma,d+1}(n; t_2)} (1 + n \mathbf{d}_{\mathbb{V}_0}((y, s), (x_2, t_2)))^{\kappa(\gamma, d+1)}} \\ & \leq c \frac{n^{d+1}}{\sqrt{\mathbf{w}_{\gamma,d}(n; s)} \sqrt{\mathbf{w}_{\gamma,d}(n; t_2)} (1 + n \mathbf{d}_{\mathbb{V}_0}((y, s), (x_2, t_2)))^{\kappa(\gamma, d+1)}}, \end{aligned}$$

where the last step follows from the inequality $n^{-1}\sqrt{s} \leq (\sqrt{t_2} + n^{-1})(\sqrt{s} + n^{-1})$. The integral that contains $\Sigma_3(v_2)$ can be estimated similarly by applying Lemma 4.11 with γ replaced by $\gamma + 1$ and using $n^{-1}\sqrt{1-s} \leq (\sqrt{1-t_2} + n^{-1})(\sqrt{1-s} + n^{-1})$. This completes the proof. \square

The first two assertions for the highly localized kernels are established for $w_{-1,\gamma}$ when $\gamma \geq -\frac{1}{2}$. The case of $p = 1$ of the following lemma establishes Assertion 3. Recall that $w_{\beta,\gamma,d}(n;t)$ is defined in (4.8).

Lemma 4.14. *Let $d \geq 2$, $\beta > -d$ and $\gamma > -1$. For $0 < p < \infty$, assume $\kappa > \frac{2d}{p} + (\gamma + \beta + \frac{d+1}{2})|\frac{1}{p} - \frac{1}{2}|$. Then for $(x,t) \in \mathbb{V}_0^{d+1}$,*

$$(4.19) \quad \int_{\mathbb{V}_0^{d+1}} \frac{w_{\beta,\gamma}(s) d\sigma(y,s)}{w_{\beta,\gamma,d}(n;s)^{\frac{p}{2}} (1 + nd_{\mathbb{V}_0}((x,t), (y,s)))^{\kappa p}} \leq cn^{-d} w_{\beta,\gamma,d}(n;t)^{1-\frac{p}{2}}.$$

Proof. Let J_p denote the left-hand side of (4.19). By Lemma 2.4, it is sufficient to estimate J_2 . Let $x = t\xi$ and $y = s\eta$. The definition of $d_{\mathbb{V}_0}(\cdot, \cdot)$ shows that it is a function of $\langle \xi, \eta \rangle$. Hence, by (4.5), we deduce that

$$\begin{aligned} J_{2,\kappa} &\leq c \int_0^1 \int_{-1}^1 \frac{s^{d-1} w_{\beta,\gamma}(s) (1-u^2)^{\frac{d-3}{2}}}{w_{\beta,\gamma,d}(n;s) (1 + n \arccos(\sqrt{ts}\sqrt{\frac{1+u}{2}} + \sqrt{1-t}\sqrt{1-s}))^{2\kappa}} du ds \\ &\leq c \int_0^1 \int_0^1 \frac{s^{d-1} w_{\beta,\gamma}(s) v^{d-2} (1-v^2)^{\frac{d-3}{2}}}{w_{\beta,\gamma,d}(n;s) (1 + n\sqrt{1-tsv} - \sqrt{1-t}\sqrt{1-s})^{2\kappa}} dv ds, \end{aligned}$$

where the second step follows from changing variable $\sqrt{\frac{1+u}{2}} \mapsto v$ and the relation $\theta \sim \sin \frac{\theta}{2} \sim \sqrt{1-\cos \theta}$. Making a further changing of variable $v \mapsto z/\sqrt{s}$ gives

$$\begin{aligned} J_2 &\leq c \int_0^1 \int_0^{\sqrt{s}} \frac{s w_{\beta,\gamma}(s) z^{d-2} (s-z^2)^{\frac{d-3}{2}}}{w_{\beta,\gamma,d}(n;s) (1 + n\sqrt{1-tz} - \sqrt{1-t}\sqrt{1-s})^{2\kappa}} dz ds \\ &\leq c \int_0^1 \int_0^{\sqrt{s}} \frac{(s-z^2)^{\frac{d-3}{2}}}{(1-s+n^{-2})^{\frac{1}{2}} (1 + n\sqrt{1-tz} - \sqrt{1-t}\sqrt{1-s})^{2\kappa}} dz ds, \end{aligned}$$

where we have used $sz^{d-2} \leq s^{\frac{d}{2}} \leq (s+n^{-2})^{\frac{d}{2}}$. One more change of variable $s \mapsto 1-w^2$ with $ds = wdw$ and $w \leq (w^2 + n^{-2})^{\frac{1}{2}}$, we obtain

$$J_2 \leq c \int_0^1 \int_0^{\sqrt{1-w^2}} \frac{(1-w^2-z^2)^{\frac{d-3}{2}}}{(1 + n\sqrt{1-tz} - \sqrt{1-t}w)^{2\kappa}} dz dw,$$

which is an integral over the positive quadrant $\{(z,w) \in \mathbb{B}^2 : w \geq 0, z \geq 0\}$ of the unit disk \mathbb{B}^2 . Setting $p = \sqrt{t}z + \sqrt{1-t}w$ and $q = -\sqrt{1-t}z + \sqrt{t}w$ in the integral, which is an orthogonal transformation, and enlarging the integral domain while taking into account that $p \geq 0$, it follows that

$$\begin{aligned} J_2 &\leq c \int_0^1 \frac{1}{(1 + n\sqrt{1-p})^{2\kappa}} \int_{-\sqrt{1-p^2}}^{\sqrt{1-p^2}} (1-p^2-q^2)^{\frac{d-3}{2}} dq dp \\ &\leq c \int_0^1 \frac{(1-p^2)^{\frac{d-2}{2}}}{(1 + n\sqrt{1-p})^{2\kappa}} \leq cn^{-d} \int_0^n \frac{r^{d-1}}{(1+r)^\kappa} dr \leq cn^{-d} \end{aligned}$$

by setting $r = n\sqrt{1-p}$ and recalling that $\kappa > d$. This completes the proof. \square

Proposition 4.15. *Let $d \geq 2$ and $\gamma \geq -\frac{1}{2}$. For $0 < p < \infty$ and $(x, t) \in \mathbb{V}_0^{d+1}$,*

$$\int_{\mathbb{V}_0^{d+1}} |\mathbb{L}_n(\mathbf{w}_{-1,\gamma}; (x, t), (y, s))|^p \mathbf{w}_{-1,\gamma}(s) d\sigma(y, s) \leq c \left(\frac{n^d}{\mathbf{w}_{\gamma,d}(n; t)} \right)^{p-1}.$$

The proof is immediate by (i) of Theorem 4.10 and Lemma 4.14.

Corollary 4.16. *For $\gamma \geq -\frac{1}{2}$, the space $(\mathbb{V}_0^{d+1}, \mathbf{w}_{-1,\gamma}, d_{\mathbb{V}_0})$ is a localizable homogeneous space.*

4.5. Maximal ε -separated sets and MZ inequality. Let $\varepsilon > 0$. An ε -separated set is defined in Definition 2.13. In this subsection, we provide a construction of some examples of maximal ε -separated points on the conic surface.

For our construction, we shall need ε -separated points on the unit sphere \mathbb{S}^{d-1} . We adopt the following notation. For $\varepsilon > 0$, we denote by $\Xi_{\mathbb{S}}(\varepsilon)$ a maximal ε -separated set on the unit sphere \mathbb{S}^{d-1} and we let $\mathbb{S}_{\xi}(\varepsilon)$ be the subsets in \mathbb{S}^{d-1} so that the collection $\{\mathbb{S}_{\xi}(\varepsilon) : \xi \in \Xi_{\mathbb{S}}(\varepsilon)\}$ is a partition of \mathbb{S}^{d-1} , and we assume

$$(4.20) \quad \mathbb{c}_{\mathbb{S}}(\xi, c_1\varepsilon) \subset \mathbb{S}_{\xi}(\varepsilon) \subset \mathbb{c}_{\mathbb{S}}(\xi, c_2\varepsilon), \quad \xi \in \Xi_{\mathbb{S}}(\varepsilon),$$

where $\mathbb{c}_{\mathbb{S}}(\xi, \varepsilon)$ denotes the spherical cap centered at ξ with radius ε , c_1 and c_2 depending only on d . Such a $\Xi_{\mathbb{S}}(\varepsilon)$ exists for all $\varepsilon > 0$, see for example [12, Section 6.4], and its cardinality satisfies

$$(4.21) \quad c'_d \varepsilon^{-d+1} \leq \#\Xi_{\mathbb{S}}(\varepsilon) \leq c_d \varepsilon^{-d+1}.$$

We now consider the subsets of points on the cone \mathbb{V}_0^{d+1} . For $\varepsilon > 0$, we denote by $\Xi_{\mathbb{V}_0} = \Xi_{\mathbb{V}_0}(\varepsilon)$ an ε -separated set and denote by $\{\mathbb{V}_0(\xi, t) : (t\xi, t) \in \Xi_{\mathbb{V}_0}\}$ a partition of \mathbb{V}_0^{d+1} . We start with an explicit construction of such $\Xi_{\mathbb{V}_0}$ and $\mathbb{V}_0(\xi, t)$.

Let $\varepsilon > 0$ and let $N = \lfloor \frac{\pi}{2\varepsilon} \rfloor$. For $1 \leq j \leq N$ we define

$$\theta_j := \frac{(2j-1)\pi}{2N}, \quad \theta_j^- := \theta_j - \frac{\pi}{2N} \quad \text{and} \quad \theta_j^+ := \theta_j + \frac{\pi}{2N}.$$

Let $t_j = \sin^2 \frac{\theta_j}{2}$ and define t_j^- and t_j^+ accordingly. In particular, $t_1^- = 0$ and $t_N^+ = 1$. Then $\theta_{j+1}^- = \theta_j^+$ and \mathbb{V}_0^{d+1} can be partitioned by

$$\mathbb{V}_0^{d+1} = \bigcup_{j=1}^N \mathbb{V}_0^{(j)}, \quad \text{where} \quad \mathbb{V}_0^{(j)} := \{(x, t) \in \mathbb{V}_0^{d+1} : t_j^- < t \leq t_j^+\}.$$

Let $\varepsilon_j := (2\sqrt{t_j})^{-1}\pi\varepsilon$. Then $\Xi_{\mathbb{S}}(\varepsilon_j)$ is the maximal ε_j -separated set of \mathbb{S}^{d-1} such that $\{\mathbb{S}_{\xi}(\varepsilon_j) : \xi \in \Xi_{\mathbb{S}}(\varepsilon_j)\}$ is a partition $\mathbb{S}^{d-1} = \bigcup_{\eta \in \Xi_{\mathbb{S}}(\varepsilon_j)} \mathbb{S}_{\eta}(\varepsilon_j)$, and

$$\#\Xi_{\mathbb{S}}(\varepsilon_j) \sim \varepsilon_j^{-d+1}.$$

For each $j = 1, \dots, N$, we decompose $\mathbb{V}_0^{(j)}$ by

$$\mathbb{V}_0^{(j)} = \bigcup_{\xi \in \Xi_{\mathbb{S}}(\varepsilon_j)} \mathbb{V}_0(\xi, t_j), \quad \text{where} \quad \mathbb{V}_0(\xi, t_j) := \{(t\xi, t) : t_j^- < t \leq t_j^+, \eta \in \Xi_{\mathbb{S}}(\varepsilon_j)\}.$$

Finally, we define the subset $\Xi_{\mathbb{V}_0}$ of \mathbb{V}_0^{d+1} by

$$\Xi_{\mathbb{V}_0} = \{(t_j\xi, t_j) : \xi \in \Xi_{\mathbb{S}}(\varepsilon_j), 1 \leq j \leq N\}.$$

Proposition 4.17. *Let $\varepsilon > 0$ and $N = \lfloor \frac{\pi}{2\varepsilon} \rfloor$. Then $\Xi_{\mathbb{V}_0}$ is a maximal ε -separated set of \mathbb{V}_0^{d+1} and $\{\mathbb{V}_0(\xi, t_j) : \xi \in \Xi_{\mathbb{S}}(\varepsilon_j), 1 \leq j \leq N\}$ is a partition*

$$\mathbb{V}_0^{d+1} = \bigcup_{j=1}^N \bigcup_{\xi \in \Xi_{\mathbb{S}}(\varepsilon_j)} \mathbb{V}_0(\xi, t_j).$$

Moreover, there are positive constants c_1 and c_2 depending only on d such that

$$(4.22) \quad \mathbf{c}((t_j\xi, t_j), c_1\varepsilon) \subset \mathbb{V}_0(\xi, t_j) \subset \mathbf{c}((t_j\xi, t_j), c_2\varepsilon),$$

and c'_d and c_d such that

$$c'_d\varepsilon^{-d} \leq \#\Xi_{\mathbb{V}_0} \leq c_d\varepsilon^{-d}.$$

Proof. Let $(t_j\xi, t_j)$ and $(t_k\eta, t_k)$ be two distinct points in $\Xi_{\mathbb{V}_0}$. If $t_j \neq t_k$, then

$$\mathbf{d}_{\mathbb{V}_0}((t_j\xi, t_j), (t_k\eta, t_k)) \geq \mathbf{d}_{[0,1]}(t_j, t_k) = \frac{1}{2}|\theta_j - \theta_k| \geq \frac{\pi}{2N} \geq \varepsilon.$$

If $j = k$, then ξ and η are both elements of $\mathbb{S}(\varepsilon_j)$, so that $\mathbf{d}_{\mathbb{S}}(\xi, \eta) \geq \varepsilon_j$. Hence, using $\frac{2}{\pi}\phi \leq \sin \phi \leq \phi$, we deduce from (4.2) that

$$\mathbf{d}_{\mathbb{V}_0}((t_j\xi, t_j), (t_j\eta, t_j)) \geq \frac{2}{\pi}\sqrt{t_j}\mathbf{d}_{\mathbb{S}}(\xi, \eta) \geq \frac{2}{\pi}\sqrt{t_j}\varepsilon_j = \varepsilon.$$

Hence, $\Xi_{\mathbb{V}_0}$ is ε -separated. Moreover, since $\#\Xi_{\mathbb{S}}(\varepsilon_j) \sim \varepsilon_j^{-d+1}$ and $\varepsilon_j \sim \varepsilon/\theta_j$, it follows that

$$\#\Xi_{\mathbb{V}_0} = \sum_{j=1}^N \#\Xi_{\mathbb{S}}(\varepsilon_j) \sim \sum_{j=1}^N \varepsilon_j^{-d+1} \sim \varepsilon^{-d+1} \sum_{j=1}^N \theta_j^{d-1} \sim \varepsilon^{-d+1} N \sim \varepsilon^{-d}.$$

For the proof of (4.22), we first consider the ball $\mathbf{c}_{[0,1]}(t_j, r) = \{s : \mathbf{d}_{[0,1]}(s, t_j) \leq r\}$ on $[0, 1]$. For $0 < \delta < \pi/2$, it is easy to see that

$$\mathbf{c}_{[0,1]}(t_j, \delta/N) \subset \{s : (s\eta, s) \in \mathbb{V}_0(\xi, t_j)\} \subset \mathbf{c}_{[0,1]}(t_j, \pi/N).$$

We further choose δ so that $4\delta(1+\delta) < \frac{1}{2}$. For $s \in \mathbf{c}_{[0,1]}(t_j, \delta/N)$, write $s = \sin^2 \frac{\phi}{2}$, then

$$|t_j - s| = \frac{1}{2}|\cos \theta - \cos \phi| = \left| \sin \frac{\theta_j - \phi}{2} \sin \frac{\theta_j + \phi}{2} \right| \leq \frac{\delta}{N} \left(2\sqrt{t_j} + \frac{\delta}{N} \right),$$

where we have used $|\sin \frac{\theta_j + \phi}{2}| \leq 2 \sin \frac{\theta_j}{2} + |\sin \frac{\theta_j - \phi}{2}| \leq 2\sqrt{t_j} + \mathbf{d}_{[0,1]}(s, t_j)$. For $j \geq 1$, $N^{-1} = \frac{2}{(2j-1)\pi} \theta_j \leq 2 \sin \frac{\theta_j}{2} = 2\sqrt{t_j}$, it follows that

$$|t_j - s| \leq 2\delta(2+\delta)t_j \leq \frac{1}{2}t_j,$$

which implies in particular that $\frac{1}{2}t_j \leq s \leq \frac{3}{2}t_j$. Furthermore, the same proof shows if $s \in \mathbf{c}_{[0,1]}(t_j, \pi/N)$, then $s \leq c_* t_j$. By definition, there are constants $b_1 > 0$ and $b_2 > 0$ such that $\mathbf{c}_{\mathbb{S}}(\xi, b_1\varepsilon_j) \subset \mathbb{S}_{\xi}(\varepsilon_j) \subset \mathbf{c}_{\mathbb{S}}(\xi, b_2\varepsilon_j)$. We claim that (4.22) holds for some $c_1 < \delta$ and some $c_2 > b_2$. Indeed, if $(y, \eta) \subset \mathbf{c}((t_j\xi, t_j), c_1\varepsilon)$, then $\mathbf{d}_{[0,1]}(s, t_j) \leq c_1\varepsilon \leq \delta/N$ so that $s \geq t_j/2$, and $(st)^{\frac{1}{4}}\mathbf{d}_{\mathbb{S}}(\xi, \eta) \leq cc_1\varepsilon$ by (4.3) so that $\mathbf{d}_{\mathbb{S}}(\xi, \eta) \leq 2^{\frac{1}{4}}cc_1\varepsilon/\sqrt{t_j} \leq b_1\varepsilon_j$ by choosing c_1 small. This establishes the left-hand side inclusion of (4.22). The right-hand side inclusion can be similarly established. The proof is completed. \square

The above construction establishes the existence of maximal ε -separated set on the conic surface. Since $(\Omega, \mathbf{w}_{-1,\gamma}, \mathbf{d}_{\mathbb{V}_0})$ is a localizable homogeneous space, we can then deduce the Marcinkiewicz-Zygmund inequality on such sets for all doubling weights on the conic surface. The weight \mathbf{w} on \mathbb{V}_0^{d+1} is in general a function of both x and t .

Theorem 4.18. *Let w be a doubling weight on \mathbb{V}_0^{d+1} . Let $\Xi_{\mathbb{V}_0}$ be a maximal $\frac{\delta}{n}$ -separated subset of \mathbb{V}_0^{d+1} and $0 < \delta \leq 1$.*

(i) *For all $0 < p < \infty$ and $f \in \Pi_m(\mathbb{V}_0^{d+1})$ with $n \leq m \leq cn$,*

$$\sum_{z \in \Xi_{\mathbb{V}_0}} \left(\max_{(x,t) \in c((z,r), \frac{\delta}{n})} |f(x,t)|^p \right) w(c((z,r), \frac{\delta}{n})) \leq c_{w,p} \|f\|_{p,w}^p$$

where $c_{w,p}$ depends on the doubling constant $L(w)$ and on p when p is close to 0.

(ii) *For $0 < r < 1$, there is a $\delta_r > 0$ such that for $\delta \leq \delta_r$, $r \leq p < \infty$ and $f \in \Pi_n(\mathbb{V}_0^{d+1})$,*

$$\|f\|_{p,w}^p \leq c_{w,r} \sum_{z \in \Xi} \left(\min_{(x,t) \in c((z,r), \frac{\delta}{n})} |f(x,t)|^p \right) w(c((z,r), \frac{\delta}{n}))$$

where $c_{w,r}$ depends only on $L(w)$ and on r when r is close to 0.

4.6. Cubature rules and localized tight frames. We now turn our attention to Assertion 4 in Subsection 2.5 and construct fast decaying polynomials on the conic surface.

Lemma 4.19. *Let $d \geq 2$. For each $(x,t) \in \mathbb{V}_0^{d+1}$, there is a polynomial $T_{x,t}$ of degree n that satisfies*

(1) *$T_{x,t}(x,t) = 1$, $T_{x,t}(y,s) \geq c > 0$ if $(y,s) \in c((x,t), \frac{\delta}{n})$, and for every $\kappa > 0$,*

$$0 \leq T_{x,t}(y,s) \leq c_\kappa (1 + d_{\mathbb{V}_0}((x,t), (y,s)))^{-\kappa}, \quad (y,s) \in \mathbb{V}_0^{d+1}.$$

(2) *there is a polynomial $q(t)$ of degree n such that $q(t)T_{x,t}$ is a polynomial of degree $2n$ in (x,t) variables and $1 \leq q_n(t) \leq c$.*

Proof. Let r be a positive integer such that $\kappa \leq 2r$. For positive integer n , let $m = \lfloor \frac{n}{r} \rfloor + 1$ and define

$$S_n(\cos \theta) = \left(\frac{\sin(m + \frac{1}{2})\frac{\theta}{2}}{(m + \frac{1}{2})\sin \frac{\theta}{2}} \right)^{2r}, \quad 0 \leq \theta \leq \pi.$$

By considering $m\theta \geq 1$ and $m\theta \leq 1$ separately if necessary, it follows readily that

$$(4.23) \quad S_n(1) = 1, \quad 0 \leq S_n(\cos \theta) \leq c(1 + n\theta)^{-2r}, \quad 0 \leq \theta \leq \pi.$$

Moreover, $S_n(z)$ is an even algebraic polynomial of degree at most $2n$. For a fixed $(x,t) \in \mathbb{V}_0^{d+1}$, we define

$$T_{(x,t)}(y,s) := \frac{S_n\left(\sqrt{\frac{\langle x,y \rangle + ts}{2}} + \sqrt{1-t}\sqrt{1-s}\right) + S_n\left(\sqrt{\frac{\langle x,y \rangle + ts}{2}} - \sqrt{1-t}\sqrt{1-s}\right)}{1 + S_n(2t-1)}.$$

As an even polynomial, $S_n(z)$ is a sum of even monomials. Since, by binomial formula, $(\sqrt{a} + \sqrt{b})^{2k} + (\sqrt{a} - \sqrt{b})^{2k}$ is a polynomial of degree k in a and b , it follows that $T_{(x,t)}$ is indeed a polynomial in (y,s) of degree at most n . Moreover, it satisfies $T_{(x,t)}(x,t) = 1$ since $\langle x,x \rangle = t^2$. Furthermore, using $\frac{2}{\pi}\theta \leq \sin \theta \leq \theta$ for $0 \leq \theta \leq \pi/2$, we see that S_n satisfies

$$S_n(\cos \theta) \geq \left(\frac{2}{\pi} \right)^{2r}, \quad \text{if } 0 \leq \theta \leq \frac{2\pi}{2m+1}.$$

Hence, since $0 \leq S_n(2t-1) \leq 1$, it follows that

$$T_{(x,t)}(y,s) \geq \frac{S_n\left(\sqrt{\frac{\langle x,y \rangle + ts}{2}} + \sqrt{1-t}\sqrt{1-s}\right)}{1 + S_n(2t-1)} \geq \frac{1}{2}S_n(\cos d_{\mathbb{V}_0}((x,t), (y,s))) \geq \frac{1}{2}\left(\frac{2}{\pi}\right)^{2r}$$

for $(y,s) \in \mathbf{c}((x,t), \frac{2\pi}{2m+1})$. Finally, we can use the fast decay (4.23) of S_n with $\theta \sim \sin \frac{\theta}{2} = \sqrt{\frac{1}{2}(1 - \cos \theta)}$ to derive an upper bound of $T_{(x,t)}$. Indeed, since evidently $1 - \sqrt{\frac{\langle x,y \rangle + ts}{2}} + \sqrt{1-t}\sqrt{1-s} \geq 1 - \sqrt{\frac{\langle x,y \rangle + ts}{2}} - \sqrt{1-t}\sqrt{1-s}$, we obtain then

$$\begin{aligned} 0 \leq T_{(x,t)}(y,s) &\leq c \frac{1}{\left(1 + n\sqrt{1 - \sqrt{\frac{\langle x,y \rangle + ts}{2}} - \sqrt{1-t}\sqrt{1-s}}\right)^{2r}} \\ &= c \frac{1}{\left(1 + n\sqrt{1 - \cos d_{\mathbb{V}_0}((x,t), (y,s))}\right)^{2r}}. \end{aligned}$$

Hence, by the definition of $d_{\mathbb{V}_0}$ and $1 - \cos \theta \sim \theta^2$, we obtain

$$(4.24) \quad 0 \leq T_{(x,t)}(y,s) \leq c \frac{1}{(1 + nd_{\mathbb{V}_0}((x,t), (y,s)))^{2r}}.$$

This completes the proof of (1). Finally, $q_n(t) = 1 + S_n(2t-1)$ is a polynomial of degree n and it satisfies $1 \leq q_n(t) \leq c$ for all $t \in [0, 1]$. Moreover, $q_n(t)T_{x,t}$ is a polynomial of degree $2n$ in (x, t) variable. This completes the proof. \square

From Propositions 2.17 and 2.18, we have established the following result.

Corollary 4.20. *Let w be a doubling weight function on \mathbb{V}_0^{d+1} . Then*

$$\lambda_n(w; (x, t)) \leq c w\left(\mathbf{c}\left((x, t), \frac{1}{n}\right)\right).$$

Moreover, for $\gamma \geq -\frac{1}{2}$,

$$\lambda_n(w_{-1,\gamma}; (x, t)) \geq c w_{-1,\gamma}\left(\mathbf{c}\left((x, t), \frac{1}{n}\right)\right) = cn^{-d} w_{-1,\gamma}(n; t).$$

This implies, in particular, that the cubature rule in Theorem 2.20 holds for all doubling weights on the conic surface.

Theorem 4.21. *Let w be a doubling weight on \mathbb{V}_0^{d+1} . Let Ξ be a maximum $\frac{\delta}{n}$ -separated subset of \mathbb{V}_0^{d+1} . There is a $\delta_0 > 0$ such that for $0 < \delta < \delta_0$ there exist positive numbers $\lambda_{z,r}$, $(z, r) \in \Xi$, so that*

$$(4.25) \quad \int_{\mathbb{V}_0^{d+1}} f(x, t) w(x, t) d\sigma(x, t) = \sum_{(z,r) \in \Xi} \lambda_{z,r} f(z, r), \quad \forall f \in \Pi_n(\mathbb{V}_0^{d+1}).$$

Moreover, $\lambda_{z,r} \sim w(\mathbf{c}((z, r), \frac{\delta}{n}))$ for all $(z, r) \in \Xi$.

We can now construct localized frame on the conic surface. Let us recapitulate necessary definitions in the setting of \mathbb{V}_0^{d+1} . For $j = 0, 1, \dots$, let Ξ_j be a maximal $\frac{\delta}{2^j}$ -separated subset in \mathbb{V}_0^{d+1} , so that

$$\int_{\mathbb{V}_0^{d+1}} f(x, t) w(x, t) d\sigma(x, t) = \sum_{(z,r) \in \Xi_j} \lambda_{(x,r),j} f(z, r), \quad f \in \Pi_{2^j}(\mathbb{V}_0^{d+1}).$$

Let \hat{a} be an admissible cut-off function satisfying (2.43). Let $\mathbf{L}_{2^j}(\mathbf{w}) * f$ denote the near best approximation operator defined by

$$\mathbf{L}_{2^j}(\mathbf{w}) * f(x) := \int_{\mathbb{V}_0^{d+1}} f(y, s) \mathbf{L}_{2^j}(\mathbf{w}; (x, t), (y, s)) \mathbf{w}(y, s) d\sigma(y, s).$$

For $j = 1, \dots$, we introduce the notation $F_j(\mathbf{w}) = \mathbf{L}_{2^j}(\mathbf{w})$ for both the kernel and the operator. More precisely, $F_0(\mathbf{w}; \cdot, \cdot) = 1$ and for $j \geq 1$,

$$F_j(\mathbf{w}; \cdot, \cdot) = \mathbf{L}_{2^{j-1}}(\mathbf{w}; \cdot, \cdot) \quad \text{and} \quad F_j(\mathbf{w}) * f = F_j(\mathbf{w}) * f$$

and define the frame elements $\psi_{(z,r),j}$ for $(z, r) \in \Xi_j$ by

$$\psi_{(z,r),j}(x, t) := \sqrt{\lambda_{(z,r),j} F_j(\mathbf{w}; (x, t), (z, r))}, \quad (x, t) \in \mathbb{V}_0^{d+1},$$

which are well defined for all doubling weight by Theorem 4.21. By Theorem 2.21, the system $\Phi = \{\psi_{(z,r),j} : (z, r) \in \Xi_j, j = 1, 2, 3, \dots\}$ is a tight frame of $L^2(\mathbb{V}_0^{d+1}, \mathbf{w})$.

Theorem 4.22. *Let \mathbf{w} be a doubling weight on \mathbb{V}_0^{d+1} . If $f \in L^2(\mathbb{V}_0^{d+1}, \mathbf{w})$, then*

$$f = \sum_{j=0}^{\infty} \sum_{(z,r) \in \Xi_j} \langle f, \psi_{(z,r),j} \rangle_{\mathbf{w}} \psi_{(z,r),j} \quad \text{in } L^2(\mathbb{V}_0^{d+1}, \mathbf{w})$$

and

$$\|f\|_{2,\mathbf{w}} = \left(\sum_{j=0}^{\infty} \sum_{(z,r) \in \Xi_j} |\langle f, \psi_{(z,r),j} \rangle_{\mathbf{w}}|^2 \right)^{1/2}.$$

For the weight function $\mathbf{w}_{-1,\gamma}$, the frame elements are highly localized.

Theorem 4.23. *For $\gamma \geq -\frac{1}{2}$, the frame for $\mathbf{w}_{-1,\gamma}$ is highly localized in the sense that, for every $\kappa > 0$, there exists a constant $c_\kappa > 0$ such that*

$$(4.26) \quad |\psi_{(z,r),j}(x, t)| \leq c_\sigma \frac{2^{jd/2}}{\sqrt{\mathbf{w}_{\gamma,d}(2^j; t) (1 + 2^j d_{\mathbb{V}_0}((x, t), (z, r)))^\kappa}}, \quad (x, t) \in \mathbb{V}_0^{d+1}.$$

Proof. The inequality (4.26) follows from the highly localized estimate of the kernel in Theorem 4.10 and $\lambda_{(z,r),j} \sim 2^{-jd} \mathbf{w}_\gamma(2^j; r)$ which holds for $\mathbf{w}_{-1,\gamma}$ according to Corollary 4.20. \square

It is worth pointing out that the localization of the frame elements is established for $\mathbf{w}_{-1,\gamma}(t) = t^{-1}(1-t)^\gamma$ but not for the Lebesgue measure on \mathbb{V}_0^{d+1} .

4.7. Characterization of best approximation. For $f \in L_p(\mathbb{V}_0^{d+1}, \mathbf{w})$, $1 \leq p < \infty$, and $f \in C(\mathbb{V}_0^{d+1})$ if $p = \infty$, the error of best approximation by polynomials of degree at most n is defined by

$$\mathbf{E}_n(f)_{p,\mathbf{w}} = \inf_{g \in \Pi_n(\mathbb{V}_0^{d+1})} \|f - g\|_{p,\mathbf{w}}, \quad 1 \leq p \leq \infty.$$

Following the study in Section 3, we can give a characterization for this quantity by the modulus of smoothness defined via the operator $\mathbf{S}_{\theta,\mathbf{w}}$ and the K -functional defined via the differential operator $\Delta_{0,\gamma}$ for $\mathbf{w}_{-1,\gamma}$.

More specifically, for $f \in L^p(\mathbb{V}_0^{d+1}, \mathbf{w}_{-1,\gamma})$ and $r > 0$, the modulus of smoothness is defined by

$$\omega_r(f; \rho)_{p,\mathbf{w}_{-1,\gamma}} = \sup_{0 \leq \theta \leq \rho} \left\| (I - \mathbf{S}_{\theta,\mathbf{w}_{-1,\gamma}})^{r/2} f \right\|_{p,\mathbf{w}_{-1,\gamma}}, \quad 1 \leq p \leq \infty,$$

where the operator $S_{\theta, \mathbf{w}_{-1, \gamma}}$ is defined by, for $n = 0, 1, 2, \dots$ and $\lambda = \gamma + d - 1$,

$$\text{proj}_n(\mathbf{w}_{-1, \gamma}; S_{\theta, \mathbf{w}_{-1, \gamma}} f) = R_n^{(\lambda - \frac{1}{2}, -\frac{1}{2})}(\cos \theta) \text{proj}_n(\mathbf{w}_{-1, \gamma}; f).$$

Moreover, in terms of the fractional differential operator $(-\Delta_{0, \gamma})^{\frac{r}{2}}$, the K -functional is defined for a weight \mathbf{w} by

$$K_r(f, t)_{p, \mathbf{w}} := \inf_{g \in W_p^r(\mathbb{V}_0^{d+1}, \mathbf{w})} \left\{ \|f - g\|_{p, \mathbf{w}} + t^r \|(-\Delta_{0, \gamma})^{\frac{r}{2}} f\|_{p, \mathbf{w}} \right\}.$$

Both these quantities are well defined as shown in Section 3. Since Assertions 1–4 hold for $\mathbf{w}_{-1, \gamma}$, we only have to verify that the Assertion 5 in Subsection 3.5 holds for the differential operator $\mathfrak{D}_{\varpi} = \Delta_{0, \gamma}$. By Theorem 4.8, the kernel $L_n^{(r)}(\varpi)$ in Assertion 5 becomes

$$\mathbb{L}_n^{(r)}(\mathbf{w}_{-1, \gamma}; (x, t), (y, s)) = \sum_{k=0}^{\infty} \hat{a}\left(\frac{k}{n}\right) (k(k + \gamma + d - 1))^{\frac{r}{2}} P_k(\mathbf{w}_{-1, \gamma}; (x, t), (y, s)).$$

Lemma 4.24. *Let $\gamma \geq -\frac{1}{2}$ and $\kappa > 0$. Then, for $r > 0$ and $(x, t), (y, s) \in \mathbb{V}_0^{d+1}$,*

$$|\mathbb{L}_n^{(r)}(\mathbf{w}_{-1, \gamma}; (x, t), (y, s))| \leq c_{\kappa} \frac{n^{d+r}}{\sqrt{\mathbf{w}_{\gamma, d}(n; t)} \sqrt{\mathbf{w}_{\gamma, d}(n; s)}} (1 + n d_{\mathbb{V}_0}((x, t), (y, s)))^{-\kappa}.$$

Proof. By (4.9), the kernel can be written as

$$\begin{aligned} \mathbb{L}_n^{(r)}(\mathbf{w}_{-1, \gamma}; (x, t), (y, s)) &= b_{\gamma, d} \int_{[-1, 1]^2} L_{n, r}(2\zeta(x, t, y, s; v)^2 - 1) \\ &\quad \times (1 - v_1^2)^{\frac{d-2}{2}-1} (1 - v_2^2)^{\gamma - \frac{1}{2}} dv, \end{aligned}$$

in which $L_{n, r}$ is defined by, with $\lambda = \gamma + d - 1$,

$$L_{n, r}(t) = \sum_{k=0}^{\infty} \hat{a}\left(\frac{k}{n}\right) (k(k + \gamma + d - 1))^{\frac{r}{2}} \frac{P_n^{(\lambda - \frac{1}{2}, -\frac{1}{2})}(1) P_n^{(\lambda - \frac{1}{2}, -\frac{1}{2})}(t)}{h^{(\lambda - \frac{1}{2}, -\frac{1}{2})}}.$$

Applying (2.18) with $\eta(t) = \hat{a}(t) (t(t + n^{-1}(\gamma + d - 1)))^{\frac{r}{2}}$ and $m = 0$, it follows that

$$|L_{n, r}(t)| \leq c n^r \frac{n^{2\lambda+1}}{(1 + n\sqrt{1-t})^{\ell}}.$$

Consequently, it follows readily that

$$\begin{aligned} |\mathbb{L}_n^{(r)}(\mathbf{w}_{-1, \gamma}; (x, t), (y, s))| &\leq c n^{2\lambda+r+1} \int_{[-1, 1]^2} \frac{1}{\left(1 + n\sqrt{1 - \zeta(x, t, y, s; v)^2}\right)^{\kappa+2\gamma+d+1}} \\ &\quad \times (1 - v_1^2)^{\frac{d-2}{2}-1} (1 - v_2^2)^{\gamma - \frac{1}{2}} dv. \end{aligned}$$

Apart from n^r , this estimate is the same as that of $L_n(\mathbf{w}_{-1, \gamma})$ appeared in the proof of Theorem 4.10, so that the desired upper bound follows from the estimate there. \square

With all assertions hold for the $\mathbf{w}_{-1, \gamma}$, the characterization of the best approximation by polynomials in Subsection 3.3 holds on the conic surface, which we state below.

Theorem 4.25. *Let $f \in L^p(\mathbb{V}_0^{d+1}, \mathbf{w})$ if $1 \leq p < \infty$ and $f \in C(\mathbb{V}_0^{d+1})$ if $p = \infty$. Let $r > 0$ and $n = 1, 2, \dots$. For $\mathbf{w} = \mathbf{w}_{-1, \gamma}$ with $\gamma \geq -\frac{1}{2}$, there holds*

(i) *direct estimate*

$$\mathbb{E}_n(f)_{p, \mathbf{w}_{-1, \gamma}} \leq c K_r(f; n^{-1})_{p, \mathbf{w}_{-1, \gamma}};$$

(ii) *inverse estimate*

$$\mathbf{K}_r(f; n^{-1})_{p, \mathbf{w}_{-1, \gamma}} \leq cn^{-r} \sum_{k=0}^n (k+1)^{r-1} \mathbf{E}_k(f)_{p, \mathbf{w}_{-1, \gamma}}.$$

Furthermore, the characterization can be given via the modulus of smoothness, since it is equivalent to the K -functional.

Theorem 4.26. *Let $\gamma \geq -\frac{1}{2}$ and $f \in L_p^r(\mathbb{V}_0^{d+1}, \mathbf{w}_{-1, \gamma})$, $1 \leq p \leq \infty$. Then for $0 < \theta \leq \pi/2$ and $r > 0$*

$$c_1 \mathbf{K}_r(f; \theta)_{p, \mathbf{w}_{-1, \gamma}} \leq \omega_r(f; \theta)_{p, \mathbf{w}_{-1, \gamma}} \mathbf{K}_r(f; \theta)_{p, \mathbf{w}_{-1, \gamma}}.$$

5. HOMOGENEOUS SPACE ON SOLID CONES

In this section we work in the setting of homogeneous space on the solid cone

$$\mathbb{V}^{d+1} = \{(x, t) : \|x\| \leq t, x \in \mathbb{B}^d, 0 \leq t \leq 1\},$$

which is bounded by \mathbb{V}_0^{d+1} and the hyperplane $t = 1$ in \mathbb{R}^{d+1} . As in the previous section, we shall verify that the framework we developed for approximation and tight frame on homogeneous space is applicable on this domain with respect to the Jacobi weight function $W_{\gamma, \mu}(x, t) = (1-t)^\gamma (t^2 - \|x\|^2)^{\mu - \frac{1}{2}}$.

The structure of the section is parallel to that of the conic surface. We again need to define and understand an intrinsic distance and doubling weights on the cone and do so in the first two subsections. The orthogonal structure with respect to the Jacobi weight is reviewed in the third subsection. The highly localized kernels are established in the fourth subsection, where the Assertions 1 and 3 are verified for $W_{\gamma, \mu}$ for all $\mu \geq 0$ but the Assertion 2 is established only for $W_{\gamma, 0}$. We also provide construction of ve -separated set on the cone and state the Marcinkiewicz-Zygmund inequality in the fifth subsection. Assertion 4 is verified in the sixth subsection, which allows us to state the positive cubature rules and the tight frame on the cone. While the tight localized frame holds for the Jacobi weight with $\mu = 0$, the characterization of the best approximation by polynomials works out for all $\mu \geq 0$ in the seventh subsection.

5.1. Distance on solid cone. The distant function $\mathbf{d}_{\mathbb{B}}(\cdot, \cdot)$ on the unit ball \mathbb{B}^d can be deduced from regarding \mathbb{B}^d as the projection of the upper hemisphere of \mathbb{S}^{d+1} , or the distant function $\mathbf{d}_{\mathbb{S}}(\cdot, \cdot)$ applied on the points $X = (x, \sqrt{1 - \|x\|^2})$ and $Y = (y, \sqrt{1 - \|y\|^2})$ with $x \in \mathbb{B}^d$. The same holds for the solid cone \mathbb{V}^{d+1} .

Definition 5.1. *For (x, t) and (y, s) on \mathbb{V}^{d+1} , let $X = (x, \sqrt{t^2 - \|x\|^2})$ and $Y = (y, \sqrt{s^2 - \|y\|^2})$. Define*

$$(5.1) \quad \mathbf{d}_{\mathbb{V}}((x, t), (y, s)) := \arccos \left(\sqrt{\frac{\langle X, Y \rangle + ts}{2}} + \sqrt{1-t} \sqrt{1-s} \right).$$

Proposition 5.2. *The function $\mathbf{d}_{\mathbb{V}}(\cdot, \cdot)$ defines a distance function on the solid cone \mathbb{V}^{d+1} .*

Proof. Since $\|X\| = t$ and $\|Y\| = s$, both (X, t) and (Y, s) are elements of \mathbb{V}_0^{d+2} . Furthermore,

$$(5.2) \quad \mathbf{d}_{\mathbb{V}^{d+1}}((x, t), (y, s)) = \mathbf{d}_{\mathbb{V}_0^{d+2}}((X, t), (Y, s)),$$

from which it is easy to see that $\mathbf{d}_{\mathbb{V}}(\cdot, \cdot)$ is a distance function on \mathbb{V}^{d+1} . \square

The distance function $d_V(\cdot, \cdot)$ is closely related to the distance function $d_{[0,1]}(\cdot, \cdot)$ of the interval $[0, 1]$ and the distance $d_{\mathbb{B}}(\cdot, \cdot)$ of the unit ball \mathbb{B}^d .

Proposition 5.3. *For $d \geq 2$ and $(x, t), (y, s) \in \mathbb{V}^{d+1}$, write $x = tx'$ and $y = sy'$ with $x', y' \in \mathbb{B}^d$. Then*

$$1 - \cos d_V((x, t), (y, s)) = 1 - \cos d_{[0,1]}(t, s) + \sqrt{t}\sqrt{s} \left[1 - \cos \left(\frac{1}{2} d_{\mathbb{B}}(x', y') \right) \right].$$

In particular,

$$c_1 d_V((x, t), (y, s)) \leq d_{[0,1]}(t, s) + (ts)^{\frac{1}{4}} d_{\mathbb{B}}(x', y') \leq c_2 d_V((x, t), (y, s)).$$

Proof. With $x = tx'$, we see that $X = t(x', \sqrt{1 - \|x'\|^2})$ and $Y = s(y', \sqrt{1 - \|y'\|^2})$. Hence, it follows as in the case of \mathbb{V}_0^{d+1} that

$$(5.3) \quad d_V((x, t), (y, s)) = \arccos \left[\sqrt{t}\sqrt{s} \cos \left(\frac{1}{2} d_{\mathbb{B}}(x', y') \right) + \sqrt{1-t}\sqrt{1-s} \right].$$

The rest of the proof follows from that of Proposition 4.3 almost verbatim. \square

Like the distance function on the surface, we see that the distance on the line segment from the apex to $(x', 1)$ on the top boundary of the cone \mathbb{V}^{d+1} is exactly the distance function on $[0, 1]$; that is, for $x' \in \mathbb{B}^d$,

$$d_V((x', t), (x', s)) = d_{[0,1]}(t, s), \quad 0 \leq t, s \leq 1.$$

The following lemma is an analog of Lemma 4.4 and will be needed in the next subsection.

Lemma 5.4. *For $(x, t), (y, s) \in \mathbb{V}^{d+1}$,*

$$|\sqrt{t} - \sqrt{s}| \leq d_V((x, t), (y, s)) \quad \text{and} \quad |\sqrt{1-t} - \sqrt{1-s}| \leq d_V((x, t), (y, s)),$$

and

$$|\sqrt{t^2 - \|x\|^2} - \sqrt{s^2 - \|y\|^2}| \leq c \max\{\sqrt{s}, \sqrt{t}\} d_V((x, t), (y, s)).$$

Proof. The first two inequalities are immediate consequence of (5.2) and Lemma 4.4. Let $x = tx'$ and $y = sy'$ with $x', y' \in \mathbb{B}^d$. We know [12, (A.1.4)] that

$$(5.4) \quad \left| \sqrt{1 - \|x'\|^2} - \sqrt{1 - \|y'\|^2} \right| \leq \sqrt{2} d_{\mathbb{B}}(x', y').$$

Without loss of generality, assuming $t \geq s$. Then $s \leq \sqrt{s}\sqrt{t}$, so that

$$\begin{aligned} \left| \sqrt{t^2 - \|x\|^2} - \sqrt{s^2 - \|y\|^2} \right| &= \left| t\sqrt{1 - \|x'\|^2} - s\sqrt{1 - \|y'\|^2} \right| \\ &\leq |t - s| \sqrt{1 - \|x'\|^2} + s \left| \sqrt{1 - \|x'\|^2} - \sqrt{1 - \|y'\|^2} \right| \\ &\leq \sqrt{2} \max\{\sqrt{s}, \sqrt{t}\} \left(|\sqrt{t} - \sqrt{s}| + (ts)^{\frac{1}{4}} d_{\mathbb{B}}(x', y') \right), \end{aligned}$$

which is bounded by $c \max\{\sqrt{s}, \sqrt{t}\} d_V((x, t), (y, s))$ by the first inequality in the statement and the inequality in Proposition 5.3. \square

5.2. A family of doubling weights. For $d \geq 1$, $\mu \geq 0$, $\beta + 2\mu > -d$ and $\gamma > -1$, we consider the Jacobi weight function defined on the solid cone by

$$W_{\beta,\gamma,\mu}(x,t) := (t^2 - \|x\|^2)^{\mu-\frac{1}{2}} t^\beta (1-t)^\gamma, \quad (x,t) \in \mathbb{V}^{d+1}.$$

Let $\mathbf{b}_{\beta,\gamma,\mu}$ be the normalization constant so that $\mathbf{b}_{\beta,\gamma,\mu} W_{\beta,\gamma,\mu}$ has unit integral on \mathbb{V}^{d+1} . Setting $x = tx'$ with $x' \in \mathbb{B}^d$, then

$$\mathbf{b}_{\beta,\gamma,\mu} = \left(\int_0^1 t^{d+\beta+2\mu-1} (1-t)^\gamma dt \int_{\mathbb{B}^d} (1-\|x'\|^2)^{\mu-\frac{1}{2}} dx' \right)^{-1} = c_{\beta+2\mu+d-1,\gamma} b_\mu^\mathbb{B},$$

where $c_{\alpha,\beta}$ is defined in (2.14) and $b_\mu^\mathbb{B}$ is the normalization constant of the classical weight function on the unit ball. The case $\beta = 0$ is of particular interest and will be denoted by $W_{\gamma,\mu}$; that is,

$$W_{\gamma,\mu}(x,t) := W_{0,\gamma,\mu}(x,t) = (t^2 - \|x\|^2)^{\mu-\frac{1}{2}} (1-t)^\gamma, \quad (x,t) \in \mathbb{V}^{d+1}.$$

We show that this is a doubling weight with respect to the distance $\mathbf{d}_\mathbb{V}$. For $r > 0$ and $(x,t) \in \mathbb{V}^{d+1}$, we denote the ball centered at (x,t) with radius r by

$$\mathbf{c}((x,t), r) := \{(y,s) \in \mathbb{V}^{d+1} : \mathbf{d}_\mathbb{V}((x,t), (y,s)) \leq r\}.$$

Proposition 5.5. *Let $r > 0$ and $(x,t) \in \mathbb{V}^{d+1}$. Then for $\beta > -d$, $\gamma > -1$ and $\mu \geq 0$,*

$$(5.5) \quad W_{\beta,\gamma,\mu}(\mathbf{c}((x,t), r)) := \int_{\mathbf{c}((x,t), r)} W_{\beta,\gamma,\mu}(y,s) dy ds \\ \sim r^{d+1} (t+r^2)^{\beta+\frac{d-1}{2}} (1-t+r^2)^{\gamma+\frac{1}{2}} (t^2 - \|x\|^2 + r^2)^\mu.$$

In particular, $W_{\beta,\gamma,\mu}$ is a doubling weight and the doubling index $\alpha(W_{\beta,\gamma,\mu})$ is give by $\alpha(W_{\beta,\gamma,\mu}) = 2\mu + d + 1 + 2\max\{0, \beta + \frac{d-1}{2}\} + 2\max\{0, \gamma + \frac{1}{2}\}$.

Proof. Let $\tau_r(t,s)$ and $\theta_r(t,s) = \arccos \tau_r(t,s)$ be as in the proof of Proposition 4.6. From $\mathbf{d}_\mathbb{V}((x,t), (y,s)) \leq r$, we obtain $\mathbf{d}_{[0,1]}(t,s) \leq r$ and, by (5.1),

$$\mathbf{d}_\mathbb{B}(x', y') \leq \arccos(2[\tau_r(t,s)]^2 - 1) = \frac{1}{2} \arccos \tau_r(t,s) = \frac{1}{2} \theta_r(t,s),$$

where $\mathbf{d}_\mathbb{B}(\cdot, \cdot)$ is the distance on the unit ball \mathbb{B}^d . Hence, it follows that

$$W_{\beta,\gamma,\mu}(\mathbf{c}((x,t), r)) = \int_{\mathbf{d}_{[0,1]}(t,s) \leq r} s^d \int_{\mathbf{d}_\mathbb{B}(x', y') \leq \frac{1}{2} \theta_r(t,s)} W_{\beta,\gamma,\mu}(sy', s) dy' ds \\ \sim \int_{\mathbf{d}_{[0,1]}(t,s) \leq r} s^{d+\beta+2\mu-1} (1-s)^\gamma \int_{\mathbf{d}_\mathbb{B}(x', y') \leq \frac{1}{2} \theta_r(t,s)} (1-\|y'\|^2)^{\mu-\frac{1}{2}} dy' ds$$

For $\mu \geq 0$ and $0 < \rho \leq 1$, it is known [38, Lemma 5.3] or [12, p. 107] that

$$(5.6) \quad \int_{\mathbf{d}_\mathbb{B}(x', y') \leq \rho} (1-\|y'\|^2)^{\mu-\frac{1}{2}} dy' \sim (1-\|x'\|^2 + \rho^2)^\mu \rho^d,$$

which can also be established by following the approach in Proposition 4.6. Consequently, using $\theta_r(t,s) \sim \sqrt{1-\tau_r(t,s)}$, we conclude that

$$W_{\beta,\gamma,\mu}(\mathbf{c}((x,t), r)) \sim \int_{\mathbf{d}_{[0,1]}(t,s) \leq r} s^{d+\beta+2\mu-1} (1-s)^\gamma \\ \times (1-\|x'\|^2 + 1-\tau_r(t,s))^\mu (1-\tau_r(t,s))^{\frac{d}{2}} ds.$$

If $t \geq 3r^2$, then $s \sim t + r^2$ and, by (4.7), it follows that

$$s^\mu (1-\|x'\|^2 + 1-\tau_r(t,s))^\mu \sim (t^2 - \|x\|^2 + r^2)^\mu.$$

With this term removed, the integral of the remaining integrand can be estimated by following the estimates of Case 1 and Case 3 of the proof of Proposition 4.6. If $t \leq 3r^2$, then $t^2 - \|x\|^2 + r^2 \sim r^2$. We use $1 - \|x'\|^2 + 1 - \tau_r(t, s) \leq 2$ for the upper bound and $1 - \tau_r(t, s) \geq 2/\pi^2$ on the subset $\mathbf{d}_{[0,1]}(t, s) \leq r/2$, proved in the Case 2 of the proof of Proposition 4.6, to remove the term $(1 - \|x'\|^2 + 1 - \tau_r(t, s))^\mu$ from the integral. The rest of the proof then follows from that of the Case 2 of the proof of Proposition 4.6. This completes the proof. \square

Corollary 5.6. *For $d \geq 2$, $\beta > -d$, $\gamma > -1$ and $\mu \geq 0$, $(\mathbb{V}^{d+1}, W_{\beta, \gamma, \mu}, \mathbf{d}_\mathbb{V})$ is a homogeneous space.*

5.3. Orthogonal polynomials on the solid cone. These polynomials are studied in [53] and they are orthogonal with respect to the inner product

$$\langle f, g \rangle_{\beta, \gamma, \mu} = \mathbf{b}_{\beta, \gamma, \mu} \int_{\mathbb{V}^{d+1}} f(x, t) g(x, t) W_{\beta, \gamma, \mu}(x, t) dx dt.$$

Let $\mathcal{V}_n(\mathbb{V}^{d+1}, W_{\beta, \gamma, \mu})$ be the space of orthogonal polynomials of degree n , which has dimension $\binom{n+d+1}{n}$. An orthogonal basis of $\mathcal{V}_n(\mathbb{V}^{d+1}, W_{\beta, \gamma, \mu})$ can be given in terms of the Jacobi polynomials and the orthogonal polynomials on the unit ball. For $m = 0, 1, 2, \dots$, let $\{P_{\mathbf{k}}^m(W_\mu) : |\mathbf{k}| = m, \mathbf{k} \in \mathbb{N}_0^d\}$ be an orthonormal basis of $\mathcal{V}_n(\mathbb{B}^d, W_\mu)$ on the unit ball. Let $\alpha := \mu + \frac{\beta+d-1}{2}$ and

$$(5.7) \quad \mathbf{J}_{m, \mathbf{k}}^n(x, t) := P_{n-m}^{(2\alpha+2m, \gamma)}(1-2t)t^m P_{\mathbf{k}}^m\left(W_\mu; \frac{x}{t}\right),$$

Then $\{\mathbf{J}_{m, \mathbf{k}}^n(x, t) : |\mathbf{k}| = m, 0 \leq m \leq n\}$ is an orthogonal basis of $\mathcal{V}_n(\mathbb{V}^{d+1}, W_{\beta, \gamma, \mu})$ and the norm of $\mathbf{J}_{m, \mathbf{k}}^n$ is given by

$$(5.8) \quad \mathbf{H}_{m, n}^{(\alpha, \gamma)} := \langle \mathbf{J}_{m, \mathbf{k}}^n, \mathbf{J}_{m, \mathbf{k}}^n \rangle_{W_{\mu, \beta, \gamma}} = \frac{c_{2\alpha, \gamma}}{c_{2\alpha+2m, \gamma}} h_{n-m}^{(2\alpha+2m, \gamma)},$$

where $c_{\alpha, \beta}$ is as in (2.14) and $h_m^{(\alpha, \gamma)}$ is the norm square of the Jacobi polynomial. We call $\mathbf{J}_{m, \mathbf{k}}^n$ the Jacobi polynomials on the cone.

The reproducing kernel of the space $\mathcal{V}_n(\mathbb{V}^{d+1}, W_{\beta, \gamma, \mu})$, denoted by $\mathbf{P}_n(W_{\beta, \gamma, \mu}; \cdot, \cdot)$, can be written in terms of the above basis,

$$\mathbf{P}_n(W_{\beta, \gamma, \mu}; (x, t), (y, s)) = \sum_{m=0}^n \sum_{|\mathbf{k}|=n} \frac{\mathbf{J}_{m, \mathbf{k}}^n(x, t) \mathbf{J}_{m, \mathbf{k}}^n(y, s)}{\mathbf{H}_{m, n}^{\beta, \gamma}}.$$

It is the kernel of the projection $\text{proj}_n(W_{\beta, \gamma, \mu}) : L^2(\mathbb{V}^{d+1}, W_{\beta, \gamma, \mu}) \rightarrow \mathcal{V}_n(\mathbb{V}^{d+1}, W_{\beta, \gamma, \mu})$,

$$\text{proj}_n(W_{\beta, \gamma, \mu}; f) = \int_{\mathbb{V}^{d+1}} f(y, s) \mathbf{P}_n(W_{\beta, \gamma, \mu}; \cdot, (y, s)) W_{\beta, \gamma, \mu}(s) dy ds.$$

In contrast to the conic surface, it is the case $\beta = 0$ that turns out to be the most interesting one. For $\beta = 0$, there is a second order differential operator that has orthogonal polynomials as eigenfunctions. Recall that $W_{0, \gamma, \mu} = W_{\gamma, \mu}$.

Theorem 5.7. *Let $\mu > -\frac{1}{2}$, $\gamma > -1$ and $n \in \mathbb{N}_0$. Define the second order differential operator $\mathfrak{D}_{\mu, \gamma}$ by*

$$\begin{aligned} \mathfrak{D}_{\gamma, \mu} := & t(1-t)\partial_t^2 + 2(1-t)\langle x, \nabla_x \rangle \partial_t + \sum_{i=1}^d (t - x_i^2) \partial_{x_i}^2 - 2 \sum_{i < j} x_i x_j \partial_{x_i} \partial_{x_j} \\ & + (2\mu + d) \partial_t - (2\mu + \gamma + d + 1) (\langle x, \nabla_x \rangle + t \partial_t), \end{aligned}$$

where ∇_x and Δ_x denote the gradient and the Laplace operator in x -variable. Then the polynomials in $\mathcal{V}_n(\mathbb{V}^{d+1}, W_{\gamma,\mu})$ are eigenfunctions of $\mathfrak{D}_{\gamma,\mu}$,

$$(5.9) \quad \mathfrak{D}_{\gamma,\mu} u = -n(n + 2\mu + \gamma + d)u, \quad \forall u \in \mathcal{V}_n(\mathbb{V}^{d+1}, W_{\gamma,\mu}).$$

The reproducing kernel enjoys an addition formula that is a mixture of the addition formula for orthogonal polynomials with respect to the classical weight function W_μ on the unit ball and the Jacobi polynomials. The addition formula is complicated and has the most elegant form when $\beta = 0$.

Theorem 5.8. *Let $d \geq 2$. For $\mu \geq 0$ and $\gamma \geq -\frac{1}{2}$, let $\alpha = \mu + \frac{d-1}{2}$; then*

$$(5.10) \quad \mathbf{P}_n(W_{\gamma,\mu}; (x, t), (y, s)) = c_{\mu,\gamma,d} \int_{[-1,1]^3} Z_{2n}^{2\alpha+\gamma+1}(\xi(x, t, y, s; u, v)) \\ \times (1 - u^2)^{\mu-1} (1 - v_1^2)^{\alpha-1} (1 - v_2^2)^{\gamma-\frac{1}{2}} du dv,$$

where $c_{\mu,\gamma,d}$ is a constant, so that $\mathbf{P}_0 = 1$ and $\xi(x, t, y, s; u, v) \in [-1, 1]$ is defined by

$$(5.11) \quad \xi(x, t, y, s; u, v) = v_1 \sqrt{\frac{1}{2} \left(ts + \langle x, y \rangle + \sqrt{t^2 - \|x\|^2} \sqrt{s^2 - \|y\|^2} u \right)} \\ + v_2 \sqrt{1-t} \sqrt{1-s}.$$

When $\mu = 0$ or $\gamma = -\frac{1}{2}$, the identity (5.10) holds under the limit (2.28). In particular,

$$(5.12) \quad \mathbf{P}_n(W_{\gamma,0}; (x, t), (y, s)) = c_{0,\gamma,d} \\ \times \int_{[-1,1]^2} \left[Z_{2n}^{\gamma+d}(\xi(x, t, y, s; 1, v)) + Z_{2n}^{\gamma+d}(\xi(x, t, y, s; -1, v)) \right] \\ \times (1 - v_1^2)^{\frac{d-1}{2}-1} (1 - v_2^2)^{\gamma-\frac{1}{2}} dv.$$

As a consequence, the orthogonal structure for the weight function $W_{\gamma,\mu}$ on the cone \mathbb{V}^{d+1} satisfies both characteristic properties specified in Definition 3.1 and Definition 3.2. In the next subsection, we use the addition formula to establish Assertions 1–3 for highly localized kernels.

5.4. Highly localized kernels. Let \hat{a} be an admissible cut-off function. For $(x, t), (y, s) \in \mathbb{V}^{d+1}$, the localized kernel $\mathbf{L}_n(W_{\gamma,\mu}; \cdot, \cdot)$ is defined by

$$\mathbf{L}_n(W_{\gamma,\mu}; (x, t), (y, s)) = \sum_{j=0}^{\infty} \hat{a}\left(\frac{j}{n}\right) \mathbf{P}_j(W_{\gamma,\mu}; (x, t), (y, s)).$$

For $\gamma > -1$ and $\mu \geq 0$, we need the function $n^{d+1} W_{\gamma,\mu}(\mathbf{c}((x, t), n^{-2}))$, which we denote by $W_{\gamma,\mu}(n; x, t)$; that is,

$$W_{\gamma,\mu}(n; x, t) := (t + n^{-2})^{\frac{d-1}{2}} (1 - t + n^{-2})^{\gamma+\frac{1}{2}} (t^2 - \|x\|^2 + n^{-2})^\mu.$$

Theorem 5.9. *Let $d \geq 2$, $\gamma \geq -\frac{1}{2}$ and $\mu \geq 0$. Then for any $k > 0$ and $(x, t), (y, s) \in \mathbb{V}^{d+1}$,*

$$|\mathbf{L}_n(W_{\gamma,\mu}; (x, t), (y, s))| \leq \frac{c_\kappa n^{d+1}}{\sqrt{W_{\gamma,\mu}(n; x, t)} \sqrt{W_{\gamma,\mu}(n; y, s)}} (1 + n d_{\mathbb{V}}((x, t), (y, s)))^{-\kappa}.$$

Proof. The proof follows along the line of the estimate on the surface of the cone, but it will be more involved as can be seen in the lower bound of $1 - \xi(x, t, y, s; u, v)$ at (5.15) below.

The kernel $\mathbf{L}_n(W_{\mu,\gamma})$ can be written in terms of the kernel of the Jacobi polynomials by the addition formula in Theorem 5.8, which gives, with $\alpha = \mu + \frac{d-1}{2}$ and $\lambda = 2\mu + \gamma + d$, that

$$\begin{aligned} \mathbf{L}_n(W_{\gamma,\mu}; (x, t), (y, s)) &= c_{\mu,\gamma,d} \int_{[-1,1]^3} L_n^{(\lambda-\frac{1}{2}, -\frac{1}{2})} (2\xi(x, t, y, s; u, v)^2 - 1) \\ &\quad \times (1-u^2)^{\mu-1} (1-v_1^2)^{\alpha-1} (1-v_2^2)^{\gamma-\frac{1}{2}} du dv. \end{aligned}$$

Then, as in the proof of Theorem 4.10, we can apply the estimate (2.17) for $L_n^{\alpha,\beta}$ with $\alpha = \lambda - 1/2$, $\beta = -1/2$ to obtain

$$\begin{aligned} |\mathbf{L}_n(W_{\gamma,\mu}; (x, t), (y, s))| &\leq cn^{2\lambda+1} \int_{[-1,1]^3} \frac{1}{\left(1 + n\sqrt{1 - \xi(x, t, y, s; u, v)^2}\right)^{\kappa+5\mu+3\gamma+\frac{3d}{2}+1}} \\ &\quad \times (1-u^2)^{\mu-1} (1-v_1^2)^{\alpha-1} (1-v_2^2)^{\gamma-\frac{1}{2}} du dv. \end{aligned}$$

We first need an lower bound for $\xi(x, t, y, s; u, v)$. We simplify the notation and write

$$\xi(x, t, y, s; u, v) = v_1 \sqrt{\eta(x, t, y, s; u)} + v_2 \sqrt{1-t} \sqrt{1-s}$$

by introducing the notation

$$\eta(x, t, y, s; u) = \frac{1}{2} \left(ts + \langle x, y \rangle + \sqrt{t^2 - \|x\|^2} \sqrt{s^2 - \|y\|^2} u \right).$$

Let $x = tx'$ and $y = sy'$, where $x', y' \in \mathbb{B}^d$. Then by $|u| \leq 1$, it is easy to verify that $0 \leq \eta(x, t, y, s; u) \leq ts$ and, consequently, $|\xi(x, t, y, s; u, v)| \leq \sqrt{ts} + \sqrt{1-t} \sqrt{1-s} \leq 1$. Furthermore, we can write

$$\begin{aligned} (5.13) \quad 1 - \xi(x, t, y, s; u, v) &= 1 - \sqrt{\eta(x, t, y, s; u)} - \sqrt{1-t} \sqrt{1-s} \\ &\quad + (1-v_1) \sqrt{\eta(x, t, y, s; u)} + (1-v_2) \sqrt{1-t} \sqrt{1-s} \\ &\geq 1 - \sqrt{\eta(x, t, y, s; 1)} - \sqrt{1-t} \sqrt{1-s} \\ &= 1 - \cos d_{\mathbb{V}}((x, t), (y, s)) = 2 \sin^2 \frac{d_{\mathbb{V}}((x, t), (y, s))}{2} \\ &\geq \frac{2}{\pi^2} [d_{\mathbb{V}}((x, t), (y, s))]^2. \end{aligned}$$

As in the case of \mathbb{V}_0^{d+1} , we use this inequality to obtain the estimate

$$|\mathbf{L}_n(W_{\gamma,\mu}; (x, t), (y, s))| \leq cn^{2\lambda+1} \frac{1}{(1 + nd_{\mathbb{V}}((x, t), (y, s)))^{\kappa+\mu+\gamma+\frac{d}{2}}} I(x, t, y, s)$$

with the integral $I(x, t, y, s)$ defined by

$$I(x, t, y, s) = c_{\gamma,\mu} \int_{[-1,1]^3} \frac{(1-u^2)^{\mu-1} (1-v_1^2)^{\alpha-1} (1-v_2^2)^{\gamma-\frac{1}{2}}}{(1 + n\sqrt{1 - \xi(x, t, y, s; u, v)^2})^{4\mu+2\gamma+d+1}} du dv,$$

where $c_{\gamma,\mu}$ is the normalization constant so that $I(x, t, y, s) = 1$ when $n = 0$. To complete the proof, we need to show that

$$\frac{I(x, t, y, s)}{(1 + nd_{\mathbb{V}}((x, t), (y, s)))^{\mu+\gamma+\frac{d}{2}}} \leq \frac{cn^{-(4\mu+2\gamma+d)}}{\sqrt{W_{\gamma,\mu}(n; x, t)} \sqrt{W_{\gamma,\mu}(n; y, s)}}.$$

This is a special case of Lemma 5.10 established below. \square

Lemma 5.10. *Let $d \geq 2$ and $\gamma \geq -\frac{1}{2}$ and $\mu \geq 0$. Then for $\beta \geq 4\mu + 2\gamma + d + 1$,*

$$(5.14) \quad c_{\gamma, \mu} \int_{[-1, 1]^3} \frac{(1-u^2)^{\mu-1} (1-v_1^2)^{\alpha-1} (1-v_2^2)^{\gamma-\frac{1}{2}}}{(1+n\sqrt{1-\xi(x, t, y, s; u, v)})^\beta} du dv$$

$$\leq \frac{cn^{-(4\mu+2\gamma+d)}}{\sqrt{W_{\gamma, \mu}(n; x, t)} \sqrt{W_{\gamma, \mu}(n; y, s)} (1+n\mathbf{d}_V((x, t), (y, s)))^{\beta-5\mu-3\gamma-\frac{3d}{2}-1}}.$$

Proof. Let $I(x, t, y, s)$ be defined as in the proof of the previous proof. Using the lower bound of ξ in (5.13), we see that the left-hand side of (5.14) is bounded by

$$\frac{1}{(1+n\mathbf{d}_V((x, t), (y, s)))^{\beta-4\mu-2\gamma-d-1}} I(x, t, y, s).$$

In order to estimate $I(x, t, y, s)$ we need a refined lower bound for $1 - \xi(x, t, y, s; u, v)$. Let $x = tx'$ and $y = sy'$ with $x', y' \in \mathbb{B}^d$. We claim that for $-1 \leq v_1, v_2, u \leq 1$

$$(5.15) \quad 1 - \xi(x, t, y, s; u, v) \geq \frac{1}{4} \sqrt{ts}(1-v_1) + \sqrt{1-s}\sqrt{1-t}(1-v_2)$$

$$+ \frac{1}{4} \sqrt{ts} \sqrt{1-\|x'\|^2} \sqrt{1-\|y'\|^2} (1-u).$$

To prove this lower bound, we write $A(x', y') = \sqrt{1-\|x'\|^2} \sqrt{1-\|y'\|^2}$ so that the notation can be further simplified as

$$\eta(x, t, y, s; u) = \frac{1}{2} ts (1 + \langle x', y' \rangle + A(x', y') u).$$

We shall use ξ and η without their variables below. Our first step is to write

$$1 - \xi = 1 - \sqrt{ts} - \sqrt{1-t}\sqrt{1-s} + \sqrt{ts} - \sqrt{\eta} + (1-v_1)\sqrt{\eta} + (1-v_2)\sqrt{1-t}\sqrt{1-s}$$

$$\geq \sqrt{ts} - \sqrt{\eta} + (1-v_1)\sqrt{\eta} + (1-v_2)\sqrt{1-t}\sqrt{1-s}.$$

Next, since $\eta(x, t, y, s; u) \leq ts$ by the Cauchy's inequality, we obtain

$$\sqrt{ts} - \sqrt{\eta} = \frac{ts - \eta}{\sqrt{ts} + \sqrt{\eta}} \geq \frac{ts}{4\sqrt{ts}} (1 - \langle x', y' \rangle - A(x', y') u)$$

$$= \frac{\sqrt{ts}}{4} (1 - \langle x', y' \rangle - A(x', y') + (1-u)A(x', y')).$$

Since $1 - \langle x', y' \rangle - A(x', y') \geq 0$ and $1 \geq 1 - v_1$, this leads to

$$1 - \xi \geq B\sqrt{ts}(1-v_1) + \sqrt{1-s}\sqrt{1-t}(1-v_2) + \frac{\sqrt{ts}}{4} A(x', y')(1-u),$$

where B is given by

$$B = \frac{1}{4} (1 - \langle x', y' \rangle - A(x', y')) + \sqrt{\frac{1}{2} (1 + \langle x', y' \rangle + A(x', y') u)},$$

which can be rewritten to give

$$B = 1 - \frac{1}{4} (1-u)A(x', y') - \frac{1}{2} \left(1 - \sqrt{\frac{1}{2} (1 + \langle x', y' \rangle + A(x', y') u)} \right)^2$$

$$\geq 1 - \frac{1}{4} - \frac{1}{2} = \frac{1}{4}.$$

This proves (5.15). We are now ready to prove (5.14). Denote its left-hand side by $J(x, t, u, v)$. Using the inequality (5.15), we obtain

$$J(x, t, y, s) \leq c \int_{[0,1]^3} \frac{(1-u^2)^{\mu-1} (1-v_1^2)^{\mu+\frac{d-1}{2}-1} (1-v_2^2)^{\gamma-\frac{1}{2}}}{\left(1+n\sqrt{B_1(1-v_1)+B_2(1-v_2)+B_3(1-u)}\right)^{4\mu+2\gamma+d+1}} du dv,$$

where $B_1 = \frac{1}{4}\sqrt{t}\sqrt{s}$, $B_2 = \sqrt{1-s}\sqrt{1-t}$ and $B_3 = \frac{1}{4}\sqrt{ts}\sqrt{1-\|x'\|^2}\sqrt{1-\|y'\|^2}$. We can estimate the integral as in the proof of Theorem 4.10 by applying (4.14), which we now need to apply three times, and we obtain

$$\begin{aligned} J(x, t, y, s) &\leq c \frac{n^{-2\mu}}{B_3^\mu} \int_{[0,1]^2} \frac{(1-v_1^2)^{\mu+\frac{d-1}{2}-1} (1-v_2^2)^{\gamma-\frac{1}{2}}}{\left(1+n\sqrt{B_1(1-v_1)+B_2(1-v_2)}\right)^{2\mu+2\gamma+d}} dv \\ &\leq c \frac{n^{-2\mu-2\gamma-1}}{B_3^\mu B_2^{\gamma+\frac{1}{2}}} \int_0^1 \frac{(1-v_1^2)^{\mu+\frac{d-1}{2}-1}}{\left(1+n\sqrt{B_1(1-v_1)}\right)^{2\mu+d-1}} dv_1 \\ &\leq c \frac{n^{-4\mu-2\gamma-d}}{B_3^\mu B_2^{\gamma+\frac{1}{2}} B_1^{\mu+\frac{d-1}{2}}}. \end{aligned}$$

Combining part of B_1 and B_3 , it follows from $\sqrt{t}\sqrt{s}B_3 = \frac{1}{4}\sqrt{t^2-\|x\|^2}\sqrt{s^2-\|y\|^2}$ that

$$\begin{aligned} J(x, t, y, s) &\leq c \frac{n^{-4\mu-2\gamma-d}}{\left(\sqrt{t^2-\|x\|^2}\sqrt{s^2-\|y\|^2}\right)^\mu \left(\sqrt{1-t}\sqrt{1-s}\right)^{\gamma+\frac{1}{2}} \left(\sqrt{t}\sqrt{s}\right)^{\frac{d-1}{2}}} \\ &\leq c \frac{n^{-4\mu-2\gamma-d}}{\left(\sqrt{t^2-\|x\|^2}\sqrt{s^2-\|y\|^2}+n^{-2}\right)^\mu \left(\sqrt{1-t}\sqrt{1-s}+n^{-2}\right)^{\gamma+\frac{1}{2}} \left(\sqrt{t}\sqrt{s}+n^{-2}\right)^{\frac{d-1}{2}}}, \end{aligned}$$

where the second inequality follows since $I(x, t, y, s) \leq 1$ holds trivially. We can now apply (4.15) three times to obtain

$$\begin{aligned} J(x, t, y, s) &\leq c \frac{n^{-4\mu-2\gamma-d}}{\sqrt{W_{\mu,\gamma}(n; x, t)}\sqrt{W_{\mu,\gamma}(n; y, s)}} (1+n|\sqrt{t^2-\|x\|^2}-\sqrt{s^2-\|y\|^2}|)^\mu \\ &\quad \times (1+n|\sqrt{t}-\sqrt{s}|)^{\frac{d-1}{2}} (1+n|\sqrt{1-t}-\sqrt{1-s}|)^{\gamma+\frac{1}{2}}. \end{aligned}$$

Finally, by Lemma 5.4, we conclude

$$J(x, t, y, s) \leq c \frac{n^{-4\mu-2\gamma-d} (1+n d_V((x, t), (y, s)))^{\mu+\gamma+\frac{d}{2}}}{\sqrt{W_{\gamma,\mu}(n; x, t)}\sqrt{W_{\gamma,\mu}(n; y, s)}}$$

which is what we need to complete the proof. \square

The theorem we just proved verifies Assertion 1 for $W_{\gamma,\mu}$. Our next theorem verifies Assertion 2 but only for $W_{\gamma,0}$. Recall that $W_{\gamma,0}$ depends only on t ,

$$W_{\gamma,0}(n; x, t) = (t+n^{-2})^{\frac{d-1}{2}} (1-t+n^{-2})^{\gamma+\frac{1}{2}},$$

Theorem 5.11. *Let $d \geq 2$, $\gamma \geq -\frac{1}{2}$. For $(x_i, t_i), (y, s) \in \mathbb{V}^{d+1}$, and $(x_1, t_1) \in \mathfrak{c}((x_2, t_2), c^* n^{-1})$ with c^* small, and any $\kappa > 0$,*

$$(5.16) \quad |\mathbf{L}_n(W_{\gamma,0}; (x_1, t_1), (y, s)) - \mathbf{L}_n(W_{\gamma,0}; (x_2, t_2), (y, s))| \\ \leq \frac{c_\kappa n^{d+2} \mathbf{d}_\mathbb{V}((x_1, t_1), (x_2, t_2))}{\sqrt{W_{\gamma,0}(n; x_2, t_2)} \sqrt{W_{\gamma,0}(n; y, s)} (1 + n \mathbf{d}_\mathbb{V}((x_2, t_2), (y, s)))^\kappa}.$$

Proof. Let K denote the left-hand side of (5.16). As in the proof of Theorem 4.13, we use the integral expression of $\mathbf{L}_n(W_{\gamma,0})$ and (5.12) to obtain $K \leq K_1 + K_{-1}$, where

$$K_u \leq 2 \int_{[-1,1]^2} \left\| \partial L_n^{\gamma+d-\frac{1}{2}, -\frac{1}{2}}(2(\cdot)^2 - 1) \right\|_{L^\infty(I_{u,v})} |\xi_1(u, v)^2 - \xi_2(u, v)^2| \\ \times (1 - v_1^2)^{\frac{d-1}{2}-1} (1 - v_2^2)^{\gamma-\frac{1}{2}} dv,$$

where $u = 1$ or -1 , $\partial f = f'$, $\xi_i(u, v) = \xi(x_i, t_i, y, s; u, v)$ and $I_{u,v}$ is the interval with end points $\xi_1(u, v)$ and $\xi_2(u, v)$. We first consider the case $u = 1$. Clearly $|\xi_1(1, v)^2 - \xi_2(1, v)^2| \leq 2|\xi_1(1, v) - \xi_2(1, v)|$. We claim that it has the upper bound

$$(5.17) \quad |\xi_1(u, v) - \xi_2(u, v)| \leq c \mathbf{d}_\mathbb{V}((x_1, t_1), (x_2, t_2)) [\Sigma_1 + \Sigma_2(v_1) + \Sigma_3(v_2)],$$

where

$$\Sigma_1 = \mathbf{d}_\mathbb{V}((x_i, t_i), (y, s)) + \mathbf{d}_\mathbb{V}((x_1, t_1), (x_2, t_2)), \\ \Sigma_2(v_1) = (1 - v_1)\sqrt{s}, \\ \Sigma_3(v_2) = (1 - v_2)\sqrt{1 - s}.$$

To see this, we follow the notation used in the proof of Lemma 5.10, writing $\xi_i = v_1 \sqrt{\eta_i(1)} + v_2 \sqrt{1 - t} \sqrt{1 - s}$, where $\eta_i(1) = \eta(x_i, t_i, s_i; 1)$. Using (5.3) and writing $x_i = t_i x'_i$ and $y = sy'$, we obtain

$$(5.18) \quad \xi_1(1, v) - \xi_2(1, v) = \xi_1(1, \mathbf{1}) - \xi_2(1, \mathbf{1}) \\ + (1 - v_1)(\sqrt{\eta_2(1)} - \sqrt{\eta_1(1)}) \\ + (1 - v_2)(\sqrt{1 - t_2} - \sqrt{1 - t_1})\sqrt{1 - s},$$

where $\mathbf{1} = (1, 1)$. For $u = 1$, as in Proposition 5.3,

$$\eta_i(1) = t_i s \cos \frac{\mathbf{d}_\mathbb{B}(x'_i, y')}{2} \quad \text{and} \quad \xi_i(1, \mathbf{1}) = \cos \mathbf{d}_\mathbb{V}((x_i, t_i), (y, s)).$$

Substituting these into (5.18), the resulted identity is similar to the corresponding expression in the proof of Theorem 4.13 and can be analogously estimated using Proposition 5.3 and Lemma 5.4, so that (5.17) follows.

As in the proof of Theorem 4.13, it follows from (2.18) with $m = 1$ that

$$\left\| \partial L_n^{\lambda-\frac{1}{2}, -\frac{1}{2}}(2(\cdot)^2 - 1) \right\|_{L^\infty(I_{1,v})} \\ \leq c \left[\frac{n^{2\lambda+3}}{(1 + n\sqrt{1 - \xi_1(1, v)^2})^\kappa} + \frac{n^{2\lambda+3}}{(1 + n\sqrt{1 - \xi_2(1, v)^2})^\kappa} \right].$$

Hence, K_1 is bounded by, with $\lambda = \gamma + d$,

$$K_1 \leq c \mathbf{d}_\mathbb{V}((x_1, t_1), (x_2, t_2)) \int_{[-1,1]^2} \left[\frac{n^{2\lambda+3}}{(1 + n\sqrt{1 - \xi_1(1, v)^2})^\kappa} + \frac{n^{2\lambda+3}}{(1 + n\sqrt{1 - \xi_2(1, v)^2})^\kappa} \right] \\ \times (\Sigma_1 + \Sigma_2(v_1) + \Sigma_3(v_2)) (1 - v_1^2)^{\frac{d-1}{2}-1} (1 - v_2^2)^{\gamma-\frac{1}{2}} dv.$$

Using the assumption that $(x_1, t_1) \in \mathbf{c}((x_2, t_2), c^* n^{-1})$, the integral that contains Σ_1 term can be handled as in the proof of Theorem 4.13, which leads to

$$\begin{aligned} & \int_{[-1,1]^2} \frac{n^{2\lambda+3}}{(1+n\sqrt{1-\xi_i(1,v)^2})^\kappa} \Sigma_1(1-v_1^2)^{\frac{d-1}{2}-1} (1-v_2^2)^{\gamma-\frac{1}{2}} dv \\ & \leq \frac{n^{2\lambda+2}}{(1+n\mathbf{d}_\mathbb{V}((x,t),(y,s)))^{\kappa-2\gamma-d-2}} \int_{[-1,1]^2} \frac{(1-v_1^2)^{\frac{d-1}{2}-1} (1-v_2^2)^{\gamma-\frac{1}{2}}}{(1+n\sqrt{1-\xi_i(1,v)^2})^{2\gamma+d+2}} dv. \end{aligned}$$

Now, the inequality (5.15) shows that

$$1 - \xi_i(1, v) \geq \frac{1}{4} \sqrt{t_i s} (1 - v_1) + \sqrt{1 - t_i} \sqrt{1 - s} (1 - v_2).$$

Hence, applying Corollary 4.12 with $d-1$ replaced by d and Lemma 5.4, we conclude that the integral in the right-hand side is bound by

$$\frac{cn^{d+2}}{\sqrt{W_{\gamma,0}(n; x_i, t_i)} \sqrt{W_{\gamma,0}(n; y, s)} (1 + n\mathbf{d}_\mathbb{V}((x_i, t_i), (y, s)))^{\kappa(\gamma, d)}},$$

where $\kappa(\gamma, d) = \kappa - 3\gamma - \frac{3d+4}{2}$, since $W_{\gamma,0}(n; x, t) = \mathbf{w}_{\gamma, d+1}(n; t)$. Putting together, we see that the integral containing Σ_1 has the desired upper bound. The two integrals containing $\Sigma_2(v_1)$ and $\Sigma_3(v_2)$ can be estimated similarly and more straightforwardly by using Corollary 4.12. Moreover, the estimate is parallel to the two corresponding terms in the proof of Theorem 4.13 and can be carried out similarly. This completes the proof of the case $u = 1$.

For $u = -1$, we observe that if we define $Y_* = (y, -\sqrt{s^2 - \|y\|^2})$ and

$$\mathbf{d}_*((x, t), (y, s)) = \arccos \left(\sqrt{\frac{\langle X, Y_* \rangle + ts}{2}} + \sqrt{1-t}\sqrt{1-s} \right),$$

then for $u = -1$ and $v = \mathbf{1} = (1, 1)$,

$$\xi(x, t, y, s; -1, \mathbf{1}) = \cos \mathbf{d}_*((x, t), (y, s)).$$

Since $(Y_*, s) \in \mathbb{V}_0^{d+2}$ and $\mathbf{d}_*((x, t), (y, s)) = \mathbf{d}_{\mathbb{V}_0^{d+2}}((X, t), (Y_*, s))$, it follows that \mathbf{d}_* is also a distance function and satisfies, in particular, the triangle inequality. As a result, we can repeat the proof for $u = 1$ to estimate K_{-1} , so that K_{-1} is bounded by the right-hand side of (5.16) with $(1 + n\mathbf{d}_*((x, t), (y, s)))^\kappa$ in place of $(1 + n\mathbf{d}_\mathbb{V}((x, t), (y, s)))^\kappa$ in the denominator. Since $\langle X, Y \rangle \geq \langle X, Y_* \rangle$, $\cos \mathbf{d}_\mathbb{V}((x, t), (y, s)) \geq \cos \mathbf{d}_*((x, t), (y, s))$ and, consequently, $\mathbf{d}_\mathbb{V}((x, t), (y, s)) \leq \mathbf{d}_*((x, t), (y, s))$. Hence, the upper bound of K_{-1} is bounded by the right-hand side of (5.16). This completes the proof. \square

Remark 5.1. It is not clear if this theorem also holds for $W_{\gamma, \mu}$ when $\mu > 0$. The main task lies in establishing an analog of (5.15), perhaps with a multiple of $1 - u$ as one more term. This turns out to be elusive, not because of lack of trying, and there appears to be a real obstacle for (x_i, t_i) around the apex of the cone.

Our next lemma is an analog of Lemma 4.14, which establishes Assertion 3 when $p = 1$ for the weight function $W_{\gamma, \mu}$ on the solid cone.

Lemma 5.12. *Let $d \geq 2$, $\gamma > -1$ and $\mu > -\frac{1}{2}$. For $0 < p < \infty$, assume $\kappa > \frac{2d+2}{p} + (\gamma + \mu + \frac{d}{2})|\frac{1}{p} - \frac{1}{2}|$. Then for $(x, t) \in \mathbb{V}^{d+1}$,*

$$(5.19) \quad \int_{\mathbb{V}^{d+1}} \frac{W_{\gamma, \mu}(s)}{W_{\gamma, \mu}(n; s)^{\frac{p}{2}} (1 + n\mathbf{d}_\mathbb{V}((x, t), (y, s)))^{\kappa p}} dy ds \leq cn^{-d-1} W_{\gamma, \mu}(n; t)^{1-\frac{p}{2}}.$$

Proof. Let J_p denote the left-hand side of (5.19). Using the doubling property of $W_{\beta,\gamma}$ and by Lemma 2.4, it is sufficient to estimate $J_{2,\kappa}$. Setting $y = sy'$, $y' \in \mathbb{B}^d$, we obtain

$$J_2 = \int_0^1 s^d \int_{\mathbb{B}^d} \frac{W_{\gamma,\mu}(sy', s)}{W_{\gamma,\mu}(n; sy', s)(1 + n\mathbf{d}_{\mathbb{V}}((x, t), (sy', s)))^{2\kappa}} dy' ds.$$

Using $w_{\gamma,d}(n; t)$ defined in (4.10), it follows readily that

$$\frac{W_{\gamma,\mu}(y, s)}{W_{\gamma,\mu}(n; y, s)} = c \frac{sw_{-1,\gamma}(s)}{(s^2 - \|y\|^2 + n^{-2})^{\frac{1}{2}} w_{\gamma,d+1}(n; s)} \leq c \frac{w_{-1,\gamma}(n; s)}{w_{\gamma,d+1}(n; s) \sqrt{1 - \|y'\|^2}},$$

which leads to

$$J_2 \leq c \int_0^1 s^d \int_{\mathbb{B}^d} \frac{w_{-1,\gamma}(n; s)}{w_{\gamma,d+1}(n; s)(1 + n\mathbf{d}_{\mathbb{V}}((x, t), (sy', s)))^{2\kappa}} \frac{dy'}{\sqrt{1 - \|y'\|^2}} ds.$$

Setting $x = tx'$, $X = (x', \sqrt{1 - \|x'\|^2})$ and $Y = (y', \sqrt{1 - \|y'\|^2})$, so that $\mathbf{d}_{\mathbb{S}^d}(X, Y)$, we use the identity

$$(5.20) \quad \int_{\mathbb{B}^d} g\left(y', \sqrt{1 - \|y'\|^2}\right) \frac{dy'}{\sqrt{1 - \|y'\|^2}} = \int_{\mathbb{S}_+^d} g(y) d\sigma(y),$$

where \mathbb{S}_+^d denotes the upper hemisphere of \mathbb{S}^d , which allows us to follow the proof of Proposition 4.15 to obtain

$$J \leq c \int_0^1 \int_{-1}^1 \frac{s^d w_{-1,\gamma}(s)(1 - u^2)^{\frac{d-2}{2}}}{w_{\gamma,d+1}(n; s) \left(1 + n \arccos\left(\sqrt{ts}\sqrt{\frac{1+u}{2}} + \sqrt{1-t}\sqrt{1-s}\right)\right)^{2\kappa}} du ds.$$

Comparing with the proof of Proposition 4.15, we see that the above integral with d replaced by $d-1$ has already appeared in that proof, so that it is bounded by cn^{-d-1} as the proof of Proposition 4.15 shows. The proof is completed. \square

Proposition 5.13. *Let $d \geq 2$, $\mu \geq 0$ and $\gamma \geq -\frac{1}{2}$. For $0 < p < \infty$ and $(x, t) \in \mathbb{V}^{d+1}$,*

$$\int_{\mathbb{V}^{d+1}} |\mathbf{L}_n(W_{\gamma,\mu}; (x, t), (y, s))|^p W_{\gamma,\mu}(y, s) dy ds \leq c \left(\frac{n^{d+1}}{W_{\gamma,\mu}(n; x, t)} \right)^{p-1}.$$

This follows immediately from applying Lemma 5.12 on (i) of Theorem 5.9.

Corollary 5.14. *For $d \geq 2$ and $\gamma \geq -\frac{1}{2}$, the space $(\mathbb{V}^{d+1}, W_{\gamma,0}, \mathbf{d}_{\mathbb{V}})$ is a localizable homogeneous space.*

5.5. Maximal ε -separated sets and MZ inequality. We provide a construction of maximal ε -separated sets on the solid cone as defined in Definition 2.13.

Our construction follows the approach in Subsection 4.5. Instead of starting with ε -separated sets on the unit sphere, we now need such sets on the unit ball \mathbb{B}^d . We adopt the following notation. For $\varepsilon > 0$, we denote by $\Xi_{\mathbb{B}}(\varepsilon)$ a maximal ε -separated set on the unit ball \mathbb{B}^d and we let $\mathbb{B}_u(\varepsilon)$ be the subsets in \mathbb{B}^d so that the collection $\{\mathbb{B}_u(\varepsilon) : u \in \Xi_{\mathbb{B}}(\varepsilon)\}$ is a partition of \mathbb{B}^d , and we assume

$$(5.21) \quad \mathbf{c}_{\mathbb{B}}(u, c_1\varepsilon) \subset \mathbb{B}_u(\varepsilon) \subset \mathbf{c}_{\mathbb{B}}(u, c_2\varepsilon), \quad u \in \Xi_{\mathbb{B}}(\varepsilon),$$

where $\mathbf{c}_{\mathbb{B}}(u, \varepsilon)$ denotes the ball centered at u with radius ε in \mathbb{B}^d , c_1 and c_2 depend only on d . It is known (see, for example, [38]) that such a $\Xi_{\mathbb{B}}(\varepsilon)$ exists for all $\varepsilon > 0$ and its cardinality satisfies

$$(5.22) \quad c'_d \varepsilon^{-d} \leq \#\Xi_{\mathbb{B}}(\varepsilon) \leq c_d \varepsilon^{-d}.$$

On the solid cone \mathbb{V}^{d+1} we denote by $\Xi_{\mathbb{V}} = \Xi_{\mathbb{V}}(\varepsilon)$ a maximum ε -separated set and denote by $\{\mathbb{V}(u, t) : (tu, t) \in \Xi_{\mathbb{V}}\}$ a partition of \mathbb{V}^{d+1} . We give a construction of these sets.

Let $\varepsilon > 0$ and let $N = \lfloor \frac{\pi}{2}\varepsilon^{-1} \rfloor$. We define $t_j = \sin^2 \frac{\theta_j}{2}$, t_j^+ and t_j^- , $1 \leq j \leq N$, as in Subsection 4.5, so that

$$\mathbb{V}^{d+1} = \bigcup_{j=1}^N \mathbb{V}^{(j)}, \quad \text{where} \quad \mathbb{V}^{(j)} := \{(x, t) \in \mathbb{V}^{d+1} : t_j^- < t \leq t_j^+\}.$$

Let $\varepsilon_j := (2\sqrt{t_j})^{-1}\pi\varepsilon$. Then $\Xi_{\mathbb{B}}(\varepsilon_j)$ is the maximal ε_j -separated set of \mathbb{B}^d such that, for each $j \geq 1$, $\{\mathbb{B}_u(\varepsilon_j) : u \in \Xi_{\mathbb{B}}(\varepsilon_j)\}$ is a partition of \mathbb{B}^d and $\#\Xi_{\mathbb{B}}(\varepsilon_j) \sim \varepsilon_j^{-d}$. For each $j = 1, \dots, N$, we decompose $\mathbb{V}^{(j)}$ by

$$\mathbb{V}^{(j)} = \bigcup_{u \in \Xi_{\mathbb{B}}(\varepsilon_j)} \mathbb{V}(u, t_j), \quad \text{where} \quad \mathbb{V}(u, t_j) := \{(tv, t) : t_j^- < t \leq t_j^+, v \in \mathbb{B}_u(\varepsilon_j)\}.$$

Finally, we define the set $\Xi_{\mathbb{V}}$ of \mathbb{V}^{d+1} by

$$\Xi_{\mathbb{V}} = \{(t_j u, t_j) : u \in \Xi_{\mathbb{B}}(\varepsilon_j), 1 \leq j \leq N\}.$$

Proposition 5.15. *Let $\varepsilon > 0$ and $N = \lfloor \frac{\pi}{2}\varepsilon^{-1} \rfloor$. Then $\Xi_{\mathbb{V}}$ is a maximal ε -separated set of \mathbb{V}^{d+1} and $\{\mathbb{V}(t_j u, t_j) : u \in \Xi_{\mathbb{B}}(\varepsilon_j), 1 \leq j \leq N\}$ is a partition*

$$\mathbb{V}^{d+1} = \bigcup_{j=1}^N \bigcup_{u \in \Xi_{\mathbb{B}}(\varepsilon_j)} \mathbb{V}(u, t_j).$$

Moreover, there are positive constants c_1 and c_2 depending only on d such that

$$(5.23) \quad \mathbf{c}((t_j u, t_j), c_1 \varepsilon) \subset \mathbb{V}(u, t_j) \subset \mathbf{c}((t_j u, t_j), c_2 \varepsilon), \quad (t_j u, t_j) \in \Xi_{\mathbb{V}},$$

and c'_d and c_d depending only on d such that

$$c'_d \varepsilon^{-d-1} \leq \#\Xi_{\mathbb{V}} \leq c_d \varepsilon^{-d-1}.$$

Proof. The proof is parallel to that of Proposition 4.17. In fact, it can be carried out almost verbatim under obvious modifications such as replacing $\mathbf{d}_{\mathbb{S}}$ by $\mathbf{d}_{\mathbb{B}}$ and \mathbb{V}_0 by \mathbb{V} . We shall omit it. \square

Even though we only established Assertion 2 for $W_{\gamma,0}$ for $\gamma \geq 0$, it is sufficient for applying Theorem 5.16 to conclude that the Marcinkiewicz-Zygmund inequality on a maximal ε -separated set holds for any doubling weight on \mathbb{V}^{d+1} .

Theorem 5.16. *Let W be a doubling weight on \mathbb{V}^{d+1} . Let $\Xi_{\mathbb{V}}$ be a maximal $\frac{\delta}{n}$ -separated subset of \mathbb{V}^{d+1} and $0 < \delta \leq 1$.*

(i) *For all $0 < p < \infty$ and $f \in \Pi_m^{d+1}$ with $n \leq m \leq cn$,*

$$\sum_{z \in \Xi_{\mathbb{V}}} \left(\max_{(x,t) \in \mathbf{c}((z,r), \frac{\delta}{n})} |f(x, t)|^p \right) W(\mathbf{c}((z, r), \frac{\delta}{n})) \leq c_{W,p} \|f\|_{p,W}^p$$

where $c_{W,p}$ depends on p when p is close to 0 and on the doubling constant $\alpha(W)$.

(ii) *For $0 < r < 1$, there is a $\delta_r > 0$ such that for $\delta \leq \delta_r$, $r \leq p < \infty$ and $f \in \Pi_n^{d+1}$,*

$$\|f\|_{p,W}^p \leq c_{W,r} \sum_{z \in \Xi} \left(\min_{(x,t) \in \mathbf{c}((z,r), \frac{\delta}{n})} |f(x, t)|^p \right) W(\mathbf{c}((z, r), \frac{\delta}{n}))$$

where $c_{W,r}$ depends only on $L(W)$ and on r when r is close to 0.

5.6. Cubature rules and localized tight frames. We first verify Assertion 4 in Subsection 2.5 by constructing fast decaying polynomials on the solid cone.

Lemma 5.17. *Let $d \geq 2$. For each $(x, t) \in \mathbb{V}^{d+1}$, there is a polynomial $\mathcal{T}_{x,t}$ of degree n that satisfies*

(1) $\mathcal{T}_{x,t}(x, t) = 1$, $\mathcal{T}_{x,t}(y, s) \geq c > 0$ if $(y, s) \in \mathbf{c}((x, t), \frac{\delta}{n})$, and for every $\kappa > 0$,

$$0 \leq \mathcal{T}_{x,t}(y, s) \leq c_\kappa (1 + \mathbf{d}_\mathbb{V}((x, t), (y, s)))^{-\kappa}, \quad (y, s) \in \mathbb{V}^{d+1}.$$

(2) there is a polynomial q of degree n such that $q(x, t)\mathcal{T}_{x,t}$ is a polynomial of degree $3n$ in (x, t) and $1 \leq q(x, t) \leq c$.

Proof. For $(x, t), (y, s) \in \mathbb{V}^{d+1}$, we introduce the notation $X = (x, \sqrt{t^2 - \|x\|^2})$ and $Y = (y, \sqrt{s^2 - \|y\|^2})$. Moreover, we also denote $Y_* = (y, -\sqrt{t^2 - \|y\|^2})$. Then $(X, t), (Y, s)$ and (Y_*, s) are all elements of \mathbb{V}_0^{d+2} . Let $T_{(X,t)}$ denote the polynomial of degree n on \mathbb{V}_0^{d+2} defined in Lemma 4.19. We now define

$$\mathcal{T}_{x,t}(y, s) := \frac{T_{(X,t)}(Y, s) + T_{(X,t)}(Y_*, s)}{1 + T_{(X,t)}(X_*, t)}.$$

Since $T_{(X,t)}(X, t) = 1$, it follows that $\mathcal{T}_{x,t}(x, t) = 1$. Since $\sqrt{\frac{\langle X_*, X \rangle + t^2}{2}} = \|x\|$, we have

$$T_{(X,t)}(X_*, t) = \frac{S_n(\|x\| + (1-t)) + S_n(\|x\| - (1-t))}{1 + S_n(2t-1)}.$$

By $0 \leq S_n(t) \leq c$, we see that $0 \leq T_{(X,t)}(X_*, s) \leq c$. In particular, it follows that

$$\mathcal{T}_{x,t}(y, s) \geq c T_{(X,t)}(Y, s) \geq c > 0, \quad (y, s) \in \mathbf{c}((x, t), \frac{\delta}{n}),$$

since $\mathbf{d}_\mathbb{V}((x, t), (y, s)) = \mathbf{d}_{\mathbb{V}_0^{d+2}}((X, t), (Y, s))$. Furthermore, since $\cos \mathbf{d}_{\mathbb{V}_0}((X, t), (Y, s)) \geq \cos \mathbf{d}_{\mathbb{V}_0}((X, t), (Y_*, s))$, we obtain

$$\mathbf{d}_{\mathbb{V}_0}((X, t), (Y_*, s)) \geq \mathbf{d}_{\mathbb{V}_0}((X, t), (Y, s)) = \mathbf{d}_\mathbb{V}((x, t), (y, s)).$$

Hence, using the estimate of $T_{(X,t)}$ in Lemma 4.19, we conclude that

$$\begin{aligned} 0 \leq \mathcal{T}_{x,t}(y, s) &\leq c \left[(1 + n \mathbf{d}_{\mathbb{V}_0}((X, t), (Y, s)))^{-\kappa} + (1 + n \mathbf{d}_{\mathbb{V}_0}((X, t), (Y_*, s)))^{-\kappa} \right] \\ &\leq c (1 + n \mathbf{d}_\mathbb{V}((x, t), (y, s)))^{-\kappa}. \end{aligned}$$

Finally, let $q(x, t) = (1 + S_n(2t-1))T_{(X,t)}(X_*, t)$. Then $q(x, t)$ is a polynomial of degree at most $2n$, so that $q(x, t)\mathcal{T}_{x,t}$ is a polynomial of degree at most $3n$ and $1 \leq q(x, t) \leq c$. This completes the proof. \square

By Propositions 2.17 and 2.18 and using Theorem 5.9, we have established the following result on the Christoffel function $\lambda_n(W)$.

Corollary 5.18. *Let W be a doubling weight function on \mathbb{V}^{d+1} . Then*

$$\lambda_n(W; (x, t)) \leq c W(\mathbf{c}((x, t), \frac{1}{n})).$$

Moreover, for $\gamma \geq -\frac{1}{2}$ and $\mu \geq 0$,

$$\lambda_n(W_{\gamma, \mu}; (x, t)) \geq c W_{\gamma, \mu}(\mathbf{c}((x, t), \frac{1}{n})) = c n^{-d-1} W_{\gamma, \mu}(n; t).$$

Since the lower bound of the Christoffel function holds for all doubling weight, we see that the cubature rules in Theorem 2.20 holds for all doubling weights on the solid cone.

Theorem 5.19. *Let W be a doubling weight on \mathbb{V}^{d+1} . Let Ξ be a maximum $\frac{\delta}{n}$ -separated subset of \mathbb{V}^{d+1} . There is a $\delta_0 > 0$ such that for $0 < \delta < \delta_0$ there exist positive numbers $\lambda_{z,r}$, $(z, r) \in \Xi$, so that*

$$(5.24) \quad \int_{\mathbb{V}^{d+1}} f(x)W(x, t)dxdt = \sum_{(z,r) \in \Xi} \lambda_{z,r}f(z, r), \quad \forall f \in \Pi_n^{d+1}.$$

Moreover, $\lambda_{z,r} \sim W(\mathbf{c}((z, r), \frac{\delta}{n}))$ for all $(z, r) \in \Xi$.

We can now state our local frame on the solid cone. For $j = 0, 1, \dots$, let Ξ_j be a maximal $\frac{\delta}{2^j}$ -separated subset in \mathbb{V}^{d+1} , so that

$$\int_{\mathbb{V}^{d+1}} f(x, t)W(x, t)dxdt = \sum_{(z,r) \in \Xi_j} \lambda_{(z,r),j}f(z, r), \quad f \in \Pi_{2^j}^{d+1}.$$

Denote by $\mathbf{L}_n(W) * f$ the near best approximation operator defined by

$$\mathbf{L}_n(W) * f(x, t) = \int_{\mathbb{V}^{d+1}} f(y, s)\mathbf{L}_n(W; (x, t), (y, s))W(y, s)dyds.$$

For $j = 0, 1, \dots$, define the operator $F_j(W)$ by

$$F_j(W) * f = \mathbf{L}_{2^{j-1}}(W) * f$$

and define the frame elements $\psi_{(z,r),j}$ for $(z, r) \in \Xi_j$ by

$$\psi_{(z,r),j}(x, t) := \sqrt{\lambda_{(z,r),j}F_j((x, t), (z, r))}, \quad (x, t) \in \mathbb{V}^{d+1}.$$

Then $\Phi = \{\psi_{(z,r),j} : (z, r) \in \Xi_j, j = 1, 2, 3, \dots\}$ is a tight frame.

Theorem 5.20. *Let W be a doubling weight on \mathbb{V}^{d+1} . If $f \in L^2(\mathbb{V}^{d+1}, W)$, then*

$$f = \sum_{j=0}^{\infty} \sum_{(z,r) \in \Xi_j} \langle f, \psi_{(z,r),j} \rangle_W \psi_{(z,r),j} \quad \text{in } L^2(\mathbb{V}^{d+1}, W)$$

and

$$\|f\|_{2,W} = \left(\sum_{j=0}^{\infty} \sum_{(z,r) \in \Xi_j} |\langle f, \psi_{(z,r),j} \rangle_W|^2 \right)^{1/2}.$$

Furthermore, for $W_{\gamma,\mu}$ with $\gamma \geq -\frac{1}{2}$ and $\mu \geq 0$, the frame is highly localized in the sense that, for every $\kappa > 0$, there exists a constant $c_\kappa > 0$ such that

$$(5.25) \quad |\psi_{(z,r),j}(x, t)| \leq c_\sigma \frac{2^{j(d+1)/2}}{\sqrt{W_{\gamma,\mu}(2^j; x, t)(1 + 2^j d_V((x, t), (z, r)))^\kappa}}, \quad (x, t) \in \mathbb{V}^{d+1}.$$

The frame elements are well defined for all doubling weight by Theorem 5.19. The decomposition is the consequence of Theorem 2.21. Moreover, the localization (5.25) follows from Theorem 5.9 and $\lambda_{(z,r),j} \sim 2^{-j(d+1)}W_{\gamma,\mu}(2^j; t)$ that holds for $W_{\gamma,\mu}$ as see from Corollary 5.18 and (5.5).

5.7. Characterization of best approximation. For $f \in L^p(\mathbb{V}^{d+1}, W)$, we denote by $\mathbf{E}_n(f)_{p,W}$ the error of best approximation to f from Π_n^{d+1} , the space of polynomials of degree at most n , in the norm $\|\cdot\|_{p,W}$,

$$\mathbf{E}_n(f)_{p,W} := \inf_{g \in \Pi_n^{d+1}} \|f - g\|_{p,W}, \quad 1 \leq p \leq \infty.$$

We give a characterization of this quantity in terms of the modulus of smoothness defined via the operator $\mathbf{S}_{\theta,W}$ and the K -functional defined via the differential operator $\mathfrak{D}_{\gamma,\mu}$ for $W_{\gamma,\mu}$.

For $f \in L^p(\mathbb{V}^{d+1}, W_{\gamma,\mu})$ and $r > 0$, the modulus of smoothness is defined by

$$\omega_r(f; \rho)_{p,W_{\gamma,\mu}} = \sup_{0 \leq \theta \leq \rho} \left\| (I - \mathbf{S}_{\theta,W_{\gamma,\mu}})^{r/2} f \right\|_{p,W_{\gamma,\mu}}, \quad 1 \leq p \leq \infty,$$

where the operator $\mathbf{S}_{\theta,W_{\gamma,\mu}}$ is defined by, for $n = 0, 1, 2, \dots$ and $\lambda = 2\mu + \gamma + d$,

$$\text{proj}_n(W_{\gamma,\mu}; \mathbf{S}_{\theta,W_{\gamma,\mu}} f) = R_n^{(\lambda-\frac{1}{2}, -\frac{1}{2})}(\cos \theta) \text{proj}_n(W_{\gamma,\mu}; f).$$

Moreover, in terms of the fractional differential operator $(-\mathfrak{D}_{\gamma,\mu})^{\frac{r}{2}}$, the K -functional is defined for a weight W on \mathbb{V}^{d+1} by

$$\mathbf{K}_r(f, \rho)_{p,W} := \inf_{g \in \mathcal{W}_p^r(\mathbb{V}^{d+1}, W)} \left\{ \|f - g\|_{p,W} + \rho^r \|(-\mathfrak{D}_{\gamma,\mu})^{\frac{r}{2}} f\|_{p,W} \right\},$$

where $\mathcal{W}_p^r(\mathbb{V}^{d+1}, W)$ denotes the Sobolev space consisting of functions in $L^p(\mathbb{V}^{d+1}, W)$ with finite $\|(-\mathfrak{D}_{\gamma,\mu})^{\frac{r}{2}} f\|_{p,W}$.

The weight function $W_{\gamma,\mu}$ admits Assertion 1 and 3 by Theorem 5.9 and Lemma 5.12. We now verify that the Assertion 5 in Subsection 3.5 holds. By Theorem 4.8, the kernel $L_n^{(r)}(\varpi)$ in Assertion 5 becomes

$$\mathbf{L}_n^{(r)}(W_{\gamma,\mu}; (x, t), (y, s)) = \sum_{k=0}^{\infty} \hat{a}\left(\frac{k}{n}\right) (k(k + 2\mu + \gamma + d))^{\frac{r}{2}} \mathbf{P}_k(W_{\gamma,\mu}; (x, t), (y, s)).$$

Lemma 5.21. *Let $\gamma \geq -\frac{1}{2}$ and $\mu \geq 0$. Let $\kappa > 0$. Then, for $r > 0$ and $(x, t), (y, s) \in \mathbb{V}^{d+1}$,*

$$|\mathbf{L}_n^{(r)}(W_{\gamma,\mu}; (x, t), (y, s))| \leq c_{\kappa} \frac{n^{r+d}}{\sqrt{W_{\gamma,\mu}(n; x, t)} \sqrt{W_{\gamma,\mu}(n; y, s)} (1 + n d_{\mathbb{V}}((x, t), (y, s)))^{\kappa}}.$$

Proof. By (5.10), the kernel can be written as

$$\begin{aligned} \mathbf{L}_n^{(r)}(W_{\gamma,\mu}; (x, t), (y, s)) &= c_{\mu,\gamma,d} \int_{[-1,1]^3} L_{n,r} (2\xi(x, t, y, s; u, v)^2 - 1) \\ &\quad \times (1 - u^2)^{\mu-1} (1 - v_1^2)^{\alpha-1} (1 - v_2^2)^{\gamma-\frac{1}{2}} du dv, \end{aligned}$$

in which $\alpha = \mu + \frac{d-1}{2}$ and $L_{n,r}$ is defined by, with $\lambda = 2\alpha + \gamma + 1$,

$$L_{n,r}(t) = \sum_{k=0}^{\infty} \hat{a}\left(\frac{k}{n}\right) (k(k + \gamma + d - 1))^{\frac{r}{2}} \frac{P_n^{(\lambda-\frac{1}{2}, -\frac{1}{2})}(1) P_n^{(\lambda-\frac{1}{2}, -\frac{1}{2})}(t)}{h^{(\lambda-\frac{1}{2}, -\frac{1}{2})}}.$$

Applying (2.18) with $\eta(t) = \hat{a}(t) (t(t + n^{-1}(2\mu + \gamma + d)))^{\frac{r}{2}}$ and $m = 0$, it follows that

$$|L_{n,r}(t)| \leq c n^r \frac{n^{2\lambda+1}}{(1 + n\sqrt{1-t})^{\ell}}.$$

Using this estimate, we can then deduce the proof to one that has already appeared for $\mathbf{L}_n(W_{\gamma,\mu})$ in the proof of Theorem 4.10. \square

With Assertions 1, 3 and 5 verified for $W_{\gamma,\mu}$, the characterization of the best approximation by polynomials in Subsection 3.3 holds on the solid cone, which we state below.

Theorem 5.22. *Let $f \in L^p(\mathbb{V}^{d+1}, W)$ if $1 \leq p < \infty$ and $f \in C(\mathbb{V}^{d+1})$ if $p = \infty$. Let $r > 0$ and $n = 1, 2, \dots$. For $W = W_{\gamma,\mu}$ with $\gamma \geq -\frac{1}{2}$ and $\mu \geq 0$, there holds*

(i) *direct estimate*

$$\mathbf{E}_n(f)_{p,W_{\gamma,\mu}} \leq c \mathbf{K}_r(f; n^{-1})_{p,W_{\gamma,\mu}}.$$

(ii) *inverse estimate, for $\mu = 0$,*

$$\mathbf{K}_r(f; n^{-1})_{p,W_{\gamma,0}} \leq cn^{-r} \sum_{k=0}^n (k+1)^{r-1} \mathbf{E}_k(f)_{p,W_{\gamma,0}}.$$

Proof. The direct estimate follows from Theorem 3.14, which requires only Assertions 1, 3, 5 and holds for $W_{\gamma,\mu}$ by Theorem 5.9 and Lemma 5.12. The inverse estimate follows from Theorem 3.12, which requires one weight function that admits all Assertions 1–3 and 5 and that holds for $W_{\gamma,0}$ on the cone. \square

Both the direct and the inverse estimates hold for the weight function $W_{\gamma,\mu}$ in the above theorem. However, it should be noted that the inverse estimate uses the K -functional

$$\mathbf{K}_r(f, \rho)_{p,W} = \inf_{g \in \mathcal{W}_p^r(\mathbb{V}^{d+1}, W)} \left\{ \|f - g\|_{p,W} + \rho^r \|(-\mathfrak{D}_{\gamma,0})^{\frac{r}{2}} f\|_{p,W} \right\}$$

defined via the operator $\mathfrak{D}_{\gamma,0}$ for the weight $W_{\gamma,0}$.

For $W = W_{\gamma,\mu}$, both direct and inverse estimates can be given via the modulus of smoothness, since it is equivalent to the K -functional.

Theorem 5.23. *Let $\gamma \geq -\frac{1}{2}$, $\mu \geq 0$ and $f \in L_p^r(\mathbb{V}^{d+1}, W_{\gamma,\mu})$, $1 \leq p \leq \infty$. Then for $0 < \theta \leq \pi/2$ and $r > 0$*

$$c_1 \mathbf{K}_r(f; \theta)_{p,W_{\gamma,\mu}} \leq \omega_r(f; \theta)_{p,W_{\gamma,\mu}} \mathbf{K}_r(f; \theta)_{p,W_{\gamma,\mu}}.$$

REFERENCES

- [1] R. Askey, *Orthogonal Polynomials and Special Functions*, Regional Conference Series in Applied Mathematics **21**, SIAM, Philadelphia, 1975.
- [2] P. Baldi, G. Kerkycharian, D. Marinucci, and D. Picard. Asymptotics for spherical needlets, *Ann. Statist.* **37** (2009), 1150–1171.
- [3] P. Baldi, G. Kerkycharian, D. Marinucci, and D. Picard. Adaptive density estimation for directional data using needlets, *Ann. Statist.* **37** (2009), 3362–3395.
- [4] H. Berens, P. L. Butzer, and S. Pawelke, Limitierungsverfahren von Reihen mehrdimensionaler Kugelfunktionen und deren Saturationsverhalten, *Publ. Res. Inst. Math. Sci. Ser. A.* **4** (1968), 201–268.
- [5] G. Brown, F. Dai, Approximation of smooth functions on compact two-point homogeneous spaces, *J. Funct. Anal.* **220** (2005), 401–423.
- [6] A. Cohen and M. Dolbeault, Optimal sampling and Christoffel functions on general domains. arXiv:2010.11040
- [7] F. Dai, Multivariate polynomial inequalities with respect to doubling weights and A_∞ weights, *J. Funct. Anal.* **235** (2006), 137–170.
- [8] F. Dai and A. Prymak, L_p -Bernstein inequalities on C^2 domains, 2020. arXiv:2010.06728
- [9] F. Dai and A. Prymak, On directional Whitney inequality, 2020. arXiv:2010.08374

- [10] F. Dai and H. Wang, Optimal cubature formulas in weighted Besov spaces with A_∞ weights on multivariate domains, *Const. Approx.* **37** (2013), 167–194.
- [11] F. Dai and Y. Xu, Moduli of smoothness and approximation on the unit sphere and the unit ball. *Adv. Math.* **224** (2010), 1233–1310.
- [12] F. Dai and Y. Xu, *Approximation theory and harmonic analysis on spheres and balls*. Springer Monographs in Mathematics, Springer, 2013.
- [13] C. F. Dunkl, Differential-difference operators associated to reflection groups. *Trans. Amer. Math. Soc.* **311** (1989), 167–183.
- [14] C. F. Dunkl and Y. Xu, *Orthogonal Polynomials of Several Variables*. Encyclopedia of Mathematics and its Applications **155**, Cambridge University Press, Cambridge, 2014.
- [15] J. Dziubański, Triebel-Lizorkin spaces associated with Laguerre and Hermite expansions, *Proc. Amer. Math. Soc.* **125** (1997), 3547–3554.
- [16] J. Dziubański, E. Hernández, Band-limited wavelets with subexponential decay, *Canad. Math. Bull.* **41** (1998), 398–403.
- [17] R. A. DeVore and G. G. Lorentz, *Constructive approximation*, Grundlehren der Mathematischen Wissenschaften, vol. 303, Springer-Verlag, Berlin, 1993.
- [18] J. Epperson, Hermite and Laguerre wave packet expansions, *Studia Math.* **126** (1997), 199–217.
- [19] J. Faraut and A. Korányi *Analysis on symmetric cones*. Oxford Mathematical Monographs. Oxford University Press, New York, 1994.
- [20] G. Kerkycharian, P. Petrushev, D. Picard and Y. Xu, Decomposition of Triebel-Lizorkin and Besov spaces in the context of Laguerre expansions, *J. Funct. Anal.* **256** (2009), 1137–1188.
- [21] H. L. Krall and I. M. Sheffer, Orthogonal polynomials in two variables, *Annali di Matema. Pura & Appl.* **76** (1967), 325–376.
- [22] A. Króo, Christoffel functions on convex and starlike domains in \mathbb{R}^d , *J. Math. Anal. Appl.* **421** (2015), 718–729.
- [23] A. Króo and D. S. Lubinsky, Christoffel functions and universality in the bulk for multivariate orthogonal polynomials, *Canadian J. Math.* **65**, 600–620.
- [24] G. Kyriazis, P. Petrushev and Y. Xu, Jacobi decomposition of weighted Triebel-Lizorkin and Besov spaces, *Studia Math.* **186** (2008), 161–202.
- [25] G. Kyriazis, P. Petrushev and Y. Xu, Decomposition of weighted Triebel-Lizorkin and Besov spaces on the ball, *Proc. London Math. Soc.* **97** (2008), 477–513.
- [26] K. Ivanov and P. Petrushev, Fast memory efficient evaluation of spherical polynomials at scattered points, *Adv. Comput. Math.* **41** (2015), 191–230.
- [27] K. Ivanov and P. Petrushev, Highly effective stable evaluation of bandlimited functions on the sphere, *Numer. Algorithms*, **71** (2016), 585–611.
- [28] K. Ivanov, P. Petrushev and Y. Xu, Sub-exponentially localized kernels and frames induced by orthogonal expansions. *Math. Z.* **264** (2010), 361–397.
- [29] K. Ivanov, P. Petrushev and Y. Xu, Decomposition of spaces of distributions induced by tensor product bases. *J. Funct. Anal.* **263** (2012), 1147–1197.
- [30] K. Ivanov and V. Totik, Fast decreasing polynomials, *Constr. Approx.* **6** (1990), 1–20.
- [31] Q. T. Le Gia, I. H. Sloan, Y. G. Wang and R. S. Womersley, Needlet approximation for isotropic random fields on the sphere. *J. Approx. Theory* **216** (2017), 86–116.
- [32] G. Mastroianni and V. Totik Weighted polynomial inequalities with doubling and A_∞ weights. *Const. Approx.* **16** (2000) 37–71.
- [33] H. N. Mhaskar, F. J. Narcowich and J. D. Ward, Spherical Marcinkiewicz-Zygmund inequalities and positive quadrature, *Math. Comp.* **70** (2001), 1113 - 1130 (Corrigendum: *Math. Comp.* **71** (2001), 453 - 454).
- [34] F. J. Narcowich, P. Petrushev and J. D. Ward, Localized tight frames on spheres, *SIAM J. Math. Anal.* **38** (2006), 574–594.
- [35] F. J. Narcowich, P. Petrushev and J. D. Ward, Decomposition of Besov and Triebel-Lizorkin spaces on the sphere, *J. Funct. Anal.* **238** (2006), 530–564.
- [36] S. Pawelke, Über Approximationsordnung bei Kugelfunktionen und algebraischen Polynomen, *Tôhoku Math. J.* **24** (1972), 473–486.
- [37] P. Petrushev and Y. Xu, Localized polynomial frames on the interval with Jacobi weights, *J. Fourier Anal. and Appl.* **11** (2005), 557–575.
- [38] P. Petrushev and Y. Xu, Localized polynomial frames on the ball, *Constr. Approx.* **27** (2008), 121–148.

- [39] P. Petrushev and Y. Xu, Decomposition of spaces of distributions induced by Hermite expansion, *J. Fourier Anal. and Appl.* **14** (2008), 372–414.
- [40] A. Prymak, Upper estimates of Christoffel function on convex domains, *J. Math. Anal. Appl.* **455** (2017), 1984–2000.
- [41] A. Prymak, Christoffel functions on planar domains with piecewise smooth boundary, *Acta Math. Hungar.* **158** (2019), 216–234.
- [42] Kh. P. Rustamov, On the approximation of functions on a sphere, (Russian), *Izv. Ross. Akad. Nauk Ser. Mat.* **57** (1993), 127–148; translation in Russian Acad. Sci. *Izv. Math.* **43** (1994), no. 2, 311–329.
- [43] E. Saff and V. Totik, *Logarithmic Potentials with External Fields*, Springer-Verlag, Berlin-Heidelberg, 1997.
- [44] E. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals* Princeton University Press, Princeton, 1993.
- [45] E. Stein and G. Weiss, *Introduction to Fourier Analysis on Euclidean Spaces*. Princeton Univ. Press, Princeton, 1971.
- [46] G. Szegő, *Orthogonal polynomials*. 4th edition, Amer. Math. Soc., Providence, RI. 1975
- [47] V. Totik, Polynomial approximation on polytopes, *Mem. Amer. Math. Soc.* **232** (2014), no. 1091, vi+112.
- [48] V. Totik, Polynomial approximation in several variables, *J. Approx. Theory* **252**: 105364 (2020).
- [49] Y. G. Wang, Q. T. Le Gia, I. H. Sloan and R. S. Womersley, Fully discrete needlet approximation on the sphere. *Appl. Comput. Harmon. Anal.* **43** (2017), 292–316.
- [50] Y. Xu, Christoffel functions and Fourier series for multivariate orthogonal polynomials. *J. Approx. Theory* **82** (1995), 205–239.
- [51] Y. Xu, Summability of Fourier orthogonal series for Jacobi weight on a ball in \mathbb{R}^d , *Trans. Amer. Math. Soc.*, **351** (1999), 2439–2458.
- [52] Y. Xu, Weighted approximation of functions on the unit sphere. *Const. Approx.* **21** (2005), 1–28.
- [53] Y. Xu, Fourier series in orthogonal polynomials on a cone of revolution, *J. Fourier Anal. Appl.*, **26** (2020), Paper No. 36, 42 pp.
- [54] Y. Xu, Orthogonal structure and orthogonal series in and on a double cone or a hyperboloid. *Trans. Amer. Math. Soc.*, 374 (2021), 3603–3657.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, OR 97403–1222, USA
 Email address: yuan@uoregon.edu