

Massless finite and infinite spin representations of Poincaré group in six dimensions

I.L. Buchbinder^{1,2}, S.A. Fedoruk³, A.P. Isaev^{3,4}, M.A. Podoinitsyn³

¹*Department of Theoretical Physics, Tomsk State Pedagogical University,
634041 Tomsk, Russia,
joseph@tspu.edu.ru*

²*National Research Tomsk State University,
634050 Tomsk, Russia*

³*Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research,
141980 Dubna, Moscow Region, Russia,
fedoruk@theor.jinr.ru, isaevap@theor.jinr.ru, mpod@theor.jinr.ru*

⁴*Physics Faculty, M.V. Lomonosov Moscow State University,
119991 Moscow, Russia,*

Abstract

We study the massless irreducible representations of the Poincaré group in the six-dimensional Minkowski space. The Casimir operators are constructed and their eigenvalues are found. It is shown that the finite spin (helicity) representation is defined by two integer or half-integer numbers while the infinite spin representation is defined by the real parameter μ^2 and one integer or half-integer number.

PACS: 11.10.Kk, 11.30.Cp, 11.30.-j

Keywords: massless representations, helicity, infinite spin representations, Casimir operators

1 Introduction

Study of the various aspects of field theory in higher dimensions attracts much attention due to the remarkable and sometimes even unexpected properties at classical and quantum levels. Many of such properties are closely related to superstring theory which may be treated as a theory of infinite number of higher spin fields in higher dimensional space-time (see e.g. [1]). In particular, the low-energy limits of superstring theory are supersymmetric gauge theories and supergravity in ten dimensions that after reduction yield to field models in dimensions from ten to four. Since the details of field theories are essentially defined by the space-time symmetry, it seems useful to focus an attention on studying the diverse specific properties of symmetry groups in the higher dimensions.

The fundamental space-time background in relativistic theory is Minkowski space where the basic symmetry is described by Poincaré group. Theory of unitary irreducible representations of Poincaré group in four dimensions was constructed in the pioneer papers [2–4]. The aspects of unitary irreducible representations in higher dimensions and their applications are considered in the papers [5], [6], [7], [8], [9] and in lectures [10] (see also the recent paper [11]). Although the generic scheme of constructing the representations of the Poincaré group in any dimension seems can be realized on the base of known method of induced representations (see e.g. [12], [13]), many specific aspects important for classical and quantum field theory deserve a separate attention and require independent study. Some of such aspects are appropriate only for each concrete dimension and can not be formulated at once for all dimensions. For example, the spinor representations of the Lie algebra of multidimensional Lorenz group are defined independently for each space-time dimension. Therefore one can expect that the structure of relativistic symmetry representations in higher dimensions is much richer and more complicated than in the four-dimensional Minkowski space.

In this letter we construct the massless finite and infinite spin irreducible representations of the Lie algebra of the Poincaré group in six-dimensional Minkowski space. Some aspects of such representations are considered in papers [14], [15] however many issues, especially the infinite spin representations, were not addressed and complete analysis was not done. Recently there was the paper [16] where the unitary irreducible massless representations of the Poincaré group in five-dimensional Minkowski space were constructed and some issues related to representations in arbitrary dimensions were briefly studied and the representations of super Poincaré group were considered. The infinite spin representations were not addressed.

The letter is organized as follows. Section 2 is devoted to Casimir operators and their properties in the six-dimensional standard massless momentum reference frame. In section 3 we describe the massless finite spin irreducible representations and show that they are described by two integer or half-integer numbers. Section 4 is devoted to infinite spin representations which are described by arbitrary real parameter and a single integer or half-integer number. Section 5 is a summary of the results.

2 Poincaré algebra and light-cone reference frame

The generators P_m and $M_{mn} = -M_{nm}$ of the Lie algebra $\mathfrak{iso}(1, D-1)$ of the Poincaré group in D -dimensional space-time have the commutators

$$[P_n, P_k] = 0, \quad [M_{mn}, P_k] = i(\eta_{mk}P_n - \eta_{nk}P_m), \quad (2.1)$$

$$[M_{mn}, M_{kl}] = i(\eta_{mk}M_{nl} + \eta_{nl}M_{mk} - \eta_{ml}M_{nk} - \eta_{nk}M_{ml}), \quad (2.2)$$

where the D -vector indices run the values $m, n = 0, 1, \dots, D-1$ and we use the space-time metric $\eta^{mn} = \text{diag}(+1, \underbrace{-1, \dots, -1}_{D-1})$. We call the Lie algebra $\mathfrak{iso}(1, D-1)$ of the Poincaré group as D -dimensional Poincaré algebra.

2.1 Casimir operators of 6-dimensional Poincaré algebra

We introduce the third rank tensor W_{mnk} and the vector Υ_m as the elements of the enveloping algebra of $\mathfrak{iso}(1, 5)$ [5, 14]

$$W_{mnk} = \varepsilon_{mnklpr} P^l M^{pr}, \quad (2.3)$$

$$\Upsilon_m = \varepsilon_{mnklpr} P^n M^{kl} M^{pr}. \quad (2.4)$$

Here we use the totally antisymmetric tensor ε_{mnklpr} and normalize it as $\varepsilon_{012345} = 1$. The operators (2.3) and (2.4) satisfy the equations

$$P^m W_{mnk} = 0, \quad [P_l, W_{mnk}] = 0, \quad (2.5)$$

$$P^m \Upsilon_m = 0, \quad [P_l, \Upsilon_m] = 0. \quad (2.6)$$

By using of these equations one can check that the operators

$$C_2 := P^m P_m, \quad (2.7)$$

$$C_4 := \frac{1}{24} W^{mnk} W_{mnk}, \quad (2.8)$$

$$C_6 := \frac{1}{64} \Upsilon^m \Upsilon_m \quad (2.9)$$

are the Casimir operators of the Poincaré algebra $\mathfrak{iso}(1, 5)$. It is clear that C_2 , C_4 and C_6 are second, fourth and sixth order operators in the Poincaré algebra generators, respectively. The six-dimensional Poincaré algebra has no other additional Casimir operators (see comments in the Appendix).

Taking into account the expressions (2.3), (2.4) we obtain explicit form of the Casimir operators (2.7), (2.8), (2.9):

$$C_2 = P^m P_m, \quad (2.10)$$

$$C_4 = \Pi^m \Pi_m - \frac{1}{2} M^{mn} M_{mn} C_2, \quad (2.11)$$

$$\begin{aligned} C_6 = & -\Pi^k M_{km} \Pi_l M^{lm} + \frac{1}{2} (M^{mn} M_{mn} - 8) C_4 \\ & + \frac{1}{8} \left[M^{kl} M_{kl} (M^{mn} M_{mn} - 8) + 2 M^{mn} M_{nk} M^{kl} M_{lm} \right] C_2, \end{aligned} \quad (2.12)$$

where we introduce new vector Π with components

$$\Pi_m := P^k M_{km} = M_{km} P^k - 5i P_m, \quad (2.13)$$

which satisfy commutation relations (cf. (2.1))

$$[\Pi_n, \Pi_k] = -i M_{nk} C_2, \quad [M_{mn}, \Pi_k] = i (\eta_{mk} \Pi_n - \eta_{nk} \Pi_m). \quad (2.14)$$

Further in this paper we consider the massless unitary representations of the algebra $\mathfrak{iso}(1, 5)$ when the quadratic Casimir operator (2.10) is fixed as following:

$$C_2 \equiv P^2 = P^m P_m = 0. \quad (2.15)$$

2.2 Standard massless momentum reference frame

Let the algebra (2.1), (2.2) acts in the representation space \mathcal{H} with basis vectors $|k, \sigma\rangle$, where σ is a set of eigenvalues of all operators commuting with P_m and $P_m |k, \sigma\rangle = k_m |k, \sigma\rangle$. We take the light-cone reference frame for massless particle momentum $k^m = (k^0, k^a, k^5) = (k, 0, 0, 0, 0, k)$ in which momentum operator (2.15) has the standard form

$$P^0 = P^5 = k, \quad P^a = 0, \quad a = 1, 2, 3, 4. \quad (2.16)$$

We stress that all operator formulas presented in this Section (and written in the light-cone frame) should be understood as a result of their action on the subspace $\mathcal{H}_k \subset \mathcal{H}$ spanned by vectors $|k, \sigma\rangle$ with fixed light-cone momentum k_m .

The transition to this light-cone reference frame is conveniently performed in the light-cone basis where any 6D vector $X^m = (X^0, X^a, X^5)$ has the light-cone coordinates $X^m = (X^+, X^-, X^a)$, where

$$X^\pm = \frac{1}{\sqrt{2}} (X^0 \pm X^5), \quad X_\pm = \frac{1}{\sqrt{2}} (X_0 \pm X_5) \quad \Rightarrow \quad X^\pm = X_\mp. \quad (2.17)$$

Then, in the light-cone basis the contraction of two 6D vectors X^m and Y^m is

$$\begin{aligned} X^m Y_m &= X^+ Y_+ + X^- Y_- + X^a Y_a \\ &= \eta^{-+} X_- Y_+ + \eta^{+-} X_+ Y_- + \eta^{ab} X_b Y_a = X_- Y_+ + X_+ Y_- - X_a Y_a, \end{aligned} \quad (2.18)$$

where we use the light-cone metric $\eta^{\pm\mp} = \eta_{\pm\mp} = 1$, $\eta^{\pm\pm} = \eta_{\pm\pm} = 0$, $\eta^{ab} = \eta_{ab} = -\delta_{ab}$. In the light-cone basis the total antisymmetric tensor ε_{mnklpr} has components

$$\varepsilon_{-+abcd} = -\varepsilon_{+-abcd} = \varepsilon^{+-abcd} = -\varepsilon^{-+abcd} = \varepsilon_{abcd},$$

and we normalize the antisymmetric tensors ε_{mnklpr} and ε_{abcd} as $\varepsilon_{012345} = 1$ and $\varepsilon_{1234} = 1$.

In the light-cone basis the standard momentum (2.16) has the components

$$P^+ = P_- = \sqrt{2}k, \quad P^- = P_+ = 0, \quad P^a = 0, \quad a = 1, 2, 3, 4. \quad (2.19)$$

Thus, in the light-cone reference frame (2.19) the Casimir operators (2.11), (2.12) take the form¹

$$\hat{C}_4 = -\hat{\Pi}_a \hat{\Pi}_a, \quad (2.20)$$

$$\hat{C}_6 = \hat{\Pi}_b M_{ba} \hat{\Pi}_c M_{ca} - \frac{1}{2} M_{bc} M_{bc} \hat{\Pi}_a \hat{\Pi}_a, \quad (2.21)$$

¹When we deduce (2.20) and (2.21) it is necessary, since we project all operator relations to the subspace \mathcal{H}_k , first move all operators P_m in the expressions (2.11) and (2.12) to the right and only then perform the substitution (2.19).

where we introduce Hermitian operators

$$\hat{\Pi}_a := \sqrt{2}k M_{+a}. \quad (2.22)$$

Formula (2.20) directly follows from (2.11), while derivation of (2.21) from (2.12) takes some efforts.

In view of (2.2) the operators $\hat{\Pi}_a$ (2.22) and M_{ab} , which generate (2.20) and (2.21), form the Lie algebra of $ISO(4)$ group

$$[\hat{\Pi}_a, \hat{\Pi}_b] = 0, \quad [\hat{\Pi}_a, M_{bc}] = i \left(\delta_{ab} \hat{\Pi}_c - \delta_{ac} \hat{\Pi}_b \right), \quad (2.23)$$

$$[M_{ab}, M_{cd}] = i (\delta_{bc} M_{ad} - \delta_{bd} M_{ac} + \delta_{ac} M_{db} - \delta_{ad} M_{cb}), \quad (2.24)$$

and therefore generate the isometries of the four-dimensional Euclidean space. As a result, the operators \hat{C}_4 and \hat{C}_6 defined in (2.20) and (2.21) are the Casimir operators of the $\mathfrak{iso}(4)$ algebra.

Six generators of rotations M_{ab} in four-dimensional Euclidean space are decomposed into the sum

$$M_{ab} = M_{ab}^{(+)} + M_{ab}^{(-)}, \quad (2.25)$$

where

$$M_{ab}^{(\pm)} := \frac{1}{2} \left(M_{ab} \pm \frac{1}{2} \epsilon_{abcd} M_{cd} \right) \quad (2.26)$$

are (anti)selfdual parts. They satisfy the identities

$$M_{ab}^{(\pm)} = \pm \frac{1}{2} \epsilon_{abcd} M_{cd}^{(\pm)}. \quad (2.27)$$

The generators (2.26) form the algebra

$$[M_{ab}^{(\pm)}, M_{cd}^{(\pm)}] = i \left(\delta_{bc} M_{ad}^{(\pm)} - \delta_{bd} M_{ac}^{(\pm)} + \delta_{ac} M_{db}^{(\pm)} - \delta_{ad} M_{cb}^{(\pm)} \right), \quad [M_{ab}^{(+)}, M_{cd}^{(-)}] = 0, \quad (2.28)$$

which is direct sum of two algebras with three generators $M_{ab}^{(+)}$ and with three generators $M_{ab}^{(-)}$ respectively. Each of these algebras, containing three generators $M_{ab}^{(+)}$ or $M_{ab}^{(-)}$, is the $\mathfrak{su}(2)$ algebra.

This becomes clear (see e.g. [13]) after using the 't Hooft symbols [17]. The 't Hooft symbols $\eta_{ab}^i = -\eta_{ba}^i$, $i = 1, 2, 3$ and $\bar{\eta}_{ab}^{i'} = -\bar{\eta}_{ba}^{i'}$, $i = 1, 2, 3$ are (anti-)selfdual tensors with respect to the $SO(4)$ indices a, b :

$$\eta_{ab}^i = \frac{1}{2} \epsilon_{abcd} \eta_{cd}^i, \quad \bar{\eta}_{ab}^{i'} = -\frac{1}{2} \epsilon_{abcd} \bar{\eta}_{cd}^{i'}. \quad (2.29)$$

Below we use the following standard representations for the 't Hooft symbols

$$\eta_{ab}^i = \begin{cases} \epsilon_{iab} & a, b = 1, 2, 3, \\ \delta_{ia} & b = 4, \end{cases} \quad \bar{\eta}_{ab}^{i'} = \begin{cases} \epsilon_{i'ab} & a, b = 1, 2, 3, \\ -\delta_{i'a} & b = 4. \end{cases} \quad (2.30)$$

Due to the properties (2.29) the 't Hooft symbols connect (anti-)selfdual $SO(4)$ tensors $M_{ab}^{(\pm)}$ (2.26) with the $SO(3)$ vectors $M_i^{(+)}$, $M_{i'}^{(-)}$ by means of the following relations

$$M_{ab}^{(+)} = -\eta_{ab}^i M_i^{(+)}, \quad M_{ab}^{(-)} = -\bar{\eta}_{ab}^{i'} M_{i'}^{(-)}. \quad (2.31)$$

Such defined operators $M_i^{(+)}$ and $M_{i'}^{(-)}$ form two $\mathfrak{su}(2)$ algebras with standard form of the commutators

$$[M_i^{(+)}, M_j^{(+)}] = i\epsilon_{ijk}M_k^{(+)}, \quad [M_{i'}^{(-)}, M_{j'}^{(-)}] = i\epsilon_{i'j'k'}M_{k'}^{(-)}, \quad [M_i^{(+)}, M_{j'}^{(-)}] = 0. \quad (2.32)$$

In term of the operators (2.31) the Casimir (2.21) takes the form (we use the equalities $\eta_{ab}^i\eta_{ab}^j = 4\delta^{ij}$, $\bar{\eta}_{ab}^{i'}\bar{\eta}_{ab}^{j'} = 4\delta^{i'j'}$ and $\eta_{ab}^i\bar{\eta}_{ab}^{j'} = 0$)

$$\hat{C}_6 = 2M_i^{(+)}M_{j'}^{(-)}\eta_{ab}^i\bar{\eta}_{ac}^{j'}\hat{\Pi}_b\hat{\Pi}_c - \left(M_i^{(+)}M_i^{(+)} + M_{i'}^{(-)}M_{i'}^{(-)}\right)\hat{\Pi}_a\hat{\Pi}_a. \quad (2.33)$$

Thus, in massless case (2.15) unitary irreducible representations are defined by the eigenvalues of the $\mathfrak{iso}(4)$ Casimir operators (2.20) and (2.21). In case of this noncompact symmetry there are two different cases defined the value of Casimir operator (2.20), i.e. square of “four-translation” generator $\hat{\Pi}_a$. So, in next sections we consider following unitary massless representations:

- **Finite spin (helicity) representations.**

In these cases the $SO(4)$ four-vector $\hat{\Pi}_a$ has zero norm:

$$\hat{\Pi}_a\hat{\Pi}_a = 0. \quad (2.34)$$

- **Infinite (continuous) spin representations.**

In case of these representations the Euclidean four-vector $\hat{\Pi}_a$ has nonzero norm:

$$\hat{\Pi}_a\hat{\Pi}_a = \mu^2 \neq 0. \quad (2.35)$$

In next sections we consider these massless representations in details.

3 Massless finite spin representations

This case is characterized by the fulfillment of condition (2.34), which implies that all components $\hat{\Pi}_a$ (since they are Hermitian operators) of the Euclidean vector are zero:

$$\hat{\Pi}_a = 0 \quad \text{at all} \quad a = 1, 2, 3, 4. \quad (3.1)$$

As result, the Casimir operators (2.20) and (2.21) are vanish in this case: $\hat{C}_4 = 0$ and $\hat{C}_6 = 0$. In passing from this light-cone reference frame to an arbitrary frame, we get that all Casimir operators (2.11), (2.12) on the massless finite spin states take zero values (see also [14]):

$$C_4 = 0, \quad C_6 = 0, \quad (3.2)$$

and in view of (2.15) we have $\Pi^k\Pi_k = 0$ and $\Pi^k M_{km}\Pi_\ell M^{\ell m} = 0$.

Due to (3.1) the Euclidean four-translations are realized trivially in case of these representations. As a result such representations of $ISO(4)$ are finite dimensional. Each such representation defines some $6D$ standard massless representation. As we saw above, such representations are induced from irreducible $SO(4)$ representations. Let us show below that the Casimir operators of the stability subgroup $SO(4)$ define the $6D$ helicity operators.

3.1 6D helicity operators

First, let us consider the vector Υ_m defined in (2.4). In the case $C_6 = 0$, according to (2.9), we have $\Upsilon_m \Upsilon^m = 0$ and, in the light-cone reference frame (2.16), (2.19), the components of 6D vector Υ are

$$\Upsilon^+ = \Lambda_1 P^+, \quad \Upsilon^- = \Upsilon_a = 0, \quad (3.3)$$

where we have

$$\Lambda_1 := \epsilon_{abcd} M_{ab} M_{cd}. \quad (3.4)$$

This operator is the Casimir operator of the $\mathfrak{so}(4)$ algebra.

The conditions (3.3) demonstrate that vectors Υ and P are collinear in the light-cone reference frame and this property is conserved in any reference frame. Namely, the relations (2.6) show that the light like vector Υ is transverse to the vector P and its components Υ_m commute with P_k . Therefore, the vector Υ_m is proportional to the vector P_m :

$$\Upsilon_m = \Lambda_1 P_m. \quad (3.5)$$

This relation was also pointed out in [14, 16]. Note that the operator (3.4) can be represented in the form

$$\Lambda_1 := \frac{\Upsilon_0}{P_0}. \quad (3.6)$$

This expression appears for the 4D helicity operator when Υ_m is replaced by W_m . Due to the relations

$$[M_{0i}, \Lambda_1] = \frac{i}{P_0} (\Upsilon_i - \Lambda_1 P_i) = 0, \quad [M_{ik}, \Lambda_1] = 0 = [P_k, \Lambda_1], \quad (i, k = 1, \dots, 5), \quad (3.7)$$

we conclude that the operator (3.6) is invariant with respect to the 6D Poincare symmetry. Therefore, the operator Λ_1 , defined in (3.6), is a 6D analog of the helicity operator and it coincides with one of $\mathfrak{so}(4)$ Casimir operators in the light-cone reference frame.

We note that irreducible $\mathfrak{so}(4)$ representations are characterized by two quadratic Casimir operators. The second Casimir operator arises as helicity operator if we use the construction proposed in [16]. Indeed, by using the prescription of [16], one can construct another (third order in generators of $\mathfrak{iso}(1, 5)$) vector with components²

$$S_m := 3M^{nk} P_{[m} M_{nk]} = M^{nk} M_{nk} P_m - 2M^{kn} M_{mn} P_k. \quad (3.8)$$

The square of this 6D vector is

$$S^m S_m = M^{nm} M_{nm} M^{lk} M_{lk} P^2 + 4 \left[\Pi^k M_{km} \Pi_l M^{lm} - M^{nm} M_{nm} (\Pi^l \Pi_l + P^2) + \Pi^l \Pi_l \right], \quad (3.9)$$

while its contraction with 6D vector momentum P_m gives

$$P^m S_m = M^{mn} M_{mn} P^2 - 2\Pi^m \Pi_m \equiv -2C_4, \quad (3.10)$$

²In the definition of antisymmetrization of n indices, we use the factor $n!$, i.e.

$$A_{[m} B_n C_k] = \frac{1}{3!} (A_m B_n C_k - A_m B_k C_n + \text{cyclic permutations}).$$

and the commutators of S_m and P_n are

$$[S_m, P_n] = 2iM_{mn}P^2 + 4i\Pi_{[m}P_{n]} . \quad (3.11)$$

For the massless finite spin representations, defined by the conditions (2.15), (3.1) and (3.2), equations (3.9), (3.10) and (3.11) are reduced to

$$S^m S_m = 0, \quad P^m S_m = 0, \quad [S_m, P_n] = 0, \quad (3.12)$$

which are the same as conditions (2.6) for light-like vectors Υ and P . So, in the case of massless finite spin representations, the vectors P_m and S_m are also proportional to each other. One can check this in the light-cone reference frame, when subject to the conditions (2.19) and (3.1) the components of the $6D$ vector (3.8) are equal to

$$S^+ = \Lambda_2 P^+, \quad S^- = S_a = 0, \quad (3.13)$$

where the operator

$$\Lambda_2 := M_{ab}M_{ab} \quad (3.14)$$

is second $\mathfrak{so}(4)$ Casimir operator.

Due to the relations (3.12) in general frame the relations (3.13) take the form

$$S_m = \Lambda_2 P_m, \quad (3.15)$$

where the operator Λ_2 (3.14) defines second helicity operator and has equivalent ‘‘covariant’’ form

$$\Lambda_2 := \frac{S_0}{P_0}. \quad (3.16)$$

So these massless representations of finite spin are characterized by the pair (λ_1, λ_2) , where real numbers $\lambda_{1,2}$ define the eigenvalue of the Casimir operators $\Lambda_{1,2}$ presented in (3.6) and (3.16), respectively.

Using (2.25) and (2.31) we represent helicity operators (3.4) and (3.14) in the form

$$\Lambda_1 = 2 \left(M_{ab}^{(+)} M_{ab}^{(+)} - M_{ab}^{(-)} M_{ab}^{(-)} \right) = 8 \left(M_i^{(+)} M_i^{(+)} - M_{i'}^{(-)} M_{i'}^{(-)} \right), \quad (3.17)$$

$$\Lambda_2 = M_{ab}^{(+)} M_{ab}^{(+)} + M_{ab}^{(-)} M_{ab}^{(-)} = 4 \left(M_i^{(+)} M_i^{(+)} + M_{i'}^{(-)} M_{i'}^{(-)} \right). \quad (3.18)$$

In case of unitary representations, the operators $M_i^{(+)} M_i^{(+)}$ and $M_{i'}^{(-)} M_{i'}^{(-)}$ equal $j_+(j_+ + 1)$ and $j_-(j_- + 1)$ respectively. Therefore, the eigenvalues of the helicity operators (3.6) and (3.16) take the values

$$\lambda_1 = 8j_+(j_+ + 1) - 8j_-(j_- + 1), \quad (3.19)$$

$$\lambda_2 = 4j_+(j_+ + 1) + 4j_-(j_- + 1), \quad (3.20)$$

where j_{\pm} are integer or half-integer numbers in case of the unitary representations.

We note that the standard $4D$ helicity operator is invariant under proper $SO(1, 3)$ rotations but changes its sign under improper $O(1, 3)$ rotations (reflections). We have the same property for Λ_1 but it is not the case for Λ_2 .

3.2 Examples

Here we will demonstrate the use of the obtained formulas for determining the helicities on the examples of some massless finite spin fields. To clarity and avoid technical complications, we will consider only bosonic integer-spin fields.

Since the irreducible massless representations of the $6D$ Poincaré group are induced by the irreducible $SO(4)$ representations in the light-cone reference frame, we will use the following procedure.

Below, in all examples of this section, we first consider a fixed irreducible $SO(4)$ representation and determine the values of the helicities. Here we will use the defining representation for the $\mathfrak{so}(4)$ generators

$$(\mathcal{M}_{ab})_{eg} = i(\delta_{ae}\delta_{bg} - \delta_{ag}\delta_{be}). \quad (3.21)$$

Then we reconstruct the corresponding $6D$ field, for which the equations of motion and gauge fixing show that the independent components are exactly those $SO(4)$ fields which were considered earlier in the Euclidean four-dimensional picture.

3.2.1 Vector field

Let us consider the $SO(4)$ vector field A_a . In this case the $\mathfrak{so}(4)$ generators coincide with (3.21):

$$(M_{ab})_{eg} = (\mathcal{M}_{ab})_{eg}. \quad (3.22)$$

Then, the $SO(4)$ Casimir operators take the form

$$\begin{aligned} (\Lambda_1)_{eg} &= \epsilon_{abcd}(M_{ab}M_{cd})_{eg} = 0, \\ (\Lambda_2)_{eg} &= (M_{ab}M_{ab})_{eg} = 6\delta_{eg}. \end{aligned} \quad (3.23)$$

When acting on the $SO(4)$ vector field A_a , the operators (3.23) give the following values of helicities:

$$\lambda_1 = 0, \quad \lambda_2 = 6; \quad j_+ = j_- = \frac{1}{2}. \quad (3.24)$$

This Euclidean vector field A_a describes physical components of the $6D$ vector gauge field A_m . In the momentum representation the $U(1)$ massless gauge field A_m is described by the equations of motion

$$P^m F_{mn} = 0, \quad (3.25)$$

where $F_{mn} = i(P_m A_n - P_n A_m)$ is the field strength, and determined up to gauge transformations

$$\delta A_m = i P_m \varphi. \quad (3.26)$$

One of the possible gauge fixing for transformations (3.26) is the light-cone gauge (see e.g. [18])

$$A^+ = 0. \quad (3.27)$$

Then in the light-cone frame (2.19), the equations of motion (3.25) give $A^- = 0$ and independent field is given by the transverse part A_a of the $6D$ gauge field A_m .

3.2.2 Second rank symmetric tensor field

Now we consider the $SO(4)$ second rank tensors. In this case the $\mathfrak{so}(4)$ generators take the matrix form

$$(M_{ab})_{e_1 e_2, g_1 g_2} = ((\mathcal{M}_{ab})_1 + (\mathcal{M}_{ab})_2)_{e_1 e_2, g_1 g_2} = (\mathcal{M}_{ab})_{e_1 g_1} \delta_{e_2 g_2} + \delta_{e_1 g_1} (\mathcal{M}_{ab})_{e_2 g_2} \quad (3.28)$$

and the $SO(4)$ Casimir operators are

$$\begin{aligned} (\Lambda_1)_{e_1 e_2, g_1 g_2} &= \epsilon_{abcd} (M_{ab} M_{cd})_{e_1 e_2, g_1 g_2} = 2 \epsilon_{abcd} ((M_{ab})_1 (M_{cd})_2)_{e_1 e_2, g_1 g_2} = 8 \epsilon_{e_1 e_2 g_1 g_2} , \\ (\Lambda_2)_{e_1 e_2, g_1 g_2} &= (M_{ab} M_{ab})_{e_1 e_2, g_1 g_2} = ((M_{ab}^2)_1 + (M_{ab}^2)_2 + 2(M_{ab})_1 (M_{ab})_2)_{e_1 e_2, g_1 g_2} = \\ &= 12 \delta_{e_1 g_1} \delta_{e_2 g_2} + 4(\delta_{e_1 g_2} \delta_{e_2 g_1} - \delta_{e_1 e_2} \delta_{g_1 g_2}) . \end{aligned} \quad (3.29)$$

First, we consider the $SO(4)$ second rank tensor \hat{h}_{ab} , which is symmetric $\hat{h}_{ab} = \hat{h}_{ba}$ and traceless \hat{h}_{aa} . On this field the helicity operators (3.29) take the values

$$\lambda_1 = 0, \quad \lambda_2 = 16; \quad j_+ = j_- = 1. \quad (3.30)$$

Let us show that this field \hat{h}_{ab} describes the physical components of the $6D$ linearized gravitational field.

The $6D$ linearized gravitational field $h^{mn} = h^{nm}$ is determined by the well known equations of motion

$$P^2 h^{mn} - P^m P_k h^{nk} - P^n P_k h^{mk} + P^m P^n h_k{}^k = 0, \quad (3.31)$$

and has gauge invariance

$$\delta h^{mn} = i P^{(m} \varphi^{n)}. \quad (3.32)$$

For the transformations (3.32) we can put again the light-cone gauge (see also [18])

$$h^{+m} = 0. \quad (3.33)$$

The equations of motion (3.31) produce $h^{-m} = 0$, $h_a{}^a = 0$ in the light-cone frame (2.19). As a result, nonvanishing physical components of the $6D$ gravity field h_{mn} are given by the traceless part \hat{h}_{ab} of its transverse components h_{ab} .

3.2.3 Third rank (anti-)selfdual antisymmetric tensor fields

Now we consider the $SO(4)$ antisymmetric tensors of the second rank $B_{ab}^{(\pm)} = -B_{ba}^{(\pm)}$, which are (anti-)selfdual

$$B_{ab}^{(\pm)} = \pm \frac{1}{2} \epsilon_{abcd} B_{cd}^{(\pm)}. \quad (3.34)$$

These tensors form the spaces of two $SO(4)$ irreducible representations which make up the $SO(4)$ reducible representation in the space of all antisymmetric rank 2 tensors associated to Young diagram $[1^2] \equiv \begin{array}{|c|} \hline \square \\ \hline \end{array}$. In this case the $\mathfrak{so}(4)$ generators M_{ab} and helicity operators Λ_1, Λ_2 have the same expressions (3.28) and (3.29). Then the eigenvalues of the operators Λ_1, Λ_2 and $(M_i^{(\pm)} M_i^{(\pm)})$ are given by numbers

$$\lambda_1 = 16, \quad \lambda_2 = 8; \quad j_+ = 1, \quad j_- = 0 \quad (3.35)$$

on the space of the selfdual fields $B_{ab}^{(+)}$, and by

$$\lambda_1 = -16, \quad \lambda_2 = 8; \quad j_+ = 0, \quad j_- = 1 \quad (3.36)$$

on the space of the anti-selfdual fields $B_{ab}^{(-)}$. It is clear that these $SO(4)$ (anti-)selfdual fields $B_{[ab]}^{(\pm)}$ are independent components of the $6D$ massless (anti-)selfdual 3-rank fields $B_{mnk}^{(\pm)}$ which satisfy the identities

$$B_{mnk}^{(\pm)} = \pm \frac{1}{3!} \varepsilon_{mnklpr} B^{(\pm)lpr}. \quad (3.37)$$

So, the equations of motion of the $6D$ massless fields $B_{mnk}^{(\pm)}$ are

$$\text{a) } P^m B_{mnk}^{(\pm)} = 0, \quad \text{b) } P_{[m} B_{nkl]}^{(\pm)} = 0, \quad \text{c) } P^2 B_{nkl}^{(\pm)} = 0. \quad (3.38)$$

Then in the light-cone frame (2.19) the equations (3.38a) give $B^{(\pm)-mn} = 0$ whereas the equations (3.38b) produce $B^{(\pm)abc} = 0$. As a result, independent fields of the $6D$ tensors $B_{mnk}^{(\pm)}$ are the $SO(4)$ (anti-)selfdual fields $B^{(\pm)+ab} \equiv B^{(\pm)ab}$ which are subjected the $SO(4)$ (anti-)selfdual conditions (3.34) due to the $6D$ (anti-)selfdual conditions (3.37).

Remark. One can generalize this example to the case of special $3n$ -rank selfdual and anti-selfdual 6-dimensional tensor fields. These fields correspond to $SO(4)$ irreducible representations in spaces of $2n$ -rank traceless selfdual and anti-selfdual tensors with components $B_{a_1 \dots a_{2n}}^{(\pm)}$

be symmetrized in accordance to the Young diagram $[n^2] \equiv \begin{array}{|c|c|c|c|} \hline & & \cdots & \\ \hline & & \cdots & \\ \hline \end{array}$. It is clear that for highest weights of such selfdual and anti-selfdual representations of $SO(4)$ we have respectively $j_+ = n, j_- = 0$ and $j_+ = 0, j_- = n$ and in view of (3.19) and (3.20) we obtain the eigenvalues of helicity operators $\lambda_1 = 8n(n+1)$, $\lambda_2 = 4n(n+1)$ and $\lambda_1 = -8n(n+1)$, $\lambda_2 = 4n(n+1)$ which is a generalization of (3.35) and (3.36).

4 Massless infinite (continuous) spin representations

In this case, when the condition (2.35) is satisfied and the Euclidean four-vector $\hat{\Pi}_a$ is nonzero. Then here the representations of the $ISO(4)$ group, which induce the $6D$ relativistic massless representations, are infinite dimensional.

In case of these representations the Casimir operator (2.20) has nonvanishing eigenvalue

$$C_4 = \hat{C}_4 = -\mu^2, \quad \mu \neq 0. \quad (4.1)$$

Moreover, for the orbits (2.35) we can take the basis with with nonzero only the fourth component:

$$\hat{\Pi}_1 = \hat{\Pi}_2 = \hat{\Pi}_3 = 0, \quad \hat{\Pi}_4 = \mu. \quad (4.2)$$

Then taking into account $\eta_{a4}^i = \delta_{ia}$ and $\bar{\eta}_{a4}^i = -\delta_{i'a}$ (see (2.30)) we obtain from (2.33) the value of the Casimir operator (2.21):

$$\hat{C}_6 = -\mu^2 J_i J_i, \quad (4.3)$$

where

$$J_i := M_i^{(+)} + M_i^{(-)} \quad (4.4)$$

are the generators of the diagonal $su(2)$ subalgebra of the $so(4) = su(2) \oplus su(2)$ stability algebra. Using (2.26) and (2.31) and explicit expressions of the 't Hooft symbols (see e.g. Sect. 3.3.3 in [13]) we find

$$J_i = -\frac{1}{2} \epsilon_{ijk} M_{jk}, \quad i = 1, 2, 3. \quad (4.5)$$

So the operators (4.4) are in fact the generators of the $SO(3)$ subgroup of the $SO(4)$ stability group. Therefore, in case of the unitary representations it is necessary to satisfy the equality

$$J^2 = s(s+1), \quad (4.6)$$

where s is fixed integer or half-integer number.

So, in case of the irreducible representations of infinite (continuous) spin, the Casimir operator (2.12) takes the value

$$C_6 = \hat{C}_6 = -\mu^2 s(s+1), \quad (4.7)$$

Such irreducible representations describe a tower of infinite number of massless states.

As a result, the massless infinite spin representations are characterized by the pair (μ, s) , where the real parameter μ defines the eigenvalue of the Casimir operator (4.1) and the (half-)integer number s defines the eigenvalue of the Casimir operator (4.7).

Let us examine in our consideration the $D = 6$ infinite integer spin system [19] which is higher dimension generalization of the $D = 4$ model [2], [3], [4]. This model [19] is described by the pair of the space-time phase operators

$$x^m, p_m, \quad [x^m, p_k] = i\delta_k^m \quad (4.8)$$

and two pairs of the additional bosonic phase vectors

$$w^m, \xi_m, \quad [w^m, \xi_k] = i\delta_k^m; \quad u^m, \zeta_m, \quad [u^m, \zeta_k] = i\delta_k^m. \quad (4.9)$$

These two pairs of vectors (4.9) are responsible for spinning degrees of freedom.

Infinite integer spin field Ψ in [19] is described by the $D = 6$ generalization of the Wigner-Bargmann equations

$$p^2 \Psi = 0, \quad (4.10)$$

$$\xi \cdot p \Psi = 0, \quad (4.11)$$

$$(w \cdot p - \mu) \Psi = 0, \quad (4.12)$$

$$(\xi \cdot \xi + 1) \Psi = 0, \quad (4.13)$$

and additional equations with vectorial operators from the second pair (4.9)

$$u \cdot p \Psi = 0, \quad (4.14)$$

$$\zeta \cdot p \Psi = 0, \quad (4.15)$$

$$\zeta \cdot \xi \Psi = 0, \quad (4.16)$$

$$\zeta \cdot \zeta \Psi = 0, \quad (4.17)$$

$$(u \cdot \zeta - s) \Psi = 0, \quad (4.18)$$

where $\xi \cdot p := \xi^m p_m$, *etc.*

Note that, in contrast to the four-dimensional case [2], [3], [4] with one pair of auxiliary variables w^m, ξ_m ,³ in the six-dimensional case it is necessary to use the second pair of auxiliary vector variables u^m, ζ_m to describe arbitrary infinite spin representations.

In the light-cone frame (2.19), *i.e.* $p^- = p_a = 0$, $p^+ = \text{const} \neq 0$, and in the representation $\xi_m = -i\partial/\partial w^m$, $\zeta_m = -i\partial/\partial u^m$ the equations (4.11)-(4.13) give the conditions

$$\frac{\partial}{\partial w^+} \Psi = 0, \quad (4.19)$$

$$(p^+ w^- - \mu) \Psi = 0, \quad (4.20)$$

$$\left(\frac{\partial}{\partial w_a} \frac{\partial}{\partial w_a} + 1 \right) \Psi = 0, \quad (4.21)$$

whereas (4.14)-(4.18) yield

$$p^+ u^- \Psi = 0, \quad (4.22)$$

$$\frac{\partial}{\partial u^+} \Psi = 0, \quad (4.23)$$

$$\frac{\partial}{\partial u_a} \frac{\partial}{\partial w_a} \Psi = 0, \quad (4.24)$$

$$\frac{\partial}{\partial u_a} \frac{\partial}{\partial u_a} \Psi = 0, \quad (4.25)$$

$$\left(u_a \frac{\partial}{\partial u_a} - s \right) \Psi = 0, \quad (4.26)$$

The solution of the equations (4.19)-(4.26) is the field

$$\Psi = \delta(p^+ w^- - \mu) \delta(p^+ u^-) \Phi(w_a, u_a), \quad (4.27)$$

where $\Phi(w_a, u_a)$ is subjected (4.21), (4.24)-(4.26) and has series expansions presented in [19].

Now we can determine the values of the Casimir operators (2.20), (2.21) on the field (4.27).

For the field (4.27) the generators of the *iso*(4) algebra (2.23), (2.24) have the form

$$M_{ab} = i \left(w_a \frac{\partial}{\partial w_b} - w_b \frac{\partial}{\partial w_a} + u_a \frac{\partial}{\partial u_b} - u_b \frac{\partial}{\partial u_a} \right), \quad \hat{\Pi}_a = -i\mu \frac{\partial}{\partial w_a}. \quad (4.28)$$

As result, due to the equation (4.21), we obtain the fulfillment of the condition (4.1) for the Casimir operator C_4 : $C_4 = \hat{C}_4 = -\mu^2$. Moreover, the representations (4.28) lead to the expression

$$\begin{aligned} \hat{C}_6 &= \mu^2 u_a \frac{\partial}{\partial u_a} \left(u_b \frac{\partial}{\partial u_b} + 1 \right) \frac{\partial}{\partial w_c} \frac{\partial}{\partial w_c} \\ &+ \mu^2 \left(u_a \frac{\partial}{\partial w_a} u_b \frac{\partial}{\partial w_b} - u_a u_a \frac{\partial}{\partial w_b} \frac{\partial}{\partial w_b} \right) \frac{\partial}{\partial u_c} \frac{\partial}{\partial u_c} \\ &+ \mu^2 \left(u_a u_a \frac{\partial}{\partial u_b} \frac{\partial}{\partial w_b} - 2u_a \frac{\partial}{\partial u_a} u_b \frac{\partial}{\partial w_b} \right) \frac{\partial}{\partial u_c} \frac{\partial}{\partial w_c} \end{aligned} \quad (4.29)$$

³Note that in the twistor formulation of the infinite spin particle [20], it was more convenient for us to use dimensional additional variables $y^m = w^m/\mu$, $q_m = \mu\xi_m$.

for the sixth order Casimir operator. So, due to the equations (4.21), (4.24)-(4.26) the operator (2.21) takes the value $C_6 = \hat{C}_6 = -\mu^2 s(s+1)$ on the field (4.27).

Thus, the infinite spin field with only one additional vector variables and obeying the Wigner-Bargmann equations (4.10)-(4.13) and additional equations (4.19)-(4.26) describes the irreducible (μ, s) infinite spin representation. The system with only one pair of auxiliary variables w^m, ξ_m in (4.9) (without using the second pair of auxiliary vector variables u^m, ζ_m) and with only the equations of motion (4.11)-(4.13) describe the infinite spin representations at $s = 0$ [19].

5 Summary and outlook

We have studied the massless irreducible representations of the Poincaré group in six-dimensional Minkowski space. The representations are described by three Casimir operators written in the form (2.7), (2.8), (2.9) or in the equivalent form (2.10), (2.11), (2.12). The properties of these operators are explored in the standard massless momentum reference frame, where it is seen that the unitary representations of $ISO(1, 5)$ group are induced from representations of $SO(4)$ and $ISO(4)$ groups and correspondingly are divided into finite spin (helicity) and infinite spin representations. Both these representations are studied in details. It is proved that the finite spin representation is described by two integer or half-integer numbers while the infinite spin representation is described by one real parameter and one integer or half-integer number. In case of half-integer spin we should introduce an additional spinor or twistor variables like in [19]. The results obtained here are in agreement with the statements of [5,10] where it was done the classification of massless representations of any dimension Poincaré group via the induced representation method.

As a continuation of this research it would be interesting to describe the massless representations with half-integer spin and massive irreducible representations of six-dimensional Poincaré group with both integer and half-integer spin. Another open problem is constructing the representations of the corresponding six-dimensional *super* Poincaré group. Also it would be useful to work out the field realizations of the massless representations considered in this paper (see, e.g., Remark at the end of Sect.3.2.3) and explore the new aspects of Lagrange formulation for these fields in six-dimensional Minkowski space including infinite spin cases.⁴ We plan to study all these problems in the forthcoming papers.

Acknowledgments

We are grateful to N. Boulanger, S.M. Kuzenko and Yu.M. Zinoviev for correspondence. Work of I.L.B, S.A.F and A.P.I was supported in part by The Ministry of Education of Russian Federation, project FEWF-2020-0003. I.L.B is grateful to RFBR grant, project No 18-02-00153. M.A.P is grateful to RFBR grant, project No 19-01-00726A.

⁴Currently, there is a fairly large literature on various aspects of infinite spin Lagrange formulation (see e.g., the recent paper [21] and the references therein).

Appendix. On the Casimir operators of the Poincaré algebra

The quantity $\varepsilon^{mnklpr}W_{mnk}W_{lpr}$ could be an additional Casimir operator for $\mathfrak{iso}(1, 5)$ algebra. But it is identically equal to zero. This fact is a special case of the property of any rank r antisymmetric tensor $W_{m_1\dots m_r}$ in $2r$ -dimensional space, when r is odd number. Indeed, in this case we have $(W, V)_\varepsilon = (-1)^r(V, W)_\varepsilon$, where $(W, V)_\varepsilon := \varepsilon^{m_1\dots m_r n_1\dots n_r}W_{m_1\dots m_r}V_{n_1\dots n_r}$ and $\varepsilon^{m_1\dots m_r n_1\dots n_r}[W_{m_1\dots m_r}, V_{n_1\dots n_r}] = 0$. Thus, for antisymmetric tensor with components

$$W_{m_1\dots m_r} = \varepsilon_{m_1\dots m_r n_1\dots n_r} P^{n_1} M^{n_2 n_3} \dots M^{n_{r-1} n_r},$$

which is defined only for odd r , we always have $(W, W)_\varepsilon = 0$. In this case a Casimir operator for $\mathfrak{iso}(1, 2r - 1)$ algebra, of the second order in W , has the unique form

$$W^2 = \frac{1}{(r+1)!} W^{m_1\dots m_r} W_{m_1\dots m_r}.$$

Whereas for even r we have antisymmetric tensor with components

$$L_{m_1\dots m_r} = \varepsilon_{m_1\dots m_r n_1\dots n_r} M^{n_1 n_2} \dots M^{n_{r-1} n_r}$$

which yields for $\mathfrak{so}(\ell, 2r - \ell)$ algebra additional to $L^2 = L^{m_1\dots m_r} L_{m_1\dots m_r}$ Casimir operator $(L, L)_\varepsilon \neq 0$ (see operator (3.4) written for the case of $\mathfrak{so}(4)$ algebra).

References

- [1] M.B. Green, J.H. Schwarz, E. Witten, *Superstring theory*, Cambridge Univ. Press, 1987.
- [2] E.P. Wigner, *On unitary representations of the inhomogeneous Lorentz group*, *Annals Math.* **40** (1939) 149.
- [3] E.P. Wigner, *Relativistische Wellengleichungen*, *Z. Physik* **124** (1947) 665.
- [4] V. Bargmann, E.P. Wigner, *Group theoretical discussion of relativistic wave equations*, *Proc. Nat. Acad. Sci. US* **34** (1948) 211.
- [5] L. Brink, A.M. Khan, P. Ramond, X.-Z. Xiong, *Continuous spin representations of the Poincare and superPoincare groups*, *J. Math. Phys.* **43** (2002) 6279, [arXiv:hep-th/0205145](#).
- [6] A.M. Khan, P. Ramond, *Continuous spin representations from group contraction*, *J. Math. Phys.* **46** (2005) 053515, [arXiv:hep-th/0410107](#).
- [7] X. Bekaert, N. Boulanger, *On geometric equations and duality for free higher spins*, *Phys. Lett.* **B561** (2003) 183, [arXiv:hep-th/0301243](#).
- [8] X. Bekaert, N. Boulanger, *Tensor gauge fields in arbitrary representations of $GL(D, R)$* , *Commun. Math. Phys.* **271** (2007) 723, [arXiv:hep-th/0606198](#).
- [9] I. Bandos, X. Bekaert, J.A. de Azcarraga, D. Sorokin, M. Tsulaia, *Dynamics of higher spin fields and tensorial space*, *JHEP* **0505** (2005) 031, [hep-th/0501113](#).

- [10] X. Bekaert, N. Boulanger, *The unitary representations of the Poincaré group in any space-time dimension*, Lectures presented at 2nd Modave Summer School in Theoretical Physics, 6-12 Aug 2006, Modave, Belgium, [arXiv:hep-th/0611263](#).
- [11] S. Weinberg, *Massless Particles in Higher Dimensions*, Phys. Rev. **D102** (2020) 095022, [arXiv:2010.05823 \[hep-th\]](#).
- [12] A.O. Barut, R. Raczka, *Theory of Group Representations and Applications*, Polish Scientific Publishing, 1977.
- [13] A.P. Isaev, V.A. Rubakov, *Theory Of Groups And Symmetries (I): Finite Groups, Lie Groups, And Lie Algebras*. World Scientific, 2019.
- [14] L. Mezincescu, A.J. Routh, P.K. Townsend, *Supertwistors and massive particles*, Annals Phys. **346** (2014) 66, [arXiv:1312.2768 \[hep-th\]](#).
- [15] A.S. Arvanitakis, L. Mezincescu, P.K. Townsend, *Pauli-Lubanski, supertwistors, and the super-spinning particle*, JHEP **1706** (2017) 151, [arXiv:1601.05294 \[hep-th\]](#).
- [16] S.M. Kuzenko, A.E. Pindur, *Massless particles in five and higher dimensions*, Phys. Lett. **B812** (2021) 136020, [arXiv:2010.07124 \[hep-th\]](#).
- [17] G. 't Hooft, *Computation of the quantum effects due to a four-dimensional pseudoparticle*, Phys. Rev. **D14** (1976) 3432.
- [18] W. Siegel, *Fields*, [arXiv:hep-th/9912205](#).
- [19] X. Bekaert, J. Mourad, *The continuous spin limit of higher spin field equations*, JHEP **0601** (2006) 115, [arXiv:hep-th/0509092](#).
- [20] I.L. Buchbinder, S. Fedoruk, A.P. Isaev, *Twistorial and space-time descriptions of massless infinite spin (super)particles and fields*, Nucl. Phys. **B945** (2019) 114660, [arXiv:1903.07947 \[hep-th\]](#).
- [21] M.V. Khabarov, Yu.M. Zinoviev, *Infinite (continuous) spin fields in the frame-like formalism*, Nucl. Phys. **B928** (2018) 182, [arXiv:1711.08223 \[hep-th\]](#).