

NEGATIVE CLUSTER CATEGORIES FROM SIMPLE MINDED COLLECTION QUADRUPLES

FRANCESCA FEDELE

ABSTRACT. Fomin and Zelevinsky’s definition of cluster algebras laid the foundation for cluster theory. The various categorifications and generalisations of the original definition led to Iyama and Yoshino’s generalised cluster categories $\mathcal{T}/\mathcal{T}^{fd}$ coming from positive-Calabi-Yau triples $(\mathcal{T}, \mathcal{T}^{fd}, \mathcal{M})$. Jin later defined simple minded collection quadruples $(\mathcal{T}, \mathcal{T}^p, \mathbb{S}, \mathcal{S})$, where the special case $\mathbb{S} = \Sigma^{-d}$ is the analogue of Iyama and Yang’s triples: negative-Calabi-Yau triples.

In this paper, we further study the quotient categories $\mathcal{T}/\mathcal{T}^p$ coming from simple minded collection quadruples. Our main result uses limits and colimits to describe Hom-spaces over $\mathcal{T}/\mathcal{T}^p$ in relation to the easier to understand Hom-spaces over \mathcal{T} . Moreover, we apply our theorem to give a different proof of a result by Jin: if we have a negative-Calabi-Yau triple, then $\mathcal{T}/\mathcal{T}^p$ is a negative cluster category.

1. INTRODUCTION

Cluster theory developed starting from the definition of cluster algebras by Fomin and Zelevinsky, see [11, Definition 2.3]. Buan, Marsh, Reineke, Reiten, and Todorov later categorified the original definition by introducing cluster categories in [5, Section 1]. Further categorifications have been widely studied in the past few years, see for example [12], [2], [13]. This led to two parallel generalisations of cluster categories coming from positive and negative Calabi-Yau triples respectively.

First, Iyama and Yang introduced generalised cluster categories, see [16], which instead of coming from dg algebras, come from d -Calabi-Yau triples of the form $(\mathcal{T}, \mathcal{T}^{fd}, \mathcal{M})$ for some positive integer. The idea behind such a triple is that, starting from a triangulated category \mathcal{T} with some extra assumptions and a triangulated subcategory \mathcal{T}^{fd} , we obtain a triangulated quotient category $\mathcal{T}/\mathcal{T}^{fd}$ which is a generalised cluster category: it is Hom-finite, $(d-1)$ -Calabi-Yau and it has a $(d-1)$ -cluster tilting object. This was first proven by Iyama and Yang in [16, Section 5] and we presented an alternative proof using more classic means such as limits, colimits and a Gap Theorem in [10].

Note that in this setup $(\mathcal{T}, \mathcal{T}^{fd})$ is relative d -Calabi-Yau in the sense that for every $X \in \mathcal{T}^{fd}$ and $Y \in \mathcal{T}$ there exists a bifunctorial isomorphism of the form

$$D\mathcal{T}(X, Y) \cong \mathcal{T}(Y, \Sigma^d X). \quad (\star)$$

An interesting question is what happens if we substitute Σ^d in (\star) with a more general “restricted Serre functor” $\mathbb{S}: \mathcal{T} \rightarrow \mathcal{T}$. In particular, in the same fashion negative-Calabi-Yau categories can be studied, we can look at the special case when we have Σ^{-d} instead of Σ^d

2020 *Mathematics Subject Classification.* 16E45, 18A30, 18G80.

Key words and phrases. Hom-spaces, limits and colimits, $(d+1)$ -simple minded system, SMC quadruple, $(-d)$ -Calabi-Yau triple, negative cluster category, quotient categories, truncation triangles.

in (\star) . This motivates the parallel generalisation of cluster categories to negative cluster categories, see [6, 9, 14].

Coelho Simões showed there is a parallel between the mutation theory for d -simple minded systems in $(-d)$ -Calabi-Yau triangulated categories and d -cluster-tilting objects in d -Calabi-Yau triangulated categories, see [7, Theorems A-D and the subsequent paragraph]. Moreover, in [17], Jin defined SMC quadruples $(\mathcal{T}, \mathcal{T}^p, \mathbb{S}, \mathcal{S})$, where the case $\mathbb{S} = \Sigma^{-d}$ is the analogue of Iyama and Yang's triples, that is $(-d)$ -Calabi-Yau triples, see Definition 2.3. Moreover, Coelho Simões, Pauksztello and Ploog then introduced negative cluster categories in [9], which is the analogue of the “classical” cluster category considered by Buan, Marsh, Reineke, Reiten, and Todorov in [5] and by Thomas in [21].

The aim of this paper is to prove that if $(\mathcal{T}, \mathcal{T}^p, \mathcal{S})$ is a $(-d)$ -Calabi-Yau triple, then $\mathcal{T}/\mathcal{T}^p$ is a negative cluster category, that is it is Hom-finite, $(-d-1)$ -Calabi-Yau and it has a $(d+1)$ -simple minded system in the sense of Definition 2.4. Note that Jin already proved this in [17, Theorem 4.5]. In the same way we reproved the corresponding result in the positive-Calabi-Yau setup in [10], we present an alternative proof of this result which uses different and more classic means, giving in this way a deeper understanding of $\mathcal{T}/\mathcal{T}^p$. In particular, we use limits and colimits to explicitly describe Hom spaces in this quotient category.

Let k be an algebraically closed field and $(\mathcal{T}, \mathcal{T}^p, \mathbb{S}, \mathcal{S})$ be an SMC quadruple. By property (RS2) of SMC quadruples, see Definition 2.3, for every integer i , the pair

$$(\mathcal{T}_{>i}, \mathcal{T}_{\leq i}) := ({}^\perp(\Sigma^{>-i-1}\mathcal{S}), {}^\perp(\Sigma^{<-i}\mathcal{S}))$$

is a co- t -structure on \mathcal{T} . Hence, for every object X in \mathcal{T} , there is a truncation triangle of the form

$$X_{>i} \xrightarrow{f_i} X \xrightarrow{g_i} X_{\leq i} \rightarrow \Sigma X_{>i},$$

where $X_{>i} \in \mathcal{T}_{>i}$ and $X_{\leq i} \in \mathcal{T}_{\leq i}$. Note that such a triangle is not unique and when we write $X_{>i}$, respectively $X_{\leq i}$, we mean an object fitting in one such triangle.

The main result of this paper gives a relation between Hom-spaces in \mathcal{T} and Hom-spaces in $\mathcal{T}/\mathcal{T}^p$ using limits and colimits. Note that our theorem recalls some results by Artin and Zhang, see [3, Propositions 2.2 and 3.13].

Theorem A (=Theorem 3.8). *Let X and Y be objects in \mathcal{T} .*

(a) *For $p \gg 0$, each direct system*

$$\mathcal{T}(Y, X) \longrightarrow \mathcal{T}(Y, X_{\leq 0}) \longrightarrow \mathcal{T}(Y, X_{\leq -1}) \longrightarrow \cdots \longrightarrow \mathcal{T}(Y, X_{\leq -p}) \longrightarrow \cdots$$

stabilizes. Moreover, we have that $\varinjlim_p \mathcal{T}(Y, X_{\leq -p}) \cong \mathcal{T}/\mathcal{T}^p(Y, X)$.

(b) *Fixing truncation triangles for X and Y , we have that*

$$\varprojlim_q \left(\varinjlim_p \mathcal{T}(Y_{\leq -q}, \Sigma X_{>-p}) \right) \cong \mathcal{T}/\mathcal{T}^p(Y, X).$$

Assume now that $\mathbb{S} = \Sigma^{-d}$ for some positive integer d . Let Y be in \mathcal{S} . Considering $\Sigma^j Y$ for any integer j , we prove that the system from part (a) of the theorem is stable for $p \geq j - d + 1$, and so

$$\mathcal{T}/\mathcal{T}^p(\Sigma^j Y, X) \cong \mathcal{T}(\Sigma^j Y, X_{\leq -j+d-1}),$$

see Corollary 4.2. If we further have $X \in \mathcal{S}$, we can say when the Hom-space over $\mathcal{T}/\mathcal{T}^p$ vanishes, see Corollary 4.3.

We apply our theorem to prove the following result saying that $\mathcal{T}/\mathcal{T}^p$ is a negative cluster category coming from a negative-Calabi-Yau triple. Note that this is the analogue of Iyama and Yang's generalised cluster category coming from a positive-Calabi-Yau triple, see [16, Section 5].

Theorem B.

- (a) *The category $\mathcal{T}/\mathcal{T}^p$ is Hom-finite and $(-d-1)$ -Calabi-Yau.*
- (b) *The subcategory $\mathcal{S} \subseteq \mathcal{T}/\mathcal{T}^p$ is a $(d+1)$ -simple minded system $((d+1)$ -SMS) in the sense of Definition 2.4.*

Parts (a) and (b) correspond to Theorems 4.5 and 4.7 respectively. Note that these results have already been proven by Jin in [17, Theorem 4.5] but we present a more efficient proof. We use Theorem A to prove both part (a) and properties (1) and (2) of Definition 2.4 and a lemma by Iyama and Yang from [15] to prove property (3). Note that Jin relies on [2] to prove part (a) and on [15] for his proof of part (b) but he applies the main theorem of Iyama and Yang's paper.

The paper is organised as follows. Section 2 presents some background material and our setup. Section 3 studies Hom spaces in $\mathcal{T}/\mathcal{T}^p$ and proves Theorem A. Finally, Section 4 applies Theorem A to prove Theorem B.

2. BACKGROUND

The definition of simple-minded collections was first introduced by Koenig and Yang in [18, Definition 3.2], but the same concept was also defined earlier by Al-Nofayee in [1] as *cohomologically Schurian set of generators*. We recall it here in the form presented in [17, Definition 2.4] by Jin. Note that the base field k is assumed to be algebraically closed throughout the paper.

Definition 2.1. Let \mathcal{T} be a k -linear, Hom-finite, Krull-Schmidt triangulated category and \mathcal{S} be a subcategory of \mathcal{T} . We say that \mathcal{S} is a *simple minded collection (SMC)* if the following hold for any objects X and Y in \mathcal{S} :

- (1) $\mathcal{T}(X, \Sigma^{<0}Y) = 0$;
- (2) $\dim_k \mathcal{T}(X, Y) = \delta_{XY}$;
- (3) $\text{thick}(\mathcal{S}) = \mathcal{T}$.

Notation 2.2. Let \mathcal{T} be a triangulated category and \mathcal{X}, \mathcal{Y} two full subcategories of \mathcal{T} . We write

$$\mathcal{X} * \mathcal{Y} := \{T \in \mathcal{T} \mid \text{there exists a triangle } X \rightarrow T \rightarrow Y \rightarrow \Sigma X \text{ in } \mathcal{T}, \text{ with } X \in \mathcal{X}, Y \in \mathcal{Y}\}.$$

When $\mathcal{T}(\mathcal{X}, \mathcal{Y}) = 0$ holds, we write $\mathcal{X} * \mathcal{Y} = \mathcal{X} \perp \mathcal{Y}$. Moreover, we write

$$\mathcal{X}^\perp = \{T \in \mathcal{T} \mid \mathcal{T}(\mathcal{X}, T) = 0\} \quad \text{and} \quad {}^\perp \mathcal{X} = \{T \in \mathcal{T} \mid \mathcal{T}(T, \mathcal{X}) = 0\}.$$

Definition 2.3 ([17, Definition 4.1]). We say that a quadruple $(\mathcal{T}, \mathcal{T}^p, \mathbb{S}, \mathcal{S})$ is a *simple minded collection (SMC) quadruple* if the following are satisfied:

- (RS0) \mathcal{T} is a k -linear, Hom-finite, Krull-Schmidt triangulated category and \mathcal{T}^p is a thick subcategory of \mathcal{T} ;
- (RS1) $\mathbb{S} : \mathcal{T} \rightarrow \mathcal{T}$ is a triangle equivalence restricting to an equivalence $\mathbb{S} : \mathcal{T}^p \rightarrow \mathcal{T}^p$ and satisfying a bifunctorial isomorphism for any $X \in \mathcal{T}^p$ and $Y \in \mathcal{T}$:

$$D\mathcal{T}(X, Y) \cong \mathcal{T}(Y, \mathbb{S}X);$$

- (RS2) \mathcal{S} is a SMC in \mathcal{T} and $\mathcal{T} = {}^\perp(\Sigma^{\geq 0}\mathcal{S}) \perp {}^\perp(\Sigma^{< 0}\mathcal{S}) = (\Sigma^{\geq 0}\mathcal{S})^\perp \perp (\Sigma^{< 0}\mathcal{S})^\perp$ are co- t -structures of \mathcal{T} satisfying ${}^\perp(\Sigma^{\geq 0}\mathcal{S}) \subset \mathcal{T}^p$ and $(\Sigma^{< 0}\mathcal{S})^\perp \subset \mathcal{T}^p$.

Moreover, if $\mathbb{S} = \Sigma^{-d}$ for some integer $d \geq 0$, we say that $(\mathcal{T}, \mathcal{T}^p, \mathcal{S})$ is a $(-d)$ -Calabi-Yau $((-d)$ -CY) triple.

The definition of n -simple minded system was first introduced in [7, Section 1] and, applying [8, Lemma 2.8] to the equivalent definition [8, Definition 2.1], one can see it is equivalent to the following.

Definition 2.4. Let \mathcal{T} be a k -linear, Hom-finite, Krull-Schmidt triangulated category and $n \geq 1$ be an integer. A collection of objects \mathcal{C} in \mathcal{T} is an n -simple minded system (n -SMS) if:

- (1) $\dim_k \mathcal{T}(X, Y) = \delta_{XY}$ for every X, Y in \mathcal{C} ;
- (2) if $n \geq 2$, $\text{Hom}(\Sigma^m X, Y) = 0$ for $1 \leq m \leq n-1$ and X, Y in \mathcal{C} ;
- (3) $\mathcal{T} = \langle \mathcal{C} \rangle * \Sigma^{-1}\langle \mathcal{C} \rangle * \dots * \Sigma^{1-n}\langle \mathcal{C} \rangle$.

Remark 2.5. Note that we are following the definition as stated in [8], while Jin applies a shift to it when defining an n -SMS in [17, Definition 2.7].

Setup 2.6. Let $(\mathcal{T}, \mathcal{T}^p, \mathbb{S}, \mathcal{S})$ be an SMC quadruple.

Notation 2.7. Let i be an integer, we set $\mathcal{T}_{>i} = \mathcal{T}_{\geq i+1} := {}^\perp(\Sigma^{>-i-1}\mathcal{S})$, $\mathcal{T}_{<i+1} = \mathcal{T}_{\leq i} := {}^\perp(\Sigma^{<-i}\mathcal{S})$, $\mathcal{T}^{>i} = \mathcal{T}^{\geq i+1} := \langle \Sigma^{<-i}\mathcal{S} \rangle$ and $\mathcal{T}^{<i} = \mathcal{T}^{\leq i-1} := \langle \Sigma^{\geq 1-i}\mathcal{S} \rangle$.

Remark 2.8. Note that $\mathcal{T} = \mathcal{T}_{>i} \perp \mathcal{T}_{\leq i}$ is a co- t -structure with $\mathcal{T}_{>i} \subset \mathcal{T}^p$. Hence $\mathcal{T}_{>i+1} \subseteq \Sigma \mathcal{T}_{>i+1} = \mathcal{T}_{>i}$ and $\mathcal{T}_{\leq i} \subseteq \Sigma^{-1} \mathcal{T}_{\leq i} = \mathcal{T}_{\leq i+1}$. As $(\mathcal{T}_{>i}, \mathcal{T}_{\leq i})$ is a torsion pair, for any $X \in \mathcal{T}$ there is a (non-unique) truncation triangle of the form

$$X_{>i} \xrightarrow{f_i} X \xrightarrow{g_i} X_{\leq i} \rightarrow \Sigma X_{>i},$$

where $X_{>i} \in \mathcal{T}_{>i}$, $X_{\leq i} \in \mathcal{T}_{\leq i}$ and we may assume that g_i is left minimal. When we write $X_{>i}$, respectively $X_{\leq i}$, we mean an object fitting in one of such truncation triangles.

Note also that $\mathcal{T} = \mathcal{T}^{\leq i} \perp \mathcal{T}^{>i}$ is a bounded t -structure and $\mathcal{T}^{\leq i} = \mathcal{T}_{\leq i}$, see [17, Top of page 12].

Lemma 2.9. For any $X \in \mathcal{T}$, fixing a set of truncation triangles as described in Remark 2.8, there is a direct system of the form

$$X \xrightarrow{g_0} X_{\leq 0} \xrightarrow{\xi_0} X_{\leq -1} \xrightarrow{\xi_{-1}} X_{\leq -2} \xrightarrow{\xi_{-2}} \dots.$$

Proof. Given X in \mathcal{T} and an integer i , then fixing truncation triangles as in Remark 2.8, there is a morphism of triangles of the form:

$$\begin{array}{ccccccc} X_{>i} & \xrightarrow{f_i} & X & \xrightarrow{g_i} & X_{\leq i} & \longrightarrow & \Sigma X_{>i} \\ \downarrow \alpha_i & & \parallel & & \downarrow \xi_i & & \downarrow \\ X_{>i-1} & \xrightarrow{f_{i-1}} & X & \xrightarrow{g_{i-1}} & X_{\leq i-1} & \longrightarrow & \Sigma X_{>i-1} \end{array}$$

where α_i such that $f_{i-1} \circ \alpha_i = f_i$ exists since $X_{>i} \in \mathcal{T}_{>i} \subset \mathcal{T}_{>i-1}$, $X_{\leq i-1} \in \mathcal{T}_{\geq i-1}$ and $(\mathcal{T}_{>i-1}, \mathcal{T}_{\leq i-1})$ is a torsion pair and ξ_i then exists by the axioms of triangulated categories. Then

$$X \xrightarrow{g_0} X_{\leq 0} \xrightarrow{\xi_0} X_{\leq -1} \xrightarrow{\xi_{-1}} X_{\leq -2} \xrightarrow{\xi_{-2}} \dots$$

is a direct system. \square

3. MORPHISMS IN $\mathcal{T}/\mathcal{T}^p$

The goal of this section is to prove Theorem 3.8. The whole section closely follows [10, Section 3] which proves the corresponding result for cluster categories coming from positive-Calabi-Yau triples. Most of the proofs in this section are the same as the proofs of the corresponding results in [10] up to using the direct system obtained from co- t -structure truncations and the Hom-vanishing coming from the simple-minded analogues in [17] of the corresponding silting results in [16]. Note that in [10] the truncation triangles come from t -structures and are hence unique. On the other hand, here they come from co- t -structures and are not unique, and so the direct system and the objects C_{-p}^X introduced later in the section depend on the choice of the truncation triangles.

Lemma 3.1. Let $X \in \mathcal{T}$. For any integer m , in the quotient category $\mathcal{T}/\mathcal{T}^p$ for each truncation triangle we have that g_m and ξ_m become isomorphisms. Moreover, $X \cong_{\mathcal{T}/\mathcal{T}^p} X_{\leq m}$.

Proof. By (RS2) and since \mathcal{T}^p is closed under integer powers of Σ , we have that $\mathcal{T}_{>m} \subset \mathcal{T}^p$ for any integer m . Then any truncation triangle

$$X_{>m} \xrightarrow{f_m} X \xrightarrow{g_m} X_{\leq m} \rightarrow \Sigma X_{>m}$$

viewed as a triangle in $\mathcal{T}/\mathcal{T}^p$ is such that $X_{>m} \cong_{\mathcal{T}/\mathcal{T}^p} 0$ and g_m is an isomorphism. Hence $X \cong_{\mathcal{T}/\mathcal{T}^p} X_{\leq m}$. Since $\xi_m \circ g_m = g_{m-1}$, while g_{m-1} and g_m become isomorphisms in $\mathcal{T}/\mathcal{T}^p$, so does ξ_m . \square

Remark 3.2. Given X and Y in \mathcal{T} , applying the functor $\mathcal{T}(Y, -)$ to a direct system as in Lemma 2.9, we obtain a direct system of the form

$$\mathcal{T}(Y, X) \longrightarrow \mathcal{T}(Y, X_{\leq 0}) \longrightarrow \mathcal{T}(Y, X_{\leq -1}) \longrightarrow \mathcal{T}(Y, X_{\leq -2}) \longrightarrow \dots$$

Moreover, passing to the quotient category $\mathcal{T}/\mathcal{T}^p$ using the quotient functor $Q : \mathcal{T} \rightarrow \mathcal{T}/\mathcal{T}^p$, we obtain another direct system and a commutative diagram of the form

$$\begin{array}{ccccccc} \mathcal{T}(Y, X) & \longrightarrow & \mathcal{T}(Y, X_{\leq 0}) & \longrightarrow & \mathcal{T}(Y, X_{\leq -1}) & \longrightarrow & \mathcal{T}(Y, X_{\leq -2}) \longrightarrow \dots \\ \downarrow Q(-) & & \downarrow Q(-) & & \downarrow Q(-) & & \downarrow Q(-) \\ \mathcal{T}/\mathcal{T}^p(Y, X) & \xrightarrow{\sim} & \mathcal{T}/\mathcal{T}^p(Y, X_{\leq 0}) & \xrightarrow{\sim} & \mathcal{T}/\mathcal{T}^p(Y, X_{\leq -1}) & \xrightarrow{\sim} & \mathcal{T}/\mathcal{T}^p(Y, X_{\leq -2}) \xrightarrow{\sim} \dots \end{array}$$

where all the arrows in the bottom row are isomorphisms by Lemma 3.1. Then, by the universal property of direct systems, there exists a unique morphism of the form

$$\Psi : \varinjlim_p \mathcal{T}(Y, X_{\leq -p}) \rightarrow \varinjlim_p \mathcal{T}/\mathcal{T}^p(Y, X_{\leq -p})$$

such that the diagram

$$\begin{array}{ccc}
 \mathcal{T}(Y, X_{\leq -q}) & \xrightarrow{Q(-)} & \mathcal{T}/\mathcal{T}^p(Y, X_{\leq -q}) \\
 \nu_{-q} \downarrow & & \bar{\nu}_{-q} \downarrow \wr \\
 \varinjlim_p \mathcal{T}(Y, X_{\leq -p}) & \xrightarrow{\Psi} & \varinjlim_p \mathcal{T}/\mathcal{T}^p(Y, X_{\leq -p})
 \end{array} \tag{1}$$

commutes for every $q \geq 0$.

Lemma 3.3. For a construction as in Remark 3.2, the morphism

$$\Psi : \varinjlim_p \mathcal{T}(Y, X_{\leq -p}) \rightarrow \varinjlim_p \mathcal{T}/\mathcal{T}^p(Y, X_{\leq -p})$$

is an isomorphism.

Proof. We first prove that Ψ is injective. Suppose that $\beta \in \varinjlim_p \mathcal{T}(Y, X_{\leq -p})$ is such that $\Psi(\beta) = 0$. By [20, Lemma 5.30(i)], we have that β comes from an element in one of the Hom-spaces of the direct system, say from the element $\alpha \in \mathcal{T}(Y, X_{\leq -q})$. Then, considering the commutative diagram (1), we have

$$\bar{\nu}_{-q} \circ Q(\alpha) = \Psi \circ \nu_{-q}(\alpha) = \Psi(\beta) = 0.$$

Since $\bar{\nu}_{-q}$ is an isomorphism, we have that $Q(\alpha) = 0$. Hence there exists a morphism $f : X_{\leq -q} \rightarrow K$ in the multiplicative system being inverted when we pass to $\mathcal{T}/\mathcal{T}^p$ such that $f \circ \alpha = 0$. Consider the triangle extending f , say

$$Z \xrightarrow{\gamma} X_{\leq -q} \xrightarrow{f} K \longrightarrow \Sigma Z,$$

where $Z \in \mathcal{T}^p$. By [17, Lemma 4.9], since $Z \in \mathcal{T}^p$, we have that $\mathcal{T}(Z, \Sigma^i \mathcal{S}) \neq 0$ for finitely many $i \in \mathbb{Z}$. Hence there is an integer j such that $\mathcal{T}(Z, \Sigma^{>j} \mathcal{S}) = 0$, that is $Z \in \mathcal{T}_{\geq -j} = {}^\perp(\mathcal{T}_{\leq -j-1})$. Then, as $X_{\leq -j-1} \in \mathcal{T}_{\leq -j-1}$, we have that $\mathcal{T}(Z, X_{\leq -j-1}) = 0$. Since $f \circ \alpha = 0$, there exists a morphism $c : Y \rightarrow Z$ such that $\gamma \circ c = \alpha$, that is such that the following commutes:

$$\begin{array}{ccccc}
 & & Y & & \\
 & \swarrow c & \downarrow \alpha & \searrow 0 & \\
 Z & \xrightarrow{\gamma} & X_{\leq -q} & \xrightarrow{f} & K \longrightarrow \Sigma Z.
 \end{array}$$

If $-j-1 > -q$, then $Z \in \mathcal{T}_{\geq -j} \subseteq \mathcal{T}_{\geq -q+1} = {}^\perp(\mathcal{T}_{\leq -q})$. So $\gamma = 0$ and $\alpha = \gamma \circ c = 0$ and $\beta = \nu_{-q}(\alpha) = 0$. In the other case, that is $-j-1 \leq -q$, the direct system gives us a morphism $\xi : X_{\leq -q} \rightarrow X_{\leq -j-1}$. Then, as $\mathcal{T}(Z, X_{\leq -j-1}) = 0$, we have that $\xi \circ \gamma = 0$ and so $\xi \circ \alpha = \xi \circ \gamma \circ c = 0$. Consider the commutative diagram

$$\begin{array}{ccc}
 \mathcal{T}(Y, X_{\leq -q}) & \xrightarrow{\xi_*} & \mathcal{T}(Y, X_{\leq -j-1}) \\
 \searrow \nu_{-q} & & \downarrow \nu_{-j-1} \\
 & & \varinjlim_p \mathcal{T}(Y, X_{\leq -p}).
 \end{array}$$

We have that $0 = \nu_{-j-1}(0) = \nu_{-j-1}(\xi \circ \alpha) = \nu_{-j-1} \circ \xi_*(\alpha) = \nu_{-q}(\alpha) = \beta$. Hence Ψ is injective.

It remains to show that Ψ is also surjective. Let γ be an element in $\varinjlim \mathcal{T}/\mathcal{T}^p(Y, X_{\leq -p})$. By [20, Lemma 5.30(i)], γ comes from an element in one of the Hom-spaces of the direct system and since these are all isomorphic, we may assume γ comes from an element of the form

$$\alpha = \left[\begin{array}{ccc} & Z & \\ f \nearrow & & \nwarrow s \\ Y & & X \end{array} \right] \in \mathcal{T}/\mathcal{T}^p(Y, X),$$

where the triangle extending s , say

$$W \longrightarrow X \xrightarrow[\sim]{s} Z \longrightarrow \Sigma W,$$

has W in \mathcal{T}^p since s is in the multiplicative system being inverted. Then, [17, Lemma 4.9] implies that there exists an integer j such that $\mathcal{T}(W, \Sigma^{>j} \mathcal{S}) = 0$, that is $W \in \mathcal{T}_{\geq -j} = {}^\perp(\mathcal{T}_{\leq -j-1})$. As $X_{\leq -j-1} \in \mathcal{T}_{\leq -j-1}$, we have that $\mathcal{T}(W, X_{\leq -j-1}) = 0$. Without loss of generality, we may assume j is positive. Then, the direct system from Lemma 2.9 gives a morphism $\xi : X \rightarrow X_{\leq -j-1}$ and we have a commutative diagram:

$$\begin{array}{ccccc} W & \xrightarrow{\quad} & X & \xrightarrow{s} & Z \longrightarrow \Sigma W. \\ & \searrow 0 & \downarrow \xi & \swarrow z & \\ & & X_{\leq -j-1} & & \end{array}$$

Then, we have that

$$\left[\begin{array}{ccc} & X_{\leq -j-1} & \\ \xi \nearrow & & \nwarrow \\ X & & X_{\leq -j-1} \end{array} \right] \circ \left[\begin{array}{ccc} & Z & \\ f \nearrow & & \nwarrow s \\ Y & & X \end{array} \right] = \left[\begin{array}{ccc} & X_{\leq -j-1} & \\ z \circ f \nearrow & & \nwarrow \\ Y & & X_{\leq -j-1} \end{array} \right] = Q(z \circ f).$$

Consider the commutative diagram

$$\begin{array}{ccc} & \mathcal{T}/\mathcal{T}^p(Y, X) & \\ & \downarrow \bar{\mu} \wr & \\ \mathcal{T}(Y, X_{\leq -j-1}) & \xrightarrow{Q(-)} & \mathcal{T}/\mathcal{T}^p(Y, X_{\leq -j-1}) \\ \downarrow \nu_{-j-1} & & \downarrow \bar{\nu}_{-j-1} \wr \\ \varinjlim_p \mathcal{T}(Y, X_{\leq -p}) & \xrightarrow[\Psi]{} & \varinjlim_p \mathcal{T}/\mathcal{T}^p(Y, X_{\leq -p}) \end{array}$$

Then, we have

$$\gamma = \bar{\nu}_{-j-1} \circ \bar{\mu}(\alpha) = \bar{\nu}_{-j-1} \circ Q(z \circ f) = \Psi \circ \nu_{-j-1}(z \circ f)$$

and so Ψ is surjective. \square

Notation 3.4. Given X in \mathcal{T} , consider a direct system as in Lemma 2.9. Then, for $i \geq 1$, the triangle in \mathcal{T} extending ξ_{-i} is

$$X_{\leq -i} \xrightarrow{\xi_{-i}} X_{\leq -i-1} \longrightarrow \Sigma C_{-i}^X \longrightarrow \Sigma X_{\leq -i}.$$

Note that the object C_{-i}^X depends not only on the object X , but also on the choice of the truncation triangles from Remark 2.8.

Lemma 3.5. Given $X \in \mathcal{T}$ and an integer $i \geq 1$, we have that each object C_{-i}^X is in $\mathcal{T}_{-i} := \mathcal{T}_{\leq -i} \cap \mathcal{T}_{> -i-1}$. In particular, $C_{-i}^X \in \mathcal{T}^p$.

Proof. Using Remark 2.8 and the octahedral axiom, we have a commutative diagram of triangles in \mathcal{T} of the form

$$\begin{array}{ccccccc}
X_{>-i} & \xrightarrow{f_{-i}} & X & \xrightarrow{g_{-i}} & X_{\leq -i} & \longrightarrow & \Sigma X_{>-i} \\
\downarrow \alpha_{-i} & & \parallel & & \downarrow \xi_{-i} & & \downarrow \\
X_{>-i-1} & \xrightarrow{f_{-i-1}} & X & \xrightarrow{g_{-i-1}} & X_{\leq -i-1} & \longrightarrow & \Sigma X_{>-i-1} \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
C_{-i}^X & \longrightarrow & 0 & \longrightarrow & \Sigma C_{-i}^X & \xlongequal{\quad} & \Sigma C_{-i}^X \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
\Sigma X_{>-i} & \longrightarrow & \Sigma X & \longrightarrow & \Sigma X_{\leq -i} & \longrightarrow & \Sigma^2 X_{>-i}.
\end{array}$$

Note that

$$X_{>-i-1} \in \mathcal{T}_{>-i-1} \text{ and } \Sigma X_{>-i} \in \Sigma \mathcal{T}_{>-i} = \mathcal{T}_{>-i-1}.$$

Since $\mathcal{T}_{>-i-1}$ is closed under extensions, we have that $C_{-i}^X \in \mathcal{T}_{>-i-1} \subseteq \mathcal{T}^p$. Moreover, note that

$$X_{\leq -i-1} \in \mathcal{T}_{\leq -i-1} \text{ and } \Sigma X_{\leq -i} \in \Sigma \mathcal{T}_{\leq -i} = \mathcal{T}_{\leq -i-1}.$$

Since $\mathcal{T}_{\leq -i-1}$ is closed under extensions, we have that $\Sigma C_{-i}^X \in \mathcal{T}_{\leq -i-1} = \Sigma \mathcal{T}_{\leq -i}$ and so $C_{-i}^X \in \mathcal{T}_{\leq -i}$. \square

Lemma 3.6. Let X and Y be objects in \mathcal{T} , then for each possible C_{-p}^X , we have that $\mathcal{T}(C_{-p}^X, Y) = 0$ for $p \gg 0$.

Proof. By Remark 2.8, $\mathcal{T} = \mathcal{T}^{\leq j} \perp \mathcal{T}^{>j}$ is a bounded t -structure for any integer j and so there is an integer i such that $Y \in \mathcal{T}^{>i}$. Pick an integer $p \geq -i$, so that $-p \leq i$. Then, by Lemma 3.5 we have that

$$C_{-p}^X \in \mathcal{T}_{\leq -p} \subseteq \mathcal{T}_{\leq i} = \mathcal{T}^{\leq i},$$

where the last equality holds by Remark 2.8. As $\mathcal{T} = (\mathcal{T}^{\leq i}, \mathcal{T}^{>i})$ is a torsion pair, we then have that $\mathcal{T}(C_{-p}^X, Y) = 0$. \square

Lemma 3.7. Let X and Y be objects in \mathcal{T} and let $q \gg 0$ be an integer. We have that $\mathcal{T}(X_{\leq -q}, Y) = 0$ for each choice of $X_{\leq -q}$ from the construction in Remark 2.8.

Proof. By Remark 2.8, $\mathcal{T} = \mathcal{T}^{\leq j} \perp \mathcal{T}^{>j}$ is a bounded t -structure for any integer j and so there is an integer i such that $Y \in \mathcal{T}^{>i}$. Pick an integer $q \geq -i$, so that $-q \leq i$. Then,

$$X_{\leq -q} \in \mathcal{T}_{\leq -q} \subseteq \mathcal{T}_{\leq i} = \mathcal{T}^{\leq i},$$

where the last equality holds by Remark 2.8. As $\mathcal{T} = (\mathcal{T}^{\leq i}, \mathcal{T}^{>i})$ is a torsion pair, we then have that $\mathcal{T}(X_{\leq -q}, Y) = 0$. \square

Theorem 3.8. Let X and Y be objects in \mathcal{T} .

(a) For $p \gg 0$, each direct system

$$\mathcal{T}(Y, X) \longrightarrow \mathcal{T}(Y, X_{\leq 0}) \longrightarrow \mathcal{T}(Y, X_{\leq -1}) \longrightarrow \cdots \longrightarrow \mathcal{T}(Y, X_{\leq -p}) \longrightarrow \cdots$$

stabilizes. Moreover, we have that $\varinjlim_p \mathcal{T}(Y, X_{\leq -p}) \cong \mathcal{T}/\mathcal{T}^p(Y, X)$.

(b) *Fixing truncation triangles for X and Y , we have that*

$$\varprojlim_q \left(\varinjlim_p \mathcal{T}(Y_{\leq -q}, \Sigma X_{> -p}) \right) \cong \mathcal{T}/\mathcal{T}^p(Y, X).$$

Proof. (a) Consider any triangle $C_{-i}^X \longrightarrow X_{\leq -i} \xrightarrow{\xi_{-i}} X_{\leq -i-1} \longrightarrow \Sigma C_{-i}^X$. Applying $\mathcal{T}(Y, -)$ to it, we obtain the exact sequence

$$\mathcal{T}(Y, C_{-i}^X) \longrightarrow \mathcal{T}(Y, X_{\leq -i}) \longrightarrow \mathcal{T}(Y, X_{\leq -i-1}) \longrightarrow \mathcal{T}(Y, \Sigma C_{-i}^X).$$

By Lemma 3.5, we have that $C_{-i}^X \in \mathcal{T}^p$, so that $\mathcal{T}(Y, C_{-i}^X) \cong D\mathcal{T}(C_{-i}^X, \mathbb{S}Y)$ and $\mathcal{T}(Y, \Sigma C_{-i}^X) \cong D\mathcal{T}(C_{-i}^X, \Sigma^{-1}\mathbb{S}Y)$ by (RS1). For $i \gg 0$, by Lemma 3.6, we have that $\mathcal{T}(Y, X_{\leq -i}) \cong \mathcal{T}(Y, X_{\leq -i-1})$. Hence the direct system stabilizes as claimed and for $i \gg 0$ we have that

$$\varinjlim_p \mathcal{T}(Y, X_{\leq -p}) \cong \mathcal{T}(Y, X_{\leq -i}).$$

Then we have

$$\varinjlim_p \mathcal{T}(Y, X_{\leq -p}) \cong \varinjlim_p \mathcal{T}/\mathcal{T}^p(Y, X_{\leq -p}) \cong \mathcal{T}/\mathcal{T}^p(Y, X),$$

where the first isomorphism holds by Lemma 3.3 and the second one holds because all the Hom-spaces in the direct system over $\mathcal{T}/\mathcal{T}^p$ are isomorphic and isomorphic to the direct limit.

(b) Consider a truncation triangle $X_{> -p} \longrightarrow X \longrightarrow X_{\leq -p} \longrightarrow \Sigma X_{> -p}$. Applying $\mathcal{T}(Y_{\leq -q}, -)$ to it, we obtain the exact sequence

$$\mathcal{T}(Y_{\leq -q}, X) \longrightarrow \mathcal{T}(Y_{\leq -q}, X_{\leq -p}) \longrightarrow \mathcal{T}(Y_{\leq -q}, \Sigma X_{> -p}) \longrightarrow \mathcal{T}(Y_{\leq -q}, \Sigma X). \quad (2)$$

Taking the direct limit of (2) with respect to p , we obtain the exact sequence

$$\mathcal{T}(Y_{\leq -q}, X) \longrightarrow \mathcal{T}/\mathcal{T}^p(Y, X) \longrightarrow \varinjlim_p \mathcal{T}(Y_{\leq -q}, \Sigma X_{> -p}) \longrightarrow \mathcal{T}(Y_{\leq -q}, \Sigma X). \quad (3)$$

In the above, the first and last terms are unchanged since they do not depend on p . Moreover, for the second term, we observed that

$$\varinjlim_p \mathcal{T}(Y_{\leq -q}, X_{\leq -p}) \cong \mathcal{T}/\mathcal{T}^p(Y_{\leq -q}, X) \cong \mathcal{T}/\mathcal{T}^p(Y, X),$$

where we used part (a) for the first isomorphism and Lemma 3.1. For $q \gg 0$, by Lemma 3.7, we have that

$$\mathcal{T}(Y_{\leq -q}, X) = 0 \text{ and } \mathcal{T}(Y_{\leq -q}, \Sigma X) = 0.$$

Hence, for $q \gg 0$, (3) becomes

$$0 \longrightarrow \mathcal{T}/\mathcal{T}^p(Y, X) \xrightarrow{\sim} \varinjlim_p \mathcal{T}(Y_{\leq -q}, \Sigma X_{> -p}) \longrightarrow 0.$$

By [20, Exercise 5.22(ii)], chopping off the tail of an inverse system, we obtain the same inverse limit. Hence, we have that

$$\varprojlim_q \left(\varinjlim_p \mathcal{T}(Y_{\leq -q}, \Sigma X_{> -p}) \right) \cong \mathcal{T}/\mathcal{T}^p(Y, X).$$

□

4. APPLICATIONS OF OUR THEOREM

In this section, we work in the following more specific setup.

Setup 4.1. Let $(\mathcal{T}, \mathcal{T}^p, \mathcal{S})$ be a $(-d)$ -CY triple.

When Y is a shift of an object in \mathcal{S} , we can say more about when the direct system in Theorem 3.8(a) stabilizes and we obtain the following result.

Corollary 4.2. Let $X \in \mathcal{T}$ and $Y \in \mathcal{S}$. Then, for any integer j and any choice of truncation triangles for X , we have that

$$\mathcal{T}/\mathcal{T}^p(\Sigma^j Y, X) \cong \mathcal{T}(\Sigma^j Y, X_{\leq -j+d-1}).$$

Proof. By Theorem 3.8(a), for $p \gg 0$, we have that $\mathcal{T}(\Sigma^j Y, X_{\leq -p}) \cong \mathcal{T}/\mathcal{T}^p(\Sigma^j Y, X)$. We show that $p = j - d + 1$ is big enough. For any integer l , consider a truncation triangle

$$X_{\leq -l} \xrightarrow{\xi_{-l}} X_{\leq -l-1} \longrightarrow \Sigma C_{-l}^X \longrightarrow \Sigma X_{\leq -l},$$

where $C_{-l}^X \in \mathcal{T}_{-l}$ by Lemma 3.5. Applying $\mathcal{T}(\Sigma^j Y, -)$ to this triangle, we obtain the exact sequence

$$\mathcal{T}(\Sigma^j Y, C_{-l}^X) \longrightarrow \mathcal{T}(\Sigma^j Y, X_{\leq -l}) \xrightarrow{\alpha} \mathcal{T}(\Sigma^j Y, X_{\leq -l-1}) \longrightarrow \mathcal{T}(\Sigma^j Y, \Sigma C_{-l}^X).$$

Since $C_{-l}^X \in \mathcal{T}^p$, we have that

$$\mathcal{T}(\Sigma^j Y, C_{-l}^X) \cong D\mathcal{T}(C_{-l}^X, \Sigma^{j-d} Y) \text{ and } \mathcal{T}(\Sigma^j Y, \Sigma C_{-l}^X) \cong D\mathcal{T}(C_{-l}^X, \Sigma^{j-d-1} Y).$$

Moreover, since $C_{-l}^X \in \mathcal{T}_{\leq -l}$, we have that $\mathcal{T}(C_{-l}^X, \Sigma^{< l} \mathcal{S}) = 0$ and so $\mathcal{T}(C_{-l}^X, \Sigma^{< l} Y) = 0$. Hence, if $l \geq j - d + 1$, the morphism α is an isomorphism. \square

Moreover, if we also fix X to be an object in \mathcal{S} , then Corollary 4.2 has the following important special case.

Corollary 4.3. Let X and Y be objects in \mathcal{S} . Then, we have that $\mathcal{T}/\mathcal{T}^p(\Sigma^j Y, X) = 0$ for $1 \leq j \leq d$.

Proof. By Corollary 4.2, we have that $\mathcal{T}/\mathcal{T}^p(\Sigma^j Y, X) \cong \mathcal{T}(\Sigma^j Y, X_{\leq -j+d-1})$. Consider a truncation triangle

$$X_{> -j+d-1} \xrightarrow{f} X \xrightarrow{g} X_{\leq -j+d-1} \longrightarrow \Sigma X_{> -j+d-1},$$

where we may assume g is left minimal, see Remark 2.8, and so f is right minimal by [19, Lemma 2.5]. If $1 \leq j \leq d-1$, then $-j+d-1 \geq 0$ and $X_{> -j+d-1} \in \mathcal{T}_{> -j+d-1} \subseteq \mathcal{T}_{> 0}$. Since $\mathcal{S} \subseteq \mathcal{T}_{\leq 0}$, $X \in \mathcal{S}$ and $(\mathcal{T}_{> 0}, \mathcal{T}_{\leq 0})$ is a torsion pair, we have that $\mathcal{T}(X_{> -j+d-1}, X) = 0$. Then $f = 0$ and, as it is right minimal, we have that $X_{> -j+d-1} = 0$ and $X \cong X_{\leq -j+d-1}$. Hence

$$\mathcal{T}/\mathcal{T}^p(\Sigma^j Y, X) \cong \mathcal{T}(\Sigma^j Y, X_{\leq -j+d-1}) \cong \mathcal{T}(\Sigma^j Y, X) \cong \mathcal{T}(Y, \Sigma^{-j} X) = 0$$

by (RS2), since $-j \leq -1$.

It remains to show that the result holds for the case $j = d$. Note that in each truncation triangle

$$X_{> -1} \xrightarrow{f} X \xrightarrow{g} X_{\leq -1} \longrightarrow \Sigma X_{> -1},$$

we have that X and $\Sigma^{-1}X_{\leq -1}$ are in $\mathcal{T}_{\leq 0}$. Then, as $\mathcal{T}_{\leq 0}$ is closed under extensions, we have that $X_{> -1} \in \mathcal{T}_{\leq 0}$. Moreover, such a truncation triangle induces the exact sequence

$$\mathcal{T}(\Sigma^d Y, X) \longrightarrow \mathcal{T}(\Sigma^d Y, X_{\leq -1}) \longrightarrow \mathcal{T}(\Sigma^d Y, \Sigma X_{> -1}).$$

Note that as $Y \in \mathcal{T}_{\leq 0}$, we have that the first term is zero. Moreover, as $\Sigma X_{> -1} \in \mathcal{T}^p$, by (RS1) we have that

$$\mathcal{T}(\Sigma^d Y, \Sigma X_{> -1}) \cong D\mathcal{T}(\Sigma^{d+1} X_{> -1}, \Sigma^d Y) \cong D\mathcal{T}(X_{> -1}, \Sigma^{-1} Y) = 0,$$

where the last equality holds as $X_{> -1} \in \mathcal{T}_{\leq 0}$ by above and $\Sigma^{-1} Y \in \Sigma^{-1} \mathcal{S}$. Hence, we have that $\mathcal{T}(\Sigma^d Y, X_{\leq -1}) = 0$ and so $\mathcal{T}(\Sigma^d Y, X) = 0$ by Corollary 4.2. \square

Lemma 4.4. We have that $\dim_k \mathcal{T}/\mathcal{T}^p(S_i, S_j) = \delta_{S_i S_j}$ for every S_i and S_j in \mathcal{S} .

Proof. By Corollary 4.2, we have that $\mathcal{T}/\mathcal{T}^p(S_i, S_j) \cong \mathcal{T}(S_i, (S_j)_{\leq d-1})$. Note that since $\mathcal{S} \subseteq \mathcal{T}_{\leq 0}$, then $S_j \in \mathcal{T}_{\leq 0}$. Consider a truncation triangle

$$(S_j)_{> d-1} \xrightarrow{f} S_j \longrightarrow (S_j)_{\leq d-1} \longrightarrow \Sigma(S_j)_{> d-1},$$

where we may assume f is right minimal. Note that $(S_j)_{> d-1} \in \mathcal{T}_{> d-1} \subseteq \mathcal{T}_{> 0}$ and as $(\mathcal{T}_{> 0}, \mathcal{T}_{\leq 0})$ is a torsion pair, we have that $f = 0$. As f is right minimal, then $(S_j)_{> d-1} = 0$ and $S_j \cong (S_j)_{\leq d-1}$. Then

$$\mathcal{T}/\mathcal{T}^p(S_i, S_j) \cong \mathcal{T}(S_i, (S_j)_{\leq d-1}) \cong \mathcal{T}(S_i, S_j).$$

The result then follows because \mathcal{S} is SMC in \mathcal{T} by (RS2). \square

Theorem 4.5. The category $\mathcal{T}/\mathcal{T}^p$ is Hom-finite and $(-d-1)$ -Calabi-Yau.

Proof. Let X and Y be in \mathcal{T} . By Theorem 3.8(a), for $p \gg 0$ we have that

$$\mathcal{T}/\mathcal{T}^p(Y, X) \cong \mathcal{T}(Y, X_{\leq -p})$$

and \mathcal{T} is Hom-finite by assumption, see (RS0). Hence $\mathcal{T}/\mathcal{T}^p$ is Hom-finite.

Now let q, p be integers. For any choice of $Y_{\leq -q}$, applying the covariant functor $D\mathcal{T}(-, Y_{\leq -q})$ to a shift of a truncation triangle

$$X_{> -p} \rightarrow X \rightarrow X_{\leq -p} \rightarrow \Sigma X_{> -p},$$

we obtain the exact sequence

$$D\mathcal{T}(\Sigma^d X_{\leq -p}, Y_{\leq -q}) \longrightarrow D\mathcal{T}(\Sigma^{d+1} X_{> -p}, Y_{\leq -q}) \longrightarrow D\mathcal{T}(\Sigma^{d+1} X, Y_{\leq -q}) \longrightarrow D\mathcal{T}(\Sigma^{d+1} X_{\leq -p}, Y_{\leq -q}).$$

Taking the direct limit of the above with respect to p , we obtain the exact sequence

$$0 \longrightarrow \varinjlim_p D\mathcal{T}(\Sigma^{d+1} X_{> -p}, Y_{\leq -q}) \longrightarrow D\mathcal{T}(\Sigma^{d+1} X, Y_{\leq -q}) \longrightarrow 0,$$

where

$$\varinjlim_p D\mathcal{T}(\Sigma^d X_{\leq -p}, Y_{\leq -q}) \cong \varinjlim_p D\mathcal{T}(X_{\leq -p}, \Sigma^{-d} Y_{\leq -q}) = 0$$

by Lemma 3.7 and similarly the last term is zero. Hence

$$D\mathcal{T}(\Sigma^{d+1} X_{> -p}, Y_{\leq -q}) \cong D\mathcal{T}(\Sigma^{d+1} X, Y_{\leq -q}). \quad (4)$$

Moreover, by Theorem 3.8(a) the inverse system

$$D\mathcal{T}(\Sigma^{d+1}X, Y) \longleftarrow D\mathcal{T}(\Sigma^{d+1}X, Y_{\leq 0}) \longleftarrow D\mathcal{T}(\Sigma^{d+1}X, Y_{\leq -1}) \longleftarrow \dots$$

stabilizes and so

$$\varprojlim_q D\mathcal{T}(\Sigma^{d+1}X, Y_{\leq -q}) \cong D(\varinjlim_q \mathcal{T}(\Sigma^{d+1}X, Y_{\leq -q})). \quad (5)$$

Hence, we have that

$$\begin{aligned} \mathcal{T}/\mathcal{T}^p(Y, X) &\cong \varprojlim_q (\varinjlim_p \mathcal{T}(Y_{\leq -q}, \Sigma X_{> -p})) \\ &\cong \varprojlim_q (\varinjlim_p D\mathcal{T}(\Sigma^{d+1}X_{> -p}, Y_{\leq -q})) \\ &\cong \varprojlim_q (D\mathcal{T}(\Sigma^{d+1}X, Y_{\leq -q})) \\ &\cong D(\varinjlim_q \mathcal{T}(\Sigma^{d+1}X, Y_{\leq -q})) \\ &\cong D\mathcal{T}/\mathcal{T}^p(\Sigma^{d+1}X, Y), \end{aligned}$$

where the first isomorphism follows by Theorem 3.8(b), the second by (RS1) with $\mathbb{S} = \Sigma^{-d}$, the third by (4), the fourth by (5) and the last one by Theorem 3.8(a). \square

To prove the following, we apply [15, Lemma 3.1]. In order to be able to do this, we first need to check that the required conditions (T0)-(T2) from [15, Section 1.2] hold. Note that these conditions have also been checked by Jin in the proof of [17, Theorem 4.5]. Even if both us and Jin rely on results by Iyama and Yang from [15], our arguments differ as Jin uses the main theorem while we apply another lemma.

Lemma 4.6. We have that

$$\mathcal{T}/\mathcal{T}^p \cong \langle \mathcal{S} \rangle * \Sigma^{-1}\langle \mathcal{S} \rangle * \dots * \Sigma^{1-d}\langle \mathcal{S} \rangle * \Sigma^{-d}\langle \mathcal{S} \rangle.$$

Proof. We check that the required conditions (T0)-(T2) from [15, Section 1.2] hold:

- (T0) \mathcal{T} is a triangulated category and \mathcal{T}^p is a thick subcategory,
- (T1) \mathcal{T}^p has a torsion pair $\mathcal{T}^p = \mathcal{X} \perp \mathcal{Y}$,
- (T2) \mathcal{T} has torsion pairs $\mathcal{T} = \mathcal{X} \perp \mathcal{X}^\perp = {}^\perp \mathcal{Y} \perp \mathcal{Y}$.

Condition (T0) holds by construction. Let $\mathcal{X} = \mathcal{T}_{>0}$, $\mathcal{Y} = \mathcal{T}_{\leq 0} \cap \mathcal{T}^p$ and $\mathcal{Z} = \mathcal{X}^\perp \cap {}^\perp(\Sigma \mathcal{Y})$. We have that $\mathcal{X}^\perp = \mathcal{T}_{>0}^\perp = \mathcal{T}_{\leq 0}$ and so $\mathcal{T} = \mathcal{X} \perp \mathcal{X}^\perp$ is a co- t -structure by Remark 2.8. Moreover, for any $X \in \mathcal{T}^p$, consider a truncation triangle of the form

$$X_{>0} \longrightarrow X \longrightarrow X_{\leq 0} \longrightarrow \Sigma X_{>0},$$

where $X_{>0} \in \mathcal{T}_{>0} \subseteq \mathcal{T}^p$ and $X_{\leq 0} \in \mathcal{T}_{\leq 0}$. Since \mathcal{T}^p is closed under extensions, we have that $X_{\leq 0} \in \mathcal{T}^p$. Hence $\mathcal{T} = \mathcal{X} \perp \mathcal{X}^\perp$ restricts to the co- t -structure

$$\mathcal{T}^p = \mathcal{X} \perp (\mathcal{X}^\perp \cap \mathcal{T}^p) = \mathcal{X} \perp (\mathcal{T}_{\leq 0} \cap \mathcal{T}^p) = \mathcal{X} \perp \mathcal{Y},$$

hence (T1) holds. It remains to show the existence of the second torsion pair from (T2). We have that $\mathcal{Y} = \mathcal{T}_{\leq 0} \cap \mathcal{T}^p = {}^\perp(\Sigma^{<0}\mathcal{S}) \cap \mathcal{T}^p$. Then, for any $Y \in \mathcal{Y}$, by (RS1) we have that

$$D\mathcal{T}(\Sigma^{<0}\mathcal{S}, \mathbb{S}Y) \cong \mathcal{T}(Y, \Sigma^{<0}\mathcal{S}) = 0.$$

Hence $\mathbb{S}Y \in (\Sigma^{<0}\mathcal{S})^\perp$ and $\mathcal{Y} \subseteq \mathbb{S}^{-1}(\Sigma^{<0}\mathcal{S})^\perp$. Moreover, we have that $(\Sigma^{<0}\mathcal{S})^\perp \subseteq \mathcal{T}^p$ by (RS2) and so, using (RS1) one can check that $\mathbb{S}^{-1}(\Sigma^{<0}\mathcal{S})^\perp \subseteq \mathcal{T}^p$. Now, for any $X \in \mathbb{S}^{-1}(\Sigma^{<0}\mathcal{S})^\perp$, we have that $\mathbb{S}X \in (\Sigma^{<0}\mathcal{S}^\perp) \subseteq \mathcal{T}^p$ and

$$\mathcal{T}(X, \Sigma^{<0}\mathcal{S}) \cong D\mathcal{T}(\Sigma^{<0}\mathcal{S}, \mathbb{S}X) = 0$$

by (RS1), so that $X \in {}^\perp(\Sigma^{<0}\mathcal{S})$ and $\mathbb{S}^{-1}(\Sigma^{<0}\mathcal{S})^\perp \subseteq {}^\perp(\Sigma^{<0}\mathcal{S})$. Hence $\mathcal{Y} = \mathbb{S}^{-1}(\Sigma^{<0}\mathcal{S})^\perp$. Using the second co- t -structure from (RS2), we then conclude that $\mathcal{T} = {}^\perp\mathcal{Y} \perp \mathcal{Y}$ is a co- t -structure and (T2) holds.

Since conditions (T0)-(T2) are satisfied, we apply [15, Lemma 3.1] and conclude that for any X in \mathcal{T} , there is an object Z in \mathcal{Z} such that $X \cong_{\mathcal{T}/\mathcal{T}^p} Z$. Note that, since $\mathbb{S} = \Sigma^{-d}$ by assumption, we have that

$${}^\perp\mathcal{Y} = \mathbb{S}^{-1}(\Sigma^{\geq 0}\mathcal{S})^\perp = \Sigma^d(\Sigma^{\geq 0}\mathcal{S})^\perp = \mathcal{T}^{>-d}.$$

Recalling that $\mathcal{T}_{\leq 0} = \mathcal{T}^{\leq 0}$ by Remark 2.8, we have that

$$\mathcal{Z} = \mathcal{X}^\perp \cap {}^\perp(\Sigma\mathcal{Y}) = \mathcal{T}^{\leq 0} \cap \mathcal{T}^{\geq -d}.$$

Moreover, by [4, Proposition 1.3.13], we have that

$$\mathcal{T}^{\leq 0} \cap \mathcal{T}^{\geq -d} = \Sigma^d\langle\mathcal{S}\rangle * \Sigma^{d-1}\langle\mathcal{S}\rangle * \cdots * \Sigma\langle\mathcal{S}\rangle * \langle\mathcal{S}\rangle.$$

Hence, we conclude that

$$\begin{aligned} \mathcal{T}/\mathcal{T}^p &\cong \mathcal{T}^{\leq 0} \cap \mathcal{T}^{\geq -d} \\ &\cong \Sigma^d\langle\mathcal{S}\rangle * \Sigma^{d-1}\langle\mathcal{S}\rangle * \cdots * \Sigma\langle\mathcal{S}\rangle * \langle\mathcal{S}\rangle \\ &\cong \langle\mathcal{S}\rangle * \Sigma^{-1}\langle\mathcal{S}\rangle * \cdots * \Sigma^{1-d}\langle\mathcal{S}\rangle * \Sigma^{-d}\langle\mathcal{S}\rangle. \end{aligned}$$

□

Theorem 4.7. *We have that \mathcal{S} is a $(d+1)$ -SMS in $\mathcal{T}/\mathcal{T}^p$.*

Proof. The three properties from Definition 2.4 correspond respectively to Lemma 4.4, Corollary 4.3 and Lemma 4.6. □

REFERENCES

- [1] S. Al-Nofayee, *Simple objects in the heart of a t -structure*, J. Pure Appl. Algebra **213** (2009), 54-59.
- [2] C. Amiot, *Cluster categories for algebras of global dimension 2 and quivers with potential*, Ann. Inst. Fourier **59** (2009), 2525-2590.
- [3] M. Artin and J. J. Zhang, *Noncommutative projective schemes*, Adv. Math. **109** (1994), 228-287.
- [4] A. A. Beilinson, J. Bernstein, and P. Deligne, *Faisceaux pervers*, Astérisque **100** (1982) (Vol. 1 of the proceedings of the conference “Analysis and topology on singular spaces”, Luminy, 1981).
- [5] A. B. Buan, R. J. Marsh, M. Reineke, I. Reiten, and G. Todorov, *Tilting theory and cluster combinatorics*, Adv. Math. **204** (2006), 572-618.
- [6] R. Coelho Simões, *Hom-configurations and noncrossing partitions*, J. Algebraic Combin. **35** (2012), 313-343.
- [7] R. Coelho Simões, *Mutations of simple-minded systems in Calabi-Yau categories generated by a spherical object*, Forum Math. **29** (2017), 1065-1081.
- [8] R. Coelho Simões and D. Pauksztello, *Simple-minded systems and reduction for negative Calabi-Yau triangulated categories*, Transactions of the American Mathematical Society **373** (2020), 2463-2498.
- [9] R. Coelho Simões and D. Pauksztello and D. Ploog and A. Zvonareva, *Functorially finite hearts, simple-minded systems in negative cluster categories, and noncrossing partitions*, Compositio Math. (2021).

- [10] F. Fedele *Properties of triangulated and quotient categories arising from n -Calabi-Yau triples*, Pacific J. Math. **310** (2021), 1-21.
- [11] S. Fomin and A. Zelevinsky, *Cluster algebras I: Foundations*, J. Amer. Math. Soc. **15** (2002), 497-529.
- [12] V. Ginzburg, *Calabi-Yau algebras*, preprint (2006). [math.AG/0612139v3](#)
- [13] L. Guo, *Cluster tilting objects in generalized higher cluster categories*, J. Pure Appl. Algebra **215** (2011), 2055-2071.
- [14] O. Iyama and H. Jin, *Positive Fuss-Catalan numbers and simple-minded systems in negative Calabi-Yau triples*, preprint (2020). [math.RT/2002.09952](#)
- [15] O. Iyama and D. Yang, *Quotients of triangulated categories and Equivalences of Buchweitz, Orlov, and Amiot-Guo-Keller*, American Journal of Mathematics **142(5)** (2020), 1641-1659.
- [16] O. Iyama and D. Yang, *Silting reduction and Calabi-Yau reduction of triangulated categories*, Trans. Amer. Math. Soc. **370** (2018), 7861-7898.
- [17] H. Jin, *Reductions of triangulated categories and simple-minded collections*, preprint (2019). [math.RT/1907.05114v2](#)
- [18] S. Koenig and D. Yang, *Silting objects, simple minded collections, t -structures and co- t -structures for finite dimensional algebras*, Documenta Math. **19** (2014), 403-438.
- [19] H. Krause, *Auslander-Reiten theory via Brown representability*, K-Theory **20** (2011), 331-344.
- [20] J. J. Rotman, "An introduction to homological algebra", Springer Science and Business Media, 2008, second edition.
- [21] H. Thomas, *Defining an m -cluster category*, J. Algebra **318** (2007), 37-46.

DIPARTIMENTO DI INFORMATICA - SETTORE DI MATEMATICA UNIVERSITA' DEGLI STUDI DI VERONA
 STRADA LE GRAZIE 15 - CA' VIGNAL, I-37134 VERONA, ITALY

Email address: francesca.fedele@univr.it