

Higgs alignment and novel CP -Violating observables in two-Higgs-doublet models

Ian Low^{a,b}, Nausheen R. Shah^c, Xiao-Ping Wang^{d,e,1}

^{1a}High Energy Physics Division, Argonne National Laboratory, Argonne, Illinois 60439, USA

^bDepartment of Physics and Astronomy, Northwestern University, Evanston, Illinois 60208, USA

^cDepartment of Physics and Astronomy, Wayne State University, Detroit, Michigan 48201, USA

^dSchool of Physics, Beihang University, Beijing 100191, China

^eBeijing Key Laboratory of Advanced Nuclear Materials and Physics, Beihang University, Beijing 100191, China

Null results from searches for new physics at the Large Hadron Collider (LHC) tend to enforce the belief that new particles must be much heavier than the weak scale. We undertake a systematic study of the interplay between Higgs alignment and CP -violation in complex two-Higgs-doublet models, which enables us to construct a CP -violating scenario where new Higgs bosons are close to the weak scale after including stringent constraints from the electric dipole moment and measurements at the LHC. In addition, we propose a novel potential signal of CP -violation in the Higgs-to-Higgs decays, $h_3 \rightarrow h_2 h_1$, where h_3, h_2 , and h_1 are the heaviest, second heaviest and the Standard Model-like neutral Higgs bosons, respectively. The decay could manifest itself in triple boson final states in $h_1 h_1 h_1$ and $h_1 h_1 Z$, which are quite distinct and provide unique venues for new measurements at the LHC.

I. INTRODUCTION

CP -violation (CPV) is a critical ingredient for the matter-antimatter asymmetry in the Universe [1] and its presence is of existential significance. However, the amount of CPV in the Standard Model (SM), via the Cabbibo-Kobayashi-Maskawa mechanism, is insufficient to generate the observed baryon asymmetry [2, 3]; new sources of CPV must be present outside of the SM. A two-Higgs-doublet model (2HDM) [4] is not only one of the simplest extensions of the SM which may provide new sources for CPV [5–7], but also the prototype employed in numerous more elaborate new physics models [8].

There is vast literature on CPV and 2HDMs. However, the majority of these studies focus on detecting a CP -even and CP -odd mixture in a mass eigenstate through angular correlations or asymmetries in kinematic distributions [9–17], which requires significant experimental resources and statistics.¹ On the other hand, there are two major results derived from data collected at the LHC: 1) null searches for new particles beyond the SM, and 2) a SM-like 125 GeV Higgs. The first result suggests that new particles, if present, should be much heavier than the weak scale, while the latter implies a dominantly CP -even 125 GeV Higgs.

In light of these considerations, it becomes clear that we must reevaluate the possibility of CPV in 2HDM under the assumption of a SM-like 125 GeV Higgs, which is dubbed the alignment limit [19–21]. Of particular interest is the “alignment without decoupling” limit, where new Higgs bosons could still be present near the weak scale [22–24]. This has been done only under limited purview in the past [16, 25, 26] but we aim to achieve a comprehensive and analytical understanding.

Specifically we emphasize there are two distinct sources of CPV in 2HDM; in the mixing and in the decay of the Higgs bosons. Kinematic distributions are only sensitive to CPV in the mixing. This realization allows us to construct a benchmark scenario where new Higgs bosons are not far above the weak scale, at around 500 GeV or lighter, and propose a novel signature of CPV, without recourse to angular correlations or electric dipole moment (EDM) signals, in the Higgs-to-Higgs decay, ($h_3 \rightarrow h_2 h_1 \rightarrow 3 h_1$), whose existence is sufficient to establish CPV in complex two-Higgs-doublet models (C2HDMs).² The presence of such an observable is non-trivial, as this decay channel vanishes in the exact alignment limit. Our benchmark survives constraints from EDMs [29–33] and collider measurements, and could be discovered at the LHC in the near future.

II. THE HIGGS BASIS

The most general potential for a 2HDM [34–36] in terms of the two hypercharge-1, $SU(2)$ doublet fields $\Phi_a = (\Phi_a^+, \Phi_a^0)^T$, $a = \{1, 2\}$, is given by

$$\begin{aligned} \mathcal{V} = & m_1^2 \Phi_1^\dagger \Phi_1 + m_2^2 \Phi_2^\dagger \Phi_2 - \left(m_{12}^2 \Phi_1^\dagger \Phi_2 + \text{H.c.} \right) \\ & + \frac{\lambda_1}{2} (\Phi_1^\dagger \Phi_1)^2 + \frac{\lambda_2}{2} (\Phi_2^\dagger \Phi_2)^2 + \lambda_3 (\Phi_1^\dagger \Phi_1) (\Phi_2^\dagger \Phi_2) \\ & + \lambda_4 (\Phi_1^\dagger \Phi_2) (\Phi_2^\dagger \Phi_1) + \left[\frac{\lambda_5}{2} (\Phi_1^\dagger \Phi_2)^2 + \lambda_6 (\Phi_1^\dagger \Phi_1) (\Phi_1^\dagger \Phi_2) \right. \\ & \left. + \lambda_7 (\Phi_2^\dagger \Phi_2) (\Phi_1^\dagger \Phi_2) + \text{H.c.} \right]. \end{aligned} \quad (1)$$

¹ An exception is Ref. [18] which proposed a combination of three different decay channels.

² In models with additional CP -even scalars beyond the 2HDM, such decays may be present without CPV [27, 28]. However, the mass spectrum in this case is different from that of 2HDM.

We assume a vacuum preserving the $U(1)_{em}$ gauge symmetry and adopt a convention where both scalar vacuum expectation values (VEVs) are real and non-negative,

$$\langle \Phi_1 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v_1 \end{pmatrix}, \quad \langle \Phi_2 \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v_2 \end{pmatrix}, \quad (2)$$

where $\sqrt{v_1^2 + v_2^2} \equiv v = 246$ GeV. It is customary to define an angle β through $\tan \beta = v_2/v_1$.

We choose to study the alignment limit [22–24] in the Higgs basis [37], which is defined by two doublet fields, $H_i, i = \{1, 2\}$, having the following property

$$\langle H_1^0 \rangle = v/\sqrt{2}, \quad \langle H_2^0 \rangle = 0. \quad (3)$$

We will parametrize the Higgs basis doublets as $H_1 = (G^+, (v + \phi_1^0 + iG^0)/\sqrt{2})^T$ and $H_2 = (H^+, (\phi_2^0 + ia^0)/\sqrt{2})^T$, where G^+ and G^0 are the Goldstone bosons. The neutral fields are ϕ_1^0 , ϕ_2^0 and a^0 , and the charged field is H^+ . Moreover, our phase convention is such that ϕ_2^0 and a^0 are the CP -even and CP -odd eigenstates, defined with respect to the fermion Yukawa couplings. There is a residual $U(1)$ redundancy in the Higgs basis, labeled by $H_2 \rightarrow e^{i\eta} H_2$, which leaves Eq. (3) invariant and motivates writing the scalar potential in terms of $\mathcal{H}_2 \equiv e^{i\eta} H_2$ [38],

$$\begin{aligned} \mathcal{V} = & Y_1 H_1^\dagger H_1 + Y_2 \mathcal{H}_2^\dagger \mathcal{H}_2 + \left(Y_3 e^{-i\eta} H_1^\dagger \mathcal{H}_2 + \text{H.c.} \right) \\ & + \frac{Z_1}{2} (H_1^\dagger H_1)^2 + \frac{Z_2}{2} (\mathcal{H}_2^\dagger \mathcal{H}_2)^2 \\ & + Z_3 (H_1^\dagger H_1) (\mathcal{H}_2^\dagger \mathcal{H}_2) + Z_4 (H_1^\dagger \mathcal{H}_2) (\mathcal{H}_2^\dagger H_1) \\ & + \left[\frac{Z_5}{2} e^{-2i\eta} (H_1^\dagger \mathcal{H}_2)^2 + Z_6 e^{-i\eta} (H_1^\dagger H_1) (H_1^\dagger \mathcal{H}_2) \right. \\ & \left. + Z_7 e^{-i\eta} (\mathcal{H}_2^\dagger \mathcal{H}_2) (H_1^\dagger \mathcal{H}_2) + \text{H.c.} \right]. \end{aligned} \quad (4)$$

In the above, different choices of parameters truly represent physically distinct theories [38]. The potentially complex parameters are $\{Y_3, Z_5, Z_6, Z_7\}$.

The minimization of the scalar potential gives $Y_1 = -Z_1/2v^2$ and $Y_3 = -Z_6 v^2/2$. The first relation can be viewed as the definition of v in the Higgs basis, while the second relation implies there are only three independent complex parameters, usually taken to be $\{Z_5, Z_6, Z_7\}$. If one can find a choice of η such that all parameters in Eq. (4) are real after imposing the minimization condition, the vacuum and the bosonic sector of the 2HDM is CP -invariant. This can happen if and only if [39]

$$\text{Im}(Z_5^* Z_6^2) = \text{Im}(Z_5^* Z_7^2) = \text{Im}(Z_6^* Z_7) = 0. \quad (5)$$

Otherwise, CP invariance is broken.

In a 2HDM the most general Higgs-fermion interactions result in tree-level flavor-changing neutral currents, which can be removed by imposing a discrete $\mathbb{Z}_2$ symmetry [40–42], $\Phi_1 \rightarrow \Phi_1$ and $\Phi_2 \rightarrow -\Phi_2$. In addition, the $\mathbb{Z}_2$ symmetry can be broken softly by mass terms,

leading to $\lambda_6 = \lambda_7 = 0$ in Eq. (1).

In the Higgs basis, the existence of a softly broken $\mathbb{Z}_2$ symmetry is guaranteed through the condition [38, 43],

$$\begin{aligned} (Z_1 - Z_2) [(Z_3 + Z_4)(Z_6 + Z_7)^* - Z_2 Z_6^* - Z_1 Z_7^* \\ + Z_5^* (Z_6 + Z_7)] - 2(Z_6 + Z_7)^* (|Z_6|^2 - |Z_7|^2) = 0. \end{aligned} \quad (6)$$

Equation. (6) assumes $Z_6 + Z_7 \neq 0$ and $Z_1 \neq Z_2$, and eliminates two real degrees of freedom. In the end there are a total of nine real parameters in a complex 2HDM.

III. THE ALIGNMENT LIMIT

The alignment limit [19–21] is defined by the limit where the scalar carrying the full VEV in the Higgs basis is aligned with the 125 GeV mass eigenstate [22–24], in which case the observed Higgs boson couples to the electroweak gauge bosons with SM strength. The mass-squared matrix $\mathcal{M}^2$ in the $\phi_1^0 - \bar{\phi}_2^0 - \bar{a}^0$ basis, where $\mathcal{H}_2 = (\bar{H}^+, (\bar{\phi}_2^0 + ia^0)/\sqrt{2})^T$, can be diagonalized by an orthogonal matrix R relating $\vec{\phi} = (\phi_1^0, \bar{\phi}_2^0, \bar{a}^0)^T$ to the mass eigenstates $\vec{h} = (h_3, h_2, h_1)^T$, $\vec{h} = R \cdot \vec{\phi}$ [38],

$$\begin{aligned} R = R_{12} R_{13} \bar{R}_{23} \\ = \begin{pmatrix} c_{12} & -s_{12} & 0 \\ s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} c_{13} & 0 & -s_{13} \\ 0 & 1 & 0 \\ s_{13} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \bar{c}_{23} & -\bar{s}_{23} \\ 0 & \bar{s}_{23} & \bar{c}_{23} \end{pmatrix}. \end{aligned} \quad (7)$$

Here we have used the notation $c_{ij} = \cos \theta_{ij}$, $s_{ij} = \sin \theta_{ij}$, $\bar{c}_{23} = \cos \bar{\theta}_{23}$ and $\bar{s}_{23} = \sin \bar{\theta}_{23}$. An important observation is that $\bar{\theta}_{23}$ [44] rotates between $\bar{\phi}_2^0$ and $\bar{a}^0$, which corresponds to the phase rotation $\mathcal{H}_2 \rightarrow e^{i\bar{\theta}_{23}} \mathcal{H}_2$. Therefore the effect of the $\bar{\theta}_{23}$ rotation is to shift the η parameter labelling the Higgs basis. In the end the combination that appears in the physical couplings is $\theta_{23} \equiv \eta + \bar{\theta}_{23}$. This motivates defining [38]

$$\begin{aligned} \widetilde{\mathcal{M}}^2 \equiv \bar{R}_{23} \mathcal{M}^2 \bar{R}_{23}^T \\ = v^2 \begin{pmatrix} Z_1 & \text{Re}[\tilde{Z}_6] & -\text{Im}[\tilde{Z}_6] \\ \text{Re}[\tilde{Z}_6] & \text{Re}[\tilde{Z}_5] + A/v^2 & -\frac{1}{2}\text{Im}[\tilde{Z}_5] \\ -\text{Im}[\tilde{Z}_6] & -\frac{1}{2}\text{Im}[\tilde{Z}_5] & A/v^2 \end{pmatrix}, \end{aligned} \quad (8)$$

where $\tilde{Z}_5 = Z_5 e^{-2i\theta_{23}}$, $\tilde{Z}_{6/7} = Z_{6/7} e^{-i\theta_{23}}$, and $A = Y_2 + v^2(Z_3 + Z_4 - \text{Re}[\tilde{Z}_5])/2$. Alignment is achieved by the conditions $\text{Re}[\tilde{Z}_6] = \text{Im}[\tilde{Z}_6] = 0$.

$\widetilde{\mathcal{M}}^2$ can be diagonalized by just two angles: $\tilde{R} \widetilde{\mathcal{M}}^2 \tilde{R}^T = \text{diag}(m_{h_3}^2, m_{h_2}^2, m_{h_1}^2)$ and

$$\tilde{R} = R_{12} R_{13} = \begin{pmatrix} c_{12} c_{13} & -s_{12} & -c_{12} s_{13} \\ s_{12} c_{13} & c_{12} & -s_{12} s_{13} \\ s_{13} & 0 & c_{13} \end{pmatrix}, \quad (9)$$

which relates the mass eigenbasis (h_1, h_2, h_3) to the CP -

eigenbasis $(\phi_1^0, \phi_2^0, a^0)$

$$\begin{pmatrix} h_3 \\ h_2 \\ h_1 \end{pmatrix} = \tilde{R} \begin{pmatrix} \phi_1^0 \\ \phi_2^0 \\ \phi_3^0 \end{pmatrix} = \tilde{R} \begin{pmatrix} \phi_1^0 \\ c_{23} \phi_2^0 - s_{23} a^0 \\ s_{23} \phi_2^0 + c_{23} a^0 \end{pmatrix}. \quad (10)$$

θ_{23} will be important when discussing CP -conservation.

Recall ϕ_1^0 carries the full SM VEV and exact alignment is when ϕ_1^0 coincides with a mass eigenstate. We choose to align ϕ_1^0 with h_1 , which can be achieved by setting $c_{13} = 0$ and $\theta_{13} = \pi/2$ in Eq. (9). We also impose the ordering, $m_{h_1} \leq m_{h_2} \leq m_{h_3}$ so that $m_{h_1} = 125$ GeV.

Small departures from alignment can be parametrized by writing $\theta_{13} = \pi/2 + \epsilon$, $\epsilon \ll 1$,

$$\tilde{R} = \begin{pmatrix} -\epsilon c_{12} & -s_{12} & -c_{12}(1 - \epsilon^2/2) \\ -\epsilon s_{12} & c_{12} & -s_{12}(1 - \epsilon^2/2) \\ 1 - \epsilon^2/2 & 0 & -\epsilon \end{pmatrix}. \quad (11)$$

Equation (6) remains the same after we changing $\{Z_5, Z_6, Z_7\}$ into $\{\tilde{Z}_5, \tilde{Z}_6, \tilde{Z}_7\}$. We can use Eq. (6) to eliminate Z_2 and $\text{Im}[\tilde{Z}_7]$ and choose nine input parameters $\{v, m_{h_1}, m_{h_2}, m_{h_3}, m_{H^\pm}, \theta_{12}, \theta_{13}, Z_3, \text{Re}[\tilde{Z}_7]\}$. Some important relations are, in the approximate alignment limit,

$$\begin{aligned} \text{Re}[\tilde{Z}_5] &= \frac{1}{v^2} [c_{2\theta_{12}} (m_{h_2}^2 - m_{h_3}^2) \\ &\quad + \epsilon^2 (m_{h_3}^2 c_{12}^2 + m_{h_2}^2 s_{12}^2 - m_{h_1}^2)] , \end{aligned} \quad (12)$$

$$\text{Im}[\tilde{Z}_5] = \frac{1}{v^2} s_{2\theta_{12}} \left(1 - \frac{\epsilon^2}{2}\right) (m_{h_2}^2 - m_{h_3}^2) , \quad (13)$$

$$\text{Re}[\tilde{Z}_6] = \frac{\epsilon}{2v^2} s_{2\theta_{12}} (m_{h_3}^2 - m_{h_2}^2) , \quad (14)$$

$$\text{Im}[\tilde{Z}_6] = \frac{\epsilon}{v^2} (m_{h_1}^2 - m_{h_2}^2 c_{12}^2 - m_{h_3}^2 s_{12}^2) , \quad (15)$$

$$g_{h_1 h_2 h_3} = \epsilon v \text{Re}[\tilde{Z}_7 e^{-2i\theta_{12}}] . \quad (16)$$

From the above we see that the mass splitting between h_3 and h_2 is determined at leading order in ϵ by $\Delta m_{23}^2 \equiv (m_{h_3}^2 - m_{h_2}^2) = v^2 |Z_5|$. Therefore, in general, an $\mathcal{O}(v^2)$ splitting can be achieved with $|Z_5| \sim \mathcal{O}(1)$. Further, the CPV coupling $g_{h_1 h_2 h_3}$ is nonzero away from exact alignment and for nonzero $\text{Re}[\tilde{Z}_7 e^{-2i\theta_{12}}]$. Hence the decay $(h_3 \rightarrow h_2 h_1)$ may be achieved for reasonable choices of parameters, which however are constrained from LHC and EDM constraints, as will be discussed later. In the $\mathbb{Z}_2$ basis, where each field in the model has a well-defined $\mathbb{Z}_2$ charge, the Yukawa interactions must also respect the $\mathbb{Z}_2$ invariance, which necessitates assigning $\mathbb{Z}_2$ charges to SM fermions as well [45, 46]. Two distinct possibilities exist in the literature, leading to type I [47, 48] and type II [48, 49] models which differ by interchanging $\tan \beta$ with $\cot \beta$ in the Yukawa couplings. Importantly $\tan \beta$ is a derived parameter [38, 50] which strongly depends on the mass spectrum for type II 2HDM. In the left panel of Fig. 2 we show contours of $\tan \beta$ in the $m_{h_2} - m_{h_3}$ plane. For our parameter region of interest, $\tan \beta \sim 1$ except

when m_{h_2} and m_{h_3} are degenerate. We focus on type II models with $\tan \beta \sim \mathcal{O}(1)$, in which region type I and type II models have similar Yukawa couplings.

IV. TWO CP -CONSERVING LIMITS

The condition for CP invariance in Eq. (5) can be re-alized as follows [6, 38]:

$$\text{CPC1} : \text{Im}[\tilde{Z}_5] = \text{Im}[\tilde{Z}_6] = \text{Im}[\tilde{Z}_7] = 0 , \quad (17)$$

$$\text{CPC2} : \text{Im}[\tilde{Z}_5] = \text{Re}[\tilde{Z}_6] = \text{Re}[\tilde{Z}_7] = 0 . \quad (18)$$

In CPC1, $\tilde{\mathcal{M}}^2$ in Eq. (8) is block-diagonal; $\tilde{\mathcal{M}}_{13}^2 = \tilde{\mathcal{M}}_{23}^2 = 0$, in which case ϕ_1^0 and $\tilde{\phi}_2^0$ defined in Eq. (10) are CP -even and can mix in general, whereas $\tilde{\phi}_3^0$ is CP -odd. This can be achieved by $\theta_{23} = 0$ so that $\tilde{\phi}_3^0 = a^0$ in Eq. (10). Further, neither of the two CP -even states can mix with the CP -odd state. From Eq. (9) we see θ_{13} controls the mixing between ϕ_1^0 and $\tilde{\phi}_3^0$, which implies $\theta_{13} = \pi/2$ in the CP -conserving limit. This coincides with the exact alignment limit $\epsilon = 0$. The mixing between $\tilde{\phi}_2^0$ and $\tilde{\phi}_3^0$ is dictated by θ_{12} and can be removed by $\theta_{12} = 0$ or $\pi/2$, which corresponds to $h_3 = a^0$ or $h_2 = a^0$, respectively. Therefore, CPC1 is reached by

$$\theta_{13} = \frac{\pi}{2}, \theta_{23} = 0, \theta_{12} = \{0, \pi/2\}, \text{Im}[\tilde{Z}_7] = \text{Im}[Z_7] = 0 . \quad (19)$$

One sees from Eqs. (13) and (15) that $\text{Im}[\tilde{Z}_5] = \text{Im}[\tilde{Z}_6] = 0$ under the choice of parameters in Eq. (19). It can be further checked that fermionic couplings of the mass eigenstates follow from their CP -property and the EDM constraints vanish as expected [51].

In CPC2, $\tilde{\mathcal{M}}_{12}^2 = \tilde{\mathcal{M}}_{23}^2 = 0$ and $\tilde{\mathcal{M}}^2$ is again block-diagonal. In this case ϕ_1^0 can mix with $\tilde{\phi}_3^0$, since they are both CP -even. The CP -odd state is ϕ_2^0 . Referring back to Eq. (10) we see that this requires $\theta_{23} = \pi/2$. In contrast to the CPC1 scenario, the mixing angle θ_{13} , which controls alignment, can now be arbitrary. Turning off mixing between $\tilde{\phi}_2^0$ and $\tilde{\phi}_3^0$ again implies $\theta_{12} = 0$ or $\pi/2$. Hence CPC2 is represented by:

$$\theta_{23} = \pi/2, \theta_{12} = \{0, \pi/2\}, \text{Re}[\tilde{Z}_7] = \text{Im}[Z_7] = 0 . \quad (20)$$

Again one can check that $\text{Im}[\tilde{Z}_5] = \text{Re}[\tilde{Z}_6] = 0$ and couplings of the mass eigenstates to the fermions behave as expected from their CP quantum numbers.

There is an important distinction between these two scenarios. In CPC1 the CP -conserving limit coincides with the alignment limit because misalignment introduces a small CP -odd component to the SM-like Higgs boson. Then the stringent EDM limits on CPV also constrain the misalignment, $\epsilon \sim \mathcal{O}(10^{-4})$, thereby forcing the 125 GeV Higgs to be almost exactly SM-like [51]. This is consistent with the findings in Refs. [25, 26, 52].




Figure 1. $\tan\beta$ contours in the $m_{h_2} - m_{h_3}$ plane. The relevant parameters are specified in Eq. (21). Stars denote our benchmark point.

To the contrary, in CPC2 the SM-like Higgs boson only contains a CP -even non-SM-like component. Therefore EDM limits do not constrain misalignment.³

Equations. (17) and (18) also make it clear that there are two sources of CPV in 2HDM; $\tilde{Z}_5$ and $\tilde{Z}_6$ enter into the scalar mass-squared matrix in Eq. (8), while $\tilde{Z}_7$ does not. When $\text{Im}[\tilde{Z}_5] = \text{Im}[\tilde{Z}_6] = 0$ or $\text{Im}[\tilde{Z}_5] = \text{Re}[\tilde{Z}_6] = 0$, there is no CPV in the scalar mixing matrix and each mass eigenstate h_i is also a CP -eigenstate: two are CP -even and one is CP -odd. In this case, CPV could still be present through nonzero $\text{Re}[\tilde{Z}_7]$ or $\text{Im}[\tilde{Z}_7]$ and will manifest in the bosonic interactions of the Higgs bosons. In light of these considerations, we construct a benchmark which interpolates between the CPC1 and CPC2 limits,

$$\begin{aligned} \{Z_3, \text{Re}[\tilde{Z}_7], \theta_{12}, \theta_{23}, \epsilon\} &= \{0.1, 3, \pi/2, 1.23, 0.1\}, \\ \{m_{h_3}, m_{h_2}, m_{H^\pm}\} &= \{420, 280, 420\} \text{ GeV}. \end{aligned} \quad (21)$$

In Fig. 1 we show the $\tan\beta$ contours on the $m_{h_3} - m_{h_2}$ plane, for the region of parameter space close to our benchmark; our benchmark has $\tan\beta \sim 2.3$. With these parameters, h_1 is mostly CP -even, while h_2 and h_3 are CP -mixed states. In our benchmark the charged Higgs and h_3 are degenerate in mass so as to be consistent with precision electroweak measurements, which include the oblique parameters S , T and U [53–55]. Conventional wisdom has it that a charged Higgs lighter than 800 GeV is constrained by $b \rightarrow s\gamma$ measurements [56, 57]. However, more recent results [58] argued that the theoretical uncertainty leaves more room for the new physics contribution. So in our analysis, we set the charged Higgs mass at 420 GeV, which is considered safe for the $b \rightarrow s\gamma$

³ We emphasize that this statement concerns the EDM constraints on the alignment. It was pointed out in Ref. [25] that the Z_2 condition in Eq. (6) would force CPV to vanish in the exact alignment limit.




Figure 2. LHC constraints on $|\epsilon|$ from Higgs couplings with gluons (κ_g), vector bosons (κ_V), fermions (κ_F) and photons (κ_γ), as well as searches for $H^+ \rightarrow tb$ (cyan), $h_{2/3} \rightarrow Zh_1$ (magenta) and $h_{2/3} \rightarrow h_1 h_1$ (orange). Stars denote our benchmark point.

measurement [33].

V. LHC/EDM CONSTRAINTS

In Fig. 2 we show the LHC constraints on $|\epsilon|$ and $\text{Re}[\tilde{Z}_7]$. For Higgs coupling measurements we use recent results from both ATLAS [59, 60] and CMS [61], which constrain $\kappa_i = g_i^{\text{measured}}/g_i^{\text{SM}}$, $i = g, V, F, \gamma$. Blue, gray, red and green shaded regions correspond to regions excluded by constraints coming from κ_g , κ_V , κ_F , and κ_γ , respectively. The cyan shaded region is excluded due to searches for $H^+ \rightarrow tb$ [62, 63], which requires $\tan\beta \geq 2$ and is satisfied by our benchmark point, $\tan\beta = 2.3$. For $m_{h_3(h_2)} = 420(280)$ GeV the experimental limit from double Higgs production [64] is shown as the orange shaded region in Fig. 2 and the limit from $h_3/h_2 \rightarrow Zh_1$ search [65, 66] is given by the magenta shaded region in Fig. 2. We also checked that LHC limits on heavy Higgs decays to $t\bar{t}$ final states [67] are not relevant for our benchmark.

In our analysis, we consider both constraints from the electron EDM (eEDM) [32, 68, 69] and neutron EDM (nEDM) [70]. The most recent constraints are

$$\begin{aligned} |d_n| &< 1.8 \times 10^{-26} \text{ e cm (90\% C.L.)} \\ |d_e| &< 1.1 \times 10^{-29} \text{ e cm (90\% C.L.)} \end{aligned} \quad (22)$$

The dominant contribution for both eEDM and nEDM are the two-loop Barr-Zee(BZ) diagrams [71–77]. The BZ diagrams to d_f ($f = e, d$, and u) and d_q^C ($q = d$ and u) contain contributions from fermion-loops, Higgs boson-loops, and gauge boson-loops [16]

$$d_f = d_f(\text{fermion}) + d_f(\text{Higgs}) + d_f(\text{gauge}), \quad (23)$$




Figure 3. Contours for eEDM (d_e) in θ_{23} vs. $|\epsilon|$ (top), and $\text{Re}[\tilde{Z}_7]$ (bottom) plane. Only regions within the dashed red lines are experimentally allowed $|d_e| < 1.1 \times 10^{-29} \text{e cm}$ (90%CL) [32]. Thick red line denotes $|d_e| = 0$. Note different scales for the left/right axes and legends. Stars denote our benchmark point.

and each contribution includes

$$d_f(X) = d_f^I(X) + d_f^Z(X) + d_f^W(X). \quad (24)$$

For the nEDM, the relevant formula for d_n [73] is

$$d_n = 0.79d_d - 0.2d_u + \frac{e}{g_s} (0.59d_d^C + 0.3d_u^C), \quad (25)$$

where g_s is the QCD gauge coupling constant. Both fermion-loops and gauge boson-loops contributions are related to the CP property of the Yukawa interactions, which are parametrized by θ_{23}, θ_{12} and ϵ . The Higgs-loops contributions are both related to the CP property of Yukawa interaction and the coupling of $g_{H^\pm H^\mp h_j}$, which not only depends on $\epsilon, \theta_{23}, \theta_{12}$, but also depends on $\text{Re}[\tilde{Z}_7]$. After consider both eEDM and nEDM, we found that the eEDM constraints are stronger than those from the neutron EDM, so we only show the relevant plots for eEDM. In Fig. 3 contours for the eEDM and the experimental constraints on the most relevant parameters are shown: θ_{23} vs ϵ (left) and $\text{Re}[\tilde{Z}_7]$ (right). The solid red line denotes $d_e = 0$, while the dashed red lines bound the experimentally allowed region $|d_e| < 1.1 \times 10^{-29} \text{e cm}$ (90%CL) [32]. We fix the

mass spectrum as for the LHC constraints, and again choose $\theta_{12} = 0$. While not shown, EDM constraints are minimized when the masses are degenerate [38]. However, regardless of the mass spectrum, eEDM constraints severely limit the CPV components of the mass eigenstates. This can be seen from the limits on d_e tracking the behavior expected from our analysis of CPC1 and CPC2. Small values of θ_{23} (CPC1 limit) can only be obtained for small values of $|\epsilon|$, but for $|\theta_{23}| \sim \pi/2$ (CPC2), ϵ is effectively unconstrained. Further, small values of $\text{Re}[\tilde{Z}_7]$ are obtained for values of $\theta_{23} \sim \pi/2$ (CPC2 limit), but larger values are allowed as θ_{23} decreases. Additionally, we see that in regions far from CPC1 and CPC2, d_e can be 0 due to cancellations between various contributions. This is the region where our benchmark resides.

VI. COLLIDER PHENOMENOLOGY

With the generically small CPV components allowed in the mass eigenstates due to experimental constraints, directly probing the CP nature of the mass eigenstates will be challenging. However, the decay ($h_3 \rightarrow h_2 h_1$) could provide a smoking gun signature for CPV in 2HDMs. If kinematically accessible, this signal is maximized for maximum possible misalignment ϵ and largest possible $\text{Re}[\tilde{Z}_7]$ [cf. Eq. (16)], as allowed from LHC and where eEDM constraints are minimized. Further, we are interested in the possibility of both additional Higgs bosons being within reach of the LHC, which motivates the benchmark presented in Eq. (21).

Fig. 4 shows the branching ratios of h_3 (top panel) and h_2 (bottom panel). Gray hatching denotes mass spectra in tension with eEDM constraints. One particular decay mode we would like to focus on is the CPV Higgs-to-Higgs decay, $h_3 \rightarrow h_2 h_1$, which in our benchmark has a branching fraction $\text{BR}(h_3 \rightarrow h_2 h_1) \sim 1.5\%$. As can be seen from Eq. (16), the coupling controlling this decay is CP -violating. Under the assumption that there are two CP -even and one CP -odd scalars in 2HDM, it is easy to see that such a decay mode is forbidden if CP is conserved.

From Fig. 4 we also see that h_2 can decay into $h_1 h_1$ and $h_1 Z$, giving rise to $h_3 \rightarrow h_1 h_1 h_1$ and $h_3 \rightarrow h_1 h_1 Z$ final states via $h_3 \rightarrow h_2 h_1$. The main production channel for both h_2 and h_3 is gluon fusion. At the $\sqrt{s} = 13 \text{ TeV}$ LHC [74]:

$$\sigma(gg \rightarrow h_2) \simeq 5.8 \text{ pb}, \quad \sigma(gg \rightarrow h_3) \simeq 2.7 \text{ pb}. \quad (26)$$

The large production rate for h_3 stems from its sizable CP -odd component. Therefore, for an integrated luminosity of $\mathcal{L} = 3000 \text{ fb}^{-1}$, we will have approximately 500 CPV triple Higgs events from $h_3 \rightarrow h_1 h_1 h_1$, which is a smoking gun signature of CP -violation in C2HDMs.

The decay modes of h_3 , include $h_3 \rightarrow h_2 Z$ and $h_3 \rightarrow h_1 h_2$, will gives rise to the final state $h_1 ZZ$ and $h_1 h_1 Z$, from $h_2 \rightarrow h_1 Z$, $h_2 \rightarrow ZZ$ and $h_2 \rightarrow h_1 h_1$. The event

- [11] C.-Y. Chen, S. Dawson, and Y. Zhang, JHEP **06**, 056 (2015), arXiv:1503.01114 [hep-ph].
- [12] D. Fontes, J. C. Romão, R. Santos, and J. a. P. Silva, JHEP **06**, 060 (2015), arXiv:1502.01720 [hep-ph].
- [13] B. Grzadkowski, O. Ogreid, and P. Osland, JHEP **05**, 025 (2016), [Erratum: JHEP 11, 002 (2017)], arXiv:1603.01388 [hep-ph].
- [14] D. Fontes, M. Mühlleitner, J. C. Romão, R. Santos, J. a. P. Silva, and J. Wittbrodt, JHEP **02**, 073 (2018), arXiv:1711.09419 [hep-ph].
- [15] K. Cheung, A. Jueid, Y.-N. Mao, and S. Moretti, (2020), arXiv:2003.04178 [hep-ph].
- [16] S. Kanemura, M. Kubota, and K. Yagyu, JHEP **08**, 026 (2020), arXiv:2004.03943 [hep-ph].
- [17] L. Bian, H. M. Lee, and C. B. Park, (2020), arXiv:2008.03629 [hep-ph].
- [18] D. Fontes, J. C. Romão, R. Santos, and J. a. P. Silva, Phys. Rev. D **92**, 055014 (2015), arXiv:1506.06755 [hep-ph].
- [19] J. F. Gunion and H. E. Haber, Phys. Rev. D **67**, 075019 (2003), arXiv:hep-ph/0207010.
- [20] A. Delgado, G. Nardini, and M. Quiros, JHEP **07**, 054 (2013), arXiv:1303.0800 [hep-ph].
- [21] N. Craig, J. Galloway, and S. Thomas, (2013), arXiv:1305.2424 [hep-ph].
- [22] M. Carena, I. Low, N. R. Shah, and C. E. M. Wagner, JHEP **04**, 015 (2014), arXiv:1310.2248 [hep-ph].
- [23] M. Carena, H. E. Haber, I. Low, N. R. Shah, and C. E. M. Wagner, Phys. Rev. **D91**, 035003 (2015), arXiv:1410.4969 [hep-ph].
- [24] M. Carena, H. E. Haber, I. Low, N. R. Shah, and C. E. M. Wagner, Phys. Rev. **D93**, 035013 (2016), arXiv:1510.09137 [hep-ph].
- [25] B. Grzadkowski, O. Ogreid, and P. Osland, JHEP **11**, 084 (2014), arXiv:1409.7265 [hep-ph].
- [26] B. Grzadkowski, H. E. Haber, O. M. Ogreid, and P. Osland, JHEP **12**, 056 (2018), arXiv:1808.01472 [hep-ph].
- [27] S. Baum and N. R. Shah, JHEP **12**, 044 (2018), arXiv:1808.02667 [hep-ph].
- [28] S. Baum and N. R. Shah, (2019), arXiv:1904.10810 [hep-ph].
- [29] C. Baker *et al.*, Phys. Rev. Lett. **97**, 131801 (2006), arXiv:hep-ex/0602020.
- [30] W. C. Griffith, M. D. Swallows, T. H. Loftus, M. V. Romalis, B. R. Heckel, and E. N. Fortson, Phys. Rev. Lett. **102**, 101601 (2009), arXiv:0901.2328 [physics.atom-ph].
- [31] C. Wang, X.-H. Guo, Y. Liu, and R.-C. Li, Eur. Phys. J. C **74**, 3140 (2014), arXiv:1408.0086 [hep-ph].
- [32] V. Andreev *et al.* (ACME), Nature **562**, 355 (2018).
- [33] W. Altmannshofer, S. Gori, N. Hamer, and H. H. Patel, (2020), arXiv:2009.01258 [hep-ph].
- [34] H. Georgi, Hadronic J. **1**, 155 (1978).
- [35] M. Carena and H. E. Haber, Prog. Part. Nucl. Phys. **50**, 63 (2003), arXiv:hep-ph/0208209.
- [36] S. Davidson and H. E. Haber, Phys. Rev. D **72**, 035004 (2005), [Erratum: Phys.Rev.D 72, 099902 (2005)], arXiv:hep-ph/0504050.
- [37] F. J. Botella and J. P. Silva, Phys. Rev. **D51**, 3870 (1995), arXiv:hep-ph/9411288 [hep-ph].
- [38] R. Boto, T. V. Fernandes, H. E. Haber, J. C. Romão, and J. P. Silva, (2020), arXiv:2001.01430 [hep-ph].
- [39] L. Lavoura and J. P. Silva, Phys. Rev. **D50**, 4619 (1994), arXiv:hep-ph/9404276 [hep-ph].
- [40] S. L. Glashow and S. Weinberg, Phys. Rev. **D15**, 1958 (1977).
- [41] E. A. Paschos, Phys. Rev. **D15**, 1966 (1977).
- [42] H. Georgi and D. V. Nanopoulos, Phys. Lett. **82B**, 95 (1979).
- [43] H. E. Haber and O. Stål, Eur. Phys. J. C **75**, 491 (2015), [Erratum: Eur.Phys.J.C 76, 312 (2016)], arXiv:1507.04281 [hep-ph].
- [44] H. E. Haber and D. O’Neil, Phys. Rev. D **74**, 015018 (2006), [Erratum: Phys.Rev.D 74, 059905 (2006)], arXiv:hep-ph/0602242.
- [45] G. C. Branco and M. N. Rebelo, Phys. Lett. **160B**, 117 (1985).
- [46] L. Lavoura, Phys. Rev. D **50**, 7089 (1994), arXiv:hep-ph/9405307.
- [47] H. E. Haber, G. L. Kane, and T. Sterling, Nucl. Phys. **B161**, 493 (1979).
- [48] L. J. Hall and M. B. Wise, Nucl. Phys. **B187**, 397 (1981).
- [49] J. F. Donoghue and L. F. Li, Phys. Rev. D **19**, 945 (1979).
- [50] V. D. Barger, J. L. Hewett, and R. J. N. Phillips, Phys. Rev. D **41**, 3421 (1990).
- [51] I. Low, N. R. Shah, and X. Wang, forthcoming (2020).
- [52] B. Li and C. E. M. Wagner, Phys. Rev. D **91**, 095019 (2015), arXiv:1502.02210 [hep-ph].
- [53] S. Hossenfelder and W. Hollik, Eur. Phys. J. C **77**, 178 (2017), arXiv:1607.04610 [hep-ph].
- [54] H. E. Haber and D. O’Neil, Phys. Rev. **D83**, 055017 (2011), arXiv:1011.6188 [hep-ph].
- [55] G. Funk, D. O’Neil, and R. M. Winters, Int. J. Mod. Phys. A **27**, 1250021 (2012), arXiv:1110.3812 [hep-ph].
- [56] M. Misiak and M. Steinhauser, Eur. Phys. J. C **77**, 201 (2017), arXiv:1702.04571 [hep-ph].
- [57] M. Misiak, A. Rehman, and M. Steinhauser, JHEP **06**, 175 (2020), arXiv:2002.01548 [hep-ph].
- [58] F. U. Bernlochner, H. Lacker, Z. Ligeti, I. W. Stewart, F. J. Tackmann, and K. Tackmann (SIMBA), Phys. Rev. Lett. **127**, 102001 (2021), arXiv:2007.04320 [hep-ph].
- [59] *Projections for measurements of Higgs boson cross sections, branching ratios, coupling parameters and mass with the ATLAS detector at the HL-LHC*, Tech. Rep. ATL-PHYS-PUB-2018-054 (CERN, Geneva, 2018).
- [60] ATLAS, Phys. Rev. D **101**, 012002 (2020).
- [61] *Combined measurements of the Higgs boson’s couplings at $\sqrt{s} = 13$ TeV*, Tech. Rep. CMS-PAS-HIG-17-031 (CERN, 2018).
- [62] A. M. Sirunyan *et al.* (CMS), JHEP **01**, 096 (2020), arXiv:1908.09206 [hep-ex].
- [63] *Search for charged Higgs bosons decaying into a top-quark and a bottom-quark at $\sqrt{s} = 13$ TeV with the ATLAS detector*, Tech. Rep. ATLAS-CONF-2020-039 (CERN, Geneva, 2020).
- [64] A. M. Sirunyan *et al.* (CMS), Phys. Rev. Lett. **122**, 121803 (2019), arXiv:1811.09689 [hep-ex].
- [65] M. Aaboud *et al.* (ATLAS), JHEP **03**, 174 (2018), [Erratum: JHEP 11, 051 (2018)], arXiv:1712.06518 [hep-ex].
- [66] *Search for a heavy pseudoscalar boson decaying to a Z boson and a Higgs boson at $\sqrt{s}=13$ TeV*, Tech. Rep. CMS-PAS-HIG-18-005 (CERN, Geneva, 2018).
- [67] A. M. Sirunyan *et al.* (CMS), JHEP **04**, 171 (2020), arXiv:1908.01115 [hep-ex].
- [68] K. Cheung, J. S. Lee, E. Senaha, and P.-Y. Tseng, JHEP **06**, 149 (2014), arXiv:1403.4775 [hep-ph].

- [69] M. Jung and A. Pich, JHEP **04**, 076 (2014), arXiv:1308.6283 [hep-ph].
- [70] C. Abel *et al.* (nEDM), Phys. Rev. Lett. **124**, 081803 (2020), arXiv:2001.11966 [hep-ex].
- [71] S. M. Barr and A. Zee, Phys. Rev. Lett. **65**, 21 (1990), [Erratum: Phys. Rev. Lett. 65, 2920 (1990)].
- [72] A. Pilaftsis and C. E. Wagner, Nucl. Phys. B **553**, 3 (1999), arXiv:hep-ph/9902371.
- [73] T. Abe, J. Hisano, T. Kitahara, and K. Tobioka, JHEP **01**, 106 (2014), [Erratum: JHEP 04, 161 (2016)], arXiv:1311.4704 [hep-ph].
- [74] J. R. Andersen *et al.* (LHC Higgs Cross Section Working Group), (2013), 10.5170/CERN-2013-004, arXiv:1307.1347 [hep-ph].
- [75] S. Inoue, M. J. Ramsey-Musolf, and Y. Zhang, Phys. Rev. **D89**, 115023 (2014), arXiv:1403.4257 [hep-ph].
- [76] L. Bian, T. Liu, and J. Shu, Phys. Rev. Lett. **115**, 021801 (2015), arXiv:1411.6695 [hep-ph].
- [77] D. Egana-Ugrinovic and S. Thomas, (2018), arXiv:1810.08631 [hep-ph].