ON INITIAL-BOUNDARY VALUE PROBLEM OF THE STOCHASTIC NAVIER–STOKES EQUATIONS IN THE HALF SPACE

TONGKEUN CHANG AND MINSUK YANG

ABSTRACT. We study the initial-boundary value problem of the stochastic Navier–Stokes equations in the half space. We prove the existence of weak solutions in the standard Besov space valued random processes when the initial data belong to the critical Besov space.

1. Introduction

In this paper we study the following stochastic Navier-Stokes equations

$$du(t,x) = (\Delta u(t,x) - \nabla p(t,x) + \operatorname{div}(u \otimes u)(t,x))dt + g(t,x)dB_t$$
(1.1)

for $(t,x) \in (0,\infty) \times \mathbb{R}^n_+$, $n \ge 2$, with the boundary condition u(t,x) = 0 for $(t,x) \in (0,\infty) \times \mathbb{R}^{n-1}$, the initial condition $u(0,x) = u_0$ for $x \in \mathbb{R}^n_+$, and the random noise $g(t,x)dB_t$. Here $\{B_t(\omega) : t \ge 0, \omega \in \Omega\}$ denotes an n-dimensional Wiener process defined on a probability space Ω . The Navier–Stokes system is considered as a reliable model both in science and engineering. In spite of huge effort done by countless mathematicians, many mathematical issues related to the Navier–Stokes equations including uniqueness and regularity remain as open questions. Moreover, one of the important issues is to understand turbulence in fluid motions. Although turbulent phenomena can be frequently observed in a lot of real world situations, it is extremely hard to define or characterize them mathematically. Because chaotic behavior in fluid motions may be interpreted as a presence of randomness, many interests on the stochastic Navier–Stokes equations have been increased.

The study of the stochastic Navier–Stokes equations started around early 70s', for example, by Bensoussan and Temam [2] and Foias [14]. Although many mathematicians made important contributions, we list here only a few of them. Capiński and Gatarek [6] obtained an existence theorem for the stochastic Navier–Stokes equations in a Hilbert space setting. Mikulevicius and Rozovskii [20] proved the existence of a global weak (martingale) solution of the stochastic Navier–Stokes equation. Kim [16] established the existence of local strong solutions to the stochastic Navier–Stokes equations in \mathbb{R}^3 when the initial data are sufficiently small and slightly more regular. Taniguchi [22] studied the energy solutions to the stochastic Navier–Stokes equations in two dimensional unbounded domains. For a recent overview, we refer the reader to the lecture notes or monographs by Flandoli [13], Kuksin and Shirikyan [19], and Breit, Feireisl, Hofmanová [4] and the references cited therein.

1

For the deterministic Navier–Stokes equations, Fujita and Kato [15] proved existence of mild solutions. Many papers continued to study such solutions with wider class of initial data, which should be in some special classes due to the scaling structure of the Navier–Stokes equations. Adapting harmonic analysis tools, for example, Littlewood–Paley theory, Fujita–Kato's result extended to the refined function space setting like Besov spaces. For this, we refer the reader the monograph by Cannone [5]. After that, Koch and Tataru [17] extends the class of initial data to BMO^{-1} . This line of development is well illustrated in the monograph by Bahouri, Chemin, and Danchin [1].

Recently, Du and Zhang [12] proved existence of solutions to the stochastic Navier–Stokes equations in the whole space. Due to a technical reason they found solutions in Chemin–Lerner type spaces with the norm defined by a weighted sum of space-time integration of Littlewood–Paley projections. Our aim of this paper is to establish existence of solutions to the stochastic Navier–Stokes equations in the half space. We found our solutions in the standard Besov spaces.

To state precise concept of solutions we need to recall some standard definitions. Let $(\Omega, \mathcal{G}, \{\mathcal{G}_t\}, \mathbb{P})$ be a probability space, where $\{\mathcal{G}_t : t \geq 0\}$ is a filtration of σ -fields $\mathcal{G}_t \subset \mathcal{G}$ with \mathcal{G}_0 containing all \mathbb{P} -null subsets of Ω . Assume that w is a one-dimensional $\{\mathcal{G}_t\}$ -adapted Wiener processes defined on $(\Omega, \mathcal{G}, \mathbb{P})$. We denote by $\mathbb{E}X$ the expectation of a random variable X. We suppress the argument $\omega \in \Omega$ of random variable $X(\omega)$ when it brings no confusion.

We note that the solution u and data (u_0,g) in (1.1) are random variables. We construct suitable spaces for them using the Besov spaces. We shall use two different types of spaces. The first type emphasizes the regularity in x whereas the second type does the joint regularity in (t,x). We can consider u and g as Banach space-valued stochastic processes. Hence $(\Omega \times (0,\infty), \mathscr{P}, \mathbb{P} \otimes \ell(0,\infty))$ is a suitable choice for their common domain, where \mathscr{P} is the predictable σ -field generated by $\{\mathscr{G}_t : t \geq 0\}$ (see, for instance, pp. 84–85 of [18]) and $\ell(0,\infty)$ is the Lebesgue measure on $(0,\infty)$.

For a Banach space X and $\alpha \in \mathbb{R}$ we define the stochastic Banach spaces $\mathcal{L}^r_{\alpha}(\Omega \times (0,T),\mathcal{P};X)$ to be the space of X-valued processes with the norm

$$||f||_{\mathscr{L}^{r}_{a}(\Omega\times(0,T),\mathscr{P};X)} = \left(\mathbb{E}\int_{0}^{T} s^{\alpha r} ||f(s,\cdot)||_{X}^{r} ds\right)^{1/r}.$$

Here is the definition of solutions.

Definition 1.1 (Local solution). Let $n \ge 2$ and $2 \le p, q \le \infty$. Assume that u_0 is a $\dot{B}_{pq0}^{-1+\frac{n}{p}}(\mathbb{R}^n_+)$ -valued \mathscr{F}_0 -measurable random variable. We say that (u, τ) is a local solution for the stochastic Navier-Stokes equations (1.1) if $u \in \mathscr{L}_q^q(\Omega \times (0, \infty), \mathscr{P}; L^p(\mathbb{R}^n_+))$ is a progressively measurable process and for any $0 \le t < \infty$ and

$$\tau(\omega) = \inf \left\{ 0 \le T \le \infty : \|u(\omega)\|_{L^q_{\alpha}(0,T;L^p(\mathbb{R}^n_+))} \ge R \right\}$$

where R is a positive number, and for almost surely, $u \in L^q_\alpha(0,\tau;L^p(\mathbb{R}^n_+))$ satisfies

$$\int_{\mathbb{R}^{n}_{+}} u(x, t \wedge \tau) \Phi(x) dx - \langle u_{0}, \Phi(\cdot, 0) \rangle$$

$$= \int_{0}^{t \wedge \tau} \int_{\mathbb{R}^{n}} (u \cdot \Delta \Phi + u \cdot (\Phi_{t} + (u \otimes u) : \nabla \Phi)) dx dt + \int_{0}^{t \wedge \tau} \langle g, \Phi \rangle dB_{t}$$

for every $\Phi \in C_0^{\infty}(\overline{\mathbb{R}^n}_+ \times [0, \tau(\omega))$ with $\operatorname{div}_x \Phi = 0$, $\Phi\big|_{x_n = 0} = 0$, where $\langle \cdot, \cdot \rangle$ denotes the duality pairing.

The exact definitions and notations will be given in the next section. Here are our main theorems.

Theorem 1.2. Let $n , <math>2 < q \le \infty$, and $2\alpha = 1 - \frac{n}{p} - \frac{2}{q} \ge 0$. Assume $u_0 \in \mathbb{B}_{pq0}^{-1 + \frac{n}{p}}(\mathbb{R}_+^n)$ with div $u_0 = 0$ and $g \in \mathcal{L}_{\alpha_2}^q(\Omega \times (0, \infty), \mathcal{P}; L^{p_2}(\mathbb{R}_+^n))$ with div g = 0 where p_2 and α_2 satisfy

$$p_2 \le p$$
, $\alpha \le \alpha_2$, $\left(\frac{n}{p_2} - \frac{n}{p}\right) + 2(\alpha_2 - \alpha) = 1$.

Given a positive number ϵ , there exists a positive number δ such that if

$$||u_0||_{\mathbb{B}_{p_00}^{-1+\frac{n}{p}}(\mathbb{R}^n_+)} + ||g||_{\mathcal{L}_{a_2}^{q_2}(\Omega\times(0,\infty),\mathscr{P};L^{p_2}(\mathbb{R}^n_+))} < \delta,$$

then there is a local unique solution (u, τ) of the initial boundary value problem (1.1) in the sense Definition 1.1 such that $\mathbb{P}(\tau > 0) = 1$ and $\mathbb{P}(\tau = \infty) \ge 1 - \epsilon$.

Theorem 1.3. In addition, if $g \in \mathcal{L}^q(\Omega \times (0, \infty), \mathcal{P}; \dot{B}_{pq0}^{-2\alpha-1}(\mathbb{R}^n_+))$, then the solution u obtained in Theorem 1.2 belongs to $L^q(0, \tau(\omega); \dot{B}_{pq}^{-2\alpha}(\mathbb{R}^n_+))$ almost surely.

Remark 1.4. (1) Both theorems contains the limiting exponent $q = \infty$.

- (2) The standard Besov spaces are used throughout the paper (cf. [12]).
- (3) The regularity exponent of the initial data is negative.
- (4) The exponents in the main theorems are rather complicated and restrictive due to the Hardy–Littlewood type embedding and scaling structure of governing equations.

The paper is organized as follows. In Section 2 we present preliminaries including a few estimates for the Newtonian potential and the Heat kernel and a representation of the Helmholtz projection operator. In Section 3 we prove the existence of the solutions to the linearized Navier–Stokes equations in half space. In Section 4 and 5 we prove Theorem 1.2 and 1.3, respectively.

2. Preliminaries

Let $\mathscr{S}(\mathbb{R}^n)$ denote the Schwartz space on \mathbb{R}^n and $\mathscr{S}'(\mathbb{R}^n)$ the space of tempered distributions, which is the dual space of $\mathscr{S}(\mathbb{R}^n)$. The Fourier transform of $f \in \mathscr{S}(\mathbb{R}^n)$ is initially defined by

$$\widehat{f}(\xi) = \int_{\mathbb{R}^n} e^{-2\pi i x \cdot \xi} f(x) dx,$$

and extended to the dual space $\mathcal{S}'(\mathbb{R}^n)$.

Definition 2.1 (Besov spaces on \mathbb{R}^n). Let $\Psi \in \mathcal{S}(\mathbb{R}^n)$ be a radial Schwartz function whose Fourier transform is nonnegative, supported in the ball $|\xi| \leq 2$, equal to 1 in the ball $|\xi| \leq 1$. Let

$$\widehat{\Phi}(\xi) = \widehat{\Psi}(\xi) - \widehat{\Psi}(2\xi).$$

We denote the Littlewood-Paley projection operator by

$$\Delta_j f(x) = \int_{\mathbb{R}^n} e^{2\pi i x \cdot \xi} \widehat{\Phi}(2^{-j}\xi) \widehat{f}(\xi) d\xi \quad \text{for } j \in \mathbb{Z}.$$

Notice that

$$\sum_{j\in\mathbb{Z}}\widehat{\Phi}(2^{-j}\xi)=1, \quad \text{for } \xi\neq 0.$$

We define for $s \in \mathbb{R}$ and $1 \le p, q \le \infty$, the homogeneous Besov space $\dot{B}^s_{pq}(\mathbb{R}^n)$, which is the set of $f \in \mathscr{S}'(\mathbb{R}^n)$ with the norm

$$||f||_{\dot{B}^{s}_{pq}(\mathbb{R}^{n})} = \begin{cases} \left(\sum_{j \in \mathbb{Z}} (2^{sj} ||\Delta_{j}f||_{L^{p}(\mathbb{R}^{n})})^{q} \right)^{1/q} < \infty & \text{for } q < \infty, \\ \sup_{j \in \mathbb{Z}} 2^{sj} ||\Delta_{j}f||_{L^{p}(\mathbb{R}^{n})} < \infty & \text{for } q = \infty. \end{cases}$$

Definition 2.2 (Besov spaces on \mathbb{R}^n_+). For $1 \le p, q \le \infty$ and $s \in \mathbb{R}$, the space

$$\dot{B}^{s}_{pq}(\mathbb{R}^{n}_{+}) = \{F|_{\mathbb{R}^{n}_{+}} : F \in \dot{B}^{s}_{pq}(\mathbb{R}^{n})\}$$

is defined as the set of restrictions of $F \in \dot{B}^{s}_{pq}(\mathbb{R}^{n})$ with the norm

$$||f||_{\dot{B}^{s}_{p,q}(\mathbb{R}^{n}_{+})} = \inf \Big\{ ||F||_{\dot{B}^{s}_{p,q}(\mathbb{R}^{n})} : F \in \dot{B}^{s}_{p,q}(\mathbb{R}^{n}), \ F|_{\mathbb{R}^{n}_{+}} = f \Big\}.$$

For $1 \leq p, q < \infty$ and $s \in \mathbb{R}$, the space $\dot{B}^s_{pq0}(\mathbb{R}^n_+)$ is the closure of $C_c^{\infty}(\mathbb{R}^n_+)$ in $\dot{B}^s_{pq}(\mathbb{R}^n_+)$.

We denote by $\mathbb{B}_{pq0}^{-1+\frac{n}{p}}(\mathbb{R}^n_+)$ the stochastic Besov space with the norm

$$||f||_{\mathbb{B}^{-1+\frac{n}{p}}_{pq0}(\mathbb{R}^n_+)} = \mathbb{E}||f||_{B^{-1+\frac{n}{p}}_{pq0}(\mathbb{R}^n_+)}$$

Remark 2.3. Let $0 \le s < \infty$, $1 \le p, q \le \infty$, and let p', q' denote the Hölder conjugates of p, q, respectively. Then

$$(\dot{B}^{s}_{p'q'}(\mathbb{R}^{n}_{+}))' = \dot{B}^{-s}_{pq0}(\mathbb{R}^{n}_{+}) \quad and \quad (\dot{B}^{s}_{p'q'0}(\mathbb{R}^{n}_{+}))' = \dot{B}^{-s}_{pq}(\mathbb{R}^{n}_{+}).$$

Definition 2.4. Let X be a Banach space and I an interval in \mathbb{R} . For $1 \le r \le \infty$ and $0 \le \alpha < \infty$, we denote by $L^r_{\alpha}(I;X)$ the weighted Bochner space with the norm

$$||f||_{L^r_{\alpha}(I;X)} = \left(\int_I s^{\alpha r} ||f(s)||_X^r ds\right)^{\frac{1}{r}}.$$

In particular, if $\alpha = 0$, then it is the same as the usual Bochner space, i.e., $L_0^r(I;X) = L^r(I;X)$.

Definition 2.5 (Newtonian potential). The fundamental solution of the Laplace equation in \mathbb{R}^n is defined by

$$N(x) = \begin{cases} \frac{1}{\sigma_n(2-n)|x|^{n-2}} & \text{if } n \ge 3, \\ \frac{1}{2\pi} \ln|x| & \text{if } n = 2. \end{cases}$$

Here σ_n denotes the surface area of the unit sphere in \mathbb{R}^n .

Lemma 2.6 (Lemma 3.3 in [8]). *If* 1*and* $<math>0 \le s < \infty$, *then*

$$\left\| \nabla_x^2 \int_{\mathbb{R}^n_+} N(\cdot - y) f(y) dy \right\|_{\dot{H}^s_p(\mathbb{R}^n_+)} \lesssim \|f\|_{\dot{H}^s_p(\mathbb{R}^n_+)},$$

where $\dot{H}^s_p(\mathbb{R}^n_+)$ is the space of all restriction of $f \in \dot{H}^s_p(\mathbb{R}^n)$ on \mathbb{R}^n_+ , which is the standard homogenous factional Sobolev space defined by the Fourier inversion. (In particular, $\dot{H}^0_p(\mathbb{R}^n_+) = L^p(\mathbb{R}^n_+)$.)

Lemma 2.7. *If* $1 , <math>1 \le q \le \infty$, and 0 < s < 1/p', then

$$\left\| \nabla_{x}^{2} \int_{\mathbb{R}^{n}_{+}} N(\cdot - y) f(y) dy \right\|_{\dot{B}^{-s}_{pq}(\mathbb{R}^{n}_{+})} \lesssim \|f\|_{\dot{B}^{-s}_{pq}(\mathbb{R}^{n}_{+})},$$

$$\left\| \nabla_{x}^{2} \int_{\mathbb{R}^{n}_{+}} N(\cdot - y^{*}) f(y) dy \right\|_{\dot{B}^{-s}_{pq}(\mathbb{R}^{n}_{+})} \lesssim \|f\|_{\dot{B}^{-s}_{pq}(\mathbb{R}^{n}_{+})}.$$

where p' is the Hölder conjugate of p, i.e., 1/p + 1/p' = 1.

Proof. Applying the real interpolation (see for example Section 2 in [9]) to Lemma 2.6 we get

$$\left\| \nabla_{x}^{2} \int_{\mathbb{R}^{n}_{+}} N(\cdot - y) f(y) dy \right\|_{\dot{B}^{s}_{p'q'}(\mathbb{R}^{n}_{+})} \lesssim \|f\|_{\dot{B}^{s}_{p'q'}(\mathbb{R}^{n}_{+})}.$$

Since the operator

$$Tf(x) := \nabla_x^2 \int_{\mathbb{R}^n_+} N(x - y) f(y) dy$$

is symmetric, we use the duality argument to obtain

$$\left\| \nabla_{x}^{2} \int_{\mathbb{R}^{n}_{+}} N(\cdot - y) f(y) dy \right\|_{\dot{B}^{-s}_{pq0}(\mathbb{R}^{n}_{+})} \lesssim \|f\|_{\dot{B}^{-s}_{pq0}(\mathbb{R}^{n}_{+})}.$$

We note that if -1/p < s < 1/p', then

$$\dot{B}_{pq0}^{-s}(\mathbb{R}^{n}_{+}) = \dot{B}_{pq}^{-s}(\mathbb{R}^{n}_{+}).$$

This proves the first estimate and the proof of the second estimate is almost the same.

Definition 2.8 (Heat kernel). The fundamental solution of the heat equation in \mathbb{R}^n is defined by

$$\Gamma_{t}(x) = \Gamma(x, t) = \begin{cases} (2\pi t)^{-\frac{n}{2}} e^{-\frac{|x|^{2}}{4t}} & \text{if } t > 0\\ 0 & \text{if } t \leq 0. \end{cases}$$

We define

$$\Gamma_t * f(x) = \int_{\mathbb{R}^n} \Gamma(x - y, t) f(y) dy,$$

$$\Gamma_t^* * f(x) = \int_{\mathbb{R}^n} \Gamma(x - y^*, t) f(y) dy.$$

The following lemma is a consequence of a pointwise estimate of the localized heat kernel, which is obtained by the standard integration by parts argument.

From now on we simply write $||f||_p$ instead of $||f||_{L^p(\mathbb{R}^n)}$.

Lemma 2.9. There exists a positive constant c such that for all $1 \le p \le \infty$, t > 0, and $j \in \mathbb{Z}$

$$\|\Delta_j(\Gamma_t * f)\|_p \lesssim \exp(-ct2^{2j})\|\Delta_j f\|_p$$

Proof. Fix t > 0. For $\lambda > 0$ we define

$$K_{\lambda}(x,t) = \int_{\mathbb{R}^n} e^{2\pi i x \cdot \xi} \widehat{\Psi}(\xi/\lambda) e^{-t|\xi|^2} d\xi.$$

By a change of variables we have

$$K_{\lambda}(x,t) = \lambda^n \int_{\mathbb{R}^n} e^{2\pi i \lambda x \cdot \xi} \widehat{\Psi}(\xi) e^{-t\lambda^2 |\xi|^2} d\xi.$$

Using the identity

$$(I - \Delta_{\varepsilon})e^{2\pi i \lambda x \cdot \xi} = (1 + 4\pi^2 |\lambda x|^2)e^{2\pi i \lambda x \cdot \xi}$$

we can carry out repeated integration by parts to get the pointwise estimate of the localized heat kernel

$$K_{\lambda}(x,t) \lesssim \frac{\lambda^n \exp(-c_1 t \lambda^2)}{(1+|\lambda x|^2)^{(n+1)/2}}$$

for some positive constant c_1 . Using a change of variable yields

$$\int_{\mathbb{R}^n} K_{\lambda}(x,t) dx \lesssim \exp(-c_1 t \lambda^2). \tag{2.1}$$

The implied constant does not depend on λ . Since $\Delta_j = (\Delta_{j-1} + \Delta_j + \Delta_{j+1})\Delta_j$ and the convolutions commute, we have

$$\Delta_i(\Gamma_t * f) = (\Delta_{i-1} + \Delta_i + \Delta_{i+1})\Gamma_t * (\Delta_i f).$$

Applying Young's inequality and (2.1) we obtain that

$$\begin{split} \left\| \Delta_{j}(\Gamma_{t} * f) \right\|_{p} &\leq \left(\|K_{2^{j-1}}\|_{1} + \|K_{2^{j}}\|_{1} + \|K_{2^{j+1}}\|_{1} \right) \left\| \Delta_{j} f \right\|_{p} \\ &\leq \left(\exp(-c_{1} t 2^{2^{j-2}}) + \exp(-c_{1} t 2^{2^{j}}) + \exp(-c_{1} t 2^{2^{j+2}}) \right) \left\| \Delta_{j} f \right\|_{p} \\ &\leq 3 \exp(-c t 2^{2^{j}}) \left\| \Delta_{j} f \right\|_{p} \end{split}$$

where
$$c = c_1/4$$
.

Lemma 2.10. *If* $1 \le p, q \le \infty$ *and* $\beta \in \mathbb{R}$ *, then*

$$\|\Gamma_t * f\|_{L^q(0,\infty;\dot{B}^{\beta}_{pq}(\mathbb{R}^n))} + \|\Gamma_t^* * f\|_{L^q(0,\infty;\dot{B}^{\beta}_{pq}(\mathbb{R}^n))} \lesssim \|f\|_{\dot{B}^{\beta-\frac{2}{q}}_{pq}(\mathbb{R}^n)}.$$

Proof. Fix $1 \le p \le \infty$. If $1 \le q < \infty$, then by Lemma 2.9

$$\begin{split} \|\Gamma_t * f\|_{L^q(0,\infty;\dot{B}^{\beta}_{pq}(\mathbb{R}^n))}^q &= \int_0^\infty \sum_{j \in \mathbb{Z}} 2^{q\beta j} \|\Delta_j (\Gamma_t * f)\|_p^q dt \\ &\lesssim \int_0^\infty \sum_{j \in \mathbb{Z}} 2^{q\beta j} \exp(-ct2^{2j}q) \|\Delta_j f\|_p^q dt \\ &= \sum_{j \in \mathbb{Z}} 2^{q\beta j} \|\Delta_j f\|_p^q \int_0^\infty \exp(-ct2^{2j}q) dt \\ &= \frac{1}{cq} \sum_{j \in \mathbb{Z}} 2^{(q\beta - 2)j} \|\Delta_j f\|_p^q \\ &= \frac{1}{cq} \|f\|_{\dot{B}^{\beta - \frac{2}{q}}(\mathbb{R}^n)}^q. \end{split}$$

Note that the operator norm does not depend on q since $\lim_{q\to\infty}(cq)^{1/q}=1$. If $q=\infty$, then by Lemma 2.9

$$\begin{split} \|\Gamma_t * f\|_{L^{\infty}(0,\infty;\dot{B}^{\beta}_{p\infty}(\mathbb{R}^n))} &= \sup_{t>0} \sup_{j \in \mathbb{Z}} 2^{\beta j} \|\Delta_j(\Gamma_t * f)\|_p \\ &\lesssim \sup_{j \in \mathbb{Z}} \sup_{t>0} 2^{\beta j} \exp(-ct2^{2j}) \|\Delta_j f\|_p \\ &= \sup_{j \in \mathbb{Z}} 2^{\beta j} \|\Delta_j f\|_p \\ &= \|f\|_{\dot{B}^{\beta}_{p\infty}(\mathbb{R}^n)}. \end{split}$$

The estimates for $\Gamma_t^* * f$ can be done by exactly the same way.

Lemma 2.11. Let $1 \le p, q \le \infty$ and $0 < \alpha < \infty$. Then

$$\|\Gamma_t * f\|_{L^q_a(0,\infty;L^p(\mathbb{R}^n))} + \|\Gamma_t^* * f\|_{L^q_a(0,\infty;L^p(\mathbb{R}^n))} \lesssim \|f\|_{\dot{B}^{-2\alpha-2/q}_{g_{\mathbb{R}^n}}}.$$

Proof. First we consider the case $q = \infty$. Using the identity $f = \sum_{j \in \mathbb{Z}} \Delta_j f$ and the triangle inequality and applying Lemma 2.9, we get

$$t^{\alpha} \|\Gamma_t * f\|_p \le t^{\alpha} \sum_{i \in \mathbb{Z}} \|\Delta_j(\Gamma_t * f)\|_p \lesssim t^{\alpha} \sum_{i \in \mathbb{Z}} \exp(-ct2^{2j}) \|\Delta_j f\|_p.$$

Since $2^{-2\alpha j} \|\Delta_j f\|_p \le \|f\|_{\dot{B}^{-2\alpha}_{p\infty}(\mathbb{R}^n)}$ for all j, we have

$$t^{\alpha} \sum_{j \in \mathbb{Z}} \exp(-ct2^{2j}) \|\Delta_{j} f\|_{p} \lesssim \|f\|_{\dot{B}^{-2\alpha}_{p\infty}(\mathbb{R}^{n})} \sum_{j \in \mathbb{Z}} (t2^{2j})^{\alpha} \exp(-ct2^{2j}) \lesssim \|f\|_{\dot{B}^{-2\alpha}_{p\infty}(\mathbb{R}^{n})}.$$

Thus, we have for all t,

$$t^{\alpha} \| \Gamma_t * f \|_p \lesssim \| f \|_{\dot{B}^{-2\alpha}_{n\infty}(\mathbb{R}^n)}.$$

This proves the lemma for the case $q = \infty$.

Now we consider the case q = 1. By the same way

$$\begin{split} \int_0^\infty t^\alpha \|\Gamma_t * f\|_p dt &\leq \int_0^\infty t^\alpha \sum_{j \in \mathbb{Z}} \left\| \Delta_j (\Gamma_t * f) \right\|_p dt \\ &\lesssim \sum_{j \in \mathbb{Z}} \left\| \Delta_j f \right\|_p \int_0^\infty t^\alpha \exp(-ct2^{2j}) dt \\ &\lesssim \sum_{j \in \mathbb{Z}} 2^{-2\alpha j - 2j} \left\| \Delta_j f \right\|_p \int_0^\infty (t2^{2j})^\alpha \exp(-ct2^{2j}) 2^{2j} dt \\ &\lesssim \|f\|_{\dot{B}_{n}^{-2\alpha - 2}(\mathbb{R}^n)} \end{split}$$

since the integral $\int_0^\infty (t2^{2j})^\alpha \exp(-ct2^{2j})2^{2j}dt$ does not depend on j by a change of variable. This proves the lemma for the case q=1. Using complex interpolation (5.6.3. Theorem in [3])

$$[L^{\infty}_{\alpha}(0,\infty;L^{p}(\mathbb{R}^{n})),L^{1}_{\alpha}(0,\infty;L^{p}(\mathbb{R}^{n}))]_{\theta}=L^{q}_{\alpha}(0,\infty;L^{p}(\mathbb{R}^{n}))$$

and (6.4.5. Theorem in [3])

$$[\dot{B}_{p\infty}^{-2\alpha}(\mathbb{R}^n), \dot{B}_{p1}^{-2\alpha-2}(\mathbb{R}^n)]_{\theta} = \dot{B}_{pq}^{-2\alpha-2/q}(\mathbb{R}^n)$$

with $1/q=1-\theta$, we get the result. The estimates for Γ_t^**f can be done by exactly the same way. \Box

Here we recall the following representation of the Helmholtz projection operator. See Section 3 of [11] for more details.

Definition 2.12 (Helmholtz projection \mathbb{P} in \mathbb{R}^n_+). Let $\mathscr{F} = (F_{kl})^n_{k,l=1}$ with

$$F_{kl} = F_{lk}$$
 and $F_{mk}|_{x_n=0} = 0$.

If $f = \text{div}\mathcal{F}$, then the Helmholtz projection operator $\mathbb{P}f$ in \mathbb{R}^n_+ is defined by

$$\mathbb{P} f = \operatorname{div} \mathscr{F}',$$

where $\mathscr{F}' = (F'_{km})^n_{k,m-1}$ is given by

$$\begin{split} F'_{nm} &= F_{nm} - \delta_{nm} F_{nn}, \ m = 1, \cdots, n, \\ F'_{\beta\gamma} &= F_{\beta\gamma} - \delta_{\beta\gamma} F_{nn} + \sum_{q=1}^{n} D_{x_{\gamma}} \int_{\mathbb{R}^{n}_{+}} D_{y_{q}} N^{+}(x, y) F_{\beta q}(y) dy \\ &+ D_{x_{\gamma}} \int_{\mathbb{R}^{n}_{+}} \left(D_{y_{n}} N^{+}(x, y) F_{n\beta}(y) - D_{y_{\beta}} N^{+}(x, y) F_{nn}(y) \right) dy, \quad \beta, \gamma \neq n, \\ F'_{\beta n} &= -\sum_{\gamma=1}^{n-1} D_{x_{\gamma}} \int_{\mathbb{R}^{n}_{+}} D_{x_{n}} N^{+}(x, y) F_{\beta\gamma}(y) dy + D_{x_{\beta}} \int_{R_{+}} D_{x_{n}} N^{+}(x, y) F_{nn}(y) dy \\ &- 2F_{\beta n}(x) - 2\sum_{\gamma=1}^{n-1} D_{x_{\gamma}} \int_{\mathbb{R}^{n}_{+}} D_{x_{\gamma}} N^{-}(x, y) F_{\beta n}(y) dy \quad \beta \neq n. \end{split}$$

Here we used shorthand notations

$$N^{+}(x, y) := N(x - y) + N(x - y^{*}),$$

$$N^{-}(x, y) := N(x - y) - N(x - y^{*}).$$

Remark 2.13. By Lemma 2.6 for 1

$$\|\mathscr{F}'\|_{L^p(\mathbb{R}^n_+)} \lesssim \|\mathscr{F}\|_{L^p(\mathbb{R}^n_+)}.\tag{2.2}$$

Lemma 2.14 (Hardy-Littilewood-Sobolev inequality [23]). Let $0 < \lambda < 1$ and define for t > 0

$$I_{\lambda}f(t) = \int_{0}^{t} (t-s)^{-\lambda}f(s)ds.$$

Then the operator $I_{\lambda}: L^p_{\alpha}(0,\infty) \to L^q_{\beta}(0,\infty)$ is bounded if $1 and <math>\beta \le \alpha$ satisfy

$$0 < \alpha p + 1 < p$$
, $0 < \beta q + 1 < q$, $1 + \frac{1}{q} + \beta = \frac{1}{p} + \lambda + \alpha$.

3. LINEARIZED NAVIER-STOKES EQUATIONS

In this section, we consider the initial-boundary value problem of the stochastic linearized Navier– Stokes equations in half space

$$du(t,x) = \left(\Delta u(t,x) - \nabla p(t,x) + \operatorname{div} \mathscr{F}(t,x)\right) dt + g(t,x) dB_t \tag{3.1}$$

in $\Omega \times (0, \infty) \times \mathbb{R}^n_+$ with the boundary condition $u|_{x_n=0}=0$ and the initial condition $u|_{t=0}=u_0$. The goal of this section is to prove the following proposition.

Proposition 3.1. Let $2 \le p < \infty$, $2 < q < \infty$, and $0 \le 2\alpha < 1 - \frac{1}{p} - \frac{2}{q}$. Assume $u_0 \in \dot{\mathbb{B}}_{pq0}^{-2\alpha - \frac{2}{q}}(\mathbb{R}^n_+)$ with $\operatorname{div} u_0 = 0$, $\mathscr{F} \in \mathcal{L}_{a_1}^{q_1}(\Omega \times (0, \infty), \mathscr{P}; L^{p_1}(\mathbb{R}^n_+))$ for some p_1 , q_1 , and α_1 satisfying

$$q_1 \le q, \quad \alpha \le \alpha_1 < 1 - \frac{1}{q_1}, \quad \left(\frac{n}{p_1} - \frac{n}{p}\right) + \left(\frac{2}{q_1} - \frac{2}{q}\right) + 2(\alpha_1 - \alpha) = 1,$$
 (3.2)

and $g\in\mathcal{L}^q_{\alpha_2}(\Omega\times(0,\infty),\mathcal{P}\,;L^{p_2}(\mathbb{R}^n_+))$ for some p_2 and α_2 satisfying

$$p_2 \le p$$
, $\alpha \le \alpha_2$, $\left(\frac{n}{p_2} - \frac{n}{p}\right) + 2(\alpha_2 - \alpha) = 1$.

Then there is at least one weak solution $u \in \mathcal{L}^q_a(\Omega \times (0, \infty), \mathcal{P}; L^p(\mathbb{R}^n_+))$ to the Stokes equations (3.1) satisfying

$$\|u\|_{\mathcal{L}^{q}_{\alpha}(\Omega\times(0,\infty),\mathcal{P};L^{p}(\mathbb{R}^{n}_{+}))} \leq C_{1}\Big(\|u_{0}\|_{\dot{\mathbb{B}}^{-2\alpha-\frac{2}{q}}_{pq_{0}}(\mathbb{R}^{n}_{+})} + \|g\|_{\mathcal{L}^{q}_{\alpha_{2}}(\Omega\times(0,\infty),\mathcal{P};L^{p_{2}}(\mathbb{R}^{n}_{+}))} + \|\mathcal{F}\|_{\mathcal{L}^{q_{1}}_{\alpha_{1}}(\Omega\times(0,\infty),\mathcal{P};L^{p_{1}}(\mathbb{R}^{n}_{+}))}\Big). \tag{3.3}$$

$$\text{Moreover, the solution } u \text{ is unique in } \mathcal{L}^{q}_{\alpha}(\Omega\times(0,\infty),\mathcal{P};L^{p}(\mathbb{R}^{n}_{+})) \text{ when } 0 \leq \alpha < 1 - \frac{1}{q}.$$

Before proving the proposition we recall the following solvability lemma for the initial-boundary value problem of the deterministic linearized Navier–Stokes equations in half space. For the proof we refer the reader to Theorem 1.2 in [7] for the case $q < \infty$ and to Theorem 1.3 in [9] for the case $q = \infty$.

Lemma 3.2. Let $1 , <math>2 < q \le \infty$, and $0 \le 2\alpha < 1 - \frac{1}{p} - \frac{2}{q}$. Assume $u_0 \in \dot{B}_{pq0}^{-2\alpha - \frac{2}{q}}(\mathbb{R}^n_+)$ with $\operatorname{div} u_0 = 0$, $\mathscr{F} \in L^{q_1}_{\alpha_1}(0,\infty;L^{p_1}(\mathbb{R}^n_+))$ for some p_1 , q_1 , and α_1 satisfying (3.2). Then there is at least one weak solution $u_1 \in L^q_{\alpha}(0,\infty;L^p(\mathbb{R}^n_+))$ to the Stokes equations

$$\partial_t u_1 = \Delta u_1 - \nabla p_1 - \operatorname{div} \mathscr{F} \quad \text{and} \quad \operatorname{div} u_1 = 0 \tag{3.4}$$

in $(0, \infty) \times \mathbb{R}^n_+$ with the boundary condition $u_1|_{x_n=0}=0$ and the initial condition $u_1|_{t=0}=u_0$. Moreover, the solution satisfies the estimate

$$||u_1||_{L^q_{\alpha}(0,\infty;L^p(\mathbb{R}^n_+))} \lesssim ||u_0||_{B^{-2\alpha-\frac{2}{q}}_{n\alpha_0}(\mathbb{R}^n_+)} + ||\mathscr{F}||_{L^{q_1}_{\alpha_1}(0,\infty;L^{p_1}(\mathbb{R}^n_+))}. \tag{3.5}$$

Moreover, the solution u_1 is unique in $L^q_\alpha(0,\infty;L^p(\mathbb{R}^n_+))$ when $0\leq \alpha<1-\frac{1}{q}$.

Proof of Proposition 3.1. For each fixed $\omega \in \Omega$, we solve the Stokes equation (3.4) with the initial data $u_0(\omega)$ to obtain the unique solution u_1 satisfying the estimate (3.5). If we solve the following stochastic Stokes problem

$$du_2(t,x) = \left(\Delta u_2(t,x) - \nabla p_2(t,x)\right)dt + g(t,x)dB_t \tag{3.6}$$

in $\Omega \times (0, \infty) \times \mathbb{R}^n_+$ with the boundary condition $u_2|_{x_n=0}=0$ and the initial condition $u_2|_{t=0}=0$, then we can take $u=u_1+u_2$. We note that the solution u_2 can be represented by

$$u_2(x,t) = \int_0^t T_K g(x,t,s) dB_s,$$

where

$$T_K g(x,t,s) = \int_{\mathbb{R}^n_+} K(x,y,t-s)g(y,s)dy$$

and the kernel $K = (K_{ij})$ is given by

$$K_{ij}(x, y, t) = \delta_{ij} \Big(\Gamma(x - y, t) - \Gamma(x - y^*, t) \Big)$$

$$+ 4(1 - \delta_{jn}) D_{x_j} \int_0^{x_n} \int_{\mathbb{R}^{n-1}} D_{x_i} N(x - z) \Gamma(z - y^*, t) dz$$
(3.7)

for $i, j = 1, \dots, n$. Here $y^* = (y', -y_n)$. (See [21] for details.) We have

$$\begin{split} \mathbb{E} \int_0^\infty t^{\alpha q} \|u_2\|_{L^p(\mathbb{R}^n_+)}^q dt &= \mathbb{E} \int_0^\infty t^{\alpha q} \left(\int_{\mathbb{R}^n_+} \left| \int_0^t T_K g(x,t,s) dB_s \right|^p dx \right)^{\frac{q}{p}} dt \\ &\lesssim \mathbb{E} \int_0^\infty t^{\alpha q} \left(\int_{\mathbb{R}^n_+} \left| \int_0^t |T_K g(x,t,s)|^2 ds \right|^{\frac{p}{2}} dx \right)^{\frac{q}{p}} dt \\ &\leq \mathbb{E} \int_0^\infty t^{\alpha q} \left(\int_0^t \left| \int_{\mathbb{R}^n_+} |T_K g(x,t,s)|^p dx \right|^{\frac{2}{p}} ds \right)^{\frac{q}{2}} dt \end{split}$$

by the Burkholder–Davis–Gundy inequality (Section 2.7 in [18]) and Minkowski's integral inequality. We now recall the following estimate

$$\int_{\mathbb{R}^{n}_{+}} |T_{K}g(x,t,s)|^{p} dx \lesssim \int_{\mathbb{R}^{n}_{+}} \left| \int_{\mathbb{R}^{n}_{+}} \Gamma_{t-s}(x-y) f(y,s) dy \right|^{p} dx$$

$$+ \int_{\mathbb{R}^{n}_{+}} \left| \int_{\mathbb{R}^{n}_{+}} \Gamma_{t-s}(x-y^{*}) f(y,s) dy \right|^{p} dx$$

in Section 3.1 in [10]. Then we obtain that, by Young's inequality and Lemma 2.14,

$$\begin{split} \mathbb{E} \int_{0}^{\infty} t^{\alpha q} \|u_{2}\|_{L^{p}(\mathbb{R}^{n}_{+})}^{q} dt &\leq \mathbb{E} \int_{0}^{\infty} t^{\alpha q} \left(\int_{0}^{t} \left| \int_{\mathbb{R}^{n}_{+}} \left| \int_{\mathbb{R}^{n}_{+}} \Gamma_{t-s}(x-y)g(y,s) dy \right|^{p} dx \right|^{\frac{2}{p}} ds \right)^{\frac{q}{2}} dt \\ &+ \mathbb{E} \int_{0}^{\infty} t^{\alpha q} \left(\int_{0}^{t} \left| \int_{\mathbb{R}^{n}_{+}} \left| \int_{\mathbb{R}^{n}_{+}} \Gamma_{t-s}(x-y^{*})g(y,s) dy \right|^{p} dx \right|^{\frac{2}{p}} ds \right)^{\frac{q}{2}} dt \\ &\leq \mathbb{E} \int_{0}^{\infty} t^{\alpha q} \left(\int_{0}^{t} (t-s)^{-\frac{n}{p_{2}} + \frac{n}{p}} \|g(s)\|_{L^{p_{2}}(\mathbb{R}^{n}_{+})}^{2} ds \right)^{\frac{q}{2}} dt \\ &\leq \mathbb{E} \int_{0}^{\infty} t^{\alpha_{2}q} \|g\|_{L^{p_{2}}(\mathbb{R}^{n}_{+})}^{q} dt, \end{split}$$

where $\left(\frac{n}{p_2} - \frac{n}{p}\right) + 2(\alpha_2 - \alpha) = 1$. Combining this together with Lemma 3.2, we obtain that

$$||u||_{\mathscr{L}_{a}^{q}(\Omega\times(0,\infty),\mathscr{P};L^{p}(\mathbb{R}^{n}_{+}))} \lesssim ||g||_{\mathscr{L}_{a_{2}}^{q}(\Omega\times(0,\infty),\mathscr{P};L^{p_{2}}(\mathbb{R}^{n}_{+}))}$$

Summing this and (3.5) we get (3.3).

4. Proof of Theorem 1.2

We divide the proof into a few steps.

Step 1) Define the Banach space ${\mathscr X}$ and its closed subset ${\mathscr C}_R$ as

$$\mathscr{X} := \mathscr{L}_{q}^{q}(\Omega \times (0, \infty), \mathscr{P}; L^{p}(\mathbb{R}^{n}_{+}))$$
 and $\mathscr{C}_{R} := \{u \in \mathscr{X}: ||u||_{\mathscr{X}} \leq R\},$

where the number $0 < R \le 1$ will be determined in later. Fix $0 \le \alpha < \frac{1}{q'}$ and an initial data $u_0 \in \dot{\mathbb{B}}_{pq0}^{-2\alpha - \frac{2}{q}}(\mathbb{R}^n_+)$ with div $u_0 = 0$. Then for each $v \in \mathscr{C}_R$ there exists a unique solution $w(\omega)$, for every $\omega \in \Omega$, of the following problem

$$dw = (\Delta w + \nabla P_v - \operatorname{div}(\chi_v v \otimes v))dt + gdB_t \quad \text{and} \quad \operatorname{div} w = 0$$
(4.1)

with the boundary condition $w|_{x_n=0}=0$ and the initial condition $w|_{t=0}=u_0$ by Proposition 3.1. Thus, we can define the solution map by

$$S(v) = w$$
.

We introduced the function $\chi_{\nu(\omega)}:[0,\infty)\to[0,1]$ defined by

$$\chi_{\nu(\omega)}(t) := \theta(\|\nu(\omega)\|_{L^{q}_{\sigma}(0,t;L^{p}(\mathbb{R}^{n}_{+}))}),$$

where $\theta:[0,\infty)\to[0,1]$ is a continuous decreasing function defined by

$$\theta(s) = \{0 \lor (2 - R^{-1}s)\} \land 1.$$

Step 2) Choose $q_1 = \frac{q}{2}$, $p_1 = \frac{p}{2}$, $\alpha_1 = 2\alpha$ so that $\left(\frac{n}{p_1} - \frac{n}{p}\right) + \left(\frac{2}{q_1} - \frac{2}{q}\right) + 2(\alpha_1 - \alpha) = 1$. Hence $-2\alpha - \frac{2}{q} = -1 + \frac{n}{p}$. For every $\omega \in \Omega$, we get by Hölder's inequality

$$\|\chi_{\nu}\nu\otimes\nu\|_{L^{q_1}_{a_1}(0,\infty;L^{p_1}(\mathbb{R}^n_+))}\leq \|\sqrt{\chi_{\nu}}\nu\|_{L^{q}_{a}(0,\infty;L^{p}(\mathbb{R}^n_+))}^2\leq 4R^2.$$

If $v \in \mathcal{C}_R$, then by using Proposition 3.1 and assumption of Theorem 1.2

$$||S(v)||_{\mathcal{X}} = ||w||_{\mathcal{X}} \le C_1(||u_0||_{\dot{\mathbb{B}}^{-2\alpha - \frac{2}{q}}_{pq0}(\mathbb{R}^n_+)} + ||g||_{\mathcal{L}^q_{a_2}(\Omega \times (0,\infty),\mathcal{P};L^{p_2}(\mathbb{R}^n_+))} + 4R^2)$$

$$\le C_1(\delta + 4R^2).$$

Now, we can take $\delta = R^2$ so that

$$||S(v)||_{\mathscr{X}} \le 5C_1 R^2.$$
 (4.2)

Step 3) We claim that

$$|\chi_u - \chi_v| \le R^{-1} ||u - v||_{L^q(0, t \cdot L^p(\mathbb{R}^n))}.$$
 (4.3)

Note first that $\chi_{\nu(\omega)}(t) = 1$ if $\|\nu(\omega)\|_{L^q_\alpha(0,t;L^p(\mathbb{R}^n_+))} \le R$ and that $\chi_{\nu(\omega)}(t) = 0$ if $\|\nu(\omega)\|_{L^q_\alpha(0,t;L^p(\mathbb{R}^n_+))} \ge 2R$. To prove the claim, we may consider the following six cases.

- (a) If $||u||_{L^q_a(0,t;L^p(\mathbb{R}^n_+))} \le ||v||_{L^q_a(0,t;L^p(\mathbb{R}^n_+))} \le R$, then $|\chi_u \chi_v| = |1-1| = 0$.
- (b) If $||u||_{L^q_{\alpha}(0,t;L^p(\mathbb{R}^n_+))} \le R \le ||v||_{L^q_{\alpha}(0,t;L^p(\mathbb{R}^n_+))} \le 2R$, then

$$\begin{split} |\chi_{u} - \chi_{v}| &= |1 - (2 - R^{-1} ||v||_{L_{\alpha}^{q}(0,t;L^{p}(\mathbb{R}^{n}_{+}))})| \\ &= R^{-1} |||v||_{L_{\alpha}^{q}(0,t;L^{p}(\mathbb{R}^{n}_{+}))} - R| \\ &\leq R^{-1} |||v||_{L_{\alpha}^{q}(0,t;L^{p}(\mathbb{R}^{n}_{+}))} - ||u||_{L_{\alpha}^{q}(0,t;L^{p}(\mathbb{R}^{n}_{+}))}| \\ &\leq R^{-1} ||u - v||_{L_{\alpha}^{q}(0,t;L^{p}(\mathbb{R}^{n}_{+}))}. \end{split}$$

(c) If $||u||_{L^q_\alpha(0,t;L^p(\mathbb{R}^n_+))} \le R \le 2R \le ||v||_{L^q_\alpha(0,t;L^p(\mathbb{R}^n_+))}$, then

$$\begin{aligned} |\chi_{u} - \chi_{v}| &= |1 - 0| \le \frac{\|v\|_{L_{\alpha}^{q}(0, t; L^{p}(\mathbb{R}^{n}_{+}))} - \|u\|_{L_{\alpha}^{q}(0, t; L^{p}(\mathbb{R}^{n}_{+}))}}{2R - R} \\ &\le R^{-1} \|u - v\|_{L_{\alpha}^{q}(0, t; L^{p}(\mathbb{R}^{n}_{+}))}. \end{aligned}$$

(d) If $R \le ||u||_{L^q_a(0,t;L^p(\mathbb{R}^n_+))} \le ||v||_{L^q_a(0,t;L^p(\mathbb{R}^n_+))} \le 2R$, then

$$\begin{aligned} |\chi_{u} - \chi_{v}| &= |(2 - R^{-1} ||u||_{L_{\alpha}^{q}(0,t;L^{p}(\mathbb{R}_{+}^{n}))}) - (2 - R^{-1} ||v||_{L_{\alpha}^{q}(0,t;L^{p}(\mathbb{R}_{+}^{n}))})| \\ &= R^{-1} |||v||_{L_{\alpha}^{q}(0,t;L^{p}(\mathbb{R}_{+}^{n}))} - ||u||_{L_{\alpha}^{q}(0,t;L^{p}(\mathbb{R}_{+}^{n}))}| \\ &\leq R^{-1} ||u - v||_{L_{\alpha}^{q}(0,t;L^{p}(\mathbb{R}_{+}^{n}))}. \end{aligned}$$

(e) If $R \leq ||u||_{L^{q}_{\alpha}(0,t;L^{p}(\mathbb{R}^{n}_{+}))} \leq 2R \leq ||v||_{L^{q}_{\alpha}(0,t;L^{p}(\mathbb{R}^{n}_{+}))}$, then

$$\begin{split} |\chi_{u} - \chi_{v}| &= |(2 - R^{-1} ||u||_{L_{\alpha}^{q}(0,t;L^{p}(\mathbb{R}^{n}_{+}))}) - 0| \\ &= R^{-1} |2R - ||u||_{L_{\alpha}^{q}(0,t;L^{p}(\mathbb{R}^{n}_{+}))}| \\ &\leq R^{-1} |||v||_{L_{\alpha}^{q}(0,t;L^{p}(\mathbb{R}^{n}_{+}))} - ||u||_{L_{\alpha}^{q}(0,t;L^{p}(\mathbb{R}^{n}_{+}))}| \\ &\leq R^{-1} ||u - v||_{L_{\alpha}^{q}(0,t;L^{p}(\mathbb{R}^{n}_{+}))}. \end{split}$$

(f) If $2R \le ||u||_{L^q_\alpha(0,t;L^p(\mathbb{R}^n_+))} \le ||v||_{L^q_\alpha(0,t;L^p(\mathbb{R}^n_+))}$, then $|\chi_u - \chi_v| = |0 - 0| = 0$. Thus, (4.3) always holds true.

Step 4) We claim that there exists $C_2 > 0$ such that

$$||S(u) - S(v)||_{\mathscr{X}} \le C_2 R ||u - v||_{\mathscr{X}}. \tag{4.4}$$

Note first that

$$|\chi_{u}u\otimes u-\chi_{v}v\otimes v|\leq |(\chi_{u}-\chi_{v})(u\otimes u)|+|\chi_{v}(u-v)\otimes u|+|\chi_{v}v\otimes (u-v)|.$$

To prove the claim, we may consider the following two cases.

(a) If $\chi_u(t) > 0$ and $\chi_v(t) > 0$, then by (4.3)

$$\begin{aligned} \|(\chi_{u} - \chi_{v})(u \otimes u)\|_{L^{\frac{q}{2}}_{2a}(0,t;L^{\frac{p}{2}}(\mathbb{R}^{n}_{+}))} &\leq R^{-1} \|u - v\|_{L^{q}_{a}(0,t;L^{p}(\mathbb{R}^{n}_{+}))} \|u\|_{L^{q}_{a}(0,t;L^{p}(\mathbb{R}^{n}_{+}))}^{2} \\ &\leq 4R \|u - v\|_{L^{q}_{a}(0,t;L^{p}(\mathbb{R}^{n}_{+}))}. \end{aligned}$$

and

$$\begin{split} &\|\chi_{\nu}(u-\nu)\otimes u\|_{L^{\frac{q}{2}}_{2\alpha}(0,\infty;L^{\frac{p}{2}}(\mathbb{R}^{n}_{+}))} + \|\chi_{\nu}\nu\otimes(u-\nu)\|_{L^{\frac{q}{2}}_{2\alpha}(0,\infty;L^{\frac{p}{2}}(\mathbb{R}^{n}_{+}))} \\ &\leq \|u-\nu\|_{L^{q}_{\alpha}(0,t;L^{p}(\mathbb{R}^{n}_{+}))} \Big(\|u\|_{L^{q}_{\alpha}(0,t;L^{p}(\mathbb{R}^{n}_{+}))} + \|\nu\|_{L^{q}_{\alpha}(0,t;L^{p}(\mathbb{R}^{n}_{+}))} \Big) \\ &\leq 4R\|u-\nu\|_{L^{q}_{\alpha}(0,t;L^{p}(\mathbb{R}^{n}_{+}))}. \end{split}$$

(b) If $\chi_u(t) > 0$ and $\chi_v(t) = 0$, then by the similar way

$$\begin{aligned} \|(\chi_{u} - \chi_{v})(u \otimes u)\|_{L^{\frac{q}{2}}_{2\alpha}(0,t;L^{\frac{p}{2}}(\mathbb{R}^{n}_{+}))} &\leq R^{-1} \|u - v\|_{L^{q}_{\alpha}(0,t;L^{p}(\mathbb{R}^{n}_{+}))} \|u\|_{L^{q}_{\alpha}(0,t;L^{p}(\mathbb{R}^{n}_{+}))}^{2} \\ &\leq 4R \|u - v\|_{L^{q}_{\alpha}(0,t;L^{p}(\mathbb{R}^{n}_{+}))}. \end{aligned}$$

and
$$\|\chi_{\nu}(u-\nu)\otimes u\|_{L^{\frac{q}{2}}_{2a}(0,\infty;L^{\frac{p}{2}}(\mathbb{R}^n_+))} = \|\chi_{\nu}\nu\otimes(u-\nu)\|_{L^{\frac{q}{2}}_{2a}(0,\infty;L^{\frac{p}{2}}(\mathbb{R}^n_+))} = 0.$$

Thus, we use Theorem 3.1 to obtain that

$$||S(u) - S(v)||_{\mathcal{L}^{q}_{a}(0,\infty;L^{p}(\mathbb{R}^{n}_{+}))} \lesssim ||\chi_{u}u \otimes u - \chi_{v}v \otimes v||_{\mathcal{L}^{\frac{q}{2}}_{2a}(0,\infty;L^{\frac{p}{2}}(\mathbb{R}^{n}_{+}))} \lesssim 4R||u - v||_{L^{q}_{a}(0,\infty;L^{p}(\mathbb{R}^{n}))}.$$

This proves the claim (4.4).

Step 5) From (4.2) $S(\mathscr{C}_R) \subset \mathscr{C}_R$ and from (4.4) the map S is contractive in the complete space \mathscr{C}_R whenever $R \leq \frac{1}{5C_1+C_2}$. Therefore we obtain a unique solution u to (4.1) in \mathscr{C}_R by the Banach fixed point theorem. Now, we define the stopping time

$$\tau(\omega) = \inf \left\{ 0 \le T \le \infty : \|u\|_{L^q_{\sigma}(0,T;L^p(\mathbb{R}^n))} \ge R \right\}.$$

If $\tau(\omega) < \infty$, then we set u(t) = 0 for $t \ge \tau(\omega)$ so that (u, τ) is the unique solution in the sense of Definition 1.1.

Notice from definition of τ that for all $h \in (0, \infty)$,

$$\{\omega \mid \tau(\omega) \leq h\} \subset \{\omega \mid ||u||_{L^{q}(0,h\wedge\tau;L^{p}(\mathbb{R}^{n}))} \geq R\}.$$

Using the Chebyshev inequality we obtain that for all $h \in (0, \infty)$,

$$\mathbb{P}(\tau=0) \le \mathbb{P}(\{\tau \le h\}) \le \frac{1}{R} \mathbb{E} \|u\|_{L^q_\alpha(0,h\wedge\tau;L^p(\mathbb{R}^n_+))}$$

Since $\|u\|_{L^q_a(0,h\wedge\tau;L^p(\mathbb{R}^n_+))} \le \|u\|_{L^q_a(0,h;L^p(\mathbb{R}^n_+))} \to 0$ as $h\to 0$ almost surely, we have

$$\mathbb{P}(\tau=0)=0$$
 and $\mathbb{P}(\tau>0)=1$.

By the definition of the stopping time τ we have

$$\mathbb{P}(\tau < \infty) = \mathbb{E}1_{\{\tau < \infty\}} \leq \mathbb{E}\left(1_{\{\tau < \infty\}} \frac{\|u\|_{L^q_a(0,T;L^p(\mathbb{R}^n_+))}}{R}\right) \leq \frac{1}{R} \mathbb{E}\left(\lim_{T \to \infty} \|u\|_{L^q_a(0,T \wedge \tau;L^p(\mathbb{R}^n_+))}\right)$$

By Fatou's Lemma and (4.2) it is bounded by

$$\liminf_{T\to\infty}\frac{1}{R}\mathbb{E}\Big(\|u\|_{L^q_\alpha(0,T\wedge\tau;L^p(\mathbb{R}^n_+))}\Big)\leq \frac{1}{R}\mathbb{E}\Big(\|u\|_{L^q_\alpha(0,\infty;L^p(\mathbb{R}^n_+))}\Big)\leq \frac{5C_1R^2}{R}=5C_1R.$$

If $R < \frac{1}{5C_1 + C_2} \wedge \frac{\epsilon}{5C_1}$, then

$$\mathbb{P}(\tau = \infty) = 1 - \mathbb{P}(\tau < \infty) \ge 1 - \epsilon$$
.

This completes the proof of Theorem 1.2.

5. Proof of Theorem 1.3

We divide the proof into a few steps.

Step 1) We decompose the solution u of (1.1) formally as the sum

$$u = v + V + u_2, (5.1)$$

where

$$v(x,t) = \sum_{i=1}^{n} \langle u_{0i}, K(x-\cdot,t) \rangle,$$

$$V(x,t) = -\int_{0}^{t} \int_{\mathbb{R}^{n}_{+}} K(x,y,t-s) \mathbb{P}(\operatorname{div}(\chi_{R}^{u}(u \times u)(y,s))) dy ds,$$

$$u_{2}(x,t) = \int_{0}^{t} \langle g(\cdot,s), K(x-\cdot,t-s) \rangle dB_{s}.$$

During the proof we shall verify that they are well-defined. Here K is defined in (3.7). We shall give estimates of v, V, and u_2 , separately.

Step 2) For ν we have the following estimate.

Lemma 5.1. For $1 , <math>1 \le q \le \infty$, and $\alpha > 0$,

$$\|\nu\|_{L^{q}(0,\infty;\dot{B}^{-2\alpha}_{pq}(\mathbb{R}^{n}_{+}))} \lesssim \|u_{0}\|_{\dot{B}^{-2\alpha-\frac{2}{q}}_{pq0}(\mathbb{R}^{n}_{+})}.$$

Proof. The zero extension of $u_0 \in \dot{B}^{-2\alpha-2/q}_{pq0}(\mathbb{R}^n_+)$ will be denoted as $\widetilde{u}_0 \in \dot{B}^{-2\alpha-2/q}_{pq}(\mathbb{R}^n)$, which is defined for $f \in \dot{B}^{2\alpha+2/q}_{p'q'}(\mathbb{R}^n)$ by

$$\langle \widetilde{u}_0, f \rangle = \langle u_0, f |_{\mathbb{R}^n} \rangle.$$

Note that $\|u_0\|_{\dot{B}^{-2\alpha-2/q}_{pq0}(\mathbb{R}^n_+)}$ is comparable to $\|\widetilde{u}_0\|_{\dot{B}^{-2\alpha-2/q}_{pq}(\mathbb{R}^n)}$. Following the proof of Theorem 4.1 in [7], we can decompose ν as the sum

$$v(x,t) = v_1(x,t) + v_2(x,t) + v_3(x,t)$$

where v_1 , v_2 and v_3 are defined by

$$\begin{split} v_{1i}(x,t) &= \Gamma_t * \widetilde{u}_{0i}(x) - \Gamma_t^* * \widetilde{u}_{0i}(x), \\ v_{2i}(x,t) &= 4 \frac{\partial}{\partial x_i} \int_{\mathbb{R}^n_+} \sum_{j=1}^{n-1} \frac{\partial}{\partial y_j} (N(x-y) - N(x-y^*)) \Gamma_t^* * u_{0j}(y) dy \\ &- 2 \frac{\partial}{\partial x_i} \int_{\mathbb{R}^n_+} \frac{\partial}{\partial y_n} (N(x-y) - N(x-y^*)) \sum_{j=1}^{n-1} R'_j \Gamma_t^* * u_{0j}(y) dy, \\ v_{3i}(x,t) &= -4 \delta_{in} \sum_{i=1}^{n-1} R'_j \Gamma_t^* * \widetilde{u}_{0j}(x). \end{split}$$

Here R'_j denotes the Riesz transform in \mathbb{R}^{n-1} . Since R'_j is an $L^p(\mathbb{R}^{n-1})$ -multiplier, R'_j is bounded in $L^p(\mathbb{R}^n)$ by Fubini's theorem. Thus, we have

$$\|\nu_{1}(t)\|_{\dot{B}^{-2\alpha}_{na}(\mathbb{R}^{n}_{+})} + \|\nu_{3}(t)\|_{\dot{B}^{-2\alpha}_{na}(\mathbb{R}^{n}_{+})} \lesssim \|\Gamma_{t} * \widetilde{u}_{0}\|_{\dot{B}^{-2\alpha}_{na}(\mathbb{R}^{n})} + \|\Gamma_{t}^{*} * \widetilde{u}_{0}\|_{\dot{B}^{-2\alpha}_{na}(\mathbb{R}^{n})}$$
(5.2)

and from Lemma 2.7

$$\|\nu_2(t)\|_{\dot{B}^{-2\alpha}_{pq}(\mathbb{R}^n_+)} \lesssim \|\Gamma_t^* * \widetilde{u}_0\|_{\dot{B}^{-2\alpha}_{pq}(\mathbb{R}^n)}. \tag{5.3}$$

Using the estimates (5.2), (5.3), and Lemma 2.10, we obtain that

$$\begin{split} \|\nu\|_{L^{q}(0,\infty;\dot{B}_{pq}^{-2\alpha}(\mathbb{R}_{+}^{n}))} &\leq \|\nu_{1}\|_{L^{q}(0,\infty;\dot{B}_{pq}^{-2\alpha}(\mathbb{R}_{+}^{n}))} + \|\nu_{2}\|_{L^{q}(0,\infty;\dot{B}_{pq}^{-2\alpha}(\mathbb{R}_{+}^{n}))} + \|\nu_{3}\|_{L^{q}(0,\infty;\dot{B}_{pq}^{-2\alpha}(\mathbb{R}_{+}^{n}))} \\ &\lesssim \|\Gamma_{t}*\tilde{u}_{0}\|_{L^{q}(0,\infty;\dot{B}_{pq}^{-2\alpha}(\mathbb{R}^{n}))} + \|\Gamma_{t}^{*}*\tilde{u}_{0}\|_{L^{q}(0,\infty;\dot{B}_{pq}^{-2\alpha}(\mathbb{R}^{n}))} \\ &\lesssim \|\tilde{u}_{0}\|_{\dot{B}_{pq}^{-2\alpha-\frac{2}{q}}(\mathbb{R}^{n})} \lesssim \|u_{0}\|_{\dot{B}_{pq}^{-2\alpha-\frac{2}{q}}(\mathbb{R}_{+}^{n})}. \end{split}$$

This completes the proof of Lemma 5.1.

Step 3) For *V* we have the following estimate.

Lemma 5.2. For $1 , <math>1 \le q \le \infty$, and $\alpha > 0$,

$$||V||_{L^{q}(0,\infty;\dot{B}_{pq}^{-2\alpha}(\mathbb{R}^{n}_{+}))} \lesssim ||u||_{L^{q}_{q}(0,\tau(\omega);L^{p}(\mathbb{R}^{n}_{+}))}^{2}.$$

Proof. We can decompose V as the sum

$$V(x,t) = V_1(x,t) + V_2(x,t) + V_3(x,t),$$

where V_1 , V_2 and V_3 are defined by

$$\begin{split} V_{1i}(x,t) &= D_{x'} \big(\mathscr{U} F'(x,t) - \mathscr{U}^* F'(x,t) \big) \\ &:= \int_0^t \int_{\mathbb{R}^n_+} D_{y'} \big(\Gamma(x-y,t-s) - \Gamma(x-y^*,t-s) \big) F'_i(y,s) dy ds, \\ V_{2i}(x,t) &= 4 \frac{\partial}{\partial x_i} \int_{\mathbb{R}^n_+} \sum_{j=1}^{n-1} \frac{\partial}{\partial y_j} \big(N(x-y) - N(x-y^*) \big) D_{y'} \mathscr{U}^* F'_j(y,t) dy \\ &- 2 \frac{\partial}{\partial x_i} \int_{\mathbb{R}^n_+} \frac{\partial}{\partial y_n} \big(N(x-y) - N(x-y^*) \big) \sum_{j=1}^{n-1} R'_j D_{y'} \mathscr{U}^* F'_j(y,t) dy, \\ V_{3i}(x,t) &= -4 \delta_{in} \sum_{j=1}^{n-1} R'_j D_{x'} \mathscr{U}^* F'_j(x,t), \end{split}$$

with $F_{ij} = \chi_R^u u_i u_j$. Notice that

$$||V_{k}(t)||_{\dot{B}_{pq}^{-2\alpha}(\mathbb{R}_{+}^{n})} \leq ||V_{1}(t)||_{\dot{B}_{pq}^{-2\alpha}(\mathbb{R}_{+}^{n})} + ||V_{2}(t)||_{\dot{B}_{pq}^{-2\alpha}(\mathbb{R}_{+}^{n})} + ||V_{3}(t)||_{\dot{B}_{pq}^{-2\alpha}(\mathbb{R}_{+}^{n})} \lesssim ||D_{x'}\mathscr{U}F'(t)||_{\dot{B}_{pq}^{-2\alpha}(\mathbb{R}_{+}^{n})} + ||D_{x'}\mathscr{U}^{*}F'(t)||_{\dot{B}_{pq}^{-2\alpha}(\mathbb{R}_{+}^{n})}.$$

$$(5.4)$$

If $p,q,\alpha,p_1,q_1,\alpha_1$ satisfy $0<\frac{1}{2}-\frac{1}{2}(\frac{n}{p_1'}-\frac{n}{p'})-\alpha<1,\ 1< q_1\leq q<\infty,\ 0\leq \alpha_1<1-\frac{1}{q_1},\ \text{and}$ $\left(\frac{n}{p_1}-\frac{n}{p}\right)+\left(\frac{2}{q_1}-\frac{2}{q}\right)+2(\alpha_1-\alpha)=1,\ \text{then}$

$$||D_{x'}\mathcal{U}F'||_{L^{q}(0,\infty;\dot{B}_{pa}^{-2a}(\mathbb{R}^{n}_{+}))} + ||D_{x'}\mathcal{U}^{*}F'||_{L^{q}(0,\infty;\dot{B}_{pa}^{-2a}(\mathbb{R}^{n}_{+}))} \lesssim ||F'||_{L^{q}_{\alpha_{1}}(0,\infty;L^{p_{1}}(\mathbb{R}^{n}_{+}))}.$$
(5.5)

Since the proof of this technical estimate (5.5) is a little bit long, we postpone its proof to the next step. Combining (5.4), (5.5), and (2.2), we obtain that

$$||V||_{L^{q}(0,\infty;\dot{B}_{pq}^{-2\alpha}(\mathbb{R}^{n}_{+}))} \lesssim ||F||_{L^{q_{1}}_{\alpha_{1}}(0,\infty;L^{p_{1}}(\mathbb{R}^{n}_{+}))}.$$

Finally, we can take $q_1 = \frac{q}{2}$, $p_1 = \frac{p}{2}$, and $\alpha_1 = 2\alpha$ so that the exponents $p, q, \alpha, p_1, q_1, \alpha_1$ satisfy every conditions on the exponents and

$$||F||_{L^{q_1}_{a_1}(0,\infty;L^{p_1}(\mathbb{R}^n_+))} \lesssim ||u||^2_{L^q(0,\tau(\omega);L^p(\mathbb{R}^n_+))}$$

This completes the proof of Lemma 5.2.

Step 4) Now we prove the technical estimate (5.5). Since the estimate of $D_{\chi'}\mathcal{U}^*F'$ can be derived by exactly the same way as that of $D_{\chi'}\mathcal{U}F'$, we focus on estimating the term $D_{\chi'}\mathcal{U}F'$, which is done

by duality argument. If $\psi \in L^{q'}(0,\infty;\dot{B}^{2\alpha}_{p'q'0}(\mathbb{R}^n_+))$, then by Fubini's theorem

$$\begin{split} &\int_0^\infty \int_{\mathbb{R}^n_+} D_{x'} \mathscr{U} F'(x,t) \psi(x,t) dx dt \\ &= \int_0^\infty \int_{\mathbb{R}^n} D_{x'} \mathscr{U} F'(x,t) \widetilde{\psi}(x,t) dx dt \\ &= \int_0^\infty \int_{\mathbb{R}^n} \left(\int_0^t \int_{\mathbb{R}^n_+} F'(y,s) D_{y'} \Gamma(x-y,t-s) dy ds \right) \widetilde{\psi}(x,t) dx dt \\ &= \int_0^\infty \int_0^t \int_{\mathbb{R}^n} F'(y,s) \int_{\mathbb{R}^n} D_{y'} \Gamma(x-y,t-s) \widetilde{\psi}(x,t) dx dy ds dt. \end{split}$$

Integrating by parts and using Hölder's inequality, we have

$$\left| \int_{\mathbb{R}^{n}_{+}} F'(y,s) \int_{\mathbb{R}^{n}} D_{y'} \Gamma(x-y,t-s) \widetilde{\psi}(x,t) dx dy \right|$$

$$= \left| \int_{\mathbb{R}^{n}_{+}} F'(y,s) \int_{\mathbb{R}^{n}} \Gamma(y-x,t-s) D_{x'} \widetilde{\psi}(x,t) dx dy \right|$$

$$= \left| \int_{\mathbb{R}^{n}_{+}} F'(y,s) \Gamma_{t-s} * D' \widetilde{\psi}(y,t) dy \right|$$

$$\leq \left\| F'(s) \right\|_{L^{p_{1}}(\mathbb{R}^{n}_{+})} \left\| \Gamma_{t-s} * D' \widetilde{\psi}(t) \right\|_{L^{p'_{1}}(\mathbb{R}^{n}_{+})}.$$

Using the semigroup property of the heat kernel and Young's convolution inequality, we get

$$\begin{split} \left\| \Gamma_{t-s} * D' \widetilde{\psi}(t) \right\|_{L^{p'_{1}}(\mathbb{R}^{n}_{+})} &\leq \left\| \Gamma_{t-s} * D' \widetilde{\psi}(t) \right\|_{p'_{1}} \\ &= \left\| \Gamma_{(t-s)/2} * \Gamma_{(t-s)/2} * D' \widetilde{\psi}(t) \right\|_{p'_{1}} \\ &\leq \left\| \Gamma_{(t-s)/2} \right\|_{L^{r}(\mathbb{R}^{n})} \left\| \Gamma_{(t-s)/2} * D' \widetilde{\psi}(t) \right\|_{p'} \\ &\lesssim (t-s)^{-\frac{n}{2}(1-\frac{1}{r})} \left\| \Gamma_{(t-s)/2} * D' \widetilde{\psi}(t) \right\|_{p'} \\ &= (t-s)^{\frac{n}{2}(\frac{1}{p'_{1}} - \frac{1}{p'})} \left\| \Gamma_{(t-s)/2} * D' \widetilde{\psi}(t) \right\|_{p'} \end{split}$$

where $1 + 1/p'_1 = 1/r + 1/p'$. Since Lemma 2.11 yields

$$\left\| \Gamma_{(t-s)/2} * D' \widetilde{\psi}(t) \right\|_{p'} \lesssim (t-s)^{-\frac{1}{2}+\alpha} \| D' \widetilde{\psi}(t) \|_{\dot{B}^{-1+2\alpha}_{p'q'}(\mathbb{R}^n)} \lesssim (t-s)^{-\frac{1}{2}+\alpha} \| \widetilde{\psi}(t) \|_{\dot{B}^{2\alpha}_{p'q'}(\mathbb{R}^n)},$$

we obtain that

$$\left\|\Gamma_{t-s}*D'\psi(t)\right\|_{L^{p'_{1}}(\mathbb{R}^{n}_{+})} \lesssim (t-s)^{\frac{n}{2}(\frac{1}{p'_{1}}-\frac{1}{p'})-\frac{1}{2}+\alpha} \|\psi(t)\|_{\dot{B}^{2\alpha}_{p'q'}(\mathbb{R}^{n}_{+})}.$$

Combining the estimate above and then using Hölder's equality, we get

$$\begin{split} & \left| \int_{0}^{\infty} \int_{\mathbb{R}^{n}_{+}} D_{x'} \mathcal{U} F'(x,t) \psi(x,t) dx dt \right| \\ & \leq \int_{0}^{\infty} \int_{0}^{t} \left\| F'(s) \right\|_{L^{p_{1}}(\mathbb{R}^{n}_{+})} (t-s)^{\frac{n}{2}(\frac{1}{p_{1}'} - \frac{1}{p'}) - \frac{1}{2} + \alpha} \|\psi(t)\|_{\dot{B}^{2\alpha}_{p'q'}(\mathbb{R}^{n}_{+})} ds dt \\ & = \int_{0}^{\infty} I_{\lambda} f(t) \|\psi(t)\|_{\dot{B}^{2\alpha}_{p'q'}(\mathbb{R}^{n}_{+})} dt \\ & \leq \|I_{\lambda} f\|_{L^{q}(0,\infty)} \|\psi\|_{L^{q'}(0,\infty;\dot{B}^{2\alpha}_{p'q'}(\mathbb{R}^{n}_{+}))}, \end{split}$$

where

$$\lambda := -\frac{n}{2} (\frac{1}{p'_1} - \frac{1}{p'}) + \frac{1}{2} - \alpha$$
 and $f(s) := ||F'(s)||_{L^{p_1}(\mathbb{R}^n_+)}$.

If $0 < \lambda < 1$, $1 < q_1 \le q < \infty$, $0 \le \alpha_1 < 1/q_1'$, and $1 + 1/q = 1/q_1 + \lambda + \alpha_1$, then by Lemma 2.14

$$||I_{\lambda}f||_{L^{q}(0,\infty)} \lesssim ||f||_{L^{q_1}_{a_1}(0,\infty)} = ||F'||_{L^{q_1}_{a_1}(0,\infty;L^{p_1}(\mathbb{R}^n_+))}.$$

Thus, we have

$$\left| \int_{0}^{\infty} \int_{\mathbb{R}^{n}} D_{x'} \mathscr{U} F'(x,t) \psi(x,t) dx dt \right| \lesssim \|F'\|_{L^{q_{1}}_{\alpha_{1}}(0,\infty;L^{p_{1}}(\mathbb{R}^{n}_{+}))} \|\psi\|_{L^{q'}(0,\infty;\dot{B}^{2\alpha}_{p'q'}(\mathbb{R}^{n}_{+}))}$$

and therefore by duality $\|D_{x'}\mathcal{U}F'\|_{L^q(0,\infty;\dot{B}^{-2\alpha}_{pq}(\mathbb{R}^n_+))} \lesssim \|F'\|_{L^{q_1}_{a_1}(0,\infty;L^{p_1}(\mathbb{R}^n_+))}$. This completes the proof of the estimate (5.5).

Step 5) For u_2 we have the following estimate.

Lemma 5.3. For $1 , <math>1 \le q \le \infty$, and $\alpha > 0$,

$$\mathbb{E} \int_{0}^{\infty} \|u_{2}(t)\|_{\dot{B}^{-2\alpha}_{pq}(\mathbb{R}^{n}_{+})}^{q} dt \lesssim \mathbb{E} \int_{0}^{\infty} \|g(t)\|_{\dot{B}^{-2\alpha-1}_{pq0}(\mathbb{R}^{n}_{+})}^{q} dt.$$

Proof. We can decompose u_2 as the sum

$$u_{2i}(x,t) = u_{21i}(x,t) + u_{22i}(x,t) + u_{23i}(x,t),$$

where u_{21i} , u_{22i} and u_{23i} are defined by

$$\begin{split} u_{21i}(x,t) &= \mathcal{U}\widetilde{g}(x,t) - \mathcal{U}^*\widetilde{g}(x,t) \\ &:= \int_0^t \Gamma_{t-s} * \widetilde{g}_i(\cdot,s)(x) - \Gamma_{t-s}^* * \widetilde{g}_i(\cdot,s)(x) dB_s, \\ u_{22i}(x,t) &= 4 \frac{\partial}{\partial x_i} \int_{\mathbb{R}^n_+} \sum_{j=1}^{n-1} \frac{\partial}{\partial y_j} (N(x-y) - N(x-y^*)) \mathcal{U}^*\widetilde{g}_j(y,s) dy \\ &- 2 \frac{\partial}{\partial x_i} \int_{\mathbb{R}^n_+} \frac{\partial}{\partial y_n} (N(x-y) - N(x-y^*)) \sum_{j=1}^{n-1} R'_j \mathcal{U}^*\widetilde{g}_j(y,s) dy \\ u_{23i}(x,t) &= -4 \delta_{in} \sum_{i=1}^{n-1} R'_j \mathcal{U}^*\widetilde{g}_j(x,t). \end{split}$$

Thus, by Lemma 2.7 we get

$$\sum_{k=1}^3 \|u_k(t)\|_{\dot{B}^{-2\alpha}_{pq}(\mathbb{R}^n_+)} \lesssim \|\mathscr{U}\widetilde{g}(t)\|_{\dot{B}^{-2\alpha}_{pq}(\mathbb{R}^n)} + \|\mathscr{U}^*\widetilde{g}(t)\|_{\dot{B}^{-2\alpha}_{pq}(\mathbb{R}^n)}.$$

If we prove that for $j \in \mathbb{N}$,

$$\mathbb{E} \int_{0}^{\infty} 2^{qj} \|\Delta_{j} \mathscr{U}\widetilde{g}(t)\|_{p}^{q} dt \lesssim \mathbb{E} \int_{0}^{\infty} \|\Delta_{j}\widetilde{g}(t)\|_{p}^{q} dt, \tag{5.6}$$

then we obtain that

$$\begin{split} \mathbb{E} \int_0^\infty \|\mathscr{U}\widetilde{g}(t)\|_{\dot{B}^{-2\alpha}_{pq}(\mathbb{R}^n)}^q dt &= \mathbb{E} \int_0^\infty \sum_{-\infty < j < \infty} 2^{-2\alpha q j} \|\Delta_j \mathscr{U}\widetilde{g}(t)\|_p^q dt \\ &\lesssim \mathbb{E} \int_0^\infty \sum_{-\infty < j < \infty} 2^{-(1+2\alpha)q j} \|\Delta_j \widetilde{g}(t)\|_p^q dt \\ &= \mathbb{E} \int_0^\infty \|\widetilde{g}(t)\|_{\dot{B}^{-2\alpha-1}_{pq0}(\mathbb{R}^n)}^q dt. \end{split}$$

Step 6) Fianlly, we prove the estimate (5.6). Notice that $p > n \ge 2$. The Burkholder-Davis-Gundy inequality (Section 2.7 in [18]) yields

$$\mathbb{E} \int_{0}^{\infty} \|\Delta_{j} \mathcal{U} \widetilde{g}(t)\|_{p}^{q} dt$$

$$= \mathbb{E} \int_{0}^{\infty} \left(\int_{\mathbb{R}^{n}} \left| \int_{0}^{t} \Delta_{j} (\Gamma_{t-s} * \widetilde{g})(x,s) dB_{s} \right|^{p} dx \right)^{\frac{q}{p}} dt$$

$$\lesssim \mathbb{E} \int_{0}^{\infty} \left(\int_{\mathbb{R}^{n}} \left(\int_{0}^{t} |\Delta_{j} (\Gamma_{t-s} * \widetilde{g})(x,s)|^{2} ds \right)^{p/2} dx \right)^{q/p} dt.$$

By Minkowski's integral inequality and Lemma 2.9 we obtain

$$\mathbb{E} \int_{0}^{\infty} \left(\int_{\mathbb{R}^{n}}^{t} \left| \Delta_{j} (\Gamma_{t-s} * \widetilde{g})(x,s) \right|^{2} ds \right)^{p/2} dx \right)^{q/p} dt$$

$$\leq \mathbb{E} \int_{0}^{\infty} \left(\int_{0}^{t} \left(\int_{\mathbb{R}^{n}} |\Delta_{j} (\Gamma_{t-s} * \widetilde{g})(x,s)|^{p} dx \right)^{2/p} ds \right)^{q/2} dt$$

$$= \mathbb{E} \int_{0}^{\infty} \left(\int_{0}^{t} \left\| \Delta_{j} (\Gamma_{t-s} * \widetilde{g})(s) \right\|_{p}^{2} ds \right)^{q/2} dt$$

$$\lesssim \mathbb{E} \int_{0}^{\infty} \left(\int_{0}^{t} \exp(-c(t-s)2^{2j}) \left\| \Delta_{j} \widetilde{g}(s) \right\|_{p}^{2} ds \right)^{q/2} dt$$

for some positive constant c. By Young's convolution inequality we have

$$\mathbb{E} \int_0^\infty \left(\int_0^t \exp(-c(t-s)2^{2j}) \|\Delta_j \widetilde{g}(s)\|_p^2 ds \right)^{q/2} dt$$

$$\leq \mathbb{E} \left(\int_0^\infty \exp(-ct2^{2j}) dt \right)^{q/2} \int_0^\infty \|\Delta_j \widetilde{g}(t)\|_p^q dt$$

$$= (c2^{2j})^{-q/2} \mathbb{E} \int_0^\infty \|\Delta_j \widetilde{g}(t)\|_p^q dt.$$

Thus, we get (5.6).

From (5.1), Lemma 5.1, Lemma 5.2, and Lemma 5.3, we get the desired result. This completes the proof of Theorem 1.3.

ACKNOWLEDGEMENT

T. C. has been supported by the National Research Foundation of Korea No. 2020R1A2C1A01102531. M. Y. has been supported by the National Research Foundation of Korea No. 2016R1C1B2015731 and No. 2015R1A5A1009350.

REFERENCES

- [1] Hajer Bahouri, Jean-Yves Chemin, and Raphaël Danchin. Fourier analysis and nonlinear partial differential equations, volume 343 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer, Heidelberg, 2011.
- [2] A. Bensoussan and R. Temam. Équations stochastiques du type Navier-Stokes. J. Functional Analysis, 13:195-222, 1973.
- [3] Jöran Bergh and Jörgen Löfström. *Interpolation spaces. An introduction*. Springer-Verlag, Berlin-New York, 1976. Grundlehren der Mathematischen Wissenschaften, No. 223.
- [4] Dominic Breit, Eduard Feireisl, and Martina Hofmanová. Stochastically forced compressible fluid flows, volume 3 of De Gruyter Series in Applied and Numerical Mathematics. De Gruyter, Berlin, 2018.
- [5] Marco Cannone. Ondelettes, paraproduits et Navier-Stokes. Diderot Editeur, Paris, 1995. With a preface by Yves Meyer.
- [6] Marek Capiński and Dariusz Gatarek. Stochastic equations in Hilbert space with application to Navier-Stokes equations in any dimension. *J. Funct. Anal.*, 126(1):26–35, 1994.
- [7] Tongkeun Chang and Bum Ja Jin. Initial and boundary values for $L^q_a(L^p)$ solution of the Navier-Stokes equations in the half-space. *J. Math. Anal. Appl.*, 439(1):70–90, 2016.
- [8] Tongkeun Chang and Bum Ja Jin. Global in time solvability of the Navier-Stokes equations in the half-space. *J. Differential Equations*, 267(7):4293–4319, 2019.
- [9] Tongkeun Chang and Bum Ja Jin. Global well-posedness of the half space problem of the navier-stokes equations in critical function spaces of limiting case. Ann. Univ. Ferrara Sez. VII Sci. Mat., 2020.
- [10] Tongkeun Chang and Kyungkeun Kang. Estimates of anisotropic Sobolev spaces with mixed norms for the Stokes system in a half-space. *Ann. Univ. Ferrara Sez. VII Sci. Mat.*, 64(1):47–82, 2018.
- [11] F. Crispo and P. Maremonti. On the (x,t) asymptotic properties of solutions of the Navier-Stokes equations in the half-space. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)*, 318(Kraev. Zadachi Mat. Fiz. i Smezh. Vopr. Teor. Funkts. 36 [35]):147–202, 311, 2004.
- [12] Lihuai Du and Ting Zhang. Local and global strong solutions to the stochastic incompressible Navier-Stokes equations in critical Besov space. *J. Math. Anal. Appl.*, 481(1):123472, 16, 2020.
- [13] Franco Flandoli. An introduction to 3D stochastic fluid dynamics. In SPDE in hydrodynamic: recent progress and prospects, volume 1942 of Lecture Notes in Math., pages 51–150. Springer, Berlin, 2008.
- [14] C. Foiaș. Statistical study of Navier-Stokes equations. I, II. Rend. Sem. Mat. Univ. Padova, 48:219–348 (1973); ibid. 49 (1973), 9–123, 1972.
- [15] Hiroshi Fujita and Tosio Kato. On the Navier-Stokes initial value problem. I. Arch. Rational Mech. Anal., 16:269-315, 1964.
- [16] Jong Uhn Kim. Strong solutions of the stochastic Navier-Stokes equations in R³. Indiana Univ. Math. J., 59(4):1417–1450, 2010.
- [17] Herbert Koch and Daniel Tataru. Well-posedness for the Navier-Stokes equations. Adv. Math., 157(1):22-35, 2001.

- [18] N. V. Krylov. An analytic approach to SPDEs. In *Stochastic partial differential equations: six perspectives*, volume 64 of *Math. Surveys Monogr.*, pages 185–242. Amer. Math. Soc., Providence, RI, 1999.
- [19] Sergei Kuksin and Armen Shirikyan. *Mathematics of two-dimensional turbulence*, volume 194 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2012.
- [20] R. Mikulevicius and B. L. Rozovskii. Global L₂-solutions of stochastic Navier-Stokes equations. Ann. Probab., 33(1):137–176, 2005.
- [21] V. A. Solonnikov. Estimates of the solution of a certain initial-boundary value problem for a linear nonstationary system of Navier-Stokes equations. Zap. Naučn. Sem. Leningrad. Otdel Mat. Inst. Steklov. (LOMI), 59:178–254, 257, 1976. Boundary value problems of mathematical physics and related questions in the theory of functions, 9.
- [22] Takeshi Taniguchi. The existence of energy solutions to 2-dimensional non-Lipschitz stochastic Navier-Stokes equations in unbounded domains. *J. Differential Equations*, 251(12):3329–3362, 2011.
- [23] Hans Triebel. *Interpolation theory, function spaces, differential operators*. Johann Ambrosius Barth, Heidelberg, second edition, 1995.

Yonsei University, Department of Mathematics, Mathematics, Yonsei University, 50 Yonseiro, Seodaemungu, Seoul, Republic of Korea

Email address: chang7357@yonsei.ac.kr, m.yang@yonsei.ac.kr