Finite totally k-closed groups

Dmitry Churikov^{1,2} and Cheryl E Praeger³

¹Sobolev Institule of Mathematics, Novosibirsk, Russia ²Novosibirsk State University, Novosibirsk, Russia ³Centre for the Matheamtics of Symmetry and Computation, University of Western Australia, Perth, WA, Australia

January 23, 2022

Abstract

For a positive integer k, a group G is said to be totally k-closed if in each of its faithful permutation representations, say on a set Ω , G is the largest subgroup of $\operatorname{Sym}(\Omega)$ which leaves invariant each of the Gorbits in the induced action on $\Omega \times \cdots \times \Omega = \Omega^k$. We prove that every abelian group G is totally (n(G) + 1)-closed, but is not totally n(G)closed, where n(G) is the number of invariant factors in the invariant factor decomposition of G. In particular, we prove that for each $k \geq 2$ and each prime p, there are infinitely many finite abelian p-groups which are totally k-closed but not totally (k - 1)-closed. This result in the special case k = 2 is due to Abdollahi and Arezoomand. We pose several open questions about total k-closure.

1 Introduction

In 1969 Wielandt [8, Definition 5.3] introduced, for each positive integer k, the concept of the k-closure of a permutation group G on a set Ω . The kclosure $G^{(k),\Omega}$ of G is the set of all $g \in \text{Sym}(\Omega)$ (permutations of Ω) such that g leaves invariant each G-orbit in the induced G-action on ordered ktuples from Ω . The k-closure $G^{(k),\Omega}$ is a subgroup of $\text{Sym}(\Omega)$ containing G [8, Theorem 5.4], and a permutation group G is said to be k-closed if $G^{(k),\Omega} = G$. Different faithful permutation representations of the same group G may have quite different k-closures. For example, the symmetric group S_3 acts faithfully and intransitively on $\{1, 2, 3, 4, 5\}$ with orbits $\{1, 2, 3\}$ and $\{4, 5\}$, and in this action its 2-closure is $S_3 \times C_2$; while S_3 is 2-closed in its natural action on $\{1, 2, 3\}$.

In 2016, D. F. Holt¹ suggested a stronger concept independent of the permutation representation, and this was studied first by Abdollahi and Arezoomand in [1] in the case k = 2. For a positive integer k, a group G is said to be *totally k-closed* if $G^{(k),\Omega} = G$ whenever G is faithfully represented as a permutation group on Ω . The only totally 1-closed group is the trivial group consisting of a single element (see Remark 2.3), while Abdollahi and Arezoomand [1, Theorem 2] showed that a finite nilpotent group is totally 2-closed if and only if it is cyclic, or it is a direct product of a generalised quaternion group and a cyclic group of odd order. Here we consider larger values of k.

For a permutation group $G \leq \text{Sym}(\Omega)$, and $k \geq 2$, Wielandt [8, Theorem 5.8] proved that

$$G \le G^{(k),\Omega} \le G^{(k-1),\Omega}.$$
(1)

Thus if G is totally (k-1)-closed, then it is automatically totally k-closed. Moreover $G = G^{(k),\Omega}$ for sufficiently large k, since by [8, Theorem 5.12], this holds whenever there exist k-1 points $\alpha_1, \ldots, \alpha_{k-1} \in \Omega$ such that the only element of G fixing each α_i is the identity. The inclusion (1) does suggest that the family of totally k-closed groups might be larger than that of totally (k-1)-closed groups. We show that this is the case, even for abelian groups.

Theorem 1.1. Let k be an integer with $k \ge 2$. Then, for each prime p, there are infinitely many finite abelian p-groups which are totally k-closed but not totally (k-1)-closed.

The result of Abdollahi and Arezoomand shows that the finite totally 2-closed abelian groups are precisely the cyclic groups. It turns out, also for larger values of k, that the total k-closure property for abelian groups is linked with the numbers of cyclic direct factors in their direct decompositions. A study of these decompositions leads to useful bounds, from which we deduce Theorem 1.1.

¹mathoverflow.net/questions/235114/2-closure-of-a-permutation-group

According to the fundamental theorem for finite abelian groups, each nontrivial finite abelian group G can be written as a direct product $G = H_1 \times \cdots \times H_n$, for some $n \ge 1$, such that each $H_i \cong \mathbb{Z}_{d_i}$, $d_1 > 1$, and $d_i | d_{i+1}$ for $1 \le i < n$. The integer n and the d_i are uniquely determined by G, up to the order of the factors. The H_i are called the *invariant factors* of G, and we write n(G) := n for the number of invariant factors. We also have the primary decomposition of G as $G = \prod_{p \in \pi(G)} G_p$, where $\pi(G)$ is the set of primes dividing |G| and G_p is the (unique) Sylow p-subgroup of G. It is straightforward to see that $n(G) = \max_{p \in \pi(G)} n(G_p)$. Our main result is the following theorem, from which we deduce Theorem 1.1.

Theorem 1.2. Let G be a finite abelian group with |G| > 1. Then G is totally (n(G) + 1)-closed, but is not totally n(G)-closed.

The following auxiliary assertion on the k-closure of the direct product abelian permutation p-groups may be of independent interest. It is proved in Section 2, and is used in Section 3 to reduce the proof of Theorem 1.2 to the case of p-groups. For its statement it is convenient to use Syl(G) to denote the set of all Sylow subgroups of a group G; if G is abelian, Syl(G)will consist of one Sylow p-subgroup for each prime $p \in \pi(G)$.

Theorem 1.3. Let G be a finite abelian permutation group on a set Ω , and k an integer, $k \geq 2$. Then $G^{(k),\Omega} = \prod_{P \in Svl(G)} P^{(k),\Omega}$.

The results in our short paper serve to raise a number of open questions, and we record a few here. The first relates to Theorem 1.2. It would be interesting to have a generalisation of the classification by Abdollahi and Arezoomand [1, Theorem 2] of nilpotent totally 2-closed nilpotent groups for larger values of k.

Problem 1. For k > 2 determine all finite nilpotent groups G that are totally k-closed.

As we noted above, the symmetric group S_3 is not totally 2-closed. Indeed, it was shown by Abdollahi, Arezoomand and Tracey [2, Theorem B] a finite soluble group is totally 2-closed if and only if it is nilpotent, hence known by [1, Theorem 2]. However it is not difficult to see that it is totally 3-closed, since in every faithful permutation representation of $G = S_3$ on a set Ω there must be a *G*-orbit of length 3 or 6, and the stabiliser in *G* of two points α, β from such an orbit is trivial. Hence by [8, Theorem 5.12], $G = G^{(3),\Omega}$. As a first step it would be interesting to know which other non-nilpotent soluble groups are totally 3-closed.

Problem 2. Determine the finite soluble groups that are totally 3-closed.

For some time it was believed that all finite totally 2-closed groups would be soluble, and it was somewhat surprising to the authors of [3] to discover that exactly six or the sporadic simple groups are totally 2-closed, namely J_1, J_3, J_4, Ly, Th, M .

Problem 3. Find all the the totally 3-closed sporadic simple groups. More generally, for each sporadic simple group G determine the least value of k such that G is totally k-closed.

The classification of the finite nonabelian simple totally 2-closed groups is still not complete, and we refer the reader to [3] for details of the status of this problem and other open questions about total 2-closure.

Acknowledgements

The authors would like to thank Natalia Maslova for her role in organising the 2020 Ural Workshop on Group Theory and Combinatorics. They are also grateful to an anonymous referee for some hepful comments on the exposition. The first author is supported by Mathematical Center in Akademgorodok under agreement No. 075-15-2019-1613 with the Ministry of Science and Higher Education of the Russian Federation. The second author is supported by the Australian Research Council Discovery Project DP190100450.

2 Preliminaries

In this section we give some background theory, and in particular we prove Theorem 1.3. First we state two results of Wielandt for convenience.

Theorem 2.1. [Wielandt, [8, Theorem 5.6]] Let $G \leq \text{Sym}(\Omega)$, let $k \geq 1$, and let $x \in \text{Sym}(\Omega)$. Then $x \in G^{(k),\Omega}$ if and only if, for all $(\alpha_1, \ldots, \alpha_k) \in \Omega^k$, there exists $g \in G$ such that $\alpha_i^x = \alpha_i^g$ for $i = 1, \ldots, k$.

Theorem 2.2. [Wielandt, [8, Theorem 5.12]] Let $G \leq \text{Sym}(\Omega)$ and $k \geq 1$, and suppose that $\alpha_1, \ldots, \alpha_k \in \Omega$ such that $G_{\alpha_1 \ldots \alpha_k} = 1$. Then $G^{(k+1),\Omega} = G$. Next we discuss total 1-closure.

Remark 2.3. Suppose that G is a finite totally 1-closed group. Consider the regular representation of G on $\Omega = G$. Since G is transitive on Ω it follows from Theorem 2.1 that $G^{(1),\Omega} = \text{Sym}(\Omega)$. Thus, since G is totally 1-closed, it follows that $\text{Sym}(\Omega) = G$ is regular, and hence $|G| \leq 2$. However, if $G = C_2$, then in the representation $G = \langle (12)(34) \rangle \leq \text{Sym}(\Omega)$ on $\Omega = \{1, 2, 3, 4\}$ we have $G^{(1),\Omega} = \langle (12), (34) \rangle \neq G$. Hence G = 1 is the only possibility.

For a prime $p \mid n$, the largest *p*-power divisor of *n* is denoted by n_p ; if π is a set prime divisors of *n*, then $n_{\pi} := \prod_{p \in \pi} n_p$ denotes the π -part of *n*. Recall that, for a finite group G, $\pi(G)$ is the set of prime divisors of |G|. For $p \in \pi(G)$, we denote by $\operatorname{Syl}_p(G)$ the set of Sylow *p*-subgroups of *G*. For a subgroup $G \leq \operatorname{Sym}(\Omega)$ we denote by $\operatorname{Orb}(G)$ the set of *G*-orbits in Ω .

The proof of Theorem 1.3 is developed using ideas from [5]. First we present separately two lemmas as they are general results about finite nilpotent groups.

Lemma 2.4. Let G be a finite nilpotent permutation group, let $p \in \pi(G)$, k be a positive integer, and $P \in \text{Syl}_p(G)$. Let $\Delta_1, \ldots, \Delta_k \in \text{Orb}(P), \Delta = \bigcup_{i=1}^k \Delta_i$, and L be the subgroup of G consisting of all elements fixing each Δ_i setwise. Then $L^{\Delta} = P^{\Delta}$.

Proof. By the definition of L, the subgroup $P \leq L$, and hence $P^{\Delta} \leq L^{\Delta}$. We now prove the converse. Since G is nilpotent, we have $G = P \times H$, where H is the Hall p'-subgroup of G. Let $g \in L$, so g = xy for some (unique) $x \in P$ and $y \in H$. Since $P \leq L$, we have $y = x^{-1}g \in L$.

We claim that $y^{\Delta} = 1$, or equivalently, that $y^{\Delta_i} = 1_{\Delta_i}$ for each $i = 1, \ldots, k$. Since $y \in H \leq C_G(P)$ it follows that, for each i, y^{Δ_i} belongs to the centralizer Z_i of the transitive group $P^{\Delta_i} \leq \text{Sym}(\Delta_i)$, which is semiregular by [7, Theorem 3.2]. In particular $|Z_i|$ divides $|\Delta_i|$ which is a *p*-power, so Z_i is a *p*-group. Consequently, y^{Δ_i} is a *p*-element. Since $y \in H$ and |H| is coprime to *p*, this implies that $y^{\Delta_i} = 1$, for each *i*, and hence that $y^{\Delta} = 1$, proving the claim. Thus, $g^{\Delta} = (xy)^{\Delta} = x^{\Delta}y^{\Delta} = x^{\Delta} \in P^{\Delta}$, as required. \Box

Lemma 2.5. [5, Lemma 2.4] Let $G \leq \text{Sym}(\Omega)$, where $n = |\Omega|$ and $\pi \subseteq \pi(G)$. Suppose that G is transitive and nilpotent, and let H be a Hall π -subgroup of G. Then

- (1) the size of every H-orbit is equal to n_{π} , and
- (2) G acts on Orb(H); moreover, the kernel of this action is equal to H.

Proof of Theorem 1.3

Let G be a finite abelian permutation group on a set Ω , and let $k \geq 2$. Then by [8, Theorem 5.8] and [8, Exercise 5.26], $G^{(k),\Omega}$ is abelian, and $\pi(G^{(k),\Omega}) = \pi(G)$. Let $p \in \pi(G)$, and let P and Q be the (unique) Sylow p-subgroups of G and $G^{(k),\Omega}$ respectively.

Claim 1. $P \leq P^{(k),\Omega} \leq Q$, and $\operatorname{Orb}(P) = \operatorname{Orb}(Q)$.

Proof of Claim 1. By [8, Theorem 5.8] and [8, Exercise 5.28], the group $P^{(k),\Omega}$ is a *p*-group, and hence $P \leq P^{(k),\Omega} \leq Q$. It remains to prove that each *P*-orbit is a *Q*-orbit. Let Δ be a *P*-orbit, and let Γ be the *G*-orbit containing Δ . By (1), $G \leq G^{(k),\Omega} \leq G^{(1),\Omega}$, and hence $G^{(k),\Omega}$ has the same orbits as *G* in Ω . Thus Γ is also a $G^{(k),\Omega}$ -orbit, and hence the *Q*-orbit Δ' containing Δ satisfies $\Delta \subseteq \Delta' \subseteq \Gamma$. The induced permutation groups G^{Γ} and $(G^{(k),\Omega})^{\Gamma}$ are both transitive and abelian, so applying Lemma 2.5 to each of these groups with Hall subgroups P^{Γ}, Q^{Γ} , respectively, yields $|\Delta| = |\Gamma|_p = |\Delta'|$. Thus $\Delta = \Delta'$, and Claim 1 is proved.

Claim 2. $P^{(k),\Omega} = Q$.

Proof of Claim 2. Let $(\alpha_1, \ldots, \alpha_k) \in \Omega^k$, and $g \in Q$. By Theorem 2.1, there exists $h \in G$ such that

$$(\alpha_1,\ldots,\alpha_k)^g = (\alpha_1,\ldots,\alpha_k)^h.$$

For each $i = 1 \dots k$, let Δ_i be the Q-orbit containing α_i . Then by Claim 1, each Δ_i is also a P-orbit. Since $P \leq G$, the group G permutes the P-orbits, and for each i, since $\alpha_i^h = \alpha_i^g \in \Delta_i$, it follows that h fixes each Δ_i setwise. Thus h lies in the subgroup L of Lemma 2.4, and setting $\Delta = \bigcup_{i=1}^k \Delta_i$, it follows from Lemma 2.4 that $h^{\Delta} = u^{\Delta}$ for some $u \in P$. Thus

$$(\alpha_1,\ldots,\alpha_k)^g = (\alpha_1,\ldots,\alpha_k)^h = (\alpha_1,\ldots,\alpha_k)^u.$$

Since such an element $u \in P$ exists for each k-tuple of points and each $g \in Q$, it follows from Theorem 2.1 that $g \in P^{(k),\Omega}$. Thus $Q \leq P^{(k),\Omega}$, and the reverse inclusion holds by Claim 1.

Now we complete the proof of Theorem 1.3. Since $G^{(k),\Omega}$ is abelian, $G^{(k),\Omega}$ is the direct product of its Sylow subgroups. Further, for each $p \in \pi(G)$ it follows from Claim 2 that the unique Sylow *p*-subgroup of $G^{(k),\Omega}$ is $P^{(k),\Omega}$, where *P* is the unique Sylow *p*-subgroup of *G*.

3 Proof of the main results

Recall the definition of n(G) given in Section 1 for a finite abelian group G. We also set $N(G) := \sum_{p \in \pi(G)} n(G_p)$. If $G \leq \text{Sym}(\Omega)$ then the base size $b(G, \Omega)$ of G is the smallest integer b for which there exist $\alpha_1, \ldots, \alpha_b \in \Omega$ such that $G_{\alpha_1 \ldots \alpha_b} = 1$. Such a set $\alpha_1, \ldots, \alpha_b$ is called a base of G. Note that, by Theorem 2.2, $G = G^{(b+1),\Omega}$, where $b = b(G, \Omega)$.

Lemma 3.1. Let G be a finite abelian group and suppose that G has a faithful permutation representation on a finite set Ω . Then $b(G, \Omega) \leq N(G)$, and equality holds for some Ω .

Proof. Let $G = \prod_{p \in \pi(G)} G_p$ with $\pi(G)$ the set of primes dividing |G|, and G_p the Sylow *p*-subgroup of G, for $p \in \pi(G)$. Then, by the definition of N(G) and the $n(G_p)$, G has a direct decomposition $G = H_1 \times \cdots \times H_n$, with each H_i nontrivial and cyclic of prime power order, and n = N(G). For each i, H_i acts regularly on $\Omega_i := H_i$ by (right) multiplication, and G acts faithfully on $\Omega := \bigcup_{i=1}^n \Omega_i$ (where H_j acts trivially on Ω_i for $i \neq j$). Thus the G-orbits in Ω are the sets Ω_i , and for each i the subgroup H_i acts nontrivially only on the orbit Ω_i . Thus each base must contain a point from each of the G-orbits. It follows that the base size equals N(G) for this faithful permutation representation of G.

Now consider an arbitrary faithful permutation representation of G, that is, suppose that $G \leq \text{Sym}(\Omega)$. We prove by induction on N(G) that Ghas base size at most N(G). Now $H_1 = \langle h_1 \rangle \cong \mathbb{Z}_{p^a}$, for some prime p and positive integer a, and as G acts faithfully on Ω there exists $\alpha \in \Omega$ which is not fixed by $h_1^{p^{a-1}}$. This implies that $G_{\alpha} \cap H_1 = 1$. If N(G) = 1 then $G = H_1$ is a cyclic p-group, and $G_{\alpha} = 1$, so $\{\alpha\}$ is a base. Assume now that $N(G) \geq 2$ and that the assertion holds for groups X with N(X) < N(G). Since $G_{\alpha} \cap H_1 = 1$, we have $G_{\alpha} \cong (G_{\alpha}H_1)/H_1 \leq G/H_1 \cong \prod_{i=2}^n H_i$ so $N(G_{\alpha}) \leq n-1 = N(G)-1$, and hence by induction, G_{α} has a base $\alpha_1, \ldots, \alpha_s$ in $\Omega \setminus \{\alpha\}$ with $s \leq N(G) - 1$. Then $\alpha_1, \ldots, \alpha_s, \alpha$ is a base for G in Ω , and the result follows by induction.

We now prove Theorem 1.2 in the case of *p*-groups. The second part of the lemma is proved using a construction developed from ideas in the book of Chen and Ponomarenko [4, Proposition 2.2.26]. An element $\tau \in \text{Sym}(\Omega)$ is called a *cycle* if it is not the identity and has exactly one cycle of length greater than 1 in its disjoint cycle representation; the length of this cycle is denoted $|\tau|$. Two cycles are said to be *independent* if the sets of points they move are disjoint.

Lemma 3.2. Let G be a finite abelian p-group with |G| > 1. Then G is totally (n(G) + 1)-closed, but is not totally n(G)-closed.

Proof. Since G is an abelian p-group, N(G) = n(G). By Lemma 3.1, if G is faithfully represented as a subgroup of $\text{Sym}(\Omega)$, then $b := b(G, \Omega) \leq n(G)$, and by Theorem 2.2, $G = G^{(b+1),\Omega}$. It follows from (1) that $G = G^{(n(G)+1),\Omega}$. Since this holds for all faithful permutation representations of G, G is totally (n(G) + 1)-closed.

As discussed in Section 1, $G \cong \mathbb{Z}_{d_1} \times \mathbb{Z}_{d_2} \times \ldots \times \mathbb{Z}_{d_n}$, with $d_1 > 1$, $d_i | d_{i+1}$ for $1 \leq i < n$, and n = n(G). Let Ω be a set of size $d_1 + \sum_{i=1}^n d_i$, and let $\tau_0, \tau_1, \ldots, \tau_n \in \text{Sym}(\Omega)$ be pairwise independent cycles on Ω such that $|\tau_0| = d_1$, and $|\tau_i| = d_i$ for $i = 1 \ldots n$. Let $H_1 = \langle \tau_0 \tau_1 \rangle$ and $H_i = \langle \tau_0^{-1} \tau_i \rangle$ for $i = 2 \ldots n$, and let

$$H = \langle H_1, \ldots, H_n \rangle.$$

We claim that $H \cong G$. Indeed, the groups H_i commute, and an easy proof by induction on n shows that

$$H_i \cap \langle H_1, \dots, H_{i-1}, H_{i+1}, \dots, H_n \rangle = 1$$
, for $i = 1 \dots n$.

Thus $H = H_1 \times \ldots \times H_n$, with $H_i \cong \mathbb{Z}_{d_i}$ for $i = 1 \ldots n$, proving the claim.

Now we will use Theorem 2.1 to show that $\tau_0 \in H^{(n),\Omega}$. Let $(\alpha_1, \ldots, \alpha_n) \in \Omega^n$, and for $i = 0, \ldots, n$, let Δ_i denote the set of points of Ω moved by τ_i , so that $\{\Delta_0, \ldots, \Delta_n\}$ is the set of *H*-orbits in Ω . Since *H* has n + 1 nontrivial orbits, there exists $k \in \{0, 1, \ldots, n\}$ such that $\Delta_k \cap \{\alpha_1, \ldots, \alpha_n\} = \emptyset$. Define a permutation τ as follows:

$$\tau = \begin{cases} 1, & \text{if } k = 0, \\ \tau_0 \tau_k^{-1}, & \text{if } 1 \le k \le n \end{cases}$$

By definition, $\tau \in H$. If $\tau = 1$, then both τ and τ_0 fix each of the α_i so $(\alpha_1, \ldots, \alpha_n)^{\tau_0} = (\alpha_1, \ldots, \alpha_n)^{\tau}$. On the other hand, if $\tau = \tau_0 \tau_k^{-1}$ for some k, then τ and τ_0 induce the same permutation on $\Omega \setminus \Delta_k$, and again we have $(\alpha_1, \ldots, \alpha_n)^{\tau_0} = (\alpha_1, \ldots, \alpha_n)^{\tau}$. Thus, by Theorem 2.1, $\tau_0 \in H^{(n),\Omega}$. By the construction, $\tau_0 \notin H$, and hence $H \neq H^{(n),\Omega}$. Thus G is not totally n-closed.

Remark 3.3. Theorem 1.1 follows from Lemma 3.2 since, for each integer $k \ge 2$, there are infinitely many finite abelian *p*-groups with k invariant factors.

Finally we prove Theorem 1.2 for an arbitrary finite abelian group G with |G| > 1. Suppose that G is faithfully represented on a set Ω . Since $n(G) = \max_{p \in \pi(G)} n(G_p)$, every Sylow subgroup G_p of G is (n(G) + 1)-closed by Lemma 3.2, and hence, by Theorem 1.3, we have $G^{(n(G)+1),\Omega} = G$. Thus G is totally (n(G) + 1)-closed.

Set n := n(G). If n = 1 then, since |G| > 1, it follows from Remark 2.3 that G is not totally 1-closed. Thus we may assume that $n \ge 2$. Now $n(G) = \max_{p \in \pi(G)} n(G_p)$, and hence we have $n = n(G_q)$ for some $q \in \pi(G)$. By Lemma 3.2, G_q is not totally *n*-closed, so there exists a set Ω_q such that G_q acts faithfully on Ω_q and $G^{(n),\Omega_q} \neq G_q$. There is nothing further to prove if $G = G_q$ so we may assume that $|\pi(G)| \ge 2$. For each $p \in \pi(G) \setminus \{q\}$, let $\Omega_p = G_p$, and consider G_p acting regularly on Ω_p by right multiplication. Thus G acts faithfully on $\Omega := \bigcup_{p \in \pi(G)} \Omega_p$. Since $n \ge 2$, it follows from Theorem 1.3 that

$$G^{(n),\Omega} = \prod_{p \in \pi(G)} (G_p)^{(n),\Omega_p} = (G_q)^{(n),\Omega_q} \times \prod_{\substack{p \in \pi(G) \\ p \neq q}} (G_p)^{(n),\Omega_p}$$

which is not equal to G, because $(G_q)^{(n),\Omega_q} > G_q$ and for every $p \in \pi(G), p \neq q$ the group G_p is *n*-closed as a regular group. Thus, G is not totally *n*-closed, and the proof of Theorem 1.2 is complete.

References

- A. Abdollahi and M. Arezoomand, Finite nilpotent groups that coincide with their 2-closures in all of their faithful permutation representations, J. Algebra Appl., 17(4) 2018, 1850065.
- [2] A. Abdollahi, M. Arezoomand and G. Tracey, On finite totally 2-closed groups, preprint 2020, arXiv:2001:09597
- [3] M. Arezoomand, M. A. Iranmanesh, C. E. Praeger, and G. Tracey, Totally 2-closed finite simple groups, in preparation.

- [4] G. Chen and I. Ponomarenko, *Coherent Configurations*, Central China Normal University Press, Wuhan (2019).
- [5] D. Churikov and I. Ponomarenko, On 2-closed abelian permutation groups, preprint 2020, arXiv:2011.12011
- [6] S. Evdokimov and I. Ponomarenko, *Two-closure of odd permutation group* in polynomial time, Discrete Math., **235**, No. 1-3, 221–232 (2001).
- [7] C. E. Praeger and C. Schneider, *Permutation groups and Cartesian de*compositions. London Mathematical Society Lecture Note Series, 449. Cambridge University Press, Cambridge, 2018.
- [8] H. W. Wielandt, Permutation groups through invariant relations and invariant functions, Lecture Notes, Ohio State University, 1969. Also published in: Wielandt, Helmut, Mathematische Werke (Mathematical works) Vol. 1. Group theory. Walter de Gruyter & Co., Berlin, 1994, pp. 237–296.